

On p -value combination of independent and frequent signals

Yusi Fang

Department of Biostatistics, University of Pittsburgh, Pittsburgh, U.S.A.

and

Chung Chang

Department of Applied Mathematics, National Sun Yat-sen University, Kaohsiung, Taiwan

and

George Tseng

Department of Biostatistics, University of Pittsburgh, Pittsburgh, U.S.A.

SUMMARY: Combining p -values to integrate multiple effects is of long-standing interest in social science and biomedical research. In this paper, we focus on revisiting a classical scenario closely related to meta-analysis, which combines a relatively small (finite and fixed) number of p -values while the sample size for generating each p -value is large (asymptotically goes to infinity). We evaluate a list of traditional and recently developed modified Fisher's methods to investigate their asymptotic efficiencies and finite-sample numerical performance. The result concludes Fisher and adaptively weighted Fisher method to have top performance and complementary advantages across different proportions of true signals. Finally, we propose an ensemble method, namely Fisher ensemble, to combine the two top-performing Fisher-related methods using a robust truncated Cauchy ensemble approach. We show that Fisher ensemble achieves asymptotic Bahadur optimality and integrates the strengths of Fisher and adaptively weighted Fisher methods in simulations. We subsequently extend Fisher ensemble to a variant with emphasized power for concordant effect size directions. A transcriptomic meta-analysis application confirms the theoretical and simulation conclusions, generates intriguing biomarker and pathway findings and demonstrates strengths and strategy of using proposed Fisher ensemble methods.

KEY WORDS: p -value combination; omnibus tests; global hypothesis testing; ensemble method.

1 Introduction

Methods for combining p -values are historically of substantial interest in statistics and in applications of many scientific fields to aggregate homogeneous or possibly heterogeneous information from multiple sources. Consider the problem of combining K p -values, $\vec{p} = (p_1, \dots, p_K)$, where p_i is p -value of testing $H_0^{(i)} : \theta_i \in \Theta_0^{(i)}$ versus $H_A^{(i)} : \theta_i \in \Theta^{(i)} - \Theta_0^{(i)}$. The global union-intersection test for detecting any signal in the K p -values is $H_0 : \bigcap_{1 \leq i \leq K} \{\theta_i \in \Theta_0^{(i)}\}$ versus $H_A : \bigcup_{1 \leq i \leq K} \{\theta_i \in \Theta^{(i)} - \Theta_0^{(i)}\}$. A general strategy is to combine the input p -values and form a new test statistic for globally testing the existence of any signal. In the literature, three major categories of methods have been developed, depending on the types of input data and signal. The first category considers combination of independent p -values, where K is small and fixed (usually 5-30). The sample size n_i ($1 \leq i \leq K$) for deriving p_i is large and can asymptotically goes to infinity. This first classical scenario is closely related to meta-analysis applications to integrate multiple small effects for increased statistical power. Traditional methods include Fisher's method $T_{\text{Fisher}} = \sum_{i=1}^K -2 \log p_i$ (Fisher et al., 1934) and Stouffer's method $T_{\text{Stouffer}} = \sum_{i=1}^K \Phi^{-1}(1 - p_i)$ (Stouffer et al., 1949) as well as many other transformation selections. The second category considers combining independent, sparse, and weak signals, where a large number of p -values are combined ($K \rightarrow \infty$) while only a small number ℓ of the K p -values ($\ell = K^\beta$ with $0 < \beta < \frac{1}{2}$) have weak signals and all remaining p -values have no signal. High criticism (Donoho and Jin, 2004, denoted as HC test hereafter) and Berk-Jones test (Berk and Jones, 1979; Li and Siegmund, 2015, denoted as BJ test hereafter) are two representative methods and are shown to be asymptotically optimal in terms of detection boundary across varying levels of signal sparsity ($0 < \beta < \frac{1}{2}$) as $K \rightarrow \infty$. In the third category, integration of K p -values with unknown correlation structure and with sparse and weak signals is considered. Liu and Xie (2020) and Wilson (2019) have proposed Cauchy test (CA) and harmonic mean test (HM), respectively. These methods

provide robustness under unknown dependency structure when inference is established under independence assumption and they attain optimal detection boundary for detecting highly sparse signals (with $s = K^\beta$, $0 < \beta < \frac{1}{4}$, but not for $\frac{1}{4} < \beta < \frac{1}{2}$) as $K \rightarrow \infty$ (Liu and Xie, 2020; Fang et al., 2021).

In this paper, we revisit methods of the first category, evaluate their asymptotic efficiencies, assess finite-sample numerical performance, and propose an ensemble method combining two complementary top performers for general applications. Despite increasing applications of complex and big data that require methods designed for the second and third categories, methods of the first meta-analytic scenario remain in high demand and present new challenges in many applications. Method development of the first category before the 1970-80s focuses on a class of methods aggregating transformed scores from the p -values: $T = \sum_{i=1}^K g(p_i) = \sum_{i=1}^K F_U^{-1}(p_i)$, where $F_U^{-1}(\cdot)$ is the inverse CDF of U . For example, U is chi-squared for Fisher test and standard normal for Stouffer test. Littell and Folks (1973) showed that Fisher's method is asymptotically optimal in terms of Bahadur relative efficiency, providing theoretical superiority of its log-transformation over the other types of transformations (see Section 2 for more details).

Despite optimal asymptotic efficiency of the Fisher's method, its finite-sample performance in terms of statistical power can be poor if only part of the K p -values have signals. In this commonly encountered situation with heterogeneous signals, modified Fisher's methods have been developed in the past ten years to adaptively aggregate signals only for the top ordered (i.e., the most significant) p -values: $T = -2 \sum_{i=1}^m \log p_{(i)}$, where $p_{(i)}$ is the i -th ordered p -value and m is data-driven, usually determined from an optimization criterion. Li and Tseng (2011) proposed an adaptive Fisher procedure using partial sum optimized by the corresponding p -values (denoted as AFp hereafter): $T_{\text{AFp}} = \max_{1 \leq j \leq K} -\log(h(\vec{p}, j))$, where $h(\vec{p}, j) = \bar{F}_{\chi_{2j}^2}(-2 \sum_{i=1}^j \log p_{(i)})$ is the corresponding p -value of the partial sum, and

$\bar{F}_{\chi_{2j}^2}(t) = 1 - F_{\chi_{2j}^2}(t)$, where $F_{\chi_{2j}^2}(t)$ denotes the CDF of chi-squared random variable with degree of freedom $2j$. Later, Song et al. (2016) developed another adaptive Fisher procedure using partial sum optimized by z -standardization similar to higher criticism (denoted as AFz hereafter): $T_{\text{AFz}} = \max_{1 \leq j \leq K} \frac{-\sum_{i=1}^j \log p_{(i)} - \sum_{i=1}^j w(j,i)}{\sqrt{\sum_{i=1}^j w^2(j,i)}}$ where $w(j,i) = \min\{1, i/j\}$. Another related strategy in the literature is to directly filter out p -values greater than a user-specified threshold $\tau \in (0, 1]$. For example, the truncated Fisher with hard-thresholding (denoted by TFhard) $T_{\text{TFhard}}(\tau) = \sum_{i=1}^K -\log(p_i) I_{\{p_i \leq \tau\}}$ (Zaykin et al., 2002), where $I_{\{\cdot\}}$ denotes the indicator function. The choice of τ is, however, not straightforward. Zhang et al. (2020) investigates the optimal choice of τ for TFhard under a theoretical setting of Gaussian mixture, where mixture probability and mean of the signals are known and $K \rightarrow \infty$. It is, however, not practical since such prior information is generally unknown in applications. To this end, they further propose truncated Fisher with soft-thresholding (TFsoft) to improve TFhard. Zhang et al. (2020) argues that the soft-thresholding scheme can lead to more stable performance with varying input p -values, as the soft-thresholding is continuous while the hard-thresholding is not. By replacing a single user-specified τ with a user-specified set of thresholds \mathcal{T} , two omnibus tests for TFhard and TFsoft are ultimately proposed, which alleviate the issue of choosing τ to some extent. However, the selection of \mathcal{T} is still user-specified.

Notwithstanding the active development of modified Fisher's methods, there is a lack of comprehensive and systematic evaluation of these popular methods' asymptotic properties and their finite-sample numerical performance in the first category. Our paper sets out to fill this gap. In Section 2, we examine asymptotic Bahadur optimality (ABO) of six methods in the first category: Fisher, Stouffer, AFp, AFz, TFhard, and TFsoft. The two adaptive Fisher methods, AFp and AFz, provide estimates of the subset of p -values contributing to the signal. Therefore, we investigate whether the estimates in these two methods consistently select the

subset of p -values representing true signals (signal selection consistency). For completeness, we also examine asymptotic efficiencies for methods developed for sparse signals, including Cauchy, Pareto family, minimum p -value (minP), BJ, and HC. In Section 3, we perform finite-sample numerical evaluations to compare the statistical power of these methods under different K , signal strength, and proportions of true signals. Results of Section 2 and 3 conclude complementary advantages of two top performers – Fisher and AFp –, especially in varying proportions of true signals. Consequently, we develop a Fisher ensemble (FE) method in Section 4 that applies a truncated Cauchy ensemble approach to combine Fisher and AFp. We prove asymptotic Bahadur optimality of the FE method (Section 4.2) and demonstrate its consistently high performance in varying simulation scenarios (Section 4.3). Section 5 develops an extension of the FE method, namely FE_{CS}, for enhanced statistical power on detecting signals with concordant effect size directions. Section 6 applies FE and FE_{CS} as well as existing methods to a transcriptomic meta-analysis on biomarker and pathway detection for aging (Zahn et al., 2007). Section 7 provides final discussion and conclusion.

2 Asymptotic efficiencies of existing methods

This section investigates the asymptotic efficiencies of existing p -value combination methods. Since our focus is for the scenarios with finite-number signals, we slightly generalize the setup proposed in Littell and Folks (1973) (differences between the original setup and ours are discussed in Remark 1), which uses the criterion of exact Bahadur relative efficiency (Bahadur et al., 1960). Under this setting, Fisher’s method is asymptotically Bahadur optimal (Littell and Folks, 1973) and shows theoretical advantages of log-transformation. Multiple modified Fisher’s methods (AFp, AFz, TFhard, and TFsoft) have been developed to improve finite-sample statistical power, but their asymptotic efficiencies have not been investigated. Section 2.1 introduces the problem setting and defines the exact slope of a hypothesis test, which

is a natural concept derived from the exact Bahadur relative efficiency. Section 2.2 presents asymptotic Bahadur optimality (ABO) results of the four modified Fisher's methods.

2.1 Bahadur relative efficiency and exact slope

We first introduce the concept of exact slope of a hypothesis test (Bahadur, 1967; Littell and Folks, 1973). Consider (x_1, x_2, \dots) an infinite sequence of independent observations of a random variable X from probability distribution P_θ with parameter $\theta \in \Theta$. Let T_n be a real-valued and continuous test statistic depending on the first n observations (x_1, \dots, x_n) , where large values of T_n are considered to reject the null hypothesis. Denote $\mathbb{P}_0(T_n < t) = \mathbb{P}_\theta(T_n < t)$ for $\theta \in \Theta_0$. Further denote $p^{(n)} = 1 - F_n(t_n)$ as the p -value of observed $T_n = t_n$, where $F_n(t) = \mathbb{P}_0(T_n < t)$. We then define the exact slope of T_n as follows.

DEFINITION 1: For the test statistic T_n with p -value $p^{(n)}$, if there is a positive valued function $c(\theta)$, such that for any $\theta \in \Theta - \Theta_0$,

$$-\frac{2}{n} \log p^{(n)} \rightarrow c(\theta) \text{ as } n \rightarrow \infty$$

with probability one. Then $c(\theta)$ is called the **exact slope** of T_n .

As a simple example, consider testing for zero mean ($\mu = 0$) with known variance under univariate Gaussian distribution and T_n is the conventional z -test, it is easily seen that $c(\mu) = \mu^2$ is the exact slope of the z -test. Exact slope of a test naturally connects to the exact Bahadur efficiency between test statistics. Consider two sequence of test statistics $\{T_n^{(1)}\}$ and $\{T_n^{(2)}\}$ testing the same null hypothesis with exact slopes $c_1(\theta)$ and $c_2(\theta)$ respectively. We define the ratio $\phi_{12}(\theta) = c_1(\theta)/c_2(\theta)$ as the *exact Bahadur relative efficiency* of $\{T_n^{(1)}\}$ relative to $\{T_n^{(2)}\}$, which compares the relative asymptotic efficiency between two test statistics. Indeed, considering any significance level $\alpha > 0$, for $i = 1, 2$, denote $N^{(i)}(\alpha)$ as the smallest sample size such that, for any $n \geq N^{(i)}(\alpha)$, the p -value of $T_n^{(i)}$ is smaller than α , one can show $\lim_{\alpha \rightarrow 0} N^{(2)}(\alpha)/N^{(1)}(\alpha) = \phi_{12}(\theta)$, which asymptotically characterizes the ratio of the

smallest sample sizes of two test statistics required to attain the same sufficiently small significant level α (Bahadur, 1967; Littell and Folks, 1973).

For $\theta \in \Theta_0$, the p -value $p^{(n)}$ follows uniform distribution $\text{Unif}(0, 1)$. The following lemma shows the analogous “exact slope” $-(2/n) \log p^{(n)}$ converges to zero with probability one.

LEMMA 1: For $\theta \in \Theta_0$, as n diverges, $-(2/n) \log p^{(n)} \rightarrow 0$ with probability one.

In this paper, we extend the definition of exact slope to the null parameter space, where $c(\theta) = 0$ for $\theta \in \Theta_0$.

To benchmark the asymptotic efficiency of a p -value combination method, we then introduce the theoretical setup adopted from the framework in Littell and Folks (1973). Suppose we have $K < \infty$ sequences of test statistics $\{T_{n_1}^{(1)}\}, \dots, \{T_{n_K}^{(K)}\}$ for testing $\theta_i \in \Theta_0^{(i)}$ for $1 \leq i \leq K$. Assume for all the sample sizes n_1, \dots, n_K , and when $\theta_i \in \Theta_0^{(i)}$ for $1 \leq i \leq K$, $\{T_{n_1}^{(1)}\}, \dots, \{T_{n_K}^{(K)}\}$ are independently distributed. Denote $p_i^{(n_i)}$ as the p -value of the i -th test statistic $T_{n_i}^{(i)}$. For each i ($1 \leq i \leq K$), assume that the sequence $\{T_{n_i}^{(i)}\}$ has exact slope $c_i(\theta_i)$ as $-(2/n_i) \log p_i^{(n_i)} \rightarrow c_i(\theta_i) \geq 0$ with probability one as $n_i \rightarrow \infty$. We further assume the sample sizes n_1, \dots, n_p satisfy $n = (1/K) \sum_i^K n_i$ and $\lim_{n \rightarrow \infty} (n_i/n) = \lambda_i$, where $\lambda_i > 0$ and $\sum_i^K \lambda_i = K$. Under the above setup, the goal of any p -value combination method is to test

$$H_0 : \bigcap_{i=1}^K \{\theta_i \in \Theta_0^{(i)}\} \text{ versus } H_1 : \bigcup_{i=1}^K \{\theta_i \in \Theta^{(i)} - \Theta_0^{(i)}\}. \quad (1)$$

For the simplicity of notations, we assume under the null

$$\lambda_1 c_1(\theta_1) \geq \lambda_2 c_2(\theta_2) \geq \dots \geq \lambda_K c_K(\theta_K) \geq 0,$$

where the first ℓ p -values have true signals (i.e., θ_i 's belong to $\Theta^{(i)} - \Theta_0^{(i)}$ for $1 \leq i \leq \ell$) with exact slopes $c_i(\theta_i) > 0$, while the remaining $\theta_i \in \Theta_0^{(i)}$ with exact slopes $c_i(\theta_i) = 0$.

REMARK 1: There are two differences between the original setup in Littell and Folks (1973) and ours. First, Littell and Folks (1973) assume that all the studies have strictly

positive exact slopes, while we allow some studies to have zero-valued exact slopes. Second, Littell and Folks (1973) consider a general parameter space Θ while we consider a product of parameter spaces $\Theta^{(1)} \times \Theta^{(2)} \times \dots \times \Theta^{(K)}$. Although there are differences, one can still establish the results in Littell and Folks (1973) by combining their arguments with Lemma 1.

Following Theorem 2 and arguments in Section 4 in Littell and Folks (1973), under the alternatives, the maximum attainable exact slope for any p -value combination method is $\sum_{i=1}^{\ell} \lambda_i c_i(\theta_i)$. Hence we define the asymptotic Bahadur optimality (ABO) of a p -value combination method as follows.

DEFINITION 2: Denote $\vec{\theta} = (\theta_1, \dots, \theta_K)$. Under the above setup, a p -value combination test $H(p_1, \dots, p_K)$ is asymptotically Bahadur optimal (ABO) if its exact slope $C_H(\vec{\theta})$ satisfies $C_H(\vec{\theta}) = \sum_{i=1}^{\ell} \lambda_i c_i(\theta_i)$.

2.2 Asymptotic Bahadur optimality property of p -value combination methods

Littell and Folks (1973) show that Fisher test is ABO while Stouffer and minP tests are not. Except for these methods, there is a lack of asymptotic efficiency analysis of the other methods. In this subsection, we focus on discussing four modified p -value methods: AFp, AFz, TFhard, and TFsoft. We additionally analyze four methods designed for combining sparse and weak signals: Cauchy, Pareto, BJ, and HC. As expected, they do not enjoy ABO property, and the proofs are outlined in the supplement. The theoretical results of ABO, exact slope, and signal selection consistency (to be discussed in Theorem 4 and remarks S3, S5, and S6) are summarized in Table 1.

[Table 1 about here.]

Recall that Fisher and the four modified Fisher methods combine p -values using the

following test statistics:

$$T_{\text{Fisher}} = \sum_{i=1}^K -2 \log p(i); \quad T_{\text{AFz}} = \max_{1 \leq j \leq K} \frac{-\sum_{i=1}^j \log p(i) - \sum_{i=1}^j w(j, i)}{\sqrt{\sum_{i=1}^j w^2(j, i)}};$$

$$T_{\text{AFp}} = \max_{1 \leq j \leq K} -\log(\bar{F}_{\chi^2_j}(-2 \sum_{i=1}^j \log p(i))); \quad T_{\text{TFhard}}(\tau) = \sum_{i=1}^K (-2 \log p_i) \mathbf{1}_{\{p_i \leq \tau\}};$$

$$T_{\text{TFsoft}}(\tau) = \sum_{i=1}^K (-2 \log p_i + 2 \log \tau)_+.$$

Here $w(j, i) = \min\{1, i/j\}$. In addition, $\tau \in (0, 1]$ is a user-specified constant for the two truncated Fisher methods and $(x)_+$ denotes $\max(x, 0)$.

All the five methods can be characterized in the form of $H(-\log p_1, \dots, -\log p_K)$ by some function $H(\cdot)$. With the log-transform on p -values as a key ingredient, the above methods potentially can achieve high asymptotic efficiency. Indeed, combined with Lemma 1 in Littell and Folks (1973), by using almost the same arguments, one can show that Fisher test attains ABO, where we present the theorem below for completeness.

THEOREM 1 (extended from Littell and Folks (1973); Fisher is ABO): *Under the setup in Section 2.1, Fisher is ABO with exact slope $C_{\text{Fisher}}(\vec{\theta}) = \sum_{i=1}^{\ell} \lambda_i c_i(\theta_i)$.*

Although achieving high asymptotic efficiency, the Fisher test has been shown to have poor performance empirically when only small part of p -values contain signals (e.g., 2 out of 10 p -values have signals); see Song et al. (2016) and Li and Tseng (2011) for more discussions. Many modified Fisher methods have been proposed to address this problem (Zaykin et al., 2002; Yu et al., 2009; Kuo and Zaykin, 2011; Zhang et al., 2020; Li and Tseng, 2011; Song et al., 2016). The idea is to filter out large p -values that are less likely to carry signals and reduce the impact of noise, while still using the log-transformation on p -values to achieve high efficiency. Particularly, AFp and AFz combine the first m smallest ordered p -values. Both methods use some optimization criterion that adaptively selects m to achieve superior finite-sample power in varying proportions of signals. Whether AFp and AFz retain the

ABO property of Fisher is an intriguing question and is investigated below. In fact, we will surprisingly find in the following two theorems that AFp is ABO, but AFz is not.

THEOREM 2 (AFp is ABO): *Under the setup in Section 2.1, T_{AFp} is similar to Fisher test to be ABO with exact slope $C_{AFp}(\vec{\theta}) = \sum_{i=1}^{\ell} \lambda_i c_i(\theta_i)$.*

THEOREM 3 (AFz is not ABO): *Under the setup in Section 2.1, consider the following test statistic $T_A = \max_{1 \leq j \leq K} \frac{-2 \sum_{i=1}^j \log p_{(i)} - A_j}{B_j}$, where $B_j > 0$ and A_j are some finite constants only depend on j and K . Assume there is no tie for $\frac{\sum_{i=1}^j \lambda_i c_i(\theta_i)}{B_j}$, $j=1, \dots, K$, and B_j is monotonic increasing. Then T_A is not ABO with exact slope*

$$C_A(\vec{\theta}) \leq \max_{1 \leq j \leq \ell} (B_1/B_j) \sum_{i=1}^j \lambda_i c_i(\theta_i),$$

where the equality holds if and only if $\operatorname{argmax}_j (1/B_j) \sum_{i=1}^j \lambda_i c_i(\theta_i) = 1$.

By taking $A_j = 2 \sum_{i=1}^j w(j, i)$ and $B_j = \sqrt{4 \sum_{i=1}^j w(j, i)}$, T_A reduces to T_{AFz} , indicating that AFz is not ABO in general (e.g., a special case that AFz is ABO is when $\ell = 1$).

AFp's better asymptotic efficiency property compared to AFz may be due to its attempt to estimate the subset of p -values with true signals. Consider the equivalent form of AFp for combining independent p -values:

$$T'_{AFp} = \min_{\vec{w}} \bar{F}_{\chi^2_{2(\sum_{i=1}^K w_i)}} (-2 \sum_{i=1}^K w_i \log p_i),$$

where $\vec{w} = (w_1, \dots, w_K) \in \{0, 1\}^K$ is the vector of binary weights that identify the candidate subset of p -values with true signals. Note that T'_{AFp} is the original form proposed in Li and Tseng (2011). Denote by $\hat{\vec{w}} = \operatorname{argmin}_{\vec{w}} \bar{F}_{\chi^2_{2(\sum_{i=1}^K w_i)}} (-2 \sum_{i=1}^K w_i \log p_i)$ and let $\vec{w}^* = \{(w_1^*, \dots, w_K^*) : w_k^* = 1 \text{ if } \theta_i \in \Theta - \Theta_0 \text{ and } w_k^* = 0 \text{ if } \theta_i \in \Theta_0\}$ be the indicators of the true signals (i.e., $\sum_{k=1}^K w_k^* = \ell$), we can show signal selection consistency of AFp in the following theorem.

THEOREM 4 (signal selection by AFp is consistent): *Under the setup in Section 2.1, $\hat{w} \rightarrow \vec{w}^*$ as $n \rightarrow \infty$ with probability one for the AFp test.*

The following Theorem 5 states that for any given value of $\tau \in (0, 1]$, TFhard and TFsoft are ABO:

THEOREM 5 (TFhard and TFsoft are ABO): *Under the setup in Section 2.1, TFhard and TFsoft are ABO with exact slopes $C_{TFhard}(\vec{\theta}) = C_{TFsoft}(\vec{\theta}) = \sum_{i=1}^{\ell} \lambda_i c_i(\theta_i)$.*

Although TFhard and TFsoft are ABO, the choice of τ may significantly impact their finite-sample performance (Zhang et al., 2020). To address this issue, Zhang et al. (2020) propose the following omnibus tests for both methods (denoted by oTFhard and oTFsoft, respectively):

$$T_{oTFhard} = \min_{\tau \in \mathcal{T}} 1 - F_{U_{TFhard}(\tau)}(T_{TFhard}(\tau)); \quad T_{oTFsoft} = \min_{\tau \in \mathcal{T}} 1 - F_{U_{TFsoft}(\tau)}(T_{TFsoft}(\tau)),$$

where $\mathcal{T} = \{\tau_1, \dots, \tau_m\}$ is a user-specified set of the candidates of τ . Here $U_{TFhard}(\tau)$ and $U_{TFsoft}(\tau)$ denote the random variables that follow the same null distributions of $T_{TFhard}(\tau)$ and $T_{TFsoft}(\tau)$, respectively. Although the omnibus tests alleviate the issue of sensitivity of the choice of τ for both TFhard and TFsoft to some extent, selection of \mathcal{T} is still user-specified and subjective. In addition, Zhang et al. (2020) derive the null distributions of both omnibus tests in an asymptotic sense as $K \rightarrow \infty$, which may not be accurate for small K with small p -value thresholds that are commonly used in applications, such as genomics studies, to handle multiplicity.

Proofs of theorems for Fisher and modified Fisher methods in this subsection can be found in Supplement Section S2.1. For completeness, we also show that methods designed for combining sparse and weak signals, such as Cauchy, Pareto, BJ and HC, are not ABO (Supplement Section S1) and the proofs are available in Supplement Section S2.3. In conclusion, Fisher, AFp, TFhard and TFsoft are the only four methods with ABO property. AFp and

AFz are two methods to provide signal selection (i.e., subset estimation of the true signal) and AFp is the only method to have consistency in the signal identification.

3 Power comparison in finite-sample simulations

Although Section 2 evaluates asymptotic efficiencies of p -value combination methods, the finite-sample statistical power of the methods under different proportions of signals has not been assessed. In this section, we evaluate six methods that are designed for relatively frequent signal setting described in Section 2: Fisher, Stouffer, AFp, AFz, TFhard, and TFsoft. Additionally, we also evaluate methods designed for combining sparse and weak signals: minimum p -value (minP), Cauchy (CA), harmonic mean (HM), Berk & Jongs (BJ), and higher criticism (HC). As TFhard and TFsoft are sensitive to the choice of tuning parameter τ , for a fair comparison, we use the corresponding omnibus tests, oTFhard and oTFsoft, instead. The tuning candidate set \mathcal{T} is set to be $\{0.01, 0.05, 0.5, 1\}$, which is used in the original paper (Zhang et al., 2020).

For better illustration, we first present results of the six methods designed for combining relatively frequent signals in Figure 1. Results comparing all eleven methods can be found in Supplement Figure S1, where modified Fisher’s methods generally dominate the other methods designed for sparse and weak signals, except when the signals are indeed sparse and weak (e.g., only $\ell = 1$ true signal out of $K = 10$ p -values). However, even for such cases, methods such as AFp and AFz still have comparable power with the top-performing methods such as minP.

We simulate $X = (X_1, \dots, X_K) \stackrel{D}{\sim} N(\vec{\mu}, I_K)$, where $\vec{\mu} = (\mu_1, \mu_2, \dots, \mu_K)$ contains ℓ non-zero signals $\mu_1 = \dots = \mu_\ell = \mu_0$ and $n - \ell$ with no signal ($\mu_{\ell+1} = \dots = \mu_K = 0$). We evaluate for a wide range of K and proportions of true signals ℓ/K : $K = 10, 20, 40, 80$, and $\ell/K = 0.1, 0.2, \dots, 0.7$. We also vary $\mu_0 = 0.5, 0.65, \dots, 5$ for a broad range of signal strength. The p -values are calculated by two-sided test $p_i = 2(1 - \Phi(|X_i|))$ for $i = 1, \dots, K$.

For each combination of parameter values, we draw 10^6 Monte Carlo samples to calculate the critical values for all the methods at significance level $\alpha = 0.01$, given that there is a lack of fast and accurate algorithm for some methods with small K (e.g., oTFsoft and oTFhard).

Figure 1 shows the empirical power of Fisher, Stouffer, and four modified Fisher methods at $\alpha = 0.01$. For a given K and proportion of signals ℓ/K , we choose the smallest μ_0 such that the best method has at least 0.5 statistical power, which allows optimized visualization and comparison of different methods in different signal settings. We first note that AFz is inferior to the other modified Fisher methods, consistent with our theoretical result that AFz is not ABO. We further note that AFp, oTFhard, and oTFsoft have comparable performance across varying proportions of signals, significantly dominating Fisher for detecting infrequent signals (e.g., when the proportion of true signals is smaller than 0.3). On the contrary, Fisher dominates all other methods for detecting frequent signals (e.g., when the proportion of true signals is greater than 0.5). In many real applications (e.g., the transcriptomic meta-analysis in Section 6), the p -value combination test is repeated many times (i.e., for each gene). It is expected that some true biomarkers are more homogeneous with frequent true signals and some with less frequent signals. The result in Figure 1 shows the need to develop an ensemble method to integrate the advantages of Fisher and one of the top-performing modified Fisher methods, which is presented in the next section.

[Figure 1 about here.]

4 Fisher ensemble to combine Fisher and AFp

As shown in Section 2 and 3, Fisher and three modified Fisher methods (AFp, TFhard, and TFsoft) are ABO, and they have complementary strength in the finite-sample evaluation of varying proportions of true signals. A natural idea to improve is to ensemble Fisher and one of the three modified Fisher methods for more stable and universally competitive performance. Since oTFhard and oTFsoft methods require an ad hoc decision of user-specified set \mathcal{T} and

their existing computing algorithms are not accurate for small K , we choose to develop an ensemble method to combine Fisher and AFp in this section. In Section 4.1, we propose an ensemble approach, namely Fisher ensemble (FE), using truncated Cauchy method (Fang et al., 2021) to combine Fisher and AFp. In section 4.2, we provide theoretical support of FE and show that FE is ABO. Section 4.3 presents simulation results similar to Section 3 to demonstrate the balanced and superior performance of FE across varying proportions of true signals.

4.1 Fisher ensemble by truncated Cauchy integration

Denote by p^{Fisher} and p^{AFp} the p -values derived from Fisher and AFp combination tests, respectively. We propose to ensemble the two methods by combining their p -values using $T_h = [h(p^{\text{Fisher}}) + h(p^{\text{AFp}})]/2$ with function h . Since p^{Fisher} and p^{AFp} can be highly dependent, if we use the Cauchy combination test $h(p) = \tan(\pi(\frac{1}{2} - p))$, theorems and simulations in Liu and Xie (2020) and Liu et al. (2019) have shown that the resulting test statistics T_h is robust to the dependence of p^{Fisher} and p^{AFp} , and the method has fast-computing algorithm with Cauchy distribution under the null hypothesis (i.e., null distribution of T_h follows standard Cauchy). This Cauchy ensemble approach is, however, problematic when either p^{Fisher} or p^{AFp} is close to 1. In this case, the Cauchy transformation generates a $-\infty$ score and the power is greatly reduced. We propose to adopt the truncated Cauchy transformation (Fang et al., 2021) in our Fisher ensemble (FE) method by

$$T_{\text{FE}} = [h_{\delta}(p^{\text{Fisher}}) + h_{\delta}(p^{\text{AFp}})]/2, \quad (2)$$

where $h_{\delta}(p) = h(p)\mathbf{I}_{\{p < 1-\delta\}} + h(1-\delta)\mathbf{I}_{\{p \geq 1-\delta\}}$ and δ is set as 0.01 as justified in the original paper. The purpose of using $h_{\delta}(p)$ instead of $h(p)$ is to avoid the large negative penalty issue of the transformation by Cauchy distribution; see Fang et al. (2021) for more details. Except for avoiding large negative penalty issue, ensemble by truncated Cauchy using $h_{\delta}(p)$ performs

almost identically to Cauchy $h(p)$. Supplement Section S3.4 provides numeric examples where the ensemble method using $h_\delta(p)$ performs better than that using $h(p)$.

In the implementation, FE is fully data-driven with fast-computing algorithms. Indeed, for $p_1, \dots, p_K \stackrel{\text{i.i.d.}}{\sim} \text{Unif}(0, 1)$, null distribution of Fisher test follows chi-squared distribution with degree of freedom $2K$. Fast p -value calculation for AFp has been previously developed in Huo et al. (2020) using importance sampling. Using the null reference distribution library for AFp pre-built by Huo et al. (2020), one can calculate the p -value of AFp in linear complexity (Bioconductor package ‘‘AWFisher’’). Finally, Proposition 3 in Fang et al. (2021) has shown that the truncated Cauchy approach proposed here has at most $(1 + \delta)^2 - 1 = (1 + 0.01)^2 - 1 = 2\%$ inflation of the type I error control if we naively use Cauchy distribution as the null distribution (see Proposition S2 in supplement). In other words, if the calculated p -value is 0.05 from null distribution using Cauchy distribution, the true p -value from null distribution using truncated Cauchy is between 0.05 and $0.05 \cdot (1 + 0.01)^2 = 0.051$. As a result, fast p -value computation for Fisher ensemble $T_{h;\delta}$ is warranted. Table S1 in Section S3.1 justifies the above fast-computing procedure, where we show the type I error control for FE with $\delta = 0.01$ is accurate for $\alpha \leq 0.05$ across a broad range of $5 \leq K \leq 100$.

4.2 Asymptotic efficiency of Fisher ensemble

In this subsection, we will show that Fisher ensemble (FE) is asymptotically Bahadur optimal (ABO). We first introduce a heavy-tailed distribution family, namely regularly-varying distribution R , where Cauchy and truncated Cauchy distributions are special cases of the family. Consider an ensemble method induced by a regularly-varying distribution (e.g., truncated Cauchy in our case) to combine multiple p -value combination methods (e.g., Fisher and AFp in our case). The ensemble method will be shown to be ABO if at least one of the p -value combination methods is ABO. Since both Fisher and AFp are ABO and truncated Cauchy is a regularly varying distribution, we conclude that Fisher ensemble is

also ABO. Below, we outline the definition of regularly-varying distribution and the theorem. The detailed proof is available in Supplement Section S2.2.

DEFINITION 3: A distribution F is said to belong to the regularly-varying tailed family with index γ (denoted by $F \in R_{-\gamma}$) if $\lim_{x \rightarrow \infty} \frac{\bar{F}(xy)}{\bar{F}(x)} = y^{-\gamma}$ for some $\gamma > 0$ and all $y > 0$.

We denote the whole family of regularly varying tailed distributions by R . For two positive functions $u(\cdot)$ and $v(\cdot)$, we write $u(t) \sim v(t)$ if $\lim_{t \rightarrow \infty} \frac{u(t)}{v(t)} = 1$. It can be shown that every distribution F belonging to $R_{-\gamma}$ can be characterized by $\bar{F}(t) \sim L(t)t^{-\gamma}$, where $\bar{F}(t) = 1 - F(t)$ and $L(t)$ is a slowly varying function. A function L is called slowly varying if $\lim_{y \rightarrow \infty} \frac{L(ty)}{L(y)} = 1$ for any $t > 0$. Regularly varying distribution is a wide class of heavy-tailed distributions, which includes Cauchy, harmonic mean, Pareto and truncated Cauchy.

Consider $L < \infty$ p -value combination test statistics T_1, \dots, T_L , each combining $\vec{p} = (p_1, \dots, p_K)$. Denote by p_{T_1}, \dots, p_{T_L} the resulting p -values of T_1, \dots, T_L . In Fisher ensemble, we have $L = 2$ and (T_1, T_2) are Fisher and AFp. Under Definition 3, consider the following ensemble method by a regularly varying tailed distribution:

$$T_{RV}(\gamma) = \sum_{i=1}^L g_{\gamma}(p_{T_i}) = \sum_{i=1}^L F_{U(\gamma)}^{-1}(1 - p_{T_i}),$$

where $F_{U(\gamma)}$ is CDF of $U(\gamma)$ and $U(\gamma) \in R_{-\gamma}$. Under the null hypothesis, the test statistic transforms all the p_{T_i} 's into regularly varying tailed random variables with index γ . The following theorem suggests that there is no loss of asymptotic efficiency in terms of Bahadur relative efficiency to ensemble T_1, \dots, T_L by regularly varying tailed distribution.

THEOREM 6: For each $i = 1, \dots, L$, let $C_i(\vec{\theta})$ be the exact slope of T_i and assume $\max_{1 \leq i \leq L} C_i(\vec{\theta}) > 0$. For $T_{RV}(\gamma)$ with $U \in R_{-\gamma}$ and $\bar{F}_{U(\gamma)}(t) \sim L(t)/t^{\gamma}$, assume $L(t)/t^{\gamma}$ is continuous at $[t_0, +\infty)$ for some $t_0 \geq 0$. Denote by $C_{RV}^{(\gamma)}(\vec{\theta})$ the exact slope of $T_{RV}(\gamma)$, then we have $C_{RV}^{(\gamma)}(\vec{\theta}) \geq \max_{1 \leq i \leq L} C_i(\vec{\theta})$.

Theorem 6 suggests that the exact slope of $T_{RV}(\gamma)$ is always greater than or equal to the

maximum of exact slopes of aggregated p -value combination methods. In other words, $T_{RV}(\gamma)$ is ABO as long as at least one of T_1, \dots, T_L methods is ABO. Consequently, Fisher ensemble is ABO since truncated Cauchy belongs to regularly-varying tailed distribution and both Fisher and AFp are ABO.

4.3 Finite-sample power comparison of Fisher ensemble

In this subsection, we evaluate the finite-sample power of FE. To illustrate that FE can take advantages of integrated methods, we also include AFp and Fisher as the baseline methods. We use the same simulation scheme in Section 3 to generate the simulated data. Figure 2 shows statistical power of FE, AFp, and Fisher at $\alpha = 0.01$. Similar to Figure 1, For a given proportion of signals ℓ/K and number of combined p -values K , we choose the smallest μ_0 that allows the best method to have power larger than 0.5. As expected, we note that FE has a stable statistical power that is comparable to the better of Fisher and AFp in different proportions of true signals. Specifically, when the proportion of signals is high, FE performs close to Fisher and is superior to AFp. When the proportion is low, FE performs close to AFp and outperforms Fisher. In Supplement Figure S2, we implement another Fisher ensemble method combining Fisher, AFp, and minP. As expected, its power for only one signal (i.e., $\ell = 1$) is improved but at the expense of a large reduction of power when signals are frequent. From the asymptotic efficiency in Section 4.2 and simulations above, we recommend using the Fisher ensemble method combining Fisher and AFp for general applications.

[Figure 2 about here.]

5 Detection of signals with concordant direction

5.1 Fisher ensemble focused on concordant signal (FE_{CS})

For all methods we have discussed so far, the global hypothesis setting is designed for two-sided tests, regardless of directions of the effects. Recall from Equation 1 that the hypothesis

testing considered is $H_0 : \bigcap_{i=1}^K \{\theta_i = 0\}$ vs $H_1 : \bigcup_{i=1}^K \{\theta_i \neq 0\}$. Consider the alternative hypothesis that only the first ℓ p -values have true signals (i.e., $\theta_i \neq 0$ for $1 \leq i \leq \ell$ and $\theta_{\ell+1} = \dots = \theta_K = 0$). The two-sided tests to obtain p_i ($1 \leq i \leq K$) cannot guarantee signals with concordant directions ($\text{sgn}(\theta_1) = \dots = \text{sgn}(\theta_\ell)$, denoted by $\text{sgn}(\cdot)$ the sign function), which is desirable in some applications. For example, when conducting meta-analysis of K transcriptomic studies believed to be relatively homogeneous, we are interested in identifying biomarkers concordantly up-regulated or down-regulated. For this problem, Owen (2009) revisits the Pearson test statistic and proposes to use

$$T_{\text{Pearson}} = \min\{\tilde{p}^{\text{Fisher},L}, \tilde{p}^{\text{Fisher},R}\},$$

where $\tilde{p}^{\text{Fisher},L}$ and $\tilde{p}^{\text{Fisher},R}$ uses Fisher to combine the left and right one-sided p -values respectively, and the Pearson test takes the more significant one as the test statistics. In this subsection, we similarly extend the Fisher ensemble method to use truncated Cauchy approach to combine the two left and right one-sided p -values of Fisher and AFp (denoted as FE_{CS}; Fisher ensemble for concordant signal):

$$T_{\text{FECS}} = \frac{h_\delta(\tilde{p}^{\text{Fisher},L}) + h_\delta(\tilde{p}^{\text{Fisher},R}) + h_\delta(\tilde{p}^{\text{AFp},L}) + h_\delta(\tilde{p}^{\text{AFp},R})}{4}.$$

REMARK 2: When combining one-sided p -values, it is very common to observe p -values close to 1 and it is even more critical to use truncated Cauchy, instead of Cauchy, to avoid $-\infty$ scores.

REMARK 3: Let $C^L(\vec{\theta})$ be the maximum attainable exact slope for any p -value combination method combining left one-sided p -values, and define $C^R(\vec{\theta})$ in a similar manner for right one-sided p -values. By Theorem 6, the exact slope of FE_{CS} is $\max\{C^L(\vec{\theta}), C^R(\vec{\theta})\}$, indicating high asymptotic efficiency for utilizing information within one-sided p -values.

For computation, similar to FE, one can use p -value calculation for standard Cauchy to calculate p -values for FE_{CS} and there is at most $(1 + 0.01)^4 - 1 = 4\%$ inflation on type I error control using the standard Cauchy when $\delta = 0.01$. This approximation procedure is justified by simulation results in Table S1 in Section S3.1 for a broad range of significance levels α and numbers of input p -values K .

5.2 Finite-sample power comparison of Fisher ensemble for concordant signal

In this subsection, we evaluate the finite-sample power of FE_{CS} with $\delta = 0.01$. To demonstrate the advantages of FE_{CS}, we also include the regular FE and Pearson as the baseline methods. We use the same simulation scheme in Section 3 to generate the simulated data. For FE_{CS} and Pearson, the one-sided p -values are generated by $\tilde{p}_i^{(L)} = 1 - \Phi(X_i)$ and $\tilde{p}_i^{(R)} = \Phi(X_i)$ ($i = 1, \dots, K$), respectively. While for the regular FE, we combine the two-sided p -values $p_i = 2(1 - \Phi(|X_i|))$ for $i = 1, \dots, K$.

Figure 3 shows the empirical power of FE_{CS}, Pearson, and the regular FE at $\alpha = 0.001$. For a given frequency of signals ℓ/K and a number of combined p -values K , we choose the smallest μ_0 that allows the best method to have power larger than 0.6. Both FE_{CS} and Pearson dominate the regular FE, indicating the latter two methods perform better for the alternatives with one-sided direction consistent effects (as $\mu_1 = \dots = \mu_s = \mu_0 > 0$ under the alternatives). We further note that FE_{CS} outperforms Pearson when the frequency of signals is low ($\ell/K \leq 0.3$), while FE_{CS} still has a comparative power with Pearson for detecting frequent signals ($\ell/K \geq 0.3$).

[Figure 3 about here.]

6 Real Application to AGEMAP data

In this section, we apply different p -value combination methods to analyze the AGEMAP study (Zahn et al., 2007). The dataset contains microarray expression of 8,932 genes in

sixteen tissues as well as age and sex variables of 618 mice subjects. We are interested in identifying age-related genes. Following the original paper, we fit the regression model below to detect age-associated genes in each tissue:

$$Y_{ijk} = \beta_{0jk} + \beta_{\text{age},jk} \text{Age}_{ijk} + \beta_{\text{sex},jk} \text{Sex}_{ijk} + \varepsilon_{ijk} \text{ for } i = 1, \dots, m_{jk},$$

where Y_{ijk} is the expression level of the i -th subject for the j -th gene and k -th tissue. We consider designs of both two-sided and one-sided tests when combining p -values across tissues. In two-sided test design, two-sided p -values (p_{j1}, \dots, p_{jK}) for their corresponding $\beta_{\text{age},jk}$ coefficients are combined using Fisher, AFp and FE methods. In this case, the association directions (positive or negative association) are not considered. By contrast, one-sided test design combines left-tailed p -values ($\tilde{p}_{j1}^L, \dots, \tilde{p}_{jK}^L$) or right-tailed p -values ($\tilde{p}_{j1}^R, \dots, \tilde{p}_{jK}^R$) respectively using FE_{CS}. Compared to FE, FE_{CS} is expected to have increased power to detect age-related biomarkers with concordant signals (all positive association or all negative association) across tissues while have reduced power for markers with heterogeneous signals (i.e., positive association in some tissues and negative association in some others). In this application, both concordant and heterogeneous age-related biomarkers are of interest. FE and FE_{CS} will serve as complementary tools for different biological objectives.

Figure 4(a) shows Fisher, AFp, and FE p -value combination results in the two-sided test design. Under q -value ≤ 0.05 , Fisher detects 576 genes (yellow color) and AFp detects 452 genes (green color), where Category II (379 genes) represents overlapped detected genes by Fisher and AFp and Category I (197 genes) and III (73 genes) represent biomarkers uniquely detected by Fisher or by AFp. The heatmap shows age-association measure defined as: $E_{jk} = -\text{sign}(\beta_{\text{age},jk}) \log(\min\{\tilde{p}_{jk}^L, \tilde{p}_{jk}^R\})$ for gene j on the rows and tissue k on the columns; i.e., the signed log-transformed (base 10) two-sided p -values. Consequently, the red color of E_{jk} represents a strong positive association while blue means a strong negative association. As expected, FE combines the strengths of Fisher and AFp to detect 598 genes (purple color)

that contain all genes in Category II and most genes in Category I and III. By counting the number of tissues with p -values $p_{jk} \leq 0.05$, Supplement Figure S4 shows that category I genes (detected by Fisher but not by AFp) are age-associated in more tissues, while Category III (detected by AFp but not by Fisher) are age-associated in fewer tissues, which is consistent to the theoretical insight and simulation result that Fisher is more powerful for detecting frequent signals and AFp is more powerful for less frequent signals.

We next perform hierarchical clustering (using 1-correlation between tissues as dissimilarity measure and complete linkage) for the sixteen tissues based on the E_{jk} values in the 598 age-related genes detected by FE, and the dendrogram is shown in Figure 4(a). By cutting the dendrogram, five clear tissue modules of similar age-association patterns are identified: (1) thymus and gonads; (2) spleen and lung; (3) eye, kidney, and heart (4) hippocampus, adrenal glands, and muscle; (5) cerebrum and spinal cord (also see Figure 4(b) for heatmap of pairwise correlations). For the first module, the thymus has long been regarded as an endocrine organ that is closely related to Gonads and sexual physiology, such as sexual maturity and reproduction. (Grossman, 1985; Leposavić and Pilipović, 2018). The spleen-lung module is consistent with the finding in Zahn et al. (2007), and many reports suggest that spleen and lung share a similar aging pattern (e.g., Schumacher et al. (2008)). For the third module, literature shows that kidney and eye share structural, developmental, physiological, and pathogenic similarities and pathways. The relationships between age-related eye, kidney, and cardiovascular diseases have been widely reported (e.g., Farrah et al. (2020)) For the fourth module, extensive literature reports the relationship between adrenal glands and hippocampal aging (e.g., Landfield et al. (1978)). For the last module, few existing studies have investigated the aging process of the spinal cord (Knight and Nigam, 2017). But it is reasonable that the cerebrum and spinal cord might share a similar aging pattern as they both belong to the central nervous system. On the other hand, the liver has intriguingly

negative correlations of aging effects with muscle, adrenal glands, and several brain regions, such as the hippocampus, cerebellum, and cerebrum (also see Figure 4(b)).

Next, we evaluate FE_{CS} for one-sided test design and compare it with FE. We calculate $S_{\text{sign},j} = \sum_{k=1}^{16} \text{sign}(\beta_{\text{age},jk}) I_{\{\min\{\tilde{p}_{jk}^L, \tilde{p}_{jk}^R\} \leq 0.05\}}$ to determine whether the detected concordant aging marker j is positively associated ($S_{\text{sign},j} > 0$) or negatively associated ($S_{\text{sign},j} \leq 0$). Similar to the previous analysis, Figure 5 shows age-associated genes detected by FE (598 genes, Category II(A), II(B) and III) and FE_{CS} (407 genes, Category I(A), I(B), II(A) and II(B)), where Category II(A) and II(B) are overlapped genes detected by FE and FE_{CS} , Category III are only detected by FE and Category I(A) and I(B) are only detected by FE_{CS} . For genes detected by FE_{CS} , Category I(A) and II(A) are concordant aging markers with positive association (mostly red) and Category I(B) and II(B) are negatively associated (mostly blue), which are visually consistent with the heatmap. In contrast, genes in Category III mostly have discordant association directions (partial red and partial blue). Supplement Figure S5 shows the numbers of positive or negative associated tissues, which confirms the above observation.

At significance level $q \leq 0.05$, FE_{CS} identifies 189 positively associated genes (Category I(A) and II(A)) and 218 negatively associated genes (Category I(B) and II(B)). We perform Ingenuity Pathway Analysis (IPA) to these two concordant age-associated gene lists. The result identifies eleven enriched pathways from the 189 positively associated genes and five enriched pathways from the 218 negatively associated genes (significance level $p \leq 0.01$). Table S2 shows details of these enriched pathways with pathway names, enrichment p -values, and supporting literature of the pathways related to aging/early development processes (see complete references in Supplement References II). The result shows the advantage of FE_{CS} to identify age-associated markers concordant across tissues and to deliver interpretable biological insights.

[Figure 4 about here.]

[Figure 5 about here.]

7 Conclusion and discussion

P-value combination is a common and effective information synthetic tool in many scientific applications. In this paper, we focus on a meta-analytic scenario, where the number of combined p -values K is relatively small and fixed in the asymptotic setting. Our contribution is three-fold. Firstly, this is the first study to comprehensively evaluate p -value combination methods for their asymptotic efficiencies in terms of asymptotic Bahadur optimality (ABO). We investigate classical methods (Fisher and Stouffer) and modified Fisher’s methods (AFp, AFz, TFhard, and TFsoft). The result shows that Fisher, AFp, TFhard, and TFsoft are ABO, but Stouffer and AFz are not. We also find an interesting consistency property for the estimation of signal contributing subset in AFp (Theorem 4). Secondly, we perform an extensive finite-sample power comparison and conclude that Fisher and AFp are two top performers with complementary advantages, where Fisher is more powerful with frequent signals and AFp is more powerful in a relatively sparse setting. Thirdly, we propose a Fisher ensemble (FE) method to combine Fisher and AFp. A one-sided test modification, FE_{CS}, is further developed for detecting concordant signal. The advantages of FE and FE_{CS} includes: (A) Both methods have high asymptotic efficiencies (FE is ABO). (B) The truncated Cauchy combination avoids the $-\infty$ score in the traditional Cauchy. (C) We numerically demonstrate their constantly high performance across varying proportions of signals. (D) Both methods have fast-computing procedures. Finally, an application to integrate 16 tissues of AGEMAP transcriptomic data verifies theoretical conclusions, demonstrates superior performance of FE and FE_{CS}, and discovers intriguing biological findings in age-association biomarkers and pathways.

Modern data science faces challenges from data heterogeneity, increased complex structure,

and the need for effective methods for new scientific hypotheses. The ensemble methods proposed in this paper, FE and FE_{CS}, have solid theoretical and numerical support for their superior performance in a wide range of signal settings. We believe the method will find impactful applications in many other scientific problems.

Acknowledgements

YF and GCT are funded by NIH R21LM012752; CC is funded by Ministry of Science and Technology of ROC 109-2118-M-110-002 and 110-2118-M-110-001. The authors thank Yongseok Park for some helpful discussions.

References

- Bahadur, R. R. (1967). Rates of convergence of estimates and test statistics. *The Annals of Mathematical Statistics* **38**, 303–324.
- Bahadur, R. R. et al. (1960). Stochastic comparison of tests. *The Annals of Mathematical Statistics* **31**, 276–295.
- Berk, R. H. and Jones, D. H. (1979). Goodness-of-fit test statistics that dominate the Kolmogorov statistics. *Probability Theory and Related Fields* **47**, 47–59.
- Donoho, D. and Jin, J. (2004). Higher criticism for detecting sparse heterogeneous mixtures. *The Annals of Statistics* **32**, 962–994.
- Fang, Y., Tseng, G. C., and Chang, C. (2021). Heavy-tailed distribution for combining dependent p -values with asymptotic robustness. *arXiv preprint arXiv:2103.12967*.
- Farrah, T. E., Dhillon, B., Keane, P. A., Webb, D. J., and Dhaun, N. (2020). The eye, the kidney, and cardiovascular disease: old concepts, better tools, and new horizons. *Kidney International* **98**, 323–342.
- Fisher, R. et al. (1934). *Statistical methods for research workers*. Oliver and Boyd, Edinburgh and London.

- Grossman, C. J. (1985). Interactions between the gonadal steroids and the immune system. *Science* **227**, 257–261.
- Huo, Z., Tang, S., Park, Y., and Tseng, G. (2020). P-value evaluation, variability index and biomarker categorization for adaptively weighted Fisher’s meta-analysis method in omics applications. *Bioinformatics* **36**, 524–532.
- Knight, J. and Nigam, Y. (2017). Anatomy and physiology of ageing 5: the nervous system. *Nursing Times* **113**, 55–58.
- Kuo, C.-L. and Zaykin, D. V. (2011). Novel rank-based approaches for discovery and replication in genome-wide association studies. *Genetics* **189**, 329–340.
- Landfield, P. W., Waymire, J., and Lynch, G. (1978). Hippocampal aging and adrenocorticoids: quantitative correlations. *Science* **202**, 1098–1102.
- Leposavić, G. M. and Pilipović, I. M. (2018). Intrinsic and extrinsic thymic adrenergic networks: sex steroid-dependent plasticity. *Frontiers in Endocrinology* **9**, 13.
- Li, J. and Siegmund, D. (2015). Higher criticism: p -values and criticism. *The Annals of Statistics* **43**, 1323–1350.
- Li, J. and Tseng, G. C. (2011). An adaptively weighted statistic for detecting differential gene expression when combining multiple transcriptomic studies. *The Annals of Applied Statistics* **5**, 994–1019.
- Littell, R. C. and Folks, J. L. (1973). Asymptotic optimality of Fisher’s method of combining independent tests II. *Journal of the American Statistical Association* **68**, 193–194.
- Liu, Y., Chen, S., Li, Z., Morrison, A. C., Boerwinkle, E., and Lin, X. (2019). Acat: a fast and powerful p -value combination method for rare-variant analysis in sequencing studies. *The American Journal of Human Genetics* **104**, 410–421.
- Liu, Y. and Xie, J. (2020). Cauchy combination test: a powerful test with analytic p -value calculation under arbitrary dependency structures. *Journal of the American Statistical*

Association **115**, 393–402.

- Owen, A. B. (2009). Karl Pearson’s meta-analysis revisited. *The Annals of Statistics* **37**, 3867–3892.
- Schumacher, B., Van Der Pluijm, I., Moorhouse, M. J., Kosteas, T., Robinson, A. R., Suh, Y., Breit, T. M., Van Steeg, H., Niedernhofer, L. J., Van Ijcken, W., et al. (2008). Delayed and accelerated aging share common longevity assurance mechanisms. *PLoS Genetics* **4**, e1000161.
- Song, C., Min, X., and Zhang, H. (2016). The screening and ranking algorithm for change-points detection in multiple samples. *The Annals of Applied Statistics* **10**, 2102.
- Stouffer, S. A., Suchman, E. A., DeVinney, L. C., Star, S. A., and Williams Jr, R. M. (1949). The American soldier: Adjustment during army life. *Studies in Social Psychology in World War II* **5**,
- Wilson, D. J. (2019). The harmonic mean p-value for combining dependent tests. *Proceedings of the National Academy of Sciences* **116**, 1195–1200.
- Yu, K., Li, Q., Bergen, A. W., Pfeiffer, R. M., Rosenberg, P. S., Caporaso, N., Kraft, P., and Chatterjee, N. (2009). Pathway analysis by adaptive combination of p-values. *Genetic Epidemiology: The Official Publication of the International Genetic Epidemiology Society* **33**, 700–709.
- Zahn, J. M., Poosala, S., Owen, A. B., Ingram, D. K., Lustig, A., Carter, A., Weeraratna, A. T., Taub, D. D., Gorospe, M., Mazan-Mamczarz, K., et al. (2007). AGEMAP: a gene expression database for aging in mice. *PLoS Genetics* **3**, e201.
- Zaykin, D. V., Zhivotovsky, L. A., Westfall, P. H., and Weir, B. S. (2002). Truncated product method for combining p-values. *Genetic Epidemiology: The Official Publication of the International Genetic Epidemiology Society* **22**, 170–185.
- Zhang, H., Tong, T., Landers, J., and Wu, Z. (2020). TFisher: A powerful truncation and

weighting procedure for combining p -values. *The Annals of Applied Statistics* **14**, 178–201.




Figure 1: Statistical power of Fisher, Stouffer, and four modified Fisher’s methods at significance level $\alpha = 0.01$ across varying frequencies of signals $\ell/K = 0.1, 0.2, \dots, 0.7$ and varying numbers of combined p -values $K = 10, 20, 40, 80$. The standard errors are negligible compared to the scale of the mean power (smaller than 0.1% of the power) and hence omitted. The results of Stouffer with power smaller than 0.25 when $\ell/K = 0.1$ or 0.2 are omitted.




Figure 2: Statistical power of FE, Fisher, and AFp at significance level $\alpha = 0.01$ across varying frequencies of signals $\ell/K = 0.1, 0.2, \dots, 0.7$ and varying numbers of combined p -values $K = 10, 20, 40, 80$. The standard errors are negligible and hence omitted.




Figure 3: Statistical power of FE, FE_{CS} , and Pearson at significance level $\alpha = 0.001$ across varying frequencies of signals $\ell/K = 0.1, 0.2, \dots, 0.7$ and varying numbers of combined p -values $K = 10, 20, 40, 80$. The standard errors are negligible and hence omitted.




Figure 4: (a) Heatmaps of age-association measure E_{jk} of significant genes ($q \leq 0.05$) detected in the two-sided test design. Category I: genes detected by Fisher but not AFp; II: genes detected by both Fisher and AFp; III: genes detected by AFp but not Fisher. (b) Heatmap of pair-wise correlations between tissues based on the detected genes by FE ($q \leq 0.05$.) in (a).




Figure 5: Heatmaps of age-association measure E_{jk} of genes detected by FE_{CS} or by FE ($q \leq 0.05$). Heatmap I(A) represents up-regulated genes detected only by FE_{CS} (43 genes); heatmap I(B) represents down-regulated genes detected only by FE_{CS} (58 genes); heatmap II(A) represents up-regulated genes detected both by FE_{CS} and FE (146 genes); heatmap II(B) represents down-regulated genes detected both by FE_{CS} and FE (160 genes); heatmap III represents genes detected only by FE (292 genes), respectively.

Table 1: Results of asymptotic properties of eleven p -value combination methods: Fisher, Stouffer, four modified Fisher (AFp, AFz, TFhard and TFsoft) and five methods designed for sparse and weak signal (Cauchy, Pareto, minP, BJ and HC).

Methods	ABO	Exact slopes	Signal selection consistency	Proofs
Fisher	Yes	$\sum_{i=1}^{\ell} \lambda_i c_i(\theta_i)$	–	Littell and Folks (1973)
Stouffer	No	$\frac{1}{K} \left[\sum_{i=1}^{\ell} (\lambda_i c_i(\theta_i))^{\frac{1}{2}} \right]^2$	–	Littell and Folks (1973)
AFp	Yes	$\sum_{i=1}^{\ell} \lambda_i c_i(\theta_i)$	Yes	Theorems 2 & 4
AFz	No	$\leq \max_j \frac{\sum_{i=1}^j \lambda_i c_i(\theta_i)}{\sqrt{\sum_{i=1}^j \min^2\{1, i/j\}}}$	No	Theorem 3
TFhard	Yes	$\sum_{i=1}^{\ell} \lambda_i c_i(\theta_i)$	–	Theorem 5
TFsoft	Yes	$\sum_{i=1}^{\ell} \lambda_i c_i(\theta_i)$	–	Theorem 5
Cauchy	No	$\max_i \lambda_i c_i(\theta_i)$	–	Theorem S1
Pareto	No	$\max_i \lambda_i c_i(\theta_i)$	–	Theorem S2
minP	No	$\max_i \lambda_i c_i(\theta_i)$	–	Littell and Folks (1973)
BJ	No	$\max_i i \lambda_i c_i(\theta_i)$	No	Theorem S3
HC	No	–	No	Proposition S1