Variations and extensions of the Gaussian concentration inequality, Part II

Daniel J. Fresen*

Abstract

We prove concentration inequalities for f(X) about its median, where X is a random vector in \mathbb{R}^n with independent heavy tailed coordinates of Weibull or power type, and $f : \mathbb{R}^n \to \mathbb{R}$ is a locally Lipschitz function. This paper is part of a series of four papers, Part I, Part II and two supporting papers. It can be read independently of Part I.

Contents

Rest	ults	1
1.2	Power tails	4
Pro	ofs	6
2.1	Outline of the general method	6
2.2	Small lemmas	6
2.3	Tools from $[5]$, $[7]$ and $[13]$	7
2.4	Proof of Theorem 1	9
2.5	Proof of Theorem 2	11
	1.1 1.2 Prod 2.1 2.2 2.3 2.4	Results 1.1 Weibull tails 1.2 Power tails 1.2 Power tails Proofs 2.1 Outline of the general method 2.2 Small lemmas 2.3 Tools from [5], [7] and [13] 2.4 Proof of Theorem 1 2.5 Proof of Theorem 2

1 Results

Let Lip(T, x) denote the local Lipschitz constant of a function around a point x,

$$Lip(T, x) = \lim_{\varepsilon \to 0^+} Lip(T|_{B(x,\varepsilon)})$$
(1)

Associated to a real valued random variable X_1 are its distribution, CDF and quantile function defined as

$$\mu_{1}(E) = \mathbb{P}\{X_{1} \in E\}, E \in \mathcal{B}(\mathbb{R})
F_{1}(t) = \mathbb{P}\{X_{1} \leq t\}, t \in \mathbb{R}
F_{1}^{-1}(s) = \inf\{t \in \mathbb{R} : F_{1}(t) \geq s\}, s \in (0, 1)$$
(2)

^{*}University of Pretoria, Department of Mathematics and Applied Mathematics, daniel.fresen@up.ac.za

where $\mathcal{B}(\mathbb{R})$ is the Borel sigma algebra. If X_1 has a continuous density $h = d\mu_1/dx$ then for all points $s \in (0, 1)$ and $t \in \mathbb{R}$,

$$Lip(F_1^{-1}, s) = \frac{1}{h(F_1^{-1}(s))} \qquad Lip(F_1^{-1} \circ \Phi, t) = \frac{\phi(t)}{h(F_1^{-1}(\Phi(t)))}$$

where we interpret $a/0 = \infty$ for a > 0,

$$\phi(t) = \frac{1}{\sqrt{2\pi}} e^{-t^2/2} \qquad and \qquad \Phi(t) = \int_{-\infty}^t \phi(u) du$$

For a differentiable function $f : \mathbb{R}^n \to \mathbb{R}$ and $1 \le s \le \infty$ set

$$Lip_{s}(f) = \sup_{x \in \mathbb{R}^{n}} \left| \nabla f(x) \right|_{s} \qquad Lip_{s}^{\sharp}(f) = \left(\sum_{i=1}^{n} \sup_{x \in \mathbb{R}^{n}} \left| \frac{\partial f}{\partial x_{i}}(x) \right|^{s} \right)^{1/s} \tag{3}$$

with the usual interpretation (involving max) when $s = \infty$. In general, $Lip_s(f) \leq Lip_s^{\sharp}(f)$. There are two special cases when $Lip_s(f) = Lip_s^{\sharp}(f)$. The first is when f is linear. The second is when $s = \infty$. By considering a path integral, and a local linear approximation, it follows that $Lip_s(f)$ is the Lipschitz constant of f with respect to the $\ell_{s^*}^n$ norm on \mathbb{R}^n , where $s^* = s/(s-1)$ when $1 < s \le \infty$ (setting $\infty/(\infty - 1) := 1$).

In both Theorems 1 and 2 below, the assumption of differentiability is a convenience rather than a necessity.

1.1 Weibull tails

Theorem 1 (Weibull type tails ~ exp $(-|t|^q)$, 0 < q < 1) Let $n \in \mathbb{N}$, 0 < q < 1, and let $(\mu_i)_1^n$ be a sequence of probability measures on \mathbb{R} , each with corresponding cumulative distribution F_i and quantile function F_i^{-1} as in (2) such that for all $s \in \mathbb{R}$,

$$Lip(F_i^{-1} \circ \Phi, s) \le (1 + |s|)^{-1 + 2/q}$$
 (4)

Let $f : \mathbb{R}^n \to \mathbb{R}$ be differentiable and let $(X_i)_1^n$ be a sequence of independent random variables, each with corresponding distribution μ_i . Then for all t > 0,

$$\mathbb{P}\left\{\left|f\left(X\right) - \mathbb{M}f\left(X\right)\right| > t\right\} \le 2\exp\left(-c_q \min\left\{\left(\frac{t}{Lip_2^{\sharp}\left(f\right)}\right)^2, \left(\frac{t}{Lip_{\infty}^{\sharp}\left(f\right)}\right)^q\right\}\right)$$
(5)

and

$$\mathbb{P}\left\{\left|f\left(X\right) - \mathbb{M}f\left(X\right)\right| > C_{q}\left(\log\left(e + \frac{n}{t^{-2+4/q}}\right)\right)^{1/q-1/2}\left(tLip_{2}\left(f\right) + t^{2/q}Lip_{\infty}\left(f\right)\right)\right\} \le 2\exp\left(-t^{2}/2\right) \tag{6}$$

Comments for Theorem 1

• Comment 1: When f is linear, say $f(x) = \sum_{i=1}^{n} a_i x_i$, for non-zero $a \in \mathbb{R}^n$, and we assume without loss of generality that $\mathbb{E}X_i = 0$, then (5) can be written as

$$\mathbb{P}\left\{\left|\sum_{i=1}^{n} aX_{i}\right| > t\right\} \le 2\exp\left(-c_{q}\min\left\{\left(\frac{t}{|a|}\right)^{2}, \left(\frac{t}{|a|_{\infty}}\right)^{q}\right\}\right)$$
(7)

This is already known and holds under the slightly weaker tail assumption

$$\mathbb{P}\left\{|X_i| \ge t\right\} \le 2\exp\left(-t^q\right), t > 0$$

There are several ways to show this. For example, one could use moment estimates of Hitczenko, Montgomery-Smith and Oleszkiewicz [9, Theorem 1.1] as modified in [4] (see also Latała's paper [10]), or one could use a modified version of the Gaussian concentration inequality (either in the style of this paper, or using a method of restricting and extending functions). We refer the reader to [4] for a more detailed discussion of this.

• Comment 2: A related nonlinear estimate was proved by Barthe, Cattiaux, and Roberto [1, Example 5.4]. If X has i.i.d. coordinates, each with distribution μ_i and corresponding density $d\mu_i/dx = (2\Gamma(1+1/q))^{-1} \exp(-|x|^q)$, then using a weak Poincaré inequality they proved that

$$\mathbb{P}\left\{\left|f\left(X\right) - \mathbb{M}f\left(X\right)\right| > t\right\} \le 2\exp\left(-c_q \min\left\{\frac{t}{Lip_2\left(f\right)}\left(\log n\right)^{1-1/q}, \left(\frac{t}{Lip_2\left(f\right)}\right)^q\right\}\right)$$

Or equivalently,

$$\mathbb{P}\left\{ |f(X) - \mathbb{M}f(X)| > C_q\left(t^2 \left(\log n\right)^{-1+1/q} + t^{2/q}\right) Lip_2(f) \right\} \le 2\exp\left(-t^2/2\right)$$
(8)

For t below a certain value (8) is superior to (6), however for large values of t (6) improves on (8) by replacing $Lip_2(f)$ with the potentially smaller quantity $Lip_{\infty}(f)$, which is the correct behaviour in the linear case.

• Comment 3: The quantity

$$\left(\log\left(e + \frac{n}{t^{-2+4/q}}\right)\right)^{1/q-1/2}$$

cannot be completely erased from (6), otherwise this could be written as

$$\mathbb{P}\left\{\left|f\left(X\right) - \mathbb{M}f\left(X\right)\right| > t\right\} \le 2\exp\left(-c_{q}\min\left\{\left(\frac{t}{Lip_{2}\left(f\right)}\right)^{2}, \left(\frac{t}{Lip_{\infty}\left(f\right)}\right)^{q}\right\}\right)$$

which would imply that $\operatorname{Var}(|X|_{\infty}) < C_q$, when in fact $\operatorname{Var}(|X|_{\infty}) \approx (\log n)^{-2+2/q}$. The current bound gives the estimate $\operatorname{Var}(|X|_{\infty}) \leq C_q (\log n)^{-1+2/q}$ which is off by a factor of $\log n$, exactly the same factor by which the classical Gaussian concentration inequality is off by in the case when q = 2 (specifically for the ℓ_{∞}^n norm). For certain functions it may be possible to decrease the exponent of the logarithmic term, say from 1/q –

1/2 to 1/q - 1, using Talagrand's L_1 - L_2 inequality (the Gaussian version as in [3]), or by applying the methodology presented here to $\eta \circ f$ for an appropriate choice of η to achieve a superconcentrated estimate, which is a trick we have exploited in connection with Gaussian concentration of the ℓ_p^n norm, see [6]. Or one could simply combine (6) with (8).

• Comment 4: In case $|\nabla f(X)|_s$ is with high probability much smaller than $Lip_s(f)$, including the case when $Lip_s(f) = \infty$ and/or when one has a high probability bound on $\partial_i f(X)$, $1 \le i \le n$, one can prove variations of Theorem 1 (using a similar proof) that take the distribution of these into account.

• Comment 5: For $q \in [1, 2]$ related estimates can be proved using results of Talagrand [14, Theorem 2.4] and Gozlan [8, Proposition 1.2], which give bounds of the form

$$\mu^{n} \left(A + tB_{2}^{n} + t^{2/q}B_{q}^{n} \right) \ge 1 - \exp\left(-Dt^{2}\right) : t \ge 0$$

assuming that μ satisfies a type of Poincaré inequality on \mathbb{R} with constant C, μ^n is the *n*-fold product of μ , and D is a constant depending only on C. This includes the case $d\mu/dx = (2\Gamma(1+1/q))^{-1} \exp(-|x|^q)$. Taking $A = \{x : f(x) \leq \mathbb{M}f(x)\}$, if $x \in$ $A + tB_2^n + t^{2/q}B_q^n$ then there exists $a \in A$, $u \in B_2^n$, $v \in B_q^n$ such that $x = a + tu + t^{2/q}v$, so $|f(x) - f(a + tu)| \leq t^{2/q}Lip_{q^*}(f)$ and $|f(a + tu) - f(a)| \leq tLip_2(f)$, which implies $f(x) \leq \mathbb{M}f(x) + tLip_2(f) + t^{2/q}Lip_{q^*}(f)$. Together with a similar lower bound,

$$\mathbb{P}\left\{\left|f\left(X\right) - \mathbb{M}f\left(X\right)\right| > t\right\} \le 2\exp\left(-c_{q}\min\left\{\left(\frac{t}{Lip_{2}\left(f\right)}\right)^{2}, \left(\frac{t}{Lip_{q^{*}}\left(f\right)}\right)^{q}\right\}\right)$$

1.2 Power tails

Theorem 2 There exists a universal constant C > 0 such that the following is true. Let $n \in \mathbb{N}, 2 < q < \infty, 2q (q-2)^{-1} < p < \infty$, and let $(\mu_i)_1^n$ be a sequence of probability measures on \mathbb{R} , each with corresponding cumulative distribution F_i and quantile function F_i^{-1} as in (2) such that for all $s \in (0, 1)$,

$$Lip(F_i^{-1}, s) \le \min\{s, 1-s\}^{-1-1/q}$$
 (9)

Let $f : \mathbb{R}^n \to \mathbb{R}$ be differentiable and let $(X_i)_1^n$ be a sequence of independent random variables, each with corresponding distribution μ_i . Then for all t > 0,

$$\mathbb{P}\left\{\left|f\left(X\right) - \mathbb{M}f\left(X\right)\right| > C_{q}t\left(Lip_{2}^{\sharp}\left(f\right) + e^{t^{2}/(2q)}Lip_{q}^{\sharp}\left(f\right)\right)\right\} \le Ce^{-t^{2}/2}$$
(10)

where $C_q > 0$ is a function of q, and

$$\mathbb{P}\left\{\left|f\left(X\right) - \mathbb{M}f\left(X\right)\right| > C_{p,q}tLip_{p}\left(f\right)\left(n^{1/2-1/p} + n^{1/q}e^{t^{2}/(2q)}\right)\right\} \le Ce^{-t^{2}/2}$$
(11)

where $C_{p,q} > 0$ is a function of (p,q).

Comments for Theorem 2

• Comment 1: The following result is due to Barthe, Cattiaux and Roberto [1, Example 5.3] (see Theorem 5.1 in their paper for a more general result).

Theorem 3 Let $\alpha > 0$, let X be a random vector in \mathbb{R}^n with probability density (with respect to Lebesgue measure) $d\mu/dx = \alpha^n 2^{-n} \prod_{i=1}^n (1+|x_i|)^{-1-\alpha}$, and let $f : \mathbb{R}^n \to \mathbb{R}$ satisfy $|f(x) - f(y)| \leq |x - y|$ for all $x, y \in \mathbb{R}^n$. Then there exists $t_0(\alpha) > e$ and $C(\alpha) > 0$ such that for all $t \geq t_0(\alpha)$,

$$\mathbb{P}\left\{\left|f\left(X\right) - \mathbb{M}f\left(X\right)\right| > tn^{1/\alpha}\right\} \le C\left(\alpha\right) \left(\frac{\log t}{t}\right)^{c}$$

Their bound can be written as

$$\mathbb{P}\left\{\left|f\left(X\right) - \mathbb{M}f\left(X\right)\right| > C_{q}t^{2}e^{t^{2}/(2q)}n^{1/q}\right\} \le C_{q}e^{-t^{2}/2}$$
(12)

We have taken the liberty of including the C_q in the deviation. For t large enough so that

$$te^{t^2/(2q)} > C_q n^{\frac{1}{2} - \frac{1}{p} - \frac{1}{q}} Lip_p(f)$$

(11) improves on (12). Note that by the assumptions on p and q, $\frac{1}{2} - \frac{1}{p} - \frac{1}{q} > 0$ and $Lip_p(f) \leq Lip_2(f)$, which in their result is at most 1. It is not unusual for $Lip_p(f)$ to be much smaller than 1.

• Comment 2: Whenever f is linear, say $f(x) = \sum_{1}^{n} a_i X_i$ for $a \neq 0$, (10) improves on (12) for all t. In this case we may replace the basic assumption (9) with $\mathbb{P}\{|X_i| > t\} \leq c_q (1+t)^{-q}$. To do this, take an i.i.d. sequence $(Y_i)_1^n$ independent of $(X_i)_1^n$ with $\mathbb{P}\{|Y_i| > t\} = (1+t)^{-q}$. One can show, as we did in the 3^{rd} arXiv version of [5], that both $(Y_i)_1^n$ and $(X_i+Y_i)_1^n$ satisfy (9). Applying the result to $\sum a_i(X_i+Y_i)$ and to $\sum a_iY_i$ gives the result for $\sum a_iX_i$. However in the linear case, using a different approach, this can be improved for certain values of t (see again [5] and references therein).

• Comment 3: Consider the special case of Theorem 2 where $f(x) = n^{-1/2} \sum_{i=1}^{n} x_i$, and (10) implies

$$\mathbb{P}\left\{ |f(X) - \mathbb{M}f(X)| > C_q t \left(1 + n^{1/q - 1/2} e^{t^2/(2q)}\right) \right\} \le C e^{-t^2/2}$$

which is sub-Gaussian up to probability $Cn^{1-q/2}$ and matches the variance up to the factor C_q . Compare this with the following non-uniform version of the Berry-Esseen bound (see [12, Chapter V §2-4 Theorems 3 and 13]).

Theorem 4 Let $r \in [3, \infty)$, and let $(X_i)_1^n$ be an i.i.d. sequence with $\mathbb{E}X_1 = 0$, $\mathbb{E}X_1^2 = 1$, and $\mathbb{E}|X_1|^r < \infty$. Then for all $x \in \mathbb{R}$,

$$\left| \Phi(x) - \mathbb{P}\left\{ \frac{1}{\sqrt{n}} \sum_{i=1}^{n} X_i \le x \right\} \right| \le C_r \left(1 + |x| \right)^{-r} \left(n^{-1/2} \mathbb{E} \left| X_1 \right|^3 + n^{-(r-2)/2} \mathbb{E} \left| X_1 \right|^r \right)$$

Assuming that $\mathbb{E} |X_1|^r < C_r$ and $n > C_r$, this gives a sub-Gaussian bound on $n^{-1/2} \sum X_i$ up to probability $C_r (\log n)^{-r/2} n^{-1/2}$. There is more that can be said here; the point is simply that as a nonlinear result Theorem 2 does reasonably well in the linear case.

Notation and conventions

M denotes median, C, c, C' etc. denote universal constants that may represent different values at each appearance, dependence on variables will usually be indicated by subscripts, C_q , c_q etc. N denotes $\{1, 2, 3, ...\}$. The term 'sub-Gaussian bound' is used loosely to mean a bound where the probability is sub-Gaussian in terms of the deviation, at least for some values of t (usually in a central region of the distribution). This differs from a sub-Gaussian random variable that would satisfy sub-Gaussian bounds for all t. Upper bounds (for probabilities) of the form $C \exp(-c_1 t^2)$ can be replaced with $2 \exp(-c_2 t^2)$ by taking c_2 sufficiently small, and we do this without further explanation.

2 Proofs

2.1 Outline of the general method

We write $X_i = F_i^{-1}(\Phi(Z_i))$ where $Z = (Z_i)_1^n$ has the standard normal distribution in \mathbb{R}^n . The underlying probability space is of no relevence, and we may assume without loss of generality that such a Z exists. Then $f(X) = \psi(Z)$ where $\psi(x) = f(F_i^{-1}(\Phi(Z_i))_1^n)$. We then apply Pisier's version of the Gaussian concentration inequality, see [13] and Proposition 10 below, to conclude that for all convex $\varphi : \mathbb{R} \to \mathbb{R}$,

$$\mathbb{E}\varphi\left(\psi(Z) - \psi(Y)\right) \le \mathbb{E}\varphi\left(Z'\left|\nabla\psi(Z)\right|\right)$$

where Y is an independent copy of Z and Z' is a random variable in \mathbb{R} with the standard normal distribution independent of Y and Z. By the theory of tail comparison inequalities under convex majorization, in particular by a result of Meilijson and Nádas [11] that is modified in [7] to suite our purposes, see Proposition 11 below, we conclude that the quantiles of $\psi(Z) - \psi(Y)$ cannot be much larger than the quantiles of $Z' |\nabla \psi(Z)|$.

This brings us to the problem of estimating the quantiles of $Z' |\nabla \psi(Z)|$. The gradient is a sum of random variables and one can use the triangle inequality, Hölder's inequality or a different duality to write it in terms of a sum of independent random variables. We then apply Propositions 7, 8 and 9 below to handle such sums.

2.2 Small lemmas

Lemma 5 Let $\delta > 0$ and let Q be a random variable taking values in $[0, \infty)$ such that for all $u, v \in [0, \infty)$,

$$\mathbb{P}\left\{Q \ge uv\right\} \ge \delta e^{-v^2/4} \mathbb{P}\left\{Q \ge u\right\}$$

Let Z' be a random variable with the standard normal distribution in \mathbb{R} . Then for all t > 0,

$$\mathbb{P}\left\{|Z'|Q \ge t\right\} \le C\delta^{-1}\mathbb{P}\left\{Q \ge ct\right\}$$

Proof.

$$\mathbb{P}\left\{ |Z'| \, Q \ge t \right\} = \frac{2}{\sqrt{2\pi}} \int_0^\infty e^{-s^2/2} \mathbb{P}\left\{ Q \ge \frac{t}{s} \right\} ds \le \frac{2}{\sqrt{2\pi}} \int_0^\infty e^{-s^2/4} \delta^{-1} \mathbb{P}\left\{ Q \ge t \right\} ds$$

The following lemma will be used implicitly several times.

Lemma 6 If $f, g : [0, \infty) \to [0, \infty)$ are continuous strictly increasing functions with $f(0) = g(0) = 0, t \in [0, \infty)$ and $s = \max \{f(t), g(t)\}$ then $t = \min \{f^{-1}(s), g^{-1}(s)\}$.

2.3 Tools from [5], [7] and [13]

Proposition 7 (see proof of [5, Lemma 8]) Let $n \in \mathbb{N}$, $\lambda \in [2, \infty)$, and let $(Y_i)_1^n$ be an *i.i.d.* sequence of non-negative random variables, each with cumulative distribution F, quantile function F^{-1} , and corresponding order statistics $(Y_{(i)})_1^n$. With probability at least $1 - 3^{-1}\pi^2 \exp(-\lambda^2/2)$, the following event holds: for all $j, k \in \mathbb{Z}$ with $0 \le j \le k < n$,

$$\sum_{i=n-k}^{n-j} Y_{(i)} \leq F^{-1} \left(1 - \frac{j+1}{n+1} e^{-1} \exp\left(\frac{-\lambda^2 - 4\log(j+1)}{2(j+1)}\right) \right) + (n+1) \int_{(j+1)/(n+1)}^{(k+1)/(n+1)} F^{-1} \left(1 - e^{-1-2/e} t \exp\left(\frac{-\lambda^2}{2(n+1)t}\right) \right) dt$$

Proof sketch. This is based on concentration of order statistics from the uniform distribution on (0,1) which in turn is based on estimates for the binomial distribution. One then transforms the order statistics in (0,1) to the order statistics $(Y_{(i)})_1^n$ using the quantile function. One then peels off the largest term and bounds $\sum_{i=n-k}^{n-j-1} Y_{(i)}$ above by a deterministic sum (with high probability) and compares the deterministic sum to an integral.

Proposition 8 ([5, Proposition 10]) Consider the setting and assumptions of Proposition 7, and assume in addition that there exists p > 0 and $T \ge 1$ such that for all $\delta, x \in (0, 1)$,

$$H^*(\delta x) \ge T^{-1} \delta^{-1/p} H^*(x)$$
 (13)

where $H^*(x) = F^{-1}(1-x)$. With probability at least $1 - 3^{-1}\pi^2 \exp(-\lambda^2/2)$, the following event holds: for all $j, k \in \mathbb{Z}$ with $0 \le j \le k < n$, $\sum_{i=n-k}^{n-j} Y_{(i)}$ is bounded above by

$$\left[1+T\lambda^{2}A\right]H^{*}\left(e^{-1-2/e}\frac{j+1}{n+1}\exp\left(\frac{-\lambda^{2}}{2(j+1)}\right)\right)+Cn\int_{\frac{j+1}{n+1}\exp\left(\frac{-\lambda^{2}}{2(j+1)}-1-2/e\right)}^{\frac{k+1}{n+1}\exp\left(\frac{-\lambda^{2}}{2(j+1)}-1-2/e\right)}H^{*}(x)dx$$

where A = 0 if $\lambda^2/2 \le j + 1$ and A equals

$$C^{1+1/p} \min\left\{p, \lambda^2 \left(\frac{1}{j+1} - \frac{1}{\min\left\{\lambda^2/2, k+1\right\}}\right)\right\} \left(p+1 + \frac{\lambda^2}{j+1}\right)^{-2} + C \min\left\{1, \log\frac{\min\left\{k+1, \lambda^2/2\right\}}{j+1}\right\} \left[\frac{\lambda^2}{2(j+1)} \exp\left(\frac{\lambda^2}{2(j+1)}\right)\right]^{-1/p} \left[1 + \frac{\lambda^2}{k+1}\right]^{-1} if \lambda^2/2 > j+1.$$

Proof sketch. One simplifies the integral in Proposition 7 by making a change of variables and proving various expressions for the resulting integrals that arise. While the details are at times mildly tedious, the ingredients are mostly standard and idea is simple.

Let $(W_i)_1^n$ be an i.i.d. sequence of non-negative random variables such that for all t > 0,

$$\mathbb{P}\left\{W_i > t\right\} = e^{q/2} (e+t)^{-q/2} \left(\ln(e+t)\right)^{q/2}$$

Proposition 9 ([5, Proposition 21]) For all $b \in [0, \infty)^n$ and all t > 0, with probability at least $1 - Ce^{-t^2/2}$,

$$\sum_{i=1}^{n} b_i W_i \le C_q \left(|b|_1 + t^2 e^{t^2/q} |b|_{q/2} \right)$$

Proof sketch. One defines a norm on \mathbb{R}^n , $[\cdot]_{\delta,W}$, which essentially measures the quantiles of $\sum_{1}^{n} |x_i| W_i$. One estimates $[x]_{\delta,W}$ for all $x \in \{0,1\}^n$, and by a duality argument (expressing this norm as the dual norm of its dual norm) one gets a bound on $[x]_{\delta,W}$ for all $x \in \mathbb{R}^n$. This estimate is good for most directions, but for certain directions becomes crude. One then compares $\sum_{1}^{n} |x_i| W_i$ to $\sum_{1}^{n} |x_{I(i)}| W_i$, where $(I(i))_{1}^{n}$ is an i.i.d. sequence of random integers in $\{1, 2, \dots, n\}$. The advantage of this is that the latter sum is a sum of i.i.d. random variables, so one can use Proposition 8. This second approach also works for most coefficient sequences and becomes crude in some cases. When we combine the first approach, which is more geometric, with the second approach, which is more combinatorial, we end up with Proposition 9.

Proposition 10 (Pisier [13]) Let Y and Z be independent random vectors in \mathbb{R}^n each with the standard normal distribution, and let Z' be a random variable with the standard normal distribution in \mathbb{R} independent of Z. Let $\psi : \mathbb{R}^n \to \mathbb{R}$ be differentiable on \mathbb{R} . For all convex functions $\varphi : \mathbb{R} \to \mathbb{R}$,

$$\mathbb{E}\varphi\left(\psi\left(Z^{*}\right)-\psi\left(Z\right)\right) \leq \mathbb{E}\varphi\left(\frac{\pi}{2}\left|\nabla\psi\left(Z\right)\right|Z'\right)$$
(14)

Proof sketch. This follows by an elementary analysis using Jensen's inequality of the corresponding path integral from Y to Z, for a carefully chosen path.

Proposition 11 ([7, Corollary 6]) Let X and Y be real valued random variables such that $\mathbb{E} \max\{0, Y\} < \infty$ and for all $\alpha \in \mathbb{R}$, $\mathbb{E} \max\{0, X - \alpha\} \leq \mathbb{E} \max\{0, Y - \alpha\}$. Suppose that $T, \gamma \geq 1$, p > 1, and $Q: (0, \infty) \to (0, \infty)$ is a function that satisfies

$$Q(t) \exp\left(\frac{-t^2}{2p}\right) \le TQ(s) \exp\left(\frac{-s^2}{2p}\right)$$

for all 0 < s < t. If $\mathbb{P}\{Y > Q(t)\} < \gamma \exp(-t^2/2)$ for all t > 0, then for all t > 0

$$\mathbb{P}\left\{X > \frac{pT}{p-1}Q(t)\right\} \le \gamma \exp\left(-t^2/2\right)$$

Proof sketch. This is based on the result of Meilijson and Nádas [11], that if X and Y are real valued random variables with $\mathbb{E} \max\{0, Y\} < \infty$ and $\mathbb{E}\varphi(X) \leq \mathbb{E}\varphi(Y)$ for all non-decreasing convex $\varphi : \mathbb{R} \to [0, \infty)$, then for all $s \in \mathbb{R}$ with $\mathbb{P}\{Y > s\} \neq 0$, $\mathbb{P}\{X \geq \mathbb{E}(Y|Y > s)\} \leq \mathbb{P}\{Y > s\}$. Unless Y has very thick tails, $\mathbb{E}(Y|Y > s)$ is the same order of magnitude as s; this is quantified in the function Q.

2.4 Proof of Theorem 1

Without loss of generality we may assume that each F_i^{-1} is differentiable. Let $Tx = ((F_i^{-1}\Phi(x_i))_{i=1}^n)$, and let Z be a random vector in \mathbb{R}^n with the standard normal distribution. f_i denotes the partial derivative of f with respect to the i^{th} coordinate. By the assumed bound on $Lip(F_i^{-1}\Phi, Z_i)$ and the triangle inequality,

$$\begin{aligned} |\nabla(f \circ T)(Z)| &= \left(\sum_{i=1}^{n} f_{i} (TZ)^{2} Lip \left(F_{i}^{-1}\Phi, Z_{i}\right)^{2}\right)^{1/2} \\ &\leq \left(\sum_{i=1}^{n} f_{i} (TZ)^{2} (1+|Z_{i}|)^{-2+4/q}\right)^{1/2} \\ &\leq \left(\sum_{i=1}^{n} \sup_{x \in \mathbb{R}^{n}} |f_{i} (x)|^{2} \mathbb{E} (1+|Z_{i}|)^{-2+4/q}\right)^{1/2} \\ &+ \left|\sum_{i=1}^{n} \sup_{x \in \mathbb{R}^{n}} |f_{i} (x)|^{2} \left\{ (1+|Z_{i}|)^{-2+4/q} - \mathbb{E} (1+|Z_{i}|)^{-2+4/q} \right\} \right|^{1/2} \end{aligned}$$
(15)

Since 0 < q < 1, for all t > 0,

$$\mathbb{P}\left\{ \left| (1+|Z_i|)^{-2+4/q} - \mathbb{E} \left(1+|Z_i|\right)^{-2+4/q} \right| > t \right\} \le 2 \exp\left(-c_q t^{q/(2-q)}\right)$$

Noting that 0 < 2/(2-q) < 1 and using (7), with probability at least $1 - 2 \exp(-t^2/2)$,

$$\left|\sum_{i=1}^{n} \sup_{x \in \mathbb{R}^{n}} |f_{i}(x)|^{2} \left\{ (1+|Z_{i}|)^{-2+4/q} - \mathbb{E} \left(1+|Z_{i}|\right)^{-2+4/q} \right\} \right|^{1/2} \le C_{q} \left(t^{1/2} Lip_{4}^{\sharp}(f) + t^{(2-q)/q} Lip_{\infty}^{\sharp}(f) \right)$$

When adding this to the remaining term of $C_q Lip_2^{\sharp}(f)$, the term involving $Lip_4^{\sharp}(f)$ can be erased since by Hölder's inequality $|\cdot|_4 \leq |\cdot|_2^{1/2} |\cdot|_{\infty}^{1/2}$ and since we may assume that $t \geq 1, t^{1/2} \leq (1)^{1/2} (t^{(2-q)/q})^{1/2}$. What we have shown is that with probability at least $1 - 2 \exp(-t^2/2)$,

 $|\nabla(f \circ T)(Z)| \le C_q Lip_2^{\sharp}(f) + C_q t^{(2-q)/q} Lip_{\infty}^{\sharp}(f)$

and therefore with probability at least $1 - C \exp(-t^2/2)$,

$$|Z'| \cdot |\nabla (f \circ T)(Z)| \le C_q t Lip_2^{\sharp}(f) + C_q t^{2/q} Lip_{\infty}^{\sharp}(f)$$

As in Section 2.1, Y is an independent copy of Z, and Z' is a real valued variable with the standard normal distribution, independent of Y and Z. It now follows from Pisier's version of the Gaussian concentration inequality that for all non-decreasing convex $\varphi : \mathbb{R} \to \mathbb{R}$,

$$\mathbb{E}\varphi\left((f\circ T)(Z)-(f\circ T)(Y)\right)\leq \mathbb{E}\varphi\left(Z'\left|\nabla(f\circ T)(Z)\right|\right)\leq \mathbb{E}\varphi\left(|Z'|\cdot\left|\nabla(f\circ T)(Z)\right|\right)$$

By [7, Proposition 6], the quantiles of $(f \circ T)(Z) - (f \circ T)(Y)$ cannot be much larger than those of $|Z'| \cdot |\nabla (f \circ T)(Z)|$ (this should not be surprising, even if the reader is not familiar with [7]). More precisely, with probability at least $C \exp(-t^2/2)$,

$$\left|\left(f\circ T\right)\left(Z\right) - \left(f\circ T\right)\left(Y\right)\right| \le C_q t Lip_2^{\sharp}\left(f\right) + C_q t^{2/q} Lip_{\infty}^{\sharp}\left(f\right)$$

Here we are also using symmetry of the distribution of $(f \circ T)(Z) - (f \circ T)(Y)$. It is well known, and an easy computation, that concentration about an independent random point implies concentration about the median by changing the constants involved (this is a more robust phenomenon than a comparison to concentration about the mean). This proves (5).

For $r \geq 1$ consider the following norm on \mathbb{R}^n ,

$$|x|_{B_1^n \cap r^{-1} B_\infty^n} = \max\{|x|_1, r \, |x|_\infty\}$$

After a brief consideration we see that the dual norm is given by

$$\begin{aligned} |y|_{conv\left(B_{\infty}^{n}, rB_{1}^{n}\right)} &= \max\left\{\sum_{i=1}^{n} x_{i}y_{i} : |x|_{1} \leq 1, |x|_{\infty} \leq r^{-1}\right\} \\ &\leq r^{-1}\sum_{i=1}^{\min\{\lceil r \rceil, n\}} y^{(i)} \leq 2 |y|_{conv\left(B_{\infty}^{n}, rB_{1}^{n}\right)} \end{aligned}$$

where $y^{(1)} \ge y^{(2)} \dots$ is the non-increasing rearrangement of $(|y_i|)_1^n$. Set $r = t^{-2+4/q}$, $k = \lceil r \rceil$ and $Y_i = (1 + |Z_i|)^{-2+4/q}$. Each Y_i has quantile function $H^*(1 - x)$, where

$$H^*(x) \le C_q \left(1 + \log \frac{1}{x}\right)^{-1+2/q}$$

By Proposition 7, with probability at least $1-C \exp(-t^2/2)$, (assuming first that $t^{-2+4/q} \le c_q n$),

$$\begin{split} \sum_{i=1}^{k} Y^{(i)} &= \sum_{i=n-k+1}^{n} Y_{(i)} \le H^* \left(\frac{c}{n} \exp\left(\frac{-t^2}{2}\right) \right) + Cn \int_{1/(n+1)}^{k/(n+1)} H^* \left(cx \exp\left(\frac{-t^2}{2(n+1)x}\right) \right) dx \\ &\le C_q \left(t^2 + \log n \right)^{-1+2/q} + C_q n \frac{k-1}{n+1} + C_q n \int_{1/(n+1)}^{k/(n+1)} \left(\log \frac{1}{x} \right)^{-1+2/q} dx \\ &+ C_q t^{-2+4/q} n^{2-2/q} \int_{1/(n+1)}^{k/(n+1)} x^{1-2/q} dx \\ &\le C_q t^{-2+4/q} + C_q \left(\log n \right)^{-1+2/q} + C_q k + C_q k \left(\log \frac{n+1}{k} \right)^{-1+2/q} + C_q t^{-2+4/q} k^{2-2/q} \end{split}$$

Recalling the relationships between t, k and n, and the fact that 3 - 2/q < 1, this is bounded above by $(n + 1)^{-1+2/q}$

$$C_q k \left(\log \frac{n+1}{k} \right)^{-1+2_j}$$

This implies

$$\left| \left(\left(1 + |Z_i| \right)^{-2+4/q} \right)_1^n \right|_{conv\left(B_\infty^n, rB_1^n\right)} \le C_q \left(\log \frac{n+1}{k} \right)^{-1+2/q}$$

For $t^{-2+4/q} > c_q n$, it follows by classical Gaussian concentration of the $\ell_{-2+4/q}^n$ norm (which is 1-Lipschitz since $-2 + 4/q \ge 2$) that with probability at least $1 - C \exp(-t^2/2)$,

$$\left| \left((1+|Z_i|)^{-2+4/q} \right)_1^n \right|_{conv\left(B_\infty^n, rB_1^n\right)} \le C_q n^{-1} \sum_{i=1}^n (1+|Z_i|)^{-2+4/q} \le C_q + C_q t^{-2+4/q} n^{-1} \le C_q$$

In either case,

$$\left| \left(\left(1 + |Z_i| \right)^{-2+4/q} \right)_1^n \right|_{conv\left(B_\infty^n, rB_1^n\right)} \le C_q \left(\log \left(e + \frac{n}{t^{-2+4/q}} \right) \right)^{-1+2/q}$$

Applying duality to the sum in (15),

$$\left|\nabla\left(f\circ T\right)(Z)\right|^{2} \leq \left|\left(f_{i}\left(TZ\right)^{2}\right)_{1}^{n}\right|_{B_{1}^{n}\cap r^{-1}B_{\infty}^{n}}\left|\left(\left(1+|Z_{i}|\right)^{-2+4/q}\right)_{1}^{n}\right|_{conv\left(B_{\infty}^{n},rB_{1}^{n}\right)}\right|_{Conv\left(B_{\infty}^{n},rB_{1}^{n}\right)}\right|_{Conv\left(B_{\infty}^{n},rB_{1}^{n}\right)}$$

 \mathbf{SO}

$$|\nabla (f \circ T) (Z)| \le C_q \left(\log \left(e + \frac{n}{t^{-2+4/q}} \right) \right)^{1/q-1/2} \left(Lip_2 (f) + t^{(2-q)/q} Lip_\infty (f) \right)$$

(6) now follows, as earlier in the proof, from Pisier's version of Gaussian concentration and applying tail comparison under convex majorization.

2.5 Proof of Theorem 2

Without loss of generality we may assume that each F_i^{-1} is differentiable, and that $Lip_2^{\sharp}(f) = 1$. Let $Tx = (F_i^{-1}\Phi(x_i))_{i=1}^n$ and let Z be a random vector in \mathbb{R}^n with the standard normal distribution. Using the estimate

$$\frac{\phi(x)}{\min\{\Phi(x), 1 - \Phi(x)\}} \le C(1 + |x|) \le C\sqrt{\log\min\{\Phi(x), 1 - \Phi(x)\}^{-1}}$$

valid for all $x \in \mathbb{R}$, which follows by integrating the tails of ϕ using log-concavity, and using

$$\begin{aligned} Lip\left(F_i^{-1} \circ \Phi, x\right) &\leq Lip\left(F_i^{-1}, \Phi(x)\right) \cdot Lip\left(\Phi, x\right) \\ &\leq \min\left\{\Phi(x), 1 - \Phi(x)\right\}^{-1/q} \frac{\phi(x)}{\min\left\{\Phi(x), 1 - \Phi(x)\right\}} \end{aligned}$$

we see that

$$|\nabla (f \circ T) (Z)| = \left(\sum_{i=1}^{n} f_i (TZ)^2 Lip \left(F_i^{-1} \Phi, Z_i\right)^2\right)^{1/2} \le \left(\sum_{i=1}^{n} f_{i,\sharp}^2 U_i\right)^{1/2}$$

where

$$f_{i,\sharp} = \sup_{x \in \mathbb{R}^n} |f_i(x)| \qquad U_i = C \min \left\{ \Phi(Z_i), 1 - \Phi(Z_i) \right\}^{-2/q} \log \min \left\{ \Phi(Z_i), 1 - \Phi(Z_i) \right\}^{-1}$$
(16)

It follows from the definition that each U_i has quantile function

$$G^{-1}(t) = \left(\frac{1-t}{2}\right)^{-2/q} \log\left(\frac{1-t}{2}\right)^{-1} : 0 < t < 1$$
(17)

By Proposition 9,

$$\mathbb{P}\left\{\sum_{i=1}^{n} f_{i,\sharp}^{2} U_{i} > C_{q} \sum_{i=1}^{n} f_{i,\sharp}^{2} + C_{q} t^{2} e^{t^{2}/q} \left(\sum_{i=1}^{n} f_{i,\sharp}^{q}\right)^{q/2}\right\} \le C e^{-t^{2}/2}$$

Recalling the expression for $|\nabla (f \circ T)(Z)|$ in terms of the U_i : with probability at least $1 - Ce^{-t^2/2}$,

$$\left|\nabla\left(f\circ T\right)\left(Z\right)\right| \le C_q \left(Lip_2^{\sharp}\left(f\right) + te^{t^2/(2q)}Lip_q^{\sharp}\left(f\right)\right)$$

This means we can write

$$\left|\nabla\left(f\circ T\right)\left(Z\right)\right| = C_q Lip_2^{\sharp}\left(f\right) + C_q G Lip_q^{\sharp}\left(f\right)$$

where G is a random variable with heavy tails (polynomial type heaviness). Upon multiplication, a random variable with sufficiently heavy tails absorbs a standard normal random variable, in that the quantiles of the product are the same order of magnitude. This is made precise, for example, in Lemma 5. If Z' is a standard normal random variable in \mathbb{R} independent of Z, then this observation means that with probability at least $1 - Ce^{-t^2/2}$,

$$|Z'| C_q Lip_2^{\sharp}(f) \le C_q t Lip_2^{\sharp}(f) \qquad \text{and} \qquad |Z'| C_q G Lip_q^{\sharp}(f) \le C_q t e^{t^2/(2q)} Lip_q^{\sharp}(f)$$

 \mathbf{SO}

$$|Z'| \cdot |\nabla (f \circ T) (Z)| \le C_q t \left(Lip_2^{\sharp} (f) + e^{t^2/(2q)} Lip_q^{\sharp} (f) \right)$$

Eq. (10) now follows, as in the proof of Theorem 1, from Gaussian concentration and tail comparison under convex majorization. Here one uses Proposition 11 with, say, $T = C_q$ and p = 1.5. We now consider Eq. (11). No longer assuming that $Lip_2^{\sharp}(f) = 1$, using Hölder's inequality for $\ell_{p/2}$ and $\ell_{p/(p-2)}$,

$$|\nabla (f \circ T) (Z)| \le Lip_p(f) \left(\sum_{i=1}^n U_i^{p/(p-2)}\right)^{(p-2)/(2p)}$$

By Proposition 8, with probability at least $1 - C \exp(-t^2/2)$,

$$\left(\sum_{i=1}^{n} U_i^{p/(p-2)}\right)^{(p-2)/(2p)} \le C_{p,q} \left(n^{1/2-1/p} + n^{1/q} t e^{t^2/(2q)}\right)$$

The result now follows, as before, by absorbing a Gaussian into a heavy tailed random variable, and applying Gaussian concentration and convex majorization.

Acknowledgements

Thanks to Gusti van Zyl and the anonymous referee for comments and suggestions. Part of this work was done while the author was a postdoctoral fellow at the Weizmann Institute of Science.

References

- [1] Barthe, F., Cattiaux, P., Roberto C.: Concentration for independent random variables with heavy tails. AMRX, Appl. Math. Res. Express (2), 39-60 (2005)
- [2] Cattiaux, P., Gozlan, N., Guillin A., Roberto, C.: Functional inequalities for heavy tailed distributions and application to isoperimetry. Electon. J. Probab. 15 (13), 346-385 (2010)
- [3] Cordero-Erausquin, D., Ledoux, M.: Hypercontractive measures, Talagrand's inequality, and influences. Geometric aspects of functional analysis, 169-189, Lecture Notes in Math., 2050, Springer (2012)
- [4] Fresen, D. J: Variations and extensions of the Gaussian concentration inequality, Part I. To appear in Quaest. Math. Published online at https://www.tandfonline.com/doi/abs/10.2989/16073606.2022.2074908. Preprint available at https://arxiv.org/abs/1812.10938
- [5] Fresen, D. J: Deviation inequalities for linear combinations of random variables with power tails. arXiv: 2207.01867 (results cited with numbering as in v3, v4 to be released soon with updated title)
- [6] Fresen, D. J: Random Euclidean embeddings in finite dimensional Lorentz spaces. To appear. arXiv:2104.11974
- [7] Fresen, D. J: Optimal tail comparison under convex majorization. arXiv: 2207.01872.
- [8] Gozlan, N.: Poincaré inequalities and dimension free concentration of measure. Ann. Inst. Henri Poincaré Probab. Stat. 46 (3), 708-739 (2010)
- [9] Hitczenko, P., Montgomery-Smith, S. J., Oleszkiewicz, K.: Moment inequalities for sums of certain independent symmetric random variables. Studia Math. 123 (1), 15-42 (1997)

- [10] Latała, R.: Estimates of moments of sums of independent real random variables. Ann. Probab. 25 (3), 1502-1513 (1997)
- [11] Meilijson, I., Nádas, A.: Convex Majorization with an Application to the Length of Critical Paths. J. Appl. Prob. 16 (3), 671-677 (1979)
- [12] Petrov, V. V.: Sums of independent random variables. Springer-Verlag 1975 (translated from Russian into English)
- [13] Pisier, G.: Probabilistic methods in the geometry of Banach spaces. CIME, Varenna, 1985. Lecture Notes in Mathematics 1206, 167-241 (1986)
- [14] Talagrand, M.: The supremum of some canonical processes. Amer. J. Math. 116 (2), 283-325 (1994)