Binomial Approximation to Locally Dependent CDO

Amit N. Kumar^{*} and P. Vellaisamy[†]

^{*}Department of Mathematical Sciences, Indian Institute of Technology (BHU), Varanasi (Uttar Pradesh) – 221005, India.

[†] Department of Mathematics, Indian Institute of Technology Bombay, Powai, Mumbai – 400076, India.

> *Email: amit.kumar2703@gmail.com †Email: pv@math.iitb.ac.in

Abstract

In this paper, we develop Stein's method for binomial approximation using the stop-loss metric that allows one to obtain a bound on the error term between the expectation of call functions. We obtain the results for a locally dependent collateralized debt obligation (CDO), under certain conditions on moments. The results are also exemplified for an independent CDO. Finally, it is shown that our bounds are sharper than the existing bounds.

Keywords: Binomial distribution; error bounds; Stein's method; CDO. MSC 2020 Subject Classifications: Primary: 62E17, 62E20; Secondary: 60F05, 60E05.

1 Introduction

The collateralized debt obligation (CDO) is a financial tool of structured asset-backed security. It is used to repackage assets into a product and sold to investors in the secondary market. These packages are in terms of debt such as auto loans, credit card debt, mortgages, and corporate debt. The assets are sliced into tranches, a set of repayment ties with different payment priorities and interest rates. The elementary tranches are senior (low risk and low return), mezzanine and equity (high risk and high return). Investors can choose a tranche of their interest to invest. For more details, see Jiao and Karoui [11], Jiao *et al.* [12], Hull and White [13], Kumar [14, 15], Neammanee and Yonghint [18], Yonghint *et al.* [23], and the reference therein.

Condider the tranche pricing with n portfolios, where each portfolio has a constant recovery rate R > 0. Then, the percentage loss at time T can be defined as (see Neammane and Yonghint [18, p. 2])

$$L(T) = \frac{(1-R)}{n} \sum_{i=1}^{n} \mathbf{1}_{\{\tau_i \le T\}},$$
(1.1)

where τ_i is the default time of the *i*-th portfolio and $\mathbf{1}_A$ denotes the indicator function of the set A. For a detachment point or an attachment point, say z^* , the main problem of CDO pricing is to evaluate the value of the mean of percentage total loss for each tranche given by (see Yonghint *et al.* [23, p. 2])

$$\mathbb{E}[(L(T) - z^*)^+] = \frac{(1-R)}{n} \mathbb{E}\left[(W_n - z)^+\right]$$

where $h(x) = x^+ = \max\{x, 0\}$ is the call function, $W_n = \sum_{i=1}^n \mathbf{1}_{\{\tau_i \leq T\}}$ and $z = (nz^*/(1-R)) > 0$. For additional details, see, for example, Kumar [14, 15], Neammanee and Yonghint [18], Yonghint *et al.* [23].

In general, the mean of percentage total loss is difficult to compute when $\mathbf{1}_{\{\tau_i \leq T\}}$, i = 1, 2, ..., n, are locally dependent random variables. Therefore, it is of interest to find a suitable approximating distribution which is close to W_n . The proximity between $\mathbb{E}[(W_n - z)^+]$ and $\mathbb{E}[(\mathbf{P}_{\lambda} - z)^+]$ is studied by Yonghint *et al.* [23], where \mathbf{P}_{λ} denotes the Poisson random variable with parameter λ . Observe that W_n is the sum of dependent Bernoulli random variables and so the binomial distribution also could a good choice to approximate W_n . See Vellaisamy and Punnen [21], where it is shown that the binomial distribution may arise as the distribution of the sum of a certain dependent Bernoulli random variables. Therefore, we choose the binomial distribution as the target distribution.

Throughout this paper, let $B_{\alpha,p}$ follow the binomial distribution with probability mass function

$$\mathbb{P}(\mathbf{B}_{\alpha,p}=k) = \binom{\alpha}{k} p^k q^{\alpha-k}, \quad \text{for } k = 0, 1, \dots, \alpha,$$
(1.2)

where α is a positive integer and 0 . We derive the result using the stop-loss metric defined by

$$d_{sl}(X,Y) = \sup_{z \in \mathbb{R}} \left| \mathbb{E}[(X-z)^+] - \mathbb{E}[(Y-z)^+] \right|,$$
(1.3)

where \mathbb{R} denotes the set of real numbers. For more details, see Bautsikas and Veggelatou [4].

The paper is organized as follows. In Section 2, we develop Stein's method for binomial distribution, under the stop-loss metric. In particular, we discuss uniform and non-uniform bounds for the solution of the Stein equation. In Section 3, we derive the error in approximation of a locally dependent CDO to a suitable binomial distribution. As a special case, we demonstrate the results for the independent CDO. Also, we give some numerical comparisons between our results and the existing results. It is shown that our bounds improve significantly over the existing bounds.

2 Stein's Method

Stein [20] proposed an elegant method to find the error in approximating the sums of real-valued random variables to a normal distribution. Then, Chen [9] adapted the technique for approximating the sums of discrete random variables to a suitable Poisson distribution. Later, numerous authors adapted or used Stein's method for several distributions and under various distance metrics. To mention a few relevant ones, Stein's method for negative binomial distribution has been studied by Brown and Phillips [5] under the total variation distance and for the Poisson distribution has been studied by Neammanee and Yonghint [18], under the stop-loss metric. For recent developments, see Barbour *et al.* [1], Brown and Xia [6], Eichelsbacher and Reinert [10] and Kumar [15].

In this section, we develop the Stein's method for the binomial distribution, under the stop-loss metric defined in (1.3). Our work is mainly focused on finding the upper bound for $|\Delta g_z(k)| = |g_z(k+1) - g_z(k)|$, where $g_z(k)$ is the solution of the Stein equation given by

$$\mathcal{A}g_{z}(k) = (k-z)^{+} - \mathbb{E}(\mathbf{B}_{\alpha,p} - z)^{+}.$$
(2.1)

Here, A denotes the Stein operator of $B_{\alpha,p}$ given by

$$\mathcal{A}g_{z}(k) = \frac{(\alpha - k)p}{q}g_{z}(k+1) - kg_{z}(k), \quad \text{for } k = 0, 1, \dots, \alpha.$$
(2.2)

Note that $\mathbb{E}(Ag_z(\mathbf{B}_{\alpha,p})) = 0$. Using (2.2), the Stein equation (2.1) leads to

$$\frac{(\alpha - k)p}{q}g_z(k+1) - kg_z(k) = (k-z)^+ - \mathbb{E}(\mathbf{B}_{\alpha,p} - z)^+.$$

From Section 2 of Kumar *et al.* [16, p. 4] with appropriate changes, it can easily be verified that the solution of the above equation is

$$g_{z}(k) = \begin{cases} 0, & \text{if } k = 0; \\ -\sum_{j=k}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{(k-1)!}{j!} \left(\frac{p}{q}\right)^{j-k} [(j-z)^{+} - \mathbb{E}(\mathbf{B}_{\alpha,p} - z)^{+}], & \text{if } 1 \le k \le \alpha. \end{cases}$$
(2.3)

The bounds for $|\Delta g_z(k)|$ may be essentially uniform for all k and z. However, the bound depends on k may be useful in practice to get a sharper bound for the stop-loss metric. So, we obtain the non-uniform upper bound for $|\Delta g_z(k)|$.

The following lemma gives the non-uniform upper bound for $|\Delta g_z(k)|$, for all $z \ge 0$.

Lemma 2.1. For $z \ge 0$, the following inequality holds:

$$|\Delta g_z(k)| \le \begin{cases} 2q^{1-\alpha} - q, & \text{for } k = 0;\\ 2q^{k-\alpha}, & \text{for } 1 \le k \le \alpha. \end{cases}$$

Proof. It can be easily verified that $\mathbb{E}[(\mathbf{B}_{\alpha,p}-z)^+] \leq \alpha p$. For k = 0, we have

$$\begin{aligned} |\Delta g_z(0)| &= |g_z(1)| \le \sum_{j=1}^{\alpha} \frac{(\alpha-1)!}{(\alpha-j)!j!} \left(\frac{p}{q}\right)^{j-1} \left[(j-z)^+ + \mathbb{E}(B_{\alpha,p}-z)^+\right] \\ &\le \sum_{j=1}^{\alpha} \left(\frac{\alpha-1}{j-1}\right) \left(\frac{p}{q}\right)^{j-1} + p \sum_{j=1}^{\alpha} \binom{\alpha}{j} \left(\frac{p}{q}\right)^{j-1} \\ &= 2q^{1-\alpha} - q. \end{aligned}$$

For $1 \le k \le \alpha - 1$, let

$$h_1(k) = \sum_{j=k}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{(k - 1)!}{j!} \left(\frac{p}{q}\right)^{j-k} (j - z)^+$$

and

$$h_2(k) = \sum_{j=k}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{(k-1)!}{j!} \left(\frac{p}{q}\right)^{j-k} \mathbb{E}(\mathbf{B}_{\alpha,p}-z)^+.$$

Then,

$$\Delta g_z(k) = g_z(k+1) - g_z(k) = [h_1(k) - h_1(k+1)] - [h_2(k) - h_2(k+1)].$$
(2.4)

First, consider

$$0 < h_1(k) = \sum_{j=k}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{(k - 1)!}{j!} \left(\frac{p}{q}\right)^{j-k} (j - z)^+$$

$$\leq \sum_{j=k}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{(k - 1)!}{(j - 1)!} \left(\frac{p}{q}\right)^{j-k}$$

$$= 1 + \sum_{j=k+1}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{1}{k(k+1)\dots(j-1)} \left(\frac{p}{q}\right)^{j-k}$$

$$\leq 1 + \sum_{j=k+1}^{\alpha} \binom{\alpha - k}{j-k} \left(\frac{p}{q}\right)^{j-k}$$

$$= 1 + \sum_{j=1}^{\alpha-k} {\alpha-k \choose j} \left(\frac{p}{q}\right)^j$$
$$= q^{k-\alpha}.$$
 (2.5)

Similarly, $0 < h_1(k+1) \le q^{k+1-\alpha} \le q^{k-\alpha}$. Now, consider

$$0 < h_{2}(k) = \sum_{j=k}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{(k-1)!}{j!} \left(\frac{p}{q}\right)^{j-k} \mathbb{E}(\mathbf{B}_{\alpha,p}-z)^{+}$$

$$\leq \frac{\alpha p}{k} \sum_{j=k}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{k!}{j!} \left(\frac{p}{q}\right)^{j-k}$$

$$\leq \frac{\alpha p}{k} \left(1 + \sum_{j=k+1}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{1}{(k+1)\dots j} \left(\frac{p}{q}\right)^{j-k}\right)$$

$$= \frac{\alpha p}{k} \left(1 + \frac{1}{\alpha-k+1} \sum_{j=k+1}^{\alpha} \left(\frac{\alpha-k+1}{j-k+1}\right) \left(\frac{p}{q}\right)^{j-k}\right)$$

$$= \frac{\alpha p}{k} \left(1 + \frac{1}{\alpha-k+1} \sum_{j=1}^{\alpha-k} \left(\frac{\alpha-k+1}{j+1}\right) \left(\frac{p}{q}\right)^{j}\right)$$

$$= \frac{\alpha p}{k} \left(1 + \frac{q^{k-\alpha} - (\alpha-k)p - 1}{(\alpha-k+1)p}\right)$$

$$\leq \frac{\alpha p q^{k-\alpha}}{(\alpha-k+1)kp}$$

$$\leq q^{k-\alpha}.$$
(2.6)

Similarly, it follows that $0 < h_2(k+1) \le q^{k+1-\alpha} \le q^{k-\alpha}$. Therefore, from (2.4), (2.5) and (2.6), the result follows for $1 \le k \le \alpha - 1$.

Finally, for $k = \alpha$, it is easy to verify that

$$|\Delta g_z(\alpha)| = |g_z(\alpha)| = \frac{(\alpha - z)^+}{\alpha} + \frac{\mathbb{E}(B_{\alpha, p} - z)^+}{\alpha} \le 2.$$

This proves the result.

Remark 2.1. (i) Observe that the uniform bound can be taken as

$$|\Delta g_z(k)| \le 2q^{1-\alpha}, \quad \text{for all } z \ge 0 \text{ and } 0 \le k \le \alpha.$$
(2.7)

(ii) From (2.3), (2.5) and (2.6), it can be easily seen that

$$|g_z(k)| \le 2q^{k-n}, \quad \text{for all } z \ge 0.$$
(2.8)

Also, the uniform bound for $|g_z(k)|$ is the same as given in (2.7).

Next, using the similar technique of the proof of Lemma 2.1, we obtain the following lemma which gives the non-uniform upper bound for $|\Delta g_z(k)|$ for all z > 1.

Lemma 2.2. For z > 1, the following inequality holds:

$$|\Delta g_z(k)| \le \begin{cases} 2\left(1 + \frac{q^{k-\alpha} - 1}{qz}\right), & \text{for } k \ge z;\\ \frac{3\left(q^{k-\alpha} - 1\right)}{pz}, & \text{for } 2 \le k < z;\\ \frac{2(\alpha - 1)pq^{1-\alpha}}{z}, & \text{for } 1 = k < z. \end{cases}$$

Proof. Let $k \ge z$. First, consider

$$0 < h_{1}(k) = \sum_{j=k}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{(k - 1)!}{j!} \left(\frac{p}{q}\right)^{j-k} (j - z)^{+}$$

$$\leq 1 + \sum_{j=k+1}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{(k - 1)!}{(j - 1)!} \left(\frac{p}{q}\right)^{j-k}$$

$$= 1 + \frac{(\alpha - k)p}{kq} + \frac{1}{k} \sum_{j=k+2}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{1}{(k + 1) \dots (j - 1)} \left(\frac{p}{q}\right)^{j-k}$$

$$\leq 1 + \frac{(\alpha - k)p}{kq} + \frac{1}{k} \sum_{j=2}^{\alpha-k} \binom{\alpha - k}{j} \left(\frac{p}{q}\right)^{j}$$

$$= 1 + \frac{(\alpha - k)p}{kq} + \frac{1}{k} \left(q^{k-\alpha} - \frac{(\alpha - k)p}{q} - 1\right)$$

$$\leq 1 + \frac{q^{k-\alpha} - 1}{z}.$$
(2.9)

Next, for $k \ge z$ and $k \ge \alpha p$, we have

$$0 < \sum_{j=k}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{(k-1)!}{j!} \left(\frac{p}{q}\right)^{j-k} = \frac{1}{k} \sum_{j=k}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{k!}{j!} \left(\frac{p}{q}\right)^{j-k} \leq \frac{1}{\alpha p} \left(1 + \frac{(\alpha-k)p}{(k+1)q} + \sum_{j=k+2}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{k!}{j!} \left(\frac{p}{q}\right)^{j-k} \right) = \frac{1}{\alpha p} \left(1 + \frac{(\alpha-k)p}{(k+1)q} + \frac{1}{(k+1)} \sum_{j=k+2}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{(k+1)!}{j!} \left(\frac{p}{q}\right)^{j-k} \right) = \frac{1}{\alpha p} \left(1 + \frac{(\alpha-k)p}{(k+1)q} + \frac{1}{(k+1)} \sum_{j=k+2}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{1}{(k+2)\dots j} \left(\frac{p}{q}\right)^{j-k} \right) \leq \frac{1}{\alpha p} \left(1 + \frac{(\alpha-k)p}{(k+1)q} + \frac{1}{(k+1)} \sum_{j=k+2}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!(j-k)!} \left(\frac{p}{q}\right)^{j-k} \right) = \frac{1}{\alpha p} \left(1 + \frac{(\alpha-k)p}{(k+1)q} + \frac{1}{(k+1)} \sum_{j=2}^{\alpha-k} \binom{\alpha-k}{j} \left(\frac{p}{q}\right)^{j} \right) = \frac{1}{\alpha p} \left(1 + \frac{(\alpha-k)p}{(k+1)q} + \frac{1}{(k+1)} \left(q^{k-\alpha} - \frac{(\alpha-k)p}{q} - 1\right) \right) \leq \frac{1}{\alpha p} \left(1 + \frac{q^{k-\alpha} - 1}{z} \right).$$

$$(2.10)$$

Also, for $k \ge z$ and $k \le \alpha p$, we have

$$0 < \sum_{j=k}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{(k-1)!}{j!} \left(\frac{p}{q}\right)^{j-k}$$

$$= \frac{1}{\alpha-k+1} \sum_{j=k}^{\alpha} \frac{(\alpha-k+1)!}{(\alpha-j)!} \frac{(k-1)!}{j!} \left(\frac{p}{q}\right)^{j-k}$$

$$= \frac{1}{\alpha-k+1} \left(\frac{\alpha-k+1}{k} + \frac{1}{k} \sum_{j=k+1}^{\alpha} \frac{(\alpha-k+1)!}{(\alpha-j)!} \frac{k!}{j!} \left(\frac{p}{q}\right)^{j-k}\right)$$

$$= \frac{1}{\alpha-k+1} \left(\frac{\alpha-k+1}{k} + \frac{1}{k} \sum_{j=k+1}^{\alpha} \frac{(\alpha-k+1)!}{(\alpha-j)!} \frac{1}{(k+1)\dots j} \left(\frac{p}{q}\right)^{j-k}\right)$$

$$\leq \frac{1}{\alpha-k+1} \left(\frac{\alpha-k+1}{k} + \frac{1}{k} \sum_{j=k+1}^{\alpha} \frac{(\alpha-k+1)!}{(\alpha-j)!(j-k+1)!} \left(\frac{p}{q}\right)^{j-k}\right)$$

$$= \frac{1}{\alpha - k + 1} \left(\frac{\alpha - k + 1}{k} + \frac{1}{k} \sum_{j=1}^{\alpha} \binom{\alpha - k + 1}{j + 1} \binom{p}{q}^{j} \right)$$

$$= \frac{1}{\alpha - k + 1} \left(\frac{\alpha - k + 1}{k} + \frac{1}{k} \left(\frac{q^{k - \alpha} - 1}{p} - (\alpha - k) \right) \right)$$

$$= \frac{1}{\alpha - k + 1} \left(\frac{1}{k} + \frac{q^{k - \alpha} - 1}{kp} \right)$$

$$\leq \frac{1}{(\alpha - k + 1)k} + \frac{q^{k - \alpha} - 1}{(\alpha - k)zp}$$

$$\leq \frac{1}{\alpha} + \frac{q^{k - \alpha} - 1}{\alpha pqz}$$

$$\leq \frac{1}{\alpha p} \left(1 + \frac{q^{k - \alpha} - 1}{qz} \right).$$
(2.11)

Therefore, form (2.10) and (2.11), for $k \ge z$, we have

$$0 < h_2(k) = \sum_{j=k}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{(k - 1)!}{j!} \left(\frac{p}{q}\right)^{j-k} \mathbb{E}(B_{\alpha, p} - z)^+ \\ \le 1 + \frac{q^{k-\alpha} - 1}{qz}.$$
(2.12)

Hence, from (2.4), (2.9) and (2.12), the result follows for $k \ge z$. Next, let $2 \le k < z$. Note that

$$0 < h_1(k) = \sum_{j=\lceil z \rceil}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{(k - 1)!}{j!} \left(\frac{p}{q}\right)^{j-k} (j - z)$$

$$\leq \frac{1}{z} \sum_{j=\lceil z \rceil}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{(k - 1)!}{(j - 1)!} \left(\frac{p}{q}\right)^{j-k} (j - z)$$

$$\leq \frac{1}{z} \sum_{j=k+1}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{(k - 1)!}{(j - 1)!} \left(\frac{p}{q}\right)^{j-k} (j - z)^+$$

$$\leq \frac{1}{z} \sum_{j=k+1}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{(k - 1)!}{(j - 2)!} \left(\frac{p}{q}\right)^{j-k}$$

$$= \frac{1}{z} \left(\frac{(\alpha - k)p}{q} + \sum_{j=k+2}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{1}{k(k+1)\dots(j-2)} \left(\frac{p}{q}\right)^{j-k}\right)$$

$$\leq \frac{1}{z} \left(\frac{(\alpha - k)p}{q} + \sum_{j=k+2}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{1}{2.3\dots(j-k)} \left(\frac{p}{q}\right)^{j-k}\right)$$

$$= \frac{1}{z} \left(\frac{(\alpha - k)p}{q} + \sum_{j=2}^{\alpha - k} {\alpha - k \choose j} \left(\frac{p}{q} \right)^j \right)$$
$$= \frac{q^{k-\alpha} - 1}{z}.$$
 (2.13)

Next, consider

$$\begin{split} 0 &< \sum_{j=k}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{(k-1)!}{j!} \left(\frac{p}{q}\right)^{j-k} \\ &= \frac{1}{k} \sum_{j=k}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{k!}{j!} \left(\frac{p}{q}\right)^{j-k} \\ &= \frac{1}{k} \left(1 + \sum_{j=k+1}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!} \frac{1}{(k+1)\dots j} \left(\frac{p}{q}\right)^{j-k}\right) \\ &\leq \frac{1}{k} \left(1 + 2\sum_{j=k+1}^{\alpha} \frac{(\alpha-k)!}{(\alpha-j)!(j-k+2)!} \left(\frac{p}{q}\right)^{j-k}\right) \\ &\leq \frac{1}{k} \left(1 + \frac{2}{(\alpha-k)(\alpha-k+1)} \sum_{j=1}^{\alpha-k} \binom{\alpha-k+2}{j+2} \left(\frac{p}{q}\right)^{j}\right) \\ &= \frac{1}{k} \left(1 + \frac{2}{(\alpha-k)(\alpha-k+1)} \left(\frac{2q^{k-\alpha}-2 - 2(\alpha-k)p - (\alpha-k)(\alpha-k+1)p^2}{2p^2}\right)\right) \right) \\ &\leq \frac{2(q^{k-\alpha}-1)}{(\alpha-k)(\alpha-k+1)kp^2} \\ &\leq \frac{2(q^{k-\alpha}-1)}{\alpha(\alpha-z)p^2}. \end{split}$$

Note that, for $k < z \leq \alpha$, we have

$$\mathbb{E}(B_{\alpha,p}-z)^{+} = \sum_{m=\lceil z \rceil}^{\alpha} (m-z) \binom{\alpha}{m} p^{m} q^{\alpha-m}$$
$$\leq (\alpha-z) \sum_{m=\lceil z \rceil}^{\alpha} \frac{\alpha!}{(\alpha-m!)m!} p^{m} q^{\alpha-m}$$
$$\leq \frac{\alpha(\alpha-z)}{z} \sum_{m=1}^{\alpha} \binom{\alpha-1}{m-1} p^{m} q^{\alpha-m}$$
$$= \frac{\alpha(\alpha-z)p}{z}$$

Therefore,

$$0 < h_2(k) = \sum_{j=k}^{\alpha} \frac{(\alpha - k)!}{(\alpha - j)!} \frac{(k - 1)!}{j!} \left(\frac{p}{q}\right)^{j-k} \mathbb{E}(B_{\alpha,p} - z)^+ \\ \leq \frac{2(q^{k-\alpha} - 1)}{pz}.$$
(2.14)

Hence, from (2.4), (2.13) and (2.14), the result follows for $2 \le k < z \le \alpha$. Next, for 1 = k < z, we have

$$0 < h_1(1) = \sum_{j=\lceil z \rceil}^{\alpha} \frac{(\alpha-1)!}{(\alpha-j)!j!} \left(\frac{p}{q}\right)^{j-1} (j-z)$$
$$\leq \frac{\alpha-1}{z} \sum_{j=2}^{\alpha} \binom{\alpha-2}{j-2} \left(\frac{p}{q}\right)^{j-1}$$
$$= \frac{(\alpha-1)pq^{1-\alpha}}{z}.$$

and

$$0 < h_2(1) = \sum_{j=1}^{\alpha} \frac{(\alpha - 1)!}{(\alpha - j)! j!} \left(\frac{p}{q}\right)^{j-1} \mathbb{E}(B_{\alpha, p} - z)^+$$
$$\leq \left(\frac{q^{1-\alpha} - q}{\alpha p}\right) \left(\frac{\alpha(\alpha - 1)p^2}{z}\right)$$
$$= \frac{(\alpha - 1)pq^{1-\alpha}}{z}.$$

This proves the result.

3 Bounds for Binomial Approximation

In this section, we obtain the error bounds for binomial approximation to locally dependent CDO. We derive the results for the stop-loss metric under certain conditions on moments. Moreover, we demonstrate the results under an independent setup. It is shown that binomial distribution is more suitable for a CDO using the numerical comparison between our bounds and the existing bound given by Neammanee and Yonghint [18].

In Yonghint *et al.* [23], it is shown that the locally dependent CDO is useful in real-life applications in various aspects. So, we consider a similar locally dependent structure that can also be used for independent setup. Let X_1, X_2, \ldots, X_n be a collection of random variables such that X_i is independent

of $X_{A_i^c}$, while X_{A_i} is independent of $X_{B_i^c}$, where $i \in A_i \subseteq B_i \subset \{1, 2, ..., n\}$, i = 1, 2, ..., n. Here, X_A denotes the collection of random variables $\{X_i, i \in A\}$, and A^c denotes the complement of the set A. Note that if $A_i = B_i = \{i\}$, then $X_1, X_2, ..., X_n$ become independent random variables. See Kumar [14, 15], Röllin [19] and Čekanavičius and Vellaisamy [7, 8] for a similar locally dependent setup.

Let τ_i is the default time of the *i*-th portfolio and X_i henceforth denote the random variable $\mathbf{1}_{\{\tau_i \leq T\}}$ with $\mathbb{P}(X_i = 1) = p_i = 1 - q_i = 1 - \mathbb{P}(X_i = 0)$. Consider

$$W_n = \sum_{i=1}^n X_i,\tag{3.1}$$

which is a key factor of percentage loss up to time T defined in (1.1). Our aim is to approximate W_n by a suitable binomial random variable.

Throughout this section, let $g_z = g$,

$$W_i = W_n - X_{A_i} = \sum_{j \notin A_i} X_i$$
 and $W_i^* = W_n - X_{B_i} = \sum_{j \notin B_i} X_i.$ (3.2)

Note that X_i is independent of W_i . Also, X_i and X_{A_i} are independent of W_i^* .

First, we choose one parameter of binomial distribution of our choice, and accordingly, another parameter can be obtained. Since α should be a positive integer (number of identical Bernoulli trials) for the binomial distribution, we choose $\alpha = n$ and let

$$p = \frac{1}{\alpha} \sum_{i=1}^{n} \mathbb{E}(X_i) = \frac{1}{n} \sum_{i=1}^{n} p_i.$$
(3.3)

The following theorem gives the error in the approximation between $B_{\alpha,p}$ and W_n .

Theorem 3.1. Let $B_{\alpha,p}$ and W_n be defined as in (1.2) and (3.1), respectively, and satisfy (3.3). Then

$$d_{sl}(W_n, \boldsymbol{B}_{n,p}) \le \frac{2}{pq^n} \sum_{i=1}^n \left[\mathbb{E}\left((X_i + p)q^{W_i} \right) - \mathbb{E}\left((p_i + qX_i)q^{W_n} \right) \right].$$
(3.4)

where d_{sl} denote the stop-loss distance defined in (1.3).

Proof. Multiplying the Stein operator (2.2) by q and taking expectation with respect to W_n , we have

$$q\mathbb{E}[\mathcal{A}g(W_n)] = \alpha p\mathbb{E}(g(W_n+1)) - p\mathbb{E}(W_ng(W_n+1)) - q\mathbb{E}(W_ng(W_n)).$$
(3.5)

Using (3.3), we get

$$q\mathbb{E}[\mathcal{A}g(W_n)] = \sum_{i=1}^n \mathbb{E}(X_i)\mathbb{E}(g(W_n+1)) - \sum_{i=1}^n \mathbb{E}(X_ig(W_n+1)) + q\sum_{i=1}^n \mathbb{E}(X_i\Delta g(W_n))$$

From (3.2), note that X_i and W_i are independent. Therefore, the above expression can be written as

$$q\mathbb{E}[\mathcal{A}g(W_n)] = \sum_{i=1}^{n} \mathbb{E}(X_i)\mathbb{E}(g(W_n+1) - g(W_i+1)) + q\sum_{i=1}^{n} \mathbb{E}(X_i\Delta g(W_n)) -\sum_{i=1}^{n} \mathbb{E}(X_i(g(W_n+1) - g(W_i+1))) = \sum_{i=1}^{n} \mathbb{E}(X_i)\mathbb{E}(g(W_i + X_{A_i} + 1) - g(W_i+1)) + q\sum_{i=1}^{n} \mathbb{E}(X_i\Delta g(W_n)) -\sum_{i=1}^{n} \mathbb{E}(X_i(g(W_i + X_{A_i} + 1) - g(W_i+1))) = \sum_{i=1}^{n} \mathbb{E}(X_i)\mathbb{E}\left(\sum_{j=1}^{X_{A_i}} \Delta g(W_i+j)\right) + q\sum_{i=1}^{n} \mathbb{E}(X_i\Delta g(W_n)) -\sum_{i=1}^{n} \mathbb{E}\left(X_i\sum_{j=1}^{X_{A_i}} \Delta g(W_i+j)\right).$$
(3.6)

Using Lemma 2.1, we get

$$\begin{aligned} |\mathbb{E}[\mathcal{A}g(W_n)]| &\leq \frac{2}{pq^n} \sum_{i=1}^n p_i \mathbb{E}\left(q^{W_i}(1-q^{X_{A_i}})\right) + \frac{2}{q^n} \sum_{i=1}^n \mathbb{E}\left(X_i q^{W_n}\right) \\ &+ \frac{2}{pq^n} \sum_{i=1}^n \mathbb{E}\left(X_i q^{W_i}\left(1-q^{X_{A_i}}\right)\right) \\ &= \frac{2}{pq^n} \sum_{i=1}^n \left[\mathbb{E}\left((X_i+p)q^{W_i}\right) - \mathbb{E}\left((p_i+qX_i)q^{W_n}\right)\right]. \end{aligned}$$

This proves the result.

Corollary 3.1. Let $B_{\alpha,p}$ and W_n be defined as in (1.2) and (3.1), respectively, and satisfy (3.3). Assume X_1, X_2, \ldots, X_n are independent random variables then

$$d_{sl}(W_n, B_{n,p}) \le \frac{2}{q^n} \sum_{i=1}^n |p - p_i| p_i \prod_{j \ne i} (1 - pp_j).$$
(3.7)

Proof. Substituting $A_i = \{i\}$ in (3.6), it can be easily verified that

$$q\mathbb{E}[\mathcal{A}g(W_n)] = \sum_{i=1}^n p_i^2 \mathbb{E}(\Delta g(W_i+1)) + q \sum_{i=1}^n p_i \mathbb{E}(\Delta g(W_i+1)) - \sum_{i=1}^n p_i \mathbb{E}(\Delta g(W_i+1)))$$
$$= \sum_{i=1}^n (p_i - p) p_i \mathbb{E}(\Delta g(W_i+1)).$$

Hence, using Lemma 2.1, we get

$$|\mathbb{E}[\mathcal{A}g(W_n)]| \le \frac{2}{q^n} \sum_{i=1}^n |p - p_i| p_i \mathbb{E}\left(q^{W_i}\right) = \frac{2}{q^n} \sum_{i=1}^n |p - p_i| p_i \prod_{j \neq i} (1 - pp_j).$$

This proves the result.

Remark 3.1. (i) Note that if $p_i = p, i = 1, 2, ..., n$, in (3.7) then $d_{sl}(W_n, B_{n,p}) = 0$, as expected.

- (ii) In Theorem 3.1 and Corollary 3.1, the bounds become shaper for sufficiently small values of p_i .
- (iii) If we choose the parameter p of our choice, then $\alpha = \frac{1}{p} \sum_{i=1}^{n} \mathbb{E}(X_i)$. In this situation, the parameter α may not be an integer. So, we can take

$$\alpha = \left\lfloor \frac{1}{p} \sum_{i=1}^{n} \mathbb{E}(X_i) \right\rfloor,\,$$

where $\lfloor x \rfloor$ is the integer part of x > 0, and use (2.8) and (3.12) to get

$$d_{sl}(W_n, \mathbf{B}_{\alpha, p}) \le \frac{2}{pq^{\alpha}} \sum_{i=1}^n \left[\mathbb{E}\left((X_i + p)q^{W_i} \right) - \mathbb{E}\left((p_i + qX_i - \delta p^2)q^{W_n} \right) \right].$$
(3.8)

Therefore, it is suggested to consider the minimum of the bounds given in (3.4) and (3.8).

If we choose $\alpha p = \mathbb{E}(W_n)$ and $\alpha pq = \operatorname{Var}(W_n)$ (the first two moments matching condition), then the choice of α may not be a positive integer. So, we choose

$$\alpha = \left\lfloor \frac{(\mathbb{E}(W_n))^2}{\mathbb{E}(W_n) - \operatorname{Var}(W_n)} \right\rfloor \quad \text{and} \quad p = \frac{\mathbb{E}(W_n) - \operatorname{Var}(W_n)}{\mathbb{E}(W_n)}.$$
(3.9)

Also, define $D(Z) := 2d_{TV}(Z, Z+1)$ and

$$\delta := \frac{(\mathbb{E}(W_n))^2}{\mathbb{E}(W_n) - \operatorname{Var}(W_n)} - \alpha.$$

Observe that $0 \le \delta < 1$,

$$\alpha p = \mathbb{E}(W_n) - \delta p \quad \text{and} \quad \alpha pq = \operatorname{Var}(W_n) - \delta pq$$
(3.10)

The next result gives the error in approximation between $B_{\alpha,p}$ and W_n satisfying the above conditions. **Theorem 3.2.** Let $B_{\alpha,p}$ and W_n be as defined in (1.2) and (3.1), respectively, and satisfy (3.9). Then

$$d_{sl}(W_{n}, \mathbf{B}_{\alpha, p}) \leq \frac{2}{p^{2}q^{\alpha}} \left\{ \sum_{i=1}^{n} \mathbb{E}(X_{i}) \mathbb{E}\left(\left(pX_{A_{i}} + q^{X_{B_{i}}} \left(1 - q^{-X_{A_{i}}} \right) \right) D(W_{i}^{*} | X_{A_{i}}, X_{B_{i}}) \right) \right. \\ \left. + \sum_{i=1}^{n} \mathbb{E}\left(X_{i} \left(pX_{A_{i}} + q^{X_{B_{i}}} \left(1 - q^{-X_{A_{i}}} \right) \right) D(W_{i}^{*} | X_{i}, X_{A_{i}}, X_{B_{i}}) \right) \right. \\ \left. + p \sum_{i=1}^{n} \mathbb{E}\left(X_{i} \left(q^{B_{i}} - q \right) D(W_{i}^{*} | X_{B_{i}}) \right) + \frac{\delta p^{3}}{q} \mathbb{E}(q^{W_{n}}) \right. \\ \left. + \frac{p}{q} \sum_{i=1}^{n} |\mathbb{E}(X_{i})\mathbb{E}(X_{A_{i}}) - \mathbb{E}(X_{i}X_{A_{i}}) + q\mathbb{E}(X_{i})|\mathbb{E}\left(\left(q^{B_{i}} - q \right) D(W_{i}^{*} | X_{B_{i}}) \right) \right\}.$$
(3.11)

Proof. From (3.5), we have

$$q\mathbb{E}[\mathcal{A}g(W_n)] = \alpha p\mathbb{E}(g(W_n+1)) - p\mathbb{E}(W_ng(W_n+1)) - q\mathbb{E}(W_ng(W_n)).$$

Using (3.10) and following the steps similar to the proof of Theorem 3.1, we get

$$q\mathbb{E}[\mathcal{A}g(W_n)] = \sum_{i=1}^n \mathbb{E}(X_i)\mathbb{E}\left(\sum_{j=1}^{X_{A_i}} \Delta g(W_i+j)\right) - \sum_{i=1}^n \mathbb{E}\left(X_i \sum_{j=1}^{X_{A_i}} \Delta g(W_i+j)\right) + q \sum_{i=1}^n \mathbb{E}(X_i \Delta g(W_n)) - \delta p\mathbb{E}(g(W_n+1)).$$
(3.12)

Using (3.2) and (3.10), the above expression leads to

$$q\mathbb{E}[\mathcal{A}g(W_n)] = \sum_{i=1}^n \mathbb{E}(X_i)\mathbb{E}\left(\sum_{j=1}^{X_{A_i}} (\Delta g(W_i+j) - \Delta g(W_i^*+1))\right)$$
$$-\sum_{i=1}^n \mathbb{E}\left(X_i \sum_{j=1}^{X_{A_i}} (\Delta g(W_i+j) - \Delta g(W_i^*+1))\right)$$
$$+ q \sum_{i=1}^n \mathbb{E}(X_i (\Delta g(W_n) - \Delta g(W_i^*+1))) - \delta p\mathbb{E}(g(W_n+1))$$

$$\begin{split} &-\sum_{i=1}^{n} [\mathbb{E}(X_{i})\mathbb{E}(X_{A_{i}}) - \mathbb{E}(X_{i}X_{A_{i}}) + q\mathbb{E}(X_{i})]\mathbb{E}(g(W_{n}+1) - g(W_{i}^{*}+1)) \\ &= \sum_{i=1}^{n} \mathbb{E}(X_{i})\mathbb{E}\left(\sum_{j=1}^{X_{A_{i}}}\sum_{\ell=1}^{X_{B_{i}\setminus A_{i}}+j-1} \Delta^{2}g(W_{i}^{*}+\ell)\right) \\ &-\sum_{i=1}^{n} \mathbb{E}\left(X_{i}\sum_{j=1}^{X_{B_{i}}}\sum_{\ell=1}^{X_{B_{i}\setminus A_{i}}+j-1} \Delta^{2}g(W_{i}^{*}+\ell)\right) \\ &+q\sum_{i=1}^{n} \mathbb{E}\left(X_{i}\sum_{\ell=1}^{X_{B_{i}}}\Delta^{2}g(W_{i}^{*}+\ell)\right) - \delta p\mathbb{E}(g(W_{n}+1)) \\ &-\sum_{i=1}^{n} [\mathbb{E}(X_{i})\mathbb{E}(X_{A_{i}}) - \mathbb{E}(X_{i}X_{A_{i}}) + q\mathbb{E}(X_{i})]\mathbb{E}\left(\sum_{\ell=1}^{X_{B_{i}}-1}\Delta^{2}g(W_{i}^{*}+\ell)\right) \\ &-\sum_{i=1}^{n} \mathbb{E}(X_{i})\mathbb{E}\left(\sum_{j=1}^{X_{A_{i}}}\sum_{\ell=1}^{X_{B_{i}\setminus A_{i}}+j-1} \mathbb{E}(\Delta^{2}g(W_{i}^{*}+\ell)|X_{A_{i}},X_{B_{i}})\right) \\ &-\sum_{i=1}^{n} \mathbb{E}\left(X_{i}\sum_{j=1}^{X_{A_{i}}}\sum_{\ell=1}^{X_{B_{i}\setminus A_{i}}+j-1} \mathbb{E}(\Delta^{2}g(W_{i}^{*}+\ell)|X_{i},X_{A_{i}},X_{B_{i}})\right) \\ &+q\sum_{i=1}^{n} \mathbb{E}\left(X_{i}\sum_{j=1}^{X_{B_{i}}-1}\mathbb{E}(\Delta^{2}g(W_{i}^{*}+\ell)|X_{B_{i}})\right) - \delta p\mathbb{E}(g(W_{n}+1)) \\ &-\sum_{i=1}^{n} [\mathbb{E}(X_{i})\mathbb{E}(X_{A_{i}}) - \mathbb{E}(X_{i}X_{A_{i}}) + q\mathbb{E}(X_{i})]\mathbb{E}\left(\sum_{\ell=1}^{X_{B_{i}}-1}\mathbb{E}(\Delta^{2}g(W_{i}^{*}+\ell)|X_{B_{i}})\right) \\ &-\sum_{i=1}^{n} [\mathbb{E}(X_{i})\mathbb{E}(X_{A_{i}}) - \mathbb{E}(X_{i}X_{A_{i}}) + q\mathbb{E}(X_{i})]\mathbb{E}\left(\sum_{\ell=1}^{X_{B_{i}}-1}\mathbb{E}(\Delta^{2}g(W_{i}^{*}+\ell)|X_{B_{i}})\right) \\ &+\sum_{i=1}^{n} \mathbb{E}\left(X_{i}\mathbb{E}(X_{A_{i}}) - \mathbb{E}(X_{i}X_{A_{i}}) + q\mathbb{E}(X_{i})]\mathbb{E}\left(\sum_{\ell=1}^{X_{B_{i}}-1}\mathbb{E}(\Delta^{2}g(W_{i}^{*}+\ell)|X_{B_{i}})\right) \\ &+\sum_{i=1}^{n} \mathbb{E}\left(X_{i}\mathbb{E}(X_{A_{i}}) - \mathbb{E}(X_{i}X_{A_{i}}) + q\mathbb{E}(X_{i})\mathbb{E}\left(\sum_{\ell=1}^{X_{B_{i}}-1}\mathbb{E}(\Delta^{2}g(W_{i}^{*}+\ell)|X_{B_{i}})\right) \\ &+\sum_{i=1}^{n} \mathbb{E}\left(X_{i}\mathbb{E}(X_{A_{i}}) - \mathbb{E}(X_{i}X_{A_{i}}) + q\mathbb{E}(X_{i})\mathbb{E}(X_{A_{i}}) + q\mathbb{E}(X_{i})\mathbb{E}(X_{A_{i}}) + q\mathbb{E}(X_{i})\mathbb{E}(X_{A_{i}}) + q\mathbb{E}(X_{i})\mathbb{E}\left(\sum_{\ell=1}^{N}\mathbb{E}(\Delta^{2}g(W_{i}^{*}+\ell)|X_{B_{i}})\right) \\ &+\sum_{i=1}^{n} \mathbb{E}\left(X_{i}\mathbb{E}(X_{A_{i}}) - \mathbb{E}(X_{i}X_{A_{i}}) + q\mathbb{E}(X_{i})\mathbb{E}\left(\sum_{\ell=1}^{N}\mathbb{E}(\Delta^{2}g(W_{i}^{*}+\ell)|X_{B_{i}})\right) \\ &+\sum_{i=1}^{n} \mathbb{E}\left(X_{i}\mathbb{E}(X_{A_{i}}) - \mathbb{E}\left(X_{i}X_{A_{i}}\right) + q\mathbb{E}\left(X_{i}\mathbb{E}\left(\sum_{\ell=1}^{N}\mathbb{E}\left(X_{i}X_{A_{i}}\right) + q\mathbb{E}\left(X_{i}X_{A_{i}}\right) + q\mathbb{E}\left(X_{i}X_{A_{i}}\right) + q\mathbb{E}\left(X_$$

Note that $\mathbb{E}(\Delta^2 g(W_i^* + \ell) | \cdot) \le 2q^{\ell-\alpha} D(W_i^* | \cdot)$. Hence, using (2.8), we get

$$\begin{aligned} |\mathbb{E}[\mathcal{A}g(W_{n})]| &\leq \frac{2}{p^{2}q^{\alpha}} \left\{ \sum_{i=1}^{n} \mathbb{E}(X_{i})\mathbb{E}\left(\left(pX_{A_{i}}+q^{X_{B_{i}}}\left(1-q^{-X_{A_{i}}}\right)\right)D(W_{i}^{*}|X_{A_{i}},X_{B_{i}})\right) \right. \\ &+ \sum_{i=1}^{n} \mathbb{E}\left(X_{i}\left(pX_{A_{i}}+q^{X_{B_{i}}}\left(1-q^{-X_{A_{i}}}\right)\right)D(W_{i}^{*}|X_{i},X_{A_{i}},X_{B_{i}})\right) \\ &+ p\sum_{i=1}^{n} \mathbb{E}\left(X_{i}\left(q^{B_{i}}-q\right)D(W_{i}^{*}|X_{B_{i}})\right) + \frac{\delta p^{3}}{q}\mathbb{E}(q^{W_{n}}) \\ &+ \frac{p}{q}\sum_{i=1}^{n} |\mathbb{E}(X_{i})\mathbb{E}(X_{A_{i}}) - \mathbb{E}(X_{i}X_{A_{i}}) + q\mathbb{E}(X_{i})|\mathbb{E}\left(\left(q^{B_{i}}-q\right)D(W_{i}^{*}|X_{B_{i}})\right)\right)\right\}.\end{aligned}$$

This proves the result.

Corollary 3.2. Let $B_{\alpha,p}$ and W_n be defined as in (1.2) and (3.9), respectively, and satisfy (3.2). If X_1, X_2, \ldots, X_n are independent random variables, then

$$d_{sl}(W_n, B_{\alpha, p}) \le \frac{2}{q^{\alpha}} \left\{ \sqrt{\frac{2}{\pi}} \left(\frac{1}{4} + \sum_{i=1}^n \gamma_i - \gamma^* \right)^{-1/2} \sum_{i=1}^n |p - p_i| p_i^2 + \delta p \prod_{i=1}^n (1 - pp_i) \right\}, \quad (3.14)$$

where $\gamma_j = \min\{\frac{1}{2}, 1 - \frac{1}{2}(q_j + |q_j - p_j|)\}$ and $\gamma^* = \max_{1 \le j \le n} \gamma_j$. *Proof.* Substituting $A_i = \{i\}$ in (3.12), it can be easily verified that

$$q\mathbb{E}[\mathcal{A}g(W_n)] = \sum_{i=1}^n (p_i - p)p_i\mathbb{E}(\Delta g(W_i + 1)) - \delta p\mathbb{E}(g(W_n + 1)).$$

Using (3.10), we get

$$q\mathbb{E}[\mathcal{A}g(W_n)] = -\sum_{i=1}^n (p-p_i)p_i\mathbb{E}(\Delta g(W_n+1) - \Delta g(W_i+1)) - \delta p\mathbb{E}(g(W_n+1)))$$

= $-\sum_{i=1}^n (p-p_i)p_i^2\mathbb{E}(\Delta^2 g(W_i+1)) - \delta p\mathbb{E}(g(W_n+1))$

Note that $|\mathbb{E}(\Delta^2 g(W_i + 1))| \leq 2\gamma/q^{\alpha-1}$, where $\gamma = 2 \max_{i \in J} d_{TV}(W_i, W_i + 1)$ (see Barbour and Xia [3], and Barbour and Čekanavičius [2, p. 517])). Also, from Corollary 1.6 of Mattner and Roos [17] (see also Remark 4.1 of Vellaisamy *et al.* [22]), we have

$$\gamma \le \sqrt{\frac{2}{\pi}} \left(\frac{1}{4} + \sum_{j=1}^{n} \gamma_j - \gamma^* \right)^{-1/2}.$$

where

$$\gamma_j = \min\left\{\frac{1}{2}, 1 - d_{TV}(X_j, X_j + 1)\right\}$$
$$= \min\left\{\frac{1}{2}, 1 - \frac{1}{2}(q_j + |q_j - p_j|)\right\}$$

and $\gamma^* = \max_{1 \le j \le n} \gamma_j$. Hence,

$$|\mathbb{E}[\mathcal{A}g(W_n)]| \le \frac{2}{q^{\alpha}} \left\{ \sqrt{\frac{2}{\pi}} \left(\frac{1}{4} + \sum_{i=1}^n \gamma_i - \gamma^* \right)^{-1/2} \sum_{i=1}^n |p - p_i| p_i^2 + \delta p \prod_{i=1}^n (1 - pp_i) \right\}.$$

This proves the result.

- **Remark 3.2.** (i) Note that W_i^* can be expressed as the conditional sum of independent random variables. Therefore, Subsections 5.3 and 5.4 of Röllin [19] and Remark 3.1(ii) of Kumar *et al.* [16] are useful to find the upper bound of $D(W_i^*|\cdot)$.
 - (ii) Observe that $|\Delta^2 g(X + \ell)| \le 4q^{X+\ell-\alpha}$. Therefore, from (3.13), we get

$$d_{sl}(W_{n}, \mathbf{B}_{\alpha, p}) \leq \frac{4}{p^{2}q^{\alpha}} \left\{ \sum_{i=1}^{n} \mathbb{E} \left((X_{i} + p_{i}) \left(pX_{A_{i}}q^{W_{i}^{*}} + q^{W_{n}} - q^{W_{i}} \right) \right) + p\sum_{i=1}^{n} \mathbb{E} \left(X_{i} \left(q^{W_{n}} - q^{W_{i}^{*}+1} \right) \right) + \frac{\delta p^{3}}{2} \mathbb{E} \left(q^{W_{n}} \right) + \frac{p}{q} \sum_{i=1}^{n} |\mathbb{E}(X_{i})\mathbb{E}(X_{A_{i}}) - \mathbb{E}(X_{i}X_{A_{i}}) + q\mathbb{E}(X_{i})|\mathbb{E} \left(q^{W_{n}} - q^{W_{i}^{*}+1} \right) \right\}. \quad (3.15)$$

Therefore, in practice, one could take the minimum of the bounds (3.11) and (3.15).

- (iii) Theorems 3.1 and 3.2 are established using Lemma 2.1 for all $z \ge 0$. Following the steps similar to the proofs of Theorems 3.1 and 3.2, the results can also be derived using Lemma 2.2 in terms of z > 1. This can be used to approximate $\mathbb{E}[(W_n z)^+]$ by $\mathbb{E}[(\mathbf{B}_{\alpha,p} z)^+]$ for z > 1.
- (iv) From Corollary 1 of Neammanee and Yonghint [18], we have

$$d_{sl}(\mathbf{P}_{\lambda}, W_n) \le (2e^{\lambda} - 1) \sum_{i=1}^n p_i^2,$$
(3.16)

where $\lambda = \sum_{i=1}^{n} p_i$. The bound given in (3.7) and (3.14) are better than the above bound, for example, let n = 100 and p_i , $1 \le i \le 100$, be defined as follows:

i	p_i	i	p_i	i	p_i	i	p_i	i	p_i
1-20	0.06	21-40	0.07	41-60	0.08	61-80	0.09	81-100	0.10

Table 1: The values of p_i

Then, the following table gives a comparison between our bounds (3.7) and (3.14), and the existing bound (3.16).

n	From (3.16) (existing bound)	From (3.7)	From (3.14)
10	0.095193	0	$7.6 imes 10^{-16}$
20	0.406097	0	6.8×10^{-15}
30	1.496990	0.109842	0.638717
40	4.407670	0.324195	1.188300
50	13.78920	1.186000	1.474570
60	39.44710	3.261280	1.676520
70	123.9500	12.78810	12.56050
80	370.6940	39.29820	13.90400
90	1227.670	136.3000	68.75740
100	3934.200	425.1760	335.1310

Table 2: Comparison of bounds.

For $1 \le n \le 20$, note that the bounds given in (3.7) are zero, as expected. Further, our bounds improve upon the existing bounds for various values of p_i . Also, for sufficiently large values of n, the bound given in (3.14) is better than the bound given in (3.7).

References

- [1] Barbour, A. D., Holst, L. and Janson, S. (1992). Poisson Approximation. Oxford University Press.
- [2] Barbour, A. D. and Čekanavičius, V. (2002). Total variation asymptotes for sum of independent integer random variables. *Ann. Prob.* **30**, 509–545.
- [3] Barbour, A. D. and Xia, A. (1999). Poisson perturbation. ESAIM Probab. Statist. 3, 131–150.
- [4] Boutsikas, M. V. and Vaggelatou, E. (2002). On the distance between convex-ordered random variables, with applications. *Adv. in Appl. Probab.* **34**, 349–374.
- [5] Brown, T. C. and Phillips, M. J. (1999). Negative binomial approximation with Stein's method. *Methodol. Comput. Appl. Probab.* 1, 407–421.
- [6] Brown, T. C. and Xia, A. (2001). Stein's method and birth-death processes. *Ann. Probab.* **29**, 1373–1403.
- [7] Čekanavičius, V. and Vellaisamy, P. (2015). Discrete approximations for sums of *m*-dependent random variables. *ALEA Lat. Am. J. Probab. Math. Stat.* **12**, 765–792.

- [8] Čekanavičius, V. and Vellaisamy, P. (2021). Compound Poisson approximations in ℓ_p -norm for sums of weakly dependent vectors. *J. Theoret. Probab.* **34**, 2241–2264.
- [9] Chen, L. H. Y. (1975). Poisson approximation for dependent trials. Ann. Probability 3, 534–545.
- [10] Eichelsbacher, P. and Reinert, G. (2008). Stein's method for discrete Gibbs measures. *Ann. Appl. Probab.* 18, 1588–1618.
- [11] El Karoui, N. and Jiao, Y. (2009). Stein's method and zero bias transformation for CDO tranche pricing. *Finance Stoch.* 13, 151–180.
- [12] El Karoui, N., Jiao, Y. and Kurtz, D. (2008). Gaussian and Poisson approximation: applications to CDOs tranche pricing. J. Comput. Finance 12, 31–58.
- [13] Hull, J. C. and White, A. D. (2004). Valuation of a CDO and an nth to default CDS without Monte Carlo simulation. J. Deriv. 12, 8–23.
- [14] Kumar, A. N. (2021). Approximations to weighted sums of random variables. Bull. Malays. Math. Sci. Soc. 44, 2447–2464.
- [15] Kumar, A. N. (2022). Bounds on negative binomial approximation to call function. To appear in *Revstat Stat. J.*
- [16] Kumar, A. N., Upadhye, N. S. and Vellaisamy, P. (2021). Approximations related to the sums of *m*-dependent random variables. To appear in *Braz. J. Probab. Stat.*
- [17] Mattner, L. and Roos, B. (2007). A shorter proof of Kanter's bessel function concentration bound. *Probab. Theory and Related Fields* 139, 191–205.
- [18] Neammanee, K. and Yonghint, N. (2020). Poisson approximation for call function via Stein-Chen method. *Bull. Malays. Math. Sci. Soc.* 43, 1135–1152.
- [19] Röllin, A. (2008). Symmetric and centered binomial approximation of sums of locally dependent random variables. *Electron. J. Probab.* **13**, 756–776.
- [20] Stein, C. (1972). A bound for the error in the normal approximation to the distribution of a sum of dependent random variables. In: *Proc. Sixth Berkeley Symp. Math. Statist. Probab. II. Probability Theory*, Univ. California Press, Berkeley, Calif., 583–602.
- [21] Vellaisamy, P. and Punnen, A. P. (2001). On the nature of the binomial distribution. J. Appl. *Probab.* **38**, 36–44.
- [22] Vellaisamy, P., Upadhye, N. S. and Čekanavičius, V. (2013). On negative binomial approximation. *Theory Probab. Appl.* 57, 97–109.
- [23] Yonghint, N., Neammanee, K. and Chaidee, N. (2022). Poisson approximation for locally dependent CDO. *Comm. Statist. Theory Methods* 51, 2073–2081.