

A Note on the Gessel Numbers

Jovan Mikić

University of Banja Luka

Faculty of Technology

Bosnia and Herzegovina

jovan.mikic@tf.unibl.org

Abstract

The Gessel number $P(n, r)$ represents the number of lattice paths in a plane with unit horizontal and vertical steps from $(0, 0)$ to $(n + r, n + r - 1)$ that never touch any of the points from the set $\{(x, x) \in \mathbb{Z}^2 : x \geq r\}$. In this paper, we use combinatorial arguments to derive a recurrence relation between $P(n, r)$ and $P(n - 1, r + 1)$. Also, we give a new proof for a well-known closed formula for $P(n, r)$. Moreover, a new combinatorial interpretation for the Gessel numbers is presented.

Keywords: Gessel numbers, Catalan numbers, central binomial coefficient, lattice paths.

2020 Mathematics Subject Classification: Primary 05A10; Secondary 11B65.

1 Introduction

Let n be a non-negative integer and let r be a positive integer. The Gessel number $P(n, r)$ counts all lattice paths [1, p. 191] in plane with $(1, 0)$ and $(0, 1)$ steps from $(0, 0)$ to $(n + r, n + r - 1)$ that never touch any of the points from the set $\{(x, x) \in \mathbb{Z}^2 : x \geq r\}$.

By using a combinatorial argument and an instance of the Pfaff-Saalschütz theorem, Gessel proved that (see [1, p. 191])

$$P(n, r) = \frac{r}{2(n+r)} \binom{2n}{n} \binom{2r}{r}. \quad (1)$$

Let $C_n = \frac{1}{n+1} \binom{2n}{n}$ denote the n th Catalan number. The Catalan numbers are well studied in the literature (e.g. see [3, 7]) and there are many counting problems in combinatorics whose solutions are given by the Catalan numbers. For example, the Catalan number C_n is the number of all paths in a plane from $(0, 0)$ to (n, n) with $(1, 0)$ and $(0, 1)$ steps such that they never rise above the line $y = x$ (see [3, Example 9.1], [6, Problem 158], and [4, Eq. (10.11)]).

By setting $r = 1$ in Equation (1), it follows that $P(n, 1) = C_n$.

Recently, it is shown [5, Cor. 4] that the sum $\sum_{k=0}^{2n} (-1)^k \binom{2n}{k}^m C_k C_{2n-k}$ is divisible by $\binom{2n}{n}$ for all non-negative integers n and for all positive integers m . The Gessel numbers play an interesting role in this proof [5, Eq. (68)].

It is known [1, p. 191] that, for a fixed positive integer r , the smallest positive integer K_r , such that $\frac{K_r}{n+r} \binom{2n}{n}$ is an integer for every n is $\frac{r}{2} \binom{2r}{r}$. Guo gave a generalization of the Gessel numbers [2, Eq. (1.10)].

In this paper, we study the Gessel numbers by establishing a close relationship between the Gessel and the Catalan numbers. After we establish such a relationship, we present a new combinatorial interpretation for the Gessel Numbers. This combinatorial interpretation is contained in our main result, Theorem 6.

2 Main results

Our first result expresses $P(n, r)$ in terms of the Catalan numbers.

Proposition 1. *Let n be a non-negative integer and let r be a positive integer. Then*

$$P(n, r) = \sum_{k=0}^{r-1} \binom{2k}{k} C_{n+r-k-1}. \quad (2)$$

Proof. Let k be a non-negative integer, and let p be a path from $(0, 0)$ to $(n+r, n+r-1)$, with $(1, 0)$ and $(0, 1)$ steps, that never touches any of the points from the set $\{(x, x) \in \mathbb{Z}^2 : x \geq r\}$. Assume that the last point point of intersection of p and the line $y = x$, when taking the path p from left to right, is the point (k, k) , where $0 \leq k \leq r-1$.

The number of “permitted” paths from $(0, 0)$ to (k, k) is $\binom{2k}{k}$, since every such path does not contain “forbidden” points. Every such path passes through the point $(k+1, k)$, as (k, k) is the last intersection point between this path and the line $y = x$. Note that the segment that connects points $(k+1, k)$ and $(n+r, n+r-1)$ is parallel with the line $y = x$.

The number of “permitted” paths from $(k+1, k)$ to $(n+r, n+r-1)$ is the same as the number of all paths from $(k+1, k)$ to $(n+r, n+r-1)$ with $(1, 0)$ and $(0, 1)$ steps that never rise above the line $y = x-1$. It follows that there are $C_{n+r-k-1}$ such paths.

Therefore, the number of all paths whose last intersection with the line $y = x$ is the point (k, k) is $\binom{2k}{k} C_{n+r-k-1}$. Since k takes values from 0 to $r-1$, we obtain Equation (2). \square

We use the previous proposition to obtain a recurrence relation between $P(n-1, r+1)$ and $P(n, r)$.

Theorem 2. *Let n and r be positive integers. Then*

$$P(n-1, r+1) - P(n, r) = \binom{2r}{r} C_{n-1}. \quad (3)$$

Proof. The number $P(n-1, r+1)$ counts all paths from $(0, 0)$ to $(n+r, n+r-1)$, with $(1, 0)$ and $(0, 1)$ steps, that never touch any of the points from the set $\{(x, x) \in \mathbb{Z}^2 : x \geq r+1\}$.

Therefore, the number $P(n-1, r+1) - P(n, r)$ counts all paths from $(0, 0)$ to $(n+r, n+r-1)$, with $(1, 0)$ and $(0, 1)$ steps, whose last point of intersection with the line $y = x$ is the point (r, r) .

The number of “permitted” paths from $(0, 0)$ to (r, r) is $\binom{2r}{r}$, since every such path does not contain “forbidden” points.

After (r, r) , every such path must pass through $(r+1, r)$. Note that the segment that connects points $(r+1, r)$ and $(n+r, n+r-1)$ is parallel with the line $y = x$. The number of “permitted” paths from $(r+1, r)$ to $(n+r, n+r-1)$ is the same as the number of all paths from $(r+1, r)$ to $(n+r, n+r-1)$, with $(1, 0)$ and $(0, 1)$ steps, that never rise above the line $y = x - 1$. By a well-known [4, Eq. (10.11)] property of the Catalan numbers, it follows that there are C_{n-1} such paths.

Therefore, the number of all paths whose last point of intersection with the line $y = x$ is the point (r, r) is $\binom{2r}{r}C_{n-1}$. \square

By using recurrence relation (3) and induction on n , we give a proof of Equation (1).

Let $S(n, r)$ denote $\frac{r}{2(n+r)} \binom{2n}{n} \binom{2r}{r}$. We will show that $P(n, r) = S(n, r)$ for all non-negative integers n and for all positive integers r . We use induction on n .

For $n = 0$, since the “final” point $(r, r-1)$ is below the first “forbidden” point (r, r) , the Gessel number $P(0, r)$ counts all paths in a plane from $(0, 0)$ to $(r, r-1)$ with $(1, 0)$ and $(0, 1)$ steps without any restrictions. Hence, $P(0, r) = \binom{2r-1}{r}$ or $P(0, r) = \frac{1}{2} \binom{2r}{r}$.

Therefore, it follows that $P(0, r) = S(0, r)$ for all positive integers r .

Let us assume that $P(n-1, r) = S(n-1, r)$ for some positive integer n and for all positive integers r .

We use a well-known [3, p. 26] recurrence relation for the central binomial coefficient:

$$\binom{2(r+1)}{r+1} = \frac{2(2r+1)}{r+1} \binom{2r}{r}. \quad (4)$$

Then we have that the following equalities hold:

$$\begin{aligned} P(n, r) &= P(n-1, r+1) - \binom{2r}{r} C_{n-1} && \text{(by Equation (3))} \\ &= S(n-1, r+1) - \binom{2r}{r} C_{n-1} && \text{(by the induction hypothesis)} \\ &= \frac{r+1}{2(n+r)} \binom{2(n-1)}{n-1} \binom{2(r+1)}{r+1} - \binom{2r}{r} C_{n-1} \\ &= \frac{r+1}{2(n+r)} \binom{2(n-1)}{n-1} \frac{2(2r+1)}{r+1} \binom{2r}{r} - \binom{2r}{r} C_{n-1} && \text{(by Equation (4))} \end{aligned}$$

$$\begin{aligned}
&= \frac{2r+1}{n+r} \binom{2(n-1)}{n-1} \binom{2r}{r} - \frac{1}{n} \binom{2r}{r} \binom{2(n-1)}{n-1} \\
&= \binom{2(n-1)}{n-1} \binom{2r}{r} \left(\frac{2r+1}{n+r} - \frac{1}{n} \right) \\
&= \binom{2(n-1)}{n-1} \binom{2r}{r} \frac{r(2n-1)}{n(n+r)} \\
&= \frac{r}{2(n+r)} \binom{2r}{r} \left(\frac{2(2n-1)}{n} \binom{2(n-1)}{n-1} \right) \\
&= \frac{r}{2(n+r)} \binom{2r}{r} \binom{2n}{n} \tag{by Equation (4)} \\
&= S(n, r).
\end{aligned}$$

This completes our proof by induction.

Definition 3. Let n be a non-negative integer and let r be a positive integer. The number $Q(n, r)$ counts all lattice paths in a plane with $(1, 0)$ and $(0, 1)$ steps from $(0, 0)$ to $(n+r, n+r-1)$ that never touch any of the points from the set $\{(x, x) \in \mathbb{Z}^2 : 1 \leq x \leq r\}$.

Proposition 4. Let n and r be positive integers. Then

$$Q(n, r) = \sum_{k=1}^n C_{r+k-1} \binom{2(n-k)}{n-k}. \tag{5}$$

Proof. Let k be a non-negative integer, and let p be a path from $(0, 0)$ to $(n+r, n+r-1)$, with $(1, 0)$ and $(0, 1)$ steps, that never touches any of the points from the set $\{(x, x) \in \mathbb{Z}^2 : 1 \leq x \leq r\}$. There are two cases to consider.

The first case: A path p intersects the line $y = x$ only at the point $(0, 0)$. In this case, a path p must begin with a $(1, 0)$ step. Note that the segment that connects the points $(1, 0)$ and $(n+r, n+r-1)$ is parallel with the line $y = x$. The number of “permitted” paths from $(1, 0)$ to $(n+r, n+r-1)$ is the same as the number of all paths from $(1, 0)$ to $(n+r, n+r-1)$, with $(1, 0)$ and $(0, 1)$ steps, that never rise above the line $y = x - 1$. It follows that there are C_{n+r-1} such paths.

The second case: A path p intersects the line $y = x$ in at least two points. Let $(r+k, r+k)$ be the first point of intersection between p and the line $y = x$ after the point $(0, 0)$. Here, $1 \leq k \leq n-1$. Note that, in this case, $n \geq 2$.

Let m be a positive integer. It is readily verified that the number of all paths in a plane from $(0, 0)$ to (m, m) , with $(1, 0)$ and $(0, 1)$ steps, that intersect the line $y = x$ only at points $(0, 0)$ and (m, m) is $2C_{m-1}$. Therefore, the number of “permitted” paths from $(0, 0)$ to $(r+k, r+k)$ is $2C_{r+k-1}$.

The number of “permitted” paths from $(r+k, r+k)$ to $(n+r, n+r-1)$ is the same as the number of all paths from $(r+k, r+k)$ to $(n+r, n+r-1)$ with $(1,0)$ and $(0,1)$ steps, since every such path does not contain “forbidden” points. It follows that there are $\frac{1}{2} \binom{2(n-k)}{n-k}$ such paths (see [4, Equation (10.3)]).

Therefore, the number of all paths that intersect the line $y=x$ at the point $(r+k, r+k)$ for the first time after the point $(0,0)$ is $C_{r+k-1} \binom{2(n-k)}{n-k}$. Since k can take values from 1 to $n-1$, it follows that there are $\sum_{k=1}^{n-1} C_{r+k-1} \binom{2(n-k)}{n-k}$ such paths.

Putting ever together, it holds:

$$\begin{aligned} Q(n, r) &= C_{n+r-1} + \sum_{k=1}^{n-1} C_{r+k-1} \binom{2(n-k)}{n-k} \\ &= \sum_{k=1}^n C_{r+k-1} \binom{2(n-k)}{n-k}. \end{aligned}$$

□

Remark 5. Note that, for $n=0$, the number $Q(0, r)$ is equal to C_{r-1} .

We now use Proposition 1 and Proposition 4 to prove our main result that gives us a new combinatorial interpretation for the Gessel numbers.

Theorem 6. *Let n and r be positive integers. Then*

$$P(n, r) = Q(r, n). \tag{6}$$

Proof. By setting $n := r$ and $r := n$ in Proposition 4, it follows that

$$Q(r, n) = \sum_{k=1}^r C_{n+k-1} \binom{2(r-k)}{r-k}. \tag{7}$$

By substituting t for $r-k$, it follows that Equation (7) becomes

$$Q(r, n) = \sum_{t=0}^{r-1} C_{n+r-t-1} \binom{2t}{t}. \tag{8}$$

By using Proposition 1 and Equation (8), it follows that $Q(r, n) = P(n, r)$. □

Theorem 6 gives a new combinatorial interpretation for Gessel numbers. By Theorem 6, the Gessel number $P(n, r)$ is the number of all lattice paths in a plane with $(1,0)$ and $(0,1)$ steps from $(0,0)$ to $(n+r, n+r-1)$ that never touch any of the points from the set $\{(x, x) \in \mathbb{Z}^2 : 1 \leq x \leq n\}$.

3 Concluding remarks

We end this paper with some formulas for the Gessel numbers.

Let n be a non-negative integer, and let r be a positive integer. By Equation (1), Theorem 6, and Remark 5, it follows that

$$Q(n, r) = \begin{cases} C_{r-1}, & \text{if } n = 0 \\ \frac{n}{2(n+r)} \binom{2n}{n} \binom{2r}{r}, & \text{if } n > 0. \end{cases} \quad (9)$$

Let n and r be positive integers. Then the following formulas are true:

$$\frac{1}{2} \binom{2n+2r}{n+r} - P(n, r) = \sum_{k=1}^n \binom{2(r+k-1)}{r+k-1} C_{n-k}, \quad (10)$$

$$\frac{1}{2} \binom{2n+2r}{n+r} - Q(n, r) = \sum_{l=1}^r \binom{2(n+r-l)}{n+r-l} C_{l-1}. \quad (11)$$

The left side of Equation (10) represents the number of all lattice paths in a plane with $(1, 0)$ and $(0, 1)$ steps from $(0, 0)$ to $(n+r, n+r-1)$ whose intersection with the set $\{(x, x) \in \mathbb{Z}^2 : r \leq x \leq n+r-1\}$ is non-empty. It is readily verified that there are $\binom{2(r+k-1)}{r+k-1} C_{n-k}$ lattice paths in a plane with $(1, 0)$ and $(0, 1)$ steps from $(0, 0)$ to $(n+r, n+r-1)$ whose last point of intersection with the set $\{(x, x) \in \mathbb{Z}^2 : r \leq x \leq n+r-1\}$ is the point $(r+k-1, r+k-1)$. Here, $1 \leq k \leq n$.

Similarly, the left side of Equation (11) represents the number of all lattice paths in a plane with $(1, 0)$ and $(0, 1)$ steps from $(0, 0)$ to $(n+r, n+r-1)$ whose intersection with the set $\{(x, x) \in \mathbb{Z}^2 : 1 \leq x \leq r\}$ is non-empty. It is readily verified that there are $\binom{2(n+r-l)}{n+r-l} C_{l-1}$ lattice paths in a plane with $(1, 0)$ and $(0, 1)$ steps from $(0, 0)$ to $(n+r, n+r-1)$ whose first point of intersection with the set $\{(x, x) \in \mathbb{Z}^2 : 1 \leq x \leq r\}$, after the point $(0, 0)$, is the point (l, l) . Here, $1 \leq l \leq r$.

Note that, by using Equations (10) and (11), one can give another proof of Theorem 6.

Remark 7. By using a combinatorial argument, Gessel proved [1, Equation (39)] the following formula:

$$\sum_{k=0}^n P(k, r) \binom{2n-2k}{n-k} = \frac{1}{2} \binom{2n+2r}{n+r}, \quad (12)$$

where n is a non-negative integer and r is a positive integer. It is known that Equation (12) uniquely determines the numbers $P(n, r)$. Gessel used Equation (12) in order to prove Equation (1).

Let n and r be positive integers. By using Equation (12) and Theorem 6, it can be proved that

$$\sum_{k=0}^{r-1} \binom{2k}{k} Q(n, r-k) = \frac{1}{2} \binom{2n+2r}{n+r} - \frac{1}{2} \binom{2n}{n} \binom{2r}{r}. \quad (13)$$

Note that, for positive integers n , Equation (13) uniquely determines the numbers $Q(n, r)$.

Acknowledgments

I want to thank professor Duško Bogdanić for valuable comments which helped to improve the article.

References

- [1] I. M. Gessel, Super ballot numbers, *J. Symbolic Comput.* **14** (1992), 179–194.
- [2] V. J. W. Guo, Proof of two divisibility properties of binomial coefficients conjectured by Z. W. Sun, *Electron. J. Combin.* **21** (2014), # P54.
- [3] T. Koshy, *Catalan numbers with applications*, Oxford University Press, 2009.
- [4] C. Krattenthaler, Lattice Path Enumeration, arxiv.org/pdf/1503.05930.pdf.
- [5] J. Mikić, On a certain sums divisible by the central binomial coefficient, *J. Integer Sequences* **23** (2020), [Article 20.1.6](#).
- [6] R. P. Stanley, *Bijjective proof problems*, 2009.
- [7] R. P. Stanley, *Catalan Numbers*, Cambridge University Press, Cambridge, 2015.