

THE SINGULAR SUPPORT OF SHEAVES IS γ -COISOTROPIC

STÉPHANE GUILLERMOU AND CLAUDE VITERBO

Monday 18th September, 2023, 00:45

CONTENTS

1. Introduction	2
2. Notation, comments and acknowledgements	3
3. Reminders on the singular support of sheaves	4
4. Quantization of Hamiltonian isotopies	6
5. Spectral invariants for sheaves and Lagrangians	8
5.1. Definition of spectral invariants	10
5.2. The case of Hamiltonian maps	12
5.3. Defining γ -coisotropic subsets	15
6. The γ_g -metric on sheaves	15
6.1. The γ_g -topology	17
6.2. Link with spectral invariants	20
6.3. γ_g -limits and colimits	23
7. Proof of Theorem 1.2	28
8. γ -coisotropic vs cone-coisotropic	29
8.1. A γ -coisotropic set is cone-coisotropic	29
8.2. A cone-coisotropic set that is not γ -coisotropic	34
9. Questions and comments	36
Appendix A. Persistence modules and Barcodes	39
Appendix B. Decomposition in the completion of $D_{lc}(\mathbb{R})$	41
B.1. Limits of constructible sheaves	41
B.2. Decomposition in $D_{lc}(\mathbb{R})$	47
Appendix C. Comparing different interleaving metrics on $D(N \times \mathbb{R})$.	50

SG: UMR 6629 du CNRS - Laboratoire de Mathématiques Jean LERAY, Nantes Université, France. Part of this paper was written as the author was a member of UMR 5582 du CNRS - Institut Fourier, Université Grenoble Alpes, France.

CV: Université Paris-Saclay, CNRS, Laboratoire de mathématiques d'Orsay, 91405, Orsay, France. Part of this paper was written as the author was a member of DMA, École Normale Supérieure, 45 Rue d'Ulm, 75230 Cedex 05, FRANCE.

Both authors acknowledge support from ANR MICROLOCAL (ANR-15-CE40-0007) and ANR COSY (ANR-21-CE40-0002). The first author also acknowledges support from the Centre Henri Lebesgue ANR-11-LABX-0020-01.

Appendix D. Local Floer cohomology	53
Appendix E. Duality for sheaf spectral invariants and connection with Floer's spectral theory	56
E.1. Not really a duality	56
E.2. Compact support	57
E.3. Duality	57
E.4. Case of Hamiltonian maps	58
E.5. Sheaf and Floer cohomology for Hamiltonian maps	58
References	62

1. INTRODUCTION

In [Vit92], a metric, denoted γ , was introduced on the set $\mathcal{L}(T^*N)$ of Lagrangians Hamiltonianly isotopic to the zero section in T^*N , where N is a compact manifold and on $\mathcal{D}\mathfrak{H}\mathfrak{a}\mathfrak{m}_c(T^*N)$ the group of Hamiltonian maps with compact support for $N = \mathbb{R}^n$ or T^n (and in [Vit06] it was extended to $\mathcal{D}\mathfrak{H}\mathfrak{a}\mathfrak{m}_c(T^*N)$ for general compact N). The metric was generalized by Schwartz and Oh to general symplectic manifolds (M, ω) using Floer cohomology for the Hamiltonian case (see [Sch00; Oh05]), and by [Lec08] for the Lagrangian case¹. The notion of γ -coisotropic sets in a symplectic manifold was defined in [Vit22]. A similar definition was due to Usher [Ush19] in the setting of Hofer distance under the name of “locally rigid”. Notice that this yields a weaker notion : γ -coisotropic implies locally rigid in the sense of Usher.

In the context of sheaves on manifolds, there is a notion of coisotropic sets in the sense of Kashiwara-Schapira (see [KS90], definition 6.5.1, p. 271 and Definition 8.2), that the authors call **involutivity**. To avoid any confusion with other notions, we shall use the term cone-coisotropic for their definition as it is defined using the contingent and paratingent cones (see [Bou32] and Section 8). With this notion, Kashiwara and Schapira proved

Theorem 1.1 ([KS90], theorem 6.5.4, p. 272). *Given a sheaf $\mathcal{F} \in D^b(N)$, its singular support $SS(\mathcal{F})$ is cone-coisotropic.*

The aim of this paper is to prove that

Theorem 1.2. *Given a sheaf $\mathcal{F} \in D^b(N)$, its singular support $SS(\mathcal{F})$ is γ -coisotropic.*

Theorem 1.2 implies Theorem 1.1, since we shall prove the following connection between the two notions

¹Assuming $[\omega]\pi_2(M, L) = 0$, $\mu_L\pi_2(M, L) = 0$, where μ_L is the Maslov class of L .

Proposition 1.3. *Let V be a closed subset in (M, ω) . If V is γ -coisotropic then it is cone-coisotropic.*

Remark 1.4. This is related to questions asked by Vichery in section 4 of [Vic13]. Our result, together with Proposition 1.3, stating that a γ -coisotropic set is cone-coisotropic, gives a more natural proof of the Kashiwara-Schapira theorem. Moreover, as opposed to the notion of cone-coisotropic, which is only invariant by C^1 symplectic diffeomorphisms, the notion of γ -coisotropic set is invariant by homeomorphisms preserving γ , in particular by symplectic homeomorphisms, i.e. homeomorphisms which are C^0 limits of symplectic diffeomorphisms. Along the way we prove a number of results relating the singular support and the spectral norm γ .

2. NOTATION, COMMENTS AND ACKNOWLEDGEMENTS

Since our results are local, we shall assume from now on that N is a compact manifold. The following definitions will turn out useful :

- The category $D(N)$ is the unbounded derived category of sheaves of k -vector spaces on N , for some given field k , but except in Appendix B the reader may assume we are dealing with its bounded version. For the operations on sheaves associated to a continuous map $f : M \rightarrow N$ that is $f^{-1}, f_*, f^!, f_!$ and the two operations $\otimes, \mathcal{H}om$ and their derived versions, $f^{-1}, Rf_*, Rf_!, f^!$ and $\otimes, R\mathcal{H}om$ on $D(N)$ we refer to [KS90].
- Choosing a real analytic structure on N we set $D_c(N)$ to be the category of constructible complexes \mathcal{F} , i.e. complexes such that, for some subanalytic stratification, the restriction of \mathcal{F} to each stratum is locally constant and of finite rank.
- For $\mathcal{F} \in D(N)$, $SS(\mathcal{F})$ is the singular support (also called micro-support) defined by Kashiwara-Schapira in [KS90]. We set for short $SS^*(\mathcal{F}) = SS(\mathcal{F}) \cap (T^*N \setminus 0_N)$.
- For an open set U in the symplectic manifold (M, ω) we denote by $\mathcal{D}\mathcal{H}am_c(U)$ the set of time one flows of Hamiltonian with compact support contained in U .

We thank Pierre Schapira for useful conversations and Nicolas Vichery for Remark 6.3. While writing this paper, we realized that some of the results in Subsection 6.3 were independently discovered by Asano and Ike in [AI22]. We thank Yuichi Ike for a careful reading of the manuscript.

3. REMINDERS ON THE SINGULAR SUPPORT OF SHEAVES

Let N be a manifold, k a field and $D(N)$ the derived category of sheaves of k -vector spaces on N . For $\mathcal{F} \in D(N)$ we recall that $SS(\mathcal{F}) \subset T^*N$ is the closed conic subset defined by Kashiwara-Schapira as the closure of the set of points $(x; \xi)$ such that $(R\Gamma_{\{f \geq 0\}} \mathcal{F})_x \neq 0$ for some function f of class C^1 such that $f(x) = 0$ and $df_x = \xi$. It is called microsupport or singular support of \mathcal{F} .

The functor $R\Gamma_{\{f \geq 0\}}$ does not commute with infinite direct sums and $(-)_x$ does not commute with infinite direct products. Here are two variations on the definition which have these commutation properties. For the statement of the lemmas we introduce the following definition.

Definition 3.1. *Let $\Omega \subset T^*N \setminus 0_N$ be an open conic subset. We call Ω -lens a locally closed subset Σ of N with the following properties: $\bar{\Sigma}$ is compact and there exists an open neighbourhood U of $\bar{\Sigma}$ and a function $g: U \times [0, 1] \rightarrow \mathbb{R}$ of class C^1 such that*

- (1) $dg_t(x) \in \Omega$ for all $(x, t) \in U \times [0, 1]$, where $g_t = g|_{U \times \{t\}}$,
- (2) $\{g_t < 0\} \subset \{g_{t'} < 0\}$ if $t \leq t'$,
- (3) the hypersurfaces $\{g_t = 0\}$ coincide on $U \setminus \bar{\Sigma}$,
- (4) $\Sigma = \{g_1 < 0\} \setminus \{g_0 < 0\}$.

We recall that $SS^*(k_{\{g_t < 0\}}) = \{(x; \lambda dg_t(x)); g_t(x) = 0, \lambda > 0\}$. Using the triangle $k_{\{g_0 < 0\}} \rightarrow k_{\{g_1 < 0\}} \rightarrow k_\Sigma \xrightarrow{+1}$ and the ‘‘triangle inequality’’ for the singular support (see [KS90]) we obtain $SS^*(k_\Sigma) \subset \Omega$. The next lemma is




FIGURE 1. A lens. The arrows represent ∇g_1 and ∇g_0 .

essentially a reformulation of the definition of the singular support.

Lemma 3.2. *Let $\mathcal{F} \in D(N)$ and let $\Omega \subset T^*N \setminus 0_N$ be an open conic subset. Then $SS(\mathcal{F}) \cap \Omega = \emptyset$ if and only if $\text{RHom}(k_\Sigma, \mathcal{F}) \simeq 0$ for any Ω -lens Σ .*

Proof. (i) We first assume $SS(\mathcal{F}) \cap \Omega = \emptyset$. We recall that $\mathrm{RHom}(k_V, \mathcal{F}) \simeq \mathrm{R}\Gamma(V; \mathcal{F})$ for any open subset V of N . Using the notations in the definition of Ω -lens we have a distinguished triangle $k_{\{g_0 < 0\}} \rightarrow k_{\{g_1 < 0\}} \rightarrow k_\Sigma \xrightarrow{+1}$. Applying $\mathrm{RHom}(-, \mathcal{F}|_U)$ to this triangle we obtain the result, using the non-characteristic deformation lemma ([KS90, Prop. 2.7.2]) and the definition of the singular support.

(ii) We assume that there exists $(x_0; \xi_0) \in SS(\mathcal{F}) \cap \Omega$. Hence we can find $(x_1; \xi_1) \in \Omega$ (close to $(x_0; \xi_0)$) and a function $f: N \rightarrow \mathbb{R}$ of class C^1 such that $f(x_1) = 0$, $df(x_1) = \xi_1$ and $(\mathrm{R}\Gamma_{\{f \geq 0\}} \mathcal{F})_{x_1} \neq 0$. Now we can find a basis of open neighbourhoods of x_1 , say U_n , $n \in \mathbb{N}$, such that $\Sigma_n := \{f \geq 0\} \cap U_n$ is an Ω -lens. Then there exists n such that $\mathrm{R}\Gamma(U_n; \mathrm{R}\Gamma_{\{f \geq 0\}} \mathcal{F}) \neq 0$ and the result follows from

$$\begin{aligned} \mathrm{R}\Gamma(U_n; \mathrm{R}\Gamma_{\{f \geq 0\}} \mathcal{F}) &\simeq \mathrm{R}\Gamma(N; \mathrm{R}\mathcal{H}\mathrm{om}(k_{U_n}, \mathrm{R}\mathcal{H}\mathrm{om}(k_{\{f \geq 0\}}, \mathcal{F}))) \\ &\simeq \mathrm{R}\Gamma(N; \mathrm{R}\mathcal{H}\mathrm{om}(k_{\Sigma_n}, \mathcal{F})) \simeq \mathrm{RHom}(k_{\Sigma_n}, \mathcal{F}) \end{aligned}$$

□

The functor $\mathrm{RHom}(k_\Sigma, -)$ commutes with direct products but not with direct sums. Here is a dual version of the previous lemma using the functor $\mathrm{R}\Gamma(N; k_\Sigma \otimes -)$ which commutes with direct sums.

Lemma 3.3. *Let $\mathcal{F} \in D(N)$ and let $\Omega \subset T^*N \setminus 0_N$ be an open conic subset. Then $SS(\mathcal{F}) \cap \Omega = \emptyset$ if and only if $\mathrm{R}\Gamma(N; k_\Sigma \otimes \mathcal{F}) \simeq 0$ for any Ω^a -lens Σ , where Ω^a is the antipodal set of Ω , that is, its image by $(x; \xi) \mapsto (x; -\xi)$.*

The first author thanks Pierre Schapira for a suggestion simplifying the next proof.

Proof. Let Σ be an Ω^a -lens. We write it as the difference of the two closed sets $\bar{\Sigma}$ and $Z = \bar{\Sigma} \setminus \Sigma$. We can find two families of open neighbourhoods of $\bar{\Sigma}$, say $\{U_n\}_{n \in \mathbb{N}}$, and Z , say $\{V_n\}_{n \in \mathbb{N}}$, such that, for all n , we have: $V_n \subset U_n$ and $U_n \setminus V_n$ is an Ω -lens. For each n we have the commutative diagram of restriction maps

$$\begin{array}{ccc} \mathrm{R}\Gamma(U_n; \mathcal{F}) & \xrightarrow{u_n} & \mathrm{R}\Gamma(V_n; \mathcal{F}) \\ \downarrow & & \downarrow \\ \mathrm{R}\Gamma(N; \mathcal{F} \otimes k_{\bar{\Sigma}}) & \xrightarrow{u} & \mathrm{R}\Gamma(N; \mathcal{F} \otimes k_Z) \end{array}$$

The cone of u is $\mathrm{R}\Gamma(N; k_\Sigma \otimes \mathcal{F})$, so we want to prove that u is an isomorphism, that is, u induces an isomorphism on all cohomology groups. The cone of u_n is $\mathrm{RHom}(k_{U_n \setminus V_n}, \mathcal{F})$ and the previous lemma says that u_n is an isomorphism. Now the result follows from $H^i(N; \mathcal{F} \otimes k_{\bar{\Sigma}}) \simeq$

$\varinjlim_n H^i(U_n; \mathcal{F})$ and $H^i(N; \mathcal{F} \otimes k_Z) \simeq \varinjlim_n H^i(V_n; F)$, for all i (see [KS90, Rem. 2.6.6]). \square

The previous two lemmas give the following proposition (see [KS90, Ex. 5.7]), away from the zero-section. To deal with the zero-section we use $SS(\mathcal{F}) \cap 0_N = \text{supp}(\mathcal{F})$.

Proposition 3.4. *Let $\mathcal{F}_n \in D(N)$, $n \in \mathbb{N}$, be given. Then $SS(\bigoplus_n \mathcal{F}_n) \subset \overline{\bigcup_n SS(\mathcal{F}_n)}$ and $SS(\prod_n \mathcal{F}_n) \subset \overline{\bigcup_n SS(\mathcal{F}_n)}$.*

4. QUANTIZATION OF HAMILTONIAN ISOTOPIES

Everything in the next subsection is from [GKS] or [KS90]. Given φ , a homogeneous Hamiltonian map from $T^*N \setminus 0_N$ to $T^*N \setminus 0_N$, Guillermou, Kashiwara and Schapira associate to φ a Kernel, that is an element of $D^b(N \times N)$, noted K_φ having the following properties². First K_φ is a *quantization* of the homogeneous Lagrangian

$$\Lambda_\varphi = \{(z, \varphi(z)) \mid z \in T^*N \setminus 0_N\} \subset \overline{T^*N} \times T^*N$$

in the sense that $SS^\bullet(K_\varphi) = \Lambda_\varphi$, where we defined, for a sheaf \mathcal{F} on a manifold N ,

$$SS^\bullet(\mathcal{F}) = SS(\mathcal{F}) \cap (T^*N \setminus 0_N)$$

Before we state the second property we recall the *composition* of sheaves:

Definition 4.1. *Let $\mathcal{F} \in D^b(M)$ and $K \in D^b(M \times N)$. We define*

$$K(\mathcal{F}) = \mathcal{F} \circ K = Rq_{N!}(K \otimes q_M^{-1}\mathcal{F}) \in D^b(N),$$

where q_M, q_N are the projections of $M \times N$ on M and N respectively.

The second property is then :

Proposition 4.2 (See Proposition 7.1.2 (ii) in [KS90] and [GKS], formula (1.12)). *We have*

$$SS^\bullet(K_\varphi(\mathcal{F})) = \varphi(SS^\bullet(\mathcal{F}))$$

Now if φ is an exact non-homogeneous symplectic map from T^*M to T^*N (i.e. $PdQ - pdq$ is exact), we replace Λ_φ by the homogeneous lift of the graph of φ , say $\widehat{\Lambda}_\varphi \subset \overline{T^*M} \times T^*N \times T^*\mathbb{R}$, given by

$$\widehat{\Lambda}_\varphi = \{(q, -\tau p, Q(q, p), \tau P(q, p), F(q, p), \tau) \mid \varphi(q, p) = (Q, P), \tau > 0\}$$

²In fact in [GKS] the authors associate to a Hamiltonian isotopy φ^s a Kernel $\mathcal{K}_\Phi \in D^b(N \times N \times [0, 1])$ such that $\mathcal{K}_{\Phi|_{N \times N \times \{s\}}} = K_{\varphi^s}$. Note that K_{φ^1} depends on Φ not only φ^1 and our notation is slightly abusive.

where $dF = PdQ - pdq$. Note that considering $\widehat{\Lambda}_\varphi$ as a correspondence in $\overline{T^*M} \times T^*N \times T^*\mathbb{R}$, if L is an exact Lagrangian in T^*M and

$$\widehat{L} = \{(q, \tau p, f_L(q, p), \tau) \mid (q, p) \in L, \tau > 0\}$$

we have $\widehat{\Lambda}_\varphi \circ \widehat{L} = \widehat{\varphi(L)}$.

From now on we assume $M = N$ and φ is a Hamiltonian map with compact support (which will always be the case in this paper), the existence of a quantization \mathcal{K}_φ for $\widehat{\Lambda}_\varphi$ follows from the homogeneous case by considering $\widehat{\Lambda}_\varphi$ as a deformation of $\widehat{\Lambda}_{\text{id}}$. However we need to take care of two points. The first one is that the natural map which realizes this deformation, $\widehat{\Phi} = \text{id}_{T^*N} \times \widehat{\varphi}$, given by

$$\widehat{\Phi}(q, p, q', p', t, \tau) = (q, p, Q(q', p'/\tau), \tau P(q', p'/\tau), t + F(q', p'/\tau), \tau)$$

is not defined everywhere outside the zero section (for $p' \neq 0$ and $\tau = 0$). The second point is that it is not enough to deform $\widehat{\Lambda}_{\text{id}}$ because the singular support of $k_{\Delta_N \times [0, +\infty[}$ is slightly bigger than desired:

$$SS^\bullet(k_{\Delta_N \times [0, +\infty[}) = \widehat{\Lambda}_{\text{id}} \cup \{(q, p, q, -p, t, 0) \mid t \geq 0\}$$

(only its intersection with $\{\tau > 0\}$ is equal to $\widehat{\Lambda}_{\text{id}}$). Taking these points into account, we can find a homogeneous map ψ acting on $T^*(N^2 \times \mathbb{R}) \setminus \mathbf{0}_{N^2 \times \mathbb{R}}$ such that ψ is the identity map near $\{\tau = 0\}$ and $\psi(\widehat{\Lambda}_{\text{id}}) = \widehat{\Lambda}_\varphi$. We set

$$\mathcal{K}_\varphi = K_\psi(k_{\Delta_N \times [0, +\infty[})$$

and we find

$$SS^\bullet(\mathcal{K}_\varphi) = \widehat{\Lambda}_\varphi \cup \{(q, p, q, -p, t, 0) \mid t \geq 0\}$$

If we consider the whole isotopy φ_s and choose t_0 big enough so that $\widehat{\Lambda}_{\varphi_s}$ is contained in $T^*(N^2 \times]-t_0, t_0[)$ for all $s \in [0, 1]$, then we see that $SS^\bullet(\mathcal{K}_{\varphi_s}) \cap T^*U$ is independent of s , where $U = N^2 \times (\mathbb{R} \setminus [-t_0, t_0])$. By [KS90, Prop. 5.4.5] we deduce that

$$\mathcal{K}_\varphi|_{N^2 \times]-\infty, -t_0[} \simeq \mathbf{0}, \quad \mathcal{K}_\varphi|_{N^2 \times]t_0, +\infty[} \simeq k_{\Delta_N \times [0, +\infty[}|_{N^2 \times]t_0, +\infty[}$$

In other words $\mathcal{K}_\varphi = \mathcal{K}_{\text{id}} = k_{\Delta_N \times [0, +\infty[}$ outside of a compact set.

Remark 4.3. We could obtain the existence of \mathcal{K}_φ when $M = N$ and φ has compact support without assuming that φ is Hamiltonian by adapting the result of [Gui12; Vit19] (see Theorem 5.2). In *loc. cit.* the existence of a quantization is proved for a Legendrian of J^1N which is the lift of a compact exact Lagrangian of T^*N . Here the graph of φ is not compact but coincides with the diagonal $T_{\Delta_N}^*N^2$ outside a compact set (if we choose $F = 0$ at infinity) so it should not be too difficult to extend the result in our situation.

Since we use \mathcal{K}_φ , defined over $N^2 \times \mathbb{R}$, instead of sheaf defined over $(N \times \mathbb{R})^2$, we change slightly the composition functor as follows, which becomes a mixture of \circ and the convolution $*$ (see Section 5 for the definition)

Definition 4.4. *Let $\mathcal{F} \in D^b(M \times \mathbb{R})$ and $\mathcal{K} \in D^b(M \times N \times \mathbb{R})$. We define*

$$\mathcal{K}^\circledast(\mathcal{F}) = \mathcal{F} \circledast \mathcal{K} = R s_! R q_{2!} (q_{1,2}^{-1} \mathcal{K} \otimes q_1^{-1} \mathcal{F}) \in D^b(N \times \mathbb{R}),$$

where the maps

$$\begin{aligned} q_{1,2}: M \times N \times \mathbb{R}^2 &\rightarrow M \times N \times \mathbb{R}, & q_1: M \times N \times \mathbb{R}^2 &\rightarrow M \times \mathbb{R}, \\ q_2: M \times N \times \mathbb{R}^2 &\rightarrow N \times \mathbb{R}^2, & s: N \times \mathbb{R}^2 &\rightarrow N \times \mathbb{R} \end{aligned}$$

are given by $q_{1,2}(x, y, t_1, t_2) = (x, y, t_1)$, $q_1(x, y, t_1, t_2) = (x, t_2)$, $q_2(x, y, t_1, t_2) = (y, t_1, t_2)$, $s(y, t_1, t_2) = (y, t_1 + t_2)$.

Now we obtain a non homogeneous version of Proposition 4.2.

Proposition 4.5. *For a compactly supported Hamiltonian map φ from T^*N to itself and any $\mathcal{F} \in D^b(N \times \mathbb{R})$ with $SS(\mathcal{F}) \subset \{\tau \geq 0\}$, we have $SS^\bullet(\mathcal{K}_\varphi^\circledast(\mathcal{F})) = \widehat{\varphi}(SS^\bullet(\mathcal{F}))$.*

In particular, for $L \in \mathcal{L}(T^*N)$ we have $SS^\bullet(\mathcal{K}_\varphi(\mathcal{F}_L)) = SS^\bullet(\mathcal{F}_{\varphi(L)})$ and, by uniqueness of the quantization (see Theorem 5.2, (4)), $\mathcal{K}_\varphi^\circledast(\mathcal{F}_L) \simeq \mathcal{F}_{\varphi(L)}$.

5. SPECTRAL INVARIANTS FOR SHEAVES AND LAGRANGIANS

For M a symplectic manifold, (here we only need $M = T^*N$), if $\Lambda(M)$ is the bundle of Lagrangian subspaces of the tangent bundle to M , with fibre the Lagrangian Grassmannian $\Lambda(T_z M) \simeq \Lambda(n)$, we denote by $\widetilde{\Lambda}(M)$ the bundle induced by the universal cover $\widetilde{\Lambda}(n) \rightarrow \Lambda(n)$.

When using coefficients others than $\mathbb{Z}/2\mathbb{Z}$ we assume we have a lifting \widetilde{G}_L of the Gauss map $G_L: L \rightarrow \Lambda(T^*N)$ given by $x \mapsto T_x L$ to $\widetilde{\Lambda}(T^*N)$. This is called a grading. Given a graded L , the canonical deck transformation of the covering induces a new grading and we denote it as $L[1]$ (or $\widetilde{L}[1]$), and its k -th iteration as $L[k]$ (or $\widetilde{L}[k]$). The grading yields an absolute grading for the Floer homology³ and hence for the complex of sheaves in the Theorem stated below. We refer to [Sei00] and [Vit22] for more details on this, but point out that we shall never mention explicitly the grading.

³Without this extra piece of information, the Floer cohomology $FH^k(L_1, L_2)$ is generally defined only up to a shift in grading. However, when $L_2 = \varphi_H^1(L_1)$, there is an absolute grading, *a priori* depending on the choice of H . However in our situation we do not assume L_1 is Hamiltonianly isotopic to L_2 , so the grading is required to have an absolute grading of the Floer cohomology.

An exact Lagrangian in T^*N is a pair (L, f_L) such that $df_L = \lambda|_L$ where $\lambda = pdq$ is the Liouville form. For an exact Lagrangian in T^*N , a grading always exists since the obstruction to its existence is given by the Maslov class, and for exact Lagrangians in T^*N the Maslov class vanishes, as was proved by Kragh and Abouzaid (see [Kra13] and also the sheaf-theoretic proof by [Gui12]). When f_L is implicit we only write L , for example 0_N means $(0_N, 0)$. A Lagrangian brane is a triple $\tilde{L} = (L, f_L, \tilde{G}_L)$, where L is a compact Lagrangian. For c a real constant, we write $T_c\tilde{L} = (L, f_L + c, \tilde{G}_L)$. We also use the notation $\tilde{L} + c$ or $L + c$ if the grading is irrelevant (as it will be most of the time).

Definition 5.1. *We set $\mathcal{L}(T^*N)$ to be the set of Lagrangian branes in T^*N .*

Let L be a Lagrangian brane in T^*N and

$$\hat{L} = \{(q, \tau p, f_L(q, p), \tau) \mid (q, p) \in L, \tau > 0\}$$

the homogenized Lagrangian in $T^*(N \times \mathbb{R})$.

Similarly for Hamiltonian maps in an exact manifold $(M, \omega = d\lambda)$ we look for pairs (φ, F) such that $\varphi^*\lambda - \lambda = dF$ on M .

To state the next result we recall the two adjoint functors, $*$ and $\mathcal{H}om^*$, on the category $D(N \times \mathbb{R})$ introduced by Tamarkin in [Tam08]. First we set $s, q_1, q_2 : N \times (\mathbb{R})^2 \rightarrow N \times \mathbb{R}$ given by $s(x, t_1, t_2) = (x, t_1 + t_2)$, $q_1(x, t_1, t_2) = (x, t_1)$, $q_2(x, t_1, t_2) = (x, t_2)$ and we define

$$\mathcal{F} * \mathcal{G} = R s_! (q_1^{-1} \mathcal{F} \otimes q_2^{-1} \mathcal{G}),$$

$$\mathcal{H}om^*(\mathcal{F}, \mathcal{G}) = R q_{1*} R \mathcal{H}om(q_2^{-1} \mathcal{F}, s^! \mathcal{G})$$

Then $\mathcal{H}om^*$ is the adjoint of $*$ in the sense that

$$\text{Mor}_{D(N \times \mathbb{R})}(\mathcal{F}, \mathcal{H}om^*(\mathcal{G}, \mathcal{H})) = \text{Mor}_{D(N \times \mathbb{R})}(\mathcal{F} * \mathcal{G}, \mathcal{H})$$

We recall some properties of $\mathcal{H}om^*$. We denote by $D_{\tau \geq 0}(N \times \mathbb{R})$ the subcategory of $D(N \times \mathbb{R})$ of objects with singular support contained⁴ in $\{\tau \geq 0\}$. A base change formula gives, for any $c \in \mathbb{R}$, and $\mathcal{F}, \mathcal{G} \in D^b(N \times \mathbb{R})$

$$(5.1) \quad \text{R}\Gamma_{N \times \{-c\}}(N \times \mathbb{R}; \mathcal{H}om^*(\mathcal{F}, \mathcal{G})) \simeq \text{RHom}(\mathcal{F}, T_{c*}(\mathcal{G})),$$

with $T_c : N \times \mathbb{R} \rightarrow N \times \mathbb{R}$, $(x, t) \mapsto (x, t + c)$. Note that $\widehat{T_c \tilde{L}} = T_c \hat{L}$.

For \mathcal{H} with singular support in $\{\tau \geq 0\}$, we have a morphism

$$\text{R}\Gamma_{N \times \{b\}}(N \times \mathbb{R}; \mathcal{H}) \rightarrow \text{R}\Gamma_{N \times [a, b]}(N \times \mathbb{R}; \mathcal{H}) \xleftarrow{\sim} \text{R}\Gamma_{N \times \{a\}}(N \times \mathbb{R}; \mathcal{H})$$

for any $a \leq b$. We remark that the cone of this morphism is $\text{R}\Gamma_{N \times [a, b]}(N \times \mathbb{R}; \mathcal{H})$. Using (5.1) we deduce a morphism, say u , from $\text{RHom}(\mathcal{F}, T_{c*}(\mathcal{G}))$

⁴This category contains the ‘‘Tamarkin category’’ which is the left orthogonal of $D_{\tau \leq 0}(N \times \mathbb{R})$. The sheaves \mathcal{F}_L belongs to the Tamarkin category. The category $D_{\tau \geq 0}(N \times \mathbb{R})$ is stable by $*$ and $\mathcal{H}om^*$. The Tamarkin category is stable by $*$ but not $\mathcal{H}om^*$: typically when N is a point we have $\mathcal{H}om^*(k_{[a, \infty[}, k_{[b, \infty[}) \simeq k_{]-\infty, b-a[}[1]$.

to $\mathrm{RHom}(\mathcal{F}, T_{d*}(\mathcal{G}))$ for any $c \leq d$. Taking $\mathcal{F} = \mathcal{G}$, $c = 0$ the image of $\mathrm{id}_{\mathcal{G}}$ gives

$$(5.2) \quad \tau_{0,d}: \mathcal{G} \rightarrow T_{d*}(\mathcal{G}) \quad \text{for any } d \geq 0$$

(we come back to this in §6). Then u is nothing but the composition with $T_{c*}(\tau_{0,d-c})$.

According to [Gui12] and [Vit19] we have

Theorem 5.2. *To each $L \in \mathcal{L}(T^*N)$ we can associate $\mathcal{F}_L \in D^b(N \times \mathbb{R})$ such that*

- (1) $SS^\bullet(\mathcal{F}_L) = \widehat{L}$.
- (2) \mathcal{F}_L is simple (cf. [KS90, Def. 7.5.4]), $\mathcal{F}_L = 0$ near $N \times \{-\infty\}$ and $\mathcal{F}_L = k_{N \times \mathbb{R}}$ near $N \times \{+\infty\}$.
- (3) We have an isomorphism

$$FH^\bullet(L_0, L_1; a, b) = H_{N \times [a, b[}^*(N \times \mathbb{R}; \mathcal{H}om^*(\mathcal{F}_{L_0}, \mathcal{F}_{L_1}))$$

- (4) \mathcal{F}_L is unique satisfying properties (1) and (2).
- (5) There is a natural product map

$$\mathcal{H}om^*(\mathcal{F}_{L_1}, \mathcal{F}_{L_2}) \otimes \mathcal{H}om^*(\mathcal{F}_{L_2}, \mathcal{F}_{L_3}) \longrightarrow \mathcal{H}om^*(\mathcal{F}_{L_1}, \mathcal{F}_{L_3})$$

inducing in cohomology a map

$$H_{N \times [\lambda, +\infty[}^*(N \times \mathbb{R}; \mathcal{H}om^*(\mathcal{F}_{L_1}, \mathcal{F}_{L_2})) \otimes H_{N \times [\mu, +\infty[}^*(N \times \mathbb{R}; \mathcal{H}om^*(\mathcal{F}_{L_2}, \mathcal{F}_{L_3}))$$

$$\downarrow \cup_*$$

$$H_{N \times [\lambda + \mu, +\infty[}^*(N \times \mathbb{R}; \mathcal{H}om^*(\mathcal{F}_{L_1}, \mathcal{F}_{L_3}))$$

that coincides through the above identifications with the triangle product in Floer cohomology.

Remarks 5.3.

- (1) Note that we have

$$\mathcal{F}_{T_c L} = T_{c*}(\mathcal{F}_L)$$

- (2) In case L has a Generating Function Quadratic at infinity, $S(x, \xi)$ defined on the vector bundle $E \xrightarrow{\pi} N$, we can take for \mathcal{F}_L the complex $R(\pi \times \mathrm{id}_{\mathbb{R}})_*(k_{U_S})$ where $U_S = \{(x, \xi, t) \mid S(x, \xi) \leq t\}$.

5.1. Definition of spectral invariants. Now we can give a sheafy definition of the spectral invariants introduced in [Vit92]. The next formulation appears in [Vic13] (see also [Vit22]).

In fact the invariants can be defined for objects \mathcal{F} of $D_{\tau \geq 0}(N \times \mathbb{R})$ satisfying:

$$(5.3) \quad \mathcal{F}|_{N \times]-\infty, -t_0[} \simeq 0, \quad \mathcal{F}|_{N \times]t_0, +\infty[} \simeq k_{N \times \mathbb{R}}|_{N \times]t_0, +\infty[} \quad \text{for some } t_0.$$

We have seen that \mathcal{F}_L satisfies (5.3) for any $L \in \mathcal{L}(T^*N)$. For $\mathcal{F}_1, \mathcal{F}_2 \in D_{\tau \geq 0}(N \times \mathbb{R})$ satisfying (5.3), $\mathcal{H}om^*(\mathcal{F}_1, \mathcal{F}_2)$ is $k_{N \times \mathbb{R}}[1]$ near $N \times \{-\infty\}$ and 0 near $N \times \{+\infty\}$. Using 5.1 and $H_{N \times \{c\}}^0(N \times \mathbb{R}; k_{N \times \mathbb{R}}[1]) \simeq k$, we deduce a natural morphism $\mathcal{F}_1 \rightarrow T_{c^*}(\mathcal{F}_2)$, for any c big enough, which restricts to $\text{id}: k_{N \times \mathbb{R}} \rightarrow k_{N \times \mathbb{R}}$ near $N \times \{+\infty\}$. More generally $H_{N \times \{c\}}^*(N \times \mathbb{R}; k_{N \times \mathbb{R}}[1]) \simeq H^*(N; k_N)$ and any class $\alpha \in H^d(N; k_N)$ yields a morphism

$$u(\alpha, \mathcal{F}_1, \mathcal{F}_2, c): \mathcal{F}_1 \rightarrow T_{c^*}(\mathcal{F}_2)[d],$$

for c large enough, which restricts to $\alpha: k_{N \times \mathbb{R}} \rightarrow k_{N \times \mathbb{R}}[d]$ near $N \times \{+\infty\}$ (using $\text{Mor}(k_N, k_N[d]) \simeq H^d(N; k_N)$). The cup product corresponds to the composition, and for c (and t_0) large enough this is summarized by the following diagram

$$\begin{array}{ccc} H^*(N, k) \simeq R\Gamma(N, k_N) & \xrightarrow{\simeq} & \text{RHom}(\mathcal{F}_1, T_{c^*}\mathcal{F}_2) \\ \downarrow \simeq & & \downarrow \simeq \\ \text{RHom}(k_{N \times]t_0, +\infty[}, k_{N \times]t_0, +\infty[}) & \xrightarrow{\simeq} & \text{RHom}(\mathcal{F}_1|_{N \times]t_0, +\infty[}, T_{c^*}\mathcal{F}_2|_{N \times]t_0, +\infty[}) \end{array}$$

Definition 5.4. For $\mathcal{F}_1, \mathcal{F}_2 \in D_{\tau \geq 0}(N \times \mathbb{R})$ satisfying (5.3) and $\alpha \in H^d(N; k_N)$ we set

$$c(\alpha, \mathcal{F}_1, \mathcal{F}_2) = -\inf\{c; \exists u: \mathcal{F}_1 \rightarrow T_{c^*}(\mathcal{F}_2)[d], u|_{N \times]t_0, +\infty[} = \alpha, \text{ for } t_0 \gg 0\}$$

(Here t_0 is big enough so that both \mathcal{F}_1 and $T_{c^*}(\mathcal{F}_2)$ satisfy (5.3)). We also set $c_-(\mathcal{F}_1, \mathcal{F}_2) = c(1, \mathcal{F}_1, \mathcal{F}_2)$ and if N is oriented with fundamental class μ_N , $c_+(\mathcal{F}_1, \mathcal{F}_2) = c(\mu_N, \mathcal{F}_1, \mathcal{F}_2)$. Finally we set $\gamma(\mathcal{F}_1, \mathcal{F}_2) = c_+(\mathcal{F}_1, \mathcal{F}_2) - c_-(\mathcal{F}_1, \mathcal{F}_2)$.

For $L_1, L_2 \in \mathcal{L}(T^*N)$ we set $c(\alpha, L_1, L_2) = c(\alpha, \mathcal{F}_{L_1}, \mathcal{F}_{L_2})$ and we define $c_{\pm}(L_1, L_2), \gamma(L_1, L_2)$ in the same way. Finally we set $c_{\pm}(L) = c_{\pm}(0_N, L)$.

It is not difficult to see that

$$c_-(\mathcal{F}_1, \mathcal{F}_2) = \inf\{c(\alpha, \mathcal{F}_1, \mathcal{F}_2); \alpha \in H^*(N; k_N)\}$$

and

$$c_+(\mathcal{F}_1, \mathcal{F}_2) = \sup\{c(\alpha, \mathcal{F}_1, \mathcal{F}_2); \alpha \in H^*(N; k_N)\}$$

Indeed, a class α induces by multiplication $v: \mathcal{F}_2 \rightarrow \mathcal{F}_2[d]$, $v|_{N \times]t_0, +\infty[} = \alpha$; now if we have $u: \mathcal{F}_1 \rightarrow T_{c^*}(\mathcal{F}_2)[0]$, $u|_{N \times]t_0, +\infty[} = 1$, then $v \circ u: \mathcal{F}_1 \rightarrow T_{c^*}(\mathcal{F}_2)[d]$ satisfies $v \circ u|_{N \times]t_0, +\infty[} = \alpha$ so $c(1, \mathcal{F}_1, \mathcal{F}_2) \leq c(\alpha, \mathcal{F}_1, \mathcal{F}_2)$.

Using the construction of the sheaf \mathcal{F}_L by using Floer cohomology as in [Vit19] it is easy to see that the present definition of $c(\alpha, \mathcal{F}_{L_1}, \mathcal{F}_{L_2})$ coincides with the definition of $c(\alpha, L_1, L_2)$ using Floer cohomology. In particular, we have that $c_+(\mathcal{F}_{L_1}, \mathcal{F}_{L_2}) = -c_-(\mathcal{F}_{L_2}, \mathcal{F}_{L_1})$ and this can be extended to the general case of $\mathcal{F}_1, \mathcal{F}_2$ using [Vit19].

We have given the definition using the right hand side of (5.1). We can also rewrite it with the left hand side as follows. Let us set $\mathcal{H} = Rt_* \mathcal{H}om^*(\mathcal{F}, \mathcal{G})$, where $t: N \times \mathbb{R} \rightarrow \mathbb{R}$ is the projection. Then \mathcal{H} contains all the information about the spectral invariants. More precisely (5.1) is isomorphic to $R\Gamma_{\{-c\}}(\mathcal{H})$ and \mathcal{H} has its singular support in $\{\tau \geq 0\}$ so we have the morphisms

$$(5.4) \quad R\Gamma(N; k_N) \simeq R\Gamma_{\{-t_0\}}(\mathbb{R}; \mathcal{H}) \xrightarrow{\sim} R\Gamma_{[-t_0, -c]}(\mathbb{R}; \mathcal{H}) \leftarrow R\Gamma_{\{-c\}}(\mathbb{R}; \mathcal{H})$$

for $t_0 \gg 0$ and any $c \leq t_0$. To find the spectral invariants we look for the infimum on the c such that a given class is in the image of the composition of this morphisms.

Remark 5.5. We can see that the infimum in Definition 5.4 is actually a minimum, at least when \mathcal{F} is of the form \mathcal{F}_L . Indeed, this follows either by identifying the spectral invariants of the Definition to those obtained by using Floer cohomology, or by appealing to the theory of persistence modules and barcodes (see Appendix A Proposition A.1).

We notice the following immediate consequence of the definitions.

Proposition 5.6. *There exists a morphism $u: \mathcal{F}_{L_1} \rightarrow \mathcal{F}_{L_2}$ such that $u = \text{Id}$ at $+\infty$ if and only if $c_-(L_1, L_2) \geq 0$. In particular there is a morphism $k_{N \times [0, +\infty[} \rightarrow \mathcal{F}$ if and only if $c_-(L) \geq 0$.*

5.2. The case of Hamiltonian maps. If φ is a Hamiltonian map of $T^*\mathbb{R}^n$ with compact support, we define the spectral invariants of φ as those of its graph, say $\Gamma_\varphi \subset \overline{T^*\mathbb{R}^n} \times T^*\mathbb{R}^n$, compactified in the following way: we choose a symplectomorphism $\psi: \overline{T^*\mathbb{R}^n} \times T^*\mathbb{R}^n \xrightarrow{\sim} T^*\Gamma_{\text{id}}$ such that $\psi(\Gamma_{\text{id}}) = 0_{\Gamma_{\text{id}}}$; then $\psi(\Gamma_\varphi)$ coincides with the zero-section outside a compact set and we can compactify it as Γ_φ^c in the cotangent bundle of the sphere. In particular $\Gamma_{\text{id}}^c = 0_{S^{2n}}$ and we set

Definition 5.7. *The invariants of φ , $c_+(\varphi)$, $c_-(\varphi)$ are defined as those of $(\Gamma_\varphi^c, 0_{S^{2n}})$.*

This is well-defined since another choice of ψ would give a pair $(\Gamma_\varphi'^c, 0_{S^{2n}})$ symplectomorphic to $(\Gamma_\varphi^c, 0_{S^{2n}})$.

On the other hand we have seen in §4 that φ has an associated sheaf $\mathcal{K}_\varphi \in D(\mathbb{R}^{2n+1})$. In the next paragraph we explain how we can turn the above definition of $c_+(\varphi)$, $c_-(\varphi)$ into Definition 5.8 which uses \mathcal{K}_φ , \mathcal{K}_{id} and not the intermediate symplectomorphism ψ .

Support in a Darboux chart. Let now $S_\varphi(q, P, \xi)$ be a generating function quadratic at infinity for Γ_φ over $\Delta_{\mathbb{R}^{2n}}$, that is,

$$\begin{cases} p - P = \frac{\partial S_\varphi}{\partial q}(q, P; \xi) \\ Q - q = \frac{\partial S_\varphi}{\partial P}(q, P; \xi) \\ \frac{\partial S_\varphi}{\partial \xi}(q, P; \xi) = 0 \end{cases}$$

Since φ is compactly supported, so is S_φ in the q, P variables: for $|q| + |P|$ large enough, $S_\varphi(q, P; \xi) = B(\xi)$ where B is a non-degenerate quadratic form. We can then extend S_φ to a generating function quadratic at infinity on a bundle over S^{2n} (see [Vit92]), and $c_\pm(\varphi)$ are defined as

$$\begin{aligned} c_+(\varphi) &= \inf\{t \mid T \cup \mu_{S^{2n}} \neq 0 \text{ in } H^*(S^t, S^{-\infty})\}, \\ c_-(\varphi) &= \inf\{t \mid T \cup 1 \neq 0 \text{ in } H^*(S^t, S^{-\infty})\}, \end{aligned}$$

where T is the Thom class associated to the negative bundle of B and $S^{-\infty}$ means S^{-c} for c large enough. Note that these coincide with the spectral invariants defined using Floer cohomology (see [Vit95]).

Now if φ is defined on T^*N but has support in a Darboux chart, it can be considered either as a compact supported Hamiltonian map on T^*N or on \mathbb{R}^{2n} and we can define the spectral invariants in two ways (using Floer cohomology). By an easy argument (see Appendix D) we can see that these two definitions of $c_\pm(\varphi)$ do coincide.

Sheaf definition of spectral invariants for a Hamiltonian map. Now we can give another formulation of the definition of $c_\pm(\varphi)$. Let N be a compact manifold and let φ be a Hamiltonian map of T^*N with compact support. We consider its homogenized graph $\widehat{\Lambda}_\varphi \subset \overline{T^*N} \times T^*N \times T^*\mathbb{R}$ as in §4. We have associated to φ its quantization \mathcal{K}_φ , with singular support $\widehat{\Lambda}_\varphi$, and it coincides with $\mathcal{K}_{\text{id}} = k_{\Delta_N \times [0, +\infty[}$ outside a compact subset of $N^2 \times \mathbb{R}$. Now $\mathcal{H}\text{om}^*(\mathcal{K}_{\text{id}}, \mathcal{K}_\varphi)$ has similar properties as $\mathcal{H}\text{om}^*(\mathcal{F}_{L_1}, \mathcal{F}_{L_2})$, namely, it is $k_{\Delta_N \times \mathbb{R}}[1]$ near $N^2 \times \{-\infty\}$ and 0 near $N^2 \times \{+\infty\}$. As in the case of compact Lagrangians we have $H_{N^2 \times \{c\}}^*(N^2 \times \mathbb{R}; k_{\Delta_N \times \mathbb{R}}[1]) \simeq H^*(N; k_N)$ and any class $\alpha \in H^d(N; k_N)$ yields a morphism

$$u(\alpha, \varphi, c): \mathcal{K}_{\text{id}} \rightarrow T_{c*}(\mathcal{K}_\varphi)[d],$$

for c big enough, which restricts to $\alpha: k_{\Delta_N \times \mathbb{R}} \rightarrow k_{\Delta_N \times \mathbb{R}}[d]$ near $N \times \{+\infty\}$ (using $\text{Hom}(k_{\Delta_N}, k_{\Delta_N}[d]) \simeq H^d(N; k_N)$).

Definition 5.8. *For a compactly supported Hamiltonian map φ of T^*N and $\alpha \in H^d(N; k_N)$ we set*

$$c(\alpha, \varphi) = -\inf\{c; \exists u: \mathcal{K}_{\text{id}} \rightarrow T_{c*}(\mathcal{K}_\varphi)[d], u|_{N^2 \times]t_0, +\infty[} = \alpha, \text{ for } t_0 \gg 0\}$$

(Here t_0 is large enough so that over $N^2 \times]t_0, +\infty[$ we have $T_{c^*}(\mathcal{K}_\varphi) \simeq k_{N \times \mathbb{R}}$.) We also set $c_-(\varphi) = c(1, \varphi)$, $c_+(\varphi) = -c_-(\varphi^{-1})$ and $\gamma(\varphi) = c_+(\varphi) - c_-(\varphi)$. If ψ is another compactly supported Hamiltonian map, we set $\gamma(\varphi, \psi) = \gamma(\varphi\psi^{-1})$.

We have $c_-(\varphi) = \inf\{c(\varphi, \alpha); \alpha \in H^*(N; k_N)\}$. Since $\Gamma(\varphi)$ coincides with the zero section outside a compact set, normalizing the primitive of λ on $\Gamma(\varphi)$ to be zero outside a compact set, we have $c_-(\varphi) \leq 0 \leq c_+(\varphi)$ as in [Vit92].

Equivalence between Definitions 5.7 and 5.8. We first remark that we don't really need to compactify Γ_{id} into the sphere to obtain the invariants. Let D_1 be the subcategory of $D_{\tau \geq 0}(\Gamma_{\text{id}} \times \mathbb{R})$ of complexes isomorphic to $k_{\Gamma_{\text{id}} \times]c, +\infty[}$ (for some c) and D_2 be the subcategory of $D_{\tau \geq 0}(S^{2n} \times \mathbb{R})$ of complexes isomorphic to $k_{S^{2n} \times]c, +\infty[}$ near $\{\infty\} \times]c, +\infty[$. Then the compactification and the restriction map induce functors $\alpha: D_1 \rightarrow D_2$ and $\beta: D_2 \rightarrow D_1$ such that $\beta \circ \alpha = \text{id}$, $\alpha(\tau_{a,b}(\mathcal{F})) = \tau_{a,b}(\alpha(\mathcal{F}))$. We deduce that $c_-(\alpha(\mathcal{F}_1), \alpha(\mathcal{F}_2)) = c_-(\mathcal{F}_1, \mathcal{F}_2)$ (we remark that c_- makes sense even in the non compact case). In particular

$$(5.5) \quad c_-(\varphi) = c_-(\mathcal{F}_{\Gamma_\varphi^c}, \mathcal{F}_{0_{S^{2n}}}) = c_-(\mathcal{F}_{\Gamma_\varphi}, \mathcal{F}_{0_{\mathbb{R}^{2n}}})$$

Now the symplectomorphism ψ induces an equivalence of categories between $\mathcal{K}_\psi^\otimes: D_{\tau \geq 0}(\mathbb{R}^{2n+1}) \xrightarrow{\sim} D_{\tau \geq 0}(\Gamma_{\text{id}} \times \mathbb{R})$ which commutes with the translation T_{c^*} in the last variables. We have $\mathcal{K}_\psi^\otimes(\mathcal{K}_\varphi) \simeq \mathcal{F}_{\Gamma_\varphi}$ and we obtain from (5.5) that the spectral invariant $c_-(\varphi)$ we have defined above coincides with the one in Definition 5.8 below. To obtain c_+ we use $c_+(\mathcal{F}_1, \mathcal{F}_2) = -c_-(\mathcal{F}_2, \mathcal{F}_1)$.

The existence of \mathcal{K}_ψ doesn't really follows from §4 because ψ is not compactly supported. But the choice of ψ is irrelevant and we can choose $\psi: T^*\mathbb{R}^n \times T^*\mathbb{R}^n \rightarrow T^*\Delta_{\mathbb{R}^{2n}}$ given by $\psi(q_1, q_2, p_1, p_2) = (q_1, p_2, p_1 + p_2, q_1 - q_2)$. In this case we can give a direct construction of \mathcal{K}_ψ as follows. Let us define $\Sigma^+ \subset \mathbb{R}^{4n+1}$ by $\Sigma^+ = \{(q_1, q_2, Q_1, Q_2, t) \mid Q_1 = q_1, t \geq (q_1 - q_2)Q_2\}$, whose boundary is the projection of the lift $\widehat{\Lambda}_\psi$ of the graph of ψ to the base. Then we can check that $\mathcal{K}_\psi = k_{\Sigma^+}$ is the quantization of ψ .

This justifies that both definitions coincide for a Hamiltonian map of T^*S^n obtained from a compactly supported map of $T^*\mathbb{R}^n$. We saw in a previous paragraph that the case of a Hamiltonian map of T^*N with support in a Darboux chart is reduced to this case.

Note the following analogue of Proposition 5.6

Lemma 5.9. *Let φ be a compact supported Hamiltonian map. Then there is a morphism $\mathcal{K}_{\text{id}} \rightarrow \mathcal{K}_\varphi$ in $D^b(N \times N \times \mathbb{R})$ equal to Id at $+\infty$ if and only*

if $c_-(\varphi) = 0$. Analogously there is a morphism $\mathcal{K}_\varphi \longrightarrow \mathcal{K}_{\text{id}}$ in $D^b(N \times N \times \mathbb{R})$ if and only if $c_+(\varphi) = 0$.

Proof. The first statement follows from Proposition 5.6. For the second one, we have by assumption a morphism $\mathcal{K}_{\text{id}} \longrightarrow \mathcal{K}_{\varphi^{-1}}$. Composing by \mathcal{K}_φ we get a morphism $\mathcal{K}_\varphi \longrightarrow \mathcal{K}_{\text{id}}$. Analogously, if we have such a morphism, composing by $\mathcal{K}_{\varphi^{-1}}$ we get $c_+(\varphi) = 0$. \square

5.3. Defining γ -coisotropic subsets. For an open set U in the symplectic manifold (M, ω) we denote by $\mathcal{D}\mathfrak{H}\text{am}_c(U)$ the set of time one flows of Hamiltonian with compact support contained in U .

Definition 5.10 (see definition 6.1 in [Vit22]). *Let V be a set in (M, ω) . We say that V is γ -coisotropic at $x \in V$ if there exists $\varepsilon > 0$ such that, for all η with $0 < \eta < \varepsilon$, there is a $\delta > 0$ such that, for all φ in $\mathcal{D}\mathfrak{H}\text{am}_c(B(x, \varepsilon))$ satisfying $\varphi(V) \cap B(x, \eta) = \emptyset$, we have $\gamma(\varphi) > \delta$.*

We shall say that V is γ -coisotropic if it is closed, non-empty and γ -coisotropic at each $x \in V$.

We refer to [Vit22], in particular section 6, for properties and examples of γ -coisotropic subsets. In particular we remind the reader of the following result from [Vit22].

Proposition 5.11. *Let φ be a homeomorphism preserving γ , for example a C^0 -limit of Hamiltonian maps (usually called Hamiltonian homeomorphisms) and V a γ -coisotropic set. Then, the image by φ of V is a γ -coisotropic set.*

Note that the completion of $\mathcal{L}(T^*N)$ for γ is denoted by $\widehat{\mathcal{L}}(T^*N)$. In [Vit22] the γ -support of an element L in $\widehat{\mathcal{L}}(T^*N)$ was defined, and it was proved that it must be a γ -coisotropic subset.

6. THE γ_g -METRIC ON SHEAVES

We shall define the analogue of the γ distance between sheaves. It is also a sheaf analogue of the barcode distance (see [KS18] or [AI20]) and relies on the morphism τ introduced by Tamarkin (see [Tam08] or [GKS, Prop. 4.8]) that we have already encountered in (5.2).

To define τ in general we recall that the subcategory $D_{\tau \geq 0}(N \times \mathbb{R})$ of $D(N \times \mathbb{R})$ can be characterized as follows (see [KS90, Prop. 5.2.3, 3.5.4]). Let P be the endofunctor of $D(N \times \mathbb{R})$ defined by

$$P(\mathcal{F}) = Rs_* (\mathcal{F} \boxtimes k_{[0, +\infty[})$$

with $s: N \times \mathbb{R}^2 \rightarrow N \times \mathbb{R}$, $(x, t_1, t_2) \mapsto (x, t_1 + t_2)$ (this is almost $\mathcal{F} * k_{N \times [0, +\infty[}$ up to replacing from $Rs_!$ to Rs_*). Then the natural morphism $k_{[0, +\infty[} \rightarrow k_{\{0\}}$ induces a morphism $P(\mathcal{F}) \rightarrow \mathcal{F}$ and $D_{\tau \geq 0}(N \times \mathbb{R})$ is the set of \mathcal{F} such

that $P(\mathcal{F}) \rightarrow \mathcal{F}$ is an isomorphism. Now we have in general $R_{s_*}(\mathcal{F} \boxtimes k_{\{c\}}) \simeq T_{c^*}(\mathcal{F})$, where $T_c(x, t) = (x, t + c)$ and we obtain in the same way, using the morphism $k_{[0, +\infty[} \rightarrow k_{\{c\}}$, a morphism $P(\mathcal{F}) \rightarrow T_{c^*}(\mathcal{F})$ for any $c \geq 0$. In particular we have a morphism of endofunctors defined on $D_{\tau \geq 0}(N \times \mathbb{R})$ for any $c \geq 0$

$$\tau_{0,c}(\mathcal{F}): \mathcal{F} \rightarrow T_{c^*}(\mathcal{F})$$

We shall call this morphism the Tamarkin morphism.

The case of $D_{\tau \geq 0}(N \times \mathbb{R})$ is in fact sufficient for this paper but we can consider a slightly more general case without modifying the proofs. Let $g: T^*N \setminus 0_N \rightarrow \mathbb{R}$ be a 1-homogeneous autonomous Hamiltonian function and let $D_{g \geq 0}(N)$ be the full subcategory of $D(N)$ formed by the \mathcal{F} with $SS^*(\mathcal{F}) \subset \{g \geq 0\}$. We let $\varphi^t = \varphi_g^t: T^*N \setminus 0_N \rightarrow T^*N \setminus 0_N$ be the corresponding homogeneous flow. We set for short $K_\varphi^a(\mathcal{F}) = \mathcal{F} \circ K_{\varphi^a}$.

We consider the lifting of the graph of the whole isotopy φ^t to Lagrangian in $T^*(N \times N \times \mathbb{R})$ given by

$$\{(q, p, Q_t(q, p), P_t(q, p), t, g(q, p))\},$$

where $\varphi^t(q, p) = (Q_t(q, p), P_t(q, p))$ and the associated kernel $K_\varphi \in D(N^2 \times \mathbb{R})$. Then $K_\varphi(\mathcal{F}) \in D(N \times \mathbb{R})$ is such that $K_\varphi(\mathcal{F})|_{N \times \{a\}} = K_\varphi^a(\mathcal{F})$. Now, if $\mathcal{F} \in D_{g \geq 0}(N)$, then $K_\varphi(\mathcal{F}) \in D_{\tau \geq 0}(N \times \mathbb{R})$. Indeed a point of $SS^*(K_\varphi(\mathcal{F}))$ is written (q, p, Q, P, t, τ) with $(q, p) \in SS^*(\mathcal{F})$ and $\tau = g(q, p)$. We thus have the Tamarkin morphism $\tau_{0,c}(K_\varphi(\mathcal{F})): K_\varphi(\mathcal{F}) \rightarrow T_{c^*}(K_\varphi(\mathcal{F}))$ for $c \geq 0$. Restricting to $N \times \{a\}$ and setting $b = a + c$ we obtain

$$(6.1) \quad \tau_{a,b}(\mathcal{F}): K_\varphi^a(\mathcal{F}) \rightarrow K_\varphi^b(\mathcal{F}), \quad \text{for } a \leq b.$$

The morphisms $\tau_{a,b}$ are invariant by K_φ^t , for any t , and compatible with composition:

$$(6.2) \quad \begin{aligned} K_\varphi^t(\tau_{a,b}(\mathcal{F})) &= \tau_{a+t, b+t}(K_\varphi^t(\mathcal{F})), \\ \tau_{a,c}(\mathcal{F}) &= \tau_{b,c}(\mathcal{F}) \circ \tau_{a,b}(\mathcal{F}), \quad \text{for } a \leq b \leq c \end{aligned}$$

Examples 6.1.

- (1) Let $N = \mathbb{R}$ and g be the norm given by the standard metric, so $g(t, \tau) = |\tau|$. Then $\varphi^a(t, \tau) = (t + a \frac{\tau}{|\tau|}, \tau)$. As a result $K_{\varphi^a}(k_{[x,y]}) = k_{[x+a, y+a]}$ and $SS(K_{\varphi^a}) = \{(t, t + a \frac{\tau}{|\tau|}, \tau, \tau)\} \subset T^*(\mathbb{R}^2)$.
- (2) In the case of $N \times \mathbb{R}$ and $g = \tau$ we recover the case we introduced in the first paragraph (and originally due to Tamarkin).
- (3) If g is non negative we have $D_{g \geq 0}(N) = D(N)$. Typically g is the norm given by a Riemannian metric and φ_g is the normalized geodesic flow. In this case, denoting by $B(x, r)$ the open ball at x of

radius r , we find

$$K_\varphi^a(k_{B(x,r)}) \simeq \begin{cases} k_{B(x,r+a)} & \text{for } -r < a < r_{inj} - r, \\ k_{\overline{B(x,-r-a)}}[-\dim N] & \text{for } -r_{inj} - r < a \leq -r, \end{cases}$$

and

$$K_\varphi^a(k_{\overline{B(x,r)}}) \simeq \begin{cases} k_{\overline{B(x,r-a)}} & \text{for } r - r_{inj} \leq a \leq r, \\ k_{B(x,a-r)}[\dim N] & \text{for } r < a < r + r_{inj}, \end{cases}$$

where r_{inj} is the injectivity radius.

6.1. The γ_g -topology. The following definition is the analogue of the spectral metric (in the case $g = \tau$) (and a sheaf analogue of the barcode distance). It is introduced in [KS18] or [AI20] (up to a small modification).

Definition 6.2. *Let $g: T^*N \setminus 0_N \rightarrow \mathbb{R}$ be a 1-homogeneous autonomous Hamiltonian function and $\varphi = \varphi_g^t$ its flow. For $\mathcal{F}, \mathcal{G} \in D_{g \geq 0}(N)$, we let $\gamma_g(\mathcal{F}, \mathcal{G}) \in [0, \infty]$ be the infimum of the $c \geq 0$ for which there exist $a, b \geq 0$ such that $a + b = c$ and such that there are morphisms*

$$u: \mathcal{F} \rightarrow K_\varphi^a(\mathcal{G}), \quad v: \mathcal{G} \rightarrow K_\varphi^b(\mathcal{F})$$

satisfying $K_\varphi^a(v) \circ u = \tau_{0,a+b}(\mathcal{F})$ and $K_\varphi^b(u) \circ v = \tau_{0,a+b}(\mathcal{G})$.

The authors thank N. Vichery for the next remark.

Remark 6.3. The case $D_{g \geq 0}(N)$ is not really more general than the case $D_{\tau \geq 0}(N \times \mathbb{R})$. Indeed the functor $K_\varphi(-): D(N) \rightarrow D(N \times \mathbb{R})$ is fully faithful and we have $T_{a^*}(K_\varphi(-)) \simeq K_\varphi(K_\varphi^a(-))$. Hence, for $\mathcal{F}, \mathcal{G} \in D_{g \geq 0}(N)$ we have $\gamma_g(\mathcal{F}, \mathcal{G}) = \gamma_\tau(\mathcal{K}_\varphi(\mathcal{F}), \mathcal{K}_\varphi(\mathcal{G}))$.

The definition of γ_g in [KS18] imposes $a = b$ which leads to a bigger distance, say γ'_g , with $\gamma_g \leq \gamma'_g \leq 2\gamma_g$. The definition in [AI20] only asks $K_\varphi^a(v) \circ u = \tau_{0,a+b}(\mathcal{F})$ and $K_\varphi^b(u') \circ v' = \tau_{0,a+b}(\mathcal{G})$, for two possibly different morphisms u', v' (see the arXiv version for another definition). This leads to a smaller distance but we don't know if it is equivalent to γ_g . Lemma 6.8 below says that γ_g is equivalent to $\inf\{\gamma_g(\mathcal{C}(u), 0); u: \mathcal{F} \rightarrow \mathcal{G}\}$, where $\mathcal{C}(u)$ denotes the cone of u .

Lemma 6.4. *For $\mathcal{F}, \mathcal{G}, \mathcal{H} \in D_{g \geq 0}(N)$ we have*

- (1) $\gamma_g(\mathcal{F}, \mathcal{G}) = \gamma_g(K_\varphi^t(\mathcal{F}), K_\varphi^t(\mathcal{G})) = \gamma_g(\mathcal{G}, \mathcal{F})$.
- (2) $\gamma_g(\mathcal{F}, \mathcal{H}) \leq \gamma_g(\mathcal{F}, \mathcal{G}) + \gamma_g(\mathcal{G}, \mathcal{H})$.
- (3) $\gamma_g(0, \mathcal{F}) = \inf\{t \geq 0; \tau_{0,t}(\mathcal{F}) = 0\}$.
- (4) *If \mathcal{F} satisfies $\gamma_g(\mathcal{F}, 0) = 0$, then $\mathcal{F} \simeq 0$.*

(5) For any distinguished triangle $\mathcal{F} \xrightarrow{u} \mathcal{G} \xrightarrow{v} \mathcal{H} \xrightarrow{+1}$, we have

$$\gamma_g(0, \mathcal{G}) \leq \gamma_g(0, \mathcal{F}) + \gamma_g(0, \mathcal{H})$$

(6) For composable morphisms u, v with cones $\mathcal{C}(u), \mathcal{C}(v), \mathcal{C}(v \circ u)$, we have $\gamma_g(0, \mathcal{C}(v \circ u)) \leq \gamma_g(0, \mathcal{C}(u)) + \gamma_g(0, \mathcal{C}(v))$.

Proof. (1) follows from (6.2) and (3) follows from the definition of γ_g . The triangular inequality (2) is proved in [AI20, Prop. 4.11].

Let us prove (4). By Remark 6.3 we can assume that $N = M \times \mathbb{R}$ and $g = \tau$. Let $x_0 \in M$ be given and let $i: \{x_0\} \times \mathbb{R} \rightarrow M \times \mathbb{R}$ be the inclusion. We have $\tau_{a,b}(i^{-1}\mathcal{F}) = i^{-1}(\tau_{a,b}(\mathcal{F}))$ and it follows that $\gamma_g(i^{-1}\mathcal{F}, i^{-1}\mathcal{G}) \leq \gamma_g(\mathcal{F}, \mathcal{G})$. If $\mathcal{F} \neq 0$, there exists x_0 such that $i^{-1}(\mathcal{F}) \neq 0$. Hence we can assume that $N = \mathbb{R}$. We can also take $g = |\tau|$ (it is differentiable on $T^*\mathbb{R} \setminus 0_{\mathbb{R}}$) instead of $g = \tau$. The corresponding τ morphisms, say $\tau_{a,b}^g$, coincide with $\tau_{a,b}$ on $D_{\tau \geq 0}(\mathbb{R})$ but they are defined on $D(\mathbb{R})$. Up to shifting \mathcal{F} in degrees and translating, we can assume that $H^0\mathcal{F}_0 \neq 0$. Hence there exists $u \in H^0(-\varepsilon, \varepsilon; \mathcal{F})$ with $u_0 \neq 0$. Interpreting u as a morphism and choosing a map $v: \mathcal{F}_0 \rightarrow k$ non-vanishing on the germ defined by u , we get that the composition of v and u ; $k_{]-\varepsilon, \varepsilon[} \xrightarrow{u} \mathcal{F} \xrightarrow{v} k_{\{0\}}$ is the canonical morphism. Now the morphisms $\tau_{0,\delta}^g(-)$, for $0 < \delta < \varepsilon$ give the commutative diagram (using Example 6.1 (3)) where the vertical arrows are given by $\tau_{0,\delta}^g$

$$\begin{array}{ccccc} k_{]-\varepsilon, \varepsilon[} & \xrightarrow{u} & \mathcal{F} & \xrightarrow{v} & k_{\{0\}} \\ \downarrow & & \downarrow & & \downarrow w \\ k_{]-\varepsilon+\delta, \varepsilon-\delta[} & \longrightarrow & T_{\delta*}(\mathcal{F}) & \longrightarrow & k_{]-\delta, \delta[}[1] \end{array}$$

We check that $w \circ v \circ u \neq 0$; this then implies that $\tau_{0,\delta}^g(\mathcal{F}) \neq 0$, as required. Since $w \neq 0$, we have a distinguished triangle $k_{]-\delta, \delta[} \rightarrow k_{]-\delta, 0[} \oplus k_{[0, \delta[} \rightarrow k_{\{0\}} \xrightarrow{w} k_{]-\delta, \delta[}[1]$ (in other words w gives a non trivial Yoneda extension). If $w \circ (v \circ u)$ were 0, then $v \circ u$ would factorize through $k_{]-\delta, 0[} \oplus k_{[0, \delta[}$. But $\text{Hom}(k_{]-\varepsilon, \varepsilon[}, k_{]-\delta, 0[}) = \text{Hom}(k_{]-\varepsilon, \varepsilon[}, k_{[0, \delta[}) = 0$ because $\varepsilon > \delta$ and we cannot have such a factorization.

Let us prove (5) (a similar but more detailed argument will be given in the proof of Lemma 6.8-(i)). If $\tau_{0,a}(\mathcal{F}) = 0$, then $\tau_{0,a}(\mathcal{G}) \circ u = K_{\varphi}^a(u) \circ \tau_{0,a}(\mathcal{F}) = 0$ and $\tau_{0,a}(\mathcal{G})$ factorizes through v . In the same way, if $\tau_{0,b}(\mathcal{H}) = 0$, then $\tau_{0,b}(\mathcal{G})$ factorizes through $K_{\varphi}^b(u)$. Hence $\tau_{0,a+b}(\mathcal{G})$ factorizes via $K_{\varphi}^b(v) \circ K_{\varphi}^b(u) = 0$. We conclude with (3).

We deduce (6) from (5) since, by the octaedron axiom, the three cones appear in a distinguished triangle. \square

If N is real analytic and \mathcal{F}, \mathcal{G} are constructible and satisfy $\gamma_g(\mathcal{F}, \mathcal{G}) = 0$, then $\mathcal{F} \simeq \mathcal{G}$ (see [PSW] and also Proposition B.8 for limits of constructible sheaves). However γ_g is only a pseudo-metric as shown by Berkouk and Ginot (see [BG18], proposition 6.9) and the following example.

Example 6.5. We define two sheaves on the real line $\mathcal{F} = \bigoplus_{x \in \mathbb{Q}} k_{[x, \infty[}$ and $\mathcal{G} = \bigoplus_{x \in \mathbb{Q}^\times} k_{[x, \infty[}$, where $\mathbb{Q}^\times = \mathbb{Q} \setminus \{0\}$. We remark that $K_\varphi^t(\mathcal{F}) \simeq T_{t*}(\mathcal{F})$, where $T_t(x) = x + t$. For a given $\varepsilon > 0$ we choose a bijection $f: \mathbb{Q} \rightarrow \mathbb{Q}^\times$ such that $x \leq f(x) \leq x + \varepsilon$, for all $x \in \mathbb{Q}$, and we define $u = \bigoplus_{x \in \mathbb{Q}} u_x: \mathcal{F} \rightarrow \mathcal{G}$, $v = \bigoplus_{x \in \mathbb{Q}^\times} v_x: \mathcal{G} \rightarrow T_{\varepsilon*} \mathcal{F}$ where $u_x: k_{[x, \infty[} \rightarrow k_{[f(x), \infty[}$ and $v_x: k_{[x, \infty[} \rightarrow k_{[f^{-1}(x) + \varepsilon, \infty[}$ are the natural morphisms. Using u, v we see that $\gamma_g(\mathcal{F}, \mathcal{G}) \leq \varepsilon$. Hence $\gamma_g(\mathcal{F}, \mathcal{G}) = 0$.

Definition 6.6. We denote by γ_g the topology associated to the pseudo-distance γ_g . In other words a sequence $(\mathcal{F}_j)_{j \geq 1}$ in $D_{g \geq 0}(N)$ γ_g -converges to \mathcal{F} if and only if $\lim_j \gamma_g(\mathcal{F}_j, \mathcal{F}) = 0$. We may write abusively $\gamma_g - \lim \mathcal{F}_j = \mathcal{F}$ although $\gamma_g - \lim \mathcal{F}_j$ is not well-defined.

Remark 6.7. In view of the Lemma 6.8, a sequence $(\mathcal{F}_n)_{n \in \mathbb{N}}$ converges to \mathcal{F} if and only if there exist a sequence of positive numbers $(\varepsilon_n)_{n \in \mathbb{N}}$ converging to 0 and morphisms $u_n: \mathcal{F}_n \rightarrow K_\varphi^{\varepsilon_n}(\mathcal{F})$ such that $\gamma_g(0, \mathcal{C}(u_n))$ converges to 0.

Part (2) in the next lemma is analogue to [AI20, Lem. 4.14].

Lemma 6.8. Let $u: \mathcal{F} \rightarrow \mathcal{F}'$ be a morphism in $D_{g \geq 0}(N)$ and let \mathcal{C} be the cone of u .

- (1) If there exists $v: \mathcal{F}' \rightarrow K_\varphi^\varepsilon(\mathcal{F})$ such that $v \circ u = \tau_{0, \varepsilon}$ and $K_\varphi^\varepsilon(u) \circ v = \tau_{0, 2\varepsilon}$, then $\gamma_g(\mathcal{C}, 0) \leq 2\varepsilon$.
- (2) If $\gamma_g(\mathcal{C}, 0) < \varepsilon$, then there exists $v: \mathcal{F}' \rightarrow K_\varphi^{2\varepsilon}(\mathcal{F})$ such that $v \circ u = \tau_{0, 2\varepsilon}$ and $K_\varphi^{2\varepsilon}(u) \circ v = \tau_{0, 4\varepsilon}$.

Proof. (1) We consider the morphism of triangles given by the functorial morphism (6.1),

$$\begin{array}{ccccccc}
 \mathcal{F} & \xrightarrow{u} & \mathcal{F}' & \xrightarrow{\alpha} & \mathcal{C} & \xrightarrow{\beta} & \mathcal{F}[1] \\
 \downarrow & & \downarrow & \searrow f & \downarrow & \searrow g & \downarrow \\
 K_\varphi^\varepsilon(\mathcal{F}) & \longrightarrow & K_\varphi^\varepsilon(\mathcal{F}') & \longrightarrow & K_\varphi^\varepsilon(\mathcal{C}) & \longrightarrow & K_\varphi^\varepsilon(\mathcal{F})[1]
 \end{array}$$

Here all small triangles commute (which defines f and g). We have $f = K_\varphi^\varepsilon(\alpha) \circ \tau_{0, \varepsilon}(\mathcal{F}') = K_\varphi^\varepsilon(\alpha \circ u) \circ v = 0$. In the same way g also vanishes. Since $\tau_{0, \varepsilon}(\mathcal{C}) \circ \alpha = f$ vanishes, $\tau_{0, \varepsilon}(\mathcal{C})$ factorizes through β as $\tau_{0, \varepsilon}(\mathcal{C}) = h \circ \beta$. Hence $\tau_{0, 2\varepsilon}(\mathcal{C}) = K_\varphi^\varepsilon(\tau_{0, \varepsilon}(\mathcal{C})) \circ \tau_{0, \varepsilon}(\mathcal{C}) = K_\varphi^\varepsilon(h) \circ K_\varphi^\varepsilon(\beta) \circ \tau_{0, \varepsilon}(\mathcal{C}) = K_\varphi^\varepsilon(h) \circ g$ vanishes, which means $\gamma_g(\mathcal{C}, 0) \leq 2\varepsilon$.

(2) Since $\tau_{0,\varepsilon}(\mathcal{C}) = 0$, both morphisms f and g vanish. Since $K_\varphi^\varepsilon(\alpha) \circ \tau_{0,\varepsilon}(\mathcal{F}') = f$ vanishes, $\tau_{0,\varepsilon}(\mathcal{F}')$ factorizes as $\tau_{0,\varepsilon}(\mathcal{F}') = K_\varphi^\varepsilon(u) \circ w$ for some w .

In the same way, the vanishing of g gives a factorization $\tau_{0,\varepsilon}(\mathcal{F}) = w' \circ u$ but we cannot say that $w = w'$.

We set $v = K_\varphi^\varepsilon(w') \circ \tau_{0,\varepsilon}(\mathcal{F}'): \mathcal{F}' \rightarrow K_\varphi^{2\varepsilon}(\mathcal{F})$. In the following computation we omit abusively to write K_φ^ε and write τ for whatever $\tau_{-,-}(-)$. A diagram chase on the diagram below gives $v \circ u = w' \circ \tau \circ u = w' \circ u \circ \tau = \tau \circ \tau = \tau$ and $u \circ v = u \circ w' \circ \tau = u \circ w' \circ u \circ w = u \circ \tau \circ w = \tau \circ u \circ w = \tau \circ \tau = \tau$, as required.

$$\begin{array}{ccc}
 \mathcal{F} & \xrightarrow{u} & \mathcal{F}' \\
 \downarrow & \swarrow w & \downarrow \\
 K_\varphi^\varepsilon(\mathcal{F}) & \xrightarrow{u} & K_\varphi^\varepsilon(\mathcal{F}') \\
 \downarrow & \swarrow w' & \downarrow \\
 K_\varphi^{2\varepsilon}(\mathcal{F}) & \xrightarrow{u} & K_\varphi^{2\varepsilon}(\mathcal{F}')
 \end{array}$$

□

6.2. Link with spectral invariants. In this paragraph we work on $N \times \mathbb{R}$ and choose $g = \tau$. So we work with the category $D_{\tau \geq 0}(N \times \mathbb{R})$ as in §5. We prove Propositions 6.11 and 6.14 which explain the relations between the γ_τ -distance and the spectral invariants.

We first give a useful variation on the Morse theorem for sheaves. The following result is mentioned in [Nad16, §6.1].

Proposition 6.9. *Let M be a manifold and I an open interval of \mathbb{R} . Let $\mathcal{F} \in D(M)$, $\mathcal{G} \in D(M \times I)$ and set $\mathcal{G}_t = \mathcal{G}|_{M \times \{t\}}$ for $t \in I$. We assume*

- (1) *there exists a compact set C of M such that $\mathcal{G}|_{(M \setminus C) \times I}$ is of the form $\mathcal{G}' \boxtimes k_I$ for some $\mathcal{G}' \in D(M \setminus C)$,*
- (2) $SS^*(\mathcal{G}) \cap (0_M \times T^*I) = \emptyset$
- (3) $(SS^*(\mathcal{F}) \times T^*I) \cap SS^*(\mathcal{G}) = \emptyset$.

Then $\mathrm{RHom}(\mathcal{F}, \mathcal{G}_t)$ does not depend on $t \in I$.

Proof. We let $i_t: M \times \{t\} \rightarrow M \times I$, $t \in I$, be the inclusion. We set $\mathcal{H} = \mathrm{R}\mathcal{H}om(\mathcal{F} \boxtimes k_I, \mathcal{G})$. By [KS90, Prop. 5.4.14], condition (3) gives $SS(\mathcal{H}) \subset \Lambda := (SS(\mathcal{F}) \times 0_I)^a + SS(\mathcal{G})$. We can see moreover that Λ is also non-characteristic for all i_t (that is, $\Lambda \cap (0_M \times T^*I) \subset 0_{M \times I}$). Using condition (2) $i_t^{-1}\mathcal{G} \simeq i_t^!\mathcal{G}[1]$ and $i_t^{-1}\mathcal{H} \simeq i_t^!\mathcal{H}[1]$ by [KS90, Prop. 5.4.13] and we deduce $\mathrm{R}\mathcal{H}om(\mathcal{F}, \mathcal{G}_t) \simeq i_t^{-1}\mathcal{H}$ by the adjunction formula $i_t^!\mathrm{R}\mathcal{H}om(-, -) \simeq \mathrm{R}\mathcal{H}om(i_t^{-1}(-), i_t^!(-))$. Hence $\mathrm{RHom}(\mathcal{F}, \mathcal{G}_t) \simeq \mathrm{R}\Gamma(M; i_t^{-1}\mathcal{H})$. By the condition (1) $\mathcal{H}|_{(M \setminus C) \times I}$ is of the form $\mathcal{H}' \boxtimes k_I$, for some $\mathcal{H}' \in D(M \setminus C)$,

and we can use the base change formula and the bound for the singular support of a direct image. Hence $\mathrm{R}\Gamma(M; i_t^{-1} \mathcal{H}) \simeq (Rq_*(\mathcal{H}))_t$, where $q: M \times I \rightarrow I$ is the projection and it is enough to check that $Rq_*(\mathcal{H})$ is constant. Since $\Lambda \cap (0_M \times T^*I) \subset 0_{M \times I}$, we have $\mathrm{SS}(Rq_*(\mathcal{H})) \subset 0_I$ and the result follows. \square

Corollary 6.10. *Let $\mathcal{F} \in D_{\tau \geq 0}(N \times \mathbb{R})$ and t_0 satisfy (5.3). We assume that $\mathrm{SS}(\mathcal{F}) \cap \mathrm{SS}(T_{c^*}(\mathcal{F})) \subset 0_{N \times \mathbb{R}}$ for any $c \neq 0$. Then the restriction map*

$\mathrm{RHom}(\mathcal{F}, T_{c^*}(\mathcal{F})) \rightarrow \mathrm{RHom}(\mathcal{F}|_U, T_{c^*}(\mathcal{F})|_U) \simeq \mathrm{RHom}(k_U, k_U) \simeq \mathrm{R}\Gamma(N; k_N)$
is an isomorphism for any $c > 0$, where $U = N \times]c + t_0, +\infty[$. In particular, if $u: \mathcal{F} \rightarrow T_{c^}(\mathcal{F})$ is id near $+\infty$, then $u = \tau_{0,c}(\mathcal{F})$.*

When $\mathrm{SS}^*(\mathcal{F})$ is the cone over a Legendrian, the hypothesis means that there is no Reeb chord. In particular the hypothesis is satisfied for the sheaves \mathcal{F}_L , $L \in \mathcal{L}(T^*N)$.

Proof. We are going to apply Proposition 6.9 with $M = N \times \mathbb{R}$, $I =]0, d]$ for $d > 0$ and $\mathcal{G} = \delta^{-1}(\mathcal{F})$, where $\delta: N \times \mathbb{R}^2 \rightarrow N \times \mathbb{R}$ is the map

$$(x, t_1, t_2) \mapsto (x, t_1 - t_2)$$

so that $\mathcal{G}|_{N \times \mathbb{R} \times \{c\}} \simeq T_{c^*}(\mathcal{F})$.

For $a < b$, \mathcal{G} is constant on $N \times]-\infty, a - t_0[\times]a, b]$ and $N \times]b + t_0, +\infty[\times]a, b]$. Hence \mathcal{G} satisfies (1) in Proposition 6.9. It is not difficult to check (2) since its singular support is contained in the conormal of $t_1 = t_2$ that is $\tau_1 + \tau_2 = 0$. For (3) it follows from the hypothesis on $\mathrm{SS}(\mathcal{F})$. The proposition gives $\mathrm{RHom}(\mathcal{F}, T_{c^*}(\mathcal{F})) \simeq \mathrm{RHom}(\mathcal{F}, T_{d^*}(\mathcal{F}))$, for any $0 < c \leq d$. \square

Proposition 6.11. *Let $L_1, L_2 \in \mathcal{L}(T^*N)$. Then*

$$\gamma_\tau(\mathcal{F}_{L_1}, \mathcal{F}_{L_2}) = \begin{cases} c_+(L_1, L_2) & \text{if } c_-(L_1, L_2) > 0, \\ -c_-(L_1, L_2) & \text{if } c_+(L_1, L_2) < 0, \\ c_+(L_1, L_2) - c_-(L_1, L_2) & \text{else.} \end{cases}$$

Proof. We recall that $\mathcal{F}_{L_i} \simeq k_{N \times \mathbb{R}}$ near $+\infty$ and that $-c_-(L_1, L_2)$ is the infimum on the a such that there exists $u: \mathcal{F}_{L_1} \rightarrow T_{a^*}(\mathcal{F}_{L_2})$ restricting to $\mathrm{id}_{k_{N \times \mathbb{R}}}$ near $+\infty$. Similarly $c_+(L_1, L_2)$ is the infimum on the b such that there exists $v: \mathcal{F}_{L_2} \rightarrow T_{b^*}(\mathcal{F}_{L_1})$ restricting to $\mathrm{id}_{k_{N \times \mathbb{R}}}$ near $+\infty$. We then have $T_{a^*}(v) \circ u = \tau_{0, a+b}(\mathcal{F}_{L_1})$ and $T_{b^*}(u) \circ v = \tau_{0, a+b}(\mathcal{F}_{L_2})$ by Corollary 6.10. Hence u, v satisfy Definition 6.2, except that here a, b may be negative. Assuming moreover $a, b \geq 0$ we find that $\gamma_\tau(\mathcal{F}_{L_1}, \mathcal{F}_{L_2})$ is less than the right hand side of the equality in the statement.

Conversely, if $\gamma_\tau(\mathcal{F}_{L_1}, \mathcal{F}_{L_2}) < c$, there exists a, b, u, v as above with $a + b = c$, $a, b \geq 0$ but u, v only satisfy the condition on the composition.

Near $+\infty$ we only know that they are $u = \lambda_u \text{id}_{k_{N \times \mathbb{R}}}$, $v = \lambda_v \text{id}_{k_{N \times \mathbb{R}}}$ (indeed $\text{Hom}(k_{N \times \mathbb{R}}, k_{N \times \mathbb{R}}) \simeq k$). Since $T_{a^*}(v) \circ u = \tau_{0, a+b}(\mathcal{F}_{L_1})$, we have $\lambda_u \lambda_v = 1$. Hence λ_u, λ_v are not zero and, replacing u, v by a non zero multiple, we can assume that $u = v = \text{id}_{k_{N \times \mathbb{R}}}$ near $+\infty$. We deduce $a \geq \max\{-c_-(L_1, L_2), 0\}$, $b \geq \max\{c_+(L_1, L_2), 0\}$. \square

Remark 6.12. Note that in [Vit22] the metric on $\mathcal{L}(T^*N)$ is defined by $c(L_1, L_2) = \max\{0, c_+(L_1, L_2)\} - \min\{0, c_-(L_1, L_2)\}$. We thus get

Proposition 6.13. *The map $L \mapsto \mathcal{F}_L$ is then an isometric embedding.*

The proof of Proposition 6.11 works exactly the same way for Hamiltonian maps, replacing the condition “ $\mathcal{F}_{L_i} \simeq k_{N \times \mathbb{R}}$ near $+\infty$ ” by “ $\mathcal{K}_\varphi \simeq k_{\Delta_N \times [0, +\infty[}$ near $+\infty$ ”. Since we know that $c_-(\varphi) \leq 0 \leq c_+(\varphi)$ by [Vit92] we obtain the following statement:

Proposition 6.14. *Let φ_1, φ_2 be compactly supported in a Darboux chart Hamiltonian maps of T^*N . Then $\gamma_\tau(\mathcal{K}_{\varphi_1}, \mathcal{K}_{\varphi_2}) = \gamma(\varphi_1, \varphi_2)$.*

We end this section by the useful result that \otimes is Lipschitz for the γ_τ -distance.

Lemma 6.15. *Let M, N, P be manifolds and $\mathcal{K}_1 \in D_{\tau \geq 0}(M \times N \times \mathbb{R})$, $\mathcal{K}_2 \in D_{\tau \geq 0}(N \times P \times \mathbb{R})$. Then, for any $a \in \mathbb{R}$, we have natural isomorphisms $T_{a^*}(\mathcal{K}_1 \otimes \mathcal{K}_2) \simeq T_{a^*}(\mathcal{K}_1) \otimes \mathcal{K}_2 \simeq \mathcal{K}_1 \otimes T_{a^*}(\mathcal{K}_2)$ and, through these isomorphisms, we have the equalities, for any $a \leq b$,*

$$\tau_{a,b}(\mathcal{K}_1 \otimes \mathcal{K}_2) = \tau_{a,b}(\mathcal{K}_1) \otimes \mathcal{K}_2 = \mathcal{K}_1 \otimes \tau_{a,b}(\mathcal{K}_2)$$

Remark 6.16. The notation $\tau_{a,b}(\mathcal{K}_1) \otimes \mathcal{K}_2$ means that we apply the functor $\otimes \mathcal{K}_2$ to the morphism $\tau_{a,b}(\mathcal{K}_1)$. This means that $(\tau_{a,b}(\mathcal{K}_1) \otimes \mathcal{K}_2)$ is a morphism

$$\tau_{a,b}(\mathcal{K}_1) \otimes \mathcal{K}_2 : T_{a^*}(\mathcal{K}_1) \otimes \mathcal{K}_2 \longrightarrow T_{b^*}(\mathcal{K}_1) \otimes \mathcal{K}_2$$

Proof. In the proof the manifolds M, N, P play an auxiliary role and we can consider that our sheaves live on \mathbb{R} to simplify the notations. Hence $\mathcal{K}_1 \otimes \mathcal{K}_2 = R\mathcal{S}!(\mathcal{K}_1 \boxtimes \mathcal{K}_2)$ where $s: \mathbb{R}^2 \rightarrow \mathbb{R}$ is the sum. The isomorphisms then follow from the equalities $T_a \circ s = s \circ (T_a \times \text{id}_{\mathbb{R}}) = s \circ (\text{id}_{\mathbb{R}} \times T_a)$. However to prove the equalities of morphisms we need to be more precise.

Let $(t_1, t_2, \tau_1, \tau_2)$ be the coordinates on $T^*\mathbb{R}^2$. Then the singular support of $\mathcal{K}_1 \boxtimes \mathcal{K}_2$ is contained in $\{\tau_1 \geq 0\} \times \{\tau_2 \geq 0\}$ and, setting $g_u = u\tau_1 + (1-u)\tau_2$, $\mathcal{K}_1 \boxtimes \mathcal{K}_2$ belongs to $D_{g_u \geq 0}(\mathbb{R}^2)$ for any $u \in [0, 1]$. Let us write $\tau_{a,b}^u$ for the morphism $\tau_{a,b}$ induced by g_u . Then $\tau_{a,b}^1$ gives $\tau_{a,b}(\mathcal{K}_1) \otimes \mathcal{K}_2$ by applying $R\mathcal{S}!$ and $\tau_{a,b}^0$ gives $\mathcal{K}_1 \otimes \tau_{a,b}(\mathcal{K}_2)$. It remains to see that $\tau_{a,b}^u$

gives the same morphism for all u . We define $\mathcal{K} = \mathcal{K}_1 \boxtimes \mathcal{K}_2 \boxtimes k_{[0,1]} \in D(\mathbb{R}^2 \times [0, 1])$ and $g = u\tau_1 + (1 - u)\tau_2$. Then $K \in D_{g \geq 0}(\mathbb{R}^2 \times [0, 1])$ (here we are cheating a little bit because the singular support is not defined on a manifold with boundary – however we can replace $[0, 1]$ by $] -\varepsilon, 1 + \varepsilon[$ and u by a smooth function with values in $[0, 1]$ which is 0 on $] -\varepsilon, 0[$ and 1 on $[1, 1 + \varepsilon[$). We thus obtain a morphism $\tau_{a,b}^g: T'_{a*}(\mathcal{K}) \rightarrow T'_{b*}(\mathcal{K})$ where $T'_c(t_1, t_2, u) = (t_1 + uc, t_2 + (1 - u)c, u)$. Let $s': \mathbb{R}^2 \times [0, 1] \rightarrow \mathbb{R} \times [0, 1]$ be the sum of the first two variables. Then $s' \circ T'_c = T'_c \times \text{id}_{[0,1]}$ and $\tau_{a,b}^g$ induces $\tau'_{a,b}: (T_{a*}(\mathcal{K}_1 \otimes \mathcal{K}_2)) \boxtimes k_{[0,1]} \rightarrow (T_{b*}(\mathcal{K}_1 \otimes \mathcal{K}_2)) \boxtimes k_{[0,1]}$ which restricts to $\tau_{a,b}^s$ along $\mathbb{R} \times \{s\}$. Now restricting along $\mathbb{R} \times \{s\}$ always induces the same isomorphism $\text{Hom}(\mathcal{F} \boxtimes k_{[0,1]}, \mathcal{G} \boxtimes k_{[0,1]}) \xrightarrow{\sim} \text{Hom}(\mathcal{F}, \mathcal{G})$, for all \mathcal{F}, \mathcal{G} , namely, the inverse of the isomorphism induced by the pull-back along the projection to $[0, 1]$. This concludes the proof. \square

We deduce an analogue of [PS21], theorem 2.4.1, in our situation.

Lemma 6.17. *Let $\mathcal{K}_1, \mathcal{K}_2 \in D_{\tau \geq 0}(M \times N \times \mathbb{R})$ and $\mathcal{F} \in D_{\tau \geq 0}(M \times \mathbb{R})$. Then $\mathcal{K}_j \otimes \mathcal{F} \in D_{\tau \geq 0}(N \times \mathbb{R})$ and we have*

$$\gamma_{\tau}(\mathcal{K}_1^{\otimes}(\mathcal{F}), \mathcal{K}_2^{\otimes}(\mathcal{F})) \leq \gamma_{\tau}(\mathcal{K}_1, \mathcal{K}_2)$$

Proof. We have $\gamma_{\tau}(\mathcal{K}_1, \mathcal{K}_2) < c$, for a given c , if and only if there exist $a, b \geq 0$ with $a + b = c$ and morphisms $u: \mathcal{K}_1 \rightarrow T_{a*}(\mathcal{K}_2)$, $v: \mathcal{K}_2 \rightarrow T_{b*}(\mathcal{K}_1)$ satisfying the conditions in Definition 6.2. The morphisms induce $u \circ \text{id}_{\mathcal{F}}$, $v \circ \text{id}_{\mathcal{F}}$ which also satisfies the conditions of the definition by Lemma 6.15. \square

6.3. γ_g -limits and colimits. In this paragraph we shall prove that any sequence $(\mathcal{F}_n)_{n \in \mathbb{N}}$ which γ_g -converges to \mathcal{F} can be turned into an inductive system whose homotopy colimit is \mathcal{F} . We recall the notion of homotopy colimit in $D(N)$ (see [BN93, Def. 2.1])

Definition 6.18. *Let $(\mathcal{F}_n, f_n)_{n \in \mathbb{N}}$, with $f_n: \mathcal{F}_n \rightarrow \mathcal{F}_{n+1}$, be an inductive system in $D(N)$; then $\text{hocolim}_n \mathcal{F}_n$ is the cone of the morphism*

$$(\text{id} - f): \bigoplus_{n \geq 0} \mathcal{F}_n \rightarrow \bigoplus_{n \geq 0} \mathcal{F}_n$$

where $f = \bigoplus_{n \geq 0} f_n$.

We thus have a distinguished triangle

$$(6.3) \quad \bigoplus_{n \geq 0} \mathcal{F}_n \xrightarrow{\text{id} - f} \bigoplus_{n \geq 0} \mathcal{F}_n \xrightarrow{p} \text{hocolim}_n \mathcal{F}_n \xrightarrow{+1}$$

However, like any cone, $\text{hocolim}_n \mathcal{F}_n$ is only defined up to (non unique) isomorphism and the morphism p itself is not uniquely defined. For a given triangle (6.3) and for any $n_0 \in \mathbb{N}$, we have a morphism $i_{n_0}: \mathcal{F}_{n_0} \rightarrow$

$\text{hocolim}_n \mathcal{F}_n$ defined as $i_{n_0} = p \circ j_{n_0}$ where j_{n_0} is the obvious morphism to the sum. For another choice of triangle, with p' instead of p , there exists an automorphism ψ of $\text{hocolim}_n \mathcal{F}_n$ such that $p' = \psi \circ p$. Then $i'_{n_0} = p' \circ j_{n_0}$ is “conjugate” to i_{n_0} in the sense $i'_{n_0} = \psi \circ i_{n_0}$.

We recall the “nine diagram” from [BBD82, Prop. 1.1.11] (see [KS06, Ex. 10.6] where it is stated precisely which arrows are given and which ones are obtained), a variant of the octaedron axiom. We assume to be given a commutative square as on the top left corner of the diagram below and four distinguished triangles (two vertical and two horizontal) given by the unmarked solid arrows. The arrows marked [1] are just translations of already defined solid arrows.

The proposition claims that the dotted arrows can be chosen to make all squares commutative, except for the one with Z^2 at its top left corner, which is anticommutative, and the new line and the new column are distinguished triangles.

$$\begin{array}{ccccccc}
 X^0 & \longrightarrow & X^1 & \longrightarrow & X^2 & \longrightarrow & X^0[1] \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow [1] \\
 Y^0 & \longrightarrow & Y^1 & \longrightarrow & Y^2 & \longrightarrow & Y^0[1] \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow [1] \\
 Z^0 & \cdots\cdots\cdots & Z^1 & \cdots\cdots\cdots & Z^2 & \cdots\cdots\cdots & Z^0[1] \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \ominus -[1] \\
 X^0[1] & \xrightarrow{[1]} & X^1[1] & \xrightarrow{[1]} & X^2[1] & \xrightarrow{-[1]} & X^0[2]
 \end{array}$$

Remark 6.19. Using the nine diagram built on the square

$$\begin{array}{ccc}
 \bigoplus_{n \geq n_0} \mathcal{F}_n & \xrightarrow{\text{id}-f} & \bigoplus_{n \geq n_0} \mathcal{F}_n \\
 \downarrow & & \downarrow \\
 \bigoplus_{n \geq 0} \mathcal{F}_n & \xrightarrow{\text{id}-f} & \bigoplus_{n \geq 0} \mathcal{F}_n
 \end{array}$$

we can check that $\text{hocolim}_{n \geq 0} \mathcal{F}_n \xrightarrow{\sim} \text{hocolim}_{n \geq n_0} \mathcal{F}_n$ is indeed an isomorphism. With the above notations, we take $X^0 = X^1 = \bigoplus_{n \geq n_0} \mathcal{F}_n$, $Y^0 = Y^1 = \bigoplus_{n \geq 0} \mathcal{F}_n$, $Z^0 = Z^1 = \bigoplus_{0 \leq n \leq n_0-1} \mathcal{F}_n$. The map from Z^0 to Z^1 is also equal to $\text{id} - f$, but as is an upper triangular finite-dimensional matrix with Id on the diagonal, so it is invertible. This implies $Z^2 = 0$. As a result the vertical arrow from X^2 to Y^2 is an isomorphism (in $D(N)$).

A more general statement of the following Lemma can be found in Lemma 13.33.4 of [Stacks, Section 0A5K]. It tells us, as we might expect, that the limit of a subsequence coincides with the limit of the original sequence.

Lemma 6.20. *Let $(\mathcal{F}_n, f_n)_{n \in \mathbb{N}}$ be an inductive system in $D(N)$ and $\sigma: \mathbb{N} \rightarrow \mathbb{N}$ an increasing map. Then the compositions of the f_n 's define an inductive system $(\mathcal{F}_{\sigma(n)}, f'_n)$ and there exists an isomorphism $j: \text{hocolim}_n \mathcal{F}_{\sigma(n)} \xrightarrow{\sim} \text{hocolim}_n \mathcal{F}_n$ such that $j \circ i'_{\sigma(n)} = i_{\sigma(n)}$, where $i'_{\sigma(n)}$ is the morphism i for $(\mathcal{F}_{\sigma(n)})$.*

A family of morphisms $u_n: \mathcal{F}_n \rightarrow \mathcal{G}$ such that $u_{n+1} \circ f_n = u_n$ induces a morphism $u: \text{hocolim}_n \mathcal{F}_n \rightarrow \mathcal{G}$ such that $u \circ i_n = u_n$ for all n . Indeed we have $\bigoplus_n u_n \circ (\text{id} - f) = 0$, hence $\bigoplus_n u_n$ factorizes as $\bigoplus_n u_n = u \circ p$. Again this morphism u is not uniquely defined.

Lemma 6.21. *Let $(\mathcal{F}_n, f_n)_{n \in \mathbb{N}}$, with $f_n: \mathcal{F}_n \rightarrow \mathcal{F}_{n+1}$, be an inductive system in $D(N)$. We let $\mathcal{C}(f_n)$ be the cone of f_n and we assume that the sum $\sum_n \gamma_g(\mathcal{C}(f_n), 0)$ is finite. Then the sequence $(\mathcal{F}_n)_{n \in \mathbb{N}}$ γ_g -converges to $\text{hocolim}_k \mathcal{F}_k$. More precisely, if we let $i_n: \mathcal{F}_n \rightarrow \text{hocolim}_k \mathcal{F}_k$ be the morphism defined after (6.3), we have $\gamma_g(\mathcal{C}(i_n), 0) \leq 2 \sum_{k=n}^{\infty} \gamma_g(\mathcal{C}(f_k), 0)$.*

Proof. Let n_0 be given. We set for short $\mathcal{S} = \bigoplus_{n \geq n_0} \mathcal{F}_n$, $\mathcal{C} = \text{hocolim}_n \mathcal{F}_n$. We checked that \mathcal{C} is the cone of $u = \text{id} - \bigoplus_{n \geq n_0} f_n: \mathcal{S} \rightarrow \mathcal{S}$. We also consider the similar sum but associated to the constant sequence $\mathcal{G}_n = \mathcal{F}_{n_0}$ for all $n \geq n_0$ and $\mathcal{T} = \bigoplus_{n \geq n_0} \mathcal{G}_n$. We define $w: \mathcal{T} \rightarrow \mathcal{T}$ as $\text{id} - s$ where $s_n: \mathcal{G}_n \rightarrow \mathcal{G}_{n+1}$ is induced by $\text{Id}_{\mathcal{F}_{n_0}}$. The cone of w is \mathcal{F}_{n_0} . We let $f_n^m: \mathcal{F}_n \rightarrow \mathcal{F}_m$ be the composition of the f_k 's and we define $F = \bigoplus_{n \geq n_0} f_{n_0}^n: \mathcal{T} \rightarrow \mathcal{S}$. Then $F \circ w = u \circ F$ and the nine diagram recalled above gives the commutative (except for one anticommuting square) diagram of distinguished triangles

$$\begin{array}{ccccccc}
\mathcal{T} & \xrightarrow{w} & \mathcal{T} & \xrightarrow{q} & \mathcal{F}_{n_0} & \xrightarrow{+1} & \\
F \downarrow & & F \downarrow & & \downarrow j_{n_0} & & \\
\mathcal{S} & \xrightarrow{u} & \mathcal{S} & \xrightarrow{p} & \mathcal{C} & \xrightarrow{+1} & \\
\downarrow & & \downarrow & & \downarrow & & \\
\mathcal{S}' & \longrightarrow & \mathcal{S}' & \longrightarrow & \mathcal{C}(j_{n_0}) & \xrightarrow{+1} & \\
\downarrow +1 & & \downarrow +1 & & \downarrow +1 & &
\end{array}$$

where $\mathcal{S}' = \bigoplus_{n \geq n_0} \mathcal{C}(f_{n_0}^n)$, j_{n_0} is conjugate to i_{n_0} (unfortunately we cannot say they are equal because of the ambiguity in the definition of the cone) and $\mathcal{C}(j_{n_0})$ is the cone of j_{n_0} .

We set $r: \mathcal{F}_{n_0} \rightarrow \mathcal{T}$ be the inclusion of the first factor. The morphisms r and w give a splitting $\mathcal{T} \simeq \mathcal{F}_{n_0} \oplus \mathcal{T}$ and it follows that $q \circ r$ is an isomorphism. Since i_{n_0} is "conjugate" to $p \circ F \circ r$ (see after (6.3)), it is also conjugate to j_{n_0} and $\mathcal{C}(j_{n_0}) \simeq \mathcal{C}(i_{n_0})$.

Now we claim that $\gamma_g(\mathcal{C}(f_n^m), 0) \leq \sum_{k=n}^{m-1} \gamma_g(\mathcal{C}(f_k), 0)$. We prove this by induction on $m-n$. When $m = n+1$ this is clear. Since $f_n^{m+1} = f_m \circ f_n^m$, we have, using Lemma 6.4 (6), $\gamma_g(\mathcal{C}(f_n^{m+1}), 0) \leq \gamma_g(\mathcal{C}(f_n^m), 0) + \gamma_g(\mathcal{C}(f_m), 0)$.

In particular $\gamma_g(\mathcal{C}(f_{n_0}^n), 0) \leq \varepsilon_{n_0} := \sum_{k=n_0}^{\infty} \gamma_g(\mathcal{C}(f_k), 0)$, for all $n \geq n_0$, and then $\gamma_g(\mathcal{S}', 0) \leq \varepsilon_{n_0}$. By the triangle in the last row of the above diagram and Lemma 6.4 (5) we deduce $\gamma_g(\mathcal{C}(j_{n_0}), 0) \leq 2\varepsilon_{n_0}$, as required. \square

Proposition 6.22. *Any Cauchy sequence in $D_{g \geq 0}(N)$ with respect to γ_g is convergent.*

Remark 6.23. The reader should be careful : the distance γ_g is degenerate, so the limit is by no means unique. However we shall see that the restriction of γ_g to the set of limits of constructible sheaves (i.e. in $D_{lc}(N)$) the distance is non-degenerate (see Appendix B).

Proof. Let $(\mathcal{F}_n)_{n \in \mathbb{N}}$ be a Cauchy sequence. Up to taking a subsequence we can assume that $\gamma_g(\mathcal{F}_n, \mathcal{F}_m) \leq 2^{-n}$ for any $n \leq m$. In particular, by Lemma 6.8, there exist $f_n: \mathcal{F}_n \rightarrow K_\varphi^{2^{-n}}(\mathcal{F}_{n+1})$ such that $\gamma_g(\mathcal{C}(f_n), 0) \leq 2^{-n+1}$. We then set $\varepsilon_n = \sum_{k \geq n} 2^{-k}$ and $\mathcal{F}'_n = K_\varphi^{-\varepsilon_n}(\mathcal{F}_n)$ so f_n induces a map $f'_n: \mathcal{F}'_n \rightarrow \mathcal{F}'_{n+1}$ and according to Lemma 6.21, $\mathcal{F}'_n \gamma_g$ converges to $\text{hocolim } \mathcal{F}'_n$. Since $\gamma_g(\mathcal{F}_n, \mathcal{F}'_n) \leq \varepsilon_n$ we have that $(\mathcal{F}_n)_{n \geq 1}$ has the same limit. \square

We just proved that to a Cauchy sequence we can associate a limit that is a homotopy colimit. In the next Proposition we prove that conversely any γ_g -limit of a sequence $(\mathcal{F}_n)_{n \in \mathbb{N}}$ is a homotopy colimit of some (slightly modified) subsequence. We first notice an easy consequence of Lemma 6.21.

Lemma 6.24. *Let $\mathcal{F} \in D(N)$ and let $\varepsilon_n > 0$ be a sequence decreasing to 0. Then we have an inductive system $(K_\varphi^{-\varepsilon_n}(\mathcal{F}), \tau_{-\varepsilon_n, -\varepsilon_{n+1}}(\mathcal{F}))$ and the maps $\tau_{-\varepsilon_n, 0}(\mathcal{F})$ induce an isomorphism $\text{hocolim } K_\varphi^{-\varepsilon_n}(\mathcal{F}) \xrightarrow{\sim} \mathcal{F}$.*

Proof. Let $\tau: \text{hocolim } K_\varphi^{-\varepsilon_n}(\mathcal{F}) \rightarrow \mathcal{F}$ be the morphism induced by the maps $\tau_{-\varepsilon_n, 0}(\mathcal{F})$. With the notation i_n of Lemma 6.21 we have $\tau \circ i_n = \tau_{-\varepsilon_n, 0}(\mathcal{F})$. By the octaedron axiom there exists a distinguished triangle relating the cones of these maps and Lemma 6.4 implies that $\gamma_g(0, \mathcal{C}(\tau)) \leq \gamma_g(0, \mathcal{C}(i_n)) + \gamma_g(0, \mathcal{C}(\tau_{-\varepsilon_n, 0}(\mathcal{F})))$. By Lemma 6.8 $\gamma_g(0, \mathcal{C}(\tau_{-\varepsilon_n, -\varepsilon_{n+1}}(\mathcal{F}))) \leq 2(\varepsilon_n - \varepsilon_{n+1})$ and we can apply Lemma 6.21. Hence $\gamma_g(0, \mathcal{C}(i_n))$ goes to 0 and $\gamma_g(0, \mathcal{C}(\tau))$ is as small as desired. So $\mathcal{C}(\tau) \simeq 0$ and τ is an isomorphism. \square

Proposition 6.25. *Let $(\mathcal{F}_n)_{n \in \mathbb{N}}$ be a sequence in $D(N)$ which γ_g -converges to \mathcal{F} . Then, up to taking a subsequence, there exist a sequence of positive numbers $(\varepsilon_n)_{n \in \mathbb{N}}$ converging to 0 and morphisms $f_n: K_\varphi^{-\varepsilon_n}(\mathcal{F}_n) \rightarrow$*

$K_\varphi^{-\varepsilon_{n+1}}(\mathcal{F}_{n+1})$, $u_n: K_\varphi^{-\varepsilon_n}(\mathcal{F}_n) \rightarrow \mathcal{F}$ such that $u_{n+1} \circ f_n = u_n$, for all n , and the morphism $\text{hocolim } K_\varphi^{-\varepsilon_n}(\mathcal{F}_n) \rightarrow \mathcal{F}$ induced by the u_n 's (see after (6.3)) is an isomorphism.

Proof. We set $\alpha_n = 2^{-n}$. Up to taking a subsequence we can assume that $\gamma_g(\mathcal{F}_n, \mathcal{F}) \leq \alpha_{n+1}$ for each n . This implies the existence of $0 \leq \beta_n \leq \alpha_{n+1}$ and morphisms $v_n: \mathcal{F} \rightarrow K_\varphi^{\beta_n}(\mathcal{F}_n)$, $w_n: K_\varphi^{\beta_n}(\mathcal{F}_n) \rightarrow K_\varphi^{\alpha_{n+1}}(\mathcal{F})$ such that $w_n \circ v_n = \tau_{0, \alpha_{n+1}}(\mathcal{F})$. Translating these morphisms by $K_\varphi^*(-)$ and alternating the \mathcal{F}_n and the $K_\varphi^{-\alpha_n}(\mathcal{F})$ we obtain an inductive system

$$\begin{aligned} \dots \rightarrow K_\varphi^{-\alpha_n}(\mathcal{F}) &\xrightarrow{K_\varphi^{-\alpha_n}(v_n)} K_\varphi^{-\varepsilon_n}(\mathcal{F}_n) \xrightarrow{K_\varphi^{-\alpha_n}(w_n)} K_\varphi^{-\alpha_{n+1}}(\mathcal{F}) \\ &\xrightarrow{K_\varphi^{-\alpha_{n+1}}(v_{n+1})} K_\varphi^{-\varepsilon_{n+1}}(\mathcal{F}_{n+1}) \xrightarrow{K_\varphi^{-\alpha_{n+1}}(w_{n+1})} K_\varphi^{-\alpha_{n+2}}(\mathcal{F}) \rightarrow \dots \end{aligned}$$

where $\varepsilon_n = \alpha_n - \beta_n$. We define $f_n = K_\varphi^{-\alpha_{n+1}}(v_{n+1}) \circ K_\varphi^{-\alpha_n}(w_n)$ and $u_n = \tau_{-\alpha_{n+1}, 0}(\mathcal{F}) \circ K_\varphi^{-\alpha_n}(w_n)$. By Lemma 6.20 we have $\text{hocolim } K_\varphi^{-\varepsilon_n}(\mathcal{F}_n) \simeq \text{hocolim } K_\varphi^{-\alpha_n}(\mathcal{F})$ and by Lemma 6.24 we have $\text{hocolim } K_\varphi^{-\alpha_n}(\mathcal{F}) \xrightarrow{\sim} \mathcal{F}$, which proves the result. \square

We now use the previous results to bound the microsupport of a γ_g -limit. For a sequence $(X_j)_{j \geq 0}$ of subsets of a topological space Z , its topological upper and lower limits are defined as

$$\begin{aligned} \limsup_j X_j &= \bigcap_n \overline{\bigcup_{j \geq n} X_j} \\ &= \left\{ x \in Z \mid \exists (x_j)_{j \geq 1}, x_j \in X_j \text{ for infinitely many } j, \lim_j x_j = x \right\} \\ \liminf_j X_j &= \left\{ x \in Z \mid \exists (x_j)_{j \geq 1}, x_j \in X_j, \lim_j x_j = x \right\} \end{aligned}$$

Obviously $\liminf_j X_j \subset \limsup_j X_j$. We have

Proposition 6.26. *Let N be a manifold and let $g: T^*N \setminus 0_N \rightarrow \mathbb{R}$ be a 1-homogeneous autonomous Hamiltonian function. If $(\mathcal{F}_j)_{j \geq 1}$ is a sequence such that $\gamma_g - \lim \mathcal{F}_j = \mathcal{F}$ then we have*

$$\text{SS}(\mathcal{F}) \subset \liminf_j \text{SS}(\mathcal{F}_j)$$

Proof. By Proposition 6.25, up to taking a subsequence, we can find some sequence (ε_j) converging to 0, such that $\text{hocolim } \mathcal{F}'_j \xrightarrow{\sim} \mathcal{F}$, where $\mathcal{F}'_j = K_{\varphi_g}^{\varepsilon_j}(\mathcal{F}_j)$. We have $\text{SS}(\mathcal{F}'_j) = \varphi_g^{\varepsilon_j}(\text{SS}(\mathcal{F}_j))$ and $\varphi_g^{\varepsilon_j}$ tends to id in C^0 norm over compact subsets. Hence $\liminf_j \text{SS}(\mathcal{F}_j) = \liminf_j \text{SS}(\mathcal{F}'_j)$. The homotopy colimit is defined by a distinguished triangle where the other two terms are $\bigoplus_{n \geq n_0} \mathcal{F}'_n$ (we can indeed restrict to $\bigoplus_{n \geq n_0}$ by Remark 6.19). By

the triangle inequality for singular supports and Proposition 3.4, this implies that $SS(\mathcal{F}) \subset \limsup_j SS(\mathcal{F}'_j)$. Now let $p = (x, \xi) \notin \liminf_j SS(\mathcal{F}'_j)$, we can find a subsequence $\sigma(j)$ such that $p \notin \limsup_j SS(\mathcal{F}'_{\sigma(j)})$; but $\mathcal{F}'_{\sigma(j)}$ has limit \mathcal{F} , so $p \notin SS(\mathcal{F})$. As a result $SS(\mathcal{F}) \subset \liminf_j SS(\mathcal{F}'_j)$. \square

7. PROOF OF THEOREM 1.2

Let $\mathcal{F} \in D(N)$. We want to prove that around a point $z_0 \in SS(\mathcal{F})$ there is no sequence $(\varphi_j)_{j \geq 1}$ in $\mathcal{D}\mathcal{H}\text{am}_c(B(z_0, \varepsilon))$ such that $\gamma\text{-}\lim_j \varphi_j = \text{Id}$ and $\varphi_j(SS(\mathcal{F})) \cap B(z_0, \eta) = \emptyset$ for some $\varepsilon > \eta > 0$. We argue by contradiction and assume the existence of such a sequence $(\varphi_j)_{j \geq 1}$ and open ball $B(z_0, \eta)$. Let \mathcal{K}_{φ_j} be the sheaf in $D^b(N \times N \times \mathbb{R})$ inducing φ_j as recalled in §4. We denote by $T_{\tau > 0}^*(N \times \mathbb{R})$ the subset of $T^*(N \times \mathbb{R})$ given by $\tau > 0$ and we set $\rho: T_{\tau > 0}^*(N \times \mathbb{R}) \rightarrow T^*N$, $(x, t, \xi, \tau) \mapsto (x, \xi/\tau)$. Then $\rho(SS(\mathcal{K}_{\varphi_j}^{\otimes}(\mathcal{F})) \cap T_{\tau > 0}^*(N \times \mathbb{R})) = \varphi_j(SS(\mathcal{F}))$ does not intersect $B(z_0, \eta)$. We want to prove that this implies that $SS(\mathcal{F}) \cap B(z_0, \eta) = \emptyset$.

We will use the following proposition.

Proposition 7.1. *Let $(\varphi_j)_{j \geq 1}$ be a sequence in $\mathcal{D}\mathcal{H}\text{am}_c(T^*N)$ such that $\gamma\text{-}\lim \varphi_j = \text{Id}$ and $\mathcal{K}_{\varphi_j}^{\otimes}$ its quantization in $D^b(N \times N \times \mathbb{R})$. Then for all $\mathcal{F} \in D^b(N)$ we have*

$$\gamma_\tau\text{-}\lim \mathcal{K}_{\varphi_j}^{\otimes}(\mathcal{F}) = \mathcal{K}_{\text{Id}}^{\otimes}(\mathcal{F}) = \mathcal{F} \boxtimes k_{[0, +\infty[} \in D^b(N \times \mathbb{R})$$

Proof. This follows from Lemma 6.17 and Proposition 6.14. \square

Remark 7.2. Note that $\varphi_j(SS(\mathcal{F}))$ is not a homogeneous manifold, and $SS(\mathcal{K}_{\varphi_j}^{\otimes}(\mathcal{F})) \subset T^*(N \times \mathbb{R})$ represents the homogenization of $\varphi_j(SS(\mathcal{F}))$.

We now claim that $SS(\mathcal{F} \boxtimes k_{[0, +\infty[}) \cap \{\tau > 0\} = SS(\mathcal{F}) \times (\{0\} \times \{\tau > 0\})$. Indeed we have the general bound $SS(\mathcal{F} \boxtimes k_{[0, +\infty[}) \subset SS(\mathcal{F}) \times SS(k_{[0, +\infty[})$ and a similar bound for $\mathcal{F} \boxtimes k_{]-\infty, 0]}$. Now we have the distinguished triangle $\mathcal{F} \boxtimes k_{\mathbb{R}} \rightarrow \mathcal{F} \boxtimes (k_{]-\infty, 0]} \oplus k_{[0, +\infty[}) \rightarrow \mathcal{F} \boxtimes k_{\{0\}} \xrightarrow{+1}$ and the equalities $SS(\mathcal{F} \boxtimes k_{\{0\}}) = SS(\mathcal{F}) \times (\{0\} \times \mathbb{R})$, $SS(\mathcal{F} \boxtimes k_{\mathbb{R}}) = SS(\mathcal{F}) \times (\mathbb{R} \times \{0\})$, by [KS90, Prop. 5.4.4, 5.4.5]. The triangular inequality for the microsupport then gives the claim.

We return to the sequence $(\varphi_j)_{j \geq 1}$ and the ball $B(z_0, \eta)$. Proposition 7.1 and Proposition 6.26 (applied with $N \times \mathbb{R}$ and $g = \tau$) give

$$SS(\mathcal{F}) \times (\{0\} \times \{\tau > 0\}) \subset SS(\mathcal{K}_{\text{Id}}^{\otimes}(\mathcal{F})) \subset \liminf_j SS(\mathcal{K}_{\varphi_j}^{\otimes}(\mathcal{F}))$$

Applying the map ρ we deduce $SS(\mathcal{F}) \subset \liminf_j \varphi_j(SS(\mathcal{F}))$. In particular $SS(\mathcal{F}) \cap B(z_0, \eta) = \emptyset$ which contradicts $z_0 \in SS(\mathcal{F})$.

8. γ -COISOTROPIC VS CONE-COISOTROPIC

8.1. A γ -coisotropic set is cone-coisotropic. Our goal in this section is to prove Proposition 1.3.

Following Bouligand (see [Bou32]), one may define for a point x of a subset V in a smooth manifold, two cones :

Definition 8.1. *Let V be a closed subset of the C^1 -manifold M and $x \in V$. We choose a C^1 chart of M at x and define*

(1) *The **paratingent cone** of V at x is*

$$C^+(x, V) = \left\{ \lim_n c_n(x_n - y_n) \mid x_n, y_n \in V, c_n \in \mathbb{R}, \lim_n x_n = \lim_n y_n = x, \lim_n c_n = +\infty \right\}$$

(2) *The **contingent cone** of V at x is*

$$C^-(x, V) = \left\{ \lim_n c_n(x_n - x) \mid x_n \in V, c_n \in \mathbb{R}, \lim_n x_n = x, \lim_n c_n = +\infty \right\}$$

Clearly $C^-(x, V) \subset C^+(x, V) \subset T_x V$. Note that $C^+(x, V)$ is invariant by $v \mapsto -v$, while it is not necessarily the case for $C^-(x, V)$.

We shall say that a cone C is closed (resp. open) if $C \cap S(x, 1)$ is closed (resp. open). Note that a closed cone is closed in the usual sense, while an open cone is not open (but becomes open if we remove x). Then the contingent cone is closed. It is also easy to see that both cones are independent from the choice of the C^1 chart. We then have the following definition, which appears under the name ‘‘involutivity’’ in the book of Kashiwara and Schapira

Definition 8.2 (Cone-coisotropic, see [KS90], definition 6.5.1 p. 271). *We shall say that V is **cone-coisotropic** if whenever a hyperplane H is such that $C^+(x, V) \subset H$ then the symplectic orthogonal of H , H^ω is contained in $C^-(x, V)$.*

Proof of Proposition 1.3. We want to prove that if V is not cone-coisotropic at $z \in V$ then it is not γ -coisotropic at z . Using a chart, we may assume $V \subset \mathbb{R}^d$ and $z = 0$. For H a hyperplane we set n_H to be the normal to H (for the usual scalar product). Let us assume that $C^+(0, V) \subset H$ and $H^\omega \not\subset C^-(0, V)$. We must prove that for all $\varepsilon > 0$ there exists a ball $B(0, \eta)$ such that for all $\delta > 0$ we can find $\varphi \in \mathcal{D}\mathfrak{H}am_c(B(x, \varepsilon))$ such that $\varphi(V) \cap B(0, \eta) = \emptyset$ and $\gamma(\varphi) < \delta$. We first prove

Lemma 8.3. *Let V be a closed set in \mathbb{R}^d containing 0 and H a hyperplane such that $C^+(0, V) \subset H$. Then there exists a continuous function $f_{H, V}$ defined in a neighbourhood of 0 in H such that $f_{H, V}(0) = 0$ and*




FIGURE 2. Coisotropic cone on the left, non-coisotropic on the right

$d_G f_{H,V}(0) = 0$, where d_G is the Gâteaux derivative, and for any open cone $C_\rho^-(0, V) := \{\lambda x; \lambda \geq 0, x \in S^{d-1}, d(x, C^-(0, V)) < \rho\}$, we have for δ small enough

$$V \cap B(0, \delta) \subset \left\{ x + f_{H,V}(x)n_H \mid x \in C_\rho^-(0, V) \right\}$$

Proof. For $x \in H$ let $V_x = V \cap (x + \mathbb{R}n_H)$. We claim that for x close enough to 0, V_x contains at most one point. Indeed, if this was not the case, we would have sequences $(y_n)_{n \geq 1}, (w_n)_{n \geq 1}$ converging to 0 such that $y_n - w_n = t_n n_H$ with $t_n \neq 0$. But then by definition of $C^+(0, V)$, we would have $n_H \in C^+(0, V)$ a contradiction.

As a result, on the closed set $\Gamma \subset H$ defined as the set of x such that V_x is non empty, we have a function f such that

$$x \in \Gamma \iff x + f(x)n_H \in V$$

Let us prove f is continuous on Γ , at least in a neighbourhood of 0. If this was not the case, there would be a sequence $(z^n)_{n \geq 1}$ converging to 0 and for each n two sequences $(x_k^n)_{k \geq 1}, (y_k^n)_{k \geq 1}$ such that $\lim_k x_k^n = \lim_k y_k^n = z^n$ and $\lim_k f(x_k^n) \neq \lim_k f(y_k^n)$. Setting $u_k^n = x_k^n + f(x_k^n)n_H, v_k^n = y_k^n + f(y_k^n)n_H$ we have

$$u_k^n - v_k^n = (x_k^n - y_k^n) + (f(x_k^n) - f(y_k^n))n_H$$

which can be normalized so that, for any n , it has a limit $(0, 1)$ when $k \rightarrow +\infty$. As a result for $k = k(n)$ large enough, setting $u_n = u_{k(n)}^n, v_n = v_{k(n)}^n$, there is a τ_n so that

$$\|\tau_n(u_n - v_n) - (0, 1)\| < 1/3$$

We can then find a subsequence of $\tau_n(u_n - v_n)$ converging to a vector w which does not belong to H . Since $w \in C^+(0, V)$ we have a contradiction. From now on we replace Γ by $\Gamma \cap B(0, r)$ with r small enough so that f is now continuous on Γ .

To show that $\Gamma \subset C_\rho^-(0, V)$ we first show that $\lim_{\substack{|x| \rightarrow 0 \\ x \in \Gamma}} \frac{f(x)}{|x|} = 0$. Indeed, if

we had a sequence $(x_k)_{k \geq 1}$ in Γ converging to 0 and such that $\lim_k \frac{f(x_k)}{|x_k|} \neq 0$, the sequence of vectors $\frac{1}{|x_k|}(x_k + f(x_k)n_H)$ would have a subsequence with a limit which does not belong to H , so $C^+(0, V) \not\subset H$, a contradiction. To prove that $\Gamma \subset C_\rho^-(0, V)$ we argue by contradiction, and assume we have a sequence $x_n \in \Gamma$ such that $\lim_n x_n = 0$ and $w = \lim_n \frac{x_n}{|x_n|} \notin C^-(0, V)$. Then there exists a sequence $x_n + f(x_n)n_H$ converging to 0 and belonging to V . As a result $\lim_n \frac{x_n + f(x_n)n_H}{|x_n|} = \lim_n \frac{x_n}{|x_n|} = w \notin C^-(0, V)$ a contradiction.

We now will extend f to $f_{H,V}$ such that

- (1) $f_{H,V}(x) = f(x)$ for $x \in \Gamma$
- (2) $f_{H,V}$ is continuous
- (3) $d_G(f_{H,V}(0)) = 0$

For this we can assume for simplicity that $r = 1$ and write $x = \rho \cdot \theta$ with $0 \leq \rho \leq 1, \theta \in S^{d-1}$. We write $g(s, \theta) = f(e^s \cdot \theta)$ with $s \in]-\infty, 0]$ and identify Γ with its image in $]-\infty, 0] \times S^{d-1}$. Now let $g_n(s, \theta)$ be a continuous function defined on $C_n =]-n-1, -n+1[\times S^{d-1}$ for $n \in \mathbb{N}$, coinciding with g on $\Gamma \cap C_n$ and having the same bound, i.e.

$$\sup \{g_n(s, \theta) \mid (s, \theta) \in C_n\} = \sup \{g(s, \theta) \mid (s, \theta) \in \Gamma \cap C_n\}$$

$$\inf \{g_n(s, \theta) \mid (s, \theta) \in C_n\} = \inf \{g(s, \theta) \mid (s, \theta) \in \Gamma \cap C_n\}$$

Such a function exists by Tietze's extension theorem.

Now let χ_n be a partition of unity subordinated to the covering of $]-\infty, 0] \times S^{d-1}$ by the C_n . We set

$$G(s, \theta) = \sum_{n \geq 0} \chi_n(s) g_n(s, \theta)$$

Note that this is well defined since whenever $g_n(s, \theta)$ is not defined, we have $\chi_n(s, \theta) = 0$, so we set $\chi_n(s) g_n(s, \theta) = 0$ outside of C_n . And since whenever $g_n(s, \theta)$ is defined and $(s, \theta) \in \Gamma$ we have $g_n(s, \theta) = g(s, \theta)$, we see that G is an extension of g . We now set

$$f_{H,V}(e^s \cdot \theta) = G(s, \theta)$$

Properties (1) and (2) are then obvious. As for the third one, we notice that

$$\begin{aligned} \sup \{f_{H,V}(x) \mid |x| \leq e^{-n}\} &= \sup \{G(s, \theta) \mid (s, \theta) \mid s \leq -n\} \leq \\ &\sup_{k \geq n} \sup \{g_k(s, \theta) \mid (s, \theta) \in C_k\} \leq \sup \{g(s, \theta) \mid (s, \theta) \in \Gamma, s \leq -n\} = \\ &\sup \{f(x) \mid x \in \Gamma, |x| \leq e^{-n}\} \end{aligned}$$

We see that since $\lim_{\substack{|x| \rightarrow 0 \\ x \in \Gamma}} \frac{f(x)}{|x|} = 0$ we have $\lim_{x \rightarrow 0} \frac{f_{H,V}(x)}{|x|} = 0$.

As a result

$$\lim_{t \rightarrow 0} \frac{1}{t} (f_{H,V}(th) - f_{H,V}(0)) = 0$$

and we may conclude that $d_G f_{H,V}(z) = 0$. \square

Lemma 8.4. *Let f be a continuous function on the hyperplane H such that $d_G f(0) = 0$ and C a closed cone in H not containing H^ω . Set*

$$\tilde{C} = \{x + f(x)n_H \mid x \in C\}$$

Then for η small enough and any positive δ , there is a Hamiltonian map φ supported in $B(0, 1)$, such that $\gamma(\varphi) \leq \delta$ and

$$\varphi(\tilde{C}) \cap B(0, \eta) = \emptyset$$

Proof. We shall assume w.l.o.g. that $H = \{p_n = 0\}$, so that $\tilde{C} = \{(q_n, \bar{q}, p_n, \bar{p}) \mid p_n = f(q_n, \bar{q}, \bar{p}) \text{ and } H^\omega \text{ is given by the direction } \frac{\partial}{\partial q_n}\}$. First let us mention that the case $f = 0$ is trivial: just take a Hamiltonian $K_0(p_n)$ such that $K_0(0) = 0$, $K_0'(0) = \frac{1}{2}$ and K_0 is C^0 small, η being chosen so that we have

$$B(0, \eta) \cap \left(\frac{1}{2} \frac{\partial}{\partial q_n} + C\right) = \emptyset$$

We may then truncate K_0 outside the ball $B(0, 1)$ so that we still have


$$B(0, \eta) \cap (\varphi_{K_0}^1(C)) = \emptyset$$

This extends to the case where f is smooth. Indeed, let us consider a hamiltonian $K(q_n, p_n, \bar{q}, \bar{p})$ to be a truncation of

$$F(q_n, p_n, \bar{q}, \bar{p}) = \frac{1}{2}(p_n - f(q_n, \bar{q}, \bar{p}))$$

$K(q_n, f(q_n, \bar{q}, \bar{p}), \bar{q}, \bar{p}) = 0$ and $\frac{\partial K}{\partial p_n}(q_n, f(q_n, \bar{q}, \bar{p}), \bar{q}, \bar{p}) = \frac{1}{2}$. So for any $(q_n, p_n, \bar{q}, \bar{p}) \in \tilde{C}$, $\varphi_K^t(q_n, p_n, \bar{q}, \bar{p}) = (q_n + \frac{t}{2}, p_n(t), \bar{q}(t), \bar{p}(t))$ and we get

$$\varphi_K^1(\tilde{C}) \cap B(0, \eta) = \emptyset$$

FIGURE 3. The cone C and its translation by $\frac{1}{2} \frac{\partial}{\partial q_n}$

Since K is constant on \tilde{C} , we can truncate the flow so that it is unchanged on the trajectory of points in $\varphi_K^t(\tilde{C}) \cap B(0, 1)$ and $\gamma(\varphi_K) \leq \delta$.

Note that if we replace \tilde{C} by the region

$$\tilde{C}_{g,\delta} = \{(q_n, p_n, \bar{q}, \bar{p}) \mid (q_n, \bar{q}, \bar{p}) \in C, g(q_n, \bar{q}, \bar{p}) \leq p_n \leq g(q_n, \bar{q}, \bar{p}) + \delta\}$$

we can choose K to be the truncation of a function F such that

- (1) $F(q_n, g(q_n, \bar{q}, \bar{p}), \bar{q}, \bar{p}) = 0$
- (2) $F(q_n, g(q_n, \bar{q}, \bar{p}) + \delta, \bar{q}, \bar{p}) = \frac{1}{2}\delta$
- (3) $\frac{\partial F}{\partial p_n}(p_n, q_n, \bar{q}, \bar{p}) = \frac{1}{2}$ for $g(q_n, \bar{q}, \bar{p}) \leq p_n \leq g(q_n, \bar{q}, \bar{p}) + \delta$

As a result the flow of K preserves $\tilde{C}_{g,\delta}$ and on $\tilde{C}_{g,\delta}$ is given by $(q_n, p_n, \bar{q}, \bar{p}) \mapsto (q_n + \frac{1}{2} \frac{\partial}{\partial q_n}, p_n, \bar{q}, \bar{p})$. Then $\varphi_K^1(\tilde{C}_{g,\delta}) \cap B(0, \eta) = \emptyset$, since the projection of $\varphi_K^1(\tilde{C}_{g,\delta})$ on H does not intersect $B(0, \eta)$. However now we cannot have $\gamma(\varphi_K)$ arbitrarily small because K has oscillation bounded from below by 2δ . Indeed we must have $\frac{\partial K}{\partial p_n} = \frac{1}{2}$ on a segment in the p_n direction of length δ . However we can assume $\gamma(\varphi_K) \leq \|K\| \leq \frac{1}{2}\delta + \delta = \frac{3\delta}{2}$.

The general case is obtained as follows. We take a sequence g_k such that $g_k \leq f \leq g_k + \frac{1}{k}$. We can then displace $\tilde{C}_{g_k,\delta}$ hence \tilde{C} by a map with $\gamma(\varphi_k) \leq \frac{3}{2k}$. This concludes our proof. \square

Let us now conclude the proof of the Proposition. According to Lemma 8.3 there is a ball $B(0, \varepsilon)$ such that $V \cap B(0, \varepsilon)$ can be written as the graph of $f_{H,V}$ over $\Gamma \subset C_\rho^-(0, V) \subset H$. By assumption $C_\rho^-(0, V)$ does not contain H^ω , so according to Lemma 8.4, where we replace $B(0, 1)$ by $B(0, \varepsilon)$, we have for any $\delta > 0$ a Hamiltonian map φ supported in $B(0, \varepsilon)$, such that $\gamma(\varphi) < \delta$ and $\varphi(V) \cap B(0, \eta) = \emptyset$. This means that V is not γ -coisotropic at 0. \square

Remark 8.5. The proof actually shows that V is locally rigid in the sense of Usher (see [Ush19]). So we have that γ -coisotropic implies locally rigid which implies cone-coisotropic.

8.2. A cone-coisotropic set that is not γ -coisotropic. Let K_a be the central Cantor set of ratio a . In other words we set

$$F_1 = [0, a] \cup [1 - a, 1]$$

$$F_2 = [0, a^2] \cup [a(1 - a), a] \cup [1 - a, 1 - a + a^2] \cup [1 - a^2, 1]$$

and $K_a = \bigcap_k F_k$.

Proposition 8.6. *For $a < 2^{-2n}$, the set $X = K_a^{2n} \subset \mathbb{R}^{2n}$ is cone-coisotropic but not γ -coisotropic.*

Proof. We claim that $C^+(z, X)$ is not contained in any hyperplane. Indeed for $x \in K = K_a$ the cone $C^+(x, K)$ is never empty since any point in K is a cluster point of K . In fact $C^-(x, K)$ is either $\mathbb{R}_-, \mathbb{R}_+$ or \mathbb{R} depending whether x is the limit of points of K less than x , the limit of points of K greater than x or both, and $C^+(x, K) = \mathbb{R}$. Even though the paratingent cone of a product does not necessarily contain the product of the paratingent cones of the factors, in the present case $C^+(z, X)$ contains the rays $\{0^{j-1}\} \times C^+(x_j, K) \times \{0^{n-j}\}$ where $z = (x_1, \dots, x_{2n})$, and $0^k = (0, \dots, 0) \in \mathbb{R}^k$. Therefore $C^+(z, X)$ is not contained in any hyperplane and X is trivially cone-coisotropic.

We now claim X is not γ -coisotropic. Pick $z \in X$ and balls $B(z, \varepsilon)$, $B(z, \eta)$, $0 < \eta < \varepsilon$. Let k big enough (to be made precise later). Since K is contained in F_k which consists of 2^k intervals of length a^k , our X is contained in a family \mathcal{S} of cubes of edge length a^k with $|\mathcal{S}| = 2^{2nk}$. The projection $(q, p) \mapsto (q_2, \dots, q_n, p_1, \dots, p_n)$ maps \mathcal{S} to a family \mathcal{S}' of cubes of \mathbb{R}^{2n-1} (again these cubes have edge length a^k and $|\mathcal{S}'| = 2^{(2n-1)k}$). We choose disjoint neighbourhoods U_i , $i \in \mathcal{S}'$, of the cubes in \mathcal{S}' . For a given $i_0 \in \mathcal{S}'$, let C_j , $j = 1, \dots, N$, be the cubes of \mathcal{S} contained in $B(z, \varepsilon) \cap (\mathbb{R} \times U_{i_0})$, ordered according to the q_1 variable. Let N_1 be the maximal index j such that C_j meets $B(z, \eta)$. We push all cubes C_2, \dots, C_{N_1} to the left (that is, decreasing their q_1 coordinates), close to C_1 , in the space between $B(z, \varepsilon)$ and $B(z, \eta)$ (if k is big enough), as follows. We move C_2 by

a translation T_2 with direction $-\partial_{q_1}$, so that $T_2(C_2)$ is close to C_1 . Then we move C_3 by a similar translation T_3 , so that $T_3(C_3)$ is close to $T_2(C_2)$. We go on until C_{N_1} . We have thus accumulated at most 2^k cubes of edge length a^k near C_1 . Since $a < 1/2$, we have room to put all these cubes in $B(z, \varepsilon) \setminus B(z, \eta)$ for k big enough.

Now the translation T_j can be realized by a Hamiltonian function of the form $h_j = h(p_1)\rho_j(q, p)$, where $h' = 1$ and ρ_j is a bump function vanishing outside $B(z, \varepsilon) \cap (\mathbb{R} \times U_{i_0})$ and over $C_1 \cup \dots \cup T_{j-1}(C_{j-1}) \cup C_{j+1} \cup \dots \cup C_{N_1}$ and equals to 1 over the path swept by C_j . We have $\|h_j\| \sim a^k$ and we apply the isotopy for a length of time less than $\varepsilon < 1$. We have to do this less than $|\mathcal{S}|$ times. Since we chose $a < 2^{-2n}$, the composition of all these isotopies has norm as small as required.




FIGURE 4. Moving the cubes..

□

Remarks 8.7. (1) We can weaken the assumption $a < 2^{-2n}$ to $a < 1/4$ at the expense of a more complicated proof.

(2) According to [AGHIV] where we prove that $SS(\widehat{Q}(L)) = \gamma\text{-supp}(L)$ and [AHV22], where we prove that the Birkhoff attractor is a γ -support, we may find a limit of constructible sheaves such that $SS(\mathcal{F}) = \gamma\text{-supp}(L)$ is an indecomposable continuum (i.e. a compact connected set that cannot be the non-trivial union of compact connected subsets).

Question 8.8. Find an example where the image of a smooth non-coisotropic submanifold C by a symplectic homeomorphism, φ , is such that $\varphi(C)$ is cone-coisotropic (but will not be γ -coisotropic by [Vit22]).

9. QUESTIONS AND COMMENTS

We gather in this section a number of questions that sprang up naturally while writing this paper.

Question 9.1. What is the γ_g -distance of k_X and k_Y for two closed sets X, Y . If X, Y do not have the same cohomology, the distance is infinite. So we can focus on the case when $Y = X_\varepsilon$ is an ε -neighbourhood of X . When is it true that $\gamma_g(k_X, k_{X_\varepsilon}) \leq C\varepsilon$? Or when do we have $\gamma_g - \lim k_{X_\varepsilon} = k_X$?

For the next four questions we choose on N a real analytic structure. We let $D_{lc}(N)$ be the category of limits of constructible sheaves considered in Appendix B.

Question 9.2. In Appendix B we describe the objects of $D_{lc}(\mathbb{R})$ with micro-support contained in $\{\tau \geq 0\}$. Is there a similar description in the general case?

Question 9.3. Using Lemma B.4 and the proof of Lemma B.2 it is probably not hard to see that $D_{lc}(N)$ is a triangulated category. If we have a map $f: N \rightarrow M$, can we find conditions so that $Rf_*(D_{lc}(N)) \subset D_{lc}(M)$, $f^{-1}(D_{lc}(M)) \subset D_{lc}(N)$?

For example this is true for a map of the type $f \times \text{id}_{\mathbb{R}}: N \times \mathbb{R} \rightarrow M \times \mathbb{R}$, if we consider the γ_τ -distance, because Rf_* and f^{-1} are Lipschitz for this distance. In general f^{-1} is not Lipschitz: take $f: \mathbb{R} \rightarrow \mathbb{R}^2$, $x \mapsto (x, 0)$, and D_ε the closed disc with center $(0, 1)$ and radius $1 - \varepsilon$; then $\gamma_g(k_{D_0}, k_{D_\varepsilon}) = \varepsilon$ but $\gamma_g(f^{-1}(k_{D_0}), f^{-1}(k_{D_\varepsilon})) = +\infty$ (here $g(q, p) = \|p\|$). Hence it is not clear that $f^{-1}(D_{lc}(M)) \subset D_{lc}(N)$.

Question 9.4. Let $D_{dom}(N)$ be the subcategory of $D(N)$ generated in the triangulated sense by the k_U , where U runs over the domains with smooth boundary. It should not be difficult to prove that $D_{dom}(N) \subset D_{lc}(N)$ by approximating domains with smooth boundary by domains with analytic boundary. Is it true that, for any subanalytic open subset V of X and any $\varepsilon > 0$, there exists a domain U with smooth boundary such that $U \subset V$ and $K_{\varphi_g}^\varepsilon(k_U) \simeq k_{U_\varepsilon}$ with U_ε open and $V \subset U_\varepsilon$? In this case we can see that $D_{lc}(N)$ coincides with the γ_g -closure of $D_{dom}(N)$, which would imply that $D_{lc}(N)$ only depends on the smooth structure of N .

Question 9.5. Let $D_{tri}(N)$ be the set of sheaves which are constructible with respect to some (non-fixed!) triangulation of N . Do the γ_g -closures of $D_{dom}(N)$ and $D_{tri}(N)$ coincide? Are they equal to $D_{lc}(N)$?

We remind the reader of the following (see [KS18], theorem 2.11, and [PS21], corollary 2.4.2)

Proposition 9.6. *We have for all 1-Lipschitz maps $f : M \rightarrow N$ the inequality*

$$\gamma_g(Rf_*(\mathcal{F}), Rf_*(\mathcal{G})) \leq \gamma_g(\mathcal{F}, \mathcal{G})$$

Question 9.7. Can we give conditions on \mathcal{F}, \mathcal{G} (for example $\gamma_g(\mathcal{F}, \mathcal{G}) < \infty$) such that the opposite inequality

$$\gamma_g(\mathcal{F}, \mathcal{G}) \leq C \sup_{f \in \text{Lip}_1(X, \mathbb{R})} \gamma_{|\tau|}(Rf_*(\mathcal{F}), Rf_*(\mathcal{G}))$$

holds for some constant C ?

*Question 9.8.*⁵ Let f be a continuous function on N . Then $\mathcal{F}_f = k_{\{(x,t)|f(x) \leq t\}}$ is a sheaf on $N \times \mathbb{R}$, and Vichery (in [Vic13]) defined a subdifferential as

$$\partial f = \rho(\text{SS}(\mathcal{F}_f) \cap \{\tau = 1\}) \subset T^*N,$$

where $\rho(x, t, \xi, \tau) = (x, \xi/\tau)$ is defined in §7. Its intersection with T_x^*N is $\partial f(x)$. On the other hand $df \in \widehat{\mathcal{L}}(T^*N)$, so we may ask whether

$$\rho(\text{SS}(\mathcal{F}_f) \cap \{\tau = 1\}) = \partial f = \gamma - \text{supp}(df)$$

This leads to more general questions. First notice the following

Proposition 9.9. *There is a functor $Q: \widehat{\mathcal{L}}(T^*N) \rightarrow D_{lc}(N \times \mathbb{R})$ extending the usual quantization from [Gui12; Vit19]. Moreover Q is an isometric embedding from $(\widehat{\mathcal{L}}(T^*N), \gamma)$ to $(D_{lc}(N \times \mathbb{R}), \gamma_\tau)$.*

Proof. Let L_n be a Cauchy sequence in $\mathcal{L}(T^*N)$. Then \mathcal{F}_{L_n} is a Cauchy sequence in $D_{lc}(N \times \mathbb{R})$. The map $L \rightarrow \mathcal{F}_L$ is an isometric embedding from the metric c to the (pseudo-) metric γ_τ . As we saw in Proposition 6.13 where c is the metric defined on $\mathcal{L}(T^*N)$ in [Vit22] by $c(\widetilde{L}_1, \widetilde{L}_2) = \max\{0, c_+(L_1, L_2)\} - \min\{0, c_-(L_1, L_2)\}$. Since according to Proposition B.8 the metric γ_τ is a bona fide metric in $D_{lc}(N \times \mathbb{R})$, limits are unique and the limit of \mathcal{F}_{L_n} is well-defined. \square

*Question 9.10.*⁶ Set $\rho(x, t; \xi, \tau) = (x; \xi/\tau)$. Given $\Lambda \in \widehat{\mathcal{L}}(T^*N)$, do we have $\gamma - \text{supp}(\Lambda) = \rho(\text{SS}^*(Q(\Lambda)))$?

A positive answer would also imply a positive answer to Question 9.8, i.e. $\partial f = \gamma - \text{supp}(df)$ by taking $\Lambda_n = df_n$ for some sequence of differentiable functions converging to f .

Question 9.11. Can one characterize the sheaves \mathcal{F} such that there exists an element $\widehat{\text{SS}}(\mathcal{F})$ in $\widehat{\mathcal{L}}(T^*N)$ so that $\text{SS}(\mathcal{F}) = \gamma - \text{supp}(\widehat{\text{SS}}(\mathcal{F}))$? In the previous question we considered the \mathcal{F} which are γ -limits of sheaves \mathcal{F}_j quantizing some exact Lagrangian.

⁵This question is answered positively in [AGHIV]

⁶This question is answered positively in [AGHIV]

Let us give an example of such a situation : let M, N be closed manifolds. For a map $f \in C^\infty(M, N)$ set Λ_f be the correspondence in $T^*M \times T^*N$ defined by

$$\Lambda_f = \{(x, p_x, y, p_y) \mid y = f(x), p_y \circ df(x) = -p_x\}$$

Now let f_n be a sequence of smooth maps C^0 -converging to $f \in C^0(M, N)$. For $\mathcal{F} \in D^b(M)$ the sequence $(R(f_n)_*\mathcal{F})_{n \geq 1}$ γ -converges to $Rf_*(\mathcal{F})$ and $SS(R(f_n)_*\mathcal{F}) \subset \Lambda_{f_n} \circ SS(\mathcal{F})$. Thus $SS(R(f_n)_*\mathcal{F})$ should γ -converge to a subset of $\Lambda_f \circ SS(\mathcal{F})$. Of course since f is only C^0 , Λ_f is not defined as a set, but it is well defined in $\widehat{\mathcal{L}}(T^*N \times T^*M)$, hence $\Lambda_f \circ SS(\mathcal{F}) \in \widehat{\mathcal{L}}(T^*N)$. Thus we expect $\widehat{SS}(Rf_*(\mathcal{F})) = \Lambda_f \circ SS(\mathcal{F})$ and $SS(Rf_*(\mathcal{F})) \subset \gamma\text{-supp}(\Lambda_f \circ SS(\mathcal{F}))$.

Question 9.12. Let \mathcal{F}_n be a sequence in $D(N)$ γ_g -converging to \mathcal{F}_∞ . We set $\Lambda_n = SS(\mathcal{F}_n)$ and we assume that Λ_n Hausdorff converges to Λ_∞ . We know that $SS(\mathcal{F}_\infty) \subset \Lambda_\infty$ but when is it true that $\Lambda_\infty = SS(\mathcal{F}_\infty)$? We can prove it if Λ_∞ is a smooth Lagrangian or more generally minimally γ -coisotropic. Indeed Proposition 6.26 shows that $SS(\mathcal{F}_\infty) \subset \Lambda_\infty$ and the involutivity of the singular support gives the equality (by an argument similar to the one in the proof of Proposition 9.13). Assuming $N = M \times \mathbb{R}$ and the \mathcal{F}_n are quantizations of exact Lagrangian, Proposition 9.13 shows that this still hold when Λ_∞ is the conification of some $\varphi(L_0)$ for L_0 smooth Lagrangian, φ a symplectic homeomorphism.

Note however that the answer cannot be positive in general: Let $f(x)$ be a smooth bounded function on \mathbb{R} , such that $\max f' = 1$, $\min f' = -1$. Let $f_n(x) = f(nx)/n$ and $\mathcal{F}_n = k_{\{t \geq f_n(x)\}}$. Then \mathcal{F}_n converges to $\mathcal{F}_\infty = k_{\mathbb{R} \times [0, \infty[}$ but $SS(\mathcal{F}_n)$ converges to a set Λ_∞ bigger than $SS(\mathcal{F}_\infty)$. In fact Λ_∞ is the cone over a ‘‘cross’’ C in $T^*\mathbb{R}$ with $C = 0_{\mathbb{R}} \cup (\{0\} \times [-1, 1])$.

Proposition 9.13. *Let $L = \varphi(L_0)$ where L_0 is smooth Lagrangian and $\varphi \in \mathcal{H}_\gamma(M, \omega) = \widehat{\mathcal{D}\mathfrak{H}\text{am}}(M, \omega) \cap \text{Homeo}(M)$. Then any closed proper subset of L is not γ -coisotropic.*

Proof. Let $L' \subsetneq L$. Since being γ -coisotropic is invariant by $\widehat{\mathcal{D}\mathfrak{H}\text{am}}(M, \omega) \cap \text{Homeo}(M)$ it is enough to prove the Proposition for L_0 i.e. in the smooth case. Now there is a ball $B(z, r)$ in L such that $\overline{B(z, r)} \cap L' = \{z'\}$. Moreover we may choose r to be arbitrarily small. Let us consider a flow on L such that $\rho^t(B(z, r)) \subset B(z', r)$. In a chart containing $B(z, 2r)$ we may assume that ρ^t is a translation. Then $\rho^1(L') \cap B(z', r) = \emptyset$. Since if X_t is the vector field generating ρ^t , the Hamiltonian written in action-angle coordinates (i.e. L is given locally by $p = 0$) can be chosen of the form $\chi(|p|)\langle p, X_t(x) \rangle$ for some bump function χ , and this can be made arbitrarily small. As a result, L' is not γ -coisotropic.




FIGURE 5. Proof of Proposition 9.13

□

Question 9.14. Let V be γ -coisotropic of dimension n . Then for $z \in V$ and ε small enough we have $V \setminus B(z, \varepsilon)$ is not coisotropic.

APPENDIX A. PERSISTENCE MODULES AND BARCODES

For this section we refer also to [Zha20]

We have already noticed after Definition 5.4 that the spectral invariants of $L_1, L_2 \in \mathcal{L}(T^*N)$ are encoded in the sheaf over \mathbb{R} given by $\mathcal{H} = R t_* \mathcal{H}om^*(\mathcal{F}, \mathcal{G})$, where $t: N \times \mathbb{R} \rightarrow \mathbb{R}$ is the projection, because of the isomorphism

$$R\Gamma_{N \times \{t\}}(N \times \mathbb{R}; \mathcal{H}om^*(\mathcal{F}_{L_1}, \mathcal{F}_{L_2})) \simeq R\Gamma_{\{t\}}(\mathbb{R}; \mathcal{H})$$

We recall that \mathcal{H} belongs to the category $D_{\tau \geq 0}(\mathbb{R})$ of sheaves with singular support in $\{\tau \geq 0\}$.

We remark that $D_{\tau \geq 0}(\mathbb{R})$ contains the derived category of *persistence modules*, where by a persistence module, we mean a constructible sheaf on (\mathbb{R}, \leq) . We refer to [Bar94; Cha+09; ELZ02; KS18; ZC05] for the theory and applications. Such a persistence module is uniquely defined by the finite-dimensional vector spaces $V_t = \mathcal{F}(\cdot - \infty, t]$ and the linear maps $r_{s,t}: V_t \rightarrow V_s$ defined for $s \leq t$, such that

- (1) for $s < t < u$ we have $r_{u,t} \circ r_{s,t} = r_{s,u}$,
- (2) $\lim_{t > s} V_t = V_s$ where the limit is that of the directed system given by the $r_{s,t}$,
- (3) $r_{t,t} = \text{Id}$.

It is a well-known result that persistence modules have a decomposition as sum of barcodes : this is Gabriel's theorem ([Gab72] on quivers in the finite case and Crawley-Boevey's theorem ([Cra15]) in the locally finite case. More precisely, using Gabriel's theorem and the fact that any complex of objects in an Abelian category of homological dimension 1 is the

sum of its cohomology, we find: any $\mathcal{F} \in D_{\tau \geq 0}(\mathbb{R})$ which is constructible for a finite stratification can be written $\mathcal{F} \simeq \bigoplus_{j \in \mathcal{J}} k_{[a_j, b_j]}[-n_j]$, where \mathcal{J} is a finite family, and this decomposition is unique up to isomorphism. (The hypothesis that the singular support is in $\{\tau \geq 0\}$ is useless here.) We remark that the stratification consists only of a finite set of points in \mathbb{R} and the connected components of its complement, where \mathcal{F} is constant. This finite set is thus the projection to \mathbb{R} of $SS^*(\mathcal{F})$. If $L \in \mathcal{L}(T^*N)$ is generic in the sense that L meets 0_N transversely, then $Rt_*(\mathcal{F}_L)$ is of the above type, where the stratification is given by the set of actions of the points in $L \cap 0_N$.

In [KS18] it is explained how to go from the case of a finite stratification to the case of a locally finite one (then \mathcal{J} is at most countable). By [Cra15] the same result holds in fact (at least in $D_{\tau \geq 0}(\mathbb{R})$) under the more general hypothesis that \mathcal{F}_x is finite dimensional for all $x \in \mathbb{R}$. However the sheaves we encounter in this paper are not necessarily of this type: if f is the function $f(x) = x^2 \sin(1/x)$ and $\mathcal{F} = Rt_*(k_{\{t \geq f(x)\}})$, then \mathcal{F}_0 is infinite dimensional. But, by a small perturbation of L , we see that all sheaves of the form $Rt_*(\mathcal{F}_L)$ are in the closure of the set of constructible sheaves. Using Corollary B.12 we deduce:

Proposition A.1. *Let $L_1, L_2 \in \mathcal{L}(T^*N)$. Then there exists an at most countable family \mathcal{J} (which is finite in the case of transverse intersection) such that*

$$(Rt)_*(\mathcal{H}om^*(\mathcal{F}_{L_1}, \mathcal{F}_{L_2})) = \bigoplus_{j \in \mathcal{J}} k_{[a_j, b_j]}[-n_j],$$

where for each $j \in \mathcal{J}$ we have $a_j \in \mathbb{R} \cup \{-\infty\}$, $b_j \in \mathbb{R}$, $n_j \in \mathbb{Z}$. Moreover there is a unique j_- such that $a_{j_-} = -\infty$ and $n_{j_-} = -1$ and then $b_{j_-} = c_-(L_1, L_2)$ and a unique j_+ such that $a_{j_+} = +\infty$ and $n_{j_+} = n - 1$ and then $b_{j_+} = c_+(L_1, L_2)$.

More generally, with the notations of the proposition we let \mathcal{J}_∞ be the set of j such that $a_j = -\infty$. Then $H_{t_0}^*((Rt)_*(\mathcal{H}om^*(\mathcal{F}_{L_1}, \mathcal{F}_{L_2}))) \simeq \bigoplus_{j \in \mathcal{J}_\infty} k[-n_j]$, for $t_0 \ll 0$, and we deduce that the spectral invariants $c(\alpha, L_1, L_2)$ are the b_j 's with $j \in \mathcal{J}_\infty$.

Note that the filtered Floer cohomology, $FH^*(L_1, L_2; -\infty, t)$ defines a persistence module equal to the one obtained from $\mathcal{H}om^*(\mathcal{F}_{L_1}, \mathcal{F}_{L_2})$ as follows from Theorem 5.2.

It will be useful to remind the reader that

Lemma A.2 (see [KS18], (1.10)).

$$\mathrm{RHom}(k_{[a, b]}, k_{[c, d]}) \simeq \begin{cases} k & \text{for } a \leq c < b \leq d, \\ k[-1] & \text{for } c < a \leq d < b, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. The formula $k_{[c,d]} \simeq \mathbf{R}\mathcal{H}\text{om}(k_{[c,d]}, k_{\mathbb{R}})$ and the adjunction between \otimes and $\mathbf{R}\mathcal{H}\text{om}$ give $\mathbf{R}\text{Hom}(k_{[a,b]}, k_{[c,d]}) \simeq \mathbf{R}\text{Hom}(k_I, k_{\mathbb{R}})$, where $I = [a, b[\cap]c, d]$. If I is half closed or empty, the result is 0. The two other cases correspond to I open or I closed. \square

APPENDIX B. DECOMPOSITION IN THE COMPLETION OF $D_{lc}(\mathbb{R})$

B.1. Limits of constructible sheaves. We assume in this section that the γ_g -distance on $D_{g \geq 0}(N)$ is given by a *real analytic* function $g: T^*N \setminus 0_N \rightarrow \mathbb{R}$. We recall that we work with coefficients in a field k . We denote by $D_c(N)$ the subcategory of $D(N)$ of constructible sheaves, that is, the sheaves \mathcal{F} for which there exists a locally finite subanalytic stratification of N such that the restriction of \mathcal{F} to each stratum is locally constant of finite rank (see [KS90, Chap. VIII]). We let $D_{lc}(N)$ be the set of objects which are limits of objects in $D_c(N) \cap D_{g \geq 0}(N)$ with respect to the γ_g -topology. Since g is analytic, $\varphi = \varphi_g$ is also analytic and $K_\varphi \in D_c(N^2)$. Hence $K_\varphi^l(-)$ preserves $D_c(N) \cap D_{g \geq 0}(N)$. A sheaf \mathcal{F} belongs to $D_{lc}(N)$ if and only if, for any $\varepsilon > 0$, there exists $\mathcal{F}' \in D_c(N) \cap D_{g \geq 0}(N)$ and morphisms $K_\varphi^{-\varepsilon}(\mathcal{F}) \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow K_\varphi^\varepsilon(\mathcal{F}') \rightarrow K_\varphi^\varepsilon(\mathcal{F})$ whose compositions are the morphisms τ (notice that $K_\varphi^\varepsilon(\mathcal{F}')$ itself belongs to $D_c(N)$).

Our aim in this section is to prove that the distance γ_g is non degenerate on $D_{lc}(N)$. For $\mathcal{F}, \mathcal{G} \in D_{lc}(N)$ with $\gamma_g(\mathcal{F}, \mathcal{G}) = 0$ we will prove the existence of an isomorphism between \mathcal{F} and \mathcal{G} as follows: we first see that $\text{Hom}(\mathcal{F}, \mathcal{G})$ is almost $\varinjlim_n \text{Hom}(\mathcal{F}, K_\varphi^{1/n}(\mathcal{G}))$ and then we use the morphisms in $\text{Hom}(\mathcal{F}, K_\varphi^{1/n}(\mathcal{G}))$ whose existence follows from the condition $\gamma_g(\mathcal{F}, \mathcal{G}) = 0$. Actually we will prove

$$\text{Hom}(\mathcal{F}, \mathcal{G}) \simeq \varinjlim_n \text{Hom}(\mathcal{F}|_{U_n}, K_\varphi^{1/n}(\mathcal{G})|_{U_n})$$

where U_n is an increasing family of open subsets with compact closures. The reason for introducing these U_n 's is that this enables us to check the Mittag-Leffler criterion by approximating \mathcal{F} and \mathcal{G} by constructible sheaves. We often write $\text{Hom}(\mathcal{F}|_U, \mathcal{G}|_U)$ for $\text{Hom}_{D(U)}(\mathcal{F}|_U, \mathcal{G}|_U)$ when the working category (here $D(U)$) is understood. We will also use the isomorphisms, for $\mathcal{F}, \mathcal{G} \in D(N)$ and $U \subset N$ open,

$$\text{Hom}(\mathcal{F}, \mathcal{G}) \simeq H^0 \mathbf{R}\text{Hom}(\mathcal{F}, \mathcal{G}), \quad \text{Hom}(\mathcal{F}|_U, \mathcal{G}|_U) \simeq \text{Hom}(\mathcal{F}_U, \mathcal{G}),$$

where $\mathcal{F}_U = \mathcal{F} \otimes k_U$.

When cutting by the open sets U_n we need to compare $K_\varphi^a(\mathcal{F}_U)$ and $K_\varphi^a(\mathcal{F})$. This is the object of the next lemma.

Lemma B.1. *Let $V \subset U$ be open subsets of N such that \overline{V} is compact and $\overline{V} \subset U$. There exists $a_0 > 0$ such that, for any $a \in [-a_0, a_0]$ and $\mathcal{F} \in D(N)$,*

the morphisms

$$(K_\varphi^a(\mathcal{F}_V))_U \rightarrow K_\varphi^a(\mathcal{F}_V), \quad (K_\varphi^a(\mathcal{F}_U))_V \rightarrow (K_\varphi^a(\mathcal{F}))_V,$$

are isomorphisms.

Proof. Set $\bar{V}_a = \bigcup_{0 \leq b \leq a} \pi(\varphi_b(\pi^{-1}(\bar{V})))$, where $\pi: T^*N \setminus 0_N \rightarrow N$ is the projection. Since \bar{V} is compact, there exists $a_0 > 0$ such that $\bar{V}_{a_0} \subset U$.

Let $\mathcal{G} \in D(N \times [-a_0, 0])$ be the sheaf obtained from \mathcal{F}_V by applying the whole isotopy, that is, $\mathcal{G}|_{N \times \{t\}} = K_{\varphi_t}(\mathcal{F}_V)$. Since $SS(\mathcal{F}_V) \subset \pi^{-1}(\bar{V})$, we have $SS(\mathcal{G}) \subset T^*\bar{V}_{a_0} \times T^*[-a_0, 0]$ outside the zero-section. Hence, for any $x \in N \setminus \bar{V}_{a_0}$, \mathcal{G} is locally constant near $x \times [-a_0, 0]$, hence 0 near $x \times [-a_0, 0]$ since $(\mathcal{G})_{(x,0)} = 0$. It follows that the support of $K_\varphi^a(\mathcal{F}_V)$ is contained in U and this gives the first isomorphism.

We also see that the support of $K_\varphi^a(\mathcal{F}_{N \setminus U})$ avoids V , which gives the second isomorphism. \square

Lemma B.2. *Let $\mathcal{F}, \mathcal{G} \in D_{lc}(N)$, let $\varepsilon > 0$ and let $V \subset U$ be two open subsets of N . We assume that \bar{V} is compact and $\bar{V} \subset U$. Then the morphism $\text{Hom}_{D(U)}(\mathcal{F}|_U, \mathcal{G}|_U) \rightarrow \text{Hom}_{D(V)}(\mathcal{F}|_V, K_\varphi^\varepsilon(\mathcal{G})|_V)$ induced by $\tau_{0,\varepsilon}(\mathcal{G})$ and the restriction from U to V has finite dimensional image.*

Proof. We recall that $\text{Hom}_{D(U)}(\mathcal{F}|_U, \mathcal{G}|_U) \simeq \text{Hom}_{D(N)}(\mathcal{F}_U, \mathcal{G})$ and our morphism is given by composing with $\mathcal{F}_V \rightarrow \mathcal{F}_U$ and $\tau_{0,\varepsilon}(\mathcal{G})$. For $0 < \eta < \varepsilon$ we have $\text{Hom}(\mathcal{F}_V, K_\varphi^\varepsilon(\mathcal{G})) \simeq \text{Hom}(K_\varphi^{-\eta}(\mathcal{F}_V), K_\varphi^{\varepsilon-\eta}(\mathcal{G}))$. We choose a subanalytic open subset W such that \bar{W} is compact and $\bar{V} \subset W \subset \bar{W} \subset U$. By Lemma B.1 and the commutative diagram

$$\begin{array}{ccccc} (K_\varphi^{-\eta}(\mathcal{F}_V))_W & \xrightarrow{\sim} & K_\varphi^{-\eta}(\mathcal{F}_V) & \xrightarrow{\tau_{-\eta,0}(\mathcal{F}_V)} & \mathcal{F}_V \\ \downarrow & & \swarrow \text{dotted} & & \downarrow \\ (K_\varphi^{-\eta}(\mathcal{F}))_W & \xrightarrow{(\tau_{-\eta,0}(\mathcal{F}))_W} & & & \mathcal{F}_W \end{array}$$

the composition $K_\varphi^{-\eta}(\mathcal{F}_V) \xrightarrow{\tau_{-\eta,0}(\mathcal{F}_V)} \mathcal{F}_V \rightarrow \mathcal{F}_W$ is factorized as $K_\varphi^{-\eta}(\mathcal{F}_V) \rightarrow (K_\varphi^{-\eta}(\mathcal{F}))_W \xrightarrow{(\tau_{-\eta,0}(\mathcal{F}))_W} \mathcal{F}_W$, for η small enough. Since $\mathcal{F}, \mathcal{G} \in D_{lc}(N)$, there exist constructible sheaves $\mathcal{F}', \mathcal{G}'$ and morphisms $K_\varphi^{-\eta} \mathcal{F} \xrightarrow{u_1} \mathcal{F}' \xrightarrow{u_2} \mathcal{F}$, $\mathcal{G} \xrightarrow{v_1} \mathcal{G}' \xrightarrow{v_2} K_\varphi^{\varepsilon-\eta}(\mathcal{G})$ such that $u_2 \circ u_1 = \tau_{-\eta,0}(\mathcal{F})$, $v_2 \circ v_1 = \tau_{0,\varepsilon-\eta}(\mathcal{G})$.

We sum up these morphisms in the diagram

$$\begin{array}{ccccccccc}
K_\varphi^{-\eta}(\mathcal{F}_V) & \longrightarrow & (K_\varphi^{-\eta}(\mathcal{F}))_W & \longrightarrow & \mathcal{F}'_W & \longrightarrow & \mathcal{F}_W & \longrightarrow & \mathcal{F}_U & \xrightarrow{\alpha} & \mathcal{G} \\
& & & & & & & & & & \downarrow \\
& & & & & & & & & & \mathcal{G}' \\
& & & & & & & & & & \downarrow \\
& & & & & & & & & & K_\varphi^{\varepsilon-\eta}(\mathcal{G})
\end{array}$$

γ (dotted arrow from $K_\varphi^{-\eta}(\mathcal{F}_V)$ to $K_\varphi^{\varepsilon-\eta}(\mathcal{G})$)
 β (dotted arrow from \mathcal{F}'_W to \mathcal{G}')

We deduce that the morphism of the lemma, which we have rewritten as

$$\mathrm{Hom}(\mathcal{F}_U, \mathcal{G}) \rightarrow \mathrm{Hom}(K_\varphi^{-\eta}(\mathcal{F}_V), K_\varphi^{\varepsilon-\eta}(\mathcal{G})) \simeq \mathrm{Hom}(\mathcal{F}_V, K_\varphi^\varepsilon(\mathcal{G}))$$

and corresponds to $\alpha \mapsto \gamma$ in the diagram, factorizes as $\alpha \mapsto \beta$ and $\beta \mapsto \gamma$ through $\mathrm{Hom}(\mathcal{F}'_W, \mathcal{G}')$ which is finite dimensional. \square

We choose an increasing family of open subsets U_n , $n \in \mathbb{N}$, of N such that $N = \bigcup_{n \in \mathbb{N}} U_n$, each $\overline{U_n}$ is compact and $\overline{U_n} \subset U_{n+1}$. For $\mathcal{F}, \mathcal{G} \in D_{lc}(N)$ we set

$$A_n(\mathcal{F}, \mathcal{G}) = \mathrm{Hom}(\mathcal{F}_{U_n}, K_\varphi^{1/n}(\mathcal{G}))$$

The morphisms $\tau_{1/n, 1/(n-1)}(\mathcal{G}): K_\varphi^{1/n}(\mathcal{G}) \rightarrow K_\varphi^{1/(n-1)}(\mathcal{G})$ and $\mathcal{F}_{U_{n-1}} \rightarrow \mathcal{F}_{U_n}$ induce $a_n: A_n(\mathcal{F}, \mathcal{G}) \rightarrow A_{n-1}(\mathcal{F}, \mathcal{G})$. Lemma B.2 implies that the projective system $(A_n(\mathcal{F}, \mathcal{G}), a_n)$, satisfies the Mittag-Leffler criterion.

By Lemma 6.24 we have an isomorphism $f: \mathrm{hocolim} K_\varphi^{-1/n}(\mathcal{F}) \xrightarrow{\sim} \mathcal{F}$. The morphisms $f_{n-1}: K_\varphi^{-1/(n-1)}(\mathcal{F}_{U_{n-1}}) \rightarrow K_\varphi^{-1/n}(\mathcal{F}_{U_n})$ give a similar injective system $(K_\varphi^{-1/n}(\mathcal{F}_{U_n}), f_n)_n$ and the morphisms $K_\varphi^{-1/n}(\mathcal{F}_{U_n}) \rightarrow \mathcal{F}$ yield

$$f': \mathrm{hocolim} K_\varphi^{-1/n}(\mathcal{F}_{U_n}) \rightarrow \mathcal{F}$$

which is also an isomorphism: indeed this can be checked locally and we can restrict to a relatively compact open subset V ; for n big enough we have $(K_\varphi^{-1/n}(\mathcal{F}))|_V \simeq (K_\varphi^{-1/n}(\mathcal{F}_{U_n}))|_V$ by Lemma B.1 and f' coincides with f . Applying $\mathrm{RHom}(-, \mathcal{G})$ we obtain

$$(B.1) \quad \mathrm{RHom}(\mathcal{F}, \mathcal{G}) \xrightarrow{\sim} \mathrm{holim}_n \mathrm{RHom}(\mathcal{F}_{U_n}, K_\varphi^{1/n}(\mathcal{G}))$$

We can deduce $H^i \mathrm{RHom}(\mathcal{F}, \mathcal{G}) \simeq \varprojlim_n H^i \mathrm{RHom}(\mathcal{F}_{U_n}, K_\varphi^{1/n}(\mathcal{G}))$ using the Mittag-Leffler criterion. Since this criterion is not often stated in the framework of homotopy limits, we recall it in the next lemma (which is a variation on the usual Mittag-Leffler condition; see for example [KS90, Prop. 1.12.4]). Here Ab denotes the category of Abelian groups and $D(\mathrm{Ab})$ its derived category.

Lemma B.3. *Let (A_n, a_n) , $a_n: A_n \rightarrow A_{n-1}$, be a projective system in $D(\text{Ab})$ and define $\text{holim}_n A_n$ by the distinguished triangle $\text{holim}_n A_n \rightarrow \prod_n A_n \xrightarrow{\alpha} \prod_n A_n \xrightarrow{+1}$ where $\alpha = \text{id} - \prod_n a_n$. We assume that $H^i A_n$ satisfies the Mittag-Leffler criterion, for some i , that is, for each n the image of $H^i A_m \rightarrow H^i A_n$ stabilizes when $m \rightarrow \infty$. Then $H^{i+1}(\text{holim}_n A_n) \simeq \varprojlim_n H^{i+1} A_n$.*

Proof. In Ab the product \prod is exact, hence H^i and \prod commute and we have a long exact sequence $\prod_n H^i A_n \xrightarrow{H^i \alpha} \prod_n H^i A_n \rightarrow H^{i+1}(\text{holim}_n A_n) \rightarrow \prod_n H^{i+1} A_n \rightarrow \prod_n H^{i+1} A_n$. Hence the statement is equivalent to the surjectivity of $H^i \alpha$. We write for short $B_n = H^i A_n$, $b_n = H^i a_n$ and $B_n^\infty = \text{im}(B_m \rightarrow B_n)$ for $m \gg n$. We let b_n^∞, \bar{b}_n be the maps induced by b_n on $B_n^\infty, B_n/B_n^\infty$. We have $H^i \alpha = \text{id} - \prod b_n$ and the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \prod B_n^\infty & \longrightarrow & \prod B_n & \longrightarrow & \prod B_n/B_n^\infty \longrightarrow 0 \\ & & \downarrow \text{id} - \prod b_n^\infty & & \downarrow \text{id} - \prod b_n & & \downarrow \text{id} - \prod \bar{b}_n \\ 0 & \longrightarrow & \prod B_n^\infty & \longrightarrow & \prod B_n & \longrightarrow & \prod B_n/B_n^\infty \longrightarrow 0 \end{array}$$

We see that $\text{id} - \prod b_n^\infty$ is surjective because the b_n^∞ 's are surjective. Let us write $t = \prod \bar{b}_n$. By the Mittag-Leffler criterion, for each n , there exists $m \geq n$ such that $\bar{b}_{n+1} \circ \bar{b}_{n+2} \circ \cdots \circ \bar{b}_m$ is the zero map. It follows that $t' = \sum_{i=0}^\infty t^i$ is well-defined. Hence $\text{id} - t$ is an isomorphism, with inverse t' . We deduce that $\text{id} - \prod b_n$ is also surjective, as required. \square

We apply this result to the homotopy limit in (B.1). By Lemma B.2 the maps $H^i \text{RHom}(\mathcal{F}_{U_{n+1}}, K_\varphi^{1/(n+1)}(\mathcal{G})) \rightarrow H^i \text{RHom}(\mathcal{F}_{U_n}, K_\varphi^{1/n}(\mathcal{G}))$ have finite dimensional image (recall that $H^i \text{RHom}(X, Y) \simeq \text{Hom}(X, Y[i])$ in any derived category). Hence we can indeed apply Lemma B.3 and we obtain

Lemma B.4. *Let $\mathcal{F}, \mathcal{G} \in D_{lc}(N)$ and let $U_n, n \in \mathbb{N}$, be a sequence of open subsets as above. We have a natural isomorphism*

$$\text{Hom}(\mathcal{F}, \mathcal{G}) \xrightarrow{\simeq} \varprojlim_n \text{Hom}(\mathcal{F}_{U_n}, K_\varphi^{1/n}(\mathcal{G}))$$

We recall that $A_n(\mathcal{F}, \mathcal{G}) = \text{Hom}(\mathcal{F}_{U_n}, K_\varphi^{1/n}(\mathcal{G}))$. We set

$$B_n(\mathcal{F}, \mathcal{G}) = \text{Im}(\text{Hom}(\mathcal{F}, \mathcal{G}) \rightarrow A_n(\mathcal{F}, \mathcal{G}))$$

which is a finite dimensional vector subspace of $A_n(\mathcal{F}, \mathcal{G})$ by Lemma B.2. For $u \in \text{Hom}(\mathcal{F}, \mathcal{G})$, we denote its image in $B_n(\mathcal{F}, \mathcal{G})$ by

$$[u]_n \in B_n(\mathcal{F}, \mathcal{G})$$

The map $a_n: A_n(\mathcal{F}, \mathcal{G}) \rightarrow A_{n-1}(\mathcal{F}, \mathcal{G})$ induces $b_n: B_n(\mathcal{F}, \mathcal{G}) \rightarrow B_{n-1}(\mathcal{F}, \mathcal{G})$. Since $A_n(\mathcal{F}, \mathcal{G})$ satisfies Mittag-Leffler, the images of the maps $A_r(\mathcal{F}, \mathcal{G}) \rightarrow$

Proof. (1) Let $r(n)$ be as in (B.2) and define $s(n)$ in the same way, with \mathcal{F} and \mathcal{G} switched. We choose $m \geq \max\{r(n), s(n)\}$. Since $\gamma_g(\mathcal{F}, \mathcal{G}) = 0$, we can find $u_1: \mathcal{F} \rightarrow K_\phi^{1/2m}(\mathcal{G})$ and $v_1: \mathcal{G} \rightarrow K_\phi^{1/2m}(\mathcal{F})$ such that $u_1 \circ K_\phi^{1/2m}(v_1)$ and $v_1 \circ K_\phi^{1/2m}(u_1)$ are the morphisms τ . We let $u_2 \in A_{2m}(\mathcal{F}, \mathcal{G})$ be the composition of $\mathcal{F}_{U_{2m}} \rightarrow \mathcal{F}$ and u_1 and we let u be the image of u_2 in $A_n(\mathcal{F}, \mathcal{G})$. Then u belongs to $B_n(\mathcal{F}, \mathcal{G})$. We define v in the same way. Since $v_1 \circ u_1 = \tau_{0,1/m}(\mathcal{F})$ we have $v \circ_n u = \tau_{0,1/n}(\mathcal{F})$. Similarly $u \circ_n v = \tau_{0,1/n}(\mathcal{G})$. Hence $(u, v) \in C_n$.

(2) We first remark that $\beta_{u,v}(u', v')$ belongs to $B_n^\times(\mathcal{F}, \mathcal{F})$: indeed the inverse of $v \circ_n u'$ is $v' \circ_n u$. Now we compute $\beta_{u,v}(\alpha_{u,v}(a)) = v \circ_n u \circ_n a = a$ and $\alpha_{u,v}(\beta_{u,v}(u', v')) = (u \circ_n v \circ_n u', v' \circ_n u \circ_n v) = (u', v')$.

(3) We pick $(u, v) \in C_m$ and let (\bar{u}, \bar{v}) be its image by the natural map $C_m \rightarrow C_n$. We obtain a commutative diagram

$$\begin{array}{ccc} B_m^\times(\mathcal{F}, \mathcal{F}) & \xrightarrow{\alpha_{u,v}} & C_m \\ q \downarrow & & \downarrow \\ B_n^\times(\mathcal{F}, \mathcal{F}) & \xrightarrow{\alpha_{\bar{u}, \bar{v}}} & C_n \end{array}$$

where the horizontal maps are bijections. The map $B_m(\mathcal{F}, \mathcal{F}) \rightarrow B_n(\mathcal{F}, \mathcal{F})$ is surjective by definition. Hence the map q is surjective by Lemma B.7 and the result follows. \square

Lemma B.7. *Let A, B be finite dimensional algebras over k and let $f: A \rightarrow B$ be a surjective algebra morphism. Then the map induced on the sets of invertible elements $f^\times: A^\times \rightarrow B^\times$ is also surjective.*

Proof. We pick $x \in A$ and assume that $f(x)$ is invertible. We have to prove that there exists $z \in \ker(f)$ such that $x + z$ is invertible. We let $A' = \langle x \rangle$ be the subalgebra of A generated by x and define similarly $B' = \langle f(x) \rangle$. The maps $u: k[X] \rightarrow A$, $X \mapsto x$, and $f \circ u$ identify A' and B' with quotients of $k[X]$. Since $k[X]$ is principal, we can write $A' = k[X]/\langle PQ \rangle$, with $x = [X]_{A'}$, and $B' = k[X]/\langle Q \rangle$, with $f(x) = [X]_{B'}$.

Since $f(x)$ is invertible in B , it is not a zero divisor and we have $X \nmid Q$. Let us write $P = X^n P'$ with $X \nmid P'$. We set $R = X + P'Q$. Then $f([R]_{A'}) = f(x)$. We claim that $[R]_{A'}$ is invertible, which concludes the proof. We have to check that $\langle R, PQ \rangle = k[X]$. Let D be a generator of $\langle R, PQ \rangle$. Then $D \mid X^n R - PQ = X^{n+1}$. Either $X \mid D$, but then $X \mid R$ and finally $X \mid P'Q$ which is false, or $D \in k^\times$ and we are done. \square

Proposition B.8. *We recall that we take coefficients in a field. Let \mathcal{F}, \mathcal{G} be elements in $D_{lc}(N)$. Assume that $\gamma_g(\mathcal{F}, \mathcal{G}) = 0$. Then $\mathcal{F} \simeq \mathcal{G}$.*

Question B.9. Is the Proposition true if we replace the field by \mathbb{Z} or a more general ring ?

Proof. If $\mathcal{F} = 0$, then $\gamma_g(0, \mathcal{G}) = 0$ and we obtain $\mathcal{G} = 0$ by Lemma 6.4. We assume $\mathcal{F} \neq 0$, hence $\tau_{0,1/n}(\mathcal{F}) \neq 0$ for n big enough. We use the notation C_n of Lemma B.6. By this lemma the inductive system $\cdots \rightarrow C_n \rightarrow C_{n-1} \rightarrow \cdots$ is made of surjective maps between non empty sets for $n \gg 0$. Hence its limit is non empty and we deduce a pair $(u, v) \in \text{Hom}(\mathcal{F}, \mathcal{G}) \times \text{Hom}(\mathcal{F}, \mathcal{G})$ such that $([u]_n, [v]_n) \in C_n$ for all n , that is, $[v \circ u]_n = \tau_{0,1/n}(\mathcal{F})$ and $[u \circ v]_n = \tau_{0,1/n}(\mathcal{G})$. Let \mathcal{C} be the cone of u . We want to prove that $\mathcal{C} \simeq 0$. This can be done locally and we can restrict to some open subset V with compact closure. For any n big enough so that $V \subset U_n$ we have the following morphism of triangles given by the morphisms $\tau_{0,1/n}(-)$ (recall that we work on V , so $\mathcal{F}_{U_n} \simeq \mathcal{F}$, $\mathcal{G}_{U_n} \simeq \mathcal{G}$):

$$\begin{array}{ccccccc} \mathcal{F} & \xrightarrow{u} & \mathcal{G} & \longrightarrow & \mathcal{C} & \longrightarrow & \mathcal{F}[1] \\ \downarrow & \swarrow [v]_n & \downarrow & & \downarrow & & \downarrow \\ K_\varphi^{1/n}(\mathcal{F}) & \longrightarrow & K_\varphi^{1/n}(\mathcal{G}) & \longrightarrow & K_\varphi^{1/n}(\mathcal{C}) & \longrightarrow & K_\varphi^{1/n}(\mathcal{F})[1], \end{array}$$

where both maps $\tau_{0,1/n}(\mathcal{F})$ and $\tau_{0,1/n}(\mathcal{G})$ factorize through $[v]_n$. Using the same argument as in the proof of Lemma 6.8-(1) we can deduce that $\tau_{0,2/n}(\mathcal{C}) = 0$. Hence $\gamma_g(0, \mathcal{C}) \leq 2/n$, for any n . Thus the cone of u vanishes and u is an isomorphism. \square

B.2. Decomposition in $D_{lc}(\mathbb{R})$. Let $\text{Mod}(\mathbb{R})$ be the category of sheaves of k -vector spaces on \mathbb{R} . We consider it as usual as the full subcategory of $D(\mathbb{R})$ of complexes concentrated in degree 0. A sheaf $\mathcal{F} \in \text{Mod}(\mathbb{R})$ which is constructible with respect to a finite stratification of \mathbb{R} is described by a quiver representation of type A_n . Hence Gabriel theorem implies that \mathcal{F} is decomposed as $\mathcal{F} \simeq \bigoplus_{I \in \mathcal{I}} k_I$ where \mathcal{I} is a finite family of intervals. In this section we extend this result to γ_g -limits of constructible sheaves under the assumption that the microsupport is contained in $\{\tau \geq 0\}$.

Let Int^+ be the set of intervals of the form $[a, b[$, $a \in \mathbb{R}$, $b \in \mathbb{R} \cup \{+\infty\}$. We endow Int^+ with the order

$$(B.3) \quad [a, b[\leq [a', b'[\quad \text{if and only if} \quad a \leq a' \text{ and } b \leq b'$$

By Lemma A.2, for $I, I' \in \text{Int}^+$ we have $\text{Hom}(k_I, k_{I'}) \simeq k$ canonically if and only if $I \cap I' \neq \emptyset$ and $I \leq I'$ and in this case we denote by $e_I^{I'}$ its generator. Otherwise $\text{Hom}(k_I, k_{I'}) \simeq 0$ and we set $e_I^{I'} = 0$. For $I \leq I' \leq I''$ we have $e_{I'}^{I''} \circ e_I^{I'} = e_I^{I''}$. For two finite families $\mathcal{I}, \mathcal{I}' \subset \text{Int}^+$ and $\mathcal{G} = \bigoplus_{I \in \mathcal{I}} k_I$, $\mathcal{G}' = \bigoplus_{J \in \mathcal{I}'} k_J$, we obtain a natural morphism

$$E = E_{\mathcal{I}'}^{\mathcal{I}} : \text{Mat}_{l_t}(\mathcal{I}' \times \mathcal{I}) \rightarrow \text{Hom}(\mathcal{G}, \mathcal{G}'), \quad (a_{I', I})_{I' \in \mathcal{I}', I \in \mathcal{I}} \mapsto \sum a_{I', I} e_{I'}^{I'}$$

where Mat_{I_t} denotes the space of lower triangular matrices ($a_{I',I} \neq 0$ implies $I \leq I'$). The map E is surjective and, for three families $\mathcal{I}, \mathcal{I}', \mathcal{I}''$, we have $E^{\mathcal{I}''}(B) \circ E^{\mathcal{I}'}(A) = E^{\mathcal{I}''}(BA)$.

Lemma B.10. *Let $\mathcal{I}, \mathcal{I}' \subset \text{Int}^+$ be two finite families and $\mathcal{G} = \bigoplus_{I \in \mathcal{I}} k_I$, $\mathcal{G}' = \bigoplus_{J \in \mathcal{I}'} k_J$. Let $\varepsilon > 0$ be given and let $u: \mathcal{G} \rightarrow \mathcal{G}'$, $v: \mathcal{G}' \rightarrow T_{\varepsilon^*}(\mathcal{G})$ be such that $v \circ u = \tau_{0,\varepsilon}(\mathcal{G})$. We assume that each $I \in \mathcal{I}$ is of length $> \varepsilon$. Then there exist an isomorphism $\phi: \mathcal{G}' \xrightarrow{\sim} \mathcal{G}'$ and an injective map $\sigma: \mathcal{I} \rightarrow \mathcal{I}'$ such that $I \leq \sigma(I)$ for all $I \in \mathcal{I}$ (for the order \leq defined in (B.3)) and $\phi \circ u: \mathcal{G} \rightarrow \mathcal{G}'$ is the sum of the natural morphisms $k_I \rightarrow k_{\sigma(I)}$.*

Proof. We set $\mathcal{I}'' = \{T_\varepsilon(I); I \in \mathcal{I}\}$. In view of the discussion after (B.3) we can find lower triangular matrices $A \in \text{Mat}_{I_t}(\mathcal{I}' \times \mathcal{I})$, $B \in \text{Mat}_{I_t}(\mathcal{I}'' \times \mathcal{I}')$ representing u, v . Then $BA \in \text{Mat}_{I_t}(\mathcal{I}'' \times \mathcal{I})$ is lower triangular. We identify \mathcal{I} and \mathcal{I}'' through T_ε and we claim that $BA \in \text{Mat}_{I_t}(\mathcal{I} \times \mathcal{I})$ is still lower triangular. We take $I_1, I_2 \in \mathcal{I}$ and assume that $0 \neq (BA)_{I_1, I_2} = \sum_{J \in \mathcal{I}'} B_{T_\varepsilon(I_1), J} A_{J, I_2}$. There exists J such that $B_{T_\varepsilon(I_1), J}, A_{J, I_2} \neq 0$, hence $I_2 \leq J \leq T_\varepsilon(I_1)$. If we assume moreover that $I_2 \not\leq I_1$ we must have $T_\varepsilon(I_1) \cap I_2 \neq \emptyset$ because I_1 and I_2 are of length $> \varepsilon$. Hence $e_{I_2}^{T_\varepsilon(I_1)} \neq 0$ and we deduce that the decomposition of $v \circ u$ on the basis of the non zero $e_I^{I''}$'s has a non zero coefficient on $e_{I_2}^{T_\varepsilon(I_1)}$. Since $v \circ u = \tau_{0,\varepsilon}(\mathcal{G})$ this means that $I_1 = I_2$, contradicting $I_2 \not\leq I_1$. Hence $I_2 \leq I_1$, as claimed.

Finally BA is lower triangular (in $\text{Mat}_{I_t}(\mathcal{I} \times \mathcal{I})$) and has entries 1 along the diagonal since it represents $\tau_{0,\varepsilon}(\mathcal{G})$. Hence BA is invertible. In particular $|\mathcal{I}| \leq |\mathcal{I}'|$, A has rank $|\mathcal{I}|$, we can decompose \mathcal{I}' as $\mathcal{I}' = \mathcal{I} \sqcup \mathcal{I}'_1$ and find an invertible matrix $A' \in \text{Mat}_{I_t}(\mathcal{I}' \times \mathcal{I}')$ such that the block $\mathcal{I} \times \mathcal{I}$ in $A'A$ is the identity matrix. Now we let σ be the map given by the partition $\mathcal{I}' = \mathcal{I} \sqcup \mathcal{I}'_1$ and ϕ be the morphism induced by A' . \square

Proposition B.11. *Let $\mathcal{F} \in \text{Mod}(\mathbb{R})$ such that $\text{SS}(\mathcal{F}) \subset \{(t; \tau) \in T^*\mathbb{R}; \tau \geq 0\}$. We assume that \mathcal{F} is a γ_τ -limit of constructible (for finite stratifications) sheaves $\mathcal{F}_n \in \text{Mod}(\mathbb{R})$ such that $\text{SS}(\mathcal{F}_n) \subset \{(t; \tau) \in T^*\mathbb{R}; \tau \geq 0\}$. Then there exists an at most countable set of intervals \mathcal{I} such that $\mathcal{F} \simeq \bigoplus_{I \in \mathcal{I}} k_I$.*

We notice that the singular support hypothesis in the proposition implies that our intervals are of the type $[a, b]$, $[a, \infty[$ or $]-\infty, b]$.

Proof. (i) Up to taking a subsequence we can assume $\gamma_\tau(\mathcal{F}, \mathcal{F}_n) \leq \varepsilon_n = 2^{-n}$. For each n there exists a finite family $\mathcal{I}_n \subset \text{Int}^+$ such that $\mathcal{F}_n := \bigoplus_{I \in \mathcal{I}_n} k_I$. By Proposition 6.25 we can assume, up to translating \mathcal{F}_n by less than ε_n , that there exist compatible morphisms $f_n: \mathcal{F}_n \rightarrow \mathcal{F}_{n+1}$ and that $\text{hocolim} \mathcal{F}_n \xrightarrow{\sim} \mathcal{F}$. Since $\mathcal{F}_n \in \text{Mod}(\mathbb{R})$, we have $\text{hocolim} \mathcal{F}_n \simeq \varinjlim \mathcal{F}_n$. We can also assume (maybe changing ε_n by $4\varepsilon_n$) that there exist morphisms $g_n: \mathcal{F}_{n+1} \rightarrow T_{\varepsilon_n^*}(\mathcal{F}_n)$ such that $g_n \circ f_n = \tau_{0,\varepsilon_n}(\mathcal{F}_n)$.

(ii) It remains to check that $\varinjlim \mathcal{F}_n$ is decomposed. For this we modify the f_n 's using Lemma B.10. We let $\mathcal{J}_n \subset \mathcal{I}_n$ be the set of intervals of length $> 2\varepsilon_n$ and we set $\widetilde{\mathcal{F}}_n = \bigoplus_{I \in \mathcal{J}_n} k_I$. We first put the f_n 's in "diagonal form", \widehat{f}_n (see the diagram (B.4)). Lemma B.10 applied with $f_0|_{\widetilde{\mathcal{F}}_0} : \widetilde{\mathcal{F}}_0 \rightarrow \mathcal{F}_1$ gives an injective map $\sigma_0 : \mathcal{J}_0 \rightarrow \mathcal{J}_1$ and an isomorphism $\phi_1 : \mathcal{F}_1 \xrightarrow{\sim} \mathcal{F}_1$ such that $\phi_1 \circ f_0|_{\widetilde{\mathcal{F}}_0}$ has a "diagonal form" (that is, it is the sum of the natural morphisms $k_I \rightarrow k_{\sigma_0(I)}$). We set $\widehat{f}_0 = \phi_1 \circ f_0$ and $f'_1 = f_1 \circ \phi_1^{-1}$. Now we apply the lemma with $f'_1|_{\widetilde{\mathcal{F}}_1} : \widetilde{\mathcal{F}}_1 \rightarrow \mathcal{F}_2$ and obtain $\phi_2 : \mathcal{F}_2 \xrightarrow{\sim} \mathcal{F}_2$ and $\sigma_1 : \mathcal{J}_1 \rightarrow \mathcal{J}_2$. We set $\widehat{f}_1 = \phi_2 \circ f'_1$ and $f'_2 = f_2 \circ \phi_2^{-1}$. We go on inductively and end up with a sequence of injective maps $\sigma_n : \mathcal{J}_n \rightarrow \mathcal{J}_{n+1}$ and morphisms $\widehat{f}_n : \widetilde{\mathcal{F}}_n \rightarrow \mathcal{F}_{n+1}$ such that \widehat{f}_n restricted to $\widetilde{\mathcal{F}}_n$ has a "diagonal form".

$$(B.4) \quad \begin{array}{ccccc} \widetilde{\mathcal{F}}_0 & \equiv & \bigoplus_{I_0 \in \mathcal{J}_0} k_{I_0} & \longrightarrow & \bigoplus_{I_0 \in \mathcal{J}_0} k_{I_0} & \equiv & \mathcal{F}_0 \\ \widetilde{f}_0 \downarrow & & \searrow & & \downarrow \widehat{f}_0 & & \downarrow f_0 \\ \widetilde{\mathcal{F}}_1 & \equiv & \bigoplus_{I_1 \in \mathcal{J}_1} k_{I_1} & \longrightarrow & \bigoplus_{I_1 \in \mathcal{J}_1} k_{I_1} & \xrightarrow[\sim]{\phi_1} & \mathcal{F}_1 \\ \widetilde{f}_1 \downarrow & & \searrow & & \downarrow \widehat{f}_1 & & \downarrow f_1 \\ \widetilde{\mathcal{F}}_2 & \equiv & \bigoplus_{I_2 \in \mathcal{J}_2} k_{I_2} & \longrightarrow & \bigoplus_{I_2 \in \mathcal{J}_2} k_{I_2} & \xrightarrow[\sim]{\phi_2} & \mathcal{F}_2 \\ \widetilde{f}_2 \downarrow & & \searrow & & \downarrow \widehat{f}_2 & & \downarrow f_2 \\ \widetilde{\mathcal{F}}_3 & \equiv & \bigoplus_{I_3 \in \mathcal{J}_3} k_{I_3} & \longrightarrow & \bigoplus_{I_3 \in \mathcal{J}_3} k_{I_3} & \xrightarrow[\sim]{\phi_3} & \mathcal{F}_3 \end{array}$$

(iii) With our definition of \mathcal{J}_n we have moreover $\sigma_n(\mathcal{J}_n) \subset \mathcal{J}_{n+1}$. Indeed, modifying g_n as $\widehat{g}_n = T_{\varepsilon_n^*}(\phi_n) \circ g_n \circ \phi_{n+1}^{-1}$, we see that $\widehat{g}_n \circ \widehat{f}_n = \tau_{0, \varepsilon_n}(\mathcal{F}_n)$. It follows that, for any $[a, b[\in \mathcal{J}_n$ and $[a', b'[= \sigma_n([a, b[$, we must have $a \leq a' \leq a + \varepsilon_n$ and $b \leq b' \leq b + \varepsilon_n$. Hence $b' - a' \geq b - a - \varepsilon_n \geq 2\varepsilon_n - \varepsilon_n = 2\varepsilon_{n+1}$, which shows that $[a', b'[\in \mathcal{J}_{n+1}$.

We can then define $\widetilde{f}_n = \widehat{f}_n|_{\widetilde{\mathcal{F}}_n} : \widetilde{\mathcal{F}}_n \rightarrow \widetilde{\mathcal{F}}_{n+1}$ and \widetilde{f}_n has a "diagonal form" with respect to the inclusion $\sigma_n|_{\mathcal{J}_n} : \mathcal{J}_n \hookrightarrow \mathcal{J}_{n+1}$. For a given $I \in \mathcal{J}_n$ we write $\sigma_k \circ \sigma_{k-1} \circ \dots \circ \sigma_n(I) = [a_k, b_k[$; the sequences (a_k) , (b_k) are non decreasing and have limits, say a_∞ , b_∞ . Then $\varinjlim k_{[a_k, b_k[} \simeq k_{[a_\infty, b_\infty[}$. We set $I_\infty = [a_\infty, b_\infty[$. We let \mathcal{J} be the increasing union of the \mathcal{J}_n 's. Then each element of \mathcal{J} corresponds to an interval I_∞ as above and we have proved $\varinjlim \widetilde{\mathcal{F}}_n \simeq \bigoplus_{I_\infty \in \mathcal{J}} k_{I_\infty}$. By Lemma 6.21 the maps $\widetilde{\mathcal{F}}_k \rightarrow \varinjlim \widetilde{\mathcal{F}}_n$ and $\mathcal{F}_k \rightarrow \varinjlim \mathcal{F}_n$ have cones which go to 0 when $k \rightarrow \infty$. It follows by Lemma 6.4 that the natural map $\varinjlim \widetilde{\mathcal{F}}_n \rightarrow \varinjlim \mathcal{F}_n$ has a cone which is arbitrarily close to 0, hence it is an isomorphism. \square

Corollary B.12. *We recall that k is a field. Let $\mathcal{F} \in D_{lc}(\mathbb{R}) \cap D_{\tau \geq 0}(\mathbb{R})$ be a limit of constructible (for finite stratifications) objects of $D_{\tau \geq 0}(\mathbb{R})$. Then there exists an at most countable set of intervals \mathcal{I} and integers d_I , $I \in \mathcal{I}$, such that $\mathcal{F} \simeq \bigoplus_{I \in \mathcal{I}} k_I[d_I]$.*

Proof. As in the case of constructible sheaves the result follows from Proposition B.11 and a decomposition of a complex as sum of its cohomology. However we didn't prove that limits of constructible sheaves have no Ext^2 so we have to be careful to decompose the cohomology before we take the limit, as follows.

There exists a sequence of constructible objects $\mathcal{F}_n \in D_{\tau \geq 0}(\mathbb{R})$ which γ_τ -converges to \mathcal{F} . Since $\text{Ext}^2(\mathcal{G}, \mathcal{G}') = 0$ for any two constructible $\mathcal{G}, \mathcal{G}' \in \text{Mod}(\mathbb{R})$, we have $\mathcal{F}_n \simeq \bigoplus_{i \in \mathbb{Z}} H^i \mathcal{F}_n[-i]$ (see [KS06, Cor. 13.1.20] for example).

In $D_{\tau \geq 0}(\mathbb{R})$ we have $K_\varphi^a(-) = T_{a^*}(-)$, where T_a is as usual the translation by a , and this functor commutes with the cohomology, that is, $H^i(T_{a^*}(\mathcal{G})) \simeq T_{a^*}(H^i(\mathcal{G}))$. It follows that $\gamma_\tau(H^i \mathcal{G}, H^i \mathcal{G}') \leq \gamma_\tau(\mathcal{G}, \mathcal{G}')$, for any $\mathcal{G}, \mathcal{G}' \in D_{\tau \geq 0}(\mathbb{R})$. Hence $H^i \mathcal{F}_n$ γ_τ -converges to $H^i \mathcal{F}$.

It follows that $\mathcal{F}_n \simeq \bigoplus_i H^i \mathcal{F}_n[-i]$ γ_τ -converges to $\bigoplus_i H^i \mathcal{F}[-i]$. By uniqueness of the limit (Proposition B.8) we have $\mathcal{F} \simeq \bigoplus_i H^i \mathcal{F}[-i]$. Now the result follows from Proposition B.11. \square

APPENDIX C. COMPARING DIFFERENT INTERLEAVING METRICS ON $D(N \times \mathbb{R})$.

To prove Theorem 1.2 we have used the distance γ_τ on sheaves over $N \times \mathbb{R}$ and the fact that its restriction to sheaves associated with Hamiltonian maps coincides with the spectral distance (see Proposition 6.14). We can also consider the distance γ_g on sheaves, where g is some Riemannian metric on N . For the sake of completeness we describe the distance γ_g from the point of view of spectral distance and we prove that the distances γ_τ and γ_g induce equivalent topologies (when restricted to sheaves associated with Hamiltonian maps).

In the following we let $H: T^*N \rightarrow \mathbb{R}$ be a function which coincides with $(q, p) \mapsto \|p\|_g$ outside a compact neighbourhood of 0_N and $\varphi = \varphi_H$ denotes its Hamiltonian flow. We will use spectral invariants and sheaves for φ^s but in §4 and §5 we only defined them for compactly supported isotopies. However we can extend them to a positive Hamiltonian H as follows. Let H_r be a truncation of H to the disc bundle of T^*N of radius r , so that H_r is an increasing sequence of Hamiltonians converging to H . Then the homogeneous lift of $s \mapsto \varphi_{H_r}^{-s} \circ \varphi_{H_{r+1}}^s$ to $T^*(N \times \mathbb{R}) \setminus 0_{N \times \mathbb{R}}$ is a non negative Hamiltonian isotopy. Hence we have a natural map $\mathcal{K}_{\text{id}} \rightarrow \mathcal{K}_{\varphi_{H_r}^{-s}} \circ \mathcal{K}_{\varphi_{H_{r+1}}^s}$, or equivalently, $\mathcal{K}_{\varphi_{H_r}^s} \rightarrow \mathcal{K}_{\varphi_{H_{r+1}}^s}$. We define \mathcal{K}_{φ_s}

as the homotopy colimit of this inductive system (say r runs over the integers). In the same way $c_+(\varphi_r^{-s}\psi)$ is a decreasing sequence, and we say that $c_+(\varphi^{-s}\psi) = 0$ if $\lim_r c_+(\varphi_r^{-s}\psi) = 0$. We shall prove that the sequence is in fact stationary, so this means $c_+(\varphi_r^{-s}\psi) = 0$ for r large enough.

Now Lemma 5.9 extends as follows: there exist morphisms $\mathcal{K}_\psi \rightarrow \mathcal{K}_{\varphi^s} \rightarrow \mathcal{K}_{\varphi^{2s} \circ \psi}$ if and only if $c_-(\varphi^s\psi) = 0$ and $c_-(\varphi^s\psi^{-1}) = 0$. Using Corollary 6.10 we obtain

Lemma C.1. *If $c_-(\varphi^s\psi) = 0$ and $c_-(\varphi^s\psi^{-1}) = 0$, the composition of the map of the map $\mathcal{K}_\psi \rightarrow \mathcal{K}_{\varphi^s}$ and $\mathcal{K}_{\varphi^s} \rightarrow \mathcal{K}_{\varphi^{2s} \circ \psi}$ coincides with the canonical map $\mathcal{K}_\psi \rightarrow \mathcal{K}_{\varphi^{2s}\psi}$.*

Now we may define the distance γ^g on $\text{Ham}(T^*N)$ as follows

Definition C.2. *We set*

$$c_-^g(\psi) = -\inf\{s \geq 0 \mid c_-(\varphi^s\psi) = 0\}$$

and

$$c_+^g(\psi) = -c_-^g(\psi^{-1}) = \inf\{s \geq 0 \mid c_+(\varphi^{-s}\psi) = 0\}$$

Finally we set

$$\gamma^g(\psi) = c_+^g(\psi) - c_-^g(\psi)$$

Proposition C.3. *We have*

- (1) *The function γ^g defines a metric by $\gamma^g(\psi_1, \psi_2) = \gamma^g(\psi_1\psi_2^{-1})$.*
- (2) *The topologies defined by γ^g and γ coincide.*

Proof. (1) Apply section 3 of [PS21] to the Kernels \mathcal{K}_{φ^s} (or the symplectic flow φ^s).

(2) We shall first prove that $c_+(\psi) \leq C_W c_+^g(\psi)$ and $\gamma(\psi) \leq C_W \gamma^g(\psi)$ for all ψ such that $\text{supp}(\psi) \subset K$. We consider $c_+(\varphi^{-s}\psi)$ and remember that φ^s is generated by a positive homogeneous Hamiltonian H . By a classical argument (see e.g. [Vit92], lemma 4.7 p. 699) the map $s \mapsto c_+(\varphi^{-s}\psi)$ is piecewise C^1 and where it is C^1 we have

$$\frac{d}{ds} c_+(\varphi^{-s}\psi)|_{s=s_0} = -K_s(z)$$

where K_s is the Hamiltonian associated to the map $\varphi^{-s}\psi$ and z is some fixed point of $\varphi^{-s}\psi$ having non-negative action. Indeed, if S_s is a Generating function Quadratic at infinity for $\varphi^{-s}\psi$ we know that $c_+(\varphi^{-s}\psi)$ is a critical value of S_s and is non-negative. It thus corresponds to fixed points of $\varphi^{-s}\psi$ with non-negative action and its derivative is $\frac{d}{ds} S_s(\xi_z)$ where ξ_z corresponds to z , one of the fixed points such that $S_s(\xi_z) = c_+(\varphi^{-s}\psi)$.

In our case, if z is outside the support of ψ , the fixed point is a fixed point of φ^{-s} and these have action 0 since H is 1-homogeneous (as on an

orbit $p\dot{q} - H(q, p) = p\frac{\partial H}{\partial p} - H(q, p) = 0$). So we only need to estimate H in the support of ψ and we set W to be a neighbourhood of this support. Then if $C_W = \sup\{H(x, p) \mid (x, p) \in W\}$ we have

$$\frac{d}{ds} c_+(\varphi^{-s}\psi)|_{s=s_0} \geq -C_W$$

so that

$$c_+(\varphi^{-s}\psi) - c_+(\psi) \geq -C_W s$$

and if $c_+(\varphi^{-s}\psi) = 0$ we have

$$C_W s \geq c_+(\psi)$$

hence

$$c_+(\psi) \leq C_W c^g(\psi)$$

Applying the same to ψ^{-1} , we get

$$\gamma(\psi) \leq C_W \gamma^g(\psi)$$

Note that we also proved that $\gamma(\varphi_r^{-s}\psi)$ is stationary for r such that $B(0, r)$ contains the support of ψ .

Conversely, let us write $\varphi^s = \xi_r^s \circ \eta_r^s$ where ξ_r^s (resp. η_r^s) is the flow of $f_r(H)$ (resp. $(1 - f_r)(H)$) where f_r is a function given by

- (1) $f_r(t) = t$ for $t \geq 2r$
- (2) $f_r = \frac{3r}{2}$ for $t \leq r$
- (3) f_r is non-decreasing

Note that the flows ξ_r^s, η_r^s are autonomous, commute, and $f_r(H)$ is bounded from below by $\frac{3r}{2}$ and η_r^s is supported in a ball of radius $K \cdot r$.

Then

$$c_+(\varphi^{-s}\psi) = c_+(\xi_r^{-s}\eta_r^{-s}\psi) \leq c_+(\eta_r^{-s}\psi) - \frac{3r}{2}s$$

as long as $c_+(\varphi^{-s}\psi) > 0$ so we have


$$\frac{3r}{2} c_+(\psi) \leq c_+(\eta_r^{-s}\psi) \leq \pi K^2 \cdot r^2 + c_+(\psi)$$

As a result if $c_+(\psi_j)$ converges to zero, let $r_j = \sqrt{c_+(\psi_j)}$. Then

$$c_+^g(\psi_j) \leq \frac{2}{3} \left(\pi K \cdot r_j + \frac{c_+(\psi_j)}{r_j} \right) \leq \frac{2}{3} (\pi K + 1) \sqrt{c_+(\psi_j)}$$

and we deduce that $\lim_j c_+^g(\psi_j) = 0$. □

Remark C.4. (1) We do not know if the metrics γ and γ^g are equivalent, only that they define the same topology.

FIGURE 6. The function f_r

- (2) Here is an analogy for the difference between γ and γ^g . Consider the following norm on $C_b^0(X, \mathbb{R})$: we first define

$$\max_f(g) = \inf \{s \geq 0 \mid \max(g - sf) \leq 0\}$$

and

$$\min_f(g) - \max_f(-g) = \inf \{s \geq 0 \mid \min(g + sf) \geq 0\}$$

Then we set $\|g\|_f = \max_f(g) - \min_f(g)$. Provided

$$0 < \inf_{x \in X} f(x) \leq \sup_{x \in X} f(x) < +\infty$$

it is obvious that $\|\cdot\|_f$ and $\|\cdot\|$ are equivalent.

APPENDIX D. LOCAL FLOER COHOMOLOGY

Our goal is to prove the following statement, which was used in Subsection 5.2.

Let φ be a Hamiltonian map generated by a Hamiltonian supported in a sufficiently small Darboux chart, V . Since $\Gamma(\varphi)$ coincides with the diagonal, $\Gamma(\text{Id})$ outside $W = V \times V$, we can consider two types of spectral numbers.

Consider φ^t as a compact supported Hamiltonian isotopy defined on \mathbb{R}^{2n} and denote by $\tilde{\varphi}^t$ this isotopy. Let $\tilde{S}(q, P, \xi)$ be a Generating Function Quadratic at Infinity for $\Gamma(\tilde{\varphi})$ that we compactify as a Lagrangian in

T^*S^{2n} . In the sequel we will shorten Generating Function Quadratic at Infinity to G.F.Q.I. and refer to [Vit92] for their properties.

Let $S : \Delta_M \times \mathbb{R}^k \rightarrow \mathbb{R}$ be a G.F.Q.I. for $\Gamma(\varphi)$ over Δ_M . Then we want to prove that

$$c(1_{S^{2n}}, \tilde{S}) = c(1_\Delta, S)$$

Since by [Vit95] $c(1_\Delta, S)$ corresponds to the $c_-(\varphi)$ obtained by using Floer cohomology (note that this is also a consequence of Theorem 5.2, (3)), this will imply

$$c_-(\tilde{\varphi}^1) = c(1_{S^{2n}}, \tilde{S}) = c(1_\Delta, S) = c_-(\varphi^1)$$

We are thus in a special case of the more general situation where L is Hamiltonianly isotopic to the zero section 0_N by a Hamiltonian isotopy supported in W

Definition D.1. Let $S(q, P, \xi)$ be a G.F.Q.I. for $\Gamma(\varphi)$ over a bounded connected codimension 0 submanifold W . We set $c_-^W(\varphi) = c(1_W, S)$ where $1_W \in H^0(W)$. the spectral number $c(1_W, S)$ was defined in [Vit21], Definition 5.1 and $c(\mu_W, S) = c_+^W(L)$ where $\mu_W \in H^n(W, \partial W)$ is the fundamental class. They are obtained by setting

(1) For $\alpha \in H^*(W)$

$$c(\alpha, S) = \inf \left\{ t \mid T \otimes \alpha \neq 0 \text{ in } H^*(S_{|W}^t, S_{|W}^{-\infty}) \right\}$$

(2) For $\alpha \in H_c^*(W) = H^*(W, \partial W)$

$$c(\alpha, S) = \inf \left\{ t \mid T \otimes \alpha \neq 0 \text{ in } H^*(S_{|W}^t, S_{|W}^{-\infty} \cup S_{|\partial W}^t) \right\}$$

Proposition D.2. Let W be a codimension zero connected submanifold of the closed manifold N . Let φ^t be an isotopy supported in $T^*W \subset T^*N$. Let $S(q; \xi)$ be a G.F.Q.I. for $L = \varphi^1(0_N)$ normalized so that outside W the critical points of S have critical value 0. Set $c_-^W(L) = c(1_W, S)$. Then we have $c_-^W(L) = c_-(L)$ and by duality $c_+^W(L) = c_+(L)$.

Proof. We prove the first equality. Since L coincides with the zero section outside of W , we know that $c_-(L) \leq 0$. Let ψ^t be flow of a fiber preserving vector field on $S^{-\varepsilon}$ where ε is some small positive real number. Let ψ^t be a flow on $N \times \mathbb{R}^k$ such that

(1) ψ^t is the flow of $-\frac{\nabla_\xi S(x, \xi)}{|\nabla_\xi S(x, \xi)|^2}$ outside of $W \times \mathbb{R}^k$

(2) $S(\psi^t(x, \xi)) \leq S(x, \xi)$ for $t \geq 0$

This is easy to construct by noticing that in $(N \setminus W) \times \mathbb{R}^k \cap S^{-\varepsilon}$ the vector field $-\frac{\nabla_\xi S(x, \xi)}{|\nabla_\xi S(x, \xi)|^2}$ is well defined, since S has no fiberwise critical points in $(N \setminus W) \times \mathbb{R}^k$. Note that ψ^t is then obtained by multiplying this vector field by a smooth function equal to 1 on $(N \setminus W) \times \mathbb{R}^k \cap S^{-\varepsilon}$ and vanishing

outside a neighbourhood of it. Remember that we denote by $S^{-\infty}$ the set S^{-c} for some large c . Then for $t < -\varepsilon$ we have

$$H^*(S^t, S^{-\infty}) = H^*(\psi^s(S^t), \psi^s(S^{-\infty}))$$

but since for s large enough

$$\psi^s(S^t) \subset S^t_{|W \times \mathbb{R}^k} \cup S^{-\infty}$$

we have by excision,

$$H^*(\psi^s(S^t), \psi^s(S^{-\infty})) \simeq H^*(\psi^s(S^t_{|W \times \mathbb{R}^k}), \psi^s(S^{-\infty}_{|W \times \mathbb{R}^k}))$$

and applying the isotopy ψ^{-s} we get that for $t < -\varepsilon$

$$H^*(S^t, S^{-\infty}) \simeq H^*(S^t_{|W \times \mathbb{R}^k}, S^{-\infty}_{|W \times \mathbb{R}^k})$$

We therefore have the following diagram for $t < -\varepsilon$

$$\begin{array}{ccc} H^{*-d}(N) & \longrightarrow & H^{*-d}(W) \\ \downarrow \simeq & & \downarrow \simeq \\ H^*(S^\infty, S^{-\infty}) & \longrightarrow & H^*(S^\infty_{|W \times \mathbb{R}^k}, S^{-\infty}_{|W \times \mathbb{R}^k}) \\ \downarrow & & \downarrow \\ H^*(S^t, S^{-\infty}) & \xrightarrow{\simeq} & H^*(S^t_{|W \times \mathbb{R}^k}, S^{-\infty}_{|W \times \mathbb{R}^k}) \end{array}$$

Since the top horizontal map sends 1_N to 1_W , we have that (when $t < -\varepsilon$) the composition of the two vertical left-hand side maps sends 1_N to zero if and only if the composition of the two vertical right-hand side maps sends 1_W to zero. In other words if $t < -\varepsilon$ and $t < c(1_N, S)$ then $t \leq c(1_W, S)$ so that $\min\{c(1_N, S), -\varepsilon\} \leq c(1_W, S)$ and conversely if $t < -\varepsilon$ and $t < c(1_W, S)$ then $t \leq c(1_N, S)$ in other words $\min\{c(1_W, S), -\varepsilon\} \leq c(1_N, S)$. It is however well-known that $c(1_W, S) \leq 0$ and $c(1_N, S) \leq 0$ and since $\varepsilon > 0$ is arbitrarily small, we get $c(1_W, S) = c(1_N, S)$. \square

From this it immediately follows

Corollary D.3. *Let U be a codimension 0 submanifold of the closed manifolds N_1, N_2 and $L_1 = \varphi_1^1(0_{N_1})$, $L_2 = \varphi_2^1(0_{N_2})$ where $\varphi_1^t = \text{Id}$ and $\varphi_2^t = \text{Id}$ outside of T^*U and $\varphi_1^t = \varphi_2^t$ on T^*U . Then*

$$c_{\pm}^1(L_1) = c_{\pm}^2(L_2)$$

In other words c_{\pm} does not depend on the extension of $\varphi_1^1(0_U) = \varphi_2^1(0_U)$.

Applying this to $\Gamma(\varphi)$ we get, provided the small Darboux chart is such that $V \times V$ is contained in a Weinstein neighbourhood of the diagonal (i.e.

a neighbourhood of the diagonal that is symplectomorphic to a neighbourhood of the zero section in $T^*\Delta$). Using the remark at the beginning of this Appendix, we get

Corollary D.4. *Let φ^t be a Hamiltonian isotopy of (M, ω) supported in a small Darboux chart and consider φ^t as a compact supported Hamiltonian isotopy defined on \mathbb{R}^{2n} . Denote by $\tilde{\varphi}^t$ this isotopy and $\tilde{S}(q, P, \xi)$ be a G.F.Q.I. for $\Gamma(\tilde{\varphi})$ that we compactify as a Lagrangian in T^*S^{2n} . Let $S : \Delta_M \times \mathbb{R}^k \rightarrow \mathbb{R}$ be a G.F.Q.I. for $\Gamma(\varphi)$ over Δ_M . Then*

$$c_-(\tilde{\varphi}^1) = c(1_{S^{2n}}, \tilde{S}) = c(1_{\Delta_M}, S) = c_-(\varphi^1)$$

We similarly get

$$c_+(\tilde{\varphi}^1) = c_+(\varphi^1)$$

APPENDIX E. DUALITY FOR SHEAF SPECTRAL INVARIANTS AND CONNECTION WITH FLOER'S SPECTRAL THEORY

Our goal in this Appendix is twofold. On one hand we look for a replacement of the duality yielding the equality $c_-(\varphi) = -c_+(\varphi^{-1})$ in the sheaf framework. As will be clear the duality involves spectral invariants associated to compact supported cohomology classes. On the other hand we shall prove in general that the Floer cohomology of a symplectic map is isomorphic to the corresponding sheaf cohomology. In particular this implies that the spectral invariants coincide, which we only proved for symplectic maps supported in a Darboux ball in Subsection 5.2.

Recall that the Tamarkin category $\mathcal{D}(N)$ is the left orthogonal of $D_{\{\tau \leq 0\}}(k_{N \times \mathbb{R}})$ and the convolution $k_{N \times [0, +\infty[} * - : D(k_{N \times \mathbb{R}}) \rightarrow D(k_{N \times \mathbb{R}})$ is a projector onto $\mathcal{D}(N)$. We can define a duality in $\mathcal{D}(N)$ as follows. We first recall the usual duality functors, for a manifold M , $D'(F) = R\mathcal{H}om(F, k_M)$ and $D(F) = R\mathcal{H}om(F, \omega_M)$, where $\omega_M \simeq or_M[d_M]$ is the dualizing sheaf. We define $i : N \times \mathbb{R} \rightarrow N \times \mathbb{R}$ by $i(x, t) = (x, -t)$. For $F \in \mathcal{D}(N)$, we set

$$D^T(F) = k_{N \times [0, +\infty[} * i^{-1} D'(F)$$

For $L \in \mathcal{L}(T^*M)$ and $F_L = Q(L)$ we then have $D^T(F_L) \simeq F_{-L}$.

E.1. Not really a duality. If we restrict to constructible objects, we also have a biduality statement $F \xrightarrow{\simeq} D^T(D^T(F))$ and $D^T(-)$ is a contravariant equivalence of categories. We deduce, for F, G constructible

$$R\text{Hom}(F, T_{c^*}(G)) \simeq R\text{Hom}(D^T(T_{c^*}(G)), D^T(F)) \simeq R\text{Hom}(D^T(G), T_{c^*}(D^T(F)))$$

It follows that $c(\alpha, F, G) = c(\alpha, D^T(G), D^T(F))$. In particular $c(\alpha, L, L') = c(\alpha, -L', -L)$.

E.2. Compact support. For $F, G \in \mathcal{D}(N)$ which coincide with $k_{N \times \mathbb{R}}$ near $N \times \{+\infty\}$ we recall that $c(\alpha, F, G)$ is obtained as follows. For $c \leq d$ and d big enough we have

$$u_c^1: \mathrm{RHom}(F, T_{c^*}(G)) \rightarrow \mathrm{RHom}(F, T_{d^*}(G)) \simeq \mathrm{R}\Gamma(N; k_N)$$

and, for $\alpha \in H^i(N; k_N)$, we can define $c(\alpha, F, G)$ as the infimum on the set of c such that α is in the image of $H^i(u_c^1)$. To express a dual statement we set for short

$$\mathrm{RHom}_c(F, G) = \mathrm{R}\Gamma_c(N \times \mathbb{R}; \mathrm{R}\mathcal{H}\mathrm{om}(F, G))[1]$$

Then $\mathrm{RHom}_c(F, T_{d^*}(G)) \simeq \mathrm{R}\Gamma_c(N; k_N)$ for $d \ll 0$. For $d \leq c$ and d small enough we have

$$u_c^2: \mathrm{R}\Gamma_c(N; k_N) \simeq \mathrm{RHom}_c(F, T_{d^*}(G)) \rightarrow \mathrm{RHom}_c(F, T_{c^*}(G))$$

and we can define spectral invariants in this way, say $c'(\alpha, F, G)$, for $\alpha \in H_c^i(N; k_N)$: we let $c'(\alpha, F, G)$ be the infimum on the set of c such that $H^i(u_c^2)(\alpha) = 0$. If N is compact, we have $c'(\alpha, F, G) = c(\alpha, F, G)$: we can check that there is a distinguished triangle

$$\mathrm{RHom}(F, T_{c^*}(G)) \xrightarrow{u_c^1} \mathrm{R}\Gamma(N; k_N) \xrightarrow{u_c^2} \mathrm{RHom}_c(F, T_{c^*}(G)) \xrightarrow{+1}$$

and, taking the long exact sequence, we see that a given $\alpha \in H^*(N)$ belongs to $\mathrm{im}(u_c^1)$ exactly when $u_c^2(\alpha) = 0$.

E.3. Duality. For F, G constructible and with disjoint microsupports we also have $D^* \mathrm{R}\mathcal{H}\mathrm{om}(F, G) \simeq \mathrm{R}\mathcal{H}\mathrm{om}(G, F)$. Tensoring with $p^{-1}\omega_N$ and using the Poincaré-Verdier duality we get $(\mathrm{RHom}(F, G))^* \simeq \mathrm{RHom}_c(G, F \otimes p^{-1}\omega_N)$. Hence for F, G constructible, with microsupports in $\{\tau > 0\}$, and for a generic c we have the commutative diagram

$$\begin{array}{ccc} \mathrm{RHom}(F, T_{c^*}(G)) & \xrightarrow{u_c^1} & \mathrm{R}\Gamma(N; k_N) \\ \left| \wr \right. & & \left| \wr \right. \\ (\mathrm{RHom}_c(G, T_{-c^*}(F) \otimes p^{-1}\omega_N))^* & \xrightarrow{(u_c^2)^*} & \mathrm{R}\Gamma_c(N; \omega_N)^* \end{array}$$

To find $c(1, F, G)$ we look for the values c such that $H^0(u_c^1)$ is surjective and, similarly, $c'(\mu_N, F, G)$ is related with the injectivity of $H^{d_N}(u_c^2)$.

If N is orientable, then $\omega_N \simeq k_N[d_N]$ and we deduce $c(1, F, G) = -c'(\mu_N, G, F)$.

If N is compact and orientable, we obtain finally

$$c(1, F, G) = -c'(\mu_N, G, F) = -c(\mu_N, G, F) = -c(\mu_N, D^T(F), D^T(G))$$

E.4. Case of Hamiltonian maps. Now we work on N^2 and our sheaves $\mathcal{K}, \dots \in \mathcal{D}(N^2)$ represent graphs of Hamiltonian maps. Hence the hypothesis “ $F \simeq k_{N \times \mathbb{R}}$ near $N \times \{+\infty\}$ ” is changed into “ $\mathcal{K} \simeq k_{\Delta_N \times \mathbb{R}}$ near $N^2 \times \{+\infty\}$ ”. Most (iso)morphisms from the above paragraphs §E.1-§E.3 have an analog in this framework. In particular we have, denoting by $\omega_{\Delta_N|N^2}$ is the relative dualizing sheaf, $\omega_{\Delta_N|N^2} \simeq or_N[-d_N]$,

$$\begin{aligned} D^T(\mathcal{K}_\varphi) &\simeq \mathcal{K}_{\varphi^{-1}} \otimes \omega_{\Delta_N|N^2} \\ c(\alpha, \mathcal{K}_\varphi, \mathcal{K}_\psi) &= c(\alpha, D^T(\mathcal{K}_\psi), D^T(\mathcal{K}_\varphi)) = c(\alpha, \mathcal{K}_{\psi^{-1}}, \mathcal{K}_{\varphi^{-1}}) \\ u_c^1: \mathrm{RHom}(\mathcal{K}, T_{c^*}(\mathcal{K}')) &\rightarrow \mathrm{RHom}(\mathcal{K}, T_{d^*}(\mathcal{K}')) \\ &\simeq \mathrm{R}\Gamma(N^2; k_{\Delta_N}) \simeq \mathrm{R}\Gamma(N; k_N) \\ u_c^2: \mathrm{R}\Gamma_c(N^2; k_{\Delta_N}) &\simeq \mathrm{R}\Gamma_c(N; k_N) \simeq \mathrm{RHom}_c(\mathcal{K}, T_{d^*}(\mathcal{K}')) \\ &\rightarrow \mathrm{RHom}_c(\mathcal{K}, T_{c^*}(\mathcal{K}')) \end{aligned}$$

But now we don't have $D' \mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{K}, \mathcal{K}') \simeq \mathrm{R}\mathcal{H}\mathrm{om}(\mathcal{K}', \mathcal{K})$, not even up to shift by d_N (the microsupports are not disjoint even after a small translation). So we don't have duality; instead we get “new” invariants. Let us set

$$\mathrm{RHom}^D(\mathcal{K}, \mathcal{K}') = \mathrm{R}\Gamma_c(N^2 \times \mathbb{R}; \mathrm{DR}\mathcal{H}\mathrm{om}(\mathcal{K}', \mathcal{K}))$$

Then $\mathrm{RHom}^D(\mathcal{K}, T_{d^*}(\mathcal{K}')) \simeq \mathrm{R}\Gamma_c(N^2; \omega_{\Delta_N}) \simeq \mathrm{R}\Gamma_c(N; \omega_N)$ for $d \ll 0$ and we can use this to define $c''(\alpha, \mathcal{K}, \mathcal{K}')$ for $\alpha \in H_c^*(N; \omega_N) \simeq (H^*(N; k_N))^*$ (in the same way as $c'(\alpha, \mathcal{K}, \mathcal{K}')$). So the complete set of invariants for $(\mathcal{K}, \mathcal{K}')$ is now $c(\alpha, \mathcal{K}, \mathcal{K}')$, $c''(\beta, \mathcal{K}, \mathcal{K}')$, for $\alpha \in H^*(N; k_N)$ and $\beta \in H_c^*(N; \omega_N)$.

We have the diagram

$$\begin{array}{ccc} \mathrm{RHom}(\mathcal{K}, T_{c^*}(\mathcal{K}')) & \xrightarrow{u_c^1} & \mathrm{R}\Gamma(N^2; k_{\Delta_N}) \\ \left| \wr \right. & & \left| \wr \right. \\ (\mathrm{RHom}^D(\mathcal{K}', T_{c^*}(\mathcal{K})))^* & \xrightarrow{(u_c^3)^*} & \mathrm{R}\Gamma_c(N^2; \omega_{\Delta_N})^* \end{array}$$

which says that $c''(\mu_N, \mathcal{K}, \mathcal{K}') = -c(1, \mathcal{K}', \mathcal{K})$. In particular $c''(\mu_N, \mathcal{K}_\varphi, \mathcal{K}_\psi) = -c(1, \mathcal{K}_\psi, \mathcal{K}_\varphi) = -c(1, \mathcal{K}_{\varphi^{-1}}, \mathcal{K}_{\psi^{-1}})$.

E.5. Sheaf and Floer cohomology for Hamiltonian maps. Let φ be a Hamiltonian map of T^*N with compact support. Let us consider the cohomology groups $H^*(N \times N \times [a, b], \mathcal{K}_\varphi, \mathcal{K}_\psi)$ by which we mean $H^*(\mathrm{R}\Gamma_{N \times N \times [a, b]}(\mathrm{RHom}(\mathcal{K}_\varphi, \mathcal{K}_\psi)))$.

Similarly we can consider the Floer cohomology $FH^*(\Gamma(\varphi), \Delta_N; a, b)$ that is the Floer cohomology filtered between action values a and b with

$-\infty < a < b < +\infty$. In [Vit19] it is proved that for L a closed Lagrangian in T^*N and F_L its quantized sheaf, we have

$$H^*(N \times [a, b[, F_{L_1}, F_{L_2}) = FH^*(L_1, L_2; a, b)$$

The issue here is that the graph of φ is not closed (it is diffeomorphic to the diagonal of $\overline{T^*N} \times T^*N$). So our goal is to prove

Theorem E.1. *We have for $\varphi \in DHam_c(T^*N)$ that*

$$FH^*(\Gamma(\varphi), \Delta_N; a, b) = H^*_{N \times N \times [a, b[}(RHom(\mathcal{K}_\varphi, \mathcal{K}_\psi))$$

Since the sheaf theoretic version of the spectral invariants $c_s(\alpha, \varphi_1, \varphi_2)$ is obtained by looking at the barcode associated to the persistence module $H^*(R\Gamma_{N \times N \times [a, b[}(RHom(\mathcal{K}_{\varphi_1}, \mathcal{K}_{\varphi_2})))$, we obtain, denoting by $c(\alpha, \varphi_1, \varphi_2)$ the Floer theoretic spectral invariants associated to the persistence module $FH^*(\Gamma(\varphi_1), \Gamma(\varphi_2); a, b)$

Corollary E.2. *We have for $\alpha \in H^*(N) \setminus \{0\}$*

$$c_s(\alpha, \varphi_1, \varphi_2) = c(\alpha, \varphi_1, \varphi_2)$$

The proof of the theorem is based on the analogous result for closed Lagrangians. Indeed, let M be a Riemannian manifold and let S be a submanifold of M , and $d(x, S)$ be the distance from x to S . Let $\chi_\rho : \mathbb{R} \rightarrow \mathbb{R}$ be a function such that

- $\chi_\rho(t) = t$ for $t \leq \rho$
- $\forall t \in \mathbb{R}, 0 \leq \chi'_\rho(t)$
- $\chi'_\rho(t) = 0$ for $t \geq 2\rho$

Finally we set $f_{\varepsilon, \rho} = \chi_\rho\left(\frac{d(x, S)^2}{\varepsilon}\right)$ and $S_{\varepsilon, \rho} = \text{graph}(df_{\varepsilon, \rho})$. Using that


$$df_{\varepsilon, \rho}(x) = \frac{1}{\varepsilon} \chi'_\rho\left(\frac{d(x, S)^2}{\varepsilon}\right) d(x, S) \nu_S(x)$$

where $\nu_S(x)$ is the normal to the level set of $d(x, S)$ through x , i.e. $\nu_S(x) = d(d(x, S)) \in T_x^*M$, we see that, for $0 < A < \sqrt{\rho/\varepsilon}$,

- $S_{\varepsilon, \rho} \cap \{(x, p) \mid |p| \leq A, d(x, S) \leq \sqrt{\varepsilon\rho}\}$ corresponds to $\{x \mid d(x, S) \leq A\varepsilon\}$, and in this region $0 \leq f_{\varepsilon, \rho}(x) \leq A^2\varepsilon$
- Over $A\varepsilon \leq d(x, S) \leq \sqrt{\rho\varepsilon}$ we have $S_{\varepsilon, \rho} \cap \{(x, p) \mid |p| \leq A\} = \emptyset$
- Over $\sqrt{\varepsilon\rho} \leq d(x, S)$ we have $f_{\varepsilon, \rho}(x) \geq \rho$

As a result for a fixed Lagrangian L contained in $|p| \leq A/2$ and such that f_L is bounded by $\rho/2$, the intersection points $L \cap S_{\varepsilon, \rho}$ are

- over $\{x \mid d(x, S) \leq A\varepsilon\}$ and $0 \leq f_{\varepsilon, \rho}(x) \leq A^2\varepsilon$
- $\sqrt{\varepsilon\rho} \leq d(x, S)$ and then $f_{\varepsilon, \rho}(x) \geq \rho$

FIGURE 7. The function $f_{\epsilon, \rho}$

Let us set $V_\epsilon = \text{graph}(df_{\epsilon, \rho}) \cap \{x \mid d(x, S) \leq A\epsilon\}$. As a result, the intersection points of $L \cap D_{\epsilon, \rho}$ contributing to $FH^*(L, D_{\epsilon, \rho}; a, b)$ are, for ϵ small enough, in one-to-one correspondence with the points (x, p) of $L \cap V_\epsilon$ such that $f_L(x, p) \in [a, b]$.

Proposition E.3. *For a, b finite and ρ sufficiently large we have*

$$\lim_{\substack{\epsilon \rightarrow 0 \\ \epsilon' \rightarrow 0}} FH^*((\varphi \times \text{id})D_{\epsilon, \rho}, D_{\epsilon', \rho}; a, b) = FH^*(\Gamma(\varphi), \Delta_{T^*N}; a, b)$$

Proof. Note that $\lim_{\epsilon \rightarrow 0} V_\epsilon = \nu^*S$, the limit being understood as C^1 -convergence on compact subsets. Indeed $df_{\epsilon, \rho}(x) = \frac{1}{\epsilon}d(x, S)\nu_S(x)$ so if $x = \exp_y(t\nu)$ where $\nu \perp T_yS$, and $t \leq A\epsilon$, $df_{\epsilon, \rho}(x) = \frac{t}{\epsilon}\nu_S(x)$ where $\nu_S(x) = d\exp(t\nu)\nu$.

In local coordinates $y \in V, \nu \in (T_yS)^\perp$ we may consider the map

$$(x, \nu) \mapsto (\exp_y(\epsilon\nu), \nu)$$

Set $D_AT^*M = \{(q, p) \in T^*M \mid |p| \leq A\}$. The above map defines a smooth diffeomorphism C^1 close to the identity sending $\nu^*S \cap D_AT^*M$ to $S_{\epsilon, \rho} \cap D_AT^*M$.

Now we set $S = \Delta_N$ and $\nu^*\Delta_N = \Delta_{T^*N}$ and $D_{\epsilon, \rho} = S_{\epsilon, \rho}$. We have to be careful with the definition of $FH^*(\Gamma(\varphi), \Delta_{T^*N}; a, b)$ since $\Gamma(\varphi)$ and Δ_{T^*N} coincide at infinity. Indeed we must then slightly perturb $\Gamma(\varphi)$ so that we have transversality. Then we slightly change the action of the intersection points at infinity. This action is zero, and we replace them with finitely

many points with arbitrarily small action. We assume $a, b \neq 0$ and the perturbation so small that no critical value crosses a or b .

It is now enough to prove that - after this perturbation- the holomorphic strips involved in the definition of $FH^*(\Gamma(\varphi), \Delta_{T^*N}; a, b)$ must remain in $D_A T^*M$. Here a, b are fixed (and finite !), and we may increase A . This follows from Lemma E.4 below.

Finally, since $(\varphi \times \text{id})D_{\varepsilon, \rho} C^1$ converges to $(\varphi \times \text{id})\Delta_{T^*N}$ and $D_{\varepsilon', \rho} C^1$ converges to Δ_{T^*N} , we may conclude our proof. \square

Lemma E.4. *Let N be a complete Riemannian manifold. Let L be a Lagrangian coinciding with 0_N outside a compact set, and f a C^1 bounded function without critical points at infinity (i.e. $10\delta_0 > |df(x)| > \delta_0 > 0$ for x large enough and some $\delta_0 > 0$). We define $FC_J^*(L, 0_N; a, b)$ to be the limit as ε goes to 0 of $FC_J^*(L, \text{gr}(\varepsilon df); a, b)$. Then the holomorphic curves defining $FC_J^*(L, \text{gr}(\varepsilon df); a, b)$ for ε small enough are contained in some bounded set independent from ε .*

Proof. We shall choose an almost complex structure such that outside a bounded set, J is analytic and in a neighbourhood

$$W = D_r(T^*N) = \{(q, p) \in T^*N \mid |p| \leq r\}$$

of 0_N , there is an antisymplectic involution such that $\tau^*J = -J$. This is always possible by taking a complex structure that is a fixed point of $J \mapsto -\tau^*(J)$ where $\tau(q, p) = (q, -p)$. Since the set of almost complex structures is a convex set, such a fixed point for the above involution must exist.

We now claim that if C_ε is a holomorphic strip with boundaries in L and $\varepsilon \text{graph}(df)$, and containing a point x_ε going to infinity as ε goes to 0, then the area of $C_\varepsilon \cap W$ is going to infinity. Indeed, the areas of the curves is bounded by $b - a$, so in the limit ε going to 0 we would get a curve C_0 with boundary in $L \cup 0_N$. Then using the reflection principle (see [Sib85], Theorem 3.5 and also [Chi83; Chi79; IS10]) we get a closed curve in W through a point x_ε and exiting from the ball of radius R_ε centered at x_ε . Here $\lim_{\varepsilon \rightarrow 0} R_\varepsilon = +\infty$ since we only need that in $B(x_\varepsilon, R_\varepsilon)$ $L = 0_N$ and J is analytic and invariant by $-\tau^*$. Then the area of such a curve goes to ∞ , by the standard monotonicity argument⁷. But then the area of C would be infinite. \square

⁷see [Whi16], p. 8 and 9. We have $\frac{\text{area}(B(x_\varepsilon, r) \cap C)}{\pi r^2}$ is increasing with r and greater than 1. A similar statement holds in the general case for r less than the injectivity radius ρ . But we may cover $B(x_\varepsilon, R_\varepsilon)$ by balls of radius ρ , and C will intersect an increasing number of these balls.

Similarly setting $U_{\varepsilon,\rho} = \{(x, t) \mid f_{\varepsilon,\rho}(x) \leq t\}$, and noticing that $SS(k_{U_{\varepsilon,\rho}}) = D_{\varepsilon,\rho}$, we prove

Proposition E.5. *We have*

$$\lim_{\varepsilon \rightarrow 0} \lim_{\varepsilon' \rightarrow 0} H_{N \times N \times [a,b]}^*(RHom(K_{\varphi} \circ k_{U_{\varepsilon,\rho}}, k_{U_{\varepsilon',\rho}})) = H_{N \times N \times [a,b]}^*(RHom(\mathcal{K}_{\varphi}, \mathcal{K}_{id}))$$

Proof. Indeed this follows from the fact that clearly

$$\lim_{\varepsilon \rightarrow 0} k_{U_{\varepsilon,\rho}}(W) = k_{V \times [0, +\infty[}(W)$$

on bounded sets (i.e. bounded in the \mathbb{R} component). But since the direct limits of sheaves are sheafification of the direct limit of the presheaves, we get

$$\lim_{\varepsilon \rightarrow 0} k_{U_{\varepsilon,\rho}} = k_{V \times [0, +\infty[}$$

and in our case

$$\lim_{\varepsilon \rightarrow 0} k_{D_{\varepsilon,\rho}} = k_{\Delta_N \times [0, +\infty[} = \mathcal{K}_{id}$$

Because direct limits is an exact functor, we get

$$\lim_{\varepsilon' \rightarrow 0} H_{N \times N \times [a,b]}^*(RHom(K_{\varphi} \circ k_{U_{\varepsilon,\rho}}, k_{U_{\varepsilon',\rho}})) = H_{N \times N \times [a,b]}^*(RHom(\mathcal{K}_{\varphi} \circ k_{U_{\varepsilon,\rho}}, \mathcal{K}_{id}))$$

and taking the limit as ε goes to 0 we get the Proposition. \square

Proof of Theorem E.1. Now since the quantization of $D_{\varepsilon,\rho}$ is $k_{U_{\varepsilon,\rho}}$ and the quantization of $(\varphi \times id)D_{\varepsilon,\rho}$ is $\mathcal{K}_{\varphi} \circ k_{U_{\varepsilon,\rho}}$ the combination of Proposition E.3 and E.5 yields Theorem E.1 \square

REFERENCES

- [AHV22] M.-C. Arnaud, V. Humilière, and C. Viterbo. *In preparation*. 2022.
- [AI22] T. Asano and Y. Ike. *Completeness of derived interleaving distances and sheaf quantization of non-smooth objects*. 2022. arXiv: 2201.02598 [math.SG].
- [AI20] T. Asano and Y. Ike. *Persistence-like distance on Tamarkin's category and symplectic displacement energy*. *J. Symp. Geometry* 18.3 (2020), pp. 613–649. DOI: DOI : 10 . 4310 / JSG . 2020 . v18 . n3 . a1. arXiv: 1712.06847.
- [AGHIV] T. Asano et al. *Comparison of microsupport and γ -support*. arXiv: 2211.13945 [math.SG].
- [Bar94] S. Barannikov. *The framed Morse complex and its invariants*. *Advances in Soviet Maths* 21 (1994), pp. 96–115. DOI: 10 . 1090 / advsov / 021 / 03.

- [BBD82] A. A. Beilinson, J. Bernstein, and P. Deligne. *Analyse et topologie sur les espaces singuliers (I)*. Vol. 100. Astérisque. Société Mathématique de France, 1982. URL: http://www.numdam.org/item/AST_1982__100__1_0/.
- [BG18] N. Berkouk and G. Ginot. *A derived isometry theorem for sheaves*. 2018. ArXiv: 1805.09694.
- [BN93] M. Bökstedt and A. Neeman. *Homotopy limits in triangulated categories*. *Compositio Mathematica* 86.2 (1993), pp. 209–234. URL: http://www.numdam.org/item/CM_1993__86_2_209_0/.
- [Bou32] G. Bouligand. *Introduction à la géométrie infinitésimale directe*. Paris: Vuibert, 1932.
- [Cha+09] F. Chazal et al. “Proximity of persistence modules and their diagrams”. In: *Proceedings of the Twenty-fifth Annual Symposium on Computational Geometry*. Vol. SCG '09. New-York, 2009, pp. 237–246. URL: <https://hal.inria.fr/inria-00292566v4>.
- [Chi79] E. Chirka. *Removable singularities of analytic sets*. *Dokl. Akad. Nauk SSSR (N.S.)* 248.1 (1979), pp. 47–50.
- [Chi83] E. Chirka. *Regularity of the boundaries of analytic sets*. *Math. USSR Sbornik* 45.3 (1983), pp. 291–355. DOI: 10.1070/SM1983v045n03ABEH001010.
- [Cra15] W. Crawley-Boevey. *Decomposition of pointwise finite-dimensional persistence modules*. *J. Algebra Appl* 14 (2015), p. 1550066. DOI: 10.1142/S0219498815500668.
- [ELZ02] H. Edelsbrunner, D. Letscher, and A. Zomorodian. *Topological Persistence and Simplification*. *Discrete & Computational Geometry* 28 (2002), pp. 511–533. DOI: 10.1007/s00454-002-2885-2.
- [Gab72] P. Gabriel. *Unzerlegbare Darstellungen I*. *Manuscripta Math* 6 (1972), pp. 71–103. DOI: 10.1007/BF01298413.
- [Gui12] S. Guillermou. *Quantization of conic Lagrangian submanifolds of cotangent bundles*. 2012. ArXiv: 1212.5818.
- [GKS] S. Guillermou, M. Kashiwara, and P. Schapira. *Sheaf quantization of Hamiltonian isotopies and applications to non-displaceability problems*. ArXiv: 1005.1517.
- [IS10] S. Ivashkovich and A. Sukhov. *Schwarz Reflection Principle, Boundary Regularity and Compactness for J -Complex Curves*. *Annales de l'Institut Fourier* 60.4 (2010), pp. 1489–1513. URL: http://aif.cedram.org/item?id=AIF_2010__60_4_1489_0.
- [KS90] M. Kashiwara and P. Schapira. *Sheaves on manifolds*. Vol. 292. Grundlehren der Math. Wissenschaften. Springer-Verlag, 1990.

- [KS06] M. Kashiwara and P. Schapira. *Categories and Sheaves*. Vol. 332. Grundlehren der Mathematische Wissenschaften. Springer-Verlag, 2006. DOI: doi.org/10.1007/3-540-27950-4.
- [KS18] M. Kashiwara and P. Schapira. *Persistent homology and microlocal sheaf theory*. 2018. arXiv: 1705.00955 [math.AT].
- [Kra13] T. Kragh. *Parametrized ring-spectra and the nearby Lagrangian conjecture*. Appendix by M. Abouzaid. *Geometry & Topology* 17 (2013), pp. 639–731. DOI: [10.2140/gt.2013.17.639](https://doi.org/10.2140/gt.2013.17.639).
- [Lec08] R. Leclercq. *Spectral invariants in Lagrangian Floer theory*. *Journal of Modern Dynamics* 2 (2008), pp. 249–286. DOI: [10.3934/jmd.2008.2.249](https://doi.org/10.3934/jmd.2008.2.249). URL: <https://www.aims sciences.org/article/doi/10.3934/jmd.2008.2.249>.
- [Nad16] D. Nadler. *Non-characteristic expansions of Legendrian singularities*. 2016. arXiv: 1507.01513 [math.SG].
- [Oh05] Y.-G. Oh. *Construction of spectral invariants of Hamiltonian paths on closed symplectic manifolds*. In: *The breadth of symplectic and Poisson geometry*. Vol. 232. Progr. Math. vol. 232. Boston: Birkhäuser, 2005, pp. 525–570.
- [PS21] F. Petit and P. Schapira. *Thickening of the diagonal and interleaving distance*. 2021. ArXiv: 2006.13150.
- [PSW] F. Petit, P. Schapira, and L. Waas. *A property of the interleaving distance for sheaves*. arXiv: 2108.13018 [math.AG].
- [Sch00] M. Schwarz. *On the action spectrum for closed symplectically aspherical manifolds*. *Pacific J. Math* 193 (2000), pp. 419–461.
- [Sei00] P. Seidel. *Graded Lagrangian submanifolds*. *Bulletin de la S. M. F* 128.1 (2000), pp. 103–149. URL: http://www.numdam.org/item?id=BSMF_2000__128_1_103_0.
- [Sib85] N. Sibony. *Quelques problèmes de prolongement de courants en analyse complexe*. *Duke Math Journal* 52.1 (1985), pp. 157–197. DOI: [10.1215/S0012-7094-85-05210-X](https://doi.org/10.1215/S0012-7094-85-05210-X).
- [Tam08] D. Tamarkin. *Microlocal condition for non-displaceability*. Ed. by M. Hitrik et al. 2008. DOI: [10.1007/978-3-030-01588-6_3](https://doi.org/10.1007/978-3-030-01588-6_3). arXiv: 0809.1584 [math.SG].
- [Stacks] The Stacks project authors. *The Stacks project*. <https://stacks.math.columbia.edu>. 2022.
- [Ush19] M. Usher. *Local rigidity, symplectic homeomorphisms, and coisotropic submanifolds*. 2019. ArXiv: 1912.13043v2 (math.SG).
- [Vic13] N. Vichery. *Homological differential calculus*. 2013. arXiv: 1310.4845v1 [math.SG].
- [Vit92] C. Viterbo. *Symplectic topology as the geometry of generating functions*. *Mathematische Annalen* 292 (1992), pp. 685–710.

- [Vit95] C. Viterbo. *Functors and Computations in Floer cohomology, II*. (revised in 2003). 1995. arXiv: 1805.01316 [math.SG]. URL: <https://arxiv.org/abs/1805.01316>.
- [Vit06] C. Viterbo. “Symplectic topology and Hamilton-Jacobi equations”. In: *Morse Theoretic Methods in Nonlinear Analysis and in Symplectic Topology. NATO Science Series II: Mathematics, Physics and Chemistry book series (NAII, volume 217*. Ed. by P. Biran, O. Cornea, and F. Lalonde. Springer-Verlag, Netherlands, 2006, pp. 439–459.
- [Vit19] C. Viterbo. *Sheaf Quantization of Lagrangians and Floer cohomology* (2019). arXiv: 1901.09440 [math.SG].
- [Vit21] C. Viterbo. *Stochastic Homogenization for variational solutions of Hamilton-Jacobi Equations*. 2021. arXiv: <https://arxiv.org/abs/2105.04445> [math.SG].
- [Vit22] C. Viterbo. *On the support of elements in the Humilière completion of smooth Lagrangians and the inverse reduction inequality*. 2022. arXiv: 2204.04133 [math.SG].
- [Whi16] B. White. *Lectures on minimal surface theory*. 2016. DOI: 10.48550/arXiv.1308.3325. arXiv: 1308.3325 [math.SG]. URL: <https://arxiv.org/pdf/1308.3325.pdf>.
- [Zha20] J. Zhang. *Quantitative Tamarkin Theory*. CRM Short courses. Springer-Verlag, 2020. DOI: 10.1007/978-3-030-37888-2.
- [ZC05] A. Zomorodian and G. Carlsson. *Computing persistent homology*. *Discrete Comput. Geom* 33 (2005), pp. 249–274. DOI: 10.1007/s00454-004-1146-y.