

SAGEX-22-05
DESY 22-032
DO-TH 22/07
RISC Report number 22-03

arXiv:2203.13015 [hep-th]

The SAGEX Review on Scattering Amplitudes

Chapter 4: Multi-loop Feynman Integrals

Johannes Blümlein

Deutsches Elektronen-Synchrotron DESY, Platanenallee 6, 15738 Zeuthen, Germany
E-mail: Johannes.Bluemlein@desy.de

Carsten Schneider

Johannes Kepler University Linz, Research Institute for Symbolic Computation
(RISC), Altenberger Straße 69, A-4040 Linz, Austria
E-mail: Carsten.Schneider@risc.jku.at

March, 2022

Abstract. The analytic integration and simplification of multi-loop Feynman integrals to special functions and constants plays an important role to perform higher order perturbative calculations in the Standard Model of elementary particles. In this survey article the most recent and relevant computer algebra and special function algorithms are presented that are currently used or that may play an important role to perform such challenging precision calculations in the future. They are discussed in the context of analytic zero, single and double scale calculations in the Quantum Field Theories of the Standard Model and effective field theories, also with classical applications. These calculations play a central role in the analysis of precision measurements at present and future colliders to obtain ultimate information for fundamental physics.

1. Introduction

The present and upcoming high luminosity results at the Large Hadron Collider (LHC) at CERN, with input of the results measured at the ep -collider HERA at DESY, yield a big amount of precision data which require further fundamental precision calculations in perturbative Quantum Chromodynamics (QCD). This also applies to projects in the Future like the EIC [1], the LHeC [2,3], the ILC [4–7] or CLIC [8–10], the FCC_ee [11], and the proton version of the FCC [11], also for Quantum Electrodynamics (QED). Many of these outstanding problems can be formulated by large expressions in terms of hundred thousands (and even millions) of sophisticated Feynman integrals at higher

arXiv:2203.13015v3 [hep-th] 10 Jan 2023

loop order of presently up to three and four loops, with no, a single and two scales, or even multiscale problems.† Up to one scale, and in much fewer cases at two scales, technologies have been designed to calculate these integrals analytically over sets of basic functions, the properties of which have been studied to a certain extent. Furthermore, numerical representations of these building blocks have been derived.

The central objects are s -fold multiple integrals of the form

$$F(n, \varepsilon) = \int_0^1 \cdots \int_0^1 f(n, \varepsilon, x_1, \dots, x_s) dx_1 \cdots dx_s \quad (1)$$

where the discrete parameter n stands for the Mellin variable, and $\varepsilon = D - 4, \varepsilon \in \mathbb{R}, |\varepsilon| \ll 1$, is the dimensional parameter. A crucial property is that the integrand f is hyperexponential‡ in each of the integration variables x_i ($1 \leq i \leq s$) and hypergeometric in the discrete parameter n . In particular, one is interested in calculating the first coefficients of their Laurent series expansion w.r.t. ε :

$$F(n, \varepsilon) = F_l(n)\varepsilon^l + F_{l+1}(n)\varepsilon^{l+1} + \cdots + F_r(n)\varepsilon^r + O(\varepsilon^{r+1}). \quad (2)$$

For m -loop Feynman integrals without infrared divergences such expansions start usually at $l = -m$. In other cases, one obtains $l = -2m$. Alternatively, one looks for such an ε -expansions for the inverse Mellin transform $f(x, \varepsilon)$ with

$$\mathbf{M}[f(x, \varepsilon)](n) = F(n, \varepsilon) = \int_0^1 x^{n-1} f(x, \varepsilon) dx \quad (3)$$

or its power series representation

$$\bar{f}(x, \varepsilon) = \sum_{n=0}^{\infty} F(n, \varepsilon) x^n. \quad (4)$$

During the last decades more and more significant methods have been derived to simplify such Feynman integrals. Based on the representation (for instance (2)–(4)), we will present important tools that are currently used to perform such challenging calculations and discuss the associated special function spaces. We further relate these aspects to different precision calculations.

It is needless to say that we had to leave out the description of a series of techniques, which are also important. This concerns a series of aspects, which have been surveyed in Ref. [12], appearing in the same volume, and has been agreed between the different authors. It concerns e.g. the use of the symbol [13] and specific Hopf algebra structures [14–16], which are omnipresent in quantum field theoretic calculations. Related to this, many methods found in algebraic and arithmetic geometry are applicable [17–24]. For mostly numerical methods in use in multi-leg calculations we refer to [25].

†For zero scale problems results are available also at the five loop level.

‡ $h(x)$ is hyperexponential (or hypergeometric) in x if $\frac{h'(x)}{h(x)}$ (or $\frac{h(x+1)}{h(x)}$) is a rational function in x .

One way to classify the emergence of new mathematical structures in quantum field theories is given by the study of their differential equations. This is first of all a practical issue at the respective loop-level, where these structures are recognized and tied up with the respective graph topologies. There is in general no all order statement possible *ab initio*. However, as has been found out recently, Calabi–Yau motives play an important role here, cf. e.g. [24]. A selection criterion on what we are focusing on in the following are key technologies for multi-loop calculations in the massive case, presently to three-loop order in the zero-, single-, and two-scale cases. These technologies do synonymously apply to the corresponding massless calculations. There, clearly simplifications can be obtained using even other technologies, unlike the case in the massive case.

In the following we discuss the following key research topics:

- *guessing methods* (see Section 2),
- *solving linear recurrences and differential equations* (see Section 3),
- *solving coupled systems* (see Section 4),
- *transformation to special integral and sum representations* (see Section 5),
- *symbolic summation* (see Section 6),
- *symbolic integration* (see Section 7),
- *the large moment method* (see Section 8),
- *special function tools* (see Section 9)
- and *concrete calculations in the Quantum Field Theories of the Standard Model and within effective field theories* (see Section 10).

We emphasize that each of the different techniques cannot be considered as a stand-alone toolbox. Contrary, they all have to be applied in non-trivial interactions. In particular, based on a concrete problem, one has to choose the best tactic among the conglomeration of tools. For further and supplementary aspects we refer also to [26–28] and Chapter 3 [12] of the SAGEX review. We will conclude this survey on multi-loop tools and multi-loop calculations in Section 11.

2. Guessing methods

Often physical quantities can be evaluated up to a certain precision and one seeks for a mathematical representation that allows one to represent the data in a more compact fashion, to gain further insight and to support further calculations that depend on these quantities. Here we will emphasize two crucial tactics: (1) to predict from a given floating point number (that approximates a real number to very high precision) an alternative representation in terms of special constants and (2) to guess from a finite set of evaluations at integer points a linear recurrence or linear differential equation that satisfies all evaluations at integer points of the physical quantity.

2.1. Guessing integer relations

Using the LLL algorithm [29], or the PSLQ algorithm introduced in [30] and substantially improved in [31, 32], one can try to solve the following problem: Given a finite set of finite floating point numbers a_1, \dots, a_n with high precision (say l fractional digits), find integers $z_1, \dots, z_n \in \mathbb{Z}$ as small as possible in its absolute value§ such that

$$z_1 a_1 + \dots + z_n a_n < 10^{-m},$$

where m is large.

In order to obtain further confidence one may apply this method with further precision (i.e., more digits l of the input) and checks if the obtained result remains the same but m gets larger. For instance, suppose that we are given the finite floating point number

$$a_1 = 5.68700407989058207630312605688168433418849655155997$$

with $l = 50$ fractional digits that approximates a real number $r_1 \in \mathbb{R}$. Then one can use, e.g., the Mathematica implementation of the PSLQ algorithm to search for an alternative representation in terms of $\zeta(2)$, $\zeta(3)$ and $\zeta(5)$. Namely, by activating `FindIntegerNullVector` $\{a_1, \text{N}[\text{Zeta}[2], 50], \text{N}[\text{Zeta}[3], 50], \text{N}[\text{Zeta}[5], 50]\}$ one obtains the result $(z_1, z_2, z_3, z_4) = (-2, 2, 5, 2)$ with $m = 49$. Thus we may conjecture that

$$r_1 = \zeta(2) + \frac{5}{2}\zeta(3) + \zeta(5)$$

holds. We remark that the PSLQ method finds this relations already with $l = 5$ fractional digits with precision $m = 4$. Of course, if one is given even more digits of r_1 , one may check if even more digits m agree. Similarly, one may apply PSLQ again for this improved data in order to check if there is even a smaller relation (with smaller z_i).

Summarizing, the method can be very efficiently used if one knows the set of numbers, by which the result of a calculation is finally spanned or if one wants to find a linear combination of such numbers. A highly non-trivial example is, e.g., the calculation of the 5-loop β -function in QCD in Ref. [33], where these guessing tools were instrumental. For a recent survey on these techniques (covering not only PSLQ but also the LLL approach) and further applications we refer to [34].

2.2. Guessing recurrences and differential equations

In Section 8 below we will introduce a method that enables one to compute many moments $F(n, \varepsilon)$ in (3) or coefficients in (4) for $n = 0, 1, 2, \dots$. More precisely, if we

§Since any finite floating point number can be written as a rational number, this problem can be always solved if the integers z_i can be arbitrarily large. Thus a solution to the problem might indicate a proper relation among the approximated real numbers if l is large but the values z_i are small.

write $F(n, \varepsilon)$ in its ε -expansion (2), we will be able to compute the moments of the first ε -coefficients, say $F_j(n)$ for $l \leq j \leq r$ with $n = 0, 1, 2, \dots, \mu$ where μ is large (e.g., $\mu = 10^4$).

Within multi-loop calculations these moments depend linearly also on special constants, such as the multiple zeta values [35], with rational coefficients. This finally leads to several finite sequences, $F(0), F(1), \dots, F(\mu)$, of rational numbers. Then given these numbers, one can try to guess a linear recurrence

$$a_0(n)F(n) + a_1(n)F(n+1) + \dots + a_\lambda(n)F(n+\lambda) = 0 \quad (5)$$

of order λ with polynomial coefficients $a_i(n) \in \mathbb{Q}[n]$ that contains this finite sequence as solution. Namely, fixing the order λ and assuming that the degrees of the polynomials $a_i(n)$ are less than or equal to δ , one searches for the $r = (\delta + 1)(\lambda + 1)$ unknown coefficients. More precisely, by setting $n = 0, \dots, r - 2$ in (5) and plugging in the rational numbers $F(0), \dots, F(r + \lambda - 2)$ one gets $r - 1$ equations in r unknowns over the rational numbers which can be solved by linear algebra. In many cases this yields solutions that do not hold for $n \geq r - 1$. Thus one usually takes an over-determined system (by more evaluations, say $0 \leq n \leq r + 100$). In this way one can exclude basically all wrong solutions. Finally, given a found solution one usually checks at many extra points if the recurrence is still valid. This gives further evidence that the guessed recurrence is reliable.

This tactic implemented, e.g., in the Maple package `gfun` [36] or the Mathematica package `GeneratingFunction` [37] is surprisingly simple and can be carried out in this naive fashion for small examples (i.e., for recurrences of small orders λ and small degree bounds δ). For large examples within QCD calculations this straightforward procedure utterly fails. Here highly efficient computer algebra technologies, such as homomorphic image calculations and rational/polynomial reconstructions, are essential [38]. Using in addition gcd-calculations to determine recurrences with minimal order, the Mathematica implementation `Guess.m` by Kauers could be utilized with about $\mu = 5000$ moments to guess all the recurrences that determine the massless unpolarized 3-loop anomalous dimensions and Wilson coefficients in deep-inelastic scattering [39–41] in Ref. [42], see also [43–47]. For even larger problems, the highly efficient Sage implementation in `ore_algebra` [48] (utilizing among other smart techniques the fast integer arithmetic of `Flint`) was instrumental to guess linear recurrences with minimal order. E.g., for the massive form factor [49, 50] about $\mu = 10000$ moments were needed to obtain recurrences up to order $\lambda = 55$ and degree $\delta = 1300$. In the case of a massive operator matrix element 8000 moments [51] could be calculated and difference equations were derived for all contributing color and ζ -value structures. Recently, also the 3-loop splitting functions [44], the anomalous dimensions from off shell operator matrix elements [44, 52, 45, 46], and lately the polarized transition matrix element $A_{gq}(N)$ [53] and the logarithmic contributions to the polarized $O(\alpha_s^3)$ asymptotic massive Wilson coefficients [54] have been derived by guessing the underlying recurrence relations.

Further we note that one can guess in a similar fashion a linear differential equation

of the power series $f(x) = \sum_{n=0}^{\infty} F(n)x^n$, say

$$a_0(x)f(x) + a_1(x)D_x f(x) + \cdots + a_\lambda(x)D_x^\lambda f(x) = 0$$

where $D_x = \frac{d}{dx}$ denotes the differentiation w.r.t. x . Both, the Mathematica implementation in `Guess.m` and the Sage implementation in [48] cover this extra feature.

Given such recurrences, one succeeds in many cases to solve the recurrences in terms of special functions that are most relevant in QCD calculations. Further details on these solving aspects will be given in the next section.

3. Solving linear recurrences and differential equations

As already motivated in Section 2.2 above and further emphasized in Sections 6–8, one can derive a linear recurrence (linear difference equation) or a linear differential equation which contains the given multi-loop Feynman integral (1) or a given physical expression in terms of such Feynman integrals as a solution. Then a natural strategy is to apply the available toolboxes to compute all solutions of the derived equations that can be represented in terms of certain classes of function spaces that will be introduced in more detail in Section 9. In the case that one finds sufficiently many (linearly independent) solutions one may obtain an alternative representation of the physical problem in terms of these solutions.

In the following we describe different algorithms that can provide solutions of linear difference and differential equations that occur in QCD calculations.

3.1. Ordinary linear equations

We start with equations in one variable, i.e., with ordinary linear difference equations of the form

$$\sum_{i=0}^{\lambda} a_i(n) F(n+i) = r(n) \tag{6}$$

and ordinary linear differential equations of the form

$$\sum_{i=0}^{\lambda} a_i(x) D_x^i f(x) = r(x) \tag{7}$$

where $D_x = \frac{d}{dx}$ denotes the differentiation w.r.t. x .

3.1.1. Ordinary linear difference equations The first major contribution for recurrence solving is elaborated in [55] and is substantially improved in [56]. Given rational functions $a_0(n), \dots, a_\lambda(n), r(n) \in \mathbb{K}(n)$ (\mathbb{K} denotes a computable field that contains the rational numbers) it finds all rational solutions $F(n) \in \mathbb{K}(x)$ of (6). More generally, using the algorithms from [57] and the more efficient version given in [58] one can compute

all hypergeometric solutions of (6), this means one can compute all solutions that can be written as hypergeometric products

$$F(n) = \prod_{k=l}^n f(k),$$

where l is an integer and $f(k)$ is a rational function in k ; here l is chosen such that the evaluation $f(k)$ for $k \in \mathbb{N}$ with $k \geq l$ has no pole and is nonzero. In particular, the solutions can be described in terms of a product of Γ -functions, Pochhammer-symbols, factorials, binomial coefficients and rational functions. Even more generally, using the algorithms described in [59, 60] one can search for all d'Alembertian solutions, i.e., all solutions that can be expressed in terms of iterative sums defined over hypergeometric products. Special cases of this class of sums are harmonic sums [61, 62], cyclotomic harmonic sums [63], generalized harmonic sums [64, 65] and finite binomial sums [66]; infinite binomial sums have been also studied in [67, 68, 66]. Further details and extra properties of such sums are presented in Section 9.

Finally, one can search in addition for all Liouvillian solutions [69] which cover in addition the interlacing of expressions in terms of iterated sums over hypergeometric products. Basically all these tools have been generalized to the setting of difference fields [70] and rings [71] (utilizing results from above and [72–76]) that allows one to find such solutions for difference equations (6) where the coefficients $a_i(n)$ and the inhomogeneous part $r(n)$ are not just rational functions but can be built again by indefinite nested sums over hypergeometric products. E.g., using the summation package **Sigma** [77–79], that contains this general toolbox, one can compute for the recurrence

$$\begin{aligned} & (1 + S_1(n) + nS_1(n))^2 (3 + 2n + 2S_1(n) + 3nS_1(n) + n^2S_1(n))^2 F(n) \\ & - (1 + n)(3 + 2n)S_1(n)(3 + 2n + 2S_1(n) + 3nS_1(n) + n^2S_1(n))^2 F(n + 1) \\ & + (1 + n)^2(2 + n)^3 S_1(n)(1 + S_1(n) + nS_1(n)) F(n + 2) = 0 \end{aligned}$$

the complete solution set

$$\left\{ c_1 S_1 n \prod_{l=1}^n S_1(l) + c_2 S_1(n)^2 \prod_{l=1}^n S_1(l) \mid c_1, c_2 \in \mathbb{K} \right\};$$

here $S_1(n) = \sum_{k=1}^n \frac{1}{k}$ denotes the n th harmonic number. Internally, the recurrence operator is factorized as much as possible into linear factors. Then each extra factor provides one extra linearly independent solution which is constructed by one extra indefinite sum. In other words, finding ν linear factors (ideally $\nu = \lambda$) yields ν linearly independent solutions where the most complicated solution is built by an iterative nested sum over hypergeometric products of nesting depth $\nu - 1$; the particular solution will lead to a nested sum of depth ν . Then a key task is to simplify these sum solutions further such that the nesting depth is minimal; further aspects on such simplifications will be given in Section 6.1. We note that all solutions of a linear recurrence can be given

in terms of d'Alembertian solutions if the linear recurrence operator factors completely into first-order linear factors. We call such a recurrence also first order factorizing. If this is not the case, i.e., if only parts of the recurrence can be factored into linear right-hand factors then it is called non-first order factorizing.

The recurrences coming from QCD calculations usually have polynomial coefficients $a_i(n)$ and the right-hand side $r(n)$ is either 0 or is built by indefinite nested sums over hypergeometric products. One of the largest homogeneous recurrences ($r(n) = 0$) that have been solved with **Sigma** were of order $\lambda = 35$ and the degree of the coefficients of $a_i(n)$ was up to 1000 and the occurring integers required up to 1400 decimal digits; for details see, e.g., [80, 42]. The largest inhomogeneous recurrences were up to order $\lambda = 12$ where $r(n)$ may be built up to hundreds of highly nested indefinite nested sums.

In most cases Feynman diagrams or physical expressions in terms of such integrals depend on the dimensional parameter ε . In particular, this parameter ε occurs in the coefficients $a_i(n)$ and the inhomogeneous part $r(n)$ of the recurrence (6). In some special cases, the solution $F(n)$ can be given in terms of indefinite nested sums over hypergeometric products where ε occurs inside of the sums and products. In such situations, the above methods implemented in **Sigma** can find the complete solution in n and ε . However, in most instances such a closed form solution does not exist and one seeks for closed form solutions of the first coefficients $F_i(n)$ (free of ε) of the ε -expansion (2). In order to accomplish this task, one can apply the algorithm from [81] implemented in **Sigma** in order to constructively decide if the coefficients $F_i(n)$ can be represented in terms of nested sums over hypergeometric products.

The more complicated multi-loop Feynman integrals are considered, the more complicated function spaces arise. Thus further techniques are extremely desirable that extend the class of indefinite nested sums over hypergeometric products. In this regard, one should mention the special case of factorial series [82–84] solutions of the form $f(n) = \sum_{k=0}^{\infty} \frac{k!}{(n+k)!} a_k$. Namely, given a linear recurrence in $f(n)$, an operator method is described in [85] and further considered in [86], to provide a linear recurrence for the sequence a_k . Precisely here one can utilize the recurrence solver of **Sigma** to decide, if a_k can be written in terms of d'Alembertian solutions. E.g., for the recurrence

$$(1+n)(2+n)(3+n)f(n) - (2+n)^2(3+n)f(1+n) + (2+n)(3+n)f(2+n) - f(3+n) = 0$$

one finds the factorial series solution $\sum_{k=0}^{\infty} \frac{n!}{(k+n)!} \sum_{i=0}^k \frac{(-1)^i}{i!}$. Furthermore, M. Petkovšek proposed new ideas in [87] to find solutions of truncated binomial sums: instead of $\frac{k!}{(n+k)!}$ one can choose certain products of binomial coefficients and the upper bound should be integer-linear in n .

3.1.2. Ordinary linear differential equations In various instances one is interested in a power series solution $f(x) = \sum_{n=0}^{\infty} F(n)x^n$ of a linear differential equation (7). If the coefficients $a_i(x)$ are rational functions in x and the inhomogeneous part $r(x)$ itself can be given in form of a power series representation, one can utilize holonomic closure

properties [36, 37, 88] as follows. Plugging the power series ansatz into the differential equation and comparing coefficients w.r.t. x^n yield a linear recurrence of the form (7) (with updated $a_i(n)$ and $r(n)$) for the desired coefficients $F(n)$. In a nutshell, one can activate the available recurrence solver introduced in Section 3.1.1 to compute closed form representations of the coefficients $F(n)$.

Alternatively, there are also direct algorithms available, similarly to the difference equation case, that can solve linear differential equations in terms of rather general classes of special functions. Namely, using the algorithms from [55] one can find all rational solutions. More generally, using [89] and, e.g., the improved versions given in [56, 90] one can find all hyperexponential functions $f(x)$. In general, the functions can be given in the form $e^{\int^x h(x)dx}$ for some rational function h and lower bound l ; special cases are, e.g., rational functions or roots over such functions. More generally, one can use these algorithms to compute all d'Alembertian [59, 60], i.e., all solutions that can be given in terms of iterated integrals over hyperexponential functions. Special cases of these integrals, are harmonic polylogarithms [91], cyclotomic polylogarithms [63], generalized multiple polylogarithms [64, 65] but also root-valued nested integrals [66]; further details are given in Section 9. As for the recurrence case the corresponding differential operator is factorized as much as possible into linear factors. Then each factor yields one extra linearly independent solution by introducing one extra indefinite integration quantifier. These d'Alembertian solutions can be computed with the package `HarmonicSums` [92]. Similarly to the recurrence case we note that all solutions of a linear differential equation can be given in terms of d'Alembertian solutions if the differential operator factors completely into first-order linear factors. We call such a differential equation also first order factorizing. If this is not the case, i.e., if only parts of the linear differential equation can be factored into first-order linear right-factors then it is called non-first order factorizing. More generally, also Liouvillian solutions [93] can be calculated partially with `HarmonicSums` by utilizing Kovacic's algorithm [94]. For instance, given

$$(11 + 20x)f'(x) + (1 + x)(35 + 134x)f''(x) + 3(1 + x)^2(4 + 37x)f^{(3)}(x) + 18x(1 + x)^3f^{(4)}(x) = 0$$

`HarmonicSums` finds the general solution

$$\left\{ c_1 + c_2 \int_0^x \frac{1}{1+\tau_1} d\tau_1 + c_3 \int_0^x \int_0^{\tau_1} \frac{\sqrt[3]{1+\sqrt{1+\tau_2}}}{(1+\tau_1)(1+\tau_2)} d\tau_2 d\tau_1 + c_4 \int_0^x \int_0^{\tau_1} \frac{\sqrt[3]{1-\sqrt{1+\tau_2}}}{(1+\tau_1)(1+\tau_2)} d\tau_2 d\tau_1 \mid c_1, c_2, c_3, c_4 \in \mathbb{K} \right\},$$

where $\sqrt{1+x}$ is hyperexponential and $\sqrt[3]{1-\sqrt{1+x}}$ is algebraic over a field generated by x and $\sqrt{1+x}$. More generally, in [93] an algorithm has been described that finds all Liouvillian solutions of a homogeneous linear differential equations, i.e., all solutions that can be given by iterated integrals over hyperexponential function and functions

that are algebraic over the extension below. More generally, an algorithm has been proposed in [95] that can find all Liouvillian solutions of linear differential equations whose coefficients are given in terms of functions that are Liouvillian. In some sense, this highly general solver can be considered as the continuous version of the recurrence solver [70] implemented within the package `Sigma`.

As already emphasized in Section 3.1.1, also the dimensional parameter ε appears in the coefficients $a_i(x)$ and the inhomogeneous part $r(x)$ of the linear differential equation (7) when one deals with Feynman integrals. In some special cases one can use the above algorithms directly where ε may arise inside of d'Alembertian and Liouvillian solutions. However, similarly to the recurrence case, this approach usually does not work and one aims at finding closed forms of the first coefficients of the ε -expansion

$$f(x, \varepsilon) = f_l(x)\varepsilon^l + f_{l+1}(x)\varepsilon^{l+1} + \dots + f_r(x)\varepsilon^r + O(\varepsilon^{r+1}).$$

In this regard, the package `HarmonicSums` can decide constructively if the first coefficients $f_i(x)$ (free of ε) can be given in terms iterated integrals over hyperexponential functions; for the underlying algorithm we refer to [96] which is based on ideas given in [81].

By looking at more and more complicated Feynman integrals also the class of Liouvillian solutions is not sufficient. For second order linear differential equations van Hoeij proposed algorithms in [97] that can find hypergeometric series solutions (${}_pF_q$'s) in terms of certain rational function arguments. These advanced tools turned out to be instrumental to deal with the ρ -parameter in [98, 99] and related quantities.

3.2. Partial linear equations

Solving partial linear difference and differential equations is a hard problem. It has been shown in [100] based on [101] that already the task to solve such equations in terms of polynomial solutions is an unsolvable problem. Recently, new methods have been introduced in [102, 103] that enable one to search for (not necessarily all) rational solutions of partial linear difference equations of the form

$$\sum_{(s_1, \dots, s_r) \in S} a_{(s_1, \dots, s_r)}(n_1, \dots, n_r) F(n_1 + s_1, \dots, n_r + s_r) = 0, \quad (8)$$

where the coefficients $a_{(s_1, \dots, s_r)}$ are rational functions in the variables n_1, \dots, n_r and $S \subset \mathbb{Z}^r$ is a finite set. In [104] further ideas coming from Section 3.1.1 have been incorporated to hunt also for solutions in terms of a given set of nested sums. E.g., suppose that we are given the partial linear difference equation

$$\begin{aligned} & - (n+1)^2 (k+n^2+2) (4k^2 - 3kn^2 + 5kn + 12k - 2n^3 - 2n^2 + 8n + 8) F(n, k+1) \\ & + (n+1)^2 (k+n^2+3) (2k^2 - 2kn^2 + 2kn + 6k - n^3 - n^2 + 4n + 4) F(n, k+2) \\ & + (n+1)^2 (k+n+1) (2k - n^2 + n + 4) (k+n^2+1) F(n, k) \\ & - (k+1)n^2(n+2)^2 (k+n^2+2n+2) F(n+1, k) \end{aligned}$$

$$+ kn^2(n+2)^2(k+n^2+2n+3)F(n+1, k+1) = 0$$

and the set $W = \{S_1(k), S_1(n+k), S_{2,1}(n+k)\}$ in terms of the harmonic numbers and the harmonic sum $S_{2,1}(n) = \sum_{k=1}^n \frac{S_1(k)}{k^2}$; compare Section 9. Then fixing the total degree bound 5 or the arising objects in the numerator, one can compute with the package `SolvePLDE` introduced in [104] the 37 solutions $\frac{p}{(1+n)^2(1+k+n^2)}$ where p is taken from the set

$$\begin{aligned} & \left\{ 1 + \frac{1}{2}nS_1(k+n), k, n, kn, kn^2, kn^3, kn^4, kS_1(n), knS_1(n), kn^2S_1(n), kn^3S_1(n), kS_1(n)^2, \right. \\ & knS_1(n)^2, kn^2S_1(n)^2, kS_1(n)^3, knS_1(n)^3, kS_1(n)^4, kS_{2,1}(n), knS_{2,1}(n), kn^2S_{2,1}(n), kn^3S_{2,1}(n), \\ & kS_1(n)S_{2,1}(n), knS_1(n)S_{2,1}(n), kn^2S_1(n)S_{2,1}(n), kS_1(n)^2S_{2,1}(n), knS_1(n)^2S_{2,1}(n), \\ & kS_1(n)^3S_{2,1}(n), kS_{2,1}(n)^2, knS_{2,1}(n)^2, kn^2S_{2,1}(n)^2, kS_1(n)S_{2,1}(n)^2, knS_1(n)S_{2,1}(n)^2, \\ & \left. kS_1(n)^2S_{2,1}(n)^2, kS_{2,1}(n)^3, knS_{2,1}(n)^3, kS_1(n)S_{2,1}(n)^3, kS_{2,1}(n)^4 \right\}. \end{aligned}$$

In particular, the method for scalar linear difference equations in [80] has been carried over in this new package to search also for closed form solutions of the first coefficients of an ε -expansion.

We emphasize that this new package enables one also to attack partial linear differential equations and to find solutions in its multivariate power series expansion $f(x_1, \dots, x_r) = \sum_{(n_1, \dots, n_r) \in \mathbb{N}^r} F(n_1, \dots, n_r) x_1^{n_1} \dots x_r^{n_r}$. Namely by plugging the power series ansatz into the partial differential equation and comparing coefficients w.r.t. $x_1^{n_1} \dots x_r^{n_r}$ produce a partial linear difference equation of the form (8). Thus one can utilize the tools described above to search for closed form representations of $F(n_1, \dots, n_r)$. In Section 4.2 this tactic will be refined further to find solutions for certain classes of coupled systems of partial linear differential equations.

4. Solving coupled systems of linear differential equations

In order to solve open problems at the forefront of elementary particle physics, millions of complicated Feynman integrals have to be tackled. As a preprocessing step one often applies integration-by-parts (IBP) methods [105–107]¶ that crunch these integrals to a few hundred (or thousand) so-called master integrals; for a recent survey, possible refinements and applications see, e.g., [107, 110, 111]. Then the main task is to simplify only these master integrals to expressions in terms of special functions and to assemble the original problem with these sub-results. Most of these master integrals $f_i(x, \varepsilon)$ can be determined as solutions of coupled systems of linear differential equations. For single-variate systems they are of the form

$$D_x \begin{pmatrix} f_1(x, \varepsilon) \\ \vdots \\ f_\lambda(x, \varepsilon) \end{pmatrix} = A \begin{pmatrix} f_1(x, \varepsilon) \\ \vdots \\ f_\lambda(x, \varepsilon) \end{pmatrix} + \begin{pmatrix} g_1(x, \varepsilon) \\ \vdots \\ g_\lambda(x, \varepsilon) \end{pmatrix}, \quad (9)$$

with A being a $\lambda \times \lambda$ matrix with entries from $\mathbb{K}(x, \varepsilon)$ where the right-hand sides are given in terms of simpler master integrals. They are either determined by other coupled

¶Here the method of syzygies [108, 109] from computational algebraic geometry helps to reduce the number of contributing scalar products.




Figure 1. Solving systems.

systems or have to be tackled by tools presented, e.g., in Sections 6 and 7. Here we elaborate the most relevant approaches. Before one considers to solve such systems, one may also analyze them further as exemplified in [112] in order gain further insight or to find further relations among them.

4.1. Uncoupling algorithm and scalar solvers

In the last years a general toolbox has been elaborated that finds all solutions that can be given in terms of iterated integrals (or sums) as follows. By uncoupling algorithms [113, 114] available, e.g., in the package `OreSys` [115], one first decouples the system (9) to a scalar linear differential equation in one of the unknowns. Using the differential equation solver in `HarmonicSums` [92] (based on [93, 94, 59]) one finds, whenever possible, a closed form representations of the unknown functions f_1, \dots, f_λ in terms of d'Alembertian (and partially of Liouvillian) solutions. Based on this strategy we recalculated the 2-loop form factors [116] and obtained first results for the 3-loop case [117]. Another fruitful approach [118] is based on recurrence solving. Here one assumes that the arising Feynman integrals can be given in the power series representations

$$f_i(x) = \sum_{n=0}^{\infty} F_i(n)x^n, \quad i = 1, \dots, \lambda. \quad (10)$$

Then the machinery proceeds as summarized in Fig. 1. After the decoupling of the system (step 1) one takes the scalar differential equation of $f_1(x)$ and calculates by means of holonomic closure properties [88] a scalar linear recurrence of $F_1(n)$ (step 2). Activating the recurrence solver of `Sigma` (based on [55, 57–59, 75, 81]) one can decide algorithmically in step 3 if $F_1(n)$ can be represented in terms of indefinite nested sums. If yes, one plugs this representation into the decoupled system and gets closed forms of the remaining $F_2(n), \dots, F_\lambda(n)$ in step (4). For advanced QCD-calculations see, e.g., [43, 119, 120, 118, 121, 44, 122, 123]

4.2. Direct solver

So far there are only few algorithms available that can compute directly (i.e., without uncoupling) the desired set of solutions of a given coupled system of the form (9). For instance, with the algorithms in [124] one can find all hyperexponential solutions of (higher order) coupled systems but so far no algorithms are available to compute all d'Alembertian or Liouvillian solutions. First steps have been elaborated in [125] within the general difference field setting.

However, in various instances it has been observed in [126] that the arising coupled systems of the form (9) can be transformed to a system of the form

$$D_x \tilde{\mathbf{f}}(x, \varepsilon) = \varepsilon \tilde{A}(x) \tilde{\mathbf{f}}(x, \varepsilon) + \tilde{\mathbf{g}}(x, \varepsilon) \quad (11)$$

for a matrix $\tilde{A}(x)$ which is free of ε and where $\tilde{\mathbf{f}}$ and $\tilde{\mathbf{g}}$ are defined by the multiplication of an invertible matrix T with $\mathbf{f} = (f_1, \dots, f_\lambda)$ and $\mathbf{g} = (g_1, \dots, g_\lambda)$, respectively. Under the assumption that such a transformed system exists, algorithms are available to compute such a transformation matrix T . Furthermore, methods exist, cf. [127–130] that can hunt for such a basis transformation for the multivariate case, i.e., for systems of partial linear differential equations. Given such a transformed system one obtains the benefit that one can read off the coefficients of the ε -expansion in terms of indefinite nested integrals. However, such a transformation does not hold for more complicated systems.

As elaborated in [104], there are other special cases that enable one to solve coupled systems of partial linear differential equations if one considers the solution as a multivariate power series solution where the coefficients satisfy a nicely coupled system of linear difference equations. For instance, take the coupled partial system

$$\begin{aligned} (x-1)yD_{xy}f(x,y) + (x(2\varepsilon + \frac{7}{2}) - \varepsilon + 1)D_x f(x,y) + (x-1)xD_x^2 f(x,y) \\ + y(2\varepsilon + 1)D_y f(x,y) + \frac{3}{2}(2\varepsilon + 1)f(x,y) = 0, \\ x(y-1)D_{xy}f(x,y) + x(4-\varepsilon)D_x f(x,y) + (y(\frac{13}{2} - \varepsilon) - \varepsilon + 1)D_y f(x,y) \\ + (y-1)yD_y^2 f(x,y) + \frac{3(4-\varepsilon)}{2}f(x,y) = 0. \end{aligned}$$

Then writing $f(x, y)$ as a multivariate power series $f(x, y) = \sum_{n,m=0}^{\infty} F(n, m)x^n y^m$ one obtains by coefficient comparison w.r.t. $x^n x^m$ the coupled system of homogeneous first-order difference equations

$$\begin{aligned} \frac{3}{2}(2\varepsilon + 1)F(n, m) - n(\varepsilon - 1)F(n + 1, m) &= 0, \\ -\frac{3}{2}(\varepsilon - 4)F(n, m) - m(\varepsilon - 1)F(n, m + 1) &= 0. \end{aligned}$$

Given such a first-order homogeneous system, it follows that its solution can be expressed in terms of hypergeometric products. Namely, using the algorithm given in [104, Sec. 4.1] (which is a simplified version of the algorithm given in [131]) and implemented in the package `HypSeries` one obtains the solution

$$F(n, m) = \left(\prod_{i=1}^n \frac{(1+2i)(3+i-\varepsilon)}{2i(-2+i+\varepsilon)} \right) \prod_{i=1}^m \frac{(1+2i+2n)(i+2\varepsilon)}{2i(-2+i+n+\varepsilon)}$$

$$= \frac{\left(\frac{3}{2}\right)_{m+n}(4-\varepsilon)_n(1+2\varepsilon)_m}{m!n!(-1+\varepsilon)_{m+n}}$$

in terms of hypergeometric products or equivalently in terms of factorial and Pochhammer symbols. As a consequence the derived solution of the original coupled system of differential equations can be given in the form

$$F(x, y) = \sum_{n,m=0}^{\infty} \frac{\left(\frac{3}{2}\right)_{m+n}(4-\varepsilon)_n(1+2\varepsilon)_m}{m!n!(-1+\varepsilon)_{m+n}} x^n y^m.$$

Using this sum representation one can deploy the summation tools in Section 6 to calculate the first coefficients of its ε -expansion. We note that these solutions (coming from homogeneous first-order difference systems) are closely related to special functions that are introduced in the next section.

5. Transformation to special integral and sum representations

In the simplest cases, the integrands of Feynman integrals (1) exhibit Euler Beta-function structures and by clever rewriting, cf. also e.g. [132], the integral can be rewritten in terms of hypergeometric functions and their generalization [133–136]. More precisely, one may rewrite simple Feynman integrals in terms of the following hierarchy of ${}_{p+1}F_p$ functions, the first of which read

$$\begin{aligned} B(a_1, a_2) &= \int_0^1 dt t^{a_1-1}(1-t)^{a_2-1} \\ {}_2F_1(a_1, a_2, b_1; x) &= \frac{\Gamma(b_1)}{\Gamma(a_2)\Gamma(b_1-a_2)} \int_0^1 dt t^{a_2-1}(1-t)^{b_1-a_2-1}(1-tx)^{-a_1} \\ {}_3F_2(a_1, a_2, b_1; x) &= \frac{\Gamma(b_2)}{\Gamma(a_3)\Gamma(b_2-a_3)} \int_0^1 dt t^{a_3-1}(1-t)^{-a_3+b_2-1} {}_2F_1(a_1, a_2, b_1; tx). \end{aligned}$$

Here the parameters a_i, b_i are such, that the corresponding integrals exists, [136].|| We note that computer algebra can be used non-trivially to explore further properties on these special functions. E.g., using symbolic summation (see also Section 6) one can compute all arising contiguous relations of a finite set of sums [139]. More generally, it is possible to represent Feynman integrals by Mellin–Barnes [140–142] representations [81]. As well-known, the Mellin–Barnes representations are also used for hypergeometric functions and their generalizations, originally in terms of Pochhammer Umlauf-integrals.

At 3-loop order, also Appell functions [143–150, 104] and their generalizations arise; see, e.g., [151, 152, 118]. For instance, the F_1 function has the integral representation

$$F_1(a, b_1, b_2, c; x, y) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \int_0^1 dt t^{a-1}(1-t)^{c-a-1}(1-xt)^{-b_1}(1-yt)^{-b_2},$$

$$\operatorname{Re}(c) > \operatorname{Re}(a) > 0.$$

||One may then perform corresponding analytic continuations, cf. [137, 138].

When one succeeds in detecting that the the given Feynman integrals can be rewritten in integral representations that can be connected to ${}_{p+1}F_p$ or Appell-like functions, one can utilize the essential property that all the ${}_{p+1}F_p$ functions have a single infinite sum representation, while the Appell-functions are represented by two infinite sums. For instance, we get

$$F_1(a; b_1, b_2; c; x, y) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(a)_{m+n} (b_1)_m (b_2)_n}{m! n! (c)_{m+n}} x^m y^n.$$

Similarly, there are also other classes of higher transcendental functions, which obey multi-sum representations [146, 147, 149, 104]. Up to the level of massless and single mass two-loop integrals, cf. [153] and in some cases in the three loop case, cf. [118], these representations are usually sufficient. For more complicated integrand-structures, however, one has to apply other techniques. Applying successively Newton's binomial theorem and Mellin–Barnes [140–142] decompositions on the integrand, implemented in different packages [154–157], enables one to carry out all integrals by introducing Mellin–Barnes integrals. Finally, carrying out the remaining Mellin–Barnes integrals with the residue theorem yields definite multiple sums

$$\sum_{k_1=1}^{L_1(n)} \cdots \sum_{k_v=1}^{L_v(n, k_1, \dots, k_{v-1})} f(n, k_1, \dots, k_v). \quad (12)$$

Here the upper bounds $L_1(n), \dots, L_v(n, k_1, \dots, k_{v-1})$ are integer linear (i.e., linear combinations of the variables over the integers) in the dependent parameters or ∞ , and f is hypergeometric in n and the summation variables k_i . For further details on this rewriting process we refer, e.g., to [81, 28].

In physical applications the dimensional parameter ε arises in the parameters a_i, b_i, c_i, \dots of the ${}_{p+1}F_p$ and Appell-type representations. Moreover, the hypergeometric summand f in (12) may also depend on ε . In all these cases one seeks for an ε -expansion where the coefficients are represented in sum representations that are as simple as possible. In order to accomplish this task, highly general summation methods introduced in Section 6 can be applied.

6. Symbolic summation

Following the strategy sketched in Section 5 one ends up at (thousands or even millions) of definite multiple sums where the summand is built by hypergeometric products and indefinite nested sums, like harmonic sums [61, 62], cyclotomic harmonic sums [63], generalized harmonic sums [64, 65]; these sums may pop up in particular if one expands the summand w.r.t. the ε -parameter (i.e., if one applies the differential operator w.r.t. ε to the hypergeometric products; for a detailed description see, e.g., [104]). Producing such sum representations without making the original problem more complicated is highly non-trivial. However, if one succeeds in getting an

appropriate sum representation, one can apply various symbolic summation algorithms to simplify these sums.

6.1. Simplification of indefinite nested sums

The simplification of indefinite nested sums defined over hypergeometric products started with Gosper's and Karr's summation algorithms [158, 72] and has been enhanced significantly within the last 20 years to a strong summation machinery based on difference field and ring theories [73, 159–163]. Using our summation package `Sigma` [77, 78] it is now possible to design completely automatically appropriate difference rings in which one can represent such indefinite nested sums fulfilling various optimality criteria: e.g., the number of nested summation quantifiers or the degrees in the denominators are minimized; see [164–167]. Furthermore, employing our contributions to a refined Galois theory of difference rings [168] (see also [69, 169–171]), the used sums do not admit any algebraic relations. As a consequence, one obtains canonical (unique) product-sum representations [79].

Furthermore, these algorithms can be accompanied with quasi-shuffle relations [172–176] for the discovery of such relations in a very efficient way; for further details we refer to Section 9.

6.2. The WZ-summation approach

The treatment of single nested definite hypergeometric sums started with Zeilberger's creative telescoping paradigm [177–181] and has been enhanced to multi-summation with the WZ-summation approach due to [182] and its refinements given, e.g., in [183–185]. Given a multiple sum $F(n, \varepsilon)$ over a hypergeometric summand, like on the left-hand side of**

$$\sum_{j=0}^{n-2} \sum_{r=0}^{j+1} \sum_{s=0}^{n-j+r-2} \frac{(-1)^r (n-j-2)! \binom{j+1}{r} r! (-1)^s \binom{n-j+r-2}{s}}{(n-j+r)! (n-s)(s+1)} = \frac{((-1)^n - 1)(n^2 + n + 1)}{n^2(n+1)^3} + \frac{S_1(n)}{(n+1)^2} - \frac{S_2(n) + 2S_{-2}(n)}{n+1}, \quad (13)$$

one can search for a recurrence/difference equation of order λ of the form

$$\sum_{i=0}^{\lambda} a_i(n, \varepsilon) F(n+i, \varepsilon) = r(n, \varepsilon)$$

with polynomials $a_i(n, \varepsilon)$ in n, ε and $r(n, \varepsilon)$ being an expression in terms of multiple sums of simpler type than $F(n, \varepsilon)$. By further tricks one can compute even a homogeneous recurrence. With the package `MultiSum` [183] one obtains, for instance, for the triple

**For $a \in \mathbb{Z} \setminus \{0\}$ we define the generalized harmonic numbers $S_a(n) = \sum_{k=1}^n \frac{(\text{sign}(a))^k}{k^{|a|}}$.

sum in (13) a homogeneous linear recurrence with polynomial coefficients in n of order $\lambda = 4$ in about 2 days.

Given this recurrence, one can utilize algorithms from [55, 57, 58, 69, 75, 70] (see Section 3.1.1) encoded in our package **Sigma** that find all d'Alembertian solutions, i.e., all solutions that can be expressed in terms of indefinite nested sums defined over hypergeometric products. More precisely, **Sigma** computes 4 linearly independent solutions (i.e., their linear span generates all solutions) where

$$\frac{-1}{4(1+n)^2} \sum_{i=3}^n \sum_{j=3}^i \frac{(8-24j+11j^2+3j^3-3j^4+j^5)}{(-2+j)^2(-1+j)^2j^2(1+j)} \sum_{k=1}^j \frac{(-1)^k(-2+k)^2(9-86k+229k^2-156k^3-26k^4+64k^5-26k^6+4k^7)}{(36-20k-26k^2+25k^3-8k^4+k^5)(8-24k+11k^2+3k^3-3k^4+k^5)} \quad (14)$$

is the most complicated sum solution. Finally, with four initial values of the triple sum one finds an alternative representation of it in terms of indefinite nested sums.

In general, these are highly nested, and the summands might consist of ugly polynomials in the denominator (like in (14)) that do not factorize nicely. However, employing our sophisticated difference ring algorithms introduced in Section 6.1, one can simplify the found representation further and obtains the right-hand side in (13). In total, the solving and simplification steps need around 10 seconds.

Summarizing, combining the WZ-approach (recurrence finding) and solving tools, one obtains a summation machinery that can transform a definite nested sum to expressions in terms of indefinite nested sums. When the input sum depends furthermore on the dimensional parameter ε , this machinery has been generalized in [81] to determine the coefficients of the ε -expansion of (2) whenever they are expressible in terms of indefinite nested sums defined over hypergeometric products. This toolbox is very general, but has a substantial drawback: it reaches already with such simple sums like in (13) its limit. With the difference ring approach described next, this situation can be improved substantially.

6.3. The difference ring approach

With the difference ring and field theories worked out in [72, 159, 165, 167, 161, 168] one can simplify not only indefinite nested sums, but one can also apply Zeilberger's creative telescoping paradigm [177]. This means that one can try to compute a linear recurrence of order λ for a definite sum, say $S(n) = \sum_{k=0}^n f(n, k)$, where $f(n, k)$ is given in terms of indefinite nested sums defined over hypergeometric products w.r.t. the summation variable k . Given such a recurrence, one can solve it in terms of indefinite nested sums defined over hypergeometric products by the algorithms given in Section 3.1.1. If one succeeds in combining the solutions accordingly (matching λ initial values), one obtains an alternative representation of $S(n)$. If this expression itself is summed over n , one can repeat this process w.r.t. another variable (over which one may sum later again). In a nutshell, one can apply the summation spiral illustrated in Fig. 2 iteratively with the goal to transform a given multi-sum from inside to outside to a representation purely in terms of indefinite nested sums.




Figure 2. Sigma's summation spiral; see [186].




Figure 3. 2-mass 3-loop diagram.

This interplay has been automated in the package `EvaluateMultiSums` [78] based on `Sigma`'s difference ring algorithms and produces the right-hand side in (13) in about 70 seconds. If a sum depends also on the dimensional parameter ε , one can first expand the summand of the multi-sum w.r.t. ε and can apply afterwards the summation quantifiers to each of the coefficients being free of ε . A clear drawback of this approach is that the summands blow up when higher ε -orders are calculated. Nevertheless, the pure difference ring approach produces simplifications that currently no other toolbox can achieve. E.g., while treating the 2-mass 3-loop integral given in Fig. 3 triple and quadruple sums between 0.4 to 1.6 GB of memory arose in [187] that could be simplified to expressions in terms of binomial sums using 8.4 MB of memory only. Further challenging calculations based on the difference ring/field approach can be found, e.g., in [118, 187–189].

6.4. The holonomic-difference ring approach

Another prominent branch of symbolic summation is the holonomic system approach which has been introduced in [190] and pushed further, e.g., in [191, 192] to determine recurrence relations. Here the summand of a definite sum is described by a system of homogeneous recurrences with polynomial coefficients. Then given such a system, one can try to compute a linear recurrence system by introducing the next definite summation quantifier. Applying these algorithms iteratively from inside to outside yields a linear recurrence in n for the input sum. However, the underlying recurrence systems may grow heavily and the holonomic approach usually fails due to time and memory limitations. In [193] a hybrid strategy has been introduced and developed further in [194, 195] that brings the holonomic and difference ring/field approach under a common umbrella. This new approach allows one to deal with recurrence systems with




Figure 4. A 3-loop ladder diagram with a central triangle [118].

inhomogeneous parts in terms of indefinite nested sums that covers the pure holonomic and difference ring approaches as special cases. So far this approach has been non-trivially applied to obtain the first computer assisted proof [196] of Stembridge’s TSPP theorem [197] and to provide the first proof of a non-trivial identity in [198] that is connected to irrationality proofs of zeta-values. In QCD-calculations this new approach has been explored further to evaluate, e.g., bubble topologies [199].

7. Symbolic integration

In the following we will present some of the most relevant tools of symbolic integration that have been used (at least in parts) in particular for multi-loop calculations in the case of a few number of external legs in elementary particle physics. Other tools suited for lower loop multi-leg calculations are described in part e.g. in [12].

7.1. The hyperlogarithm approach

If a Feynman diagram of the form (1) has no pole terms in (2) (i.e., $l = 0$) or can be made finite by certain transformations splitting off its pole terms [200], it can be calculated under certain conditions by using the method of hyperlogarithms [201]. Since here the denominator of the integral (1) is a multinomial in the Feynman parameters $x_i \in [0, 1]$, one may seek a sequence of integrations, such that the denominator is always a linear function in the integration variable. In this case the Feynman integral can be found as a linear combination of Kummer-Poincaré iterated integrals (also known as Goncharov iterated integrals), [202–206]. The method has been first devised for massless scalar integrals in [201], for a corresponding code see [207], and it has been generalized to massive diagrams [208], dealing even with cases with no thorough multi-linearity, which is an extension to [201, 207].

7.2. The multivariate Almkvist-Zeilberger approach

Similar to the WZ summation approach its continuous version, the multivariate Almkvist-Zeilberger algorithm [185], can compute a linear recurrence/difference equation for a Feynman integral of the form (1). Likewise, if the Feynman integral depends on a continuous parameter x and the integrand is hyperexponential in x , one can search for a linear differential equation of the form

$$\sum_{i=0}^{\lambda} a_i(x, \varepsilon) D_x^i F(x, \varepsilon) = r(x, \varepsilon).$$

A refined and improved method for the input class of Feynman integrals has been developed [209, 118, 96] which can hunt efficiently for homogeneous recurrences or differential equations. E.g., a recurrence in n of order 5 can be calculated in about 8 hours for the master integral

$$\int_0^1 \int_0^1 \int_0^1 \int_0^{1-u} \frac{(x+y-1)^N x^{\varepsilon/2} (1-x)^{\varepsilon/2} y^{\varepsilon/2} (1-y)^{\varepsilon/2} (1-u-v)^N \left(1-u \frac{x}{x-1} - v \frac{y}{y-1}\right)^{-1+3/2\varepsilon}}{u^{1+\varepsilon/2} v^{1+\varepsilon/2}} dx dy du dv$$

that arose in the context to tackle the highly non-trivial 3-loop Feynman diagram given in Fig. 4; see [118]. Then using the the linear difference equation solver of **Sigma** one can compute the first coefficients of the ε -expansion (2) in terms of harmonic sums and generalized harmonic sums. More generally, one can utilize the algorithms from [81, 96] to solve linear difference and differential equations in terms iterated sums and integrals; for further details see Section 3.

7.3. The differential field and holonomic approach

Risch's algorithm [210] for indefinite integration (for details see [211]) allows as input an integrand from the class of elementary functions (they are recursively built by compositions of algebraic, exponential or logarithmic functions and the standard operations $+$, $-$, \times , $/$) and one can decide, if the indefinite integral defined over the input function can be written again in terms of elementary functions. Inspired by this result many further extensions have been derived. In particular, with [212, 211] it is possible to deal with special classes of Liouvillian integrands (recursively built by indefinite integrals and hyperexponentials). In this regard, e.g., the package **Integrator** [213] enables one to treat not only indefinite integration problems, but also to compute difference/differential equations if the integrand depends on a discrete/continuous parameter. These tools have been exploited, e.g., to study root-valued integrals in [66] that arise within massive 3-loop Feynman integral calculations.

In particular, the holonomic system approach [190–192] can be applied not only to multi-sums, but also to multi-integrals of the form (1) to determine a linear recurrence in a discrete parameter n or a linear differential equation in a continuous parameter x . Analogously to the sum case, one can compute stepwise systems of linear differential/difference equations working from inside to outside of the multi-integral and ending up at a scalar equation of the free parameter n or x . First examples have been elaborated in [192, 214] using the package **HolonomicFunctions** that illustrate further possibilities in QCD-calculations.

8. The method of arbitrarily large moments

One is often interested in the calculation of a certain number of moments in the Mellin variable, say $n = 0, 1, 2, \dots, \mu$, to predict extra properties of physical quantities in terms of Feynman integrals. Standard procedures, like **Mincer** [215] or **MATAD** [216],




Figure 5. The large moment engine.

allow the calculation of a comparable small number of Mellin moments, e.g., $\mu = 20$. Recently, a new method has been worked out in Ref. [217] and implemented within the package `SolveCoupledSystem` [218, 50, 219] to compute thousands of such moments.

In general, this new method assumes that we are given a coupled system (9) with (10) where already μ moments for the inhomogeneous part in (9) are computed (by applying this method recursively). Then given such an input, one follows the calculation steps in Fig. 1 but instead of solving the recurrence in step (3), one uses the recurrence together with a small number of initial values of $F_1(n)$ (bounded by the order of the recurrence) to compute in linear time the moments $F_1(0), \dots, F_1(\mu)$, and finally the corresponding moments for $F_2(n), \dots, F_\lambda(n)$. If the $F_i(n)$ depend also on ε , one can calculate the moments of the coefficients of the ε -expansions by exploiting refined ideas from [81].

More generally, using IBP methods [105, 106], we suppose that a physical expression $\bar{f}(x)$ is given in terms of master integrals that are described in terms of recursively defined coupled systems of differential equations. Then using the large moment method iteratively one can calculate for a very large μ the moments of the master integrals. Assembling all the building blocks in the physical expression $\bar{f}(x)$, one finally derives at the coefficients $F(0), \dots, F(\mu)$ of its power series (4) in terms of rational numbers (if ζ -values and other constants arise linearly, they are separated accordingly).

While in traditional solving methods very complicated function spaces might arise in intermediate steps, the large moment method deals simply with rational numbers and one can represent physical quantities with such sequences without entering any structural challenges.

Given a large number of moments, one may follow various strategies illustrated in Fig. 5. First, one can try to obtain interpolation expressions, e.g., by using orthogonal polynomials [220] that provide numerical data of sufficient high precision being relevant for the experiments at the LHC and other future colliders.

Second, one can apply the guessing methods from Section 2.2 in order to produce linear recurrences with minimal order for the physical quantities. In short, analyzing this quantity amounts precisely to the exploration of the computed recurrence.

Next, one can try to solve the recurrences in terms of special functions by using the tools from Section 3. This strategy is particularly successful if the final result (but not necessarily the intermediate results) can be given in terms of indefinite nested sums over hypergeometric products. As demonstrated in [80], we could calculate from about $\mu = 5000$ moments all the recurrences that determine the massless unpolarized 3-loop anomalous dimensions and Wilson coefficients in deep-inelastic scattering [39–41] by solving the recurrences. Similarly, we could calculate, e.g., the 3-loop splitting functions [44], the massive 2- and 3-loop form factor [49, 50], the anomalous dimensions from off shell operator matrix elements [45, 52, 46], the polarized transition matrix element $A_{gq}(N)$ [53] and others, the logarithmic contributions to the polarized $O(\alpha_s^3)$ asymptotic massive Wilson coefficients [54], and the two-loop massless off-shell QCD operator matrix elements [47].

We remark that the found recurrences may also contribute substantially in the case that one fails to find closed form solutions. For instance, one may extract the asymptotic behavior of the physical quantities by using methods described in [221, 222].

9. Special functions and their algorithms

The representation of the results of calculations in QCD and QED are characterized by special constants and functions. The former ones appear in zero scale calculations and as boundary conditions in single and more scale problems. Since in particular in QCD and QED the Mellin transform (see (3) where in the following x is replaced by z) relates nested sums at the one hand to nested integrals at the other hand, and vice versa, two principle classes of special single scale functions emerge: indefinitely nested sums over hypergeometric products and iterated integrals over certain alphabets of letters. Both in the limit $n \rightarrow \infty$ of the sums and at $z = 1$ for the iterated integrals special numbers are obtained. Examples on different classes of functions are given in Tab. 1. All these function spaces obey (quasi) shuffle relations, cf. [172, 173], implying algebraic relations, which allow to reduce to the respective algebraic bases [173, 176].

Historically, most of the Feynman diagram calculations in the time before 1998 were performed using z -space representations leading to classical polylogarithms and Nielsen integrals [223–228], partly with involved arguments.

A systematic description in terms of harmonic sums started in 1998 with [61, 62]. They are defined by

$$S_{b,\bar{a}}(n) = \sum_{k=1}^n \frac{(\text{sign}(b))^k}{k^{|b|}} S_{\bar{a}}(k), \quad S_{\emptyset} = 1, \quad b, a_i \in \mathbb{Z} \setminus \{0\}, n \in \mathbb{N} \setminus \{0\}. \quad (15)$$

Related to that, the iterative integrals are the harmonic polylogarithms,

$$H_{b,\bar{a}}(z) = \int_0^z dy f_b(y) H_{\bar{a}}(y), \quad H_{\emptyset} = 1, \quad b, a_i \in \{0, 1, -1\}, \quad (16)$$

Nested sums	Nested integrals	Special numbers
Harmonic Sums $\sum_{k=1}^n \frac{1}{k} \sum_{l=1}^k \frac{(-1)^l}{l^3}$	Harmonic Polylogarithms $\int_0^z \frac{dy}{y} \int_0^y \frac{dx}{1+x}$	multiple zeta values $\int_0^1 dx \frac{\text{Li}_3(x)}{1+x} = -2\text{Li}_4(1/2) + \dots$
gen. Harmonic Sums $\sum_{k=1}^n \frac{(1/2)^k}{k} \sum_{l=1}^k \frac{(-1)^l}{l^3}$	gen. Harmonic Polylogarithms $\int_0^z \frac{dy}{y} \int_0^y \frac{dx}{x-3}$	gen. multiple zeta values $\int_0^1 dx \frac{\ln(x+2)}{x-3/2} = \text{Li}_2(1/3) + \dots$
Cycl. Harmonic Sums $\sum_{k=1}^n \frac{1}{(2k+1)} \sum_{l=1}^k \frac{(-1)^l}{l^3}$	Cycl. Harmonic Polylogarithms $\int_0^z \frac{dy}{1+y^2} \int_0^y \frac{dx}{1-x+x^2}$	cycl. multiple zeta values $\mathbf{C} = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)^2}$
Binomial Sums $\sum_{k=1}^n \frac{1}{k^2} \binom{2k}{k} (-1)^k$	root-valued iterated integrals $\int_0^z \frac{dy}{y} \int_0^y \frac{dx}{x\sqrt{1+x}}$	associated numbers $\text{H}_{8,w_3} = 2\arccot(\sqrt{7})^2$
	iterated integrals on ${}_2F_1$'s $\int_0^z \frac{\ln(x)}{1+x} {}_2F_1\left[\frac{4}{3}, \frac{5}{3}; \frac{x^2(x^2-9)^2}{(x^2+3)^3}\right] dx$	associated numbers $\int_0^1 {}_2F_1\left[\frac{4}{3}, \frac{5}{3}; \frac{x^2(x^2-9)^2}{(x^2+3)^3}\right] dx$

Table 1. Special functions and numbers.

with the alphabet

$$\mathfrak{A} = \left\{ f_0(z) = \frac{1}{z}, f_1(z) = \frac{1}{1-z}, f_{-1}(z) = \frac{1}{1+z} \right\}, \quad (17)$$

[91]. A special definition is required for the case $\text{H}_{\vec{a}}(z), \forall a_i = 0, i = 1 \dots n$, which has no integral representation, but is defined as $\ln^n(z)/n!$, for completeness. In the case of infinite sums we also allow for the symbol $\sigma_{\infty} := \sum_{k=1}^{\infty} (1/k)$, which is not a number, but simplifies various algebraic relations and is therefore useful.

The special numbers are multiple zeta values in both cases. Their representations at high order can be found in [35].

At the next level, generalized harmonic sums [64, 65] contribute, e.g. in the case of the pure singlet 3-loop massive Wilson coefficients in the asymptotic region [43]. These quantities are given by

$$S_{b,\vec{a}}(\{c, \vec{d}\}; n) = \sum_{k=1}^n \frac{c^k}{k^b} S_{\vec{a}}(\{\vec{d}\}; k), \quad S_{\emptyset} = 1, \\ b, a_i \in \mathbb{N} \setminus \{0\}, c, d_i \in \mathbb{C} \setminus \{0\}, n \in \mathbb{N} \setminus \{0\}. \quad (18)$$

The corresponding iterated integrals are also called Kummer–Poincaré iterated integrals [202–206] and are given by

$$\text{H}_{b,\vec{a}}(z) = \int_0^z dy f_b(y) \text{H}_{\vec{a}}(y), \quad \text{H}_{\emptyset} = 1, \quad (19)$$

with the alphabet

$$\mathfrak{A} = \left\{ f_{c_i}(z) = \frac{1}{z - c_i}, \quad c_i \in \mathbb{C} \right\}. \quad (20)$$

Further, cyclotomic harmonic sums and polylogarithms [63] contribute. The letters of the alphabet forming the iterated integrals are those of the harmonic polylogarithms extended with letters of the type

$$f_{k,a}^{\text{cycl.}}(z) = \frac{z^a}{P_k(z)}, \quad k \geq 3, \quad (21)$$

with k labeling the cyclotomic polynomials and $a \in [0, \varphi(k)]$, and $\varphi(k)$ is Euler's totient function. The associated cyclotomic harmonic sums iterate monomials of the type

$$\frac{s^k}{(ak+b)^c}, \quad a, c \in \mathbb{N}_+, b \in \mathbb{N}, s \in \mathbb{C} \setminus \{0\}. \quad (22)$$

Finite binomial sums [66] contribute for a series of topologies in the massive OMEs $A_{gg}^{(3)}$ [229] and $A_{Qg}^{(3)}$ [118]. The corresponding sums are generalized sums with an additional factor of $\binom{2k}{k}$ in the numerator or denominator. The associated iterated integrals, obtained by a Mellin inversion, are formed out of letters containing square root valued structures, as e.g. shown in Tab. 1. Another example is

$$\begin{aligned} \sum_{i=1}^n \frac{1}{\binom{2i}{i} i^3} &= \sum_{i=1}^{\infty} \frac{1}{\binom{2i}{i} i^3} \\ &+ \frac{1}{4^n} \int_0^1 z^n \left(\frac{3 \ln(z)^2 - 12 \ln(2) \ln(z) + 12 \ln(2)^2 - \pi^2}{6(-4+z)} + \frac{\int_0^z \frac{1}{\tau_1} \int_0^{\tau_1} \frac{\sqrt{1-\tau_2}-1}{\tau_2} d\tau_2 d\tau_1 - 2 \int_0^z \frac{\sqrt{1-\tau}-1}{\tau} d\tau}{-4+z} \right) dz. \end{aligned}$$

In particular, iterative application of integration by parts yield the asymptotic expansion

$$\sum_{i=1}^n \frac{1}{\binom{2i}{i} i^3} \sim 2^{-2n} \sqrt{n} \sqrt{\pi} \left(-\frac{34924547}{884736n^7} + \frac{91999}{9216n^6} - \frac{10537}{3456n^5} + \frac{77}{72n^4} - \frac{1}{3n^3} + O\left(\frac{1}{n^8}\right) \right) + \sum_{i=1}^{\infty} \frac{1}{\binom{2i}{i} i^3}. \quad (23)$$

Such expansions are extremely useful for limit calculations and for analyzing the expression behavior for large values of n . Moreover, the sum and integral representations equipped with their shuffle and quasi-shuffle algebras [172–175] give rise to algebraic relations of infinite sums. In particular, attaching special constants to sums that cannot be simplified further, one can discover evaluations such as

$$\sum_{i=1}^{\infty} \frac{2^i S_1(i)}{i \binom{2i}{i}} = 2C - \frac{\pi \log(2)}{2} + \frac{3}{4} \zeta(2)$$

where $C = \sum_{i=1}^{\infty} \frac{(-1)^i}{(2i+1)^2}$ denotes the Catalan constant; for further details see [230, 231]. For general classes, like nested binomial sums, more flexible methods were developed recently to map between n - and z -space, cf. [187]: given a recurrence of $F(n)$ (resp. a differential equation of $f(z)$), compute a differential equation for $f(z)$ (resp. a recurrence for $F(n)$). In particular, using the introduced solvers from Section 3.1, one can check, if the Mellin transform (resp. inverse Mellin transform) can be given in terms of indefinite nested sums (resp. integrals). Infinite (inverse) binomial sums have been also studied in [67, 68]. For the simpler cases efficient rewrite rules have been developed to switch between the sum and integral representations via the (inverse) Mellin transform.

In more general cases, in particular in two-scale problems, the so called G -functions appear, which are iterated integrals over larger alphabets, partly with root-valued letters. They are given by

$$G(f_a(x), f_{b_1}(x), \dots, f_{b_n}(x)) = \int_0^x dy f_a(y) G(f_{b_1}(y), \dots, f_{b_n}(y)). \quad (24)$$

Actually $f_c(x)$ even denotes in general a differentiable function, up to regularizations in special cases.

At higher and higher orders in perturbation theory, new building blocks arise that cannot be represented in terms of indefinite nested sums or iterated integrals. In particular, one ends up at linear difference/differential equations, that cannot be solved completely in terms of d'Alembertian/Liouvillian solutions. For this reason, the class of iterative non-iterative integrals have been introduced in 2016 [232].

Probably the first case in which complete elliptic integrals emerged in a quantum field theoretic calculation has been the fourth order spectral functions for the electron propagator by Sabry 1962 [233]. For complete physical processes more recently elliptic integrals were needed. This is the case in massive three-loop calculations for the QCD corrections of the ρ parameter [98, 99] 2017^{††} and in massless three-three loop calculations 2018 and later [235, 236]. There is a series of well-known examples of individual integrals of a certain structure in the literature as the sun-rise integral, cf. e.g. [237–239] and the kite-integral [233, 240, 241]; for a collection of recent surveys see Ref. [242].^{‡‡} In general, these classes of integrals form iterative non-iterative integrals, cf. [98]. Beyond this level one has Abel-integrals [246] and Calabi–Yau structures, cf. [247, 24, 248]. Even more involved structures will occur at higher topologies. In Mellin space they have the common characteristics of difference equations with rational coefficients which are not factorizing at first order. Any of the corresponding solutions also needs efficient numerical representations, as e.g. [249–251], for phenomenological and experimental applications. This also applies to Mellin space representations for $n \in \mathbb{C}$, [252–254, 174, 175]. Feynman integrals will imply a multitude of new function spaces in the future.

10. Calculations in Quantum Field Theory

Our major topic concerns analytic Feynman diagram calculations. As has been shown, this is deeply rooted in solving large systems of differential or difference equations. The single scale cases are mathematically widely understood and one may project to the zero scale case, i.e. to special numbers. However, just by this one will probably not be able to find all relations between these special numbers by using, e.g., the techniques described in Section 2.1, beginning at a certain level of complexity, which requires further advanced methods in these cases. On the other side, as experience shows, certain two-scale problems can still be solved analytically, as we will discuss in Section 10.4. But already starting at that level, one has to deal with partial differential and difference

^{††}The same differential equations rule the non first-order factorizing cases in the calculation of the massive three-loop operator matrix element A_{Qg} in the single mass case [234], found together with the calculation [99].

^{‡‡}Very naturally, as now the technical aspects on complete elliptic integrals are very well known in particle physics, many applications find these contributions, cf. e.g. [243–245], the reason being the occurrence of the corresponding Heun and ${}_2F_1$ -type differential equations, cf. e.g. [98].

equations, on which is much less known, cf. [255, 256]. In the single scale case, going to higher and higher orders, one will face non-first order factorizing differential and difference equations of higher and higher order [257, 258, 246]§§, for which only the properties of very few concrete classes have been studied so far and a very wide field of future mathematical investigation is opening up.

For more scales, one probably will have to rely on using numerical precision methods in the first place, because of the wide variety of structures [25]. Computational Quantum Field Theory (QFT) is urged to invest much more efforts to obtain fast and highly reliable methods in this direction to be able to cope with the challenges in future precision measurements. Developments of this kind may take quite a long time and need intense collaboration with experts in the field of numerical mathematics.

10.1. Zero Scale Calculations

Zero scale quantities in QFTs, as QED and QCD, are characterized by color factors, rational coefficients and special numbers like multiple zeta values [35]. Examples are fixed moments for massive three-loop OMEs [259] and massless four-loop anomalous dimensions [260]. Particularly for massive problems more special numbers contribute, as those related to generalized harmonic sums [65], cyclotomic harmonic sums [63], binomial sums [66], and those related to elliptic integrals [261], see also Tab. 1. More and more different sets will emerge including even higher topologies. One also may calculate moments of single scale quantities, which depend on an integer parameter n , by obtaining sequences of rational numbers. These numbers incorporate thus an essential part of the more involved single scale dependence for general values of n . It is sometimes of advantage to first work with these moments, despite the fact that the general n relation is determined by a difference equation, which does not factorize at first order. This is often the case for master integrals in the massive case. However, the corresponding recurrences for anomalous dimensions are factorizing at first order. One inserts first the master integrals for fixed moments and then determines the difference equation for the anomalous dimension, see [44, 52].

10.2. Massless Single Scale Calculations

These quantities are the anomalous dimensions, currently known to three-loop order [39, 40, 262, 43, 44, 52, 45, 46], the massless Wilson coefficients for deep-inelastic scattering [263] up to $O(\alpha_s^3)$ [41], the Drell-Yan process and Higgs production to two-loop order [264–267]. All these quantities can be expressed by harmonic sums [62, 61] in Mellin space and by harmonic polylogarithms in z -space [91]. For Higgs production and the Drell-Yan process at three-loop order [235, 236] also elliptic integrals contribute. It is generally expected that a further nesting in the Feynman diagram topologies leads to

§§For more literature on elliptic integrals and modular forms see Refs. [26, 98, 242].

new mathematical structures also in the massless case in higher orders of the coupling constant.

For massless single scale calculations one may very efficiently apply the method of arbitrarily high moments [217], together with guessing to obtain the recurrences, which may either be solved or reduced, by factoring of the first order factors, using `Sigma`. This also applies to the case of massive single scale calculations, to which we turn now.

10.3. Massive Single Scale Calculations

The method of massive OMEs [268] allowed to calculate single scale quantities, such as the heavy flavor Wilson coefficients to three-loop order in the asymptotic region, obtaining all logarithmic contributions [269, 54] and also the constant term. The method has also been applied to problems in QED, cf. [270–272], see Section 10.3.1. In some cases even full results have been obtained at two-loop order [268, 273–275, 270], cf. Section 10.3.2.

10.3.1. Massive Single Scale Calculations: logarithmic and constant corrections The asymptotic heavy flavor Wilson coefficients of deep-inelastic scattering contain single scale logarithmic and constant contributions. At two-loop order all contributions are known [268, 276, 277, 153, 278, 279]. At three-loop order all but the massive OME $A_{Qg}^{(3)}$ have been calculated analytically in complete form both in the unpolarized and polarized case. In Mellin n space they can be expressed by harmonic sums for all N_F -terms [280, 122], and for $A_{qq,Q}^{(3),NS}$, $A_{qq,Q}^{(3),PS}$, $A_{qg,Q}^{(3)}$ and $A_{gg,Q}^{(3)}$ [281, 269, 123, 188, 54]. Generalized harmonic sums contribute in the pure singlet case $A_{Qq}^{(3),PS}$ [43, 282] and finite binomial sums for $A_{gg,Q}^{(3)}$ [229]. Finally, $A_{Qg}^{(3)}$ receives also contributions by complete elliptic integrals [234].

Another case belonging to this class of integrals are the contributions to the massive form factor at three-loop order. The first order factorizing contributions can be expressed by harmonic and cyclotomic harmonic polylogarithms in the variable x

$$\frac{q^2}{m^2} = -\frac{(1-x)^2}{x}, \quad (25)$$

with q^2 the virtuality and m the heavy quark mass, [49, 283, 117, 284, 285, 50]

The method of massive OMEs has also been applied for the calculation of the initial state radiation to the process $e^+e^- \rightarrow Z^*/\gamma^*$. These corrections are of importance for planned future high luminosity measurements at the ILC, CLIC, and FCC-ee. The results at $O(\alpha^2)$ [270] showed disagreement with an earlier direct calculation [286]. The only way to find the correct results has been a complete diagrammatic calculation, without expansion in the small parameter $\rho = m_e^2/s$, with m_e the electron mass and s the cms energy. This has been performed in Ref. [287]. Furthermore, we expanded in ρ through different steps, controlled by high precision numerics, and confirmed the results of [270]. The fermionic integrals could be represented using iterated Kummer-elliptic integrals over larger alphabets. Numerical results were presented in [288, 289].

The method of massive OMEs has then been extended to calculate the first three logarithmic series up to $O(\alpha^6 L^5)$, where $L = \ln(s/m_e^2)$ in [271]. Here in Mellin space also generalized harmonic sums contribute. Higher order corrections for the forward–backward asymmetry were calculated in [272], where also cyclotomic harmonic polylogarithms contribute to the radiators. ¶¶

10.3.2. Massive Single Scale Calculations: including also power corrections In some cases, massive two–loop problems can be integrated analytically in the whole kinematic region. This applies to the flavor non–singlet contributions [268,273] and the pure singlet contributions [274,275]. While in the non–singlet case classical polylogarithms with root–valued arguments suffice for the representation, in the pure singlet case iterated integrals over alphabets containing Kummer–elliptic letters are necessary. Part of them integrates to incomplete elliptic integrals, which do not destroy the iterated integral structure, unlike the case in the iterative non–iterative integrals [291]. In establishing the contributing alphabet also rationalization of roots is performed as far as possible; for other investigations see also [292]. Examples for these letters are, cf. [274]

$$f_{w_{11}}(t) = \frac{t}{\sqrt{1-t^2}\sqrt{1-k^2t^2}} \quad (26)$$

$$f_{w_{12}}(t) = \frac{t}{\sqrt{1-t^2}\sqrt{1-k^2t^2}(k^2(1-t^2(1-z^2)) - z^2)}, \quad (27)$$

with $k = \sqrt{z}/\sqrt{1-(1-z)\beta^2}$, $\beta = \sqrt{1-4m^2z/(Q^2(1-z))}$ and z is the momentum fraction variable. Depth-three iterated integrals over the contributing integrals emerge.

The fact that one finds analytic integral representations in these cases is related to the tree–like structure of the contributing diagrams. At higher orders or for other processes, correspondingly, one has to perform corresponding expansions in m^2/Q^2 , to successively obtain analytic results, improving the logarithmic and constant orders obtained in the region $Q^2 \gg m^2$. The possibility to analytically calculate the pure–singlet corrections, conjectured by van Neerven and J.B. around 2000, turned out to be correct, however, the necessary technologies for this became only available with [274] later and Nielsen integrals with whatsoever complicated argument are not sufficient to represent the final result.

10.4. Massive Double Scale Calculations

In the case of deep–inelastic scattering from the level of three loop onward, diagrams contribute, which contain charm and bottom quarks. This leads to a double scale problem, as similarly also in the case of the massive from factor and for other processes. In the following we will consider the case of deep–inelastic scattering. In all cases but the massive OMEs $A_{Qg}^{(3)}$ and its polarized counterpart $\Delta A_{Qg}^{(3)}$, complete analytic solutions are possible either in Mellin n or momentum fraction z –space. The different OMEs

¶¶For a recent survey on the QED corrections see [290].

have been calculated in Refs. [293, 294, 187, 295, 189, 188]. They can be expressed by G -functions, cf. (24). One example is [187]

$$\begin{aligned}
G_{37} &= G\left(\left\{\frac{1}{1-x+\eta x}, \sqrt{1-x}\sqrt{1-x+\eta x}\right\}, z\right) \\
&= -\frac{\eta^2}{(1-\eta)^{5/2}}\left\{\frac{1}{16}\ln\left(2-\eta+2\sqrt{1-\eta}\right)+\frac{1}{4}\arcsin^2\left(\frac{1}{\sqrt{\eta}}\right)\right. \\
&\quad +\frac{i}{2}\left[-\ln\left(1-\sqrt{1-\eta}\right)+\ln(\eta)-\ln(2)\right]\arcsin\left(\frac{1}{\sqrt{\eta}}\right) \\
&\quad +\frac{1}{4}\text{Li}_2\left(\frac{\eta+2\sqrt{1-\eta}-2}{\eta}\right)+\frac{i}{2}\arcsin\left(\frac{\sqrt{1-z+\eta z}}{\sqrt{\eta}}\right)\ln(1+\chi) \\
&\quad -\frac{1}{8}\ln\left(\sqrt{1-\eta}\sqrt{1-z}+\sqrt{1-z+\eta z}\right)-\frac{1}{4}\arcsin^2\left(\frac{\sqrt{1-z+\eta z}}{\sqrt{\eta}}\right) \\
&\quad +\frac{1}{4}\ln(1-z+\eta z)\left[-\ln\left(1-\eta+\sqrt{1-\eta}\right)+\frac{\ln(\eta)}{2}+\frac{1}{2}\ln(1-\eta)-\frac{i\pi}{2}\right] \\
&\quad \left.-\frac{1}{4}\text{Li}_2(-\chi)\right\}+\frac{3\eta-2}{8(1-\eta)^2}-\frac{(2-\eta)}{4(1-\eta)^2}\ln(1-z+\eta z) \\
&\quad +\sqrt{1-z}\sqrt{1-z+\eta z}\frac{(2-3\eta+2\eta z-2z)}{8(1-\eta)^2}, \tag{28}
\end{aligned}$$

where $\eta = m_1^2/m_2^2$ and $\chi = (1/\eta)(\sqrt{1-\eta}\sqrt{1-z}-\sqrt{1-z+\eta z})^2$. In the case of the pure singlet two-mass contributions [294, 189] we work in z -space by using Mellin-Barnes integrals [140–142]. One also obtains G -functions and in part integrals over them, with a different support than usual, expressed by Heaviside functions. These problems cannot be solved in Mellin n space.

10.5. Classical Gravity

The classical kinematics of massive astrophysical objects, such as black holes and neutron stars, can be calculated by using methods of effective field theory developed for Quantum Field Theory. Concepts like the path integral [296] and Feynman diagrams are also applicable at the classical level. This is an enormous bonus to the field of general relativity and classical gravity, since very advanced computation technologies, starting from Feynman diagram generation [297], effective performance of Lorentz algebra [298, 299], integration-by-parts reduction [107], and the calculation of master integrals already exist. One expands Einstein–Hilbert gravity in terms of auxiliary fields [300]. Furthermore, one performs the classical limit using the method of expansion by regions, cf. [301, 302], where only the potential and radiation modes are contributing. These methods can be applied for the inspiraling process of the massive objects [303–308], as well as for their scattering process, cf. [309, 310]. The main challenge is here to deal with the ever growing effective vertex structures and the integration by parts reduction, which can be performed by packages like *Crusher*, cf. [107]. The bound

state kinematics is described by the post-Newtonian (PN) approach, having reached now 6PN order [311, 308, 306] and the scattering process by the post-Minkowskian approach, now available at $O(G_N^4)$, where G_N denotes Newton's constant [309, 310]. After expanding the potential contributions of the post-Minkowskian results, a part of the PN results is re-obtained, which has been shown to 6PN in [311, 306]. This applies to the potential contributions. In principle, the method of guessing could be used obtaining post-Minkowskian results, again from potential contributions, [312]. While agreement has been reached for the 4PN level between various approaches, the level of 5PN is still under discussion because of the non-potential contributions. For their description the different methods proposed in the literature do not lead to consistent results as of yet, requiring both more clear theoretical foundations and also more work to obtain the final result for bound state problems.

11. Conclusion

For less than the last quarter century, a technological revolution has happened in the field of perturbative analytic calculations in renormalizable Quantum Field Theories, which is accompanying this field since. Considering single scale Feynman diagrams, in the time until 1998, the analytic integration of these amplitudes has been an art based on hypergeometric function structures and maximally Nielsen integrals dealing with sets of up to $O(50)$ Feynman diagrams mostly to two-loop order. Before about this time computer algebra has been inspired and motivated often by its own discipline or by other mathematical research areas, such as combinatorics, number theory, or special functions. In this survey article we introduced recent methods from both communities and showed how they can be combined in non-trivial ways to new methods that may be instrumental for current and future precision calculations in particle physics. For a graphical summary of the different interactions presented in this article we refer to Fig. 6.

Elements of the revolution in Quantum Field Theories were also (quasi)shuffle algebras and the discovery of a hierarchy of function spaces both in Mellin n and momentum fraction z space. For all quantities considered one may find recurrences by applying the methods of arbitrary large moments and guessing, which delivers closed form equations in the first place. In the moment technology presented in Section 8 one simply deals with rational numbers ignoring completely the possible function spaces that may arise there. At the end of the day, one can apply the computer algebra tools introduced above and obtains from this data numerical representations or even symbolic representations of the final physical problem. In general, we feel that such new strategies will be crucial for future calculations and we are curious to see how these techniques can be developed further or can be outperformed with new ideas and strategies.

Systematic mathematical methods, like difference ring theory, allowed to reveal various new structures. Nowadays first order factorizing difference and differential equations (or systems thereof) are fully understood. Non-first order factorizing systems,




Figure 6. The computer algebra and special function tools in interaction

leading to ${}_2F_1$ -solutions, complete elliptic integrals and modular forms are understood as well and steps in the direction of equations related to Calabi–Yau manifolds are done. Yet these are rather special systems only and Feynman diagrams can in principle cause more general, yet unknown structures also belonging to non–first order factorizing equations. They are fascinating as such and their complex analysis is a highly interesting topic. One may intend to derive general characteristics for these quantities [313]. Many of the present massless and massive three-loop problems of single and double scales could be solved by the technologies described in the present survey and new structures challenge further innovative mathematical solutions and efficient computer-algebraic implementations. All present achievements have in various instances been achieved by sophisticated computer algebra algorithms. Another challenge comes from the experimental possibilities at future colliders, operating at high luminosity, with which the theoretical results have to cope. All methods described do not only apply to relativistic renormalizable Quantum Field Theories, but also to effective field theories, e.g. dealing with (non–linear) Einstein general relativity in post–Newtonian and post–Minkowskian expansions at the classical level and various applications more, e.g. also to solid state physics. Problems with more scales do still escape complete analytic solutions at present and require more research in the future.

The close collaboration of theoretical physicist, mathematicians and researchers in the field of computer algebra led both to the use of known algorithms from quite different fields in Quantum Field Theory, but have also triggered new mathematical and

algorithmic research. The success reached has only been possible due to this symbiosis. This process will continue in full strength in the future.

Acknowledgments

We would like to thank D. Kreimer for a discussion. This work was supported by the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 764850 “SAGEX”. We also acknowledge support from the Austrian Science Fund (FWF) grants SFB F50 (F5009-N15) and P33530. The Feynman diagrams have been drawn using `Axodraw` [314].

- [1] Boer D *et al.* 2011 *Gluons and the quark sea at high energies: Distributions, polarization, tomography* (Preprint 1108.1713)
- [2] Agostini P *et al.* (LHeC, FCC-he Study Group) 2021 *J. Phys. G* **48** 110501 (Preprint 2007.14491)
- [3] Abelleira Fernandez J L *et al.* (LHeC Study Group) 2012 *J. Phys. G* **39** 075001 (Preprint 1206.2913)
- [4] Accomando E *et al.* (ECFA/DESY LC Physics Working Group) 1998 *Phys. Rept.* **299** 1–78 (Preprint hep-ph/9705442)
- [5] Aguilar-Saavedra J A *et al.* (ECFA/DESY LC Physics Working Group) 2001 TESLA: The Superconducting electron positron linear collider with an integrated x-ray laser laboratory. Technical design report. Part 3. Physics at an e^+e^- linear collider (Preprint hep-ph/0106315)
- [6] Aarons G *et al.* (ILC) 2007 International Linear Collider Reference Design Report Volume 2: Physics at the ILC ed Djouadi A, Lykken J, Mönig K, Okada Y, Oreglia M and Yamashita S (Preprint 0709.1893)
- [7] Aihara H *et al.* (ILC) 2019 The International Linear Collider. A Global Project (Preprint 1901.09829)
- [8] van der Meer S 1988 *The CLIC project and design for an e^+e^- collider INFN International School on Electromagnetic Radiation and Particle Beams Acceleration: Physics and Applications*
- [9] Accomando E *et al.* (CLIC Physics Working Group) 2004 Physics at the CLIC multi-TeV linear collider *11th International Conference on Hadron Spectroscopy CERN Yellow Reports: Monographs* (Preprint hep-ph/0412251)
- [10] Roloff P *et al.* (CLIC, CLICdp) 2018 The Compact Linear e^+e^- Collider (CLIC): Physics Potential ed Roloff P, Franceschini R, Schnoor U and Wulzer A (Preprint 1812.07986)
- [11] Abada A *et al.* (FCC) 2019 *Eur. Phys. J. ST* **228** 261–623
- [12] Abreu S, Britto R and Duhr C 2022 *J. Phys. A: Math. Theor.* **55** 443004 (Preprint 2203.13014)
- [13] Duhr C, Gangl H and Rhodes J R 2012 *JHEP* **10** 075 (Preprint 1110.0458)
- [14] Connes A and Kreimer D 1998 *Commun. Math. Phys.* **199** 203–242 (Preprint hep-th/9808042)
- [15] Connes A and Kreimer D 2000 *Commun. Math. Phys.* **210** 249–273 (Preprint hep-th/9912092)
- [16] Connes A and Kreimer D 2001 *Commun. Math. Phys.* **216** 215–241 (Preprint hep-th/0003188)
- [17] Manin Y I 2005 *Iterated integrals of modular forms and noncommutative modular symbols*, in Algebraic geometry and number theory, vol. 253 of Progr. Math., (Boston), pp. 565–597, Birkhäuser Boston, 2006. (Preprint math/0502576)
- [18] Bloch S, Esnault H and Kreimer D 2006 *Commun. Math. Phys.* **267** 181–225 (Preprint math/0510011)
- [19] Bloch S and Kreimer D 2010 *Commun. Num. Theor. Phys.* **4** 709–753 (Preprint 1007.0338)
- [20] Brown F 2011 *On the decomposition of motivic multiple zeta values Galois-Teichmüller theory and arithmetic geometry* (Adv. Studies in Pure Math. vol 68) (Math. Soc. Japan) pp 31–58 (Preprint 1102.1310)

- [21] Golden J, Goncharov A B, Spradlin M, Vergu C and Volovich A 2014 *JHEP* **01** 091 (*Preprint* 1305.1617)
- [22] Broadhurst D and Schnetz O 2014 *PoS* **LL2014** 078 (*Preprint* 1409.5570)
- [23] Brown F 2017 *Commun. Num. Theor. Phys.* **11** 453–556 (*Preprint* 1512.06409)
- [24] Bönisch K, Duhr C, Fischbach F, Klemm A and Nega C 2021 Feynman Integrals in Dimensional Regularization and Extensions of Calabi-Yau Motives (*Preprint* 2108.05310)
- [25] Heinrich G 2021 *Phys. Rept.* **922** 1–69 (*Preprint* 2009.00516)
- [26] Blümlein J and Schneider C 2018 *Int. J. Mod. Phys. A* **33** 1830015 (*Preprint* 1809.02889)
- [27] Blümlein J 2021 Analytic Integration Methods in Quantum Field Theory: An Introduction *Anti-Differentiation and the Calculation of Feynman Amplitudes* ed Blümlein J and Schneider C (Cham: Springer International Publishing) pp 1–33 (*Preprint* 2103.10652)
- [28] Weinzierl S 2022 *Feynman Integrals* arXiv:2201.03593 [hep-th]
- [29] Lenstra A K, Lenstra H W and Lovasz L 1982 *MATH. ANN* **261** 515–534
- [30] Ferguson H R P and Forcade R W 1979 *Bulletin of the American Mathematical Society* **1** 912–914
- [31] Ferguson H R P and Bailey D H 1991 *A Polynomial Time, Numerically Stable Integer Relation Algorithm* Tech. rep. RNR-91-032
- [32] Bailey D H and Broadhurst D J 1999 *Mathematics of Computation* **70** 1719–1736
- [33] Luthe T, Maier A, Marquard P and Schröder Y 2017 *JHEP* **10** 166
- [34] Acres K and Broadhurst D 2021 Empirical determinations of Feynman integrals using integer relation algorithms *Anti-Differentiation and the Calculation of Feynman Amplitudes* ed Blümlein J and Schneider C (Cham: Springer International Publishing) pp 63–82
- [35] Blümlein J, Broadhurst D J and Vermaseren J A M 2010 *Comput. Phys. Commun.* **181** 582–625 (*Preprint* 0907.2557)
- [36] Salvy B and Zimmermann P 1994 *ACM Trans. Math. Software* **20** 163–177
- [37] Mallinger C 1996 *Algorithmic Manipulations and Transformations of Univariate Holonomic Functions and Sequences* Master’s thesis RISC, J. Kepler University, Linz
- [38] Kauers M 2008 *Nucl. Phys. B (Proc. Suppl.)* **183** 245–250
- [39] Moch S, Vermaseren J A M and Vogt A 2004 *Nucl. Phys. B* **688** 101–134 (*Preprint* hep-ph/0403192)
- [40] Vogt A, Moch S and Vermaseren J 2004 *Nucl. Phys. B* **691** 129–181
- [41] Vermaseren J, Vogt A and Moch S 2005 *Nucl. Phys. B* **724** 3–182
- [42] Blümlein J, Kauers M, Klein S and Schneider C 2008 *PoS* **ACAT08** 106 (*Preprint* 0902.4095)
- [43] Ablinger J, Behring A, Blümlein J, De Freitas A, von Manteuffel A and Schneider C 2014 *Nucl. Phys. B* **890** 48–151 (*Preprint* 1409.1135)
- [44] Ablinger J, Behring A, Blümlein J, De Freitas A, von Manteuffel A and Schneider C 2017 *Nucl. Phys. B* **922** 1–40 (*Preprint* 1705.01508)
- [45] Blümlein, J and Marquard, P and Schneider, C and Schönwald, K 2021 *Nucl. Phys. B* **971** 115542 (*Preprint* 2107.06267)
- [46] Blümlein J, Marquard P, Schneider C and Schönwald K 2022 *JHEP* **01** 193 (*Preprint* 2111.12401)
- [47] Blümlein J, Marquard P, Schneider C and Schönwald K 2022 *Nucl. Phys. B* **980** 115794 (*Preprint* 2202.03216)
- [48] Kauers M, Jaroschek M and Johansson F 2014 Ore polynomials in Sage *Computer Algebra and Polynomials* Lecture Notes in Computer Science ed Gutierrez J, Schicho J and Weimann M pp 105–125
- [49] Ablinger J, Blümlein J, Marquard P, Rana N and Schneider C 2018 *Phys. Lett. B* **782** 528–532 (*Preprint* 1804.07313)
- [50] Blümlein J, Marquard P, Rana N and Schneider C 2019 *Nucl. Phys. B* **949** 114751 (*Preprint* 1908.00357)
- [51] Ablinger J, Behring A, Blümlein J, De Freitas A, von Manteuffel A and Schneider C 2017 Heavy flavor Wilson coefficients in deep-inelastic scattering: Recent results *PoS(QCDEV2017)031* ed Gamberg L, Prokudin A, Qiu J and Radyushkin A pp 1–10

- [52] Behring A, Blümlein J, De Freitas A, Goedicke A, Klein S, von Manteuffel A, Schneider C and Schönwald K 2019 *Nucl. Phys. B* **948** 114753 (*Preprint* 1908.03779)
- [53] Behring A, Blümlein J, De Freitas A, von Manteuffel A, Schönwald K and Schneider C 2021 *Nucl. Phys. B* **964** 115331 (*Preprint* 2101.05733)
- [54] Blümlein J, De Freitas A, Saragnese M, Schneider C and Schönwald K 2021 *Phys. Rev. D* **104** 034030 (*Preprint* 2105.09572)
- [55] Abramov S 1989 *U.S.S.R. Comput. Math. Math. Phys.* **29** 7–12
- [56] Van Hoeij M 1997 *J. Symb. Comput.* **24** 537–561
- [57] Petkovšek M 1992 *J. Symbolic Comput.* **14** 243–264
- [58] van Hoeij M 1999 *J. Pure Appl. Algebra* **139** 109–131
- [59] Abramov S and Petkovšek M 1994 D’Alembertian solutions of linear differential and difference equations *Proc. ISSAC’94* ed von zur Gathen J (ACM Press) pp 169–174
- [60] Abramov S and Zima E V 1996 D’Alembertian solutions of inhomogeneous linear equations (differential, difference, and some other) *Proc. ISSAC’96* (ACM Press) pp 232–240
- [61] Blümlein J and Kurth S 1999 *Phys. Rev. D* **60** 014018 (*Preprint* hep-ph/9810241)
- [62] Vermaseren J 1999 *Int. J. Mod. Phys. A* **14** 2037–2976
- [63] Ablinger J, Blümlein J and Schneider C 2011 *J. Math. Phys.* **52** 102301 (*Preprint* 1105.6063)
- [64] Moch S, Uwer P and Weinzierl S 2002 *J. Math. Phys.* **6** 3363–3386
- [65] Ablinger J, Blümlein J and Schneider C 2013 *J. Math. Phys.* **54** 082301 (*Preprint* 1302.0378)
- [66] Ablinger J, Blümlein J, Raab C G and Schneider C 2014 *J. Math. Phys.* **55** 112301 (*Preprint* 1407.1822)
- [67] Davydychev A I and Kalmykov M Yu 2004 *Nucl. Phys. B* **699** 3–64
- [68] Weinzierl S 2004 *J. Math. Phys.* **45** 2656–2673
- [69] Hendriks P and Singer M 1999 *J. Symbolic Comput.* **27** 239–259
- [70] Abramov S, Bronstein M, Petkovšek M and Schneider C 2021 *J. Symb. Comput.* **107** 23–66
- [71] Ablinger J and Schneider C 2021 Solving linear difference equations with coefficients in rings with idempotent representations *Proceedings of the 2021 International Symposium on Symbolic and Algebraic Computation (Proc. ISSAC 21)* ed Mezzarobba M pp 27–34
- [72] Karr M 1981 *J. ACM* **28** 305–350
- [73] Bronstein M 2000 *J. Symbolic Comput.* **29** 841–877
- [74] Schneider C 2004 *An. Univ. Timișoara Ser. Mat.-Inform.* **42** 163–179 extended version of Proc. SYNASC’04
- [75] Schneider C 2005 *J. Differ. Equations Appl.* **11** 799–821
- [76] Schneider C 2005 *Appl. Algebra Engrg. Comm. Comput.* **16** 1–32
- [77] Schneider C 2007 *Sém. Lothar. Combin.* **56** 1–36 article B56b
- [78] Schneider C 2014 Modern summation methods for loop integrals in quantum field theory: The packages Sigma, EvaluateMultiSums and SumProduction *Proc. ACAT 2013 (J. Phys.: Conf. Ser. vol 523)* pp 1–17
- [79] Schneider C 2021 Term Algebras, Canonical Representations and Difference Ring Theory for Symbolic Summation *Anti-Differentiation and the Calculation of Feynman Amplitudes* ed Blümlein J and Schneider C (Cham: Springer International Publishing) pp 423–485
- [80] Blümlein J, Kauers M, Klein S and Schneider C 2009 *Comput. Phys. Commun.* **180** 2143–2165 (*Preprint* 0902.4091)
- [81] Blümlein J, Klein S, Schneider C and Stan F 2012 *J. Symb. Comput.* **47** 1267–1289 (*Preprint* 1011.2656)
- [82] Nielsen N 1906 *Die Gammafunktion* (Leipzig: Teubner)
- [83] Landau E 1906 *Sitzungsber. Kgl. Bayer. Akademie der Wissenschaften* **XXXVI** 151–221
- [84] Nørlund N E 1924 *Vorlesungen über Differenzenrechnung* (Berlin: Springer)
- [85] Milne-Thomson L 1932 *Math. Proc. of the Cambridge Phil. Soc.* **28** 311–318
- [86] Laporta S 2001 *Phys. Lett. B* **504** 188–194 (*Preprint* hep-ph/0102032)
- [87] Jiménez-Pastor A and Petkovšek M 2022 Definite Sums as Solutions of Linear Recurrences With

- Polynomial Coefficients (*Preprint 2202.05550*)
- [88] Kauers M and Paule P 2011 *The Concrete Tetrahedron* 1st ed Text and Monographs in Symbolic Computation (Springer Wien)
- [89] Bronstein M 1992 Linear ordinary differential equations: Breaking through the order 2 barrier *Papers from the International Symposium on Symbolic and Algebraic Computation ISSAC '92* (New York, NY, USA: Association for Computing Machinery) pp 42–48
- [90] Johansson F, Kauers M and Mezzarobba M 2013 Finding hyperexponential solutions of linear odes by numerical evaluation *Proceedings of the 38th International Symposium on Symbolic and Algebraic Computation ISSAC '13* (New York, NY, USA: Association for Computing Machinery) pp 211–218
- [91] Remiddi E and Vermaseren J 2000 *Int. J. Mod. Phys. A* **15** 725–754
- [92] Ablinger J 2017 Computing the inverse Mellin transform of holonomic sequences using Kovacic's algorithm *PoS (RADCOR2017) 069* ed Hoang A and Schneider C pp 1–8
- [93] Singer M 1981 *Amer. J. Math.* **103** 661–682
- [94] Kovacic J J 1986 *J. Symb. Computation* **2** 3–43
- [95] Singer M F 1991 *J. Symb. Computation* **11** 251–273
- [96] Ablinger J 2021 Extensions of the AZ-algorithm and the Package MultiIntegrate *Anti-Differentiation and the Calculation of Feynman Amplitudes* ed Blümlein J and Schneider C (Cham: Springer International Publishing) pp 35–61
- [97] Imamoglu E and van Hoeij M 2017 *J. Symbolic Comput.* **83** 245–271
- [98] Ablinger J, Blümlein J, De Freitas A, van Hoeij M, Imamoglu E, Raab C G, Radu C S and Schneider C 2018 *J. Math. Phys.* **59** 062305 (*Preprint 1706.01299*)
- [99] Blümlein J, De Freitas A, Van Hoeij M, Imamoglu E, Marquard P and Schneider C 2018 *PoS LL2018* 017 (*Preprint 1807.05287*)
- [100] Abramov S A and Petkovsek M 2012 On polynomial solutions of linear partial differential and (q-)difference equations *Computer Algebra in Scientific Computing - 14th International Workshop, CASC 2012, Maribor, Slovenia, September 3-6, 2012. Proceedings (Lecture Notes in Computer Science vol 7442)* ed Gerdt V P, Koepf W, Mayr E W and Vorozhtsov E V (Springer) pp 1–11
- [101] Matiyasevich Y V 1993 *Hilbert's tenth problem. With a foreword by Martin Davis* (Cambridge, MA: MIT Press)
- [102] Kauers M and Schneider C 2010 Partial denominator bounds for partial linear difference equations *Proc. ISSAC'10* ed Watt S pp 211–218
- [103] Kauers M and Schneider C 2011 A refined denominator bounding algorithm for multivariate linear difference equations *Proc. ISSAC 2011* ed Leykin A pp 201–208
- [104] Blümlein J, Saragnese M and Schneider C 2023 Hypergeometric Structures in Feynman Integrals *Annals of Mathematics and Artificial Intelligence, in press* (*Preprint 2111.15501*)
- [105] Chetyrkin K and Tkachov F 1981 *Nucl. Phys. B* **192** 159–204
- [106] Laporta S 2000 *Int. J. Mod. Phys. A* **15** 5087–5159
- [107] Marquard P 2021 Integration-by-parts: A survey *Anti-Differentiation and the Calculation of Feynman Amplitudes* ed Blümlein J and Schneider C (Cham: Springer International Publishing) pp 313–320
- [108] Hilbert D 1890 *Math. Ann.* **36** 473–534
- [109] Gluza J, Kajda K and Kosower D A 2011 *Phys. Rev. D* **83** 045012 (*Preprint 1009.0472*)
- [110] Vermaseren J A M 2021 Some steps towards improving ibp calculations and related topics *Anti-Differentiation and the Calculation of Feynman Amplitudes* ed Blümlein J and Schneider C (Cham: Springer International Publishing) pp 501–518
- [111] Kotikov A V 2021 Differential equations and feynman integrals *Anti-Differentiation and the Calculation of Feynman Amplitudes* ed Blümlein J and Schneider C (Cham: Springer International Publishing) pp 235–259
- [112] Dreyfus T and Weil J A 2021 Differential galois theory and integration *Anti-Differentiation and*

- the Calculation of Feynman Amplitudes* ed Blümlein J and Schneider C (Cham: Springer International Publishing) pp 145–171
- [113] Zürcher B 1994 *Rationale Normalformen von pseudo-linearen Abbildungen* Ph.D. thesis Mathematik, ETH Zürich
- [114] Bostan A, Chyzak F and de Panafieu É 2013 Complexity estimates for two uncoupling algorithms *Proc. ISSAC'13* (Boston)
- [115] Gerhold S 2002 *Uncoupling Systems of Linear Ore Operator Equations* Master's thesis RISC, J. Kepler University Linz
- [116] Ablinger J, Behring A, Blümlein J, Falcioni G, De Freitas A, Marquard P, Rana N and Schneider C 2018 *Phys. Rev. D* **97** 094022 (*Preprint* 1712.09889)
- [117] Ablinger J, Blümlein J, Marquard P, Rana N and Schneider C 2019 *Nucl. Phys. B* **939** 253–291 (*Preprint* 1810.12261)
- [118] Ablinger J, Behring A, Blümlein J, De Freitas A, von Manteuffel A and Schneider C 2016 *Comput. Phys. Commun.* **202** 33–112 (*Preprint* 1509.08324)
- [119] Behring A, Blümlein J, De Freitas A, von Manteuffel A and Schneider C 2015 *Nucl. Phys. B* **897** 612–644 (*Preprint* 1504.08217)
- [120] Behring A, Blümlein J, De Freitas A, Hasselhuhn A, von Manteuffel A and Schneider C 2015 *Phys. Rev. D* **92** 114005 (*Preprint* 1508.01449)
- [121] Behring A, Blümlein J, Falcioni G, De Freitas A, von Manteuffel A and Schneider C 2016 *Phys. Rev. D* **94** 114006 (*Preprint* 1609.06255)
- [122] Blümlein J, Hasselhuhn A, Klein S and Schneider C 2013 *Nucl. Phys. B* **866** 196–211 (*Preprint* 1205.4184)
- [123] Ablinger J, Blümlein J, De Freitas A, Hasselhuhn A, von Manteuffel A, Round M, Schneider C and Wißbrock F 2014 *Nucl. Phys. B* **882** 263–288 (*Preprint* 1402.0359)
- [124] Barkatou M A and Raab C G 2012 Solving linear ordinary differential systems in hyperexponential extensions *Proceedings of the 37th International Symposium on Symbolic and Algebraic Computation ISSAC '12* (New York, NY, USA: Association for Computing Machinery) pp 51–58
- [125] Middeke J and Schneider C 2018 Denominator bounds for systems of recurrence equations using $\Pi\Sigma$ -extensions *Advances in Computer Algebra. WWCA 2016. (Springer Proceedings in Mathematics & Statistics* vol 226) ed Schneider C and Zima E (Springer) pp 149–173
- [126] Henn J 2013 *Phys. Rev. Lett.* **110** 251601
- [127] Lee R N 2015 *JHEP* **04** 108 (*Preprint* 1411.0911)
- [128] Gituliar O and Magerya V 2017 *Comput. Phys. Commun.* **219** 329–338 (*Preprint* 1701.04269)
- [129] Prausa M 2017 *Comput. Phys. Commun.* **219** 361–376 (*Preprint* 1701.00725)
- [130] Meyer C 2018 *Comput. Phys. Commun.* **222** 295–312
- [131] Abramov S and Petkovsek M 2002 *Adv. Appl. Math.* **29** 386–411
- [132] Hamberg R 1991 *Second order gluonic contributions to physical quantities* Ph.D. thesis University of Leiden
- [133] Klein F 1933 *Vorlesungen über die hypergeometrische Funktion* (Berlin: Springer)
- [134] Bailey W N 1935 *Generalized Hypergeometric Series* (Cambridge: Cambridge University Press)
- [135] Slater L J 1966 *Generalized hypergeometric functions* (Cambridge: Cambridge University Press)
- [136] Andrews G, Askey R and Roy R 1999 *Special functions (Encyclopedia of Mathematics and its Applications* vol 71) (Cambridge University Press, Cambridge)
- [137] Whittaker E T and Watson G N 1927 *A Course of Modern Analysis* (Cambridge: Cambridge University Press)
- [138] Kratzer A and Franz W 1963 *Transzendente Funktionen* (Leipzig: Akademische Verlagsgesellschaft Geest & Portig)
- [139] Paule P 2021 Contiguous relations and creative telescoping *Anti-Differentiation and the Calculation of Feynman Amplitudes* ed Blümlein J and Schneider C (Cham: Springer International Publishing) pp 335–394

- [140] Mellin H 1910 *Math. Ann* **68** 305–337
- [141] Barnes E W 1908 *Proc. Lond. Math. Soc.* **6** 14–177
- [142] Barnes E W 1910 *Quarterly Journal of Mathematics* **41** 136–140
- [143] Appell P and Kampé de Fériet J 1926 *Fonctions hypergéométriques et hypersphériques. Polynômes d’Hermite*. Paris: Gauthier-Villars. vii, 434 p. (1926).
- [144] Appell P 1925 *Sur les fonctions hypergéométriques de plusieurs variables, les polynômes d’Hermite et autres fonctions sphériques dans l’hyperespace*. IV + 76 p. Paris, Gauthier-Villars (Mémorial des sciences mathématiques, fasc. 3) (1925).
- [145] Kampé de Fériet J 1937 *La fonction hypergéométrique* Mém. Sci. Math. 85. Paris: Gauthier-Villars. 87 p. (1937).
- [146] Exton H 1976 *Multiple hypergeometric functions and applications* Mathematics and Its Applications. Chichester: Ellis Horwood Limited, Publisher; New York etc.: Halsted Press, a division of John Wiley & Sons, Inc. 312 p. (1976).
- [147] Exton H 1978 *Handbook of hypergeometric integrals. Theory, applications, tables, computer programs* Mathematics & its Applications. Chichester: Ellis Horwood Limited Publishers. 316 p. (1978).
- [148] Schlosser M J 2013 *Multiple hypergeometric series: Appell series and beyond* Computer algebra in quantum field theory. Integration, summation and special functions (Wien: Springer) pp 305–324
- [149] Srivastava H M and Karlsson P W 1985 *Multiple Gaussian hypergeometric series* Ellis Horwood Series in Mathematics and Its Applications. Chichester: Ellis Horwood Limited; New York etc.: Halsted Press: a division of John Wiley & Sons. 425 p. (1985).
- [150] Kalmykov M, Bytev V, Kniehl B A, Moch S O, Ward B F L and Yost S A 2021 Hypergeometric functions and feynman diagrams *Anti-Differentiation and the Calculation of Feynman Amplitudes* ed Blümlein J and Schneider C (Cham: Springer International Publishing) pp 189–234
- [151] Anastasiou C, Glover E W N and Oleari C 2000 *Nucl. Phys.* **B572** 307–360
- [152] Ablinger J, Blümlein J, Hasselhuhn A, Klein S, Schneider C and Wißbrock F 2012 *Nucl. Phys. B* **864** 52–84 (*Preprint* 1206.2252)
- [153] Bierenbaum I, Blümlein J and Klein S 2007 *Nucl. Phys. B* **780** 40–75 (*Preprint* hep-ph/0703285)
- [154] Czakon M 2006 *Comput. Phys. Commun.* **175** 559–571
- [155] Smirnov A and Smirnov V 2009 *Eur. Phys. J.* **C62** 445–449
- [156] Gluza J, Kajda K and Riemann T 2007 *Comput. Phys. Commun.* **177** 879–893
- [157] Ananthanarayan B, Banik S, Friot S and Ghosh S 2021 *Phys. Rev. Lett.* **127**(15) 151601 (*Preprint* hep-ph/2012.15108)
- [158] Gosper R W 1978 *Proc. Nat. Acad. Sci. U.S.A.* **75** 40–42
- [159] Schneider C 2001 Symbolic summation in difference fields Tech. Rep. 01-17 RISC-Linz, J. Kepler University PhD Thesis
- [160] Schneider C 2014 A streamlined difference ring theory: Indefinite nested sums, the alternating sign and the parameterized telescoping problem *Symbolic and Numeric Algorithms for Scientific Computing (SYNASC), 2014 15th International Symposium* ed Winkler F, Negru V, Ida T, Jebelean T, Petcu D, Watt S and Zaharie D (IEEE Computer Society) pp 26–33
- [161] Schneider C 2016 *J. Symb. Comput.* **72** 82–127
- [162] Ocansey E and Schneider C 2018 Representing (q-)hypergeometric products and mixed versions in difference rings *Advances in Computer Algebra. WWCA 2016. (Springer Proceedings in Mathematics & Statistics vol 226)* ed Schneider C and Zima E (Springer) pp 175–213
- [163] Ocansey E D and Schneider C 2020 Representation of hypergeometric products of higher nesting depths in difference rings RISC Report Series 20-19 Research Institute for Symbolic Computation (RISC), Johannes Kepler University Linz Schloss Hagenberg, 4232 Hagenberg, Austria arXiv:2011.08775
- [164] Schneider C 2007 *J. Algebra Appl.* **6** 415–441

- [165] Schneider C 2008 *J. Symbolic Comput.* **43** 611–644
- [166] Schneider C 2010 *Appl. Algebra Engrg. Comm. Comput.* **21** 1–32
- [167] Schneider C 2015 Fast algorithms for refined parameterized telescoping in difference fields
Computer Algebra and Polynomials (Lecture Notes in Computer Science (LNCS) no 8942)
ed Guitierrez J, Schicho J and Weimann M (Springer) pp 157–191
- [168] Schneider C 2017 *J. Symb. Comput.* **80** 616–664
- [169] Kauers M and Zimmermann B 2008 *Journal of Symbolic Computation* **43** 787–803
- [170] Hardouin C and Singer M 2008 *Math. Ann.* **342** 333–377
- [171] Schneider C 2010 *Ann. Comb.* **14** 533–552 appeared also in FPSAC’07
- [172] Hoffman M 2000 *J. Algebraic Combin.* **11** 49–68
- [173] Blümlein J 2004 *Comput. Phys. Commun.* **159** 19–54 (*Preprint hep-ph/0311046*)
- [174] Blümlein J 2009 *Comput. Phys. Commun.* **180** 2218–2249 (*Preprint 0901.3106*)
- [175] Blümlein J 2010 *Clay Math. Proc.* **12** 167–188 (*Preprint 0901.0837*)
- [176] Ablinger J and Schneider C 2018 *Annals of Combinatorics* **22** 213–244
- [177] Zeilberger D 1991 *J. Symbolic Comput.* **11** 195–204
- [178] Paule P and Schorn M 1995 *J. Symbolic Comput.* **20** 673–698
- [179] Paule P 1995 *J. Symbolic Comput.* **20** 235–268
- [180] Chen S and Kauers M 2012 Order-Degree Curves for Hypergeometric Creative Telescoping *Proc. ISSAC 2012* ed Hoeven J and van Hoeij M pp 122–129
- [181] Bostan A, Dumont L and Salvy B 2016 Efficient algorithms for mixed creative telescoping
ISSAC’16—Proceedings of the 2016 ACM International Symposium on Symbolic and Algebraic Computation (ACM Press) pp 127–134
- [182] Wilf H and Zeilberger D 1992 *Invent. Math.* **108** 575–633
- [183] Wegschaider K 1997 *Computer generated proofs of binomial multi-sum identities* Master’s thesis
RISC, J. Kepler University
- [184] Lyons R, Paule P and Riese A 2002 *Appl. Algebra Engrg. Comm. Comput.* **13** 327–333
- [185] Apagodu M and Zeilberger D 2006 *Adv. Appl. Math.* **37** 139–152
- [186] Schneider C 2004 *Discrete Math. Theor. Comput. Sci.* **6** 365–386
- [187] Ablinger J, Blümlein J, De Freitas A, Goedicke A, Schneider C and Schönwald K 2018 *Nucl. Phys. B* **932** 129–240 (*Preprint 1804.02226*)
- [188] Ablinger J, Blümlein J, De Freitas A, Goedicke A, Saragnese M, Schneider C and Schönwald K 2020 *Nucl. Phys. B* **955** 115059 (*Preprint 2004.08916*)
- [189] Ablinger J, Blümlein J, De Freitas A, Saragnese M, Schneider C and Schönwald K 2020 *Nucl. Phys. B* **952** 114916 (*Preprint 1911.11630*)
- [190] Zeilberger D 1990 *J. Comput. Appl. Math.* **32** 321–368
- [191] Chyzak F 2000 *Discrete Math.* **217** 115–134
- [192] Koutschan C 2013 Creative telescoping for holonomic functions *Computer Algebra in Quantum Field Theory* Texts and Monographs in Symbolic Computation ed Schneider C and Blümlein J (Springer) pp 171–194
- [193] Schneider C 2005 *Adv. in Appl. Math.* **34** 740–767
- [194] Ablinger J, Blümlein J, Round M and Schneider C 2012 *PoS LL2012* 050 (*Preprint 1210.1685*)
- [195] Blümlein J, Round M and Schneider C 2018 Refined holonomic summation algorithms in particle physics *Advances in Computer Algebra. WWCA 2016. (Springer Proceedings in Mathematics & Statistics vol 226)* ed Schneider C and Zima E (Springer) pp 51–91 (*Preprint 1706.03677*)
- [196] Andrews G, Paule P and Schneider C 2005 *Adv. in Appl. Math.* **34** 709–739
- [197] Stembridge J 1995 *Advances in Mathematics* **111** 227 – 243
- [198] Schneider C and Zudilin W 2021 A case study for $\zeta(4)$ *Transcendence in Algebra, Combinatorics, Geometry and Number Theory. TRANS 2019 (Proceedings in Mathematics & Statistics vol 373)* ed Bostan A and Raschel K (Springer) pp 421–435
- [199] Behring A *et al.* 2013 *PoS RADCOR2013* 058 (*Preprint 1312.0124*)
- [200] van Manteuffel A, Panzer E and Schabinger R 2015 *JHEP* **02** 120

- [201] Brown F 2009 *Commun. Math. Phys.* **287** 925–958
- [202] Kummer E E 1840 *J. Reine Angew. Math. (Crelle)* **21** 74–90, 193–225, 328–371
- [203] Poincaré H 1884 *Acta Math. (Crelle)* **4** 201–312
- [204] Lappo-Danilevsky J A 1953 *Mémoires sur la Théorie des Systèmes Différentielles Linéaires* (New York: Chelsea Pub. Co.)
- [205] Chen K 1971 *A.M.S.* **156** 359–379
- [206] Goncharov A 1998 *Math. Res. Lett.* **5** 497–516
- [207] Panzer E 2015 *Comput. Phys. Commun.* **188** 148–166
- [208] Ablinger J, Blümlein J, Raab C, Schneider C and Wißbrock F 2014 *Nucl. Phys. B* **885** 409–447 (Preprint 1706.03678)
- [209] Ablinger J 2012 *Computer algebra algorithms for special functions in particle physics* Ph.D. thesis J. Kepler University Linz
- [210] Risch R 1969 *Trans. Amer. Math. Soc.* **139** 167–189
- [211] Bronstein M 1997 *Symbolic Integration I, Transcendental functions* (Springer)
- [212] Singer M, Saunders B and Caviness B 1985 *SIAM J. Comput.* **14** 966–990 ISSN 0097-5397
- [213] Raab C 2012 *Definite integration in differential fields* Ph.D. thesis Johannes Kepler Universität Linz, Austria
- [214] Koutschan C 2021 Holonomic anti-differentiation and feynman amplitudes *Anti-Differentiation and the Calculation of Feynman Amplitudes* ed Blümlein J and Schneider C (Cham: Springer International Publishing) pp 261–277
- [215] Larin S, Tkachov F and Vermaseren J 1991 The FORM version of Mincer Tech. Rep. NIKHEF-H-91-18 NIKHEF, Netherlands
- [216] Steinhauser M 2001 *Comput. Phys. Commun.* **134** 335–364
- [217] Blümlein J and Schneider C 2017 *Phys. Lett.* **B771** 31–36 (Preprint 1706.03677)
- [218] Ablinger J, Schneider C, Behring A, Blümlein J and De Freitas A 2016 *PoS LL2016* 005 (Preprint 1608.05376)
- [219] Blümlein J, Marquard P and Schneider C 2019 *PoS RADCOR2019* 46 (Preprint 1912.04390)
- [220] Furmanski W and Petronzio R 1982 *Nucl. Phys. B* **195** 237–261
- [221] Wimp J and Zeilberger D 1985 *J. Math. Anal. Appl. v. 111, 162-177 (1985)* **111**
- [222] Kauers M 2011 A Mathematica package for computing asymptotic expansions of solutions of p-finite recurrence equations RISC Report Series 11-04 Johannes Kepler University Linz
- [223] Nielsen N 1909 *Nova Acta Leopold.* **XC** 125–211
- [224] Kölbig K S, Mignoco J A and Remiddi E 1970 *BIT* **10** 38–74
- [225] Kölbig K S 1986 *SIAM J. Math. Anal.* **17** 1232–1258
- [226] Lewin L 1958 *Dilogarithms and associated functions* (London: Macdonald)
- [227] Lewin L 1981 *Polylogarithms and associated functions* (New York: North Holland)
- [228] Devoto A and Duke D W 1984 *Riv. Nuovo Cim.* **7N6** 1–39
- [229] Ablinger J, Blümlein J, De Freitas A, Hasselhuhn A, von Manteuffel A, Round M and Schneider C 2014 *Nucl. Phys. B* **885** 280–317 (Preprint 1405.4259)
- [230] Ablinger J 2019 *Experimental Mathematics* 1–15
- [231] Ablinger J 2020 Proving Two Conjectural Series for $\zeta(7)$ and Discovering More Series for $\zeta(7)$ *Mathematical Aspects of Computer and Information Science* ed Slamanig D, Tsigaridas E and Zafeirakopoulos Z (Springer International Publishing) pp 42–47
- [232] Blümlein J 2016 Heavy Flavor Wilson Coefficients in Deep-Inelastic Scattering: Recent Results, The 5th International Congress on Mathematical Software ZIB Berlin from July 11 to July 14, 2016, Session: Symbolic computation and elementary particle physics, <https://www.risc.jku.at/conferences/ICMS2016/>, 2016 The 5th International Congress on Mathematical Software ZIB Berlin from July 11 to July 14, 2016, Session: Symbolic computation and elementary particle physics, <https://www.risc.jku.at/conferences/ICMS2016/>
- [233] Sabry A 1962 *Nucl. Phys. B* **33** 401–430
- [234] Blümlein, Johannes and Ablinger, Jakob and Behring, Arnd and De Freitas, Abilio and von

- Manteuffel, Andreas and Schneider, Carsten and Schneider, C 2017 *PoS QCDEV2017* 031 (Preprint 1711.07957)
- [235] Mistlberger B 2018 *JHEP* **05** 028 (Preprint 1802.00833)
- [236] Duhr C, Dulat F and Mistlberger B 2020 *Phys. Rev. Lett.* **125** 172001 (Preprint 2001.07717)
- [237] Broadhurst D, Fleischer J and Tarasov O 1993 *Z. Phys.* **C60** 287–302
- [238] Bloch S and Vanhove P 2015 *J. Number Theor.* **148** 328–364
- [239] Adams L, Bogner C and Weinzierl S 2016 *J. Math. Phys.* **57** 032304
- [240] Remiddi E and Tancredi L 2016 *Nucl. Phys. B* **907** 400–444
- [241] Adams L, Bogner C, Schweitzer A and Weinzierl S 2016 *J. Math. Phys.* **57** 122302
- [242] Blümlein J, Schneider C and Paule P (eds) 2019 *Proceedings, KMPB Conference: Elliptic Integrals, Elliptic Functions and Modular Forms in Quantum Field Theory: Zeuthen, Germany, October 23-26, 2017* (Wien: Springer)
- [243] Henn J, Mistlberger B, Smirnov V A and Wasser P 2020 *JHEP* **04** 167 (Preprint 2002.09492)
- [244] Bern Z, Parra-Martinez J, Roiban R, Ruf M S, Shen C H, Solon M P and Zeng M 2021 *Phys. Rev. Lett.* **126** 171601 (Preprint 2101.07254)
- [245] Bargiela P, Caola F, von Manteuffel A and Tancredi L 2022 *JHEP* **02** 153 (Preprint 2111.13595)
- [246] Neumann C 1884 *Vorlesungen über Riemann's Theorie der Abel'schen Integrale* (Leipzig: Teubner)
- [247] Brown F and Schnetz O 2012 *Duke Math. J.* **161** 1817–1862 (Preprint 1006.4064)
- [248] Kreimer D 2022 Bananas: multi-edge graphs and their Feynman integrals (Preprint 2202.05490)
- [249] Gehrmann T and Remiddi E 2001 *Comput. Phys. Commun.* **141** 296–312 (Preprint hep-ph/0107173)
- [250] Vollinga J and Weinzierl S 2005 *Comput. Phys. Commun.* **167** 177 (Preprint hep-ph/0410259)
- [251] Ablinger J, Blümlein J, Round M and Schneider C 2019 *Comput. Phys. Commun.* **240** 189–201 (Preprint 1809.07084)
- [252] Blümlein J 2000 *Comput. Phys. Commun.* **133** 76–104 (Preprint hep-ph/0003100)
- [253] Blümlein J and Moch S O 2005 *Phys. Lett. B* **614** 53–61 (Preprint hep-ph/0503188)
- [254] Kotikov A V and Velizhanin V N 2005 Analytic continuation of the Mellin moments of deep inelastic structure functions (Preprint hep-ph/0501274)
- [255] Lie S 1891 *Vorlesungen über Differentialgleichungen mit bekannten infinitesimalen Transformationen* (Leipzig: Teubner)
- [256] Schwarz F 2008 *Algorithmic Lie Theory for Solving Ordinary Differential Equations* (Boca Raton: Chapman & Hall/CRC)
- [257] Tricomi F 1948 *Elliptische Funktionen* (Leipzig: Akademische Verlagsgesellschaft Geest & Portig)
- [258] Cohen H and Strömberg F 2017 *Modular Forms: A Classical Approach* (Providence,RI: AMS)
- [259] Bierenbaum I, Blümlein J and Klein S 2009 *Nucl. Phys. B* **820** 417–482 (Preprint 0904.3563)
- [260] Moch S, Ruijl B, Ueda T, Vermaseren J A M and Vogt A 2022 *Phys. Lett. B* **825** 136853 (Preprint 2111.15561)
- [261] Laporta S 2017 *Phys. Lett. B* **772** 232–238 (Preprint 1704.06996)
- [262] Moch S, Vermaseren J A M and Vogt A 2014 *Nucl. Phys. B* **889** 351–400 (Preprint 1409.5131)
- [263] Blümlein, J 2013 *Prog. Part. Nucl. Phys.* **69** 28–84 (Preprint 1208.6087)
- [264] Hamberg R, van Neerven W L and Matsuura T 1991 *Nucl. Phys. B* **359** 343–405 [Erratum: Nucl.Phys.B 644, 403–404 (2002)]
- [265] Harlander R V and Kilgore W B 2002 *Phys. Rev. Lett.* **88** 201801 (Preprint hep-ph/0201206)
- [266] Ravindran V, Smith J and van Neerven W L 2003 *Nucl. Phys. B* **665** 325–366 (Preprint hep-ph/0302135)
- [267] Blümlein J and Ravindran V 2005 *Nucl. Phys. B* **716** 128–172 (Preprint hep-ph/0501178)
- [268] Buza M, Matiounine Y, Smith J, Migneron R and van Neerven W L 1996 *Nucl. Phys. B* **472** 611–658 (Preprint hep-ph/9601302)
- [269] Behring A, Bierenbaum I, Blümlein J, De Freitas A, Klein S and Wißbrock F 2014 *Eur. Phys. J. C* **74** 3033 (Preprint 1403.6356)

- [270] Blümlein J, De Freitas A and van Neerven W 2012 *Nucl. Phys. B* **855** 508–569 (*Preprint* 1107.4638)
- [271] Ablinger J, Blümlein J, De Freitas A and Schönwald K 2020 *Nucl. Phys. B* **955** 115045 (*Preprint* 2004.04287)
- [272] Blümlein J, De Freitas A and Schönwald K 2021 *Phys. Lett. B* **816** 136250 (*Preprint* 2102.12237)
- [273] Blümlein J, Falcioni G and De Freitas A 2016 *Nucl. Phys. B* **910** 568–617 (*Preprint* 1605.05541)
- [274] Blümlein J, De Freitas A, Raab C G and Schönwald K 2019 *Nucl. Phys. B* **945** 114659 (*Preprint* 1903.06155)
- [275] Blümlein J, Raab C and Schönwald K 2019 *Nucl. Phys. B* **948** 114736 (*Preprint* 1904.08911)
- [276] Buza M, Matiounine Y, Smith J and van Neerven W L 1997 *Nucl. Phys. B* **485** 420–456 (*Preprint* hep-ph/9608342)
- [277] Buza M, Matiounine Y, Smith J and van Neerven W L 1998 *Eur. Phys. J. C* **1** 301–320 (*Preprint* hep-ph/9612398)
- [278] Bierenbaum I, Blümlein J, Klein S and Schneider C 2008 *Nucl. Phys. B* **803** 1–41 (*Preprint* 0803.0273)
- [279] Bierenbaum I, Blümlein J and Klein S 2009 *Phys. Lett. B* **672** 401–406 (*Preprint* 0901.0669)
- [280] Ablinger J, Blümlein J, Klein S, Schneider C and Wißbrock F 2011 *Nucl. Phys. B* **844** 26–54 (*Preprint* 1008.3347)
- [281] Ablinger J, Behring A, Blümlein J, De Freitas A, Hasselhuhn A, von Manteuffel A, Round M, Schneider C and Wißbrock F 2014 *Nucl. Phys. B* **886** 733–823
- [282] Ablinger J, Behring A, Blümlein J, De Freitas A, von Manteuffel A, Schneider C and Schönwald K 2020 *Nucl. Phys. B* **953** 114945 (*Preprint* 1912.02536)
- [283] Blümlein J, Marquard P and Rana N 2019 *Phys. Rev. D* **99** 016013 (*Preprint* 1810.08943)
- [284] Lee R N, Smirnov A V, Smirnov V A and Steinhauser M 2018 *JHEP* **05** 187 (*Preprint* 1804.07310)
- [285] Lee R N, Smirnov A V, Smirnov V A and Steinhauser M 2018 *JHEP* **03** 136 (*Preprint* 1801.08151)
- [286] Berends F A, van Neerven W L and Burgers G J H 1988 *Nucl. Phys. B* **297** 429 [Erratum: *Nucl.Phys.B* 304, 921 (1988)]
- [287] Blümlein J, De Freitas A, Raab C and Schönwald K 2020 *Nucl. Phys. B* **956** 115055 (*Preprint* 2003.14289)
- [288] Blümlein J, De Freitas A, Raab C G and Schönwald K 2019 *Phys. Lett. B* **791** 206–209 (*Preprint* 1901.08018)
- [289] Blümlein J, De Freitas A, Raab C G and Schönwald K 2020 *Phys. Lett. B* **801** 135196 (*Preprint* 1910.05759)
- [290] Blümlein J and Schönwald K 2022 *Mod. Phys. Lett. A* **37** 2230004 (*Preprint* 2202.08476)
- [291] Blümlein J, Ablinger J, Behring A, De Freitas A, Imamoglu E, van Hoeij M, von Manteuffel A, Raab C G, Radu C S and Schneider C 2017 *PoS RADCOR2017* 069 (*Preprint* 1711.09742)
- [292] Besier M, Wasser P and Weinzierl S 2020 *Comput. Phys. Commun.* **253** 107197 (*Preprint* 1910.13251)
- [293] Ablinger J, Blümlein J, De Freitas A, Hasselhuhn A, Schneider C and Wißbrock F 2017 *Nucl. Phys. B* **921** 585–688 (*Preprint* 1705.07030)
- [294] Ablinger J, Blümlein J, De Freitas A, Schneider C and Schönwald K 2018 *Nucl. Phys. B* **927** 339–367 (*Preprint* 1711.06717)
- [295] Blümlein J, De Freitas A, Schneider C and Schönwald K 2018 *Phys. Lett. B* **782** 362–366 (*Preprint* 1804.03129)
- [296] Feynman R P and Hibbs A R 2005 *Quantum Mechanics and Path Integral* (New York: McGraw-Hill)
- [297] Nogueira P 1993 *J. Comput. Phys.* **105** 279–289
- [298] Vermaseren J A M 2000 New features of FORM (*Preprint* math-ph/0010025)
- [299] Tentyukov M and Vermaseren J A M 2010 *Comput. Phys. Commun.* **181** 1419–1427 (*Preprint*

- hep-ph/0702279)
- [300] Kol B and Smolkin M 2008 *Class. Quant. Grav.* **25** 145011 (*Preprint* 0712.4116)
 - [301] Jantzen B 2011 *JHEP* **12** 076 (*Preprint* 1111.2589)
 - [302] Blümlein J, Maier A, Marquard P and Schäfer G 2021 *Nucl. Phys. B* **965** 115352 (*Preprint* 2010.13672)
 - [303] Foffa S, Mastrolia P, Sturani R, Sturm C and Torres Bobadilla W J 2019 *Phys. Rev. Lett.* **122** 241605 (*Preprint* 1902.10571)
 - [304] Blümlein J, Maier A and Marquard P 2020 *Phys. Lett. B* **800** 135100 (*Preprint* 1902.11180)
 - [305] Blümlein J, Maier A, Marquard P and Schäfer G 2020 *Nucl. Phys. B* **955** 115041 (*Preprint* 2003.01692)
 - [306] Blümlein J, Maier A, Marquard P and Schäfer G 2021 *Phys. Lett. B* **816** 136260 (*Preprint* 2101.08630)
 - [307] Blümlein J, Maier A, Marquard P and Schäfer G 2021 The fifth-order post-Newtonian Hamiltonian dynamics of two-body systems from an effective field theory approach (*Preprint* 2110.13822)
 - [308] Bini D, Damour T and Geralico A 2020 *Phys. Rev. D* **102** 084047 (*Preprint* 2007.11239)
 - [309] Bern Z, Parra-Martinez J, Roiban R, Ruf M S, Shen C H, Solon M P and Zeng M 2021 Scattering Amplitudes, the Tail Effect, and Conservative Binary Dynamics at $O(G^4)$ (*Preprint* 2112.10750)
 - [310] Dlapa C, Kälin G, Liu Z and Porto R A 2021 Conservative Dynamics of Binary Systems at Fourth Post-Minkowskian Order in the Large-eccentricity Expansion (*Preprint* 2112.11296)
 - [311] Blümlein J, Maier A, Marquard P and Schäfer G 2020 *Phys. Lett. B* **807** 135496 (*Preprint* 2003.07145)
 - [312] Blümlein J, Maier A, Marquard P, Schäfer G and Schneider C 2020 *Phys. Lett. B* **801** 135157 (*Preprint* 1911.04411)
 - [313] Ablinger J, Blümlein J and Schneider C 2022 in preparation
 - [314] Vermaseren J 1994 *Comput. Phys. Commun.* **83** 45–58