ON LELONG NUMBERS OF GENERALIZED MONGE-AMPERE PRODUCTS `

MARTIN L. SERA

In memory of Jean-Pierre Demailly

ABSTRACT. We consider generalized (mixed) Monge-Ampère products of quasiplurisubharmonic functions (with and without analytic singularities) as they were introduced and studied in several articles written by subsets of M. Andersson, E. Wulcan, Z. Blocki, R. Lärkäng, H. Raufi, J. Ruppenthal, and the author. We continue these studies and present estimates for the Lelong numbers of pushforwards of such products by proper holomorphic submersions. Furthermore, we apply these estimates to Chern and Segre currents of pseudoeffective vector bundles. Among other corollaries, we obtain the following generalization of a recent result by X. Wu. If the non-nef locus of a pseudoeffective vector bundle E on a Kähler manifold is contained in a countable union of k -codimensional analytic sets, and if the k -power of the first Chern class of E is trivial, then E is nef.

1. Introduction

For a plurisubharmonic (psh) function u on a domain $D \subset \mathbb{C}^n$, the Lelong number in $x_0 \in D$ is defined by

$$
\nu(u, x_0) := \liminf_{x \to x_0} \frac{u(x)}{\log ||x - x_0||}.
$$

This can be seen as a generalization of the vanishing order (of e^u) since $\nu(\log|f|,0)$ = ord_{x₀} f for a holomorphic function $f: D \to \mathbb{C}$. Since its introduction in [\[Lel57,](#page-32-0) [Lel68\]](#page-32-1), Lelong numbers of psh functions and (its generalized version) of closed positive currents have proven to be very crucial tools in many areas beside of pluripotential theory. Many of its applications involve estimates on Lelong numbers. Let us recall a particular one which gives estimates on Lelong numbers of (mixed) Monge-Ampère products by the Lelong numbers of their potentials from below, see [\[Dem85,](#page-32-2) [Dem93\]](#page-32-3).

Generalizing a fundamental result from Bedford and Taylor in [\[BT76,](#page-32-4) [BT82\]](#page-32-5) defining the Monge-Ampère products of bounded psh functions (cited as [Theorem 2.6\)](#page-6-0), Demailly proved in [\[Dem85,](#page-32-2) [Dem93\]](#page-32-3) that for a closed positive current T of bidegree (p, p) and a plurisubharmonic function u such that its unbounded locus $L(u)$ is contained in an analytic set of codimension greater than or equal to $p + k$,

$$
(dd^c u)^k \wedge T := \lim_{\kappa \to \infty} (dd^c u_\kappa)^k \wedge T
$$

is a closed positive current whereby $d^c := \frac{i}{2\pi}(\overline{\partial} - \partial)$ and u_{κ} is a sequence of smooth psh functions decreasing pointwise to u (cited as [Theorem 2.8\)](#page-6-1). To shorten the notation, we will use BTD for the reference to such Monge-Ampère (MA) products.

Date: 29th June 2023.

²⁰¹⁰ Mathematics Subject Classification. 32W20, 32U25, 32U05, 32U40 (14C17).

With BTD, we can calculate the Lelong number of a closed positive (p, p) -current T by

$$
\nu(T, x_0) = \int \mathbb{1}_{x_0} (dd^c \log ||x - x_0||)^{n-p} \wedge T.
$$

We get that $\nu(dd^cu,x_0) = \nu(u,x_0)$. Demailly's second comparison formula for Lelong numbers (see [\[Dem85,](#page-32-2) [Dem93\]](#page-32-3)) implies

$$
(1.1) \qquad \nu((dd^c u)^k \wedge T, x_0) \ge \nu(u, x_0)^k \cdot \nu(T, x_0)
$$

for $k + p$ as above. The purpose of this work is to show that such estimates hold for more general Monge-Ampère products as explained in the following.

Let X and Y be complex manifolds, and let $\pi: X \to Y$ be a proper holomorphic submersion with m-dimensional fibres. Let q be a quasipsh function, let α be a closed real $(1, 1)$ -form, and let T be a closed positive (p, p) -current, all defined on X. Furthermore, we assume that for any small enough open $V \subset Y$, there is a closed positive¹ (1, 1)-form γ on $U := \pi^{-1}(V)$ such that q is $(\alpha + \gamma)$ -psh on U, i.e. $dd^c q + \alpha \geq -\gamma$. This is always the case if a priori $dd^c q + \alpha$ is assumed to be positive, or if π is Kähler, see [Remark 3.2.](#page-8-0) Let q_{κ} be a sequence of $(\alpha+\gamma)$ -psh functions which is decreasing pointwise to q. Following [\[LRRS18\]](#page-32-6), we obtain that for all k such that $\pi(L(q))$ is contained in an analytic set of codimension $\geq k+p-m$, the current

(1.2)
$$
\pi_*([dd^c q + \alpha]^k \wedge T) := \lim_{\kappa \to \infty} \pi_*((dd^c q_\kappa + \alpha)^k \wedge T)
$$

is well-defined and locally the difference of two closed positive currents which is independent of the choice of α and q_{κ} , see [Proposition 3.1.](#page-8-1) This generalizes the definition of MA products by BTD as the $L(q)$ is allowed to be in an analytic set of codimension strictly greater than $k + p$.

The first main result of the present work gives an estimate on the Lelong numbers of currents defined by (1.2) which generalizes (1.1) .

Theorem 1.1. Let $\pi: X \to Y$ be a proper holomorphic submersion between complex manifolds X and Y with m-dimensional fibres. Fix a point $y \in Y$. Let $\theta_1, \ldots, \theta_t$ be positive $(1,1)$ -currents on X such that each θ_i is in a Kähler class on a neighbourhood of $\pi^{-1}(y)$. Then, there exist positive constants δ_i , for $i = 1, \ldots, t$, (which only depend of the Kähler class represented by θ_i on a neighbourhood of $\pi^{-1}(y)$) such that the following statements are correct.

(i) If the union of all images $\pi(L(\theta_i))$ of the unbounded loci of local dd^c-potentials of θ_i is contained in an analytic A with codim $A \geq k_1 + \cdots + k_t - m$, then

(1.3)
$$
\nu\left(\pi_*\left([\theta_t]^{k_t} \wedge \cdots \wedge [\theta_1]^{k_1}\right), y\right) \geq \prod_{i=1}^t \min\{\nu(\theta_i, x), \delta_i\}^{k_i}
$$

for all points $x \in \pi^{-1}(y)$ as long the LHS does not vanish due to the degree of the current. (ii) If $\bigcup_i \pi(L(\theta_i))$ is contained in an analytic A with codim $A \geq k_1 + \cdots + k_t - t \cdot m$, then

(1.4)
$$
\nu\left(\pi_*\left([\theta_t]^{k_t}\right)\wedge\cdots\wedge\pi_*\left([\theta_1]^{k_1}\right),y\right)\geq \prod_{i=1}^t \min\{\nu(\theta_i,x),\delta_i\}^{k_i}
$$

for all points $x \in \pi^{-1}(y)$ as long the LHS does not vanish due to the degree of the current.

¹Following the notation usual for currents, we call a (1, 1)-form α positive if $\alpha \geq 0$.

Remark 1.2. (a) The bound δ_i is determined by whether for each $x \in \pi^{-1}(y)$, there exists a closed positive current which is in the same cohomology class as θ_i near $\pi^{-1}(y)$, smooth everywhere except in isolated points, and whose Lelong number is greater than or equal to δ_i in x; see [Lemma 6.1.](#page-24-0) For example, if π is projective and θ_i is in the class induced by the Fubini-Study metric, then we can select $\delta_i = 1$.

(b) If we consider proper holomorphic submersions $\pi_i: X_i \to Y$, $i = 1, ..., t$, and if each θ_i is defined on X_i with analogous properties as in [Theorem 1.1,](#page-1-2) there are positive constants δ_i such that

$$
(1.5) \qquad \nu\Big(\pi_{t,*}\big([\theta_t]^{k_t}\big) \wedge \cdots \wedge \pi_{1,*}\big([\theta_1]^{k_1}\big),y\Big) \geq \prod_{i=1}^t \min\{\nu(\theta_i,x_i),\delta_i\}^{k_i}
$$

for all points $x_1 \in \pi_1^{-1}(y), \ldots, x_t \in \pi_t^{-1}(y)$ and suitable k_1, \ldots, k_t .

Let us sketch the proof of [Theorem 1.1](#page-1-2) (i). We will use generalized Monge-Ampère products for currents with analytic singularities which were introduced and studied in $[And05, AW14, ASWY17, ABW19, LRSW22, B1019, LSW20]$ and the present work as explained as follows.

If a quasipsh function q equals locally $c \log ||F||^2 + b$ for a positive constant c, holomorphic tuple F and bounded function b, q is said to have analytic singularities in ${F = 0}$ (given as reduced analytic set). If moreover b is smooth, q has so-called neat analytic singularities. We say that a closed quasipos $(1, 1)$ -current θ has (neat) analytic singularities in Z if it is locally given by $\theta = dd^c q$ for a quasipsh q with (neat) analytic singularities in Z. For $i = 1, \ldots, t$, let θ_i be closed quasipos (1, 1)-currents with analytic singularities in Z_i , and let α_i be (closed) real (1, 1)-forms. Then, the *generalized Monge-Ampère* product $[\theta_t]_{\alpha_t}^{k_t} \wedge \cdots \wedge [\theta_1]_{\alpha_1}^{k_1}$ for any k_1, \ldots, k_t is defined recursively by

(1.6)
$$
[\theta_i]_{\alpha_i} \wedge T := \theta_i \wedge \mathbb{1}_{Z^c} T + \alpha_i \wedge \mathbb{1}_Z T.
$$

This definition works also if we replace α_i by quasipositive (1, 1)-currents η_i with neat analytic singularities in isolated points since the last term in (1.6) is defined by BTD, see [Definition 4.6.](#page-15-0) This will be useful to get estimates for the Lelong number of such MA products, see [Proposition 5.8](#page-22-0) and [Remark 5.9.](#page-22-1) Although the generalized Monge-Ampère product lacks some properties of a product, it is a suitable extension of the BTD Monge-Ampère product to arbitrary degrees independent of the unbounded locus of q . By [Theorem 4.7](#page-16-0) (cf. [\[LRSW22,](#page-32-7) Thm 1.2]), we get that if η_i are in the same $\partial \partial$ -class as θ_i , and if $\bigcup_i \pi(L(\theta_i))$ is contained in an analytic A with codim $A \geq k_1 + \cdots + k_t - m$, then

(1.7)
$$
\pi_*([\theta_t]^{k_t} \wedge \cdots \wedge [\theta_1]^{k_1}) = \pi_*([\theta_t]^{k_t}_{\eta_t} \wedge \cdots \wedge [\theta_1]^{k_1}_{\eta_1}).
$$

Therefore, the idea behind the proof of [Theorem 1.1](#page-1-2) is to use Demailly's regularization [\[Dem92,](#page-32-9) Main Thm 1.1] (see also [Theorem 2.10\)](#page-7-0) which gives us that the (quasi-) positive $(1, 1)$ -currents θ_i can be approximated by sequences of quasipositive $(1, 1)$ -currents θ_i _K with analytic singularities (and other suitable properties). By the monotone continuity result [Theorem 3.5,](#page-10-0) we get

(1.8)
$$
\pi_*([\theta_t]^{k_t} \wedge \cdots \wedge [\theta_1]^{k_1}) := \lim_{\kappa \to \infty} \pi_*([\theta_{t,\kappa}]^{k_t} \wedge \cdots \wedge [\theta_{1,\kappa}]^{k_1})
$$

(actually for any approximation). We would like to add that [Theorem 3.5](#page-10-0) implies Corollary 3.6 which asserts that for certain k_1, \ldots, k_t ,

$$
\pi_*[\theta_t]^{k_t} \wedge \cdots \wedge \pi_*[\theta_1]^{k_1} := \lim_{\kappa \to \infty} \pi_*[\theta_{t,\kappa}]^{k_t} \wedge \cdots \wedge \pi_*[\theta_{1,\kappa}]^{k_1}.
$$

We obtain that [Theorem 1.1](#page-1-2) is implied by (1.8) , (1.7) and the following theorem, which is the second main result of the present work.

Theorem 1.3. Let $\pi: X \to Y$ be a proper holomorphic submersion between complex manifolds X and Y with m-dimensional fibres. Fix a point $y \in Y$. Let $\theta_1, \ldots, \theta_t$ be positive $(1, 1)$ -currents with analytic singularities on X such that each θ_i is in a Kähler class on a neighbourhood of $\pi^{-1}(y)$. Then, there exist positive constants δ_i , $i = 1,...,t$, (which only depend of the Kähler class represented by θ_i on a neighbourhood of $\pi^{-1}(y)$) such that the following is correct.

(i) Existence of η_i : For every $x \in \pi^{-1}(y)$, there exist closed quasipos (1,1)-currents $\eta_i = \eta_{i,x} \in \{\theta_i\}_{\partial \overline{\partial}},$ positive near $\pi^{-1}(y)$, with neat analytic singularities only in $\{x\}$ (and with $\nu(\eta_i, x) = \delta_i$) such that

(1.9)
$$
\nu\Big(\pi_*\big([\theta_t]_{\eta_t}^{k_t} \wedge \cdots \wedge [\theta_1]_{\eta_1}^{k_1}\big), y\Big) \geq \prod_{i=1}^t \min\{\nu(\theta_i, x), \delta_i\}^{k_i}
$$

for all $m \leq k_1 + \ldots + k_t \leq \dim X$. Thereby, $\eta_i = \eta_{i,x}$ depends only of the Kähler class of θ_i near $\pi^{-1}(y)$ and x.

(ii) Independency of η_i : For $i = 1, \ldots, t$, let $\eta_i \in \{\theta_i\}_{\partial \overline{\partial}}$ be closed quasipos $(1, 1)$ -currents with neat analytic singularities in isolated points on X such that there is a point $x \in X$ with $L(\eta_i) \cap \pi^{-1}(y) = \{x\}$ and $\nu(\eta_i, x) \leq \delta_i$. Then,

$$
(1.10) \qquad \nu\Big(\pi_*\big([\theta_t]_{\eta_t}^{k_t} \wedge \cdots \wedge [\theta_1]_{\eta_1}^{k_1}\big),y\Big) \geq \prod_{i=1}^t \min\{\nu(\theta_i,x),\nu(\eta_i,x)\}^{k_i}
$$

for all $m \leq k_1 + \ldots + k_t \leq \dim X$.

[Remark 1.2](#page-2-3) (a) applies to [Theorem 1.3,](#page-3-0) as well.

The idea to consider η_i with singularities in isolated points is inspired by X. Wu's approach to choose specific non-smooth approximations of Segre currents to obtain Lelong number estimates of them, see [\[Wu22,](#page-32-10) Proof of Thm 2].

Following the arguments in the proof of [\[LRSW22,](#page-32-7) Thm 1.1 (3)], the Lelong numbers of $\pi_*(\theta_t]_{\alpha_t}^{k_t} \wedge \cdots \wedge \theta_1]_{\alpha_1}^{k_1}$ are independent of α_i among closed forms in the same class as θ_i . Unfortunately, this is not correct if we replace α_i by η_i with different kind of isolated singularities, see [Remark 5.12.](#page-23-0) By [Proposition 5.13,](#page-23-1) we get at least that $\pi_*(\theta_t]_{\eta_t}^{k_t} \wedge \cdots \wedge$ $[\theta_1]_{\eta_1}^{k_1}$'s Lelong numbers are independent of η_i among currents with the same kind of singularities in all points.

By applying [Theorem 1.3](#page-3-0) to Segre currents defined by [\(7.3\)](#page-29-0) as in [\[LRSW22,](#page-32-7) Thm 1.1], we obtain the following corollary; see [Remark/Def. 7.7.](#page-29-1)

Corollary 1.4. Let X be a complex manifold, let $E \to X$ be a (pseudoeffective) holomorphic vector bundle on X of rank r, and let $e^{-\varphi}$ be a semipositive singular metric on $\mathcal{O}_{\mathbb{P}(E)}(1)$ with analytic singularities, i.e. the weights φ are psh with analytic singularities. Then, for every $x \in X$, $\xi \in \pi^{-1}(x)$ and every $k \leq \dim X$, there exists a closed real (k, k) -current $S = S_{\xi}$ in the Segre class $(-1)^{k} s_{k}(E)$ such that (i) locally S is the difference of two closed

positive currents, (ii) S is positive close to x, and (iii) its Lelong number can be estimated from below by

$$
\nu(S, x) \ge \min{\{\nu(\varphi, \xi), 1\}}^{k+tr-t}.
$$

Analogously, [Theorem 1.1](#page-1-2) gives us estimates for the Lelong numbers of Segre currents defined by [Theorem 7.2,](#page-28-0) following [\[LRRS18,](#page-32-6) Prop. 4.6].

Corollary 1.5. Let X be a complex manifold of dimension n, let $E \to X$ be a (pseudoeffective) holomorphic vector bundle on X of rank r, and let $e^{-\varphi}$ be a semipositive singular metric on $\mathcal{O}_{\mathbb{P}(E)}(1)$ such that $\pi(L(\varphi))$ is contained in an analytic set of codimension s. Then, for all partitions $k_1 + \ldots + k_t = k \leq s$, all $x \in X$ and all $\xi \in \pi^{-1}(x)$, we get

 $(-1)^k \nu(s_{k_t}(E, \varphi) \wedge \cdots \wedge s_{k_1}(E, \varphi), x) \ge \min{\{\nu(\varphi, \xi), 1\}}^{k+tr-t}.$

Furthermore, [Corollary 1.5](#page-4-0) implies the following sufficient condition for E being nef.

Corollary 1.6. Let X be a compact complex manifold of dimension n, let $E \rightarrow X$ be a (pseudoeffective) holomorphic vector bundle on X, and let $e^{-\varphi}$ be a semipositive singular metric on $\mathcal{O}_{\mathbb{P}(E)}(1)$ such that $\pi(L(\varphi))$ is contained in an analytic set of codimension s. If there are k_1, \ldots, k_t with $k_1 + \ldots + k_t \leq s$ and $s_{k_1}(E) \cdots s_{k_t}(E) = 0$, then E is nef.

Let us point out that X does not need to be Kähler in [Corollary 1.6.](#page-4-1) Yet, the condition on $\pi(L(\varphi))$, which gives us the existence of the positive Segre currents, can be seen as quite strong. Assuming that X is Kähler, we can weaken this condition by using the non-nef locus which is defined by

$$
L_{\text{nnf}}(E) := \pi \big(L_{\text{nnf}}(\mathcal{O}_{\mathbb{P}(E)}(1)) \big) = \pi \big(\bigcup_{\delta > 0} \bigcap_{\theta} E_{+}(\theta) \big)
$$

where the intersection runs over all closed $\delta\omega_{\mathbb{P}(E)}$ -positive $(1,1)$ -currents $\theta \in {\{\omega_{\mathbb{P}(E)}\}}$ = $\mathcal{O}_{\mathbb{P}(E)}(1)$ for a Kähler form $\omega_{\mathbb{P}(E)}$ on $\mathbb{P}(E)$, and $E_+(\theta) := \{\xi \in \mathbb{P}(E) : \nu(\theta,\xi) > 0\}$. We obtain the following generalization of X. Wu's main result in [\[Wu22\]](#page-32-10).

Theorem 1.7. Let (X, ω) be a compact Kähler manifold, and let E be a pseudoeffective vector bundle on X such that $L_{nnf}(E)$ is contained in a countable union of analytic sets of codimension s. If there is a $k \leq s$ with $c_1(E)^k = (-s_1(E))^k = 0$, then E is nef.

If X is strongly pseudoeffective, i.e. $L_{\text{nnf}}(E) \neq X$ (see [\[BDPP13,](#page-31-5) Def. 7.1]), then the assumption on $L_{nnf}(E)$ is satisfied for $k = 1$. Hence, we obtain X. Wu's original result that all strongly pseudoeffective vector bundles with trivial first Chern class are nef. To prove [Theorem 1.7,](#page-4-2) we will follow X. Wu's argumentation. Yet, instead of the analytical construction of closed positive currents tailored to prove his result (we will see that these coincide with the one defined above, see [Remark 7.4\)](#page-28-1), we will use [Theorem 1.1](#page-1-2) which is based on the calculus of generalized Monge-Ampère products, see Sections [4](#page-13-0) and [5.](#page-18-0)

The present work is organized as follows. In [Section 2,](#page-5-0) we recall and elaborate some basic properties which are later used. We continue with the study of currents as they are defined by [\(1.2\)](#page-1-0) in [Section 3.](#page-8-2) In particular, we prove the mentioned monotone continuity results [Theorem 3.5](#page-10-0) and [Corollary 3.6](#page-12-0) there. In [Section 4,](#page-13-0) we introduce so-called currents with analytic singularities and study their properties. We continue with a section about

6 M. SERA

Lelong numbers of closed real currents, see [Section 5.](#page-18-0) [Section 6](#page-24-1) is dedicated to prove the main results, Theorems [1.1](#page-1-2) and [1.3.](#page-3-0) We conclude the article with a short discussion on Segre currents of (pseudoeffective) vector bundles and proving [Corollary 1.6](#page-4-1) and [Theorem 1.7](#page-4-2) in [Section 7.](#page-27-0)

Acknowledgements. The author is very grateful to Elizabeth Wulcan and the unknown referee for their valuable comments and suggestions which helped to improve the article a lot. The author was supported by JSPS KAKENHI Grant Number JP20K14319.

2. Preliminaries

In this section, we will introduce some notations and collect results from the literature. Considering the wedge product of pushforwards, the following tool is very helpful.

Lemma 2.1. (Lem. 6.3 in [\[LRSW22\]](#page-32-7)) Let Y be a complex manifold. For $i = 1,...,t$, let X_i be complex manifolds, let $\pi_i: X_i \to Y$ be proper holomorphic submersions, and let α_i be forms on X_i . Let $\tilde{X} := X_t \times_Y \cdots \times_Y X_1 \stackrel{\varpi}{\longrightarrow} Y$ be the fibre product of $\pi_t, ..., \pi_1$, let $\text{pr}_i \colon \tilde{X} \to X_i$ denote its projections on the ith component, and let T be a current on X_1 . Then,

 $\pi_{t,*}\alpha_t \wedge \cdots \wedge \pi_{2,*}\alpha_2 \wedge \pi_{1,*}T = \varpi_*(\mathrm{pr}_t^*\alpha_t \wedge \cdots \wedge \mathrm{pr}_2^*\alpha_2 \wedge \mathrm{pr}_1^*T).$

2.1. Closed positive currents. In this subsection, we recall some basic facts about closed positive currents. Throughout it, let X denote a complex manifold of dimension n . In [\[LRRS18,](#page-32-6) Def. 4.2], we introduced the following notion.

Definition 2.2. A strongly positive (k, k) -form β is called bump form at a point $x \in X$ if for a (or equivalently for any) Kähler form ω defined near x, there exists a constant $\delta > 0$ such that $\beta - \delta \omega^k$ is a strongly positive form in a neighbourhood of x.

Remark 2.3 (Lem. 7.2 in [\[LRSW22\]](#page-32-7)). As presented in the proof of [\[LRRS18,](#page-32-6) Lem. 4.3], for every analytic A with dim $A \leq k$ and every point $x \in A$, there exists a (k, k) -bump form β with arbitrarily small support s.t. the support of $dd^c\beta$ is in the complement of A.

The notion of bump form turns out to be very useful to obtain the uniqueness of the extensions of closed positive currents across analytic sets as follows.

Lemma 2.4 (Lem. 4.5 in [\[LRRS18\]](#page-32-6)). Let T and S be two closed positive (p, p) -currents on X such that $T = S$ on $X \setminus A$ for an analytic set A with codim $A \geq p$. If for every point $x \in A$, there is an $(n-p, n-p)$ -bump form β at x with arbitrarily small support such that $\int T \wedge \beta = \int S \wedge \beta$, then $T = S$ on X.

Proposition 2.5 (cf. Rem. 4.7 in [\[LRRS18\]](#page-32-6)). Let A be an analytic subset of X with codim A $\geq p$, and let T_{κ} be a sequence of closed positive (p, p) -currents on X. We assume (i) the sequence T_{κ} converges weakly to a closed positive current T' on $X \setminus A$, and (ii) $\int T_{\kappa} \wedge \beta$ converges to numbers T_β for any $(n-p, n-p)$ -bump form β with arbitrarily small support as $\kappa \to \infty$. Then, T_{κ} converges weakly to a closed positive (p, p) -current T on X which is uniquely defined by T' and $T_\beta = \int T \wedge \beta$.

Proof. We follow the argumentation of the proof of Prop. 4.6 in [\[LRRS18\]](#page-32-6).

Due to the assumptions (i) and (ii), we obtain that the trace measures of T_{κ} are (locally) uniformly bounded in κ . Therefore, the Banach-Alaoglu theorem implies that there is a subsequence $\kappa_{\lambda} \to \infty$ such that $T_{\kappa_{\lambda}}$ converges weakly to a closed positive current T. Let us assume that there is another subsequence $T_{\kappa'_\lambda}$ which does not converge to T. By passing to a subsequence, we may assume that $T_{\kappa'_\lambda}$ converges to a closed positive current S different from T. By the assumption (i), we get that S equals T on $X \setminus A$. Furthermore, the assumption (ii) implies that $\int T \wedge \beta = T_{\beta} = \int S \wedge \beta$ for any $(n-p, n-p)$ -bump form with arbitrarily small support. By [Lemma 2.4,](#page-5-1) we get $T = S$ on X. This is a contradiction. \Box

Next we recall the fundamental result of Bedford-Taylor defining MA products for bounded psh functions and its generalization by Demailly.

Theorem 2.6 (cf. [\[BT76,](#page-32-4)[BT82\]](#page-32-5)). Let u_1, \ldots, u_t be bounded psh functions, and let T be a closed positive (p, p) -current on T. Then, there exists a well-defined closed positive $(t + p, t + p)$ current

$$
dd^c u_t \wedge \cdots \wedge dd^c u_1 \wedge T,
$$

locally given as the limit of $dd^c u_{t,\kappa} \wedge \cdots \wedge dd^c u_{1,\kappa} \wedge T$ for sequences of smooth psh functions $u_{i,\kappa}$ decreasing pointwise to u_i . Furthermore, we obtain that this is monotone continuous: Let $v_{i,k}$ be sequences of (not necessarily smooth) psh functions decreasing pointwise to u_i . Then,

$$
u_t dd^c u_{t-1} \wedge \cdots \wedge dd^c u_1 \wedge T = \lim_{\kappa \to \infty} v_t dd^c v_{t-1,\kappa} \wedge \cdots \wedge dd^c v_{1,\kappa} \wedge T, \quad and
$$

$$
dd^c u_t \wedge \cdots \wedge dd^c u_1 \wedge T = \lim_{\kappa \to \infty} dd^c v_{t,\kappa} \wedge \cdots \wedge dd^c v_{1,\kappa} \wedge T.
$$

Definition 2.7. For a psh function u on X, the unbounded locus $L(u)$ is defined as the closed set of all points $x \in X$ such that u is unbounded near x. In general, $L(u)$ is strictly larger than the pole set of u.

Theorem 2.8 (cf. [\[Dem85,](#page-32-2) [Dem93\]](#page-32-3); shortly BTD). Let u be a psh function, and let T be a closed positive (p, p) -current. If $L(u) \cap \text{supp } T$ is contained in an analytic A with codim $A \geq$ $k+p$, then there exists a well-defined closed positive $(k+p, k+p)$ -current $(dd^c u)^k\wedge T$ (locally) given as the limit of

$$
(dd^c u_\kappa)^k \wedge T
$$

for any sequence of smooth psh functions u_{κ} decreasing pointwise to u. Furthermore, we obtain that this is monotone continuous: Let u_1, \ldots, u_t be psh functions with $L(u_i) \cap \text{supp } T$ is contained in an analytic A with codim $A \geq t+p$, and $v_{i,\kappa}$ be sequences of (not necessarily smooth) psh functions decreasing pointwise to u_i . Then,

$$
dd^{c}u_{t}\wedge\cdots\wedge dd^{c}u_{1}\wedge T=\lim_{\kappa\to\infty}dd^{c}v_{t,\kappa}\wedge\cdots\wedge dd^{c}v_{1,\kappa}\wedge T.
$$

Considering the mixed Monge-Ampère product (with different potentials in each factor), the codimension condition can be weaken by considering the (Hausdorff) dimension of the intersections of the unbounded loci, see Thm 4.5 and Prop. 4.9 in [\[Dem12,](#page-32-11) Chp. III].

2.2. Closed quasipositive currents. Let X be a complex manifold of dimension n , and let γ be a (positive) (1, 1)-form on X. We call a function q on X γ -psh if $dd^c q + \gamma$ is positive. This is equivalent to q is quasipsh with $dd^c q + \gamma \geq 0$ in the sense of currents. Let θ be a closed quasipositive (1, 1)-current on X, i.e. locally, $\theta = dd^c q$ for a quasipsh function q (shortly quasipos; also called almost positive). θ is called γ -positive (γ -pos) if $\theta + \gamma > 0$ in the sense of currents, i.e. locally, $\theta = dd^c q$ for a γ -psh function q. Let us pick a closed real (1, 1)-form α which is in the same $\partial \overline{\partial}$ -cohomology class as the γ -pos θ , i.e. $\theta - \alpha = dd^c q$ for a $(\gamma + \alpha)$ -psh function q on X. We define $L(\theta) := L(q)$ which is independent of the choice of q and α . Let q_{κ} be a sequence of (smooth) $(\gamma + \alpha)$ -psh functions which is decreasing pointwise to q. Set $\theta_{\kappa} := dd^c q_{\kappa} + \alpha$ which are (smooth) γ -positive currents. Obviously, the sequence θ_{κ} converges weakly to θ . Let us use the following notation improperly by saying that the sequence θ_{κ} is *decreasing pointwise to* θ *.* This is motivated by the following. If u is a psh functions on a small enough open set with $dd^c u = \theta + \gamma \geq 0$ (assuming that γ is closed), then $u_{\kappa} := q_{\kappa} - q + u$ are (smooth) psh functions with $dd^c u_{\kappa} = \theta_{\kappa} + \gamma \geq 0$ such that the sequence u_{κ} is decreasing pointwise to u.

Locally, we may assume that γ is closed and positive: On any small enough open set, there is a Kähler form ω and a constant C such that $\gamma \leq C \omega$. In particular, $\theta + C \omega \geq$ $\theta + \gamma \geq 0$ (and $\theta_{\kappa} + C\omega \geq 0$). Therefore, we may replace γ by $C\omega$ on any small enough open set.

Let T be a closed positive (p, p) -current, and assume that θ_{κ} are smooth. By BTD, for all k such that $L(\theta)$ is contained in an analytic set of codimension $\geq k+p$, (assuming γ is closed, see above)

$$
(\theta + \gamma)^k \wedge T \stackrel{\text{def}}{=} \lim_{\kappa \to \infty} (\theta_\kappa + \gamma)^k \wedge T \stackrel{\text{loc}}{=} \lim_{\kappa \to \infty} (dd^c u_\kappa)^k \wedge T
$$

is a closed positive current which is independent of the choice of q_{κ} / θ_{κ} / u_{κ} . We can extend this definition to (wedge) powers of the quasipositive θ by

(2.1)
$$
\theta^k \wedge T = (\theta + \gamma - \gamma)^k \wedge T := \sum_{i=0}^k (-1)^i {k \choose i} \gamma^i \wedge (\theta + \gamma)^{k-i} \wedge T
$$

$$
\stackrel{\text{def}}{=} \lim_{\kappa \to \infty} \sum_{i=0}^k (-1)^i {k \choose i} \gamma^i \wedge (\theta_{\kappa} + \gamma)^{k-i} \wedge T = \lim_{\kappa \to \infty} \theta_{\kappa}^k \wedge T.
$$

Due to the last equation, we see that the definition is independent of the choice of γ . $\theta^k \wedge T$ is a closed real $(k+p, k+p)$ -current which is locally the difference of two closed positive currents.

At the end of this section, we recall following versions of Demailly's regularization of quasipos (1, 1)-currents fitting to our applications.

Theorem 2.9 ([\[Dem82\]](#page-32-12), Main Thm 1.1 in [\[Dem92\]](#page-32-9)). Let X be a compact complex manifold, and fix a Hermitian form ω and a real $(1,1)$ -form γ on X. Let θ be a γ -positive $(1,1)$ current on X. Then, there are a constant C only dependent of (X,ω) , continuous functions λ_{κ} and $(\gamma+C\lambda_{\kappa}\omega)$ -pos $(1,1)$ -forms $\theta_{\kappa}\in\{\theta\}_{\partial\overline{\partial}}$ such that as $\kappa\to\infty$, (i) the sequence θ_{κ} is decreasing pointwise to θ , and (ii) $\lambda_{\kappa}(x) \searrow \nu(\theta, x)$ for all $x \in X$.

Theorem 2.10 (Main Thm 1.1 in [\[Dem92\]](#page-32-9)). Let X be a complex manifold, and fix a Hermitian form ω and a real (1, 1)-form γ on X. Let θ be a γ -positive (1, 1)-current on X.

Then, for every relatively compact $U \\\in X$, there are positive constants ε_{κ} and $(\gamma + \varepsilon_{\kappa} \omega)$ -pos $(1, 1)$ -currents $\theta_{\kappa} \in {\theta}_{\partial \overline{\partial}}$ with analytic singularities on U such that as $\kappa \to \infty$, (i) the sequence θ_{κ} is decreasing pointwise to θ , (ii) $\nu(\theta_{\kappa}, x) \nearrow \nu(\theta, x)$ for all $x \in U$ (uniformly), and (iii) $\varepsilon_{\kappa} \to 0$.

In its original version, X is assumed to be compact and θ_{κ} may not have analytic singularities (instead $\nu(\theta_{\kappa},x) = \nu(\theta,x)$). Nevertheless, the proof also works in the relatively compact setting as explained in the paragraph after [\[Dem92,](#page-32-9) Main Thm 1.1]. Furthermore, [\[Dem92,](#page-32-9) Rem. 5.15] ensures the existence of θ_{κ} with analytic singularities (see [Definition 4.1\)](#page-13-1) by weakening the condition on θ_{κ} 's Lelong numbers satisfying (ii).

3. Pushforwards of MA products by submersions

Following [\[LRRS18,](#page-32-6) Sec. $4 \&$ Prop. 5.3], we can define Monge-Ampère products for higher degrees than in BTD if one considers the pushforwards by proper holomorphic submersions.

Proposition 3.1. Let $\pi: X \to Y$ be a proper holomorphic submersion between complex manifolds X and Y with m-dimensional fibres, let T be a closed positive (p, p) -current on X, and let θ be a closed γ -pos $(1,1)$ -current for a (positive) $(1,1)$ -form γ such that for any small enough open $V \subset Y$, there is a closed positive $(1,1)$ -form γ_+ on $\pi^{-1}(V)$ with $\gamma \leq \gamma_+$ (for example, π is Kähler, see [Remark 3.2\)](#page-8-0). Let $\theta_{\kappa} \in {\{\theta\}}_{\partial \overline{\partial}}$ be closed γ -positive $(1, 1)$ -forms such that the sequence θ_{κ} is decreasing pointwise to θ , i.e. locally there are γ -psh potentials of θ_{κ} decreasing pointwise to a γ -psh potential of θ ; and let k be a positive integer. If $\pi(L(\theta))$ is contained in analytic A with codim $A \geq k + p - m$, then

$$
S:=\pi_*\big([\theta]^k\wedge T\big):=\lim_{\kappa\to\infty}\pi_*\big(\theta^k_\kappa\wedge T\big)
$$

is a closed real $(k+p, k+p)$ -current on Y, independent of the choice of θ_{κ} , and locally² the difference of two closed positive currents. If θ is positive, then S is positive.

We would like to stress that in general, $[\theta]^k \wedge T$ is not defined as a current on X for codim $A - p < k \leq$ codim $A - p + m$ in the setting above. Later, we will see that it can be defined for currents with analytic singularities, see [Section 4.](#page-13-0)

Remark 3.2. Following [\[Fuj78,](#page-32-13) Def. 4.1], a holomorphic function $\pi \colon X \to Y$ is called Kähler if there is an open covering $\{U_{\kappa}\}\$ of X and plurisubharmonic functions φ_{κ} on U_{κ} s.t. φ_{κ} is strictly plurisubharmonic on $\pi^{-1}(y) \cap U_\kappa$ and $\varphi_\kappa - \varphi_\lambda$ is pluriharmonic on $\pi^{-1}(y) \cap U_\kappa \cap U_\lambda$ for all $y \in Y$. In case of that π is a locally trivial proper holomorphic submersion (which is equivalent to all fibres of π are biholomorphic to each other, cf. [\[FG65\]](#page-32-14)), we get π is Kähler if its fibres are Kähler. In general, deformations of Kähler manifolds are not necessarily Kähler morphisms, see [\[Bin83,](#page-31-6) Exp. 3.9] due to Deligne.

By [\[Fuj78,](#page-32-13) Lem. 4.4], the preimages $U = \pi^{-1}(V)$ of a proper Kähler $\pi \colon X \to Y$ are Kähler for any small enough open $V \subset Y$, i. e. there is a Kähler form ω_U on U. Moreover, for any $(1,1)$ -form γ on X and any small enough open V, there exists a constant C such

 2 to be more precise, on all V from before

that $\gamma_+ := C \omega_U \geq \gamma$ with closed positive form γ_+ on $U = \pi^{-1}(V)$. Analogously, every closed (1, 1)-form α can be decomposed as $\alpha = \alpha_+ - \alpha_-$ with closed positive forms α_{\pm} on $\pi^{-1}(V)$.

Proof of [Proposition 3.1.](#page-8-1) By transferring the argumentation in [\[LRRS18,](#page-32-6) Sec. 4] to our setting, we first prove the positive case as follows.

We may define S local. Therefore, we can shrink Y such that γ_+ is defined on $X =$ $\pi^{-1}(Y)$. Set $\vartheta := \theta + \gamma_+ \geq 0$ and $\vartheta_{\kappa} = \theta_{\kappa} + \gamma_+ \geq 0$. Let A denote the analytic set of codimension $k + p - m$ such that $\pi(L(\theta)) = \pi(L(\theta)) \subset A$. On $X \setminus \pi^{-1}(A)$, Bedford-Taylor (see [Theorem 2.6\)](#page-6-0) implies that we can define $\vartheta^k \wedge T := \lim_{\kappa \to \infty} \vartheta^k \wedge T$. The direct image $S' := \pi_*(\vartheta^k \wedge T)$ is a closed positive current on $Y \setminus A$ which is the weak limit of the closed positive currents $S_{\kappa} := \pi_*(\vartheta_{\kappa}^k \wedge T)$. We set $s := \dim Y - k - p + m = n - k - p$. As A is of codimension $k + p - m$, there is an (s, s) -bump form β at y with arbitrarily small support such that $A \cap \text{supp} dd^c \beta = \emptyset$, see [Remark 2.3.](#page-5-2) Let α be a closed real $(1, 1)$ -form in the same $\partial\overline{\partial}$ -class of ϑ (and ϑ_{κ}). Since $\vartheta_{\kappa} = \theta_{\kappa} + \gamma_{+}$ is decreasing pointwise to $\vartheta = \theta + \gamma_{+}$, there is a sequence of α -psh functions q_{κ} with $dd^c q_{\kappa} = \vartheta_{\kappa} - \alpha$ which is decreasing pointwise to an α -psh function q with $dd^c q = \vartheta - \alpha$. We get

$$
\int S_{\kappa} \wedge \beta = \int \vartheta_{\kappa}^{k} \wedge T \wedge \pi^{*} \beta
$$
\n
$$
= \int \left(\alpha^{k} + \sum_{i=1}^{k} {k \choose i} \alpha^{k-i} \wedge (dd^{c} q_{\kappa})^{i} \right) \wedge T \wedge \pi^{*} \beta
$$
\n
$$
= \int \left(\alpha^{k} \wedge \pi^{*} \beta + q_{\kappa} \cdot \pi^{*} dd^{c} \beta \wedge \sum_{i=1}^{k} {k \choose i} \alpha^{k-i} \wedge (dd^{c} q_{\kappa})^{i-1} \right) \wedge T
$$
\n
$$
\xrightarrow{\kappa \to \infty} \int \left(\alpha^{k} \wedge \pi^{*} \beta + q \cdot \pi^{*} dd^{c} \beta \wedge \sum_{i=1}^{k} {k \choose i} \alpha^{k-i} \wedge (dd^{c} q)^{i-1} \right) \wedge T =: S_{\beta}
$$

whereby the convergence follows from Bedford-Taylor as $dd^c\beta$ has no support in A, see [Theorem 2.6.](#page-6-0) By [Proposition 2.5,](#page-5-3) we get that S' uniquely extends to a closed positive current S which is the weak limit of S_{κ} .

The quasipositive case is now a direct consequence of the following equation, cf. (2.1) . (3.1)

$$
\pi_*([\theta]^k \wedge T) = \pi_*([\theta + \gamma_+ - \gamma_+]^k \wedge T) := \sum_{i=0}^k (-1)^i {k \choose i} \pi_*([\theta + \gamma_+]^{k-i} \wedge \gamma_+^i \wedge T)
$$

\n
$$
\stackrel{\text{def}}{=} \lim_{\kappa \to \infty} \sum_{i=0}^k (-1)^i {k \choose i} \pi_*((\theta_{\kappa} + \gamma_+)^{k-i} \wedge \gamma_+^i \wedge T) = \lim_{\kappa \to \infty} \pi_*(\theta_{\kappa}^k \wedge T).
$$

Thereby, the terms in the second line of (3.1) are well-defined by the first part of this proof. Due to the last equation in [\(3.1\)](#page-9-0), this definition is independent of the choice of γ_+ .

Remark 3.3. If θ_{κ} is a sequence of γ -pos currents (not necessarily smooth) which is decreasing pointwise to θ , then $L(\theta_{\kappa}) \subset L(\theta)$. Therefore, $\pi_*([\theta_{\kappa}]^k \wedge T)$ are also defined by [Proposition 3.1](#page-8-1) for all κ . Moreover, the proof above works also for non-smooth θ_{κ} such that $\pi_*([\theta]^k \wedge T) = \lim_{\kappa \to \infty} \pi_*([\theta_\kappa]^k \wedge T)$. In other words, the conclusions of Proposition 3.1 [are also correct under the exactly same assumptions within the propositio](#page-8-1)n beside of that θ_{κ} are allowed to be currents instead of forms. This is worth mentioning since we do not need the extra assumption on $\{\theta\}_{\partial\overline{\partial}}$ which will be assumed to obtain the (full) monotone continuity result, [Theorem 3.5](#page-10-0) below.

Remark 3.4. If there is a closed (positive) $(1, 1)$ -form ω on Y such that θ is $\pi^*\omega$ -positive, then we obtain as in [\(3.1\)](#page-9-0) that

$$
\pi_*([\theta]^k \wedge T) = \sum_{i=0}^k (-1)^i {k \choose i} \pi_* \left([\theta + \pi^* \omega]^{k-i} \wedge \pi^* \omega^i \wedge T \right)
$$

$$
= \sum_{i=0}^k (-1)^i {k \choose i} \pi_* \left([\theta + \pi^* \omega]^{k-i} \wedge T \right) \wedge \omega^i
$$

where the last equation follows from the projection formula and that π 's fibres are of dimension m. For $k + p = m + 1$, we get that

$$
\pi_*([\theta]^{m-p+1}\wedge T)=\pi_*([\theta+\pi^*\omega]^{m-p+1}\wedge T)-(m-p+1)\omega\wedge \pi_*([\theta+\pi^*\omega]^{m-p}\wedge T).
$$

If $T = \pi^*S$ for a closed positive current S on Y, then

$$
\pi_*([\theta]^k) \wedge S = \sum_{i=0}^{k-m} (-1)^i {k \choose i} \pi_*([\theta + \pi^* \omega]^{k-i}) \wedge \omega^i \wedge S \text{ and}
$$

$$
\pi_*([\theta]^{m+1}) \wedge S = \pi_*([\theta + \pi^* \omega]^{m+1}) \wedge S - (m+1) \omega \wedge \pi_*([\theta + \pi^* \omega]^m) \wedge S.
$$

We get the following monotone continuity result for the extended BTD MA products defined by [Proposition 3.1.](#page-8-1)

Theorem 3.5. Let $\pi: X \to Y$ be a proper holomorphic submersion between complex manifolds X and Y with m-dimensional fibres, and let T be a closed positive (p, p) -current on X. For $i = 1, \ldots, t$, let θ_i be closed γ -pos $(1, 1)$ -currents on X for a (pos) $(1, 1)$ -form γ such that for any small enough open $V \subset Y$, there are closed positive $(1, 1)$ -forms γ_+ and $\alpha_{i, \pm}$ with $\gamma \leq \gamma_+$ and $\alpha_{i,+} - \alpha_{i,-} \in {\theta_i}_{\partial \overline{\partial}}$ on $\pi^{-1}(V)$ (for example, π is Kähler). Let $\theta_{i,\kappa} \in {\theta_i}_{\partial \overline{\partial}}$ be closed γ -positive (1, 1)-currents decreasing pointwise to θ_i , i.e. there are local γ -psh potentials of $\theta_{i,k}$ decreasing pointwise to a γ -psh potential of θ_i ; and let k_i be positive integers. If $\bigcup_{i=1}^t \pi(L(\theta_i))$ is contained in an analytic A with codim $A \geq k_1 + \ldots + k_t - m + p$, then

(3.2)
$$
\pi_*([\theta_t]^{k_t} \wedge \cdots \wedge [\theta_1]^{k_1} \wedge T) = \lim_{\kappa \to \infty} \pi_*([\theta_{t,\kappa}]^{k_t} \wedge \cdots \wedge [\theta_{1,\kappa}]^{k_1} \wedge T).
$$

Proof. Without loss of generality, we may assume that $k_1 = \ldots = k_t = 1$. The assumption on the codimension of A reads then codim $A \ge t + p - m$. By a recursive application of [Proposition 3.1,](#page-8-1) we obtain that $\pi_*([\theta_t]\wedge \cdots \wedge [\theta_1]\wedge T)$ and (in case that $\theta_{i,\kappa}$ is not smooth) $\pi_*([\theta_{t,\kappa}] \wedge \cdots \wedge [\theta_{1,\kappa}] \wedge T)$ are well-defined currents for all κ .

As the statement is local with respect to Y, we may shrink Y and $X = \pi^{-1}(Y)$ such that γ_+ and α_{\pm} are defined on X. Let us set $\vartheta_i := \theta_i + \gamma_+$ and $\vartheta_{i,\kappa} := \theta_{i,\kappa} + \gamma_+$. By assumption, these are closed positive currents, and we get that the sequence $\vartheta_{i,\kappa}$ is decreasing pointwise to ϑ_i in the sense defined above. To infer the statement in the quasipos case from the positive one, we are going to prove

(3.3)
$$
\pi_*([\theta_t] \wedge \cdots \wedge [\theta_{i+1}] \wedge [\vartheta_i] \wedge \cdots \wedge [\vartheta_1] \wedge T)
$$

$$
= \lim_{\kappa \to \infty} \pi_*([\theta_{t,\kappa}] \wedge \cdots \wedge [\theta_{i+1,\kappa}] \wedge [\vartheta_{i,\kappa}] \wedge \cdots \wedge [\vartheta_{1,\kappa}] \wedge T)
$$

by induction over the number of θ_j -factors, $t-i$.

First, we proof the induction step, $t-i \mapsto t-i+1$ for $1 \leq i \leq t$ (i.e. $i \mapsto i-1$). By induction assumption, we have [\(3.3\)](#page-10-1). Yet, the induction assumption also implies that

$$
\pi_*([\theta_t] \wedge \cdots \wedge [\theta_{i+1}] \wedge \gamma_+ \wedge [\vartheta_{i-1}] \wedge \cdots \wedge [\vartheta_1] \wedge T) =
$$

$$
\lim_{\kappa \to \infty} \pi_*([\theta_{t,\kappa}] \wedge \cdots \wedge [\theta_{i+1,\kappa}] \wedge \gamma_+ \wedge [\vartheta_{i,\kappa}] \wedge \cdots \wedge [\vartheta_{1,\kappa}] \wedge T)
$$

as there are $t-1-(i-1)=t-i$ of θ_i -factors. Thereby, we use that γ_+ is positive and closed. Summing up [\(3.3\)](#page-10-1) and [\(3.4\)](#page-11-0), we obtain (since $\vartheta_i = \theta_i + \gamma_+$ and $\vartheta_{i,\kappa} = \theta_{i,\kappa} + \gamma_+$)

$$
\pi_*([\theta_t] \wedge \cdots \wedge [\theta_i] \wedge [\vartheta_{i-1}] \wedge \cdots \wedge [\vartheta_1] \wedge T) = \lim_{\kappa \to \infty} \pi_*([\theta_{t,\kappa}] \wedge \cdots \wedge [\theta_{i,\kappa}] \wedge [\vartheta_{i-1,\kappa}] \wedge \cdots \wedge [\vartheta_{1,\kappa}] \wedge T)
$$

which is the case $t - (i - 1) = (t - i) + 1$. This concludes the induction step.

Next, we prove the base case $t-i = 0$, i.e. there is not any θ_i -factor and all factors are positive. We obtain that the currents $S = \pi_*([\vartheta_t]\wedge \cdots \wedge [\vartheta_1]\wedge T)$ and $S_{\kappa} := \pi_*([\vartheta_{t,\kappa}]\wedge \cdots \wedge$ $[\vartheta_{1,\kappa}]\wedge T$ are closed positive $(t+p-m,t+p-m)$ -currents. Therefore, the weak convergence of S_{κ} to S follows from [Proposition 2.5](#page-5-3) when its assumptions (i) and (ii) are satisfied which we show as follows.

(i) Since $\vartheta_{i,k}$ is decreasing pointwise to ϑ_i , we get that $L(\vartheta_{i,k}) \subset L(\vartheta_i)$ for all i and κ . By [Theorem 2.6,](#page-6-0) we obtain that

$$
\vartheta_t \wedge \cdots \wedge \vartheta_1 \wedge T = \lim_{\kappa \to \infty} \vartheta_{t,\kappa} \wedge \cdots \wedge \vartheta_{1,\kappa} \wedge T
$$

on $X \setminus \pi^{-1}(A) \subset X \setminus \bigcup_i \pi(L(\vartheta_i))$, and

$$
S = \pi_* \big(\vartheta_t \wedge \cdots \wedge \vartheta_1 \wedge T \big) = \lim_{\kappa \to \infty} \pi_* \big(\vartheta_{t,\kappa} \wedge \cdots \wedge \vartheta_{1,\kappa} \wedge T \big) = \lim_{\kappa \to \infty} S_{\kappa}
$$

on $Y \setminus A$.

(ii) Since codim $A \geq s := t + p - m$, for each point $x \in A$, there is an $(n-s, n-s)$ -bump form β with arbitrarily small support and $dd^c \beta \cap A = \emptyset$, see [Remark 2.3.](#page-5-2) By assumption, $\alpha := \alpha_{t,+} + \gamma_+ - \alpha_{t,-}$ is the difference of two closed positive forms and in the same $\partial \overline{\partial}$ -class as $\vartheta_t = \theta_t + \gamma_+$, i.e. there is a quasipsh function q on X such that $\vartheta_t - \alpha = dd^c q$. By the assumption on $\vartheta_{t,\kappa}$, we get quasipsh functions q_{κ} on X decreasing pointwise to q s.t. $\vartheta_{t,\kappa} - \alpha = dd^c q_{\kappa}$. We get

$$
\int \pi_*([\vartheta_t] \wedge \cdots \wedge [\vartheta_1] \wedge T) \wedge \beta = \int \pi_*([\mathrm{d}d^c q] \wedge [\vartheta_{t-1}] \wedge \cdots \wedge [\vartheta_1] \wedge T) \wedge \beta
$$

$$
+ \int \pi_*([\vartheta_{t-1}] \wedge \cdots \wedge [\vartheta_1] \wedge \alpha \wedge T) \wedge \beta
$$

where the last term is defined since α is the difference of two closed positive forms. Let $q^{(\lambda)}$ be smooth α -psh functions decreasing pointwise to q, and let $\vartheta_i^{(\lambda)}$ $i^{(\lambda)}$ be positive forms decreasing pointwise to ϑ_i for $i = 1, \ldots, t-1$. Since recursively defined by [Proposition 3.1,](#page-8-1) we obtain

$$
\int \pi_*([dd^c q] \wedge [\vartheta_{t-1}] \wedge \cdots \wedge [\vartheta_1] \wedge T) \wedge \beta
$$

= $\lim_{\lambda_t \to \infty} \cdots \lim_{\lambda_1 \to \infty} \int dd^c q^{(\lambda_t)} \wedge \vartheta_{t-1}^{(\lambda_{t-1})} \wedge \cdots \wedge \vartheta_1^{(\lambda_1)} \wedge T \wedge \pi^* \beta$.

Since $q^{(\lambda)}$ are (globally defined) functions on X, we can apply integration by parts (as the definition of d and d^c on currents) in the limit-term of the RHS and get

$$
\int dd^c q^{(\lambda_t)} \wedge \vartheta_{t-1}^{(\lambda_{t-1})} \wedge \cdots \wedge \vartheta_1^{(\lambda_1)} \wedge T \wedge \pi^* \beta
$$
\n
$$
= \int q^{(\lambda_t)} \cdot \vartheta_{t-1}^{(\lambda_{t-1})} \wedge \cdots \wedge \vartheta_1^{(\lambda_1)} \wedge T \wedge dd^c \pi^* \beta
$$
\n
$$
= \int_{X \backslash U} q^{(\lambda_t)} \cdot \vartheta_{t-1}^{(\lambda_{t-1})} \wedge \cdots \wedge \vartheta_1^{(\lambda_1)} \wedge T \wedge dd^c \pi^* \tilde{\beta}
$$
\n
$$
= \int_{X \backslash U} dd^c q^{(\lambda_t)} \wedge \vartheta_{t-1}^{(\lambda_{t-1})} \wedge \cdots \wedge \vartheta_1^{(\lambda_1)} \wedge T \wedge \pi^* \tilde{\beta}
$$
\n
$$
\xrightarrow{\lambda_1 \to \infty} \cdots \xrightarrow{\lambda_t \to \infty} \int_X dd^c q \wedge \vartheta_{t-1} \wedge \cdots \wedge \vartheta_1 \wedge T \wedge \pi^* \tilde{\beta}
$$

where $U = U_{\delta} = \pi^{-1}(V_{\delta})$ for a neighbourhood $V = V_{\delta}$ of A in X (so small that $V_{\delta} \cap$ supp $dd^c \beta = \emptyset$) and $\tilde{\beta} = \chi \beta$ for a smooth cutoff function χ with support on $Y \setminus U$ and $\chi|_{\text{supp} dd^c\beta} = 1$. As $L(\vartheta_i) = L(\theta_i) \subset U$, the convergence is implied by Bedford-Taylor, see [Theorem 2.6.](#page-6-0) Since $dd^c q$ equals $\vartheta_t - \alpha$, we get

(3.5)
$$
\int \pi_*([\vartheta_t] \wedge \cdots \wedge [\vartheta_1] \wedge T) \wedge \beta = \int_{X \setminus U} \vartheta_t \wedge \cdots \wedge \vartheta_1 \wedge T \wedge \pi^* \tilde{\beta} + \int_U \pi_*([\vartheta_{t-1}] \wedge \cdots \wedge [\vartheta_1] \wedge \alpha \wedge T) \wedge \beta.
$$

Repeating exactly the same arguments in the lines above replacing ϑ_i by $\vartheta_{i,\kappa}$ and q by q_{κ} , we obtain the same formula for all κ :

(3.6)
$$
\int \pi_*([\vartheta_{t,\kappa}] \wedge \cdots \wedge [\vartheta_{1,\kappa}] \wedge T) \wedge \beta = \int_{X \setminus U} \vartheta_{t,\kappa} \wedge \cdots \wedge \vartheta_{1,\kappa} \wedge T \wedge \pi^* \tilde{\beta} + \int_U \pi_*([\vartheta_{t-1,\kappa}] \wedge \cdots \wedge [\vartheta_{1,\kappa}] \wedge \alpha \wedge T) \wedge \beta.
$$

By [Theorem 2.6,](#page-6-0) we get that the first term on the RHS in [\(3.6\)](#page-12-1) converges to the first one on the RHS in [\(3.5\)](#page-12-2). We obtain the same for the second term as we may assume that the proposition is already proven for $(t-1)$ -factors (i.e. by induction over t) since α is the difference of two closed positive forms. Hence, we get

$$
\int \pi_*([\vartheta_t] \wedge \cdots \wedge [\vartheta_1] \wedge T) \wedge \beta = \lim_{\kappa \to \infty} \int \pi_*([\vartheta_{t,\kappa}] \wedge \cdots \wedge [\vartheta_{1,\kappa}] \wedge T) \wedge \beta.
$$

Using the idea of [\[LRSW22\]](#page-32-7) to define the wedge product of pushforwards currents with the fibre product, [Theorem 3.5](#page-10-0) implies a monotone continuity result for the recursively defined mixed products:

Corollary 3.6. For $i = 1, \ldots, t$, let $\pi_i: X_i \to Y$ be proper holomorphic submersions between complex manifolds X_i and Y with m_i -dimensional fibres, let θ_i be closed γ_i -positive (1, 1)currents on X_i for (positive) (1, 1)-forms γ_i on X_i such that for any small enough open $V \subset Y$, there are closed positive $(1,1)$ -forms $\gamma_{i,+}$ and $\alpha_{i,\pm}$ on $\pi_i^{-1}(V)$ with $\gamma_i \leq \gamma_{i,+}$ and $\alpha_{i,+}-\alpha_{i,-}\in\{\theta_i\}_{\partial\overline{\partial}}$ (for example, π_i are Kähler). Let $\theta_{i,\kappa}\in\{\theta_i\}_{\partial\overline{\partial}}$ be closed γ_i -positive $(1,1)$ -currents decreasing pointwise to θ_i , let T be a closed pos (p, p) -current on X_1 , and let k_i be positive integers.

If $\bigcup_{i=1}^t \pi_i(L(\theta_i))$ is contained in an analytic A with codim $A \geq k_1 + \ldots + k_t - m_1 \ldots - m_t + p$, then

$$
(3.7) \quad \pi_{t,*}\big([\theta_t]^{k_t}\big) \wedge \cdots \wedge \pi_{1,*}\big([\theta_1]^{k_1} \wedge T\big) = \lim_{\kappa \to \infty} \pi_{t,*}\big([\theta_{t,\kappa}]^{k_t}\big) \wedge \cdots \wedge \pi_{1,*}\big([\theta_{1,\kappa}]^{k_1} \wedge T\big).
$$

Proof. Let $\varpi: \tilde{X} \to Y$ be the fibre product of $\tilde{X} = X_t \times_Y \cdots \times_Y X_1$, let $\text{pr}_i: \tilde{X} \to X_i$ be its projection on the *i*th component, set $\tilde{\theta}_i := \text{pr}_i^* \theta_i$, set $\tilde{\theta}_{i,\kappa} := \text{pr}_i^* \theta_{i,\kappa}$, and set $\gamma :=$ $\sum_i \operatorname{pr}_i^* \gamma_i$ (such that $\gamma \leq \sum_i \operatorname{pr}_i^* \gamma_{i,+}$ on $\varpi^{-1}(V)$). Let $\theta_i^{(\lambda)}$ $i^{(\lambda)}$ be γ_i -pos $(1, 1)$ -forms decreasing pointwise to θ_i , and set $\tilde{\theta}_i^{(\lambda)}$ $\hat{\theta}_i^{(\lambda)}:=\operatorname{pr}_i^*\theta_i^{(\lambda)}$ $i^{(0)}$. As recursively defined by [Proposition 3.1](#page-8-1) (using the projection formula, as well), we get

(3.8)
\n
$$
\pi_{t,*}\left([\theta_t]^{k_t}\right) \wedge \cdots \wedge \pi_{1,*}\left([\theta_1]^{k_1} \wedge T\right)
$$
\n
$$
\stackrel{\text{def}}{=} \lim_{\lambda_t \to 0} \cdots \lim_{\lambda_1 \to 0} \pi_{t,*}\left((\theta_t^{(\lambda_t)})^{k_t}\right) \wedge \cdots \wedge \pi_{1,*}\left((\theta_1^{(\lambda_1)})^{k_1} \wedge T\right)
$$
\n
$$
= \lim_{\lambda_t \to 0} \cdots \lim_{\lambda_1 \to 0} \varpi_*\left((\tilde{\theta}_t^{(\lambda_t)})^{k_t} \wedge \cdots \wedge (\tilde{\theta}_1^{(\lambda_1)})^{k_1} \wedge \text{pr}_1^* T\right)
$$
\n
$$
\stackrel{\text{def}}{=} \varpi_*\left([\tilde{\theta}_t]^{k_t} \wedge \cdots \wedge [\tilde{\theta}_1]^{k_1} \wedge \text{pr}_1^* T\right)
$$

where the second equation follows from [Lemma 2.1.](#page-5-4) Repeating these arguments for $\theta_{i,k}$ instead of θ_i , we obtain also

$$
(3.9) \qquad \pi_{t,*}\big([\theta_{t,\kappa}]^{k_t}\big) \wedge \cdots \wedge \pi_{1,*}\big([\theta_{1,\kappa}]^{k_1} \wedge T\big) = \varpi_*\big([\tilde{\theta}_{t,\kappa}]^{k_t} \wedge \cdots \wedge [\tilde{\theta}_{1,\kappa}]^{k_1} \wedge \mathrm{pr}_1^* T\big)
$$

for all κ . By [Theorem 3.5,](#page-10-0) the RHS of [\(3.9\)](#page-13-2) converges weakly to the right side of [\(3.8\)](#page-13-3) as $\kappa \to \infty$. Hence, the same applies to the LHS of the equations what proves the claimed. \square

4. Currents with analytic singularities

Throughout this section, let X be a complex manifold of dimension n. The main objective of this section is the study of wedge products of currents with analytic singularities.

Definition 4.1. We call q a quasipsh function with analytic singularities in Z if locally $q = u + \text{smth}$ whereby $u = c \log ||F||^2 + b$ is a psh function with analytic singularities in the (reduced) analytic set $\{F = 0\}$ for a positive constant c, a tuple F of holomorphic functions and bounded b. Moreover, we call q a quasipsh function with *neat* analytic singularities if b can be chosen to be smooth, i.e. locally $q = c \log ||F||^2 + \text{smth}$.

We say that a closed positive θ has (neat) analytic singularities when its local dd^cpotential is psh with (neat) analytic singularities. θ is called *quasipositive* (shortly *quasi*pos) with (neat) analytic singularities if it is locally the sum of a closed positive current with (neat) analytic singularities and a closed real $(1, 1)$ -form. This is equivalent to that locally $\theta = dd^c q$ for a quasipsh function q with (neat) analytic singularities.

Let u_1, \ldots, u_k be psh functions with analytic singularities in Z_1, \ldots, Z_k , resp. For constructible sets W_2, \ldots, W_k such that $W_j \subset Z_j^c := X \setminus Z_j$, we can define the MA product

(4.1)
$$
dd^c u_k \wedge \mathbb{1}_{W_k} dd^c u_{k-1} \wedge \mathbb{1}_{W_{k-1}} \cdots dd^c u_2 \wedge \mathbb{1}_{W_2} dd^c u_1
$$

recursively (from right to left) as a closed positive current, see [\[LRSW22,](#page-32-7) Prop. 3.2]. In case of $W_j = X \setminus Z_j$, we call [\(4.1\)](#page-13-4) the (mixed) Andersson-Wulcan (AW) MA product of $dd^c u_k, \ldots, dd^c u_1$. For a psh function u with analytic singularities in Z, $(dd^c u)^k_{\text{AW}}$ denotes the k-power $dd^c u \wedge \mathbb{1}_{Z^c} \cdots \mathbb{1}_{Z^c} dd^c u$ which was introduced in [\[AW14\]](#page-31-1).

Please note there is no restriction on k . Moreover, this definition of (mixed) Monge-Ampère products is an extension of BTD MA, see [\[LRSW22,](#page-32-7) Prop. 3.4], and (mixed) products of Bochner-Martinelli currents $dd^c \log ||F_k||^2$ for holomorphic tuples F_k as it was introduced in [\[ASWY17,](#page-31-2) Sec. 5]. Also, we would like to point out that [\(4.1\)](#page-13-4) can be defined for quasipsh potentials, as well, see $[Bb19, \text{LSW20}]$ and also $[LRSW22, \text{Lem. } 3.5]$ (i. p. for γ -psh potentials for a (pos) form γ).

As in [\[LRSW22,](#page-32-7) Sec. 3] and [\[LSW20,](#page-32-8) Sec. 2], we consider the following class of currents.

Definition 4.2. We call a (closed) real (p, p) -current T a (closed) current with analytic singularities if it is locally of the following form:

$$
(4.2) \qquad \sum_{i} \alpha_i \wedge \mathbb{1}_{W_{i,k_i+1}} dd^c u_{i,k_i} \wedge \mathbb{1}_{W_{i,k_i}} dd^c u_{i,k_i-1} \mathbb{1}_{W_{i,k_i-1}} \cdots \wedge \mathbb{1}_{W_{i,2}} dd^c u_{i,1} \mathbb{1}_{W_{i,1}}
$$

with smooth closed real forms α_i of degree $(p-k_i, p-k_i)$, psh $u_{i,j}$ with analytic singularities, and constructible sets $W_{i,j}$ such that $L(u_{i,j}) \subset W_{i,j}^c$ for $j \leq 2$. As $\mathbb{1}_{\{\text{pt}\}}T = 0$ for (p, p) currents with $p < n$, we can omit the last condition on $W_{i,j}$ if $L(u_{i,j})$ is a set of isolated points.

If furthermore $L(u_{i,1}) \subset W_{i,1}^c$, $T \wedge \cdot$ denotes an operator on closed currents with analytic singularities (which is given recursively and well-defined due to the condition on $W_{i,1}$).

In particular, the AW MA product of closed quasipositive $(1, 1)$ -currents with analytic singularities (see (4.1)) is a closed current with analytic singularities. Let us use the following more specific notation for such currents.

Definition 4.3. A closed (p, p) -current T with analytic singularities is called *quasipositive* (shortly quasipos) if T is locally of the following form:

$$
(4.3) \qquad \sum_{i} c_i \mathbb{1}_{W_{i,p+1}} dd^c q_{i,p} \wedge \mathbb{1}_{W_{i,p}} dd^c q_{i,p-1} \mathbb{1}_{W_{i,p-1}} \cdots \wedge \mathbb{1}_{W_{i,2}} dd^c q_{i,1} \mathbb{1}_{W_{i,1}}
$$

with positive constants c_i , quasipsh $q_{i,j}$ with analytic singularities, and constructible sets $W_{i,j}$ such that $L(q_{i,j}) \subset W_{i,j}^c$ for $j \geq 2$.

Obviously, a closed (p, p) -current T with analytic singularities is quasipos if and only if it is locally of the form as in [\(4.2\)](#page-14-0) with α_i is either a positive constant or the wedge product of closed real $(1, 1)$ -forms. In particular, locally T is the difference of two closed positive currents.

Remark 4.4. Let $\pi: X \to Y$ be a proper holomorphic submersion, and let γ be a (pos) $(1, 1)$ -form such that for all small enough open $V \subset Y$, there is a *closed* positive $(1, 1)$ form γ_+ with $\gamma_+ \geq \gamma$ on $U = \pi^{-1}(V)$, e.g. π is Kähler, see [Remark 3.2.](#page-8-0) Let $\theta_1, \ldots, \theta_p$ be closed γ -pos (1, 1) currents with analytic singularities, and let W_2, \ldots, W_p be constructible sets with $L(\theta_j) \subset W_j^c$ for $j \geq 2$. Then, $T = \theta_p \wedge \mathbb{1}_{W_p} \cdots \theta_2 \wedge \mathbb{1}_{W_2} \theta_1$ can be decomposed into the difference $T_+ - T_-$ of two closed positive currents T_{\pm} with analytic singularities

on $U = \pi^{-1}(V)$ for all small enough V (to be more precise, on all U on which the γ_+ from above exists).

Let us prove this by induction over p. The base case $p = 1$ is trivial. For the induction step, we may assume that there are two closed positive currents S_{\pm} with analytic singularities such that $\theta_{p-1} \wedge 1_{W_{p-1}} \cdots \theta_2 \wedge 1_{W_2} \theta_1 = S_+ - S_-$. Then,

$$
T = \theta_p \wedge \mathbb{1}_{W_p}(S_+ - S_-) = ((\theta_p + \gamma_+) \wedge \mathbb{1}_{W_p} S_+ + \gamma_+ \wedge \mathbb{1}_{W_p} S_-) - ((\theta_p + \gamma_+) \wedge \mathbb{1}_{W_p} S_- + \gamma_+ \wedge \mathbb{1}_{W_p} S_+)
$$

where minuend and subtracted are closed and positive

where minuend and subtrahend are closed and positive.

We obtain two classes of currents which are closed under the following operations:

(1) wedge products with closed real forms (of bidegree $(1, 1)$);

(2) $\mathbb{1}_W$ for any constructible set W; and

(3) $dd^c q \wedge \mathbb{1}_{Z^c}$ for any quasipsh function q with analytic singularities in Z.

If T and S are (quasipos) currents with analytic singularities, then $S \wedge T$ is well-defined and a (quasipos) current with analytic singularities where $S \wedge$ denotes the operator as defined in [Definition 4.2.](#page-14-1)

In this sense, the AW MA product has proven to be very useful, e. g. we can define the Monge-Ampère operator for arbitrary degrees. Yet, it lacks certain properties. For example, the product is neither commutative nor distributive, see [\[LRSW22,](#page-32-7) Exp. 3.1]. Furthermore, although it coincides with the BTD MA product whenever this defined (see [\[LRSW22,](#page-32-7) Prop. 3.4]), the pushforward of an AW MA product is in general different form the (natural) extension of BTD defined by [Proposition 3.1](#page-8-1) as the following example shows; cf. [\[LRSW22,](#page-32-7) Exp. 8.3].

Example 4.5. Let X be $\mathbb{C}_z \times \mathbb{P}^1$, let π be the projection on $Y = \mathbb{C}_z$, let u denote the psh function $\log |z|^2$ on X, and let ω_{FS} be the pullback of the Fubini-Study form on \mathbb{P}^1 . Then, $\pi_*((\omega_{\text{FS}}+dd^c u)_{\text{AW}}^2) = \pi_*(\omega_{\text{FS}} \wedge [z=0]) = [z=0]$. On the other side, if u_{κ} is a sequence of smooth psh functions decreasing pointwise to u, then $(\omega_{\text{FS}} + dd^c u_{\kappa})^2$ = $(dd^c u_{\kappa})^2 + 2\omega_{\rm FS} \wedge dd^c u_{\kappa} + 0$ such that $\pi_*((\omega_{\rm FS} + dd^c u_{\kappa})^2) = 2\pi_*(\omega_{\rm FS} \wedge dd^c u_{\kappa})$ which converges to $2[z=0]$, i.e.

$$
\pi_*\big([\omega_{\rm FS} + dd^c u]^2\big) = 2[z=0] \neq \pi_*\big((\omega_{\rm FS} + dd^c u)^2_{\rm AW}\big).
$$

Roughly speaking, the AW MA product removes too much. Inspired by [\[ABW19,](#page-31-3) Thm 1.2], Lärkäng, Raufi, Wulcan and the author give the following definition in [\[LRSW22,](#page-32-7) [LSW20\]](#page-32-8) which keeps track of the removed part by adding an extra term to the AW MA product.

Definition 4.6. For a closed quasipositive $(1, 1)$ -current θ with analytic singularities in Z, closed real (1, 1)-form α , and closed current T with analytic singularities, we get that the so-called *generalized Monge-Ampère product* of θ and T defined by

(4.4)
$$
[\theta]_{\alpha} \wedge T := \theta \wedge \mathbb{1}_{Z^c} T + \alpha \wedge \mathbb{1}_Z T
$$

is a well-defined closed real current with analytic singularities which is quasipos when T is. This is also defined and a (quasipos) current with analytic singularities for α replaced by a quasipos $(1, 1)$ -current η with analytic singularities in isolated points due to BTD.

Generalizing [\[LRSW22,](#page-32-7) Thm 1.2], we obtain the following correlation between the extended BTD MA product defined by [Proposition 3.1](#page-8-1) and pushforwards of the generalized MA products defined by [\(4.4\)](#page-15-1).

Theorem 4.7. Let $\pi: X \to Y$ be a proper holomorphic submersion between complex manifolds with m-dimensional fibres, let γ be a (pos) (1,1)-form, and let T be a closed real (p, p) -current such that for any small enough $V \subset Y$, there are a closed positive form γ_+ with $\gamma \leq \gamma_+$ and closed positive currents T_+ with analytic singularities and $T = T_+ - T_$ on $\pi^{-1}(V)$ (cf. [Remark 4.4\)](#page-14-2). Let θ be a γ -pos $(1,1)$ -current with analytic singularities in Z, and let $\eta \in \{\theta\}_{\partial \overline{\partial}}$ be a closed γ -pos $(1,1)$ -current with neat analytic singularities in isolated points. Then, we get

$$
\pi_*([\theta]^k \wedge T) = \pi_*([\theta]^k_\eta \wedge T)
$$

for all $k + p \leq \operatorname{codim} \pi(Z) + m$. In particular, it is independent of η .

Let us point out in contrast to the general case, $S := \pi_*([\theta]^k \wedge T)$ is actually given as a direct image of a current on X for θ with analytic singularities. We will utilize this fact to estimate the Lelong numbers of S, see [Corollary 5.10.](#page-23-2)

Remark 4.8. By [Remark 4.4,](#page-14-2) we can apply [Theorem 4.7](#page-16-0) recursively. In particular, we get

(4.5)
$$
\pi_*([\theta_t]^{k_t} \wedge \cdots \wedge [\theta_1]^{k_1}) = \pi_*([\theta_t]_{\eta_t}^{k_t} \wedge \cdots \wedge [\theta_1]_{\eta_1}^{k_1})
$$

for closed γ -pos (1, 1)-currents $\theta_1, ..., \theta_t$ with analytic singularities in Z_i , closed γ -pos (1, 1)currents η_1, \ldots, η_t with neat analytic singularities in isolated points such that $\eta_i \in \{\theta_i\}_{\partial \overline{\partial}}$ and $k_1 + \ldots + k_t \leq \operatorname{codim} \bigcup_{i=1}^t \pi(Z_i) + m$ (γ as above).

Remark 4.9. Using the idea from [\[LRSW22\]](#page-32-7) (motivated by [Lemma 2.1\)](#page-5-4), we can use the fibre product to define the wedge product of pushforwards of currents with analytic singularities in the following setting. For $i = 1, \ldots, t$, let $\pi_i : X_i \to Y$ be proper holomorphic submersions with m_i -dimensional fibres, let θ_i be closed γ_i -pos (1, 1)-currents with analytic singularities in Z_i , and let $\alpha_i \in \{\theta_i\}_{\partial \overline{\partial}}$ be closed γ_i -pos $(1, 1)$ -forms where γ_i are (pos) $(1, 1)$ -forms, each on X_i . Following [\[LRSW22,](#page-32-7) Thm 1.1], we define

$$
(4.6) \qquad \qquad \pi_{t,*}\big([\theta_t]_{\alpha_t}^{k_t}\big) \wedge \cdots \wedge \pi_{1,*}\big([\theta_1]_{\alpha_1}^{k_1}\big) := \varpi_*\big([\tilde{\theta}_t]_{\tilde{\alpha}_t}^{k_t} \wedge \cdots \wedge [\tilde{\theta}_1]_{\tilde{\alpha}_1}^{k_1}\big)
$$

where $\varpi: \tilde{X} \to Y$ is the fibre product, $\tilde{X} = X_t \times_Y \cdots \times_Y X_1$, $\mathrm{pr}_i: \tilde{X} \to X_i$ denotes the projection on the *i*th component, $\tilde{\theta}_i := \mathrm{pr}_i^* \theta_i$ and $\tilde{\alpha}_i := \mathrm{pr}_i^* \alpha_i$. In [\(4.6\)](#page-16-1), we cannot replace α_i by currents η_i with neat analytic singularities in isolated points since $\mathrm{pr}_i^* \eta_i$ would have singularities in sets of higher codimension than points. If for any small enough open $V \subset Y$, there are closed positive (1, 1)-forms $\gamma_{i,+}$ with $\gamma_i \leq \gamma_{i,+}$ on $\pi_i^{-1}(V)$, then [\(4.5\)](#page-16-2) implies

$$
(4.7) \qquad \pi_{t,*}([\theta_t]^{k_t}) \wedge \cdots \wedge \pi_{1,*}([\theta_1]^{k_1}) = \pi_{t,*}([\theta_t]_{\alpha_t}^{k_t}) \wedge \cdots \wedge \pi_{1,*}([\theta_1]_{\alpha_1}^{k_1})
$$

for all $k_1 + \ldots + k_t \leq \operatorname{codim} \bigcup_i \pi(Z_i) + m_1 + \ldots + m_t$ (cf. proof of [Corollary 3.6\)](#page-12-0).

Proof of [Theorem 4.7.](#page-16-0) Let α be in the same $\partial\overline{\partial}$ -class as θ (and η), i.e. there exist a quasipsh q with analytic singularities and a quasipsh v with neat analytic singularities in

isolated points such that $\theta = dd^c q + \alpha$ and $\eta = dd^c v + \alpha$. By the definition of the general-ized MA product [\(4.4\)](#page-15-1), we get $[\theta]_{\eta} \wedge T = [dd^c q + \alpha]_{dd^c v + \alpha} \wedge T = ([dd^c q]_{dd^c v} + \alpha) \wedge T$. We set $q_{\kappa} := \max\{q, v - \kappa\}.$ As locally, there is a constant C s.t. $q + C||z||^2$ and $v + C||z||^2$ are psh (where z denotes coordinates of X), we get $q_{\kappa} \stackrel{\text{loc}}{=} \max\{q + C||z||^2, v - \kappa + C||z||^2\} - C||z||^2$ are quasipsh. Moreover, q_{κ} have neat analytic singularities in isolated points and the sequence is decreasing pointwise to q, i.p. $\theta_{\kappa} := dd^c q_{\kappa} + \alpha \longrightarrow \theta$ weakly as $\kappa \to \infty$. On any $U := \pi^{-1}(V)$ for small enough open $V \subset Y$ where $T = T_+ - T_-$ by the assumption, [Lemma 4.11](#page-17-0) below implies

$$
[\theta]_\eta \wedge T_{\pm} = ([dd^c q]_{dd^c v} + \alpha) \wedge T_{\pm} = \lim_{\kappa \to \infty} (dd^c q_\kappa + \alpha) \wedge T_{\pm} = \lim_{\kappa \to \infty} \theta_\kappa \wedge T_{\pm}.
$$

By taking the pushforwards of both sides of the equation, we get the claimed as in the proof of [Proposition 3.1,](#page-8-1) see [Remark 3.3.](#page-9-1)

Lemma 4.10. Let T be a closed current with analytic singularities, and let q be a quasipsh function with analytic singularities in Z. If $\rho_{\kappa} : \mathbb{R} \to \mathbb{R}$ is a sequence of non-decreasing convex and bounded from below functions such that ρ_{κ} is decreasing pointwise to the identity as $\kappa \to \infty$, then

(4.8)
$$
dd^c \rho_\kappa \circ q \wedge T \stackrel{\kappa \to \infty}{\longrightarrow} dd^c q \wedge \mathbb{1}_{Z^c} T.
$$

In particular, if v is a smooth function, then

(4.9)
$$
[dd^c q]_{dd^c v} \wedge T = \lim_{\kappa \to \infty} dd^c(\rho_\kappa \circ (q - v) + v) \wedge T.
$$

Proof. [\(4.8\)](#page-17-1) is a direct consequence of the definition of the AW MA product, see [\[LRSW22,](#page-32-7) Prop. 3.2]. Now, [\(4.8\)](#page-17-1) implies

$$
dd^c \rho_{\kappa} \circ (q - v) \wedge T \stackrel{\kappa \to \infty}{\longrightarrow} dd^c(q - v) \wedge \mathbb{1}_{Z^c} T
$$

which is equivalent to [\(4.9\)](#page-17-2) since we can add $dd^c v \wedge T$ on both sides of the equation. \Box

We get the following variant for v with neat analytic singularities in isolated points.

Lemma 4.11. Let T be a closed quasipos (p, p) -current with analytic singularities, let q be a quasipsh function with analytic singularities in Z , and let v be a quasipsh function with neat analytic singularities in isolated points. We set $q_{\kappa} := \max\{q, v - \kappa\} = \max\{q - v, -\kappa\} + v$ such that q_{κ} are quasipsh functions with neat analytic singularities in isolated points and q_{κ} is decreasing pointwise to q. Then,

(4.10)
$$
[dd^c q]_{dd^c v} \wedge T = \lim_{\kappa \to \infty} dd^c q_\kappa \wedge T.
$$

Proof. As explained in the comment after [Definition 4.3,](#page-14-3) T is locally the difference of two closed positive currents (with analytic singularities). By proving the convergence (4.10) for the minuend and subtrahend separately, we may assume that T is positive. As $\rho_{\kappa} = \max\{\cdot, -\kappa\}$ are non-decreasing convex functions decreasing pointwise to the identity as $\kappa \to \infty$, [Lemma 4.10](#page-17-4) implies the weak convergence [\(4.10\)](#page-17-3) on the complement of $L(v)$. Fix an $x \in L(v)$. Let β be an $(n-p-1, n-p-1)$ -bump form at x such that $\text{supp } dd^c \beta \subset L(v)^c$, see [Remark 2.3.](#page-5-2) Fix a constant C such that $\tilde{v} := \max_{\varepsilon} \{v, -C\}$ equals v on supp $dd^c\beta$ (max_{ϵ} denotes the regularized max for a small enough ε). We set

 $\tilde{q}_{\kappa} := \max\{q, \tilde{v} - \kappa\} = \rho_{\kappa}(q - \tilde{v}) + \tilde{v}$. By [Lemma 4.10,](#page-17-4) we get that

$$
dd^c \tilde{q}_{\kappa} \wedge T \stackrel{\kappa \to \infty}{\longrightarrow} (dd^c q \wedge \mathbb{1}_{Z^c} + dd^c \tilde{v} \wedge \mathbb{1}_Z)T.
$$

Furthermore, \tilde{q}_κ equals q_κ on supp $dd^c\beta$. By applying Stokes and the facts above, we get

$$
\int \beta \wedge dd^c q_{\kappa} \wedge T = \int dd^c \beta \wedge q_{\kappa} T = \int dd^c \beta \wedge \tilde{q}_{\kappa} T = \int \beta \wedge dd^c \tilde{q}_{\kappa} \wedge T
$$
\n
$$
(4.11)
$$
\n
$$
\xrightarrow{\kappa \to \infty} \int \beta \wedge (dd^c q \wedge 1_{Z^c} + dd^c \tilde{v} \wedge 1_Z) T = \int dd^c \beta \wedge (q1_{Z^c} + \tilde{v}1_Z) T
$$
\n
$$
= \int dd^c \beta \wedge (q1_{Z^c} + v1_Z) T = \int \beta \wedge (dd^c q \wedge 1_{Z^c} + dd^c v \wedge 1_Z) T.
$$

Since locally there is a closed positive (1, 1)-form α such that q, v and q_{κ} are α -psh, $S_{\kappa} := (dd^c q_{\kappa} + \alpha) \wedge T$ are positive, and we can apply [Proposition 2.5,](#page-5-3) i.e. S_{κ} is converging weakly to a closed positive current S with S equals $[d d^c q]_{dd^c v} \wedge T + \alpha \wedge T$ on the complement of $L(v)$. In particular, we proved the claimed by using [Lemma 2.4.](#page-5-1)

Let us conclude this section with the following variant of [Lemma 2.4.](#page-5-1)

Lemma 4.12. Let T be a closed quasipos (p, p) -current with analytic singularities such that T has support in an analytic set A with codim $A \geq p$. If for all points $x \in A$, there is an $(n-p, n-p)$ -bump form β such that $\int \beta \wedge T = 0$, then T equals zero.

Proof. By [Definition 4.2,](#page-14-1) we get locally $T = \sum_i T_i$ where

$$
T_i=\alpha_i\wedge \mathbbm{1}_{W_{i,k_i+1}} dd^c u_{i,k_i}\wedge \mathbbm{1}_{W_{i,k_i}}\cdots \wedge \mathbbm{1}_{W_{i,2}} dd^c u_{i,1}\mathbbm{1}_{W_{i,1}}
$$

with smooth closed real forms α_i of degree $(p-k_i, p-k_i)$, psh $u_{i,j}$ with analytic singularities, and constructible sets $W_{i,j}$ such that $L(u_{i,j}) \subset W_{i,j}^c$ for $j \geq 2$. If codim $A > k_i$, then we may assume that T_i vanishes due to T's support (independently of the bump form assumption). So, all terms in T vanish except of T_i with $k_i = \text{codim } A = p$, i.e.

$$
T = \sum_{i} c_i \mathbb{1}_{W_{i,p+1}} dd^c u_{i,p} \wedge \mathbb{1}_{W_{i,p}} \cdots \wedge \mathbb{1}_{W_{i,2}} dd^c u_{i,1} \mathbb{1}_{W_{i,1}}.
$$

As T is quasipos, $c_i \geq 0$, see comment in [Definition 4.3.](#page-14-3) We conclude T is positive. Therefore, [Lemma 2.4](#page-5-1) implies $T = 0$.

5. Lelong numbers of closed real currents

In this section, we are going to study Lelong numbers of closed real currents which are not necessarily positive. To be more precise, we consider currents which are locally the difference of two closed positive ones or currents with analytic singularities, see [Definition 4.2.](#page-14-1) We will recall and generalize basic properties of Lelong numbers following the notation as in the Sections 5ff of [\[Dem12,](#page-32-11) Chp. III]; see also [\[Dem85,](#page-32-2) [Dem93\]](#page-32-3).

Let X be a complex manifold of dimension n, and fix a point $x_0 \in X$ and coordinates z in a neighbourhood of x_0 such shat $z(x_0) = 0$. If u is a (negative) psh function on X, the Lelong number of u in x_0 is defined by

$$
\nu(u, x_0) := \liminf_{x \to x_0} \frac{u(x)}{\log ||z||}.
$$

This definition trivially extends to quasipsh functions q. The following lemmata are direct consequences of this definition.

Lemma 5.1. If $S \subset X$ is a submanifold s.t. $q|S \neq -\infty$, then $\nu(q|S, x_0) \geq \nu(q, x_0)$.

Lemma 5.2. If $\pi: Y \to X$ is a holom. submersion, then $\nu(q \circ \pi, y_0) = \nu(q, \pi(y_0)).$

Proof. As the statement is local with respect to y_0 , we may assume that $X \subset \mathbb{C}^n$, $Y =$ $X \times D$ for $D \subset \mathbb{C}^m$, $\pi: Y \to X$ is the projection and $y_0 = (0, 0)$. Then, for any sequence x_k in X converging to 0, we get $\frac{q(x_k)}{\log \|x_k\|} = \frac{q(\pi(x_k,0))}{\log \| (x_k,0) \|}$ $\frac{q(\pi(x_k,0))}{\log ||(x_k,0)||}$. Therefore, $\nu(q,0) \geq \nu(q \circ \pi,(0,0))$. If $y_k = (x_k, a_k)$ is a sequence in Y converging to $(0, 0)$, then $\frac{q(\pi(x_k, a_k))}{\log ||(x_k, a_k)||} \geq \frac{q(x_k)}{\log ||x_k||}$ $\frac{q(x_k)}{\log \|x_k\|}$ for almost all k. This implies the other inequality $\nu(q, 0) \leq \nu(q \circ \pi, (0, 0)).$

If T is a closed positive (p, p) -current, then the Lelong number of T in x_0 is given by

(5.1)
$$
\nu(T, x_0) = \int \mathbb{1}_{\{x_0\}} \big(dd^c \log ||z|| \big)^{n-p} \wedge T.
$$

Thereby, we use that $L(d d^c \log ||z||) = \{x_0\}$ is of codimension n such that the current $(dd^c\log ||z||^2)^{n-p} \wedge T$ is defined in the sense of BTD; see e.g. [\[Dem12,](#page-32-11) Def. 5.4]. As introduced in [\[LRSW22,](#page-32-7) Sec. 6.4], we can extend the definition of Lelong numbers to currents which locally are given as differences of two closed positive currents as follows.

Lemma 5.3. Let T be a closed real current on X such that in a neighbourhood of a point $x_0, T = T_+ - T_-$ for two closed positive currents T_{\pm} . Then, the Lelong number of T in x_0

$$
\nu(T, x_0) := \nu(T_+, x_0) - \nu(T_-, x_0)
$$

is well-defined since it is independent of the decomposition.

We call $\nu(T, x_0)$ the Lelong number of T in x_0 . By this extension, $\nu(\cdot, x_0)$ is a linear operator on the currents which are given as the difference of two closed positive currents in a neighbourhood of x_0 .

Proof. Let T equal $T_+ - T_-$ and $S_+ - S_-$ for closed positive currents T_{\pm} and S_{\pm} in a neighbourhood of x_0 . Then, we get $T_+ + S_- = S_+ + T_-$ is a closed positive current. As the Lelong number of the sum of closed positive currents is the sum of the Lelong numbers of the summands, we get

$$
\nu(T_+,x_0) - \nu(T_-,x_0) = \nu(S_+ + T_+,x_0) - \nu(S_+ + T_-,x_0)
$$

= $\nu(S_+ + T_+,x_0) - \nu(S_- + T_+,x_0) = \nu(S_+,x_0) - \nu(S_-,x_0).$

By Prop. 5.12 in [\[Dem12,](#page-32-11) Chp. III], we have the following semicontinuity property.

Proposition 5.4. Let T be a closed current such that $T = T_+ - T_-$ for closed positive T_{\pm} in a neighbourhood of x_0 . T_k be a sequence of closed currents which converges weakly to T such that $T_{\kappa,+} := T_{\kappa} + T_{-}$ are closed positive currents for (almost) all κ . Then,

$$
\limsup_{\kappa \to \infty} \nu(T_{\kappa}, x_0) \le \nu(T, x_0).
$$

For proper holomorphic submersions, we get the following estimates between the Lelong numbers of closed positive currents and their pushforwards or pullbacks.

Proposition 5.5. Let $\pi: X \to Y$ be a proper holomorphic submersion between complex manifolds X and Y, with m-dimensional fibres, and let T be a closed positive (p, p) -current on X with $p \geq m$. Then, for all $x_0 \in X$, we get

$$
\nu(T, x_0) \leq \nu(\pi_*T, \pi(x_0)).
$$

Proof. Let n be the dimension of X. We may assume that Y is a small enough neighbourhood of $y_0 := \pi(x_0)$ such that $Y \subset \mathbb{C}^{n-m}$ with $z' = (z_1, \ldots, z_{n-m})$ as coordinates of Y and $z'(y_0) = 0$. By the assumption on π , $F := \pi^{-1}(y_0)$ is a smooth complex manifold of dimension m. We pick some coordinates $z'' = (z_{n-m+1},...,z_n)$ of F in a neighbourhood D of x_0 such that $z''(x_0) = 0$. We may assume that $U = Y \times D$ is a subset of X and π is the projection on Y and obtain that $z = (z', z'')$ are coordinates for U with $\pi \circ z = z'$.

We set $\varphi(z) := \gamma \log ||z||$ for constant $\gamma > 0$ and $\psi(z') := \log ||z'||$ which are both semiexhaustive psh functions defining the Lelong number in x_0 and y_0 , respectively, by using Demailly's generalized Lelong numbers, see [\[Dem12,](#page-32-11) §5 in Chp. III], i. e.

$$
\gamma^{n-p}\nu(T,x_0) = \nu(T,\varphi) := \lim_{r \to -\infty} \nu(T,\varphi,r), \ \nu(T,\varphi,r) := \int_{\{\varphi < r\}} T \wedge (dd^c \varphi)^{n-p} \quad \text{and}
$$

$$
\nu(\pi_*T,y_0) = \nu(\pi_*T,\psi) := \lim_{r \to -\infty} \nu(\pi_*T,\psi,r), \ \nu(\pi_*T,\psi,r) := \int_{\{\psi < r\}} \pi_*T \wedge (dd^c \psi)^{n-p}.
$$

We define $\psi_{\varepsilon} := \frac{1}{2} \log (||z'||^2 + \varepsilon ||z''||^2)$ whose sequence is decreasing pointwise to ψ as $\varepsilon \to 0$ and $\psi_{\varepsilon,s} := \max\{\psi_{\varepsilon,s}\}\$ whose sequence is decreasing pointwise to $\psi_s := \max\{\psi, s\}$ as $\varepsilon \to 0$ and to ψ_{ε} as $s \to -\infty$.

Since $||z'||^2 + \varepsilon ||z''||^2 \le ||z||^2$, we obtain $\gamma \psi_{\varepsilon} \le \varphi$, i.e. $\frac{\varphi(z)}{\psi_{\varepsilon}(z)} \le \gamma$ close to x. Moreover, we have

$$
\gamma \geq \limsup_{\psi_{\varepsilon}(z) \to -\infty} \frac{\varphi(z)}{\psi_{\varepsilon}(z)} \geq \limsup_{z' \to 0} \frac{\varphi(z',0)}{\psi_{\varepsilon}(z',0)} = \gamma.
$$

By Demailly's second comparison theorem for Lelong numbers (cf. Thm 7.8 in [\[Dem12,](#page-32-11) Chp. III]), we get that

$$
\nu(T \wedge (dd^c \varphi)^{n-p}, x_0) \le \gamma^{n-p} \nu(T \wedge (dd^c \psi_{\varepsilon})^{n-p}, x_0) < \nu(T \wedge (dd^c \psi_{\varepsilon})^{n-p}, \varphi),
$$

i.e. for all constants R there is an $r < R$ such that

(5.2)
$$
\int_{B_r} T \wedge (dd^c \varphi)^{n-p} \leq \int_{B_R} T \wedge (dd^c \psi_{\varepsilon})^{n-p}
$$

where $B_r = \{ \varphi < r \}$ and $B_R = \{ \varphi < R \}$ are balls with radii e^r and e^R , resp. As $\psi_{\varepsilon,s} \searrow \psi_{\varepsilon,s}$ we get

(5.3)
$$
\int_{B_R} T \wedge (dd^c \psi_{\varepsilon})^{n-p} \leq \int_{B_R} T \wedge (dd^c \psi_{\varepsilon,s})^{n-p}
$$

for all s small enough, see for example proof of Thm 3.7 in [\[Dem12,](#page-32-11) Chp. III]. Furthermore,

(5.4)

$$
\lim_{\varepsilon \to 0} \int_{B_R} T \wedge (dd^c \psi_{\varepsilon, s})^{n-p} = \int_{B_R} T \wedge (dd^c \psi_s \circ \pi)^{n-p}
$$

$$
\leq \int_{\{\psi \circ \pi < R\}} T \wedge (dd^c \psi_s \circ \pi)^{n-p} = \int_{\{\psi < R\}} \pi_* T \wedge (dd^c \psi_s)^{n-p}
$$

$$
= \int_{\{\psi < R\}} \pi_* T \wedge (dd^c \psi)^{n-p} = \nu(\pi_* T, \psi, R)
$$

where the second last equation follows from Stokes, cf. Prop. 9.3 in [\[Dem12,](#page-32-11) Chp. III]. [\(5.2\)](#page-20-0), [\(5.3\)](#page-20-1) and [\(5.4\)](#page-21-0) combined imply for every R there is an $r < R$ such that $\nu(T, \varphi, r) \leq$ $\nu(\pi_*T, \psi, R)$. Since the RHS converges to $\nu(\pi_*T, y_0)$ as $R \to -\infty$ and the LHS converges to $\gamma^{n-p}\nu(T,x_0)$ as $r \to -\infty$ for every $\gamma < 1$, we proved the claimed.

Proposition 5.6 (Prop. 5 in [\[Meo96\]](#page-32-15)). Let $\pi: X \to Y$ be a proper holomorphic submersion between complex manifolds X and Y, and let T be a closed positive (p, p) -current on Y. Then, for all $x_0 \in X$, we get

$$
\nu(\pi^*T, x_0) \ge \nu(T, \pi(x_0)).
$$

To be more precise, Meo proved that for every holomorphic $f: D_1 \to D_2$ with $D_1 \subset \mathbb{C}^n$ and $D_2 \subset \mathbb{C}^m$, and every closed positive (p, p) -current $T = \lim_{\kappa \to \infty} \theta_{\kappa}$ on D_2 with closed positive (p, p) -forms θ_{κ} such that $f^*T := \lim_{\kappa \to \infty} f^* \theta_{\kappa}$ is a closed positive current on D_1 , we get $\nu(f^*T, x_0) \ge \nu(T, f(x_0)).$

Lelong numbers of currents with analytic singularities.

Remark/Def. 5.7. If T is a closed positive (p, p) -current and α a closed strongly positive (k, k) -form $(k > 0)$, then $\alpha \wedge T$ is a closed positive current by definition. For every point $x \in X$, we get that

$$
\nu(\alpha \wedge T, x) = \int \mathbb{1}_{\{x\}} (dd^c \log ||z||)^{n-p-k} \wedge \alpha \wedge T
$$

$$
= \int \alpha \wedge \mathbb{1}_{\{x\}} (dd^c \log ||z||)^{n-p-k} \wedge T = 0
$$

where z denotes coordinates of X with $z(x) = 0$. The last equation follows from that a closed positive (s, s) -current with support in an analytic set of codimension strictly greater than s vanishes.

Therefore, if α is real (not necessary strongly positive), we may trivially extend the definition of Lelong numbers to currents as $\alpha \wedge T$, and their linear combinations. In particular, this defines the Lelong number for every closed current T with analytic singularities. We get

$$
\nu(T, x) = \nu \Big(\sum_{i} c_{i} \cdot \mathbb{1}_{W_{i, p+1}} dd^{c} u_{i, p} \wedge \mathbb{1}_{W_{i, p}} \cdots \wedge \mathbb{1}_{W_{i, 2}} dd^{c} u_{i, 1}, x \Big)
$$

=
$$
\sum_{i} c_{i} \cdot \nu \big(\mathbb{1}_{W_{i, p+1}} dd^{c} u_{i, p} \wedge \mathbb{1}_{W_{i, p}} \cdots \wedge \mathbb{1}_{W_{i, 2}} dd^{c} u_{i, 1}, x \big)
$$

for constants c_i , psh functions $u_{i,1},...,u_{i,p}$ with analytic singularities and (suitable) constructible sets $W_{i,2},...,W_{i,p+1}$ (defined in a neighbourhood of x). This is clear since the Lelong numbers of other terms in the local form of T given by (4.2) vanish.

The following is a special variant of Cor. 7.9 in [\[Dem12,](#page-32-11) Chp. III] inferred from Demailly's second comparison theorem.

Proposition 5.8. Let X be a complex manifold of dimension n, let T be a closed quasipos (p, p) -current with analytic singularities for $p < n$, and let q be a quasipsh function with analytic singularities in Z. For every point $x \in X$, we get

(5.5)
$$
\nu(dd^cq\wedge 1_{Z^c}T,x)\geq \nu(q,x)\cdot\nu(1_{Z^c}T,x).
$$

If η is a closed quasipos (1,1)-current with analytic singularities in isolated points, then

(5.6)
$$
\nu([dd^c q]_\eta \wedge T, x) \ge \min{\{\nu(q, x), \nu(\eta, x)\}} \cdot \nu(T, x).
$$

Remark 5.9. The cutoff $\mathbb{1}_{Z^c}$ in the RHS of [\(5.5\)](#page-22-2) cannot be omitted as the following example shows. Let f be holomorphic, let σ be smooth, and set $q := \log |f|^2 + \sigma$. Then

$$
\nu((dd^c q)_{AW}^2, 0) = \nu(dd^c q \wedge \mathbb{1}_{Z^c} dd^c q, 0) = \nu(\left([f = 0] + dd^c \sigma\right) \wedge dd^c \sigma, 0) = 0
$$

while $\nu(dd^c q,0) = \nu([f = 0],0) \neq 0.$

Proof of [Proposition 5.8.](#page-22-0) The proof follows Demailly's strategy adapted to our setting.

Without loss of generality, we may assume that T is positive since the Lelong number of each non-positive term in the local form of T given by (4.2) vanishes, see [Remark/Def. 5.7.](#page-21-1)

Let us assume that we work on a coordinate patch of X with $x = 0$ and z denoting the coordinates, and that $q = u + \sigma$ for a psh u with analytic singularities and smooth function σ . Due to the homogeneity of the Lelong number, we may assume that $\nu(u, 0) =$ $\nu(q, 0) = 1 - \varepsilon$ by rescaling q, and to prove [\(5.5\)](#page-22-2), it is enough to show

(5.7)
$$
\nu(\mathbb{1}_{Z^c}T,0) \leq \nu(dd^c u \wedge \mathbb{1}_{Z^c}T,0) = \nu(dd^c q \wedge \mathbb{1}_{Z^c}T,0).
$$

We set $v := \log ||z||$ and $u_{\kappa} := \max\{u, v - \kappa\} = \max\{u-v, -\kappa\} + v$. Since $\nu(u, 0) < \nu(v, 0)$, there is an $r_0 > 0$ such that $v - \kappa \geq u$, i.e. $u_{\kappa} = v - \kappa$ on $\{|z| < r_0\}$. In particular, $\nu(u_{\kappa}, 0) = \nu(v, 0) = 1$ and

$$
\nu(dd^cu_\kappa\wedge T,0)=\nu(dd^cv\wedge T,0),
$$

whereby the wedge product is well-defined due to BTD. On the other side, $u_{\kappa} \searrow u$ pointwise as $\kappa \to \infty$ such that by [Lemma 4.11,](#page-17-0) we get the weak convergence

$$
dd^c u_{\kappa} \wedge T \stackrel{\kappa \longrightarrow \infty}{\longrightarrow} dd^c u \wedge \mathbb{1}_{Z^c} T + dd^c v \wedge \mathbb{1}_Z T.
$$

As the Lelong number is upper semicontinuous in its first argument (see [Proposition 5.4\)](#page-19-0), we get

$$
\nu(dd^cv\wedge T,0)=\nu(dd^cu_\kappa\wedge T,0)\leq\nu(dd^cu\wedge\mathbb{1}_{Z^c}T,0)+\nu(dd^cv\wedge\mathbb{1}_Z T,0).
$$

By subtracting $\nu(dd^c v\wedge \mathbb{1}_Z T,0)$ on both sides, we get [\(5.7\)](#page-22-3) since $\nu(\mathbb{1}_{Z^c}T,0)=\nu(dd^c v\wedge$ $\mathbb{1}_{Z^c}T$, 0), see [\(5.1\)](#page-19-1).

As $[dd^c q]_\eta \wedge T = dd^c q \wedge \mathbb{1}_{Z^c} T + \eta \wedge \mathbb{1}_Z T$ and the Lelong number is linear (in its first argument), [\(5.6\)](#page-22-4) follows from [\(5.5\)](#page-22-2) and

$$
\nu(\eta \wedge \mathbb{1}_Z T, 0) \ge \nu(\eta, 0) \cdot \nu(\mathbb{1}_Z T, 0).
$$

which also follows from [\(5.5\)](#page-22-2) since $\nu(\mathbb{1}_{X\setminus0}\mathbb{1}_ZT,0) = \nu(\mathbb{1}_ZT,0)$ as $p < n$.

Combining [Proposition 5.5](#page-20-2) with [Proposition 5.8,](#page-22-0) we get the following corollary.

Corollary 5.10. Let X be a complex manifold of dimension n, and let $\pi: X \rightarrow Y$ be a proper holomorphic submersion with m-dimensional fibres. Let T, θ and η be closed positive currents with analytic singularities on X, such that T is of bidegree (p, p) , θ and η are of bidegree (1, 1) and in the same $\partial \overline{\partial}$ -class, $L(\eta)$ is a set of isolated points. Then, for all k with $m \leq k + p \leq n$, and for all $x \in X$,

(5.8)
$$
\nu\big(\pi_*\big([\theta]_\eta^k \wedge T\big), \pi(x)\big) \ge \min\{\nu(\theta, x), \nu(\eta, x)\}^k \cdot \nu(T, x).
$$

Remark 5.11. In general, [\(5.8\)](#page-23-3) cannot hold if we consider quasipos θ which are not positive near $\pi^{-1}(\pi(x))$. This is obvious since the RHS in [\(5.8\)](#page-23-3) is always positive for such θ while the LHS can be negative as in the following simple example.

Let Y be $\mathbb C$ with coordinate z, let F be $\mathbb P^1$ with the coordinates $[\xi_1 : \xi_2]$, let X be $\mathbb C \times \mathbb P^1$, let $\pi: X \to Y$ be the projection on the first component, and let ω_{FS} be the pullback to X of the Fubini-Study form on \mathbb{P}^1 . Then, $\theta = [z = 0] - \omega_{FS}$ is quasipos and $\pi_*([\theta]_{{\eta}}^2) =$ $\pi_*([\theta]^2) = \pi_*(-\omega_{\text{FS}}) \wedge [z=0] = -[z=0]$ (for any η) such that $\nu(\pi_*([\theta]^2), 0) = -1$.

Remark 5.12. If $\alpha_1, \alpha_2 \in \{\theta\}_{\partial \overline{\partial}}$ are smooth, then $\nu(\pi_*(\theta_{\alpha_1}^k \wedge T), y) = \nu(\pi_*(\theta_{\alpha_2}^k \wedge T), y)$ following the argumentation in the proof of [\[LRSW22,](#page-32-7) Thm 1.1 (3)]. As the following counter example shows, this is not correct if one of them has neat analytic singularities in isolated points.

Let Y be $\mathbb C$ with the coordinate z, let X be $\mathbb C \times \mathbb P^1$ with the coordinate $[\xi_1 : \xi_2]$, and let $\pi: X \to Y$ be the projection on the first component. Consider the current $\theta = [\xi_2 = 0]$ which has analytic singularities in $Z = \{\xi_2 = 0\}, \eta := \frac{1}{2}dd^c \log (|z|^2 |\xi_1|^2 + |\xi_2|^2)$ which has only one neat analytic singularity in $(0, [1:0])$, and $\alpha = \omega_{\text{FS}} = \frac{1}{2}$ $\frac{1}{2}dd^c \log (|\xi_1|^2 + |\xi_2|^2)$ the Fubini-Study form on \mathbb{P}^1 . Then, $\eta - \alpha = dd^c \log \frac{|z|^2 |\xi_1|^2 + |\xi_2|^2}{|\xi_1|^2 + |\xi_2|^2}$ $\frac{|2|\xi_1|^2+|\xi_2|^2}{|\xi_1|^2+|\xi_2|^2}$. For any ball $B \subset \mathbb{C}$, we get that

$$
\int_{B} \pi_{*}([\theta]_{\eta}^{2}) - \int_{B} \pi_{*}([\theta]_{\alpha}^{2}) = \int_{B} \pi_{*}((\eta - \alpha) \wedge \mathbb{1}_{Z}[\xi_{2} = 0]) = \int_{B \times \{[1:0]\}} (\eta - \alpha) = \int_{B} dd^{c} \log |z|.
$$

Therefore, $\nu(\pi_{*}([\theta]_{\eta}^{2}), 0) - \nu(\pi_{*}([\theta]_{\alpha}^{2}), 0) = \nu(\log |z|, 0) = 1.$

At least, we obtain that the Lelong numbers of currents $\pi_*(\theta|_{\eta} \wedge T)$ are independent of η among all quasipos (1, 1)-currents with the same type of singularities as follows.

Proposition 5.13. Let $\pi: X \to Y$ be a proper holomorphic submersion, let T and θ be closed quasipos current with analytic singularities of bidegree (p, p) and $(1, 1)$, resp. Let η_1, η_2 be two quasipos (1, 1)-currents with neat analytic singularities in isolated points both in the same $\partial \overline{\partial}$ -class as θ such that $\nu(\eta_1, x) = \nu(\eta_2, x)$ for all $x \in X$. Then, for all $y \in Y$, we get

$$
\nu(\pi_*([\theta]^k_{\eta_1}\wedge T),y)=\nu(\pi_*([\theta]^k_{\eta_2}\wedge T),y).
$$

Proof. As η_1 and η_2 are in the same $\partial\overline{\partial}$ -class, we get that $\eta_1 - \eta_2 = dd^c q$ for a function q which is locally the difference of two quasipsh q_1 and q_2 function with neat analytic singularities in isolated points (given by $\eta_1 = dd^c q_1$ and $\eta_2 = dd^c q_2$). As $\nu(q_1, x) = \nu(q_2, x)$, we obtain that $q_1 - q_2 = q$ is smooth. Now, we can repeat the argumentation in the proof of [\[LRSW22,](#page-32-7) Thm 1.1 (3)] to prove the claimed. \square

6. Proofs of the main results

Before proving the second main result of the present work, let us observe the following.

Lemma 6.1. Let (X, ω) be a Kähler manifold. For each relatively compact $U \subset X$, there exists a constant $\delta = \delta(\omega)$ such that for every $x \in U$ and positive $\tilde{\delta} \leq \delta$, there is a closed positive (1, 1)-current $\eta_x \in {\omega}$ with neat analytic singularities only in x and $\nu(\eta_x, x) = \tilde{\delta}$. If $X \subset B \times \mathbb{P}^m$ for $B \subset \mathbb{C}^n$ and if ω is given as restriction of the product of the Euclidean and the Fubini-Study metric, then we can take $\delta = 1$.

Proof. Set $n := \dim X$. For every $x \in U$, let $B \subset U$ be a neighbourhood of x with coordinates $z = (z_1, \ldots, z_n)$ such that $B \subset \mathbb{C}^n$ with $z(x) = 0$, and let χ be a cutoff function on B which is identically 1 near x. The function $q_x(z) := \chi(z) \log ||z||$ is a quasipsh function on B which extends trivially to U. There is a constant $\delta_x > 0$ such that $\delta_x dd^c q_x + \omega \geq 0$. We can choose δ_x continuously depending of x. As U is relatively compact in X, we get $\delta := \min_{x \in U} \delta_x > 0$. Then, $\eta_x := \tilde{\delta} dd^c q_x + \omega$ has the claimed properties for all $\tilde{\delta} \leq \delta$.

For $X = B \times \mathbb{P}^m$ with the coordinates z' for B and $[\xi_1 : \ldots : \xi_{m+1}]$ for \mathbb{P}^m , we may assume that $x = (0, [1:0:\dots:0])$. Then, $u_x(z', [\xi]) = \frac{1}{2} \log (||z'||^2 |\xi_1|^2 + |\xi_2|^2 + \dots + |\xi_{m+1}|^2) + ||z'||^2$ is psh. So, $dd^c u_x$ is a well-defined positive current on X, in the same cohomology class as $ω$, and $ν(dd^cu_x, x) = 1$. In particular, $η_x = δdd^cu_x + (1 - δ)ω$ has the claimed properties.

For $X \subset B \times \mathbb{P}^m$, we set $\tilde{u}_x = (\text{pr}_2^* u_x)|_X$ where $\text{pr}_2: B \times \mathbb{P}^m \to \mathbb{P}^m$ denotes the projection on \mathbb{P}^m . Then, $\eta_x = \tilde{\delta}\mu \cdot dd^c \tilde{u}_x + (1 - \tilde{\delta}\mu)\omega$ has the claimed properties for $\mu =$ $\nu(u_x, x)/\nu(\tilde{u}_x, x) \le 1$ (for the last inequality, see [Lemma 5.1](#page-19-2) and [Lemma 5.2\)](#page-19-3).

Theorem [1.3.](#page-3-0) Let $\pi: X \to Y$ be a proper holomorphic submersion between complex manifolds X and Y with m-dimensional fibres. Fix a point $y \in Y$. Let $\theta_1, \ldots, \theta_t$ be positive $(1, 1)$ -currents with analytic singularities on X such that each θ_i is in a Kähler class on a neighbourhood of $\pi^{-1}(y)$. Then, there exist positive constants δ_i , $i = 1,...,t$, (which only depend of the Kähler class represented by θ_i on a neighbourhood of $\pi^{-1}(y)$) such that the following is correct.

(i) Existence of η_i : For every $x \in \pi^{-1}(y)$, there exist closed quasipos (1,1)-currents $\eta_i = \eta_{i,x} \in \{\theta_i\}_{\partial \overline{\partial}},$ positive near $\pi^{-1}(y)$, with neat analytic singularities only in $\{x\}$ (and with $\nu(\eta_i, x) = \delta_i$) such that

(6.1)
$$
\nu\Big(\pi_*\big([\theta_t]_{\eta_t}^{k_t} \wedge \cdots \wedge [\theta_1]_{\eta_1}^{k_1}\big),y\Big) \geq \prod_{i=1}^t \min\{\nu(\theta_i,x),\delta_i\}^{k_i}
$$

for all $m \leq k_1 + \ldots + k_t \leq \dim X$. Thereby, $\eta_i = \eta_{i,x}$ depends only of the Kähler class of θ_i near $\pi^{-1}(y)$ and x.

(ii) Independency of η_i : For $i = 1, \ldots, t$, let $\eta_i \in {\theta_i}_{\partial \overline{\partial}}$ be closed quasipos $(1, 1)$ -currents with neat analytic singularities in isolated points on X such that there is a point $x \in X$ with $L(\eta_i) \cap \pi^{-1}(y) = \{x\}$ and $\nu(\eta_i, x) \leq \delta_i$. Then,

(6.2)
$$
\nu\Big(\pi_*\big([\theta_t]_{\eta_t}^{k_t} \wedge \cdots \wedge [\theta_1]_{\eta_1}^{k_1}\big),y\Big) \geq \prod_{i=1}^t \min\{\nu(\theta_i,x),\nu(\eta_i,x)\}^{k_i}
$$

for all $m \leq k_1 + \ldots + k_t \leq \dim X$.

Proof. By the assumptions, there are Kähler forms ω_i on $U = \pi^{-1}(V)$ such that θ_i and ω_i are in the same cohomology class on U . Therefore, [Lemma 6.1](#page-24-0) gives us positive constants δ_i and, for every $x \in \pi^{-1}(y)$, closed positive $(1, 1)$ -currents $\tilde{\eta}_i \in {\{\omega_i\}}_{\partial \overline{\partial}} = {\{\theta_i\}}_{\partial \overline{\partial}}$ with neat analytic singularities only in x on U such that $\nu(\tilde{\eta}_i, x) = \delta_i$. In particular, δ_i and $\tilde{\eta}_i$ only depend on the cohomology class represented by ω_i .

Let us extend $\tilde{\eta}_i$ to the whole X. Pick a closed real (1, 1)-form α_i on X in the same $\partial\partial$ -class as θ_i . In particular, there is a quasipsh function q_i with neat analytic singularities in isolated points on U such that $dd^c q_i = \tilde{\eta}_i - \alpha_i$. Let χ be a cutoff function with support on V and identically 1 in a neighbourhood of y. We define $\eta_i := dd^c(\chi \circ \pi \cdot q_i) + \alpha_i$ which is in the same $\partial \overline{\partial}$ -class as θ_i and $\eta_i = \tilde{\eta}_i$ on a neighbourhood of $\pi^{-1}(y)$. Furthermore, η_i is positive there and $\nu(\eta_i, x) = \delta_i$.

Following [Definition 4.3,](#page-14-3) we get that $T := [\theta_1]_{\eta_t}^{k_t} \wedge \cdots \wedge [\theta_1]_{\eta_1}^{k_1}$ is a closed quasipos current with analytic singularities which is positive in a neighbourhood of $\pi^{-1}(y)$. In particular, [Proposition 5.5](#page-20-2) implies $\nu(\pi_*T, y) \geq \nu(T, x)$ as long $m \leq k_1 + \ldots + k_t \leq n$. By [Proposition 5.8,](#page-22-0) we get that $\nu(T, x) \ge \prod_i \min\{\nu(\theta_i, x), \delta\}^{k_i}$. This proves (i).

(ii) is now a direct consequence of [Proposition 5.13](#page-23-1) (recursively applied): We can exchange the quasipositive η_i in the current on the LHS of [\(6.2\)](#page-25-0) by a positive $\tilde{\eta}_i$ with the same Lelong number in x which is given by [Lemma 6.1.](#page-24-0)

Let us continue with the proof of the first main result of the present work.

Theorem [1.1.](#page-1-2) Let $\pi: X \to Y$ be a proper holomorphic submersion between complex manifolds X and Y with m-dimensional fibres. Fix a point $y \in Y$. Let $\theta_1, \ldots, \theta_t$ be positive $(1,1)$ -currents on X such that each θ_i is in a Kähler class on a neighbourhood of $\pi^{-1}(y)$. Then, there exist positive constants δ_i , for $i = 1, \ldots, t$, (which only depend of the Kähler class represented by θ_i on a neighbourhood of $\pi^{-1}(y)$) such that the following statements are correct.

(i) If the union of all images $\pi(L(\theta_i))$ of the unbounded loci of local dd^c-potentials of θ_i is contained in an analytic A with codim $A \geq k_1 + \cdots + k_t - m$, then

(6.3)
$$
\nu\left(\pi_*\left([\theta_t]^{k_t} \wedge \cdots \wedge [\theta_1]^{k_1}\right), y\right) \geq \prod_{i=1}^t \min\{\nu(\theta_i, x), \delta_i\}^{k_i}
$$

for all points $x \in \pi^{-1}(y)$ as long the LHS does not vanish due to the degree of the current. (ii) If $\bigcup_i \pi(L(\theta_i))$ is contained in an analytic A with codim $A \geq k_1 + \cdots + k_t - t \cdot m$, then

(6.4)
$$
\nu\left(\pi_*\left([\theta_t]^{k_t}\right)\wedge\cdots\wedge\pi_*\left([\theta_1]^{k_1}\right),y\right)\geq \prod_{i=1}^t \min\{\nu(\theta_i,x),\delta_i\}^{k_i}
$$

for all points $x \in \pi^{-1}(y)$ as long the LHS does not vanish due to the degree of the current.

Proof. Since the statement is local with respect to neighbourhoods of $y \in Y$, and since θ_i is in a Kähler class near $\pi^{-1}(y)$ by assumption, we may assume that X is Kähler by shrinking Y and $X = \pi^{-1}(Y)$. Let ω denote a Kähler form on X. By [Lemma 6.1,](#page-24-0) there are constants $\delta_i > 0$ and for each $x \in \pi^{-1}(y)$, closed positive $(1, 1)$ -currents $\eta_i = \eta_{i,x} \in {\{\tilde{\theta}_i\}}_{\partial \overline{\partial}}$ with neat analytic singularities only in x .

(i) By shrinking Y and $X = \pi^{-1}(Y)$ again, Demailly's regularization result (see Theorem 2.10) implies that there are positive constants ε_{κ} and closed $\varepsilon_{\kappa}\omega$ -positive (1, 1)currents $\theta_{i,k} \in {\theta_i}_{\partial \overline{\partial}}$ with analytic singularities such that $\theta_{i,k}$ is decreasing pointwise to θ_i , $\nu(\theta_{i,\kappa},x) \to \nu(\theta_i,x)$ and $\varepsilon_{\kappa} \to 0$ as $\kappa \to \infty$. We set $\tilde{\theta}_{i,\kappa} := \theta_{i,\kappa} + \varepsilon_{\kappa} \omega$ and $\tilde{\eta}_{i,\kappa} := \eta_i + \varepsilon_\kappa \omega \in \{ \tilde{\theta}_{i,\kappa} \}_{\partial \overline{\partial}}$. As $\tilde{\theta}_{i,\kappa}$ are positive, we can apply [Theorem 1.3](#page-3-0) such that

(6.5)
$$
\nu\Big(\pi_*\big([\tilde{\theta}_{t,\kappa}]_{\tilde{\eta}_{t,\kappa}}^{k_t} \wedge \cdots \wedge [\tilde{\theta}_{1,\kappa}]_{\tilde{\eta}_{1,\kappa}}^{k_1}\big),y\Big) \ge \prod_{i=1}^t \min\{\nu(\tilde{\theta}_{i,\kappa},x),\nu(\tilde{\eta}_{i,\kappa},x)\}^{k_i} = \prod_{i=1}^t \min\{\nu(\theta_{i,\kappa},x),\delta_i\}^{k_i}.
$$

By the Theorems [3.5](#page-10-0) and [4.7,](#page-16-0) we obtain that

$$
\pi_*\big([\theta_t]^{k_t} \wedge \cdots \wedge [\theta_1]^{k_1}\big) = \lim_{\kappa \to \infty} \pi_*\big([\theta_{t,\kappa}]^{k_t}_{\eta_t} \wedge \cdots \wedge [\theta_{1,\kappa}]^{k_1}_{\eta_1}\big).
$$

Since $[\tilde{\theta}_{i,\kappa}]_{\tilde{\eta}_{i,\kappa}} \wedge \cdot = ([\theta_{i,\kappa}]_{\eta_i} + \varepsilon_{\kappa} \omega) \wedge \cdot$, we get

$$
\begin{aligned} [\tilde{\theta}_{i,\kappa}]_{\tilde{\eta}_{i,\kappa}}^{k_i} \wedge T &= \sum_{\ell=0}^{k_i} (-1)^{\ell} \binom{k_i}{\ell} [\theta_{i,\kappa}]_{\eta_i}^{k_i-\ell} \wedge \varepsilon_{\kappa}^{\ell} \omega^{\ell} \wedge T \\ &= [\theta_{i,\kappa}]_{\eta_i}^{k_i} \wedge T + \sum_{\ell=1}^{k_i} (-1)^{\ell} \binom{k_i}{\ell} [\theta_{i,\kappa}]_{\eta_i}^{k_i-\ell} \wedge \varepsilon_{\kappa}^{\ell} \omega^{\ell} \wedge T \end{aligned}
$$

for any closed current T with analytic singularities. By applying this recursively to each factor in $\left[\tilde{\theta}_{t,\kappa}\right]_{\tilde{n}_{t,\kappa}}^{k_{t-1}}$ $_{\tilde{\eta}_{t,\kappa}}^{k_{t-1}} \wedge \cdots \wedge [\tilde{\theta}_{1,\kappa}]_{\tilde{\eta}_1}^{k_1}$ $_{\tilde{\eta}_{1,\kappa}}^{\kappa_1}$, we get

$$
\pi_*\big([\tilde{\theta}_{t,\kappa}]_{\tilde{\eta}_{t,\kappa}}^{k_t} \wedge \cdots \wedge [\tilde{\theta}_{1,\kappa}]_{\tilde{\eta}_{1,\kappa}}^{k_1}\big) = \pi_*\big([\theta_{t,\kappa}]_{\eta_t}^{k_t} \wedge \cdots \wedge [\theta_{1,\kappa}]_{\eta_1}^{k_1}\big) + \sum_j \varepsilon_{\kappa}^{\ell_j} T_{j,\kappa}
$$

for positive integers $\ell_j > 0$ and currents $T_{j,k}$ which are pushforwards of quasipos currents with analytic singularities. Moreover, we can apply [Theorem 3.5](#page-10-0) for each term and get

$$
\pi_*\big([\theta_t]^{k_t} \wedge \cdots \wedge [\theta_1]^{k_1}\big) = \lim_{\kappa \to \infty} \pi_*\big([\tilde{\theta}_{t,\kappa}]^{k_t}_{\tilde{\eta}_{t,\kappa}} \wedge \cdots \wedge [\tilde{\theta}_{1,\kappa}]^{k_1}_{\tilde{\eta}_{1,\kappa}}\big)
$$

.

As all currents are positive, we can estimate the Lelong number of the LHS from below with the limit superior of the Lelong number of the RHS, see [Proposition 5.4.](#page-19-0) This and (6.5) imply

$$
\nu\Big(\pi_*\big([\theta_t]^{k_t}\wedge\cdots\wedge[\theta_1]^{k_1}\big),y\Big)\geq \lim_{\kappa\to\infty}\prod_{i=1}^t\min\{\nu(\theta_{i,\kappa},x),\delta_i\}^{k_i}=\prod_{i=1}^t\min\{\nu(\theta_i,x),\delta_i\}^{k_i}
$$

where the last equation follows from the choice of $\theta_{i,\kappa}$, cf. [Theorem 2.10](#page-7-0) (ii).

(ii) We prove [\(6.4\)](#page-25-1) by induction over t. The case $t = 1$ follows from (i). For the induction step, we will prove

(6.6)
$$
\nu\big(\pi_*\big([\theta_t]^{k_t}\big) \wedge T, y\big) \geq \min\{\nu(\theta_t, x), \delta_t\}^{k_t} \cdot \nu(T, y)
$$

for the closed positive current $T := \pi_*([\theta_{t-1}]^{k_{t-1}}) \wedge \cdots \wedge \pi_*([\theta_1]^{k_1}).$

Pick a closed real (1, 1)-form $\alpha \in {\theta_t}_{\partial \overline{\partial}} = {\eta_t}_{\partial \overline{\partial}}$. There are α -psh functions q and v (the latter with neat analytic singularities only in x) such that $\theta_t = dd^c q + \alpha$ and $\eta_t = dd^c v + \alpha$. We set $q^{(\lambda)} := \max\{q, \frac{\delta}{\delta_t} v - \lambda\}$ with $\delta < \nu(q, x) = \nu(\theta_t, x)$ and $\delta \leq \delta_t$. We

get $q^{(\lambda)}$ is α -psh and decreasing pointwise to q as $\lambda \to \infty$. As in the proof of [Proposition 5.8,](#page-22-0) $q^{(\lambda)}$ has neat analytic singularities only in x and $\nu(q^{(\lambda)}, x) = \delta$. We set $\theta_t^{(\lambda)} = dd^c q^{(\lambda)}$ and get

$$
\pi_*\big([\theta_t]^{k_t}\big) \wedge T = \lim_{\lambda \to \infty} \pi_*\Big([\theta_t^{(\lambda)}]^{k_t}\Big) \wedge T = \lim_{\lambda \to \infty} \pi_*\Big((\theta_t^{(\lambda)})^{k_t} \wedge \pi^*T\Big)
$$

where the first equation follows from [Theorem 3.5](#page-10-0) and the second from [Proposition 3.1](#page-8-1) and BTD (by selecting a smooth approximation of $\theta_t^{(\lambda)}$ $t^{(\lambda)}$ for each λ). Semicontinuity of the Lelong number, see [Proposition 5.4,](#page-19-0) implies

$$
\nu(\pi_*([\theta_t]^{k_t}) \wedge T, y) \geq \limsup_{\lambda \to \infty} \nu\Big(\pi_*\Big((\theta_t^{(\lambda)})^{k_t} \wedge \pi^*T\Big), y\Big).
$$

By the Propositions [5.5](#page-20-2) and [5.6,](#page-21-2) we get

$$
\nu\Big(\pi_*\Big(\big(\theta_t^{(\lambda)}\big)^{k_t}\wedge \pi^*T\Big),y\Big)\geq \nu\big(\big(\theta_t^{(\lambda)}\big),x\big)^{k_t}\cdot \nu(\pi^*T,x)\geq \delta^{k_t}\cdot \nu(T,y).
$$

Since this holds for all $\delta \leq \delta_t$ with $\delta < \nu(\theta_t, x)$, [\(6.6\)](#page-26-1) is proven.

7. Segre currents

In [\[LRRS18\]](#page-32-6), Lärkäng, Raufi, Ruppenthal and the author define so-called Chern and Segre currents for singular Hermitian metrics on vector bundles which are positively curved in the sense of Griffiths³. These currents represent the Chern and Segre classes of the vector bundles. In this section, we are going to present the construction of these currents in a (slightly) more general setting.

Setting 7.1. Let X be a complex manifold of dimension n , let E be a holomorphic vector bundle on X of rank r, let $\pi: \mathbb{P}(E) \to X$ denote the projectivization of hyperplanes of E (i.e. $\pi^{-1}(x) = \mathbb{P}(E_x^*)$ for all $x \in X$), let $L := \mathcal{O}_{\mathbb{P}(E)}(1) \to \mathbb{P}(E)$ denote the hyperplane bundle of E, and let $e^{-\varphi}$ be a singular metric on L such that its curvature current $\theta := dd^c \varphi$ is quasipositive, i.e. the weights φ are quasipsh functions. Let $L(\varphi)$ denote the unbounded locus of φ , i. e. $L(\varphi) = L(\theta)$. There is a (pos) (1, 1)-form γ on $\mathbb{P}(E)$ such that $\theta + \gamma \geq 0$, i. e. θ is γ-pos. For any small enough open set $U \subset X$, we may pick γ such that γ is closed and positive on $\pi^{-1}(U)$, see [Remark 3.2.](#page-8-0) Let $e^{-\varphi_{\kappa}}$ be a sequence of smooth/singular metrics on L such that φ_{κ} is decreasing pointwise to φ and $dd^c \varphi_{\kappa}$ is γ -positive. Let T be a closed real (p, p) -current on X which is locally given as difference of two closed positive currents.

If H is a singular Hermitian metric on E, then it induces a singular metric $e^{-\psi}$ on L whose dual metric is given by $e^{\psi} = \pi^* H^*|_{L^*}$ since $L^* = \mathcal{O}_{\mathbb{P}(E)}(-1)$ is a subbundle of π^*E^* . $e^{-\psi}$ has a quite special geometric behaviour on each fibre $\pi^{-1}(x)$ coming from the Hermitian structure of H_x which turns out to be unnecessary for the definition of Chern and Segre currents. To keep the setting more general, we consider any singular metric $e^{-\varphi}$ on L . There is a one-to-one correspondence between singular metrics on L and singular metrics on E^* considering so-called singular Finsler metrics⁴.

 3_{as} defined by [\[BP08,](#page-31-7) Def. 3.2]; also called singularly (semi-) positive in the sense of Griffiths

⁴A singular Finsler metric h on E^* is given by $E^* \to [0, \infty]$, $(p, \xi) \mapsto ||\xi||_{h(p)}$ with $||\lambda \xi||_{h(p)} = |\lambda| \cdot ||\xi||_{h(p)}$.

If the weights of φ are psh, i.e. $dd^c \varphi$ is positive, then $L = \mathcal{O}_{\mathbb{P}(E)}(1)$ is pseudoeffective, and so is E by definition⁵. Moreover, we get that E is strongly pseudoeffective if $\pi(L(\varphi))$ does not equal X by definition, see [\[BDPP13,](#page-31-5) Def. 7.1]. In particular, the following results give Segre and Chern currents for strongly pseudoeffective vector bundles and not only vector bundles positively curved in the sense of Griffiths. Let us stress that we work in the even more general [Setting 7.1](#page-27-1) where the weights φ are just supposed to be quasipsh.

As in [\[LRRS18,](#page-32-6) Prop. 4.6], [Proposition 3.1](#page-8-1) implies the following result.

Theorem 7.2. In the [Setting 7.1](#page-27-1) with φ_{κ} smooth, if $\pi(L(\varphi))$ is contained in an analytic set of codimension $\geq k+p$, then

$$
(7.1) \ s_k(E,\varphi) \wedge T := (-1)^k \pi_* \left([dd^c \varphi]^{k+r-1} \wedge \pi^* T \right) \stackrel{\text{def}}{=} (-1)^k \lim_{\kappa \to \infty} \pi_* \left((dd^c \varphi_\kappa)^{k+r-1} \wedge \pi^* T \right)
$$

is a well-defined closed real current which is locally the difference of two closed positive currents, and independent of the choice of the smooth approximation φ_{κ} .

For a smooth Hermitian metric H on E, the total Segre form $s(E, H)$ is (classically) defined as the inverse of the total Chern form $c(E, H) = 1 + c_1(E, H) + \ldots + c_n(E, H)$. Then, its (k, k) -component $s_k(E, H)$ can be calculated using the projectivization of the vector bundle, $s_k(E, H) = (-1)^k \pi_*(dd^c \psi)^{k+r-1}$ where $e^{\psi} = \pi^* H^*|_{L^*}$. Therefore, the Segre currents $s_k(E, \varphi)$ defined above are naturally extending the concept of Segre forms to singular metrics on $\mathcal{O}_{\mathbb{P}(E)}(1)$. Moreover, the so-called Chern currents

(7.2)
$$
c_k(E,\varphi) := (-1)^t \sum_{k_1+\ldots+k_t=k} s_{k_t}(E,\varphi) \wedge \cdots \wedge s_{k_1}(E,\varphi)
$$

given by a recursive application of [Theorem 7.2](#page-28-0) are naturally extending the concept of Chern forms.

[Corollary 3.6](#page-12-0) implies the following monotone continuity result generalizing [\[LRRS18,](#page-32-6) Thm 1.5].

Theorem 7.3. In the [Setting 7.1,](#page-27-1) if $\pi(L(\varphi))$ is contained in an analytic set of codimension $\geq k_1 + \ldots + k_t$, then

$$
s_{k_t}(E,\varphi)\wedge\cdots\wedge s_{k_1}(E,\varphi)=\lim_{\kappa\to\infty}\pi_*\big((dd^c\varphi_\kappa)^{k_t+r-1}\big)\wedge\cdots\wedge\pi_*\big((dd^c\varphi_\kappa)^{k_1+r-1}\big)
$$

for any (not necessarily smooth) approximation φ_{κ} decreasing pointwise to φ .

Remark 7.4. In particular, the currents given by [\[Wu22,](#page-32-10) Thm 2] coincide with the currents defined in [\[LRRS18\]](#page-32-6).

Remark 7.5. [Theorem 7.3](#page-28-2) generalizes [\[LRRS18,](#page-32-6) Thm 1.5] in various ways. Among these generalizations, let us highlight that we do not need to assume that on $X \setminus \pi(L(\varphi))$, φ is continuous and φ_{κ} converges locally uniformly to φ . This is particularly interesting since the well-known calculus for Chern and Segre forms can be also applied to Chern and Segre currents defined by [\(7.1\)](#page-28-3) and [\(7.2\)](#page-28-4) in this general setting, cf. [\[LRRS18,](#page-32-6) Cor. 1.9].

Furthermore, we obtain the following cohomology result.

⁵In the literature, sometimes called weakly to distinguish it from strongly pseudoeffective.

Theorem 7.6. In [Setting 7.1](#page-27-1) with compact X, if $L(\varphi)$ is contained in an analytic set of $codimension \geq k_1 + \ldots + k_t$, then we get

$$
s_{k_t}(E, \varphi) \wedge \cdots \wedge s_{k_1}(E, \varphi) \in s_{k_t}(E) \cdots s_{k_1}(E) \quad and
$$

$$
c_{k_t}(E, \varphi) \wedge \cdots \wedge c_{k_1}(E, \varphi) \in c_{k_t}(E) \cdots c_{k_1}(E).
$$

Proof. The argumentation is exactly the same as in the proof of [\[LRRS18,](#page-32-6) Thm 1.13] by applying Demailly's regularization, see [Theorem 2.9.](#page-7-2)

Remark/Def. 7.7. Let us assume that the metric φ on $\mathcal{O}_{\mathbb{P}(E)}(1)$ has analytic singularities. Using the generalized Monge-Ampère products, the Segre currents

(7.3)
$$
s_k(E,\varphi,\alpha) := (-1)^k \pi_* \left([dd^c \varphi]_{\alpha}^{k+r-1} \right)
$$

and their wedge products $s_{k_t}(E,\varphi,\alpha) \wedge \cdots \wedge s_{k_1}(E,\varphi,\alpha)$ can be defined for arbitrary degrees following [\[LRSW22,](#page-32-7) Thm 1.1]. If α is a closed $(1,1)$ -form representing $\mathcal{O}_{\mathbb{P}(E)}(1)$, we get that these currents represent their corresponding Segre classes. $s_k(E, \varphi, \eta)$ can be defined for η with analytic singularities in isolated points, as well. For every $\xi \in \mathbb{P}(E)$, [Corollary 5.10](#page-23-2) implies

(7.4)
$$
\nu(s_k(E,\varphi,\eta),\pi(\xi)) \ge \min{\{\nu(\varphi,\xi),\nu(\eta,\xi)\}}^{k+r-1}
$$

if $k \leq n$ and if $dd^c \varphi$ and η are positive near $\pi^{-1}(\pi(\xi))$. By [Lemma 6.1,](#page-24-0) we can always find such an $\eta = \eta_{\xi}$ with $\nu(\eta, \xi) = 1$ and neat analytic singularities, see proof of [Theorem 1.3.](#page-3-0)

For wedge products of Segre currents, the situation is more complicated. Let ϖ : $P :=$ $\mathbb{P}(E) \times_X \cdots \times_X \mathbb{P}(E) \to X$ be the t-fibre power of $\pi \colon \mathbb{P}(E) \to X$, and let φ_i denote the pullback metric $pr_i^* \varphi$ on $pr_i^* \mathcal{O}_{\mathbb{P}(E)}(1)$ where $pr_i: P \to \mathbb{P}(E)$ denotes the projection on the *i*th component. For all $k = k_1 + \ldots + k_t$, [\[LRSW22\]](#page-32-7) defines

$$
(7.5) \quad s_{k_t}(E,\varphi,\alpha) \wedge \cdots \wedge s_{k_1}(E,\varphi,\alpha) := (-1)^k \varpi_* \big([dd^c \varphi_t]_{\alpha_t}^{k_t + r - 1} \wedge \cdots \wedge [dd^c \varphi_1]_{\alpha_1}^{k_1 + r - 1} \big)
$$

where $\alpha_i = \text{pr}_i^* \alpha$ [for a closed real \(1](#page-16-3), 1)-form α on $\mathbb{P}(E)$ representing $\mathcal{O}_{\mathbb{P}(E)}(1)$; cf. Remark 4.9. This cannot be extended straightforwardly to η with (neat) analytic singularities in isolated points since the pullbacks of such η have singularities in sets of lower codimension than points and BTD cannot be applied in general. We guess that it might be possible to still define these products as the singularities of $pr_i^* \eta$ are transversal to each others. An alternative approach could be the following. For every $\xi \in \mathbb{P}(E)$, there is (exactly) one $p \in \bigcap_{i=1}^t \operatorname{pr}_i^{-1}(\xi)$. Let η_i be a closed quasipositive $(1, 1)$ -currents with analytic singularities in isolated points including p such that η_i is representing $pr_i^* \mathcal{O}_{\mathbb{P}(E)}(1)$. Then,

$$
S := \varpi_*\big([dd^c \varphi_t]_{\eta_t}^{k_t + r - 1} \wedge \dots \wedge [dd^c \varphi_1]_{\eta_1}^{k_1 + r - 1} \big)
$$

is locally the difference of two closed positive currents and the pushforward of a closed quasipos (k, k) -current with analytic singularities, and $S \in (-1)^k s_{k_1}(E) \cdot \ldots \cdot s_{k_t}(E)$. By [Corollary 5.10,](#page-23-2) we obtain

$$
\nu(S, \pi(\xi)) \ge \min{\{\nu(\varphi, \xi), \nu(\eta_i, p)\}}^{k+tr-r}
$$

if $k \leq n$ and if $dd^c \varphi$ positive near $\pi^{-1}(\pi(\xi))$ and η_i near $\varpi^{-1}(\pi(\xi))$ since $\nu(\varphi, \xi) = \nu(\varphi_i, p)$ by [Lemma 5.2.](#page-19-3) Unfortunately, it is not clear whether such η_i exist as $\mathrm{pr}_i^* \mathcal{O}_{\mathbb{P}(E)}(1)$ is not Kähler near $\varpi^{-1}(\pi(\xi))$ in general.

Let us conclude this section with the proofs of [Corollary 1.6](#page-4-1) and [Theorem 1.7.](#page-4-2)

Corollary [1.6.](#page-4-1) Let X be a compact complex manifold of dimension n, let $E \rightarrow X$ be a (pseudoeffective) holomorphic vector bundle on X, and let $e^{-\varphi}$ be a semipositive singular metric on $\mathcal{O}_{\mathbb{P}(E)}(1)$ such that $\pi(L(\varphi))$ is contained in an analytic set of codimension s. If there are k_1, \ldots, k_t with $k_1 + \ldots + k_t \leq s$ and $s_{k_1}(E) \cdots s_{k_t}(E) = 0$, then E is nef.

Proof. We set $k = k_1 + \ldots + k_t$. Since $S := (-1)^k s_{k_t}(E, \varphi) \wedge \cdots \wedge s_{k_1}(E, \varphi)$ is a closed positive current, and since

$$
S \in (-1)^{k} s_{k_1}(E) \cdots s_{k_t}(E) = 0,
$$

S must vanish. Therefore, $\nu(S, \pi(\xi)) = 0$, and by [Corollary 1.5,](#page-4-0) $\nu(\varphi, \xi) = 0$ for all $\xi \in$ $\mathbb{P}(E)$. Let $\omega_{\mathbb{P}(E)}$ denote the Hermitian form on $\mathbb{P}(E)$ (which is not necessarily closed). For every $\varepsilon > 0$, there is a singular Hermitian metric φ_{ε} of $\mathcal{O}_{\mathbb{P}(E)}(1)$ with analytic singularities such that $dd^c \varphi_{\varepsilon} \geq -2\varepsilon \omega_{\mathbb{P}(E)}$ and $\nu(\varphi_{\varepsilon}, y) \leq \nu(\varphi, y) = 0$, see [\[Dem92,](#page-32-9) Prop. 3.7] or [Theorem 2.10.](#page-7-0) As the Lelong number in an analytic singularity would be positive, we obtain that $L(\varphi_{\varepsilon}) = \emptyset$. By Richberg's approximation [\[Ric68\]](#page-32-16), using the version [\[Dem92,](#page-32-9) Lem. 2.15], we can get a smooth approximation $\tilde{\varphi}_{\varepsilon}$ of φ_{ε} with $dd^c \tilde{\varphi}_{\varepsilon} \geq -\varepsilon \omega_{\mathbb{P}(E)}$. As we get these smooth metrics for all $\varepsilon > 0$, we conclude $\mathcal{O}_{\mathbb{P}(E)}(1)$ is nef, i.e. E is nef. See also $[$ Bou04, Prop. 3.2 (i)].

Theorem [1.7.](#page-4-2) Let (X, ω) be a compact Kähler manifold, and let E be a pseudoeffective vector bundle on X such that $L_{nn}(E)$ is contained in a countable union of analytic sets of codimension s. If there is a $k \leq s$ with $c_1(E)^k = (-s_1(E))^k = 0$, then E is nef.

Proof. As X is Kähler, so is $\mathbb{P}(E)$. Let $\omega_{\mathbb{P}(E)}$ denote its Kähler form, which is in $\mathcal{O}_{\mathbb{P}(E)}(1)$, and let r denote the rank of E .

Fix $\varepsilon > 0$. Let H be a smooth Hermitian metric on E, and let $e^{-\sigma}$ denote the induced metric on $\mathcal{O}_{\mathbb{P}(E)}(1)$, i.e. $e^{\sigma} = \pi^* H|_{\mathcal{O}_{\mathbb{P}(E)}(-1)}$. As $dd^c \sigma$ is positive along fibres of π , there is a small enough δ such that $dd^c \sigma \ge \delta \omega_{\mathbb{P}(E)} - \delta^{-1} \pi^* \omega$. By Demailly's regularization, see [Theorem 2.10,](#page-7-0) there exist a singular Hermitian metric φ_{ε} with analytic singularities on $\mathcal{O}_{\mathbb{P}(E)}(1)$ such that $dd^c \varphi_{\varepsilon} \geq -\delta^2 \varepsilon \omega_{\mathbb{P}(E)}$ and $\delta_{\varepsilon} > 0$ (with $\delta_{\varepsilon} \to 0$ as $\varepsilon \to 0$) such that

$$
\nu(\varphi,\xi)-\delta_{\varepsilon}\leq \nu(\varphi_{\varepsilon},\xi)\leq \nu(\varphi,\xi)\quad \forall \xi\in\mathbb{P}(E).
$$

As in the proof of [\[DPS94,](#page-32-17) Thm 11.12], we can modify φ_{ε} such that $dd^c \varphi_{\varepsilon} \ge -\varepsilon \pi^* \omega$: Replace φ_{ε} by the barycentre of φ_{ε} and σ given by $(1 - \delta \varepsilon) \varphi_{\varepsilon} + \delta \varepsilon \cdot \sigma$ with

$$
dd^{c}((1 - \delta \varepsilon)\varphi_{\varepsilon} + \delta \varepsilon \sigma) \geq (\delta \varepsilon - 1)\delta^{2} \varepsilon \omega_{\mathbb{P}(E)} + \delta \varepsilon (\delta \omega_{\mathbb{P}(E)} - \delta^{-1} \pi^{*} \omega)
$$

= $(\delta^{3} \varepsilon^{2} - \delta^{2} \varepsilon + \delta^{2} \varepsilon) \omega_{\mathbb{P}(E)} - \varepsilon \pi^{*} \omega$
 $\geq -\varepsilon \pi^{*} \omega.$

Since φ_{ε} has analytic singularities, we get that $L(\varphi_{\varepsilon})$ is an analytic set. Moreover, $\xi \in$ $L(\varphi_{\varepsilon})$ if and only if $0 < \nu(\varphi_{\varepsilon}, \xi) \leq \nu(\varphi, \xi)$ such that codim $\pi(L(\varphi_{\varepsilon})) \geq k$ by assumption. This implies

$$
(c_1(E,\varphi_{\varepsilon}))^k = (-s_1(E,\varphi_{\varepsilon}))^k = \left(\pi_*([dd^c \varphi_{\varepsilon}]^r)\right)^k
$$

is a closed real (k, k) -current in $c_1(E)^k$. As calculated in [Remark 3.4,](#page-10-2) we get

$$
\pi_*\big([dd^c \varphi_\varepsilon]^r \big) \wedge T = \big(\pi_*[dd^c \varphi_\varepsilon + \varepsilon \pi^* \omega]^{r-1} - r\varepsilon \omega \big) \wedge T.
$$

Furthermore,

$$
S_{\varepsilon} := \left(\pi_* \left([dd^c \varphi_{\varepsilon} + \varepsilon \pi^* \omega]^r) \right)^k\right)
$$

is a closed positive (k, k) -current in $(c_1(E) + r\varepsilon {\{\omega\}})^k$. So, [Theorem 1.1](#page-1-2) implies

(7.6)
$$
\nu(S_{\varepsilon}, \pi(\xi)) \ge (\nu(dd^c \varphi_{\varepsilon} + \varepsilon \pi^* \omega, \xi))^{kr} = (\nu(\varphi_{\varepsilon}, \xi))^{kr}
$$

for all $\xi \in \mathbb{P}(E)$.

By weak compactness, there is a subsequence ε_{λ} such that $S_{\varepsilon_{\lambda}}$ converges weakly to a closed positive current S in $(c_1(E))^k$. As assumed, this cohomology vanishes, and so does the closed positive S . By [Proposition 5.4,](#page-19-0) we get that

$$
\lim_{\lambda \to \infty} \nu(S_{\varepsilon_\lambda}, x) \le \nu(S, 0) = 0.
$$

This in combination with [\(7.6\)](#page-31-9) and the choice of φ_{ε} implies

$$
\nu(\varphi,\xi) = \lim_{\lambda \to \infty} \nu(\varphi_{\varepsilon_{\lambda}},\xi) = 0.
$$

Repeating the arguments in the proof of [Corollary 1.6](#page-4-1) (or by [\[Bou04,](#page-31-8) Prop. 3.2 (i)]), we obtain that $\mathcal{O}_{\mathbb{P}(E)}(1)$ is nef.

References

- [And05] M. Andersson, *Residues of holomorphic sections and Lelong currents*, Ark. Mat. 43 (2005), no. 2, 201–219.
- [ABW19] M. Andersson, Z. Blocki, and E. Wulcan, On a Monge-Ampère operator for plurisubharmonic functions with analytic singularities, Indiana Univ. Math. J. 68 (2019), no. 4, 1217–1231.
- [ASWY17] M. Andersson, H. Samuelsson Kalm, E. Wulcan, and A. Yger, Segre numbers, a generalized King formula, and local intersections, J. Reine Angew. Math. 728 (2017), 105–136.
	- [AW14] M. Andersson and E. Wulcan, Green functions, Segre numbers, and King's formula, Annales de l'Institut Fourier 64 (2014), no. 6, 2639–2657.
	- [Bin83] J. Bingener, On deformations of Kähler spaces. II., Arch. Math. 41 (1983), 517–530.
	- [Bło19] Z. Błocki, On the complex Monge-Ampère operator for quasi-plurisubharmonic functions with analytic singularities, Bull. Lond. Math. Soc. 51 (2019), no. 3, 431–435.
	- [Bou04] S. Boucksom, Divisorial Zariski decompositions on compact complex manifolds, Ann. Sci. Ec. ´ Norm. Supér. (4) 37 (2004), no. 1, 45-76.
- [BDPP13] S. Boucksom, J.-P. Demailly, M. Păun, and T. Peternell, The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension, J. Algebr. Geom. 22 (2013), no. 2, 201–248.
	- [BP08] B. Berndtsson and M. Păun, Bergman kernels and the pseudoeffectivity of relative canonical bundles, Duke Math. J. 145 (2008), no. 2, 341–378.
- [BT76] E. Bedford and B. A. Taylor, The Dirichlet problem for a complex Monge-Ampère equation, Invent. Math. 37 (1976), no. 1, 1–44.
- [BT82] , A new capacity for plurisubharmonic functions, Acta Math. 149 (1982), no. 1-2, 1-40.
- [Dem82] J.-P. Demailly, Estimations L^2 pour l'opérateur $\overline{\partial}$ d'un fibré vectoriel holomorphe semi-positif au-dessus d'une variété kählérienne complète, Ann. Sci. École Norm. Sup. (4) 15 (1982), no. 3, 457–511.
- [Dem85] , Mesures de Monge-Ampère et caractérisation géométrique des variétés algébriques affines, Mémoires de la Société Mathématique de France, Société mathématique de France, 1985.
- [Dem92] , Regularization of closed positive currents and intersection theory, J. Algebr. Geom. 1 (1992), no. 3, 361–409.
- [Dem93] $_____\$ Monge-Ampère operators, Lelong numbers and intersection theory, In: Complex analysis and geometry, Plenum Press., 1993, pp. 115–193.
- [Dem12] , Complex Analytic and Differential Geometry, 2012. Institut Fourier, Université de Grenoble I. OpenContent AG-Book.
- [DPS94] J.-P. Demailly, T. Peternell, and M. Schneider, Compact complex manifolds with numerically effective tangent bundles, J. Algebr. Geom. 3 (1994), no. 2, 295–345.
- [FG65] W. Fischer and H. Grauert, Lokal-triviale Familien kompakter komplexer Mannigfaltigkeiten, Nachr. Akad. Wiss. Gött., II. Math.-Phys. Kl. 1965 (1965), 89-94.
- [Fuj78] A. Fujiki, Closedness of the Douady spaces of compact Kähler spaces, Publ. Res. Inst. Math. Sci. 14 (1978), 1–52.
- [Lel57] P. Lelong, *Intégration sur un ensemble analytique complexe*, Bulletin de la Société Math-´ematique de France 85 (1957), 239–262.
- [Lel68] , Fonctions plurisousharmoniques et formes différentielles positives, Gordon & Breach, Paris-London-New York (distributed by Dunod Éditeur, Paris), 1968.
- [LRRS18] R. Lärkäng, H. Raufi, J. Ruppenthal, and M. Sera, Chern forms of singular metrics on vector bundles, Adv. Math. 326 (2018), 465–489.
- [LRSW22] R. Lärkäng, H. Raufi, M. Sera, and E. Wulcan, Chern forms of Hermitian metrics with analytic singularities on vector bundles, Indiana Univ. Math. J. 71 (2022), no. 1, 153–189.
- [LSW20] R. Lärkäng, M. Sera, and E. Wulcan, On a mixed Monge-Ampère operator for quasiplurisubharmonic functions with analytic singularities, Bull. Lond. Math. Soc. 52 (2020), no. 1, 77–93.
- [Meo96] M. Meo, Image inverse d'un courant positif fermé par une application analytique surjective, C. R. Acad. Sci. Paris Sér. I Math. 322 (1996), no. 12, 1141-1144.
- [Ric68] R. Richberg, Stetige streng pseudokonvexe Funktionen, Math. Ann. 175 (1968), 257–286.
- [Wu22] X. Wu, Pseudo-effective and numerically flat reflexive sheaves, J. Geom. Anal. 32 (2022), no. 4, article 124.

M. Sera, Faculty of Engineering, Kyoto University of Advanced Science, Kyoto 615-8577, Japan

Email address: sera.martin@kuas.ac.jp