# A more flexible counterpart of a Huang-Kotz's copula-type

H. M. Barakat<sup>\*</sup>, M. A. Alawady<sup>\*</sup>, I. A. Husseiny<sup>\*</sup> and M. A. Abd Elgawad<sup>†</sup>

#### Abstract

We propose a more flexible symmetric counterpart of the Huang-Kotz's copula of the 1st type. Both the counterpart and Huang-Kotz's copula of the 1st type provide the same improvement of the correlation level. Moreover, the proposed copula includes special cases of many other extensions of the Farlie-Gumbel-Morgenstern (FGM) copula.

AMS 2010 Subject Classification: Primary 62H05; Secondary 62H20.

**Key Words:** FGM copula; Huang-Kotz FGM copula; iterated FGM copula; Spearman's Rho; Kendall's Tau.

## 1 Introduction

A bivariate copula is a bivariate distribution function defined on the unit square  $I^2 = [0, 1]^2$ , which has uniform marginals on [0, 1]. Copula is mainly used to describe the dependence between random variables (RVs). Any bivariate function, C(u, v), is a copula if and only if (cf. [20])

$$C(u,0) = C(0,v) = 0, \ C(u,1) = u, \ C(u,v) \le 1, \ C(1,v) = v, \ \forall \ (u,v) \in I^2,$$
(1.1)

and 
$$C(u_2, v_2) - C(u_2, v_1) - C(u_1, v_2) + C(u_1, v_1) \ge 0,$$
 (1.2)

$$\forall (u_i, v_i) \in I^2, i = 1, 2, u_1 \leq u_2, v_1 \leq v_2.$$

The twice differentiable 2-increasing property (1.2) can be replaced by the condition

$$c(u,v) \ge 0,\tag{1.3}$$

where c(u, v) is called copula density.

The FGM copula is defined by

$$C_{FGM}(u,v;a) = uv\{1 + a(1-u)(1-v)\}, \ a \in [-1,1], \ (u,v) \in I^2.$$

$$(1.4)$$

While the copula (1.4) is a flexible family and valuable in many applications, a well-known limitation of this copula is the low dependence level it permits between RVs, Spearman's Rho  $\rho \in [-0.333, 0.333]$  and Kendall's Tau  $\tau \in [-0.222, 0.222]$ . Several extensions to the FGM copula have been introduced in the literature, where most of these extensions aimed to improve the correlation level. Some examples

<sup>\*</sup>Department of Mathematics, Faculty of Science, Zagazig University, Zagazig 44519, Egypt.

<sup>&</sup>lt;sup>†</sup>Department of Mathematics, Faculty of Science, Benha University, Benha 13518, Benha, Egypt. The corresponding author: M. A. Abd Elgawad, E-mail: mohamed.abdelgwad@fsc.bu.edu.eg; mohamed\_salem240@yahoo.com

of these important extensions are [8, 14, 15, 16]. Recently, most of these extensions were studied in different important aspects by Abd Elgawad et al. [2, 3, 4] and Alawady et al. [6, 7].

Huang and Kotz [19] proposed two analogous extensions, the 1st HK-FGM and 2nd HK-FGM types which are defined respectively by

$$C_{HK-FGM1}(u,v;a,b) = uv \left[ 1 + a \left( 1 - u^b \right) \left( 1 - v^b \right) \right], \ b > 0, a \in \left[ -\frac{1}{b^2}, \frac{1}{b} \right],$$
(1.5)

$$C_{HK-FGM2}(u,v;a,b) = uv \left[ 1 + a(1-u)^b (1-v)^b \right], \ b > 1, a \in \left[ -1, \left( \frac{b+1}{b-1} \right)^{b-1} \right], \quad (1.6)$$

with  $\rho \in [-0.333, 0.375]$  and  $\rho \in [-0.333, 0.391]$ , respectively. The difference between the maximal positive correlations of the two types (1.5) and (1.6) is marginal. For more details about the Huang-Kotz copulas, see the recent works [1, 12].

Ebaid et al. [17] proposed the following new extended FGM copula

$$C(u,v) = uv \left[1 + a(1-u)(1-v)(1-bu)(1-bv)\right], \ b \in [0,1).$$
(1.7)

Ebaid et al. [17] claimed that, the admissible range of the parameters is

$$a \in \left[-1, \frac{1}{1-b}\right]$$
 and  $b \in [0, 1).$  (1.8)

Moreover,  $\rho \in [-0.333, 1]$  and  $\tau \in [-0.222, 1]$ . Barakat et al. [10] proved that the admissible range (1.8) and the claim about the correlations are wrong. Moreover, Barakat et al. [10] showed that the copula (1.7), whenever  $b \in [0, 1)$ , does not increase the maximum positive correlations for the FGM copula (1.4).

In this paper, we modify the copula (1.7) by elongating the range of the parameter b to  $[-\infty, \infty]$ , where  $\pm \infty$  is interpreted as  $\lim_{b \to \pm \infty}$ . The modified copula would be considered as a counterpart of the Huang and Kotz's copula of the 1st type defined by (1.5), in the sense that both have two shape parameters and provide the same improvement of the positive correlation between the dependent variables compared with the FGM copula. However, the modified copula has an evident preference than the copula (1.5) because the new copula has a simpler functional form than the copula (1.5), for being that in order to get it, as an extension of the classical FGM copula (1.3), we used an extra shape parameter as a multiplicative factor, while to get the copula (1.5), the extra shape parameter is used as an exponent. Besides this evident motivation, the modified copula includes some special cases of other extensions of the FGM copula.

Moreover, Ebaid et al. [17] used the copula (1.7), with a wrong admissible range (cf. [10]), to estimate the reliability in a dependent stress-strength model with an application to the Egyptian finance system. There is no doubt that this important application can not be benefited from as long as the admissible range of the copula (1.7) has not been determined or was specified in a wrong way. Therefore, the result of this paper enables us to use this application.

#### 2 The main result

The suggested modified copula of (1.7) is defined by

$$C(u, v; a, b) = uv \{1 + a(1 - u)(1 - v)(1 + bu)(1 + bv)\}, \quad (u, v) \in I^2, \ b \in [-\infty, \infty].$$
(2.1)

Clearly, the Spearman's Rho and Kendall's Tau of the copula (2.1) can be determined by reversing the sign of b in (17) and (18) in Ebaid et al. [17] as  $\rho = \frac{a(2+b)^2}{12}$  and  $\tau = \frac{a(2+b)^2}{18}$ , respectively.

In the following theorem, we determine a subset  $\Omega$  of the admissible range of the copula (2.1), on which the copula provides an improvement of the correlation coefficients  $\rho$  and  $\tau$ .

**Theorem 2.1.** The set  $\Omega$  is given by  $\Omega = \Omega^+ \cup \Omega^-$ , where

$$\Omega^{+} = \left\{ (a,b) : 0 \le b \le 1, -\frac{1}{(1+b)^{2}} \le a \le \frac{1}{(1+b)}; \text{ or } b > 1, -\frac{1}{(1+b)^{2}} \le a \le \frac{1}{(1+b)^{2}} \right\}, 
\Omega^{-} = \left\{ (a,b) : -2 \le b \le 0, -1 \le a \le 0; \text{ or } b < -2, -\frac{1}{(1+b)^{2}} \le a \le \frac{1}{(1+b)^{2}} \right\}.$$
(2.2)

*Proof.* The admissible range  $\Omega$  will be determined relying on the conditions (1.1) and (1.3). The corresponding copula density of (2.1) can be written in the form

$$c(u, v; a, b) = 1 + af(u, b)f(v, b),$$
(2.3)

where  $f(u,b) = 3bu^2 + 2u(1-b) - 1$ . Clearly, the condition (1.1) is satisfied, when 0 < b < 1. Furthermore, when 0 < b < 1, in view of (2.3), we get f(0,b) = -1 and f(1,b) = 3b + 2(1-b) - 1 = b + 1 > 0. On the other hand,  $f'(u,b) = \frac{df(u,b)}{du} = 6bu + 2(1-b) > 0$ . Thus, f(u,b) is strictly increasing. Moreover,  $\min_{0 \le u \le 1} f(u,b) = -1$  and  $\max_{0 \le u \le 1} f(u,b) = 1 + b > 0$ . Therefore, in order that, the condition (1.3) is satisfied, i.e.  $c(u,v;a,b) \ge 0$ , we must have

$$1 + a \min f(u, b) \cdot \min f(v, b) \ge 0 \Longrightarrow a \ge -1,$$
  
$$1 + a \min f(u, b) \cdot \max f(v, b) \ge 0 \Longrightarrow a \le \frac{1}{1 + b},$$
  
$$1 + a \max f(u, b) \cdot \max f(v, b) \ge 0 \Longrightarrow a \ge \frac{-1}{(1 + b)^2}$$

The above restrictions on a imply that  $a \in \left[\frac{-1}{(1+b)^2}, \frac{1}{1+b}\right]$ . Bearing in mind that,  $C(u, v; a, 0) = C_{FGM}(u, v; a), -1 \le a \le 1$  and  $C(u, v; a, 1) = C_{HK-FGM1}(u, v; a, 2), -\frac{1}{4} \le a \le \frac{1}{2}$ , we get  $(a, b) \in \Omega^+$ , for  $0 \le b \le 1$ , and  $a \in \left[\frac{-1}{(1+b)^2}, \frac{1}{1+b}\right]$ . Now, consider the case b > 1. Clearly, while for this case the condition (1.3) is still satisfied, but the condition (1.1) is not, especially for large values of b unless we alter the upper bound of a to be  $\frac{1}{(1+b)^2}$ , note that in this case  $\lim_{b\to\infty} C(u, v; \frac{\alpha}{(1+b)^2}, b) = uv[1 + \alpha uv(1-u)(1-v)] := C_{IFGM}(u, v; 0, \alpha)$ , where  $-1 \le \alpha \le 1$ , and  $C_{IFGM}(u, v; a, b)$  is a single iterated FGM copula, see Proposition 1. Thus, we proved that  $\Omega^+ \subset \Omega$ . We turn now to the case b < 0. When  $-2 \le b \le 0$ , by proceeding as we have done in the case b > 0, we can check that the condition (1.3) is satisfied if  $a \in \left[-1, \frac{1}{1+b}\right]$ , but the condition  $a \le \frac{1}{1+b}$  makes C(u, v; a, b) > 1, for some values of b (when the value of b is equal -1, or close to it). In order to fix the upper bound of a, we invoke the general relation  $C(u, v; a, b) \le 1 + a \le 1$ , which enables us to alter the upper bound of  $\frac{1}{1+b}$  by 0. Thus,  $(a, b) \in \Omega^-$ , if  $-2 \le b \le 0$ , and  $-1 \le a \le 0$ . Finally, when b < -2, we can show that the condition (1.3) is satisfied if  $a \in \left[\frac{-1}{(1+b)^2}, \frac{1}{1+b}\right]$ . On the other hand, in order that the condition (1.1) is satisfied (especially for large |b|), we can alter the admissible range of a to be  $\left[\frac{-1}{(1+b)^2}, \frac{1}{(1+b)^2}\right]$ . This proves that  $\Omega^- \subset \Omega$ , as required to complete the proof.

**Remark 1.** We can easily check that the improvement of the correlation attains only on  $\Omega^+$ . Namely, at  $b = 1, a = \frac{1}{2}$ , we get  $\rho = 0.375$ . On the other hand, no improvement can be gained on  $\Omega^-$ , which coincides with the remark of Barakat et al. [10] that the copula (1.7) does not improve the correlation.

**Proposition 1.** It is easily to check the validity of the following relations and properties:

$$1. \ C(u, v; a, 0) = C_{FGM}(u, v; a), -1 \le a \le 1,$$

$$2. \ C(u, v; a, 1) = C_{HK-FGM1}(u, v; a, 2), -\frac{1}{4} \le a \le \frac{1}{2},$$

$$3. \ C(u, v; a, -1) = C_{HK-FGM2}(u, v; a, 2), -1 \le a \le 3,$$

$$4. \ \lim_{b \to \pm \infty} C(u, v; \frac{\alpha}{(1+b)^2}, b) = C_{IFGM}(u, v; 0, \alpha), \text{ where } -1 \le \alpha \le 1 \text{ and}$$

$$C_{IFGM}(u, v; a, b) = uv \{1 + a(1 - u)(1 - v) + buv(1 - u)(1 - v)\},$$

$$- 1 \le a \le 1, a + b \ge -1, b \le \frac{3 - a + \sqrt{9 - 6a - 3a^2}}{2},$$

is the single iterated FGM copula, which was introduced by Huang and Kotz [18] and further studied by Alawady et al. [5], Barakat and Husseiny [9], and Barakat et al. [11, 13].

### References

- Abd Elgawad, M. A., Alawady, M. A., Barakat, H. M. and Shengwu Xiong. (2020a). Concomitants of generalized order statistics from Huang-Kotz Farlie-Gumbel-Morgenstern bivariate distribution: some information measures. Bull. Malays. Math. Sci. Soc., 43: 2627-2645.
- [2] Abd Elgawad, M. A., Barakat, H. M. and Alawady, M. A. (2020b). Concomitants of generalized order statistics under the generalization of Farlie-Gumbel-Morgenstern type bivariate distributions. Bull. Iran. Math. Soc., 47: 1045-1068. https://doi.org/10.1007/s41980-020-00427-0.
- [3] Abd Elgawad, M. A., Barakat, H. M. and Alawady, M. A. (2021a). Concomitants of generalized order statistics from bivariate Cambanis family: Some information measures. Bull. Iranian. Math. Soc. https://doi.org/10.1007/s41980-021-00532-8.
- [4] Abd Elgawad, M. A., Barakat, H. M., Xiong, S., and Alyami, S. A. (2021b). Information measures for generalized order statistics and their concomitants under general framework from Huang-Kotz FGM bivariate distribution. Entropy, 23(3), 335. https://doi.org/10.3390/e23030335.
- [5] Alawady, M. A., Barakat, H. M., Shengwu Xiong and Abd Elgawad, M. A. (2020). Concomitants of generalized order statistics from iterated Farlie–Gumbel–Morgenstern type bivariate distribution. Comm. Statist.-Theory Meth., to appear. https://doi.org/10.1080/03610926.2020.1842452.
- [6] Alawady, M. A., Barakat, H. M., Shengwu Xiong and Abd Elgawad, M. A. (2021a). On concomitants of dual generalized order statistics from Bairamov-Kotz-Becki Farlie-Gumbel-Morgenstern bivariate distributions. Asian-European J. Math. https://doi.org/10.1142/S1793557121501850.

- [7] Alawady, M. A., Barakat, H. M. and Abd Elgawad, M. A. (2021b). Concomitants of generalized order statistics from bivariate Cambanis family of distributions under a general setting. Bull. Malays. Math. Sci. Soc., to appear. https://doi.org/10.1080/03610926.2020.1842452.
- [8] Bairamov, I. and Kotz, S. (2002). Dependence structure and symmetry of Huang-Kotz FGM distributions and their extensions. Metrika, 56 (1): 55-72. https://doi:10.1007/s001840100158.
- [9] Barakat, H. M. and Husseiny, I. A. (2021). Some information measures in concomitants of generalized order statistics under iterated Farlie-Gumbel-Morgenstern bivariate type. Quaestiones Math., 44(5): 581-598. https://doi.org/10.2989/16073606.2020.1729271.
- [10] Barakat, H. M. Alawady, M. A. and Abd Elgawad, M. A. (2021a). Correction to the paper "A new extension of the FGM copula with an application in reliability" by Rasha Ebaid, Walid Elbadawy, Essam Ahmed and Abdalla Abdelghaly. Comm. Statist.-Theory Meth., to appear. https://doi.org/10.1080/03610926.2021.1879864
- [11] Barakat, H. M., Nigm, E. M. and Husseiny, I. A. (2020). Measures of information in order statistics and their concomitants for the single iterated Farlie-Gumbel-Morgenstern bivariate distribution. Math. Popul. Studies, to appear. https://doi.org/10.1080/08898480.2020.1767926.
- [12] Barakat, H. M., Nigm, E. M. and Syam, A. H. (2019). Concomitants of ordered variables from Huang-Kotz FGM type bivariate-generalized exponential distribution. Bull. Malays. Math. Sci. Soc., 42(1): 337-353. https://doi.org/10.1007/s40840-017-0489-5.
- [13] Barakat, H. M., Nigm, E. M., Alawady, M. A. and Husseiny, I. A. (2021b). Concomitants of order statistics and record values from iterated FGM type bivariate-generalized exponential distribution. REVSTAT, 19(2): 291-307.
- [14] Bekrizadeh, H. and Jamshidi, B. (2017). A new class of bivariate copulas: Dependence measures and properties. Metron 75 (1): 31-50. DOI: 10.1007/s40300-017-0107-1.
- [15] Bekrizadeh, H., Parham, G. A. and Zadkarmi, M. R. (2012). The new generalization of Farlie-Gumbel-Morgenstern copulas. App. Math. Sci., 6 (71): 3527-3533.
- [16] Durante, F. (2006). A new class of symmetric bivariate copulas. J. Nonparametric Statisti., 18 (7-8): 499-510. DOI: 10.1080/10485250701262242.
- [17] Ebaid, R., Elbadawy, W., Ahmed, E. and Abdelghaly, A. (2020). A new extension of the FGM copula with an application in reliability. Comm. Statist. Theory and Meth., to appear. https://doi.org/10.1080/03610926.2020.1785501.
- [18] Huang, J. S. and Kotz, S. (1984). Correlation structure in iterated Farlie-Gumbel-Morgenstern distributions. Biometrika, 71(3): 633-636. https://doi.org/10.2307/2336577.
- [19] Huang, J. S. and Kotz, S. (1999). Modifications of the Farlie-Gumbel-Morgenstern distributions. A tough hill to climb. Metrika 49(2): 135-145. https://doi.org/10.1007/s001840050030.
- [20] Nelsen, R. B. (2006). An Introduction to Copulas. 2nd ed. New York: Springer.