COMMENTS ON A NOTE BY M. WALDSCHMIDT

IGOR V. NIKOLAEV¹

ABSTRACT. This note is the follow up to a paper by M. Waldschmidt.

In [3] it was argued that the value of $\mu = \log \varepsilon$ in [2, Theorem 1] must be equal to 1. As a consequence, [2, Theorem 1] is claimed to be false. In this note we explain why $\mu = \log \varepsilon$ cannot be 1, thus upholding [2, Theorem 1]. Our comments touch [3, Second paragraph], since [3, First paragraph] is covered in [2, Remark 2].

The β -KMS (Kubo–Martin–Schwinger) state on a C^* -algebra \mathcal{A} is a real number β , such that

$$\omega(x\sigma_{i\beta}(y)) = \omega(yx), \quad \forall x, y \in \mathcal{A}, \tag{1}$$

where σ_t is the 1-parameter group of automorphisms of \mathcal{A} and ω is a weight on \mathcal{A} . Such a state is the fundamental invariant of the algebra \mathcal{A} known in the quantum statistical mechanics as an inverse temperature. If \mathcal{A} is an AF (Approximate Finite) C^* -algebra given by a constant incidence matrix $A \in GL_n(\mathbb{Z})$, the inverse temperature was calculated in [1, Chapter 4 and Fig. 1-5] by the remarkable formula

$$\beta = \log \left(\lambda_A\right),\tag{2}$$

where $\lambda_A > 1$ is the Perron-Frobenius eigenvalue of the matrix A. Moreover, the KMS state (2) is unique, *ibid*.

In our case n = 2 and the Pimsner-Voiculescu theorem says that the noncommutative torus \mathcal{A}_{θ} embeds into an AF-algebra preserving the respective K_0 groups. In particular, the incidence matrix A is nothing but the matrix form of the fundamental unit $\epsilon > 1$ of the number field $\mathbb{Q}(\theta)$ and therefore the inverse temperature of \mathcal{A}_{θ}

$$\beta = \log (\epsilon). \tag{3}$$

Since μ in [2, Lemma 3] is known to parametrize the state space of the C^* -algebra \mathcal{A}_{θ} , we identify it with β keeping the notation in below.

Assuming $\beta = 1$ [3, Second paragraph] would simply mean that a representation of the Sklyanin *-algebra $\{S(q) \mid q = \beta e^{2\pi i \theta}\}$ by the linear operators on a Hilbert space does not exist, since there is no scalar product to apply the standard GNS (Gelfand–Naimark–Segal) construction. The same is true for all other values of β , unless $\beta = \log(\epsilon)$ in which case the GNS construction gives a representation of the Sklyanin algebra. Such a representation is critical for the validity of [2, Lemma 2].

References

^{1.} O. Bratteli, P. E. T. Jorgensen and V. Ostrovsky1, Representation Theory and Numerical AF-invariants. The representations and centralizers of certain states on \mathcal{O}_d , Memoirs of the A. M. S. 168 (2004), No. 797

²⁰¹⁰ Mathematics Subject Classification. Primary 11J81, 11R35.

Key words and phrases. KMS states, inverse temperature.

NIKOLAEV

- 2. I. Nikolaev On algebraic values of function exp $(2\pi i x + \log \log y),$ Ramanujan J. 47 (2018), 417–425.
- 3. M. Waldschmidt, On a paper by Nikolaev, Ramanujan J. 57 (2022), 1517-1518.

 1 Department of Mathematics and Computer Science, St. John's University, 8000 Utopia Parkway, New York, NY 11439, United States.

Email address: igor.v.nikolaev@gmail.com

 $\mathbf{2}$