Realization of zero-divisor graphs of finite commutative rings as threshold graphs

Rameez Raja¹ and Samir Ahmad Wagay²

March 14, 2023

Abstract

Let R be a finite commutative ring with unity, and let G = (V, E) be a simple graph. The zero-divisor graph, denoted by $\Gamma(R)$ is a simple graph with vertex set as R, and two vertices $x, y \in R$ are adjacent in $\Gamma(R)$ if and only if xy = 0. In [10], the authors have studied the Laplacian eigenvalues of the graph $\Gamma(\mathbb{Z}_n)$ and for distinct proper divisors d_1, d_2, \dots, d_k of n, they defined the sets as, $\mathcal{A}_{d_i} = \{x \in \mathbb{Z}_n : (x, n) = d_i\}$, where (x, n) denotes the greatest common divisor of x and n. In this paper, we show that the sets $\mathcal{A}_{d_i}, 1 \leq i \leq k$ are actually orbits of the group action: $Aut(\Gamma(R)) \times R \longrightarrow R$, where $Aut(\Gamma(R))$ denotes the automorphism group of $\Gamma(R)$. Our main objective is to determine new classes of threshold graphs, since these graphs play an important role in several applied areas. For a reduced ring R, we prove that $\Gamma(R)$ is a connected threshold graph if and only if $R \cong F_q$ or $R \cong F_2 \times F_q$. We provide classes of threshold graphs realized by some classes of local rings. Finally, we characterize all finite commutative rings with unity of which zero-divisor graphs are not threshold.

Keywords: Group action, orbits, zero-divisor, zero-divisor graph, threshold graph.

2020 AMS Classification Code: 13A70, 05C25, 05C50.

1 Introduction

The zero-divisor graph of a commutative ring R with unity was introduced by Beck [3]. The graph $\Gamma(R)$ is defined to be the graph with vertex set as R, and two distinct vertices $x, y \in R$ are adjacent in $\Gamma(R)$ if and only if xy = 0 in R. The notion of zero-divisor graphs was further studied by Anderson and Livingston in [1], who restricted vertices of the graph $\Gamma(R)$ to non-zero zero-divisors

 $^{^{1,2}}$ Department of Mathematics, National Institute of Technology Srinagar, Jammu and Kashmir, India. Email: rameeznaqash@nitsri.ac.in, samir_03phd20@nitsri.net, Corresponding author:¹

of the ring R. The main objective of the investigation of associating a graph to R is to study the interplay of graph theoretic properties of $\Gamma(R)$ and ring theoretic properties of R. This interplay between different properties of R and $\Gamma(R)$ was thoroughly investigated in [4, 11, 12, 13, 14]. For basic definitions from graph theory, we refer to [10], and for ring theory we refer to [8].

A vertex in a graph G is called *dominating* if it is adjacent to every other vertex of G. A graph **G** is called a *threshold* graph if it is obtained by the following procedure.

Start with K_1 , a single vertex, and use any of the following steps, in any order, an arbitrary number of times.

(i) Add an isolated vertex.

(ii) Add a dominating vertex, that is, add a new vertex and make it adjacent to each existing vertex. The family of threshold graphs represent an important class of simple graphs.

The concept of threshold graphs was first established by the authors Chvátal and Hammer in [6] and Henderson and Zalcstein in [7]. There are many different generalizations of threshold graphs. These graphs can be viewed as special cases of some wider classes of graphs like cographs, split graphs and interval graphs.

By the definition of Beck's zero-divisor graph $\Gamma(R)$, it appears that the vertex 0 in $\Gamma(R)$ is adjacent to every other vertex of $\Gamma(R)$. Therefore, $\Gamma(R)$ is a simple connected graph, that is, $\Gamma(R)$ is without parallel edges and self loops. The distance between any two vertices in $\Gamma(R)$ is at least 1, which is the minimum eccentricity of $\Gamma(R)$ and at most 2, which is the maximum eccentricity of $\Gamma(R)$.

A graph G = (V, E) is said to be complete if every pair of distinct vertices of G are adjacent in G. Recall that a *clique* of G is a complete subgraph of G. A subset S of a vertex set of G is said to be an *independent set* if no two members of S are adjacent in G. A graph G is said to be *bipartite* if V can be partitioned into non-empty subsets V_1 and V_2 such that each edge of G has one end in V_1 and other in V_2 . A bipartite graph is said to be complete if every element of V_1 is adjacent to every element of V_2 . A complete bipartite is called a *star graph* if either $|V_1| = 1$ or $|V_2| = 1$. If $R \cong F_q$, where F_q is a finite field, then it is easy verify that $\Gamma(R)$ is a star graph

An element $r \in R$ is said to be *nilpotent* if $r^n = 0$, for some $n \in \mathbb{Z}_{>0}$. A ring R is called *reduced* if it has no non-zero nilpotent element and it is called *local* if R has a unique maximal ideal.

This research article is organised as follows. In Section 2, we present some preliminary information related to the zero-divisor graph $\Gamma(\mathbb{Z}_n)$. For $R \cong \mathbb{Z}_{p^{\alpha}}$, we show that the sets $\mathcal{A}_{d_i} = \{x \in \mathbb{Z}_n : (x, n) = d_i\}$ defined in [10] are actually orbits of the group action $Aut(\Gamma(R)) \times R \longrightarrow R$. In Section 3, we show that for a reduced ring R, $\Gamma(R)$ is a connected threshold graph if and only if $R \cong F_q$ or $R \cong F_2 \times F_q$. We discuss classes of threshold graphs realized by some classes of local rings and finally we characterize all finite commutative rings of which zero-divisor graphs are not threshold.

2 Preliminaries

It is always interesting to determine new classes of threshold graphs, since a threshold graph **G** with n vertices can be represented by a binary code of the type $b = (b_1b_2\cdots b_n) = 0^{s_1}1^{t_1}0^{s_2}1^{t_2}\cdots 0^{s_k}1^{t_k}$, where s_i and t_i are positive integers for all $1 \le i \le k$. In the binary code b, $b_i = 0$ if a vertex v_i is added as an isolated vertex in **G** and $b_i = 1$ if v_i is added as a dominating vertex in **G**. These codes realized by some class of graphs (threshold graphs) are known as the creation sequences of these graphs.

We partition the vertex set V of **G** as V_1, V_2, \dots, V_k . The equivalent form of an equitable partition matrix $\mathcal{M} = (a_{ij})$ of **G** is defined as,

$$a_{ij} = \begin{cases} |V_i| - 1 & \text{if } i = j \text{ and } V_i \text{ induces a complete subgraph in } \mathbf{G}, \\ |V_j| & \text{if there exists } v_i \in V_i \text{ and } v_j \in V_j \text{ such that } v_i \text{ is adjacent to } v_j, \\ 0 & \text{otherwise.} \end{cases}$$
(1)

The following example illustrates the equitable partition matrix of a threshold graph.

Let **G** be a threshold graph of order 10 represented by a binary code $b = (0000111001) = 0^4 1^3 0^2 1$ as shown in the Figure 1. Then the equitable partition matrix of **G** is,

$$\mathcal{M} = \begin{pmatrix} 0 & 3 & 0 & 1 \\ 4 & 2 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 4 & 3 & 2 & 0 \end{pmatrix}$$

Figure 1: Threshold graph (0000111001)

The multiplicities of eigenvalues 0 and -1 are 4 and 2, which can be verified from [2]. The

remaining eigenvalues of **G** can be computed from the equitable partition matrix defined in (1). In fact, the remaining eigenvalues of **G** are roots of the characteristic polynomial $x^4 - 2x^3 - 21x^2 - 12x + 24$.

Thus constructing some new classes of threshold graphs provides us a way of developing some new binary codes from graphs and a theory which involves spectral analysis related to graphs.

Recall that the graph obtained from the union of two graphs G_1 and G_2 by adding new edges from each vertex of G_1 to every vertex of G_2 is called join of G_1 and G_2 , denoted by $G_1 \vee G_2$. The generalised join graph which is also called generalised composition graph in [15] is defined as follows.

Definition 1. Let G be a graph with vertex set $V(G) = \{v_1, v_2, \dots, v_k\}$, and let G_1, G_2, \dots, G_k be k pairwise disjoint graphs. The G-generalized join graph $G_{\vee} = [G_1, G_2, \dots, G_k]$ of G_1, G_2, \dots, G_k is the graph obtained by replacing each vertex v_i of G with the graph G_i and two vertices G_i and G_j are adjacent in G_{\vee} if and only if every vertex v_i of G_i is adjacent to every vertex v_j of G_j in G.

Let Aut(G) denotes the automorphism group of a graph G, consider the group action Aut(G)acting on V(G) by some permutation of Aut(G), that is, $Aut(G) \times V(G) \to V(G)$ is given as, $\sigma(v) = u$, where $\sigma \in Aut(G)$ and $v, u \in V(G)$ are any two vertices of G.

The graph $\Gamma(\mathbb{Z}_n)$ is simple and its vertices are all elements of the ring \mathbb{Z}_n and two distinct vertices are adjacent in $\Gamma(\mathbb{Z}_n)$ if and only if there product is zero in \mathbb{Z}_n . The canonical decomposition of n is $p_1^{n_1}p_2^{n_2}\cdots p_r^{n_r}$ and total number of divisors of n is $\prod_{i=1}^r (n_i+1)$.

The authors in [10] defined Υ_n as a simple graph whose vertices are the proper divisors of n, and two distinct vertices d_i and d_j are adjacent in Υ_n if and only if $d_i d_j$ is a multiple of n. Thus, $V(\Upsilon_n) = \{d_1, d_2, \dots, d_r\}$ and $E(\Upsilon_n) = \{(d_i, d_j) : d_i d_j \equiv 0 \pmod{n}\}.$

Let $n = p^{\alpha}$, where p is a prime number and $\alpha \geq 1$. If $R \cong \mathbb{Z}_{p^{\alpha}}$, then orbits of the group action $Aut(\Gamma(R)) \times R \longrightarrow R$ are given as, $\mathcal{O}_{\alpha,p^{i}} = \{p^{i}b(\text{mod } p^{\alpha}) \mid b \in \mathbb{Z}, (b,p) = 1\}, \mathcal{O}_{\alpha,p^{i}} \text{ is the } p^{i}\text{-th}$ orbit of R, where $i \in [0, \alpha - 1]$.

For $0 \leq i < j \leq \alpha - 1$, $p^i b \equiv p^j b' \pmod{p^{\alpha}}$, where (b, p) = 1 and (b', p) = 1. Therefore, $p^{(j-i)}b' \equiv b \pmod{p^{(\alpha-i)}}$, which is a contradiction. Thus, for $i \neq j$, $\mathcal{O}_{\alpha,p^i} \cap \mathcal{O}_{\alpha,p^j} = \emptyset$.

Any element $a \in \mathbb{Z}_{p^{\alpha}}$ can be expressed as,

$$a \equiv p^{\alpha-1}b_1 + p^{\alpha-2}b_2 + \dots + pb_{\alpha-1} + b_{\alpha} \pmod{p^{\alpha}},$$

where $b_i \in [0, p-1]$. If $a \in \mathcal{O}_{\alpha,1}$, then $b_{\alpha} \neq 0$. So, $|\mathcal{O}_{\alpha,1}| = p^{\alpha-1}(p-1) = \phi(p^{\alpha})$. If $a' \in \mathcal{O}_{\alpha,p}$, then for some $a \in \mathcal{O}_{\alpha,1}$, a' = pa, that is, $b_{\alpha} \neq 0$, so $|\mathcal{O}_{\alpha,p}| = \phi(\frac{p^{\alpha}}{p})$. Similarly, for $i \in [0, \alpha - 1]$, we have $|\mathcal{O}_{\alpha,p^i}| = \phi(\frac{p^{\alpha}}{p^i})$ and $\mathcal{O}_{\alpha,p^{\alpha}} = \{0\}$. It follows that the size of the sets $\mathcal{O}_{\alpha,p^i} = \phi(\frac{p^{\alpha}}{p^i})$ for $i \in [0, \alpha]$. Therefore, total number of elements in all the sets $\mathcal{O}_{\alpha,p^i} = \sum_{i=0}^{\alpha} \phi(\frac{p^{\alpha}}{p^i}) = |\mathbb{Z}_{p^{\alpha}}| = p^{\alpha}$

Note that $1, p, \dots, p^{\alpha}$ are the representatives of orbits $\mathcal{O}_{\alpha,1}, \mathcal{O}_{\alpha,p}, \dots, \mathcal{O}_{\alpha,p^{\alpha}}$ and divisors of the order of R. It follows that the vertices of the graph Υ_n are elements of sets $\mathcal{O}_{\alpha,p}, \dots, \mathcal{O}_{\alpha,p^{\alpha-1}}$. Moreover, by definition of the graph $\Gamma(R)$, an element 0 of R is adjacent to all vertices in $\Gamma(R)$, elements of the orbit $\mathcal{O}_{\alpha,1}$ are adjacent to 0 only in $\Gamma(R)$. Also, elements of the orbit $\mathcal{O}_{\alpha,p}$ are adjacent to 0 and elements of the orbit $\mathcal{O}_{\alpha,p^{\alpha-1}}$. The elements of the orbit \mathcal{O}_{α,p^2} are adjacent to 0 and elements of orbits $\mathcal{O}_{\alpha,p^{\alpha-1}}, \mathcal{O}_{\alpha,p^{\alpha-2}}$. This, for $k \geq 1$, elements of the orbit \mathcal{O}_{α,p^k} are adjacent to 0 and elements of orbits $\mathcal{O}_{\alpha,p^{\alpha-k}}, \mathcal{O}_{\alpha,p^{\alpha-k+1}}, \dots, \mathcal{O}_{\alpha,p^{\alpha-1}}$.

Thus, the graph Υ_n can be obtained as a subgraph of $\Gamma(R)$ with vertices as elements of orbits $\mathcal{O}_{\alpha,p}, \cdots, \mathcal{O}_{\alpha,p^{\alpha-1}}$, and two vertices x and y are adjacent in Υ_n if and only xy = 0.

There is an advantage for knowing the orbits of this group action $Aut(\Gamma(R)) \times R \to R$, since we get some structural information about some elements of R from $\Gamma(R)$. As a consequence, we do not consider all elements of R to decode the symmetry of $\Gamma(R)$. We explore this information to reveal some interesting spectral properties of $\Gamma(R)$.

From the structure of $\Gamma(R)$, the subgraphs induced by elements of orbits $\bigcup_{i=0}^{k-1} \mathcal{O}_{\alpha,p^i}$, are independent parts in $\Gamma(R)$, whereas the subgraphs induced by elements of orbits $\bigcup_{j=0}^{k-1} \mathcal{O}_{\alpha,p^{k+j}}$ are cliques in $\Gamma(R)$. Also the orbit $\mathcal{O}_{\alpha,p^{\alpha}} = \{0\}$ is connected to every element of $\Gamma(R)$. The following Lemma shows adjacency of vertices in $\Gamma(\mathbb{Z}_{p^{\alpha}})$.

Lemma 2. For $i, j \in \{0, 1, \dots, \alpha - 1\}$, two vertices $x \in \mathcal{O}_{\alpha, p^i}$, $y \in \mathcal{O}_{\alpha, p^j}$ of $\Gamma(R)$ are adjacent in $\Gamma(R)$ if and only if $i + j \ge \alpha$.

Proof. Since $x \in \mathcal{O}_{\alpha,p^i}$ and $y \in \mathcal{O}_{\alpha,p^j}$, therefore, $x = p^i b$ and $y = p^j b'$ for some integers b and b' with (p,b) = (p,b') = 1. The vertices x and y are adjacent in $\Gamma(R)$ if and only if p^{α} divides xy. This implies x and y are adjacent in $\Gamma(R)$ if and only if $i + j \ge \alpha$.

Lemma 3. Let $\Gamma(R)$ be the graph realized by $R \cong \mathbb{Z}_{p^{\alpha}}$. Then the induced subgraph $\Gamma(\mathcal{O}_{\alpha,p^{i}})$ of $\Gamma(R)$ on the vertex set $\mathcal{O}_{\alpha,p^{i}}$ for $i \in \{0, 1, 2, \cdots, \alpha - 1\}$ is either the complete graph $K_{\phi(\frac{p^{\alpha}}{p^{i}})}$ or its complement graph $\overline{K}_{\phi(\frac{p^{\alpha}}{p^{i}})}$. Furthermore, for $\alpha = 2k$ or 2k - 1, $\Gamma(\mathcal{O}_{\alpha,p^{i}}) = K_{\phi(\frac{p^{\alpha}}{p^{i}})}$ if and only if $i \geq k$ and $\Gamma(\mathcal{O}_{\alpha,p^{\alpha}}) = K_{1}$.

Proof. The subgraph $\Gamma(\mathcal{O}_{\alpha,p^i})$ has $\phi(\frac{p^{\alpha}}{p^i})$ vertices for $i \in [0, \alpha]$. By taking i = j in Lemma (2), we get the result.

Observation 2.1. If $\Gamma(\mathcal{O}_{\alpha,p^i})$ is the induced subgraph of $\Gamma(R)$ on vertices \mathcal{O}_{α,p^i} for $i \in \{0, 1, 2, \cdots, \alpha\}$, then $\Gamma(R)_{\vee} = [\Gamma(\mathcal{O}_{\alpha,1}), \Gamma(\mathcal{O}_{\alpha,p}), \cdots, \Gamma(\mathcal{O}_{\alpha,p^{\alpha}})].$

Proof. Proof follows from the adjacency relations in $\Gamma(R)$, since orbits are adjacent in $\Gamma(R)_{\vee}$ if and only if their representatives are adjacent in $\Gamma(R)$.

We conclude this section with the following example which illustrates the zero-divisor graph realized by \mathbb{Z}_{3^3} with its orbits.

Example 4. Let $R = \mathbb{Z}_{3^3}$. Consider the group action $Aut(\Gamma(\mathbb{Z}_{3^3})) \times \mathbb{Z}_{3^3} \to \mathbb{Z}_{3^3}$. The orbits of this action are: $\mathcal{O}_{3,1} = \{1, 2, 4, \dots, 25, 26\}$, $\mathcal{O}_{3,3} = \{3, 6, 12, 15, 21, 24\}$, $\mathcal{O}_{3,3^2} = \{9, 18\}$ and $\mathcal{O}_{3,3^3} = \{0\}$. Therefore, by observation (2.1),

$$\Gamma(\mathbb{Z}_{3^3})_{\vee} = [\Gamma(\mathcal{O}_{3,1}), \Gamma(\mathcal{O}_{3,3}), \Gamma(\mathcal{O}_{3,3^2}), \Gamma(\mathcal{O}_{3,3^3})],$$

where $\Gamma(\mathcal{O}_{3,1}) = \overline{K}_{18}$, $\Gamma(\mathcal{O}_{3,3}) = \overline{K}_6$, $\Gamma(\mathcal{O}_{3,3^2}) = K_2$, and $\Gamma(\mathcal{O}_{3,3^3}) = K_1$ as shown in the Figure 2.



Figure 2: Generalised join graph $\Gamma(\mathbb{Z}_{3^3})_{\vee}$

3 Threshold graphs realized by reduced, local and mixed rings

The Artinian decomposition of $R = R_1 \times \cdots \times R_r \times F_1 \times \cdots \times F_s$, where each R_i is a finite commutative local ring with unity and F_j is a field, $1 \le i \le r$ and $1 \le j \le s$. A ring R is said to be product of local rings if s = 0 and $r \ge 2$. A ring R is said to be a *mixed ring* if $r \ge 1$ and $s \ge 1$. A ring R is

said to be a *reduced ring* if r = 0 and $s \ge 1$.

This section is dedicated to study the realization of zero-divisor graphs as threshold graphs for reduced rings, product of local rings and mixed rings.

An alternating 4 - cycle of a graph G = (V, E) is a configuration consisting of four distinct vertices a, b, c, d such that $(a, b), (c, d) \in E$ and $(a, c), (b, d) \notin E$. By considering the presence or absence of edges (a, d) and (b, c), we see that the vertices of an alternating 4-cycle induce a path P_4 , a cycle C_4 , or a matching $2K_2$.

Since threshold graphs has been characterised by many ways. One of the characterizations of a threshold graph [[9], Theorem 1.2.4] is presented in the following result.

Theorem 5. For a graph G = (V, E), the following are equivalent,
1. G is a threshold graph,
2. G does not have an alternating 4 - cycle.

Let $R \cong F_{q_1} \times F_{q_2} \times \cdots \times F_{q_n}$ be a reduced ring. In the following characterization, we discuss the realization of $\Gamma(R)$ as a threshold graph.

Theorem 6. Let $R \cong \mathbb{F}_{q_1} \times F_{q_2} \times \cdots \times F_{q_n}$. Then $\Gamma(R)$ is a connected threshold graph if and only if $R \cong F_{q_i}$, or $R \cong F_{q_i} \times F_{q_j}$, where $F_{q_i} \cong \mathbb{Z}_2$ for any $i, j \in \{1, 2, \cdots, n\}$.

Proof. Let R be a ring such that $R \cong F_{q_i}$ for some $i \in \{1, 2, \dots, n\}$. Then $\Gamma(R)$ is a complete bipartite graph isomorphic to K_{1,q_i-1} . This implies, $\Gamma(R)$ is a star graph. So, $\Gamma(R)$ is a connected threshold graph.

Let R be a ring such that $R \cong F_{q_i} \times F_{q_j}$ for some $i, j \in \{1, 2, \dots, n\}$ with $F_{q_i} \cong \mathbb{Z}_2$. Then $|V(\Gamma(R))| = 2q_i$. Let $V(\Gamma(R)) = X \bigcup Y$, where X represents the independent part of the graph whose vertices are of the form $(0, a_k)$ and $(1, a_k)$, $k \in \{1, 2, \dots, q_j - 1\}$. The set Y represents complete part of the graph which consists of only two vertices (1, 0) and (0, 0).

To prove $\Gamma(R)$ is threshold, it is suffices to show that $\Gamma(R)$ does not contain an alternating 4-cycle, that is, the graph $\Gamma(R)$ has no induced subgraph of four vertices isomorphic to P_4 , C_4 or $2K_2$.

Let a, b, c, d be any four distinct vertices of $\Gamma(R)$. We consider the following cases.

Case - 1. If $a, b, c, d \in X$, then they are all isolated vertices. This implies, $\Gamma(R)$ has no alternating 4-cycle. Therefore, the result follows.

Case - 2. If $a, b, c \in X$ and $d \in Y$, then the only possibilities are as follows,



but none of them is isomorphic to P_4 , C_4 or $2K_2$. Therefore, $\Gamma(R)$ is a connected threshold graph.

Case - 3. If $a, b \in X$ and $c, d \in Y$, where c = (1, 0) and d = (0, 0). By definition, the vertex (0, 0) is adjacent to three other vertices. Consequently, no two disjoint edges exists and therefore no alternating 4 - cycle is possible. Thus, $\Gamma(R)$ is threshold.

Conversely, suppose that $\Gamma(R)$ is a connected threshold graph, we show that $R \cong F_{q_i}$, or $R \cong F_{q_i} \times F_{q_i}$, where $F_{q_i} \cong \mathbb{Z}_2$ for any $i, j \in \{1, 2, \dots, n\}$. We consider the following cases.

Case - 1. Suppose $R \cong F_{q_1} \times F_{q_2}$, $2 < q_1 \leq q_2$. Then, $|V(\Gamma(R))| = q_1q_2$. Since (a, b), (a, c), (c, d), $(b, d) \in E(\Gamma(R))$, where a = (1, 0), b = (0, y), c = (0, 1) and $d = (x, 0), x \in F_{q_1} \setminus \{0, 1\}$, $y \in F_{q_2} \setminus \{0, 1\}$. This implies, $\Gamma(R)$ has an induced subgraph of order 4 isomorphic to C_4 , a contradiction. Thus, $R \ncong F_{q_1} \times F_{q_2}$, $2 < q_1 \leq q_2$.

Case - 2. Suppose $R \cong F_{q_1} \times F_{q_2} \times \cdots \times F_{q_n}$ with $q_1 \leq q_2 \leq \cdots \leq q_n$ and $n \geq 3$. Then, $\Gamma(R)$ always contains an induced subgraph of the form $P_4 = (1, 0, 1, 0, \cdots, 0) \sim (0, 1, 0, 0, \cdots, 0) \sim (1, 0, 0, 0, \cdots, 0) \sim (0, 1, 1, 0, \cdots, 0)$, or $2K_2 = \{(1, 0, 1, 0, \cdots, 0) \sim (0, 1, 0, 1, \cdots, 0), (0, 1, 1, 0, \cdots, 0) \sim (1, 0, 0, 1, \cdots, 0)\}$ or $C_4 = (a, b), (a, c), (c, d), (b, d)$, where $a = (1, 0, 0, \cdots, 0), b = (0, y, 0, \cdots, 0), c = (0, 1, 0, \cdots, 0)$ and $d = (x, 0, 0, \cdots, 0), x \in F_{q_1} \setminus \{0, 1\}, y \in F_{q_2} \setminus \{0, 1\}$, a contradiction. This implies, $R \neq F_{q_1} \times F_{q_2} \times \cdots \times F_{q_n}$, for $q_1 \leq q_2 \leq \cdots \leq q_n$.

Therefore, $\Gamma(R)$ fails to be threshold for cases 1 and 2. We conclude that either $R \cong F_{q_i}$ or $R \cong F_{q_i} \times F_{q_j}$.

Now, we consider the case of local rings. We show that if R is a local ring, then $\Gamma(R)$ is not always a threshold graph. Let $R \cong \mathbb{Z}_8[x]/(x^2, 2x), \mathbb{Z}_2[x]/(x^3, 2x^2, 2x)$ or $\mathbb{Z}_2[x, y]/(x^3, xy, y^2)$. Then the zero-divisor graph realized by $\mathbb{Z}_8[x]/(x^2, 2x)$ is shown in the Figure 3. If we relabel the vertices of $\Gamma(\mathbb{Z}_8[x]/(x^2, 2x))$ by elements of rings $\mathbb{Z}_2[x]/(x^3, 2x^2, 2x)$ and $\mathbb{Z}_2[x, y]/(x^3, xy, y^2)$, then we obtain isomorphic copies of $\Gamma(\mathbb{Z}_8[x]/(x^2, 2x))$ as zero-divisor graphs of $\mathbb{Z}_2[x]/(x^3, 2x^2, 2x)$ and $\mathbb{Z}_2[x, y]/(x^3, xy, y^2)$. It can be easily verified from the Figure 3, that $\Gamma(\mathbb{Z}_8[x]/(x^2, 2x))$ is a threshold graph.



Figure 3: $\Gamma(\mathbb{Z}_8[x]/(x^2, 2x))$

On the other hand, if $R \cong \mathbb{Z}_4[x]/(x^2)$, $\mathbb{Z}_2[x, y]/(x^2, y^2)$ or $\mathbb{Z}_4[x, y]/(x^2, y^2, xy - 2, 2x, 2y)$, then the graph realized by these rings are isomorphic to each other. The graph realized by the ring $\mathbb{Z}_4[x]/(x^2)$) is shown in the Figure 4. The graph $\Gamma(\mathbb{Z}_4[x]/(x^2))$ fails to be threshold, since it contains an induced subgraph isomorphic to $2K_2$, that is the edges (x, 3x) and (2, 2 + 2x) are disjoint in $\Gamma(\mathbb{Z}_4[x]/(x^2))$.



Thus from preceding examples it is hard to characterize local rings such that their zero-divisor graphs are threshold. Below, we present some classes of local rings of which associated zero-divisor

graphs are threshold.

Theorem 7. For each prime p and positive integer α , the zero-divisor graph realized by the ring $\mathbb{Z}_{p^{\alpha}}[x]/(x^2, px)$ is a connected threshold graph.

Proof. Let $R \cong \mathbb{Z}_{p^{\alpha}}[x]/(x^2, px)$. Then orbits of the group action $Aut(\Gamma(R)) \times R \longrightarrow R$ are as

follows,

$$\mathcal{O}_{\alpha,p^i} = \{p^i a + bx : 0 \neq a \in \mathbb{Z}_{p^{\alpha}}, b \in \mathbb{Z}_p \text{ and } (a,p) = 1\}, \text{ where } 0 \le i \le \alpha - 2, \text{ and}$$

$$\mathcal{O}_{\alpha, p^{\alpha-1}} = \{ p^{\alpha-1}a + bx \neq 0 : a \in \mathbb{Z}_{p^{\alpha}}, b \in \mathbb{Z}_{p} \text{ and } (a, p) = 1 \}.$$

Moreover, for $0 \le i \le \alpha - 2$, $|\mathcal{O}_{\alpha,p^i}| = \phi(\frac{p^{\alpha+1}}{p^i})$ and $|\mathcal{O}_{\alpha,p^{\alpha-1}}| = \phi(\frac{p^{\alpha+1}}{p^{\alpha-1}}) + p - 1 = p^2 - 1$. Note that $\mathcal{O}_{\alpha,p^{\alpha}} = \{0\}$. The vertex set $V(\Gamma(R)) = (\bigcup_{i=0}^{\alpha} \mathcal{O}_{\alpha,p^i})$. To prove the desired result, we consider the following cases.

Case - 1. $\alpha = 2k$ for some $k \ge 1$. Then the subgraphs associated with orbits $X = \left(\bigcup_{i=0}^{k-1} \mathcal{O}_{\alpha,p^{k+i}}\right)$ represents complete part of the graph $\Gamma(R)$ and the subgraphs associated with orbits $Y = \left(\bigcup_{i=0}^{k-1} \mathcal{O}_{\alpha,p^i}\right)$ represents the independent part of $\Gamma(R)$ and each element of X and Y are connected to $\mathcal{O}_{\alpha,p^{\alpha}} = \{0\}$. Furthermore, if $r \in \mathcal{O}_{\alpha,p^{k+i}}$ and $s \in \mathcal{O}_{\alpha,p^i}$ for some *i*, then *r* and *s* are adjacent in the graph if and only if p^{α} divides p^{2i+k} . This implies,

$$|V(\Gamma(R))| = \sum_{i=0}^{k-1} |\mathcal{O}_{\alpha,p^i}| + \sum_{i=0}^{k-2} |\mathcal{O}_{\alpha,p^{k+i}}| + (|\mathcal{O}_{\alpha,p^{2k-1}}| + (p-1)) + 1 = p^{\alpha+1}.$$

To prove that $\Gamma(R)$ is a threshold graph it suffices to show that $\Gamma(R)$ has no induced subgraph of the form C_4 , P_4 or $2K_2$, that is, it has no alternating 4-cycle.

Let a, b, c, d be four distinct vertices of $\Gamma(R)$.

Subcase - 1. If a = 0 and $b, c, d \in X \cup Y$, then $(b, 0), (c, 0), (d, 0) \in E(\Gamma(R))$. Thus no two disjoint edges are present and therefore no alternating 4 - cycle is possible.

Subcase - 2. If all $a, b, c, d \in X \cup Y$, that is, no one among a, b, c, d represents the vertex 0, then we have the following cases .

Subcase - 2.1. Suppose that all four vertices $a, b, c, d \in X$. Then they are all adjacent to each other and hence no alternating 4 - cycle is possible.

Subcase - 2.2. Suppose that all four vertices $a, b, c, d \in Y$. Then these vertices are all isolated and therefore no alternating 4 - cycle is possible.

Subcase - 2.3. Suppose $a \in X$ and $b, c, d \in Y$. Then in $\Gamma(R)$, there may not be any edge from a to b, c, d. Without of loss of generality, the other possibilities are either $(a, d) \in E(\Gamma(R))$ or $(a, b), (a, d) \in E(\Gamma(R))$ or $(a, b), (a, c), (a, d) \in E(\Gamma(R))$ as follows.



Thus no one among the above graphs is isomorphic to C_4 , P_4 or $2K_2$. Therefore the result follows. **Subcase - 2.4.** Suppose $a, b \in X$ and $c, d \in Y$, then the only possibilities are as follows.



Subcase - 2.5. Suppose $a, b, c \in X$ and $d \in Y$, then the only possibilities are as follows.

$$a \rightarrow b \\ c \rightarrow d$$
 $a \rightarrow b \\ c \rightarrow d$ $a \rightarrow b \\ c \rightarrow d$ $a \rightarrow b \\ c \rightarrow d$ $a \rightarrow b \\ c \rightarrow d$

Thus it follows that the graph $\Gamma(R)$ has no alternating 4 - cycle. Therefore, $\Gamma(R)$ is a connected threshold graph.

Case - 2. $\alpha = 2k - 1$ for some $k \ge 1$. Then the subgraphs associated with orbits $X = \begin{pmatrix} \bigcup_{i=0}^{k-2} \mathcal{O}_{\alpha,p^{k+i}} \end{pmatrix}$ represents complete part of the graph $\Gamma(R)$ and the subgraphs associated with orbits $Y = \begin{pmatrix} \bigcup_{i=0}^{k-1} \mathcal{O}_{\alpha,p^i} \end{pmatrix}$ represents the independent part of $\Gamma(R)$ and each element of X and Y are connected to $\mathcal{O}_{\alpha,p^{\alpha}} = \{0\}$. This implies,

$$|V(\Gamma(R))| = \sum_{i=0}^{k-1} |\mathcal{O}_{\alpha,p^i}| + \sum_{i=0}^{k-3} |\mathcal{O}_{\alpha,p^{k+i}}| + (|\mathcal{O}_{\alpha,p^{2k-2}}| + (p-1)) + 1 = p^{\alpha+1}.$$

We can prove that $\Gamma(R)$ does not have an alternating 4-cycle by considering four distinct vertices a, b, c, d and using the same argument as above for the case $\alpha = 2k$.

Theorem 8. For each prime p, zero-divisor graphs realized by rings $\mathbb{Z}_p[x]/(x^p)$, $\mathbb{Z}_p[x,y]/(x^3, xy, y^2)$, and $\mathbb{Z}_{p^2}[x]/(px, x^2 - p)$ are connected threshold graphs.

Proof. Let $R \cong \mathbb{Z}_p[x]/(x^p)$. Then orbits of the group action $Aut(\Gamma(R)) \times R \longrightarrow R$ are as follows,

$$\mathcal{O}_{p,x^i} = \{ x^i (a_0 + a_1 x + \dots + a_{p-(1+i)} x^{p-(1+i)}) : a_i \in \mathbb{Z}_p, \text{ and } a_0 \neq 0 \},\$$

where $0 \leq i \leq p-1$. For $0 \leq i \leq p-1$, $|\mathcal{O}_{p,x^i}| = \phi(\frac{p^p}{p^i})$ and $|\mathcal{O}_{p,x^i}| = \{0\}$. The vertex set $V(\Gamma(R)) = X \bigcup Y$, where $X = \begin{pmatrix} \bigcup_{i=0}^{\frac{p-1}{2}} \mathcal{O}_{p,x^{i+\frac{p+1}{2}}} \end{pmatrix}$ and $Y = \begin{pmatrix} \bigcup_{i=0}^{\frac{p-1}{2}} \mathcal{O}_{p,x^i} \end{pmatrix}$, represents complete and independent parts of the graph $\Gamma(R)$, and $\mathcal{O}_{p,x^p} = \{0\}$. Also, two distinct vertices $r \in \mathcal{O}_{p,x^i}$ and $s \in \mathcal{O}_{p,x^j}$ for $i, j \in \{0, 1, \cdots, \frac{p-1}{2}\}$ are adjacent in $\Gamma(R)$ if and only if $i+j \geq p$. Therefore,

$$|V(\Gamma(R))| = \sum_{i=0}^{\frac{p-1}{2}} |\mathcal{O}_{p,x^{i+\frac{p+1}{2}}}| + \sum_{i=0}^{\frac{p-1}{2}} |\mathcal{O}_{p,x^{i}}| = p^{p}.$$

Moreover, for $R \cong \mathbb{Z}_p[x,y]/(x^3,xy,y^2)$ and for each $a_0, a_1, a_2, b_1 \in \mathbb{Z}_p$, the orbits of the group

action $Aut(\Gamma(R)) \times R \longrightarrow R$ are as follows,

$$\begin{array}{rcl} \mathcal{O}_{p,x} &=& \{a_1x + a_2x^2 + b_1y : & a_1 \neq 0\}, \\ \\ \mathcal{O}_{p,y} &=& \{a_2x^2 + b_1y \neq 0\}, \\ \\ \mathcal{O}_{p,1} &=& \{a_0 + a_1x + a_2x^2 + b_1y : & a_0 \neq 0\} \\ \\ \mathcal{O}_{p,p} &=& \{0\}, \end{array}$$

where x, y, p and 1 are representatives of orbits $\mathcal{O}_{p,x}$, $\mathcal{O}_{p,y}$, $\mathcal{O}_{p,1}$ and $\mathcal{O}_{p,p}$ with $|\mathcal{O}_{p,x}| = p^3 - p^2$, $|\mathcal{O}_{p,y}| = p^2 - 1$, $|\mathcal{O}_{p,1}| = p^4 - p^3$ and $|\mathcal{O}_{p,p}| = 1$. The vertex set $V(\Gamma(R)) = X \bigcup Y$, where $X = (\mathcal{O}_{p,y})$ and $Y = (\mathcal{O}_{p,x} \cup \mathcal{O}_{p,1})$ represents complete and independent parts of the graph $\Gamma(R)$, and $\mathcal{O}_{p,p} = \{0\}$. Furthermore, every vertex in $\mathcal{O}_{p,y}$ is adjacent to each vertex in $\mathcal{O}_{p,x}$ and the vertex 0 is adjacent to each vertex of $\Gamma(R)$. Therefore, $|V(\Gamma(R))| = p^4$.

Finally, for $R \cong \mathbb{Z}_{p^2}[x]/(px, x^2 - p)$, the orbits of the group action $Aut(\Gamma(R)) \times R \longrightarrow R$ are as follows,

$$\begin{array}{rcl} \mathcal{O}_{p,x} &=& \{pa+bx: \ 0 \neq b \in \mathbb{Z}_p, \ a \in \mathbb{Z}_p\}, \\ \mathcal{O}_{p,p} &=& \{pa: \ 0 \neq a \in \mathbb{Z}_p\}, \\ \mathcal{O}_{p,1} &=& \{a+bx: \ 0 \neq a \in \mathbb{Z}_{p^2}, \ b \in \mathbb{Z}_p \ and \ (a,p)=1\} \\ \mathcal{O}_{p,p^2} &=& \{0\}, \end{array}$$

where $|\mathcal{O}_{p,x}| = p^2 - p$, $|\mathcal{O}_{p,p}| = p - 1$, $|\mathcal{O}_{p,1}| = p^3 - p^2$ and $|\mathcal{O}_{p,p^2}| = 1$. The vertex set $V(\Gamma(R)) = X \bigcup Y$, where $X = (\mathcal{O}_{p,p})$ and $Y = (\mathcal{O}_{p,x} \cup \mathcal{O}_{p,1})$ represents complete and independent parts of the graph $\Gamma(R)$ and $\mathcal{O}_{p,p^2} = \{0\}$. From the structure of the graph, every vertex in $\mathcal{O}_{p,1}$ is adjacent to the vertex 0 only, whereas every vertex in $\mathcal{O}_{p,p}$ is adjacent to each vertex of $\mathcal{O}_{p,x}$ and to the vertex 0. Thus, $|V(\Gamma(R))| = p^3$.

By taking any four distinct vertices a, b, c and d from vertex set of each of the graphs realized by rings $(\mathbb{Z}_p[x]/(x^p))$, $\mathbb{Z}_p[x,y]/(x^3,xy,y^2)$, $\mathbb{Z}_{p^2}[x]/(px,x^2-p)$ and use the similar approach as in Theorem (7), it can be shown that zero divisor graphs realized by above mentioned rings does not contain an alternating 4 - cycle and therefore the result follows.

Corollary 9. For each prime p and positive integer α , the zero-divisor graph realized by a local ring $\mathbb{Z}_{p^{\alpha}}$ is a connected threshold graph.

Problem. Characterise all finite local rings such that $\Gamma(R)$ is a threshold graph.

In the following result, we consider the Artinian decomposition of R for $r \ge 2$ and s = 0, that is, we investigate the structure of $\Gamma(R)$ when R is a product of local rings only. **Lemma 10.** If $R \cong R_1 \times R_2 \times \cdots \times R_r$, where $r \ge 2$ and each $R_i, 1 \le i \le r$ is a finite local ring (not a field), then $\Gamma(R)$ is not a threshold graph.

Proof. Since R is a product of at least two finite local rings, therefore $\Gamma(R)$ always has an induced subgraph either of the form P_4 or C_4 or $2K_2$ described as follows.



Note that $a_i \in Z^*(R_1)$ and $b_i \in Z^*(R_2)$. where $Z^*(R_1)$ and $Z^*(R_2)$ denotes the set of non-zero zero-divisors of rings R_1 and R_2 .

In the following result, we classify all finite commutative rings R such that $\Gamma(R)$ is not a threshold graph.

Theorem 11. If $R \cong R_1 \times R_2 \times \cdots \times R_r \times F_1 \times F_2 \times \cdots \times F_s$, where r and s are positive integers, each R_i is a finite commutative local ring (not a field), and each F_i is a finite field, then $\Gamma(R)$ is not threshold.

Proof. The proof follows from Theorem (6) and Lemma (10).

Conclusion: In this research article, we presented some new classes of threshold graphs. As discussed above, the construction of new classes of threshold graphs are effective in a sense that it assists to advance in coding theory and spectral analysis. We obtained the results related to reduced and local rings. In fact, we characterized all reduced rings of which zero-divisor graphs are realized as threshold graphs. Furthermore, we obtained classes of threshold graphs realized by some classes of local rings and provided an example which illustrates that a zero-divisor graph associated with some local ring is not always a threshold graph. So, it is interesting to determine all classes of local rings of which zero-divisor graphs are always threshold. This investigation has been included in the paper as a problem for future research work.

Acknowledgement

We thank the anonymous reviewers for their careful reading of the manuscript and their many

insightful comments and suggestions. The second author's research is supported by the University Grants Commission, Govt. of India under UGC-Ref. No. 1337/(CSIR-UGC NET JUNE 2019).

Declaration of competing interest

There is no conflict of interest to declare.

Data Availability.

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

References

- D. F. Anderson and P. S. Livingston, The zero-divisor graph of a commutative ring, J. Algebra 217 (1999) 434 - 447.
- [2] R. B. Bapat, Graphs and Matrices, Springer/Hindustan Book Agency, London/New Delhi 2010.
- [3] I. Beck, Coloring of commutative rings, J. Algebra 116 (1988) 208 226.
- [4] Bilal A. Rather, S. Pirzada, T. A. Naikoo, Y. Shang, On Laplacian eigenvalues of the zero-divisor graph associated to the ring of integers modulo n, Math. 9 (5) (2021) 482.
- [5] S. Chattopadhyay, K. L. Patra and B. K. Sahoo, Laplacian eigenvalues of the zero divisor graph of the ring \mathbb{Z}_n , L. Algeb. Appl. 584 (2020) 267 286.
- [6] V. Chvátal, P. L. Hammer, Aggregation of Inequalities in Integer Programming, Ann. Disc. Math. 1 (1977) 145 - 162.
- [7] P. B. Henderson and Y. Zalcstein, Synchronization problems solvable by generalized pv systems, J. ACM. 27 (1) (1980) 60 - 71.
- [8] I. Kaplansky, Commutative Rings, rev. edn. University of Chicago Press, Chicago, 1974.
- [9] N. V. R. Mahadev, U.N. Peled, Threshold Graphs and Related Topics, Ann. Disc. Math. 56, (1995).
- [10] S. Pirzada, An Introduction to Graph Theory, Universities Press, OrientBlackSwan, Hyderabad, 2012.
- [11] S. Pirzada, Bilal A. Rather and T. A. Chishti, On distance Laplacian spectrum of zero

divisor graphs of \mathbb{Z}_n , Carpathian Mathematical Publications, 13 (1) (2021) 48 - 57.

- [12] S. Pirzada, Bilal A. Wani, A. Somasundaram, On the eigenvalues of zero divisor graph associated to finite commutative ring $\mathbb{Z}_{p^Mq^N}$. AKCE International J. Graphs Combinatorics, 18 (1) (2021) 1 16.
- [13] R. Raja, Total perfect codes in graphs realized by commutative rings, Transactions of Comb. 11 (4) (2022) 295 - 307.
- [14] S. P. Redmond, On zero-divisor graphs of small finite commutative rings, Disc. Math. 307 (9-10) (2007) 1155 - 1166.
- [15] A.J. Schwenk, Computing the characteristic polynomial of a graph, Graphs and Comb. Springer, Berlin, Heidelberg (1974) 153 - 172.