ON MACKEY DECOMPOSITION FOR LOCALLY PROFINITE GROUPS

YUKI YAMAMOTO

Abstract. To study induced representation of some class of groups, Mackey's theory is very useful. In this paper, we consider some generalization of Mackey's theory for locally profinite groups. In particular, we give conditions on groups under which we have the Mackey decomposition and some examples such that we do not have the Mackey decomposition in some sense.

CONTENTS

1. INTRODUCTION

In representation theory for groups G , the induction Ind is fundamental as a method to obtain representations of G from representations of its subgroups. To study induced representations, Mackey's theory is very useful. Mackey [\[2\]](#page-8-1) showed that for any finite group G and its subgroups H and K , we have the Mackey decomposition

$$
\operatorname{Res}^G_K\operatorname{Ind}^G_H\rho\cong\bigoplus_{g\in K\backslash G/H}\operatorname{Ind}^K_{K\cap^gH}\operatorname{Res}^{^gH}_{K\cap^gH} {^g\rho}
$$

for any H-representation ρ over some field, where for $g \in G$ we put $^g H := g H g^{-1}$ and $g \circ \rho$ is a gH-representation defined as $g \circ \rho(ghg^{-1}) = \rho(h)$ for any $h \in H$. By Frobenius reciprocity, we can also examine intertwining operators between induced representations from the Mackey decomposition.

There exist generalizations of Mackey's theory for some class of groups. The goal of this paper is to give a generalization of Mackey's theory for smooth representations of locally profinite groups, based on previous researches [\[1\]](#page-8-2), [\[3\]](#page-8-3).

In smooth representation theory for locally profinite groups, there two functors treated as induction, "induction" Ind and "compact induction" c-Ind. We give conditions on groups under which we have the Mackey decomposition for Ind and c-Ind. The main theorem in this paper is the following.

Theorem 1.1 (Theorem [2.5,](#page-4-1) Example [2.6\)](#page-5-0). Let G be a locally profinite group. Let H and K be closed subgroups in G. Let ρ be a smooth representation of H over some commutative ring R with unit.

(1) If either H or K is open in G, then we have the Mackey decomposition

$$
\mathrm{Res}^G_K\mathrm{c}\text{-}\mathrm{Ind}^G_H\rho\cong \bigoplus_{g\in K\backslash G/H}\mathrm{c}\text{-}\mathrm{Ind}^K_{K\cap^gH}\mathrm{Res}^{^gH}_{K\cap^gH} {^g}\rho.
$$

(2) If K is open in G, then we have the Mackey decomposition

$$
\operatorname{Res}^G_K\operatorname{Ind}^G_H\rho\cong \left(\prod_{g\in K\backslash G/H}\operatorname{Ind}^K_{K\cap^gH}\operatorname{Res}^{^gH}_{K\cap^gH} {^g\rho}\right)^{\infty}
$$

where for a K-representation τ , we denote by τ^{∞} the K-smooth part of τ . (3) If we omit the assumption that K is open, there exists an example such that H is open in G and the isomorphism in [\(2\)](#page-1-2) does not hold.

,

The example in [\(3\)](#page-1-3) shows that the condition that HgK is open and closed in G for any $g \in G$ is not enough to have the Mackey decomposition in general. (cf. [\[3,](#page-8-3) I.5.5])

As a corollary of this theorem, we also consider intertwining operators between induced representations, which are studied by Kutzko [\[1\]](#page-8-2) for some special cases.

Acknowledgment I am deeply grateful to my supervisor Naoki Imai for checking my draft carefully and giving helpful comments to improve the draft. I also thank Marie-France Vignéras and Masao Oi for many help, comments and suggestions. I am supported by the FMSP program at Graduate School of Mathematical Sciences, the University of Tokyo. I was also supported by JSPS KAKENHI Grant Number JP21J13751.

2. Mackey decomposition

In this section, we collect knowledge on the generalization of Mackey theory [\[2\]](#page-8-1) to locally profinite groups, based on [\[3,](#page-8-3) I.5].

2.1. Induction and compact induction. Let G be a locally profinite group. Let H be a closed subgroup in G, and let ρ be a smooth H-representation over some (commutative) ring R (with unit). Then we can consider the induction $\text{Ind}_{H}^{G} \rho$ and the compact induction c-Ind $_G^G \rho$ of ρ .

We recall the definition of $\text{Ind}_{H}^{G} \rho$ and c-Ind $_{H}^{G} \rho$. Let V be the representation space of ρ . We denote by $\text{IND}_{H}^{G}(\rho)$ the set of functions $f: G \to V$ such that $f(hg) = \rho(h)f(g)$ for any $g \in G$ and $h \in H$. The set $\text{IND}_{H}^{G}(\rho)$ can be equipped with a canonical G-representation structure by right translation. Let $\text{Ind}_{H}^{G} \rho$ be the smooth part of $\text{IND}_{H}^{G}(\rho)$. Moreover, we also denote by c-Ind $_{H}^{G} \rho$ the set of elements $f \in \text{Ind}_{H}^{G} \rho$ such that supp f is compact modulo H. Then we can consider functors Ind_H^G and c-Ind_H^G from the category $\mathcal{M}_R(H)$ of smooth H-representations over R to the category $\mathcal{M}_R(G)$ of smooth G-representations over R.

Let K be another closed subgroup in G. Then we can consider functors $\mathrm{Res}^G_K \mathrm{Ind}^G_H$ and Res_{K}^{G} c-Ind $_{H}^{G}$ from $\mathcal{M}_{R}(H)$ to $\mathcal{M}_{R}(K)$. We also can define other functors $\mathcal{M}_R(H) \to \mathcal{M}_R(K)$ as in [\[3,](#page-8-3) I.5.4] as we recall in the following. First, for $g \in G$ and

an H-representation (ρ, V) , we can construct K-representations $\text{Ind}_{K\cap^{g}H}^{K} \text{Res}_{K\cap^{g}H}^{g} g$ and c-Ind K_{Γ} _{K∩gH} Res_{K∩gH} g _P, where $^gH := gHg^{-1}$ and $^g\rho$ is a gH -representation defined as $^g \rho(h') := \rho(g^{-1}h'g)$ for $h' \in {}^g H$. On the other hand, let $I(H, g, K; \rho)$ be the set of locally constant functions $f: HgK \to V$ such that $f(hx) = \rho(h)f(x)$ for $h \in H$ and $x \in HgK$. The set $I(H, g, K; \rho)$ can be equipped with a canonical K-representation structure by right translation. Let $I^{\infty}(H, g, K; \rho)$ be the smooth part of $I(H, g, K; \rho)$. Moreover, we also denote by $I_c^{\infty}(H, g, K; \rho)$ the set of elements $f \in I^{\infty}(H, g, K; \rho)$ such that supp f is compact modulo H. Then $\rho \mapsto I^{\infty}(H, g, K; \rho)$ and $\rho \mapsto I_c^{\infty}(H, g, K; \rho)$ are functors $\mathcal{M}_R(H) \to \mathcal{M}_R(K)$.

The Mackey decomposition is a theorem which relates these functors when H and K are "nice". The following proposition is proved by arguments in $[3, 1.5.4]$.

Proposition 2.1. Let G be a locally profinite group, and let H and K be closed subgroups in G. Suppose HgK is open in G for any $g \in G$. Moreover, let R be a commutative ring with unit, and let ρ be an H-representation over R.

- (1) Let $\phi_{H,g,K} : \text{IND}_H^G(\rho) \to I(H,g,K;\rho)$ be a map defined by the restriction of functions $f \mapsto f|_{HgK}$. Then the map $\phi := \prod_{g \in H \backslash G/K} \phi_{H,g,K}$: IND $_H^G(\rho) \rightarrow$ $\prod_{g\in H\backslash G/K}I(H,g,K;\rho)$ induces the following inclusions of K -representations:
	- (a) $\operatorname{Res}_{K}^{G} \operatorname{Ind}_{H}^{G} \rho \hookrightarrow \prod_{g \in H \backslash G/K} I^{\infty}(H, g, K; \rho);$
	- (b) Res^G_K c-Ind $^G_H \rho \hookrightarrow \bigoplus_{g \in H \backslash G/K} I_c^{\infty}(H, g, K; \rho)$.
- (2) For $g \in G$, the map $I(H, g, K; \rho) \to \text{IND}_{g^{-1}H \cap K}^{K} \text{Res}_{g^{-1}H \cap K}^{g^{-1}H} g^{-1} \rho$ defined as $f \mapsto [k \mapsto f(gk)]$ induces an inclusion of K-representations

$$
I^{\infty}(H,g,K;\rho)\hookrightarrow \text{Ind}_{K\cap {{}^{g^{-1}}H}}^{K}\text{Res}^{{}^{g^{-1}}H}_{K\cap {{}^{g^{-1}}H}}\,g^{{}^{-1}}\rho.
$$

(3) We have a canonical inclusion of K-representations

$$
\text{Res}^G_K\text{Ind}^G_H\rho\hookrightarrow \prod_{g\in K\backslash G/H}\text{Ind}^K_{K\cap^gH}\text{Res}^{^gH}_{K\cap^gH}^{\ g}\rho.
$$

Next, we consider these inclusions with some assumption on topological groups. We apply the assumption in the case as in Proposition [2](#page-2-0).1. In particular, the set HgK is open in G for any $g \in G$.

Assumption 2.2. For any open subgroup H' in H , any open subgroup K' in K and $g \in G$, the set $H'gK'$ is open in G.

Proposition [2](#page-2-0).3. Let G , H, K, R and ρ be as in Proposition 2.1. Moreover, suppose that Assumption [2](#page-2-1).2 holds.

- (1) The inclusion in Proposition 2.[1](#page-2-0)[\(2\)](#page-2-2) is isomorphic.
- (2) The isomorphism [\(1\)](#page-2-3) induces a K-isomorphism

$$
I_c^{\infty}(H,g,K;\rho)\cong\operatorname{c-Ind}_{K\cap {^{g^{-1}}H}}^K\operatorname{Res}_{K\cap {^{g^{-1}}H}}^{{^{g^{-1}}H}}\rho.
$$

(3) The inclusion in Proposition 2.[1\(](#page-2-0)[3\)](#page-2-4) induces a canonical inclusion of smooth K-representations

$$
\mathrm{Res}^G_K\operatorname{c-Ind}^G_H \rho \hookrightarrow \bigoplus_{g\in K\backslash G/H}\operatorname{c-Ind}^K_{K\cap^g H} \mathrm{Res}^{^g H}_{K\cap^g H} {^g \rho}.
$$

Proof. The claim [\(3\)](#page-2-5) follows from [\(2\)](#page-2-6) and Proposition [2.1](#page-2-0) [\(1b\)](#page-2-7).

We consider [\(1\)](#page-2-3). Let V be the representation space for ρ . For a map $f: K \to V$ such that $f(hk) = g^{-1} \rho(h) f(k)$ for any $h \in g^{-1} H g \cap K$ and $k \in K$, we can obtain a well-defined map \hat{f} : $HgK \to V$ as $\hat{f}(hgk) = \rho(h)f(k)$ for $h \in H$ and $k \in K$. The map \hat{f} is the unique map $HgK \to V$ such that $\hat{f}(gk) = f(k)$ and $\hat{f}(hx) = \rho(h)\hat{f}(x)$ for any $h \in H, k \in K$ and $x \in HgK$. Moreover, if f is K-smooth, then \hat{f} is also Ksmooth. Then, to show the inclusion in Proposition [2.1](#page-2-0) [\(2\)](#page-2-2) is surjective, it suffices to show that $\hat{f} \in I^{\infty}(H, g, K; \rho)$ for all $f \in \text{Ind}_{K\cap g^{-1}H}^{K} \text{Res}_{K\cap g^{-1}H}^{g^{-1}H} g^{-1} \rho$, that is, the map \hat{f} is locally constant.

Let $x \in HgK$. Then we have $x = hgk$ for some $h \in H$ and $k \in K$, and so $\hat{f}(x) = \rho(h)f(k)$ by the definition of \hat{f} . Since ρ is smooth, there exists an open subgroup H' in H that fixes $f(k)$. On the other hand, the map f is invariant under the right translation by some open subgroup K'' in K. When we put $K' = kK''k^{-1}$, the subgroup K' is also open in K. By Assumption [2.2,](#page-2-1) the set $H'gK'$ is open in G. Then $hH'gkK'' = hH'gK'k$ is an open neighborhood of $h g k = x$ in G. Moreover, for $h' \in H'$ and $k'' \in K''$, we have $\hat{f}(hh'gkk'') = \rho(hh')f(kk'') = \rho(h)\rho(h')f(k) =$ $\rho(h)f(k) = \hat{f}(x)$. Therefore, the map \hat{f} is constant on $hH'gkK''$, and so \hat{f} is locally constant.

To show [\(2\)](#page-2-6), we consider the continuous bijection $\overline{\iota}: (g^{-1}Hg \cap K)\backslash K \to H\backslash HgK$ induced by $\iota: K \to HgK$; $k \mapsto gk$. By Assumption [2.2,](#page-2-1) for any open subgroup K' in K, the set HgK' is open in G. In particular, $H\backslash HgK'$ is open in HgK , which implies that $\overline{\iota}$ is an open map and so $\overline{\iota}$ is homeomorphic.

Here, let $f \in \text{Ind}_{K \cap g^{-1}H}^{K} \text{Res}_{K \cap g^{-1}H}^{g^{-1}H}$ g^{-1} . If we prove $Hg \text{supp } f = \text{supp }\hat{f}$ for any f, we have $\bar{\iota}((g^{-1}Hg \cap K) \setminus \mathrm{supp} f) = H \setminus Hg \mathrm{supp} f = H \setminus \mathrm{supp} \hat{f}$. Since $\bar{\iota}$ is homeomorphic, the set $(g^{-1}Hg \cap K)\backslash K$ is compact if and only if $H\backslash$ supp \hat{f} is compact, which implies [\(2\)](#page-2-6). Then, it suffices to show Hg supp $f = \sup \hat{f}$.

Let $x \in Hg$ supp f. Then we have $x = hgk$ for some $h \in H$ and $k \in \text{supp } f$. Since f is locally constant, we have $f(k) \neq 0$. Therefore $\hat{f}(x) = \rho(h)f(k) \neq 0$ since $\rho(h)$ is isomorphic, and so $x \in \text{supp} \hat{f}$. On the other hand, let $x \in \text{supp} \hat{f}$. Since $x \in \text{supp} \hat{f} \subset HgK$, there exist $h \in H$ and $k \in K$ such that $x = hgk$. We have $\hat{f}(x) \neq 0$ since \hat{f} is locally constant. Moreover, we also have $\hat{f}(x) = \hat{f}(h g k)$ $\rho(h)f(k)$ by definition of \hat{f} . Since $\rho(h)$ is isomorphic, we obtain $f(k) \neq 0$, and so $k \in \text{supp } f$. Therefore we have $x \in Hg$ supp f, which completes the proof. \Box

We consider sufficient conditions to satisfy the statement of Assumption [2.2.](#page-2-1)

Lemma [2](#page-2-0).4. Let G, H, K be as in Proposition 2.1. Moreover, suppose that (G, H, K) satisfies either of the following:

- (1) either H or K is open in G ;
- (2) the group G is second-countable.

Then Assumption [2](#page-2-1).2 holds. In particular, if G is the group of F-rational points of some linear algebraic group over a non-archimedean local field F, then there exist the inclusions and isomorphisms as in Proposition [2](#page-2-8).3.

Proof. Suppose H is open in G . Then any open subgroup H' in H is also open in G, and so $H'gK'$ is open in G for any subset K' in K and $g \in G$. We can also prove the lemma for the case that K is open in G similarly.

Suppose G is second-countable. Then H, K and HqK are also second-countable. Let H' (resp. K') be an open subgroup in H (resp. K). Since H, K and HgK are closed in G , the set HgK is locally compact, Hausdorff and totally disconnected. Then there exists a compact open subgroup H'' (resp. K'') in H' (resp. K'). If $H''gK''$ is open in HgK , then $H'gK'$, which is a union of open subsets in G which are of the form $h'H''gK''k'$ for some $h' \in H'$ and $k' \in K'$, is also open in HgK . Since HgK is open in G, the subset $H'gK'$ is also open in G. Therefore, we may assume H' and K' are compact.

Since H and K are second-countable, we have $H = \bigcup_{i \in \mathbb{N}} h_i H'$ and $K = \bigcup_{j \in \mathbb{N}} K' k_j$ for some $h_i \in H$ and $k_j \in K$. Then $HgK = \bigcup_{i,j} h_i H'gK'k_j$. Since $H'gK'$ is compact, the subset $H'gK'$ is closed in HgK as HgK is Hausdorff. Therefore, by the Baire category theorem, there exist $(i, j) \in \mathbb{N}^2$ and a nonempty open subset U in HgK such that $U \subset h_i H' gK' k_j$. Then $U' := h_i^{-1} U k_j^{-1}$ is a nonempty subset in $H'gK'$ which is open in G. Since $H'gK' = H'U'K'$, the set $H'gK'$ is open in HgK and G .

2.2. Mackey decomposition and some counterexamples. In general, the inclusion [\(3\)](#page-2-4) in Proposition [2](#page-2-0).1 is not necessarily isomorphic, as shown later. Our main proposition gives us some sufficient conditions to make the inclusions in Proposition 2.[1](#page-2-0) and Proposition 2.[3](#page-2-8) isomorphic.

Proposition 2.5 (Mackey decomposition). Let G, H, K, R and ρ be as in Proposition [2](#page-2-0).1.

(1) Suppose K is open. Then we have

$$
\operatorname{Res}^G_K\operatorname{Ind}^G_H \rho \cong \left(\prod_{g\in K\backslash G/H}\operatorname{Ind}^K_{K\cap^g H} \operatorname{Res}^{^g H}_{K\cap^g H} {^g \rho}\right)^{\infty},
$$

where for a K-representation τ , the representation τ^{∞} is the K-smooth part of τ.

(2) Suppose either H or K is open in G. Then we have the Mackey decomposition

$$
\operatorname{Res}^G_K \operatorname{c-Ind}^G_H \rho \cong \bigoplus_{g \in K \backslash G/H} \operatorname{c-Ind}^K_{K \cap^g H} \operatorname{Res}^g_{K \cap^g H}^g \rho.
$$

Proof. We show [\(1\)](#page-4-2). Since G-smooth representations are also K-smooth, the inclusion [\(1a\)](#page-2-9) in Proposition [2.1](#page-2-0) induces an inclusion

$$
\operatorname{Res}^G_K\operatorname{Ind}^G_H\rho\hookrightarrow \left(\prod_{g\in H\backslash G/K}I^\infty(H,g,K;\rho)\right)^\infty
$$

.

By [\(1\)](#page-2-3) in Proposition [2.3](#page-2-8) and Lemma [2.4,](#page-3-0) it is enough to show that the above inclusion is surjective. Let $(f_g)_{g\in H\backslash G/K}\in \left(\prod_{g\in H\backslash G/K}I^{\infty}(H,g,K;\rho)\right)^{\infty}$. Then there exists an open subgroup K' in K such that K' fixes f_g for all $g \in H \backslash G/K$. We define a map $f: G \to V$ as $f(x) = f_g(x)$ for $x \in HgK$. Then $f \in \text{IND}_{H}^{G} \rho$ and $\phi(f) = (f_g)_{g \in H \backslash G/K}$. Therefore, it is enough to show that $f \in \text{Ind}_{H}^{G} \rho$, that is, the element f is G -smooth.

6 YUKI YAMAMOTO

Since K is open in G , the open subgroup K' in K is also open in G. We show that K' fixes f, which implies f is G-smooth. Let $x \in G$. Then $x \in HgK$ for some double coset HgK . For $k' \in K'$, we have $f(xk') = f_g(xk')$ as $xk' \in HgKk' = HgK$. Since f_g is fixed by K', we also have $f_g(xk') = f_g(x) = f(x)$.

We show [\(2\)](#page-4-3). It is enough to show that the inclusion [\(1b\)](#page-2-7) in Proposition [2.1](#page-2-0) is surjective. Let $g \in G$, and let $f \in I_c^{\infty}(H, g, K; \rho)$. Let V be the representation space of ρ . We define a map $\tilde{f}: G \to V$ as

$$
\tilde{f}(x) = \begin{cases} f(x) & (x \in HgK) \\ 0 & (x \notin HgK). \end{cases}
$$

Then $\tilde{f} \in \text{IND}_{H}^{G} \rho$ and $\phi(\tilde{f}) = f$. Moreover, we have supp $\tilde{f} = \text{supp} f$, where the latter is compact modulo H. Therefore, it suffices to show that \tilde{f} is a smooth element in $\text{IND}_{H}^{G}\rho$, that is, there exists an open subgroup J in G such that \tilde{f} is J-invariant.

Suppose K is open. Then any open subgroup K' in K which fixes f is also open in G. By the assumption of K', we have $\tilde{f}(x k') = f(x k') = f(x) = \tilde{f}(x)$ for any $x \in HgK$ and $k' \in K'$. Moreover, for this K', we have $(G \setminus HgK) \cdot K' = G \setminus HgK$, which implies $\tilde{f}(x k') = 0 = \tilde{f}(x)$ for any $x \in G \setminus HgK$ and $k' \in K'$. Therefore it is enough to put $J = K'$.

Suppose H is open. Then the set $H \setminus \text{supp } f = H \setminus \text{supp } f$ is discrete. Since $H\setminus \text{supp } f$ is also compact, the set $H\setminus \text{supp } \tilde{f}$ is finite. Therefore, there exist finitely many elements $x_1, \ldots, x_n \in G$ such that supp \tilde{f} is the disjoint union $\bigcup_{i=1}^n Hx_i$. For any *i*, there exists a compact open subgroup H_i in H which fixes $f(x_i)$. We put $J := \bigcap_{i=1}^n x_i^{-1} H_i x_i$, which is an open subgroup in G.

We will show \hat{f} is J-invariant. First, we will show supp $\hat{f} \cdot j = \sup \hat{f}$ for any $j \in J$. We have supp $\tilde{f} \cdot j = \bigcup_{i=1}^n Hx_i j$. For any i, there exists $h'_i \in H_i$ such that $j = x_i^{-1} h_i x_i$ since $J = \bigcap_{i=1}^n x_i^{-1} H_i x_i$. Then we obtain $\bigcup_{i=1}^n H x_i j = \bigcup_i H x_i$. $x_i^{-1}h_ix_i = \bigcup_i Hx_i = \text{supp }\tilde{f}$. In particular, we also have $(G\sup p \tilde{f})\cdot J = G\sup p \tilde{f}$, which implies that $\tilde{f}(xj) = 0 = \tilde{f}(x)$ for any $x \in G\$ supp \tilde{f} and $j \in J$. On the other hand, let $x \in \text{supp }\tilde{f}$ and $j \in J$. Then there exists i such that $x = hx_i$ for some $h \in H$ and $j = x_i^{-1}h_ix_i$ for some $h_i \in H_i$. Therefore, we have $\tilde{f}(xj) = \tilde{f}(hh_ix_i)$ $\rho(hh_i)\tilde{f}(x_i)$. Here, we have $\rho(hh_i)\tilde{f}(x_i) = \rho(h)(\rho(h_i)f(x_i)) = \rho(h)f(x_i)$ since $h_i \in H_i$ fixes $f(x_i)$. Then we obtain $\tilde{f}(x_j) = \rho(h)f(x_i) = f(hx_i) = f(x) = \tilde{f}(x)$, which completes the proof. \Box

In the following, we construct some example such that the inclusion [\(1a\)](#page-2-9) in Proposition [2.1](#page-2-0) is not surjective.

Example 2.6. Let F be a non-archimedean local field, and let E/F be a ramified quadratic extension of fields. We denote by \mathfrak{o}_F (resp. \mathfrak{o}_E) the ring of integers in F (resp. E.) We also denote by \mathfrak{p}_F (resp. \mathfrak{p}_E) the maximal ideal in \mathfrak{o}_F (resp. \mathfrak{o}_E .) We put $G = GL_2(E)$ and $K = GL_2(F)$. We also put $H = \begin{pmatrix} 1 + \mathfrak{p}_E & \mathfrak{p}_E \\ \mathfrak{p}_E & 1 + \mathfrak{p}_E \end{pmatrix}$ $\mathfrak{p}_{E} \qquad 1+\mathfrak{p}_{E}$ $\bigg),$ which is a compact open subgroup in G. Let (ρ, R) be the trivial representation of H for some commutative ring R with unit. Then G, H, K, R and ρ satisfies the condition in Proposition [2](#page-2-0).1.

We define $f: HK \to R$ as $f(x) = 1$ for $x \in HK$. Then $f \in I^{\infty}(H, 1_G, K; \rho) \subset$ $\left(\prod_{g\in H\backslash G/K}I^{\infty}(H,g,K;\rho)\right)^{\infty}$, and we can define $\tilde{f}\in \text{IND}_{H}^{G}\rho$ such that $\phi(\tilde{f})=f$

as in the proof of Proposition [2](#page-4-1).5. Here, by the definition of \tilde{f} , the function \tilde{f} is the characteristic function of HK in G .

In the following, we will show that $\tilde{f} \notin \text{Ind}_{H}^{G} \rho$, that is, the function \tilde{f} is not fixed by any open subgroup U in G. Since \tilde{f} is the characteristic function of HK in G, it is enough to show that $HKU \neq HK$ for any open subgroup U in G. Let U be an open subgroup in G. We put $u_n = \begin{pmatrix} 1 & \varpi_E^{2n+1} \\ 0 & 1 \end{pmatrix}$ for $n \in \mathbb{N}$, where we fix a uniformizer ϖ_E of E. Since $\lim_{n\to\infty} u_n = 1_G$, we have $u_n \in U$ for some $n \in \mathbb{N}$. Then $\begin{pmatrix} 1 & 0 \\ 0 & -n \end{pmatrix}$ 0 ϖ_F^{n+1} $\bigg\}\bigg\|u_n\in KU\subset HKU$, where we also fix a uniformizer ϖ_F of F. Here, suppose that $HKU = HK$. Then we have $\begin{pmatrix} 1 & \varpi_F^{-(n+1)} \varpi_E^{2n+1} \\ 0 & 1 \end{pmatrix} =$ $(1 \ 0)$ $0 \quad \varpi_F^{n+1}$ $\bigg\}\,u_n\cdot\left(\begin{array}{cc} 1 & 0 \\ 0 & -\frac{1}{2}\end{array}\right)$ $\begin{pmatrix} 1 & 0 \ 0 & \varpi_F^{-(n+1)} \end{pmatrix} \in HKU \cdot K = HK.$ Therefore, there exists F $h = \begin{pmatrix} 1+a & b \\ 0 & 1 \end{pmatrix}$ $c \quad 1+d$ $\left(\begin{array}{cc} 1 & \overline{\omega}_F^{-(n+1)}\overline{\omega}_E^{2n+1} \\ 0 & 1 \end{array}\right) \in K.$ However, we have $h\begin{pmatrix} 1 & \varpi_F^{-(n+1)}\varpi_E^{2n+1} \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} x' & (1+a)\varpi_F^{-(n+1)}\varpi_E^{2n+1} + b \\ z' & w' \end{pmatrix}$ z' w' $\bigg)$ for some $x', z', w' \in E$. When we define v_E as the normalized valuation $E \to \mathbb{Z} \cup \{\infty\},$ we have $v_E\left((1+a)\varpi_F^{-(n+1)}\varpi_E^{2n+1}+b\right)=-1$ since E/F is ramified quadratic and $a, b \in \mathfrak{p}_E$. Since $-1 \notin 2\mathbb{Z} \cup {\infty} = v_E(F)$, we have $(1+a)\varpi_F^{-(n+1)}\varpi_E^{2n+1} + b \notin F$, which is a contradiction.

The Mackey decomposition leads to the following corollary on intertwining operators.

Corollary 2.7. Let G be a locally profinite group, H be a closed subgroup in G , and let K be an open subgroup in G. Let σ (resp. τ) be a smooth representation of K (resp. H) over some commutative ring R with unit.

- (1) We have $\text{Hom}_G(\text{c-Ind}_K^G \sigma, \text{Ind}_H^G \tau) \cong \prod_{g \in K \backslash G/H} \text{Hom}_{K \cap^g H}(\sigma,^g \tau).$
- (2) Moreover, we further assume that $H\backslash HgK$ is compact for any $g \in G$ and σ is finitely generated as a K-representation. Then we also have $\text{Hom}_G(\text{c-Ind}_K^G \sigma, \text{c-Ind}_H^G \tau) \cong \bigoplus_{g \in K \backslash G/H} \text{Hom}_{K \cap^g H}(\sigma,^g \tau).$

Proof. We show [\(1\)](#page-6-0). Since K is open in G, we have $\text{Hom}_G(c\text{-}\text{Ind}_K^G \sigma, \text{Ind}_H^G \tau) \cong$ $\text{Hom}_K(\sigma, \text{Res}_K^G \text{Ind}_H^G \tau)$ by Frobenius reciprocity for compact induction. By Propo-sition [2.5](#page-4-1) [\(1\)](#page-4-2), we also have $\text{Res}_{K}^{G} \text{Ind}_{H}^{G} \tau \cong \left(\prod_{g \in K \setminus G/H} \text{Ind}_{K \cap^{g} H}^{K} \text{Res}_{K \cap^{g} H}^{g} g \tau\right)^{\infty}$. Since σ is smooth, we obtain

$$
\text{Hom}_{K}(\sigma, \text{Res}_{K}^{G} \text{Ind}_{H}^{G} \tau) \cong \text{Hom}_{K} \left(\sigma, \left(\prod_{g \in K \backslash G/H} \text{Ind}_{K \cap^{g} H}^{K} \text{Res}_{K \cap^{g} H}^{g} \sigma \right)^{\infty} \right)
$$
\n
$$
\cong \text{Hom}_{K} \left(\sigma, \prod_{g \in K \backslash G/H} \text{Ind}_{K \cap^{g} H}^{K} \text{Res}_{K \cap^{g} H}^{g} \sigma \right)
$$
\n
$$
\cong \prod_{g \in K \backslash G/H} \text{Hom}_{K}(\sigma, \text{Ind}_{K \cap^{g} H}^{K} \text{Res}_{K \cap^{g} H}^{g} \sigma \tau).
$$

By Frobenius reciprocity for induction, we obtain

$$
\text{Hom}_{K}(\sigma, \text{Ind}_{K\cap^{g}H}^{K}\text{Res}_{K\cap^{g}H}^{g}{}_{H})\cong \text{Hom}_{K\cap^{g}H}(\sigma, {^g\tau})
$$

and complete the proof.

We show [\(2\)](#page-6-1). We have $\text{Hom}_G(c\text{-}\text{Ind}_K^G \sigma, c\text{-}\text{Ind}_H^G \tau) \cong \text{Hom}_K(\sigma, \text{Res}_K^G c\text{-}\text{Ind}_H^G \tau)$ as [\(1\)](#page-6-0). Here, we also have Res_{K}^{G} c-Ind $_{H}^{G} \tau \cong \bigoplus_{g \in K \backslash G/H} C$ -Ind $_{K \cap g}^{K}$ Res $_{K \cap g}^{g}$ H Proposition [2.5](#page-4-1) [\(2\)](#page-4-3). Since σ is finitely generated, we obtain

$$
\text{Hom}_G(\text{c-Ind}_{K}^G \sigma, \text{c-Ind}_{H}^G \tau) \cong \text{Hom}_K \left(\sigma, \bigoplus_{g \in K \backslash G/H} \text{c-Ind}_{K \cap {}^g H}^K \text{Res}_{K \cap {}^g H}^g \tau \right)
$$

$$
\cong \bigoplus_{g \in K \backslash G/H} \text{Hom}_K \left(\sigma, \text{c-Ind}_{K \cap {}^g H}^K \text{Res}_{K \cap {}^g H}^g \tau \right).
$$

Here, since K is open, the continuous bijection $(K \cap {}^g H) \backslash K \to g H g^{-1} \backslash g H g^{-1} K \cong$ $H\backslash Hg^{-1}K$ is homeomorphic. Then, by the assumption that $H\backslash Hg^{-1}K$ is compact, the set $(K \cap {}^{g}H)\backslash K$ is also compact, and so the functor c-Ind ${}_{K\cap {}^{g}H}^{K}$ is equal to Ind $K_{\bigcap^g H}$. Therefore we obtain

$$
\text{Hom}_{K}\left(\sigma, \text{c-Ind}_{K\cap^{g}H}^{K} \text{Res}_{K\cap^{g}H}^{g} g_{\tau}\right) = \text{Hom}_{K}\left(\sigma, \text{Ind}_{K\cap^{g}H}^{K} \text{Res}_{K\cap^{g}H}^{g} g_{\tau}\right)
$$

$$
\cong \text{Hom}_{K\cap^{g}H}(\sigma, g_{\tau}),
$$

where the last equation follows from Frobenius reciprocity for induction. \Box

Remark 2.8. (1) When $R = \mathbb{C}$ and σ is a finite-dimensional representation of K, this corollary is proved by Kutzko $[1,$ Theorem 1.1].

(2) In the statement of [\[1,](#page-8-2) Theorem 1.1], line 8, the word " $I(\sigma_c^G, \tau^G)$ " seems to be a typographical error and seems to need to be replaced with " $I(\sigma_c^G, \tau_c^G)$ " based on the proof. Even though this typo is corrected, the assumption that the image of K in $H\backslash G$ is compact is not sufficient to make the theorem hold. The following example is a counterexample for the theorem.

Example 2.9. Let $\varphi: \mathbb{Z}/2\mathbb{Z} \to \text{Aut}(\mathbb{Z}^2)$ be the group homomorphism defined as $\varphi(1+2\mathbb{Z})(n_1,n_2)=(n_2,n_1)$ for $(n_1,n_2)\in\mathbb{Z}^2$. We put $G=\mathbb{Z}^2\rtimes_{\varphi}(\mathbb{Z}/2\mathbb{Z})$ with the discrete topology. In particular, G is a locally profinite group. We put $H =$ $K = (\mathbb{Z} \times \{0\}) \rtimes \{0\}.$ We also put $g_n := ((0, n), 0) \in G$ for $n \in \mathbb{Z}$ and $g' :=$ $((0,0), 1+2\mathbb{Z}) \in G$. Moreover, let χ_1, χ_2 be distinct characters $H \to \mathbb{C}^{\times}$.

We check that the tuple $(G, H, K, R = \mathbb{C}, \sigma = \chi_1, \tau = \chi_2)$ satisfies the condition which is assumed in Kutzko's statement. First, H is a closed subgroup in G and $K = H$ contains a compact open subgroup $\{1\}$ in G since G is discrete. Next, $\sigma = \chi_1$ and $\tau = \chi_2$ are smooth, finite-dimensional representations of $K = H$. Moreover, the image of K to $H\backslash G$ is $H\backslash HK = H\backslash H$, which is compact.

Here, we consider the set of double cosets $K\backslash G/H$. For $n \in \mathbb{Z}$, we have $Kg_nH =$ $(\mathbb{Z} \times \{n\}) \rtimes \{0\}$. On the other hand, we also have $Kg'H = \mathbb{Z}^2 \rtimes \{1+2\mathbb{Z}\}$. Therefore, the set G is the disjoint union of Kg_nH for $n \in \mathbb{Z}$ and $Kg'H$, that is, the set ${g_n | n \in \mathbb{Z}} \cup {g'}$ is a representative of $K\backslash G/H$.

We show $\text{Hom}_G(\text{c-Ind}_K^G \sigma, \text{c-Ind}_H^G \tau) \not\cong \bigoplus_{g \in K \backslash G/H} \text{Hom}_{K \cap^g H}(\sigma,^g \tau)$. On the left-hand side, we have

$$
\text{Hom}_G(\text{c-Ind}_{K}^G \sigma, \text{c-Ind}_{H}^G \tau) \cong \bigoplus_{g \in K \backslash G / H} \text{Hom}_K(\sigma, \text{c-Ind}_{K \cap g}^K \text{Res}_{K \cap g}^{g} H^{g} \tau)
$$

as in the proof of Corollary 2.[7](#page-6-2)[\(2\)](#page-6-1). If $g \in Kg_nH$, then g centralizes H, and so $^gH =$ H and ${}^g\tau = \tau$. Then we have $\text{Hom}_K(\sigma, c\text{-}\text{Ind}_{K\cap {}^g H}^K \text{Res}_{K\cap {}^g H}^{\mathcal{I} H} {}^{g}\tau) = \text{Hom}_K(\sigma, \tau) =$ 0. If $g \in Kg'H$, then $^gH = (\{0\} \times \mathbb{Z}) \rtimes \{0\}$, and so $K \cap {}^gH = \{1_G\}$. Therefore we $\mathit{have} \ \operatorname{c-Ind}_{K\cap^{g}H}^{\check{K}}\operatorname{Res}^{{}^gH}_{K\cap^{g}H}$ $^g\tau=\operatorname{c-Ind}_{\{1_G\}}^{\check{K}}\mathbf{1}\cong \operatorname{c-Ind}_{\{*\}}^{\mathbb{Z}}\mathbf{1}=\{f\colon \mathbb{Z}\to \mathbb{C}\mid |\operatorname{supp} f|<\infty\}$ $\{\infty\}$. Then to show Hom_K(σ , c-Ind $K \cap H \otimes_{K \cap H} H^{\circ}$ σ) = 0, it suffices to show that c-Ind $\mathbb{Z}_{\{*\}}$ 1 does not have any nontrivial, finite-dimensional subrepresentation of \mathbb{Z} . Suppose W is a nonzero, finite-dimensional subspace in c-Ind $\mathbb{Z}_{\{*\}}$ 1. Then the min- $\lim_{n_0} n_0 := \min\{n \in \mathbb{Z} \mid f(n) \neq 0 \text{ for some } f \in W\}$ exists. Let $f \in W$ such that $f(n_0) \neq 0$. Then, we have $((c\text{-Ind}_{\{*\}}^{\mathbb{Z}} 1(1)) f)(n_0 - 1) = f((n_0 - 1) + 1) =$ $f(n_0) \neq 0$. By the definition of n_0 , we obtain $\left(\text{c-Ind}_{\{*\}}^{\mathbb{Z}} \mathbf{1}(1)\right) f \notin W$. In particular, W is not invariant by the action of \mathbb{Z} , that is, c-Ind $\tilde{\ell}_{\ast}^{\sharp}$ 1 does not have any nontrivial, finite-dimensional subrepresentations of Z. Therefore we obtain $\text{Hom}_G(\text{c-Ind}_K^G \sigma, \text{c-Ind}_H^G \tau) = 0.$

On the right-hand side, let $g \in Kg_nH$ for some $n \in \mathbb{Z}$. Then, as the left-hand side, we have $K \cap {}^g H = H$, ${}^g \tau = \tau$ and so $\text{Hom}_{K \cap {}^g H}(\sigma, {}^g \tau) = \text{Hom}_H(\sigma, \tau) =$ 0. On the other hand, suppose that $g \in Kg'H$. Then $K \cap {}^{g}H = \{1_G\}$, and so $\text{Hom}_{K\cap^{g}H}(\sigma,{}^{g}\tau) = \text{Hom}_{\mathbb{C}}(\mathbb{C},\mathbb{C}) \cong \mathbb{C}.$ Therefore, $\bigoplus_{g\in K\backslash G/H} \text{Hom}_{K\cap^{g}H}(\sigma,{}^{g}\tau) \cong$ $\mathbb{C} \neq 0$.

REFERENCES

- [1] P. C. Kutzko, *Mackey's theorem for nonunitary representations*, Proc. Amer. Math. Soc. 64 (1977), 173-175.
- [2] G. W. Mackey, On induced representations of groups, Amer. J. Math. 73 (1951), 576-592. MR 13, 106.
- [3] M.-F. Vignéras, Représentations ℓ -modulaires d'un groupe réductif p-adique avec $\ell \neq p$, Progress in mathematics 137. Birkhauser 1996.

Yuki Yamamoto

Graduate School of Mathematical Sciences, The University of Tokyo, 3–8–1 Komaba, Meguro-ku, Tokyo, 153–8914, Japan

E-mail address: yukiymmt@ms.u-tokyo.ac.jp