Robust and Efficient Bayesian Inference for Non-Probability Samples

Ali Rafei¹, Michael R. Elliott^{*1}, and Carol A. C. Flannagan²

¹Survey Methodology Program, University of Michigan ²University of Michigan Transportation Research Institute

Abstract: The declining response rates in probability surveys along with the widespread availability of unstructured data has led to growing research into non-probability samples. Existing robust approaches are not well-developed for non-Gaussian outcomes and may perform poorly in presence of influential pseudo-weights. Furthermore, their variance estimator lacks a unified framework and rely often on asymptotic theory. To address these gaps, we propose an alternative Bayesian approach using a partially linear Gaussian process regression that utilizes a prediction model with a flexible function of the pseudo-inclusion probabilities to impute the outcome variable for the reference survey. By efficiency, we mean not only computational scalability but also superiority with respect to variance. We also show that Gaussian process regression behaves as a kernel matching technique based on the estimated propensity scores, which yields double robustness and lowers sensitivity to influential pseudo-weights. Using the simulated posterior predictive distribution, one can directly quantify the uncertainty of the proposed estimator and derive associated 95% credible intervals. We assess the repeated sampling properties of our method in two simulation studies. The application of this study deals with modeling count data with varying exposures under a non-probability sample setting.

Keywords and phrases: Non-probability sample, doubly robust, semi-parametric Bayesian modeling, Gaussian process regression, count data.

1. Introduction

The declining response rates in probability surveys increasingly challenge the validity of this long-standing touchstone for finite population inference (Groves, 2011; Johnson and Smith, 2017; Miller, 2017; Neyman, 1934). According to a report by Pew Research Center, the average response rate in telephone surveys has dropped by 75% over the past two decades (Keeter et al., 2017). Unless additional effort and cost is expended, a similar trend is expected to hold for in-person household surveys. Researchers speculate that multiple factors, including the rising response burden from a multitude of surveys with lengthy and sophisticated instruments, busier-than-ever lifestyles, and increased privacy concerns, contribute to this downward trend (Brick and Williams, 2013; Williams and Brick, 2018). It is perhaps because of this issue that pollsters increasingly fail to predict the outcome of the political elections in the U.S. (Forsberg, 2020; Vittert et al., 2020). Although the emergence of responsive and adaptive survey designs has paved novel routes to maximize the response propensity by design (Brick and Tourangeau, 2017; Groves and Heeringa, 2006; Tourangeau et al., 2017), these approaches may not remain efficient forever as the cost of refusal conversion continues to rise (Luiten et al., 2020).

At the same time, use of non-probability sample "Big Data" is gaining popularity increasingly in various domains. These massive unstructured data are often accumulated naturally as a byproduct of human activities. Examples include but not limited to administrative and transactional records, social media, and sensor data (Johnson and Smith, 2017). Being cheaper and faster to collect than probability sample data has led to growing interest in using this wide range of data for producing official statistics (Beresewicz et al., 2018; Groves, 2011). However, the organic nature of their data-generating process, mainly due to self-selection, makes big-data-based inferences prone to selection bias (Johnson and Smith, 2017; Kreuter

^{*} Corresponding author; address: 426 Thompson St. Ann Arbor, MI 48109. Rm 4068 ISR, email: mrelliot@umich.edu.

and Peng, 2014). For a biased sample with an unknown selection mechanism, larger data volume even amplifies the relative contribution of bias to total error, which extremely reduces the effective sample size (Keiding and Louis, 2016; Meng et al., 2018). This heightens the need for novel approaches that optimally calibrate such large-scale unstructured data for finite population inference.

As the motivating application, we are interested in estimating crash rates per distance unit driven for a subpopulation of American drivers. The current estimates are based on a ratio of the annual total police-reported crashes and annual total miles driven obtained from the General Estimates System (GES) (Administration et al., 2014) and the American Driving Survey (ADS), respectively (Tefft, 2017). The denominator, however, can be widely subject to systematic measurement error as it relies on respondents' guess/estimate about their annual driven miles over the telephone (Kim et al., 2019; Prieger, 2004). In contrast, naturalistic driving studies (NDS) are capable of capturing these two quantities simultaneously for each participant by monitoring traffic incidents and kinematic indices continuously via a series of in-vehicle sensors and cameras (Guo et al., 2009). However, because of the high administrative and technical costs, participants of NDS are usually recruited via convenience samples from limited geographical areas. Therefore, naive inference based on such a non-probabilistic sample may suffer from selection bias (Antin et al., 2015; Rafei, 2021).

Consider a two-sample setup where a well-designed probability sample is available with a common set of auxiliary variables, also known as a "reference survey". Under certain assumptions, two general classes of adjustment methods can be followed: (1) quasi-randomization (QR), where the unknown selection probabilities are estimated for units of the non-probability sample (Lee, 2006; Lee and Valliant, 2009; Valliant and Dever, 2011), and (2) prediction modeling (PM), where the analytic variable is predicted for units of the reference survey (Kim et al., 2018; Kim and Rao, 2012; Rivers, 2007; Wang et al., 2015). In either case, design-based approaches can then be utilized to compute point and interval estimates. To further protect against model misspecification, Chen et al. (2019) reconciles the QR approach with the PM method using the idea of augmented inverse propensity weighting (AIPW) (Robins et al., 1994). This method is doubly robust (DR) in the sense that the estimator is consistent if either model holds (Scharfstein et al., 1999).

Although the QR, PM, and AIPW methods all involve modeling, the ultimate form of their estimator is design-based and therefore, are subject to the general drawbacks of the design-based methods. For instance, a presence of outlying pseudo-weights may yield extremely inefficient estimates, especially if the sample size is small (Chen et al., 2017). Furthermore, design-based approaches lack a unified framework for quantifying all sources of uncertainty in the point estimates (Zangeneh, 2012). Existing methods rely on multiple assumptions about the design of the two samples and the distribution of the outcome variable that hold often asymptotically (Chen et al., 2019; Kim et al., 2018). As another limitation, the AIPW method proposed by Chen et al. (2019) requires the sets of variables governing the selection mechanism and response surface to be identical, while this may not be the case in many situations. Furthermore, a unique solution may not necessarily exist for the joint estimating equation systems associated with the two underlying models.

To minimize these limitations, Zheng and Little (2003) propose an alternative class of inferential methods for the probability surveys with a probability proportional-to-size (PPS) design, which is called Penalized Spline of Propensity Prediction (PSPP). Unlike the previously discussed methods, PSPP is fully model-based, in which the outcome variable is predicted for the entire non-sampled units of the population. This method borrows the idea of Linear-in-Weight Prediction (LWP), in which estimated pseudo-weights are specified as a predictor in the outcome model (Zhang and Little, 2009). Bang and Robins (2005) show that an LWP estimator is equivalent to an AIPW estimator in terms of double robustness. In situations where auxiliary variables are missing for the non-sampled units, Little and Zheng (2007) recommend synthesizing multiple populations via finite population Bayesian bootstrapping (FPBB). An and Little (2008) extend this approach to a missing data imputation scenario, where measures of size are replaced by the estimated propensity scores (PS) of being observed, and demonstrate its DR property in a simulation study.

As a likelihood-based method, the PSPP can be implemented under a fully Bayesian framework. This allows for direct estimation of the variance by simulating the posterior predictive distribution of the population parameters. Zangeneh and Little (2015) expands the Bayesian PSPP under a PPS design for situations where the totals of the measures of size are known from external data and where there is evidence of heteroscedasticity with respect to the estimated PS. Further extensions to probability samples with unequal selection probabilities are proposed by Chen et al. (2012). The PSPP is also suitable for situations where the design of the reference survey is complex. For instance, Zhou et al. (2016) develop a synthetic population approach based on a multi-stage cluster survey by undoing the sampling steps through a weighted Pólya posterior distribution. More recently, Tan et al. (2019) compare PSPP with AIPW to make inference for incomplete data, where PS is predicted using Bayesian Additive Regression Trees (BART) and find that the former outperforms in terms of the mean squares error.

While the use of a more flexible non-parametric function of the estimated PS improves when influential pseudo-weights are present (Zhang and Little, 2011), the theoretical rationale for using a penalized spline model among a wider class of smoothers is not quite clear. Saarela et al. (2016) argue that the convergence of the posterior sampling to any well-defined joint distribution of the outcome and PS may be hard to achieve. Alternatively, in this study, we propose to use Gaussian Process (GP) priors to link the PS to the outcome conditional mean (Si et al., 2015). GP is a powerful non-parametric Bayesian tool for functional regression that assigns prior distributions over multidimensional non-linear functions. Because of its flexibility and generalizability, GP is gaining popularity in statistics and machine learning (Kaufman et al., 2010; Neal, 1997; Oakley and O'Hagan, 2004; Shi and Choi, 2011; Wang and Xu, 2019; Williams and Rasmussen, 2006; Yi et al., 2011).

While the correspondence between splines and GP has long been understood (Kimeldorf and Wahba, 1970; Seeger, 2000), the latter can exploit a kernel with infinite basis functions (Williams and Rasmussen, 2006). In this regard, GP may outdo the spline in terms of flexibility while depending on no arbitrary tuning parameters. In a regular spline regression, one has to determine the polynomial order as well as the frequency and location of the knots empirically. More importantly, Huang et al. (2019) demonstrate that a stationary isotropic covariance matrix in GP behaves as a non-parametric matching technique using the estimated PS as a measure of similarity. In a more *ad hoc* manner, Rivers (2007) suggests matching units of a web non-probability survey to those from a parallel reference survey. Very recently, a kernel weighting approach has been proposed by Wang et al. (2020), where the weighted estimator is proved to be consistent under a weak exchangeability condition. To further weaken the modeling assumptions, Kern et al. (2020) propose to use algorithmic tree-based methods, including random forests and gradient tree boosting, for estimating the PS in kernel weighting.

To eliminate the need to account for the sampling weights in the PS model, one possible solution is to generate synthetic populations using the weights in the reference survey (An and Little, 2008; Dong et al., 2014; Zangeneh and Little, 2015). However, fitting Bayesian joint models on a synthetic population can be computationally demanding if not impossible (Mercer, 2018). In addition, one has to rely on Rubin's combining rules to derive the final point and interval estimates. Therefore, direct simulation of the posterior predictive distribution is not possible for the unknown population quantity in this approach. To augment the prediction model in the GPPP estimator while avoiding this technique, we propose an alternative approach that is inspired by Si et al. (2015). In this method, we limit the prediction of outcome to the reference survey, instead of the whole non-sampled units of the population. Having the sampling weights and other design features known in the reference survey, we utilize the posterior predictive draws of the outcome to derive the final point and variance estimates based on Rubin's combining rules (Rubin, 2004).

The rest of the article is organized as follows: In Section 2, we start by describing the proposed method through rigorous mathematical notations. Section 3 assesses the repeated sampling properties of the proposed method and compares its performance with the AIPW estimator through a simulation study. Section 4 involves application on the sensor-based data from Strategic Highway Research Program 2 (SHRP2) where we deal with modeling count data with varying exposures. All the statistical analyses have been performed using R/Stan with annotated codes accessible publicly at

Section 5 reviews the strengths and weaknesses of the s

https://github.com/arafei/GPPP. Finally, Section 5 reviews the strengths and weaknesses of the study in more detail and suggests some future research directions. Supplemental information, including proofs, additional theory, and preliminary descriptive results, are provided in Appendix 8.

2. Methods

2.1. Notation

Consider U to be a finite population of unknown size N. For each i = 1, ..., N, we denote y_i to be the realized values of a scalar outcome variable, Y, in U, and $x_i = [x_{i1}, x_{i2}, ..., x_{ip}]^T$ the values of a p-dimensional set of relevant auxiliary variables, X. Let S_A be a non-probability sample selected from U with y_i being observed, and n_A being the sample size. The main objective of descriptive inference is to learn about an unknown finite population quantity that is a function of Y, e.g. the population mean, $Q(y) = \overline{y}_U = \sum_{i=1}^N y_i/N$. Suppose $\delta_i^A = I(i \in S_A)$ represents the inclusion indicator variable of S_A for $i \in U$, and x_i is the vector of variables associated with the selection mechanism of S_A . The unknown π_i^A for $i \in S_A$ is the biggest challenge to the analyst, and unbiased inference about Q(y) cannot be drawn without imposing a set of strong conditions. Therefore, we consider the following assumptions:

- C1. Positivity—The non-probability sample S_A actually does have a probabilistic sampling mechanism, albeit unknown. That means $p(\delta_i^A = 1|x_i) > 0$ for all possible values of x_i in U.
- **C2. Ignorability**—the selection mechanism of S_A is fully governed by x, which implies that $Y \perp \delta^A | X$. Then, for $i \in U$, the pseudo-inclusion probability associated with S_A is defined as $\pi_i^A = p(\delta_i^A = 1 | x_i)$.
- **C3.** Independence—units in S_A are selected independently given x, i.e. $\delta_i^A \perp \delta_j^A | x_i, x_j$ for $i \neq j \in U$. Note that this assumption is made to avoid unnecessary complications; otherwise extensions can be given for situations where S_A is a clustered sample.

Now, suppose S_R is a parallel "reference survey" of size n_R , for which the same set of covariates, X, has been measured. Note that in a non-probability sample setting, y_i has to be unobserved for $i \in S_R$; otherwise, inference could be directly drawn based on S_R . Also, let $\delta_i^R = I(i \in S_R)$ denotes the inclusion indicator variable associated with S_R for $i \in U$. Units of S_R may be selected independently or through a stratified multistage clustered sampling design. Being a full probability sample implies that the selection mechanism in S_R is ignorable given its design features, i.e. $p(\delta_i^R|y_i, d_i) = p(\delta_i^R|d_i)$ for $i \in U$, where d_i denotes a set of associated design variables. Thus, one can define the inclusion probabilities in S_R as $\pi_i^R = p(\delta_i^R = 1|d_i)$ for $i \in U$. This study deals with reference surveys whose sampling design involves an independent selection of population units but with unequal inclusion probabilities. A common example of such a design is PPS, under which d is regarded as a scalar variable representing measures of size in U, and $\pi_i^R \propto n^R \overline{d}/N\overline{D}$ according to Zangeneh and Little (2015).

Normally, probability surveys come with a set of sampling weights in their public-use data which are inversely proportional to the selection probabilities, i.e. $w_i^R \propto 1/\pi_i^R$. While w_i^{R} 's may comprise of post-survey adjustments for non-response and non-coverage errors (Valliant et al., 2013), not all the auxiliary information used for the construction of weights are necessarily supplied to the analyst. In addition, public-use survey data may lack a detailed guideline on how the sampling weights are calculated. To simplify the problem under such situations, Si et al. (2015) assume that weights with identical values represent a unique post-stratum in the population. Therefore, one can identify d as the indicator of Junique post-strata in U, and consider $w_j^R \propto N_j/n_j^R$ (j = 1, 2, ..., J), where N_j and n_j^R are the j-th post-stratum size in U and S_R , respectively. For instance, in RDD telephone surveys or mail surveys, whose design involves equiprobability sampling, the inequality in weights may arise exclusively from non-response adjustment or post-stratification.

Now, we combine the two samples and define $S_C = S_A \cup S_R$ with $n_C = n_A + n_R$ being the total sample size. While X and D may overlap or correlate, in addition to the aforementioned conditions, we assume

that, conditional on [X, D], S_R and S_A are selected independently, i.e. $\delta^A \perp \delta^R | X, D$. We denote this condition as **C4**. Considering **C1-C4**, the joint density of y_i , δ^A_i and δ^R_i can be factorized as below:

$$p(y_i, \delta_i^A, \delta_i^R | x_i, d_i; \theta, \beta) = p(y_i | x_i, d_i; \theta) p(\delta_i^A | x_i; \beta) p(\delta_i^R | d_i), \quad \forall i \in U$$

$$(2.1)$$

where $\eta = (\theta, \beta)$ are some unknown parameters indexing the conditional distribution of Y|X, D and $\delta^A|X$, respectively. The conditional density $p(y_i|x_i;\theta)$ denotes the underlying model that governs the response surface structure of a superpopulation from which U has been selected. Also, $p(\delta_i^A|x_i;\beta)$ and $p(\delta_i^R|d_i)$ denote the randomization distributions associated with the design of S_A and S_R , respectively. Note that the latter does not depend on any unknown parameter as S_R is a probability sample with a known sampling design. A QR approach involves modeling $p(\delta_i^A|x_i;\beta)$, whereas a PM approach deals with modeling $p(y_i|x_i, d_i;\theta)$. In the following subsection, we describe a fully model-based strategy for Bayesian inference based on non-probability samples by jointly modeling the PS and the outcome variable.

2.2. Bayesian model-based inference

As discussed in the introduction, under a fully model-based approach, the inference paradigm is viewed as imputing the unobserved outcome, Y, for the non-sampled units of the population with respect to S_A , i.e. $\overline{S_A} = U - S_A$. Apparently, one can directly estimate the population unknown quantity as soon as y_i is known for $i \in U$. For the population mean, such an estimator, also known as a *prediction* estimator, is given by

$$\widehat{\overline{y}}_{U} = \left(\sum_{i \in S_{A}} y_{i} + \sum_{i \in \overline{S}_{A}} \widehat{y}_{i}\right) / N$$

$$= \left(\sum_{i \in S_{A}} (y_{i} - \widehat{y}_{i}) + \widehat{y}_{U}\right) / N$$
(2.2)

where \hat{y}_i is the prediction of y_i for $i \in U$, and $\hat{y}_U = \sum_{i \in U} \hat{y}_i$. The last expression in Eq. 2.2 is also known as a "generalized difference estimator" (Wu and Sitter, 2001).

A fully Bayesian approach specifies a model to the joint distribution of (y_i, δ_i^A) across the units of U, which can be formulated by

$$p(y_i, \delta_i^A | x_i, d_i; \theta, \beta) = p(y_i | x_i, d_i, \delta_i^A; \theta) p(\delta_i^A | x_i; \beta), \quad i \in U$$

$$(2.3)$$

For brevity, to show that a variable, say x, is indexed across units of U, S_A , S_R , or S_C , we denote them by x_U , x_A , x_R , or x_C , respectively. Then, the likelihood function for (θ, β) based on the observed data is given by

$$L(\beta, \theta | y_A, \delta_U^A, x_U, d_U) \propto p(y_A, \delta_U^A | x_U, d_U, \theta, \beta) = \int p(y_U, \delta_U^A | x_U, d_U, \theta, \beta) dy_U$$
(2.4)

Under a Bayesian approach, model parameters are assigned prior distributions $p(\theta, \beta | x_U, d_U)$, and analytical inference is drawn based on the posterior distribution as below:

$$p(\beta, \theta | y_A, \delta_U^A, x_U, d_U) \propto p(\theta, \beta | x_U, d_U) L(\beta, \theta | y_A, \delta_U^A, x_U, d_U)$$
(2.5)

Note that in a Bayesian setting, to preserve the ignorable assumption, **C2**, in S_A , it is essential to specify independent priors, i.e. $p(\theta, \beta | x_U, d_U) = p(\theta | x_U, d_U) p(\beta | x_U, d_U)$ (Little and Zheng, 2007).

Descriptive inference about \overline{y}_U requires deriving the posterior predictive distribution conditional on the observed data, which is given by

$$p(\overline{y}_U|y_A, \delta_U^A, x_U, d_U) = \int \int p(\overline{y}_U|y_A, \delta_U^A, x_U, d_U, \theta, \beta) p(\theta, \beta|y_A, \delta_U^A, x_U, d_U) d\theta d\beta$$
(2.6)

We will further expand this formula in the following subsections and show how one can jointly estimate (y_U, π_U^A) in deriving the posterior predictive distribution of \overline{y}_U .

Obviously, estimating \hat{y}_U in Eq. 2.2 requires (X, D) to be observed for the entire population units, while it is not in practice, and most often, the measurement of auxiliary information is confined to the pooled sample, S_C . One way to tackle this issue is to generate a finite set of synthetic populations, say M, by undoing the sampling mechanism in S_R , which can be performed non-parametrically through the idea of finite population Bayesian bootstrapping (FPBB) (Dong et al., 2014; Little and Zheng, 2007). Given a synthetic population, attempts are then made to imputed the outcome variable for the entire non-sampled units.

However, considering the limited resources of high-performance computing, it is computationally expensive, if not infeasible, to fit Bayesian joint models repeatedly on large synthetic populations and simulate the posterior predictive distribution for all population units based on a custom posterior sampler (Mercer, 2018; Savitsky et al., 2016). In addition, the two-step algorithm proposed by Zangeneh and Little (2015) may not be fully implementable on the existing Bayesian platforms such as Stan (Carpenter et al., 2017), and therefore, the authors propose to combine the estimates across synthetic populations through Rubin's combining rules (Rubin, 1976). This may not be ideal especially when the posterior predictive distribution of the target population quantity tends to be heavily skewed and consequently, a symmetric confidence interval may fail to properly approximate the direct credible intervals of the posterior predictive distribution.

2.3. Proposed computationally tractable method

To reduce the computational burden and to be able to directly simulate the posterior predictive distribution of \overline{y}_U via a unified algorithm that is implementable in Stan, we limit the imputation of the outcome, y_i , to units of the combined survey, i.e. $i \in S_C$, and use the following estimator, as defined by Si et al. (2015), to multiply impute \hat{y}_U in Eq. 2.2 *M* times as below:

$$\widehat{y}_{U}^{(m)} = \sum_{j=1}^{J} N_{j}^{(m)} \widehat{\overline{y}}_{j}^{(m)}
= \sum_{j=1}^{J} \sum_{i=1}^{n_{j}} \frac{N_{j[i]}^{(m)}}{n_{j[i]}^{R}} \widehat{y}_{j[i]}^{(m)}$$
(2.7)

where $\left[N_{j[i]}^{(m)}, \hat{y}_{j[i]}^{(m)}\right]$ is the *m*-th draw of the joint posterior predictive distribution of the *j*-th post-stratum size and *i*-th outcome within *j*-th post-stratum. Therefore, the *m*-th posterior predictive draw of \overline{y}_U is given by

$$\widehat{y}_U^{(m)} = \left(\sum_{i=1}^{n_A} \left(y_i - \widehat{y}_i^{(m)}\right) + \widehat{y}_U^{(m)}\right) / N$$
(2.8)

To this end, we are interested in modeling the joint distribution of (y_i, δ_i^A, n_i^R) for $i \in S_C$ as below:

$$p(y_A, \delta_C^A, w_R, n_R | x_C, d_C, \theta, \beta, \xi) = p(n_R | w_R, \xi) p(y_A, \delta_C^A, w_R | x_C, d_C, \theta, \beta)$$
(2.9)

where $n_R = [n_1^R, n_2^R, ..., n_J^R]^T$ is the sizes of post-strata in S_R , and ξ is a *J*-dimensional vector of parameters associated with non-parametric modeling of $n_R|w_R$. While we thoroughly discuss each components of the rightmost expression of Eq. 2.9 later, one can derive the final estimate of \overline{y}_U by

$$\widehat{\bar{y}}_{U} = \frac{1}{M} \sum_{m=1}^{M} \widehat{\bar{y}}_{U}^{(m)}$$
(2.10)

and the associated $100(1 - \alpha)\%$ credible interval can be constructed by sorting $(\hat{\overline{y}}_U^{(1)}, \hat{\overline{y}}_U^{(2)}, ..., \hat{\overline{y}}_U^{(M)})$ ascendingly, and finding the $\alpha/2$ and $1 - \alpha/2$ percentiles of this ordered sequence that correspond to lower and upper limits of the credible interval, respectively.

2.3.1. Finite population Bayesian bootstrapping for modeling $p(n_R|w_R,\xi)$

We begin by modeling $p(n^R|w^R, \xi^R)$ non-parametrically via Bayesian bootstrapping (BB), with the aim to simulate the posterior predictive distribution of the N_j 's. The idea of BB operates quite similar to the regular bootstrap approach (Efron, 1981), except for the fact that BB simulates the posterior predictive distribution of a given population parameter instead of the sample distribution of the statistic estimating that parameter (Rubin, 1981). In a finite population Bayesian bootstrap (FPBB) setting, the goal is to derive the posterior predictive distribution of the post-strata sizes for the non-sampled population units, i.e. \overline{S}_R . Although FPBB imposes no parametric assumptions, it is assumed that all the existing post-strata in U are limited to those observed in the collected sample (exchangeability).

Under a simple random sample, Ghosh and Meeden (1983) propose to use a Polýa Urn Scheme, in which a Dirichlet-multinomial conjugate model is considered to expand the sample to the population. Cohen (1997) generalizes this approach to a weighted sample with independent draws, and the attributed Polýa posterior distribution for the non-sampled units of U given the observed sampling weights is formulated by Dong et al. (2014). Little and Zheng (2007) propose a modified FPBB method to generate synthetic populations based on the samples with a PPS design. Further extension based on a constrained BB is provided by Zangeneh and Little (2015) for situations where totals are known for auxiliary variables at the population level.

In the present article, we modify the FPBB method proposed by Little and Zheng (2007) by letting $v^R = \{v_1^R, v_2^R, ..., v_J^R\}$ represent the set of J distinct values of the sampling weights in S_R , and $\xi^R = \{\xi_1^R, \xi_2^R, ..., \xi_J^R\}$ denote the vector of conditional probabilities that $p(w^R = v_j^R | \delta^R = 1) = \xi_j^R$ for j = 1, 2, ..., J, where $\sum_{j=1}^J \xi_j^R = 1$. Now, suppose n_j^R and r_j^R are the frequencies of w^R taking the value v_j^R in S_R and \overline{S}_R , respectively, for j = 1, 2, ..., J. It is clear that $\sum_{j=1}^K n_j^R = n_R$, and $\sum_{j=1}^K r_j^R = N - n_R$. Considering a Dirichlet prior, i.e. $\xi^R \sim Dirichlet(\alpha_{J\times 1}), \ \alpha \in \mathbb{R}^{J>0}$, with a multinomial likelihood function of $p(n_1^R, n_2^R, ..., n_J^R | \xi) \propto \prod_{j=1}^J (\xi_j^R)^{n_j^R}$, the posterior distribution of ξ^R is given by $(\xi^R | n_1^R, n_2^R, ..., n_J^R) \sim Dirichlet(n_1^R + \alpha_1 - 1, n_2^R + \alpha_2 - 1, ..., n_J^R + \alpha_J - 1)$. Using Bayes' rule, Little and Zheng (2007) show that

$$\begin{split} \xi_{j}^{\overline{R}} &= p(w_{i}^{R} = v_{j}^{R} | \delta_{i}^{R} = 0) \\ &= p(\delta_{i}^{R} = 0 | w_{i}^{R} = v_{j}^{R}) \frac{p(w_{i}^{R} = v_{j}^{R})}{p(\delta_{i}^{R} = 0)} \\ &= p(\delta_{i}^{R} = 0 | w_{i}^{R} = v_{j}^{R}) \frac{p(w_{i}^{R} = v_{j}^{R} | \delta_{i}^{R} = 0) p(\delta_{i}^{R} = 0) + p(w_{i}^{R} = v_{j}^{R} | \delta_{i}^{R} = 1) p(\delta_{i}^{R} = 1)}{p(\delta_{i}^{R} = 0)} \\ &= p(\delta_{i}^{R} = 0 | w_{i}^{R} = v_{j}^{R}) \left\{ \xi^{\overline{R}} + \xi^{R} \frac{p(\delta_{i}^{R} = 1)}{p(\delta_{i}^{R} = 0)} \right\} \end{split}$$
(2.11)

Since $p(\delta_i^R = 0 | w_i^R = v_j^R) = 1 - \pi_j^R$, and $p(\delta_i^R = 1) / p(\delta_i^R = 0)$ can be treated as a normalizing constant,

$$\xi_j^{\overline{R}} \propto \xi_j^R \frac{1 - \pi_j^R}{\pi_j^R} \tag{2.12}$$

After normalizing $\xi^{\overline{R}}$ such that $\sum_{j=1}^{J} \xi_{j}^{\overline{R}} = 1$, the posterior predictive distribution of r^{R} is given by

$$p(r_1^R, r_2^R, ..., r_J^R | n_1^R, n_2^R, ..., n_J^R, \xi^R) = \binom{N - n_R}{r_1, r_2, ..., r_J} \prod_{j=1}^J \left[c\xi_j^R (1 - \pi_j^R) / \pi_j^R \right]^{r_j^R}$$
(2.13)

where c is the normalizing constant. The m-th posterior predictive draw of the size of post-stratum j in the population is $N_j^{(m)} = n_j^R + r_j^{R(m)}$, (m = 1, 2, ..., M).

2.3.2. Modeling the joint distribution of (y_i, δ_i^A) given the combined sample

As discussed earlier, the goal of PM in this study is to model $p(y_i|x_i;\theta)$ in order to obtain the posterior predictive distribution of y_i for $i \in S_R$, i.e. $p(y_R|y_A, x_C) \propto \int p(y_R|y_A, x_C; \theta) p(\theta|y_A, x_C) d\theta$. Although θ is a parameter defined in U, the ignorable assumption guarantees a consistent estimate of θ by fitting $p(y|x;\theta)$ on S_A , because

$$p(y_A|x_A;\theta) = p(y_U|\delta_U^A = 1, x_U, d_U;\theta) = \frac{p(\delta_U^A = 1|y_U, x_U;\theta)}{p(\delta_U^A = 1|x_U;\theta)} p(y_U|x_U, d_U;\theta) = p(y_U|x_U, d_U;\theta)$$
(2.14)

If π_i^A was known for $i \in S_C$, one could augment the PM by incorporating π_i^A as a predictor into the PM, e.g. $p(y_i|x_i, f(\pi_i^A); \theta)$. A robust estimator is achieved by choosing a flexible f(.), as detailed later.

While a non-probability sample is characterized by its unknown selection mechanism, given the conditions **C1-C4**, π_i^A can be estimated by modeling $p(\delta_U^A|x_U;\beta)$. Assuming that S_A is selected by a Poisson sampling, one can formulate the likelihood of β given δ_U^A as:

$$L(\beta|\delta_U^A, x_U) = \prod_{i=1}^N p(\delta_i^A = 1|x_i, \beta)^{\delta_i^A} \left[1 - p(\delta_i^A = 1|x_i, \beta)\right]^{1-\delta_i^A}$$
(2.15)

Under a logistic regression model,

$$\pi_i^A = p(\delta_i^A = 1 | x_i; \beta) = \frac{exp\{x_i^T\beta\}}{1 + exp\{x_i^T\beta\}}$$
(2.16)

By assigning appropriate prior distributions to β , one can simulate the posterior distribution of π_i^A for $i \in U$ through the Hamiltonian Monte Carlo (HMC) algorithm.

One major issue with Eq. 2.15 is that the observed (δ_i^A, x_i) is restricted to S_C . Although there exist several approaches restricting the estimation of β to S_C (Chen et al., 2019; Elliott and Valliant, 2017; Valliant et al., 2018; Wang et al., 2020), the majority rely on a pseudo-maximum likelihood estimation (PMLE) idea to account for unequal w_i^R 's, which necessitates solving a set of estimating equations. A corresponding method in a Bayesian setting is called pseudo-Bayesian. While such a method guarantees consistency in point estimates, the uncertainty tends to be underestimated in the posterior distribution of parameters (Gunawan et al., 2020; Savitsky et al., 2016; Williams and Savitsky, 2021). To avoid this problem, we employ a two-step pseudo-weighting approach proposed by Elliott and Valliant (2017). Assuming that $p(\delta_i^A + \delta_i^R = 2) \approx 0$, i.e. S_A and S_R have no overlap, one can show that

$$p(\delta_U^A = 1|x_U; \beta) = p(\delta_U^R = 1|x_U; \gamma) \frac{p(\delta_C^A = 1|x_C; \phi)}{1 - p(\delta_C^A = 1|x_C; \phi)}$$
(2.17)

where $\beta = (\gamma, \phi)^T$ is the associated model parameters. Rafei et al. (2020) call this approach propensity-adjusted probability prediction (PAPP) and prove the asymptotic properties of a pseudo-weighted estimate based on this method including consistency and variance estimation. As can be seen, this approach reduces the modeling of $p(\delta_U^A = 1|x_U)$ to the modeling of $p(\delta_C^A = 1|x_C)$ with an additional step, which is modeling $p(\delta_U^R|x_U)$. Treating π_i^R as a random variable for $i \in S_A$ conditional on x_i , one can estimate this probability by regressing the π_i^R 's on the x_i 's in U (Pfeffermann and Sverchkov, 2009), because

$$p(\delta_{U}^{R} = 1 | x_{U}; \gamma) = \int_{0}^{1} p(\delta_{U}^{R} = 1 | \pi_{U}^{R}, x_{U}; \gamma) p(\pi_{U}^{R} | x_{U}; \gamma) d\pi_{U}^{R}$$

$$= \int_{0}^{1} \pi_{U}^{R} p(\pi_{U}^{R} | x_{U}; \gamma) d\pi_{U}^{R}$$

$$= E(\pi_{U}^{R} | x_{U}; \gamma)$$
(2.18)

Pfeffermann and Sverchkov (1999) demonstrate that $E(\pi_U^R|x_U) = E^{-1}(w_R|x_R)$ where w_R are the sampling weights in S_R . Since π_i^R is only observed in S_R , then, the sample estimator of π_i^R is given by

$$p(\delta_C^A = 1 | x_C; \gamma, \phi) = E^{-1}(w_R | x_C; \gamma) \frac{p(\delta_C^A = 1 | x_C; \phi)}{1 - p(\delta_C^A = 1 | x_C; \phi)}$$
(2.19)

 $E(w_R|x)$ is modeled using a GLM with a log link function, as the distribution of the w_i^R 's tends to be right-skewed in the actual survey data. In addition, we know that the sampling weights are usually a multiplicative factor of selection probabilities×non-response adjustment×post-stratification. Therefore, given the posterior distribution of $p(\gamma, \beta|x_C, w_R)$, one can obtain the posterior distribution of π_i^A for $i \in S_C$ by

$$p(\delta_C^A = 1 | x_C; \gamma, \phi) = exp\{x_C^T(\phi - \gamma)\}$$
(2.20)

Note that modeling w_i^R is not required if w_i^R is known for $i \in S_A$ (Rafei et al., 2021). The joint distribution of (y_A, δ_C^A, w_R) can be written as:

$$p(y_A, \delta_C^A, w_R | x_C) = \int p(y_R, y_A | f(\pi^A [x_C, \delta_C^A, w_R; \gamma, \phi]), x_C; \theta) p(\delta_C^A | x_C; \phi) p(w_R | x_R; \gamma) dy_R$$
(2.21)

where $\pi^A[x_C, \delta^A_C, w_R; \gamma, \phi] = exp\{x^T_C(\phi - \gamma)\}$ according to Eq. 2.20. The corresponding posterior predictive distribution of y_R is given by

$$p(y_R|y_A, \delta_C^A, w_R, x_C, \delta_C^A, \pi_R^R) = \int \int \int \int p(y_R|y_A, f(\pi^A[x_C, \delta_C^A, \pi_R^R; \gamma, \phi]), x_C; \theta) \times p(\phi|\delta_C^A, x_C) p(\gamma|w_R, x_R) d\theta d\phi d\gamma$$
(2.22)

Although Zigler (2016) argues that such a factorization of the joint distribution of (y_i, δ_i^A, w^R) does not correspond to a valid use of the Bayes' theorem, for certain reasons, it has been advocated by several studies. First, Little (2004) highlights the fact that Bayesian joint modeling can result in better repeated sampling properties. It has been well-understood that the performance of the alternative two-step Bayesian methods with respect to frequentist properties depends on the choice of priors (Kaplan and Chen, 2012). Furthermore, having both π_i^A and x_i as predictors in the PM cuts the notorious feedback between the QR and PM models, which leads to incorrect estimation of the PS posterior distribution (Zigler et al., 2013).

However, what matters most in this study is the double robustness property that the likelihood factorization in Eq. 2.21 offers. For instance, by choosing a parametric form $f(\pi_i^A) = \theta^*/\pi_i^A$, where θ^* is an unknown scalar parameter, this factorization leads to a linear-in-weight Prediction (LWP) model. Scharfstein et al. (1999) and Bang and Robins (2005) identified the correspondence between LWP and AIPW estimators. In the causal inference context, this has been termed a clever covariate by Rose and van der Laan (2008) as it characterizes the correct relationship between the propensity scores and the outcome model. In the context of item-missing data imputation, Little and An (2004) suggest that the use of a more flexible non-parametric function can improve the efficiency of the adjusted estimator, especially when there are extreme values in the estimated PS. The authors propose to use a penalized spline model, which is piecewise continuous polynomials of the estimated PS, paired with a mathematical penalization to find the best fit of PM to the data (Fahrmeir et al., 2011; Ruppert et al., 2003). Alternatively, McCandless et al. (2009) suggest categorizing propensity scores into quantiles and using them as dummy variables to augment the PM.

In the current study, we extend the PSPP idea to a non-probability sample setting while using Gaussian process (GP) regression instead of a penalized spline model. As a flexible non-parametric Bayesian approach, GP can automatically capture non-linear associations as well as multi-way interactions (Neal, 1997; Rusmassen and Williams, 2005). Having $\pi_i^A = p(\delta_i^A = 1|x_i, w^R; \gamma, \phi)$ estimated for $i \in S_C$, for a continuous outcome variable, we fit a semiparametric model on S_A as below:

$$y_i|x_i, d_i, \widehat{\pi}_i, \theta = \theta_0 + \sum_{j=1}^p \theta_j x_{ij} + \sum_{j=p+1}^{p+q} \theta_j d_{ij} + f\left(\widehat{\pi}_i^A\right) + \epsilon_i$$
(2.23)

where θ denotes a (p+q+1)-dimensional vector of the PM parameters, and $\epsilon_i \sim N(0, \sigma^2)$ with σ^2 being unknown. Eq. 2.23 involves two parts: a linear regression parameterized by θ and a GP denoted by f(.).

A GP $\{f(u) : u \in \mathbb{R}^N\}$ is a set of random variables, any finite number of which jointly follow a multivariate Gaussian distribution. In a full-ranked GP, f(.) is a priori defined by its mean and covariance functions as below:

$$f(u) \sim GP\left(\mu(u), K\left(u, u'\right)\right) \tag{2.24}$$

where $\mu(u)$ is the mean vector and K(u, u') is the covariance matrix. The latter encompasses all our prior beliefs about the functional association between x and y, including continuity, smoothness, periodicity and scale properties (Riutort-Mayol et al., 2020). For notational simplicity, we set $\mu(u) = 0$, though it is not necessary. It is worth noting that the LWP model can be viewed as a specific type of GP with a dot product covariance matrix as $\alpha^2 [1 + ((\pi_i^A)^T \pi_j^A)^{-1}]$ if the regression coefficient is specified a prior of $N(0, \alpha^2)$ (Rusmassen and Williams, 2005). While literature suggests a variety of covariance functions for GP, the most common type is the squared exponential (SE) covariance matrix whose elements take the following form:

$$k(u, u') = \alpha^2 exp \left\{ -\frac{||u - u'||^2}{2\rho} \right\}$$
(2.25)

where ρ is called a *length-scale* parameter, and α is known as the *marginal standard error*. One can show that the SE covariance structure represents a kernel with an infinite number of basis functions (Rusmassen and Williams, 2005).

From a weight-space viewpoint, Huang et al. (2019) show that with a stationary isotropic kernel, where $K(\pi_i^A, \pi_j^A) = f(||\pi_i^A - \pi_j^A||)$, GP acts as a non-parametric matching technique. Wang et al. (2020) prove the consistency of a kernel-weighted estimator under certain regularity conditions. Refer to Appendix 8.1 to see the connection between GP and kernel weighting. In our non-probability sample setting, one can view it as matching units of S_A to units of S_R based on the estimated propensity scores, π_i^A 's (Rivers, 2007). Further theoretical properties of kernel optimal matching, such as consistency, can be found in Kallus et al. (2018). Although the SE covariance has desirable properties, empirical results show that it is not a strong fit for the real-world data as it is infinitely differentiable (Rusmassen and Williams, 2005). Therefore, we propose to use a Matérn kernel added to an inhomogeneous standardized polynomial kernel of order p as below:

$$K(u_i, u_j) = \alpha^2 \frac{2^{1-\nu}}{\Gamma(\nu)} \left(\sqrt{2\nu} \frac{||u_i - u_j||}{\rho}\right)^{\nu} K_{\nu} \left(\sqrt{2\nu} \frac{||u_i - u_j||}{\rho}\right) + \left(\frac{\tau^2 + u_i^T u_j}{\sqrt{\tau^2 + u_i^T u_i}}\sqrt{\tau^2 + u_j^T u_j}\right)^p$$
(2.26)

where $\Gamma(.)$ denotes the gamma function, and $K_{\nu}(.)$ is a modified Bessel function of the second kind. This combination of two kernels ensures capturing both local variations and long-range discrepancies in the estimated propensity scores (Vegetabile, 2018). Note that for $\nu \to \infty$, Matérn covariance will converge to the SE covariance, and the sum of two valid kernels is still a valid kernel.

In this chapter, we set $\nu = 3/2$ and p = 1 throughout the simulation and empirical studies, which yields the following covariance function:

$$K(u_i, u_j) = \alpha^2 \left(1 + \frac{\sqrt{3}||u_i - u_j||}{\rho} \right) exp\left(-\frac{\sqrt{3}||u_i - u_j||}{\rho} \right) + \frac{\tau^2 + u_i u_j}{\sqrt{\tau^2 + u_i^2}\sqrt{\tau^2 + u_j^2}}$$
(2.27)

In addition, we propose to use a log transformation of the $\hat{\pi}_i^A$'s as GP input, i.e. $u_i = log(\hat{\pi}_i^A)$. This is because the input of GP will become a linear combination, i.e. $x_C^T(\phi - \gamma)$, and given normal priors assigned to β , this linear combination is expected to follow a Gaussian distribution (Si et al., 2015).

Fully Bayesian inference using GP comes with computational issues even for a moderate n_A as one has to invert the covariance matrix at each posterior sampling step that needs $O(n_A^3)$ computations. The problem becomes even more severe when the joint posterior distribution of (π_i^A, y_i) has to be simulated, we propose to use a low-ranked sparse GP based on the Laplace eigenvectors approximation (Riutort-Mayol et al., 2020; Solin and Särkkä, 2020). Such a method reduces the computational complexity up to $O(n_A l^2)$ where $l << n_A$ is the reduced rank of the covariance matrix.

Under a standard Bayesian framework, a set of independent prior distributions are assigned to the model parameters, and conditional on the observed data through a joint likelihood function, the associated posterior distributions are obtained. To this end, we use the "black box" solver Stan (Carpenter et al., 2017), which employs an HMC technique to simulate the posterior predictive distribution of the parameters. In the following, we show the structure of our proposed method in Stan.

STEP 1: Priors

$$\begin{split} \theta, \gamma, \phi &\sim t\text{-}student(3,0,1) \\ \lambda, \alpha, \sigma &\sim t\text{-}student^+(3,0,1) \\ \rho &\sim GIG(0,1,2) \\ \xi^R &\sim Dirichlet(1,1,...,1) \end{split}$$

STEP 2: Joint likelihood

 \mathbf{R}_{\perp}

$$w^{r}[x_{R}, \gamma, \lambda \sim N \ (exp\{x_{R}\gamma\}, \lambda^{-}) \\ \delta^{A}_{C}[x_{C}, \phi \sim Bernoulli \ (logit^{-1}\{x_{C}^{T}\phi\}) \\ y_{A}[x_{A}, d_{A}, \theta, \sigma \sim Normal \ (\theta_{0} + x_{A}^{T}\theta_{1} + d_{A}^{T}\theta_{2} + f \ (x_{A}^{T}(\phi - \gamma), \alpha, \rho, \tau), \sigma^{2}) \\ n^{R}[\xi^{R} \sim Multinomial(n_{R}, \xi)$$

STEP 3: Posteriors

$$\begin{aligned} \widehat{y}_R | y_A, x_R, d_R, \theta, \sigma &\sim Normal \left(\theta_0 + x_R^T \theta_1 + d_R^T \theta_2 + f \left(x_R^T (\phi - \gamma), \alpha, \rho, \tau \right), \sigma^2 \right) \\ \widehat{N} | \pi^R, \xi^R &\sim Multinomial \left(N - n_R, c\xi^R (1 - \pi^R) / \pi^R \right) \\ \widehat{\overline{y}}_U &= \left\{ \sum_{j=1}^J \frac{\widehat{N}_j}{n_j^R} \sum_{i=1}^{n_j^R} \widehat{y}_{j[i]} + \sum_{i=1}^{n_A} \{ y_i - \widehat{y}_i \} \right\} / N \end{aligned}$$

where t-student⁺ denotes a half t-student and GIG stands for the Generalized Inverse Gaussian distribution, which is recommended in Stan User's Guide (Stan Development Team, 2019) for the length-scale parameter of a partially linear GP regression. Also, f(.) denotes a low-ranked GP approximation with l = 10 and a boundary condition factor of c = 1.25, where the covariance function is given by Eq. 2.3.2. Throughout the analysis, we simulate the posterior predictive distribution of \hat{y}_U in Stan using M = 500 HMC draws after discarding the first 500 draws as the burn-in period.

3. Simulation study

Two simulations are presented in this section, in which we compare the performance of our proposed GPPP method with those of LWP, AIPW, and PAPP with respect to the bias magnitude, efficiency, and accuracy of the variance estimator. All of the competing methods are DR, except for the PAPP method, which is an inverse PS weighted estimate of the observed y_i for $i \in S_A$ with PS estimated from Eq. 2.17. The GPPP and LWP methods are fully implemented under a Bayesian setting, whereas AIPW and PAPP estimates are obtained under a frequentist method (Rafei et al., 2021). Therefore, for the earlier class of methods, we are able to compute 95% credible intervals (95% CIs) while for the latter, a bootstrap method with B = 100 replications is employed to estimate the variance and 95% confidence intervals.

Various scenarios are considered with different assumptions about the functional form of the relationship among variables. For both studies, S_A and S_R are given a random selection mechanism with unequal inclusion probabilities. Note that units of both samples are selected independently with no clustering or stratification. Once S_A and S_R are drawn from U, we assume that π_i^A for $i \in S_C$ and y_j for $j \in S_R$ are unobserved, and the aim is to adjust for the selection bias in S_A based on the combined sample, S_C . The simulation is then iterated K = 216 times (which is a multiple of 36 as conducted parallel computing using 36 cores), where the bias-adjusted point estimates, SE and associated 95% credible/confidence interval (CI) for \overline{y}_U are estimated in each iteration.

To evaluate the repeated sampling properties of the competing method, relative bias (rBias), relative root mean square error (rMSE), the nominal coverage rate of 95% CIs (crCI), relative length of 95% CIs (rlCI) and SE ratio (rSE) are calculated as below:

$$rbias\left(\widehat{\overline{y}}_{U}\right) = 100 \times \frac{1}{K} \sum_{k=1}^{K} \left(\widehat{\overline{y}}_{U}^{(k)} - \overline{y}_{U}\right) / \overline{y}_{U}$$
(3.1)

$$rMSE\left(\widehat{\overline{y}}_{U}\right) = 100 \times \sqrt{\frac{1}{K} \sum_{k=1}^{K} \left(\widehat{\overline{y}}_{U}^{(k)} - \overline{\overline{y}}_{U}\right)^{2} / \overline{\overline{y}}_{U}}$$
(3.2)

$$crCI\left(\widehat{\overline{y}}_{U}\right) = 100 \times \frac{1}{K} \sum_{k=1}^{K} I\left(\left|\widehat{\overline{y}}_{U}^{(k)} - \overline{y}_{U}\right| < z_{0.975} \sqrt{var\left(\widehat{\overline{y}}_{U}^{(k)}\right)}\right)$$
(3.3)

$$rlCI\left(\widehat{\overline{y}}_{U}\right) = 100 \times \frac{2}{K} \sum_{k=1}^{K} z_{0.975} \sqrt{var\left(\widehat{\overline{y}}_{U}^{(k)}\right)}$$
(3.4)

$$rSE\left(\widehat{\overline{y}}_{U}\right) = \frac{1}{K} \sum_{k=1}^{K} \sqrt{var(\widehat{\overline{y}}_{U}^{(k)})} / \sqrt{\frac{1}{K-1} \sum_{k=1}^{K} \left(\widehat{\overline{y}}_{U}^{(k)} - \overline{\overline{y}}_{U}\right)^{2}}$$
(3.5)

where $\hat{\overline{y}}_U^{(k)}$ denotes the adjusted sample mean from iteration k, $\overline{\overline{y}}_U = \sum_{k=1}^K \hat{\overline{y}}_U^{(k)} / K$, \overline{y}_U is the finite population true mean, and var(.) represents the variance estimate of the adjusted mean based on the sample. Finally, to test the DR property of the proposed methods, we investigate different scenarios regarding whether models for QR and PM are correctly specified or not.

3.1. Simulation I

3.1.1. Design

S:3.1.1 The design of our first study is based on the simulation implemented in Chen et al. (2019). Consider a finite population of size $N = 10^5$ with $z = \{z_1, z_2, z_3, z_4\}$ being a set of auxiliary variables generated as follows:

$$z_1 \sim Ber(p=0.5)$$
 $z_2 \sim U(0,2)$ $z_3 \sim Eup(\mu=1)$ $z_4 \sim \chi^2_{(4)}$ (3.6)

and $x = \{x_1, x_2, x_3, x_4\}$ is subsequently defined as a linear function of z as below:

$$x_1 = z_1 \qquad x_2 = z_2 + 0.3z_1 \qquad x_3 = z_3 + 0.2(x_1 + x_2) \qquad x_4 = z_4 + 0.1(x_1 + x_2 + x_3) \qquad (3.7)$$

Given x, a continuous outcome variable y is constructed by

$$y_i = 2 + x_{1i} + x_{2i} + x_{3i} + x_{4i} + \sigma\epsilon_i \tag{3.8}$$

where $\epsilon_i \sim N(0, 1)$, and σ is defined such that the correlation between y_i and $\sum_{k=1}^4 x_{ki}$ equals $\rho = 0.8$. Further, associated with the design of S_A , a set of selection probabilities are assigned to the population units through the following logistic model:

$$\log\left(\frac{\pi_i^A}{1-\pi_i^A}\right) = \gamma_0 + 0.1x_{1i} + 0.2x_{2i} + 0.1x_{3i} + 0.2x_{4i}$$
(3.9)

where γ_0 is determined such that $\sum_{i=1}^N \pi_i^A = n_A$. For the selection probabilities in S_R , we assume that $\pi_i^R \propto \gamma_1 + z_{3i}$, where γ_1 is obtained such that $max\{\pi_i^R\}/min\{\pi_i^R\} = 50$. It is important to note that in this simulation study π_i^R is assumed to be known for $i \in S_A$ as z_3 is observed in S_A .

Using these measures of size, we repeatedly draw pairs of samples corresponding to S_A and S_R from U through a Poisson sampling design. The simulation is then repeated for different pairs of expected sample sizes, i.e. $(n_A, n_R) = (500, 500), (n_A, n_R) = (1, 000, 500)$ and also $(n_A, n_R) = (500, 1, 000)$. (Note that the actual sample size is a random variable under a Poisson sampling design.) Both Y and π^A are associated with a linear combination of X in this simulation study. Finally, in order to misspecify a model, we omit x_4 from the predictors of the working model.

3.1.2. Results

Table 1 summarizes the numerical results of the first simulation study across different sample size scenarios for $\rho = 0.8$. As illustrated, naive estimates of the population mean are biased in both S_R and S_A while weighting fully corrects for the bias in both samples. For the non-robust method, PAPP, estimates are unbiased as long as the QR model is correct. The DR methods produce unbiased estimates when either the QR model or PM holds, though there is evidence of residual bias for the LWP method when the QR model holds but the PM is misspecified. In terms of rMSE, all the methods perform similarly, except for the LWP method with correct and incorrect models specified the QR and PM, respectively, which shows higher degrees of rMSE compared to the alternative methods.

TABLE 1 Comparing the performance of the bias adjustment methods in the first simulation study for $\rho = 0.8$

$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$			$n_A = 50$	$00, n_R$	= 500		1	$n_A = 1, 0$	$000, n_{E}$	r = 500		1	$n_A = 500$), $n_{R} =$	= 1,000	
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	Measure	rBias	rMSE	crCI	rlCI	rSE	rBias	rMSE	crCI	rlCI	rSE	rBias	rMSE	crCI	rlCI	rSE
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Probabili	ty samp	le (S_R)													
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	UW	8.866	9.093	0.926	0.724	0.982	8.866	9.093	0.926	0.724	0.982	8.819	8.942	0.000	0.513	0.953
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	FW	0.150	2.322	93.981	0.844	0.998	0.150	2.322	93.981	0.844	0.998	0.030	1.692	94.907	0.598	0.969
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Non-prob	ability s	sample ((S_A)												
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	UW	30.675	30.794	0.000	0.940	0.950	29.958	30.006	0.000	0.657	1.063	30.675	30.794	0.000	0.940	0.950
	FW	-0.038	2.354	93.519	0.811	0.944	-0.044	1.618	93.056	0.570	0.965	-0.038	2.354	93.519	0.811	0.944
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Model spe	cification	: QR-Tr	ue, PM-'	True											
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	GPPP	0.054	2.473	96.759	0.935	1.035	0.014	2.171	95.370	0.862	1.087	0.003	2.039	95.370	0.750	1.007
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	LWP	0.034	2.586	97.222	1.502	1.591	-0.027	2.193	94.907	0.860	1.074	-0.071	2.107	94.907	0.763	0.992
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	AIPW	-0.042	2.528	93.981	0.867	0.940	0.001	2.166	93.519	0.772	0.976	-0.094	2.086	93.981	0.709	0.932
	PAPP	0.899	2.641	90.741	0.873	0.963	0.646	2.118	93.519	0.735	0.998	1.119	2.455	88.426	0.766	0.960
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Model spe	cification	: QR-Tr	ue, PM-l	False											
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	GPPP	0.025	2.465	96.759	0.934	1.038	0.003	2.177	93.981	0.856	1.078	-0.007	2.027	95.833	0.752	1.016
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	LWP	0.022	2.462	96.759	0.935	1.041	0.007	2.161	94.907	0.858	1.087	-0.028	2.036	95.833	0.749	1.007
	AIPW	-0.002	2.452	92.593	0.844	0.943	0.003	2.150	93.981	0.757	0.964	-0.068	2.019	93.519	0.697	0.947
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Model spe	cification	: QR-Fa	lse, PM-	True											
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	GPPP	0.945	2.855	95.370	0.983	0.999	0.798	2.453	96.759	0.898	1.061	0.990	2.461	91.204	0.792	0.963
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	LWP	3.989	5.740	76.852	1.413	0.937	4.233	5.511	75.463	1.278	0.992	3.959	5.245	71.759	1.136	0.904
	AIPW	0.215	2.532	93.056	0.855	0.929	0.092	2.068	93.981	0.744	0.987	0.173	2.138	93.519	0.752	0.967
GPPP 27.303 27.460 0.000 1.591 1.485 26.513 26.590 0.000 1.443 1.962 27.308 27.451 0.000 1.291 1.263 LWP 27.132 27.295 0.000 1.600 1.470 26.437 26.514 0.000 1.441 1.955 27.194 27.341 0.000 1.307 1.262 AIPW 27.162 27.322 0.000 0.986 0.914 26.453 26.531 0.000 0.725 0.978 27.114 27.252 0.000 0.947 0.934 PAPP 27.946 28.097 0.000 0.976 0.918 26.969 27.070 0.000 0.715 0.979 28.166 28.303 0.000 0.954 0.941	Model spe	cification	: QR–Fa	lse, PM-	False											
	GPPP	27.303	27.460	0.000	1.591	1.485	26.513	26.590	0.000	1.443	1.962	27.308	27.451	0.000	1.291	1.263
AIPW 27.162 27.322 0.000 0.986 0.914 26.453 26.531 0.000 0.725 0.978 27.110 27.252 0.000 0.947 0.934 PAPP 27.946 28.097 0.000 0.976 0.918 26.996 27.070 0.000 0.715 0.979 28.166 28.303 0.000 0.954 0.941	LWP	27.132	27.295	0.000	1.600	1.470	26.437	26.514	0.000	1.441	1.955	27.194	27.341	0.000	1.307	1.262
PAPP 27.946 28.097 0.000 0.976 0.918 26.996 27.070 0.000 0.715 0.979 28.166 28.303 0.000 0.954 0.941	AIPW	27.162	27.322	0.000	0.986	0.914	26.453	26.531	0.000	0.725	0.978	27.110	27.252	0.000	0.947	0.934
	PAPP	27.946	28.097	0.000	0.976	0.918	26.996	27.070	0.000	0.715	0.979	28.166	28.303	0.000	0.954	0.941

NOTE 1: GPPP: Gaussian Process of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting; PAPP: Propensity-adjusted Probability Prediction. NOTE 2: The PAPP and AIPW methods have been implemented through a bootstrap method. NOTE 3: The PAPP method is non-robust, while the rest of the methods, i.e. GPPP, LWP, and AIPW, are doubly robust.

AIPW and PAPP have slightly narrower CIs than the Bayesian methods, GPPP and LWP. The LWP performs poorly with respect to efficiency when the PM is incorrectly specified. Generally, the values of rSE suggest that variance estimation is unbiased across different model specification scenarios with a slight overestimation and underestimation in the Bayesian and bootstrap methods, respectively. Under the situations where the working model for QR is correct while that for PM is incorrect, LWP tends to underestimate the variance. The coverage rates of 95% CIs are also close to the nominal value when at least one of the QR and PM models is correctly specified. However, we observe that 95% CIs based on the frequentist methods tend to undercover the true population mean to some degrees, and the poorest result of crCI belongs to the LWP method when the PM is wrongly specified. These findings are generalizable to all other sample size combinations, and to the other extensions of the simulation for $\rho = 0.3, 0.5$, whose tables are displayed in Appendix 8.2.1.

3.2. Simulation II

3.2.1. Design

In the previous simulation study, the ignorable assumption was violated to misspecify the working model by dropping a key auxiliary variable. Now, we focus on a situation where models misspecified with respect to the functional form of their conditional means. To this end, we consider (non-)linear associations and two-way interactions in construction of the outcome variables. In addition, to build a more realistic situation, two separate sets of auxiliary variables are generated, D associated with the design of S_A , and X associated with the design of S_R . However, we allow the two variables to be correlated through a bivariate Gaussian distribution as below:

$$\begin{pmatrix} d \\ x \end{pmatrix} \sim MVN\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 4 & 2\rho \\ 2\rho & 1 \end{pmatrix}\right)$$
(3.10)

Note that ρ controls how strongly the sampling design of S_R is associated with that of S_A . Primarily, we set $\rho = 0.5$, but later we check other values ranging from 0 to 0.9 as well.

We then generate a continuous outcome variable (y_i^c) and a binary outcome variable (y_i^b) for $i \in U$ as below: $y_i^c = 3 + f_i(x_i) + d_i + 0.2x_i d_i + \sigma c_i$

$$y_i^{-} = 3 + f_k(x_i) + a_i + 0.2x_i a_i + \sigma \epsilon_i$$

$$p(y_i^{b} = 1 | x_i, d_i) = \frac{exp\{-1 + f_k(x_i) + d_i + 0.2x_i d_i\}}{1 + exp\{-1 + f_k(x_i) + d_i + 0.2x_i d_i\}}$$
(3.11)

where $\epsilon_i \sim N(0,1)$, and σ is determined such that the correlation between y_i^c and $f_k(x_i) + d_i + 0.2x_i d_i$ equals 0.8 for $i \in U$. The function $f_k(.)$ is assumed to take one of the following forms:

$$LIN: f_1(x) = x CUB: f_2(x) = (x/3)^3 (3.12)$$

$$EXP: f_3(x) = exp(x/2)/5 SIN: f_4(x) = 5sin(\pi x/3)$$

Figure 1 depicts the relationships between y^c and π^A , and between y^c and $w^A = 1/\pi^A$.



Fig 1: The proposed relationships between the outcome variable Y and $log(\pi^A)$ in U for (a) LIN, (b) CUB, (c) EXP and (d) SIN scenarios, and between the outcome Y and sampling weights w^A for (e) LIN, (f) CUB, (g) EXP and (h) SIN scenarios.

We then consider an informative sampling strategy with unequal probabilities of inclusion, where the selection mechanism of S_A and S_R depends on x and d, respectively. Thus, each $i \in U$ is assigned

two values within (0, 1) corresponding to the probabilities of selection in S_R and S_A through a *logistic* function as below:

$$\pi^{R}(d_{i}) = p(\delta_{i}^{R} = 1|d_{i}) = \frac{exp\{\gamma_{0} - 0.4d_{i}\}}{1 + exp\{\gamma_{0} - 0.4d_{i}\}}$$
(3.13)

$$\pi^{A}(x_{i}) = p(\delta^{A}_{i} = 1 | x_{i}) = \frac{exp\{\gamma_{1} + \gamma_{2}x_{i}\}}{1 + exp\{\gamma_{1} + \gamma_{2}x_{i}\}}$$
(3.14)

where δ_i^R and δ_i^A are the indicators of being selected in S_R and S_A , respectively, for $i \in U$, we initially set $\gamma_2 = 0.3$, which yields PS with a normal range. To assess how the adjustments behave in presence of influential weights, later we set $\gamma_2 = 0.6$, which yields relatively extreme weights.

Associated with S_R and S_A , independent samples of expected sizes $n_R = 1,000$ and $n_A = 500$ are selected randomly from U with a Poisson sampling design. We choose $n_A < n_R$ as is the case in the two applications of this study. The model intercepts, γ_0 and γ_1 in 3.13, are obtained such that $\sum_{i=1}^N \pi_i^R = n_R$ and $\sum_{i=1}^N \pi_i^A = n_A$, respectively. The rest of the simulation design is similar to that defined in Simulation I, except for the way we specify a working model. A QR model is misspecified by replacing x_i with x_i^2 , and a PM model is misspecified by replacing $f_k(x_i)$ with x_i^2 and d_i with d_i^2 , and also by dropping the interaction term $x_i d_i$.

3.2.2. Results

Figure 2 compares the relative bias (rBias) magnitude and efficiency of the competing methods for the continuous outcome variable, y^c , across different scenarios of model specification while $\gamma_2 = 0.3$. Note that the error bars reflect the relative length of 95% CIs (rlCI). As illustrated, point estimates from both S_R and S_A are biased if the sampling true weights are ignored. At the first glance, one can infer that for all f_k , k = 1, 2, 3, 4, the magnitude of rBias is close to zero as long as either QR or PM model is valid. However, in situations where π^A is non-linearly associated with y^c , i.e. plots (b), (c), and (d), the AIPW and PAPP estimators are biased when the PM is misspecified, but the QR model is valid. In contrast, the LWP method yields slightly biased estimates in all plots when the QR model is misspecified, but the Scenarios with respect to model specification and functional form of the PM. We did not observe consistent results across the adjustment methods with respect to efficiency. However, the GPPP method consistently shows high efficiency compared to the other methods across all the studied scenarios.

We summarize the simulation results for the binary outcome, y^b , with $\gamma_2 = 0.3$ in Figure 3. Again, adjusted estimates are unbiased if the working model for either QR or PM holds. Exceptions are seen for the PAPP and AIPW methods with residual bias in the plots related to (c) EXP, and (d) SIN when the PM is incorrectly specified. Unlike the simulation results for the continuous variable, the LWP consistently produces unbiased estimates for the binary outcome when the working model for QR fails. However, the magnitude of bias seems to be much larger in the LWP method when both underlying models for QR and PM are misspecified. Again, as for the continuous outcome, the proposed GPPP method consistently gives unbiased and efficient estimates. The lowest efficiency is associated with the AIPW and PAPP methods in the EXP scenario when the PM is misspecified.

Figure 4 displays the results of crCI and rSE for the continuous outcomes where $\gamma_2 = 0.3$. According to the rSE values, all methods perform well in variance estimation except for the LWP method which consistently underestimates the variance. A similar problem appears in the PAPP and AIPW methods for the EXP scenario when the outcome model is invalid. Generally, the Bayesian methods, i.e. GPPP and LWP, tend to slightly overestimate the variance. The values of crCI seem to be close to the nominal level for all the methods across almost all the scenarios, as long as at least one of the underlying models holds. For the non-linear associations, i.e. (b) CUB, (c) EXP and (d) SIN, the 95% CIs associated with frequentist methods, i.e. AIPW and PAPP, tend to undercover the population mean when the outcome model is false. Figure 5 depicts similar results for the binary outcome when $\gamma_2 = 0.3$. Overall, the results look analogous to those obtained for the continuous outcome. However, the degree of overestimation of variance by the Bayesian methods seems to be larger in the binary outcome than the continuous outcome.

Extensions of the simulation for other sample size combinations, i.e. $(n_A, n_R) = (500, 500)$ and $(n_A, n_R) = (1,000,500)$ and also for $\gamma_2 = 0.6$, which creates extreme sampling weights in S_A , are included in Appendix 8.2. While we observe no major discrepancy in the simulation results for other sample size scenarios than $(n_A, n_R) = (500, 1, 000)$, having influential weights presented in S_A leads to a larger magnitude of bias and lower efficiency in the estimates of PAPP, AIPW when the PM is incorrectly specified, but the QR model is valid. However, the GPPP method seems to be least affected by the presence of extreme weights.



Fig 2: Comparing the performance of the adjusted estimators under different model-specification scenarios for the continuous outcome variable with $\gamma_2 = 0.3$ under (a) LIN, (b) CUB, (c) EXP, and (d) SIN scenarios. The error bars have been drawn based on the 2.5% and 97.5% percentiles of the empirical distribution of bias over the simulation iterations. UW: unweighted; FW: Fully weighted; PAPP: Propensity-adjusted Probability Prediction; GPPP: Gaussian Processes of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting

In Figures 4 and 5, we depict the measures associated with the accuracy of the variance methods for GPPP/AIPW estimators. One can immediately infer that for both employed methods, the variance estimator is approximately unbiased when at least one of the underlying models holds. However, in situations where both models are invalid, according to the rSE values, the AIPW estimator tends to underestimate/overestimate the variance to a significant extent, while the variance estimator under GPPP shows more robustness across the model specification scenarios as well as outcome variables. Last but not least, the proximity of the crCI values to 95% for the GPPP methods, especially when both underlying models are wrong, reflects the accuracy of both point and variance estimates under the GPPP method.

So far, the results we discussed were limited to a case where $\rho = 0.5$. As the final step, we replicate the simulation for different values of ρ ranging from 0 to 0.9 to show how stable the competing methods perform in terms of rbias and rMSE. Figure 6 depicts changes in the values of rBias and rMSE in the continuous outcome, y^c , for different adjustment methods and across different model specification scenarios as the value of ρ increases. Generally, it seems that the values of rBias and rMSE decline for all competing methods with an increase in ρ . In addition, for all values of ρ , it is evident that the GPPP method outperforms the PAPP, AIPW, LWP methods when the outcome model is wrong. This strength in GPPP is more evident when the association between the outcome and the PS is non-linear, i.e. in (b)



Fig 3: Comparing the performance of the adjusted estimators under different model-specification scenarios for the binary outcome variable with $\gamma_2 = 0.3$ under (a) LIN, (b) CUB, (c) EXP, and (d) SIN scenarios. The error bars have been drawn based on the 2.5% and 97.5% percentiles of the empirical distribution of bias over the simulation iterations. UW: unweighted; FW: Fully weighted; PAPP: Propensity-adjusted Probability Prediction; GPPP: Gaussian Processes of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting



Fig 4: Comparing the 95% CI coverage rates (crCI) of the DR adjusted means for the *continuous* outcome variable with $\gamma_2 = 0.3$ under (a) LIN, (b) CUB, (c) EXP, and (d) SIN scenarios, and SE ratios (rSE) under (e) LIN, (f) CUB, (g) EXP, and (h) SIN scenarios, across different DR methods under different model specification scenarios. UW: unweighted; FW: Fully weighted; PAPP: Propensity-adjusted Probability Prediction; GPPP: Gaussian Processes of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting

CUB, (c) EXP, and (d) SIN. In Figure 7, we display corresponding comparisons for the binary outcome. The results are similar to those based on the continuous outcome, with a difference in that the values of



Fig 5: Comparing the 95% CI coverage rates (crCI) of the DR adjusted means for the *binary* outcome variable with $\gamma_2 = 0.3$ under (a) LIN, (b) CUB, (c) EXP, and (d) SIN scenarios, and SE ratios (rSE) under (e) LIN, (f) CUB, (g) EXP, and (h) SIN scenarios, across different DR methods under different model specification scenarios. UW: unweighted; FW: Fully weighted; PAPP: Propensity-adjusted Probability Prediction; GPPP: Gaussian Processes of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting

rMSE increase with an increase in the value of ρ . Detailed numerical results of Simulation II is available in Appendix 8.2.



Fig 6: Comparing the magnitude of rBias of the DR adjusted means for the *continuous* outcome variable with $\gamma_2 = 0.3$ under (a) LIN, (b) CUB, (c) EXP, and (d) SIN, and rMSE under (e) LIN, (f) CUB, (g) EXP, and (h) SIN across different model specification scenarios and different values of ρ . UW: unweighted; FW: Fully weighted; PAPP: Propensity-adjusted Probability Prediction; GPPP: Gaussian Processes of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting

4. Applications

As the application part, we conduct an empirical study involving inference for a non-probability sample. Our goal is to estimate police-reportable crash rates per 100M miles driven using the sensor-based data



Fig 7: Comparing the magnitude of rBias of the DR adjusted means for the *binary* outcome variable with $\gamma_2 = 0.3$ under (a) LIN, (b) CUB, (c) EXP, and (d) SIN, and rMSE under (e) LIN, (f) CUB, (g) EXP, and (h) SIN across different model specification scenarios and different values of ρ . UW: unweighted; FW: Fully weighted; PAPP: Propensity-adjusted Probability Prediction; GPPP: Gaussian Processes of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting

from the second phase of the Strategic Highway Research Program (SHRP2). To this end, we consider the National Household Travel Survey (NHTS) 2017 as the reference survey to adjust for the potential selection bias in crash rates. The following two subsections briefly describe the design of these samples and discuss the findings in detail.

4.1. Strategic Highway Research Program 2

SHRP2 is the largest naturalistic driving study (NDS) conducted to date, with the primary aim to assess how people interact with their vehicle and traffic conditions while driving (SHRP2, 2013). About A = 3,140 drivers aged 16 – 95 years were recruited from six geographically dispersed sites across the United States, and over five million trips and 50 million driven miles have been recorded during their participation time. The average follow-up time per person was 440 days. A quasi-random approach was initially employed to select samples by random cold calling from a pool of 17,000 pre-registered volunteers. However, because of the low success rate along with budgetary constraints, the investigators later chose to pursue voluntary recruitment. Sites were assigned one of three pre-determined sample sizes according to their population density (Campbell, 2012). The youngest and eldest age groups were oversampled in the sense that crash risk is expected to be higher among those subgroups. Thus, one can conclude that the selection mechanism in SHRP2 is a combination of convenience and quota sampling methods. Further description of the study design and recruitment process can be found in Antin et al. (2015).

SHRP2 data are collected in multiple stages. Selected participants are initially asked to complete multiple assessment tests, including executive function and cognition, visual perception, visual-cognitive, physical and psychomotor capabilities, personality factors, sleep-related factors, general medical condition, driving knowledge, etc. In addition, demographic information such as age, gender, household income, education level, and marital status as well as vehicle characteristics such as vehicle type, model year, manufacturer, and annual mileage are gathered at the screening stage. A trip in SHRP2 is defined as the time interval during which the vehicle is operating. The in-vehicle sensors start recording kinematic information, the driver's behaviors, and traffic events continuously as soon as the vehicle is switched on. Encrypted data are stored in a removable hard drive, and participants are asked to provide access to the vehicle every four to six months, so that hard drives with accumulated data are removed and replaced. Then, trip-related information such as average speed, duration, distance, and GPS trajectory coordinates are obtained by aggregating the sensor records at the trip level (Antin et al., 2019; Campbell, 2012).

4.2. National Household Travel Survey 2017

In the present study, we use data from the eighth round of the NHTS conducted from March 2016 through May 2017 as the reference survey. The NHTS is a nationally representative survey, repeated cross-sectionally almost every seven years. It is aimed at characterizing personal travel behaviors among the civilian, non-institutionalized population of the United States. The 2017 NHTS was a mixed-mode survey, in which households were initially recruited by mailing through an address-based sampling (ABS) technique. Within the selected households, all eligible individuals aged ≥ 5 years were requested to report the trips they made on a randomly assigned weekday through a web-based travel log. Proxy interviews were requested for younger household members who were ≤ 15 years old.

The overall sample size was 129,696, of which roughly 20% was used for national representativity and the remaining 80% was regarded as add-ons for the state-level analysis. The recruitment response rate was 30.4%, of which 51.4% reported their trips via the travel logs (Santos et al., 2011). In NHTS, a travel day is defined from 4 : 00 AM of the assigned day to 3 : 59 AM of the following day on a typical weekday. A trip is defined as that made by one person using any mode of transportation. While trip distance was measured by online geocoding, the rest of the trip-related information was based on self-reporting. A total of 264, 234 eligible individuals aged ≥ 5 took part in the study, for which 923, 572 trips were recorded (McGuckin and Fucci, 2018).

4.3. Auxiliary variables and analysis plan

Our focus here was on inference at the participant level, so both SHRP2 and NHTS data were aggregated. Considering this, we calculated the total distance driven and total number of police-reported crashes for each participant of SHRP2. The total of these quantities by all SHRP2 participants were 28M miles and 210, respectively. To make the two datasets more comparable, we also filtered out all the subjects in NHTS who were not drivers or were younger than 16 years old or used public transportation or transportation modes other than cars, SUVs, vans, or light pickup trucks. In addition, we restricted the NHTS sample to those respondents who reside in the SHRP2-specific six states, so our inferences are only generalizable to those six states. The final sample sizes of the complete datasets were $n_A = 2,862$ and $n_R = 29,572$ in SHRP2 and NHTS, respectively.

To address the expressed objective of the present study, we set the outcome variable to be the frequency of police-reported crashes by SHRP2 participants throughout their follow-up time. In addition, we utilize the total miles driven by each SHRP2 participant as the model offset to obtain the rates by a driven mile. Particular attention was paid to identify as many relevant common auxiliary variables as possible in the combined sample that are expected to govern both selection mechanism and response surface in SHRP2. Two distinct sets of variables were considered: (i) demographic and socio-economic information of the drivers including sex, age groups, race, ethnicity, birth country, education level, household size, number of owned vehicles, and state of residence, and (ii) vehicle characteristics including vehicle age, vehicle manufacturer, vehicle type and fuel type.

We chose to use a Bayesian negative binomial (NB) regression for modeling the response surface because the outcome variable was count data and effects of overdispersion were present. The *log* of the total miles driven by SHRP2 participants is included as offset in the model such that crash rates can be predicted by unit of distance driven, conditional on x_i and $\hat{\pi}_i^i$, the outcome model is given by

$$y_i|x_i, t_i, \theta, \phi, \gamma, \alpha, \tau, \sigma \sim NB\left(exp\{\theta_0 + x_i^T\theta_1 + f\left(x_A^T(\phi - \gamma), \alpha, \rho, \tau\right) + log(t_i)\}, 1/\sigma\right)$$
(4.1)

where t_i is the total distance driven by respondent *i*, and $\sigma \sim Cauchy^+(0,3)$. Note that we also checked and found no evidence of zero-inflation in the distribution of the outcome by comparing the observed zeros with the expected number of zeros under the proposed NB model.

21

4.3.1. Results

According to Figure 8, one can visually infer that the largest discrepancies between the sample distribution of auxiliary variables in SHRP2 and that in the population stem from participants' age, race, and population size of the residential area as well as vehicles' age and vehicles' type. The youngest and oldest age groups are overrepresented as are Whites and non-Hispanics. In addition, we found that the proportion of urban dwellers is higher in SHRP2 than that in the NHTS. In terms of vehicle characteristics, SHRP2 participants tend to own passenger cars more than the population average, whereas individuals with other vehicle types were underrepresented in SHRP2.

Before any attempt for bias adjustment, we check the positivity assumption as well as the existence of influential pseudo-weights. To this end, we estimate the pseudo-selection probabilities for the units of the SHRP2 sample using the PAPP method as well as the PMLE method by Wang et al. (2020). Figure 9a compares the distribution of estimated PS in log scale between the SHRP2 and NHTS samples. As illustrated, there is a slight lack of common support in the distribution of PS, which may lead to extreme weights. The box-plot on the right side (Figure 9b) confirms the presence of outlying pseudo-weights based on the PAPP method. However, it seems no outliers exist in the pseudo-weights based on the PMLE method. Figure 10 compares the distribution of auxiliary variables between the two samples after (pseudo-)weighting. As illustrated, pseudo-weighting obviates most of the previously seen discrepancies in the distribution of common covariates.

Figure 11 displays the adjusted estimates of police-reported crash rates per 100M miles driven and associated 95% CIs using the LWP and GPPP methods by age groups. The plot also compares the adjusted estimates in SHRP2/NHTS data with the naive estimate using SHRP2-only data and that based on the GES/ADS data, which is here considered as the benchmark Tefft (2017). Note that the latter represents the entire population of American drivers while our adjusted estimates represent the SHRP2 target population. As illustrated, for most of the age groups, adjustments shift the unweighted crash rates to the true population value, and the associated 95% CIs overlap, except for the last age group, i.e 80+ years old. In particular, the unweighted crash rate for the age group 50-59 years seems to be severely biased while adjusted estimates are desirably close to the true population value. While we observe no significant differences in the performance of the GPPP and LWP methods, it is evident that GPPP offers more efficient estimates than the LWP method, as the length of 95% CIs is consistently lower in GPPP than LWP. Finally, one can infer from Figure 11 that the risk of traffic accidents is higher among young and elder people.

In Figure 12, we assess the adjusted rates of police-reportable crashes across levels of auxiliary variables. The major associations we observe are as follows: Whites, more educated drivers, and those in middle-income families are at lower risk of traffic accidents. In addition, there is a positive relationship between the crash risk and household size. There is also evidence of higher crash rates among Vans, European, and gas/diesel vehicles. Numerical values associated with this plot have been provided in Table 16 of Appendix 8.3.







Fig 9: Comparing the empirical density of (a) estimated propensity scores between SHRP2 and NHTS and (b) estimated pseudo-weights in SHRP2 across the applied quasi-randomization methods







Fig 11: Comparing the performance of adjustment methods for estimating crash rates per 100M miles and associated 95% CIs in SHRP2/NHTS with native estimates and those based on CES/ADS as benchmark across age groups. UW: unweighted; FW: Fully weighted; PAPP: Propensity-adjusted Probability Prediction; GPPP: Gaussian Processes of Propensity Prediction; LWP: Linear-in-weight Prediction.



Fig 12: Comparing the performance of adjustment methods for estimating crash rates per 100M miles and associated 95% CIs in SHRP2/NHTS with native estimates across levels of (a) sex, (b) race, (c) education, (d) household income, (e) household size, (f) vehicle make, (g) vehicle type, and (d) fuel type. UW: unweighted; GPPP: Gaussian Process of Propensity Prediction; LWP: Linear-in-weight Prediction.

26

5. Discussion

In the present article, we proposed alternative methods for inference based non-probability samples with the goal to fill some of the gaps in the current literature. To our knowledge, this was the first study proposing a fully Bayesian method in a non-probability sample setting that jointly estimates the PS and outcome, and allows one to directly simulate the posterior predictive distribution of the population quantity under a non-probability sample setting. Bayesian approaches provide a unified framework for deriving the variance of the point estimator by simulating the posterior predictive distribution of the population unknown parameters. A *well-calibrated* Bayesian method can appropriately capture all sources of uncertainty, and therefore, yields desirable frequentist repeated sampling properties (Dawid, 1982). Unlike the Bayesian two-step methods (Kaplan and Chen, 2012; Rafei et al., 2021), it is well-understood that joint modeling of the PS and the outcome, as was the case in this article, results in accurate variance estimation (Little, 2004).

The alternative design-based approaches, such as the AIPW estimator, are sensitive to the presence of influential pseudo-weights if the outcome model is invalid. In addition, the variance estimator proposed by Chen et al. (2019) relies on asymptotic theory, and there is no guarantee that simultaneously solving the estimating leads to a unique set of solutions. As another major limitation, such a method works only when the auxiliary variables are identical in the QR and PM models. According to the likelihood we factorized in Eq. 2.1, the dimension of the auxiliary variables may vary across the QR and PM methods in a non-probability sample setting ($\{X, D\}$ vs X), which makes it impossible to use Chen's AIPW method in practice. On the other hand, employing a fully model-based approach can be extremely expensive computationally, as one has to fit the propensity model on a synthesized population, and predict the outcome variable for its entire non-sampled units Little and Zheng (2007); Mercer (2018). However, the method we proposed requires fitting the model only on the combined sample, which makes it computationally more parsimonious, especially in a Big Data setting.

The results of our simulation studies reveal that the proposed GPPP method is doubly robust in terms of both point and variance estimates. Furthermore, estimates based on the GPPP method were more efficient than those based on the LWP method especially when outlying pseudo-weights are present and the PM is misspecified. We also showed that the use of GP with a stationary isotropic covariance structure provides a stronger rationale than Spline in the PSPP method as it is equivalent to a non-parametric matching technique based on the estimated PS (Huang et al., 2019). While Bayesian joint modeling demonstrates good frequentist properties, feedback occurs between the two models (Zigler et al., 2013). This can be controversial in the sense that PS estimates should not be informed by the outcome model (Rubin, 2007).

It is worth noting that although our proposed method limited computations to the combined samples, Bayesian joint modeling can still turn out computationally very expensive, even using the low-rank approximation techniques. To conduct the simulation studies of this research, we had to hire high-performance computing servers to be able to do parallel processing, but these resources are costly to hire and may not be available for every researcher. Finally, we want to pinpoint that our proposed estimator still contains a design-based term, and therefore, adjusted estimates can still be inefficient if there are extreme values in the sampling weights of S_R . However, this should not be of a big concern compared to the pseudo-weights of S_A as S_R is supposed to be a well-designed probability sample. As discussed earlier, a fully model-based estimator will require generating synthetic populations which cannot be fully implemented on the current Bayesian platforms.

6. Acknowledgement

The present study was part of the doctoral research of the first author of this article at the Michigan Program in Survey and Data Science. Therefore, we would like to thank the respected members of the dissertation committee, Professors Brady T. West, Roderick J. Little, and Philip S. Boonstra at the University of Michigan, who have continuously supported this research with their excellent comments and critical feedback. Our gratitude also goes to Professors Katharine Abraham, Stanley Presser, and Joseph Sedarski at the University of Maryland who have significantly contributed to the development of the main idea of this paper with their valuable comments and feedback over the doctoral seminar course. Last but not least, the authors would like to thank all the researchers and staff who have been involved in collecting the data of SHRP2 and NHTS.

7. Conflict of Interest

The authors declare that there was no conflict of interest in the current research.

References

- Administration, N. H. T. S. et al. (2014). National automotive sampling system (nass) general estimates system (ges) analytical user's manual 1988-2012 (dot publication no. dot hs 811 853).
- An, H. and R. J. Little (2008). Robust model-based inference for incomplete data via penalized spline propensity prediction. Communications in Statistics-Simulation and Computation 37(9), 1718–1731.
- Antin, J., K. Stulce, L. Eichelberger, and J. Hankey (2015). Naturalistic driving study: descriptive comparison of the study sample with national data. Technical report.
- Antin, J. F., S. Lee, M. A. Perez, T. A. Dingus, J. M. Hankey, and A. Brach (2019). Second strategic highway research program naturalistic driving study methods. *Safety Science* 119, 2–10.
- Bang, H. and J. M. Robins (2005). Doubly robust estimation in missing data and causal inference models. *Biometrics* 61(4), 962–973.
- Beresewicz, M., R. Lehtonen, F. Reis, L. Di Consiglio, and M. Karlberg (2018). An overview of methods for treating selectivity in big data sources.
- Brick, J. M. and R. Tourangeau (2017). Responsive survey designs for reducing nonresponse bias. *Journal* of Official Statistics 33(3), 735–752.
- Brick, J. M. and D. Williams (2013). Explaining rising nonresponse rates in cross-sectional surveys. The ANNALS of the American academy of political and social science 645(1), 36–59.
- Campbell, K. L. (2012). The shrp 2 naturalistic driving study: Addressing driver performance and behavior in traffic safety. *Tr News* (282).
- Carpenter, B., A. Gelman, M. D. Hoffman, D. Lee, B. Goodrich, M. Betancourt, M. A. Brubaker, J. Guo, P. Li, and A. Riddell (2017). Stan: a probabilistic programming language. *Grantee Submission* 76(1), 1–32.
- Chen, Q., M. R. Elliott, D. Haziza, Y. Yang, M. Ghosh, R. J. Little, J. Sedransk, M. Thompson, et al. (2017). Approaches to improving survey-weighted estimates. *Statistical Science* 32(2), 227–248.
- Chen, Q., M. R. Elliott, and R. J. Little (2012). Bayesian inference for finite population quantiles from unequal probability samples. *Survey methodology* 38(2), 203.
- Chen, Y., P. Li, and C. Wu (2019). Doubly robust inference with nonprobability survey samples. *Journal* of the American Statistical Association, 1–11.
- Cohen, M. P. (1997). The bayesian bootstrap and multiple imputation for unequal probability sample designs. In Proceedings of the Survey Research Methods Section, American Statistical Association, pp. 635–638.
- Dawid, A. P. (1982). The well-calibrated bayesian. Journal of the American Statistical Association 77(379), 605–610.
- Dong, Q., M. R. Elliott, and T. E. Raghunathan (2014). A nonparametric method to generate synthetic populations to adjust for complex sampling design features. Survey methodology 40(1), 29.
- Efron, B. (1981). Nonparametric estimates of standard error: the jackknife, the bootstrap and other methods. *Biometrika* 68(3), 589–599.
- Elliott, M. R. and R. Valliant (2017). Inference for nonprobability samples. *Statistical science* 32(2), 249–264.
- Fahrmeir, L., T. Kneib, et al. (2011). Bayesian smoothing and regression for longitudinal, spatial and event history data. *OUP Catalogue*.
- Forsberg, O. J. (2020). Polls and the us presidential election: real or fake? Significance 17(5), 6–7.

- Ghosh, M. and G. Meeden (1983). Estimation of the variance in finite population sampling. Sankhyā: The Indian Journal of Statistics, Series B, 362–375.
- Groves, R. M. (2011). Three eras of survey research. Public Opinion Quarterly 75(5), 861–871.
- Groves, R. M. and S. G. Heeringa (2006). Responsive design for household surveys: tools for actively controlling survey errors and costs. *Journal of the Royal Statistical Society: Series A (Statistics in Society)* 169(3), 439–457.
- Gunawan, D., A. Panagiotelis, W. Griffiths, and D. Chotikapanich (2020). Bayesian weighted inference from surveys. Australian & New Zealand Journal of Statistics 62(1), 71–94.
- Guo, F., J. M. Hankey, et al. (2009). Modeling 100-car safety events: A case-based approach for analyzing naturalistic driving data. Technical report, Virginia Tech. Virginia Tech Transportation Institute.
- Huang, B., C. Chen, and J. Liu (2019). Gpmatch: A bayesian doubly robust approach to causal inference with gaussian process covariance function as a matching tool. arXiv preprint arXiv:1901.10359.
- Johnson, T. P. and T. W. Smith (2017). Big data and survey research: Supplement or substitute? In Seeing Cities Through Big Data, pp. 113–125. Springer.
- Kallus, N., B. Pennicooke, and M. Santacatterina (2018). More robust estimation of sample average treatment effects using kernel optimal matching in an observational study of spine surgical interventions. arXiv preprint arXiv:1811.04274.
- Kaplan, D. and J. Chen (2012). A two-step bayesian approach for propensity score analysis: Simulations and case study. *Psychometrika* 77(3), 581–609.
- Kaufman, C. G., S. R. Sain, et al. (2010). Bayesian functional {ANOVA} modeling using gaussian process prior distributions. *Bayesian Analysis* 5(1), 123–149.
- Keeter, S., N. Hatley, C. Kennedy, and A. Lau (2017). What low response rates mean for telephone surveys. *Pew Research Center* 15, 1–39.
- Keiding, N. and T. A. Louis (2016). Perils and potentials of self-selected entry to epidemiological studies and surveys. Journal of the Royal Statistical Society: Series A (Statistics in Society) 179(2), 319–376.
- Kern, C., Y. Li, and L. Wang (2020). Boosted kernel weighting-using statistical learning to improve inference from nonprobability samples. *Journal of Survey Statistics and Methodology* 00, 1–26.
- Kim, J. K., S. Park, Y. Chen, and C. Wu (2018). Combining non-probability and probability survey samples through mass imputation. arXiv preprint arXiv:1812.10694.
- Kim, J. K. and J. N. Rao (2012). Combining data from two independent surveys: a model-assisted approach. *Biometrika* 99(1), 85–100.
- Kim, W., V. Anorve, and B. Tefft (2019). American driving survey, 2014–2017.
- Kimeldorf, G. S. and G. Wahba (1970). A correspondence between bayesian estimation on stochastic processes and smoothing by splines. The Annals of Mathematical Statistics 41(2), 495–502.
- Kreuter, F. and R. D. Peng (2014). 12 extracting information from big data: Issues of measurement, inference and linkage. *Privacy, Big Data, and the Public Good: Frameworks for Engagement*, 257.
- Lee, S. (2006). Propensity score adjustment as a weighting scheme for volunteer panel web surveys. Journal of official statistics 22(2), 329.
- Lee, S. and R. Valliant (2009). Estimation for volunteer panel web surveys using propensity score adjustment and calibration adjustment. Sociological Methods & Research 37(3), 319–343.
- Little, R. and H. An (2004). Robust likelihood-based analysis of multivariate data with missing values. Statistica Sinica, 949–968.
- Little, R. J. (2004). To model or not to model? competing modes of inference for finite population sampling. *Journal of the American Statistical Association* 99(466), 546–556.
- Little, R. J. and H. Zheng (2007). The bayesian approach to the analysis of finite population surveys. Bayesian Statistics 8(1), 1–20.
- Luiten, A., J. Hox, and E. de Leeuw (2020). Survey nonresponse trends and fieldwork effort in the 21st century: Results of an international study across countries and surveys. *Journal of Official Statistics* 36(3), 469–487.
- McCandless, L. C., P. Gustafson, and P. C. Austin (2009). Bayesian propensity score analysis for observational data. *Statistics in medicine* 28(1), 94–112.
- McGuckin, N. and A. Fucci (2018). Summary of travel trends: 2017 national household travel survey (report fhwa-pl-18-019). Washington, DC: Federal Highway Administration, US Department of

Transportation.

- Meng, X.-L. et al. (2018). Statistical paradises and paradoxes in big data (i): Law of large populations, big data paradox, and the 2016 us presidential election. The Annals of Applied Statistics 12(2), 685–726.
- Mercer, A. W. (2018). Selection Bias in Nonprobability Surveys: A Causal Inference Approach. Ph. D. thesis.
- Miller, P. V. (2017). Is there a future for surveys? Public Opinion Quarterly 81(S1), 205–212.
- Neal, R. M. (1997). Monte carlo implementation of gaussian process models for bayesian regression and classification. arXiv preprint physics/9701026.
- Neyman, J. (1934). On the two different aspects of the representative method: the method of stratified sampling and the method of purposive selection. *Journal of the Royal Statistical Society* 97(4), 558–625.
- Oakley, J. E. and A. O'Hagan (2004). Probabilistic sensitivity analysis of complex models: a bayesian approach. Journal of the Royal Statistical Society: Series B (Statistical Methodology) 66(3), 751–769.
- Pfeffermann, D. and M. Sverchkov (1999). Parametric and semi-parametric estimation of regression models fitted to survey data. Sankhyā: The Indian Journal of Statistics, Series B, 166–186.
- Pfeffermann, D. and M. Sverchkov (2009). Inference under informative sampling. In *Handbook of statistics*, Volume 29, pp. 455–487. Elsevier.
- Prieger, J. E. (2004). An empirical investigation of biased survey data. UC Davis Department of Economics Working Paper (04-4).
- Rafei, A. (2021). Robust and Efficient Bayesian Inference for Large-scale Non-probability Samples. Ph. D. thesis.
- Rafei, A., C. A. C. Flannagan, and M. R. Elliott (2020). Big data for finite population inference: Applying quasi-random approaches to naturalistic driving data using bayesian additive regression trees. *Journal* of Survey Statistics and Methodology In press.
- Rafei, A., C. A. C. Flannagan, B. T. West, and M. R. Elliott (2021). Robust bayesian inference for big data: Combining sensor-based records with traditional survey data.
- Riutort-Mayol, G., P.-C. Bürkner, M. R. Andersen, A. Solin, and A. Vehtari (2020). Practical hilbert space approximate bayesian gaussian processes for probabilistic programming. arXiv preprint arXiv:2004.11408.
- Rivers, D. (2007). Sampling for web surveys. In *Joint Statistical Meetings*.
- Robins, J. M., A. Rotnitzky, and L. P. Zhao (1994). Estimation of regression coefficients when some regressors are not always observed. *Journal of the American statistical Association* 89(427), 846–866.
- Rose, S. and M. J. van der Laan (2008). Simple optimal weighting of cases and controls in case-control studies. The International Journal of Biostatistics 4(1).
- Rubin, D. B. (1976). Inference and missing data. *Biometrika* 63(3), 581–592.
- Rubin, D. B. (1981). The bayesian bootstrap. The annals of statistics, 130–134.
- Rubin, D. B. (2004). Multiple imputation for nonresponse in surveys, Volume 81. John Wiley & Sons.
- Rubin, D. B. (2007). The design versus the analysis of observational studies for causal effects: parallels with the design of randomized trials. *Statistics in Medicine* 26(1), 20–36.
- Ruppert, D., M. P. Wand, and R. J. Carroll (2003). Semiparametric regression. Number 12. Cambridge university press.
- Rusmassen, C. and C. Williams (2005). Gaussian process for machine learning.
- Saarela, O., L. R. Belzile, and D. A. Stephens (2016). A bayesian view of doubly robust causal inference. Biometrika 103(3), 667–681.
- Santos, A., N. McGuckin, H. Y. Nakamoto, D. Gray, and S. Liss (2011). Summary of travel trends: 2009 national household travel survey. Technical report.
- Savitsky, T. D., D. Toth, et al. (2016). Bayesian estimation under informative sampling. *Electronic Journal of Statistics* 10(1), 1677–1708.
- Scharfstein, D. O., A. Rotnitzky, and J. M. Robins (1999). Adjusting for nonignorable drop-out using semiparametric nonresponse models. *Journal of the American Statistical Association* 94 (448), 1096–1120.
- Seeger, M. (2000). Relationships between gaussian processes, support vector machines and smoothing splines. Machine Learning.
- Shi, J. Q. and T. Choi (2011). Gaussian process regression analysis for functional data. CRC Press.

- SHRP2 (2013). The 2nd Strategic Highway Research Program Naturalistic Driving Study Dataset. Transportation Research Board National Academy of Sciences.
- Si, Y., N. S. Pillai, A. Gelman, et al. (2015). Bayesian nonparametric weighted sampling inference. Bayesian Analysis 10(3), 605–625.
- Solin, A. and S. Särkkä (2020). Hilbert space methods for reduced-rank gaussian process regression. Statistics and Computing 30(2), 419–446.
- Tan, Y. V., C. A. Flannagan, and M. R. Elliott (2019). "robust-squared" imputation models using bart. Journal of Survey Statistics and Methodology 7(4), 465–497.
- Tefft, B. (2017). Rates of motor vehicle crashes, injuries and deaths in relation to driver age, united states, 2014-2015. AAA Foundation for Traffic Safety..
- Tourangeau, R., J. M. Brick, S. Lohr, and J. Li (2017). Adaptive and responsive survey designs: A review and assessment. Journal of the Royal Statistical Society: Series A (Statistics in Society) 1(180), 203–223.
- Valliant, R. and J. A. Dever (2011). Estimating propensity adjustments for volunteer web surveys. Sociological Methods & Research 40(1), 105–137.
- Valliant, R., J. A. Dever, and F. Kreuter (2013). Practical tools for designing and weighting survey samples. Springer.
- Valliant, R., J. A. Dever, and F. Kreuter (2018). Nonprobability Sampling, pp. 565–603. Springer International Publishing.
- Vegetabile, B. G. (2018). Methods for Optimal Covariate Balance in Observational Studies for Causal Inference. Ph. D. thesis, UC Irvine.
- Vittert, L., R. D. Enos, and S. Ansolabehere (2020). Predicting the 2020 presidential election. Harvard Data Science Review 2(4).
- Wang, B. and A. Xu (2019). Gaussian process methods for nonparametric functional regression with mixed predictors. *Computational Statistics & Data Analysis 131*, 80–90.
- Wang, L., B. I. Graubard, H. A. Katki, and Y. Li (2020). Improving external validity of epidemiologic cohort analyses: a kernel weighting approach. *Journal of the Royal Statistical Society: Series A* (Statistics in Society) 183(3), 1293–1311.
- Wang, L., R. Valliant, and Y. Li (2020). Adjusted logistic propensity weighting methods for population inference using nonprobability volunteer-based epidemiologic cohorts. arXiv preprint arXiv:2007.02476.
- Wang, W., D. Rothschild, S. Goel, and A. Gelman (2015). Forecasting elections with non-representative polls. International Journal of Forecasting 31 (3), 980–991.
- Williams, C. K. and C. E. Rasmussen (2006). Gaussian processes for machine learning, Volume 2. MIT press Cambridge, MA.
- Williams, D. and J. M. Brick (2018). Trends in us face-to-face household survey nonresponse and level of effort. Journal of Survey Statistics and Methodology 6(2), 186–211.
- Williams, M. R. and T. D. Savitsky (2021). Uncertainty estimation for pseudo-bayesian inference under complex sampling. *International Statistical Review* 89(1), 72–107.
- Wu, C. and R. R. Sitter (2001). A model-calibration approach to using complete auxiliary information from survey data. Journal of the American Statistical Association 96(453), 185–193.
- Yi, G., J. Shi, and T. Choi (2011). Penalized gaussian process regression and classification for high-dimensional nonlinear data. *Biometrics* 67(4), 1285–1294.
- Zangeneh, S. Z. (2012). Model-based methods for robust finite population inference in the presence of external information.
- Zangeneh, S. Z. and R. J. Little (2015). Bayesian inference for the finite population total from a heteroscedastic probability proportional to size sample. Journal of Survey Statistics and Methodology 3(2), 162–192.
- Zhang, G. and R. Little (2009). Extensions of the penalized spline of propensity prediction method of imputation. *Biometrics* 65(3), 911–918.
- Zhang, G. and R. Little (2011). A comparative study of doubly robust estimators of the mean with missing data. *Journal of Statistical Computation and Simulation* 81(12), 2039–2058.
- Zheng, H. and R. J. Little (2003). Penalized spline model-based estimation of the finite

populations total from probability-proportional-to-size samples. JOURNAL OF OFFICIAL STATISTICS-STOCKHOLM- 19(2), 99–118.

- Zhou, H., M. R. Elliott, and T. E. Raghunathan (2016). Synthetic multiple-imputation procedure for multistage complex samples.
- Zigler, C. M. (2016). The central role of bayes' theorem for joint estimation of causal effects and propensity scores. The American Statistician 70(1), 47–54.
- Zigler, C. M., K. Watts, R. W. Yeh, Y. Wang, B. A. Coull, and F. Dominici (2013). Model feedback in bayesian propensity score estimation. *Biometrics* 69(1), 263–273.

8. Appendix

8.1. Gaussian Processes and kernel weighting

Suppose $\hat{\pi}_i^A$ is the estimated PS for $i \in S_A$ based on a pseudo-weighting approach. Consider the following Gaussian Process (GP) regression model:

$$y_i = f(u_i) + \epsilon_i \tag{8.1}$$

where $u_i = log(\widehat{\pi}_i^A)$, and $f \sim Gp(0, K)$ with $K(u_i, u_j; \alpha, \rho) = Cov(f(u_i), f(u_j))$. From a weight-space viewpoint, one can show that the model 8.1 predicts y_i for $i \in S_R$ using a weighted sum of observed y_i in S_A as below:

$$\widehat{y}_i = \sum_{j=1}^{n_A} \widetilde{w}_{ij} y_j \tag{8.2}$$

where

$$\widetilde{w}_{ij} = \frac{k_{ij}}{\sum_{j=1}^{n_A} k_{ij}} \quad and \quad k_{ij} = k^T (u_j) \Sigma^{-1}$$
(8.3)

with $k(\hat{\pi}_j^A) = k(\hat{\pi}_j^A, \hat{\pi}_i^A)_{n_A \times 1}$. According to Huang et al. (2019), \hat{y}_i can be regarded as the Nadaraya-Watson estimator of the observed outcome and selection indicator in the population.

Considering an isotropic covariance structure, which is a function of $||u_j - u_i||$, k_{ij} quite resembles the kernel weights Wang et al. (2020), with the bandwidth h equivalent to the GP length-scale parameter ρ . Since the kernel weights obtained by GP is used in the PM estimator, it is clear that the final weights will be multiplied by w^R , i.e.

$$\widehat{w}_j = \sum_{i=1}^{n_R} k_{ij} w_i^R \tag{8.4}$$

One can show that the major kernel-related condition determined by Wang et al. (2020) to obtain consistency in the kernel-weighted estimates holds for a Matérn family covariance structure, i.e. K(u), $\int K(u)du = 1$, $Sup_u|K(u)| < \infty$, and $\lim_{|u|\to\infty} |u||K(u)| = 0$.

8.2. Further extensions of the simulation study

8.2.1. Simulation study I

This subsection provides additional results associated with Simulation I. Table 2 and Table 3 summarize the findings of the simulation in Section 3.1 for $\rho = 0.5$ and $\rho = 0.3$, respectively.

			_		Mo	h.	_		Mc	h.	_		Mc		ħ.	_		Mo			z	_		$\mathbf{P}_{\mathbf{r}}$	Me		
GPPP: Weighti NOTE:	PAPP	AIPW	LWP	GPPP	odel spe	AIPW	LWP	GPPP	odel spe	AIPW	LWP	GPPP	odel spe	PAPP	AIPW	LWP	GPPP	odel spe	FW	UW	on-prob	FW	UW	obabili	easure		
Gaussia ng; PA The P	27.912	27.115	27.075	27.400	cificatio	0.040	4.052	1.317	cificatio	-0.056	-0.007	0.146	cificatio	0.832	-0.101	-0.063	0.129	cificatio	-0.134	30.708	ability	0.260	8.866	ty sam	rBias		
an Proc PP: Pro APP an	28.191	27.402	27.361	27.678	n: QR–]	3.987	6.752	4.321	n: QR–J	3.916	3.876	3.829	n: QR-	4.012	4.035	4.200	3.876	n: QR-	3.763	30.926	sampl	3.770	9.445	ple $(S_F$	rMSE	$n_A =$	
ess of Pi opensity	0.000	0.000	0.000	3 0.000	False, Pl	95.83	89.81	99.07	False, Pl	94.90'	99.53'	99.53'	Γrue, PΝ	93.05	93.98	99.53'	99.53'	Γrue, PN	93.98	0.000	$e(S_A)$	93.050	21.75	2)	crCI	$500, n_{I}$	
ropensit -adjuste	1.354	1.366	2.247	2.239	M–False	3 1.392	5 2.236	4 1.934	M-True	7 1.358	7 1.910	7 1.903	I–False	5 1.406	1 1.406	7 2.516	7 1.912	I -True	1 1.363	1.304		5 1.363	9 1.149		lCI	r = 500	
y Predic ed Prob	0.936	0.944	1.558	1.564		0.954	1.132	1.285		0.948	1.348	1.360		0.980	0.953	1.638	1.350		0.991	0.974		0.991	0.965		rSE		
ction; LV ability I	27.02	26.43	26.43	26.59		-0.010	4.147	1.146		-0.08	-0.12	-0.01		0.547	-0.098	-0.14	0.000		-0.140	30.00		0.260	8.866		rBias		
VP: Line redictic	4 27.15	2 26.57	4 26.57	5 26.73		3.025	6.026	3.477		7 3.061	1 3.084	7 3.068		2.998	3.139	5 3.140	3.055		2.659	8 30.10		3.770	9.445		rMSE	$n_A = 1$	
ear-in-w	8 0.000	0.000	0.000	3 0.000		94.44	92.13	99.53		95.83	99.53	99.53		93.98	94.90	99.53	99.53		95.83	4 0.000		93.05	21.75		crCI	,000, r	
eight pro	0.968	0.981	2.056	2.042		4 1.074	0 2.004	7 1.748		3 1.070	7 1.730	7 1.720		1 1.078	7 1.107	7 1.737	7 1.724		3 0.957	0.916		6 1.363	9 1.149		lCI	$u_R = 500$	
ediction	0.983	0.987	2.053	2.055		0.971	1.253	1.456		0.957	1.535	1.533		1.000	0.965	1.514	1.544		0.986	1.039		0.991	0.965		rSE		
; AIPW:	28.12	27.05	27.12'	27.43		0.073	4.090	1.314		-0.180	-0.015	0.161		0.984	-0.232	-0.135	0.092		-0.134	30.70		0.080	8.810		rBias		
Augmei	1 28.381	27.328	7 27.395) 27.692		3.716	6.299	4.001		3.590	3.575	3.555		3.844	3.760	3.742	3.567		3.763	30.926		2.736	9.115		rMSE	$n_A = 5$	
nted Inv	1 0.000	3 0.000	0.000	2 0.000		93.519	87.03	96.296		93.98	98.148	98.148		95.370	94.90	97.68	98.148		93.98	0.000		95.83	2.778		crCI	$00, n_R$	
erse Pro	1.329	1.334	1.837	1.814) 1.338	7 1.850	3 1.607		1.272	3 1.585	3 1.575) 1.339	7 1.311	5 1.613	3 1.576		1.363	1.304		3 0.967	0.816		lCI	= 1,000	
pensity	0.950	0.949	1.314	1.306		0.984	1.056	1.163		0.970	1.213	1.213		0.985	0.955	1.179	1.209		0.991	0.974		0.967	0.954		rSE		

Ц
a,
Ы
Φ
N
~
2
Ħ
q,
aı
H
ω.
÷
зe
Ч
ē
f
Ĕ
В
ar
ī
ð
of
£
he
~
Ĕ
S
ð
<u>e</u> .
n.
Ξ
Ц
'n

В
et
þ
õ.
Ś
ii
4
þ
to the second se
fr
st
S
in
ц
la
ti
10
5
stı
pr
Y
fc
\mathbf{r}
ρ
Ш
0
С

NOTE: The PAPP and AIP w ше ę IIdve 0 en implemented through a bootstrap method.

÷
0
θ
for
dy
stu
tion
ılat
simı
first
the
in
spc
thc
me
ent
stm
suljus
bias a
he
of t
e e
nanc
for
per
the
ng 1
arii
mp
So
ä
le
[ab

		$n_A = 5$	$00, n_B$	= 500			$i_A = 1, 0$	$100, n_R$	= 500		ı	$n_A = 500$	$n_{B} = 1$: 1,000	
Measure	rBias	rMSE	crCI	ICI	rSE	rBias	rMSE	crCI	ICI	rSE	rBias	$\mathbf{r}MSE$	crCI	ICI	rSE
Probabil	ity saml	ple (S_R)													
ΜŊ	8.867	10.406	58.796	1.927	0.965	8.867	10.406	58.796	1.927	0.965	8.797	9.612	38.889	1.369	0.964
FW	0.424	6.361	92.593	2.303	0.989	0.424	6.361	92.593	2.303	0.989	0.154	4.580	95.370	1.633	0.973
Non-prol	bility	sample	(S_A)												
ΠM	30.759	31.251	0.000	2.037	1.006	30.082	30.322	0.000	1.435	1.028	30.759	31.251	0.000	2.037	1.006
FW	-0.277	6.284	95.833	2.335	1.014	-0.298	4.529	93.519	1.639	0.989	-0.277	6.284	95.833	2.335	1.014
Model spe	cification	n: QR-Th	rue, PM–	True											
GPPP	0.438	6.297	100.000	3.413	1.482	0.061	4.754	99.537	3.094	1.775	0.489	6.052	99.074	2.821	1.275
LWP	-0.342	6.811	100.000	4.061	1.628	-0.266	4.917	99.537	3.131	1.739	-0.232	6.429	98.148	2.932	1.244
AIPW	-0.390	6.666	95.370	2.378	0.975	-0.296	4.911	94.444	1.737	0.966	-0.395	6.337	95.370	2.328	1.004
PAPP	0.574	6.537	95.370	2.366	0.991	0.360	4.696	95.370	1.709	0.995	0.873	6.291	94.907	2.334	1.022
Model spe	cification	1: QR-Th	rue, PM-	False											
GPPP	0.553	6.241	100.000	3.391	1.488	0.077	4.733	99.537	3.080	1.775	0.585	6.033	98.611	2.817	1.280
LWP	-0.115	6.302	100.000	3.464	1.499	-0.255	4.748	99.537	3.105	1.786	-0.020	6.114	99.074	2.879	1.284
AIPW	-0.271	6.301	95.370	2.312	1.002	-0.275	4.789	94.444	1.688	0.963	-0.287	6.042	94.907	2.233	1.009
Model spe	cification	a: QR-Fa	alse, PM-	-True											
GPPP	1.891	6.750	99.537	3.409	1.435	1.372	5.092	99.537	3.097	1.722	1.981	6.536	97.685	2.837	1.242
LWP	4.179	8.638	98.611	3.692	1.332	4.373	7.326	99.537	3.309	1.535	4.277	8.280	95.833	3.098	1.192
AIPW	-0.146	6.594	94.907	2.377	0.983	-0.187	4.794	94.444	1.708	0.973	-0.125	6.319	95.370	2.333	1.007
Model spe	cification	n: QR-Fa	alse, PM-	-False											
GPPP	27.642	28.234	5.093	3.483	1.652	26.734	27.051	2.315	3.212	2.121	27.792	28.364	0.463	2.792	1.344
LWP	26.915	27.545	11.111	3.542	1.649	26.278	26.601	2.315	3.242	2.142	27.022	27.636	1.852	2.900	1.366
AIPW	26.987	27.603	0.000	2.120	0.997	26.405	26.734	0.000	1.528	0.996	27.022	27.630	0.000	2.076	0.982
PAPP	27.881	28.473	0.000	2.122	1.002	26.994	27.300	0.000	1.501	1.004	28.146	28.716	0.000	2.104	1.008
	Council	Duces	or of Duc	- tipoo o	Ducklink	Const VA/E	. T incom	aion oi	benne td.	liotion.	A TDXX7. A	+0000000	ad Inno	Duor	
	Caussia	un rroce	SS UL LTU	pensity	Freulu.	IOII, LVVF	. TUTTEAL	-III-weig	ur pret	IICUIOII; -	ALF VV : A	ungmenu	ed IIIver	se rru	ensuy

Weighting; PAPP: Propensity-adjusted Probability Prediction NOTE: The PAPP and AIPW methods have been implemented through a bootstrap method.

8.2.2. Simulation study II

Table 4 exhibits the numerical results associated with Figure 2 and Figure 3.

Table 4: Comparing the performance of the bias adjustment methods in the second simulation study for the *continuous* outcome with $(n_A, n_R) = (500, 1, 000)$ and $\gamma_1 = 0.3$

ч U	⊳ t	50	Moo	A	L,	G	Mod	A	Ľ	G	Mod	A	Ľ	G	Mod	Dou	P	Moc	P	Mod	Nor	দ	U	Nor	দ	U	\mathbf{Pro}	Mea		
W: Ur	IPW	NP -	lel spec	1PW	WΡ	PPP	lel spec	IPW	WΡ	PPP	lel spec	IPW	WΡ	PPP	lel spec	ubly re	APP	lel spec	APP	lel spec	1-robu	W	W	1-prob	W	W	babilit	asure		
ıweighte	36.146	36 049	af afo	0.546	-0.294	0.523	ification	-0.033	10.099	0.099	ification	0.000	-0.038	0.025	ification	bust m	35.346	ification	0.066	ification	st meth	-0.178	48.795	ability s	-0.180	-23.870	;y samp	rBias		and 771 -
d; FW:	36.468	36 361	26 676	6.979	4.923	4.628	QR-Fal	4.608	4.776	4.467	QR-Tru	4.439	4.478	4.439	QR-Tru	ethods	35.660	QR-Fal	4.771	QR-Tru	od	6.184	49.086	ample (3.921	24.078	le (S_R)	rMSE		= U.J
Fully w	0.000	0 000	n nnn	94.444	96.296	97.685	se, PM–	94.907	91.667	97.685	ie, PM–	97.222	97.222	98.148	1e, PM-'		0.000	se	95.370	ıe		91.204	0.000	(S_A)	93.519	0.000		crCI	LIN	
veighted	17.448	22.546	Palse	29.210	20.356	20.143	True	18.233	17.809	18.732	False	18.266	18.960	18.703	True		17.238		19.982			22.458	19.644		15.087	12.411		ICI		
; GPPP	0.919	1 219	1 917	1.069	1.064	1.128		1.007	0.100	1.078		1.047	1.088	1.081			0.929		1.066			0.925	0.936		0.980	1.001		rSE		
: Gaussi	18.293	18 306	18 999	-15.872	0.231	1.310		-0.109	7.636	-0.061		0.250	-0.005	-0.060			17.490		-7.545			-0.218	30.073		-0.156	-17.120		rBias		
an Proc	18.583	18 690	18 59/	46.090	6.026	3.829		3.198	3.382	3.136		4.060	3.133	3.150			17.760		24.919			4.249	30.366		2.749	17.243		rMSE		
ess of P	0.000	0 000	0 0 0	87.500	93.981	95.370		95.833	88.889	97.222		98.148	98.148	96.759			0.000		91.667			92.593	0.000		94.907	0.000		crCI	CUB	
ropensity	12.436	15 611	1л дод	102.060	15.853	15.280		12.461	12.078	12.796		16.137	13.227	13.124			11.878		53.569			15.612	15.989		10.799	8.442		ICI		
y Predic	0.967	1 213	1 105	0.600	0.677	1.093		0.992	0.087	1.047		1.014	1.084	1.067			0.980		0.574			0.936	0.966		1.001	1.045		rSE		
tion; LW	10.587	10.390	10 /09	6.885	0.509	0.575		-0.048	5.845	0.005		-0.022	-2.254	-0.027			10.111		2.269			0.004	18.146		-0.091	-12.097		rBias		
/P: Line	10.848	10.100	10 758	7.391	5.817	2.531		2.188	17.130	2.140		2.194	4.458	2.164			10.362		3.182			2.282	18.363		1.947	12.179		rMSE		
ar-in-wei	0.000	0.926	960 0	26.389	96.759	96.296		95.370	89.815	96.296		93.981	95.833	97.222			0.000		83.333			93.981	0.000		94.444	0.000		crCI	EXP	
ight pre	8.968	11 005	10 0/1	10.549	11.564	11.294		8.400	8.171	8.654		8.482	9.065	8.823			8.707		8.794			8.639	10.710		7.392	5.427		lCI		
diction;	0.965	1 185	1 1 2 9	0.999	0.512	1.175		0.977	0.078	1.037		0.984	0.080	1.048			0.977		1.003			0.964	0.969		0.967	0.976		rSE		
AIPW:	30.664	30 650	30.056	14.490	-1.060	0.819		0.245	7.886	0.439		0.157	0.299	0.301			30.831		11.642			-0.034	27.948		0.015	-18.475		rBias		
Augment	31.483	31 450	21 75N	15.884	14.048	6.523		6.221	14.740	6.189		6.179	6.335	6.250			31.610		13.063			8.276	28.716		5.549	19.159		rMSE		
ted Inve	0.000	2 778	2 9/1	35.648	92.130	95.833		94.907	92.130	95.833		94.444	96.296	96.296			0.463		46.759			93.519	0.926		93.519	3.241		crCI	$_{NIS}$	
rse Prop	26.807	35 040	33 560	24.548	37.060	27.890		23.752	24.236	25.041		23.568	26.621	26.369			26.668		23.010			31.425	25.807		20.503	18.535		lCI		
pensity	0.956	1 276	1 999	0.96	0.679	1.108		0.972	0.175	1.043		0.971	1.076	1.084			0.973		0.988			0.966	0.995		0.94	0.93		rSE		

Weighting. NOTE: The PAPP and AIPW methods have been implemented through a bootstrap method.

Table 5: Comparing the performance of the bias adjustment methods in the second simulation study for the *continuous* outcome with $(n_A, n_B) = (1, 000, 500)$

ņ
\circ
Ш
γ_1
and

			LIN					GILB					E V D					CIN		
sure	rBias	rMSE	crCI	ICI	rSE	rBias	rMSE	crCI	ICI	rSE	rBias	$\mathbf{r}\mathbf{MSE}$	crCl	ICI	rSE	rBias	rMSE	crCI	ICI	rSE
babilit	y samp	de (S_R)																		
MM	-23.868 0.118	24.297 5.477	$0.000 \\ 93.519$	17.587 21.395	0.985 0.994	-17.158 0.016	17.440 3.998	$\frac{0.000}{94.444}$	$\frac{11.959}{15.257}$	$0.974 \\ 0.971$	-12.107 0.015	$12.274 \\ 2.673$	$\frac{0.000}{95.833}$	7.683 10.454	$0.967 \\ 0.995$	-18.706 -0.081	19.966 7.162	22.222 96.296	26.219 29.064	$0.956 \\ 1.033$
n-prob	ability s	sample	(S_A)																	
M	48.581	48.727	0.000	13.878 15 070	0.935	29.971 0.107	30.129	0.000	11.296	0.933	18.042	18.159 1 596	0.000	7.569	0.935	27.647 0.907	28.037 = 417	0.000	18.285	0.999
n-robus	st meth	po	000.00	710.01	0.000	10110	7007	100.10	00000	0100	0000	000-T	007.00	1710	100.0	07.0-	112-0	00000	007.77	0=0.1
lel spec	ification	: QR-Tr	ue																	
APP	0.504	4.495	96.296	18.667	1.064	-16.241	32.851	98.611	74.882	0.667	0.403	1.995	95.833	7.859	1.024	11.569	13.086	53.241	24.302	1.011
del spec	ification	: QR-Fa	lse																	
APP	35.317	35.521	0.000	14.654	0.982	17.616	17.838	0.000	10.819	0.982	10.136	10.324	0.000	7.535	0.977	30.431	30.802	0.000	18.895	1.008
ubly ro	bust m	ethods																		
del spec	ification	: QR-Tr	ue, PM-	-True																
PPP	0.326	4.758	98.611	21.123	1.142	0.169	3.505	96.759	14.875	1.091	0.120	2.338	96.759	10.083	1.108	0.291	6.396	96.759	29.612	1.186
WP	0.285	4.746	98.611	21.279	1.152	-1.313	3.504	96.759	16.475	0.196	-0.655	8.930	96.296	10.952	0.316	0.246	6.403	97.685	29.663	1.19
MdI	0.287	4.808	97.222	20.161	1.069	0.767	5.589	97.222	20.610	0.947	0.140	2.361	96.296	9.838	1.062	0.292	6.293	95.370	26.295	1.065
lel spec	ification	: QR-Tr	ue, PM-	-False																
PPP	0.397	4.731	97.685	21.125	1.151	0.123	3.530	97.222	14.813	1.072	0.096	2.337	98.148	10.056	1.097	0.340	6.358	97.685	29.112	1.175
WP	21.964	5.322	84.259	19.456	0.077	16.925	4.571	81.019	13.776	0.071	12.837	31.570	78.241	9.029	0.061	17.658	14.123	85.648	26.438	0.131
IPW	0.430	4.877	97.685	20.009	1.048	0.153	3.531	95.833	14.218	1.026	0.108	2.359	96.759	9.660	1.043	0.357	6.324	95.370	26.422	1.065
del spec	ification	: QR-Fa	lse, PM-	-True																
PPP	0.679	4.644	98.611	22.199	1.242	1.028	3.562	98.611	16.651	1.251	0.538	2.275	99.074	12.101	1.401	0.734	6.435	98.148	30.583	1.228
WP	0.107	4.712	99.074	22.396	1.223	-1.191	3.593	97.222	17.681	0.335	0.118	2.437	99.537	12.407	1.310	-2.162	8.254	96.759	38.983	1.248
ΠΡW	1.102	6.334	95.833	25.788	1.052	-33.598	67.410	95.833	153.677	0.669	6.226	6.774	39.815	10.775	1.028	14.128	15.372	36.111	24.900	1.046
del spec	ification	: QR-Fa	lse, PM-	-False																
PPP	36.360	36.574	0.000	24.865	1.615	18.538	18.801	0.000	17.376	1.427	10.648	10.868	0.463	12.232	1.443	30.417	30.823	1.852	36.965	1.893
WP	36.137	36.354	0.000	24.980	1.612	18.636	18.902	0.000	17.596	1.426	10.702	10.920	0.926	12.309	1.452	30.259	30.656	0.926	38.086	1.988
IPW	36.248	36.471	0.000	15.439	0.976	18.546	18.800	0.000	12.121	1.003	10.730	10.951	0.000	8.629	1.005	30.357	30.749	0.000	19.215	0.999
JW: Ur	ıweighte	d; FW:	Fully v	weighted;	GPPP:	Gaussiaı	1 Proces	ss of Pro	opensity	Predicti	ion; LWF	: Linea	-in-weig	tht Pred	liction; /	AIPW: 4	Augment	ted Inver	se Prop	ensity

Weighting. NOTE: The PAPP and AIPW methods have been implemented through a bootstrap method.

NC	UU UU	AII	LW	GP	Mode	AII	LW	GP	Mode	AII	LW	GP	Mode	AII	LW	GP	Mode	Doub	PA.	Mode	PA	Mode	Non-	FW	WU	Non-	FW	MD	Prob	Meas		
ighting TE: T	/: Unv	W 3	Р 3	PP 3	specif	⁹ W 1	Р -	PP 0	l specif	0 Me	P 1	PP 0	l specif	0 Me	P 0	PP 0	l specif	ly rob	PP 3	l specif	PP 0	specif	robust	L	4	probal	, 0	۲ ۱	ability	ure		
he PA	veighte	6.291	6.191	6.537	ication	.226	1.094	.879	ication	.291	0.150	.402	ication	.292	.293	.359	ication	ust m	5.408	ication	.487	ication	meth	0.178	8.795	oility s	.118	23.868	samp	Bias		
PP and	d; FW:	36.644	36.544	36.882	: QR–Fa	7.554	5.826	5.451	: QR–Fa	5.475	5.659	5.359	: QR-Tr	5.410	5.384	5.406	: QR–Tr	ethods	35.754	: QR–Fa	5.489	: QR-Tr	od	6.184	49.086	ample	5.477	24.297	le (S_R)	rMSE		
AIPW	Fully	0.000	0.000	0.000	lse, PM	94.444	97.685	97.685	lse, PM	95.370	91.204	96.759	ue, PM-	96.296	97.222	97.222	ue, PM-		0.000	lse	94.907	ue.		91.204	0.000	(S_A)	93.519	0.000		crCI	LIN	
method	weightee	19.219	27.447	27.271	–False	31.434	24.830	24.471	-True	22.112	22.008	23.118	-False	22.264	23.243	23.039	-True		18.412		22.317			22.458	19.644		21.395	17.587		ICI		
s have b	i; GPPI	0.964	1.391	1.392		1.073	0.530	1.169		1.029	0.126	1.110		1.049	1.108	1.098			0.945		1.039			0.925	0.936		0.994	0.985		rSE		
een impl	: Gaussi	18.489	18.536	18.359		-20.691	-1.064	1.589		0.136	7.552	0.149		0.570	0.130	0.147			17.598		-10.305			-0.218	30.073		0.016	-17.158		rBias		
emented	ian Proc	18.836	18.894	18.731		1 52.415	4.293	4.338		3.982	4.184	3.926		4.891	3.860	3.855			17.930		5 26.492			4.249	30.366		3.998	3 17.440		rMSE		
l through	cess of P	0.000	0.000	0.000		92.130	96.759	95.833		95.833	88.426	95.370		97.222	96.759	96.296			0.000		94.444			92.593	0.000		94.444	0.000		crCI	CUB	
ı a boots	ropensit	14.274	19.235	19.077		121.585	19.139	18.450		15.500	14.928	15.959		20.056	16.270	16.208			13.143		62.285			15.612	15.989		15.257	11.959		ICI		
trap met	y Predic	1.010	1.350	1.320		0.643	0.535	1.172		0.991	0.110	1.039		1.051	1.083	1.080			0.974		0.650			0.936	0.966		0.971	0.974		rSE		
shod.	tion; LW	10.726	10.681	10.586		6.562	0.132	0.715		0.086	5.872	0.137		0.100	0.060	0.112			10.135		1.287			0.004	18.146		0.015	-12.107		rBias		
	P: Linea	11.039	10.998	10.905		7.243	2.775	2.775		2.662	23.999	2.580		2.610	2.595	2.546			10.422		2.842			2.282	18.363		2.673	12.274		rMSE		
	ar-in-wei	0.463	7.870	8.333		43.981	98.148	96.759		95.370	88.426	96.759		96.759	97.222	97.685			0.000		91.667			93.981	0.000		95.833	0.000		crCI	EXP	
	ght Pre	10.142	13.431	13.442		12.248	13.657	13.472		10.677	10.056	10.813		10.631	11.233	10.919			9.452		9.682			8.639	10.710		10.454	7.683		ICI		
	diction;	0.989	1.314	1.317		1.017	1.262	1.290		1.022	0.098	1.077		1.037	1.111	1.105			0.991		0.973			0.964	0.969		0.995	0.967		rSE		
	AIPW:	30.643	30.488	30.765		14.329	-1.868	0.827		0.156	7.821	0.389		0.311	0.306	0.235			30.805		11.663			-0.034	27.948		-0.081	-18.706		rBias		
	Augmen	31.452	31.305	31.605		16.096	11.807	7.547		7.197	17.007	7.230		7.290	7.389	7.323			31.598		13.627			8.276	28.716		7.162	19.966		rMSE		
	ted Inve	0.000	9.259	9.259		50.000	94.444	96.296		94.907	91.204	97.685		96.296	96.296	96.296			0.463		60.648			93.519	0.926		96.296	22.222		crCI	NIS	
	rse Prop	26.975	42.309	40.674		28.830	43.580	33.826		28.904	29.704	31.411		29.106	32.653	32.247			26.762		27.601			31.425	25.807		29.064	26.219		ICI		
	pensity	0.968	1.529	1.443		1.001	0.96	1.156		1.022	0.214	1.112		1.017	1.133	1.129			0.968		0.997			0.966	0.995		1.033	0.956		rSE		

	Tabl
	e 6:
and	Con
$\gamma_1 =$	ıpari
$^{-}0.3$	ng tl
	ıe p∈
	erfori
	manc
	e of
	the l
	oias a
	adjus
	stme
	at m
	etho
	ds in
	the :
	secoi
	ıd siı
	mula
	tion
	stud
	y for
	$_{\mathrm{the}}$
	cont
	inuo
	us ou
	itcor
	ne wi
	ith (1
	n_A, n
	(R)
	11
	=(50)

/Robust and Efficient Bayesian Inference for Non-Probability Samples

	/F	cob	u_{S}
(0			rS
0, 1, 00			ICI
) = (50)		SIN	crCI
(n_A,n_R)			rMSE
me with			rBias
<i>us</i> outco			rSE
ontinuo			ICI
r the cc		EXP	crCI
study fo			rMSE
ulation			rBias
cond sin			rSE
1 the se			ICI
hods ir		CUB	crCI
nent met			rMSE
s adjustr			rBias
of the bia			rSE
nance c			ICI
perforn		LIN	crCI
ring the $= 0.6$			rMSE
Compa: and γ_1			rBias
Table 7: (Measure ⁻
-		-	

Ŕ	ob	$u_{\rm SI}^{\rm st}$	an	0.98	Ef 0.0	fic	1.038	1.027	B	ay	0.935	ar	0.979	nf	er	1.067	1.088	0.974	r.	0.964	0.034	₹6.0	rot	1.091	0.236	0.948	S_{i}	1.0 4	0.288	0.938	ghting.
		ICI		18.535	20.503		25.372	49.982			22.936		28.544			33.426	39.758	25.116		28.027	27.053	25.796		39.581	92.284	28.031		31.475	28.693	28.899	sity Wei
	SIN	crCI		3.241	93.519		0.000	94.907			0.926		0.000			93.519	94.444	95.370		93.519	61.111	92.593		94.444	56.481	0.000		0.000	0.000	0.000	Propens
		rMSE		19.159	5.549		37.411	12.399			28.131		55.056			8.500	9.842	6.698		7.465	7.895	7.136		9.473	46.251	35.388		55.704	125.351	55.217	d Inverse
		rBias		-18.475	0.015		36.869	-0.451			27.459		54.554			0.083	-0.139	-0.070		0.252	68.011	0.111		-0.452	36.851	34.579		55.161	166.531	54.681	Augmente
		$^{\mathrm{rSE}}$		0.976	0.967		0.978	0.947			0.969		0.928			1.011	1.008	0.927		0.985	0.074	0.893		0.971	0.202	0.972		1.060	0.260	0.936	AIPW: ∤
		ICI		5.427	7.392		13.875	13.283			10.631		9.782			11.306	12.962	9.824		9.677	21.390	9.389		16.145	52.561	11.967		11.227	25.490	9.978	liction;
	EXP	crCI		0.000	94.444		0.000	92.593			49.537		0.000			95.833	96.296	93.981		95.370	60.185	95.833		90.278	82.407	3.704		0.000	0.000	0.000	ght Prec
		rMSE		12.179	1.947		45.091	3.570			5.987		15.440			2.871	3.301	2.696		2.529	28.166	2.678		4.765	19.000	11.978		14.437	113.140	14.575	ar-in-wei
		rBias		-12.097	-0.091		44.946	-0.070			5.295		15.205			0.017	-0.076	-0.055		-0.118	52.013	-0.124		2.127	32.100	11.560		14.179	146.864	14.321	WP: Line od.
		rSE		1.045	1.001		0.992	0.782			0.435		0.928			1.012	1.035	0.853		1.038	0.062	0.912		0.874	0.145	0.365		1.053	0.250	0.944	iction; L p methe
		ICI		8.442	10.799		21.935	27.339			46.558		13.272			16.700	19.394	19.238		14.732	26.171	13.978		21.919	68.192	92.613		15.244	28.728	13.586	ty Predi pootstra
	CUB	crCI		0.000	94.907		0.000	91.667			48.148		0.000			97.685	96.759	93.056		94.444	58.796	92.593		81.944	78.704	48.611		0.000	0.463	0.000	Propensi ough a l
	-	rMSE		17.243	2.749		74.114	8.917			27.294		24.516			4.219	4.808	5.745		3.652	4.030	3.903		8.273	22.233	64.920		23.242	118.013	23.115	ocess of I ented thr
		rBias		-17.120	-0.156		73.900	-0.515			1.511		24.244			-0.014	-0.168	-0.214		-0.227	77.157	-0.148		5.206	45.933	-5.904		22.943	177.674	22.823	ussian Pr n implem
		rSE		1.001	0.980		0.929	0.955			1.065		0.921			0.995	1.061	0.989		1.027	0.065	0.890		1.039	0.166	0.967		1.069	0.255	0.924	PPP: Ga lave bee
		ICI		12.411	15.087		20.820	40.191			36.302		18.790		eue	23.490	28.647	24.890	ilse	24.644	36.944	22.724	rue	27.203	66.505	53.458	alse	20.925	28.177	18.450	hted; Gl ethods h
	LIN	crCI		0.000	93.519	S_A)	0.000	92.130		Ð	94.444	e	0.000		e, PM-T	95.833	96.759	93.981	e, PM–Fa	95.370	61.111	92.593	e, PM-T	94.907	62.037	90.278	e, PM-F	0.000	0.000	0.000	ully weig AIPW m
		rMSE	le (S_R)	24.078	3.921	ample (100.399	10.721	po	QR-Tru	8.677	QR-Fals	53.104	ethods	QR-Tru	6.062	6.959	6.410	QR-Tru	6.177	6.228	6.499	QR-Fals	6.880	24.018	14.115	QR-Fals	51.589	96.444	51.947	l; FW: F PP and A
		rBias	ty sampl	-23.870	-0.180	ability s	100.237	-0.428	ist meth	cification:	0.267	cification:	52.850	obust me	cification:	0.023	-0.400	-0.244	cification:	0.578	102.755	0.108	cification:	1.435	38.447	1.156	cification:	51.343	195.557	51.698	nweighted The PAI
		Measure	Probabili	MD	FW	Non-prob	MN	FW	Non-robu	Model spec	PAPP	Model spec	PAPP	Doubly rc	Model spec	GPPP	LWP	AIPW	Model spec	GPPP	LWP	AIPW	Model spec	GPPP	LWP	AIPW	Model spec	GPPP	LWP	AIPW	UW: UI NOTE:

 $\overline{7}$

UW: NOT	AIPV	LWP	GPPI	Model s	AIPV	LWP	GPPI	Model s	AIPV	LWP	GPPI	Model s	AIPV	LWP	GPPI	Model s	Doubly	PAPI	Model s	PAPI	Model s	Non-ro	FW	WU	Non-pr	FW	WU	Probab	Measu		
Unweigh E: The F	V 51.84:	186.6	51.23	pecification	V = 1.481	-3.714	2 1.462	pecification	V = 0.764	160.6	0.847	pecification	V = 0.328	0.696	0.467	pecification	' robust	51.93	pecification	0.837	pecification	bust me	-0.355	98.62	obability	0.118	-23.86	ility san	re rBias		and
ted; FW: APP and	3 52.02	28 124.6	4 51.40.	on: QR–F	12.22	1 21.45	5.582	on: QR–F	5.471	57 5.868	5.217	on: QR–T	5.304	9.836	5.136	on: QR–T	methods	8 52.10	on: QR–F	6.971	on: QR–T	thod	5 8.567	9 98.70	y sample	5.477	38 24.29	aple (S_R)	rMSE		1 - 0.0
Fully w I AIPW	1 0.000	70 0.000	4 0.000	alse, PM-	1 88.88	4 78.70	97.68	alse, PM-	96.75	44.44	96.29	rue, PM-	96.29	97.22	97.68	rue, PM-	-	2 0.000	alse	93.98	rue		90.27	9 0.000	(S_A)	93.51	7 0.000)	crCI	LIN	
eighted; methods) 16.91	36.04	22.46	–False	39 44.39)4 31.84	35 25.71	-True	59 22.25	14 57.99	6 23.74	-False	6 23.71	22 25.61	35 23.45	-True) 16.10		31 28.49			78 29.83) 14.60		[9 21.39]) 17.58		ICI		
GPPP: (s have be	8 1.001	8 0.281	4 1.382		4 0.931	9 0.335	3 1.227		4 1.046	5 0.103	0 1.191		2 1.140	7 0.670	7 1.177			6 0.990		6 1.048			5 0.887	6 0.936		5 0.994	7 0.985		rSE		
Gaussian en implei	23.384	168.28(23.626		-35.926	1.510	4.015		0.188	121.45!	-0.091		0.323	0.169	0.310			23.574		-15.335			-0.483	71.503		0.016	-17.158		rBias		
Process o mented th	23.626) 123.557	23.875		i 106.302	13.601	6.286		3.787	5 15.991	3.657		9.295	3.997	3.766			23.768		47.984			7.679	71.609		3.998	3 17.440		rMSE		
f Propen 1rough a	0.000	7 0.000	0.000		2 75.926	91.667	89.352		95.370	38.889	97.222		96.759	96.759	97.222			0.000		81.944			88.889	0.000		94.444	0.000		crCI	CUB	
sity Prec bootstra	13.378	36.124	16.807		175.824	23.485	20.044		14.917	50.917	15.703		26.822	17.628	16.493			11.833		80.081			21.564	15.102		15.257	11.959		lCI		
p metho	1.008	0.283	1.249		0.447	0.307	1.062		1.004	0.120	1.102		0.735	1.134	1.126			0.993		0.448			0.716	0.988		0.971	0.974		rSE		
WP: Line d.	14.678	138.694	14.541		10.829	1.474	1.752		0.103	79.393	-0.035		0.236	0.129	0.237			14.805		2.217			-0.008	43.622		0.015	-12.107		rBias		
ear-in-wei	14.877	108.095	14.741		11.221	13.319	3.622		2.524	20.870	2.398		2.504	2.695	2.519			14.960		3.140			2.547	43.693		2.673	12.274		rMSE		
ght Pred	0.000	0.000	0.000		5.093	92.593	97.685		97.222	42.130	97.222		96.296	98.611	96.759			0.000		86.574			92.130	0.000		95.833	0.000		crCI	EXP	
liction; A	9.732	32.502	12.211		12.150	19.695	14.822		10.169	40.288	10.482		10.389	12.199	11.192			8.330		9.040			9.577	9.597		10.454	7.683		lCI		
JPW: A	1.020	0.287	1.293		1.051	0.383	1.197		1.026	0.139	1.121		1.061	1.161	1.145			0.987		1.035			0.957	0.980		0.995	0.967		rSE		
ugmente	55.491	158.745	54.802		35.268	-7.084	0.278		0.301	99.141	0.301		0.292	0.240	0.247			55.512		27.350			-0.590	37.338		-0.081	-18.706		rBias		
d Inverse	55.756	122.172	55.140		35.750	31.041	7.406		6.422	6.863	6.461		6.326	7.532	6.833			55.750		27.937			9.229	37.581		7.162	19.966		rMSE		
) Propens	0.000	0.000	0.000		0.000	68.981	97.222		94.907	45.370	97.685		95.833	97.685	98.148			0.000		0.463			93.519	0.000		96.296	22.222		crCI	$_{NIS}$	
sity Wei	21.258	34.886	33.025		26.074	51.814	36.447		27.431	34.496	30.374		27.065	35.788	32.759			20.689		23.160			36.092	17.907		29.064	26.219		lCI		
inting.	0.997	0.292	1.393		1.134	0.44	1.265		1.088	0.092	1.209		1.09	1.216	1.234			1.022		1.035			0.997	1.067		1.033	0.956		rSE		

Table 8: Comparing the performance of the bias adjustment methods in the second simulation study for the *continuous* outcome with $(n_A, n_R) = (1, 000, 500)$ and $\gamma_1 = 0.6$

	/F	lob	₿t	
0,500)			ICI	
) = (50)		SIN	crCI	
(n_A, n_R)			rMSE	
ome with			rBias	
ous outco			\mathbf{rSE}	
ontinuc			ICI	
\mathbf{r} the c		EXP	crCI	
study fo			rMSE	
nulation			rBias	
econd sin			\mathbf{rSE}	
n the se			ICI	
thods i		CUB	crCI	
ment me			rMSE	
as adjusti			rBias	
of the bia			\mathbf{rSE}	
nance (ICI	
perfor		LIN	crCI	
ring the $= 0.6$			rMSE	
Compa and γ_1			rBias	
Table 9:			Measure	

'F	cob	E tr	ar	0.956	1 1 1 3 3 3 3	fic	1.618	1.627	B	ay	0.963	iar	$0.\overline{9}65$	nf	er	1.855	0.843	1. B 05	r.	1.637	0.001	0.985	roł	1. 1 7	0.382	1,012	S	$1. \frac{1}{2} 99$	0.622	н С
		lCI		26.219	29.064		25.372	49.982			26.590		28.872			38.202	43.943	30.253		33.641	33.876	30.870		43.960	74.878	31.247		37.361	36.164	29.376
	SIN	crCI		22.22	96.296		0.000	94.907			1.852		0.000			96.759	97.222	94.907		95.370	62.037	94.907		95.370	68.519	0.926		0.000	0.000	0.000
		$\mathbf{r}MSE$		19.966	7.162		37.411	12.399			28.332		55.253			9.290	10.462	7.659		8.338	8.286	7.974		10.124	41.854	35.418		55.521	116.824	55 414
		rBias		-18.706	-0.081		36.869	-0.451			27.447		54.727			0.017	-2.658	-0.011		0.150	67.796	-0.031		-0.476	-4.843	34.536		54.945	163.575	54 858
		rSE		0.967	0.995		0.978	0.947			0.961		0.949			1.044	0.987	0.973		1.017	0.093	0.942		1.057	0.222	0.988		1.153	0.231	0.978
		lCI		7.683	10.454		13.875	13.283			11.529		10.533			12.998	14.712	11.765		11.661	26.690	11.341		17.944	30.616	13.911		13.478	32.503	11 295
	EXP	crCI		0.000	95.833		0.000	92.593			76.852		0.000			96.759	94.907	94.444		95.833	58.333	94.444		94.907	88.426	12.500		0.926	0.000	0 000
		\mathbf{rMSE}		12.274	2.673		45.091	3.570			4.758		15.131			3.192	3.837	3.079		2.943	18.153	3.065		4.919	17.395	11.832		14.585	104.426	14 8 14
		rBias		-12.107	0.015		44.946	-0.070			3.650		14.865			0.134	0.019	0.098		-0.015	51.800	-0.042		2.304	8.852	11.277		14.274	143.469	14.519
		rSE		0.974	0.971		0.992	0.782			0.469		0.942			1.037	1.063	0.871		1.036	0.085	0.912		0.939	0.187	0.420		1.139	0.224	0.978
		ICI		11.959	15.257		21.935	27.339			56.091		14.584			19.240	21.564	24.196		17.596	36.575	16.579		24.260	38.325	116.080		18.473	35.981	15.597
	CUB	crCI		0.000	94.444		0.000	91.667			66.667		0.000			95.370	95.370	94.444		95.833	59.722	93.981		86.111	87.963	60.185		0.000	0.000	0 0 0 0
		rMSE		17.440	3.998		74.114	8.917			30.594		24.077			4.755	5.204	7.074		4.357	4.802	4.629		8.660	16.870	71.384		23.478	112.649	$23 \ 448$
		rBias		-17.158	0.016		73.900	-0.515			-3.022		23.753			0.154	0.034	-0.103		-0.061	77.847	-0.024		5.574	10.563	-11.650		23.106	173.883	23 094
		rSE		0.985	0.994		0.929	0.955			1.042		0.943			1.035	1.049	1.027		1.060	0.078	0.920		1.097	0.157	0.943		1.197	0.226	0.957
		lCI		17.587	21.395		20.820	40.191			37.207		20.075		rue	27.157	32.102	28.099	alse	27.893	44.391	25.904	rue	30.532	46.601	54.945	alse	25.043	35.914	20.474
	LIN	crCI		0.000	93.519	S_A)	0.000	92.130		0	95.833	e	0.000		e, PM-T	94.444	95.833	95.370	e, PM-Fa	96.759	62.963	91.667	e, PM-T	94.444	74.074	92.593	e, PM-F	0.000	0.000	0.000
		$\mathbf{r}MSE$	e (S _R)	24.297	5.477	ample (2	100.399	10.721	p	QR-True	9.134	QR-Fals	52.495	$_{\mathrm{thods}}$	QR-True	6.749	7.864	6.969	QR-True	6.819	6.510	7.184	QR-Fals	7.386	20.713	14.943	QR-Fals	51.753	108.237	51 0.43
		rBias	y sample	-23.868	0.118	ubility se	100.237	-0.428	it metho	fication:	0.893	fication:	52.215	bust me	fication:	0.319	0.090	0.180	fication:	0.876	103.450	0.490	fication:	1.843	8.759	1.835	fication:	51.474	191.826	51 656
		Measure _	Probability	UW.	FW	Non-proba	ΜŊ	FW	Non-robus	Model speci	PAPP	Model speci	PAPP ,	Doubly rol	Model speci	GPPP	LWP	AIPW	Model speci	GPPP	LWP	AIPW	Model speci	GPPP	LWP	AIPW	Model speci	GPPP	LWP	ATPW

NOTE: The PAPP and AIPW methods have been implemented through a bootstrap method.

9

UW:	AIPW	LWP	GPPI	Model s	AIPW	LWP	GPPI	Model s	AIPW	LWP	GPPI	Model s	AIPW	LWP	GPPI	Model s	Doubly	PAPF	Model s	PAPF	Model s	Non-ro	FW	UW	Non-pr	FW	UW	Probab	Measur		
Unweigh	38.32	38.289	38.289	pecificatio	/ 0.295	0.089	0.623	pecificatio	/ 0.216	-0.193	-0.286	pecificatio	/ -0.028	-0.243	-0.345	pecificatio	robust	38.049	pecificatio	-0.038	pecificatio	bust me	-0.043	49.180	obability	-0.042	-24.16	ility san	e rBias		
ited; FW	1 38.770	38.70:	38.70	on: QR–F	6.609	4.790	4.813	on: QR–F	4.997	4.850	4.831	on: QR-T	4.938	4.852	4.843	on: QR–T	methods	38.53	on: QR–F	3 5.120	on: QR–T	thod	6.460	49.55	√ sample	4.761	0 24.480	$ple (S_R)$	rMSE		
: Fully v	0.000	3 0.000	3 0.000	alse, PM	95.370	98.611	98.148	alse, PM	97.685	98.611	99.074	'rue, PM-	97.685	98.611	98.611	'rue, PM-	92	3 0.000	alse	96.759	rue		93.056	3 0.000	(S_A)	94.907	0.000		crCI	LIN	
weighted	22.934	29.233	29.233	–False	28.014	24.379	24.745	-True	21.435	23.389	23.492	-False	21.216	23.554	23.428	-True		23.509		21.921			24.387	22.648		17.832	14.690		ICI		
; GPPP:	0.992	1.333	1.333		1.080	1.310	1.331		1.093	1.241	1.249		1.094	1.249	1.244			0.983		1.090			0.961	0.950		0.953	0.948		rSE		
Gaussia	25.373	26.017	26.017		-2.126	1.007	2.439		0.137	0.968	0.996		0.248	0.511	0.454			24.934		3.926			0.091	37.359		-0.400	-23.859		rBias		
un Proces	26.267	26.829	26.829		22.250	6.884	7.249		6.371	6.336	6.370		6.688	6.541	6.387			25.843		11.150			6.891	38.019		4.855	24.179		rMSE		
ss of Pro	4.167	6.481	6.481		85.648	97.222	97.685		94.907	97.685	98.148		93.981	97.685	98.148			3.704		87.037			94.444	0.000		96.759	0.000		crCI	CUB	
opensity	26.143	32.772	32.772		70.680	32.125	32.279		25.029	29.999	30.231		25.328	30.906	30.398			25.873		39.172			27.157	25.871		19.943	16.035		lCI		
Predicti	0.979	1.284	1.284		0.812	1.213	1.215		1.000	1.231	1.234		0.964	1.218	1.229			0.969		0.955			1.003	0.934		1.049	1.040		rSE		
ion; LW	18.557	18.980	18.980		11.808	1.128	1.942		0.772	1.480	1.659		0.792	0.764	0.770			18.198		6.575			0.532	27.292		-0.706	-20.146		rBias		
P: Linea	19.503	19.864	19.864		13.407	6.633	6.670		6.335	6.150	6.255		6.236	6.407	6.202			19.176		9.060			6.855	27.979		4.262	20.452		rMSE		
r-in-weig	11.111	24.074	24.074		53.241	99.074	98.611		94.444	98.148	98.611		94.444	99.074	98.611			14.352		81.944			94.907	0.000		94.907	0.000		crCI	EXP	
ght Pred	22.737	28.967	28.967		24.334	30.109	29.934		23.511	28.234	28.479		23.674	29.130	28.665			22.897		24.125			25.353	22.450		17.407	14.709		ICI		
liction; /	0.964	1.270	1.270		0.975	1.179	1.205		0.952	1.213	1.210		0.974	1.172	1.196			0.964		0.985			0.944	0.928		1.054	1.061		rSE		
AIPW: /	22.449	22.805	22.805		9.860	0.225	0.517		-0.115	0.543	0.485		-0.032	0.327	0.345			22.508		6.183			-0.300	19.208		0.169	-11.189		rBias		
Augment	23.014	23.317	23.317		10.779	5.787	4.457		4.158	4.156	4.095		4.149	4.356	4.177			23.055		7.285			5.720	19.788		3.866	11.695		rMSE		
ed Inver	0.463	2.315	2.315		42.130	96.759	96.296		93.519	92.593	92.130		94.444	95.833	94.907			0.000		71.759			94.444	1.852		94.444	10.185		crCI	SIN	
'se Prop	20.115	25.525	25.525		17.942	24.201	18.057		16.704	16.206	16.104		16.568	17.307	16.881			20.223		16.975			22.909	19.537		15.066	13.552		lCI		
ensity	1.01	1.345	1.345		1.048	1.075	1.045		1.023	1.007	1.016		1.016	1.024	1.044			1.031		1.121			1.021	1.046		0.993	1.013		rSE		

Weighting. NOTE: The PAPP and AIPW methods have been implemented through a bootstrap method.

Table 10: Comparing the performance of the bias adjustment methods in the second simulation study for the binary outcome with $(n_A, n_R) = (500, 1, 000)$ and $\gamma_1 = 0.3$

10

/Robust and Efficient Bayesian Inference for Non-Probability Samples

Table 11: Comparing the performance of the bias adjustment methods in the second simulation study for the *binary* outcome with $(n_A, n_R) = (1, 000, 500)$ and $\gamma_1 = 0.3$

Ro	bus	t a	nd	Ę	ſſ	icier	rt.	₿а	<u>y</u> e	si	$\frac{n}{2}$	Ir	ıfe	re	nc	e f	or	Λ	I oi	<i>1</i> − <i>I</i>	Pro	bo	ıbi	lit	y,	Şa	m_j	ple.	f Þ
	rSE		1.032	1.099		1.176 1.149			1.061		1.175			1.072	0.936	1.077		1.087	1.053	1.085		1.075	1.43	1.08		2.019	2.019	1.141	pensit
	ICI		19.157	21.354		13.818 16.151			18.605		14.329			19.836	20.030	19.630		19.554	19.504	19.795		20.459	26.385	19.173		27.540	27.540	14.343	rse Proi
SIN	crCI		35.648	96.759		0.000 96.759			78.241		0.000			96.296	94.907	95.370		94.907	95.833	95.833		95.833	99.074	50.000		0.463	0.463	0.000	ted Inve
	rMSE		12.367	4.947		19.164 3.624			7.462		22.420			4.796	5.585	4.639		4.710	4.850	4.651		5.008	4.765	10.465		23.046	23.046	22.284	Auoment
	rBias		-11.430	-0.014		18.929 - 0.574			5.980		22.204			0.701	0.952	-0.109		0.915	1.008	-0.091		1.026	0.445	9.440		22.778	22.778	22.053	ATPW:
	rSE		1.015	1.041		0.970 0.965			0.987		0.981			1.539	1.517	1.016		1.560	1.596	1.020		1.706	1.685	1.003		1.886	1.886	1.001	iction:
	ICI		20.818	24.676		15.876 17.877			19.074		17.047			30.631	30.879	20.331		30.723	30.677	20.202		31.759	31.708	20.331		31.709	31.709	17.500	sht Pred
EXP	crCI		4.167	96.759		$0.000 \\ 94.444$			91.667		1.852			99.074	99.537	95.833		99.537	100.000	95.370		100.000	100.000	53.241		22.222	22.222	2.315	r-in-weid
	rMSE		20.549	6.035		$27.149 \\ 4.728$			5.794		18.407			5.279	5.421	5.128		5.363	5.205	5.074		5.370	5.186	11.185		19.977	19.977	18.940	P. Lines
	rBias		-19.875	-0.219		26.828 0.380			3.062		17.868			1.324	1.427	0.594		1.777	1.679	0.565		2.438	1.868	9.924		19.504	19.504	18.409	I.W.
	rSE		1.003	1.026		$0.932 \\ 1.007$			0.896		0.999			1.509	1.501	1.004		1.505	1.520	1.022		1.578	1.559	0.825		1.863	1.863	1.005	Predicti
	ICI		22.713	28.280		18.292 19.145			42.145		19.860			33.701	33.705	23.158		33.405	33.435	22.900		34.562	34.399	90.189		35.974	35.974	20.612	nensity
CUB	crCI		0.926	97.222		0.000 94.444			95.370		0.000			99.074	99.074	93.981		99.537	99.537	95.370		100.000	100.000	94.444		3.704	3.704	0.000	s of Pro
	rMSE		24.144	7.020		37.324 4.840			12.151		25.263			5.831	5.860	5.875		5.859	5.811	5.703		6.303	6.031	30.760		27.438	27.438	26.004	n Proces
	rBias		-23.445	0.153		36.988 -0.060			-2.054		24.751			1.049	1.057	0.199		1.361	1.348	0.163		2.832	2.032	-13.115		26.986	26.986	25.475	Gaussia
	rSE		1.014	1.038		0.970			1.049		0.979			1.330	1.324	1.050		1.313	1.311	1.059		1.472	1.466	1.014		1.813	1.813	0.995	GPPP.
	ICI		20.785	25.283		16.025 17.199			20.326		18.098		rue	26.705	26.714	22.128	alse	26.629	26.602	21.900	Drue	27.464	27.407	24.931	alse	31.845	31.845	18.348	eichted.
LIN	crCI		0.926	95.833	(S_A)	0.000 94.907		le	95.370	se	0.000		ie, PM-T	98.148	98.148	96.759	ie, PM–F	98.611	98.148	95.833	se, PM-7	99.537	100.000	93.519	se, PM–I	0.000	0.000	0.000	Eully w
	rMSE	le (S_R)	24.606	6.211	sample (48.836 4.474	po	: QR-Tri	4.935	: QR-Fal	37.873	ethods	: QR-Tri	5.172	5.178	5.365	: QR-Tru	5.215	5.209	5.271	: QR-Fal	4.852	4.836	6.290	: QR-Fal	38.545	38.545	38.302	d. FW.
	rBias	ty samp	-24.046	0.329	ability s	48.655 -0.155	ist meth	cification	0.236	cification	37.580	obust m	cification	-0.350	-0.260	0.121	cification	-0.304	-0.221	0.251	cification	0.703	0.519	0.619	cification	38.279	38.279	38.013	nweighte
	Ieasure	robabili	UW MU	FW	Ion-prob	UW FW	Ion-robu	fodel spe	PAPP	fodel spe	PAPP	Joubly r	fodel spe	GPPP	LWP	AIPW	fodel spe	GPPP	LWP	AIPW	fodel spe	GPPP	LWP	AIPW	fodel spe	GPPP	LWP	AIPW	11 · /M11
	2	д			Z		Z	PZ		2		р	Z				Z				2				PZ				

Weighting. NOTE: The PAPP and AIPW methods have been implemented through a bootstrap method.

	UW	AIP	GPP	Model	AIP	LWF	GPP	Model	AIP	LWF	GPP	Model	AIP	LWF	GPP	Model	Doubl	PAP	Model	PAP	Model	Non-r	FW	UW	Non-p	FW	UW	Proba	Measu		
ghting. FE: The	: Unwei	W 38.	Р 38.0	specifica	W 0.9;	0.5	P 1.0	specifica	W 0.50	0.2	P 0.1:	specifica	W 0.3:	0.1	P 0.0	specifica	y robus	P 38.	specifica	P 0.4	specifica	obust n	-0.0	49	robabil	0.3	-24	bility s	re rBi		
PAPP	ghted;	532 39	301 30 301 30	ution: Q	39 7.	51 5.	36 5.	ution: Q)9 5.	55	33 5.	ution: Q	29 5.	19 5.	 41 5. 	ution: Q	st meth	110 38	ution: Q	18 6.	ution: Q	nethod	143 6.	180 49	ity san	29 6.	.046 24	ample	as rN		
and /	FW:	9.009	0.056	R-Fals	448	573	654	R-Fals	932	782	778	R-Tru	941	848	778	R-Tru	ods	3.636	R-Fals	024	R-Tru		460	.553	ple (,	211	1.606	(S_R)	ASE		
AIPW	Fully v	0.463 0.000	0.463	зе, PM-	97.222	99.537	99.537	e, PM−	96.296	98.148	98.611	e, PM–	94.444	97.685	98.148	e, PM-		0.000	3e	93.981	e		93.056	0.000	$S_A)$	95.833	0.926		crCI	LIN	
methods	veighted	$\frac{33.223}{24.171}$	35.225	-False	30.643	30.016	30.395	-True	25.084	28.810	29.108	False	25.256	29.079	29.040	True		24.026		24.516			24.387	22.648		25.283	20.785		lCI		
s have b	; GPPP	$1.324 \\ 1.011$	1.524		1.056	1.392	1.408		1.080	1.285	1.294		1.084	1.279	1.290			0.963		1.039			0.961	0.950		1.038	1.014		rSE		
een imple	: Gaussia	20.372 25.585	26.372		-6.206	1.474	2.959		0.439	1.365	1.387		0.566	0.923	0.907			24.997		1.301			0.091	37.359		0.153	-23.445		rBias		
emented	an Proce	27.203 26.540	27.263		26.540	7.532	7.837		7.218	7.102	7.130		7.494	7.407	7.182			25.965		12.124			6.891	38.019		7.020	24.144		rMSE		
through	ess of P	10.007 4.630	16.667		92.593	99.537	99.537		93.056	98.611	99.537		93.056	98.148	98.611			6.944		93.981			94.444	0.000		97.222	0.926		crCI	CUB	
a boots	opensity	39.480 26.995	39.480		84.822	38.708	38.926		27.731	36.637	36.942		28.540	37.189	37.212			26.728		44.579			27.157	25.871		28.280	22.713		lCI		
trap me	/ Predic	$\begin{array}{c} 1.400\\ 0.974 \end{array}$	1.465		0.837	1.346	1.378		0.980	1.356	1.358		0.972	1.304	1.342			0.968		0.941			1.003	0.934		1.026	1.003		rSE		
thod.	tion; LW	19.248 18.697	19.248		10.963	1.462	2.388		0.910	1.755	1.909		0.795	1.167	1.092			18.257		4.807			0.532	27.292		-0.219	-19.875		rBias		
	/P: Line	$\frac{20.128}{19.645}$	20.128		12.774	6.800	6.928		6.627	6.472	6.569		6.444	6.632	6.455			19.247		8.095			6.855	27.979		6.035	20.549		rMSE		
	ar-in-we	$\frac{30.374}{11.111}$	36.574		60.648	100.000	100.000		95.370	100.000	99.537		94.444	100.000	99.537			15.741		88.426			94.907	0.000		96.759	4.167		crCI	EXP	
	ight Pre	$\frac{53.137}{23.391}$	35.137		26.141	35.656	35.583		25.364	33.880	34.123		25.518	34.385	34.268			23.253		25.182			25.353	22.450		24.676	20.818		lCI		
	diction;	1.333 0.987	1.535		1.015	1.377	1.406		0.983	1.396	1.395		1.016	1.356	1.382			0.971		0.984			0.944	0.928		1.041	1.015		rSE		
	AIPW:	22.752 22.342	22.752		9.697	0.175	0.495		-0.118	0.512	0.740		-0.091	0.372	0.340			22.514		6.097			-0.300	19.208		-0.014	-11.430		rBias		
	Augmen	$\frac{23.322}{22.938}$	23.322		10.937	6.160	5.498		5.034	4.994	6.331		5.004	5.302	5.187			23.073		7.752			5.720	19.788		4.947	12.367		rMSE		
	ted Inve	$\frac{8.333}{0.926}$	8.333		57.407	98.611	95.370		95.370	94.907	95.370		95.833	97.685	95.833			0.000		78.241			94.444	1.852		96.759	35.648		crCI	NIS	
	rse Prop	$\frac{30.490}{20.299}$	30.490		21.505	29.437	22.425		21.062	20.754	20.815		21.217	21.684	21.357			20.041		20.696			22.909	19.537		21.354	19.157		lCI		
	pensity	0.995	1.528		1.082	1.226	1.057		1.065	1.075	0.85		1.079	1.054	1.06			1.01		1.1			1.021	1.046		1.099	1.032		rSE		

Table 12: Comparing the performance of the bias adjustment methods in the second simulation study for the binary outcome with $(n_A, n_R) = (500, 500)$ and $\gamma_1 = 0.3$

Table 13: Comparing the performance of the bias adjustment methods in the second simulation study for the *continuous* outcome with $(n_A, n_R) = (500, 1, 000)$ and $\gamma_1 = 0.6$

õ	
$\vec{2}$	
and	

	rSE		1.013	0.993		1.069	1.013			1.033		1.02			1.069	0.879	1.043		1.061	1.033	1.039		0.941	0.731	1.027		1.24	1.24	1.015	pensity
	lCI		13.552	15.066		19.491	37.557			17.300		21.652			21.096	23.998	17.162		17.033	17.391	17.622		23.803	29.606	21.309		25.637	25.637	22.047	rse Proj
NIS	crCI		10.185	94.444		0.000	94.444			6.944		0.000			95.370	91.667	94.444		93.981	93.519	93.981		92.593	83.796	0.000		0.000	0.000	0.000	ed Inve
	rMSE		11.695	3.866		22.962	9.445			15.234		41.367			5.066	7.087	4.192		4.147	4.355	4.317		6.565	10.385	27.089		41.104	41.104	41.769	Augment
	rBias		-11.189	0.169		22.488	-0.444			14.626		41.013			-0.179	-0.367	0.211		0.338	0.526	0.051		-0.932	-0.156	26.569		40.756	40.756	41.401	AIPW: A
	rSE		1.061	1.054		0.945	0.909			0.979		0.969			1.087	0.803	0.976		1.183	0.024	0.988		0.986	0.871	0.938		1.171	1.171	0.945	liction; /
	ICI		14.709	17.407		21.829	40.607			29.468		26.075			37.261	39.277	29.438		35.445		28.708		39.155	38.376	28.560		31.078	31.078	26.079	ght Pred
EXP	crCI		0.000	94.907		0.000	89.352			66.667		1.852			93.056	89.815	92.130		96.759	96.759	94.444		93.056	90.278	26.389		7.407	7.407	3.704	r-in-weig
	$\mathbf{r}MSE$		20.452	4.262		58.307	11.375			14.469		28.525			8.926	12.671	7.700		7.965		7.440		10.529	11.326	20.900		25.879	25.879	26.799	^o : Linea
	rBias		-20.146	-0.706		58.010	-0.125			12.275		27.691			-1.181	0.601	0.596		2.018		0.783		2.203	0.000	19.411		24.967	24.967	25.863	on; LWF
	rSE		1.040	1.049		0.937	0.976			0.737		0.982			1.155	1.011	0.981		1.230	0.584	1.018		1.105	1.032	0.571		1.163	1.163	0.956	Predicti
	ICI		16.035	19.943		25.341	41.622			50.263		29.931			37.554	39.769	29.165		36.271	33.546	29.316		43.539	41.871	81.350		34.849	34.849	29.653	pensity
CUB	crCI		0.000	96.759		0.000	93.981			56.019		0.000			96.296	93.981	95.370		98.611	98.148	94.444		96.296	93.519	53.241		2.778	2.778	0.463	s of Pro
	rMSE		24.179	4.855		79.811	10.904			22.143		38.028			8.776	10.318	7.571		7.570	14.769	7.330		10.786	10.418	37.241		33.276	33.276	34.796	Process
	rBias		-23.859	-0.400		79.514	-1.068			13.742		37.229			-2.760	-2.168	-0.311		0.141	0.028	0.007		3.623	-0.705	8.524		32.374	32.374	33.889	Gaussian
	rSE		0.948	0.953		0.982	0.956			1.060		1.012			1.201	0.941	1.054		0.802	0.098	1.074		1.257	1.160	1.044		1.263	1.263	1.047	GPPP:
	ICI		14.690	17.832		20.067	37.625			33.477		26.975		rue	27.586	30.108	25.367	alse	28.818	47.857	25.426	rue	31.009	27.687	50.214	alse	31.071	31.071	26.339	ighted;
LIN	crCI		0.000	94.907	$S_A)$	0.000	91.667		e	95.833	se	0.000		e, PM-T	96.296	94.907	95.833	e, PM–F	96.759	95.833	95.370	se, PM–T	97.685	96.759	94.907	se, PM–F	0.000	0.000	0.000	Fully we
	rMSE	$e(S_R)$	24.480	4.761	ample (,	93.380	10.053	pq	QR-Tru	8.039	QR-Fals	60.092	ethods	QR-Tru	6.355	8.425	6.143	QR-Tru	9.372	133.689	6.036	QR-Fals	6.315	6.534	12.269	QR-Fals	53.677	53.677	56.049	d; FW:
	rBias	y sampl	-24.160	-0.042	ability s	93.235	-0.792	st metho	ification:	-0.227	ification:	59.708	bust me	ification:	-2.388	-1.908	-0.429	ification:	-1.367	-12.060	0.315	ification:	-0.218	-2.276	0.857	ification:	53.302	53.302	55.682	weighted
	Measure ⁻	Probabilit	UW	FW	Non-prob	NN	FW	Non-robus	Model speci	PAPP	Model speci	PAPP	Doubly ro	Model speci	GPPP	LWP	AIPW	Model speci	GPPP	LWP	AIPW	Model speci	GPPP	LWP	AIPW	Model speci	GPPP	LWP	AIPW	UW: Un

Weighting. NOTE: The PAPP and AIPW methods have been implemented through a bootstrap method.

AIPW	LWP	GPPP	Model speci	AIPW	LWP .	GPPP	Model speci	AIPW	LWP	GPPP	Model speci	AIPW	LWP	GPPP .	Model speci	Doubly ro	PAPP	Model speci	PAPP	Model speci	Non-robus	FW	$_{ m UW}$	Non-proba	FW	UW	Probability	Measure		
55.890	51.913	51.913	fication:	1.373	-2.976	-1.790	fication:	0.392	-4.364	-3.729	fication:	-0.017	-3.647	-3.903	fication:	bust m	58.629	fication:	0.544	ification:	st meth	-0.489	92.334	ability s	0.329	-24.046	y samp	rBias		CIICI / 1
56.136	52.162	52.162	QR-Fa	9.986	6.166	5.718	QR-Fa	5.823	7.228	6.809	QR-Tr	5.864	6.913	6.864	QR-Tr	ethods	58.883	: QR–Fa	6.512	QR-Tr	od	7.243	92.413	ample	6.211	24.606	le (S_R)	rMSE		0.0
0.000	0.000	0.000	lse, PM-	94.444	99.074	100.000	lse, PM-	94.907	94.444	96.759	ue, PM–I	94.444	96.296	95.833	ue, PM-7		0.000	lse	94.907	ue		92.593	0.000	(S_A)	95.833	0.926		crCI	LIN	
20.741	32.167	32.167	False	39.167	29.028	30.256	True	24.088	28.283	28.835	alse	24.464	29.888	28.565	Frue		20.647		26.660			26.827	14.252		25.283	20.785		ICI		
1.006	1.621	1.621		1.008	1.380	1.428		1.055	1.261	1.297		1.062	1.304	1.295			0.962		1.046			0.945	0.948		1.038	1.014		rSE		
34.495	32.802	32.802		-13.701	0.070	3.429		0.428	-1.685	-0.979		0.362	-2.105	-2.421			36.297		-0.222			0.076	78.359		0.153	-23.445		rBias		
34.980	33.285	33.285		54.515	7.947	8.509		6.352	6.469	6.280		6.715	7.801	7.354			36.745		21.896			7.212	78.506		7.020	24.144		rMSE		
0.000	0.926	0.926		73.611	96.296	98.611		95.370	99.074	99.074		95.370	97.222	98.611			0.000		81.019			93.519	0.000		97.222	0.926		crCI	CUB	
23.541	36.718	36.718		128.401	39.401	40.538		25.181	35.185	36.689		26.837	38.341	37.021			23.009		62.841			29.956	17.952		28.280	22.713		lCI		
1.033	1.668	1.668		0.619	1.271	1.339		1.011	1.447	1.521		1.019	1.312	1.372			1.025		0.731			1.057	0.951		1.026	1.003		rSE		
25.807	25.110	25.110		17.160		1.870		0.772	-0.398	0.846		0.738	-0.166	-0.993			26.921		6.897			0.340	56.892		-0.219	-19.875		rBias		
26.291	25.585	25.585		18.219		7.991		6.207	5.941	6.314		6.379	8.573	7.089			27.371		9.340			7.973	57.036		6.035	20.549		rMSE		
0.000	3.241	3.241		19.444	98.148	99.074		96.296	99.074	99.537		96.759	97.685	99.537			0.000		76.852			93.519	0.000		96.759	4.167		crCI	EXP	
20.174	32.362	32.362		23.805		36.386		23.403	32.773	34.586		24.283	36.597	35.230			19.501		23.256			29.311	15.470		24.676	20.818		ICI		
1.023	1.697	1.697		0.989	0.074	1.207		0.967	1.415	1.425		0.975	1.097	1.293			1.005		0.940			0.936	0.973		1.041	1.015		rSE		
41.565	40.375	40.375		26.793	0.056	-0.683		0.005	-128.960	0.275		0.158	0.242	-0.285			41.494		14.493			-1.071	22.907		-0.014	-11.430		rBias		
41.743	40.592	40.592		27.156	9.039	5.880		4.710	1104.35	4.658		4.599	8.279	5.107			41.660		15.104			6.592	23.094		4.947	12.367		rMSE		
0.000	0.000	0.000		0.000	89.35	94.44		95.83	53 92.59	96.29		96.75	93.05	96.75			0.000		11.57			96.29	0.000		96.75	35.64		crCI	NIS	
16.317	26.482	26.482		19.895	2 28.784	4 23.425		3 20.270	3 307.22	6 19.899		9 19.998	6 23.532	9 21.765			15.831		4 18.44C			6 26.831	13.777		9 21.354	8 19.157		lCI		
1.077	1.623	1.623		1.145	0.816	1.029		1.095	5 0.071	1.1		1.107	0.734	1.097			1.084		1.104			1.05	1.195		1.099	1.032		rSE		

Table 14: Comparing the performance of the bias adjustment methods in the second simulation study for the binary outcome with $(n_A, n_R) = (1, 000, 500)$ and $\gamma_1 = 0.6$

UW: Unweighted; FW: Fully weighted; GPPP: Gaussian Process of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting. NOTE: The PAPP and AIPW methods have been implemented through a bootstrap method.

Table 15: Comparing the performance of the bias adjustment methods in the second simulation study for the *binary* outcome with $(n_A, n_R) = (500, 500)$ and

	?		
	2		
I			
5	_		

LSE		1.032 1.099		1.069	1.013			1.06	510	1.008		IJе	1.067	0.91	1.094		1.069	0.076	1.063		0.977	0.661	1.063		1.374	1.374	1.003	ensity
ICI		19.157 21.354		19.491	37.557			20.750		21.907			24.740	27.358	21.700		21.606	46.573	21.870		27.402	33.602	23.873		30.304	30.304	22.470	se Prop
SIN crCI	010	35.648 96.759		0.000	94.444			22.222		0.000			95.833	91.204	96.296		96.759	95.370	96.296		94.444	85.648	0.463		0.000	0.000	0.000	ed Inver
rMSE	1000	12.367 4.947		22.962	9.445			15.439		41.593			5.952	7.747	5.051		5.216	162.050	5.236		7.270	13.095	27.090		41.023	41.023	41.875	Augment
rBias	007	-11.430 -0.014		22.488	-0.444			14.580		41.224			-0.203	-0.293	0.128		0.354	-11.862	-0.006		-0.911	1.071	26.480		40.628	40.628	41.485	AIPW: A
SE	h T	1.015 . 1.041 .		0.945	. 609.0			0.981		1.002			1.180 .	. 898	0.985		1.346	1.308	1.004		. 960.1	0.988	0.972		1.366	1.366	0.969	liction; /
ICI		20.818 24.676		21.829 (40.607 (31.136 (26.766			41.719	44.049 (31.509 (39.760	37.732	30.408		43.869	42.761 (30.913 (36.607	36.607	27.008 (ght Pred
EXP crCl	1	4.167 96.759		0.000	89.352			78.704		2.315			94.907	92.593	93.056		99.537	98.611	93.056		97.222	92.593	36.574		13.426	13.426	5.556	tr-in-weig
rMSE		20.549 6.035		58.307	11.375			12.558		28.064			9.124	12.656	8.193		7.892	7.418	7.761		10.628	11.133	19.986		26.197	26.197	26.866	P: Linea
rBias		-19.875 -0.219		58.010	-0.125			9.612		27.229			-0.828	0.746	0.891		2.102	0.531	0.865		2.647	0.254	18.273		25.278	25.278	25.913	ion; LW
rSE	d d d d	1.003 1.026		0.937	0.976			0.746		0.968			1.203	1.157	0.980		1.307	1.274	0.988		1.172	1.143	0.626		1.322	1.322	0.951	Predict
ICI		22.713 28.280		25.341	41.622			57.366		30.541			42.778	45.327	32.175		42.119	39.763	31.529		48.346	46.784	101.955		41.353	41.353	30.859	opensity
CUB crCI	0000	0.926 97.222		0.000	93.981			76.389		0.000			98.148	96.759	93.056		97.685	98.611	94.444		96.759	98.148	65.741		4.630	4.630	1.389	ss of Pr
rMSE		24.144 7.020		79.811	10.904			21.288		37.351			9.423	10.156	8.356		8.277	8.034	8.127		11.390	10.511	41.495		33.628	33.628	35.144	n Proce
rBias	2 F 7 00	-23.445 0.153		79.514	-1.068			8.356		36.477			-2.381	-1.526	-0.122		0.499	-0.384	0.208		4.130	-0.516	2.173		32.656	32.656	34.159	Gaussia
rSE	1	$1.014 \\ 1.038$		0.982	0.956			1.033		0.995			1.253	1.126	1.082		1.279	1.229	1.065		1.338	1.276	1.011		1.413	1.413	1.037	GPPP:
ICI		20.785 25.283		20.067	37.625			34.586		27.586		I rue	32.515	33.629	28.899	false	33.459	31.969	28.536	True	35.644	32.806	51.268	False	36.316	36.316	27.460	reighted;
LIN crCI		0.926 95.833	(S_A)	0.000	91.667		ue	98.148	lse	0.000		ue, PM- ⁷	97.222	96.759	95.833	ue, PM-I	98.148	96.296	97.222	lse, PM ⁻¹	99.537	97.685	95.370	lse, PM–	0.000	0.000	0.000	Fully w
rMSE	le (S_R)	24.606 6.211	sample (93.380	10.053	pod	: QR-Tri	8.528	: QR-Fai	59.464	ethods	: QR-Tri	6.915	7.877	6.802	: QR-Tri	6.874	7.164	6.852	: QR-Fa	6.819	6.836	12.999	: QR-Fai	54.003	54.003	56.209	»d; FW:
rBias	ty samp	-24.046 0.329	ability :	93.235	-0.792	st meth	cification	0.291	cification	59.044	obust m	cification	-1.904	-1.841	-0.147	cification	-1.465	-2.607	0.646	cification	0.154	-1.833	1.505	cification	53.595	53.595	55.803	nweight ϵ
Measure	Probabili	ЪW	Non-prob	ΩM	FW	Non-robu	Model spec	PAPP	Model spe	PAPP	Doubly r	Model spe	GPPP	LWP	AIPW	Model spe	GPPP	LWP	AIPW	Model spe	GPPP	LWP	AIPW	Model spe	GPPP	LWP	AIPW	UW: U

Weighting. NOTE: The PAPP and AIPW methods have been implemented through a bootstrap method.

8.3. Supplemental results on SHRP2/NHTS data application

Adjusted police-reportable crash rates per 100M miles and associated 95% CIs by covariates

Covariate	n	Unweighted (95% CI)	GPPP (95% CI)	LWP (95% CI)	AIPW (95% CI)
Total	2,862	$1430.59\ (1417.66, 1443.52)$	461.29(296,718.88)	534.2(270.47, 1055.11)	464.58(294.06,734)
Gender					
Male	1,357	1778.61 (1740.38, 1816.84)	457.56(293.54,713.22)	543.75(270.88,1091.48)	464.59(293.9,734.42)
Female	1,505	1116.79(1107.96, 1125.62)	465.4(298.14,726.49)	524.11(268.95, 1021.35)	464.45 (293.01,736.18)
Age group	,				
16-19	453	2621.22(2565.47.2676.97)	1532.82(976.35.2406.45)	1468.28 (836.11.2578.44)	1535.73 (970.36.2430.52)
20-24	671	$1357\ 97\ (1334\ 13\ 1381\ 82)$	860 25 (533 43 1387 32)	829 97 (383 85 1794 61)	879 91 (534 14 1449 52)
25-29	254	$1058\ 64\ (1017\ 15\ 1100\ 13)$	788 27 (508 57 1221 8)	854 11 (488 08 1494 64)	794 45 (499 52 1263 49)
30-39	237	331 76 (313 61 349 9)	261 38 (168 1 406 41)	322.84(160.73.648.47)	265(1686141648)
40-49	214	290.26(273.98.306.54)	$358\ 91\ (225\ 32\ 571\ 7)$	<i>AAA</i> 7 (222 7 887 98)	361 39 (220 82 591 43)
50 50	214	4294.60(2815.74.4822.64)	200.07(256.67.620.47)	500.72 (260.20.007.82)	402.82 (252.22,601.43)
60.60	200	524.09(5015.74,4055.04)	561.02(259.07,020.47)	509.75(200.39,997.02)	544.86(242.45.864.20)
70.70	210	329.89 (309.18, 350.0)	406 44 (364 44 634 60)	405 47 (210 40 781 06)	417.80(343.45,804.59)
10-19	340 177	450.48 (455.74,407.25)	400.44 (204.44,024.09) 1020 05 (750 04 0045 60)	405.47 (210.49,781.00)	417.69(272.45,040.90) 1948 19(796 07 9119 91)
-00+	111	1514.88 (1450.84,1598.91)	1238.85 (750.24,2045.08)	1204.41 (045.22,2246.24)	1248.12 (750.97,2115.81)
Race					
White	2,530	1461.22(1445.75, 1476.7)	440.54 (281.63,689.11)	502.06(252.37,998.76)	446.8 (282.12,707.59)
Black	150	910.16 (860.84, 959.49)	521.84(334.41,814.31)	683.96(342.54, 1365.71)	511.11(323.81,806.74)
Asian	96	2197.74(2017.8, 2377.68)	521.55 (330.77, 822.36)	560.17 (311.8, 1006.42)	513.48 (313.78, 840.28)
Other	86	580.72(517.11,644.34)	632.26 (420.17, 951.43)	810.56 (443.05, 1482.92)	$634.01 \ (403.55, 996.07)$
Gender					
Non-Hisp	2,754	1442.75 (1429.07, 1456.44)	434.32(277.78,679.07)	490.06(247.28,971.23)	434.68(274.25,688.97)
Hispanic	108	1120.45 (1053.82, 1187.08)	684.24 (472.5,990.85)	1023.74 (563.45, 1860.07)	$716.38 \ (471.67, 1088.05)$
Ethnicity					
<high school<="" td=""><td>213</td><td>3659.76(3497.14,3822.38)</td><td>1169.05(730.81, 1870.09)</td><td>1329.65 (646.48, 2734.8)</td><td>1158.09(703.44, 1906.6)</td></high>	213	3659.76(3497.14,3822.38)	1169.05(730.81, 1870.09)	1329.65 (646.48, 2734.8)	1158.09(703.44, 1906.6)
HS comp	279	1606.27 (1554.61, 1657.93)	478.5 (304.26,752.52)	692.02 (356.97,1341.54)	472.07 (295.39,754.42)
College	837	1248.89 (1231.23,1266.54)	473.56 (304.61,736.21)	561.22(272.75,1154.81)	483.62 (303.5,770.65)
Graduate	1.068	603.63 (597.39.609.87)	370.9(238.69.576.34)	385.55(198.1.750.37)	374.99(238.43.589.77)
Post-grad	465	2530.49 (2347.36.2713.63)	475.47(302.42.747.56)	509.52(261.3.993.53)	473.87(299.47.749.84)
HH income		(,,,,,,,,		((
0-49	1 164	$1179\ 27\ (1167\ 12\ 1191\ 42)$	499 59 (315 31 791 56)	$594\ 45\ (288\ 18\ 1226\ 2)$	497 9 (308 84 802 71)
150-99	1 049	709.89 (702.91.716.88)	375 32 (243 18 579 25)	$421\ 01\ (215\ 49\ 822\ 54)$	$376\ 41\ (242\ 88\ 583\ 35)$
100-149	1/12	1658 96 (1605 49 1712 43)	442.6 (286.32.684.19)	506.46 (266.98.960.73)	455 88 (289 48 717 9)
150-145	207	6008.42 (5391.28 6625.55)	$676\ 53\ (425\ 78\ 1074\ 96)$	824 7 (417 97 1627 23)	685 64 (410 65 1144 77)
HH sizo	201	0000.12 (0001.20,0020.00)	010.00 (120.10,1011.00)	021.1 (111.01,1021.20)	000.01 (110.00,1111.11)
1	508	1155.7(1198.91.1183.18)	132 75 (273 84 683 87)	468 57 (231 47 048 5)	430 0 (272 26 710 74)
1	067	608.78(600.227,1103.10)	452.75(215.64,005.07) 452.72(202.66,701.06)	516.95(250.7,1026.92)	439.9(212.20,110.14)
2	507	152675(1402.82,101.34)	453.75(295.00,701.00)	510.25 (259.7, 1020.25)	442.13 (203.37,009.34)
3	510	1330.75(1493.83,1579.07)	403.80(293.02,729.33)	540.51(281.55,1000.12)	470.79(293.94,754.04)
4	075	3045.14(2885.96,3204.33)	481.49 (301.8,768.17)	561.53(291.39,1082.1)	480.84(300.07,789.85)
<u>0+</u>	275	1398.78 (1353.14,1444.42)	483.9 (314.29,745.04)	626.53 (314.87,1246.67)	512.19 (323.83,810.11)
Vehicle make	1 0 15				
American	1,045	2058.42 (2003.06,2113.78)	407.92(260.59,638.55)	496.88(246.37,1002.13)	414.98(260.96,659.91)
Asian	1,745	1034.41 (1025.7, 1043.12)	475.22(306.94,735.75)	521.23(268.96,1010.09)	478.85(305.11,751.51)
European	72	1920.11 (1707.15,2133.08)	726.7 (448.99,1176.18)	963.13(476.84,1945.38)	690.15 (409.89, 1162.04)
Vehicle type					
Car	2,061	1736.09(1715.31, 1756.86)	611.19(392.09,952.73)	667.49(342.48, 1300.92)	607.53 (381.97, 966.28)
Van	109	629.14 (581.73, 676.55)	682.49 (439.71, 1059.32)	835.22 (446.81, 1561.3)	754.96 (480.49, 1186.19)
SUV	551	724.36(696.5,752.21)	320.33(203.8,503.48)	376.21 (182.91,773.77)	325.25(208.23,508.02)
Pickup	141	344.53 (311.67, 377.4)	233.9(151.03, 362.22)	336.55 (169.77, 667.17)	238.51 (148.96, 381.9)
Vehicle age					
0-4	320	2821.77(2738.31,2905.24)	511.45(319.47,818.8)	631.43 (316.66, 1259.1)	536.52(324.72,886.47)
5-9	742	$838.31 \ (826.17, 850.46)$	483.2 (313.51, 744.74)	555.03(293.54, 1049.46)	478.57 (305.12, 750.62)
10-14	905	$977.22 \ (968.18, 986.25)$	438.34 (281.64, 682.23)	489.77 (238.5, 1005.77)	442.87 (279.4,701.98)
15-19	382	607.65(592.55,622.76)	412.86 (264.47,644.5)	480.04 (248.96,925.61)	404.66(256.76,637.75)
20-24	197	5119.19 (4543.74,5694.65)	433.61 (279.84,671.86)	478.1 (219.52,1041.29)	436.13 (281.22,676.37)
25-29	108	545.61 (515.12,576.09)	418.01 (257.83,677.71)	469.24 (214.37,1027.12)	429.71 (265.65,695.09)
30+	178	2030.45 (1897.07,2163.84)	466.84 (277.69,784.84)	580.56 (283.85,1187.42)	486.75 (288.21,822.08)
Fuel type		× / /			
Gas/Diesel	2,641	1526.32 (1511.74,1540.9)	461.79 (296.55,719.13)	535.61 (270.59,1060.18)	465.34 (294.5,735.28)
Other	221	286.58 (273.29,299.87)	439.6 (267.26,723.09)	476.92 (262.85,865.33)	432.82 (262.46,713.76)