# A proof of Toponogov's theorem in Alexandrov geometry<sup>[1](#page-0-0)</sup>

Shengqi Hu, Xiaole Su, Yusheng Wang[2](#page-0-1)

Abstract. This paper aims to give an elementary proof for Toponogov's theorem in Alexandrov geometry with lower curvature bound. The idea of the proof comes from the fact that, in Riemannian geometry, sectional curvature can be embodied in the second variation formula.

Key words. Toponogov's theorem, Alexandrov geometry Mathematics Subject Classification (2000): 53-C20.

### 0 Introduction

An Alexandrov space X with curvature  $\geq k$  is roughly defined to be a locally complete intrinsic metric space on which Toponogov's type theorem holds locally ([BGP]). In fact, if  $X$  is complete, then such a type theorem holds globally on  $X$ , which is called Toponogov's theorem in Alexandrov geometry. In its proofs ([BGP], [AKP]), to simplify the arguments, it is always assumed that there exists a minimal geodesic (shortest path) between any two points in  $X<sup>3</sup>$  $X<sup>3</sup>$  $X<sup>3</sup>$ . For any two minimal geodesics  $[pq]$  and  $[qr]$  in X, we can define the angle  $\angle pqr$  naturally; and then, Toponogov's theorem in Alexandrov geometry can be formulated as follows ([BGP]).

**Theorem A** Let X be a complete Alexandrov space with curvature  $\geq k$ . Then for any triangle  $\triangle pqr \subset X$  (a union of three minimal geodesics  $[pq], [pr], [qr]$ ), we have that

$$
\angle pqr \ge \tilde{\angle}_k pqr. \tag{0.1}
$$

In Theorem A,  $\tilde{\angle}_k pqr$  denotes the angle  $\angle \tilde{p}\tilde{q}\tilde{r}$  in the comparison triangle  $\triangle \tilde{p}\tilde{q}\tilde{r} \subset \mathbb{S}^2_k$ of  $\triangle pqr$  (i.e.  $|\tilde{p}\tilde{q}| = |pq|, |\tilde{p}\tilde{r}| = |pr|, |\tilde{q}\tilde{r}| = |qr|$ ), where  $\mathbb{S}^2_k$  denotes the complete and simply connected 2-dimensional space form with constant curvature  $k$ . And in a proof of Theorem A, one need only to consider the case where  $|pq| + |pr| + |qr| < \frac{2\pi}{\sqrt{k}}$  $\frac{\pi}{k}$  (and thus each of  $|pq|, |pr|, |qr|$  is less than  $\frac{\pi}{\sqrt{k}}$  if  $k > 0$  ([BGP])<sup>[4](#page-0-3)</sup>.

So far, there are several proofs for Theorem A ([BGP], [Pl], [Sh], [Wa]), each of which is skillful and the one in [Pl] is quite beautiful. In Riemannian geometry, sectional curvature can be seen just from geodesic variations (the second variation formula). However, all known proofs of Theorem A have no direct relation with such an idea. The main goal of this paper is to provide a proof of such an idea for Theorem A.

<sup>1</sup> Supported by NSFC 11971057 and BNSF Z190003.

<span id="page-0-0"></span><sup>2</sup>The corresponding author (E-mail: wyusheng@bnu.edu.cn).

<span id="page-0-2"></span><span id="page-0-1"></span> ${}^{3}$ If X is both complete and locally compact, there is a shortest path between any two points in it.

<span id="page-0-3"></span><sup>&</sup>lt;sup>4</sup>Once Theorem A has been proven, it can be shown that  $|pq| \leq \frac{\pi}{\sqrt{k}}$  and  $|pq| + |pr| + |qr| \leq \frac{2\pi}{\sqrt{k}}$  for all  $p, q, r \in X$  ([BGP]). Moreover, if  $|pq| = \frac{\pi}{\sqrt{k}}$ , it is a convention that  $\tilde{\angle}_k pqr = 0$ .

Our strategy is: If  $\angle pqr < \tilde{\angle}_k pqr$  for some  $\triangle pqr \subset X$ , we can split  $\triangle pqr$  into two triangles along some  $[ps]$  with  $s \in [qr]^\circ$  (the interior part of  $[qr]$ ) so that  $\angle psq < \tilde{\angle}_kpsq$ or  $\angle psr < \tilde{\angle}_kpsr$ . By repeating this process, we can locate an  $o \in [qr]$  such that there is information of "curvature  $\geq k$ " around some [po], i.e. there is  $\triangle pr_1r_2$  with  $r_i$  sufficiently close to *o* such that  $\angle pr_1r_2 < \tilde{\angle}_k pr_1r_2$ . Note that Theorem A holds on a small ball  $B(o, \delta)$  (i.e. (0.1) is true for triangles in  $B(o, \delta)$ ) because X is of curvature  $\geq k$ . Then via Alexandrov's Lemma (Lemma 1.3 below) on  $\triangle pr_1r_2$ , we can locate another  $o' \in B(o, \delta)$  with  $|po'| \leq |po| - \frac{\delta}{3}$  such that there is still information of "curvature  $\geq k$ " around some  $[p\sigma]$ . Step by step, such information can be transmitted to a small neighbourhood of  $p$ , a contradiction (because Theorem A holds around  $p$ ).

In the rest of the paper, X always denotes the space in Theorem A.

#### 1 Tools of the proof

Note that a proof of Theorem A has to depend only on its local version, where we have the following basic property on angles ([BGP]).

**Lemma 1.1** Let  $[pq], [rr'] \subset X$  with  $r \in [pq]$ <sup>o</sup>. Then  $\angle prr' + \angle qrr' = \pi$ .

Furthermore, we have the following easy observation (cf. [AKP], [Wa]).

**Lemma 1.2** Let  $[pq], [qr] \subset X$ . Then for  $q_i \in [qr]$  with  $q_i \to q$  as  $i \to \infty$ ,

$$
|pq_i| \le |pq| - \cos \angle pqr \cdot |qq_i| + o(|qq_i|)^{5}.
$$
 (1.1)

*Proof.* Since Theorem A holds around q, there is  $\bar{p} \in [pq]$  near q such that

$$
|\bar{p}q_i| \leq |\bar{p}q| - \cos \angle pqr \cdot |qq_i| + o(|qq_i|).
$$

Then (1.1) follows from that  $|pq_i| \leq |p\bar{p}| + |\bar{p}q_i|$ |.

Besides Lemmas 1.1 and 1.2, we will also use Alexandrov's lemma ([BGP]).

**Lemma 1.3** Let  $\triangle pqr$ ,  $\triangle pqs$ ,  $\triangle abc \subseteq \mathbb{S}^2_k$  (where  $\triangle pqr$  and  $\triangle pqs$  are joined to each other in an exterior way along  $[pq]$ ) such that  $|ab| = |pr|$ ,  $|ac| = |ps|$ ,  $|bc| = |qr| + |qs|$ , and  $|ab| + |ac| + |bc| < \frac{2\pi}{\sqrt{b}}$  $\frac{\pi}{k}$  if  $k > 0$ . Then  $\angle pqr + \angle pqs \leq \pi$  (resp.  $\geq \pi$ ) if and only if ∠prq  $\geq$  ∠abc and ∠psq  $\geq$  ∠acb (resp. ∠prq  $\leq$  ∠abc and ∠psq  $\leq$  ∠acb).

### 2 Proof of Theorem A

Due to the similarity of proofs for  $k < 1, \leq k \leq 0$ , we only consider the case where  $k = 0$ . And for simpleness, we denote by  $\tilde{\angle} pqr$  the angle  $\tilde{\angle}_0 pqr$ .

In the proof, we will argue by contradiction, and say that an angle  $\angle pqr$  is bad if  $\angle pqr < \angle pqr$ . If each angle of a triangle is not bad, we call the triangle a good one.

<span id="page-1-0"></span><sup>&</sup>lt;sup>5</sup>In addition, if X is locally compact, then we can select  $[pq]$  such that (1.1) is an equality (the first variation formula) once Theorem A has been proven ([BGP], [AKP]).

First of all, we have the following observation about a 'bad' angle just via Lemmas 1.1 and 1.2 (cf. [Wa], [SSW]).

**Lemma 2.1** For a triangle  $\triangle pr_1r_2 \subset X$ , if  $\angle pr_1r_2$  is bad, then there is  $s_0 \in [r_1r_2]^\circ$ such that for any  $[ps_0]$ 

$$
\angle ps_0r_1 \text{ or } \angle ps_0r_2 \text{ is bad};\tag{2.1}
$$

in particular, for each i,

$$
if |r_i s_0| \le |r_i p|, then \angle ps_0 r_i \text{ is bad.} \tag{2.2}
$$

Moreover, we have that

<span id="page-2-2"></span>
$$
|ps_0| < \max_{i=1,2} \{|pr_i|\}.\tag{2.3}
$$

*Proof.* Let  $\Delta \tilde{p} \tilde{r}_1 \tilde{r}_2 \subset \mathbb{R}^2$  be the comparison triangle of  $\Delta p r_1 r_2$ . By Lemma 1.2 (and the first variation formula on  $\mathbb{R}^2$ ), the badness of  $\angle pr_1r_2$  implies that the function  $|ps| - |\tilde{p}\tilde{s}|$  with  $s \in [r_1r_2], \tilde{s} \in [\tilde{r}_1\tilde{r}_2]$  and  $|r_i s| = |\tilde{r}_i\tilde{s}|$  attains a negative minimum at some  $s_0 \in [r_1 r_2]^\circ$ . By Lemma 1.2 again, for any  $[ps_0]$ , we have that

$$
\angle ps_0r_i \ge \angle \tilde{p}\tilde{s}_0\tilde{r}_i, \quad i=1,2.
$$

It then has to hold that

$$
\angle ps_0r_i = \angle \tilde{p}\tilde{s}_0\tilde{r}_i, \quad i = 1, 2
$$

because  $\angle ps_0r_1 + \angle ps_0r_2 = \pi$  by Lemma 1.1. On the other hand, since  $|ps_0| < |\tilde{ps}_0|$ , there is  $\tilde{p}' \in [\tilde{p}\tilde{s}_0]^\circ$  such that  $|\tilde{p}'\tilde{s}_0| = |ps_0|$ . It is clear that  $|\tilde{p}'\tilde{r}_i| < |\tilde{p}\tilde{r}_i|$  for at least one of *i*, which implies  $\angle \tilde{p} \tilde{s}_0 \tilde{r}_i < \tilde{\angle} p s_0 r_i$ , i.e. (2.1) holds. Especially, it is easy to see that  $|\tilde{p}'\tilde{r}_i| < |\tilde{p}\tilde{r}_i|$  if  $|r_i s_0| \leq |r_i p|^{6}$  $|r_i s_0| \leq |r_i p|^{6}$  $|r_i s_0| \leq |r_i p|^{6}$ ; and thus (2.2) follows. Moreover, it is clear that  $|\tilde{p}\tilde{s}_0| < \max_{i=1,2} \{|pr_i|\}^7$  $|\tilde{p}\tilde{s}_0| < \max_{i=1,2} \{|pr_i|\}^7$ , which implies [\(2.3\)](#page-2-2).

Corollary 2.2 For  $\triangle pr_1r_2 \subset X$ , if  $\angle pr_1r_2$  is bad, then there is  $\bar{s} \in [r_1r_2]$  such that

$$
|p\bar{s}| \le \max_{i=1,2} \{|pr_i|\},\tag{2.4}
$$

and for any small  $\delta > 0$  there exist  $s_1, s_2 \in [r_1r_2] \cap B(\bar{s}, \delta)$  such that for any  $[ps_i]$ 

$$
\angle ps_1s_2 \text{ or } \angle ps_2s_1 \text{ is bad.} \tag{2.5}
$$

*Proof.* By Lemma 2.1, there is  $[ps_0]$  with  $s_0 \in [r_1r_2]^\circ$  such that  $\angle ps_0r_1$  or  $\angle ps_0r_2$  is bad, say ∠ps<sub>0</sub>r<sub>1</sub>. Then we can apply Lemma 2.1 to  $\triangle pso_{r_1}$  again. By repeating this infinite times, we can locate a desired  $\bar{s}$  if we take into account (2.2) and (2.3).

<span id="page-2-0"></span><sup>&</sup>lt;sup>6</sup>For the case where  $k > 0$  and  $|r_i p| \geq \frac{\pi}{2\sqrt{k}}$ , it needs  $\angle \tilde{r}_i \tilde{p} \tilde{s}_0 < \frac{\pi}{2}$  besides  $|r_i s_0| \leq |r_i p|$  (note that  $\Delta \tilde{p}\tilde{r}_1\tilde{r}_2 \subset \mathbb{S}_k^2$  and  $|r_i p| < \frac{\pi}{\sqrt{k}}$ , cf. the comments following Theorem A). In fact, ' $\angle \tilde{r}_i \tilde{p}\tilde{s}_0 < \frac{\pi}{2}$ ' holds obviously if  $|r_i s_0| \ll |r_i p|$  (here, it needs only ' $|r_i s_0| \leq |r_i p|$ ' for other cases including  $k \leq 0$ ).

<span id="page-2-1"></span>The case where  $k > 0$  and  $\max_{i=1,2} \{|pr_i|\} \ge \frac{\pi}{2\sqrt{k}}$ , it needs to be modified to  $\left|\tilde{p}\tilde{s}_0\right| < \max_{i=1,2} \{|pr_i|\}$  +  $\tau(|r_1r_2|)'$ , where  $\tau(|r_1r_2|)/|r_1r_2| \to 0$  as  $|r_1r_2| \to 0$ ; moreover, the function  $\tau(\cdot)$  can be chosen to be the same one (i.e. not depending on  $|pr_i|$ ) if  $\max_{i=1,2} \{|pr_i|\} \leq c < \frac{\pi}{\sqrt{k}}$  for some constant c.

In our proof of Theorem A, Lemmas 1.3 and 2.1 shall be the mere keys. For simpleness, we first consider the case where  $X$  is, in addition, locally compact.

#### Proof of Theorem A where X is locally compact.

We argue by contradiction. Assume that there is a  $\triangle pqr$  such that  $\angle pqr < \angle pqr$ . Then by Corollary 2.2 (see  $(2.5)$ ), we can consider the nonempty set

$$
\mathcal{S} \triangleq \{x \in X \mid \forall \delta > 0, \exists \triangle pr_1r_2 \text{ with } r_i \in B(x,\delta) \text{ s.t. } \angle pr_1r_2 \text{ is bad}\}.
$$

It is clear that  $p \notin \mathcal{S}$  because Theorem A holds around p, and that  $\mathcal{S}$  is closed. Note that a closed and bounded subset in  $X$  is compact because  $X$  is complete and locally compact (cf. Chapter 2 in [BBI]). So, there is  $\overline{o} \in \mathcal{S}$  such that  $|\overline{po}| = \min_{o \in \mathcal{S}} { |po| } > 0.$ 

On the other hand, Theorem A holds around any  $o \in \mathcal{S}$ , and thus we can define a positive function  $\delta(o) \triangleq \min\{\frac{|po|}{2}, \delta_o\}$ , where  $\delta_o$  is the maximal number such that

$$
any \triangle xyz \subset B(o, \delta_o) \text{ is good.} \tag{2.6}
$$

**Claim**: In  $B(o, \delta(o))$ , there is another point  $o' \in S$  such that  $|po'| \le |po| - \frac{\delta(o)}{3}$ .

It is obvious that the claim contradicts the existence of  $\bar{o}$ . Thereby, we just need to verify the claim. Due to  $o \in \mathcal{S}$ , there is a  $\triangle pr_1r_2$  with  $r_i \in B(o, \delta(o))$  such that

$$
|or_i| \ll \delta(o), \text{ and } \angle pr_1r_2 \text{ is bad.}
$$

Let  $\bar{r}_1 \in [pr_1]$  with  $|p\bar{r}_1| = |po| - \frac{\delta(o)}{3}$ . Note that for any  $[\bar{r}_1r_2]$ ,  $\triangle r_2r_1\bar{r}_1$  is good because it is contained in  $B(o, \delta(o))$ ; so via Lemmas 1.3 and 1.1 on  $\triangle pr_1r_2$ , the badness of  $\angle pr_1r_2$  implies

## $\angle p\bar{r}_1r_2$  is bad.

Then we can apply (2.2) in Lemma 2.1 (note that  $|\bar{r}_1r_2| < \delta(o) < |p\bar{r}_1|$  because  $\delta(o) \le$  $\frac{|p_0|}{2}$  to  $\triangle p\bar{r}_1r_2$  to locate an  $s \in [\bar{r}_1r_2]^\circ$  such that

 $\angle ps\bar{r}_1$  is bad for any [ps].

Put  $\bar{r}_2 \triangleq s$  if  $|ps| \leq |po| - \frac{\delta(s)}{3}$ ; otherwise, similar to  $\bar{r}_1$ , we can select  $\bar{r}_2 \in [ps]$  with

$$
|p\bar{r}_2| = |po| - \frac{\delta(o)}{3}, \text{ and } \angle p\bar{r}_2\bar{r}_1 \text{ is bad for any } [\bar{r}_1\bar{r}_2].
$$

(Here,  $\bar{r}_1, s, \bar{r}_2$  all lie in  $B(o, \epsilon)$  with  $\epsilon$  very close to  $\frac{\delta(o)}{3}$ . Note that  $|r_2x| + |xp|$  with  $x = \bar{r}_1, s, \bar{r}_2$  is very close to  $|po|$  because  $|or_i| \ll \delta(o)$ .) Then by Corollary 2.2, the badness of  $\angle p\bar{r}_2\bar{r}_1$  enables us to locate a point  $o' \in [\bar{r}_1\bar{r}_2] \cap S$ , which lies in  $B(o, \delta(o))$ , such that  $|po'| \le |po| - \frac{\delta(o)}{3}$  (see (2.4) and note that  $|p\bar{r}_i| \le |po| - \frac{\delta(o)}{3}$ <sup>[8](#page-3-0)</sup>  $\Box$ 

Eventually, we prove Theorem A for general  $X$ , i.e.  $X$  might not be locally compact. Proof of Theorem A where X is not locally compact.

<span id="page-3-0"></span><sup>&</sup>lt;sup>8</sup>For the case where  $k > 0$  and  $\max_{i=1,2} \{|p\bar{r}_i|\} \ge \frac{\pi}{2\sqrt{k}}$ ,  $\delta(o)$  should be additionally so small that  $\tau(\delta(o)) \ll \delta(o)$  (e.g.  $\tau(\delta(o)) < 0.01\delta(o)$ ), where  $\tau(\cdot)$  is just the function in Footnote 7.

When X is not locally compact, there might be no  $\bar{o} \in \mathcal{S}$  with  $|p\bar{o}| = \min_{o \in \mathcal{S}} \{|po|\}$ in the proof right above. However, starting with an  $o_1 \in \mathcal{S}$  and by the claim below  $(2.6)$ , we can step by step obtain  $\{o_i\}_{i=1}^{\infty} \subset S$  such that

$$
|po_{i+1}| \le |po_i| - \frac{\delta(o_i)}{3}
$$
 and  $|o_i o_{i+1}| < \delta(o_i)$ .

It follows that

$$
|po_{i+1}| \le |po_1| - \sum_{j=1}^i \frac{\delta(o_j)}{3}
$$
, and thus  $\sum_{j=1}^i \frac{\delta(o_j)}{3} \le |po_1|$ .

This implies that lim i→∞  $\delta(o_i) = 0$ , and that  $\{o_i\}_{i=1}^{\infty}$  is a Cauchy sequence (note that  $|o_i o_{i+1}| < \delta(o_i)$ ). Then by the completeness of X and the closedness of S,  $\{o_i\}_{i=1}^{\infty}$  has a limit point  $\bar{o} \in \mathcal{S}$ . However, it follows that lim  $i \rightarrow \infty$  $\delta(o_i) = \delta(\bar{o}) > 0$ , a contradiction.  $\Box$ 

# References

- [AKP] S. Alexander, V. Kapovitch and A. Petrunin, Alexandrov Geometry, 2010.
- [BBI] D. Burago, Y. Burago and S. Ivanov, A Course in Metric Geometry, Graduate Studies in Mathematics Vol. 33, AMS, 2001.
- [BGP] Y. Burago, M. Gromov and G. Perel'man, A.D. Aleksandrov spaces with curvature bounded below, Uspeckhi Mat. Nank 47(2): 3-51, 1992.
- [Pl] C. Plaut, Spaces of Wald-Berestovskii curvature bounded below, J. Geom. Anal. 6: 113-134, 1996.
- [Sh] K. Shiohama, An introduction to the geometry of Alexandrov spaces, 1992.
- [Wa] Y. Wang, A Schur-Toponogov theorem in Riemannian geometry & a new proof of Toponogov's theorem in Alexandrov geometry, [arXiv:1809.09818.](http://arxiv.org/abs/1809.09818)
- [SSW] X. Su, H. Sun and Y. Wang, Pythagorean Theorem & curvature with lower or upper bound, to appear in Chinese Ann. of Math. (B), 2022.

School of Mathematical Sciences (and Lab. math. Com. Sys.), Beijing Normal University, Beijing, 100875 P.R.C.

E-mail: 13716456647@163.com; suxiaole@bnu.edu.cn; wyusheng@bnu.edu.cn