On the classification of low-degree ovoids of Q(4, q)

Daniele Bartoli^{*} and Nicola Durante[†]

Abstract

Ovoids of the non-degenerate quadric Q(4,q) of PG(4,q) have been studied since the end of the '80s. They are rare objects and, beside the classical example given by an elliptic quadric, only three classes are known for q odd, one class for q even, and a sporadic example for $q = 3^5$. It is well known that to any ovoid of Q(4,q) a bivariate polynomial f(x,y) can be associated. In this paper we classify ovoids of Q(4,q) whose corresponding polynomial f(x,y) has "low degree" compared with q, in particular $deg(f) < (\frac{1}{6.3}q)^{\frac{3}{13}} - 1$. Finally, as an application, two classes of permutation polynomials in characteristic 3 are obtained.

MSC: 05B25; 11T06; 51E20.

1 Introduction

Let \mathbb{F}_{q^n} , $q = p^h$, be the finite field with q^n elements. Denote by $tr(x) = x + x^p + \cdots x^{p^{hn-1}}$ the absolute trace and by $N(x) = N_{\mathbb{F}_{q^n}/\mathbb{F}_q}(x) = x^{\frac{q^n-1}{q-1}}$ the norm of an element x of \mathbb{F}_{q^n} . For every $i \in \mathbb{F}_q$, let $N_i = \{x \in \mathbb{F}_{q^n} | N(x) = i\}$.

Denote by \mathbb{P} a finite classical polar space, i.e. the set of absolute points of either a polarity or a non-singular quadratic form of a projective space PG(m,q). The maximal dimensional projective subspaces contained in \mathbb{P} are the generators of \mathbb{P} .

Definition 1.1. Let \mathbb{P} be a finite classical polar space of PG(m, q). An ovoid of \mathbb{P} is a set of points of \mathbb{P} that has exactly one point in common with each generator of \mathbb{P} . An ovoid O of \mathbb{P} is a translation ovoid with respect to a point $P \in O$ if there is a collineation group of \mathbb{P} fixing P linewise (i.e. stabilizing all lines through P) and acting sharply transitively on the points of $O \setminus \{P\}$.

Regarding ovoids of the parabolic quadric Q(m,q) of PG(m,q), m even, it is known that there are no ovoids if m > 6 (see [18,36]). Hence for parabolic quadrics the only open

 $^{^* {\}rm Department}$ of Mathematics and Informatics, University of Perugia, Perugia, Italy. Email address: daniele.bartoli@unipg.it

[†]Department of Mathematics and applications R. Caccioppoli, University of Naples Federico II, Naples, Italy. Email address: ndurante@unina.it

problems concern m = 4 and m = 6. For m = 6, there are no ovoids if q is a prime $(q \neq 3)$ and there are two families of ovoids for $Q(6, 3^h)$; see [5, 9, 31, 35, 36, 38].

For more results on ovoids of finite classical polar spaces see e.g. [12].

In this paper we will focus on ovoids of Q(4, q). Denote by $(X_0, X_1, X_2, X_3, X_4)$ the homogenous coordinates of PG(4, q) and by H_{∞} the hyperplane at infinity $X_0 = 0$.

Up to collineation, in PG(4, q) there is a unique non-degenerate quadric, the parabolic quadric Q(4, q). In the sequel we fix the following parabolic quadric:

$$\mathbf{Q}: X_0 X_4 + X_1 X_3 + X_2^2 = 0.$$

Two points of Q(4,q) are non-collinear if they are not contained in any generator of Q(4,q). It can be easily shown that an ovoid of Q(4,q) is a set of $q^2 + 1$ pairwise non-collinear points on the quadric.

We would like also to underline that ovoids of Q(4, q) are related to many relevant objects in finite projective spaces such as ovoids of PG(3, q) (if q is even) see e.g. [3], [38], semifield flocks of a quadratic cone of PG(3, q), eggs of finite projective spaces, translation generalized quadrangles and C_F^{σ} -sets of PG(2, q) (if q is odd) (see e.g. [25], [29], [14]). Also, ovoids of Q(4, q) are equivalent to symplectic spreads of PG(3, q) via the isomorphism between Q(4, q)and the dual of the symplectic generalized quadrangle W(3, q) see [34].

Since the collineation group of Q(4,q) acts transitive on pairs non-collinear points, we can always assume that an ovoid O_4 of Q(4,q) contains the points (1,0,0,0,0) and (0,0,0,0,1) and hence it can be written in the following form:

$$O_4(f) = \{(1, x, y, f(x, y), -y^2 - xf(x, y))\}_{x, y \in \mathbb{F}_q} \cup \{(0, 0, 0, 0, 1)\},$$
(1)

for some function $f : \mathbb{F}_q^2 \longrightarrow \mathbb{F}_q$ with f(0,0) = 0. The set $O_4(f)$ is an ovoid if and only if

$$\langle (1, x_1, y_1, f(x_1, y_1), -y_1^2 - x_1 f(x_1, y_1), (1, x_2, y_2, f(x_2, y_2), -y_2^2 - x_2 f(x_2, y_2) \rangle \neq 0, \quad (2)$$

for every $(x_1, y_1) \neq (x_2, y_2)$ in \mathbb{F}_q^2 , where $\langle \cdot, \cdot \rangle$ is the symmetric for q odd (or alternating for q even) form associated to the quadratic form of the quadric Q(4, q).

Note that Condition (2) reads

$$(y_1 - y_2)^2 + (x_1 - x_2)(f(x_1, y_1) - f(x_2, y_2)) \neq 0,$$
(3)

for every $(x_1, y_1) \neq (x_2, y_2)$ in \mathbb{F}_q^2 .

The only known classification theorems for ovoids of Q(4,q) are the following.

Theorem 1.2. [5] If O is an ovoid of Q(4,q), q odd prime, then O is an elliptic quadric.

Theorem 1.3. [4, 26] Let f(X,Y) = g(X) + h(Y), with both g and $h \mathbb{F}_{q'}$ -linear maps, $q = (q')^n$, q odd. If $q' \ge 4n^2 - 8n + 2$ or q' is a prime and $q' > 2n^2 - (4 - 2\sqrt{3})n + (3 - 2\sqrt{3})$, then $O_4(f)$ is either an elliptic quadric or a Kantor ovoid.

Also, since ovoids of Q(4, q) with f as in Theorem 1.3 are equivalent to rank 2 commutative semifields the following also is known.

Theorem 1.4. [27] Let f(X,Y) = g(X) + h(Y), with both g and $h \mathbb{F}_{q'}$ -linear maps, $q = (q')^n$, q odd. If either $n \leq 4$ or (q', n) = (3, 5), then $O_4(f)$ is either an elliptic quadric or a Kantor ovoid or a Thas-Payne ovoid or a Penttila-Williams ovoid.

In Section 2, using (3), a hypersurface S_f of PG(4,q) will be attached to $O_4(f)$. A similar connection between ovoids in PG(2,q) (hyperovals), q even, and algebraic curves over finite fields was used in [11,21] to classify hyperovals corresponding to o-polynomials of small degree. Investigating the existence of absolutely irreducible components defined over \mathbb{F}_q in S_f , we will provide the following classification result, which is also the main achievement of our paper.

Main Theorem. Let $f(X,Y) = \sum_{ij} a_{i,j} X^i Y^j$, with $a_{0,1} = 0$ if p > 2. Suppose that $q > 6.3(d+1)^{13/3}$, where $\deg(f) = d$. If $O_4(f)$ is an ovoid of Q then one of the following holds:

- 1. p = 2 and $f(X, Y) = a_{1,0}X + a_{0,1}Y$. Hence $O_4(f)$ is an elliptic quadric.
- 2. p > 2 and $f(X, Y) = a_{p^j,0} X^{p^j}$, $j \ge 0$. Hence $O_4(f)$ is either an elliptic quadric or a Kantor ovoid.

The only known ovoids of Q(4,q) are listed in Table 1, where *n* is a non-square of \mathbb{F}_q , $q = p^h$ odd, $\sigma \neq 1$ is an automorphism of \mathbb{F}_q , and $a \neq 1$ is an element with tr(a) = 1 of \mathbb{F}_q , q even. All but the last two examples of ovoids in Table 1 are translation ovoids of Q(4,q) with respect to their point at infinity.

Name	f(x,y)	Restrictions	Ref.
Elliptic quadric	-nx	q odd	
Elliptic quadric	ax + y	q even	
Kantor	$-nx^{\sigma}$	q odd, h > 1	[23]
Penttila-Williams	$-x^9 - y^{81}$	p = 3, h = 5	[32]
Thas-Payne	$-nx - (n^{-1}x)^{1/9} - y^{1/3}$	p = 3, h > 2	[37]
Ree-Tits slice	$-x^{2\sigma+3}-y^{\sigma}$	$p = 3, h > 1, h \text{ odd}, \sigma = \sqrt{3q}$	[23]
Tits	$x^{\sigma+1} + y^{\sigma}$	$p = 2, h > 1, h \text{ odd}, \sigma = \sqrt{2q}$	[38]

Table 1: Known ovoids of Q(4, q).

Note that the Ree-Tits slice is obtained from slicing the Ree-Tits ovoid of Q(6,q); see [23].

In the sequel we briefly discuss the connection between these examples and Theorem 2.10.

In the even characteristic case, Main Theorem does not apply to the Tits ovoid

$$f_2(x,y) = x^{2^{h+1}+1} + y^{2^{h+1}}, \qquad q = 2^{2h+1},$$

since $\deg(f_2) \simeq \sqrt{q}$.

In the odd characteristic case, more examples of ovoids are known. Apart from the case $f_3(x, y) = -nx^{p^r}$, $q = p^h$, (Elliptic quadric and Kantor) which satisfies Condition 2 in Main Theorem, the two infinite families (Ree-Tits slice and Thas-Payne in characteristic 3) arise from $f_4(x, y) = -x^{3^{2\cdot 3^{h+1}+3}} - y^{3^{h+1}}$, when $q = 3^{2h+1}$, and $f_5(x, y) = -nx - (n^{-1}x)^{1/9} - y^{1/3}$, $q = 3^h$, h > 2. Note that in these cases, Theorem 2.10 does not apply to such examples, since deg $(f_4) \simeq \sqrt{q}$ and deg $(f_5) > \sqrt[3]{q}$.

Finally, observe that for the Penttila-Williams sporadic example in \mathbb{F}_{3^5} arising from $f_6(x,y) = -x^9 - y^{81}$ the hypothesis of Main Theorem does not hold since deg (f_6) is too high with respect to $q = 3^5$.

This paper is organized as follows. The main result is proved in Section 2. Finally, in Section 3 two families of permutation polynomials connected with Pentilla-William and Thas-Payne ovoids are obtained.

2 Proof of the main theorem

The aim of this section is to provide a partial classification of the ovoids of Q(4, q) via the investigation of specific algebraic varieties; see [6] for a survey on links between algebraic varieties over finite fields and relevant combinatorial objects.

An algebraic hypersurface S is an algebraic variety that may be defined by a single implicit equation. An algebraic hypersurface defined over a field \mathbb{K} is *absolutely irreducible* if the associated polynomial is irreducible over every algebraic extension of \mathbb{K} . An absolutely irreducible \mathbb{K} -rational component of a hypersurface S, defined by the polynomial F, is simply an absolutely irreducible hypersurface such that the associated polynomial has coefficients in \mathbb{K} and it is a factor of F. As in the previous sections, we will make use of the homogeneous equations for algebraic varieties. For a deeper introduction to algebraic varieties we refer the interested reader to [19].

As mentioned in Section 1 an ovoid of $Q: X_0X_4 + X_1X_3 + X_2^2 = 0$ can be written in the form $O_4(f) = \{(1, x, y, f(x, y), -y^2 - xf(x, y)\}_{x,y \in \mathbb{F}_q} \cup \{(0, 0, 0, 0, 1)\}, \text{ for some function} f \text{ with } f(0, 0) = 0 \text{ and } (y_1 - y_2)^2 + (x_1 - x_2)(f(x_1, y_1) - f(x_2, y_2)) \neq 0 \text{ for every } (x_1, y_1) \neq (x_2, y_2) \text{ in } \mathbb{F}_q^2$. Let $\tilde{f}(X, Y, T)$ be the homogenization of f(X, Y) and let $d = \deg(f)$. In order to obtain non-existence results and partial classifications for the ovoids of Q, we will consider the hypersurface $S_f \subset PG(4, q)$ defined by $F(X_0, X_1, X_2, X_3, X_4) = 0$, where

$$F(X_0, X_1, X_2, X_3, X_4) = (X_2 - X_4)^2 X_0^{d-1} + (X_1 - X_3) \Big(\widetilde{f}(X_1, X_2, X_0) - \widetilde{f}(X_3, X_4, X_0) \Big).$$
(4)

Note that $O_4(f)$ is an ovoid of Q if and only if S_f contains no affine \mathbb{F}_q -rational points off the plane $X_1 - X_3 = 0 = X_2 - X_4$.

A fundamental tool to determine the existence of rational points in an algebraic variety over finite fields is the following result by Lang and Weil [24] in 1954, which can be seen as a generalization of the Hasse-Weil bound.

Theorem 2.1. [Lang-Weil Theorem] Let $\mathcal{V} \subset \mathbb{P}^N(\mathbb{F}_q)$ be an absolutely irreducible variety of dimension n and degree d. Then there exists a constant C depending only on N, n, and d such that

$$\left| \# \mathcal{V}(\mathbb{F}_q) - \sum_{i=0}^n q^i \right| \le (d-1)(d-2)q^{n-1/2} + Cq^{n-1}.$$
 (5)

Although the constant C was not computed in [24], explicit estimates have been provided for instance in [8,10,16,17,28,33] and they have the general shape C = f(d) provided that q > g(n,d), where f and g are polynomials of (usually) small degree. We refer to [10] for a survey on these bounds. Excellent surveys on Hasse-Weil and Lang-Weil type theorems are [15,16]. We include here the following result by Cafure and Matera [10].

Theorem 2.2. [10, Theorem 7.1] Let $\mathcal{V} \subset AG(n,q)$ be an absolutely irreducible \mathbb{F}_q -variety of dimension r > 0 and degree δ . If $q > 2(r+1)\delta^2$, then the following estimate holds:

$$|\#(\mathcal{V} \cap \mathrm{AG}(n,q)) - q^r| \le (\delta - 1)(\delta - 2)q^{r-1/2} + 5\delta^{13/3}q^{r-1}.$$

For our purposes, the existence of an absolutely irreducible \mathbb{F}_q -rational component in \mathcal{S}_f is enough to provide asymptotic non-existence results.

Theorem 2.3. Let $S_f : F(X_0, X_1, X_2, X_3, X_4) = 0$, where F is defined as in (4). Suppose that $q > 6.3(d+1)^{13/3}$, $\deg(f) = d$, and S_f contains an absolutely irreducible component \mathcal{V} defined over \mathbb{F}_q . Then $O_4(f)$ is not an ovoid of Q.

Proof. Since $q > 6.3(d+1)^{13/3}$, by Theorem 2.2, with $\delta = d+1$ and r = 3, it is readily seen that \mathcal{V} contains more than q^2 affine \mathbb{F}_q -rational points. This yields the existence of at least one affine \mathbb{F}_q -rational point (a, b, c, d) in $\mathcal{V} \subset \mathcal{S}_f$ such that $a \neq c$ or $b \neq d$ and therefore $O_4(f)$ is not an ovoid of Q.

We also include, for sake of completeness, the following results which will be useful in the sequel.

Lemma 2.4. [2, Lemma 2.1] Let $\mathcal{H}, \mathcal{S} \subset PG(n,q)$ be two projective hypersurfaces. If $\mathcal{H} \cap \mathcal{S}$ has a reduced absolutely irreducible component defined over \mathbb{F}_q then \mathcal{S} has an absolutely irreducible component defined over \mathbb{F}_q .

Lemma 2.5. [7, Lemma 2.9] Let S be a hypersurface containing O = (0, 0, ..., 0) of affine equation $F(X_1, ..., X_n) = 0$, where $F(X_1, ..., X_n) = F_d(X_1, ..., X_n) + F_{d+1}(X_1, ..., X_n) + ...,$ with F_i the homogeneous part of degree i of $F(X_1, ..., X_n)$ for i = d, d + 1, ... Let Pbe an \mathbb{F}_q -rational simple point of the variety $F_d(X_1, ..., X_n) = 0$. Then there exists an \mathbb{F}_q rational plane π through the line ℓ joining O and P such that $\pi \cap S$ has ℓ as a non-repeated tangent \mathbb{F}_q -rational line at the origin and $\pi \cap S$ has a non-repeated absolutely irreducible \mathbb{F}_q -rational component. In what follows, $f(X,Y) = \sum_{i,j} a_{i,j} X^i Y^j$, with $a_{i,j} \in \mathbb{F}_q$ for all i, j, and $d = \deg(f) = \max\{i+j : a_{i,j} \neq 0\}$. So,

$$F(X_0, X_1, X_2, X_3, X_4) = (X_2 - X_4)^2 X_0^{d-1} + (X_1 - X_3) \left[\sum_{i,j} a_{ij} \left(X_1^i X_2^j - X_3^i X_4^j \right) X_0^{d-i-j} \right].$$
(6)

In view of Theorem 2.3, our goal is to find conditions for which S_f possesses an absolutely irreducible component defined over \mathbb{F}_q . Recall that f(0,0) = 0 and therefore $a_{0,0} = 0$. Also, if p > 2, since S_f is equivalent to $S_{f'} : F(X_0, X_1, X_2 - a_{0,1}X_1/2, X_3, X_4 - a_{0,1}X_3/2) = 0$, where $a'_{0,1} = 0$, without loss of generality we can assume $a_{0,1} = 0$.

Proposition 2.6. Let S_f : $F(X_0, X_1, X_2, X_3, X_4) = 0$, with F as in (6). Let d > 1. If there exists $i \neq d$ such that $a_{i,d-i} \neq 0$ then S_f contains an absolutely irreducible component defined over \mathbb{F}_q .

Proof. Let $\mathcal{S}' = \mathcal{S}_f \cap H_\infty$, where $H_\infty : X_0 = 0$. Now, the equation of \mathcal{S}' reads

$$(X_1 - X_3) \sum_{i \le d} a_{i,d-i} \left(X_1^i X_2^{d-i} - X_3^i X_4^{d-i} \right) = 0.$$

Note that $(X_1 - X_3) | G(X_1, X_2, X_3, X_4) := \sum_{i \le d} a_{i,d-i} \left(X_1^i X_2^{d-i} - X_3^i X_4^{d-i} \right)$ if and only if $G(X_1, X_2, X_1, X_4) = \sum_{i \le d} a_{i,d-i} X_1^i \left(X_2^{d-i} - X_4^{d-i} \right)$ is the zero polynomial. This would be possible if and only if $G(X_1, X_2, X_3, X_4) = a_{d,0} \left(X_1^d - X_3^d \right)$, that is $a_{i,d-i} = 0$ for all $i \ne d$, a contradiction.

Suppose now that there exists an $i \neq d$ such that $a_{i,d-i} \neq 0$. Since $(X_1 - X_3)$ does not divide $G(X_1, X_2, X_3, X_4)$, $(X_1 - X_3)$ is a non-repeated absolutely irreducible component of \mathcal{S}' defined over \mathbb{F}_q . By Lemma 2.4, \mathcal{S} contains an absolutely irreducible component defined over \mathbb{F}_q (passing through $H_{\infty} \cap (X_1 = X_3)$).

In what follows we denote by \overline{j} the nonnegative integer max $\{j : a_{i,j} \neq 0\}$.

Proposition 2.7. Let $S_f : F(X_0, X_1, X_2, X_3, X_4) = 0$, with F as in (6). If $\overline{j} > 1$ then S_f contains an absolutely irreducible component defined over \mathbb{F}_q .

Proof. The polynomial $F(X_4, X_1, X_2, X_3, X_0)$ reads

$$(X_2 - X_0)^2 X_4^{d-1} + (X_1 - X_3) \left[\sum_{i,j} a_{ij} \left(X_1^i X_2^j - X_3^i X_0^j \right) X_4^{d-i-j} \right]$$

and the surface $\widetilde{\mathcal{S}}$: $F(X_4, X_1, X_2, X_3, X_0) = 0$ is projectively equivalent to \mathcal{S}_f .

We study now the tangent cone Ω of \widetilde{S} at the origin, whose equation depends on \overline{j} . We distinguish three cases.

- (i) $\overline{j} \geq 3$. The tangent cone Ω reads $(X_1 X_3) \sum_i a_{i,\overline{j}} X_3^i X_4^{d-i-\overline{j}}$.
- (ii) $\overline{j} = 2$. Then Ω reads $X_4^{d-1} (X_1 X_3) \sum_i a_{i,2} X_3^i X_4^{d-i-2}$.
- (iii) $\overline{j} \leq 1$. The tangent cone reads X_4^{d-1} .

If case (i) holds, then $X_1 - X_3 = 0$ is a non-repeated \mathbb{F}_q -rational component of such a tangent cone. Also, there exists a non-singular point P for Ω . By Lemma 2.5 there exists an \mathbb{F}_q -rational plane π through the line ℓ joining the origin and P such that $\pi \cap \widetilde{S}$ has ℓ as a non-repeated tangent \mathbb{F}_q -rational line at the origin and $\pi \cap \widetilde{S}$ has a non-repeated absolutely irreducible \mathbb{F}_q -rational component. Using twice Lemma 2.4, one deduces the existence of an absolutely irreducible \mathbb{F}_q -rational component in \widetilde{S} and therefore in S_f .

If case (ii) holds, then Ω is a rational surface $(X_1 \text{ has degree 1})$. This means that Ω contains an absolutely irreducible \mathbb{F}_q -rational component of degree 1 in X_1 . It is readily seen that affine singular points of Ω satisfy $X_4 = 0$ and therefore there exists at least a non-singular point P for Ω . The same conclusion as in case (i) holds.

Proposition 2.8. Let $S_f : F(X_0, X_1, X_2, X_3, X_4) = 0$, with F as in (6), and p = 2. If there exists an $a_{i,j} \neq 0$, with $(i,j) \notin \{(1,0), (0,1)\}$, then S_f contains an absolutely irreducible component defined over \mathbb{F}_q .

Proof. From Proposition 2.7, if $\overline{j} > 1$ then S_f contains an absolutely irreducible component defined over \mathbb{F}_q . Therefore we can suppose that $\overline{j} \leq 1$. Suppose that d > 1.

Consider again Equation (4) and let $\mathcal{W} = \mathcal{S}_f \cap (X_2 = X_4)$ be defined by

$$(X_1 + X_3) \left[\sum_{i,j} a_{ij} X_2^j X_0^{d-i-j} \left(X_1^i + X_3^i \right) \right]$$

= $(X_1 + X_3) \left(X_2 \sum_{0 < i \le d-2} a_{i,1} (X_1^i + X_3^i) X_0^{d-i-1} + \sum_{0 < i \le d} a_{i,0} (X_1^i + X_3^i) X_0^{d-i} \right).$

If some of the $a_{i,1}$'s, $0 < i \leq d-2$, is nonzero then \mathcal{W} contains a rational component of degree 1 in X_2 and therefore \mathcal{S}_f contains an absolutely irreducible component defined over \mathbb{F}_q through it.

So, $a_{i,1} = 0$ for i > 0 and $F(X_0, X_1, X_2, X_3, X_4)$ reads

$$(X_2 + X_4)^2 X_0^{d-1} + a_{0,1} (X_1 + X_3) (X_2 + X_4) X_0^{d-1} + (X_1 + X_3) \sum_{0 < i \le d} a_{i,0} \left(X_1^i + X_3^i \right) X_0^{d-i}.$$

If $a_{0,1} = 0$ then S_f is clearly a rational surface (in $Z = (X_2 + X_4)^2$) and therefore it contains an absolutely irreducible component defined over \mathbb{F}_q . Suppose now $a_{0,1} \neq 0$ and consider now the affine part of $S_f : F(1, X_1, X_2, X_3, X_4) = 0$. It can be written as

$$Z^{2} + Z + \frac{\sum_{0 < i \le d} a_{i,0} \left(X_{1}^{i} + X_{3}^{i} \right)}{a_{0,1}^{2} (X_{1} + X_{3})},$$
(7)

where $Z = (X_2 + X_4)/(a_{0,1}(X_1 + X_3)).$

In what follows we will prove that the surface \mathcal{Y} defined by the affine equation (7) is absolutely irreducible.

Note that the surface $Z^2 + Z + \frac{\sum_{0 \le i \le d} a_{i,0}(X_1^i + X_3^i)}{a_{0,1}^2(X_1 + X_3)} = 0$ is reducible only if there exists $\overline{Z}(X_1, X_3) = \frac{\overline{Z}_1(X_1, X_3)}{\overline{Z}_2(X_1, X_3)}$, with $\overline{Z}_i(X_1, X_3) \in \overline{\mathbb{F}}_q[X_1, X_3]$, such that $(\overline{Z}(X_1, X_3))^2 + \overline{Z}(X_1, X_3) + \frac{\sum_{0 \le i \le d} a_{i,0}(X_1^i + X_3^i)}{a_{0,1}^2(X_1 + X_3)} = 0$. This condition reads

$$(\overline{Z}_1(X_1, X_3))^2 + \overline{Z}_1(X_1, X_3)\overline{Z}_2(X_1, X_3) + (\overline{Z}_2(X_1, X_3))^2 \frac{\sum_{0 < i \le d} a_{i,0} \left(X_1^i + X_3^i\right)}{a_{0,1}^2 (X_1 + X_3)} = 0.$$
(8)

Now, $\deg(\overline{Z}_1) > \deg(\overline{Z}_2)$, otherwise $2\deg(\overline{Z}_1) \le \deg(\overline{Z}_1) + \deg(\overline{Z}_2) < 2\deg(\overline{Z}_2) + d - 1$ and (8) cannot be satisfied.

Also, the same argument shows that $2 \deg(\overline{Z}_1) = 2 \deg(\overline{Z}_2) + d - 1$ and therefore the highest homogeneous parts in $(\overline{Z}_1(X_1, X_3))^2$ and in $(\overline{Z}_2(X_1, X_3))^2 \frac{\sum_{0 \le i \le d} a_{i,0}(X_1^i + X_3^i)}{a_{0,1}^2(X_1 + X_3)}$ must be equal.

Write $d = 2^{j}k$, with $j \ge 0$ and k odd. The above argument yields that, in particular,

$$L_d(X_1, X_3) = \frac{\left(X_1^d + X_3^d\right)}{\left(X_1 + X_3\right)} = \frac{\left(X_1^k + X_3^k\right)^{2^j}}{\left(X_1 + X_3\right)}$$

the highest homogenous term in $\frac{\sum_{0 < i \leq d} a_{i,0}(X_1^i + X_3^i)}{a_{0,1}^2(X_1 + X_3)}$, is a square in $\overline{\mathbb{F}}_q[X_1, X_3]$.

- If j > 0 then $(X_1 + X_3)^{2^j 1}$ is the highest power of $(X_1 + X_3)$ dividing $L_d(X_1, X_3)$. Since $2^j - 1$ is odd, $L_d(X_1, X_3)$ is not a square in $\overline{\mathbb{F}}_q[X_1, X_3]$ and the surface \mathcal{Y} is irreducible.
- If j = 0, following the same argument as above, if \mathcal{Y} is reducible then k = 1, a contradiction to d > 1.

This shows that if d > 1 then \mathcal{Y} and therefore \mathcal{S}_f is absolutely irreducible or contains an absolutely irreducible component defined over \mathbb{F}_q . The claim follows.

Proposition 2.9. Let S_f : $F(X_0, X_1, X_2, X_3, X_4) = 0$, with F as in (6), p > 2, and $a_{0,1} = 0$. If $d \neq p^j$, with $j \geq 0$, or $a_{i,j} = 0$ for some $(i, j) \neq (d, 0)$, then S_f contains an absolutely irreducible component defined over \mathbb{F}_q .

Proof. From Proposition 2.7, if $\overline{j} > 1$ then S_f contains an absolutely irreducible component defined over \mathbb{F}_q . Therefore we can suppose that $\overline{j} \leq 1$.

Suppose that $\overline{j} = 1$. Consider $\widetilde{\mathcal{S}} : F(X_4, X_1, X_2, X_3, 1) = 0$ defined by

$$(X_2 - 1)^2 X_4^{d-1} + (X_1 - X_3) \left[\sum_i a_{i,1} \left(X_1^i X_2 - X_3^i \right) X_4^{d-i-1} \right] + (X_1 - X_3) \left[\sum_i a_{i,0} \left(X_1^i - X_3^i \right) X_4^{d-i} \right] = 0$$

Let $\mathcal{W} = \widetilde{\mathcal{S}} \cap (X_1 = 0)$ be defined by

$$(X_2 - 1)^2 X_4^{d-1} + X_3 \left[\sum_{i>0} a_{i,1} X_3^i X_4^{d-i-1} \right]$$
$$-a_{0,1} X_3 (X_2 - 1) X_4^{d-1} + X_3 \left[\sum_i a_{i,0} X_3^i X_4^{d-i} \right] = 0.$$

The above surface is of degree two in (X_2-1) and therefore, since p is odd, it is reducible if and only if

$$a_{0,1}^2 X_3^2 X_4^{2d-2} - 4X_4^{d-1} X_3 \left[\sum_{i>0} a_{i,1} X_3^i X_4^{d-i-1} + \sum_i a_{i,0} X_3^i X_4^{d-i} \right] = X_3^2 X_4^{d-1} \left(a_{0,1}^2 X_4^{d-1} - 4 \left[\sum_{i>0} a_{i,1} X_3^{i-1} X_4^{d-i-1} + \sum_i a_{i,0} X_3^{i-1} X_4^{d-i} \right] \right)$$

is a square in $\overline{\mathbb{F}}_q[X_3, X_4]$. If $a_{i,1} \neq 0$ for some i > 0 then the degrees of the monomials in $G(X_3, X_4) = a_{0,1}^2 X_4^{d-1} - 4 \left[\sum_{i>0} a_{i,1} X_3^{i-1} X_4^{d-i-1} + \sum_i a_{i,0} X_3^{i-1} X_4^{d-i} \right]$ are d-2 and d-1 and both the values are assumed. In this case, $G(X_3, X_4)$ cannot be a square in $\overline{\mathbb{F}}_q[X_3, X_4]$. Therefore \mathcal{W} is absolutely irreducible and, by Lemma 2.4, $\widetilde{\mathcal{S}}$ (and therefore \mathcal{S}) contains an absolutely irreducible component defined over \mathbb{F}_q .

Suppose now that $a_{i,1} = 0$ for all i > 0. So, $F(X_0, X_1, X_2, X_3, X_4)$ equals

$$(X_2 - X_4)^2 X_0^{d-1} + (X_1 - X_3) \left[\sum_i a_{i,0} \left(X_1^i - X_3^i \right) X_0^{d-i} \right].$$

In this case, consider the affine part of S_f defined by $F(1, X_1, X_2, X_3, X_4) = 0$. Note that, as already observed, S_f is reducible if and only if

$$(X_1 - X_3) \left[\sum_i a_{i,0} \left(X_1^i - X_3^i \right) \right]$$

is a square in $\overline{\mathbb{F}}_q[X_3, X_4]$.

In what follows we will show that if $L(X_3, X_4) = (X_1 - X_3) \left[\sum_i a_{i,0} \left(X_1^i - X_3^i \right) \right]$ is a square in $\overline{\mathbb{F}}_q[X_3, X_4]$ then there exists no *i* with 0 < i < d such that $a_{i,0} \neq 0$.

Suppose that $L(X_3, X_4)$ is a square in $\overline{\mathbb{F}}_q[X_3, X_4]$. Then, the highest homogenous term $L_d(X_3, X_4) = a_{d,0}(X_1 - X_3)(X_1^d - X_3^d)$ is a square in $\overline{\mathbb{F}}_q[X_3, X_4]$ too. Write $d = kp^j$, with $p \nmid k$. Then $L_d(X_3, X_4) = a_{d,0}(X_1 - X_3)(X_1^k - X_3^k)^{p^j}$.

- If k > 1 then there are other factors $(X_3 \eta X_4)$, $\eta^k = 1$, $\eta \neq 1$, such that $(X_3 \eta X_4)^{p^j}$ is the highest power of $(X_3 \eta X_4)$ dividing $L_d(X_3, X_4)$, a contradiction to $L_d(X_3, X_4)$ being a square.
- Let now k = 1 and $d = p^j$ for some $j \ge 0$. Suppose that there exists another 0 < i < d such that $a_{i,0} \ne 0$ and let $\overline{i} = \max\{i : a_{i,0} \ne 0, 0 < i < d\}$. Write

$$(X_1 - X_3) \left[\sum_i a_{i,0} \left(X_1^i - X_3^i \right) \right] = L_d(X_3, X_4) + L_{\overline{i}}(X_3, X_4) + \cdots$$
$$= \left((X_1 - X_3)^{(p^j + 1)/2} + M(X_3, X_4) + \cdots \right)^2,$$

where $M(X_3, X_4)$ has degree $\overline{i} - (p^j + 1)/2$. Now

$$2(X_1 - X_3)^{(p^j + 1)/2} M(X_3, X_4) = L_{\overline{i}}(X_3, X_4) = a_{\overline{i},0}(X_1 - X_3)(X_1^{\overline{i}} - X_3^{\overline{i}})$$

and then $(X_1 - X_3)^{(p^j - 1)/2} | (X_1^{\overline{i}} - X_3^{\overline{i}})$. Write $\overline{i} = k_1 p^{j_1}$, so that $(X_1^{\overline{i}} - X_3^{\overline{i}}) = (X_1^{k_1} - X_3^{k_1})^{p^{j_1}}$. Since $(X_1 - X_3)^{p^{j_1}}$ is the highest power of $(X_1 - X_3)$ dividing $(X_1^{k_1} - X_3^{k_1})^{p^{j_1}}$, we must have $(p^j - 1)/2 \le p^{j_1}$. Since $j_1 \le j - 1$,

$$p^j - 1 \le 2p^{j_1} \le 2p^j/p,$$

and therefore $p^j \leq \frac{p}{p-2}$. This yields $p^j = 3$, that is j = 1 and p = 3. Now, $(X_1 - X_3)^4 + \alpha(X_1 - X_3)(X_1^2 - X_3^2) + \beta(X_1 - X_3)^2$, $\alpha, \beta \in \mathbb{F}_q$, is a square in $\overline{\mathbb{F}_q}[X_1, X_3]$ only if $(X_1 - X_3)^2 + \alpha(X_1 + X_3) + \beta$ is a square. By direct checking, this is possible only if $\alpha = \beta = 0$.

This shows that if there exists an i, 1 < i < d, such that $a_{i,0} \neq 0$ then $L(X_3, X_4)$ is not a square in $\overline{\mathbb{F}}_q[X_3, X_4]$ and therefore \mathcal{S}_f is absolutely irreducible.

We are now in position to prove the main result of this paper.

Theorem 2.10. Let $f(X,Y) = \sum_{ij} a_{i,j} X^i Y^j$, with $a_{0,1} = 0$ if p > 2. Suppose that $q > 6.3(d+1)^{13/3}$, where $\deg(f) = d$. If $O_4(f)$ is an ovoid of Q then one of the following holds:

- 1. p = 2 and $f(X, Y) = a_{1,0}X + a_{0,1}Y$. Hence $O_4(f)$ is an elliptic quadric.
- 2. p > 2 and $f(X,Y) = a_{p^j,0}X^{p^j}$, $j \ge 0$. Hence $O_4(f)$ is either and elliptic quadric or a Kantor ovoid.

Proof. By combining Propositions 2.8 and 2.9 with Theorem 2.3 we obtain that either p = 2 and $f(X,Y) = a_{1,0}X + a_{0,1}Y$, or p > 2 and $f(X,Y) = a_{p^j,0}X^{p^j}$, $j \ge 0$. The claim then follows by applying Condition (3).

Note that the ovoids corresponding to the elliptic quadric

$$f_1(x,y) = ax + y, \qquad tr(a) = 1,$$

and the Tits ovoid

$$f_2(x,y) = x^{2^{h+1}+1} + y^{2^{h+1}}, \qquad q = 2^{2h+1}$$

yield the following equation for S_f

$$(X_2 + X_4)^2 + (X_1 + X_3)(aX_1 + X_2 + aX_3 + X_4) = 0$$
(9)

and

$$(X_2 + X_4)^2 X_0^{2^{h+1}} + (X_1 + X_3)(X_1^{2^{h+1}+1} + X_2^{2^{h+1}}X_0 + X_3^{2^{h+1}+1} + X_4^{2^{h+1}}X_0) = 0.$$
(10)

In particular, (9) defines a reducible quadric splitting into two linear factors defined over \mathbb{F}_{q^2} . It has been already observed that Main Theorem does not apply to $f_2(x, y)$, since $\deg(f_2) \simeq \sqrt{q}$. In this case, it is possible to prove that \mathcal{S}_f is absolutely irreducible.

3 Two classes of permutation polynomials in characteristic three

As a by-product of our investigation on ovoids of Q(4,q), we provide in this section two families of permutation polynomials in characteristic three. A polynomial $f(x) \in \mathbb{F}_q[x]$ is a *permutation polynomial* (PP) if it is a bijection of the finite field \mathbb{F}_q into itself. Permutation polynomials were first studied by Hermite and Dickson; see [13, 20].Particular, simple structures, or additional extraordinary properties are usually required by applications of PPs in other areas of mathematics and engineering, such as cryptography, coding theory, or combinatorial designs. In this case, permutation polynomials meeting these criteria are usually difficult to find. For a deeper introduction to the connections of PPs with other fields of mathematics we refer to [1,22,30,39] and the references therein.

In this section we will assume that $f(x,y) = \sum_i a_i x^{p^i} + \sum_j b_j y^{p^i}$ is the sum of two \mathbb{F}_p -linear functions the first one only in x and the other one only in y. Note that all known examples of ovoids $O_4(f)$, apart from the Tits and Ree-Tits slice, arise from functions f(x,y) of this type. We may assume that there exist \overline{i} such that $a_{\overline{i}} \neq 0$ and \overline{j} such that $b_{\overline{j}} \neq 0$. By using the same notation as in previous section we have that in this case the hypersurface S_f of PG(4, q) has equation $F(X_0, X_1, X_2, X_3, X_4) = 0$, where $F(X_0, X_1, X_2, X_3, X_4)$ equals

$$X_0^{d-2}(X_2 - X_4)^2 + (X_1 - X_3) \left[\sum_i a_i X_0^{d-1-p^i} (X_1 - X_3)^{p^i} + \sum_j b_j X_0^{d-1-p^j} (X_2 - X_4)^{p^j} \right].$$

Put $X_0 = 1$, $L := X_2 - X_4$ and $M := X_1 - X_3$. Then $F(X_0, X_1, X_2, X_3, X_4) = 0$ reads

$$L^{2} + \sum_{i} a_{i} M^{p^{i}+1} + M \sum_{j} b_{j} L^{p^{j}} = 0.$$
(11)

Every examples with $f(x,y) = \sum_{i} a_i x^{p^i} + \sum_{j} b_j y^{p^i}$ gives an equation in L and M of this type that has solution if and only if L = M = 0. Fixing a basis $\{\xi, \xi^p, \xi^{p^2}, \dots, \xi^{p^{h-1}}\}$ of \mathbb{F}_q over \mathbb{F}_p , every element $a \in \mathbb{F}_q$ can be written as $a = a_0\xi + a_1\xi^p + a_2\xi^{p^2} + \dots + a_{h-1}\xi^{p^{h-1}}$ for some $a_i \in \mathbb{F}_p$. Moreover, by putting $L_i := L^{p^i}$ and $M_i := M^{p^i}$, Equation (11) reads

$$L_0^2 + M_0\left(\sum_i a_i M_i + \sum_j b_j L_j\right) = 0.$$

In the projective space PG(2h-1,q) with homogeneous coordinates

 $(L_0, L_1, \ldots, L_h, M_0, M_1, \ldots, M_h)$ the previous equation gives a quadric that is always a cone with vertex a (2h-1)-dimensional subspace V given by:

$$V: L_0 = M_0 = \sum_i a_i M_i + \sum_j b_j L_j = 0$$
(12)

and projecting a non-degenerate conic $\Gamma: L_0^2 + a_{\overline{i}}M_0M_{\overline{i}} = 0$ in a plane π given by:

$$\pi: L_j = 0, j = 1, \dots, h - 1, M_i = 0, i = 1, \dots, h - 1 \text{ with } i \neq i.$$
(13)

Every example of ovoid of Q(4,q) gives a corresponding cone that is skew with the subgeometry $\Sigma = PG(2h - 1, p)$ of PG(2h - 1, q) given by

$$\Sigma = \{(a, a^p, \dots, a^{p^{h-1}}, b, b^p, \dots, b^{p^{h-1}})\}_{a,b \in \mathbb{F}_q} \text{ with } (a,b) \neq (0,0).$$
(14)

The condition that the cone is disjoint from the subgeometry Σ reads as follows: The following equation in x should have no solution unless x = 0 for every $(l_0, m_0) \neq (0, 0)$

$$\sum_{j} b_{j} m_{0} \ell_{0}^{p^{j}} x^{p^{j}} + \sum_{i} a_{i} m_{0}^{p^{i}+1} x^{p^{i}} + \ell_{0}^{2} x = 0.$$

From the previous arguments we obtain the following

Proposition 3.1. There exists an ovoid of Q(4,q), $q = p^h$, as in (1) with f(x,y) = $\sum_{i} a_{i}x^{p^{i}} + \sum_{j} b_{j}y^{p^{j}}$ if and only if in PG(2h-1,q) there exists a quadratic cone with vertex a (2h-4)-dimensional subspace V as in (12) and base a non-degenerate conic Γ as in (13) skew with the subgeometry $\Sigma \cong PG(2h-1,p)$ as in (14).

Finally, as a byproduct of the above proposition, we obtain two families of \mathbb{F}_3 -linearized permutation polynomials from the Pentilla-William ovoid of $Q(4,3^5)$ and from the Thas-Payne ovoids of $Q(4, 3^h)$, h > 2.

Proposition 3.2. Let $q = 3^5$, for every $(\ell_0, m_0) \in \mathbb{F}_q^2 \setminus \{(0, 0)\}$ the following are \mathbb{F}_3 -linearized permutation polynomials:

$$m_0\ell_0^{81}x^{81} + m_0^{10}x^9 - \ell_0^2x = 0.$$

Proposition 3.3. Let $q = 3^n$, n > 2 and let m be a fixed non-square in \mathbb{F}_q . For every $(\ell_0, m_0) \in \mathbb{F}_q^2 \setminus \{(0, 0)\}$ the following are \mathbb{F}_3 -linearized permutation polynomials:

$$(mm_0^2 - \ell_0^2)^9 x^9 + m_0^9 \ell_0^3 x^3 + (m^{-1})m_0^{10} x$$

Acknowledgments

The research of both Daniele Bartoli and Nicola Durante was supported by the Italian National Group for Algebraic and Geometric Structures and their Applications (GNSAGA - INdAM). We thank the anonymous reviewers for their careful reading of our manuscript and their comments and suggestions.

References

- A. Akbary, D. Ghioca, and Q. Wang. On permutation polynomials of prescribed shape. Finite Fields Appl., 15(2):195–206, 2009.
- [2] Y. Aubry, G. McGuire, F. Rodier. A few more functions that are not APN infinitely often, finite fields theory and applications. *Contemporary Math.* **518**, 23–31 (2010).
- [3] S. Ball. On ovoids of PG(3,q). Adv. Geom. 6(2), 177–198 (2006).
- [4] S. Ball, A. Blockhuis, M. Lavrauw. On the classification of semifield flocks. Adv. Math. 180(1), 104–111 (2003).
- [5] S. Ball, P. Govaerts, L. Storme. On ovoids of parabolic quadrics. Des. Codes Cryptogr. 38, 131–145 (2006).
- [6] D. Bartoli. Hasse-Weil type theorems and relevant classes of polynomial functions In: London Mathematical Society Lecture Note Series, Proceedings of 28th British Combinatorial Conference, Cambridge University Press, pp. 43–102 (2021).
- [7] D. Bartoli, Y. Zhou. Asymptotics of Moore exponent sets. J. Combin. Theory Ser. A 175, 105281 (2020).
- [8] E. Bombieri. Counting points on curves over finite fields (d'après S.A. Stepanov). Lecture Notes in Math. 383, 234–241 (1974).
- [9] P.J. Cameron, J.A. Thas, S.E. Payne. Polarities of generalized hexagons and perfect codes. *Geom. Dedicata* 5, 525–528 (1976).

- [10] A. Cafure, G. Matera. Improved explicit estimates on the number of solutions of equations over a finite field. *Finite Fields Appl.* **12**, 155–185 (2006).
- [11] F. Caullery, K.-U. Schmidt. On the classification of hyperovals. Adv. Math. 283, 195–203 (2015).
- [12] J. De Beule, A. Klein, K. Metsch. Substructures of finite classical polar spaces. In book: Current research topics in Galois geometry. NOVA Sci. Publ. Editors: Jan De Beule, Leo Storme, 2012.
- [13] L. E. Dickson. The analytic representation of substitutions on a power of a prime number of letters with a discussion of the linear group. Ann. of Math., 11(1-6):65–120, 1896/97.
- [14] N. Durante. Geometry of sesquilinear forms. Notes of a course given at the Summer School "Finite Geometry and friends." Bruxelles, June 2019. OnLine August 31th, 2019.
- [15] G. van der Geer. Counting curves over finite fields. *Finite Fields Appl.* 32, 207–232 (2015).
- [16] S. Ghorpade, G. Lachaud. Number of solutions of equations over finite fields and a conjecture of Lang and Weil. In: *Number Theory and Discrete Mathematics*, 269–291 (2002).
- [17] S. Ghorpade, G. Lachaud. Étale cohomology, Lefschetz theorem and number of points of singular varieties over finite fields. *Mosc. Math. J.* 2, 589–631 (2002).
- [18] A. Gunawardena, G.E. Moorhouse. The non-existence of ovoids in $O_9(q)$, European J. Combin. 18, 171–173 (1997).
- [19] R. Hartshorne. Algebraic Geometry. Graduate Texts in Mathematics, Springer-Verlag New York (1977).
- [20] C. Hermite. Sur les fonctions de sept lettres. Académie des sciences, 1863.
- [21] F. Hernando, G. McGuire. Proof of a conjecture of Segre and Bartocci on monomial hyperovals in projective planes. *Des. Codes Cryptogr.* 65, 275–289 (2012).
- [22] X.-d. Hou. Permutation polynomials over finite fields—a survey of recent advances. *Finite Fields Appl.*, 32:82–119, 2015.
- [23] W.M. Kantor. Ovoids and translation planes. Canadian J. Math. 34, 1195–1207 (1982).
- [24] S. Lang, A. Weil. Number of points of varieties in finite fields. Amer. J. Math. 76, 819–827 (1954).

- [25] M. Lavrauw. Semifield flocks, eggs and ovoids of Q(4,q). Adv. Geom. 5(3), 333–345 (2005).
- [26] M. Lavrauw. Sublines of prime order contained in the set of internal points of a conic. Des. Codes Cryptogr. 38(1), 113–123 (2006).
- [27] M. Lavrauw, M. Rodgers. Classification of 8-dimensional rank two commutative semifields. Adv. Geom. 19(1), 57–64 (2019).
- [28] R. Lidl, H. Niederreiter. Finite fields. Addison-Wesley, Reading, MA, (1983).
- [29] G. Lunardon. Normal spreads. Geom. Dedicata 75(3), 245–261 (1999).
- [30] G. L. Mullen, editor. Handbook of finite fields. Discrete Mathematics and its Applications (Boca Raton). CRC Press, Boca Raton, FL, 2013.
- [31] C.M. O'Keefe, J.A. Thas. Ovoids of the quadric Q(2n,q). European J. Combin. 16, 87–92 (1995).
- [32] T. Penttila, B. Williams. Ovoids of parabolic spaces. *Geom. Dedicata* 82, 1–19 (2000).
- [33] W. Schmidt. Equations over finite fields: an elementary approach. Lecture Notes in Math. 536, Springer, New York, 1976.
- [34] J.A. Thas Ovoidal Translation Planes. Archiv. Math. 23, 110–112 (1972).
- [35] J.A. Thas Polar spaces, generalized hexagons and perfect codes. J. Combin. Theory Ser. A. 29, 87–93 (1980).
- [36] J.A. Thas. Ovoids and spreads of finite classical polar spaces. Geom. Dedicata 10, 135–143 (1981).
- [37] J.A. Thas, S.E. Payne Spreads and ovoids in finite generalised quadrangles Geom. Dedicata 52, 227–253 (1994).
- [38] J. Tits. Les groupes simples de Suzuki et de Ree. In: Sem. Bourbaki (années 1960/61), exposés 205–222, Séminaire Bourbaki, no. 6 (1961), Exposé no. 210, 18 p.
- [39] Q. Wang. 15. polynomials over finite fields: an index approach. In Combinatorics and Finite Fields, pages 319–348. De Gruyter, 2019.
- [40] J. Zahid. Nonsingular points on hypersurfaces over \mathbb{F}_q . J. Math. Sci. 171(6), 731–735 (2010).