EXACTNESS OF LIMITS AND COLIMITS IN ABELIAN CATEGORIES REVISITED

ALEJANDRO ARGUDÍN-MONROY AND CARLOS E. PARRA

ABSTRACT. Let Σ be a small category and \mathcal{A} be a Σ -co-complete (resp. Σ complete) abelian category. It is a well-known fact that the category $\operatorname{Fun}(\Sigma, \mathcal{A})$ of functors of Σ in \mathcal{A} is an abelian category, and that the functor $\operatorname{colim}_{\Sigma}(-)$: $\operatorname{Fun}(\Sigma, \mathcal{A}) \to \mathcal{A}$ (resp. $\lim_{\Sigma}(-)$: $\operatorname{Fun}(\Sigma, \mathcal{A}) \to \mathcal{A}$) is left (resp. right) adjoint to $\kappa^{\Sigma} : \mathcal{A} \to \operatorname{Fun}(\Sigma, \mathcal{A})$, where κ^{Σ} is the associated constant diagram functor. In this paper we will show that the functor $\operatorname{colim}_{\Sigma}(-)$ (resp. $\lim_{\Sigma}(-)$) is exact if and only if the pair of functors ($\operatorname{colim}_{\Sigma}(-), \kappa^{\Sigma}$) (resp. $(\kappa^{\Sigma}, \lim_{\Sigma}(-))$) is Ext-adjoint. As an application of our findings, we will give new proofs of known results on the exactness of limits and colimits in abelian categories.

Contents

1.	Introduction	1
2.	Preliminaries	3
3.	A one-point extension of a small category	4
4.	Main results	6
References		13

1. INTRODUCTION

Let \mathcal{A} be an abelian category. Recall that \mathcal{A} is Ab3 if arbitrary coproducts exist; it is Ab4 if it is Ab3 and every coproduct of a set of monomorphisms is a monomorphism; and it is Ab5 if it is Ab3 and every colimit of a direct system of monomorphisms is a monomorphism. The corresponding dual notions are known as Ab3^{*}, Ab4^{*}, and Ab5^{*}. This classification of abelian categories was introduced by Grothendieck in [3] in order to axiomatise the constructions of the homological algebra coming from categories of modules.

Given $X, Y \in \mathcal{A}$, consider the 'big' group of extensions $\operatorname{Ext}^{1}_{\mathcal{A}}(X, Y)$ (see [5, Chapter VII]). To get a glimpse of the essence of these groups, recall that $\operatorname{Ext}^{1}_{\mathcal{A}}(X, Y)$ is made up of the equivalence classes of the short exact sequences of the form $Y \hookrightarrow E \twoheadrightarrow X$. So that, for every exact sequence $\epsilon : Y \hookrightarrow E \twoheadrightarrow X$ in \mathcal{A} , we will denote as $\overline{\epsilon}$ the extension in $\operatorname{Ext}^{1}_{\mathcal{A}}(X, Y)$ representing ϵ . Now, recall that morphisms in

²⁰²⁰ Mathematics Subject Classification. 18A25, 18A30, 18A35, 18A40, 18E10.

Key words and phrases. Abelian category, colimits, limits.

The first named author was supported by a postdoctoral fellowship from Programa de Desarrollo de las Ciencias Básicas, Ministerio de educación y cultura, Universidad de la República, Uruguay.

The second named author was supported by CONICYT/FONDECYT/REGULAR/1200090.

 \mathcal{A} induce morphisms between extension groups via pullback and pushout. Namely, every $f \in \operatorname{Hom}_{\mathcal{A}}(X', X)$ induces the additive maps $\operatorname{Ext}^{1}_{\mathcal{A}}(f, Y) : \operatorname{Ext}^{1}_{\mathcal{A}}(X, Y) \to$ $\operatorname{Ext}^{1}_{\mathcal{A}}(X', Y), \overline{\eta} \mapsto \overline{\eta} \cdot f$, and $\operatorname{Ext}^{1}_{\mathcal{A}}(Y, f) : \operatorname{Ext}^{1}_{\mathcal{A}}(Y, X') \to \operatorname{Ext}^{1}_{\mathcal{A}}(Y, X), \overline{\eta} \mapsto f \cdot \overline{\eta}$, for every $Y \in \mathcal{A}$.

Now, for every $A \in \mathcal{A}$ and for every coproduct $\coprod_{i \in X} B_i$ (resp. product $\prod_{i \in X} B_i$) in \mathcal{A} , consider the map $\Psi : \operatorname{Ext}^1_{\mathcal{A}}(\coprod_{i \in X} B_i, A) \to \prod_{i \in X} \operatorname{Ext}^1_{\mathcal{A}}(B_i, A), \overline{\eta} \mapsto (\overline{\eta} \cdot \mu_i)_{i \in X}$ (resp. $\Phi : \operatorname{Ext}^1_{\mathcal{A}}(A, \prod_{i \in X} B_i) \to \prod_{i \in X} \operatorname{Ext}^1_{\mathcal{A}}(A, B_i), \overline{\eta} \mapsto (\pi_i \cdot \overline{\eta})_{i \in X}$), where $\mu_i : B_i \to \coprod_{i \in X} B_i$ is the *i*-th canonical inclusion (resp. $\pi_i : \prod_{i \in X} B_i \to B_i$ the *i*-th projection). In [2], it was shown that the Ab4 (resp. Ab4*) categories are exactly the Ab3 (resp. Ab3*) categories where the maps Ψ (resp. Φ) are bijective. A key point of this work is that, for every set of exact sequences $H = \left\{\eta_i : A \stackrel{f_i}{\hookrightarrow} E_i \twoheadrightarrow B_i\right\}_{i \in X}$

(resp. $H = \left\{ \eta_i : B_i \hookrightarrow E_i \xrightarrow{f_i} A \right\}_{i \in X}$), it can be built an exact sequence

$$\Theta(H): A \to E \twoheadrightarrow \prod_{i \in X} A_i \qquad (\text{resp. } \Theta(H): \prod_{i \in X} B_i \hookrightarrow E \to A)$$

where E is the colimit (resp. limit) of a diagram defined through the family of morphisms $\{f_i\}_{i \in X}$ (see [2, Section 4]). It turns out that, under the right conditions, the inverse map of Ψ (resp. Φ) can be exhibited using Θ .

Let Σ be a small category and Fun (Σ, \mathcal{A}) be the category of functors $\Sigma \to \mathcal{A}$. Recall that \mathcal{A} is Σ -co-complete (resp. Σ -complete) if every $F \in \text{Fun}(\Sigma, \mathcal{A})$ admits a colimit (resp. limit) in \mathcal{A} . In this context it is immediate to see that one has an adjoint pair $(\operatorname{colim}_{\Sigma}(-), \kappa^{\Sigma})$ (resp. $(\kappa^{\Sigma}, \lim_{\Sigma}(-))$), where $\operatorname{colim}_{\Sigma}(-) : \operatorname{Fun}(\Sigma, \mathcal{A}) \to \mathcal{A}$ (resp. $\lim_{\Sigma}(-) : \operatorname{Fun}(\Sigma, \mathcal{A}) \to \mathcal{A}$) is the colimit (resp. limit) functor, and where $\kappa^{\Sigma} : \mathcal{A} \to \operatorname{Fun}(\Sigma, \mathcal{A})$ is the functor that assigns to each $A \in \mathcal{A}$ a constant functor in $\operatorname{Fun}(\Sigma, \mathcal{A})$.

In this article, motivated mainly by the sequence Θ , we will revisit the techniques used in [2] to study the exactness of Σ -colimits (resp. Σ -limits) in Σ -co-complete (resp. Σ -complete) abelian categories, for all small category Σ . Namely, we will consider the canonical map

$$\Psi_{F,A}^{\Sigma} : \operatorname{Ext}_{\mathcal{A}}^{1}(\operatorname{colim}_{\Sigma}(F), A) \to \operatorname{Ext}_{\operatorname{Fun}(\Sigma, \mathcal{A})}^{1}(F, \kappa^{\Sigma}(A))$$

(resp. $\Phi_{F,A}^{\Sigma} : \operatorname{Ext}_{\mathcal{A}}^{1}(A, \operatorname{lim}_{\Sigma}(F)) \to \operatorname{Ext}_{\operatorname{Fun}(\Sigma, \mathcal{A})}^{1}(\kappa^{\Sigma}(A), F)$)

for every $A \in \mathcal{A}$ and $F \in \operatorname{Fun}(\Sigma, \mathcal{A})$ (see Remark 4.1 and Corollary 4.4). Looking for sufficient conditions for $\Psi_{F,A}^{\Sigma}$ (resp. $\Phi_{F,A}^{\Sigma}$) to be bijective, we will see that every exact sequence $\eta : \kappa^{\Sigma}(A) \hookrightarrow E \twoheadrightarrow F$ (resp. $\eta : F \hookrightarrow E \twoheadrightarrow \kappa^{\Sigma}(A)$) in $\operatorname{Fun}(\Sigma, \mathcal{A})$ induces an exact sequence $\kappa^{\Sigma_*}(A) \xrightarrow{f_{\eta}} E_{\eta} \twoheadrightarrow F'$ (resp. $F' \hookrightarrow E_{\eta} \xrightarrow{f_{\eta}} \kappa^{\Sigma_*}(A)$) in $\operatorname{Fun}(\Sigma_*, \mathcal{A})$, where Σ_* is the one-point extension of Σ that adds a source (resp. sink) point * (see Section 3), and F' is a functor such that $F'|_{\Sigma} = F$ and F'(*) = 0.

Our main result will be that, in a Σ -co-complete (resp. Σ -complete) category \mathcal{A} , $\operatorname{colim}_{\Sigma}(-)$ (resp. $\lim_{\Sigma}(-)$) is exact if and only if f_{η} is a monomorphism (resp. epimorphism) $\forall \overline{\eta} \in \operatorname{Ext}^{1}_{\operatorname{Fun}(\Sigma,\mathcal{A})}(\kappa^{\Sigma}(A),F)$ (resp. $\forall \overline{\eta} \in \operatorname{Ext}^{1}_{\operatorname{Fun}(\Sigma,\mathcal{A})}(F,\kappa^{\Sigma}(A)))$), which in turn will be equivalent to $\Psi^{\Sigma}_{F,A}$ (resp. $\Phi^{\Sigma}_{F,A}$) being bijective $\forall A \in \mathcal{A}$ and $\forall F \in \operatorname{Fun}(\Sigma,\mathcal{A})$. Moreover, in such case, it can be seen that $\Psi^{\Sigma}_{-,?}$ (resp. $\Phi^{\Sigma}_{-,?}$) is a natural isomorphism between the respective functors. This phenomenon is what

we call an Ext-adjoint pair. An application of our results consists in reproducing with new proofs the following known results in abelian categories.

- [6, Corollary 3.2.9] Let A be an Ab3 (resp. Ab3*) abelian category. Then, A is Ab4 (resp. Ab4*) whenever A has enough injectives (resp. projectives).
- (2) [2, Theorems 4.9 and 4.11] Let A be an Ab3 (resp. Ab3*) abelian category. Then, A is Ab4 (resp. Ab4*) if, and only if, the canonical map

$$\operatorname{Ext}^{1}_{\mathcal{A}}(\coprod_{i\in I} A_{i}, A) \to \prod_{i\in I} \operatorname{Ext}^{1}_{\mathcal{A}}(A_{i}, A)$$

$$\left(\operatorname{resp.} \operatorname{Ext}^{1}_{\mathcal{B}}(B, \prod_{i\in I} B_{i}) \to \prod_{i\in I} \operatorname{Ext}^{1}_{\mathcal{B}}(B, B_{i})\right)$$

is always an isomorphism of 'big' abelian groups.

The article is organised as follows. In Section 2 we recall the preliminaries on abelian categories and (co)limits in abelian categories that we will need throughout the article. In Section 3 we show how the one-point extension of a small category Σ is related to the behaviour of the colimits on the category Σ . Finally, Section 4 contains our main results.

2. Preliminaries

In this section we recall some definitions and results, and we fix some notation that we will use throughout the paper. For more precise details, the reader is referred to [6, 7, 4].

2.1. Limits and colimits. Recall that a category is small when the isomorphism classes of its objects form a set. In this sense, every set Σ can be viewed as a small category whose objects are the elements of Σ and the only morphisms are the identity morphisms. Also, if Σ is a directed (resp. codirected) set, then Σ can be viewed as a small category whose objects are the elements of Σ and there is a unique morphism $\alpha \to \beta$ exactly when $\alpha \leq \beta$ (resp. $\beta \leq \alpha$).

In what follows Σ will denote a small category. If C is a category, then a functor $\Sigma \to C$ will be called a Σ -diagram on C. In this setting, the Σ -diagrams on C together with the respective natural transformations form a category, which will be denoted by Fun(Σ, C). In particular, if C is an abelian category, then Fun(Σ, C) is an abelian category in the obvious way (see [7, Chapter IV, Section 7]).

Observe that the assignment $C \mapsto \kappa^{\Sigma}(C)$ gives a functor $\kappa^{\Sigma} : \mathcal{C} \to \operatorname{Fun}(\Sigma, \mathcal{C})$, where $\kappa^{\Sigma}(C)$ is the Σ -diagram on \mathcal{C} such that $i \mapsto C$ and $\lambda \mapsto 1_C$, for each object iand morphism λ of Σ ; and, for every morphism $f : A \to B$ in $\mathcal{C}, \kappa^{\Sigma}(f) : \kappa^{\Sigma}(A) \to \kappa^{\Sigma}(B)$ is the natural transformation κ_f^{Σ} given by the family of morphisms $(\kappa_{f,s}^{\Sigma})_{s \in \Sigma}$ with $\kappa_{f,s}^{\Sigma} = f$ for all $s \in \Sigma$. Such functor is called the **constant diagram functor**.

Let $F \in \operatorname{Fun}(\Sigma, \mathcal{C})$, $C, L \in \mathcal{C}$, and $\rho : F \to \kappa^{\Sigma}(C)$, $\varrho : \kappa^{\Sigma}(L) \to F$ be natural transformations. Then, we will say that the pair (C, ρ) (resp. (L, ϱ)) is a Σ -colimit (resp. Σ -limit) of F when the following condition holds: for each natural transformation $\tau : F \to \kappa^{\Sigma}(D)$ (resp. $\tau : \kappa^{\Sigma}(D) \to F$), where D is an object in \mathcal{C} , there is a unique morphism $f : C \to D$ (resp. $f : D \to L$) such that the following diagram commutes.



If (C, ρ) (resp. (L, ϱ)) is a Σ -colimit (resp. Σ -limit) of F, we use the notation $C = \operatorname{colim}_{\Sigma}(F)$ (resp. $L = \lim_{\Sigma}(F)$). Now, if each Σ -diagram on C has a Σ -colimit (resp. Σ -limit), we say that C admits Σ -colimits (resp. Σ -limits) or that it is Σ -co-complete (resp. Σ -complete). In this case, the assignment $F \mapsto \operatorname{colim}_{\Sigma}(F)$ (resp. $F \mapsto \lim_{\Sigma}(F)$) gives rise to a functor $\operatorname{colim}_{\Sigma} : \operatorname{Fun}(\Sigma, C) \to C$ (resp. $\lim_{\Sigma} : \operatorname{Fun}(\Sigma, C) \to C$) which is the left (resp. right) adjoint to the associated constant diagram functor. The category C is called co-complete (resp. complete) when it admits Σ -colimits (resp. Σ -limits), for every small category Σ . In this context, for every functor $F : \Sigma \to C$, we denote by $C_F := \operatorname{colim}_{\Sigma}(F)$ (resp. $L_F := \lim_{\Sigma}(F)$) and by $\rho^F : F \to \kappa^{\Sigma}(C_F)$ (resp. $\varrho^F : \kappa^{\Sigma}(L_F) \to F$) the associated natural transformation.

Lastly, if \mathcal{C} admits Σ -colimits (resp. Σ -limits), observe that for each $A \in \mathcal{A}$ there is a unique morphism $\nabla^A : C_{\kappa^{\Sigma}(A)} \to A$ (resp. $\Delta^A : A \to L_{\kappa^{\Sigma}(A)}$) such that $\kappa_{\nabla^A}^{\Sigma} \circ \rho^{\kappa^{\Sigma}(A)} = \mathbb{1}_{\kappa^{\Sigma}(A)}$ (resp. $\varrho^{\kappa^{\Sigma}(A)} \circ \kappa_{\Delta^A}^{\Sigma} = \mathbb{1}_{\kappa^{\Sigma}(A)}$). We refer to such morphism as the **co-diagonal morphism** (resp. **diagonal morphism**).

2.2. Abelian categories. In what follows \mathcal{A} will denote an abelian category. Recall the following hierarchy among abelian categories (introduced by Grothendieck in [3]): we say that \mathcal{A} is

- Ab3 if all set-indexed coproducts exist in \mathcal{A} (equivalently, if it is co-complete);
- Ab4 if it is Ab3 and the functors colim_Σ are exact, for each set Σ viewed as a small category;
- Ab5 if it is Ab3 and the functors colim_Σ are exact, for each directed set Σ viewed as a small category.

We will denote by **Abn*** to the dual definition of **Abn** for each n in $\{3, 4, 5\}$.

3. A ONE-POINT EXTENSION OF A SMALL CATEGORY

In this section, Σ will denote a small category and \mathcal{A} will denote a Σ -co-complete abelian category. Now, from Σ we can construct a new category Σ_* which essentially consists of adding a source point to Σ . That is, Σ_* is the category consisting of the objects of Σ plus a new object *, and the morphisms of Σ_* are the morphisms of Σ plus a single morphism from * to any other point of Σ_* . More precisely, we define the category Σ_* as follows:

- $Ob(\Sigma_*) := Ob(\Sigma) \cup \{*\};$
- Let $(\alpha_i)_{i \in Ob(\Sigma)}$ be an $Ob(\Sigma)$ -indexed family of symbols, where $\alpha_i = \alpha_j$ with $i, j \in Ob(\Sigma)$, when *i* is isomorphic to *j* in Σ . In this setting, we put $\Sigma_*(i, *) := \emptyset$, $\Sigma_*(*, i) := \{\alpha_i\}$, $\Sigma_*(*, *) := \{1_*\}$ and $\Sigma_*(i, j) = \Sigma(i, j)$ for every *i* and *j* objects in Σ .

• The composition is given by the extension of the composition in the category Σ following the rule: $1_* \circ 1_* := 1_*$, $\alpha_i \circ 1_* := \alpha_i$ and $\lambda \circ \alpha_i := \alpha_j$, for all i, j and $\lambda : i \to j$, objects and morphism in Σ , respectively.

3.1. A one-point extension of a Σ -diagram. Let $\eta : \kappa^{\Sigma}(A) \stackrel{\phi}{\hookrightarrow} F \stackrel{\psi}{\twoheadrightarrow} G$ be an exact sequence in Fun (Σ, \mathcal{A}) . Define the functor $F_{\eta} : \Sigma_* \to \mathcal{A}$ as follows:

- For each $i \in Ob(\Sigma)$ we assign the object F(i), and to the object * we assign the object A.
- For each morphism λ in Σ we assign the morphism $F(\lambda)$ and, for each α_i we assign the morphism ϕ_i . Lastly, for the morphism 1_* we assign the morphism 1_A .

It can be shown that there is an exact sequence $\eta' : \kappa^{\Sigma_*}(A) \stackrel{\phi'}{\hookrightarrow} F_{\eta} \stackrel{\psi'}{\twoheadrightarrow} G'$ in $\operatorname{Fun}(\Sigma_*, \mathcal{A})$, where G' is a functor that extends G such that G'(*) = 0. In order to find the colimit of such exact sequence, we proceed as follows.

By the right exactness of the functor $\operatorname{colim}_{\Sigma}$: Fun $(\Sigma, \mathcal{A}) \to \mathcal{A}$ (see [7, Chapter V, Section 1]), there is an exact sequence in \mathcal{A} :

$$C_{\kappa^{\Sigma}(A)} \xrightarrow{\operatorname{colim}_{\Sigma}(\phi)} C_{F} \xrightarrow{\operatorname{colim}_{\Sigma}(\psi)} C_{G} \xrightarrow{} 0.$$

Consider the following commutative diagram in \mathcal{A} with exact rows, where the left square is the pushout of $\operatorname{colim}_{\Sigma}(\phi)$ and the co-diagonal morphism.

$$\begin{array}{c|c} C_{\kappa^{\Sigma}(A)} & \xrightarrow{\operatorname{colim}_{\Sigma}(\phi)} & C_{F} & \xrightarrow{\operatorname{colim}_{\Sigma}(\psi)} & C_{G} \longrightarrow 0 \\ \hline & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & \\ & & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & \\ & & & \\ & & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & &$$

Lastly, consider the morphism $\xi^{\eta}: F_{\eta} \to \kappa^{\Sigma_*}(Z_{\eta})$ in Fun (Σ_*, \mathcal{A}) given by $\xi^{\eta}_i := \mu_{\eta} \circ \rho^F_i$ for each $i \in Ob(\Sigma)$ and $\xi^{\eta}_* := f_{\eta}$. Notice that ξ^{η} is well-defined since $\mu_{\eta} \circ \rho^F_j \circ F_{\eta}(\lambda) = \mu_{\eta} \circ \rho^F_j \circ F(\lambda) = \mu_{\eta} \circ \rho^F_i$ for each $\lambda : i \to j$ morphism in Σ , and the following diagram in \mathcal{A} is commutative for all $i \in Ob(\Sigma)$:



Indeed, such claim follows from the following equalities of morphisms.

$$\mu_{\eta} \circ (\rho_i^F \circ \phi_i) = \mu_{\eta} \circ (\operatorname{colim}_{\Sigma}(\phi) \circ \rho_i^{\kappa^{\Sigma}(A)}) \quad \text{(by definition of } \operatorname{colim}_{\Sigma}(\phi))$$
$$= (f_{\eta} \circ \nabla^A) \circ \rho_i^{\kappa^{\Sigma}(A)} \qquad (By (1.1))$$
$$= f_{\eta} \circ 1_A = f_{\eta} = 1_{Z_{\eta}} \circ f_{\eta}$$

The following result shows that the morphism f_{η} coincide with the left morphism in the sequence $\Theta(\eta)$ described in the introduction (see [2]) when Σ is a set. **Lemma 3.1.** Let $\eta : \kappa^{\Sigma}(A) \xrightarrow{\phi} F \xrightarrow{\psi} G$ be an exact sequence in Fun (Σ, \mathcal{A}) , where Σ is a small category and A is an object in \mathcal{A} , and let Z_{η} be as in the diagram (1.1). Then, the pair (Z_{η}, ξ^{η}) is the Σ_{*} -colimit of F_{η} . In particular, $Z_{\eta} = \operatorname{colim}_{\Sigma_{*}}(F_{\eta})$.

Proof. Let C be an object in \mathcal{A} and let $\varphi : F_{\eta} \to \kappa^{\Sigma_*}(C)$ be a natural transformation in Fun (Σ_*, \mathcal{A}) . By definition, we have that $\varphi_j \circ F(\lambda) = \varphi_j \circ F_{\eta}(\lambda) = \varphi_i$, for every morphism $\lambda : i \to j$ in Σ , and the following commutative diagram in \mathcal{A} , for all $i \in I$:

In particular, we get that the family of morphisms $(\varphi_i)_{i \in Ob(\Sigma)} := \varphi|_{Ob(\Sigma)}$ is a natural transformation in Fun (Σ, \mathcal{A}) . Now, from the universal property of colimits, we obtain a unique morphism $\pi : C_F \to C$ in \mathcal{A} such that the equality $\kappa_{\pi}^{\Sigma} \circ \rho^F = \varphi|_{Ob(\Sigma)}$ holds in Fun (Σ, \mathcal{A}) . In particular, we get

$$\begin{aligned} (\varphi_* \circ \nabla^A) \circ \rho_i^{\kappa^{\Sigma}(A)} &= \varphi_* \circ 1_A \\ &= \varphi_i \circ \phi_i \\ &= (\pi \circ \rho_i^F) \circ \phi_i \\ &= (\pi \circ \operatorname{colim}_{\Sigma}(\phi)) \circ \rho_i^{\kappa^{\Sigma}(A)} \end{aligned}$$

for all $i \in Ob(\Sigma)$. Using once again the universal property of colimits we deduce that $\varphi_* \circ \nabla^A = \pi \circ \operatorname{colim}_{\Sigma}(\phi)$. Now, from the universal property of pushouts, we know that there is a unique morphism $\sigma : Z_{\eta} \to C$ such that $\sigma \circ f_{\eta} = \varphi_*$ and $\sigma \circ \mu_{\eta} = \pi$. And hence, it follows that the natural transformation $\kappa_{\sigma}^{\Sigma_*}$ satisfy the following equation in the category $\operatorname{Fun}(\Sigma_*, \mathcal{A})$:

$$\kappa_{\sigma}^{\Sigma_*} \circ \xi^{\eta} = \varphi$$

It remains to prove the uniqueness of σ . Let $\sigma': Z_{\eta} \to C$ be a morphism in \mathcal{A} such that $\kappa_{\sigma'}^{\Sigma_*} \circ \xi^{\eta} = \varphi$. Then, $\varphi_* = (\kappa_{\sigma'}^{\Sigma_*})_* \circ \xi^{\eta}_* = \sigma' \circ f_{\eta}$ and $\varphi_i = (\kappa_{\sigma'}^{\Sigma_*})_i \circ \xi^{\eta}_i = \sigma' \circ (\mu_{\eta} \circ \rho_i^F) = (\sigma' \circ \mu_{\eta}) \circ \rho_i^F$, for all $i \in Ob(\Sigma)$. Thus, $\kappa_{\sigma' \circ \mu_{\eta}}^{\Sigma} \circ \rho^F = \varphi|_{Ob(\Sigma)}$. And hence, from the universal property of colimits, $\sigma' \circ \mu_{\eta} = \pi$. Finally, by the universal property of pushouts, we have $\sigma = \sigma'$ as desired.

4. Main results

In this section we will use the previous results to obtain a characterisation for the exactness of the functors $\operatorname{colim}_{\Sigma}(-)$: Fun $(\Sigma, \mathcal{A}) \to \mathcal{A}$ and $\lim_{\Sigma}(-)$: Fun $(\Sigma, \mathcal{B}) \to \mathcal{B}$, when \mathcal{A} and \mathcal{B} are Σ -co-complete and Σ -complete abelian categories, respectively, for all small category Σ .

Theorem 4.1. Let Σ be a small category and let \mathcal{A} be a Σ -co-complete abelian category. Then, the following conditions are equivalent:

- (1) the functor $\operatorname{colim}_{\Sigma} : \operatorname{Fun}(\Sigma, \mathcal{A}) \to \mathcal{A}$ is exact;
- (2) f_{η} is a monomorphism, for every exact sequence $\eta : \kappa^{\Sigma}(A) \stackrel{\phi}{\hookrightarrow} F \stackrel{\psi}{\twoheadrightarrow} G$ in Fun (Σ, \mathcal{A}) , where A is an object in \mathcal{A} .

Proof. (a) \Rightarrow (b). It is clear by properties of the pushouts (see (1.1)).

(b) \Rightarrow (a). Let $H \stackrel{\zeta}{\hookrightarrow} N \stackrel{\chi}{\twoheadrightarrow} G$ be an exact sequence in Fun (Σ, \mathcal{A}) . Consider the following commutative diagram in Fun (Σ, \mathcal{A}) with exact rows, where the left square is the pushout of ζ and ρ^H

$$\begin{array}{cccc} 0 & \longrightarrow H & \xrightarrow{\zeta} & N & \xrightarrow{\chi} & G & \longrightarrow & 0 \\ & & & & & & & & \\ \rho^{H} & & & & & & & \\ \rho^{H} & & & & & & & & \\ & & & & & & & & & \\ 0 & \longrightarrow & \kappa^{\Sigma}(C_{H}) & \xrightarrow{\phi} & F & \xrightarrow{\psi} & G & \longrightarrow & 0 \end{array}$$

From the universal property of pushouts, we obtain a unique natural transformation $v: F \to \kappa^{\Sigma}(C_N)$ such that $v \circ \omega = \rho^N$ and $v \circ \phi = \kappa^{\Sigma}_{\operatorname{colim}_{\Sigma}(\zeta)}$ since $\rho^N \circ \zeta = \kappa^{\Sigma}_{\operatorname{colim}_{\Sigma}(\zeta)} \circ \rho^H$. Note that there is a unique morphism $\gamma: C_F \to C_N$ such that $\kappa^{\Sigma}_{\gamma} \circ \rho^F = v$ by the universal property of colimits.

Now, set $\eta := \kappa^{\Sigma}(C_H) \xrightarrow{\phi} F \xrightarrow{\psi} G$ and, for each $i \in Ob(\Sigma)$, consider the following commutative diagram in \mathcal{A} with exact rows, where the upper and lower left squares are pushouts (here f_{η} is a monomorphism by the assumption (b)):



In particular, we have the following equalities

$$\begin{aligned} (\gamma \circ \operatorname{colim}_{\Sigma}(\phi)) \circ \rho_{i}^{\kappa^{\Sigma}(C_{H})} &= \gamma \circ (\rho_{i}^{F} \circ \phi_{i}) \\ &= \upsilon_{i} \circ \phi_{i} \\ &= \operatorname{colim}_{\Sigma}(\zeta) \\ &= \operatorname{colim}_{\Sigma}(\zeta) \circ 1_{C_{H}} \\ &= (\operatorname{colim}_{\Sigma}(\zeta) \circ \nabla^{C_{H}}) \circ \rho_{i}^{\kappa^{\Sigma}(C_{H})} \end{aligned}$$

for all $i \in Ob(\Sigma)$. So that $\kappa_{\gamma \circ \operatorname{colim}_{\Sigma}(\phi)}^{\Sigma} \circ \rho^{\kappa^{\Sigma}(C_{H})} = \kappa_{\operatorname{colim}_{\Sigma}(\zeta) \circ \nabla^{C_{H}}}^{\Sigma} \circ \rho^{\kappa^{\Sigma}(C_{H})}$, and hence $\operatorname{colim}_{\Sigma}(\zeta) \circ \nabla^{C_{H}} = \gamma \circ \operatorname{colim}_{\Sigma}(\phi)$. Therefore, there is a unique morphism $\theta : Z_{\eta} \to C_{N}$ such that $\theta \circ \mu_{\eta} = \gamma$ and $\theta \circ f_{\eta} = \operatorname{colim}_{\Sigma}(\zeta)$. We claim that the pair $(Z_{\eta}, \kappa_{\mu_{\eta}}^{\Sigma} \circ (\rho^{F} \circ \omega) : N \to \kappa^{\Sigma}(Z_{\eta}))$ is a colimit of N. Indeed, by the universal property of colimits, there is a unique morphism $m : C_{N} \to Z_{\eta}$ in \mathcal{A} such that $\kappa_{m}^{\Sigma} \circ \rho^{N} = \kappa_{\mu_{\eta}}^{\Sigma} \circ (\rho^{F} \circ \omega)$. Therefore, the claim holds once we show that m is an isomorphism in \mathcal{A} . For this, observe that, on the one hand:

$$\begin{aligned} (\theta \circ m) \circ \rho_i^N &= \theta \circ (\mu_\eta \circ (\rho_i^F \circ \omega_i)) \\ &= \gamma \circ (\rho_i^F \circ \omega_i) \\ &= \upsilon_i \circ \omega_i \\ &= \rho_i^N = \mathbf{1}_{C_N} \circ \rho_i^N \end{aligned}$$

for all $i \in Ob(\Sigma)$. And hence, $\theta \circ m = 1_{C_N}$ by the universal property of colimits. On the other hand, from the above diagram we have that

$$f_{\eta} = f_{\eta} \circ 1_{C_H} = f_{\eta} \circ \nabla^{C_H} \circ \rho_i^{\kappa^{\Sigma}(C_H)} = \mu_{\eta} \circ \rho_i^F \circ \phi_i,$$

for all $i \in Ob(\Sigma)$. In particular, for every $i \in Ob(\Sigma)$, we get $f_{\eta} \circ \rho_i^H = \mu_{\eta} \circ \rho_i^F \circ \phi_i \circ \rho_i^H = \mu_{\eta} \circ \rho_i^F \circ \omega_i \circ \zeta_i = m \circ \rho_i^N \circ \zeta_i = m \circ \operatorname{colim}_{\Sigma}(\zeta) \circ \rho_i^H$. Therefore, $f_{\eta} = m \circ \operatorname{colim}_{\Sigma}(\zeta) = \mu_{\eta} \circ \rho_i^F \circ \phi_i$ for all $i \in Ob(\Sigma)$. Now, for each $i \in Ob(\Sigma)$, we get the following commutative diagram in \mathcal{A} (recall that $\mu_{\eta} \circ (\rho_i^F \circ \omega_i) = m \circ \rho_i^N$):



From the universal property of the upper pushout in the diagram, we obtain that $\mu_{\eta} \circ \rho_i^F = m \circ v_i$, for all $i \in Ob(\Sigma)$. But, each v_i coincide with the composition $\gamma \circ \rho_i^F$, and hence $\mu_{\eta} = m \circ \gamma$. Now, notice that the composition $m \circ \theta$ and $1_{Z_{\eta}}$ complete the following diagram in \mathcal{A} and, therefore, such morphisms coincide and our claim holds (since $m \circ \theta = 1_{Z_{\eta}}$ and $\theta \circ m = 1_{C_N}$).



Finally, from the equality $\theta \circ f_{\eta} = \operatorname{colim}_{\Sigma}(\zeta)$ we get that the morphism $\operatorname{colim}_{\Sigma}(\zeta)$ is a monomorphism as desired.

Remark 4.1. In the setting of the previous theorem, we get that for each $F \in$ Fun (Σ, \mathcal{A}) and $A \in Ob(\mathcal{A})$ there is a natural map

$$\Psi_{F,A}^{\Sigma} : Ext^{1}_{\mathcal{A}}(C_{F}, A) \to Ext^{1}_{\operatorname{Fun}(\Sigma, \mathcal{A})}(F, \kappa^{\Sigma}(A))$$

both in A and in F, defined as $\Psi_{F,A}^{\Sigma}(\overline{\epsilon}) := \overline{\kappa^{\Sigma}(\epsilon) \cdot \rho^{F}}$, where $\kappa^{\Sigma}(\epsilon) \cdot \rho^{F}$ denote the upper exact sequence in the following commutative diagram in Fun(Σ, \mathcal{A}) with exact

9

rows (here the right square is a pullback of the respective morphisms):



Lemma 4.2. Let Σ be a small category and let \mathcal{A} be a Σ -co-complete abelian category. Then, the map $\Psi_{F,A}^{\Sigma}$ is injective, for all $F \in \operatorname{Fun}(\Sigma, \mathcal{A})$ and $A \in Ob(\mathcal{A})$.

Proof. Let ϵ : $A \xrightarrow{f} B \xrightarrow{g} C_F$ be an extension in \mathcal{A} such that $\Psi_{F,A}^{\Sigma}(\epsilon)$ is the trivial extension in $\operatorname{Ext}^{1}_{\operatorname{Fun}(\Sigma,\mathcal{A})}(\kappa^{\Sigma}(A), F)$. Thus, there is a natural transformation $\tau: F \to \kappa^{\Sigma}(B)$ such that $\rho^F = \kappa^{\Sigma}(g) \circ \tau$. Now, from the universal property of colimits, we obtain a unique morphism $h: C_F \to B$ such that $\tau = \kappa_h^{\Sigma} \circ \rho^{\vec{F}}$. Using once again the universal property of colimits, we deduce that $g \circ h = 1_{C_F}$ and hence $\overline{\epsilon}$ is the trivial extension in $\operatorname{Ext}^{1}_{\mathcal{A}}(C_{F}, A)$. \square

The following is the second main result of the paper.

Theorem 4.2. Let Σ be a small category and let \mathcal{A} be a Σ -co-complete abelian category. Then, the following conditions are equivalent:

- (1) the functor $\operatorname{colim}_{\Sigma} : \operatorname{Fun}(\Sigma, \mathcal{A}) \to \mathcal{A}$ is exact;
- (2) the map $\Psi_{F,A}^{\Sigma}$ is bijective, for every $F \in \operatorname{Fun}(\Sigma, \mathcal{A})$ and $A \in Ob(\mathcal{A})$; (3) the bifunctors $\operatorname{Ext}^{1}_{\mathcal{A}}(\operatorname{colim}_{\Sigma}(?), -)$ and $\operatorname{Ext}^{1}_{\operatorname{Fun}(\Sigma, \mathcal{A})}(?, \kappa^{\Sigma}(-))$ are naturally isomorphic in the obvious way.

Proof. (b) \Leftrightarrow (c) Is clear by definition of the maps $\Psi_{F,A}^{\Sigma}$ (see Remark 4.1).

(a) \Rightarrow (b) Let F be a functor in Fun (Σ, \mathcal{A}) and let A be an object in \mathcal{A} . By Lemma 4.2 our task is reduced to check that $\Psi_{F,A}^{\Sigma}$ is surjective. For this, consider an arbitrary exact sequence $\eta : \kappa^{\Sigma}(A) \xrightarrow{\phi} G \xrightarrow{\psi} F$ in Fun (Σ, \mathcal{A}) . Now, using the exactness of the functor $\operatorname{colim}_{\Sigma}$, we obtain the following commutative diagram in \mathcal{A} with exact rows, where the left square is a pushout (see (1.1)).

$$\begin{array}{c|c} 0 \longrightarrow C_{\kappa^{\Sigma}(A)} \xrightarrow{\operatorname{colim}_{\Sigma}(\phi)} C_{G} \xrightarrow{\operatorname{colim}_{\Sigma}(\psi)} C_{F} \longrightarrow 0 \\ & & & \\ & &$$

Let ϵ be the lower exact sequence in the above diagram. We claim that $\Psi_{F,A}^{\Sigma}(\overline{\epsilon}) = \overline{\eta}$. To show this, notice that we have the following commutative diagram in Fun (Σ, \mathcal{A}) :

Using the fact that $\kappa_{\nabla^A}^{\Sigma} \circ \rho^{\kappa^{\Sigma}(A)} = \mathbb{1}_{\kappa^{\Sigma}(A)}$, we obtain that the right outer rectangle in the above diagram is a pullback (see [6, dual of Lemma 5.2, p.35]). And, hence $\Psi_{F,A}^{\Sigma}(\bar{\epsilon}) = \bar{\eta}$.

(b) \Rightarrow (a) By Theorem 4.1, it is enough to check that f_{η} is a monomorphism, for every exact sequence $\eta : \kappa^{\Sigma}(A) \xrightarrow{\phi} G \xrightarrow{\psi} F$ in Fun (Σ, \mathcal{A}) , where A is an object in \mathcal{A} . Observe that, by hypothesis, there exists an exact sequence $\epsilon : A \xrightarrow{f} B \xrightarrow{g} C_F$ in \mathcal{A} such that $\Psi_{F,A}^{\Sigma}(\overline{\epsilon}) = \overline{\eta}$. So that we have the following commutative diagram in Fun (Σ, \mathcal{A}) .

$$\begin{split} \Psi^{\Sigma}_{F,A}(\epsilon): & 0 \longrightarrow \kappa^{\Sigma}(A) \xrightarrow{\phi} G \xrightarrow{\psi} F \longrightarrow 0 \\ & & \downarrow^{\mathfrak{o}} & \downarrow^{\rho^{F}} \\ & & \kappa^{\Sigma}(\epsilon): & 0 \longrightarrow \kappa^{\Sigma}(A) \xrightarrow{\kappa^{\Sigma}(f)} \kappa^{\Sigma}(B) \xrightarrow{\kappa^{\Sigma}(g)} \kappa^{\Sigma}(C_{F}) \longrightarrow 0 \end{split}$$

Now, from the universal property of colimits, there is a unique morphism $h: C_G \to B$ such that $\kappa_h^{\Sigma} \circ \rho^G = u$. Using the fact that $\kappa^{\Sigma}(f) = u \circ \phi$ and $\kappa^{\Sigma}(\operatorname{colim}_{\Sigma}(\phi)) \circ \rho^{\kappa^{\Sigma}(A)} = \rho^G \circ \phi$, we deduce that

$$\begin{split} \kappa^{\Sigma}(h \circ \operatorname{colim}_{\Sigma}(\phi)) \circ \rho^{\kappa^{\Sigma}(A)} &= \kappa_{h}^{\Sigma} \circ \kappa^{\Sigma}(\operatorname{colim}_{\Sigma}(\phi)) \circ \rho^{\kappa^{\Sigma}(A)} \\ &= \kappa_{h}^{\Sigma} \circ \rho^{G} \circ \phi \\ &= u \circ \phi \\ &= \kappa^{\Sigma}(f) \\ &= \kappa^{\Sigma}(f) \circ 1_{\kappa^{\Sigma}(A)} \\ &= \kappa^{\Sigma}(f) \circ \kappa_{\nabla^{A}}^{\Sigma} \circ \rho^{\kappa^{\Sigma}(A)} \\ &= \kappa^{\Sigma}(f \circ \nabla^{A}) \circ \rho^{\kappa^{\Sigma}(A)}. \end{split}$$

Then, by the universal property of colimits, $h \circ \operatorname{colim}_{\Sigma}(\phi) = f \circ \nabla^{A}$. Now, from the universal property of pushouts, we get a unique morphism $\gamma : Z_{\eta} \to B$ such that $\gamma \circ f_{\eta} = f$ (see (1.1)). Hence, f_{η} is a monomorphism.

Remark 4.3. Let \mathcal{X} and \mathcal{Y} be abelian categories, and $F: \mathcal{X} \to \mathcal{Y}$ and $G: \mathcal{Y} \to \mathcal{X}$ be functors such that (F, G) is an adjoint pair. We will say that (F, G) is an **Extadjoint pair** if there is a natural isomorphism $\operatorname{Ext}^{1}_{\mathcal{X}}(-, G(?)) \cong \operatorname{Ext}^{1}_{\mathcal{Y}}(F(-), ?)$. Note that the statement (c) of the above theorem can be restated by saying that $(\operatorname{colim}_{\Sigma}, \kappa^{\Sigma})$ is an Ext-adjoint pair. It should be noted that not every adjoint pair is Ext-adjoint. The reader can find counterexamples of this in $[1, \operatorname{pp.29-30}]$.

By duality we get the following straightforward corollary.

Corollary 4.4. Let Σ be a small category and let \mathcal{B} be a Σ -complete abelian category. Then, the following conditions are equivalent:

- (1) the functor \lim_{Σ} : Fun $(\Sigma, \mathcal{B}) \to \mathcal{B}$ is exact;
- (2) the maps $\Phi_{B,F}^{\Sigma} : Ext_{\mathcal{B}}^{1}(B, L_{F}) \to Ext_{\operatorname{Fun}(\Sigma,\mathcal{B})}^{1}(\kappa^{\Sigma}(B), F)$ defined as $\Phi_{B,F}^{\Sigma}(\overline{\epsilon}) = \overline{\varrho^{F} \cdot \kappa^{\Sigma}(\epsilon)}$, where $\varrho^{F} \circ \kappa^{\Sigma}(\epsilon)$ denotes the lower exact sequence in the below commutative diagram in $\operatorname{Fun}(\Sigma, \mathcal{B})$ with exact rows (here the left square is

a pushout), are natural isomorphisms both in B and in F.



(3) the bifunctors $Ext^{1}_{\mathcal{B}}(?, \lim_{\Sigma}(-))$ and $Ext^{1}_{\operatorname{Fun}(\Sigma, \mathcal{B})}(\kappa^{\Sigma}(?), -)$ are naturally isomorphic in the obvious way.

Now, we give a new proof of the following result, which is a particular case of the dual of [6, Theorem 3.2.8].

Corollary 4.5. In the setting of the previous corollary. If \mathcal{B} has enough projectives, then the functor \lim_{Σ} is exact if and only if the functor κ^{Σ} preserves projective objects.

Proof. (\Rightarrow) It follows by Corollary 4.4(c).

 (\Leftarrow) Let *B* be an object in \mathcal{B} and let *F* be a functor in Fun (Σ, \mathcal{B}) . Consider an exact sequence in \mathcal{B} of the form:

 $\epsilon: 0 \longrightarrow A \longrightarrow P \longrightarrow B \longrightarrow 0 ,$

where P is a projective object of \mathcal{B} . Applying the functors $\Gamma := \operatorname{Hom}_{\mathcal{B}}(-, \lim_{\Sigma}(F))$ and $\Upsilon := \operatorname{Hom}_{\operatorname{Fun}(\Sigma,\mathcal{B})}(-,F)$ on ϵ and $\kappa^{\Sigma}(\epsilon)$, respectively, we get the commutative diagram with exact rows

where Γ^1 and Υ^1 denote the functors $\operatorname{Ext}^1_{\mathcal{B}}(-, \lim_{\Sigma}(F))$ and $\operatorname{Ext}^1_{\operatorname{Fun}(\Sigma,\mathcal{B})}(-,F)$ respectively, and the morphisms f_i 's are the natural isomorphisms given by the adjoint pair $(\kappa^{\Sigma}, \lim_{\Sigma})$. Using the hypothesis on κ^{Σ} , we know that $\Upsilon^1(\kappa^{\Sigma}(P)) = 0$ and hence $\Phi^{\Sigma}_{B,F}$ is an isomorphism by the Five Lemma. Thus, this implication follows by Corollary 4.4(b).

The following result is a key for a direct consequence of the above corollary.

Lemma 4.6. Let \mathcal{A} be an abelian category. Then, the functor κ^{Σ} preserve injective and projective objects, for every set Σ viewed as a small category.

Proof. We show that the functor κ^{Σ} preserves projective objects. The rest of the statement can be proved by similar arguments. Let P be a projective object in \mathcal{A} and let ϵ : $F \stackrel{\psi}{\hookrightarrow} G \stackrel{\phi}{\twoheadrightarrow} \kappa^{\Sigma}(P)$ be an exact sequence in Fun (Σ, \mathcal{A}) . In such case, ϕ_i is a split epimorphism, for all $i \in Ob(\Sigma) = \Sigma$. Thus, for each $i \in \Sigma$, we can take a morphism $\xi_i : \kappa^{\Sigma}(P)(i) = P \to G(i)$ such that $\phi_i \circ \xi_i = 1_{\kappa^{\Sigma}(P)(i)}$. Now, using the fact that the only morphisms in Σ are the identity morphisms, we obtain that the family of morphisms $\xi := (\xi_i)_{i \in \Sigma}$ is a natural transformation from the functor $\kappa^{\Sigma}(P)$ to the functor G. Finally, note that $\xi \circ \phi = 1_{\kappa^{\Sigma}(P)}$ and hence $\overline{\epsilon}$ is the trivial extension in $\operatorname{Ext}^1_{\operatorname{Fun}(\Sigma, \mathcal{A})}(\kappa^{\Sigma}(P), F)$. Therefore, $\kappa^{\Sigma}(P)$ is a projective object in Fun (Σ, \mathcal{A}) as desired.

From Corollary 4.5 and its dual, together with the Lemma 4.6, we deduce the Corollary 3.2.9 in [6].

Corollary 4.7. Let \mathcal{A} be an Ab3 abelian category and let \mathcal{B} be an Ab3^{*} abelian category. Then, the following assertions hold:

- (1) \mathcal{A} is Ab4, whenever \mathcal{A} has enough injectives;
- (2) \mathcal{B} is Ab4*, whenever \mathcal{A} has enough projectives.

The following result is straightforward.

Lemma 4.8. Let \mathcal{A} be an abelian category and let Σ be a set. Given an exact sequence $\epsilon : H \xrightarrow{\phi} G \xrightarrow{\psi} F$ in Fun (Σ, \mathcal{A}) , consider the exact sequence $\epsilon_i : H(i) \xrightarrow{\phi_i} G(i) \xrightarrow{\psi_i} F(i) \quad \forall i \in \Sigma$. Then, the assignment $\epsilon \mapsto \prod_{i \in \Sigma} \epsilon_i$ give rise a family of isomorphisms

$$\Xi_{F,A}^{\Sigma} : Ext_{\operatorname{Fun}(\Sigma,\mathcal{A})}^{1}(F,\kappa^{\Sigma}(A)) \longrightarrow \prod_{i \in \Sigma} Ext_{\mathcal{A}}^{1}(F(i),A)$$

$$\Theta_{F,A}^{\Sigma} : Ext^{1}_{\operatorname{Fun}(\Sigma,\mathcal{A})}(\kappa^{\Sigma}(A),F) \longrightarrow \prod_{i \in \Sigma} Ext^{1}_{\mathcal{A}}(A,F(i))$$

for all F in Fun(Σ, \mathcal{A}) and A in $Ob(\mathcal{A})$.

Let \mathcal{A} be an Ab3 abelian category and Σ be a set viewed as a small category. In this context, $\operatorname{colim}_{\Sigma}(F) = \coprod_{i \in \Sigma} F(i)$ for all F in $\operatorname{Fun}(\Sigma, \mathcal{A})$. We can use this to get as a corollary [2, Theorems 4.9 and 4.11].

Corollary 4.9. Let \mathcal{A} be an Ab3 abelian category and let \mathcal{B} be an Ab3^{*} abelian category. Then, the following assertions hold:

(1) A is Ab4 if, and only if, the canonical map

$$Ext^{1}_{\mathcal{A}}(\coprod_{i \in I} A_{i}, A) \to \prod_{i \in I} Ext^{1}_{\mathcal{A}}(A_{i}, A)$$

is an isomorphism, for all A and $(A_i)_{i \in I}$ object and family of objects in \mathcal{A} , respectively.

(2) \mathcal{B} is Ab4* if, and only if, the canonical map

$$Ext^{1}_{\mathcal{B}}(B,\prod_{i\in I}B_{i})\to\prod_{i\in I}Ext^{1}_{\mathcal{B}}(B,B_{i})$$

is an isomorphism, for all B and $(B_i)_{i \in I}$ object and family of objects in \mathcal{B} , respectively.

Proof. We prove statement (a), statement (b) follows by duality. Let I be a set and let $(A_i)_{i \in I}$ be a family of objects in \mathcal{A} . Then, the assignment $i \mapsto A_i$ give rise to a functor $F: I \to \mathcal{A}$. Notice that for each A in Ob(A), we have that the map in (a) coincide with the compositions $\Xi^{I}_{F,A} \circ \Psi^{I}_{F,A}$ (see Theorem 4.2 and Lemma 4.8). On the other hand, every functor in Fun (I, \mathcal{A}) can be viewed as an I-indexed family of objects in \mathcal{A} . Then, the result follows by Theorem 4.2 and Lemma 4.8. \Box

12

References

- [1] W. W. Adams and M. A. Rieffel. Adjoint functors and derived functors with an application to the cohomology of semigroups. Journal of Algebra, 7(1):25-34, 1967.
- [2] A, Argundín-Monroy. The Yoneda Ext and arbitrary coproducts in abelian categories. Glasgow Mathematical Journal, pages 1–15, 2021.
- [3] A. Grothendieck. Sur quelques points d'Algèbre Homologique. Tohoku Math. J. 9(2) (1957), 119-221.
- [4] S. Mac Lane. Categories for the working mathematician, volume 5. Springer Verlag Graduate texts in Mathematics, 1998.
- [5] B. Mitchell. Theory of Categories. Academic Press, 1965.
- [6] N. Popescu. Abelian categories with applications to rings and modules, volume 3. Academic Press, London, 1973.
- [7] B. Stenströn. Rings of quotients. Springer-Verlag, Berlin Heidelberg, 1975.

Instituto de Matemática y Estadística Rafael Laguardia, Facultad de Ingeniería, UNIVERSIDAD DE LA REPÚBLICA, JULIO HERRERA Y REISSIG 565, MONTEVIDEO, URUGUAY Email address: argudin@ciencias.unam.mx

INSTITUTO DE CIENCIAS FÍSICAS Y MATEMÁTICAS, EDIFICIO EMILIO PUGIN, CAMPUS ISLA TEJA, UNIVERSIDAD AUSTRAL DE CHILE, 5090000 VALDIVIA, CHILE

Email address: carlos.parra@uach.cl