On Convergence of General Truncation-Augmentation Schemes for Approximating Stationary Distributions of Markov Chains

Alex Infanger*

Peter W. Glynn^{*†}

Yuanyuan Liu[‡]

Abstract

In the analysis of Markov chains and processes, it is sometimes convenient to replace an unbounded state space with a "truncated" bounded state space. When such a replacement is made, one often wants to know whether the equilibrium behavior of the truncated chain or process is close to that of the untruncated system. For example, such questions arise naturally when considering numerical methods for computing stationary distributions on unbounded state space. In this paper, we study general truncation-augmentation schemes, in which the substochastic truncated "northwest corner" of the transition matrix or kernel is stochasticized (or augmented) arbitrarily. In the presence of a Lyapunov condition involving a coercive function, we show that such schemes are generally convergent in countable state space, provided that the truncation is chosen as a sublevel set of the Lyapunov function. For stochastically monotone Markov chains on \mathbb{Z}_+ , we prove that we can always choose the truncation sets to be of the form $\{0, 1, ..., n\}$. We then provide sufficient conditions for weakly continuous Markov chains under which general truncation-augmentation schemes converge weakly in continuous state space. Finally, we briefly discuss the extension of the theory to continuous time Markov jump processes.

Key words: Markov chains, stationary distributions, numerical methods, Lyapunov functions.

1 Introduction

Let $X = (X_n : n \ge 0)$ be a positive recurrent Markov chain taking values in an unbounded state space S. In many settings, one is interested in producing a

^{*}Institute for Computational & Mathematical Engineering, Stanford University, USA.

[†]Department of Management Science & Engineering, Stanford University, USA. [‡]School of Mathematics and Statistics, Central South University, China.

positive recurrent Markov chain approximation to X with bounded state space, perhaps motivated by numerical considerations. For example, in analyzing an irreducible positive recurrent Markov chain X on countable state space S, numerical computation of the stationary distribution π for X requires truncating its state space to a finite subset A_n having n elements. Since the "northwest corner" matrix $(P(x, y) : x, y \in A_n)$ is necessarily substochastic, it is natural to then "augment" the probabilities P(x, y) so as to create a stochastic matrix $P_n = (P_n(x, y) : x, y \in A_n)$. One then hopes that the stationary distribution π_n of P_n will be a good approximation to π . However, it is well known that universal convergence (as n tends to ∞) for such truncation-augmentation schemes is not guaranteed in general; see, for example, (2.5) in Wolf (1980) and Example 1 of this paper.

However, when one constructs the augmentation via a "fixed state" augmentation, Wolf (1980) showed that convergence of π_n to π is guaranteed when X is an irreducible positive recurrent countable state space Markov chain. In Infanger and Glynn (2022), we show that such convergence can be generally validated when the underlying Markov chain or process is suitably regenerative. This allows us to develop a convergence theory, in a suitably chosen weighted total variation norm, for irreducible positive recurrent Markov chains on countable state spaces and, more generally, for Harris recurrent Markov chains on general state spaces.

In this paper, we seek conditions under which one can be assured that any truncation-augmentation scheme is convergent. In the countable state space setting, one of us developed such results in Liu (2010). However, as we shall discuss in Section 2, the argument given there relies on results from Borovkov (1998) that are not correctly stated. Section 2 therefore provides new sufficient conditions guaranteeing universal convergence. Our sufficient conditions require that the truncation set be chosen as a sublevel set of a coercive Lyapunov function. This result suggests that a Lyapunov function for the Markov chain X can be useful in designing convergent truncation-augmentation schemes. We also use our approach to recover a convergence result of Gibson and Seneta (1987b) and Tweedie (1998), for stochastically monotone Markov chains; see Theorem 2.

In Section 3, we study this convergence question in the setting of Markov chains taking values in a complete separable metric space. We generalize the results of Section 2 by establishing weak convergence of π_n to π (in contrast to the total variation convergence of Infanger and Glynn (2022)), when X has suitably continuous transition probabilities. The arguments in Sections 2 and 3 do not assume Harris recurrence. Rather, they rely on Prohorov's theorem and related tightness ideas, and the methods establish convergence for arbitrary augmentations. Section 4 uses an argument based on regeneration to establish that general augmentations are valid for strongly uniformly recurrent Markov chains on general state space, thereby generalizing the known theory for Markov matrices on discrete state space to the continuous setting.

This paper concludes, in Section 5, with a brief discussion of the related convergence theory for continuous time Markov jump processes.

2 Truncation-Augmentation for Markov Chains with Countably Infinite State Space

In this section, we consider an irreducible positive recurrent Markov chain $X = (X_n : n \ge 0)$ taking values in a countably infinite state space S. We let $P = (P(x,y) : x, y \in S)$ be the one-step transition matrix of X, and we denote the (unique) stationary distribution of X by $\pi = (\pi(x) : x \in S)$. Let $(A_n : n \ge 0)$ be a strictly increasing sequence of subsets of S satisfying $A_0 \subset A_1 \subset A_2 \subset ...$ such that $\bigcap_{n=0}^{\infty} A_n^c = \emptyset$. For the *n*'th truncation A_n , let $B_n = (B_n(x,y) : x, y \in A_n)$ be the corresponding "northwest corner" truncation of P in which $B_n(x,y) = P(x,y)$ for $x, y \in A_n$. The irreducibility of P then guarantees that there exists at least one row of B_n with a row sum strictly less than 1.

We say that P_n is a general augmentation of B_n if $P_n = (P_n(x, y) : x, y \in A_n)$ is a stochastic matrix for which

$$P_n(x,y) \ge B_n(x,y) = P(x,y)$$

for $x, y \in A_n$. If there exists a probability distribution $\nu_n = (\nu_n(x) : x \in A_n)$ for which

$$P_n(x,y) = P(x,y) + \sum_{z \in A_n^c} P(x,z)\nu_n(y),$$
(2.1)

then P_n is said to be a *linear augmentation* of B_n . If there exists a probability $\nu = (\nu(x) : x \in S)$ for which

$$\nu_n(x) = \frac{\nu(x)}{\sum_{y \in A_n} \nu(y)}$$

for $x \in A_n$, then we say that P_n is formed from a fixed linear augmentation ν . If $\nu = \delta_y$ for a fixed $y \in S$, where $\delta_y = (\delta_y(x) : x \in S)$ is a unit point mass distribution at y, then we say that P_n is a fixed state augmentation of B_n . When $S = \mathbb{Z}_+$ and $A_n = \{0, 1, ..., n\}$, setting $\nu = \delta_0$ is called first state augmentation, whereas the use of $\nu_n = \delta_n$ is called last state augmentation.

Let Π_n be the set of stationary distributions on A_n associated with P_n . Since $|A_n|$ is finite, Π_n is always non-empty. It is easily seen that when P_n is a general augmentation, Π_n may not be a singleton. However, for linear augmentations (whether fixed or not), Π_n is always guaranteed to consist of a single unique stationary distribution π_n for P_n . This follows because the irreducibility of P guarantees that the probability of an exit to A_n^c is positive from every $x \in A_n$. Once X attempts to exit to A_n^c , the chain is re-distributed on A_n according to ν_n ; see (2.1). Hence, every state $y \in A_n$ for which $\nu_n(y) > 0$ is reachable from every $x \in A_n$, as are all the states reachable from such states $y \in A_n$. Consequently, P_n has exactly one closed communicating class, so that P_n has a unique stationary distribution π_n ; see also p. 261 of Seneta (1980).

To make our discussion as self-contained as possible, we now provide an example showing that even when X is very well-behaved, π_n may fail to converge to π as $n \to \infty$.

Example 1. Suppose that $S = \mathbb{Z}_+$ with P(2i, 2i + 1) = 1/2 = P(2i, 0) for $i \in \mathbb{Z}_+$, and P(2i + 1, 2i + 2) = 1 for $i \in \mathbb{Z}_+$. Then,

$$P^{2}(2i,0) \ge P(2i,0)P(0,0) = \frac{1}{4},$$

for $i \in \mathbb{Z}_+$ whereas,

$$P^{2}(2i+1,0) \ge P(2i+1,2i+2)P(2i+2,0) = \frac{1}{2}$$

for $i \in \mathbb{Z}_+$. Hence $P^2(x,0) \ge 1/4$ for $x \in \mathbb{Z}_+$, so that the two-step transition matrix is a Markov matrix, and X is uniformly ergodic.

Suppose we use last state augmentation. Then, when $A_n = \{0, 1, ..., n\}$ with n odd, state n is absorbing, and the single closed communicating class corresponding to P_n is just $\{n\}$. It follows that $\pi_n = \delta_n$ for n odd, so that π_n fails to converge to the stationary distribution π of X, despite the fact that X is uniformly ergodic.

A significant literature has developed over the years, focused on obtaining conditions under which various augmentations are guaranteed to converge. As noted in the Introduction, Wolf (1980) proved that fixed state augmentation is always convergent. This supplemented earlier work of Golub and Seneta (1974), in which it was shown that last state augmentation converges for upper Hessenberg transition matrices. Gibson and Seneta (1987a) showed that general augmentations always converge when P is either a Markov matrix or when it is upper Hessenberg. They also showed that when P is lower Hessenberg and P_n is a linear augmentation for which the sequence $(\nu_n : n \ge 0)$ is chosen to be tight, then π_n is guaranteed to converge to π . Gibson and Seneta (1987b) showed that convergence also holds for general augmentations when the underlying Markov chain is stochastically monotone. Tweedie (1998) proved convergence for the special case of last state augmentation when P corresponds to a stochastically monotone Markov chain (although most of the paper, as with much of the subsequent literature, focuses on identifying computable rates of convergence of π_n to π).

In Liu (2010), convergence of π_n to π for general augmentations is discussed, in the presence of a Lyapunov condition denoted there as D1(V, b, C). The arguments given there rely on the following result (Theorem 5.1) from Borovkov (1998) (suitably re-stated for our current exposition).

Suppose that $(P_n : n \ge 1)$ is a family of one-step transition matrices defined on a countable state space S for which there exists a finite subset $C \subseteq S$, $\lambda > 0$, $c < \infty$, and a probability $\phi = (\phi(x) : x \in S)$ such that:

- i) Under P_n , X is guaranteed to hit C from any state $x \in S$;
- ii) $P_n(x,y) \ge \lambda \phi(y)$ for $x \in C, y \in S$;
- iii) $\max_{x \in C} E_n[\tau(C)|X_0 = x] \leq c$, where $E_n(\cdot)$ is the expectation on the path-space of X associated with P_n and $\tau(C) = \inf\{n \geq 1: X_n \in C\}$ is the first return time to C.

Then, P_n has a unique stationary distribution π_n , and

$$\sup_{n \ge 1} \sum_{y \in S} |P_n^m(x, y) - \pi_n(y)| \to 0$$
 (2.2)

as $m \to \infty$.

Unfortunately, this result is not valid as stated, as made clear by the following counter-example. (The above statement is also missing the obvious aperiodicity requirement, but the aperiodic example below focuses on the more subtle flaw in the statement.)

Example 2. Suppose that $S = \mathbb{Z}_+$ with $P_n(i, i-1) = 1$ for $i \ge 1$ and $n \ge 1$. We put $P_n(0,0) = 1 - (1/(n+1))$ and $P_n(0,n) = 1/(n+1)$ for $n \ge 1$. Then, we let $C = \{0\}, \lambda = 1/2$, and $\phi = \delta_0$. Note that

$$E_n[\tau(C)|X_0=0] = 1 \cdot \left(1 - \frac{1}{n+1}\right) + (n+1) \cdot \frac{1}{n+1} \le 2,$$

so that conditions i), ii), and iii) are all in force, uniformly in $n \ge 1$. For this example,

$$\pi_n(0) = \left(2 - \frac{1}{n+1}\right)^{-1}$$

with

$$\pi_n(i) = \frac{1}{(n+1)} \left(2 - \frac{1}{n+1}\right)^{-1}$$

for $1 \leq i \leq n$. Also,

$$P_n^m(0,0) \ge \left(1 - \frac{1}{n+1}\right)^m$$

 \mathbf{SO}

$$\sup_{n \ge 1} \left[P_n^m(0,0) - \pi_n(0) \right] \ge \sup_{n \ge 1} \left[\left(1 - \frac{1}{n+1} \right)^m - \left(2 - \frac{1}{n+1} \right)^{-1} \right] = \frac{1}{2},$$

so (2.2) fails to hold.

In view of this example, we now provide here a new discussion of the convergence theory of general augmentations that does not rely on this mis-stated result from Borovkov (1998), and that is of interest in its own right. We start by recalling that a function $f: S \to \mathbb{R}_+$ is said to be *coercive* if the sublevel set $K_m(f) \stackrel{\Delta}{=} \{x \in S : f(x) \le m\}$ has finite cardinality for each $m \ge 1$. Our theory relies on the following Lyapunov condition assumption:

A1. There exists $g: S \to \mathbb{R}_+$, $b < \infty$, and a coercive function r such that

$$(Pg)(x) \le g(x) - r(x) + b$$
 (2.3)

for $x \in S$.

Remark 1. We refer to the function g appearing in (2.3) as a (stochastic) Lyapunov function.

We note that because g is non-negative, $(Pg)(\cdot)$ is non-negative, so that $r(x) \leq g(x) + b$ for $x \in S$. Hence, if $g(x) \leq m$, it follows that $r(x) \leq m + b$, so $K_m(g) \subseteq K_{m+b}(r)$. Consequently, the sublevel sets of the Lyapunov function g have finite cardinality, so g is also coercive.

We now assume that we construct our truncation sets based on A1. In particular, we use the Lyapunov function g to design our truncation sequence $(A_n : n \ge 1)$, specifically putting $A_n = K_n(g)$.

Theorem 1. Assume A1 and suppose that $A_n = K_n(g)$ for $n \ge 1$. Then, for any general augmentation sequence $(P_n : n \ge 1)$ associated with the A_n 's, and for any $\pi_n \in \Pi_n$,

$$\sum_{x \in A_n} |\pi_n(x) - \pi(x)| \to 0$$
 (2.4)

as $n \to \infty$.

Proof. We first show that the Lyapunov bound A1 for P can be extended (uniformly) to the P_n 's associated with a sequence of general augmentations. In particular, we observe that for $x \in A_n$,

$$\begin{split} \sum_{y \in A_n} P_n(x, y) g(y) &= \sum_{y \in A_n} P(x, y) g(y) + \sum_{y \in A_n} (P_n(x, y) - P(x, y)) g(y) \\ &\leq \sum_{y \in A_n} P(x, y) g(y) + \sum_{y \in A_n} (P_n(x, y) - P(x, y)) \sup_{z \in A_n} g(z) \\ &\leq \sum_{y \in A_n} P(x, y) g(y) + \sum_{y \in A_n^c} P(x, y) \inf_{z \in A_n^c} g(z) \end{split}$$

$$\leq \sum_{y \in A_n} P(x, y)g(y) + \sum_{y \in A_n^c} P(x, y)g(y)$$

= $(Pg)(x)$
 $\leq g(x) - r(x) + b,$ (2.5)

which, of course, is the desired uniform version of (2.3). (Note that the fact that A_n is chosen as the sublevel set $K_n(g)$ is used in a critical way in our second inequality above.)

We can now apply Corollary 4 of Glynn and Zeevi (2008), p. 202, to (2.5) to conclude that

$$\sum_{x \in A_n} \pi_n(x) r(x) \le b$$

for $n \ge 1$. For $\epsilon > 0$, choose $m = m(\epsilon)$ large enough that $b/m < \epsilon$. Then, Markov's inequality guarantees that

$$\sum_{\substack{x \notin K_m(r)\\x \in A_n}} \pi_n(x) \le b/m < \epsilon$$

and hence

$$\sum_{x \in K_m(r)} \pi_n(x) \ge \sum_{x \in K_m(r) \cap A_n} \pi_n(x) \ge 1 - \epsilon$$

uniformly in $n \ge 1$. It follows that $(\pi_n : n \ge 1)$ is a tight sequence of probabilities on S. Consequently, Prohorov's theorem implies that every subsequence $(\pi_{n_k} : k \ge 1)$ contains a further subsequence $(\pi_{n'_k} : k \ge 1)$ for which

$$\pi_{n'_k}(x) \to \pi'(x) \tag{2.6}$$

at each $x \in S$ as $k \to \infty$, where $\pi' = (\pi'(x) : x \in S)$ is a probability on S; see, for example, Billingsley (1968). Then, for $y \in S$,

$$\pi'(y) = \lim_{k \to \infty} \pi_{n'_k}(y)$$

$$= \lim_{k \to \infty} \sum_{x \in A_n} \pi_{n'_k}(y) P_{n'_k}(x, y)$$

$$\geq \sum_{x \in S} \lim_{k \to \infty} \pi_{n'_k}(x) P_{n'_k}(x, y)$$

$$= \sum_{x \in S} \pi'(x) P(x, y), \qquad (2.7)$$

as a result of Fatou's lemma. Since,

$$1 = \sum_{y \in S} \pi'(y) = \sum_{y \in S} \sum_{x \in S} \pi'(x) P(x, y)$$
(2.8)

it is evident that (2.7) must hold with equality. Since π is the unique stationary distribution of P, we find that $\pi = \pi'$. Since every convergent subsequence $(\pi_{n'_k}: k \ge 0)$ must have the same limit π , we conclude that for each $x \in S$,

$$\pi_n(x) \to \pi(x) \tag{2.9}$$

as $n \to \infty$. Since S is countably infinite, this easily implies (2.4).

Remark 2. The novel element in the above proof is establishing tightness. Seneta (1980) proves that tightness implies convergence; our argument of (2.6) through (2.9) is provided in order to make the argument as self-contained as possible. The argument also makes clear that if $A_n \nearrow S$ is chosen so that $\sup_{z \in A_n} g(z) \leq \inf_{z \in A_n^c} g(z)$ for $n \geq 1$, the proof continues to be valid. This observation is used in the proof of Theorem 2.

Remark 3. For Example 1, we note that if $g(2i) = e^{\theta i}$ and $g(2i+1) = e^{\theta(i+3/2)}$ for $i \ge 0$, then

$$(Pg)(2i) = \frac{1}{2}g(0) + \frac{1}{2}g(2i+1)$$

= $g(2i) - (1 - \frac{1}{2}e^{3\theta/2})e^{\theta i} + \frac{1}{2},$

while

$$(Pg)(2i+1) = g(2i+2)$$

= $g(2i+1) - (e^{3\theta/2} - e^{\theta})e^{\theta i}.$

Hence, if we choose $\theta > 0$ so that $e^{3\theta/2} < 2$, we find that A1 holds with r coercive. In this case, we note that the sublevel sets of g can only take the form $\{0, 1, ..., 2i\}$ for $i \in \mathbb{Z}_+$, so that the bad A_n 's of Example 1 are precluded by our choice of Lyapunov function.

Remark 4. Note that Theorem 1 establishes that no strictly increasing Lyapunov function g can be constructed for Example 1, for otherwise we would be guaranteed convergence along the sequence of truncation sets given by $A_n = \{0, 1, ..., n\}$.

Remark 5. For Example 1, the uniform ergodicity of X implies that one can find a bounded non-negative g for which

$$(Pg)(x) \le g(x) - 1 \tag{2.10}$$

for $x \ge 1$. (In particular, we can take g(x) as the expected hitting time of $\{0\}$, starting from x.) Inequality (2.10) is closely related to condition D1(V, b, C) of Liu (2010). Note that the sublevel sets of this g form truncation sets that either equal S (when the level is chosen higher than the bound) or take the same form as in Remark 3. Thus, in this example, Theorem 1 holds, even though the function r associated with (2.10) is non-coercive. In particular, we do not

know whether Theorem 1 continues to hold when one chooses $A_n = K_n(g)$ in the presence of (2.10) holding outside some finite subset $C \subseteq S$. (Of course, we would want to add the requirement that g be coercive, in order to ensure that $K_n(g)$ has finite cardinality for each $n \geq 1$.)

Remark 6. When A1 holds, the coerciveness of r guarantees that

$$(Pg)(x) \le g(x) - 1$$
 (2.11)

for x outside some finite subset C (say). This implies that the expected hitting time of C from $x \in C^c$ is bounded above by g(x); see, for example, p. 344 in Meyn and Tweedie (2009). Hence, A1 implies that C can be reached with positive probability from each $x \in C^c$. On the other hand, the irreducibility of P and finiteness of C ensure that each $y \in C$ can be reached from each $x \in C$ by a path lying entirely within $K_n(g)$ when n is chosen sufficiently large. Hence A1 guarantees that for n large, Π_n is guaranteed to have a single (unique) element.

We conclude this section by showing that the general augmentation result for stochastically monotone Markov chains due to Gibson and Seneta (1987b) is a special case of Theorem 1, as is the last state augmentation result of Tweedie (1998). We recall that a Markov chain is *stochastically monotone* on \mathbb{Z}_+ if $\sum_{w>y} P(x, w)$ is non-decreasing in x for each y.

Theorem 2. Suppose that X is an irreducible and positive recurrent stochastically monotone Markov chain on \mathbb{Z}_+ with A_n chosen to be $A_n = \{0, 1, ..., n\}$. Then, for any general augmentation sequence $(P_n : n \ge 1)$ associated with the A_n 's, and for any $\pi_n \in \Pi_n$,

$$\sum_{x} |\pi_n(x) - \pi(x)| \to 0$$

as $n \to \infty$, where $\pi = (\pi(x) : x \in S)$ is the unique stationary distribution of X. Proof. Put

$$\bar{P}(x) = \sum_{y \ge x} \pi(y)$$

and set

$$r(x) = \bar{P}(x)^{-\frac{1}{2}}$$

for $x \ge 0$. Clearly, r is non-negative, non-decreasing, and coercive. Furthermore,

$$0 \le \sum_{k=0}^{n} r(k)\pi(k) = \sum_{k=0}^{n} \bar{P}(k)^{-1/2} (\bar{P}(k) - \bar{P}(k+1))$$
$$= \sum_{k=0}^{n} \int_{\bar{P}(k+1)}^{\bar{P}(k)} \bar{P}(k)^{-1/2} du$$

$$\leq \sum_{k=0}^{n} \int_{\bar{P}(k+1)}^{\bar{P}(k)} u^{-1/2} du$$
$$= \int_{\bar{P}(n+1)}^{1} u^{-1/2} du \leq \int_{0}^{1} u^{-1/2} du = 2 < \infty.$$
(2.12)

Hence, (2.12) shows that

$$\alpha \stackrel{\Delta}{=} \sum_{k=0}^{\infty} r(k)\pi(k) < \infty.$$

As a consequence, we can put $r_c(x) = r(x) - \alpha$ and consider Poisson's equation

$$(P-I)\tilde{g} = -r_c. \tag{2.13}$$

It is shown in Glynn and Infanger (2022) that because X is stochastically monotone and r_c is non-decreasing, (2.13) has a finite-valued non-decreasing solution \tilde{g} . Also, we note that $g(x) = \tilde{g}(x) - \tilde{g}(0)$ is guaranteed to be nonnegative. Hence,

$$(Pg)(x) = g(x) - r(x) + \alpha$$

for $x \ge 0$, where r is non-negative and coercive and g is non-negative. We can therefore apply Theorem 1. Because g is non-decreasing, the sublevel sets $K_n(g)$ take the form $\{0, 1, 2, ..., g^{-1}(n)\} = A_{g^{-1}(n)}$, so that Theorem 1 and Remark 2 yield the desired result.

The key special feature of a stochastically monotone Markov chain is that we may always choose A_n to be an "interval" of the form $\{0, 1, ..., n\}$, and yet retain convergence for general augmentations.

3 Convergence of General Augmentations for Continuous State Space Markov Chains

In this section, we assume that $X = (X_n : n \ge 0)$ is a Markov chain taking values in a complete separable metric space X. For $x, y \in S$, let

$$P(x, dy) = P(X_{n+1} \in dy | X_n = x)$$

and let $P = (P(x, dy) : x, y \in S)$ be the one-step transition kernel of X. We assume that X has a unique stationary distribution $\pi = (\pi(dx) : x \in S)$ satisfying the equation

$$\pi(dy) = \int_{S} \pi(dx) P(x, dy)$$

for $y \in S$.

As in Section 2, our theory requires a Lyapunov function assumption. In this setting, we say that a function $f: S \to \mathbb{R}_+$ is *coercive* if for each $n \ge 1$, $K_n(f) = \{x \in S : f(x) \le n\}$ is either empty or compact in S.

A2. There exists a coercive function r, a non-negative function $g: S \to \mathbb{R}_+$, and $b < \infty$ for which

$$\int_{S} P(x, dy)g(y) \le g(x) - r(x) + b \tag{3.1}$$

for $x \in S$.

As in Section 2, $K_m(g) \subseteq K_{m+b}(r)$. If g is continuous, the fact that [0, m] is closed in \mathbb{R}_+ implies that $K_m(g)$ is closed. Also, $K_m(g)$ is a closed subset of the compact set $K_{m+b}(r)$, and so $K_m(g)$ is then compact; see p. 73 of Copson (1968). In other words, if g is continuous, A2 guarantees that g is coercive.

Let $A_n = K_n(g)$, and put $B_n = (B_n(x, dy) : x, y \in A_n)$, where $B_n(x, dy) = P(x, dy)$ for $x, y \in A_n$. Let $P_n = (P_n(x, dy) : x, y \in A_n)$ be a general augmentation for B_n , so that it is a stochastic kernel on A_n for which

$$P_n(x, dy) \ge B_n(x, dy) = P(x, dy)$$

for $x, y \in A_n$. Suppose that Π_n is the set of stationary distributions for P_n . Given our assumptions up to this point, Π_n may be either empty, consist of a singleton, or contain a multiplicity of elements.

Theorem 3. Suppose A2 holds and that Π_n is non-empty for $n \ge 1$. If $\pi_n \in \Pi_n$, then $(\pi_n : n \ge 1)$ is a tight sequence of probabilities on S.

Proof. Because A_n is chosen as a sublevel set of g, the same argument as used in Theorem 1 proves that

$$\int_{A_n} P_n(x, dy)g(y) \le g(x) - r(x) + b$$

for $x \in A_n$. Again, Corollary 4 of Glynn and Zeevi (2008) proves that

$$\int_{A_n} \pi_n(dx) r(x) \le b$$

for $n \ge 1$. For $\epsilon > 0$, choose $m = m(\epsilon)$ so that $b/m < \epsilon$. Then, Markov's inequality implies that $\pi_n(K_m(r)^c) < \epsilon$, and hence

$$\pi_n(K_m(r)) \ge 1 - \epsilon$$

uniformly in $n \ge 1$. Since $K_m(r)$ is compact, this establishes the tightness of $(\pi_n : n \ge 1)$.

Remark 7. As in the proof of Theorem 2, it suffices that $A_n \nearrow S$ with $\sup_{z \in A_n} g(z) \leq \inf_{z \in A_n^c} g(z)$, in order that the result be valid.

We now wish to argue that any weak limit point of $(\pi_n : n \ge 1)$ must equal π . For this purpose, we let bC be the space of real-valued bounded continuous functions with domain S, and assume:

A3. If
$$f \in bC$$
, then

$$(Pf)(\cdot) \stackrel{\Delta}{=} \int_{S} f(y) P(\cdot, dy)$$

also lies in bC.

A transition kernel satisfying A3 is said to be *weakly continuous* (or equivalently, *Feller continuous*).

Theorem 4. Suppose A2 and A3 hold with g continuous, and that Π_n is nonempty for $n \ge 1$. If $\pi_n \in \Pi_n$, then

$$\pi_n \Rightarrow \pi$$

as $n \to \infty$, where \Rightarrow denotes weak convergence in S.

Proof. In view of Theorem 3, let $(\pi_{n'_k} : k \ge 1)$ be a weakly convergent subsequence of $(\pi_n : n \ge 1)$, so that there exists a probability π' on S for which

$$\pi_{n'_k} \Rightarrow \pi' \tag{3.2}$$

as $k \to \infty$. For $f \in bC$, it follows that

$$\int_{A_{n'_k}} \pi_{n'_k}(dx) f(x) = \int_S \pi_{n'_k}(dx) f(x) \to \int_S \pi'(dx) f(x)$$
(3.3)

as $k \to \infty$. Since $\pi_{n'_k} \in \prod_{n'_k}$,

$$\int_{S} \pi_{n'_{k}}(dx)(P_{n'_{k}}f)(x) = \int_{S} \pi_{n'_{k}}(dx)f(x).$$
(3.4)

For $f \in bC$, let $||f|| = \sup\{|f(x)| : x \in S\}$. Also, for $\epsilon > 0$, the tightness of the sequence $(\pi_n : n \ge 1)$ guarantees the existence of a compact set $K = K(\epsilon)$ for which $\pi_n(K) \ge 1 - \epsilon$ uniformly in $n \ge 1$. We can then write

$$\begin{aligned} \left| \int_{S} \pi_{n}(dx)(P_{n}f)(x) - \int_{S} \pi_{n}(dx)(Pf)(x) \right| \\ &\leq \left| \int_{K} \pi_{n}(dx)\left((P_{n}f)(x) - (Pf)(x)\right) \right| + 2\epsilon \|f\| \\ &= \left| \int_{K\cap A_{n}} \pi_{n}(dx) \left[\int_{A_{n}} f(y)(P_{n}(x,dy) - P(x,dy)) - \int_{A_{n}^{c}} f(y)P(x,dy) \right] \right| + \epsilon \|f\| \\ &\leq \left| \int_{K\cap A_{n}} \pi_{n}(dx) \left[\int_{A_{n}} f(y)(P_{n}(x,dy) - P(x,dy)) - \int_{A_{n}^{c}} f(y)P(x,dy) \right] \right| + \epsilon \|f\| \\ &\leq \left| \int_{K\cap A_{n}} \pi_{n}(dx) \left[\int_{A_{n}} f(y)(P_{n}(x,dy) - P(x,dy)) - \int_{A_{n}^{c}} f(y)P(x,dy) \right] \right| + \epsilon \|f\| \\ &\leq \left| \int_{K\cap A_{n}} \pi_{n}(dx) \left[\int_{A_{n}} f(y)(P_{n}(x,dy) - P(x,dy)) - \int_{A_{n}^{c}} f(y)P(x,dy) \right] \right| + \epsilon \|f\| \\ &\leq \left| \int_{K\cap A_{n}} \pi_{n}(dx) \left[\int_{A_{n}} f(y)(P_{n}(x,dy) - P(x,dy)) - \int_{A_{n}^{c}} f(y)P(x,dy) \right] \right| + \epsilon \|f\| \\ &\leq \left| \int_{K\cap A_{n}} \pi_{n}(dx) \left[\int_{A_{n}} f(y)(P_{n}(x,dy) - P(x,dy)) - \int_{A_{n}^{c}} f(y)P(x,dy) \right] \right| \\ &\leq \left| \int_{K\cap A_{n}} \pi_{n}(dx) \left[\int_{A_{n}} f(y)(P_{n}(x,dy) - P(x,dy)) - \int_{A_{n}^{c}} f(y)P(x,dy) \right] \right|$$

$$2\epsilon \|f\| \le \|f\| \int_{K \cap A_n} \pi_n(dx) |P_n(x, A_n) - P(x, A_n)| + \int_{K \cap A_n} \pi_n(dx) \int_{A_n^c} |f(y)| P(x, dy) + 2\epsilon \|f\| \le \|f\| \int_{K \cap A_n} \pi_n(dx) |P_n(x, A_n^c) - P(x, A_n^c)| + \|f\| \int_{K \cap A_n} \pi_n(dx) P(x, A_n^c) + 2\epsilon \|f\| \le 2 \|f\| \int_K \pi_n(dx) P(x, A_n^c) + 2\epsilon \|f\|.$$

$$(3.5)$$

We claim that

$$\sup_{x \in K} P(x, A_n^c) \to 0 \tag{3.6}$$

as $n \to \infty$. To prove this, suppose that (3.6) does not hold. Then, there exists $\delta > 0$ and a sequence $(x_n : n \ge 1)$ such that $x_n \in K$ and

$$P(x_n, A_n^c) \ge \delta \tag{3.7}$$

for $n \geq 1$. If $P_x(\cdot) \stackrel{\Delta}{=} P(\cdot|X_0 = x)$, (3.7) is equivalent to requiring that

$$P_{x_n}(X_1 \in A_n^c) \ge \delta$$

or, in other words,

$$P_{x_n}(g(X_1) > n) \ge \delta \tag{3.8}$$

for $n \geq 1$. Because K is compact, we can extract a subsequence $(x_{n_k} : k \geq 1)$ of $(x_n : n \geq 1)$ and $x_{\infty} \in K$ for which $x_{n_k} \to x_{\infty}$ and (3.8) is in force along the subsequence.

Choose $r \in \mathbb{Z}_+$ large enough that

$$P_{x_{\infty}}(g(X_1) < r) \ge 1 - \frac{\delta}{2}.$$
 (3.9)

For $h \in bC$, the continuity of Ph (due to A3) implies that $(Ph)(x_{n_k}) \to (Ph)(x_{\infty})$ as $k \to \infty$. It follows that

$$P_{x_{n_k}}(X_1 \in \cdot) \Rightarrow P_{x_{\infty}}(X_1 \in \cdot) \tag{3.10}$$

as $k \to \infty$. The continuity of g implies that $\{x : g(x) < r\}$ is open. The weak convergence statement (3.10) implies that

$$\lim_{k \to \infty} P_{x_{n_k}}(g(X_1) < r) \ge P_{x_{\infty}}(g(X_1) < r) \ge 1 - \frac{\delta}{2}.$$
(3.11)

Hence, for $n_k > r$,

$$P_{x_{n_k}}(X_1 \in A_{n_k}^c) = P_{x_{n_k}}(g(X_1) > n_k)$$

$$\leq P_{x_{n_k}}(g(X_1) \ge r)$$

$$= 1 - P_{x_{n_k}}(g(X_1) < r).$$
(3.12)

As a consequence of (3.11) and (3.12), we find that

$$\overline{\lim_{k \to \infty}} P_{x_{n_k}}(X_1 \in A_{n_k}^c) \le 1 - \underline{\lim_{k \to \infty}} P_{x_{n_k}}(g(X_1) < r)$$
$$\le 1 - (1 - \frac{\delta}{2}) \le \frac{\delta}{2},$$

contradicting (3.8) and proving (3.6).

With (3.6) in hand, we find from (3.5) that

$$\overline{\lim_{n \to \infty}} \left| \int_{S} \pi_n(dx) (P_n f)(x) - \int_{S} \pi_n(dx) (Pf)(x) \right| \le 2\epsilon \|f\|$$

whenever $f \in bC$. Since ϵ was arbitrary, we conclude that

$$\int_{S} \pi_n(dx)(P_n f)(x) - \int_{S} \pi_n(dx)(Pf)(x) \to 0$$
(3.13)

as $n \to \infty$. Because of A3, $Pf \in bC$, so (3.2) implies that

$$\int_{S} \pi_{n'_{k}}(dx)(Pf)(x) \to \int_{S} \pi'(dx)(Pf)(x)$$
(3.14)

as $k \to \infty$. In view of (3.3), (3.4), (3.13), and (3.14), we conclude that

$$\int_{S} \pi'(dy) f(y) = \int_{S} \int_{S} \pi'(dx) P(x, dy) f(y)$$

for each $f \in bC$, which implies that

$$\pi'(dy) = \int_S \pi'(dx) P(x, dy).$$

So, π' is therefore a stationary distribution of P, and must coincide with π (due to the assumed uniqueness of π). So, $(\pi_n : n \ge 1)$ can only have one (weak) limit point, namely π , proving the theorem.

The question of when Π_n is non-empty (at least for *n* sufficiently large) must be settled separately. One approach is to impose sufficient conditions on the augmentation P_n so as to guarantee that the Markov chain having the one-step transition kernel P_n is positive Harris recurrent on A_n .

The other obvious approach is to leverage ideas related to A3. In particular, it is known that if P_n is weakly continuous as a one-step transition kernel on A_n , then the compactness of the state space A_n guarantees that Π_n is non-empty; see Karr (1975). However, if P_n fails to be weakly continuous, then Π_n may be empty, even when the augmentation is chosen to preserve as much continuity as possible. **Example 3.** Suppose that S = [0, 2], A = [0, 1], and

$$P(x,dy) = \begin{cases} \frac{1}{2}\delta_{1+x}(dy) + \frac{1}{2}\delta_2(dy), & 0 \le x \le \frac{1}{2}, \\ \frac{1}{2}\delta_{2-x}(dy) + \frac{1}{2}\delta_2(dy), & \frac{1}{2} \le x \le 2, \end{cases}$$

where $\delta_z(\cdot)$ is a unit point mass distribution at $z \in S$. Then, P is weakly continuous on S.

Of course,

$$P_x(X_1 \notin A) = \begin{cases} \frac{1}{2}, & x \in \{0, 1\} \\ 1, & 0 < x < 1 \end{cases}$$

for $x \in A$. Suppose that our augmentation takes the form

$$\tilde{P}(x,dy) = P(x,dy) + P_x(X_1 \notin A)G(x,dy)$$
(3.15)

for $x, y \in A$, where $G = (G(x, dy) : x, y \in A)$ is chosen to be weakly continuous. We claim that even when G is so chosen, \tilde{P} may fail to have a stationary distribution (despite the compactness of its state space). In particular, choose

$$G(x, dy) = \delta_{\frac{x}{2}}(dy)$$

for $x, y \in A$. Then,

$$\tilde{P}(x,dy) = \begin{cases} \frac{1}{2}\delta_1(dy) + \frac{1}{2}\delta_0(dy), & x = 0; \\ \delta_{\frac{x}{2}}(dy), & 0 < x < 1; \\ \frac{1}{2}\delta_1(dy) + \frac{1}{2}\delta_{\frac{1}{2}}(dy), & x = 1. \end{cases}$$

At any initial point $x \in A$, the Markov chain clearly converges weakly to δ_0 . As a consequence, for each $f \in bC$, $x \in A$,

$$\int_{A} f(y)\tilde{P}^{n}(x,dy) \to \int_{A} f(y)\delta_{0}(dy) = f(0)$$

as $n \to \infty$. Hence, if \tilde{P} has a stationary distribution $\tilde{\pi}$, then for $f \in bC$,

$$\int \tilde{\pi}(dy)f(y) = \int_{A} \tilde{\pi}(dx) \int_{A} \tilde{P}^{n}(x, dy)f(y) \to f(0)$$

(by the Bounded Convergence Theorem). So, the only possible choice for $\tilde{\pi}$ is δ_0 . But

$$\int_{A} \delta_0(dx) \tilde{P}(x, dy) = \tilde{P}(0, dy) \neq \delta_0,$$

so δ_0 is not a stationary distribution for \tilde{P} . This example proves that even when augmentations are constructed via (3.15), the augmentation may fail to have a

stationary distribution.

Hence, if one is relying on the compactness of A_n to directly imply that $\Pi_n \neq \emptyset$ (rather than to use, for example, Harris recurrence), one must exercise care in ensuring that P_n is weakly continuous, even when starting with a weakly continuous P.

We finish this section with a generalization of Theorem 2. For $S = \mathbb{R}_+$, we say that X is stochastically monotone if $P_x(X_1 > y)$ is non-decreasing in x for each $y \ge 0$.

Theorem 5. Let X be an \mathbb{R}_+ -valued stochastically monotone Markov chain for which $F(\cdot, y)$ is a continuous function for each $y \ge 0$, where $F(x, y) \stackrel{\Delta}{=} P_x(X_1 \le y)$. Suppose X has a unique stationary distribution $\pi = (\pi(dx) : x \ge 0)$ for which $\pi[a, b) > 0$ for all a, b for which $0 \le a < b < \infty$. Then, if $A_n = [0, n]$, Π_n is non-empty for $n \ge 1$, and $\pi_n \in \Pi_n$, we have that

$$\pi_n \Rightarrow \pi$$

as $n \to \infty$.

Proof. We start by recognizing that we can apply the argument of Theorem 2 to establish the existence of a strictly increasing sequence $(r(k) : k \ge 1)$ converging to infinity such that

$$\sum_{k=0}^{\infty} r(k+1)\pi([k,k+1)) < \infty.$$

By setting r(0) = 0 and defining $r(\cdot)$ between consecutive integers via linear interpolation, we construct a continuous function $r(\cdot)$ such that

$$\int_{\mathbb{R}_+} r(x)\pi(dx) < \infty.$$

We next take advantage of the uniqueness of π to guarantee that the shift operator defined by $\theta \circ X = (X_{1+n} : n \ge 0)$ is not only measure-preserving when X_0 has distribution π but is also ergodic; see p. 141 of Ash and Gardner (1975). Let $T_0 = 0$ and $T_{i+1} = \inf\{n > T_i : X_n \le 1\}$ for $i \ge 0$. The ergodic theorem for stationary sequences implies that

$$\frac{1}{T_n} \sum_{j=0}^{T_n-1} r(X_j) \to \int_{\mathbb{R}_+} r(x)\pi(dx) \qquad \text{a.s.}$$
(3.16)

and

$$\frac{\sum_{j=0}^{n-1} \sum_{i=T_j}^{T_{j+1}-1} r(X_i)}{\sum_{j=0}^{n-1} (T_{j+1} - T_j)} \to \frac{\int_{[0,1]} \pi(dx) E_x \sum_{i=0}^{T_1-1} r(X_i)}{\int_{[0,1]} \pi(dx) E_x T_1} \qquad \text{a.s.}$$

as $n \to \infty$, so that we may conclude that

$$\int_{\mathbb{R}_{+}} r(x)\pi(dx) = \frac{\int_{[0,1]} \pi(dx) E_x \sum_{j=0}^{T_1-1} r(X_j)}{\int_{[0,1]} \pi(dx) E_x T_1}.$$

In view of (3.16), we find that

$$\int_{[0,1]} \pi(dx) k(x) < \infty$$
 (3.17)

where

$$k(x) \stackrel{\Delta}{=} E_x \sum_{j=0}^{T_1-1} r(X_j).$$

Let $\beta_n = \inf\{j \ge 0 : X_j > n\}$ and observe that the ergodic theorem implies that $\beta_n < \infty$ a.s. when X_0 has distribution π . It is therefore evident that the set of x-values in [0, 1] for which $P_x(\beta_n < T_1) > 0$ must have positive π -probability and the set of x-values for which $k(x) < \infty$ must have π -probability one, so that there exists $w \le 1$ for which $P_w(\beta_n < T_1) > 0$ and $k(w) < \infty$. Hence,

$$\int_{(n,\infty)} P_w(X_{\beta_n} \in dy, \beta_n < T_1)k(y) \le k(w) < \infty,$$

so that there exists $y_n > n$ for which $k(y_n) < \infty$.

The stochastic monotonicity of X and the monotonicity of r imply that $k(\cdot)$ is a non-decreasing function, so that $k(z) < \infty$ for $z \leq y_n$, and hence $k(z) < \infty$ for all $z \in \mathbb{R}_+$. Furthermore, the continuity of r and weak continuity of P ensure that k is continuous on $(1, \infty)$; see Glynn and Infanger (2022) for related arguments.

Conditioning on X_1 establishes that for x > 1,

$$k(x) = r(x) + \int_{(1,\infty)} P_x(X_1 \in dy)k(y).$$

So, if we set $\tilde{k}(x) = k(x)$ for x > 1 and $\tilde{k}(x) = k(1+)$ for $x \le 1$, we find that

$$(P\tilde{k})(x) = k(1+)P_x(X_1 \le 1) + \int_{(1,\infty)} P_x(X_1 \in dy)k(y)$$
$$\le k(1+) + \int_{(1,\infty)} P_{1+}(X_1 \in dy)k(y)$$
$$= k(1+) + k(1+) - r(1)$$
$$= k(1+) + \tilde{k}(x) - r(1)$$

for $x \in [0, 1]$. As a result, A2 is satisfied with a continuous \tilde{k} , so that we can apply Theorems 3 and 4. Finally, observe that because \tilde{k} is non-decreasing, it is evident that when $A_n = [0, n]$,

$$\sup_{x \in A_n} \tilde{k}(x) \le \inf_{x \in A_n^c} \tilde{k}(x),$$

so that A_n 's of this special form are legitimate truncation sets.

This result generalizes Gibson and Seneta (1987b) and Tweedie (1998) to continuous state space. In particular, for stochastically monotone Markov chains, general augmentations yield convergent approximations when the truncation sets A_n are chosen to be of the form $A_n = [0, n]$.

4 Convergence for Strongly Uniformly Recurrent Markov Chains

In this section, we use the theory of regeneration to establish convergence of general augmentation schemes for a class of general state space Markov chains that generalize the theory developed for Markov chains having Markov transition matrices. In contrast to the earlier sections, the theory developed here pertains to arbitrary truncation sequences, so that $(A_n : n \ge 1)$ can be any sequence for which $\emptyset \neq A_1 \subseteq A_2 \subseteq ...$ for which $\bigcup_{n=1}^{\infty} A_n = S$.

We say that $X = (X_n : n \ge 0)$ is strongly uniformly recurrent if there exists $\lambda > 0$ and a probability ϕ such that

$$P(x, dy) \ge \lambda \phi(dy) \tag{4.1}$$

for $x, y \in S$.

Remark 8. Uniform recurrence requires the existence of $m \ge 1$ such that

$$P_x(X_m \in dy) \ge \lambda \phi(dy) \tag{4.2}$$

for $x, y \in S$, so that (4.1) is clearly a strong version of (4.2). We note that Example 1 satisfies (4.2) with m = 2, and yet presents a setting in which general augmentation schemes can fail to converge. So, some condition (like strong uniform recurrence) is needed beyond uniform recurrence in order to guarantee convergence for general augmentations.

Without any real loss of generality, we may assume that ϕ is supported on A_1 (at the possible cost of needing to reduce λ in (4.1)). As in our earlier sections, we allow a general augmentation $P_n = (P_n(x, dy) : x, y \in A_n)$ associated with A_n to take the form

$$P_n(x, dy) = P(x, dy) + R_n(x, dy)$$

for $x, y \in A_n$, where P_n is a stochastic kernel, and R_n is a non-negative kernel.

In the presence of (4.1), Athreya and Ney (1978) and Nummelin (1978) observed that the transition kernel P can be put in the form

$$P(x, dy) = \lambda \phi(dy) + (1 - \lambda)H(x, dy)$$
(4.3)

for $x, y \in S$, where $H = (H(x, dy) : x, y \in S)$ is a stochastic kernel. (Note that H is defined so as to make (4.3) valid.) With the "splitting representation" (4.3)

in hand, we can see that the right-hand side is a mixture of two distributions. Consequently, we have a probabilistic mechanism for how to envision the transitions of the Markov chain that evolves according to P. In particular, if $X_n = x$, we first flip a λ -coin having probability of "heads" given by λ . If the coin comes up heads, we distribute X_{n+1} according to ϕ , and X regenerates at that time. Otherwise, we distribute X_{n+1} according to $H(X_n, \cdot)$. If we let $P_{\phi}(\cdot)$ and $E_{\phi}(\cdot)$ denote the probability and expectation under which X_0 has distribution ϕ , the theory of regenerative processes (see, for example, Smith (1955)) asserts that the unique stationary distribution π associated with P is given by

$$\pi(\cdot) = \frac{E_{\phi} \sum_{j=0}^{\tau-1} I(X_j \in \cdot)}{E_{\phi} \tau}$$

$$\tag{4.4}$$

where τ is the first time at which X distributes itself according to ϕ .

Because ϕ is supported on A_1 , we note that

$$P_n(x,dy) = \lambda \phi(dy) + q_n(x)H(x,dy) + r_n(x)R_n(x,dy)$$

$$(4.5)$$

where $\lambda + q_n(x) + r_n(x) = 1$ for $x \in A_n$, $q_n(x)$, $r_n(x) \ge 0$ and $R_n = (R_n(x, dy) : x, y \in A_n)$ is a stochastic kernel. In view of (4.5), we can view the transitions of X under P_n as being implemented through a more complex randomization. In particular, if $X_m = x$, then with probability λ , X_{m+1} distributes itself according to ϕ , and X regenerates. On the other hand, with probability $q_n(X_m)$, X_{m+1} distributes itself according to $H(X_m, \cdot)$, while with probability $r_n(X_m)$, X_{m+1} distributes itself according to $R_n(X_m, \cdot)$. As in the discussion of the dynamics of X under the transition kernel P, we let τ be the first time at which X regenerates and distributes itself according to ϕ . Also, let $\beta_n - 1$ be the first time at which X_{m+1} is drawn from the distribution $R_n(X_m, \cdot)$, so that the conditional distribution of β_n has probability mass function

$$P(\beta_n = k+1|X_0, X_1, ..., X_k) = \prod_{j=0}^{k-1} (1 - r_n(X_j))r_n(X_k).$$
(4.6)

A final key observation is that

$$P_{\phi}((X_0, X_1, ..., X_k) \in \cdot, \tau \land \beta_n > k) = P_{\phi}^n((X_0, ..., X_k) \in \cdot, \tau \land \beta_n > k) \quad (4.7)$$

for $k \geq 0$, where $P_{\phi}^{n}(\cdot)$ is the probability under which X_{0} has distribution ϕ and X evolves under P_{n} .

We are now ready to state our main result.

Theorem 6. Suppose that X is strongly uniformly recurrent under P with a unique stationary distribution π . Then, X is strongly uniformly recurrent under P_n with a unique stationary distribution π_n , and

$$\sup_{A\subseteq S} |\pi_n(A) - \pi(A)| \to 0$$

as $n \to \infty$.

Note that π_n converges to π in total variation norm (rather than in the sense of weak convergence used elsewhere in this paper), regardless of how the A_n 's are chosen, and regardless of how the sequence of augmentations is defined.

Proof. We first recognize that since X regenerates at time τ under P_{ϕ}^{n} , π_{n} can be expressed in terms of regenerative cycle quantities (as in (4.4)) as

$$\pi_n(\cdot) = \frac{E_{\phi}^n \sum_{j=0}^{\tau-1} I(X_j \in \cdot)}{E_{\phi}^n \tau},$$

where $E_{\phi}^{n}(\cdot)$ is the expectation associated with P_{ϕ}^{n} . We now couple the dynamics of X under P_{ϕ}^{n} to its evolution under P_{ϕ} by drawing X_{m} from $H(X_{m-1}, \cdot)$ under P_{ϕ} whenever we draw X_{m} from $R_{n}(X_{m-1}, \cdot)$ under P_{ϕ}^{n} . Then, on account of (4.7),

$$\begin{aligned} |E_{\phi}^{n} \sum_{j=0}^{\tau-1} I(X_{j} \in \cdot) - E_{\phi} \sum_{j=0}^{\tau-1} I(X_{j} \in \cdot)| \\ &= |\sum_{j=0}^{\infty} P_{\phi}^{n}(X_{j} \in \cdot, \beta_{n} \wedge \tau > j) + E_{\phi}^{n} \sum_{j=\beta_{n}}^{\tau-1} I(X_{j} \in \cdot, \tau > \beta_{n}) \\ &- \sum_{j=0}^{\infty} P_{\phi}(X_{j} \in \cdot, \beta_{n} \wedge \tau > j) - E_{\phi} \sum_{j=\beta_{n}}^{\tau-1} I(X_{j} \in \cdot, \tau > \beta_{n})| \\ &\leq E_{\phi}^{n}(\tau - \beta_{n})I(\tau > \beta_{n}) + E_{\phi}(\tau - \beta_{n})I(\tau > \beta_{n}) \\ &\leq E_{\phi}^{n}\tau I(\tau > \beta_{n}) + E_{\phi}\tau I(\tau > \beta_{n}) \\ &\leq (E_{\phi}^{n}\tau^{2})^{1/2}P_{\phi}^{n}(\tau > \beta_{n})^{1/2} + (E_{\phi}\tau^{2})^{1/2}P_{\phi}(\tau > \beta_{n})^{1/2}. \end{aligned}$$

Since τ is geometric with parameter λ under both P_{ϕ} and P_{ϕ}^{n} , $E_{\phi}^{n}\tau^{2} = E_{\phi}\tau^{2} \leq 2/\lambda^{2}$. On the other hand,

$$P_{\phi}^{n}(\tau > \beta_{n}) = \sum_{j=1}^{\infty} P_{\phi}^{n}(\beta_{n} = j, \tau > j)$$
$$= \sum_{j=1}^{\infty} E_{\phi}^{n} \prod_{k=0}^{j-2} q_{n}(X_{k})r_{n}(X_{j-1})$$
$$= \sum_{j=1}^{\infty} E_{\phi} \prod_{k=0}^{j-2} q_{n}(X_{k})r_{n}(X_{j-1})$$
$$= P_{\phi}(\tau > \beta_{n}).$$

Since $q_n(X_k) \leq 1 - \lambda$, it follows that

$$\prod_{k=0}^{j-1} q_n(X_k) r_n(X_k) \le (1-\lambda)^j.$$

But $r_n(X_j) \downarrow 0$ a.s. as $n \to \infty$, so the Dominated Convergence Theorem implies that $P_{\phi}(\tau > \beta_n) = P_{\phi}^n(\tau > \beta_n) \to 0$ as $n \to \infty$. Consequently,

$$\sup_{B} |E_{\phi}^{n} \sum_{j=0}^{\tau-1} I(X_{j} \in B) - E_{\phi} \sum_{j=0}^{\tau-1} I(X_{j} \in B)| \to 0$$

and this implies that $E_{\phi}^{n}\tau \to E_{\phi}\tau$ as $n \to \infty$, thereby proving the theorem. \Box

5 Convergence of General Augmentations for Markov Jump Processes

We now briefly describe the extension of our discrete time theory to the setting of Markov jump processes. In particular, suppose that S is a finite or countably infinite state space. We say that $Q = (Q(x,y) : x, y \in S)$ is a *rate matrix* if $Q(x, y) \ge 0$ for $x \ne y$,

$$\lambda(x) \stackrel{\Delta}{=} \sum_{y \neq x} Q(x, y) < \infty$$

and

$$\lambda(x) = -Q(x, x)$$

for $x \in S$. We shall assume that the associated Markov jump process X is non-explosive; Meyn and Tweedie (1993) provide a Lyapunov condition that guarantees non-explosiveness. In the presence of non-explosiveness, $X = (X(t) : t \ge 0)$ can be realized as a stochastic process having piecewise constant paths that are right continuous. Miller (1963) shows that when X is non-explosive, then positive recurrence is equivalent to finding a stationary distribution $\pi = (\pi(x) : x \in S)$ satisfying

$$\pi Q = 0, \tag{5.1}$$

in which case π is the stationary distribution of X.

For a given truncation $A_n \subseteq S$, we say that Q_n is an *augmentation* of Q if Q_n is a rate matrix on A_n for which $Q_n(x,y) \ge Q(x,y)$ for all $x \ne y$ with $x, y \in A_n$.

A4. There exist non-negative coercive functions $r = (r(x) : x \in S)$ and $g = (g(x) : x \in S)$, and $b < \infty$ for which

$$(Qg)(x) \le -r(x) + b$$

for $x \in S$.

Assumption A4 is the continuous-time analog of A1. The continuous-time analog of Theorem 1 is our final result.

Theorem 7. Suppose X is an irreducible non-explosive Markov jump process with rate matrix Q and probability π satisfying (5.1). If A4 is in force, then for any augmentation sequence $(Q_n : n \ge 1)$ for which $A_n = K_n(g)$, we have that

 $\pi_n \Rightarrow \pi$

as $n \to \infty$, provided that π_n is a probability satisfying $\pi_n Q_n = 0$.

The proof of Theorem 7 is essentially identical to that of Theorem 1. The key observation is that $\pi_n r \leq b$ for $n \geq 1$, (from Glynn and Zeevi (2008)) so that $(\pi_n : n \geq 1)$ is again tight. Continuous-time analogs to all our other main results can be similarly derived.

References

- Ash, R. B. and Gardner, M. F. (1975). Topics in Stochastic Processes. Probability and Mathematical Statistics. Academic Press, New York.
- Athreya, K. B. and Ney, P. (1978). A new approach to the limit theory of recurrent Markov chains. *Transactions of the American Mathematical Society*, 245:493–501.
- Billingsley, P. (1968). Convergence of Probability Measures. John Wiley & Sons, New York.
- Borovkov, A. A. (1998). Ergodicity and Stability of Stochastic Processes. J. Wiley, Chichester.
- Copson, E. (1968). Metric Spaces. Cambridge University Press, Cambridge.
- Gibson, D. and Seneta, E. (1987a). Augmented truncations of infinite stochastic matrices. Journal of Applied Probability, 24(3):600–608.
- Gibson, D. and Seneta, E. (1987b). Monotone infinite stochastic matrices and their augmented truncations. *Stochastic Processes and their Applications*, 24(2):287–292.
- Glynn, P. W. and Infanger, A. (2022). Solutions of Poisson's equation for stochastically monotone Markov chains. arXiv:2202.10578 [math.PR]. To appear in Queueing Models and Service Management.
- Glynn, P. W. and Zeevi, A. (2008). Bounding stationary expectations of Markov processes. In *Markov processes and related topics: a Festschrift for Thomas G. Kurtz*, pages 195–214. Institute of Mathematical Statistics.
- Golub, G. H. and Seneta, E. (1974). Computation of the stationary distribution of an infinite stochastic matrix of special form. Bulletin of the Australian Mathematical Society, 10(2):255–261.

- Infanger, A. and Glynn, P. W. (2022). On convergence of a truncation scheme for approximating stationary distributions of continuous state space Markov chains and processes. *In preparation*.
- Karr, A. F. (1975). Weak convergence of a sequence of Markov chains. Zeitschrift fuer Wahrscheinlichkeitstheorie und Verwandte Gebiete, 33(1):41–48.
- Liu, Y. (2010). Augmented truncation approximations of discrete-time Markov chains. Operations Research Letters, 38(3):218–222.
- Meyn, S. P. and Tweedie, R. L. (1993). Stability of Markovian processes III: Foster–Lyapunov criteria for continuous-time processes. Advances in Applied Probability, 25(3):518–548.
- Meyn, S. P. and Tweedie, R. L. (2009). *Markov Chains and Stochastic Stability*. Cambridge University Press, Cambridge, second edition.
- Miller, R. G. (1963). Stationarity equations in continuous time Markov chains. Transactions of the American Mathematical Society, 109(1):35–44.
- Nummelin, E. (1978). A splitting technique for Harris recurrent Markov chains. Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete, 43(4):309–318.
- Seneta, E. (1980). Computing the stationary distribution for infinite Markov chains. *Linear Algebra and Its Applications*, 34:259–267.
- Smith, W. L. (1955). Regenerative stochastic processes. Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences, 232(1188):6–31.
- Tweedie, R. L. (1998). Truncation approximations of invariant measures for Markov chains. Journal of Applied Probability, pages 517–536.
- Wolf, D. (1980). Approximation of the invariant probability measure of an infinite stochastic matrix. Advances in Applied Probability, 12(3):710–726.