Characterizations of graph classes via convex geometries: a survey

Mitre C. Dourado

Instituto de Computação, Universidade Federal do Rio de Janeiro, Brazil

Marisa Gutierrez

CMaLP, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, Argentina

Fábio Protti

Instituto de Computação, Universidade Federal Fluminense, Brazil

Rudini Sampaio

Departamento de Computação, Universidade Federal do Ceará, Brazil

Silvia Tondato

CMaLP, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, Argentina

Abstract

Graph convexity has been used as an important tool to better understand the structure of classes of graphs. Many studies are devoted to determine if a graph equipped with a convexity is a *convex geometry*. In this work, we survey results on characterizations of well-known classes of graphs via convex geometries. We also give some contributions to this subject.

Keywords: Convex geometry, Convexity, Convex hull, Graph convexity

Email addresses: mitre@ic.ufrj.br (Mitre C. Dourado), marisa@mate.unlp.edu.ar (Marisa Gutierrez), fabio@ic.uff.br (Fábio Protti), rudini@dc.ufc.br (Rudini Sampaio), tondato@mate.unlp.edu.ar (Silvia Tondato)

Mitre C. Dourado is partially supported by CNPq, Brazil (305404/2020-2 and 403601/2023-1) and FAPERJ, Brazil (SEI-260003/015314/2021); Fábio Protti is partially supported by CNPq (304117/2019-6) and FAPERJ (E-26/201.083/2021); Rudini Sampaio is partially supported by CNPq (404479/2023-5 and 311070/2022-1) and CAPES (88881.712024/2022-01).

1. Introduction

A convexity on a nonempty set V is a family \mathscr{C} of subsets of V (called convex sets) such that: (a) \emptyset and V are convex sets; (b) the intersection of convex sets is a convex set. If V = V(G) for some graph G, then \mathscr{C} is a graph convexity of G. The structure of a convexity is based on the way convex sets are defined. In the context of graph convexities, there are many examples in the literature where convex sets are defined over a path system, typically according to the following rule: fix a family \mathscr{P}_G of paths (or, more generally, walks) of a graph G, and say that a set $S \subseteq V(G)$ is convex if no vertex $x \notin S$ lies in a member of \mathscr{P}_G that starts and ends at two vertices of S. The most natural example is the so-called geodesic convexity [38], for which \mathcal{P}_G is the family of all the *shortest paths* of G. Other important examples are: the monophonic convexity [17, 20], the m^3 -convexity [19], the l^k -convexity [27], the toll convexity [2], and the weakly toll convexity [28], defined over induced paths, induced paths of length at least three, induced paths of length at most k, tolled walks, and weakly toll walks, respectively. Other rules to define convex sets, not based on path systems, have also been investigated; an important example comes from the so-called *Steiner convexity*, introduced in [9]. In this case, a set $S \subseteq V(G)$ is Steiner convex if, for every $S' \subseteq S$, the vertices of any Steiner tree with terminal set S' belong to S.

Graph convexities have been studied in many contexts. A strong direction of research has focused on determining convexity invariants, such as the *hull number*, the *interval number*, and the *convexity number*, among others. A major reference work by Pelayo [38] gives an extensive overview on convexity invariants, applied to the case of the geodesic convexity.

Other studies are devoted to determine if a graph convexity is a *convex geometry*. This line of research is closely linked to the original idea of defining a combinatorial abstraction of convex sets in Geometry. The convex sets in the plane are those subsets S such that no point outside S lies on a line segment with endpoints in S. The *convex hull* of a set of points A in the plane is the smallest convex set that contains A. A point x of a convex set S is an *extreme point* of S if $S \setminus \{x\}$ is also a convex set, that is, if there is no line segment with endpoints in $S \setminus \{x\}$ containing x. For instance, an n-polygon (considered as a closed region) is clearly a convex set in the plane,

and its extreme points are precisely its n vertices². In addition, the polygon is the convex hull of its n extreme points. Another example is a circle (closed circular region); it is clearly a convex set in the plane, and all the points lying on the circle border are its extreme points. Again, the circle is the convex hull of such points.

The above concepts can be transferred to the combinatorial field in a natural way. We refer the reader to [23]. Let G be a graph and let $\mathscr E$ be a convexity of G. Given a set $S \subseteq V(G)$, the smallest set $H \in \mathscr{C}$ containing S is called the convex hull of S. A vertex x of a convex set S is an extreme vertex of S if $S \setminus \{x\}$ is also convex. The convexity \mathscr{C} is a convex geometry if it satisfies the Krein-Milman (or Minkowski-Krein-Milman) property [34, 37]: Every convex set is the convex hull of its extreme vertices. The main question dealt with in this survey is: by fixing a rule r to define the convex sets (e.g., a rule based on some path system), determine the class of graphs whose rconvexities are convex geometries. For instance, by fixing induced paths, we can obtain the following characterization: a graph G is chordal if and only if the monophonic convexity of G is a convex geometry [23]. Ptolemaic graphs, interval graphs, proper interval graphs, weak bipolarizable graphs, and 3fan-free chordal graphs can also be characterized in this way by considering, respectively, the geodesic convexity [23], the toll covexity [2], the weakly toll convexity [28], the m^3 -convexity [19], and the Steiner convexity [9]. In [27], a characterization of graphs with l^k -convexities that are convex geometries are studied. Section 3 discusses in detail most of these characterizations.

Convex geometries are equivalent, by complementation, to antimatroids (see [32]). An antimatroid consists of a finite family \mathcal{F} of finite sets (called feasible sets) such that: (a) \mathcal{F} is closed under unions; (b) if S is a nonempty feasible set, then there is $x \in S$ for which $S \setminus \{x\}$ is also a feasible set (that is, \mathcal{F} is an accessible set system). If U is the union of the sets in \mathcal{F} , then the family of complementary sets $\mathscr{C} = \{U \setminus S \mid S \in \mathcal{F}\}$ is a convex geometry. It is not difficult to see that \mathscr{C} is a convexity; in addition, it satisfies the anti-exchange property [21, 32], which is complementary to the accessibility property of antimatroids (property (b) above). In Section 2, we present in more detail the anti-exchange property, which will be useful to derive some results in Section 4.

This survey is organized as follows. Section 2 provides the necessary

²Here, the term *vertices* is used in the geometric sense.

background. In Section 3, we review the main results in the literature on characterizations of graph classes via convex geometries; chordal, Ptolemaic, strongly chordal, interval, proper interval graphs, among other classes, are dealt with. In Section 4 we present some contributions; namely, we show that forests, forests of stars, cographs, bipartite graphs, planar graphs, and triangle-free graphs can be well characterized via convex geometries. Our concluding remarks are presented in Section 5.

2. Preliminaries

All the graphs in this work are connected, unless otherwise stated. A path (resp., cycle) in a graph G is induced if there is no edge of G linking two nonconsecutive vertices of it. We denote by P_n (resp., C_n) the induced path (resp., cycle) with n vertices. For a vertex $v \in V(G)$, let N(v) denote the open neighborhood (or simply neighborhood) of v in G. In addition, let N[v] denote the closed neighborhood of v in G, where $N[v] = N(v) \cup \{v\}$. If N[v] is a clique, then v is said to be a simplicial vertex. For a set $S \subseteq V(G)$, we denote by G[S] the subgraph of G induced by S, defined as follows: V(G[S]) = S and $E(G[S]) = \{xy \in E(G) \mid x,y \in S\}$. For a graph F, we say that G is F-free if G does not contain F as an induced subgraph. For a family $\mathcal{F} = \{G_1, G_2, \ldots, G_k\}$ of prefixed graphs, we say that G is (G_1, G_2, \ldots, G_k) -free (or simply \mathcal{F} -free) if G is G_i -free for $1 \leq i \leq k$. For $\mathcal{C} = \{C_k \mid k \geq 4\}$, G is said to be a chordal graph if G is C-free.

Let \mathscr{P} be a function (called *path system*) that maps each connected graph G to a collection \mathscr{P}_G of walks of G. For instance, \mathscr{P} can map G to its shortest paths, induced paths, tolled walks etc. The members of \mathscr{P}_G are generically called p-walks of G. All the path systems dealt with in this paper satisfy the following property: For every G, all the walks of G formed by a single vertex or two adjacent vertices are p-walks.

The interval I(u, v) of $u, v \in V(G)$ is the set of all vertices that lie in a p-walk from u to v. For $S \subseteq V(G)$, we define $I(S) = \bigcup_{u,v \in S} I(u,v)$. A set $S \subseteq V(G)$ is convex if I(S) = S. Let $\mathscr{C} = \{S \subseteq V(G) : I(S) = S\}$. Clearly, \mathscr{C} is a convexity of G, defined over the collection \mathscr{P}_G of p-walks.

A vertex x of a convex set $S \subseteq V(G)$ is an extreme vertex of S if $S \setminus \{x\}$ is also a convex set. This definition implies that: (a) no p-walk in G[S] contains x as an internal vertex; (b) $x \notin I(u,v)$ for any pair of distinct vertices $u, v \in S$. The set of extreme vertices of S is denoted by ext(S). The convex hull of a set $S \subseteq V(G)$, denoted by H(S), is the smallest set

convex set of G containing S. It can be shown that H(S) can be iteratively constructed by successive applications of the interval operator, as follows: starting with a set $S \subseteq V(G)$, define $S_0 = S$ and $S_i = I(S_{i-1}), i \ge 1$. Then, there exists $k \ge 1$ such that $S_k = S_{k-1} = H(S)$.

The concept of interval is the combinatorial analog of a line segment between two points in the plane. Just as the geometric convex hull of a set of points in the plane can be built using line segments, the convex hull of a set S of vertices in a graph uses the notion of interval for the iterative construction described above.

We say that a convexity \mathscr{C} of a graph G is a convex geometry (or that it is geometric) if it satisfies the following property: for every $S \in \mathscr{C}$, it holds that S = H(ext(S)). This is called the Krein-Milman property [34]. Alternatively, the convexity \mathscr{C} is a convex geometry if the following axiom is satisfied:

Anti-exchange property [21, 32] If $y, z \notin H(S)$ and $z \in H(S \cup \{y\})$, then $y \notin H(S \cup \{z\})$.

The anti-exchange property is a combinatorial abstraction of the usual convex hull operator in the plane: for two points y and z not in the convex hull of S, if z is in the convex hull of $S \cup \{z\}$. See Figure 1.




Figure 1: Anti-exchange property applied to $S = \{a, b, c, d\}$.

3. An overview of graph classes and convex geometries

In this section, we briefly review the main existing results on convex geometries in the literature on graph convexity. Chordal graphs play in important role in this section. For basic facts on chordal graphs, see [26].

3.1. Monophonic convexity

Let \mathscr{P}^m be the path system that maps each graph G to its induced paths. The convexity defined over the induced paths of G is called the *monophonic convexity* of G. It is easy to see that v is an extreme vertex of a (monophonically) convex set $S \subseteq V(G)$ if and only if v is a simplicial vertex of G[S].

Suppose that G contains an induced cycle C with at least four vertices. Clearly, S = H(V(C)) is convex and contains no simplicial vertices, i.e., $ext(S) = \emptyset$. Thus, the Krein-Milman property fails in this case. In other words, if the monophonic convexity of G is a convex geometry, then G is chordal. On the other hand, if G is chordal, then every nonsimplicial vertex of G lies on an induced path between two simplicial vertices [23]. As chordality is an hereditary property, this implies that every convex subset of vertices of G is indeed the convex hull of its extreme vertices. Hence:

Theorem 1. [23] A graph G is chordal if and only if the monophonic convexity of G is a convex geometry.

3.2. Geodesic convexity

A path P between two vertices u and v in a graph G is a shortest path if the length of P is equal to the distance between u and v in G. Let \mathscr{P}^g be the path system that maps each graph G to its shortest paths. The convexity defined over the shortest paths of G is the geodesic convexity of G.

As in the previous subsection, it is easy to see that v is an extreme vertex of a (geodesically) convex set $S \subseteq V(G)$ if and only if v is a simplicial vertex of G[S]. In addition, if the geodesic convexity of G is a convex geometry, then G must be clearly chordal. However, it is not true that, for every chordal graph G, the geodesic convexity of G is a convex geometry. For instance, consider the graph G' (called gem) with $V(G') = \{a, b, c, d, e\}$ and $E(G') = \{ab, bc, cd, de, ae, be, ce, de\}$. See Figure 2. Assume that G' is an induced subgraph of a chordal graph G. Clearly, the extreme vertices of Y = H(V(G')) are in $\{a, d\}$. However, $H(\{a, d\}) \neq Y$.

A graph G is Ptolemaic if it is chordal and gem-free. The arguments in the previous paragraph tell us that if the geodesic convexity of G is a convex geometry, then G is a Ptolemaic graph. Conversely, if G is a Ptolemaic graph, then it can be proved that every induced path of G is a shortest path [30], i.e., the monophonic and the geodesic convexities of G coincide in this case. Therefore, in a Ptolemaic graph, every nonsimplicial vertex of G




Figure 2: A gem G' with $V(G') = \{a, b, c, d, e\}$.

lies on a *shortest* path between two simplicial vertices. Consequently, every convex subset of vertices of G is indeed the convex hull of its extreme vertices. Hence:

Theorem 2. [23] A graph G is Ptolemaic if and only the geodesic convexity of G is a convex geometry.

3.3. Strong convexity

We say that a path $P = u_0 u_1 \dots u_n$ is even-chorded if it has no odd chord (an edge that connects two vertices that are an odd distance d > 1 apart from each other in a path or cycle) and, in addition, neither u_0 nor u_n lies in a chord of P. Let \mathscr{P}^s be the path system that maps a graph G to its even-chorded paths. The convexity defined over such paths is called the strong convexity of G, and the associated convex sets are called strong convex sets.

An even cycle is a cycle with an even number of vertices. A graph is strongly chordal [22] if it is chordal and every of its even cycles with at least six vertices has an odd chord. A vertex of a graph is simple if the neighborhoods of its neighbors form a nested family of sets. Note that a simple vertex is simplicial, but not conversely. In [22], the following characterization of strongly chordal graphs is given:

Theorem 3. [22] A graph G is a strongly chordal graph if and only if every induced subgraph of G contains a simple vertex.

Suppose that the strong convexity of G is a convex geometry. Since induced paths are even-chorded, an extreme vertex v of an strong convex set S must be simplicial in G[S]. This implies that G is chordal, and, in this case, we can prove the following lemma:

Lemma 4. [23] Let G be a chordal graph. Then, a vertex $v \in V(G)$ is an extreme vertex of an strong convex set S if and only if v is a simple vertex in G[S].

By Theorem 3 and Lemma 4, G is a strongly chordal graph. Conversely, suppose that G is a strongly chordal graph. In [23], the following result is proved:

Lemma 5. [23] In a strongly chordal graph, every nonsimple vertex lies in an even-chorded path between simple vertices.

By Lemma 5, V(G) is indeed the convex hull of its extreme vertices. But since strongly chordal graphs are hereditary, we have:

Theorem 6. [23] A graph G is strongly chordal if and only if the strong convexity of G is a convex geometry.

3.4. m^3 -convexity

Let $\mathscr{P}^{\geq 3}$ be the path system that maps a graph G to its induced paths of length at least three. The convexity defined over such paths is called the m^3 -convexity of G, and the associated convex sets are called m^3 -convex sets. Note that an m^3 -convex set does not necessarily induce a connected subgraph.

An alternative definition of "simplicial" is: v is a simplicial vertex iff it is not a midpoint of an induced P_3 . In [31], this concept is relaxed as follows: a vertex is semisimplicial if it is not an internal vertex of an induced P_4 , and nonsemisimplicial otherwise. Clearly, a vertex v is an extreme vertex of an m^3 -convex set $S \subseteq V(G)$ if and only if v is semisimplicial in G[S].

A graph G is weak bipolarizable [36] if G contains no hole, house, domino, or A as an induced subgraph. See Figure 3. Weak bipolarizable graphs are also called HHDA-free graphs.




Figure 3: From left to right: hole, house, domino, A.

Using arguments similar to those presented in the preceding subsections, it can be shown if the m^3 -convexity of G is a convex geometry, then G is a weak bipolarizable graph. Suppose the m^3 -convexity of G is a convex geometry and G contains, for instance, a house G' as an induced subgraph. Then, the only possible semisimplicial vertex in the subgraph of G induced by Y = H(V(G')) is vertex e, i.e., $ext(Y) \subseteq \{e\}$. This implies $H(ext(Y)) \neq Y$, a contradiction. Analogously, G cannot contain a hole, a domino, or an G as an induced subgraph. Thus, G is weak bipolarizable. Conversely, if G is weak bipolarizable, it holds that every nonsemisimplicial vertex lies on an induced path of length at least three between semisimplicial vertices [19]. We then have the following theorem:

Theorem 7. [19] A graph G is weak bipolarizable if and only if the m^3 -convexity of G is a convex geometry.

3.5. Steiner convexity

Let $S \subseteq V(G)$. A Steiner tree for S is a connected subgraph T of G such that $S \subseteq V(T)$ and |E(T)| is minimum. The Steiner interval I(S) is the set formed by all vertices of G that lie in some Steiner tree for S. We say that S is Steiner convex if, for every $S' \subseteq S$, it holds that $I(S') \subseteq S$. The Steiner convexity of G is the family formed by the Steiner convex sets of G. Important references on the Steiner convexity are [9, 16].

In [9], the authors prove that: (a) the extreme vertices of a Steiner convex set S of G are the simplicial vertices of G[S]; (b) in a (house,hole,domino)-free graph G, the Steiner convexity and the geodesic convexity coincide. Such facts are the main ingredients to prove the following characterization:

Theorem 8. [9] A graph G is Ptolemaic if and only if the Steiner convexity of G is a convex geometry.

Proof. Suppose that G is Ptolemaic, i.e., G is chordal and gem-free. Then G is clearly (house,hole,domino)-free, so the Steiner convexity of G coincides with the geodesic convexity of G, which is a convex geometry by Theorem 2.

Conversely, suppose that the Steiner convexity of G is a convex geometry. We proceed with arguments that we have used before. If G contains an induced gem G', then, clearly, the extreme vertices of Y = H(V(G')) are in $\{a,d\}$ (see Figure 2). However, $H(\{a,d\}) \neq Y$. Thus, G is gem-free. Likewise, if G contains an induced cycle G with at least four vertices then G is Steiner convex and contains no simplicial vertices, i.e., G is chordal. G

In [10], an alternative way of defining Steiner convexity is proposed, by looking at subsets with a prescribed size. We say that S is k-Steiner convex or g_k -convex if, for every $S' \subseteq S$ with |S'| = k, it holds that $I(S') \subseteq S$. The convexity associated with the g_k -convex sets of G is called the k-Steiner convexity of G. Note that the 2-Steiner convexity of G is precisely the geodesic convexity of G.

Let us study the case k = 3. In [10], the extreme vertices of g_3 -convex sets are characterized: a vertex x is an extreme vertex of a g_3 -convex set S of G if and only if x is not the central vertex of an induced claw, paw, or P_4 contained in G[S]. See Figure 4.




Figure 4: From left to right: claw, paw, P_4 . Central vertices are indicated by arrows.

Suppose that the 3-Steiner convexity of G is a convex geometry. Suppose also that G contains an induced graph P isomorphic to a $P_4 = abcd$, where a and d have degree one in P. Observe that a and d are the only extreme vertices of P. Thus, ext(V(P)) is a set $C \subseteq \{a,d\}$. However, $H(C) = C \neq H(\{a,b,c,d\})$, that is, $H(\{a,b,c,d\})$ is a 3-Steiner convex set that is not the convex hull of its extreme points. This is a contradiction. Hence, G is P_4 -free.


Now consider the four graphs G_1, G_2, G_3, G_4 depicted in Figure 5. Note that $ext(G_i) = \emptyset, 1 \le i \le 4$. For instance, it can be checked by inspection that every vertex in G_1 is a central vertex of a claw. A similar behavior occurs for G_2, G_3 , and G_4 . This implies that $Y = H(G_i)$ is a 3-Steiner convex set and $ext(Y) = \emptyset$, a contradiction. Hence, G is (G_1, G_2, G_3, G_4) -free.

On the other hand, in [10] the authors show that if G is $(P_4, G_1, G_2, G_3, G_4)$ -free then the 3-Steiner convexity of G is a convex geometry; this proof is a bit laborious, and so it is omitted here.

Theorem 9. [10] The 3-Steiner convexity of G is a convex geometry if and only if G is $(P_4, G_1, G_2, G_3, G_4)$ -free.

Let $\mathcal{R} = \{k_1, k_2, \dots, k_t\}$ be a set of positive integers with $k_1 \leq \dots \leq k_t$ and $k_1 \in \{2, 3\}$. Say that a set $S \subseteq V(G)$ is \mathcal{R} -Steiner convex if S is k_i -Steiner convex for $1 \leq i \leq t$. The convexity associated with \mathcal{R} -Steiner convex










Figure 5: Graphs G_1, G_2, G_3, G_4 .

sets is the \mathcal{R} -Steiner convexity. Interestingly, the class of graphs for which the \mathcal{R} -Steiner convexity is a convex geometry is exactly the same as the class for which the k_1 -Steiner convexity is a convex geometry:

Theorem 10. [10] The following statements are true:

- 1. If k = 2, then the R-Steiner convexity is a convex geometry if and only if G is Ptolemaic.
- 2. If k = 3, then the \mathcal{R} -Steiner convexity is a convex geometry if and only if G is $(P_4, G_1, G_2, G_3, G_4)$ -free.

To conclude this section, we briefly discuss a further convexity notion, also presented in [10], that combines the 3-Steiner convexity and the so-called g^3 -convexity. A set S of vertices in a graph G is g^3 -convex if, for every pair $u, v \in S$ such that u and v are at distance at least 3 in G, $I[u, v] \subseteq S$. (Here, the interval is taken over shortest paths.) A set S is g_3^3 -convex if it is both g_3 - and g^3 -convex. The convexity associated with g_3^3 -convex sets is the g_3^3 -Steiner convexity.

Theorem 11. [10] The g_3^3 -Steiner convexity of a graph G is a convex geometry if and only if G is (house, hole, domino, A, gem, G_1 , G_2 , G_3 , G_4 , T_1 , T_2)-free. See Figures 2,3,5,6.

3.6. Toll convexity

A walk $u_0u_1 ldots u_{k-1}u_k$ is a tolled walk if $u_0u_i \in E(G)$ implies i = 1 and $u_ju_k \in E(G)$ implies j = k - 1. In [1], tolled walks were used to characterize interval graphs. A graph G is an interval graph if its vertices can be associated with intervals on the real line such that two vertices are adjacent if and only




Figure 6: Graphs T_1 and T_2 .

if the associated intervals intersect. The key references on the toll convexity are [2, 15].

Let \mathscr{P}^t be the path system that maps a graph G to its tolled walks. The convexity defined over such walks is called the *toll convexity* of G, and the associated convex sets are called *toll convex sets*.

If I is an interval on the real line, let R(I) and L(I) be, respectively, the right and left endpoints of I. Given a family of intervals (or interval model) $\mathscr{I} = \{I_v\}_{v \in V(G)}$, associated with an interval graph G, we say that I_a is an end interval if $L(I_a) = Min \bigcup I_v$ or $R(I_a) = Max \bigcup I_v$, i.e., I_a appears as the first or the last interval in \mathscr{I} . A vertex $a \in V(G)$ is an end vertex if there exists some interval model of G where a is associated with an end interval. In addition, a is an end simplicial vertex if a is an end vertex and is simplicial. In [2], two facts on extreme vertices in toll convex sets are given.

Lemma 12. [2] If v is an extreme vertex of a toll convex set S of a graph G, then v is simplicial in G[S].

Lemma 13. [2] A vertex v of an interval graph G is an extreme vertex of a toll convex set $S \subseteq V(G)$ if and only if v is an end simplicial vertex of G[S].

In order to characterize the graphs with toll convexities that are convex geometries, we need to resort to a well-known characterization of interval graphs. Three vertices of a graph form an *asteroidal triple* if between any pair of them there exists a path that avoids the neighborhood of the third vertex.

Theorem 14. [33] A graph G is an interval graph if and only if G is chordal and contains no asteroidal triple.

Suppose that the toll convexity of G is a convex geometry. Let us show first that G is chordal. By the assumption, V(G) (and every of its toll convex

subsets) is the convex hull of its extreme vertices. Let v be an extreme vertex of V(G). By definition of extreme vertex, $V(G) \setminus \{v\}$ is toll convex in G. Thus, the toll convexity of G - v is a convex geometry (note that any subset of $V(G) \setminus \{v\}$ is toll convex in G - v iff it is toll convex in G). Using induction, G - v is chordal. Since by Lemma 12 v is a simplicial vertex of G, it follows that G is chordal as well (recall that a graph is chordal iff there is an ordering v_1, \ldots, v_n of its vertices such that v_i is simplicial in $G[\{v_i, v_{i+1}, \ldots, v_n\}], 1 \le i \le n$ [26]).

Now, let us show that G contains no asteroidal triple. Suppose by contradiction that vertices a, b, c form an asteroidal triple in G. Consider the walk W from a to c formed by the concatenation of induced paths P_{ab} , from a to b, and P_{bc} , from b to c, such that P_{ab} avoids N[c] and P_{bc} avoids N[a]. Note that a is adjacent only to one vertex of P_{ab} and to no vertices of P_{bc} . Likewise, c is adjacent only to one vertex of P_{bc} and to no vertices of P_{ab} . Thus, W is a tolled walk from a to c passing through b. Let $Y = H(\{a, b, c\})$. Since $V(W) \subseteq Y$, b is not an extreme vertex of Y. Analogously, a and c are not extreme vertices of Y. Since no other vertices of Y are extreme vertices, this implies that Y has no extreme vertices, a contradiction.

The precedent paragraphs tell us that if the toll convexity of G is a convex geometry, then, by Theorem 14, G is an interval graph. To prove the converse, we use the following useful lemma:

Lemma 15. [2] Let G be an interval graph. Then, every vertex that is not an end simplicial vertex lies in a tolled walk between two end simplicial vertices of G.

Using Lemma 13, Lemma 15, and the fact that interval graphs are hereditary, we have:

Theorem 16. [2] A graph G is an interval graph if and only if the toll convexity of G is a convex geometry.

3.7. Weakly toll convexity

A walk $u_0u_1 u_{k-1}u_k$ is a weakly toll walk if $u_0u_i \in E(G)$ implies $u_i = u_1$ and $u_ju_k \in E(G)$ implies $u_j = u_{k-1}$. The concept of weakly toll walk is a relaxation of the concept of tolled walk (in the sense that every tolled walk is a weakly toll walk). In the previous subsection, tolled walks have been used to characterize interval graphs via convex geometries. Similarly, weakly toll walks are used as a tool to characterize proper interval graphs. A proper

interval graph is an interval graph that admits an interval model in which no interval properly contains another, or, equivalently, an interval model in which all the intervals have the same length. Roberts [40] proved that proper interval graphs are exactly the interval graphs containing no $K_{1,3}$ as an induced subgraph. The graph $K_{1,3}$ consists of four vertices a, b, c, d and three edges ab, ac, ad.

Let \mathscr{P}^w be the path system that maps a graph G to its weakly toll walks. The convexity defined over such walks is called the *weakly toll convexity* of G, and the associated convex sets are called *weakly toll convex sets*. The basic references on the weakly toll convexity are [28, 29].

Lemmas 12 and 13 have similar versions for weakly toll convex sets:

Lemma 17. [28] If v is an extreme vertex of a weakly toll convex set S of a graph G, then v is simplicial in G[S].

Lemma 18. [28] A vertex v of a proper interval graph G is an extreme vertex of a weakly toll convex set $S \subseteq V(G)$ if and only if v is an end simplicial vertex of G[S].

Using arguments similar to those used in the previous section, one can prove that if the weakly toll convexity of G is a convex geometry, then G is chordal and cannot contain asteroidal triples and induced subgraphs isomorphic to $K_{1,3}$. Hence, using the characterization of proper interval graphs in [40], G is a proper interval graph. Conversely, assume that G is a proper interval graph. Then, it is an interval graph. This means that, by Lemma 15, every vertex of G that is not an end simplicial vertex lies in a tolled walk between two end simplicial vertices. Since every tolled walk is a weakly toll walk and, by Lemma 18, end simplicial vertices of a proper interval graph are extreme vertices, we have:

Lemma 19. [28] Let G be a proper interval graph. Then, every vertex that is not an end simplicial vertex lies in a weakly toll walk between two end simplicial vertices of G.

Theorem 20. [28] A graph G is a proper interval graph if and only if the weakly toll convexity of G is a convex geometry.

3.8. l^k -convexity

Let $\mathscr{P}^{\leq k}$ be the path system that maps a graph G to its induced paths of length at most k. The convexity defined over such paths is called the

 l^k -convexity of G, and the associated convex sets are called l^k -convex sets. The l^2 -convexity is also called P_3^* -convexity [6].

Suppose that the l^2 -convexity of G is a convex geometry. It is easy to see that the extreme vertices of the l^2 -convex set V(G) are the simplicial vertices of G. Thus, if G contains an induced cycle C with at least four vertices, then H(V(C)) does not have simplicial (extreme) vertices, a contradiction. This means that G is chordal. Now, we prove that G is also a cograph (a graph that contains no P_4 as an induced subgraph – see [13] for details). Let $P = x_0, \ldots, x_p$ be a maximum induced path of G, between two simplicial vertices of G. Such a path exists because G is chordal [14]. Assume $p \geq 3$. The set H(V(P)) is an l^2 -convex set of G whose extreme vertices are x_0 and x_p . If there exists $z \in H(V(P)) \setminus V(P)$ that lies in an induced path of length two between x_0 and x_p , z must be adjacent to every x_i ; otherwise, P plus the edges x_0z and zx_p forms an induced cycle of G with at least four vertices. Since $N[x_0]$ and $N[x_p]$ are cliques in H(V(P)),

$$H(V(P)) = (N[x_0] \cap N[x_p]) \cup \{x_0, x_p\}.$$

This implies $H(V(P)) \neq H(\{x_0, x_p\})$, a contradiction. Thus G is a chordal cograph.

Conversely, let G be a chordal cograph, and let $S \subseteq V(G)$ be an l^2 -convex set of G. Since G[S] is chordal, every vertex in S lies in an induced path between two simplicial vertices of G[S]. Such a path must have length two, otherwise G contains an induced path of length at least four as an induced subgraph. Hence,

Theorem 21. [7, 27] A graph G is a chordal cograph if and only if the l^2 -convexity of G is a convex geometry.

We now analyze graphs with l^3 -convexities that are convex geometries. A vertex u is a universal vertex in a graph G if u is adjacent to every other vertex of G. An n-gem is a graph G_n such that: (i) $V(G_n) = \{x_0, \ldots, x_n, u_n\}$ $(n \ge 4)$; (ii) x_0, \ldots, x_n is an induced path; (iii) u_n is a universal vertex. See Figure 7.

Suppose that G contains an n-gem G_n as an induced subgraph. We say that G_n is *solved* if there exists in G a P_4 connecting x_0 and x_n that avoids u_n . In [27], graphs with l^3 -convexities that are convex geometries are characterized as follows:




Figure 7: An n-gem.

Theorem 22. [27] Let G be a graph. The l^3 -convexity of G is a convex geometry if and only if the following conditions hold:

- 1. G is chordal;
- 2. diam(G) < 3;
- 3. every induced n-gem $(n \ge 4)$ contained in G is solved.

The proof of the above theorem contains many details and will be omitted. The "only if" part can be generalized to prove the following result:

Theorem 23. [27] Let G be a graph and let $k \geq 2$. If the l^k -convexity of G is a convex geometry, then G is chordal and $diam(G) \leq k$.

Despite the above result, a complete characterization of graphs with l^k -convexities that are convex geometries, for arbitrary k, remains still as an open question.

A surprising fact is that the class of graphs characterized in Theorem 22 is not hereditary. The following example is elucidative:

Example 24. Let G be the graph depicted in Figure 8. Note that the l^3 -convexity of G is a convex geometry. However, for $x \in \{2, 5\}$, the l^3 -convexity of G - x is not a convex geometry, since:

- V(G-x) is trivially an l^3 -convex set of G-x,
- $ext(V(G-x)) = \{1, 7\}$, and
- $H(\{1,7\}) = \{1,7\} \neq V(G-x)$.

This shows that the class of graphs with l^3 -convexities that are convex geometries is not hereditary. As far as the authors know, this is the only practical example of a nonhereditary class of graphs with convex geometries of a certain type.




Figure 8: The l^3 -convexity of G is a convex geometry.

4. Other convexities

In this section, we present some contributions to the study of convex geometries, some of which not associated with path systems.

4.1. P_3 convexity

Consider the path system that maps a graph G to its paths of length two (not necessarily induced). The convexity defined over such paths is called the P_3 convexity of G [11], and the associated convex sets are called P_3 convex sets. Note that S is a P_3 convex set if and only if every $x \notin S$ has at most one neighbor in S. Thus, the determination of the convex hull of a set in the P_3 convexity may be viewed as an infection process where noninfected vertices having two or more infected neighbors become infected (for more details, see [11] and references therein).

Theorem 25. The P_3 convexity of G is a convex geometry if and only if G is a forest of stars.

Proof. First, suppose that the P_3 convexity of G is a convex geometry. If there are vertices $a, b, c \in V(G)$ inducing a K_3 , then $a \in H(\{b, c\})$ and $b \in H(\{a, c\})$, which contradicts the anti-exchange property, since $\{c\}$ is a convex set. Therefore, G is K_3 -free and the endpoints of any edge of G form a convex set. We also have that G is C_4 -free, because if there is an induced $C_4 = abcda$ in G, then $S = \{c, d\}$ is a convex set, $a \in H(S \cup \{b\})$, and $b \in H(S \cup \{a\})$, which contradicts the anti-exchange property. Now, suppose that G contains an induced $C_5 = abcdea$ in G. Since $\{c, d\}$ is a convex set and G is C_4 -free, we have $H(\{c, e\}) = \{c, d, e\}$, because if there is $v \in H(\{c, e\}) \setminus \{c, d, e\}$, then $\{c, d, e, v\}$ would be an induced C_4 . Since $a \in H(S \cup \{b\})$ and $b \in H(S \cup \{a\})$, the anti-exchange property does not hold. Therefore, G is C_5 -free. Next, suppose that G contains an induced

 $P_4 = abcd$. We have that $S = \{a, d\}$ is a convex set, because if there is a $P_3 = aed$ in G, then we would have that $be, ce \notin E(G)$ (since G is K_3 -free) and that abcde is an induced C_5 , which is not possible. But then the anti-exchange property does not hold because $b \in H(\{a\} \cup S)$ and $c \in H(\{b\} \cup S)$. Thus, G is P_4 -free.

Since G is (K_3, C_4, P_4) -free, we can say that G is a cograph having no triangles or induced cycles with 4 vertices. We know that every nontrivial connected cograph G is the join³ of two graphs G_1 and G_2 . Since G is K_3 -free, we have that the vertices of G_i for every $i \in \{1, 2\}$ induce an independent set; and since G is C_4 -free, we have that $|V(G_i)| = 1$ for at least one $i \in \{1, 2\}$. Therefore, every connected component of G is a star.

Conversely, consider that G is a forest of stars. Since a star has at most one vertex with degree at least 2, G satisfies the anti-exchange property. Thus the P_3 convexity of G is a convex geometry.

4.2. Triangle path convexity

We say that a path P in graph G is a triangle path if every two vertices of P with distance greater than 2 in P are non-adjacent in G. A set $S \subseteq V(G)$ is tp-convex if for any $x, y \in S$, every vertex belonging to some triangle path between x and y belongs to S. The convexity associated with the tp-convex sets of G is the triangle path convexity of G [12, 18].

Theorem 26. The triangle path convexity of G is a convex geometry if and only if G is a forest.

Proof. First, suppose that the triangle path convexity of G is a convex geometry. If there are vertices $a, b, c \in V(G)$ inducing a K_3 , then $a \in H(\{b, c\})$ and $b \in H(\{a, c\})$, which contradicts the anti-exchange property, since $\{c\}$ is a convex set. Therefore, G is K_3 -free and the endpoints of any edge of G form a convex set. We also have that G is C_k -free for $k \geq 4$, because if there is an induced $C_k = v_1 \dots v_k v_1$ in G ($k \geq 4$), then $S = \{v_2, v_3\}$ is a convex set, $v_1 \in H(S \cup \{v_4\})$, and $v_4 \in H(S \cup \{v_1\})$, which contradicts the anti-exchange property. Hence, G is acyclic.

Conversely, suppose that G is a forest. As G is K_3 -free, the triangle path and the monophonic convexities of G are identical. Since G is a forest, G

The *join* of two graphs G_1 and G_2 is the graph G with $V(G) = V(G_1) \cup V(G_2)$ and $E(G) = E(G_1) \cup E(G_2) \cup \{xy \mid x \in V(G_1), y \in V(G_2)\}.$

is a chordal graph, which means, by Theorem 1, that both the monophonic and triangle path convexities of G are convex geometries.

4.3. All-path convexity

A set $S \subseteq V(G)$ is ap-convex if, for any $x, y \in S$, every vertex lying in a path between x and y belongs to S. The convexity associated with the ap-convex sets of G is the all-path convexity of G. A study on the all-path convexity can be found in [39].

It is easy to see that v is an extreme vertex of an ap-convex set S with |S| > 1 if and only if v is a pendant vertex (a vertex of degree one) in G[S].

Theorem 27. The all-path convexity of a graph G is a convex geometry if and only if G is a forest.

Proof. Suppose that the all-path convexity of G is a convex geometry and G contains a cycle C. Then, the vertices in V(C) are non-pendant and H(V(C)) contains no extreme vertices, a contradiction. Thus G is a forest. Conversely, if G is a forest, then every non-pendant vertex of G lies in a path between two pendant vertices, and this property holds for every induced subgraph of G. This means that every ap-convex set of G is the convex hull of its extreme vertices.

4.4. Cycle convexity

We say that a set $S \subseteq V(G)$ is *c-convex* if no vertex $u \notin S$ is adjacent to distinct $x, y \in S$ such that there is a path from x to y in G[S]; in other words, no $u \notin S$ is contained in a cycle C with $V(C) \setminus S = \{u\}$. The convexity associated with the c-convex sets of G is the *cycle convexity* of G. See [4, 5] for important results on the cycle convexity.

Clearly, x is an extreme vertex of a c-convex set S if and only if there is no cycle C in G[S] containing x.

Theorem 28. The cycle convexity of a graph G is a convex geometry if and only if G is a forest.

Proof. If G contains a cycle C, then H(V(C)) contains no extreme vertices. Thus, if the cycle convexity of G is a convex geometry, then G is a forest. Conversely, if G is a forest, then every $S \subseteq V(G)$ is a c-convex set of G and satisfies ext(S) = S, i.e., it is the convex hull of its extreme vertices.

It is interesting to note that Theorems 26, 27, and 28 all apply to the class of forests. However, the triangle path, all-path, and cycle convexities do not coincide in general. If $G = C_n$ $(n \ge 4)$, the families of tp-convex sets, ap-convex sets, and c-convex sets of G are distinct.

4.5. Δ -convexity

A set $S \subseteq V(G)$ is Δ -convex if no vertex $u \notin S$ has two adjacent neighbors in S, that is, forms a triangle with two vertices in S. The Δ -convexity of G is the family formed by the Δ -convex sets of G (see [3] for more details). It is easy to see that x is an extreme vertex of a Δ -convex set S if and only if N(x) is an independent set of G[S].

Theorem 29. The Δ -convexity of a graph G is a convex geometry if and only if G is triangle-free.

Proof. Suppose that the Δ -convexity of G is a convex geometry. If G contains a triangle abc, then $S = H(\{a, b, c\})$ contains no extreme vertices, since N(x) is not an independent set for every vertex x in G[S]. This is a contradiction. Conversely, if G is triangle-free, then every $S \subseteq V(G)$ is a Δ -convex set of G and satisfies ext(S) = S, i.e., it is the convex hull of its extreme vertices. \square

4.6. F-free convexities

We now introduce the concept of \mathcal{F} -free convexity [7]. Let H be a non-trivial graph. Given a graph G, we say that $S \subseteq V(G)$ is H-free convex if for every $S' \subseteq S$ with |S'| = |V(H)| - 1 and $x \in V(G)$, if the subgraph of G induced by $S' \cup \{x\}$ is isomorphic to H, then $x \in S$. Given a family \mathcal{F} of nontrivial graphs, we say that $S \subseteq V(G)$ is \mathcal{F} -free convex if S is H-free convex for every $H \in \mathcal{F}$. The convexity associated with the \mathcal{F} -free convex sets of G is the \mathcal{F} -free convexity of G.

Theorem 30. The \mathcal{F} -free convexity of G is a convex geometry if and only if G is \mathcal{F} -free.

Proof. First, suppose that G is \mathcal{F} -free. By definition, every subset of V(G) is \mathcal{F} -free convex, which means that the anti-exchange property is valid. Therefore, the \mathcal{F} -free convexity of G is a convex geometry.

Conversely, suppose that the \mathcal{F} -free convexity of G is a convex geometry and G contains a graph of \mathcal{F} as an induced subgraph. Let $S \subseteq V(G)$ with minimum size such that G[S] is isomorphic to a graph $H \in \mathcal{F}$. Let x, y

be distinct vertices of S and write $S' = S \setminus \{x, y\}$. By the minimality of S, we have that S' is \mathcal{F} -free convex. Note also that $x \in H(S' \cup \{y\})$ and $y \in H(S \cup \{x\})$, which means that the anti-exchange property does not hold. This is a contradiction. Therefore, G is \mathcal{F} -free. \square

Corollary 31. Let \mathcal{F} be the family of odd cycles. Then, the \mathcal{F} -free convexity of G is a convex geometry if and only if G is bipartite.

Corollary 32. Let \mathcal{F} be the family of graphs that can be obtained from K_5 or $K_{3,3}$ by subdivision of edges. Then, the \mathcal{F} -free convexity of G is a convex geometry if and only if G is planar.

The \mathcal{F} -free convexity defined by $\mathcal{F} = \{P_4\}$ is called P_4^+ -convexity. Recall that a cograph is a P_4 -free graph. Given a graph G, we say that $S \subseteq V(G)$ is P_4^+ -convex if, for every induced $P_4 = abcd$ in G, if $a, b, d \in S$, then $c \in S$. The convexity associated with the P_4^+ -convex sets of G is the P_4^+ -convexity of G.

Corollary 33. The P_4^+ -convexity of G is a convex geometry if and only if G is a cograph.

To conclude this section, we remark that the cycle convexity and the Δ -convexity are \mathcal{F} -free convexities defined by taking $\mathcal{F} = \{C_k \mid k \geq 3\}$ and $\mathcal{F} = \{C_3\}$, respectively. Hence, Theorems 28 and 29 can be viewed as corollaries of Theorem 30.

5. Conclusions

In this survey, we presented characterizations of several classes of graphs via convex geometries. The characterizations follow the following structure: A graph G is in class $\mathscr G$ if and only if the $\mathscr C$ -convexity of G is a convex geometry. The table below summarizes the results.

An interesting question is to increase the table by including other important classes, such as distance-hereditary graphs and split graphs. Another line of research is to investigate convex geometries defined on digraphs. Finally, as far as the authors now, the g^3 -Steiner convexities that are convex geometries have not been characterized yet.

\mathscr{G}	\mathscr{C}
chordal	monophonic
Ptolemaic	geodesic
strongly chordal	strong
weak bipolarizable	m^3
Ptolemaic	Steiner
$(P_4, G_1, G_2, G_3, G_4)$ -free	3-Steiner
(Theorem 10)	$\mathcal{R} ext{-Steiner}$
(Theorem 11)	g_3^3 -Steiner
interval	toll
proper interval	weakly toll
chordal cograph	l^2
(Theorem 18)	l^3
forest of stars	P_3
forest	triangle path, all path, cycle
triangle-free	Δ
${\mathcal F} ext{-free}$	\mathcal{F} -free convexity
bipartite	(Corollary 24)
planar	(Corollary 25)
cograph	P_4^+

References

- [1] L. Alcón, A note on path domination, Discussiones Mathematicae Graph Theory 36 (2016) 1021–1034.
- [2] L. Alcón, B. Bešar, T. Gologranc, M. Gutierrez, T. Kraner Šumenjak, I. Peterin, A. Tepeh, Toll convexity, *European Journal of Combinatorics* 46 (2015) 161–175.
- [3] B. S. Anand, A. Anil, M. Changat, M. C. Dourado, S. S. Ramla, Computing the hull number in Δ -convexity, *Theoretical Computer Science* 844 (2020) 217–226.
- [4] J. Araújo, C. Campos, D. Girão, J. Nogueira, A. Salgueiro, A. Silva, On the hull number on cycle convexity of graphs, *Information Processing Letters* 183 (2024) 106420.

- [5] J. Araújo, G. Ducoffe, N. Nisse, K. Suchan, On interval number in cycle convexity, *Discrete Mathematics and Theoretical Computer Science* 20 (2018), no.1, 1–28.
- [6] R. T. Araújo, R. Sampaio, J. L. Szwarcfiter, The convexity of induced paths of order three, *Discrete Applied Mathematics* 237 (2018) 33–42.
- [7] R. T. Araújo, *Path Convexities and Convex Geometries*, M.Sc. Thesis, Universidade Federal do Ceará, 2013. In Portuguese.
- [8] L. M. Batten, Geodesic subgraphs, J. Graph Theory 7 (1983) 159–163.
- [9] J. Cáceres, A. Marquez, M. L. Puertas, Steiner distance and convexity in graphs. *European J. Combin.* 29 (2008) 726–736.
- [10] J. Cáceres, O. R. Oellermann. On 3-Steiner simplicial orderings. *Discrete Mathematics* 309 (2009) 5828–5833.
- [11] V. Campos, R. M. Sampaio, A. Silva, J. L. Szwarcfiter, Graphs with few P₄'s under the convexity of paths of order three, *Discrete Applied Mathematics* 192 (2015) 28–39.
- [12] M. Changat, J. Mathew, On triangle path convexity in graphs, *Discrete Mathematics* 206 (1999) 91–95.
- [13] D. G. Corneil, Y. Perl, L. K. Stewart, A linear recognition algorithm for cographs, SIAM Journal on Computing 14 (1985) 926–934.
- [14] G. A. Dirac, On rigid circuit graphs, Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg, 25 (1–2) 7–76, 1961.
- [15] M. C. Dourado, Computing the hull number in toll convexity, *Ann Oper Res* 315 (2022) 121–140.
- [16] M. C. Dourado, R. A. Oliveira, F. Protti, Algorithmic aspects of Steiner convexity and enumeration of Steiner trees. Annals of Operations Research 223 (2014) 155–171.
- [17] M. C. Dourado, F. Protti, J. L. Szwarcfiter, Complexity results related to monophonic convexity, *Discrete Applied Mathematics* 158 (2010) 1268-1274.

- [18] M. C. Dourado, R. M. Sampaio, Complexity aspects of the triangle path convexity, *Discrete Applied Mathematics* 206 (2016) 39–47.
- [19] F. Dragan, F. Nicolai, A. Brandstädt, Convexity and HHD-free graphs, SIAM J. Discrete Math 12 (1999) 119–135.
- [20] P. Duchet, Convex sets in graphs II: Minimal path convexity, *J. Combin. Theory Ser. B* 44 (1988), 307–316.
- [21] P. H. Edelman and R. E. Jamison, The theory of convex geometries, *Geometriae Dedicata* 19 (1985) 247–270.
- [22] M. Farber, Characterizations of strongly chordal graphs, *Disc. Math.* 43 (1983) 173–189.
- [23] M. Farber, R. E. Jamison, Convexity in graphs and hypergraphs, SIAM J. Algebraic Discrete Mathematics 7 (1986) 433–444.
- [24] M. R. Garey, D. S. Johnson, Computers and Intractability: A Guide to the Theory of NP-Completeness, W. H. Freeman, New York, 1979.
- [25] M. C. Golumbic, Trivially perfect graphs, *Discrete Mathematics* 24 (1978) 105–107.
- [26] M. C. Golumbic, Algorithmic Graph Theory and Perfect Graphs, Academic Press, New York (1980).
- [27] M. Gutierrez, F. Protti, S. Tondato, Convex geometries over induced paths with bounded length, *Discrete Mathematics* 346 (2023) 113133.
- [28] M. C. Dourado, M. Gutierrez, F. Protti, S. Tondato, Weakly toll convexity and proper interval graphs, *Discrete Mathematics and Theoretical Computer Science* 26:2 (2024) no.4, pp. 1–16.
- [29] M. C. Dourado, M. Gutierrez, F. Protti, S. Tondato, Computing the hull and interval numbers in the weakly toll convexity, *Theoretical Computer Science* 996 (2024) 114501.
- [30] E. Howorka, A characterization of Ptolemaic graphs, *J. Graph Theory* 5 (1981) 323–331.

- [31] B. Jamison, S. Olariu, On the semi-perfect elimination, Adv. Appl. Math. 9 (1988) 364–376.
- [32] B. Korte, L. Lovász, R. Schrader, Greedoids, Springer-Verlag, pp. 19–43, 1991.
- [33] C. G. Lekkerkerker, J. C. Boland, Representation of finite graphs by a set of intervals on the real line, *Fund. Math.* 51 (1962) 45–64.
- [34] M. Krein, D. Milman, On extreme points of regular convex sets, *Studia Mathematica* 9 (1940) 133–138.
- [35] M. H. Nielsen, O. R. Oellermann, Steiner trees and convex geometries, SIAM J. Discrete Math. 23(2) (2009) 680–693.
- [36] S. Olariu, Weak bipolarizable graphs, Discrete Mathematics 74 (1989), 159–171.
- [37] I. M. Pelayo, Generalizing the Krein-Milman property in graph convexity spaces: a short survey, *Ramanujan Math. Soc. Lect. Notes Ser.* 5 (2008) 131–142.
- [38] I. M. Pelayo, Geodesic Convexity in Graphs, Springer, New York (2013).
- [39] F. Protti and J. V. C. Thompson, All-path convexity: combinatorial and complexity aspects, *Ars Combinatoria* 148 (2020) 77–87. Also available at https://arxiv.org/abs/2303.18029.
- [40] F. S. Roberts, "Indifference graphs", in F. Harary (ed.), Proof Techniques in Graph Theory, New York, NY: Academic Press, pp. 139–146 (1969).
- [41] D. Rose, G. Lueker, R. E. Tarjan, Algorithmic aspects of vertex elimination on graphs, SIAM Journal on Computing 5 (1976) 266–283.
- [42] M. L. J. van de Vel, *Theory of Convex Structures*, North-Holland, Amsterdam (1993).