## The Alon-Tarsi number of planar graphs - a simple proof

Yangyan Gu<sup>\*</sup> Xuding Zhu<sup>†</sup>

March 31, 2022

## Abstract

This paper gives a simple proof of the result that every planar graph G has Alon-Tarsi number at most 5, and has a matching M such that G - M has Alon-Tarsi number at most 4.

Keywords: planar graph; list colouring; Alon-Tarsi number.

Assume D is an orientation of a graph G. For a subset H of arcs of D, D[H] denotes the sub-digraph induced by H. We say D[H] is an Eulerian sub-digraph if  $d^+_{D[H]}(v) = d^-_{D[H]}(v)$  for each vertex v. Let

 $\mathcal{E}(D) = \{H : D[H] \text{ is an Eulerian sub-digraph}\},\$ 

 $\mathcal{E}_e(D) = \{H \in \mathcal{E}(D) : |H| \text{ is even}\}, \ \mathcal{E}_o(D) = \{H \in \mathcal{E}(D) : |H| \text{ is odd}\}, \text{diff}(D) = |\mathcal{E}_e(D)| - |\mathcal{E}_o(D)|.$ 

We say D is an Alon-Tarsi orientation if  $diff(D) \neq 0$ .

Let  $\mathbb{N} = \{0, 1, 2, ...\}$  and  $\mathbb{N}^G = \{f : V \to \mathbb{N}\}$ . For  $f, g \in \mathbb{N}^G$ , we write  $g \leq f$  if  $g(v) \leq f(v)$  for all  $v \in V(G)$ . For  $f \in \mathbb{N}^G$ , an *f*-Alon-Tarsi orientation (an *f*-AT orientation, for short) of *G* is an Alon-Tarsi orientation *D* of *G* with  $d_D^+(v) \leq f(v) - 1$  for each vertex *v*. We say *G* is *f*-AT if *G* has an *f*-AT orientation. The Alon-Tarsi number AT(G) of *G* is the minimum *k* such that *G* is *k*-AT (i.e., *f*-AT for the constant function f(v) = k for all *v*). Let ch(G) be the choice number of *G*, and  $\chi_P(G)$  be the paint number of *G*. It is well-known [7] that for any graph *G*,

$$ch(G) \le \chi_P(G) \le AT(G).$$

It is known that for any planar graph G,  $ch(G) \leq 5$  [5],  $\chi_P(G) \leq 5$  [4],  $AT(G) \leq 5$  [6], and G has a matching M such that  $AT(G-M) \leq 4$  [2]. This note gives a simple proof of the last two results. Note that  $AT(G) \leq 5$  implies that  $ch(G) \leq \chi_P(G) \leq 5$ .

For  $f \in \mathbb{N}^G$  and  $X \subseteq V(G)$ , let  $f_{[X,-1]} \in \mathbb{N}^G$  be defined as

$$f_{[X,-1]}(v) = \begin{cases} f(v) - 1, & \text{if } v \in X, \\ f(v), & \text{otherwise.} \end{cases}$$

We write  $f_{[v,-1]}$  for  $f_{[\{v\},-1]}$ .

The following lemma is folklore (cf. Proof of Lemma 3.1.7 in [7]).

<sup>\*</sup>Department of Mathematics, Zhejiang Normal University, China. E-mail: yangyan@zjnu.edu.cn.

<sup>&</sup>lt;sup>†</sup>Department of Mathematics, Zhejiang Normal University, China. E-mail: xdzhu@zjnu.edu.cn. NSFC 11971438,U20A2068, ZJNSFC LD19A010001.

**Lemma 1** If G is f-AT and  $e = uv \in E(G)$ , then G - e is  $f_{[u,-1]}$ -AT or  $f_{[v,-1]}$ -AT.

**Observation 2** If G is f-AT, and D is an f-AT orientation of G, then  $f(x) \ge d_D^+(x) + 1 \ge 1$ for each vertex x. If f(x) = 1, then  $d_D^+(x) = 0$ , and hence any neighbour y of x has  $d_D^+(y) \ge 1$ and  $f(y) \ge 2$ . Therefore the following hold:

- (A) If G is f-AT,  $e = uv \in E(G)$ , and f(v) = 1, then G e is  $f_{[u,-1]}$ -AT.
- (B) If G is f-AT,  $e = uv, e' = vw \in E(G)$ , and f(v) = 2, f(w) = 1, then G e is  $f_{[u,-1]}$ -AT.

**Corollary 3** Assume  $d_G(w) = 2$ ,  $N_G(w) = \{u, v\}$ ,  $f: V(G) \to \mathbb{N}$  and f(w) = 2. If G is f-AT, then G - w is  $f_{[u,-1]}$ -AT or  $f_{[v,-1]}$ -AT.

**Proof.** By Lemma 1, G - wu is  $f_{[u,-1]}$ -AT or  $f_{[w,-1]}$ -AT. If G - wu is  $f_{[u,-1]}$ -AT, then G - w is  $f_{[u,-1]}$ -AT. If G - wu is  $f_{[w,-1]}$ -AT, then since  $f_{[w,-1]}(w) = 1$ ,  $G - w = G - \{wu, wv\}$  is  $f_{[v,-1]}$ -AT.

The following lemma is also easy and well-known (cf. [7]).

**Lemma 4** Assume D is an orientation of G = (V, E),  $X \subseteq V$  and all arcs between X and V - X are oriented from X to V - X. Then diff $(D) = diff(D[X]) \times diff(D[V - X])$ , where D[X], D[V - X] are sub-digraphs induced by X and V - X.

**Definition 5** Assume G is a 2-connected plane graph and  $e_1 = v_1v_2$  is a boundary edge. Assume M is a matching of G containing  $e_1$ . Let  $f_{G,v_1v_2}, f_{G,v_1v_2,M} \in \mathbb{N}^G$  be defined as

$$f_{G,v_1v_2}(x) = \begin{cases} 1, & \text{if } x \in \{v_1, v_2\}, \\ 3, & \text{if } x \in B(G) - \{v_1, v_2\}, \\ 5, & \text{if } x \in V(G) - B(G). \end{cases}$$
$$f_{G,v_1v_2,M}(x) = \begin{cases} 1, & \text{if } x \in \{v_1, v_2\}, \\ 3 - d_M(x), & \text{if } x \in B(G) - \{v_1, v_2\}, \\ 4, & \text{if } x \in V(G) - B(G). \end{cases}$$

The result that stated in the abstract follows from the following more technical theorem.

**Theorem 6** Assume G is a 2-connected plane graph and  $e_1 = v_1v_2$  is a boundary edge. Then  $G-e_1$  is  $f_{G,v_1v_2}$ -AT, and G has a matching M containing  $e_1$  such that G-M is  $f_{G,v_1v_2,M}$ -AT.

**Proof.** The proof is by induction on |V(G)|. If G is a triangle  $(v_1, v_2, v_3)$ , then let D be the orientation of  $G - e_1$  with  $v_3$  be a source. Then D is an  $f_{G,v_1v_2}$ -AT orientation of  $G - e_1$ , and also an  $f_{G,v_1v_2,M}$ -AT orientation G - M with  $M = \{e_1\}$ .

Assume  $|V(G)| \ge 4$  and Theorem 6 is true for smaller plane graphs.

4,

**Case 1** B(G) has a chord e = xy.

Let  $G_1, G_2$  be the subgraphs of G separated by e (i.e.,  $V(G_1) \cap V(G_2) = \{x, y\}$  and  $V(G_1) \cup$  $V(G_2) = V(G)$ , and assume  $v_1v_2 \in E(G_1)$ . By induction hypothesis,  $G_1 - e_1$  has an  $f_{G_1,v_1v_2}$ -AT orientation  $D_1$  and  $G_2 - xy$  has  $f_{G_2,xy}$ -AT orientation  $D_2$ . Let  $D = D_1 \cup D_2$ . It follows from Lemma 4 that  $\operatorname{diff}(D) = \operatorname{diff}(D_1) \times \operatorname{diff}(D_2) \neq 0$ . Hence D is an  $f_{G,v_1v_2}$ -AT orientation of  $G - e_1$ .

Also by induction hypothesis,  $G_1$  has a matching  $M_1$  containing  $e_1$  so that  $G_1 - M_1$  has an  $f_{G_1,v_1v_2,M_1}$ -AT orientation  $D'_1$ , and  $G_2$  has a matching  $M_2$  containing e = xy so that  $G_2 - M_2$  has an  $f_{G_2,xy,M_2}$ -AT orientation  $D'_2$ . Let  $M = M_1 \cup (M_2 - \{e\})$  and let  $D' = D'_1 \cup D'_2$ . Then M is a matching of G containing  $e_1$ . By Lemma 4, diff $(D') = diff(D'_1) \times diff(D'_2) \neq 0$ , and hence D' is an  $f_{G,v_1v_2,M}$ -AT orientation of G - M.

**Case 2** B(G) has no chord.

Assume  $B(G) = (v_1, v_2, \ldots, v_n)$  and let  $G' = G - v_n$ . Assume  $N_G(v_n) = \{v_1, u_1, u_2, \ldots, u_k, v_{n-1}\}$ . Let  $U = \{u_1, u_2, \ldots, u_k\}$ . By induction hypothesis,  $G' - e_1$  has an  $f_{G', v_1 v_2}$ -AT orientation  $D_1$ . Let G'' be obtained from G by adding paths  $P_1, P_2, \ldots, P_k$ , where  $P_i = (u_i, w_i, v_n)$ , and  $w_i$  are new vertices. Let  $b_1 = (v_n, v_{n-1}), b_2 = (v_n, v_1)$  and for  $i = 1, 2, \ldots, k, a_i = (u_i, v_n), a'_i = (u_i, w_i), a''_i = (w_i, v_n)$ . Let

$$Z = \{a_i, a'_i, a''_i : i = 1, 2, \dots, k\} \cup \{b_1, b_2\}.$$

Let  $D_2$  be the orientation of  $G'' - e_1$  obtained from  $D_1$  by adding arcs in Z. For an arc a of  $D_2$ , let

$$\mathcal{E}(D_2, a) = \{ H \in \mathcal{E}(D_2) : a \in H \}.$$

It is easy to see that

$$\mathcal{E}(D_2) = \mathcal{E}(D_1) \cup (\cup_{i=1}^k (\mathcal{E}(D_2, a_i) \cup \mathcal{E}(D_2, a'_i))),$$

where the unions are disjoint unions. The map  $\phi : \mathcal{E}(D_2, a_i) \to \mathcal{E}(D_2, a'_i)$  defined as  $\phi(H) = (H - a_i) \cup \{a'_i, a''_i\}$  is a one-to-one correspondence, and H and  $\phi(H)$  have opposite parities. So

$$|\mathcal{E}_{e}(D_{2}) \cap (\cup_{i=1}^{k} (\mathcal{E}(D_{2}, a_{i}) \cup \mathcal{E}(D_{2}, a_{i}')))| = |\mathcal{E}_{o}(D_{2}) \cap (\cup_{i=1}^{k} (\mathcal{E}(D_{2}, a_{i}) \cup \mathcal{E}(D_{2}, a_{i}')))|$$

and hence diff $(D_2) = \text{diff}(D_1) \neq 0$ . Let  $g(v) = d_{D_2}^+(v) + 1$  for  $v \in V(G'')$ . Then  $G'' - e_1$  is g-AT, and by Lemma 1,  $G - e_1$  is g-AT. It is easy to see that  $g \leq f_{G,v_1v_2}$ . Hence  $G - e_1$  is  $f_{G,v_1v_2}$ -AT.

Similarly, by induction hypothesis, G' has a matching  $M_1$  containing  $e_1$  so that  $G' - M_1$  has an  $f_{G',v_1v_2,M_1}$ -AT orientation  $D'_1$ . Let  $D'_2$  be the orientation of  $G'' - M_1$  obtained from  $D'_1$  by adding arcs in Z. For the same reason as above, we have diff $(D'_2) = \text{diff}(D'_1) \neq 0$ . Thus  $D'_2$  is an Alon-Tarsi orientation of  $G'' - M_1$ .

Let  $g(v) = d_{D'_2}^+(v) + 1$ . Then  $G'' - M_1$  is g-AT. It follows from the construction of  $D'_2$  that  $g(v) \leq f_{G,v_1v_2,M_1}(v)$  for  $v \in V(G) - U$ . For i = 1, 2, ..., k, if  $d_{D'_1}^+(u_i) \leq 1$ , then we also have  $g(u_i) = d_{D'_1}^+(u_i) + 2 + 1 \leq 4 = f_{G,v_1v_2,M_1}(u_i)$ . In particular, if  $u_i$  is covered by the matching  $M_1$ , then  $g(u_i) \leq f_{G,v_1v_2,M_1}(u_i)$ . However, if  $d_{D'_1}^+(u_i) = 2$ , then  $g(u_i) = 5 = f_{G,v_1v_2,M_1}(u_i) + 1$ .

Let  $G_k = G''$  and for i = 0, 1, ..., k - 1, let  $G_i = G_{i+1} - w_{i+1}$ . Apply Corollary 3 to  $G_{k-1}, G_{k-2}, ..., G_0$  one by one, we conclude that for  $i = k - 1, ..., 0, G_i - M_1$  is  $g_i$ -AT, where  $g_k = g$  and for i = k - 1, ..., 0, either (i)  $g_i = g_{i+1, [u_{i+1}, -1]}$ , or (ii)  $g_i = g_{i+1, [v_n, -1]}$ .

Note that  $g(v_1) = 1$  and  $g(v_n) = 3$ . By (B) of Observation 2,  $g_i(v_n) \ge 2$  for all *i*. Hence (ii) occurs at most once. If (ii) never occurs, then  $G_0 - M_1 = G - M_1$  is  $g_{[U,-1]}$ -AT. As  $g_{[U,-1]} \le f_{G,v_1v_2,M_1}$ , we are done.

Assume (ii) occurs exactly once at some *i*. Then  $G_0 - M_1 = G - M_1$  is  $g_{[(U-\{u_i\})\cup\{v_n\},-1]}$ -AT. If  $u_i$  is covered by  $M_1$ , then we have  $g_{[(U-\{u_i\})\cup\{v_n\},-1]} \leq f_{G,v_1v_2,M_1}$ , we are done. Otherwise, let  $M = M_1 \cup \{u_iv_n\}$ . Note that if D is a  $g_{[(U-\{u_i\})\cup\{v_n\},-1]}$ -orientation of  $G - M_1$ , then  $v_n$  has only one out-neighbour  $v_1$ , which is a sink. So any arc in D incident to  $v_n$  is not contained in any Eulerian sub-digraph of D. Hence  $D - (u_i, v_n)$  is an AT-orientation of G - M. So G - M is  $g_{[U\cup\{v_n\},-1]}$ -AT. As  $g_{[U\cup\{v_n\},-1]} \leq f_{G,v_1v_2,M}$ , we have G - M is  $f_{G,v_1v_2,M}$ -AT.

## References

- N. Alon, M. Tarsi, Colorings and orientations of graphs, Combinatorica 12 (2) (1992) 125–134.
- [2] J. Grytczuk and X. Zhu, The Alon-Tarsi number of a planar graph minus a matching, J. Combin. Theory Ser. B 145 (2020), 511-520.
- [3] D. Hefetz, On two generalizations of the Alon-Tarsi polynomial method, Journal of Combinaotrial Theory Ser. B, 101 (2011) 403 –414.
- [4] U. Schauz, Mr. Paint and Mrs. Correct, Electron. J. Combin. 16 (1) (2009) R77.
- [5] C. Thomassen, Every planar graph is 5-choosable, Journal of Combinatorial Theory Ser. B, 62(1) (1994):180–181.
- [6] X. Zhu, The Alon-Tarsi number of planar graphs, J. Combin. Theory Ser. B 134 (2019), 354-358.
- [7] X. Zhu and R. Balakrishnan, Combinatorial Nullstellensatz: with Applications to Graph Colouring. Chapman and Hall/CRC, 2021. isbn: 9780367686949.