INDEPENDENCE COMPLEXES OF $(n \times 4)$ AND $(n \times 5)$ -GRID GRAPHS

TAKAHIRO MATSUSHITA AND SHUN WAKATSUKI

ABSTRACT. We determine the homotopy types of the independence complexes of $(n \times 4)$ and $(n \times 5)$ -square grid graphs. In fact, they are homotopy equivalent to wedges of spheres.

1. INTRODUCTION

Let G = (V, E) be a simple graph. A subset σ of V is *independent* if no two vertices in σ are not adjacent in G. The family of independent sets in G forms a simplicial complex, and we call it the *independence complex* I(G) of G.

The independence complex is one of the most principal constructions of simplicial complexes. In fact, a simplicial complex K is isomorphic to some independence complex if and only if K is a clique complex. A *clique complex* is a simplicial complex such that the dimension of every minimal non-face is at most one, and has appeared in several branches of mathematics. The order complexes of posets, appearing in combinatorics [K2], and the Vietoris–Rips complexes, appearing in topological data analysis and geometric group theory [PRSZ], are typical examples of clique complexes. The matching complex of a graph G is the independence complex of the line graph of G, and its precise homotopy type has been recently studied by several authors (see [AGS], [BH], [BGJM], [M], and [W]). Since the barycentric subdivision of every simplicial complex is a clique complex, and hence for every simplicial complex K, there exists an independence complex homeomorphic to K.

In general, it is quite difficult to determine the homotopy type of the independence complex I(G) even though the graph G is simply described. In this paper, we study the homotopy types of the independence complexes of grid graphs $\Gamma_{n,k}$. Here, for a pair n and k of positive integers, define the $(n \times k)$ -grid graph $\Gamma_{n,k}$ by

$$V(\Gamma_{n,k}) = \{(x,y) \in \mathbb{Z}^2 \mid 1 \le x \le n, \ 1 \le y \le k\},\$$
$$E(\Gamma_{n,k}) = \{\{(x,y), (z,w)\} \mid (x,y), (z,w) \in V(\Gamma_{n,k}), \ |x-z| + |y-w| = 1\}.$$

The goal of this paper is to determine the precise homotopy types of $I(\Gamma_{n,4})$ and $I(\Gamma_{n,5})$.

We now review known results related to $I(\Gamma_{n,k})$. When k = 1, the graph $\Gamma_{n,1}$ is the path graph P_n consisting of n vertices, and the homotopy types of $I(\Gamma_{n,1})$ is determined by Kozlov [K1]. The homotopy types of $I(\Gamma_{n,2})$ and $I(\Gamma_{n,3})$ are determined by Adamaszek [A2].

²⁰²⁰ Mathematics Subject Classification. 55P10; 05C69.

Key words and phrases. independence complexes; square grid graphs.

In these cases, $I(\Gamma_{n,k})$ is contractible or homotopy equivalent to a sphere. For $k \geq 4$, the homotopy types of $I(\Gamma_{n,k})$ are not completely determined. However, Okura [O] determined the Euler characteristic of $I(\Gamma_{n,4})$, and it is an unbounded function with respect to n. His computation implies that for large n, $I(\Gamma_{n,4})$ is neither contractible nor homotopy equivalent to a sphere. Bousquet-Mélou, Linusson, and Nevo [BMLN] studied the independence complexes of several types of square grid graphs.

Our grid graph $\Gamma_{n,k}$ is the cartesian product $P_n \times P_k$ of the path graphs P_n and P_k . The homotopy types of independence complexes of $C_n \times P_k$ and $C_n \times C_k$ have also been studied by several authors, motivated by statistical physics. A celebrated result by Jonsson [J1] asserts that if n and k are coprime, then the reduced Euler characteristic of $I(C_n \times C_k)$ is -1. His computation showed that the homotopy types of $I(P_n \times P_k)$ and $I(C_n \times C_k)$ seems to be very different since the absolute value of the Euler characteristic of $I(P_n \times P_4) = I(\Gamma_{n,4})$ tends to infinity (see [O] or Theorem 1.1). See [A2], [I], [J2], [J3], and [T] for more information.

Now we state the main results in this paper.

Theorem 1.1. For $n \ge 1$, there is a following homotopy equivalence:

$$I(\Gamma_{n,4}) \simeq \begin{cases} \bigvee_{2k} S^{n-1} & (n = 6k + 1) \\ \bigvee_{2k+2} S^{n-1} & (n = 6k + 4) \\ \bigvee_{2k+1} S^{n-1} & (n = 6k + l \text{ with } l = 0, 2, 3, 5). \end{cases}$$

Theorem 1.2. There are following homotopy equivalences

$$I(\Gamma_{1,5}) \simeq S^1, \ I(\Gamma_{4,5}) \simeq S^4, \ I(\Gamma_{5,5}) \simeq S^5, \ I(\Gamma_{9,5}) \simeq S^{10}.$$

Assume that n is a positive integer which is not equal to 1,4,5,9. Let t, l, and ε be nonnegative integers satisfying $n = 20t+4l+\varepsilon$, $l \in \{0,1,2,3,4\}$, and $\varepsilon \in \{0,1,2,3\}$. Define $\nu = \lfloor \epsilon/2 \rfloor \in \{0,1\}$ (i.e., the greatest integer less than or equal to $\varepsilon/2$) and $n' = 25t+5l+\varepsilon-1+\nu$. Then the following hold:

(1) Assume that $n = 0, 4, 5, 9, 10, 14, 15, 19 \mod 20$. Then

$$I(\Gamma_{n,5}) \simeq \bigvee_{3-2\nu} S^{n'} \lor \left(\bigvee_{n'-t-\nu < i < n'} \left(\bigvee_{4} S^{i}\right)\right) \lor \bigvee_{2} S^{n'-t-\nu}$$

(2) Assume that n = 2, 3, 6, 7, 8, 11, 12, 13, 16, 17 modulo 20. Then

$$I(\Gamma_{n,5}) \simeq \bigvee_{3-2\nu} S^{n'} \lor \Big(\bigvee_{n'-t \le i < n'} \Big(\bigvee_{4} S^{i}\Big)\Big).$$

(3) Assume that n = 1, 18 modulo 20. Then

$$I(\Gamma_{n,5}) \simeq \bigvee_{3-2\nu} S^{n'} \lor \left(\bigvee_{n'-t-2\nu < i < n'} \left(\bigvee_{4} S^{i}\right)\right).$$

Corollary 1.3. The Euler characteristic of $I(\Gamma_{n,5})$ is a bounded function whose Euler characteristic is even and whose absolute value is at most 4.

Discrete Morse theory is a commonly used method to determine the homotopy types of independence complexes (see [BMLN] and [T] for example). However, it is very difficult to use discrete Morse theory to determine the homotopy type of a simplicial complex which is not a wedge of spheres of the same dimension. For other methods to determine the homotopy types of independence complexes, see [A1], [B], and [EH]. Our method is an elementary method using links of simplicial complexes, reviewed in Section 2.

Finally, we pose a problem related to the Euler characteristic of $I(\Gamma_{n,k})$, inspired by Corollary 1.3.

Problem 1.4. Given a positive integer k, determine whether the function assigning $\chi(I(\Gamma_{n,k}))$ to n is a bounded function.

The rest of this paper is organized as follows. In Section 2, we review definitions and facts related to independence complexes. Section 3 and Section 4 are devoted to the proofs of Theorem 1.1 and Theorem 1.2, respectively.

2. Preliminaries

In this section, we review several definitions and facts related to independence complexes which we need in the subsequent sections. For a comprehensive introduction to simplicial complexes, we refer to [K2]. In the latter of this section, we provide an alternative proof of determining the homotopy type of $I(\Gamma_{n,3})$.

A graph is a pair (V, E) such that V is a finite set and E is a subset of the family $\binom{V}{2}$ of 2-element subsets of V. A subset S of V is *independent* or *stable* if there are no $v, w \in S$ such that $\{v, w\} \in E$. The independence complex of a graph G is the simplicial complex whose vertex set is V and whose simplices are independent sets in G, and is denoted by I(G).

A subgraph of G = (V, E) is a graph G' = (V', E') such that $V' \subset V$ and $E' \subset E$. The subgraph G' is said to be *induced* if $E' = V' \cap E$. For a subset S of V, we write G - S to mean the induced subgraph of G whose vertex set is V - S. Note that if G' is an induced subgraph then there is a natural inclusion from I(G') to I(G).

If G is a disjoint union of two subgraphs G_1 and G_2 , then I(G) coincides with the join $I(G_1) * I(G_2)$. In particular, if G has an isolated vertex, then I(G) is contractible. Since $I(K_2) = S^0$, we have $I(K_2 \sqcup G) \simeq \Sigma I(G)$. Here, Σ denotes the suspension.

Let K be an (abstract) simplicial complex and S a subset of V(K). Let K - S denote the simplicial complex consisting of the simplices of K disjoint from S. For a face σ of K, define the star star(σ) and the link link(σ) by

$$\operatorname{star}(\sigma) = \{\tau \in K \mid \sigma \cup \tau \in K\}, \ \operatorname{link}(\sigma) = \operatorname{star}(\sigma) - \sigma.$$

For $v \in V$, we write star(v) and link(v) instead of star($\{v\}$) and link($\{v\}$), respectively.

Lemma 2.1. Let K be a simplicial complex and v a vertex of K. Then K is homeomorphic to the mapping cone of the inclusion $link(v) \hookrightarrow K - v$, and hence there is a cofiber sequence

$$link(v) \to K - v \to K.$$

In particular, if the inclusion $link(v) \hookrightarrow K - v$ is null-homotopic, then $K \simeq (K - v) \lor \Sigma link(v)$.

Let G be a graph and let $v \in V(G)$. Let N(v) be the set of vertices of G adjacent to v, and set $N[v] = N(v) \cup \{v\}$. Then it is straightforward to see I(G) - v = I(G - v) and link(v) = I(G - N[v]). Applying the above lemma to K = I(G), we have the following:

Theorem 2.2 (See [A1]). Let G be a graph and v be a vertex of G. Then there is a cofiber sequence

$$I(G - N[v]) \to I(G - v) \to I(G).$$

In particular, if the inclusion $I(G - N[v]) \rightarrow I(G - v)$ is null-homotopic, then $I(G) \simeq I(G - v) \vee \Sigma I(G - N[v])$.

This theorem implies the following simple argument, which will be frequently used.

Corollary 2.3 (Fold lemma [E]). Let G be a graph, and v, w vertices of G. Assume that $v \neq w$ and $N(v) \subset N(w)$. Then the inclusion $I(G-w) \hookrightarrow I(G)$ is a homotopy equivalence.

Proof. We write the proof for the reader's convenience. By Theorem 2.2, it suffices to see that v is an isolated vertex of G - N[w]. If $v \in N[w]$, then $v \neq w$ means that $v \in N(w)$. This means $w \in N(v) \subset N(w)$. This is a contradiction. Hence $v \in V(G - N[w])$. Since $N(v) \subset N(w)$, we have that v is an isolated vertex in G - N[w]. This completes the proof.

The fold lemma immediately determines the homotopy types of $I(\Gamma_{n,1})$ and $I(\Gamma_{n,2})$: **Theorem 2.4** (Kozlov [K1]).

$$I(\Gamma_{n,1}) = I(P_n) \simeq \begin{cases} S^{k-1} & (n = 3k) \\ \text{pt} & (n = 3k+1) \\ S^k & (n = 3k+2) \end{cases}$$

Theorem 2.5 (Adamaszek [A2]).

$$I(\Gamma_{n,2}) \simeq \begin{cases} S^{k-1} & (n=2k) \\ S^k & (n=2k+1) \end{cases}$$

We need the homotopy type of $I(C_n)$, where C_n denotes the *n*-cycle graph. **Theorem 2.6** (Kozlov [K1]). For $n \ge 3$,

$$I(C_n) \simeq \begin{cases} S^{k-1} \lor S^{k-1} & (n = 3k) \\ S^{k-1} & (n = 3k+1) \\ S^k & (n = 3k+2). \end{cases}$$



FIGURE 1.

2.1. $I(\Gamma_{n,3})$. In this subsection, we provide an alternative proof of the following theorem because our proofs in the subsequent sections are further developments of this proof:

Theorem 2.7 (Adamaszek [A2]). For l = 0, 1, 2, 3, we have

 $I(\Gamma_{4k+l,3}) \simeq S^{3k+l-1}$

Throughout this subsection, we write Γ_n to mean $\Gamma_{n,3}$, and set $X_n = \Gamma_n - (n,2)$ and $Y_n = \Gamma_n - \{(n,1), (n,3)\}$. We first determine the homotopy type of $I(X_n)$. This notation is only used in this subsection, and in the subsequent sections we use the symbols X_n and Y_n to indicate other graphs.

By the fold lemma (Corollary 2.3), we have the following homotopy equivalences:

(2.1) $I(X_n) \simeq I(K_2 \sqcup K_2 \sqcup Y_{n-2}) = \Sigma^2 I(Y_{n-2}), I(Y_n) \simeq I(K_2 \sqcup X_{n-2}) = \Sigma I(X_n).$

Hence $I(X_n)$ is recursively determined by $I(X_1)$, $I(X_2)$, $I(Y_1)$, and $I(Y_2)$. Using the fold lemma, we have

(2.2)
$$I(X_1) \simeq \text{pt}, I(X_2) \simeq S^1, I(Y_1) = \text{pt}, I(Y_2) \simeq S^0.$$

Lemma 2.8.

$$I(X_n) \simeq \begin{cases} \text{pt} & (n \text{ is odd}) \\ S^{3k-1} & (n = 4k) \\ S^{3k+1} & (n = 4k+2) \end{cases}$$

Proof. Using equation (2.1), we have that $I(X_n) \simeq \Sigma^3 I(X_{n-4})$ for n > 4, $I(X_3) \simeq \Sigma^2 I(Y_1) = \text{pt}$, and $I(X_4) \simeq \Sigma^2 I(Y_2) \simeq S^1$ by equations (2.1) and (2.2). This completes the proof.

Proof of Theorem 2.7. Applying Theorem 2.2 to the vertex v = (n, 2) in $\Gamma_n = \Gamma_{n,3}$, we have a cofiber sequence

$$I(X_{n-1}) \to I(X_n) \to I(\Gamma_n).$$

By Lemma 2.8, we have that $I(X_{n-1}) \to I(X_n)$ is null-homotopic since either $I(X_{n-1})$ or $I(X_n)$ is contractible. Hence we have

$$I(\Gamma_n) \simeq I(X_n) \lor \Sigma I(X_{n-1}).$$



FIGURE 2.

Lemma 2.8 completes the proof.

3. $I(\Gamma_{n,4})$

The goal of this section is to prove Theorem 1.1, which determines the precise homotopy type of $I(\Gamma_{n,4})$. Throughout this section, we write Γ_n to mean $\Gamma_{n,4}$.

First, define the graphs X_n and Y_n by $X_n = \Gamma_n - \{(n,2), (n,3)\}$ and $Y_n = \Gamma_n - \{(n,1), (n,4)\}$ (see Figure 1). The fold lemma implies

(3.1)
$$I(X_n) \simeq \Sigma^2 I(Y_{n-2}), \ I(Y_n) \simeq \Sigma I(X_{n-1}).$$

See Figures 2 and 3. We first determine the homotopy type of $I(X_n)$.

Lemma 3.1. There is a following homotopy equivalence:

$$I(X_n) \simeq \begin{cases} S^{n-1} & (n = 3k, 3k+2) \\ * & (n = 3k+1). \end{cases}$$

In particular, $I(X_n)$ is homotopy equivalent to a wedge of S^{n-1} .

Proof. The fold lemma implies $I(X_1) \simeq *$, $I(X_2) \simeq S^1$, and $I(X_3) \simeq \Sigma^2 I(Y_1) \simeq S^2$. For $n \ge 4$, we have $I(X_n) \simeq \Sigma^2 I(Y_{n-2}) \simeq \Sigma^3 I(X_{n-3})$. This completes the proof. \Box

Lemma 3.2. The following hold:



FIGURE 3.



FIGURE 4.

- (1) $I(\Gamma_n)$ is homotopy equivalent to a wedge of S^{n-1} .
- (2) For $n \geq 3$, there is a homotopy equivalence $I(\Gamma_n) \simeq \Sigma I(X_{n-1}) \vee \Sigma^2 I(\Gamma_{n-2})$.

Proof. We simultaneously prove both (1) and (2) by induction on n. The fold lemma implies $I(\Gamma_1) \simeq \text{pt}$ and $I(\Gamma_2) \simeq S^1$. Hence (1) holds for n = 1, 2.

Suppose $n \ge 3$. Set $G_n = \Gamma_n - \{(n-1,2), (n-1,3)\}$ for $n \ge 1$. The fold lemma implies $I(\Gamma_n) \simeq I(G_n)$. Set v = (n,4). Then we have $I(G_n - v) \simeq \Sigma I(X_{n-1})$ (Figure 4) and $I(G_n - N[v]) \simeq \Sigma I(\Gamma_{n-2})$ for $n \ge 3$ (see Figure 5). Hence Theorem 2.2 shows that there is a following cofiber sequence:

(3.2)
$$\Sigma I(\Gamma_{n-2}) \to \Sigma I(X_{n-1}) \to I(\Gamma_n)$$

Lemma 3.1 implies that $\Sigma I(X_{n-1})$ is a wedge of (n-1)-spheres, and $\Sigma I(\Gamma_{n-2})$ is a wedge of (n-2)-spheres by the inductive hypothesis. Hence the map $\Sigma I(\Gamma_{n-2}) \to I(X_{n-1})$ is null-homotopic, and we have $I(\Gamma_n) \simeq \Sigma I(X_{n-1}) \vee \Sigma^2 I(\Gamma_{n-2})$. This completes the proof of (2) in the case of n. Since $\Sigma I(X_{n-1})$ is a wedge of (n-1)-spheres and $\Sigma I(\Gamma_{n-2})$ is a wedge of (n-2)-spheres, the cofiber sequence (3.2) implies that $I(\Gamma_n)$ is a wedge of (n-1)-spheres. This completes the proof of (1) in the case of n.

Lemma 3.3. For $n \ge 7$, there is a homotopy equivalence

$$I(\Gamma_n) \simeq S^{n-1} \vee S^{n-1} \vee \Sigma^6 I(\Gamma_{n-6}).$$



FIGURE 5.

Proof. By (2) of Lemma 3.2, we have

$$I(\Gamma_n) \simeq \Sigma I(X_{n-1}) \vee \Sigma^2 I(\Gamma_{n-2})$$

$$\simeq \Sigma I(X_{n-1}) \vee \Sigma^3 I(X_{n-3}) \vee \Sigma^4 I(\Gamma_{n-4})$$

$$\simeq \Sigma I(X_{n-1}) \vee \Sigma^3 I(X_{n-3}) \vee \Sigma^5 I(X_{n-5}) \vee \Sigma^6 I(\Gamma_{n-6})$$

$$\simeq S^{n-1} \vee S^{n-1} \vee \Sigma^6 I(\Gamma_{n-6}).$$

The last homotopy equivalence follows from Lemma 3.1.

Proof of Theorem 1.1. Lemma 3.3 implies that the homotopy type of $I(\Gamma_n)$ is recursively determined by $I(\Gamma_1), \dots, I(\Gamma_6)$. Now we determine the homotopy types of these complexes. The fold lemma (Corollary 2.3) implies that

$$I(\Gamma_1) \simeq \text{pt}, \ I(\Gamma_2) \simeq S^1.$$

For $n \geq 3$, Lemma 3.1 and (2) of Lemma 3.2 implies

$$I(\Gamma_3) \simeq \Sigma I(X_2) \vee \Sigma^2 I(\Gamma_1) \simeq S^2, \ I(\Gamma_4) \simeq \Sigma I(X_3) \vee \Sigma^2 I(\Gamma_2) \simeq S^3 \vee S^3,$$

$$I(\Gamma_5) \simeq \Sigma I(X_4) \vee \Sigma^2 I(\Gamma_3) \simeq S^4, \ I(\Gamma_6) \simeq \Sigma I(X_5) \vee \Sigma^2 I(\Gamma_4) \simeq S^5 \vee S^5 \vee S^5.$$

Lemma 3.3 completes the proof.

4.
$$I(\Gamma_{n,5})$$

The goal of this section is to prove Theorem 1.2, which determines the homotopy type of $I(\Gamma_{n,5})$. Throughout this section, we write Γ_n to mean $\Gamma_{n,5}$. Since the proof is fairly intricate, we first explain its outline.

4.1. **Outline of the proof.** We introduce the graph $A_{n,k}$ recursively by k as follows. First, set $A_{n,0} = \Gamma_n$. Suppose that $A_{n,k}$ has been defined as an induced subgraph of Γ_{n+5k} . Then $A_{n,k+1}$ is defined as the induced subgraph of Γ_{n+5k+5} whose vertex set is

 $V(A_{n,k}) \cup (\{n''+1, n''+2, n''+5\} \times \{1,3\}) \cup (\{n''+2, n''+3, n''+4, n''+5\} \times \{1,5\}) \cup \{(n''+5,3)\},$



FIGURE 6.

For $n \geq 3$, let v be the vertex (n-2,3) of $A_{n,k}$. Theorem 2.2 implies that there is a cofiber sequence

(4.1)
$$I(A_{n,k} - N[v]) \to I(A_{n,k} - v) \to I(A_{n,k}).$$

We use this cofiber sequence to determine the homotopy type of $I(A_{n,k})$ inductively. We first show the following proposition, which will be proved in Subsection 4.4.

Proposition 4.1. Assume $n \ge 6$. Then there is a following homotopy equivalence:

$$I(A_{n,k} - N[v]) \simeq I(A_{n-5,k+1}).$$

By Proposition 4.1 and cofiber sequence (4.1), we have a following cofiber sequence

(4.2)
$$I(A_{n-5,k+1}) \to I(A_{n,k} - v) \to I(A_{n,k}),$$

The following proposition, which will be proved in Subsection 4.5, is a key to the whole proof:

Proposition 4.2. For $n \ge 6$, the inclusion $I(A_{n-5,k+1}) \simeq I(A_{n,k} - N[v]) \rightarrow I(A_{n,k} - v)$ is null-homotopic. Hence

$$I(A_{n,k}) \simeq I(A_{n,k} - v) \lor \Sigma I(A_{n-5,k+1}).$$

Hence to determine the homotopy types of $A_{n,k}$, it suffices to determine the homotopy types of $I(A_{n,k} - v)$ for every pair n and k, and $I(A_{n,k})$ for $n = 1, \dots, 5$. The following two propositions, which will be proved in Subsections 4.2 and 4.3, assert that we can determine the homotopy types of these complexes.

Proposition 4.3. The following hold:

(1) Assume k = 0, 1. Then

$$I(A_{n,k} - v) \simeq \begin{cases} \text{pt} & (n \text{ is odd}) \\ S^{5k+5l-1} & (n = 4l) \\ S^{5k+5l+2} & (n = 4l+2). \end{cases}$$

(2) Assume $k \geq 2$. Then

$$I(A_{n,k} - v) \simeq \begin{cases} \text{pt} & (n \text{ is odd}) \\ S^{5k+5l-1} \lor S^{5k+5l-1} & (n = 4l) \\ S^{5k+5l+2} \lor S^{5k+5l+2} & (n = 4l+2). \end{cases}$$

Proposition 4.4. The following hold:

(1)

$$I(A_{1,k}) \simeq \begin{cases} S^1 & (k=0) \\ \text{pt} & (k \ge 1) \end{cases}$$



FIGURE 7.

(2)

$$I(A_{2,k}) \simeq \begin{cases} S^{5k+2} & (k=0,1) \\ S^{5k+2} \lor S^{5k+2} & (k \ge 2) \end{cases}$$

(3)

$$I(A_{3,k}) \simeq \begin{cases} S^{5k+3} & (k=0) \\ S^{5k+3} \lor S^{5k+3} & (k \ge 1) \end{cases}$$

(4)

$$I(A_{4,k}) \simeq \begin{cases} S^{5k+4} & (k=0,1) \\ S^{5k+4} \lor S^{5k+4} & (k \ge 2) \end{cases}$$

(5)

$$I(A_{5,k}) \simeq \begin{cases} S^{5k+5} & (k=0) \\ S^{5k+5} \lor S^{5k+5} & (k \ge 1) \end{cases}$$

The rest of this section is organized as follows. Subsections 4.2, 4.3, 4.4, and 4.5 are devoted to the proofs of Propositions 4.3, 4.4, 4.1, and 4.2, respectively. In Subsection 4.6, we complete the proof of Theorem 1.2.

4.2. **Proposition 4.3.** The goal of this subsection is to show Proposition 4.3, which determines the homotopy type of $I(A_{n,k}-v)$. Before proceeding the proof, we need to determine the homotopy types of independence complexes of graphs X_n (see Lemma 4.5) and $B_{n,k}$ (see Lemma 4.6).

For $n \ge 1$, let X_n be the graph $\Gamma_n - \{(n, 2), (n, 4)\}$ and Y_n the graph $\Gamma_n - \{(n, 1), (n, 3), (n, 5)\}$ (see Figure 7). We first determine the homotopy type of $I(X_n)$. By the fold lemma (Corollary 2.3), it is clear that

(4.3)
$$I(X_n) \simeq \Sigma^3 I(Y_{n-2}), \ I(Y_n) \simeq \Sigma^2 I(X_{n-2})$$

for $n \ge 3$. Hence for $n \ge 5$, we have

(4.4)
$$I(X_n) \simeq \Sigma^5 I(X_{n-4}).$$



FIGURE 8.

Thus the homotopy type of $I(X_n)$ is determined by the homotopy types of $I(X_1), \dots, I(X_4)$. The fold lemma implies

$$I(X_1) \simeq \operatorname{pt}, I(X_2) \simeq S^2, \ I(Y_1) \simeq \operatorname{pt}, I(Y_2) \simeq S^1.$$

Hence we have

$$I(X_3) \simeq \Sigma^3 I(Y_1) \simeq \text{pt}, \ I(X_4) \simeq \Sigma^3 I(Y_2) \simeq S^4.$$

Thus (4.4) implies the following lemma:

Lemma 4.5.
$$I(X_n) \simeq \begin{cases} \text{pt} & (n \text{ is odd}) \\ S^{5l-1} & (n = 4l) \\ S^{5l+2} & (n = 4l+2) \end{cases}$$

Next we determine the homotopy types of the independence complexes of other families of graphs $B_{n,k}$ and $B'_{n,k}$. For $k \ge 1$, define the graph $B_{n,k}$ to be the induced subgraph of $A_{n,k}$ whose vertex set is $V(A_{n,k}) - V(\Gamma_n)$. For $k \ge 1$, define $B'_{n,k}$ to be the induced subgraph of $B_{n,k}$ whose vertex set is

$$(V(B_{n,k}) - V(\Gamma_{n+5})) \cup \{(n+5,2), (n+5,3), (n+5,4)\}.$$



FIGURE 9.



FIGURE 10.

Figures 8 and 9 depict the graphs $B_{n,3}$ and $B'_{n,3}$, respectively. Note that the isomorphism types of graphs $B_{n,k}$ and $B'_{n,k}$ only depend on k.

Lemma 4.6. The following hold:

(1)

$$I(B'_{n,k}) \simeq \begin{cases} S^0 & (k=1) \\ S^{5k-5} \lor S^{5k-5} & (k \ge 2) \end{cases}$$

(2)

$$I(B_{n,k}) \simeq \begin{cases} S^4 & (k=1) \\ S^{5k-1} \lor S^{5k-1} & (k \ge 2) \end{cases}$$

Proof. For $k \ge 1$, the fold lemma implies $I(B_{n,k}) \simeq \Sigma^4 I(B'_{n,k})$. Hence (1) implies (2). Thus it suffices to show (1).

Since $B'_{n,1}$ is isomorphic to P_3 , we have $I(B'_{n,1}) \simeq I(P_3) \simeq S^0$ by Theorem 2.4. Now we assume $k \geq 2$. Set x = (n + 5, 4). Then the fold lemma shows that $I(B'_{n,k} - x) \simeq$ $\Sigma^5 I(B'_{n,k-1} - x)$ (see Figure 10), and the induction on k implies $I(B'_{n,k} - x) \simeq S^{5k-5}$. On the other hand, $I(B'_{n,k} - N[x]) \simeq \Sigma^4 I(B'_{n,k-1} - x) \simeq S^{5k-6}$ (see Figure 11). Hence the inclusion $I(B'_{n,k} - N[x]) \to I(B'_{n,k} - x)$ is null-homotopic. Thus we have

$$I(B'_{n,k}) \simeq I(B'_{n,k} - x) \lor \Sigma I(B'_{n,k} - N[x]) \simeq S^{5k-5} \lor S^{5k-5}.$$

This completes the proof.

Proof of Proposition 4.3. By the fold lemma, one can show that $I(A_{n,0} - v) \simeq I(X_n)$ and $I(A_{n,k} - v) \simeq I(X_n \sqcup B_{n,k}) = I(X_n) * I(B_{n,k})$ (see Figure 12). The proof is completed by Lemmas 4.5 and 4.6.

4.3. **Proposition 4.4.** The goal of this subsection is to prove Proposition 4.4, which determines the homotopy types of $I(A_{n,k})$ for $n = 1, \dots, 5$.

Proof of (1) of Proposition 4.4. When k = 0, $A_{1,k}$ is isomorphic to $\Gamma_{5,1}$ and hence $I(A_{1,0}) \simeq S^1$ follows from Theorem 2.4.

Suppose $k \geq 1$. The fold lemma implies

$$I(A_{1,k}) \simeq I(A_{1,k} - \{(2,2), (2,4), (4,1), (4,5), (6,2), (6,4)\}).$$

Since (6,3) is an isolated vertex of $A_{1,k} - \{(2,2), (2,4), (4,1), (4,5), (6,2), (6,4)\}$, the right of the above is contractible. This completes the proof of (1).

Proof of (2) of Proposition 4.4. The case k = 0 follows from Theorem 2.5. For $k \ge 1$, the fold lemma implies

$$I(A_{2,k}) \simeq I(K_2 \sqcup K_2 \sqcup K_2 \sqcup B_{n,k}) = \Sigma^3 I(B_{n,k}).$$

Thus the proof is completed by Lemma 4.6.

Proof of (3) of Proposition 4.4. Recall v = (1,3) in this case. The fold lemma implies that

$$I(A_{3,k} - v) \simeq I(A_{3,k} - \{(1,3), (2,2), (2,4), (3,2), (1,1)\}),$$

and $A_{3,k} - \{(1,3), (2,2), (2,4), (3,2), (1,1)\}$ has isolated vertex (1,2). Hence $I(A_{3,k} - v)$ is contractible and hence $I(A_{3,k}) \simeq \Sigma I(A_{3,k} - N[v])$. The fold lemma implies that $I(A_{3,k} - N[v]) \simeq \Sigma^2 I(B'_{n,k+1})$ for $k \ge 0$. Hence the proof is completed by Lemma 4.6.

Proof of (4) of Proposition 4.4. The fold lemma implies that

$$I(A_{n,k} - N[v]) \simeq I(A_{4,k} - (N[v] \cup \{(2,1), (2,5), (4,2), (4,4)\}))$$

Since (4,3) is an isolated vertex of $A_{4,k} - (N[v] \cup \{(2,1), (2,5), (4,2), (4,4)\}$, this means that $I(A_{4,k} - N[v])$ is contractible. Thus Theorem 2.2 implies that $I(A_{4,k}) \simeq I(A_{4,k} - v)$. It follows from the fold lemma that $I(A_{4,0} - v) \simeq S^4$ and $I(A_{4,k}) \simeq \Sigma^5 I(B_{4,k})$ for $k \ge 1$. The proof is completed by Lemma 4.6.





FIGURE 12.

Proof of (5) of Proposition 4.4. Since $I(A_{5,k} - v) \simeq pt$, we have $I(A_{n,k}) \simeq \Sigma I(A_{5,k} - N[v])$. The fold lemma implies that $I(A_{5,k} - N[v])$ is isomorphic to the graph W_k depicted in Figure 13, and hence

(4.5)
$$I(A_{5,k}) \simeq \Sigma I(W_k)$$

Set u = (1, 3). Then

 $I(W_k - N[v]) \simeq I(W_k - (N[v] \cup \{(3,1), (3,5), (5,2), (5,3)\})).$

Since (5,3) is an isolated vertex in $W_k - (N[v] \cup \{(3,1), (3,5), (5,2), (5,3)\})$, we have $I(W_k - N[v]) \simeq \text{pt.}$ This shows

$$I(W_k) \simeq I(W_k - u).$$

The fold lemma shows that $I(W_k - u) \simeq \Sigma^4 I(B'_{5,k+1})$. Lemma 4.6 completes the proof. \Box

4.4. **Proposition 4.1.** The goal of this subsection is to prove Proposition 4.1, which states that $I(A_{n,k} - N[v]) \simeq I(A_{n-5,k+1})$ for $n \ge 6$.

Proof of Proposition 4.1. Set w = (n - 4, 3). The fold lemma implies that $I(A_{n,k} - N[v])$ is homotopy equivalent to the independence complex of the induced subgraph A' of $A_{n,k}$ whose vertex set is $V(A_{n-5,k+1}) \cup \{w\}$ (see Figure 14). Since $A' - w = A_{n-5,k+1}$, it suffices to show that I(A' - N[w]) is contractible (Theorem 2.2).





FIGURE 14.

The fold lemma implies

$$I(A' - N[w]) \simeq I(A' - (N[w] \cup \{(n-2,5), (n,4), (n,1), (n-3,1)\}))$$

Since (n-3,2) is an isolated vertex of $A' - (N[w] \cup \{(n-2,5), (n,4), (n,1), (n-3,1)\}$, we conclude that I(A' - N[w]) is contractible (see Figure 15).

4.5. **Proposition 4.2.** In this subsection, we prove Proposition 4.2, which states that the map $I(A_{n-5,k+1}) \simeq I(A_{n,k} - N[v]) \hookrightarrow I(A_{n,k} - v)$ is null-homotopic for $n \ge 6$. Note that $I(A_{n,k} - v) \simeq$ pt when n is odd, and that in this case $I(A_{n,k} - N[v]) \to I(A_{n,k} - v)$ is null-homotopic. Hence we prove only in the case that n is even.

Proposition 4.7. Assume that n is even. Then the following hold:

- (1) If n = 4l, $I(A_{4l,k})$ is homotopy equivalent to a wedge of spheres whose dimension is at most 5k + 5l 1. If n = 4l + 2, then $I(A_{4l+2,k})$ is homotopy equivalent to a wedge of spheres whose dimension is at most 5k + 5l + 2.
- (2) The inclusion $I(A_{n,k} N[v]) \rightarrow I(A_{n,k} v)$ is null-homotopic.

Proof. We simultaneously prove (1) and (2) by induction on n. Since the cases n = 6, 8, 10 can be directly proved by Propositions 4.1, 4.3, and 4.4, we assume that $n \ge 12$. Then we



FIGURE 15.

have a cofiber sequence

$$I(A_{n-5,k+1}) \to I(A_{n,k} - v) \to I(A_{n,k})$$

and

$$I(A_{n-10,k+2}) \to I(A_{n-5,k+1} - v) \to I(A_{n-5,k+1}).$$

Since n-5 is odd, we have $I(A_{n-5,k+1}-v) \simeq pt$, and hence we have $I(A_{n-5,k+1}) \simeq \Sigma I(A_{n-10,k+2})$. To prove (2), it suffices to see that every map from $\Sigma I(A_{n-10,k+2})$ to $I(A_{n,k}-v)$ is null-homotopic.

Suppose that n = 4l. Then Proposition 4.3 implies that $I(A_{n,k} - v) = I(A_{4l,k} - v)$ is a wedge of spheres whose dimension is 5k + 5l - 1. It follows from the inductive hypothesis that $\Sigma I(A_{n-10,k+2}) = \Sigma I(A_{4(l-3)+2,k+2})$ is a wedge of spheres whose dimension is at most

$$5(l-3) + 5(k+2) + 2 + 1 = 5k + 5l - 2$$

Hence every map from $\Sigma I(A_{4l-10,k+2})$ to $I(A_{4l,k}-v)$ is null-homotopic, and we have proved (2) in the case n = 4l. Since $I(A_{4l,k}) \simeq I(A_{4l,k}-v) \vee \Sigma^2 I(A_{4l-10,k+2})$, we conclude that $I(A_{4l,k})$ is a wedge of spheres whose dimension is at most 5k + 5l - 1. This completes the proof of (1) in the case n = 4l.

Next suppose that n = 4l+2. Then Proposition 4.3 implies that $I(A_{n,k}-v) = I(A_{4l+2,k})$ is a wedge of spheres whose dimension is 5k+5l+2. It follows from the inductive hypothesis that $\Sigma I(A_{n-10,k+2}) = \Sigma I(A_{4(l-2),k+2})$ is a wedge of spheres whose dimension is at most

$$5(l-2) + 5(k+2) - 1 + 1 = 5k + 5l.$$

Hence we have proved (2) in the case n = 4l + 2. Since

$$I(A_{4l+2,k}) \simeq I(A_{4l+2,k} - v) \vee \Sigma^2 I(A_{4l-8,k+2}),$$

we conclude that $I(A_{4l+2,k})$ is a wedge of spheres whose dimension is at most 5k + 5l + 2. This completes the proof of (1) in the case n = 4l + 2, and we have completed the proof of this proposition.

Proof of Proposition 4.2. The proof is completed by Propositions 4.1 and 4.7.

4.6. **Proof of the main theorem.** In this subsection we complete the proof of Theorem 1.2, using the results in the previous subsections.

Lemma 4.8. The following hold:

- (1) If n is odd and $n \ge 7$, then $I(A_{n,k}) \simeq \Sigma I(A_{n-5,k+1})$.
- (2) If *n* is even and $n \ge 12$, then $I(A_{n,k}) \simeq I(A_{n,k} v) \vee \Sigma^2 I(A_{n-10,k+2})$.

Proof. Suppose *n* is odd. It follows from Proposition 4.3 that $I(A_{n,k} - v)$ is contractible. Hence Theorem 2.2 and Proposition 4.1 imply $I(A_{n,k}) \simeq \Sigma I(A_{n,k} - N[v]) \simeq \Sigma I(A_{n-5,k+1})$. This completes the proof of (1).

Suppose that n is even and $n \ge 12$. By Proposition 4.2 we have

$$I(A_{n,k}) \simeq I(A_{n,k} - v) \vee \Sigma I(A_{n-5,k+1})$$

$$\simeq I(A_{n,k} - v) \vee \Sigma^2 I(A_{n-10,k+2})$$

Here the second homotopy equivalence follows from (1). This completes the proof. $\hfill \Box$

Corollary 4.9. There exists a following homotopy equivalence

$$I(A_{4l,k}) \simeq \begin{cases} \left(\bigvee_{4} S^{5k+5l-1}\right) \vee \Sigma^4 I(A_{4(l-5),k+4}) & (k=0,1) \\ \left(\bigvee_{4}^3 S^{5k+5l-1}\right) \vee \Sigma^4 I(A_{4(l-5),k+4}) & (k \ge 2). \end{cases}$$

Proof. Lemma 4.8 implies

$$I(A_{4l,k}) \simeq I(A_{4l,k} - v) \vee \Sigma^2 I(A_{4l-10,k+2})$$

$$\simeq I(A_{4l,k} - v) \vee \Sigma^2 I(A_{4l-10,k+2} - v) \vee \Sigma^4 I(A_{4l-20,k+4}).$$

Proposition 4.3 completes the proof.

Lemma 4.10. The following hold:

(1)

$$I(A_{4,k}) \simeq \begin{cases} S^{5k+4} & (k=0,1) \\ \bigvee_2 S^{5k+4} & (k \ge 2). \end{cases}$$

(2) For l = 2, 3, 4, there exists a following homotopy equivalence:

$$I(A_{4l,k}) \simeq \begin{cases} \bigvee_3 S^{5k+5l-1} & (k=0,1) \\ \bigvee_4 S^{5k+5l-1} & (k \ge 2). \end{cases}$$

(3)

$$I(A_{20,k}) \simeq \begin{cases} \bigvee_3 S^{5k+24} \lor \bigvee_2 S^{5k+23} & (k=0,1) \\ \bigvee_4 S^{5k+24} \lor \bigvee_2 S^{5k+23} & (k \ge 2). \end{cases}$$

Proof. We have already proved (1) in Proposition 4.4. To prove (2) and (3), we use the following homotopy equivalences

$$I(A_{8,k}) \simeq I(A_{8,k} - v) \lor \Sigma I(A_{3,k+1}),$$

$$I(A_{12,k}) \simeq I(A_{12,k} - v) \lor \Sigma^2 I(A_{2,k+2}),$$

$$I(A_{16,k}) \simeq I(A_{16,k} - v) \lor \Sigma^2 I(A_{6,k+2}) \simeq I(A_{16,k} - v) \lor \Sigma^2 I(A_{6,k+2} - v) \lor \Sigma^3 I(A_{1,k+3}),$$

$$I(A_{20,k}) \simeq I(A_{20,k} - v) \lor \Sigma^2 I(A_{10,k+2}) \simeq I(A_{20,k} - v) \lor \Sigma^2 I(A_{10,k+2} - v) \lor \Sigma^3 I(A_{5,k+3}).$$

Here we use Proposition 4.2 and Lemma 4.8 to derive these homotopy equivalences. Then Propositions 4.3 and 4.4 complete the proof. $\hfill \Box$

17

When n is even, the homotopy type of $I(A_{n,k})$ is determined by the following two propositions. Here we write $n = 20t + 4l + \varepsilon$ with $l \in \{0, 1, 2, 3, 4\}$ and $\varepsilon \in \{0, 1, 2, 3\}$. Define $\kappa \in \{0, 1\}$ by 0 when k = 0, 1 and by 1 when $k \ge 2$.

Proposition 4.11. Consider the case $\varepsilon = 0$ and assume $n \ge 8$. Then the following hold: (1) For l = 0, 1 (and hence $t \ge 1$), we have

$$I(A_{20t+4l,k}) \simeq \bigvee_{3+\kappa} S^{25t+5l+5k-1} \lor \left(\bigvee_{i=24t+5l+5k}^{25t+5l+5k-2} \left(\bigvee_{4}^{j} S^{i}\right)\right) \lor \bigvee_{2} S^{24t+5l+5k-1}.$$

(2) For l = 2, 3, 4, we have

$$I(A_{20t+4l,k}) \simeq \bigvee_{3+\kappa} S^{25t+5l+5k-1} \lor \Big(\bigvee_{i=24t+5l+5k-1}^{25t+5l+5k-2} \Big(\bigvee_{4}^{k} S^{i}\Big)\Big).$$

Proof. Corollary 4.9 implies

$$I(A_{4l+20t,k}) \simeq \begin{cases} \left(\bigvee_{3} S^{25t+5l+5k-1}\right) \vee \Sigma^{4} I(A_{20(t-1)+4l,k+4}) & (k=0,1) \\ \left(\bigvee_{4} S^{25t+5l+5k-1}\right) \vee \Sigma^{4} I(A_{20(t-1)+4l,k+4}) & (k \ge 2). \end{cases}$$

Lemma 4.10 completes the proof.

Proposition 4.12. Consider the case $\varepsilon = 2$. For every $n \ge 2$ (i.e. $t, l \ge 0$), the following hold:

(1) For l = 0, 1, we have

$$I(A_{20t+4l+2,k}) \simeq \bigvee_{1+\kappa} S^{25t+5l+5k+2} \lor \Big(\bigvee_{i=24t+5l+5k+2}^{25t+5l+5k+1} \Big(\bigvee_{4} S^i\Big)\Big).$$

(2) For l = 2, 3, we have

$$I(A_{20t+4l+2,k}) \simeq \bigvee_{1+\kappa} S^{25t+5l+5k+2} \lor \left(\bigvee_{i=24t+5l+5k+2}^{25t+5l+5k+1} \left(\bigvee_{4} S^{i}\right)\right) \lor \bigvee_{2} S^{24t+5l+5k+1}.$$

(3) For l = 4, we have

$$I(A_{20t+4l+2,k}) \simeq \bigvee_{1+\kappa} S^{25t+5l+5k+2} \lor \Big(\bigvee_{i=24t+5l+5k+1}^{25t+5l+5k+1} \Big(\bigvee_{4} S^{i}\Big)\Big).$$

Proof. We have already determined the homotopy type of $I(A_{2,k})$ in Proposition 4.4. Proposition 4.2 and Lemma 4.8 imply

$$I(A_{6,k}) \simeq I(A_{6,k} - v) \lor \Sigma I(A_{1,k+1}),$$

$$I(A_{10,k}) \simeq I(A_{10,k} - v) \lor \Sigma I(A_{5,k+1}),$$

$$I(A_{14,k}) \simeq I(A_{14,k} - v) \lor \Sigma^2 I(A_{4,k+2}).$$

Thus the homotopy types of $I(A_{6,k})$, $I(A_{10,k})$ and $I(A_{14,k})$ are determined by Propositions 4.3 and 4.4. For $n \ge 18$, Lemma 4.8 implies $I(A_{n,k}) \simeq I(A_{n,k} - v) \lor \Sigma^2 I(A_{n-10,k+2})$. Its homotopy type is determined in Proposition 4.11 since $n - 10 \ge 8$.

Proof of Theorem 1.2. Since $\Gamma_n = A_{n,0}$, the homotopy type of $I(\Gamma_n)$ for even n is determined by Propositions 4.11 and 4.12. The homotopy types of $I(\Gamma_1)$, $I(\Gamma_3)$, and $I(\Gamma_5)$ are determined in Proposition 4.4. For an odd number n with n > 5, (1) of Lemma 4.8 implies

$$I(\Gamma_n) \simeq \Sigma I(A_{n-5,1}),$$

and the homotopy type of $I(A_{n-5,1})$ is determined by Propositions 4.11 and 4.12. This completes the proof.

Acknowledgment

The first author is partially supported by JSPS KAKENHI Grant Number JP19K14536. The second author is supported by JSPS KAKENHI Grant Number JP20J00404.

References

- [A1] M. Adamaszek; Splittings of independence complexes and the powers of cycles, J. Comb. Theory Ser. A, 119 (2012) 1031-1047.
- [A2] M. Adamaszek, Hard squares on cylinders revisited arXiv:1202.1655
- [AGS] A. Singh, S. Goyal, S. Shukla, Matching complexes of 3 × n-grid graphs, Electr. J. Combin. (4) 28 # P4.16, 2021.
- [B] J. A. Barmak; Star clusters in independence complexes of graphs, Adv. Math. 241 (2013) 33-57.
- [BGJM] M. Bayer, B. Goeckner, M. Jelić Milutinović, Manifold matching complexes, Mathematika 66 (2020) 973-1002.
- [BH] B. Braun, W. K. Hough. Matching and independence complexes related to small grids, Electron. J. Combin., 18(1), 2011.
- [BMLN] M. Bousquet-Mélou, S. Linusson, E. Nevo, On the independence complex of square grids, J. Algebr. Comb. 2008, 27: 423-450
- [EH] R. Ehrenborg, G. Hetyei; The topology of independence complex, Eur. J. Comb. 27 (2006) 906-923.
- [E] A. Engström; Complexes of directed trees and independence complexes, Discrete Math. 309 (2009) 3299-3309.
- K. Iriye, On the homotopy types of the independence complexes of grid graphs with cylindrical identification, Kyoto J. Math., 52(3): 479-501 (2012)
- [J1] J. Jonsson, Hard squares with negative activity and rhombus tilings of the plane, Electr. J. Comb. 13 (1) (2006), #R67
- [J2] J. Jonsson, Hard Squares with negative activity on cylinders with odd circumference, Electr. J. Comb. 16 (2) (2009), #R5
- [J3] J. Jonsson, Certain homology cycles of the independence complex of grids, Discrete & Computational Geometry 43(4): 927-950 (2010)
- [K1] D.N. Kozlov, Complexes of directed trees, J. Comb. Theory Ser. A, 88 (1999) 112-122.
- [K2] D.N. Kozlov, Combinatorial algebraic topology, Springer, Berlin, Algorithms and Computation in Mathematics, Vol. 21, 2008.
- [M] T. Matsushita, Matching complexes of small grids, Electron. J. Combin. Volume 26, Issue 3, (July, 2019)
- [O] K. Okura, Simple homotopy types of independence complexes of graphs involving grid graphs, arXiv:1908.09356
- [PRSZ] L. Polterovich, D. Rosen, K. Samvelyan, J. Zhang, Topological Persistence in Geometry and Analysis, University Lecture Series, American Mathematical Society, 2020,

T. MATSUSHITA AND S. WAKATSUKI

- [T] J. Thapper, Independence complexes of cylinders constructed from square and hexagonal grid graphs, arXiv:0812.1165
- [W] M. L. Wachs. Topology of matching, chessboard, and general bounded degree graph complexes, Algebra Universalis, 49 (4):345-385, 2003.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF THE RYUKYUS, NISHIHARA-CHO, OKINAWA 903-0213, JAPAN

Email address: mtst@sci.u-ryukyu.ac.jp

Department of Mathematical Sciences, Faculty of Science, Shinshu Universigy, Matsumoto, Nagano 390-8621, Japan

Email address: swaka@shinshu-u.ac.jp