WEIGHTED HOMOMORPHISMS BETWEEN C*-ALGEBRAS

EUSEBIO GARDELLA AND HANNES THIEL

ABSTRACT. We show that a bounded, linear map between C^* -algebras is a weighted ∗-homomorphism (the central compression of a ∗-homomorphism) if and only if it preserves zero-products, range-orthogonality, and domainorthogonality. It follows that a self-adjoint, bounded, linear map is a weighted ∗-homomorphism if and only if it preserves zero-products.

As an application we show that a linear map between C^* -algebras is completely positive, order zero in the sense of Winter-Zacharias if and only if it is positive and preserves zero-products.

1. INTRODUCTION

The study of bounded, linear maps between C^* -algebras that preserve various types of orthogonality has a long history. One of the earliest results, due to Arendt [\[Are83\]](#page-16-0), is the description of such maps between unital, commutative C^* -algebras: every bounded, linear map $\varphi: C(X) \to C(Y)$ that preserves zero-products (such maps are also called *disjointness preserving*, or *separating*) is spatially implemented, that is, setting $h := \varphi(1) \in C(Y)$ and $U := \{y \in Y : h(y) \neq 0\}$, there exists a continuous map $p: U \to X$ such that $\varphi(f)(y) = h(y)f(p(y))$ for every $y \in Y$. Note that p induces a *-homomorphism $\pi: C(X) \to C_b(U)$, $f \mapsto f \circ p$. By embedding $C(Y)$ and $C_b(U)$ into a suitably larger (commutative) C^* -algebra – the canonical choice is $C(Y)^{**}$ – we obtain that $\varphi(f) = h\pi(f) = \pi(f)h$ for all $f \in C(X)$. In particular, φ is the compression of the ∗-homomorphism π by h. For this description of φ as the compression of a *-homomorphism, it is crucial to allow for an enlarged target algebra. Indeed, an example of Wolff [\[Wol94\]](#page-17-0), shows that the image of π is not in general contained in $C(Y)$.

Generalizing this idea to the noncommutative setting, we define:

Definition A. We say that a map $\varphi: A \to B$ between C^* -algebras is a *weighted* **-homomorphism* if there exist a C^{*}-algebra D containg B, a *-homomorphism $\pi: A \to D$ and an element $h \in D$ such that $\varphi(a) = \pi(a)h = h\pi(a)$ for all $a \in A$.

By the above-mentioned result of Arendt, it follows that a bounded, linear map between unital, commutative C^* -algebras is a weighted $*$ -homomorphism if and only if it preserves zero-products. The main result of this paper is a generalization of this characterization to the noncommutative setting; see [Theorem 4.2.](#page-13-0) In this case, we also need to consider the following types of orthogonality: two elements a and b in a C^* -algebra are said to be *range-orthogonal* if $a^*b = 0$; they are *domainorthogonal* if $ab^* = 0$.

 $Date: 1$ April 2022.

²⁰¹⁰ Mathematics Subject Classification. Primary 47A65, 47B49; Secondary 46L05, 47A20, 47B65.

Key words and phrases. C∗-algebras, weighted homomorphisms, zero-products, rangeorthogonality, domain-orthogonality.

The first named author was partially supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through an Eigene Stelle, and by the Swedish Research Council Grant 2021-04561. The second named author was partially supported by the ERC Consolidator Grant No. 681207.

Theorem B [\(4.2\)](#page-13-0). A bounded, linear map between C^* -algebras is a weighted ∗-homomorphism if and only if it preserves range-orthogonality, domain-orthogonality and zero-products.

Note that two elements in a *commutative* C^* -algebra are range-orthogonal if and only if they are domain-orthogonal, and if and only if they have zero-product. In particular, our [Theorem B](#page-1-0) generalizes the description of bounded, linear mpas between unital, commutative C^* -algebras that preserve zero-products.

Bounded, linear maps between (noncommutative) C^* -algebras that preserve zero-products have been extensively studied by Wong and coauthors, [\[CKLW03,](#page-16-1) [KLW04,](#page-16-2) [Won07\]](#page-17-1), and a general description of the structure of such maps seems out of reach at the moment. However, a zero-product preserving linear map which is additionally self-adjoint (equivalently, ∗-preserving) automatically preserves domain and range orthogonality, and hence we obtain:

Corollary C (4.4) . A self-adjoint, bounded, linear map between C^* -algebras is a weighted ∗-homomorphism if and only if it preserves zero-products.

The above corollary is the analog of a result of Wolff, [\[Wol94\]](#page-17-0), who showed that a self-adjoint, bounded linear map between C^* -algebras that preserves zero-products *among self-adjoint elements* is a weigthed Jordan ∗-homomorphisms. (Wolff showed the result under the assumption that the domain is unital. This condition can be removed using [\[Won07,](#page-17-1) Lemma 2.2].)

In [\[WZ09\]](#page-17-2), Winter and Zacharias introduced completely positive maps of *order* $zero$ as those completely positive maps between C^* -algebras that preserve zeroproducts among positive elements (equivalently, among self-adjoint elements). The main result of [\[WZ09\]](#page-17-2) is a characterization of completely positive maps of order zero as the *positively* weighted ∗-homomorphisms. These maps play an important role in the fine structure theory of nuclear C^* -algebras; see [\[HKW12,](#page-16-3) [WZ10\]](#page-17-3).

Using that a positive, linear map between C^* -algebras is automatically bounded and self-adjoint, we obtain the following new characterization of completely positive, order zero maps:

Corollary D (4.8) . A map between C^* -algebras is completely positive, order zero if and only if it is positive and preserves zero-products.

In particular, every positive, zero-product preserving map between C^* -algebras is completely positive (and order zero).

In [Section 2,](#page-1-1) we study bounded, linear maps between C^* -algebras that preserve zero-products, range-orthogonality, or domain-orthogonality and we provide characterizations of these properties in terms of algebraic identities. For example, a bounded, linear map φ preserves zero-products if and only if $\varphi(ab)\varphi(c) = \varphi(a)\varphi(bc)$ for all a, b, c in the domain; see [Theorem 2.2.](#page-2-0) Using this, we show that preserving these types of orthogonality passes to bitransposes and tensor products.

We also consider maps preserving zero-TRO-products or 'usual' orthogonality. In [Section 3,](#page-6-0) we consider a bounded, linear map $\varphi: A \to B$ between C^* -algebras, and a ∗-representation $B \subseteq \mathcal{B}(H)$ of B on some Hilbert space H. Assuming that π preserves zero-products, range-orthogonality, and domain-orthogonality, we construct a (canonical) *-homomorphism $\pi_{\varphi} : A \to \mathcal{B}(H)$ that takes image in the bicommutant $B'' \subseteq \mathcal{B}(H)$ and that satisfies $\pi_{\varphi}(a)\varphi(b) = \varphi(ab)$ and $\varphi(a)\pi_{\varphi}(b) =$ $\varphi(ab)$ for all $a, b \in A$. This is the main technical step in the proof of our main result, which we complete in [Section 4.](#page-12-0)

2. Maps preserving different notions of orthogonality

In this section, we study bounded linear maps between C^* -algebras that preserve certain notions of orthogonality. We begin by defining these notions.

Definition 2.1. Let A be a C^* -algebra, and let $a, b, c \in A$. We say that

- a and b have *zero-product* if $ab = 0$,
- *a* and *b* are *range-orthogonal* if $a^*b = 0$;
- a and b are *domain-orthogonal* if $ab^* = 0$;
- *a* and *b* are *orthogonal* if $a^*b = ab^* = 0$;
- *a*, *b* and *c* have *zero-TRO-product* if $ab^*c = 0$.

We will be interested in maps preserving certain combinations of the above notions. More explicitly, we say that a map $\varphi: A \to B$ between C^* -algebras preserves:

- (1) zero-products if $ab = 0$ implies $\varphi(a)\varphi(b) = 0$
- (2) range-orthogonality if $a^*b = 0$ implies $\varphi(a)^*\varphi(b) = 0$;
- (3) domain-orthogonality if $ab^* = 0$ implies $\varphi(a)\varphi(b)^* = 0$;
- (4) orthogonality if $ab^* = a^*b = 0$ implies $\varphi(a)\varphi(b)^* = \varphi(a)^*\varphi(b) = 0;$
- (5) zero-TRO-products if $ab^*c = 0$ implies $\varphi(a)\varphi(b)^*\varphi(c) = 0$.

We provide characterizations of (1) , (2) , (3) and (5) in terms of algebraic equations [\(Theorem 2.2\)](#page-2-0), which we then use to show that each of those properties passes to the bitranspose [\(Proposition 2.3\)](#page-4-0) and to tensor products [\(Proposition 2.4\)](#page-5-0). In particular, if a bounded linear map $A \rightarrow B$ preserves zero-products, then so does every amplification $M_n(A) \to M_n(B)$ – and analogously for maps preserving rangeorthogonality, preserving domain-orthogonality, or preserving zero-TRO-products; see [Corollary 2.5.](#page-6-1) In particular, a zero-product preserving map is automatically completely zero-product preserving (and similarly for range-orthogonality, domainorthogonality, and zero-TRO-products).

The situation for orthogonality-preserving maps is different: As noted in [\[Gar20,](#page-16-4) Section 4, the transpose map τ on $M_2(\mathbb{C})$ preserves orthogonality, but its amplification $\tau^{(2)}$ does not. Thus, orthogonality-preservation does not pass to tensor products. Nevertheless, using the structure result for orthogonality-preserving maps obtained in $[BFPG^+08]$, we show that the bitranspose of an orthogonality-preserving map is again orthogonality-preserving; see [Proposition 2.3.](#page-4-0)

Part (1) of the next result follows from [\[ABEV09,](#page-16-6) Lemma 3.4]. (Note that by Examples 1.3(2) and Theorem 2.11 in [\[ABEV09\]](#page-16-6), every C^* -algebra has property ($\mathbb B$) introduced in [\[ABEV09\]](#page-16-6).) The proof in [\[ABEV09,](#page-16-6) Lemma 3.4] is based on the fact that every element in a unital C^* -algebra is a linear combination of unitaries and that unitaries are doubly power-bounded operators. For maps between von Neumann algebras (and more generally, C^* -algebras of real rank zero), Part (1) of the next result has also been obtained in [\[CKLW03,](#page-16-1) Theorem 4.1], using that the linear span of projections is dense.

Our proof is inspired by [\[CKLW03,](#page-16-1) Lemma 4.4], and uses methods that go back to Wolff, [\[Wol94\]](#page-17-0). These methods allow us to also characterize maps that preserve range-orthogonality or domain-orthogonality.

Theorem 2.2. Let $\varphi: A \to B$ be a bounded, linear map between C^* -algebras. *Then:*

- (1) φ preserves zero-products if and only if $\varphi(ab)\varphi(c) = \varphi(a)\varphi(bc)$ for all $a, b, c \in A$.
- (2) φ preserves range-orthogonality if and only if $\varphi(b^*a)^*\varphi(c) = \varphi(a)^*\varphi(bc)$ for *all* $a, b, c \in A$.
- (3) φ preserves domain-orthogonality if and only if $\varphi(ab)\varphi(c)^* = \varphi(a)\varphi(cb^*)^*$ *for all* $a, b, c \in A$ *.*
- (4) φ preserves zero-TRO-products if and only if the equality $\varphi(ab)\varphi(c)^{*}\varphi(de)$ $\varphi(a)\varphi(d^*cb^*)^*\varphi(e)$ holds for all $a, b, c, d, e \in A$.

Proof. We only prove statement (2). Statements (1), (3) and (4) are shown analogously.

Backward implication: Let $a, b \in A$ satisfy $a^*b = 0$. We need to verify that $\varphi(a)^*\varphi(b) = 0$. Let $\varepsilon > 0$. Using an approximate identity in A, we can choose $c \in A$ satisfying $||b - bc|| < \varepsilon$. Using that $b^*a = 0$ and applying the assumption at the first step, we get

$$
\varphi(a)^{*}\varphi(bc) = \varphi(b^{*}a)^{*}\varphi(c) = 0
$$

and therefore

$$
\|\varphi(a)^*\varphi(b)\|=\|\varphi(a)^*\varphi(b)-\varphi(a)^*\varphi(bc)\|\leq \|\varphi(a)^*\|\|\varphi\|\|b-bc\|\leq \varepsilon \|\varphi(a)^*\|\|\varphi\|.
$$

Since $\varepsilon > 0$ was arbitrary, we get $\varphi(a)^{*}\varphi(b) = 0$, as desired.

Forward implication: We assume that φ preserves range-orthogonality. Given $a, b, c \in A$, we need to show that

$$
\varphi(b^*a)^*\varphi(c) = \varphi(a)^*\varphi(bc).
$$

Using linearity of φ , we may assume that b is self-adjoint and satisfies $||b|| < 1$.

For $k \geq 2$, let $f_k: \mathbb{R} \to [0,1]$ be the continuous function that takes the value 0 on $(-\infty, \frac{1}{k}] \cup [1 + \frac{1}{k}, \infty)$, that takes the value 1 on $[\frac{2}{k}, 1]$, and that is affine on $[\frac{1}{k}, \frac{2}{k}]$ and on $[1, 1 + \frac{1}{k}]$. Let $m \ge 1$. For each $j \in \{-m, \ldots, m-1\}$ and $k \ge 2$, we define $f_{j,k} : \mathbb{R} \to [0,1]$ by

$$
f_{j,k}(t) := f_k(mt - j)
$$

for $t \in \mathbb{R}$. Note that the support of $f_{j,k}$ is $\left(\frac{j}{m} + \frac{1}{mk}, \frac{j+1}{m} + \frac{1}{mk}\right)$. In particular, $f_{j,k}$ and $f_{j',k'}$ are orthogonal if $k \neq k'$. Moreover, the sequence $(f_{j,k})_k$ converges pointwise to the characteristic function of $(\frac{j}{m}, \frac{j+1}{m}]$.

Let $\Phi := \varphi^{**} : A^{**} \to B^{**}$ denote the bitranspose of φ . In A^{**} , the sequence $(f_{j,k}(b))_k$ converges weak* to a projection e_j .

Claim 1: Let $j > j'$. Then $\Phi(e_j^*a)^* \Phi(e_{j'}c) = 0$. To prove the claim, let $k \geq 2$. Given k' with $k \leq k'$, we have $(f_{j,k}(b)^*a)^*(f_{j',k'}(b)c) = 0$ since $f_{j,k}f_{j',k'} = 0$, and thus

$$
\varphi(f_{j,k}(b)^*a)^*\varphi(f_{j',k'}(b)c) = 0.
$$

Using that Φ is weak*-continuous, and that multiplication in B^{**} is separately weak*-continuous, we get

$$
\varphi(f_{j,k}(b)^*a)^*\Phi(e_{j'}c)=\operatorname{wk}^*\text{-}\lim_{k'}\varphi(f_{j,k}(b)^*a)^*\varphi(f_{j',k'}(b)c)=0.
$$

Since this holds for every $k \geq 2$, we get

$$
\Phi(e_j^*a)^*\Phi(e_{j'}c) = \text{wk*-}\lim_k \Phi(f_{j,k}(b)^*a)^*\varphi(e_{j'}c) = 0.
$$

This proves the claim.

Claim 2: Let $j < j'$. Then $\Phi(e_j^*a)^* \Phi(e_{j'}c) = 0$. The proof is analogous to that of the previous claim. For fixed $k' \geq 2$, and $k \geq k'$, we have $(f_{j,k}(b)^*a)^*(f_{j',k'}(b)c)$ 0, and thus $\varphi(f_{j,k}(b)^*a)^*\varphi(f_{j',k'}(b)c) = 0$. We then get

$$
\Phi(e_j^*a)^*\varphi(f_{j',k'}(b)c) = \text{wk*-}\lim_{k'}\varphi(f_{j,k}(b)^*a)^*\varphi(f_{j',k'}(b)c) = 0.
$$

Since this holds for every $k' \geq 2$, we get

$$
\Phi(e_j^*a)^*\Phi(e_{j'}c) = \operatorname{wk}^*\text{-}\lim_k \varphi(f_{j,k}(b)^*a)^*\Phi(e_{j'}c) = 0.
$$

This proves the claim.

Using that b is selfadjoint and $||b|| < 1$, we have

$$
\left\|b - \sum_{j=-m}^{m-1} \frac{j}{m} e_j\right\| \le \frac{1}{m}.
$$

Moreover, we have $\sum_j e_j = 1$ and therefore $a = \sum_j e_j^* a$ and $c = \sum_j e_j c$. Set $K := ||\varphi||^2 ||a|| ||c||.$ We write $x \approx_{\varepsilon} y$ to mean $||x - y|| \leq \varepsilon$. Using Claims 1 and 2 at the second and third steps, we get

$$
\varphi(b^*a)^*\varphi(c) \approx_{\frac{K}{m}} \Phi\left(\sum_{j=-m}^{m-1} \frac{j}{m}e_j^*a\right)^*\Phi\left(\sum_{j=-m}^{m-1} e_jc\right)
$$

$$
= \sum_{j=-m}^{m-1} \frac{j}{m}\Phi(e_j^*a)^*\Phi(e_jc)
$$

$$
= \Phi\left(\sum_{j=-m}^{m-1} e_j^*a\right)^*\Phi\left(\sum_{j=-m}^{m-1} \frac{j}{m}e_jc\right)
$$

$$
\approx_{\frac{K}{m}} \varphi(a)^*\varphi(bc).
$$

Since this holds for every $m \geq 1$, we get the desired equality.

Peralta showed in [\[Per15,](#page-16-7) Proposition 3.7] that if $\varphi: A \to B$ preserves zeroproducts, then so does the restriction of the bitranspose φ^{**} : $A^{**} \to B^{**}$ to the multiplier algebra $M(A)$. We generalize this by showing that φ^{**} preserves in fact all zero-products in A^{**} . We obtain similar results for range-orthogonality, domainorthogonality and zero-TRO-products. Using the structure result for orthogonalitypreserving maps from $[BFPG^{+}08]$, we show that orthogonality-preservation also passes to bitransposes.

Proposition 2.3. Let $\varphi: A \to B$ be a bounded, linear map between C^* -algebras. *If* φ *preserves zero-products (range-orthogonality, domain-orthogonality, zero-TROproducts, orthogonality), then so does the bitranspose* φ^{**} : $A^{**} \to B^{**}$.

Proof. Set $\Phi = \varphi^{**} : A^{**} \to B^{**}$. We first show the result for the case that φ preserves zero-products. By [Theorem 2.2\(](#page-2-0)1), we have $\varphi(ab)\varphi(c) = \varphi(a)\varphi(bc)$ for every $a, b, c \in A$. For $a, b, c \in A^{**}$, we denote by $E(a, b, c)$ the (potentially false) identity

$$
E(a, b, c) \qquad \qquad \Phi(ab)\Phi(c) = \Phi(a)\Phi(bc).
$$

By assumption, $E(a, b, c)$ holds whenever $a, b, c \in A$. In three steps, we will show that it holds for all $a, b, c \in A^{**}$.

Let $a \in A^{**}$ and $b, c \in A$ be given. Choose a net $(a_{\lambda})_{\lambda}$ in A that converges weak* to a, and note that $E(a_\lambda, b, c)$ is true. Using at the first step that Φ is weak*continuous and that the multiplication operations in A^{**} and B^{**} are separately weak*-continuous, and using $E(a_\lambda, b, c)$ at the second step, we get

$$
\Phi(ab)\Phi(c) = \mathrm{wk}^*\text{-}\lim_{\lambda} \Phi(a_\lambda b)\Phi(c) = \mathrm{wk}^*\text{-}\lim_{\lambda} \Phi(a_\lambda)\Phi(bc) = \Phi(a)\Phi(bc),
$$

so $E(a, b, c)$ holds whenever $a \in A^{**}$ and $b, c \in A$.

Given $a, b \in A^{**}$ and $c \in A$, we choose as above a net $(b_{\lambda})_{\lambda}$ in A converging weak^{*} to b. By the previous paragraph, $E(a, b_\lambda, c)$ is true for all λ . Using weak^{*}continuity of Φ and the fact that the multiplication operations in A^{**} and B^{**} are separately weak*-continuous, we deduce that $(E(a, b, c))$ $(E(a, b, c))$ $(E(a, b, c))$ holds for the triple (a, b, c) . The general case for arbitrary $a, b, c \in A^{**}$ is proved similarly. We deduce that Φ preserves zero-products.

Using the characterizations of maps preserving range-orthogonality, domainorthogonality or zero-TRO-products from [Theorem 2.2,](#page-2-0) an analogous argument shows that Φ preserves range-orthogonality, domain-orthogonality, or zero-TROproducts, whenever φ does.

Lastly, assume that φ preserves orthogonality. Recall that the triple product in a C^* -algebra is defined as $\{a, b, c\} = \frac{1}{2}(ab^*c + cb^*a)$, and that a bounded linear map θ between C^{*}-algebras is a *triple homomorphism* if $\theta({a, b, c}) = {\theta(a), \theta(b), \theta(c)}$ for all a, b, c in the domain of θ . Equivalently, θ satisfies

$$
\theta(ab^*c + cb^*a) = \theta(a)\theta(b)^*\theta(c) + \theta(c)\theta(b)^*\theta(a)
$$

for all a, b, c . Using an argument similar to the one above, it follows that the bitranspose θ^{**} is a triple homomorphism as well.

By [\[BFPG](#page-16-5)⁺08, Lemma 1], two elements a, b in a C^* -algebra are orthogonal if and only if $\{a, a, b\} = 0$. It follows that every triple homomorphism preserves orthogonality.

Set $h := \Phi(1) \in B^{**}$. Let $h = v|h|$ be the polar decomposition of h. The partial isometry v is also called the *range tripotent* of h, and it is denoted by $r(h)$ in $[BFPG⁺08]$. By $[BFPG⁺08]$, Theorem 17], there exists a triple homomorphism $\pi: A \to B^{**}$ such that

$$
\varphi(a)=hv^*\pi(a)=\pi(a)v^*h
$$

for all $a \in A$. Set $\Pi := \pi^{**} : A^{**} \to B^{**}$, which is readily checked to be a triple homomorphism It follows that

$$
\Phi(a) = hv^* \Pi(a) = \Pi(a)v^*h
$$

for all $a \in A^{**}$.

To show that Φ preserves orthogonality, let $a, b \in A^{**}$ be orthogonal, that is, $ab^* = a^*b = 0$. Since Π is a triple homomorphism and therefore preserves orthogonality, we get $\Pi(a)\Pi(b)^* = \Pi(a)^*\Pi(b) = 0$. Using this at the last steps, we get

$$
\Phi(a)\Phi(b)^* = (hv^* \Pi(a))(hv^* \Pi(b))^* = hv^* \Pi(a)\Pi(b)^* vh^* = 0
$$

and

$$
\Phi(a)^*\Phi(b) = (\Pi(a)v^*h)^*(\Pi(b)v^*h) = h^*v\Pi(a)^*\Pi(b)v^*h = 0,
$$

as desired. \Box

In the next result, we use \otimes to denote the minimal tensor product of C^* -algebras. We note that [Proposition 2.4](#page-5-0) also holds for maximal tensor products, with essentially the same proof.

Proposition 2.4. Let $\varphi: A \to B$ and $\psi: C \to D$ be bounded, linear maps be t ween C^* -algebras. If φ and ψ preserve zero-products (range-orthogonality, domain*orthogonality, zero-TRO-products), then so does the tensor product map* $\varphi \otimes \psi$: A⊗ $C \rightarrow B \otimes D$.

Proof. We prove the results for zero-product preserving maps. The statements for range-orthogonality and domain-orthogonality preserving maps are shown analogously.

Assume that φ and ψ preserve zero-products and set $\alpha := \varphi \otimes \psi : A \odot B \to C \odot D$. Using that φ and ψ satisfy the formula from [Theorem 2.2\(](#page-2-0)1), we verify that α satisfies the formula as well. Let

$$
a = \sum_j u_j \otimes v_j
$$
, $b = \sum_k w_k \otimes x_k$, and $c = \sum_l y_l \otimes z_l$

be finite sums of simple tensors in $A \otimes B$. Then

$$
\alpha(ab)\alpha(c) = \left(\varphi \otimes \psi\Big(\sum_{j,k} u_j w_k \otimes v_j x_k\Big)\right) \left(\varphi \otimes \psi\Big(\sum_l y_l \otimes z_l\Big)\right)
$$

=
$$
\sum_{j,k,l} \varphi(u_j w_k) \varphi(y_l) \otimes \psi(v_j x_k) \psi(z_l)
$$

=
$$
\sum_{j,k,l} \varphi(u_j) \varphi(w_k y_l) \otimes \psi(v_j) \psi(x_k z_l) = \alpha(a)\alpha(bc).
$$

Using that finite sums of simple tensors are dense in $A \otimes B$, and that α is continuous, it follows that $\alpha(ab)\alpha(c) = \alpha(a)\alpha(bc)$ for all $a, b, c \in A \otimes B$. Applying [Theorem 2.2\(](#page-2-0)1), it follows that α preserves zero-products.

The next result shows that every zero-product preserving map is automatically 'completely zero-product preserving' (and similarly for range-orthogonality preserving, domain-orthogonality preserving and zero-TRO-product preserving). For orthogonality-preserving maps, this is not the case: the transpose map on $M_2(\mathbb{C})$ is orthogonality-preserving, but its amplifications are not.

Corollary 2.5. Let $\varphi: A \to B$ be a bounded, linear map between C^* -algebras. *If* φ preserves zero-products (range-orthogonality, domain-orthogonality, zero-TRO*product*), then so does the amplification $\varphi^{(n)}$: $M_n(A) \to M_n(B)$ for every $n \geq 1$.

Proof. This follows by taking $C = D = M_n$ and $\psi = id_{M_n}$ in [Proposition 2.4.](#page-5-0) \Box

Some of the statements in this section do not involve the adjoint operation in a C^* -algebra, and therefore make sense in more general settings. It would thus be interesting to find other classes of Banach algebras for which part (1) of [Theorem 2.2](#page-2-0) or [Corollary 2.5](#page-6-1) are true. For example, one could explore these questions in the context of L^p -operator algebras [\[Gar21\]](#page-16-8), or at least for the well-behaved class arising from groups as in [\[GT15,](#page-16-9) [GT19\]](#page-16-10).

3. Constructing the support ∗-homomorphisms

Throughout this section, we assume that $\varphi: A \to B$ is a bounded, linear map between C^* -algebras that preserves zero-products, range-orthogonality, and also domain-orthogonality. By [Theorem 2.2,](#page-2-0) for all $a, b, c \in A$ we have

(3.1)
$$
\varphi(ab)\varphi(c) = \varphi(a)\varphi(bc);
$$

(3.2)
$$
\varphi(b^*a)^*\varphi(c) = \varphi(a)^*\varphi(bc);
$$

(3.3)
$$
\varphi(ab)\varphi(c)^* = \varphi(a)\varphi(cb^*)^*.
$$

For later use, we observe that the *adjoint* φ^* : A \rightarrow B of φ , defined by $\varphi^*(a)$ = $\varphi(a^*)^*$ for all $a \in A$, also preserves zero-products, range-orthogonality, and also domain-orthogonality. We fix a representation $B \subseteq \mathcal{B}(H)$ of B on some Hilbert space. Our goal is to construct a $*$ -homomorphism $\pi_{\varphi}: A \to B'' \subseteq \mathcal{B}(H)$ such that

$$
\pi_{\varphi}(a)\varphi(b) = \varphi(ab) = \varphi(a)\pi_{\varphi}(b)
$$

for all $a, b \in A$. Using this, we will show in [Theorem 4.2](#page-13-0) that φ is a weighted ∗-homomorphism in the sense of [Definition A.](#page-0-0)

We begin the construction by setting

$$
H_0 := \text{span}\left(\left\{\varphi(a)\xi : a \in A, \xi \in H\right\} \cup \left\{\varphi(a)^* \xi : a \in A, \xi \in H\right\}\right).
$$

Given $a \in A$, we first show that there exists a unique operator $\pi_{\varphi}(a) \in \mathcal{B}(H)$ such that

(3.4) $\pi_{\varphi}(a)\varphi(b)\xi = \varphi(ab)\xi$, and $\pi_{\varphi}(a)\varphi(c)^{*}\eta = \varphi(ca^{*})^{*}\eta$

for all $b, c \in A$ and $\xi, \eta \in H$, and such that $\pi_{\varphi}(a) \zeta = 0$ for all $\zeta \in H_0^{\perp}$.

Lemma 3.1. *Let J and K be finite index sets, let* $b_j \in A$ *and* $\xi_j \in H$ *for* $j \in J$ *, and let* $c_k \in A$ *and* $\eta_k \in H$ *for* $k \in K$ *. Then*

$$
\left\| \sum_{j \in J} \varphi(ab_j) \xi_j + \sum_{k \in K} \varphi(c_k a^*)^* \eta_k \right\| \leq \|a\| \left\| \sum_{j \in J} \varphi(b_j) \xi_j + \sum_{k \in K} \varphi(c_k)^* \eta_k \right\|
$$

for all $a \in A$ *.*

Proof. We first establish the following **Claim 1:** *Given* $a \in A$ *, we have*

$$
\Big\|\sum_{j}\varphi(ab_j)\xi_j + \sum_{k}\varphi(c_ka^*)^*\eta_k\Big\|^2
$$

$$
\leq \Big\|\sum_{j}\varphi(a^*ab_j)\xi_j + \sum_{k}\varphi(c_ka^*a)^*\eta_k\Big\|\Big\|\sum_{j}\varphi(b_j)\xi_j + \sum_{k}\varphi(c_k)^*\eta_k\Big\|.
$$

To prove the claim, let $j \in J$ and $k' \in K$. Then

$$
\langle \varphi(ab_j)\xi_j, \varphi(c_{k'}a^*)^*\eta_{k'} \rangle = \langle \varphi(c_{k'}a^*)\varphi(ab_j)\xi_j, \eta_{k'} \rangle
$$

$$
\stackrel{(3.1)}{=} \langle \varphi(c_{k'})\varphi(a^*ab_j)\xi_j, \eta_{k'} \rangle
$$

$$
= \langle \varphi(a^*ab_j)\xi_j, \varphi(c_{k'})^*\eta_{k'} \rangle.
$$

Similarly, for all $j, j' \in J$ we obtain

$$
\langle \varphi(ab_j)\xi_j, \varphi(ab_{j'})\xi_{j'} \rangle = \langle \varphi(ab_{j'})^* \varphi(ab_j)\xi_j, \xi_{j'} \rangle
$$

$$
\stackrel{(3.2)}{=} \langle \varphi(b_{j'})^* \varphi(a^*ab_j)\xi_j, \xi_{j'} \rangle
$$

$$
= \langle \varphi(a^*ab_j)\xi_j, \varphi(b_{j'})\xi_{j'} \rangle.
$$

Also, for all $k, k' \in K$ we have

$$
\langle \varphi(c_k a^*)^* \eta_k, \varphi(c_{k'} a^*)^* \eta_{k'} \rangle = \langle \varphi(c_{k'} a^*) \varphi(c_k a^*)^* \eta_k, \eta_{k'} \rangle
$$

$$
\stackrel{(3.3)}{=} \langle \varphi(c_{k'}) \varphi(c_k a^* a)^* \eta_k, \eta_{k'} \rangle
$$

$$
= \langle \varphi(c_k a^* a)^* \eta_k, \varphi(c_{k'})^* \eta_{k'} \rangle.
$$

Further, for each $k \in K$ and $j' \in J$ we get

$$
\big\langle \varphi(c_k a^*)^* \eta_k, \varphi(ab_{j'}) \xi_{j'} \big\rangle = \big\langle \varphi(c_k a^* a)^* \eta_k, \varphi(b_{j'}) \xi_{j'} \big\rangle.
$$

Using all of these equalities at the third step, and using the Cauchy-Schwarz inequality at the last step, we get

$$
\left\| \sum_{j} \varphi(ab_{j})\xi_{j} + \sum_{k} \varphi(c_{k}a^{*})^{*} \eta_{k} \right\|^{2}
$$
\n
$$
= \left\langle \sum_{j} \varphi(ab_{j})\xi_{j} + \sum_{k} \varphi(c_{k}a^{*})^{*} \eta_{k}, \sum_{j'} \varphi(ab_{j'})\xi_{j'} + \sum_{k'} \varphi(c_{k'}a^{*})^{*} \eta_{k'} \right\rangle
$$
\n
$$
= \sum_{j,j'} \left\langle \varphi(ab_{j})\xi_{j}, \varphi(ab_{j'})\xi_{j'} \right\rangle + \sum_{j,k'} \left\langle \varphi(ab_{j})\xi_{j}, \varphi(c_{k'}a^{*})^{*} \eta_{k'} \right\rangle
$$
\n
$$
+ \sum_{k,j'} \left\langle \varphi(c_{k}a^{*})^{*} \eta_{k}, \varphi(ab_{j'})\xi_{j'} \right\rangle + \sum_{k,k'} \left\langle \varphi(c_{k}a^{*})^{*} \eta_{k}, \varphi(c_{k'}a^{*})^{*} \eta_{k'} \right\rangle
$$
\n
$$
= \sum_{j,j'} \left\langle \varphi(a^{*}ab_{j})\xi_{j}, \varphi(b_{j'})\xi_{j'} \right\rangle + \sum_{j,k'} \left\langle \varphi(a^{*}ab_{j})\xi_{j}, \varphi(c_{k'})^{*} \eta_{k'} \right\rangle
$$
\n
$$
+ \sum_{k,j'} \left\langle \varphi(c_{k}a^{*}a)^{*} \eta_{k}, \varphi(b_{j'})\xi_{j'} \right\rangle + \sum_{k,k'} \left\langle \varphi(c_{k}a^{*}a)^{*} \eta_{k}, \varphi(c_{k'})^{*} \eta_{k'} \right\rangle
$$
\n
$$
= \left\langle \sum_{j} \varphi(a^{*}ab_{j})\xi_{j} + \sum_{k} \varphi(c_{k}a^{*}a)^{*} \eta_{k}, \sum_{j'} \varphi(b_{j'})\xi_{j'} + \sum_{k'} \varphi(c_{k'})^{*} \eta_{k'} \right\rangle
$$
\n
$$
\leq \left\| \sum_{j} \varphi(a^{*}ab_{j})\xi_{j} + \sum_{k} \varphi(c_{k}a^{*}a)^{*} \eta_{k} \right\| \left\| \sum_{j} \varphi(b_{j})\xi_{j} + \sum
$$

This proves the claim.

Fix $a \in A$ for the rest of the proof. We set

$$
C = \Big\|\sum_{j} \varphi(ab_j)\xi_j + \sum_{k} \varphi(c_k a^*)^* \eta_k \Big\| \text{ and } D = \Big\|\sum_{j} \varphi(b_j)\xi_j + \sum_{k} \varphi(c_k)^* \eta_k \Big\|.
$$

For $n \in \mathbb{N}$, we denote by $I(n)$ the (potentially false) expression

$$
I(n) \t C2 \leq \Big\|\sum_{j} \varphi((a^*a)^{2^n}b_j)\xi_j + \sum_{k} \varphi(c_k a^*a)^* \eta_k \Big\|^{\frac{1}{2^n}} D^{2-\frac{1}{2^n}}.
$$

Claim 2: $I(n)$ *is true for all* $n \in \mathbb{N}$. We will prove the claim by induction. Applying Claim 1, we get

$$
C^{2} = \left\| \sum_{j} \varphi(ab_{j})\xi_{j} + \sum_{k} \varphi(c_{k}a^{*})^{*} \eta_{k} \right\|^{2}
$$

\n
$$
\leq \left\| \sum_{j} \varphi(a^{*}ab_{j})\xi_{j} + \sum_{k} \varphi(c_{k}a^{*}a)^{*} \eta_{k} \right\| \left\| \sum_{j} \varphi(b_{j})\xi_{j} + \sum_{k} \varphi(c_{k})^{*} \eta_{k} \right\|
$$

\n
$$
= \left\| \sum_{j} \varphi(a^{*}ab_{j})\xi_{j} + \sum_{k} \varphi(c_{k}a^{*}a)^{*} \eta_{k} \right\|^{2} D^{2-\frac{1}{20}}.
$$

In other words, $I(0)$ holds. For the induction step, assume that $I(n)$ is true for some $n \geq 0$. Applying Claim 1 with the self-adjoint element $(a^*a)^{2^n}$ in place of a, we obtain

$$
\left\| \sum_{j} \varphi((a^*a)^{2^n} b_j) \xi_j + \sum_{k} \varphi(c_k(a^*a)^{2^n})^* \eta_k \right\|
$$

$$
\leq \left\| \sum_{j} \varphi((a^*a)^{2^{n+1}} b_j) \xi_j + \sum_{k} \varphi(c_k(a^*a)^{2^{n+1}})^* \eta_k \right\|^{\frac{1}{2}} D^{\frac{1}{2}}.
$$

Using this at the second step, we obtain

$$
C^{2} \stackrel{I(n)}{\leq} \left\| \sum_{j} \varphi((a^{*}a)^{2^{n}}b_{j})\xi_{j} + \sum_{k} \varphi(c_{k}(a^{*}a)^{2^{n}})^{*} \eta_{k} \right\|^{2} D^{2-\frac{1}{2^{n}}}
$$

$$
\leq \left\| \sum_{j} \varphi((a^{*}a)^{2^{n+1}}b_{j})\xi_{j} + \sum_{k} \varphi(c_{k}(a^{*}a)^{2^{n+1}})^{*} \eta_{k} \right\|^{2} D^{2-\frac{1}{2^{n}}}
$$

which shows that $I(n + 1)$ is true. This completes the induction and proves the claim.

Now, for every $n \geq 0$, we get

$$
C^2 \stackrel{I(n)}{\leq} \left\| \sum_j \varphi((a^*a)^{2^n} b_j) \xi_j + \sum_k \varphi(c_k a^* a)^* \eta_k \right\|^{1 \over 2^n} D^{2 - \frac{1}{2^n}} \n\leq \left(\sum_j \|\varphi\| \|\xi_j\| + \sum_k \|\varphi\| \|\eta_k\| \right)^{1 \over 2^n} \| (a^* a)^{2^n} \|^{1 \over 2^n} D^{2 - \frac{1}{2^n}}.
$$

Using that $\|(a^*a)^{2^n}\|^{\frac{1}{2^n}} = \|a\|^2$, and taking to the limit as $n \to \infty$, we deduce that $C^2 \le ||a||^2 D^2$, and so $C \le ||a||D$, as desired.

Given $a \in A$, it follows from [Lemma 3.1](#page-7-0) that the map $H_0 \to H_0$, given by

$$
\sum_{j} \varphi(b_j) \xi_j + \sum_{k} \varphi(c_k)^* \eta_k \mapsto \sum_{j} \varphi(ab_j) \xi_j + \sum_{k} \varphi(c_k a^*)^* \eta_k
$$

is well-defined, bounded and linear, and therefore extends to a bounded, linear map $\pi_{\varphi}^{(0)}(a): \overline{H_0} \to \overline{H_0}$ satisfying $\|\pi_{\varphi}^{(0)}(a)\| \leq \|a\|$ for all $a \in A$. We define $\pi_{\varphi}(a) \in \mathcal{B}(H)$ by $\pi_{\varphi}(a)\xi := \pi_{\varphi}^{(0)}(a)\xi$ for $\xi \in \overline{H_0}$ and $\pi_{\varphi}(a)\eta := 0$ for $\eta \in \overline{H_0}^{\perp}$. We note that π_{φ} is determined by

(3.5)
$$
\pi_{\varphi}(a)\varphi(b)\xi = \varphi(ab)\xi \text{ and } \pi_{\varphi}(a)\varphi(b)^{*}\xi = \varphi(ba^{*})^{*}\xi
$$

for all $a, b \in A$ and all $\xi \in H$. Moreover, one readily checks that $\pi_{\varphi} = \pi_{\varphi^*}$.

Proposition 3.2. *Let* A and B be C^* -algebras with $B \subseteq \mathcal{B}(H)$, and let $\varphi: A \to B$ *be a bounded, linear map between* C ∗ *-algebras that preserves zero-products, rangeorthogonality, and also domain-orthogonality. Denote by* π_{φ} : $A \to \mathcal{B}(H)$ *the canonical bounded, linear map defined in the preceding comments. Then:*

- *(1)* The map π_{φ} *is a* ∗*-homomorphism.*
- *(2)* The image of π_{φ} is contained in $B'' \subseteq \mathcal{B}(H)$.
- *(3)* For all $a, b \in A$, we have

$$
\pi_{\varphi}(a)\varphi(b) = \varphi(ab) = \varphi(a)\pi_{\varphi}(b) \quad and \quad \pi_{\varphi}(a)\varphi(b)^{*} = \varphi(ba^{*})^{*} = \varphi(a^{*})^{*}\pi_{\varphi}(b^{*}).
$$

Proof. (1) We first show that π_{φ} , which we will abbreviate to π throughout in the proof of this proposition, is linear and multiplicative. For a_1, a_2 and $\lambda \in \mathbb{C}$, we have

$$
\pi(a_1 + \lambda a_2)\varphi(b)\xi = \varphi((a_1 + \lambda a_2)b)\xi
$$

= $\varphi(a_1b)\xi + \lambda\varphi(a_2b)\xi = (\pi(a_1) + \lambda\pi(a_2))\varphi(b)\xi$

and

$$
\pi(a_1)\pi(a_2)\varphi(b)\xi = \pi(a_1)\varphi(a_2b)\xi = \varphi(a_1a_2b)\xi = \pi(a_1a_2)\varphi(b)\xi
$$

for all $b \in A$ and $\xi \in H$. Similarly, we have

$$
\pi(a_1 + \lambda a_2)\varphi(c)^*\eta = \varphi(c(a_1 + \lambda a_2)^*)^*\eta
$$

= $\varphi(ca_1^*)^*\eta + \lambda\varphi(ca_2^*)^*\eta = (\pi(a_1) + \lambda\pi(a_2))\varphi(c)^*\eta$

and

$$
\pi(a_1)\pi(a_2)\varphi(c)^*\eta = \pi(a_1)\varphi(ca_2^*)^*\eta = \varphi(ba_2^*a_1^*)^*\eta = \pi(a_1a_2)\varphi(c)^*\eta
$$

for all $c \in A$ and $\eta \in H$. Using linearity and continuity of $\pi(a_1 + \lambda a_2)$ and $\pi(a_1) + \lambda \pi(a_2)$, it follows that these operators agree on $\overline{H_0}$. Since both operators vanish on $\overline{H_0}^{\perp}$, it follows that $\pi(a_1 + \lambda a_2) = \pi(a_1) + \lambda \pi(a_2)$. Similarly, we obtain $\pi(a_1)\pi(a_2) = \pi(a_1a_2)$. Thus, π is linear and multiplicative.

To show that π preserves adjoints, let $a \in A$. Given $b, b' \in A$ and $\xi, \xi' \in H$, using [\(3.2\)](#page-6-3) at the fourth step, we get

$$
\langle \pi(a)^* \varphi(b)\xi, \varphi(b')\xi' \rangle = \langle \varphi(b)\xi, \pi(a)\varphi(b')\xi' \rangle = \langle \varphi(b)\xi, \varphi(ab')\xi' \rangle
$$

=
$$
\langle \varphi(ab')^* \varphi(b)\xi, \xi' \rangle = \langle \varphi(b')^* \varphi(a^*b)\xi, \xi' \rangle
$$

=
$$
\langle \varphi(a^*b)\xi, \varphi(b')\xi' \rangle = \langle \pi(a^*) \varphi(b)\xi, \varphi(b')\xi' \rangle.
$$

Similarly, given $c, c' \in A$ and $\eta, \eta' \in H$, using [\(3.3\)](#page-6-4) at the fourth step, we get

$$
\langle \pi(a)^* \varphi(c)^* \eta, \varphi(c')^* \eta' \rangle = \langle \varphi(c)^* \eta, \pi(a) \varphi(c')^* \eta' \rangle = \langle \varphi(c)^* \eta, \varphi(c'a^*)^* \eta' \rangle
$$

\n
$$
= \langle \varphi(c'a^*) \varphi(c)^* \eta, \eta' \rangle = \langle \varphi(c') \varphi(ca)^* \eta, \eta' \rangle
$$

\n
$$
= \langle \varphi(ca)^* \eta, \varphi(c')^* \eta' \rangle = \langle \pi(a^*) \varphi(c)^* \eta, \varphi(c')^* \eta' \rangle.
$$

\n
$$
= \langle \varphi(ca)^* \eta, \varphi(c')^* \eta' \rangle = \langle \pi(a^*) \varphi(c)^* \eta, \varphi(c')^* \eta' \rangle.
$$

Further, given
$$
b, c \in A
$$
 and $\xi, \eta \in H$, using (3.1) at the fourth step, we get

$$
\langle \pi(a)^* \varphi(b)\xi, \varphi(c)^* \eta \rangle = \langle \varphi(b)\xi, \pi(a)\varphi(c)^* \eta \rangle = \langle \varphi(b)\xi, \varphi(ca^*)^* \eta \rangle
$$

$$
= \langle \varphi(ca^*) \varphi(b)\xi, \eta \rangle = \langle \varphi(c)^* \varphi(a^*b)\xi, \eta \rangle
$$

$$
= \langle \varphi(a^*b)\xi, \varphi(c)^* \eta \rangle = \langle \pi(a^*) \varphi(b)\xi, \varphi(c)^* \eta \rangle.
$$

and analogously

$$
\langle \pi(a)^* \varphi(c)^* \eta, \varphi(b)\xi \rangle = \langle \pi(a^*) \varphi(c)^* \eta, \varphi(b)\xi \rangle.
$$

Using linearity and continuity of $\pi(a)^*$ and $\pi(a^*)$, we get

$$
\langle \pi(a)^* \alpha, \beta \rangle = \langle \pi(a^*) \alpha, \beta \rangle
$$

for all $\alpha, \beta \in \overline{H_0}$ and consequently $\pi(a)^* = \pi(a^*)$, as desired.

(2) Let $a \in A$, and let $x \in B'$. We need to show that $\pi(a)x = x\pi(a)$. Given $b \in B$ and $\xi \in H$, using at the first and third step that x commutes with $\varphi(b)$ and $\varphi(ab)$, we have

$$
\pi(a)x\varphi(b)\xi = \pi(a)\varphi(b)x\xi = \varphi(ab)x\xi = x\varphi(ab)\xi = x\pi(a)\varphi(b)\xi.
$$

Similarly, given $c \in B$ and $\xi \in H$, using that $\varphi(c)^*$ and $\varphi(ca^*)^*$ belong to B and that they therefore commute with x , we get

$$
\pi(a)x\varphi(c)^{*}\xi = \pi(a)\varphi(c)^{*}x\xi = \varphi(ca^{*})^{*}x\xi = x\varphi(ca^{*})^{*}\xi = x\pi(a)\varphi(c)^{*}\xi.
$$

Using linearity and continuity of $\pi(a)x$ and $x\pi(a)$, it follows that $\pi(a)x\xi = x\pi(a)\xi$ for all $\xi \in \overline{H_0}$.

Using that x commutes with $\varphi(b)$ and $\varphi(c)^*$ for all $b, c \in A$, one readily checks that $x(\overline{H_0}) \subseteq \overline{H_0}$. Similarly, since x^* commutes with $\varphi(b)$ and $\varphi(c)^*$ for all $b, c \in A$, we get $x^*(\overline{H_0}) \subseteq \overline{H_0}$. It follows that x leaves $\overline{H_0}^{\perp}$ invariant, and thus

$$
\pi(a)x\eta = 0 = x\pi(a)\eta
$$

for all $\eta \in \overline{H_0}^{\perp}$. In conclusion, we obtain $\pi(a)x = x\pi(a)$, as desired. (3) Let $a, b \in A$. We have

$$
\pi(a)\varphi(b)\xi\stackrel{(3.5)}{=}\varphi(ab)\xi
$$

for all $\xi \in H$, and therefore $\pi(a)\varphi(b) = \varphi(ab)$.

To check that $\varphi(a)\pi(b) = \varphi(ab)$, we prove that these operators agree both on $\overline{H_0}$ and on its orthogonal complement. Given $c \in A$, we have

(3.6)
$$
\varphi(a)\pi(b)\varphi(c)\xi \stackrel{(3.5)}{=} \varphi(a)\varphi(bc)\xi \stackrel{(3.1)}{=} \varphi(ab)\varphi(c)\xi.
$$

Similarly,

(3.7)
$$
\varphi(a)\pi(b)\varphi(c)^{*}\xi \stackrel{(3.5)}{=} \varphi(a)\varphi(cb^{*})^{*}\xi \stackrel{(3.3)}{=} \varphi(ab)\varphi(c)^{*}\xi.
$$

It follows from [\(3.6\)](#page-11-0) and [\(3.7\)](#page-11-1) that the linear maps $\varphi(a)\pi(b)$ and $\varphi(ab)$ agree on $\overline{H_0}$. Let $\eta \in \overline{H_0}^{\perp}$. Then $\pi(b)\eta = 0$. On the other hand, we have

$$
\langle \varphi(ab)\eta, \varphi(ab)\eta \rangle = \langle \eta, \varphi(ab)^* \varphi(ab)\eta \rangle = 0,
$$

where at the last step we use that $\varphi(ab)^*\varphi(ab)\eta$ belongs to $\overline{H_0}$. This shows that $\varphi(ab)$ and $\varphi(a)\pi(b)$ agree on $\overline{H_0}^{\perp}$, and hence $\varphi(a)\pi(b) = \varphi(ab)$.

The last two equalities are proved analogously. (They also follow by replacing φ with φ^* , which satisfies the same assumptions as φ and has $\pi_{\varphi} = \pi_{\varphi^*}$.

Notation 3.3. Let A and B be C^{*}-algebras and let $\varphi: A \to B \subseteq \mathcal{B}(H)$ be a bounded, linear map that preserves zero-products, range-orthogonality, and also domain-orthogonality. Denote by $\pi_{\varphi} : A \to B''$ the canonical *-homomorphism provided by [Proposition 3.2.](#page-9-1) We denote by $\Phi, \Pi_{\varphi} : A^{**} \to \mathcal{B}(H)$ the (unique) extensions of $\varphi, \pi_{\varphi} : A \to \mathcal{B}(H)$ to weak*-continuous, bounded linear maps, which are explicitly constructed as follows. Let $S_1(H)$ denote the space of trace-class of operators on H , which we naturally identify with the (unique) isometric predual of $\mathcal{B}(H)$. Let $\kappa: S_1(H) \to S_1(H)^{**}$ be the natural inclusion of $S_1(H)$ into its bidual. Then the transpose map $\kappa^* \colon \mathcal{B}(H)^{**} \cong S_1(H)^{***} \to S_1(H)^* \cong \mathcal{B}(H)$ is a weak*-continuous ∗-homomorphism, and

$$
\Phi = \kappa^* \circ \varphi^{**} : A^{**} \to \mathcal{B}(H), \text{ and } \Pi_{\varphi} = \kappa^* \circ \pi_{\varphi}^{**} : A^{**} \to \mathcal{B}(H).
$$

By [Proposition 2.3,](#page-4-0) φ^{**} preserves range-orthogonality, domain-orthogonality and zero-products, and hence so does Φ . Similarly, Π_{φ} is a *-homomorphism. Note that the images of Φ and Π_{φ} are contained in $B'' \subseteq \mathcal{B}(H)$.

Proposition 3.4. *Let* A and B *be* C^* -algebras and let $\varphi: A \to B \subseteq \mathcal{B}(H)$ *be a bounded, linear map that preserves zero-products, range-orthogonality, and also domain-orthogonality.* Denote by $\pi_{\varphi} \colon A \to B''$ the canonical *-homomorphism *provided by [Proposition 3.2,](#page-9-1) and let* Φ , Π_{φ} : $A^{**} \to B''$ *the maps from [Notation 3.3.](#page-11-2)*

(1) For all $a, b \in A$, we have

 $\Pi_{\varphi}(a)\Phi(b) = \Phi(ab) = \Phi(a)\Pi_{\varphi}(b)$ *and* $\Pi_{\varphi}(a)\Phi(b)^* = \Phi(ba^*)^* = \Phi(a^*)^*\Pi_{\varphi}(b^*).$

Let C *be the (not necessarily self-adjoint) closed subalgebra of* B *generated by the image of* φ *, and let* $D := C^*(\varphi(A))$ *be the sub-C*^{*}-algebra of B generated by $\varphi(A)$ *.*

- *(2) Let* $a \in M(A) \subseteq A^{**}$. Then $\Phi(a)$ and $\Pi_{\varphi}(a)$ normalize C and D, that *is,* $\Phi(a)x, x\Phi(a), \Pi_{\varphi}(a)x,$ and $x\Pi_{\varphi}(a)$ belong to C (respectively, to D) for *every* $x \in C$ *(respectively,* $x \in D$ *).*
- *(3)* If a *belongs to the center of* $M(A)$ *, then* $\Phi(a)$ *belongs to* C' *.*

Proof. (1) This follows from part (3) of [Proposition 3.2](#page-9-1) by applying the same ar-gument as in the proof of [Proposition 2.3,](#page-4-0) using twice that multiplication on $\mathcal{B}(H)$ is separately weak*-continuous.

 (2) We only show the normalization for D. The proof for C is similar, but easier. Set

$$
G := \{ \varphi(b) \colon b \in A \} \cup \{ \varphi(c)^* \colon c \in A \}.
$$

Then the set of finite linear sums of finite products of elements in G is dense in D.

We first show that $\Pi_{\varphi}(a)$ is a left normalizer of D, that is, $\Pi_{\varphi}(a)D \subseteq D$. Using that $\Pi_{\varphi}(a)$ is a bounded, linear operator, it suffices to show that $\Pi_{\varphi}(a)x \in D$ whenever $x \in G^n$ is a finite product of elements in G. Let $x = gy$ with $g \in G$ and $y \in G^n$, for some $n \in \mathbb{N}$. If $g = \varphi(b)$ for some $b \in A$, then applying part (1) of [Proposition 3.4,](#page-11-3) and using that $ab \in A$, we obtain

$$
\Pi_{\varphi}(a)x = \Pi_{\varphi}(a)gy = \Pi_{\varphi}(a)\Phi(b)y = \Phi(ab)y = \varphi(ab)y \in G^{n+1} \subseteq D.
$$

Similarly, if $g = \varphi(c)^*$ for some $c \in A$, using that $ca^* \in A$, we obtain

$$
\Pi_{\varphi}(a)x = \Pi_{\varphi}(a)gy = \Pi_{\varphi}(a)\Phi(c)^{*}y = \Phi(ca^{*})^{*}y = \varphi(ca^{*})^{*}y \in G^{n+1} \subseteq D.
$$

Analogously, one verifies that $\Pi_{\varphi}(a)$ is a right normalizer of D, that is, we have $D\Pi_{\varphi}(a) \subseteq D.$

Next, we show that $\Phi(a)$ is a left normalizer of D. Again, it suffices to verify that $\Phi(a)x \in D$ for $x = gy$ with $g \in G$ and $y \in G^n$ a product of n elements in G. If $g = \varphi(b)$ for some $b \in A$, then choose $b_1, b_2 \in A$ with $b = b_1b_2$. Using that Φ preserves zero-products and therefore satisfies the formula in [Theorem 2.2\(](#page-2-0)1), and using that $ab_1 \in A$, we obtain

$$
\Phi(a)x = \Phi(a)gy = \Phi(a)\Phi(b_1b_2)y = \Phi(ab_1)\Phi(b_2)y = \varphi(ab_1)\varphi(b_2)y \in G^{n+2} \subseteq D.
$$

Similarly, if $g = \varphi(c)^*$ for some $c \in A$, then choose $c_1, c_2 \in A$ with $c = c_1 c_2$. Using that Φ preserves domain-orthogonality and therefore satisfies the formula in part (3) of [Theorem 2.2,](#page-2-0) and using that $ac_2^* \in A$, we obtain

$$
\Phi(a)x = \Phi(a)gy = \Phi(a)\Phi(c_1c_2)^*y = \Phi(ac_2^*)\Phi(c_1)^*y = \varphi(ac_2^*)\varphi(c_1)^*y \in G^{n+2} \subseteq D.
$$

Analogously, one verifies that $\Phi(a)$ is a left normalizer for D.

(3) Let $a \in M(A)$ be central. Let $b \in A$ and denote by 1_A the unit of $M(A)$. Using that Φ preserves zero-products and therefore satisfies the formula in Theorem $2.2(1)$, we get

$$
\Phi(a)\varphi(b) = \Phi(1_A a)\Phi(b) = \Phi(1_A)\Phi(ab) = \Phi(1_A)\Phi(ba) = \Phi(1_A b)\Phi(a) = \varphi(b)\Phi(a).
$$

We deduce that $\Phi(a)$ commutes with every finite linear combination of products of finitely many elements in $\varphi(A)$. Since such elements are dense in C, we have $\Phi(a) \in C'$.

4. Weighted ∗-homomorphisms

In this section, we prove the main result of the paper [\(Theorem 4.2\)](#page-13-0), where we obtain an intrinsic and algebraic characterization of weighted ∗-homomorphisms between C^{*}-algebras. Our characterization simplifies considerably for self-adjoint and positive maps; see Corollaries [4.4](#page-14-0) and [4.8.](#page-15-0)

Lemma 4.1. Let $\varphi: A \to B$ be a bounded, linear map between C^* -algebras that *preserves range-orthogonality, domain-orthogonality and zero-products. Assume that* $B = C^*(\varphi(A))$ *, that is,* $\varphi(A)$ *is not contained in a proper sub-C*^{*}-algebra *of* B. Let C be the closed subalgebra of B generated by the image of φ . Set $h := \varphi^{**}(1) \in B^{**}.$

Then there exists a canonical **-homomorphism* π_{φ} : $A \to B^{**}$ *such that* $\varphi(a)$ = $h\pi_{\varphi}(a) = \pi_{\varphi}(a)h$ *for all* $a \in A$ *, and such that* $\pi_{\varphi}(a)$ *normalizes both* B *and* C *for every* a ∈ A*. Further,* h *normalizes* B *and* C*, and commutes with every element of* C. In particular, we may view π_{φ} as a ∗-homomorphism $\pi_{\varphi} : A \to M(B) \cap \{h\}'$, *and h belongs* to $M(B) \cap C'$ *.*

Proof. Use [\[Bla06,](#page-16-11) Section III.5.2] to choose a (universal) representation $B \subseteq \mathcal{B}(H)$ such that, with $\kappa^* : \mathcal{B}(H)^{**} \to \mathcal{B}(H)$ denoting the *-homomorphism described in [Notation 3.3,](#page-11-2) the restriction of κ^* to B^{**} is an isomorphism onto B'' . The situation is shown in the following commutative diagram:

By [Proposition 3.2,](#page-9-1) there is a canonical *-homomorphism $\pi_{\varphi} : A \to B''$ such that

$$
\pi_{\varphi}(a)\varphi(b) = \varphi(ab) = \varphi(a)\pi_{\varphi}(b)
$$

for all $a, b \in A$. Let $\Phi = \varphi^{**} : A^{**} \to B^{**} \subseteq \mathcal{B}(H)$ be the unique extension of φ to a weak-^{*} continuous map. After identifying B^{**} with B'' , the map Φ is simply the bitranspose of φ . Hence, applying part (1) of [Proposition 3.4](#page-11-3) at the second steps, we obtain

$$
\varphi(a) = \Phi(1a) = \Phi(1)\pi_{\varphi}(a) = h\pi_{\varphi}(a), \text{ and}
$$

$$
\varphi(a) = \Phi(a1) = \pi_{\varphi}(a)\Phi(1) = \pi_{\varphi}(a)h
$$

for every $a \in A$.

It follows from part (2) of [Proposition 3.4](#page-11-3) that $\pi_{\varphi}(a)$ normalizes both B and C, for each $a \in A$. Moreover, $h = \Phi(1)$ normalizes both B and C by part (3) of [Proposition 3.4,](#page-11-3) and clearly belongs to C' .

The following theorem characterizes weighted ∗-homomorphisms [\(Definition A\)](#page-0-0) in terms of orthogonality-preservation properties. Note that part (2) gives canonical choices for the algebra D and $h \in D$.

Theorem 4.2. *Let* $\varphi: A \to B$ *be a bounded, linear map between* C^* -algebras. Then *the following are equivalent:*

- (1) φ *is a weighted* *-homomorphism, namely: there exist a C^{*}-algebra D with $B \subseteq D$, a **-homomorphism* $\pi: A \rightarrow D$ and $h \in D$ *such that* $\varphi(a) =$ $h\pi(a) = \pi(a)h$ *for all* $a \in A$;
- *(2) there exists a (canonical)* *-*homomorphism* $\pi_{\varphi} : A \to B^{**}$ *such that we have* $\varphi(a) = \varphi^{**}(1)\pi_{\varphi}(a) = \pi_{\varphi}(a)\varphi^{**}(1)$ *for all* $a \in A$;
- (3) φ preserves range-orthogonality, domain-orthogonality and zero-products.

Proof. It is clear that (2) implies (1). To show that (1) implies (3), assume that there exist a C^* -algebra D with $B \subseteq D$, a *-homomorphism $\pi: A \to D$ and $h \in D$ such that $\varphi(a) = h\pi(a) = \pi(a)h$ for all $a \in A$. To verify that φ preserves rangeorthogonality, let $a, b \in A$ satisfy $a^*b = 0$. Then

$$
\varphi(a)^{*}\varphi(b) = (\pi(a)h)^{*}(\pi(b)h) = h^{*}\pi(a)^{*}\pi(b)h = h^{*}\pi(a^{*}b)h = 0.
$$

Similarly, one verifies that φ preserves domain-orthogonality and zero-products.

To show that (3) implies (2), assume that φ preserves range-orthogonality, domain-orthogonality and zero-products. Let $B_0 := C^*(\varphi(A))$ be the sub-C^{*}algebra of B generated by the image of φ . Let $\varphi_0: A \to B_0$ be the corestriction, and set $h_0 := \varphi_0^{**}(1) \in B_0^{**}$. Applying [Lemma 4.1,](#page-13-1) we obtain a *-homomorphism $\pi_0: A \to B_0^{**}$ such that $\varphi_0(a) = h_0 \pi_0(a) = \pi_0(a) h_0$ for all $a \in A$. The inclusion $\iota: B_0 \hookrightarrow B$ induces a natural inclusion $\iota^{**}: B_0^{**} \hookrightarrow B^{**}$ that identifies h_0 with $\varphi^{**}(1)$. Then the *-homomorphism $\pi_{\varphi} := \iota^{**} \circ \pi_0 \colon A \to B^{**}$ has the desired properties.

Remark 4.3. Let us clarify the relationship between our notion of 'weighted ∗ homomorphism' from [Definition A](#page-0-0) and the concept of 'weighted homomorphism' used in the theory of (Banach) algebras.

A map $\varphi: A \to B$ between Banach algebras is said to be a *weighted homomorphism* if there exists a homomorphism $\pi: A \rightarrow B$ and an invertible centralizer W on B such that $\varphi = W\pi$. Here, a *centralizer* on B is a linear map $W: B \to B$ satisfying $W(ab) = aW(b) = W(a)b$ for all $a, b \in B$. We use $\Gamma(B)$ to denote the algebra of centralizers on B.

If B is *faithful* (that is, every element $b \in B$ satisfying $bB = \{0\}$ or $Bb = \{0\}$ is zero), then centralizers correspond to central multipliers: A *multiplier* (also called a *double centralizer*) on B is a pair (L, R) of linear maps $L, R: B \to B$ such that $aL(b) = R(a)b$ for all $a, b \in B$. We obtain a natural map from $\Gamma(B)$ to the multiplier algebra $M(B)$ given by $W \mapsto (W, W)$. If B is faithful, then this map defines an isomorphism between $\Gamma(B)$ and $Z(M(B))$, the center of the multiplier algebra.

Now let $\varphi: A \to B$ be a weighted *-homomorphism between C^* -algebras. Let $B_0 := C^*(\varphi(A))$ be the sub-C^{*}-algebra of B generated by the image of φ , and let C denote the closed subalgebra of B generated by $\varphi(A)$. By [Lemma 4.1,](#page-13-1) there exists a *-homomorphism $\pi: A \to M(B_0)$ and a multiplier $h \in M(B_0) \cap C'$ such that $\varphi(a) = h\pi(a) = \pi(a)h$ for all $a \in A$. However, while h commutes with elements in the image of φ (and as a consequence belongs to C'), it may not commute with the *adjoints* of the elements in $\varphi(A)$, and therefore does not necessarily belong to $Z(M(B_0))$.

If h is normal (which is automatically the case if φ is self-adjoint), then Fuglede's theorem implies that h also commutes with the adjoints of elements in $\varphi(A)$, and consequently belongs to $M(B_0) \cap B'_0$, and therefore to $Z(M(B_0))$. Similarly, if $\varphi(A)$ is a self-adjoint subset of B (which is automatically the case if φ is surjective), then h belongs to $Z(M(B_0))$. In both cases, we see that the map $\varphi: A \to B_0 \subseteq M(B_0)$ is a weighted homomorphism in the algebraic sense.

On the other hand, we may view φ as a map from A to the Banach algebra C. Note that multiplication by h defines a centralizer on C, and that $\pi(a)$ defines a multiplier on C for each $a \in A$. However, C can be very pathological. For example, if $h^2 = 0$, then multiplication by h defines the zero centralizer on C, the product of any two elements in C is zero, C is not faithful, and the canonical map $C \to M(C)$ is the zero map.

Corollary 4.4. Let $\varphi: A \to B$ be a self-adjoint, bounded, linear map between C^* -algebras. Then φ *is a weighted* *-homomorphism if and only if φ preserves *zero-products.*

Proof. Assume that φ preserves zero-products. For $a, b \in A$ with $a^*b = 0$, we have

$$
\varphi(a)^{*}\varphi(b) = \varphi(a^{*})\varphi(b) = 0.
$$

That is, φ preserves range-orthogonality. One similarly shows that φ preserves domain-orthogonality, and the statement thus follows from [Theorem 4.2.](#page-13-0) \Box

The above corollary motivates the following problem.

Problem 4.5. Characterize weighted ∗-homomorphisms with normal weights.

The next result is a generalization to the setting of self-adjoint, zero-product preserving maps of the structure theorem of Winter and Zacharias [\[WZ09,](#page-17-2) Theorem 3.3] for completely positive, order zero maps.

Corollary 4.6. Let $\varphi: A \to B$ be a self-adjoint, bounded, linear map between C^* *algebras that preserves zero-products. Set* $C := C^*(\varphi(A)) \subseteq B$ *. Then* $h := \varphi^{**}(1)$ *belongs to* $M(C) \cap C' \subseteq C^{**} \subseteq B^{**}$, and there exists a canonical *-homomorphism $\pi_{\varphi} \colon A \to M(C) \cap \{h\}'$ such that $\varphi(a) = h \pi_{\varphi}(a)$ for all $a \in A$.

Proof. By [Corollary 4.4,](#page-14-0) φ is a weighted $*$ -homomorphism and thus preserves range-orthogonality, domain-orthogonality and zero-products by [Theorem 4.2.](#page-13-0) Since φ is self-adjoint, it follows that C agrees with the closed subalgebra of B generated by $\varphi(A)$. Now it follows from [Lemma 4.1](#page-13-1) that there exists a exists a ∗-homomorphism $\pi_{\varphi}: A \to M(C) \cap \{h\}'$ such that $\varphi(a) = h \pi_{\varphi}(a)$ for all $a \in A$, and such that h belongs to $M(C) \cap C'$.

We stress the fact that the map π_{φ} that we obtain in the previous corollary is natural. This has the following consequence:

Remark 4.7. Adopt the notation and assumptions of [Corollary 4.6.](#page-15-1) Let G be a topological group, let $\alpha: G \to \text{Aut}(A)$ and $\beta: G \to \text{Aut}(B)$ be continuous actions, and suppose that the map $\varphi: A \to B$ is equivariant. Then h is G-invariant, and there is a canonical continuous action of G on C , and hence there is a (not necessarily continuous) action $\gamma: G \to \text{Aut}(M(C) \cap \{h\}')$. Using naturality of π_{φ} at the first and third steps, and equivariance of φ at the second step, we get

$$
\pi_{\varphi} \circ \alpha_g = \pi_{\varphi \circ \alpha_g} = \pi_{\beta_g \circ \varphi} = \gamma_g \circ \pi_{\varphi}.
$$

In other words, the *-homomorphism $\pi_{\varphi} : A \to M(C) \cap \{h\}'$ is equivariant. Since the action on A is assumed to be continuous, the range of this map is contained in the γ -*continuous part* of $M(C) \cap \{h\}'$, namely

 $M(C)_{\gamma} \cap \{h\}' = \{x \in M(C) : xh = hx \text{ and } g \mapsto \gamma_g(x) \text{ is norm-continuous}\}.$

Recall a linear map $\varphi: A \to B$ between C^{*}-algebras is *positive* if $\varphi(A_+) \subseteq B_+$; it is *n*-positive if the amplification $\varphi^{(n)} = \varphi \otimes id_{M_n} : M_n(A) \to M_n(B)$ is positive; and it is *completely positive* if it is *n*-positive for every $n \in \mathbb{N}$.

Note that a weighted $*$ -homomorphism with weight h is self-adjoint (positive) if and only if h is self-adjoint (positive). It is easy to see that a positively weighted ∗-homomorphism is even completely positive. Hence, we obtain the following characterization of completely positive, order zero maps:

Corollary 4.8. *A map between* C ∗ *-algebras is completely positive, order zero if and only if it is positive and preserves zero-products. In particular, every positive, zero*product preserving map between C^* -algebras is automatically completely positive.

Remark 4.9. Following Sato, [\[Sat21\]](#page-17-4), we say that a positive (but not necessarily completely positive) map between C ∗ -algebras is *order zero* if it preserves zeroproducts of positive elements. By [\[Sat21,](#page-17-4) Corollary 3.7], every 2-positive, order zero map is completely positive. On the other hand, not every positive, order zero map is automatically completely positive, and the transpose map on $M_2(\mathbb{C})$ is a counterexample.

An alternative proof of [\[Sat21,](#page-17-4) Corollary 3.7] can be obtained from [Corollary 4.8](#page-15-0) by noting that every 2-positive, order zero map preserves zero-products. Indeed, we first note that elements a, b in a C^{*}-algebra satisfy $ab = 0$ if and only if $(a^*a)(bb^*) =$ 0. Now, let $\varphi: A \to B$ be a 2-positive, order zero map. Then φ satisfies the Kadison inequality $\varphi(a)^*\varphi(a) \leq ||\varphi||\varphi(a^*a)$ for all $a \in A$. Hence, if $a, b \in A$ satisfy $ab = 0$, then $(a^*a)(bb^*)=0$ and therefore

$$
\varphi(a^*a)\varphi(bb^*) = 0, \quad \varphi(a)^*\varphi(a) \le ||\varphi||\varphi(a^*a), \quad \text{and} \quad \varphi(b)\varphi(b)^* \le ||\varphi||\varphi(bb^*).
$$

This implies $\varphi(a)^*\varphi(a)\varphi(b)\varphi(b)^* = 0$, and thus $\varphi(a)\varphi(b) = 0$, as desired.

Proposition 4.10. *Let* $\varphi: A \to B$ *be a bounded, linear map between* C^* -algebras. *If* φ preserves range-orthogonality, domain-orthogonality and zero-products, then φ *is completely bounded with* $\|\varphi\|_{cb} = \|\varphi\| = \|\varphi^{**}(1)\|$ *.*

Proof. Set $h := \varphi^{**}(1) \in B^{**}$. By [Theorem 4.2,](#page-13-0) there exists a *-homomorphism $\pi: A \to B^{**}$ such that $\varphi(a) = h\pi(a)$ for all $a \in A$. We have

$$
\|\varphi^{**}(1)\| \le \|\varphi^{**}\| = \|\varphi\| \le \|\varphi\|_{cb}.
$$

Given $n \geq 1$, set $h^{(n)} = 1_{M_n} \otimes h \in M_n(B^{**})$, which is the diagonal matrix with diagonal entries all equal to h. It follows that the amplification $\varphi^{(n)}$ satisfies $\varphi^{(n)}(x) = h^{(n)} \pi^{(n)}(x)$ for all $x \in M_n(A)$. Using that $\pi^{(n)}$ is a *-homomorphism, we deduce that

$$
\|\varphi^{(n)}(x)\| = \|h^{(n)}\pi^{(n)}(x)\| \le \|h^{(n)}\| \|\pi^{(n)}(x)\| \le \|h\| \|x\|
$$

and thus $\|\varphi^{(n)}\| \leq \|h\|$. Since this holds for every n, we obtain $\|\varphi\|_{cb} \leq \|h\|$, as desired. \Box

Remark 4.11. In [\[GT22\]](#page-16-12), we show that [Proposition 4.10](#page-16-13) also holds for bounded linear maps that only preserve range-orthogonality or domain-orthogonality. In particular, a bounded, range-orthogonality preserving map is automatically completely bounded. We also show that a range-orthogonality preserving map from a unital C^* -algebra that has no one-dimensional irreducible representations is automatically bounded, and hence completely bounded.

REFERENCES

- [ABEV09] J. ALAMINOS, M. BREŠAR, J. EXTREMERA, and A. R. VILLENA, Maps preserving zero products, Studia Math. 193 (2009), 131–159.
- [Are83] W. ARENDT, Spectral properties of Lamperti operators, Indiana Univ. Math. J. 32 (1983), 199–215.
- [Bla06] B. Blackadar, Operator algebras, Encyclopaedia of Mathematical Sciences 122, Springer-Verlag, Berlin, 2006, Theory of C∗-algebras and von Neumann algebras, Operator Algebras and Non-commutative Geometry, III.
- [BFPG⁺08] M. Burgos, F. J. Fernández-Polo, J. J. Garcés, J. Martínez Moreno, and A. M. PERALTA, Orthogonality preservers in C[∗]-algebras, JB[∗]-algebras and JB[∗]-triples, J. Math. Anal. Appl. 348 (2008), 220–233.
- [CKLW03] M. A. Chebotar, W.-F. Ke, P.-H. Lee, and N.-C. Wong, Mappings preserving zero products, Studia Math. 155 (2003), 77–94.
- [Gar20] J. J. GARCÉS, Complete orthogonality preservers between C^* -algebras, J. Math. Anal. Appl. 483 (2020), 123596, 16.
- [Gar21] E. GARDELLA, A modern look at algebras of operators on L^p -spaces, *Expo. Math.* 39 (2021), 420–453.
- [GT15] E. GARDELLA and H. THIEL, Group algebras acting on L^p -spaces, J. Fourier Anal. Appl. 21 (2015), 1310–1343.
- [GT19] E. GARDELLA and H. THIEL, Representations of p-convolution algebras on L^q -spaces, Trans. Amer. Math. Soc. 371 (2019), 2207–2236.
- [GT22] E. GARDELLA and H. THIEL, Automatic continuity for orthogonality-preserving maps between C^* -algebras, in preparation, 2022.
- [HKW12] I. HIRSHBERG, E. KIRCHBERG, and S. WHITE, Decomposable approximations of nuclear C^* -algebras, Adv. Math. 230 (2012), 1029-1039.
- [KLW04] W.-F. Ke, B.-R. Li, and N.-C. Wong, Zero product preserving maps of operatorvalued functions, Proc. Amer. Math. Soc. 132 (2004), 1979–1985.
- [Per15] A. M. PERALTA, A note on 2-local representations of C[∗]-algebras, Oper. Matrices 9 (2015), 343–358.
- [Sat21] Y. SATO, 2-positive almost order zero maps and decomposition rank, J. Operator Theory 85 (2021), 505–526.
- [WZ09] W. WINTER and J. ZACHARIAS, Completely positive maps of order zero, Münster J. Math. 2 (2009), 311–324.
- [WZ10] W. WINTER and J. ZACHARIAS, The nuclear dimension of C[∗]-algebras, Adv. Math. 224 (2010), 461–498.
- [Wol94] M. WOLFF, Disjointness preserving operators on C^{*}-algebras, Arch. Math. (Basel) 62 (1994), 248–253.
- [Won07] N.-C. WONG, Zero product preservers of C^* -algebras, in Function spaces, Contemp. Math. 435, Amer. Math. Soc., Providence, RI, 2007, pp. 377–380.

Eusebio Gardella, Department of Mathematical Sciences, Chalmers University of Technology and University of Gothenburg, Gothenburg SE-412 96, Sweden.

Email address: gardella@chalmers.se

 $URL:$ www.math.chalmers.se/~gardella

Hannes Thiel, Department of Mathematics, Kiel University, 24118 Kiel, Germany Email address: hannes.thiel@math.uni-kiel.de URL: www.hannesthiel.org