

TORUS ORBIT CLOSURES IN THE FLAG VARIETY

EUNJEONG LEE, MIKIYA MASUDA, AND SEONJEONG PARK

ABSTRACT. The study of torus orbit closures in the (complete) flag variety was initiated by Klyachko and Gelfand–Serganova in the mid-1980s. In this chapter, we present some of the work by Klyachko and Gelfand–Serganova and our recent work on the topology, geometry, and combinatorics of torus orbit closures in the flag variety.

CONTENTS

| | | |
|----------|---|-----------|
| 1 | Introduction | 2 |
| 2 | Torus orbit closures (general) | 4 |
| 2.1 | Torus actions on flag varieties | 5 |
| 2.2 | Moment map | 5 |
| 2.3 | Torus orbit closures and their moment polytopes | 8 |
| 2.4 | Coxeter matroids and Gelfand–Serganova Theorem | 9 |
| 2.5 | Description of the fan of a torus orbit closure | 13 |
| 2.6 | Retractions and metric on finite Coxeter groups | 16 |
| 3 | Generic torus orbit closures in the flag variety | 19 |
| 3.1 | Faces of the permutohedron Π_n | 20 |
| 3.2 | Eulerian polynomial | 21 |
| 3.3 | Klyachko’s result | 23 |
| 3.4 | Relation to Hessenberg varieties | 24 |
| 4 | Generic torus orbit closures in Schubert varieties | 25 |
| 4.1 | Generic torus orbit closures in Schubert varieties | 25 |
| 4.2 | Generalized Eulerian polynomials | 28 |
| 4.3 | Toric Schubert varieties | 31 |
| 4.4 | Schubert varieties of complexity one | 34 |
| 5 | Generic torus orbit closures in Richardson varieties | 36 |
| 5.1 | Richardson varieties and Bruhat interval polytopes | 37 |
| 5.2 | Generic torus orbit closures in Richardson varieties | 37 |
| 5.3 | Toric Bruhat interval polytopes | 38 |
| 5.4 | Conditions on v and w for Q_w^v to be a cube | 40 |
| 5.5 | Toric varieties of Catalan type | 41 |
| 5.6 | Smooth toric Richardson varieties of Catalan type | 44 |
| 6 | Problems | 46 |
| 6.1 | Poincaré polynomial of Y_w^v | 46 |
| 6.2 | Combinatorics of Q_w^v | 46 |

Date: May 30, 2024.

2020 *Mathematics Subject Classification*. Primary: 14M25, 14M15, 57S12; Secondary: 05A05.

Key words and phrases. torus action, flag variety, moment map, toric variety, Coxeter matroid, permutohedron, Schubert variety, Richardson variety, Hessenberg variety, Bruhat interval polytope, Bott manifold.

Lee was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT) (RS-2022-00165641 and RS-2023-00239947), and supported by POSCO Science Fellowship of POSCO TJ Park Foundation. Masuda was supported in part by the HSE University Basic Research Program. Park was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Government of Korea (NRF-2020R1A2C1A01011045). This work was partly supported by Osaka City University Advanced Mathematical Institute (MEXT Joint Usage/Research Center on Mathematics and Theoretical Physics JPMXP0619217849).

| | | |
|-------------------|------------------------|-----------|
| Appendix A | Toric varieties | 47 |
| A.1 | Toric varieties | 47 |
| A.2 | Fano Bott manifolds | 49 |
| References | | 50 |

1. INTRODUCTION

The study of torus orbit closures in the flag variety was initiated by Klyachko [57] and Gelfand–Serganova [46]. Since then, there are some works on the subject but those are mainly about generic orbits or the normality or smoothness of orbit closures (see [38], [29], [73], [25], [26]). There have also been combinatorial studies on polytopes related to torus orbit closure in the flag variety (see [6], [58], [86], [12]). Recently, the authors investigated the topology and geometry of (not necessarily generic) torus orbit closures in connection with combinatorics. In this chapter, we present some of the work by Klyachko and Gelfand–Serganova and our recent works (see [63], [68], [67], [65], [69]) on the topology, geometry, and combinatorics of torus orbit closures in the flag variety. We mainly focus on type A case although some of the arguments work for other Lie types. We give comments for other Lie types when necessary.

Let $\text{Fl}(n)$ be the flag variety defined by

$$\text{Fl}(n) := \{(\{0\} \subset V_1 \subset V_2 \subset \cdots \subset V_n = \mathbb{C}^n) \mid \dim_{\mathbb{C}} V_i = i \text{ for all } i = 1, \dots, n\},$$

where each V_i is a linear subspace of \mathbb{C}^n . Let S_n be the symmetric group on $[n] = \{1, \dots, n\}$. An element of S_n determines a permutation flag. The natural action of the general linear group $\text{GL}_n(\mathbb{C})$ on \mathbb{C}^n induces an action of $\text{GL}_n(\mathbb{C})$ on $\text{Fl}(n)$. Let T be the maximal torus of $\text{GL}_n(\mathbb{C})$ consisting of diagonal matrices, which is isomorphic to $(\mathbb{C}^*)^n$ where $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$. Then the set of permutation flags is exactly the T -fixed point set $\text{Fl}(n)^T$ in $\text{Fl}(n)$ and we identify $\text{Fl}(n)^T$ with S_n . A key tool to study the T -action on $\text{Fl}(n)$ is a moment map

$$\mu: \text{Fl}(n) \rightarrow \mathbb{R}^n$$

which satisfies the following properties:

- $\mu(w) = (w^{-1}(1), \dots, w^{-1}(n))$ for $w \in \text{Fl}(n)^T = S_n$,
- $\mu(\text{Fl}(n))$ is the convex hull of $n!$ points $\mu(\text{Fl}(n)^T)$ in \mathbb{R}^n , that is, the permutohedron Π_n .

We will explore T -varieties (i.e., an algebraic variety with an algebraic T -action) in $\text{Fl}(n)$ with respect to the action of the maximal torus T of $\text{GL}_n(\mathbb{C})$. With this understanding, we will explain the content of each section.

The content of Section 2. The closure of a T -orbit in $\text{Fl}(n)$, denoted by Y , is a T -variety and has an open dense T -orbit, so Y is a toric variety.¹ Since scalar matrices in T act trivially on $\text{Fl}(n)$, the (complex) dimension of Y is at most $n - 1$. By the convexity theorem due to Atiyah [8, Theorem 2] (or Guillemin–Sternberg [50]), the image $\mu(Y)$ is the convex hull of the points $\mu(Y^T)$ in \mathbb{R}^n . The fan of Y is the normal fan of the polytope $\mu(Y)$, so $\mu(Y)$ determines Y as a variety, up to isomorphism. Then two fundamental questions arise:

- (Q1) Characterize the polytopes which arise as $\mu(Y)$ for some T -orbit closure Y .
(Q2) Describe the fan of Y explicitly.

Gelfand–Serganova [46] show that any edge of $\mu(Y)$ must be parallel to an edge of the permutohedron Π_n . Since the edges of Π_n are parallel to roots $\Phi = \{\mathbf{e}_i - \mathbf{e}_j \mid i \neq j \in [n]\}$, where $\mathbf{e}_1, \dots, \mathbf{e}_n$ denote the standard basis of \mathbb{R}^n , a polytope with edges parallel to roots is called a Φ -polytope. Therefore $\mu(Y)$ is a Φ -polytope. Unfortunately, the converse is not true and the complete solution to (Q1) above is unknown. We note that the notion of a Φ -polytope is known as generalized permutohedron [77]. They were originally defined by Edmonds [33], and they are also called M -convex sets (see [74]), and polymatroids (see [81]).

¹ Y is normal in type A but not necessarily normal for other Lie types (see [25]).

Φ -polytopes are closely related to Coxeter matroids. In our context, a Coxeter matroid is a subset \mathcal{M} of S_n which satisfies the Maximality (or Minimality) Condition (see Definition 2.11). The fixed point set Y^T regarded as a subset of S_n is a Coxeter matroid. One can define a retraction

$$\mathcal{R}_{\mathcal{M}}^m : S_n \rightarrow \mathcal{M} \subset S_n$$

for a Coxeter matroid \mathcal{M} and describe the fan of Y explicitly using the retraction to Y^T (see Corollary 2.22), which answers (Q2) above. Here, the suffix m in $\mathcal{R}_{\mathcal{M}}^m$ shows that this retraction is defined using the matroid structure on \mathcal{M} , so that one can distinguish it from the two more retractions introduced later (see Subsection 2.6).

The retraction $\mathcal{R}_{\mathcal{M}}^m$ has two other interpretations, one is geometric and the other is algebraic (see Subsection 2.6). It also has the following interesting property. The 1-skeleton of the permutohedron Π_n is a graph with S_n as vertices. If \mathcal{M} is a Coxeter matroid, then for each $u \in S_n$, a vertex in \mathcal{M} closest to u (with respect to the graph metric) is unique and the closest vertex in \mathcal{M} is given by $\mathcal{R}_{\mathcal{M}}^m(u)$ (see Proposition 2.28).

The content of Section 3. We discuss the topology of a *generic* T -orbit closure Y in $\text{Fl}(n)$, where Y is called generic if

$$Y^T = \text{Fl}(n)^T = S_n,$$

in other words, $\mu(Y) = \Pi_n$. Since Π_n is simple, Y is smooth and of complex dimension $n - 1$. In fact, Y is isomorphic to the permutohedral variety Perm_n which is a toric variety whose fan consists of Weyl chambers. So we may think of the generic T -orbit closure Y as Perm_n . The cohomology of Perm_n can be explicitly described and its Poincaré polynomial agrees with $A_n(t^2)$ where $A_n(t)$ denotes the n th Eulerian polynomial. The fan of Perm_n has the symmetry of S_n which induces an action of S_n on the cohomology of Perm_n . Klyachko [57] shows that the image of the restriction map

$$\iota^* : H^*(\text{Fl}(n); \mathbb{Q}) \rightarrow H^*(\text{Perm}_n; \mathbb{Q}),$$

where $\iota : \text{Perm}_n \rightarrow \text{Fl}(n)$ is the inclusion map, is the ring of S_n -invariants $H^*(\text{Perm}_n; \mathbb{Q})^{S_n}$. This fact is generalized to the setting of regular semisimple Hessenberg varieties. Here, regular semisimple Hessenberg varieties are subvarieties of $\text{Fl}(n)$ satisfying certain conditions (see Subsection 3.4) and Perm_n is the only $(n - 1)$ -dimensional regular semisimple Hessenberg variety that is toric with respect to the maximal torus of $\text{GL}_n(\mathbb{C})$ (see, for example, [31, Theorem 11]).

The content of Section 4. The Schubert variety X_w ($w \in S_n$) is a subvariety of $\text{Fl}(n)$ invariant under the T -action on $\text{Fl}(n)$ and the moment map image of X_w is the polytope

$$\mathbf{Q}_w := \text{Conv}\{(u^{-1}(1), \dots, u^{-1}(n)) \in \mathbb{R}^n \mid u \leq w\},$$

where \leq denotes the Bruhat order² on S_n . We consider a *generic* T -orbit closure, denoted by Y_w , in the Schubert variety X_w , where Y_w is called generic in X_w if

$$Y_w^T = X_w^T = \{u \in S_n \mid u \leq w\}.$$

Since $X_{w_0} = \text{Fl}(n)$ for the longest element $w_0 \in S_n$, Y_{w_0} is isomorphic to the permutohedral variety Perm_n as mentioned above. Although Y_{w_0} is smooth, Y_w is not necessarily smooth. For each $u \in S_n$ with $u \leq w$, we introduce a graph $\Gamma_w(u)$ such that Y_w is smooth at $u \in Y_w^T$ if and only if $\Gamma_w(u)$ is a forest. It seems that the graph $\Gamma_w(w)$ is most complicated among graphs $\Gamma_w(u)$ with $u \leq w$. To be more precise, we posed the following as a conjecture in [63] and it was affirmatively answered in [44] while preparing this chapter.

Theorem ([63, 44]). The graph $\Gamma_w(u)$ is a forest for any $u \leq w$ if $\Gamma_w(w)$ is a forest. (This is equivalent to saying that Y_w is smooth if it is smooth at w , in other words, \mathbf{Q}_w is simple if \mathbf{Q}_w is simple at the vertex $\mu(w)$.)

Indeed, Y_w is always smooth at the identity e and the theorem above says that Y_w is entirely smooth if it is smooth at w . In contrast to this, the Schubert variety X_w is smooth at w and

²The *Bruhat order* (or the *Bruhat-Chevalley order*) on S_n is defined as follows. Let s_i ($i = 1, \dots, n - 1$) denote the adjacent transposition interchanging i and $i + 1$. For $w \in S_n$ with a reduced expression $w = s_{i_1} s_{i_2} \cdots s_{i_q}$ and for $u \in S_n$, we denote $u \leq w$ if there is a reduced expression $u = s_{i_{j_1}} s_{i_{j_2}} \cdots s_{i_{j_k}}$ for $1 \leq j_1 < \cdots < j_k \leq q$.

entirely smooth if it is smooth at the identity e (see [14, p.208]). Interestingly, the graph $\Gamma_w(w)$ appears in a different context, i.e., it is shown in [91] that $\Gamma_w(w)$ is a forest if and only if X_w is locally factorial. Moreover, permutations w for which $\Gamma_w(w)$ is forest are characterized in terms of pattern avoidance (see [18]).

We introduce a polynomial $A_w(t)$ for $w \in S_n$ purely combinatorially by looking at ascents. The polynomial $A_w(t)$ is the Eulerian polynomial $A_n(t)$ when $w = w_0$. As is well-known, $A_{w_0}(t^2)$ agrees with the Poincaré polynomial of the permutohedral variety Perm_n , which is isomorphic to Y_{w_0} . This fact is generalized in such a way that $A_w(t^2)$ agrees with the Poincaré polynomial of Y_w for any $w \in S_n$ (see Theorem 4.17).

We set

$$c(w) := \dim_{\mathbb{C}} X_w - \dim_{\mathbb{C}} Y_w.$$

When $c(w) = 0$, we have $X_w = Y_w$ which means that X_w is a toric variety with the T -action. There are several equivalent conditions for X_w to be a toric variety (see Theorem 4.20), e.g., a reduced decomposition of w is a product of distinct adjacent transpositions. We describe the fan of a toric Schubert variety X_w explicitly in terms of w (see Theorem 4.22). This implies that X_w is a weak Fano Bott manifold, see Appendix A.2 for details of Bott manifolds. We also present a classification result of toric Schubert varieties X_w for Coxeter elements w (see Theorem 4.28).

The case where $c(w) = 1$ is also studied. In this case, X_w is not necessarily smooth. We present several equivalent conditions for $c(w) = 1$ depending on whether X_w is smooth or singular (see Theorems 4.32 and 4.33).

The content of Section 5. For a pair (v, w) of permutations with $v \leq w$, the intersection of a Schubert variety X_w and an opposite Schubert variety $w_0 X_{w_0 v}$ is non-empty and irreducible. The intersection is denoted by X_w^v and called a *Richardson variety*. It is invariant under the T -action on $\text{Fl}(n)$. The moment map image of X_w^v is the *Bruhat interval polytope*

$$Q_w^v := \text{Conv}\{(u^{-1}(1), \dots, u^{-1}(n)) \in \mathbb{R}^n \mid v \leq u \leq w\},$$

introduced by Kodama–Williams [58]. Note that $Q_w^e = Q_w$. The vertex $\mu(v)$ of Q_w^v is not necessarily simple in Q_w^v while it is simple when $v = e$. A natural generalization of the conjecture mentioned above is that Q_w^v is simple if the vertices $\mu(v)$ and $\mu(w)$ are both simple in Q_w^v .

We study the case when X_w^v is a toric variety with the T -action. A toric Richardson variety X_w^v is not necessarily smooth while every toric Schubert variety X_w is smooth. It turns out that X_w^v is a smooth toric variety if and only if Q_w^v is combinatorially equivalent to a cube (see Theorem 5.7). Although pairs (v, w) for which Q_w^v is combinatorially equivalent to a cube are not completely understood, there are many such pairs. Especially, it is shown in [51] that this is the case when

$$w = s_{n-1} s_{n-1} \cdots s_1 v \quad (\text{or } w = s_1 s_2 \cdots s_{n-1} v) \quad \text{and} \quad \ell(w) - \ell(v) = n - 1.$$

Toric Richardson varieties X_w^v for the pairs (v, w) above also arise from polygon triangulations, so we call such toric Richardson varieties of *Catalan type*. They can be classified up to isomorphism and the Wedderburn–Etherington numbers which count *unordered* binary trees appear in enumerating the isomorphism classes (see Corollary 5.23).

Finally, in Section 6 we pose several problems related to the discussion developed in this chapter. In Appendix A, we briefly review the theory of toric varieties and explain Bott manifolds in detail.

2. TORUS ORBIT CLOSURES (GENERAL)

In this section, we review the torus actions on flag varieties and the closure of a torus orbit. Moreover, we consider properties of the moment map images of torus orbit closures. Indeed, the fixed point set of a torus orbit closure becomes a Coxeter matroid. Finally, we discuss how to describe the fan of a torus orbit closure.

2.1. Torus actions on flag varieties. Let G be the general linear group $\mathrm{GL}_n(\mathbb{C})$ over \mathbb{C} , $B \subset G$ the set of upper triangular matrices, and $T \subset G$ the set of diagonal matrices. Let $B^- \subset G$ be the set of lower triangular matrices. We denote by S_n the symmetric group on $[n] := \{1, \dots, n\}$. Then $T = B \cap B^- \cong (\mathbb{C}^*)^n$ and $B^- = w_0 B w_0$, where w_0 is the longest element $[n, n-1, \dots, 1] \in S_n$. The homogeneous space G/B can be identified with the *flag variety* $\mathrm{Fl}(n)$ defined by

$$\mathrm{Fl}(n) := \{(\{0\} \subset V_1 \subset V_2 \subset \dots \subset V_n = \mathbb{C}^n) \mid \dim_{\mathbb{C}} V_i = i \text{ for all } i = 1, \dots, n\},$$

where \mathbb{C}^n is considered as the complex vector space consisting of *column* vectors. For $g \in G$, let $\mathbf{c}_1, \dots, \mathbf{c}_n$ be the column vectors of the matrix g , that is, $g = [\mathbf{c}_1 \ \dots \ \mathbf{c}_n]$. Then $gB \in G/B$ corresponds to the flag whose i th vector space is spanned by the first i columns $\mathbf{c}_1, \dots, \mathbf{c}_i$ of g .

For an element $w \in S_n$, we use the same letter w for the permutation matrix $[\mathbf{e}_{w(1)} \ \dots \ \mathbf{e}_{w(n)}]$ in $\mathrm{GL}_n(\mathbb{C})$ to simplify notation, where $\mathbf{e}_1, \dots, \mathbf{e}_n$ are the standard basis vectors in \mathbb{C}^n . The left multiplication by T on G induces the T -action on G/B .

Lemma 2.1 ([42, §10.5]). *The set of T -fixed points in G/B bijectively corresponds to the symmetric group S_n such that each $u \in S_n$ corresponds to $uB \in G/B$.*

For each $u \in S_n$, there is a T -invariant local chart U_u given by

$$(2.1) \quad U_u := \left\{ (x_{ij})B \in G/B \mid x_{ij} = \begin{cases} 1 & \text{if } j = u^{-1}(i), \\ 0 & \text{if } j > u^{-1}(i) \end{cases} \right\},$$

which is isomorphic to $\mathbb{C}^{n(n-1)/2}$ since x_{ij} with $j < u^{-1}(i)$ are arbitrary complex numbers. The T -action on G/B restricted to U_u is given by

$$(2.2) \quad (t_1, \dots, t_n) \cdot (x_{ij})B = (t_i t_{u(j)}^{-1} x_{ij})B.$$

Here, for $(t_1, \dots, t_n) \in (\mathbb{C}^*)^n$, we associate a diagonal matrix $\mathrm{diag}(t_1, \dots, t_n) \in T$. Since we have the 1's on the $(i, u^{-1}(i))$ -entries of $(x_{ij}) \in U_u$ (which is equivalent to saying that we have the 1's on the $(u(j), j)$ -entries), by considering the multiplication of an element $\mathrm{diag}(t_{u(1)}^{-1}, \dots, t_{u(n)}^{-1})$ of B from the right, we obtain the above computation. We demonstrate this action for $u = 312$ in Example 2.2. Therefore, uB is a unique T -fixed point in U_u , which corresponds to the origin in $\mathbb{C}^{n(n-1)/2}$ and a 1-dimensional T -orbit in U_u corresponds to a 1-dimensional T -eigenspace in $\mathbb{C}^{n(n-1)/2}$.

Example 2.2. When $u = 312$ in one-line notation,

$$U_{312} = \left\{ \left(\begin{array}{ccc} x_{11} & 1 & 0 \\ x_{21} & x_{22} & 1 \\ 1 & 0 & 0 \end{array} \right) B \mid x_{ij} \in \mathbb{C} \right\} \cong \mathbb{C}^3,$$

and the T -action on U_{312} is given by

$$\begin{aligned} (t_1, t_2, t_3) \cdot \left(\begin{array}{ccc} x_{11} & 1 & 0 \\ x_{21} & x_{22} & 1 \\ 1 & 0 & 0 \end{array} \right) B &= \left(\begin{array}{ccc} t_1 x_{11} & t_1 & 0 \\ t_2 x_{21} & t_2 x_{22} & t_2 \\ t_3 & 0 & 0 \end{array} \right) B \\ &= \left(\begin{array}{ccc} t_1 x_{11} & t_1 & 0 \\ t_2 x_{21} & t_2 x_{22} & t_2 \\ t_3 & 0 & 0 \end{array} \right) \begin{pmatrix} t_3^{-1} & 0 & 0 \\ 0 & t_1^{-1} & 0 \\ 0 & 0 & t_2^{-1} \end{pmatrix} B \\ &= \left(\begin{array}{ccc} t_1 t_3^{-1} x_{11} & 1 & 0 \\ t_2 t_3^{-1} x_{21} & t_2 t_1^{-1} x_{22} & 1 \\ 1 & 0 & 0 \end{array} \right) B. \end{aligned}$$

2.2. Moment map. We describe a moment map $\mu: G/B \rightarrow \mathbb{R}^n$ explicitly using the Plücker coordinates with respect to the action of the (maximal) compact torus $T_{\mathbb{R}}$ in T . Note that $T \cong (S^1)^n$. Define the set

$$I_{d,n} := \{\mathbf{i} = (i_1, \dots, i_d) \in \mathbb{Z}^d \mid 1 \leq i_1 < \dots < i_d \leq n\}.$$

For an element $x = (x_{ij}) \in G = \mathrm{GL}_n(\mathbb{C})$, the \mathbf{i} th Plücker coordinate $p_{\mathbf{i}}(x)$ of x is given by the $d \times d$ minor of x , with row indices i_1, \dots, i_d and the column indices $1, \dots, d$ for $\mathbf{i} = (i_1, \dots, i_d) \in I_{d,n}$. The Plücker embedding is defined to be

$$(2.3) \quad \psi: G/B \rightarrow \prod_{d=1}^{n-1} \mathbb{C}P^{(d)-1}, \quad xB \mapsto \prod_{d=1}^{n-1} [p_{\mathbf{i}}(x)]_{\mathbf{i} \in I_{d,n}}.$$

The map ψ is well-defined because $p_{\mathbf{i}}(xb) = (\prod_{j=1}^d b_{jj})p_{\mathbf{i}}(x)$ for $b = (b_{ij}) \in B$ and $\mathbf{i} \in I_{d,n}$. The map ψ is T -equivariant with respect to the action of T on $\prod_{d=1}^{n-1} \mathbb{C}P^{(d)-1}$ given by

$$(t_1, \dots, t_n) \cdot [p_{\mathbf{i}}]_{\mathbf{i} \in I_{d,n}} := [t_{i_1} \cdots t_{i_d} \cdot p_{\mathbf{i}}]_{\mathbf{i} \in I_{d,n}}$$

for $(t_1, \dots, t_n) \in T$ and $\mathbf{i} = (i_1, \dots, i_d) \in I_{d,n}$. Here, with the abuse of notation, we are denoting by $p_{\mathbf{i}}$ both a function $p_{\mathbf{i}}: G \rightarrow \mathbb{C}$ and the \mathbf{i} th coordinate of $\mathbb{C}P^{(d)-1}$.

Let ω_{FS} be the Fubini–Study form on a complex projective space $\mathbb{C}P^{m-1}$. With respect to the standard action of the compact torus $(S^1)^m$ on $\mathbb{C}P^{m-1}$ defined by

$$(t_1, \dots, t_m) \cdot [z_1, \dots, z_m] = [t_1 z_1, \dots, t_m z_m],$$

where each t_i is a complex number with unit length, the moment map of $(\mathbb{C}P^{m-1}, \omega_{FS}, (S^1)^m)$ is given by

$$(2.4) \quad [z_1, \dots, z_m] \mapsto \frac{-1}{2 \sum_{i=1}^m |z_i|^2} (|z_1|^2, \dots, |z_m|^2) \quad \text{up to translation.}$$

See, for example, [9, Example IV.1.2].³ Because the action of $(S^1)^n$ on the factor $\mathbb{C}P^{(d)-1}$ in (2.3) is given through the homeomorphism $\phi_d: T_{\mathbb{R}} = (S^1)^n \rightarrow (S^1)^{(d)}$ sending (t_1, \dots, t_n) to $(t_{i_1} \cdots t_{i_d})_{\mathbf{i} \in I_{d,n}}$, the moment map of $(\mathbb{C}P^{(d)-1}, 2\omega_{FS}, T_{\mathbb{R}})$ is given by

$$[p_{\mathbf{i}}]_{\mathbf{i} \in I_{d,n}} \mapsto \frac{-1}{\sum_{\mathbf{i} \in I_{d,n}} |p_{\mathbf{i}}|^2} \left(\sum_{\mathbf{1} \in I_{d,n}} |p_{\mathbf{i}}|^2, \dots, \sum_{\mathbf{n} \in I_{d,n}} |p_{\mathbf{i}}|^2 \right) \quad \text{up to translation.}$$

We note that for symplectic manifolds M_1 and M_2 having Hamiltonian actions of the same Lie group G with moment maps $\mu_1: M_1 \rightarrow \mathrm{Lie}(G)^*$ and $\mu_2: M_2 \rightarrow \mathrm{Lie}(G)^*$, respectively, the diagonal action of G on $M_1 \times M_2$ is Hamiltonian. Moreover, the moment map $\mu: M_1 \times M_2 \rightarrow \mathrm{Lie}(G)^*$ is given by the sum of moment maps μ_1 and μ_2 . Indeed, $\mu(x_1, x_2) = \mu_1(x_1) + \mu_2(x_2)$ (see, for instance, [9, Exercise III.3]). By considering a symplectic form ω on $\prod_{d=1}^{n-1} \mathbb{C}P^{(d)-1}$ given by two times of the Fubini–Study form on each complex projective space and the diagonal action on the product, the moment map $\tilde{\mu}: \prod_{d=1}^{n-1} \mathbb{C}P^{(d)-1} \rightarrow \mathbb{R}^n$ is given by

$$(2.5) \quad \prod_{d=1}^{n-1} [p_{\mathbf{i}}]_{\mathbf{i} \in I_{d,n}} \mapsto - \sum_{d=1}^{n-1} \left\{ \frac{1}{\sum_{\mathbf{i} \in I_{d,n}} |p_{\mathbf{i}}|^2} \left(\sum_{\mathbf{1} \in I_{d,n}} |p_{\mathbf{i}}|^2, \dots, \sum_{\mathbf{n} \in I_{d,n}} |p_{\mathbf{i}}|^2 \right) \right\} + \mathbf{c},$$

where \mathbf{c} is a constant vector in \mathbb{R}^n , and the action of $(S^1)^n$ on each factor $\mathbb{C}P^{(d)-1}$ is given through the homomorphism ϕ_d above. We take $\mathbf{c} = (n, \dots, n)$ in (2.5) and define

$$(2.6) \quad \mu := \tilde{\mu} \circ \psi,$$

which is a moment map of $(G/B, \psi^*\omega, T_{\mathbb{R}})$.

Before computing the moment image of the fixed point $uB \in G/B$, we introduce one notation. For a subset $S \subset [n]$, we denote by $S \uparrow$ the ordered tuple obtained from S by sorting its elements in ascending order. For instance, if $S = \{5, 2, 1, 6\} \subset [8]$, then $S \uparrow = (1, 2, 5, 6)$.

Lemma 2.3. *The moment map μ sends the fixed point $uB \in G/B$ to $(u^{-1}(1), \dots, u^{-1}(n)) \in \mathbb{R}^n$.*

³Here, we use a different sign convention from that in [9]. That is, our moment map $\mu: (M, \omega, (S^1)^n) \rightarrow \mathrm{Lie}((S^1)^n)^*$ satisfies the following: For each $X \in \mathrm{Lie}((S^1)^n)$, $d\mu^X = \iota_{X^\#} \omega$, where $\mu^X(p) = \langle \mu(p), X \rangle$ and $X^\#$ is the vector field on M generated by the one-parameter subgroup $\{\exp tX \mid t \in \mathbb{R}\} \subset (S^1)^n$.

Proof. For a permutation $u \in S_n$, the Plücker coordinates $(p_{\mathbf{i}})_{\mathbf{i} \in I_{d,n}}$ of uB are given as follows:

$$(2.7) \quad p_{\mathbf{i}} = \begin{cases} \pm 1 & \text{if } \mathbf{i} = \{u(1), \dots, u(d)\} \uparrow, \\ 0 & \text{otherwise,} \end{cases}$$

for each $\mathbf{i} \in I_{d,n}$. Therefore, one can see that for a fixed $d \in [n-1]$, $\sum_{\mathbf{i} \in I_{d,n}} |p_{\mathbf{i}}|^2 = 1$ and the entries of the vector

$$\left(\sum_{\mathbf{i} \in I_{d,n}} |p_{\mathbf{i}}|^2, \dots, \sum_{\mathbf{i} \in I_{d,n}} |p_{\mathbf{i}}|^2 \right)$$

are 1 for coordinates in $\{u(1), \dots, u(d)\}$ and 0 otherwise. Hence the summation

$$- \sum_{d=1}^{n-1} \left\{ \frac{1}{\sum_{\mathbf{i} \in I_{d,n}} |p_{\mathbf{i}}|^2} \left(\sum_{\mathbf{i} \in I_{d,n}} |p_{\mathbf{i}}|^2, \dots, \sum_{\mathbf{i} \in I_{d,n}} |p_{\mathbf{i}}|^2 \right) \right\}$$

is an integer vector such that the $u(k)$ -entry is $-(n-k)$. Therefore, $\mu(uB)$ is an integer vector whose $u(k)$ -entry is k since $\mathbf{c} = (n, \dots, n)$ in (2.5). This implies that $\mu(uB) = (u^{-1}(1), \dots, u^{-1}(n))$ since $u^{-1}(u(k)) = k$ for all k . \square

Example 2.4. When $G = \mathrm{GL}_3(\mathbb{C})$, for $x = (x_{ij}) \in \mathrm{GL}_3(\mathbb{C})$, the Plücker embedding $\psi: G/B \rightarrow \mathbb{C}P^{(3)}_{(1)}^{-1} \times \mathbb{C}P^{(3)}_{(2)}^{-1}$ maps an element $xB \in G/B$ to

$$\begin{aligned} & ([p_1(x), p_2(x), p_3(x)], [p_{1,2}(x), p_{1,3}(x), p_{2,3}(x)]) \\ & = ([x_{11}, x_{21}, x_{31}], [x_{11}x_{22} - x_{21}x_{12}, x_{11}x_{32} - x_{31}x_{12}, x_{21}x_{32} - x_{31}x_{22}]). \end{aligned}$$

Since the action of T on $\mathrm{GL}_3(\mathbb{C})$ is given by

$$(t_1, t_2, t_3) \cdot \begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{pmatrix} = \begin{pmatrix} t_1 x_{11} & t_1 x_{12} & t_1 x_{13} \\ t_2 x_{21} & t_2 x_{22} & t_2 x_{23} \\ t_3 x_{31} & t_3 x_{32} & t_3 x_{33} \end{pmatrix}$$

for $t = (t_1, t_2, t_3) \in T$, we have

$$\begin{aligned} \psi(t \cdot xB) & = ([t_1 x_{11}, t_2 x_{21}, t_3 x_{31}], [t_1 t_2 (x_{11}x_{22} - x_{21}x_{12}), t_1 t_3 (x_{11}x_{32} - x_{31}x_{12}), t_2 t_3 (x_{21}x_{32} - x_{31}x_{22})]) \\ & = t \cdot \psi(xB) \end{aligned}$$

so the map ψ is T -equivariant. The moment map

$$\tilde{\mu}: \mathbb{C}P^{(3)}_{(1)}^{-1} \times \mathbb{C}P^{(3)}_{(2)}^{-1} \rightarrow \mathbb{R}^3$$

is given by

$$\begin{aligned} & ([p_1, p_2, p_3], [p_{1,2}, p_{1,3}, p_{2,3}]) \\ & \mapsto - \frac{1}{|p_1|^2 + |p_2|^2 + |p_3|^2} (|p_1|^2, |p_2|^2, |p_3|^2) \\ & \quad - \frac{1}{|p_{1,2}|^2 + |p_{1,3}|^2 + |p_{2,3}|^2} (|p_{1,2}|^2 + |p_{1,3}|^2, |p_{1,2}|^2 + |p_{2,3}|^2, |p_{1,3}|^2 + |p_{2,3}|^2) \\ & \quad + (3, 3, 3). \end{aligned}$$

For $312 \in S_3$, we have

$$\mu(312B) = \tilde{\mu} \circ \psi(312B) = \tilde{\mu}([0, 0, 1], [0, -1, 0]) = -(0, 0, 1) - (1, 0, 1) + (3, 3, 3) = (2, 3, 1).$$

Remark 2.5. The Plücker embedding can be generalized to partial flag varieties and other Lie types (see, for example, [46] and [17]).

2.3. Torus orbit closures and their moment polytopes. For a point $x \in G/B$, the closure $\overline{T \cdot x}$ of the T -orbit $T \cdot x$ is a toric variety in G/B . The moment map image $\mu(\overline{T \cdot x})$ is a convex polytope in \mathbb{R}^n with vertices $\mu((\overline{T \cdot x})^T)$ by the convexity theorem (Atiyah [8, Theorem 2], Guillemin–Sternberg [50] for symplectic case).

On the other hand, $(\overline{T \cdot x})^T$ can be found as follows. Choose a representative $\tilde{x} \in \mathrm{GL}_n(\mathbb{C})$ of x . The non-vanishing of $p_{\mathbf{i}}(\tilde{x})$ is independent of the choice of the representative \tilde{x} of x and we define

$$I_d(x) := \{\mathbf{i} = (i_1, \dots, i_d) \in I_{d,n} \mid p_{\mathbf{i}}(\tilde{x}) \neq 0\}, \quad 1 \leq d \leq n.$$

Proposition 2.6 ([46, Proposition 1 in §5.2]). *For an element $x \in \mathrm{GL}_n(\mathbb{C})/B$, we have*

$$(\overline{T \cdot x})^T = \{wB \mid w \in S_n, \{w(1), \dots, w(d)\} \uparrow \in I_d(x) \text{ for all } 1 \leq d \leq n\}.$$

Proof. We note that for $\mathbf{i} \in I_{d,n}$, if $p_{\mathbf{i}}(\tilde{x}) = 0$, then $p_{\mathbf{i}}(\tilde{y}) = 0$ for any $y \in \overline{T \cdot x}$. Here, we denote by $\tilde{y} \in \mathrm{GL}_n(\mathbb{C})$ a representative of y . Therefore we have

$$\psi((\overline{T \cdot x})^T) \subset \left\{ [p_{\mathbf{i}}]_{\mathbf{i} \in I_{d,n}} \in \prod_{d=1}^{n-1} \mathbb{C}P^{\binom{n}{d}-1} \mid p_{\mathbf{i}} \neq 0 \text{ if } \mathbf{i} \in I_d(x) \right\}.$$

This provides

$$(\overline{T \cdot x})^T \subset \{wB \mid w \in S_n, \{w(1), \dots, w(d)\} \uparrow \in I_d(x) \text{ for all } 1 \leq d \leq n\}$$

since for $\mathbf{i} \in I_{d,n}$, we have $p_{\mathbf{i}}(wB) \neq 0$ if and only if $\{w(1), \dots, w(d)\} \uparrow = \mathbf{i}$ (see (2.7)).

Now we consider the opposite inclusion. Recall that the group of homomorphisms from T to \mathbb{C}^* is naturally isomorphic to \mathbb{Z}^n . For each $u = (u_1, \dots, u_n) \in \mathbb{Z}^n$, we have the one-parameter subgroup $\lambda_u(t) = (t^{u_1}, \dots, t^{u_n})$ of T , where $t \in \mathbb{C}^*$. Suppose that $w \in S_n$ satisfies $\{w(1), \dots, w(d)\} \uparrow \in I_d(x)$ for all $1 \leq d \leq n$. Take $u = (u_1, u_2, \dots, u_n) \in \mathbb{Z}^n$ such that $u_{w(1)} < u_{w(2)} < \dots < u_{w(n)}$. Then we have

$$\lim_{t \rightarrow 0} \lambda_u(t) \cdot x = wB.$$

This proves the desired inclusion so we are done. \square


Example 2.7. Let $x = \begin{pmatrix} \alpha & 1 & 0 \\ \beta & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} B \in \mathrm{GL}_3(\mathbb{C})/B$ with $\alpha, \beta \in \mathbb{C}^*$. Then we have

$$\begin{aligned} I_1(x) &= \{(1), (2), (3)\}, & I_2(x) &= \{(1, 2), (1, 3)\}, & I_3(x) &= \{(1, 2, 3)\}, \text{ and} \\ (\overline{T \cdot x})^T &= \{wB \mid w \in S_3, \{w(1), \dots, w(d)\} \uparrow \in I_d(x) \text{ for all } 1 \leq d \leq 3\} \\ &= \{123B, 132B, 213B, 312B\}. \end{aligned}$$

Since

$$(2.8) \quad \mu(123B) = (1, 2, 3), \quad \mu(132B) = (1, 3, 2), \quad \mu(213B) = (2, 1, 3), \quad \mu(312B) = (2, 3, 1)$$

by Lemma 2.3, $\mu(\overline{T \cdot x})$ is the convex hull of the four points in (2.8), see Figure 1. Now we label each vertex $\mu(uB)$ with u (using a different font) for simplicity.



(1) Vertices are labeled by coordinates (2) Vertices are labeled by permutations

FIGURE 1. Moment polytope $\mu(\overline{T \cdot x})$

In general, the moment polytope $\mu(\overline{T \cdot x})$ has the following property.

Proposition 2.8. *If two vertices $\mu(uB)$ and $\mu(vB)$ of $\mu(\overline{T \cdot x})$ are joined by an edge of $\mu(\overline{T \cdot x})$ for $u, v \in S_n$, then $tu = v$ for some transposition t of S_n , in other words, an edge of $\mu(\overline{T \cdot x})$ is parallel to $\mathbf{e}_i - \mathbf{e}_j$ for some $1 \leq i < j \leq n$.*

Proof. By the convexity theorem, an edge of $\mu(\overline{T \cdot x})$ is $\mu(\overline{O})$ for some 1-dimensional T -orbit O in $\overline{T \cdot x}$. Since $\overline{T \cdot x}$ is invariant under the T -action on G/B , O is also a 1-dimensional T -orbit in G/B . Note that O is isomorphic to \mathbb{C}^* , \overline{O} is $\mathbb{C}P^1$, and $\overline{O} \setminus O$ consists of two T -fixed points, say uB and vB , which map to the two end points of the edge $\mu(\overline{O})$ by the moment map μ .

Let U_u be the T -invariant chart of G/B centered at uB in (2.1). Then, as noted before, (2.2) implies that the T -orbit O is of the form

$$(u + cE_{ij})B \quad \text{for some } i, j \text{ with } j < u^{-1}(i),$$

where $c \in \mathbb{C}^*$ and E_{ij} is the $n \times n$ matrix with 1 at the (i, j) entry and 0 otherwise. Suppose that $i < u(j)$ (the essentially same argument works when $i > u(j)$). Then we apply the following elementary transformations to $(u + cE_{ij})B$:

- (1) multiply the j th column by $1/c$,
- (2) subtract the j th column from the $u^{-1}(i)$ th column,
- (3) multiply the $u^{-1}(i)$ th column by $-c$,

and then approach c to ∞ :

$$\begin{pmatrix} c & 1 \\ 1 & 0 \end{pmatrix} B \stackrel{(1)}{=} \begin{pmatrix} 1 & 1 \\ 1/c & 0 \end{pmatrix} B \stackrel{(2)}{=} \begin{pmatrix} 1 & 0 \\ 1/c & -1/c \end{pmatrix} B \stackrel{(3)}{=} \begin{pmatrix} 1 & 0 \\ 1/c & 1 \end{pmatrix} B \xrightarrow{c \rightarrow \infty} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} B,$$

where the 2×2 matrices above are intersections of the j th and $u^{-1}(i)$ th columns ($j < u^{-1}(i)$) and the i th and $u(j)$ th rows ($i < u(j)$). The other entries remain unchanged by the elementary transformations above. This shows that when c approaches ∞ , the point $(u + cE_{ij})B$ approaches the T -fixed point $t_{i,u(j)}uB$, where $t_{i,u(j)}$ is the transposition interchanging i and $u(j)$. Since $vB = t_{i,u(j)}uB$, this implies the proposition. \square

Since the root system of type A_{n-1} is $\Phi = \{\pm(\mathbf{e}_i - \mathbf{e}_j) \mid 1 \leq i < j \leq n\}$, Proposition 2.8 says that an edge vector of the moment polytope $\mu(\overline{T \cdot x})$ is parallel to a root in Φ . This fact holds in any Lie type.

Theorem 2.9 ([17, Theorem on page xii]). *Let G be a semisimple algebraic group over \mathbb{C} , B a Borel subgroup in G and $T \subset B$ a maximal torus. Let Φ be the associated root system and $\mu: G/B \rightarrow \mathfrak{t}_{\mathbb{R}}^*$ a moment map, where $\mathfrak{t}_{\mathbb{R}}^*$ denotes the vector space dual to the Lie algebra $\mathfrak{t}_{\mathbb{R}}$ of the maximal compact torus of T . Then, for any point $x \in G/B$, an edge of the moment polytope $\mu(\overline{T \cdot x})$ is parallel to a root in Φ (such a polytope is called a Φ -polytope, or a generalized permutohedron, see Definition 2.14).*

Remark 2.10. Partial flag varieties also have torus actions and one may consider moment maps. We refer the reader to [45] for the combinatorial properties of moment maps of torus orbit closures in Grassmannian.

2.4. Coxeter matroids and Gelfand–Serganova Theorem. To study the combinatorial properties of torus orbit closures, Gelfand and Serganova introduced the notion of Coxeter matroids. In this subsection, we recall the definition of Coxeter matroids and the characterization of Coxeter matroids in terms of polytopes by Gelfand–Serganova. It turns out that the permutations indexing the T -fixed points of a T -orbit closure in the flag variety G/B is a Coxeter matroid.

Let W be a finite Coxeter group, so generators of W are prescribed. The symmetric group S_n on $[n]$ with adjacent transpositions as generators is a typical example of a Coxeter group. For $u \in W$, let \leq^u denote the u -shifted order, that is, $v \leq^u w$ means $u^{-1}v \leq u^{-1}w$ in the Bruhat order on W . Note that \leq^u is a partial order on W with u as the smallest element.

Definition 2.11. A subset \mathcal{M} of a finite Coxeter group W is called a *Coxeter matroid* if it satisfies the *Maximality Property*: for any $u \in W$, there is a unique element $v \in \mathcal{M}$ such that $w \leq^u v$ for all $w \in \mathcal{M}$.

We note that for a finite Coxeter group W in type A , Coxeter matroids are also called *flag matroids*. See [89]. For more details on flag matroids and flag matroid polytopes, we refer the reader to [6, 23, 12].

Remark 2.12. (1) Since the multiplication by the longest element of W reverses the Bruhat order on W , the Maximality Property is equivalent to the *Minimality Property*: for any $u \in W$, there exists a unique element $v \in \mathcal{M}$ such that $v \leq^u w$ for all $w \in \mathcal{M}$.
 (2) In fact, a Coxeter matroid is defined more generally in [17]. Let W_P be a parabolic subgroup of a finite Coxeter group W . Then the Bruhat order on W induces a partial order on W/W_P and a subset \mathcal{M} of W/W_P is called a Coxeter matroid if it satisfies the Maximality Property above. A Coxeter matroid in Definition 2.11 is the case where W_P is the identity subgroup and an ordinary matroid can be regarded as the case where $W = S_n$ and W_P is a maximal parabolic subgroup (see [17, Theorem 1.3.1]).

Example 2.13 ([67, Example 2.2]). Let $\mathcal{M} = \{213, 132\}$ be a subset of S_3 . Since $213 \not\leq 132$ and $132 \not\leq 213$, there is no element $v \in \mathcal{M}$ such that $w \leq^{123} v$ for all $w \in \mathcal{M}$. Hence \mathcal{M} is not a Coxeter matroid. However, one can check that $\{231, 321\}$ is a Coxeter matroid of S_3 .

We recall the characterization of Coxeter matroids in terms of polytopes by Gelfand–Serganova. As is well-known, a finite Coxeter group W can be regarded as a reflection group on a vector space V , where the generators of W act on V as reflections (see [54, Section 5.3]). Let Φ be the set of roots of W .

Definition 2.14. A convex polytope Δ in V is called a Φ -polytope, or a *generalized permutohedron*, if every edge of Δ is parallel to a root in Φ .

We note that the notion of a Φ -polytope is known as generalized permutohedron [77]. They were originally defined by Edmonds [33], and they are also called M -convex sets (see [74]), and polymatroids (see [81]).

Choose a point $\nu \in V$ which is not fixed by any reflection in W . For a subset \mathcal{M} of W , we define $\Delta_{\mathcal{M}}$ to be the convex hull of the \mathcal{M} -orbit $\{w \cdot \nu \mid w \in \mathcal{M}\}$ of the point ν in V . When $\mathcal{M} = W$, Δ_W is called the W -permutohedron (see [39, Section 2.4], [52], and [7]). Two vertices $v \cdot \nu$ and $w \cdot \nu$ of Δ_W are joined by an edge of Δ_W if and only if $v^{-1}w$ is a generator of W (see [39, Lemma 2.13]). Thus any edge of Δ_W is parallel to a root and vice versa. Therefore, the W -permutohedron Δ_W is a Φ -polytope and we may say that a convex polytope Δ in V is a Φ -polytope if every edge of Δ is parallel to an edge of Δ_W .

However, $\Delta_{\mathcal{M}}$ is not necessarily a Φ -polytope unless $\mathcal{M} = W$. The following is a part of the Gelfand–Serganova Theorem [46] (also see [17, Theorem 6.3.1]).

Theorem 2.15 (Gelfand–Serganova Theorem). *A subset \mathcal{M} of a finite Coxeter group W is a Coxeter matroid if and only if $\Delta_{\mathcal{M}}$ is a Φ -polytope. Therefore, the permutations indexing the T -fixed points of a T -orbit closure in the flag variety G/B is a Coxeter matroid by Theorem 2.9.*

We provide the proof for the case $W = S_n$. We refer the reader to [17] for more details. Before giving it, we prepare two lemmas. Suppose that

$$V = \{(a_1, \dots, a_n) \in \mathbb{R}^n \mid a_1 + \dots + a_n = 0\}$$

and S_n acts on V as permuting coordinates, i.e., $u \cdot (a_1, \dots, a_n) = (a_{u^{-1}(1)}, \dots, a_{u^{-1}(n)})$, in other words,

$$u \cdot \left(\sum_{i=1}^n a_i \mathbf{e}_i \right) = \sum_{i=1}^n a_i \mathbf{e}_{u(i)}.$$

We define the ordering \preceq^u on V by putting $x \preceq^u y$ for $x, y \in V$ if there exists non-negative constants $c_i \geq 0$ such that

$$y - x = \sum_{i=1}^{n-1} c_i (\mathbf{e}_{u(i)} - \mathbf{e}_{u(i+1)}),$$

which is equivalent to saying $u^{-1} \cdot y - u^{-1} \cdot x = \sum_{i=1}^{n-1} c_i(\mathbf{e}_i - \mathbf{e}_{i+1})$ with the same coefficients c_i . This means

$$(2.9) \quad x \preceq^u y \iff u^{-1} \cdot x \preceq^e u^{-1} \cdot y.$$

Lemma 2.16 ([17, Lemma 6.2.3]). *Suppose that $\nu = (\nu_1, \nu_2, \dots, \nu_n) \in V$ satisfies $\nu_1 < \nu_2 < \dots < \nu_n$. If $v \leq^u w$, then $v \cdot \nu \preceq^u w \cdot \nu$.*

Proof. Note that $v \leq^u w$ if and only if $u^{-1}v \leq u^{-1}w$, and $v \cdot \nu \preceq^u w \cdot \nu$ if and only if $u^{-1}v \cdot \nu \preceq^e u^{-1}w \cdot \nu$ (see (2.9)). Hence it suffices to prove the statement in the case $u = e$. Suppose $v \leq w$. Then we have $v^{-1} \leq w^{-1}$ and hence $\{v^{-1}(1), \dots, v^{-1}(d)\} \uparrow \leq \{w^{-1}(1), \dots, w^{-1}(d)\} \uparrow$ for $1 \leq d \leq n$ (cf. [14, §3.2]). This implies $\sum_{i=1}^d v^{-1}(i) \leq \sum_{i=1}^d w^{-1}(i)$ for $1 \leq d \leq n$, so we get

$$(2.10) \quad \sum_{i=1}^d \nu_{v^{-1}(i)} \leq \sum_{i=1}^d \nu_{w^{-1}(i)} \quad \text{for } 1 \leq d \leq n.$$

On the other hand, we have

$$\begin{aligned} w \cdot \nu - v \cdot \nu &= (\nu_{w^{-1}(1)}, \dots, \nu_{w^{-1}(n)}) - (\nu_{v^{-1}(1)}, \dots, \nu_{v^{-1}(n)}) \\ &= \sum_{d=1}^{n-1} \left(\sum_{i=1}^d \nu_{w^{-1}(i)} - \sum_{i=1}^d \nu_{v^{-1}(i)} \right) (\mathbf{e}_d - \mathbf{e}_{d+1}). \end{aligned}$$

Accordingly, by (2.10), we obtain $v \cdot \nu \preceq^e w \cdot \nu$ and we are done. \square

The converse in Lemma 2.16 does not hold in general. For instance, if $v = 1432$ and $w = 4123$, then $v \not\leq w$ but since

$$\begin{aligned} w \cdot \nu - v \cdot \nu &= (\nu_2, \nu_3, \nu_4, \nu_1) - (\nu_1, \nu_4, \nu_3, \nu_2) \\ &= (\nu_2 - \nu_1)(\mathbf{e}_1 - \mathbf{e}_2) + (\nu_2 + \nu_3 - \nu_1 - \nu_4)(\mathbf{e}_2 - \mathbf{e}_3) + (\nu_2 - \nu_1)(\mathbf{e}_3 - \mathbf{e}_4), \end{aligned}$$

we have $v \cdot \nu \preceq^e w \cdot \nu$ if $\nu_2 + \nu_3 \geq 0$ (note that $\nu_1 + \nu_2 + \nu_3 + \nu_4 = 0$ because $\nu \in V$).

However, the converse holds in the following special case.

Lemma 2.17 ([17, Lemma 6.2.5]). *Let ν be as in Lemma 2.16. Suppose that $w \cdot \nu - v \cdot \nu$ is parallel to a root in Φ . Then $v \leq^u w$ if $v \cdot \nu \preceq^u w \cdot \nu$.*

Proof. As noted in the proof of Lemma 2.16, we may assume $u = e$. If $v \cdot \nu \preceq^e w \cdot \nu$, then

$$w \cdot \nu - v \cdot \nu = c(\mathbf{e}_i - \mathbf{e}_j)$$

for some $c > 0$ and $i < j$. This implies that $w^{-1} = v^{-1}t_{i,j}$ and $v^{-1}(i) < v^{-1}(j)$ since $c > 0$, where $t_{i,j}$ denotes the transposition interchanging i and j . Therefore $v^{-1} \leq w^{-1}$ and hence $v \leq w$. \square

Proof of Theorem 2.15 in type A. We take $\nu = (\nu_1, \nu_2, \dots, \nu_n) \in V$ which satisfies $\nu_1 < \nu_2 < \dots < \nu_n$. We will prove that if $\mathcal{M} \subset S_n$ is a Coxeter matroid, then $\Delta_{\mathcal{M}}$ is a Φ -polytope. Assume on the contrary that $\Delta_{\mathcal{M}}$ is not a Φ -polytope. Then there exists an edge l with vertices $v_1 \cdot \nu$ and $v_2 \cdot \nu$ that is not parallel to any root. Consider a linear function $f: V \rightarrow \mathbb{R}$ that is constant on l and takes smaller values on the other points of $\Delta_{\mathcal{M}}$.

Since the edge l is not parallel to any root, we may assume that f is not vanishing on any root in Φ . Accordingly, there is a unique simple system of roots $\tilde{\alpha}_1, \dots, \tilde{\alpha}_{n-1}$ such that $f(\tilde{\alpha}_i) > 0$ for $i = 1, \dots, n-1$. Moreover, since the group S_n acts transitively on the set of all simple root systems, there exists an element $u \in S_n$ which sends $\{\mathbf{e}_i - \mathbf{e}_{i+1} \mid i = 1, \dots, n-1\}$ to $\{\tilde{\alpha}_1, \dots, \tilde{\alpha}_{n-1}\}$.

For any $w \in \mathcal{M} \setminus \{v_1\}$ we have $f(w \cdot \nu) \leq f(v_1 \cdot \nu)$ and the vector $w \cdot \nu - v_1 \cdot \nu$ has at least one negative coefficient with respect to $\{\tilde{\alpha}_1, \dots, \tilde{\alpha}_{n-1}\} = \{\mathbf{e}_{u(i)} - \mathbf{e}_{u(i+1)} \mid i = 1, \dots, n-1\}$. Therefore, we have $v_1 \not\leq^u w$ and this implies $v_1 \not\leq^u w$ by Lemma 2.16. Accordingly, by the Maximality Property, v_1 must be the maximal element with respect to the u -shifted order. However, by the similar argument, one can show that v_2 is also the maximal element with respect to the u -shifted order. This contradicts the Maximality Property.

Now we prove that if $\Delta_{\mathcal{M}}$ is a Φ -polytope, then $\mathcal{M} \subset S_n$ is a Coxeter matroid. Assume on the contrary that \mathcal{M} is not a Coxeter matroid. Then there are (at least) two maximal elements v_1 and

v_2 in \mathcal{M} with respect to \leq^u for some $u \in S_n$. Let $x_1 \cdot \nu, \dots, x_\ell \cdot \nu$ be the vertices of $\Delta_{\mathcal{M}}$ adjacent to the vertex $v_1 \cdot \nu$ and $y_1 \cdot \nu, \dots, y_m \cdot \nu$ the vertices adjacent to the vertex $v_2 \cdot \nu$. Set

$$(2.11) \quad \beta_p = x_p \cdot \nu - v_1 \cdot \nu \quad (p = 1, \dots, \ell) \quad \text{and} \quad \gamma_q = y_q \cdot \nu - v_2 \cdot \nu \quad (q = 1, \dots, m).$$

Since $\Delta_{\mathcal{M}}$ is a Φ -polytope, each β_p or γ_q is parallel to a root in Φ , and therefore either $\preceq^u \mathbf{0}$ or $\succeq^u \mathbf{0}$.

Suppose that $\beta_p \preceq^u \mathbf{0}$ for all p . Then $\Delta_{\mathcal{M}}$ is contained in the cone $\{\chi \in V \mid \chi \preceq^u v_1 \cdot \nu\}$. Therefore $z \cdot \nu \preceq^u v_1 \cdot \nu$ for all $z \in \mathcal{M}$, in particular, $v_2 \cdot \nu \preceq^u v_1 \cdot \nu$. Similarly, if $\gamma_q \preceq^u \mathbf{0}$ for all q , then $v_1 \cdot \nu \preceq^u v_2 \cdot \nu$. Hence, if $\beta_p \preceq^u \mathbf{0}$ and $\gamma_q \preceq^u \mathbf{0}$ for all p and q , then $v_1 \cdot \nu = v_2 \cdot \nu$ and hence $v_1 = v_2$, a contradiction.

Thus, some β_p or γ_q is $\succeq^u \mathbf{0}$. Suppose that $\beta_p \succeq^u \mathbf{0}$ for some p , i.e.,

$$x_p \cdot \nu \succeq^u v_1 \cdot \nu.$$

Then $v_1 \leq^u x_p$ by Lemma 2.17. This contradicts the maximality of v_1 because $v_1 \neq x_p$ and $x_p \in \mathcal{M}$. The same argument works when $\gamma_q \succeq^u \mathbf{0}$ for some q , so we deduce a contradiction in any case. Thus \mathcal{M} is a Coxeter matroid. \square

Example 2.18. Using Theorem 2.15, one can check that subsets $\{123, 213, 132, 312\}$ and $\{231, 321\}$ of S_3 are Coxeter matroids while a subset $\{213, 132\}$ of S_3 is not a Coxeter matroid since the edge joining the vertices $s_1 \cdot \nu$ and $s_2 \cdot \nu$ is not parallel to any root in $\Phi = \{\pm\alpha_1, \pm\alpha_2, \pm(\alpha_1 + \alpha_2)\}$ (see Figure 2 and Example 2.13).




FIGURE 2. Examples of $\Delta_{\mathcal{M}}$

By Theorem 2.15, torus orbit closures provide Coxeter matroids. A Coxeter matroid \mathcal{M} of W is said to be *representable* (over \mathbb{C}) if \mathcal{M} can be realized as the T -fixed point set of a T -orbit closure in the flag variety G/B , that is, there exists a point $x \in G/B$ such that $\mathcal{M} = \overline{(T \cdot x)}^T$, where $(G/B)^T$ is identified with W . See [17, §1.7.5, §3.6.2, §3.10.3]. A computer check shows that any Coxeter matroid of S_n is representable when $n \leq 4$. On the other hand, there exists a non-representable Coxeter matroid of S_n when $n = 7$, which we explain in the following.

Let

$$A = \{\{1, 2, 4\}, \{1, 3, 5\}, \{1, 6, 7\}, \{2, 3, 6\}, \{2, 5, 7\}, \{3, 4, 7\}, \{4, 5, 6\}\}$$

be a collection of subsets of $\{1, 2, \dots, 7\}$. Each subset corresponds to one of the six lines or the circle in Figure 3. Define a subset \mathcal{M} of S_7 by

$$\mathcal{M} = \{w \in S_7 \mid \{w(1), w(2), w(3)\} \notin A\}.$$

Then \mathcal{M} is a Coxeter matroid (see [67]).




FIGURE 3. The Fano plane

Suppose that \mathcal{M} is the T -fixed point set of a T -orbit closure in $\mathrm{GL}_7(\mathbb{C})/B$. Then, by Proposition 2.6, there is an element x in $\mathrm{GL}_7(\mathbb{C})/B$ such that

$$\mathcal{M} = \{w \in S_7 \mid \{w(1), \dots, w(d)\} \uparrow \in I_d(x) \text{ for all } 1 \leq d \leq 7\}.$$

However, it is shown in [90, §16] and easy to check that there is no 7×3 matrix of rank 3 whose three rows $v_{j_1}, v_{j_2}, v_{j_3}$ are linearly independent if and only if $\{j_1, j_2, j_3\} \notin A$. Therefore, \mathcal{M} cannot be obtained as the T -fixed point set of any T -orbit closure in $\mathrm{GL}_7(\mathbb{C})/B$.

2.5. Description of the fan of a torus orbit closure. For a point $x \in G/B$, the closure $Y := \overline{T \cdot x}$ of its T -orbit $T \cdot x$ is a (possibly non-normal) toric variety in G/B . Although Proposition 2.6 provides how to find the fixed point set Y^T and the convexity theorem leads us to the moment polytope $\mu(Y)$, it is hard to describe the fan of Y from $\mu(Y)$. In this subsection and the next subsection, we identify $(G/B)^T = W$ for simplicity, so $Y^T \subset W$. We define a retraction \mathcal{R}_Y^g (called geometric retraction) of the Weyl group W of G onto the T -fixed point set $Y^T \subset W$ of Y by using the Orbit-Cone correspondence in toric variety and describe the fan of Y using the retraction.

We think of the Lie algebra $\mathfrak{t}_{\mathbb{R}}$ of the maximal compact torus of T as $\mathrm{Hom}(\mathbb{C}^*, T) \otimes \mathbb{R}$ and the lattice $\mathfrak{t}_{\mathbb{Z}}$ of $\mathfrak{t}_{\mathbb{R}}$ as $\mathrm{Hom}(\mathbb{C}^*, T)$, where $\mathrm{Hom}(\mathbb{C}^*, T)$ denotes the group of algebraic homomorphisms from $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ to T , i.e., the one parameter subgroup of T (see [28, §1]). The vector space $\mathfrak{t}_{\mathbb{R}}^*$ dual to $\mathfrak{t}_{\mathbb{R}}$ can be thought of as $\mathrm{Hom}(T, \mathbb{C}^*) \otimes \mathbb{R}$ and the set Φ of roots of G is a finite subset of $\mathrm{Hom}(T, \mathbb{C}^*) = \mathfrak{t}_{\mathbb{Z}}^*$. The Weyl group W of G acts on $\mathfrak{t}_{\mathbb{R}}$ as the adjoint action and on its dual space $\mathfrak{t}_{\mathbb{R}}^*$ as the coadjoint action, i.e.,

$$(w \cdot f)(x) := f(\mathrm{Ad}_{w^{-1}}(x)) \quad \text{for } w \in W, f \in \mathfrak{t}_{\mathbb{R}}^* \text{ and } x \in \mathfrak{t}_{\mathbb{R}}.$$

The Borel subgroup B determines the set Φ^+ of positive roots and the tangent space $T_{uB}(G/B)$ of G/B at a T -fixed point $u \in W$ decomposes as follows:

$$(2.12) \quad T_{uB}(G/B) = \bigoplus_{\alpha \in \Phi^+} \mathfrak{g}_{-u(\alpha)} \quad \text{as } T\text{-modules for } u \in W,$$

where \mathfrak{g}_{α} denotes the root space of α , which is the eigenspace of \mathfrak{g} for $\alpha \in \Phi$ (see, for example, [49, §3]). We recall that for $G = \mathrm{GL}_n(\mathbb{C})$, the set Φ of roots is given by

$$\Phi = \{\mathbf{e}_i - \mathbf{e}_j \mid 1 \leq i \neq j \leq n\}.$$

Here, $\mathbf{e}_i \in \mathfrak{t}_{\mathbb{R}}^*$ is the linear functional sending $(a_1, \dots, a_n) \in \mathfrak{t}_{\mathbb{R}}$ to a_i for $i = 1, \dots, n$. For each root $\mathbf{e}_i - \mathbf{e}_j$, the root space $\mathfrak{g}_{\mathbf{e}_i - \mathbf{e}_j}$ is spanned by the matrix E_{ij} , which is the $n \times n$ matrix with 1 at the (i, j) entry and 0 otherwise. Moreover, the set $\Phi^+ = \{\mathbf{e}_i - \mathbf{e}_j \mid 1 \leq i < j \leq n\}$ of positive roots corresponds to the Borel subgroup B consisting of upper triangular matrices in G . The set Δ of simple roots is $\Delta = \{\mathbf{e}_i - \mathbf{e}_{i+1} \mid 1 \leq i \leq n-1\}$. See, for example, [43] for more details.

For each $u \in W$, define

$$C(u) := \{\lambda \in \mathfrak{t}_{\mathbb{R}} \mid \langle u(\alpha), \lambda \rangle \leq 0 \text{ for all simple roots } \alpha\}$$

where $\langle \cdot, \cdot \rangle$ denotes the natural pairing between $\mathfrak{t}_{\mathbb{R}}^*$ and $\mathfrak{t}_{\mathbb{R}}$. The interiors $\mathrm{Int}(C(u))$ of the cones $C(u)$ above form the Weyl chambers. The identity in (2.12) implies that T -weights on the tangent

spaces at fixed points are given by roots. This implies that if $\lambda \in \mathfrak{t}_{\mathbb{Z}}$ satisfies $\langle \alpha, \lambda \rangle \neq 0$ for any $\alpha \in \Phi$, then

$$(2.13) \quad (G/B)^{\lambda(\mathbb{C}^*)} = (G/B)^T.$$

Example 2.19. Let $G = \mathrm{GL}_3(\mathbb{C})$ and $u = 231 \in S_3$. Following the convention that we adopt, we have two simple roots, $\alpha_1 = \mathbf{e}_1 - \mathbf{e}_2$ and $\alpha_2 = \mathbf{e}_2 - \mathbf{e}_3$. Then we have

$$u(\alpha_1) = \mathbf{e}_{u(1)} - \mathbf{e}_{u(2)} = \mathbf{e}_2 - \mathbf{e}_3, \quad u(\alpha_2) = \mathbf{e}_{u(2)} - \mathbf{e}_{u(3)} = \mathbf{e}_3 - \mathbf{e}_1.$$

Accordingly, we obtain

$$\begin{aligned} C(u) &= \{\lambda = (\lambda_1, \lambda_2, \lambda_3) \mid \langle \mathbf{e}_2 - \mathbf{e}_3, \lambda \rangle \leq 0, \langle \mathbf{e}_3 - \mathbf{e}_1, \lambda \rangle \leq 0\} \\ &= \{\lambda = (\lambda_1, \lambda_2, \lambda_3) \mid \lambda_2 \leq \lambda_3 \leq \lambda_1\}. \end{aligned}$$

For each $w \in W = (G/B)^T$, we choose an element $\lambda_w \in \mathrm{Int}(C(w)) \cap \mathfrak{t}_{\mathbb{Z}}$ and define

$$S_w := \left\{ x \in G/B \mid \lim_{t \rightarrow 0} \lambda_w(t) \cdot x = w \right\},$$

which is independent of the choice of λ_w . Then S_w is a T -invariant affine open subset of G/B and isomorphic to $T_{wB}(G/B)$ as a T -variety (see [13]).

Proposition 2.20 ([67, Proposition 3.1]). *Let x be a point of G/B and $Y = \overline{T \cdot x}$. For any $u \in W$ and $\lambda_u \in \mathrm{Int}(C(u)) \cap \mathfrak{t}_{\mathbb{Z}}$, the limit point $\lim_{t \rightarrow 0} \lambda_u(t) \cdot x$ is an element of Y^T depending only on u and Y . Furthermore, if $u \in Y^T$, then $\lim_{t \rightarrow 0} \lambda_u(t) \cdot x = u$.*

Proof. Since Y is closed, the limit point $\lim_{t \rightarrow 0} \lambda_u(t) \cdot x$ belongs to Y and clearly remains fixed under the action of $\lambda_u(\mathbb{C}^*)$. Therefore, the limit point is indeed in Y^T by (2.13). Denote the limit point by w . Since $\lambda_u(t) \cdot x \in S_w$ and S_w is T -invariant, x belongs to S_w . Moreover, because S_w is isomorphic to $T_{wB}(G/B)$ as a T -variety, it follows from (2.12) that $\lim_{t \rightarrow 0} \lambda_u(t) \cdot x$ is independent of the choice of $\lambda_u \in \mathrm{Int}(C(u)) \cap \mathfrak{t}_{\mathbb{Z}}$.

If $u \in Y^T$, then x belongs to S_u because otherwise $Y = \overline{T \cdot x}$ does not contain u (note that S_u is a T -invariant open subset of G/B). Therefore, we obtain $\lim_{t \rightarrow 0} \lambda_u(t) \cdot x = u$. \square

By Proposition 2.20, the map $\mathcal{R}_Y^g: W \rightarrow Y^T \subset W$ defined by

$$(2.14) \quad \mathcal{R}_Y^g(u) := \lim_{t \rightarrow 0} \lambda_u(t) \cdot x$$

is a retraction of W onto Y^T , which we call a *geometric retraction*.

Example 2.21. Take a point $x = \begin{pmatrix} \alpha & 1 & 0 \\ \beta & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} B \in \mathrm{GL}_3(\mathbb{C})/B$, where $\alpha, \beta \in \mathbb{C}^*$. For $Y = \overline{T \cdot x}$, we have $Y^T = \{123, 132, 213, 312\}$ as observed in Example 2.7. Choose an element $\lambda = (\lambda_1, \lambda_2, \lambda_3) \in \mathrm{Int}(C(231)) \cap \mathfrak{t}_{\mathbb{Z}}$, that is, $\lambda_2 < \lambda_3 < \lambda_1$. Since

$$(2.15) \quad x = \begin{pmatrix} \alpha\beta^{-1} & 1 & 0 \\ 1 & 0 & 1 \\ \beta^{-1} & 0 & 0 \end{pmatrix} B = \begin{pmatrix} \alpha\beta^{-1} & 1 & -\alpha\beta^{-1} \\ 1 & 0 & 0 \\ \beta^{-1} & 0 & -\beta^{-1} \end{pmatrix} B = \begin{pmatrix} \alpha\beta^{-1} & 1 & 0 \\ 1 & 0 & 0 \\ \beta^{-1} & 0 & 1 \end{pmatrix} B,$$

we have

$$\begin{aligned} \lambda(t) \cdot x &= \begin{pmatrix} t^{\lambda_1} & 0 & 0 \\ 0 & t^{\lambda_2} & 0 \\ 0 & 0 & t^{\lambda_3} \end{pmatrix} \cdot \begin{pmatrix} \alpha\beta^{-1} & 1 & 0 \\ 1 & 0 & 0 \\ \beta^{-1} & 0 & 1 \end{pmatrix} B \\ &= \begin{pmatrix} t^{\lambda_1} \alpha\beta^{-1} & t^{\lambda_1} & 0 \\ t^{\lambda_2} & 0 & 0 \\ t^{\lambda_3} \beta^{-1} & 0 & t^{\lambda_3} \end{pmatrix} B \\ &= \begin{pmatrix} t^{\lambda_1 - \lambda_2} \alpha\beta^{-1} & 1 & 0 \\ 1 & 0 & 0 \\ t^{\lambda_3 - \lambda_2} \beta^{-1} & 0 & 1 \end{pmatrix} B \xrightarrow{t \rightarrow 0} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} B \quad (\because \lambda_2 < \lambda_3 < \lambda_1). \end{aligned}$$

Therefore, we get $\mathcal{R}_Y^g(231) = 213$. By a similar computation, we obtain Table 1.

| u | 123 | 213 | 231 | 321 | 312 | 132 |
|------------------------------|-----|-----|-----|-----|-----|-----|
| $\mathcal{R}_Y^g(u)$ for Y | 123 | 213 | 213 | 312 | 312 | 132 |

 TABLE 1. $\mathcal{R}_Y^g(u)$ for $Y = \overline{T \cdot x}$ in Example 2.21.

The following corollary follows from Proposition 2.20 and the Orbit-Cone correspondence of toric varieties (see [28, Proposition 3.2.2 & Theorem 3.A.5]).

Corollary 2.22 ([63, Corollary 3.7]). *The maximal cone $C_Y(y)$ corresponding to $y \in Y^T$ in the fan of (the normalization of) Y is given by $\bigcup_{u \in (\mathcal{R}_Y^g)^{-1}(y)} C(u)$.*



Remark 2.23 ([67, Remark 3.4]). (1) Since the action of T on Y is not effective, the ambient space of the fan of Y is the quotient of $\mathfrak{t}_{\mathbb{R}}$ by the subspace $\text{Hom}(\mathbb{C}^*, T_Y) \otimes \mathbb{R}$, where T_Y is the toral subgroup of T which fixes Y pointwise. Therefore, to be precise, we need to project the cones $C_Y(y)$ to this quotient space in the corollary above.

(2) When G is of type A_n , D_4 , or B_2 , every T -orbit closure in G/B is normal ([25, Proposition 4.8]) while when $G = G_2$, non-normal torus orbit closures exist (see [25, Example 6.1] and references therein).

Example 2.24. Let Y be the torus orbit closure in Example 2.21. Corollary 2.22 together with Table 1 shows that the fan of Y consists of four maximal cones:

$$C(123), \quad C(132), \quad C(213) \cup C(231), \quad C(312) \cup C(321).$$

See Figure 5. Here, the ambient space of the fan of Y is the quotient space $\mathbb{R}^3/\mathbb{R}(1, 1, 1)$ and the identification $\mathbb{R}^3/\mathbb{R}(1, 1, 1) \xrightarrow{\cong} \mathbb{R}^2$ given by $[a_1, a_2, a_3] \mapsto (a_1 - a_2, a_2 - a_3)$ is used in Figures 4 and 5. For instance, the cone $C(213)$ consists of points $[a_1, a_2, a_3]$ satisfying $a_2 \leq a_1 \leq a_3$, so that it corresponds to the set of points $\{(b_1, b_2) \in \mathbb{R}^2 \mid b_1 \geq 0, b_1 + b_2 \leq 0\}$ under the identification.


 FIGURE 4. Cones $C(v)$ for $v \in S_3$

 FIGURE 5. The fan of Y

Now we reformulate the geometric retraction \mathcal{R}_Y^g using a Bruhat decomposition of G/B . It makes the meaning of the geometric retraction more transparent.

For $u \in W$, we set $B_u := uB^-u^{-1}$, where B^- is the opposite Borel subgroup w_0Bw_0 and w_0 is the longest element of W . The Lie algebra of the Borel subgroup B_u is given as follows:

$$\text{Lie}(B_u) = \mathfrak{t} \oplus \bigoplus_{\alpha \in \Phi^+} \mathfrak{g}_{-u(\alpha)}.$$

With respect to B_u , we obtain the following Bruhat decomposition:

$$G/B = \bigsqcup_{w \in W} B_u \cdot wB/B,$$

which follows from the Bruhat decomposition $G/B = \bigsqcup_{w \in W} BwB/B$ with respect to B (see [82, §8.3] for Bruhat decompositions). Indeed, since $G = uw_0G$ and $G/B = \bigsqcup_{w \in W} Bw_0u^{-1}wB/B$, we

have

$$G/B = uw_0G/B = \bigsqcup_{w \in W} uw_0Bw_0u^{-1}wB/B = \bigsqcup_{w \in W} B_uwB/B.$$

We set

$$(2.16) \quad A_w^u := B_u \cdot wB/B = u \cdot B^-u^{-1}wB/B.$$

Note that $A_w^{w_0}$ is the Schubert cell BwB/B and A_w^e is the opposite Schubert cell B^-wB/B , where e denotes the identity element of W . Similarly to (opposite) Schubert cells, we have

$$(2.17) \quad \overline{A_w^u} = \bigsqcup_{w \leq^u v \leq^u uw_0} A_v^u \quad \text{and hence} \quad (\overline{A_w^u})^T = \{v \in W \mid w \leq^u v \leq^u uw_0\}.$$

Indeed, since $\overline{B^-wB/B}$ is the disjoint union of opposite Schubert cells indexed by elements $z \in W$ satisfying $w \leq z \leq w_0$, it follows from (2.16) that

$$\begin{aligned} \overline{A_w^u} &= u \cdot \overline{B^-u^{-1}wB/B} \\ &= u \cdot \bigsqcup_{u^{-1}w \leq z \leq w_0} B^-zB/B \\ &= \bigsqcup_{u^{-1}w \leq u^{-1}(uz) \leq u^{-1}(uw_0)} u \cdot B^-u^{-1}(uz)B/B \\ &= \bigsqcup_{w \leq^u v \leq^u uw_0} A_v^u, \end{aligned}$$

which shows (2.17).

Proposition 2.25 ([67, Proposition 3.5]). *Let x be a point of G/B and $Y = \overline{T \cdot x}$. Then $x \in A_w^u$ if and only if $\mathcal{R}_Y^g(u) = w$.*

Example 2.26. Take $u = 231 \in S_3$. Then the corresponding Borel subgroup B_u is given by

$$B_u = uB^-u^{-1} = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} \star & 0 & 0 \\ \star & \star & 0 \\ \star & \star & \star \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} = \begin{pmatrix} \star & \star & \star \\ 0 & \star & 0 \\ 0 & \star & \star \end{pmatrix}.$$

Here, $\star \in \mathbb{C}$ and $\star \in \mathbb{C}^*$. For $w = 213 \in S_3$, the cell $A_w^u = A_{213}^{231}$ is given by

$$B_uwB/B = \begin{pmatrix} \star & \star & \star \\ 0 & \star & 0 \\ 0 & \star & \star \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} B = \begin{pmatrix} \star & \star & \star \\ \star & 0 & 0 \\ \star & 0 & \star \end{pmatrix} B = \begin{pmatrix} \star & 1 & 0 \\ 1 & 0 & 0 \\ \star & 0 & 1 \end{pmatrix} B.$$

Therefore, $x = \begin{pmatrix} \alpha & 1 & 0 \\ \beta & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} B \in \text{GL}_3(\mathbb{C})/B$ in Example 2.7 is contained in A_{213}^{231} by (2.15). On

the other hand, $\mathcal{R}_Y^g(231) = 213$ for $Y = \overline{T \cdot x}$ as observed in Example 2.7.

2.6. Retractions and metric on finite Coxeter groups. For a T -orbit closure $Y = \overline{T \cdot x}$ in the flag variety G/B , we defined the geometric retraction

$$\mathcal{R}_Y^g: W \rightarrow Y^T \subset W$$

in (2.14) by looking at the limit points of the trajectory of x by 1-parameter subgroups. Note that Y^T is a Coxeter matroid by Theorem 2.15.

In general, for a Coxeter matroid \mathcal{M} of a Coxeter group W , there is a unique \leq^u -minimal element in \mathcal{M} for any element u of W (see Remark 2.12). If we denote the minimal element by $\mathcal{R}_{\mathcal{M}}^m(u)$, then we obtain a map

$$\mathcal{R}_{\mathcal{M}}^m: W \rightarrow \mathcal{M} \subset W.$$

One can easily check that $\mathcal{R}_{\mathcal{M}}^m$ is a retraction on W and we call $\mathcal{R}_{\mathcal{M}}^m$ a *matroid retraction*.⁴

⁴In [17], a map $\mu: W \rightarrow W$ satisfying the inequality $\mu(u) \leq^v \mu(v)$ for all $u, v \in \mathcal{M}$ is called a *matroid map*. For a Coxeter matroid \mathcal{M} , the map $W \rightarrow W$ sending u to the \leq^u -maximal element of \mathcal{M} is a matroid map. Note that our matroid retraction satisfies the opposite inequality $\mathcal{R}_{\mathcal{M}}^m(u) \geq^v \mathcal{R}_{\mathcal{M}}^m(v)$ for all $u, v \in \mathcal{M}$.

Theorem 2.27 ([67, Theorem 3.7]). *Let G be a semisimple algebraic group over \mathbb{C} , B a Borel subgroup of G , and T a maximal torus of G contained in B . Then $\mathcal{R}_Y^g = \mathcal{R}_{Y^T}^m$ for any T -orbit closure Y in G/B .*

Proof. Let x be a point of G/B such that $Y = \overline{T \cdot x}$. For $u \in W$, let $\mathcal{R}_Y^g(u) = w$. By Proposition 2.25, $T \cdot x \subseteq A_w^u$ and hence $Y = \overline{T \cdot x} \subseteq \overline{A_w^u}$. Therefore, we have

$$Y^T \subset \overline{A_w^u}^T = \{v \in W \mid w \leq^u v \leq^u uw_0\},$$

where the equation above follows from (2.17). Hence, w is the unique \leq^u -minimal element in Y^T , proving $\mathcal{R}_Y^g = \mathcal{R}_{Y^T}^m$. \square

A finite Coxeter group W has a metric d defined by

$$d(v, w) := \ell(v^{-1}w) = \ell(w^{-1}v) \quad \text{for } v, w \in W$$

where $\ell(\cdot)$ denotes the length function on W . Note that the metric d is invariant under the left multiplication of W . For a subset \mathcal{M} of W , we define

$$d(v, \mathcal{M}) := \min\{d(v, w) \mid w \in \mathcal{M}\}.$$

The metric d can be interpreted in terms of the W -permutohedron Δ_W as follows. As mentioned in Subsection 2.4, the W -permutohedron Δ_W is the convex hull of the W -orbit of a generic point ν in the vector space V , and two vertices $v \cdot \nu$ and $w \cdot \nu$ are joined by an edge in Δ_W if and only if $v^{-1}w$ is a generator of W . Therefore, if we identify $w \cdot \nu$ with $w \in W$, then the distance $d(v, w)$ can be thought of as the minimum length of the paths in Δ_W connecting v and w through edges of Δ_W . In other words, the metric d is the graph metric on the graph obtained as the 1-skeleton of Δ_W . For example, if $W = S_4$ and $(v, w) = (1243, 3214)$, then $v^{-1}w = 4213$ and hence $d(v, w) = \ell(4213) = 4$. Figure 6 shows a minimal-length path joining v and w in Δ_{S_4} . In Figure 6, the vertex $v \cdot \nu$ is labeled by v for each $v \in S_4$.




FIGURE 6. A minimal-length path between 1243 and 3214 in Δ_{S_4}

For every subset \mathcal{M} of W and every $u \in W$, there exists an element $y \in \mathcal{M}$ such that $d(u, y) = d(u, \mathcal{M})$ since \mathcal{M} is a finite set, but such an element y may not be unique. However, the following proposition says that such an element y is unique if \mathcal{M} is a Coxeter matroid.

Proposition 2.28 ([67, Proposition 2.6]). *If \mathcal{M} is a Coxeter matroid of a finite Coxeter group W , then $q = \mathcal{R}_{\mathcal{M}}^m(u)$ is the unique element satisfying $d(u, q) = d(u, \mathcal{M})$.*

Proof. First we remark that if $v <^u w$, then $d(u, v) < d(u, w)$ for $u, v, w \in W$. Indeed, this is shown by the following observation:

$$\begin{aligned} v <^u w &\iff u^{-1}v < u^{-1}w \\ &\implies d(e, u^{-1}v) < d(e, u^{-1}w) \\ &\iff d(u, v) < d(u, w), \end{aligned}$$

where e denotes the identity element of W and the last equivalence follows from the invariance of the metric d under the left multiplication of W .

Since $\mathcal{R}_{\mathcal{M}}^m(u)$ is the unique \leq^u -minimal element in \mathcal{M} , it follows from the above observation that $d(u, \mathcal{R}_{\mathcal{M}}^m(u)) \leq d(u, w)$ for every $w \in \mathcal{M}$ and the equality holds only when $w = \mathcal{R}_{\mathcal{M}}^m(u)$. Hence $\mathcal{R}_{\mathcal{M}}^m(u)$ is the unique element in W closest to u . This proves the proposition. \square

There is an algorithm to find $\mathcal{R}_{\mathcal{M}}^m(u)$ when W is a Weyl group of classical Lie type. The Weyl group W of classical Lie type is of the following form:

$$W = \begin{cases} S_n & \text{if } W \text{ is of type } A_{n-1}, \\ (\mathbb{Z}/2\mathbb{Z})^n \rtimes S_n & \text{if } W \text{ is of type } B_n \text{ or } C_n, \\ (\mathbb{Z}/2\mathbb{Z})^{n-1} \rtimes S_n & \text{if } W \text{ is of type } D_n. \end{cases}$$

We denote the set $\{\bar{1}, \dots, \bar{n}\}$ by $[\bar{n}]$ and regard $\bar{i} = i$. In each type, we will use one-line notation for $u \in W$, i.e.,

$$u = u(1)u(2) \cdots u(n)$$

where $u(i) \in [n] \cup [\bar{n}]$ and $u(1)u(2) \cdots u(n)$ is a permutation on $[n]$ if we forget the bars. There is no bar in type A and the number of bars in $u(1), \dots, u(n)$ is even (possibly zero) in type D . In types B , C and D , we have $u(\bar{i}) = \overline{u(i)}$.

For $u \in W$, we define a linear order \prec^u on the set $[n] \cup [\bar{n}]$ by

$$u(1) \prec^u \cdots \prec^u u(n) \prec^u u(\bar{n}) \prec^u \cdots \prec^u u(\bar{1}).$$

This induces a u -lexicographic order \prec_{lex}^u on the set of words of length n in the alphabet $[n] \cup [\bar{n}]$. Then we obtain a linear order \prec^u on W , where $v \prec^u w$ if and only if $v(1) \cdots v(n) \prec_{\text{lex}}^u w(1) \cdots w(n)$. Note that u is the minimal element of W with respect to \prec^u .

Definition 2.29. Let W be a Weyl group of classical Lie type and \mathcal{M} an *arbitrary* subset of W . For each $u \in W$, we define $\mathcal{R}_{\mathcal{M}}^a(u)$ as the u -minimal element of \mathcal{M} with respect to the order \prec^u . Then the map

$$\mathcal{R}_{\mathcal{M}}^a : W \rightarrow \mathcal{M} (\subset W)$$

is a retraction of W onto \mathcal{M} , which we call an *algebraic retraction*.

Example 2.30. (1) We take a subset $\mathcal{M} = \{123, 132, 213, 312\}$ of S_3 . For $u = 231$, we have a linear order $2 \prec^u 3 \prec^u 1$. Since

$$213 \prec_{\text{lex}}^u 312 \prec_{\text{lex}}^u 123 \prec_{\text{lex}}^u 132,$$

we have $\mathcal{R}_{\mathcal{M}}^a(231) = 213$.

(2) We take a subset $\mathcal{M} = \{1\bar{4}23, 14\bar{3}\bar{2}, 2413, \bar{3}\bar{4}1\bar{2}\}$ of $(\mathbb{Z}/2\mathbb{Z})^4 \rtimes S_4$. For $u = \bar{2}3\bar{1}4$, we have a linear order

$$\bar{2} \prec^u 3 \prec^u \bar{1} \prec^u 4 \prec^u \bar{4} \prec^u 1 \prec^u \bar{3} \prec^u 2.$$

Since

$$14\bar{3}\bar{2} \prec_{\text{lex}}^u 1\bar{4}23 \prec_{\text{lex}}^u \bar{3}\bar{4}1\bar{2} \prec_{\text{lex}}^u 2413,$$

we have $\mathcal{R}_{\mathcal{M}}^a(\bar{2}3\bar{1}4) = 14\bar{3}\bar{2}$.

The following theorem together with Proposition 2.28 shows that if \mathcal{M} is a Coxeter matroid of a Weyl group W of classical Lie type, then $\mathcal{R}_{\mathcal{M}}^a(u)$ for $u \in W$ provides the point in \mathcal{M} closest to u .

Theorem 2.31 ([67, Theorem 4.7]). *If \mathcal{M} is a Coxeter matroid of a Weyl group W of classical Lie type, then $\mathcal{R}_{\mathcal{M}}^a = \mathcal{R}_{\mathcal{M}}^m$.*

Unless \mathcal{M} is a Coxeter matroid, $\mathcal{R}_{\mathcal{M}}^a(u)$ does not necessarily provide a point in \mathcal{M} closest to u as shown in the following example.

Example 2.32. Let $W = S_4$ and $\mathcal{M} = \{1423, 2134\}$. Then \mathcal{M} is not a Coxeter matroid because the convex hull $\Delta_{\mathcal{M}}$ of \mathcal{M} , the dotted red line in Figure 7, is not a Φ -polytope since the dotted red line is not parallel to any edge of the S_4 -permutohedron. On the other hand, one can check that \mathcal{M} has the following property: for each $u \in S_4$, there is a unique element in \mathcal{M} closest to u . However,




FIGURE 7. $\mathcal{R}_{\mathcal{M}}^{\alpha}(1324)$ is not the element of \mathcal{M} closest to 1324.

if we take $u = 1324$ for instance, then $d(u, 2134) = 2$ and $d(u, 1423) = 3$, but $\mathcal{R}_{\mathcal{M}}^{\alpha}(u) = 1423$, see Figure 7. Therefore, $\mathcal{R}_{\mathcal{M}}^{\alpha}(1324)$ is not an element of \mathcal{M} closest to 1324.

As mentioned in Proposition 2.28, the following is a necessary condition for $\mathcal{M} \subset W$ to be a Coxeter matroid.

(*) For each $u \in W$, there is a *unique* $q \in \mathcal{M}$ such that $d(u, q) = d(u, \mathcal{M})$.

However, it is not a sufficient condition as is shown in Example 2.32. On the other hand, Proposition 2.28 and Theorem 2.31 show that if \mathcal{M} is a Coxeter matroid of W , then the unique element q in (*) above must be given by $\mathcal{R}_{\mathcal{M}}^{\alpha}(u)$. We ask whether these two necessary conditions are sufficient:

Problem 2.33 ([67]). Let W be a Weyl group of classical Lie type. Suppose that a subset \mathcal{M} of W satisfies the following two conditions:

- (1) for each $u \in W$, there is a unique $q \in \mathcal{M}$ such that $d(u, q) = d(u, \mathcal{M})$, and
- (2) $q = \mathcal{R}_{\mathcal{M}}^{\alpha}(u)$.

Then, is \mathcal{M} a Coxeter matroid?

For $\mathcal{M} \subset S_n$, the answer of the above problem is yes when $|\mathcal{M}| = 2$ (see [67, Proposition 4.9]) or $n \leq 4$ (by a computer check), but we do not know the answer for an arbitrary subset \mathcal{M} of S_n with $n \geq 5$.

3. GENERIC TORUS ORBIT CLOSURES IN THE FLAG VARIETY

In this section, we observe that generic torus orbit closures in the flag variety $\text{Fl}(n)$ are in fact permutohedral varieties and then discuss their topology. Their Poincaré polynomials turn out to be Eulerian polynomials. We also discuss Klyachko's theorem which describes the restriction image of the cohomology of $\text{Fl}(n)$ to that of the generic torus orbit closure in $\text{Fl}(n)$. Finally we discuss a generalization of Klyachko's theorem to Hessenberg varieties.

The image of the moment map

$$\mu: \text{Fl}(n) = G/B \rightarrow \mathbb{R}^n$$

defined in (2.6) is the permutohedron

$$\Pi_n = \text{Conv}\{(w(1), \dots, w(n)) \mid w \in S_n\}.$$

Note that Π_n is the S_n -permutohedron in Subsection 2.4. The permutohedron Π_n lies on the hyperplane

$$(3.1) \quad \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_1 + \dots + x_n = n(n+1)/2\}$$

and one can easily see that Π_n is of dimension $n - 1$.

Definition 3.1. We say that a T -orbit O in $\text{Fl}(n)$ is *generic* if $\mu(\overline{O}) = \Pi_n$, in other words if $\overline{O}^T = S_n$.

3.1. Faces of the permutohedron Π_n . We recall some facts on the faces of the permutohedron Π_n . We refer the reader to [93, Theorem 6.1], [77], [78], or [61, Section 5.A] for more detail.

The permutohedron Π_n is contained in the half space of \mathbb{R}^n defined by

$$\left\{ (x_1, \dots, x_n) \in \mathbb{R}^n \left| \sum_{i=1}^k x_i \geq \sum_{i=1}^k i \right. \right\} \quad (k = 1, \dots, n-1)$$

and the intersection of its boundary with Π_n , that is,

$$(3.2) \quad \left\{ (x_1, \dots, x_n) \in \Pi_n \left| \sum_{i=1}^k x_i = \sum_{i=1}^k i \right. \right\} \quad (k = 1, \dots, n-1)$$

is a facet of Π_n that contains the vertex $(1, \dots, n)$. This provides all the facets of Π_n which meet at the vertex $(1, \dots, n)$. Since Π_n is invariant under permutations of the coordinates of \mathbb{R}^n , it follows that Π_n is a simple polytope of dimension $n-1$ and the facets of Π_n meeting at the vertex $(w(1), \dots, w(n))$ of Π_n ($w \in S_n$) are written as

$$(3.3) \quad \left\{ (x_1, \dots, x_n) \in \Pi_n \left| \sum_{i=1}^k x_{w^{-1}(i)} = \sum_{i=1}^k i \right. \right\} \quad (k = 1, \dots, n-1).$$

The facet in (3.3) is determined by the subset $\{w^{-1}(1), \dots, w^{-1}(k)\}$ of $[n]$ and the observation above shows that there is a bijective correspondence between non-empty proper subsets of $[n]$ and facets of Π_n . Indeed, the facet of Π_n associated to a non-empty proper subset A of $[n]$ is defined by

$$(3.4) \quad F(A) := \left\{ (x_1, \dots, x_n) \in \Pi_n \left| \sum_{a \in A} x_a = \frac{|A|(|A|+1)}{2} \right. \right\}.$$

Therefore, there are $2^n - 2$ facets in Π_n and (3.3) shows that the $n-1$ facets of Π_n meeting at the vertex $(w(1), \dots, w(n))$ are $F(A)$'s with $w(A) = \{1, \dots, |A|\}$. See Figure 8.




FIGURE 8. Facets of the permutohedron Π_3

The following lemma can easily be proved.

Lemma 3.2. Let A and B be non-empty proper subsets of $[n]$. Then $F(A) \cap F(B) \neq \emptyset$ if and only if $A \subset B$ or $B \subset A$. Therefore, a codimension k face of Π_n is $\bigcap_{i=1}^d F(A_i)$ with $A_1 \subsetneq \dots \subsetneq A_k$.

Since Π_n lies on the hyperplane in (3.1) which is perpendicular to the vector $(1, \dots, 1)$ in \mathbb{R}^n , the normal fan $\Sigma(\Pi_n)$ of Π_n is defined in the quotient space $N_{\mathbb{R}} := \mathbb{R}^n / \mathbb{R}(1, \dots, 1)$, where the lattice N of $N_{\mathbb{R}}$ is the one induced from the standard lattice \mathbb{Z}^n of \mathbb{R}^n . By (3.4), the primitive (inward) normal vector δ_A to the hyperplane in (3.4) defining the facet $F(A)$ is the $(0, 1)$ -vector with 1 in the a th coordinate for each $a \in A$ and 0 otherwise. Therefore we have the following.

Lemma 3.3. *The quotient image of the vector δ_A on $N_{\mathbb{R}} = \mathbb{R}^n / \mathbb{R}(1, \dots, 1)$ is the primitive ray vector in the normal fan $\Sigma(\Pi_n)$ corresponding to the facet $F(A)$.*

The maximal cone in the fan $\Sigma(\Pi_n)$ corresponding to the vertex $(1, \dots, n)$ is spanned by the ray vectors corresponding to the $n - 1$ facets in (3.2) and those ray vectors are the projection image of the following $n - 1$ vectors on $N_{\mathbb{R}}$:

$$(1, 0, \dots, 0), \quad (1, 1, 0, \dots, 0), \quad \dots, \quad (1, \dots, 1, 0).$$

The other maximal cones in $\Sigma(\Pi_n)$ are obtained by permuting the above cone. This shows that the collection of maximal cones coincides with the (closures of) Weyl chambers in type A . The maximal cones determine the fan $\Sigma(\Pi_n)$ and the permutohedral variety Perm_n is the toric variety whose fan has the (closure of) Weyl chambers as maximal cones. Therefore, we obtain the following.

Proposition 3.4. *The maximal cones in the fan of a generic torus orbit closure in the flag variety $\text{Fl}(n)$ are the (closures of) Weyl chambers. Therefore, the closure of a generic torus orbit in the flag variety $\text{Fl}(n)$ is isomorphic to the permutohedral variety Perm_n .*

The cohomology ring of a compact smooth toric variety is explicitly described in terms of the associated fan. Applying the general result to our setting together with Lemmas 3.2 and 3.3, we obtain the following theorem.

Theorem 3.5. *The cohomology ring of the permutohedral variety Perm_n has the following presentation:*

$$H^*(\text{Perm}_n; \mathbb{Z}) = \mathbb{Z}[\tau_A \mid \emptyset \subsetneq A \subsetneq [n]] / \mathcal{I}$$

where $\deg \tau_A = 2$ and \mathcal{I} is the ideal generated by the following two types of elements:

- (1) $\tau_A \tau_B$ for $A \not\subset B$ or $B \not\subset A$,
- (2) $\sum_{p \in A} \tau_A - \sum_{q \in B} \tau_B$ for $p, q \in [n]$.

Example 3.6. Take $n = 3$. Then $H^*(\text{Perm}_3; \mathbb{Z})$ is generated by six elements

$$\tau_{\{1\}}, \tau_{\{2\}}, \tau_{\{3\}}, \tau_{\{1,2\}}, \tau_{\{1,3\}}, \tau_{\{2,3\}}$$

with relations

- (1) $\tau_{\{1\}}\tau_{\{2\}} = \tau_{\{1\}}\tau_{\{3\}} = \tau_{\{2\}}\tau_{\{3\}} = \tau_{\{1\}}\tau_{\{2,3\}} = \tau_{\{2\}}\tau_{\{1,3\}} = \tau_{\{3\}}\tau_{\{1,2\}}$
 $= \tau_{\{1,2\}}\tau_{\{1,3\}} = \tau_{\{1,3\}}\tau_{\{2,3\}} = \tau_{\{1,2\}}\tau_{\{2,3\}} = 0,$
- (2) $\tau_{\{1\}} + \tau_{\{1,2\}} + \tau_{\{1,3\}} = \tau_{\{2\}} + \tau_{\{1,2\}} + \tau_{\{2,3\}} = \tau_{\{3\}} + \tau_{\{1,3\}} + \tau_{\{2,3\}}.$

It follows that $H^2(\text{Perm}_3)$ is freely generated by four classes, say $\tau_{\{1\}}, \tau_{\{2\}}, \tau_{\{1,2\}}, \tau_{\{2,3\}}$, and $H^4(\text{Perm}_3)$ is freely generated by one class, say $\tau_{\{1\}}\tau_{\{1,2\}}$. Therefore, the Poincaré polynomial of Perm_3 is given by

$$(3.5) \quad \text{Poin}(\text{Perm}_3, t) = 1 + 4t^2 + t^4.$$

In fact, since the fan of Perm_3 is isomorphic to the fan of the toric variety obtained by blowing up $\mathbb{C}P^2$ with the standard T^2 -action at the three fixed points, Perm_3 is isomorphic to $\mathbb{C}P^2 \# 3\overline{\mathbb{C}P^2}$, the equivariant connected sum of $\mathbb{C}P^2$ with three copies of $\overline{\mathbb{C}P^2}$, where $\overline{\mathbb{C}P^2}$ stands for $\mathbb{C}P^2$ with the opposite orientation.

3.2. Eulerian polynomial. The set of *ascents* $\text{asc}(w)$ of $w \in S_n$ is defined by

$$(3.6) \quad \text{asc}(w) := \#\{i \in [n - 1] \mid w(i) < w(i + 1)\}$$

and the Eulerian polynomial $A_n(t)$ is defined by

$$A_n(t) := \sum_{w \in S_n} t^{\text{asc}(w)}.$$

Here, for a set A , we denote by $\#A$ the cardinality of A . Note that $0 \leq \text{asc}(w) \leq n-1$ for any $w \in S_n$, where the former equality is attained only when $w = [n, n-1, \dots, 1]$ and the latter equality is attained only when w is the identity. Therefore, the constant term of $A_n(t)$ is 1 and the highest degree term is t^{n-1} . Eulerian polynomials $A_n(t)$ for $n \leq 5$ are given as follows:

$$(3.7) \quad \begin{aligned} A_1(t) &= 1, & A_2(t) &= 1 + t, & A_3(t) &= 1 + 4t + t^2, \\ A_4(t) &= 1 + 11t + 11t^2 + t^3, & A_5(t) &= 1 + 26t + 66t^2 + 26t^3 + t^4. \end{aligned}$$

Remark 3.7. (1) The *descent* $\text{des}(w)$ of $w \in S_n$ is defined by

$$\text{des}(w) := \#\{i \in [n-1] \mid w(i) > w(i+1)\}$$

and it holds that

$$\sum_{w \in S_n} t^{\text{asc}(w)} = \sum_{w \in S_n} t^{\text{des}(w)}.$$

Therefore, we may define the Eulerian polynomial $A_n(t)$ using the descents.

(2) The Eulerian polynomial $A_n(t)$ was originally defined by Euler as the polynomial which appears in the numerator of the following formula:

$$\sum_{k=1}^{\infty} k^n t^k = \frac{t A_n(t)}{(1-t)^{n+1}}.$$

As easily checked, the Eulerian polynomials defined this way have the following recurrence relation

$$A_{n+1}(t) = (nt+1)A_n(t) - t(t-1)\frac{dA_n(t)}{dt}$$

while one can see that the Eulerian polynomials defined in (3.6) satisfy the same recurrence relation as above. Since $A_1(t) = 1$ in both definitions, they define the same family of polynomials.

It follows from (3.5) and (3.7) that $\text{Poin}(\text{Perm}_3, t) = A_3(t^2)$. This is not a coincidence.

Theorem 3.8 ([57, Theorem 2]). $\text{Poin}(\text{Perm}_n, t) = A_n(t^2)$ for any $n \geq 1$.

Remark 3.9. The polynomials $A_n(t)$ for $n \leq 5$ in (3.7) are palindromic and unimodal. Indeed, this holds for any $A_n(t)$ since the Poincaré duality implies the palindromicity and the hard Lefschetz theorem implies the unimodality because Perm_n is a smooth projective toric variety.

In order to give a proof of Theorem 3.8, we recall a fact on edges of the permutohedron Π_n .

Lemma 3.10. *Two vertices $(u(1), \dots, u(n))$ and $(v(1), \dots, v(n))$ of Π_n ($u, v \in S_n$) are joined by an edge of Π_n if and only if there is an adjacent transposition $s_i = (i, i+1)$ for $i \in [n-1]$ such that $v = s_i u$.*

Proof. We note that

$$(3.8) \quad (v(1), \dots, v(n)) - (1, \dots, n) = \sum_{i=1}^{n-1} \left(\sum_{k=1}^i v(k) - \sum_{k=1}^i k \right) (\mathbf{e}_i - \mathbf{e}_{i+1})$$

where $\mathbf{e}_1, \dots, \mathbf{e}_n$ denote the standard basis vectors of \mathbb{R}^n . The coefficient of $\mathbf{e}_i - \mathbf{e}_{i+1}$ in (3.8) is non-negative for any i and $v \in S_n$. Moreover, when $v = s_i$, the right hand side at (3.8) is $\mathbf{e}_i - \mathbf{e}_{i+1}$. This implies the lemma when u is the identity. The general case follows from this special case and the invariance of Π_n under permutations of the coordinates of \mathbb{R}^n . \square

Proof of Theorem 3.8. Equation (3.8) is generalized to

$$(3.9) \quad (v(1), \dots, v(n)) - (u(1), \dots, u(n)) = \sum_{i=1}^{n-1} \left(\sum_{k=1}^i v(k) - \sum_{k=1}^i u(k) \right) (\mathbf{e}_i - \mathbf{e}_{i+1}).$$

By Lemma 3.10, the endpoints of the edges emanating from the vertex $(u(1), \dots, u(n))$ are $(s_i u(1), \dots, s_i u(n))$ for $i \in [n-1]$. Set $p = u^{-1}(i)$ and $q = u^{-1}(i+1)$. Then, since $u(p) = i$ and $u(q) = i+1$ and the action of s_i from the left interchanges i and $i+1$, (3.9) reduces to

$$(3.10) \quad (s_i u(1), \dots, s_i u(n)) - (u(1), \dots, u(n)) = \mathbf{e}_p - \mathbf{e}_q = \mathbf{e}_{u^{-1}(i)} - \mathbf{e}_{u^{-1}(i+1)}.$$

Take real numbers $a_1 > \dots > a_n$ and consider a linear function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ defined by

$$f(x_1, \dots, x_n) = a_1 x_1 + \dots + a_n x_n.$$

It follows from (3.10) that

$$f(s_i u(1), \dots, s_i u(n)) - f(u(1), \dots, u(n)) = a_{u^{-1}(i)} - a_{u^{-1}(i+1)} \begin{cases} > 0 & \text{if } u^{-1}(i) < u^{-1}(i+1), \\ < 0 & \text{if } u^{-1}(i) > u^{-1}(i+1). \end{cases}$$

Recall that $\mu(wB) = (w^{-1}(1), \dots, w^{-1}(n))$ for the moment map $\mu: \text{Fl}(n) \rightarrow \mathbb{R}^n$ and we labeled the vertex $(w^{-1}(1), \dots, w^{-1}(n))$ as \mathbf{w} . Then, the observation above shows that $\text{asc}(w)$ is the number of edges emanating from the vertex \mathbf{w} and increasing with respect to the function f . Therefore, discrete Morse theory applied to the function f on the permutohedron Π_n produces a decomposition

$$\Pi_n = \bigsqcup_{w \in S_n} P_w, \quad P_w \approx (\mathbb{R}_{\geq 0})^{\text{asc}(w)}$$

where \approx means homeomorphic as a manifold with corners. Moreover, one sees that $\mu^{-1}(P_w) \approx \mathbb{C}^{\text{asc}(w)}$ for μ restricted to Perm_n and the decomposition

$$\text{Perm}_n = \mu^{-1}(\Pi_n) = \bigsqcup_{w \in S_n} \mu^{-1}(P_w)$$


gives a cell decomposition of Perm_n . Since the dimension of the cell $\mu^{-1}(P_w)$ is $2\text{asc}(w)$, we obtain

$$\mathcal{Poin}(\text{Perm}_n, t) = \sum_{w \in S_n} t^{2\text{asc}(w)} = A_n(t^2),$$

proving the theorem. □

Example 3.11. Take $n = 3$. Consider a linear function $f(x_1, x_2, x_3) = 5x_1 + 4x_2 + x_3$. The values $f(\mu(wB))$ and $\text{asc}(w)$ are given as follows. Note that the vertex $\mu(wB)$ of Π_3 is labeled by \mathbf{w} as before.

| w | 123 | 213 | 132 | 231 | 312 | 321 |
|-----------------|-----------|-----------|-----------|-----------|-----------|-----------|
| $\mu(wB)$ | (1, 2, 3) | (2, 1, 3) | (1, 3, 2) | (3, 1, 2) | (2, 3, 1) | (3, 2, 1) |
| $f(\mu(wB))$ | 16 | 17 | 19 | 21 | 23 | 24 |
| $\text{asc}(w)$ | 2 | 1 | 1 | 1 | 1 | 0 |



Accordingly, we have

$$\mathcal{Poin}(\text{Perm}_3, t) = A_3(t^2) = 1 + 4t^2 + t^4.$$

3.3. Klyachko's result.

$$\text{Fl}(n) = \{V_\bullet = (\{0\} \subset V_1 \subset V_2 \subset \dots \subset V_n = \mathbb{C}^n) \mid \dim_{\mathbb{C}} V_i = i \text{ for all } i = 1, \dots, n\},$$

there is a sequence of tautological complex vector bundles on $\text{Fl}(n)$:

$$E_1 \subset E_2 \subset \dots \subset E_n = \text{Fl}(n) \times \mathbb{C}^n$$

where

$$E_i := \{(V_\bullet, v) \in \text{Fl}(n) \times \mathbb{C}^n \mid v \in V_i\} \quad (i = 1, \dots, n).$$

We define

$$x_i := c_1(E_i/E_{i-1}) \quad \text{for } i = 1, \dots, n,$$

where $E_0 = \text{Fl}(n)$, i.e., the 0-dimensional vector bundle over $\text{Fl}(n)$ and $c_1(E)$ denotes the first Chern class of a vector bundle E . Since

$$E_n \cong \bigoplus_{i=1}^n E_i/E_{i-1}$$

and E_n is the trivial vector bundle $\text{Fl}(n) \times \mathbb{C}^n$, we have

$$1 = c(E_n) = \prod_{i=1}^n c(E_i/E_{i-1}) = \prod_{i=1}^n (1 + x_i).$$

Here, $c(E)$ denotes the total Chern class of a vector bundle E . Therefore, the i th elementary symmetric polynomial $e_i(x)$ in x_1, \dots, x_n vanishes in $H^*(\text{Fl}(n); \mathbb{Z})$ for $i = 1, \dots, n$. The following theorem shows that $H^*(\text{Fl}(n); \mathbb{Z})$ is generated by x_1, \dots, x_n as a ring and any polynomial in x_1, \dots, x_n which vanishes in $H^*(\text{Fl}(n); \mathbb{Z})$ is generated by $e_i(x)$'s.

Theorem 3.12 (Borel [16]). *Let x_1, \dots, x_n be as above. Then*

$$H^*(\text{Fl}(n); \mathbb{Z}) = \mathbb{Z}[x_1, \dots, x_n]/(e_1(x), \dots, e_n(x)) \quad \text{as a ring.}$$

We denote by Perm_n a generic torus orbit closure in the flag variety $\text{Fl}(n)$ because it is isomorphic to the permutohedral variety Perm_n . The natural action of S_n on the fan of Perm_n induces an action of S_n on Perm_n and hence on its cohomology ring.

Theorem 3.13 (Klyachko [57]). *Let $\iota: \text{Perm}_n \rightarrow \text{Fl}(n)$ be the inclusion map. Then the image of the restriction map*

$$\iota^*: H^*(\text{Fl}(n); \mathbb{Q}) \rightarrow H^*(\text{Perm}_n; \mathbb{Q})$$

agrees with the ring of S_n -invariants $H^(\text{Perm}_n; \mathbb{Q})^{S_n}$ and*

$$(3.11) \quad H^*(\text{Perm}_n; \mathbb{Q})^{S_n} = \mathbb{Q}[\omega_1, \dots, \omega_{n-1}]/(\omega_i(\omega_{i-1} - 2\omega_i + \omega_{i+1}) \mid i = 1, \dots, n-1)$$

where $\omega_i = x_1 + \dots + x_i$ and $\omega_0 = \omega_n = 0$.

Remark 3.14. (1) The cohomology ring of Perm_n is explicitly described in Theorem 3.5. The S_n -orbit of τ_A with $|A| = i$ is $\sum_{|B|=i} \tau_B$ which is equal to $x_i - x_{i+1}$.

(2) If we set $\alpha_i := x_i - x_{i+1}$, then $\omega_{i-1} - 2\omega_i + \omega_{i+1} = -\alpha_i$. Therefore, we may express the ideal in (3.11) as $(\omega_i \alpha_i \mid i = 1, \dots, n-1)$. Note that the ω_i 's correspond to the fundamental weights while α_i 's correspond to the roots in type A .

3.4. Relation to Hessenberg varieties. Theorem 3.13 is generalized to the setting of Hessenberg varieties. Given a square matrix A of size n with complex entries and a function $h: [n] \rightarrow [n]$ (called a *Hessenberg function*) satisfying

$$h(1) \leq h(2) \leq \dots \leq h(n) \quad \text{and} \quad h(j) \geq j \quad \text{for all } j \in [n],$$

the Hessenberg variety $\text{Hess}(A, h)$ is defined as

$$\text{Hess}(A, h) := \{V_\bullet \in \text{Fl}(n) \mid A(V_j) \subset V_{h(j)} \text{ for all } j \in [n]\},$$

where the matrix A is regarded as a linear transformation on \mathbb{C}^n . We notice that $\text{Hess}(A, h)$ is isomorphic to $\text{Hess}(gAg^{-1}, h)$ via the action of G on $\text{Fl}(n)$. Accordingly, for a fixed Hessenberg function h , the geometry of Hessenberg variety $\text{Hess}(A, h)$ depends only on the conjugacy class of A . We often express the Hessenberg function h as a vector $(h(1), \dots, h(n))$ by listing the values of h . When $h = (n, \dots, n)$, it is obvious from the definition that $\text{Hess}(A, h)$ is the flag variety $\text{Fl}(n)$ regardless of A .

The Hessenberg variety $\text{Hess}(S, h)$ for a square matrix S of size n with distinct eigenvalues is called *regular semisimple*. It is known that

- (1) $\text{Hess}(S, h)$ is smooth,
- (2) $\dim_{\mathbb{C}} \text{Hess}(S, h) = \sum_{j=1}^n (h(j) - j)$,
- (3) $\text{Hess}(S, h)$ is connected if and only if $h(j) \geq j + 1$ for all $j \in [n - 1]$.

The topology of $\text{Hess}(S, h)$ is independent of the choice of S and we take S to be a diagonal matrix with distinct diagonal entries in the following. Indeed, for regular semisimple matrices S and S' , the corresponding Hessenberg varieties $\text{Hess}(S, h)$ and $\text{Hess}(S', h)$ are diffeomorphic.⁵ Since S commutes with the diagonal torus T of $\text{GL}_n(\mathbb{C})$, the restricted action of T on $\text{Fl}(n)$ leaves $\text{Hess}(S, h)$ invariant. One sees that

$$(3.12) \quad \text{Hess}(S, h)^T = \text{Fl}(n)^T = S_n.$$

When $h = (2, 3, \dots, n, n)$, it follows from (2) and (3) above that $\dim_{\mathbb{C}} \text{Hess}(S, h) = n - 1$ and $\text{Hess}(S, h)$ is connected. Although $\dim_{\mathbb{C}} T = n$, the subgroup D of T consisting of scalar matrices acts on $\text{Fl}(n)$ trivially and the induced action of T/D on $\text{Hess}(S, h)$ is effective. Therefore, $\text{Hess}(S, h)$ for $h = (2, 3, \dots, n, n)$ is a smooth toric variety and hence it is a torus orbit closure in $\text{Fl}(n)$. Moreover, the orbit is generic by (3.12). Therefore, $\text{Hess}(S, h)$ for $h = (2, 3, \dots, n, n)$ is isomorphic to the permutohedral variety Perm_n .

Using the T -action on $\text{Hess}(S, h)$, Tymoczko [87] constructed an action of S_n (called *dot action*) on $H^*(\text{Hess}(S, h); \mathbb{Z})$ and when $h = (2, 3, \dots, n, n)$, this action agrees with that on $H^*(\text{Perm}_n; \mathbb{Z})$ induced from the action of S_n on Perm_n . Therefore, the following theorem is a generalization of the former part of Theorem 3.13.

Theorem 3.15 ([1, Theorem B]). *Let $\iota: \text{Hess}(S, h) \rightarrow \text{Fl}(n)$ be the inclusion map. Then the image of the restriction map*

$$\iota^*: H^*(\text{Fl}(n); \mathbb{Q}) \rightarrow H^*(\text{Hess}(S, h); \mathbb{Q})$$

agrees with the ring of S_n -invariants $H^(\text{Hess}(S, h); \mathbb{Q})^{S_n}$ for any Hessenberg function h .*

In fact, it is known from [21] that

$$(3.13) \quad H^*(\text{Hess}(S, h); \mathbb{Q})^{S_n} \cong H^*(\text{Hess}(N, h); \mathbb{Q}),$$

where N is conjugate to a nilpotent matrix with one Jordan block ($\text{Hess}(N, h)$ is called *regular nilpotent*) and an explicit ring presentation of $H^*(\text{Hess}(N, h); \mathbb{Q})$, which reduces to (3.11) when $h = (2, 3, \dots, n, n)$, is known. Therefore, Theorem 3.13 is completely generalized to the setting of Hessenberg varieties, see [1, 2, 4, 53] for more details.

Remark 3.16. It is shown in [3] that when $h = (2, 3, \dots, n)$ (so that $\text{Hess}(S, h) = \text{Perm}_n$), Theorem 3.15 and (3.13) hold with \mathbb{Z} -coefficient but (3.11) does not.

4. GENERIC TORUS ORBIT CLOSURES IN SCHUBERT VARIETIES

In this section, we consider geometric and topological properties of generic torus orbit closures in Schubert varieties with respect to the T -action from the introduction. A generic torus orbit closure Y_w in a Schubert variety X_w is not smooth in general and we discuss how to determine the smoothness of Y_w at a fixed point by considering a certain graph. We study the Poincaré polynomial of Y_w which is a generalization of the Eulerian polynomial. Moreover, we study the fan of a toric Schubert variety and discuss the classification of toric Schubert varieties. Finally, we summarize the properties of Schubert varieties of complexity one.

4.1. Generic torus orbit closures in Schubert varieties.

Definition 4.1. For $w \in S_n$, the *Schubert variety* X_w is a subvariety of G/B defined by $X_w := \overline{BwB/B}$.

The action of T on G/B leaves X_w invariant.

Definition 4.2. We say that a T -orbit O in X_w is *generic* if $\mu(\overline{O}) = \mu(X_w)$, in other words, if $\overline{O}^T = X_w^T$. We denote by Y_w the closure of a generic torus orbit in the Schubert variety X_w .

⁵For regular semisimple matrices S and S' , there is a path connecting them in the space of regular semisimple matrices and the family of regular semisimple Hessenberg varieties are fiber bundle over the path where the Hessenberg function h is fixed. Since a path is contractible, the fiber bundle is trivial, which implies the desired result.

Since $X_{w_0} = G/B$, the generic torus orbit closure Y_{w_0} in X_{w_0} is the permutohedral variety Perm_n considered in Section 3 (cf. Definition 3.1), in other words, $Y_{w_0} = \text{Perm}_n$.

We notice that a fixed point uB is contained in the Schubert variety X_w if and only if $u \leq w$ in the Bruhat order; uB is contained in the opposite Schubert variety X^w if and only if $u \geq w$ in the Bruhat order. By Lemma 2.3, we have

$$\mu(X_w) = \text{Conv}\{(u^{-1}(1), \dots, u^{-1}(n)) \mid u \leq w\} \subset \mathbb{R}^n.$$

Motivated by this fact, we provide the following definition.

Definition 4.3. For $w \in S_n$, we define a polytope \mathbf{Q}_w by

$$\mathbf{Q}_w := \text{Conv}\{(u^{-1}(1), \dots, u^{-1}(n)) \mid u \leq w\}$$

and label the vertex $(u^{-1}(1), \dots, u^{-1}(n))$ as u as before.

Therefore we have

$$\mu(X_w) = \mu(Y_w) = \mathbf{Q}_w.$$

We depict the Bruhat interval $[e, 3412] = \{u \in S_4 \mid u \leq 3412\}$ and the polytope \mathbf{Q}_{3412} in Figure 9.




FIGURE 9. The Bruhat interval $[e, 3412]$ and the polytope \mathbf{Q}_{3412}

Remark 4.4. Note that the polytope \mathbf{Q}_w is the Φ -polytope $\Delta_{[e, w^{-1}]}$ in Subsection 2.4, where $W = S_n$ and $\nu = (1, 2, \dots, n)$.

By definition, Y_w is the toric variety whose fan is the normal fan of the polytope \mathbf{Q}_w . The polytope \mathbf{Q}_w is not simple in general. For example, the polytope \mathbf{Q}_{3412} is 3-dimensional, but there are four facets meeting at the vertex 3412 (see Figure 9). At each vertex of \mathbf{Q}_w , the primitive direction vectors of the edges emanating from the vertex are root vectors, i.e., of the form $\mathbf{e}_i - \mathbf{e}_j$. Therefore, we obtain the following.

Proposition 4.5 ([63, §8]). *For $w \in S_n$, the toric variety Y_w is smooth if and only if the polytope \mathbf{Q}_w is simple.*

To each vertex of \mathbf{Q}_w we associate a graph which detects the simpleness of the vertex. For $u \leq w$, we set


$$\tilde{E}_w(u) = \{(u(i), u(j)) \mid i < j, t_{u(i), u(j)}u \leq w, |\ell(u) - \ell(t_{u(i), u(j)}u)| = 1\},$$

where $t_{a,b}$ denotes the transposition interchanging a and b .⁶ Then the digraph $\tilde{\Gamma}_w(u)$ defined by the vertex set $[n]$ and the edge set $\tilde{E}_w(u)$ is acyclic by [86, Theorem 4.19], so it has a unique transitive reduction (see [5]). Here, a *transitive reduction* of a digraph is a digraph with the same vertices and as fewer edges as possible, such that if there is a directed path connecting two vertices, then there is also such a path in the reduction. We denote the transitive reduction of $\tilde{\Gamma}_w(u)$ by $\Gamma_w(u)$ and the edge set of $\Gamma_w(u)$ by $E_w(u)$.

Example 4.6 ([69, Example 3.2]). Take $w = 3412$ and $u = 2143$. Then we have

$$\tilde{E}_w(u) = \{(1, 4), (2, 3), (2, 1), (4, 3)\}.$$

The corresponding digraph $\tilde{\Gamma}_w(u)$ and its transitive reduction $\Gamma_w(u)$ are presented as follows.



As a result, we have $E_w(u) = \{(1, 4), (2, 1), (4, 3)\}$.

Example 4.7. For the longest element w_0 , the condition $t_{u(i),u(j)}u \leq w_0$ is always satisfied. Therefore, we have

$$\tilde{E}_{w_0}(u) = \{(u(i), u(j)) \mid i < j, |\ell(u) - \ell(t_{u(i),u(j)}u)| = 1\}.$$

One can easily check that

$$\{(u(i), u(i+1)) \mid i = 1, \dots, n-1\} \subset \tilde{E}_{w_0}(u).$$

Indeed, $E_{w_0}(u) = \{(u(i), u(i+1)) \mid i = 1, \dots, n-1\}$.

Proposition 4.8 ([63, Proposition 7.7]). *For $w \in S_n$, two vertices u and v of Q_w are joined by an edge if and only if*

$$v = t_{u(i),u(j)}u (= ut_{i,j}) \quad \text{for } (u(i), u(j)) \in E_w(u).$$

The proposition above says that the elements in $E_w(u)$ correspond to the edges of Q_w emanating from the vertex u . Therefore, one can read from the graph $\Gamma_w(u)$ the number of edges emanating from the vertex u in the polytope Q_w .

Theorem 4.9 ([63, Theorem 1.2]). *The generic torus orbit closure Y_w in a Schubert variety X_w is smooth at a fixed point uB in Y_w if and only if the graph $\Gamma_w(u)$ is a forest. Therefore, Y_w is smooth if and only if $\Gamma_w(u)$ is a forest for every $u \leq w$.*

The graph $\Gamma_w(w)$ appears in a slightly different viewpoint. Indeed, it is shown in [91] that the Schubert variety X_w is locally factorial if and only if $\Gamma_w(w)$ is a forest. This motivated the authors in [18] to study the graph and they provided a necessary and sufficient condition for $\Gamma_w(w)$ to be a forest in terms of the pattern avoidance.⁷ Combining the theorem above with [18, Theorem 1.1], we obtain the following.

Proposition 4.10. *The following statements are equivalent.*

- (1) $\Gamma_w(w)$ is a forest.
- (2) w avoids the patterns 4231 and 45 $\bar{3}$ 12.
- (3) Y_w is smooth at wB .
- (4) Q_w is simple at w .

About the smoothness of Y_w , it is enough to check the smoothness at the fixed point wB as is shown in the following theorem. It was originally conjectured in [63].

⁶There is a relation between this set and the T -weights of $T_{uB}(G/B)$ which are described in [14].

⁷We say that a permutation $w \in S_n$ contains the pattern $q = q_1q_2 \cdots q_k$ if there is a k -element set of indices $1 \leq i_1 < i_2 < \cdots < i_k \leq n$ so that $w(i_r) < w(i_s)$ if and only if $q_r < q_s$ for $1 \leq r < s \leq k$. If a permutation w does not contain q , then we say that w avoids the pattern q . Similarly, w avoids the pattern 45 $\bar{3}$ 12 if every occurrence of the pattern 4512 is a subsequence of an occurrence of 45312.

Theorem 4.11 ([44, Theorem B]). *The graph $\Gamma_w(u)$ is a forest for any $u \leq w$ if $\Gamma_w(w)$ is a forest. (This is equivalent to saying that the generic torus orbit closure Y_w in the Schubert variety X_w is smooth if it is smooth at the fixed point wB , in other words, Q_w is simple if Q_w is simple at the vertex w .)*

4.2. Generalized Eulerian polynomials. In this subsection, we introduce a polynomial $A_w(t)$ for each $w \in S_n$, which agrees with the Eulerian polynomial $A_n(t)$ when $w = w_0$, and explain that the Poincaré polynomial of Y_w is given by $A_w(t^2)$. The tool we use to compute the Poincaré polynomial of Y_w is discrete Morse theory. We have the moment map

$$\mu := Y_w \rightarrow Q_w \subset \mathbb{R}^n.$$

The polytope Q_w is not necessarily simple, but we can find a linear function on \mathbb{R}^n by which we decompose Q_w into pieces homeomorphic to orthants.

Let P be a polytope in \mathbb{R}^n and $h: \mathbb{R}^n \rightarrow \mathbb{R}$ a linear function satisfying that $h(\mathbf{u}) \neq h(\mathbf{v})$ if two vertices \mathbf{u} and \mathbf{v} of P are joined by an edge in P . Then the function h gives an orientation on each edge of P , namely, we give an orientation of the edge connecting two vertices \mathbf{u} and \mathbf{v} of P by $\mathbf{u} \rightarrow \mathbf{v}$ if $h(\mathbf{u}) < h(\mathbf{v})$. For each vertex \mathbf{u} of P , we define

$$\text{asc}(\mathbf{u}) := \#\{\mathbf{u} \rightarrow \mathbf{v}\}.$$

The following is well-known for simple polytopes and is a consequence of discrete Morse theory.

Theorem 4.12 (see [69, Theorem 2.7] for example). *Let $P \subset \mathbb{R}^n$ be a lattice polytope. Suppose that there exists a linear function $h: \mathbb{R}^n \rightarrow \mathbb{R}$ such that for each vertex \mathbf{u} of P , the direction vectors of ascending edges emanating from \mathbf{u} are linearly independent and form a face $F(\mathbf{u})$ of P , that is, these edges form the set of edges emanating from \mathbf{u} in $F(\mathbf{u})$. Then the Poincaré polynomial $\text{Poin}(X_P, t)$ of the toric variety X_P whose fan is the normal fan of P is given by*

$$\text{Poin}(X_P, t) = \sum_{\mathbf{u} \in V(P)} t^{2\text{asc}(\mathbf{u})},$$

where $V(P)$ is the set of all vertices of P .

Remark 4.13. The vertex \mathbf{u} may not be simple in P but has to be simple in $F(\mathbf{u})$. Therefore the number of k -dimensional faces of P which are contained in $F(\mathbf{u})$ and contain \mathbf{u} is $\binom{\text{asc}(\mathbf{u})}{k}$. This implies that

$$\sum_{\mathbf{u} \in V(P)} (1+t)^{2\text{asc}(\mathbf{u})} = \sum_{k=0}^n f_k(P) t^k =: f_P(t),$$

where $n = \dim P$, $f_k(P)$ denotes the number of k -dimensional faces of P , and $f_n(P) = 1$. The polynomial $f_P(t)$ is called the *f-polynomial* of P and the polynomial $h_P(t) := f_P(t-1)$ is called the *h-polynomial* of P . Theorem 4.12 can be restated as

$$(4.1) \quad \text{Poin}(X_P, t) = h_P(t^2)$$

if there exists a linear function h in Theorem 4.12. When P is simple, such a linear function h always exists and (4.1) is well-known in this case. However, it is not true that the formula (4.1) holds for every polytope P . For example, if P is an octahedron, then its *f-polynomial* is

$$f_P(t) = 6 + 12t + 8t^2 + t^3.$$

Therefore

$$h_P(t^2) = 6 + 12(t^2 - 1) + 8(t^2 - 1)^2 + (t^2 - 1)^3 = 1 - t^2 + 5t^4 + t^6$$

which cannot be the Poincaré polynomial of X_P because it has a negative coefficient. In fact, the polynomial $h_P(t^2)$ is the *virtual Poincaré polynomial* of X_P . It is known that the virtual Poincaré polynomial agrees with the ordinary Poincaré polynomial for compact smooth toric varieties, see [41, Section 4.5], but Theorem 4.17 below implies that they agree for Y_w although Y_w is not necessarily smooth.

Example 4.14. Let P be the following pyramid with five vertices:

$$P = \text{Conv}\{(1, 0, 0), (0, 1, 0), (-1, 0, 0), (0, 0, -1), (0, 0, 1)\}.$$

Choose a linear function $h: \mathbb{R}^3 \rightarrow \mathbb{R}$ defined by

$$h(x_1, x_2, x_3) = -2x_1 - x_2 + 3x_3.$$

The function h gives an orientation on edges of P as displayed in Figure 10.




FIGURE 10. The pyramid with an orientation

Then, for each vertex u of P , the corresponding face $F(u)$ and the number of ascending edges are as in Table 2. Therefore, the Poincaré polynomial $\mathcal{Poin}(X_P, t)$ of X_P is given by

$$\mathcal{Poin}(X_P, t) = 1 + t^2 + 2t^4 + t^6.$$

| | | | | | |
|-----------------|----|----|---|---|---|
| $F(u)$ | | | | | |
| $h(u)$ | -2 | -1 | 1 | 2 | 3 |
| $\text{asc}(u)$ | 3 | 2 | 2 | 1 | 0 |

TABLE 2. Faces $F(u)$ and $\text{asc}(u)$ for the pyramid

On the other hand, since $(f_0(P), f_1(P), f_2(P), f_3(P)) = (5, 8, 5, 1)$ for the pyramid P , we have

$$h_P(t^2) = 5 + 8(t^2 - 1) + 5(t^2 - 1)^2 + (t^2 - 1)^3 = 1 + t^2 + 2t^4 + t^6.$$

Therefore the formula (4.1) certainly holds in this case.

We shall apply Theorem 4.12 to Q_w to find the Poincaré polynomial of Y_w . What we have to do is to find a linear function h on \mathbb{R}^n which satisfies the conditions in Theorem 4.12.

Lemma 4.15 ([69, Lemma 3.4]). *Let $h: \mathbb{R}^n \rightarrow \mathbb{R}$ be a linear function defined by the inner product with a vector $(a_1, \dots, a_n) \in \mathbb{R}^n$ with $a_1 > a_2 > \dots > a_n$. Then, for $(u(i), u(j)) \in E_w(u)$, the edge emanating from the vertex u to the vertex v in Q_w , where $v = t_{u(i), u(j)}u$, is ascending with respect to the function h if and only if $u(i) < u(j)$. Here, we consider orientations on edges of P according to the function h .*

Motivated by Lemma 4.15, we define

$$E_w(u)^+ = \{(u(i), u(j)) \in E_w(u) \mid u(i) < u(j)\},$$

$$E_w(u)^- = \{(u(i), u(j)) \in E_w(u) \mid u(i) > u(j)\}.$$

Then

$$E_w(u) = E_w(u)^+ \sqcup E_w(u)^-.$$

Note that $E_w(u)^+$ (resp. $E_w(u)^-$) corresponds to the ascending (resp. descending) edges emanating from u by Lemma 4.15 with respect to the function h . Now we set

$$a_w(u) := |E_w(u)^+|$$

and define

$$A_w(t) := \sum_{u \leq w} t^{a_w(u)}.$$

Example 4.16. By Example 4.7, we have

$$\begin{aligned} E_{w_0}(u)^+ &= \{(u(i), u(j)) \in E_{w_0}(u) \mid u(i) < u(j)\} \\ &= \{(u(i), u(i+1)) \mid u(i) < u(i+1) \text{ for } i = 1, \dots, n-1\}. \end{aligned}$$

Accordingly, $a_{w_0}(u)$ is the number of ascents in u and

$$A_{w_0}(t) = A_n(t).$$

On the other hand, $X_{w_0} = \text{Fl}(n)$ and Y_{w_0} is the permutohedral variety Perm_n . Therefore, we have

$$\text{Poin}(Y_{w_0}, t) = A_{w_0}(t^2)$$

by Theorem 3.8.

One can check that the linear function h in Lemma 4.15 satisfies the conditions in Theorem 4.12 so that we obtain the following theorem extending Theorem 3.8.

Theorem 4.17 ([69, Theorem 3.6]). *Let Y_w be the generic torus orbit closure in the Schubert variety X_w for $w \in S_n$. Then the Poincaré polynomial of Y_w is given by*

$$\text{Poin}(Y_w, t) = A_w(t^2).$$

We close this subsection by presenting the Poincaré polynomials of singular toric varieties Y_{4231} and Y_{3412} using Theorem 4.17. The moment polytopes of Y_{4231} and Y_{3412} are non-simple polytopes Q_{4231} and Q_{3412} , respectively, see Figure 11. In both polytopes, we take $h: \mathbb{R}^4 \rightarrow \mathbb{R}$ by the inner product with a vector $(12, 2, -1, -2)$. For instance, the image of the vertex $4132 = (2, 4, 3, 1)$ under h is given by

$$h(4132) = \langle (2, 4, 3, 1), (12, 2, -1, -2) \rangle = 2 \times 12 + 4 \times 2 - 3 - 2 = 27.$$

We number each vertex u as $h(\mu(uB))$. Therefore, the vertex $4132 = (2, 4, 3, 1)$ is numbered as 27 (see Figure 11(1)).




FIGURE 11. Polytopes Q_{4231} and Q_{3412}

Using this function h , we get the Poincaré polynomials of these singular generic torus orbit closures as follows:

$$\text{Poin}(Y_{4231}, t) = 1 + 7t^2 + 11t^4 + t^6,$$

$$\text{Poin}(Y_{3412}, t) = 1 + 5t^2 + 7t^4 + t^6.$$

One can easily confirm the formula (4.1) in these two cases. Indeed, the formula (4.1) holds for every Y_w , which implies

$$A_w(t) = h_{Q_w}(t) \quad \text{for every } w \in S_n.$$

Remark 4.18. (1) For a rational polytope P allowing a *retraction sequence*, the Poincaré polynomial of the toric variety X_P is the polynomial $h_P(t^2)$. We refer the reader to [10] for a precise definition of retraction sequences. It is shown in [76] that a linear function h as in Lemma 4.15 defines a retraction sequence on the polytope Q_w . However, it is not known whether all retraction sequences come from linear functions.

(2) In [77, Section 4.1], a generalized Eulerian polynomial is defined for a *simple* generalized permutohedron P and it agrees with the h -polynomial of P (see [77, Theorem 4.2]). On the other hand, our generalized Eulerian polynomial $A_w(t)$ for $w \in S_n$ agrees with the h -polynomial of a Burhat interval polytope Q_w as mentioned above, where Q_w is a generalized permutohedron but not necessarily simple.

4.3. Toric Schubert varieties. The complexity of a T -variety X is the codimension of a maximal dimensional T -orbit in X . A toric variety is of complexity 0 and in this case there is a nice relation between geometry and combinatorics (see Appendix A.1). Indeed, a toric variety has a fruitful symmetry given by the T -action and we can extract the essential geometric information (e.g., smoothness) from the corresponding combinatorial object. One can expect a similar nice relation when the complexity is small enough. In this section, we present such kind of results for Schubert varieties of complexity 0 or 1.

Definition 4.19. For $w \in S_n$, we define

$$c(w) := \dim_{\mathbb{C}} X_w - \dim_{\mathbb{C}} Y_w = \dim_{\mathbb{C}} X_w - \dim_{\mathbb{R}} Q_w = \ell(w) - \dim_{\mathbb{R}} Q_w,$$

and call $c(w)$ the *complexity* of the Schubert variety X_w (or the *complexity* of w).

When $c(w) = 0$, we have $X_w = Y_w$; so the Schubert variety X_w is a toric variety. Let $w = s_{i_1} \cdots s_{i_m}$ be a reduced decomposition of w . Then it is known from [37, 56] that the Schubert variety X_w is a toric variety if and only if i_1, \dots, i_m are distinct. Moreover, all toric Schubert varieties are Bott manifolds, which are smooth projective toric varieties whose moment map images are combinatorially equivalent to cubes. We refer the reader to Appendix A for more details on Bott manifolds. Furthermore, all toric Schubert varieties are Bott–Samelson varieties which are smooth projective varieties. We will explain them in detail in Section 4.4.

There are several equivalent conditions characterizing toric Schubert varieties as follows.

Theorem 4.20. [37, 56, 84, 85, 66] *The following statements are equivalent:*

- (0) X_w is a toric variety (i.e., of complexity zero).
- (1) X_w is a smooth toric variety.
- (2) w avoids the patterns 321 and 3412.
- (3) A reduced decomposition of w consists of distinct letters.
- (4) X_w is isomorphic to a Bott–Samelson variety.
- (5) The Bruhat interval $[e, w]$ is isomorphic to the Boolean algebra $\mathfrak{B}_{\ell(w)}$ of rank $\ell(w)$ as posets.
- (6) Q_w is combinatorially equivalent to the cube of dimension $\ell(w)$.

Here, we say that a poset P is a *Boolean algebra* if there is a set X such that P is isomorphic to the set of all subsets of X , partially ordered by inclusion.

The fan of a toric Schubert variety X_w is the normal fan of the polytope Q_w . Since Q_w is combinatorially equivalent to a cube as stated in Theorem 4.20, we shall find its normal fan by investigating the primitive direction vectors of the edges emanating from the vertices e and w of Q_w . Let $w = s_{i_1} \cdots s_{i_m}$ be a reduced decomposition of $w \in S_n$ and we assume that i_1, \dots, i_m are distinct. Then $w^{-1} = s_{i_m} \cdots s_{i_1}$ and the Bruhat interval $[e, w^{-1}]$ has m many atoms and coatoms:

$$(4.2) \quad \begin{aligned} \text{atoms: } & s_{i_k} \quad (k = 1, \dots, m), \\ \text{coatoms: } & s_{i_m} \cdots \hat{s}_{i_k} \cdots s_{i_1} \quad (k = 1, \dots, m). \end{aligned}$$

Lemma 4.21. *Let $w = s_{i_1} \cdots s_{i_m}$. For $1 \leq k \leq m$, we set*

$$a_k = s_{i_1} s_{i_2} \cdots s_{i_{k-1}}(i_k), \quad b_k = s_{i_1} s_{i_2} \cdots s_{i_{k-1}}(i_k + 1)$$

where we understand $a_1 = i_1$ and $b_1 = i_1 + 1$. Then we have

$$s_{i_m} \cdots \hat{s}_{i_k} \cdots s_{i_1} = w^{-1} t_{a_k, b_k}.$$

Proof. For any $v \in S_n$ and transposition $t_{p,q}$, we have

$$(4.3) \quad v t_{p,q} v^{-1} = t_{v(p), v(q)}.$$

Noting that $s_{i_k} = t_{i_k, i_k+1}$ and $s_j^{-1} = s_j$, we apply (4.3) to the right hand side of the equation

$$s_{i_m} \cdots \hat{s}_{i_k} \cdots s_{i_1} = w^{-1} s_{i_1} s_{i_2} \cdots s_{i_{k-1}} s_{i_k} s_{i_{k-1}} \cdots s_{i_2} s_{i_1}.$$

Then the lemma immediately follows. \square

The *Cartan integers* $c_{i,j}$, which are entries of the Cartan matrix, are given by

$$(4.4) \quad c_{i,j} = \begin{cases} 2 & \text{if } i = j, \\ -1 & \text{if } |i - j| = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Theorem 4.22. *Let $w = s_{i_1} \cdots s_{i_m}$ be a reduced decomposition of $w \in S_n$. Assume that i_1, \dots, i_m are distinct. Then the fan of the toric Schubert variety X_w is isomorphic to the fan in \mathbb{R}^m such that the primitive ray vectors are the $2m$ column vectors of the following matrix and a subset of the column vectors forms a cone if and only if it does not contain both the i th column vectors in the left submatrix and the right submatrix for each $i = 1, \dots, m$:*

$$\left[\begin{array}{cccc|cccc} 1 & 0 & \cdots & 0 & -1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 & & -1 & \cdots & 0 \\ \vdots & & \ddots & & & & \ddots & \\ 0 & 0 & \cdots & 1 & & a_{j,k} & \ddots & \\ & & & & & & & -1 \end{array} \right],$$

where $a_{j,k} = -c_{i_j, i_k}$ for $1 \leq k < j \leq m$. Here, $c_{i,j}$ are Cartan integers (see (4.4)).

Proof. Since the fan of X_w is the normal fan of \mathbf{Q}_w and \mathbf{Q}_w is combinatorially equivalent to an m -dimensional cube by Theorem 4.20, it is enough to consider the edge vectors emanating from the vertices \mathbf{e} and \mathbf{w} to find the fan of X_w as below.

Let $\mathbf{e}_1, \dots, \mathbf{e}_n$ be the standard basis vectors of \mathbb{R}^n and we set

$$(4.5) \quad \mathbf{v}_i := \mathbf{e}_i - \mathbf{e}_{i+1} \quad \text{for } 1 \leq i \leq n-1.$$

The dual vector space to the subspace of \mathbb{R}^n spanned by $\mathbf{v}_1, \dots, \mathbf{v}_{n-1}$ is the quotient space $\mathbb{R}^n / \mathbb{R}(1, \dots, 1)$ and

$$\mathbf{v}_i^* := \mathbf{e}_1 + \cdots + \mathbf{e}_i \quad \text{for } 1 \leq i \leq n-1$$

form the basis of $\mathbb{R}^n / \mathbb{R}(1, \dots, 1)$ dual to (4.5) through the standard scalar product on \mathbb{R}^n .

At the vertex \mathbf{e} , the primitive vectors of outgoing edges are $\mathbf{v}_{i_1}, \dots, \mathbf{v}_{i_m}$ because the atoms of the Bruhat interval $[e, w^{-1}]$ are s_{i_k} 's by (4.2). Hence the primitive facet normal vectors at \mathbf{e} are $\mathbf{v}_{i_1}^*, \dots, \mathbf{v}_{i_m}^*$. On the other hand, in order to find the facet normal vectors at the vertex \mathbf{w} , we consider the following two elements by (4.2) and Lemma 4.21:

$$\begin{aligned} w^{-1} t_{a_k, b_k} &= w^{-1}(1) w^{-1}(2) \cdots w^{-1}(b_k) \cdots w^{-1}(a_k) \cdots w^{-1}(n), \\ w^{-1} &= w^{-1}(1) w^{-1}(2) \cdots w^{-1}(a_k) \cdots w^{-1}(b_k) \cdots w^{-1}(n). \end{aligned}$$

Since $w^{-1} > w^{-1} t_{a_k, b_k}$, we have $w^{-1}(a_k) > w^{-1}(b_k)$. Moreover, we get $a_k < b_k$ because $a_k \leq i_k$ and $b_k \geq i_k + 1$ by the definition of a_k and b_k in Lemma 4.21. Hence the primitive vector of the outgoing edge at \mathbf{w} to the vertex corresponding to $w^{-1} t_{a_k, b_k}$ is $-\mathbf{e}_{a_k} + \mathbf{e}_{b_k}$, which is the same as

$$-s_{i_1} s_{i_2} \cdots s_{i_{k-1}} (\mathbf{e}_{i_k} - \mathbf{e}_{i_k+1}) = -s_{i_1} s_{i_2} \cdots s_{i_{k-1}} (\mathbf{v}_{i_k}).$$

Here, we regard s_i as the reflection in \mathbb{R}^n which interchanges the i th and the $(i+1)$ st coordinates, namely $s_i(\mathbf{v}) = \mathbf{v} - (\mathbf{v}, \mathbf{e}_i - \mathbf{e}_{i+1})(\mathbf{e}_i - \mathbf{e}_{i+1})$ for $\mathbf{v} \in \mathbb{R}^n$, where (\cdot, \cdot) denotes the standard scalar product on \mathbb{R}^n . Now we set

$$\mathbf{w}_k := -s_{i_1} s_{i_2} \cdots s_{i_{k-1}}(\mathbf{v}_{i_k})$$

for $1 \leq k \leq m$. Then, since $c_{i,j} = (\mathbf{e}_i - \mathbf{e}_{i+1}, \mathbf{e}_j - \mathbf{e}_{j+1})$, we have

$$\begin{aligned} \mathbf{w}_k &= -s_{i_1} s_{i_2} \cdots s_{i_{k-1}}(\mathbf{v}_{i_k}) \\ &= -s_{i_1} s_{i_2} \cdots s_{i_{k-2}}(\mathbf{v}_{i_k} - c_{i_k, i_{k-1}} \mathbf{v}_{i_{k-1}}) \\ &= -s_{i_1} s_{i_2} \cdots s_{i_{k-3}}(\mathbf{v}_{i_k} - c_{i_k, i_{k-2}} \mathbf{v}_{i_{k-2}}) - c_{i_k, i_{k-1}} \mathbf{w}_{k-1} \\ &\quad \vdots \\ &= -\mathbf{v}_{i_k} - c_{i_k, i_1} \mathbf{w}_1 - c_{i_k, i_2} \mathbf{w}_2 - \cdots - c_{i_k, i_{k-2}} \mathbf{w}_{k-2} - c_{i_k, i_{k-1}} \mathbf{w}_{k-1}. \end{aligned}$$

The facet normal vectors at w are the dual basis $\mathbf{w}_1^*, \dots, \mathbf{w}_m^*$ to $\mathbf{w}_1, \dots, \mathbf{w}_m$, and the above computation implies that

$$\mathbf{w}_k^* = -\mathbf{v}_{i_k}^* - c_{i_{k+1}, i_k} \mathbf{v}_{i_{k+1}}^* - \cdots - c_{i_m, i_k} \mathbf{v}_{i_m}^*.$$

This proves the theorem. \square

We call the right half of the matrix in Theorem 4.22 the *reduced characteristic matrix* of X_w .

Example 4.23. For $w = s_1 s_2 s_3 s_4$, the reduced characteristic matrix of X_w is

$$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & -1 \end{bmatrix}$$

In general, the reduced characteristic matrix of X_w for $w = s_1 s_2 \cdots s_{n-1}$ (or $w = s_{n-1} s_{n-2} \cdots s_1$) has 1's just below the diagonal and 0 at the other off diagonal entries. The transpose inverse of this matrix is the upper triangular matrix with -1 on the diagonal and above the diagonal, which is the reduced characteristic of the bounded flag manifold BF_{n-1} mentioned in [22, Proposition 7.7.3]. Since the isomorphism class of the variety associated to a reduced characteristic matrix does not change by taking inverse and transpose of the reduced characteristic matrix, X_w for the w above is isomorphic to the bounded flag manifold BF_{n-1} .

A reduced characteristic matrix of a toric Schubert variety X_w depends on the choice of a reduced decomposition of w as is seen in Example 4.24, but it determines X_w up to isomorphism.

Example 4.24. For $w = s_1 s_3 s_2 s_4$, the reduced characteristic matrix of X_w is

$$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 1 & -1 & 0 \\ 0 & 1 & 0 & -1 \end{bmatrix}$$

There are four more reduced decompositions: $w = s_3 s_1 s_2 s_4 = s_1 s_3 s_4 s_2 = s_3 s_1 s_4 s_2 = s_3 s_4 s_1 s_2$. For each case, we have the following reduced characteristic matrices.

$$\begin{array}{cccc} s_3 s_1 s_2 s_4 & s_1 s_3 s_4 s_2 & s_3 s_1 s_4 s_2 & s_3 s_4 s_1 s_2 \\ \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 1 & -1 & 0 \\ 1 & 0 & 0 & -1 \end{bmatrix} & \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 1 & 1 & 0 & -1 \end{bmatrix} & \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & -1 & 0 \\ 1 & 1 & 0 & -1 \end{bmatrix} & \begin{bmatrix} -1 & 0 & 0 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 1 & 0 & 1 & -1 \end{bmatrix} \end{array}$$

Applying Proposition A.8 in Appendix A to the above five matrices, one can easily see that all of them determine the same toric variety up to isomorphism.

A reduced decomposition $w = s_{i_1} \cdots s_{i_m}$ defines a sequence $\mathbf{i} = (i_1, \dots, i_m)$ and vice versa. So we also call the sequence a reduced decomposition of w . We introduce a digraph \mathcal{G}_i associated with a sequence \mathbf{i} , which does not depend on the choice of a reduced decomposition of w , i.e., depends only on w .

Definition 4.25. For $\mathbf{i} = (i_1, \dots, i_m)$, the vertex set and the edge set of a digraph $\mathcal{G}_{\mathbf{i}}$ are defined by

- $V(\mathcal{G}_{\mathbf{i}}) = [m]$;
- $(k, j) \in E(\mathcal{G}_{\mathbf{i}})$ if and only if $|i_k - i_j| = 1$ for $1 \leq k < j \leq m$.

Example 4.26. For $\mathbf{i} = (1, 2, 3, 4)$ and $\mathbf{i}' = (1, 3, 2, 4)$ corresponding to Examples 4.23 and 4.24, we have the following digraphs:



For $\mathbf{i} = (1, 2, 4, 5)$, the digraph $\mathcal{G}_{\mathbf{i}}$ is not connected as follows:



A smooth projective variety X is called *Fano* (resp. *weak Fano*) if the anti-canonical divisor $-K_X$ is ample (resp. nef and big). Fano varieties play an important role as building blocks in the classification problem in algebraic geometry: the minimal model program. We refer the readers to [59, 60].

We consider the Fano or weak Fano characterization of toric Schubert varieties in terms of the digraphs. The digraphs in Example 4.26 are unions of directed path graphs so that each vertex has at most two outgoing edges. This always holds when \mathbf{i} has distinct components and it implies the following.

Theorem 4.27. *Let $\mathbf{i} = (i_1, \dots, i_m)$ be a reduced decomposition of $w \in S_n$. Suppose that i_1, \dots, i_m are distinct. Then the toric Schubert variety X_w is weak Fano. Moreover, it is Fano if and only if each vertex of the graph $\mathcal{G}_{\mathbf{i}}$ has at most one outgoing edge.*

Proof. We apply the criterion of toric Fano or toric weak Fano by Batyrev to our case, see Proposition A.7 for the criterion. It follows from Theorem 4.22 that a primitive collection of the fan of X_w consists of the k th column in the left matrix and the k th column in the right matrix in Theorem 4.22 for each $k = 1, \dots, m$. Then the criterion by Batyrev says that X_w is Fano if and only if 1 appears in the k th column at most once for each k in the reduced characteristic matrix of X_w . Moreover, X_w is weak Fano if and only if 1 appears in the k th column at most twice for each k in the reduced characteristic matrix of X_w . This implies the theorem. \square

We assume $n \geq 3$ in the following. An element of S_n is called a *Coxeter element* if it can be written as a product of all adjacent transpositions s_1, \dots, s_{n-1} . Let Cox_n denote the set of all Coxeter elements in S_n .

Theorem 4.28 ([64]). *Let $w, w' \in \text{Cox}_n$ and let \mathbf{i}, \mathbf{i}' be reduced decompositions of w, w' , respectively. The following statements are equivalent:*

- (1) X_w and $X_{w'}$ are isomorphic as toric varieties (with respect to the T -action on G/B).
- (2) $H^*(X_w; \mathbb{Z}) \cong H^*(X_{w'}; \mathbb{Z})$ as graded rings.
- (3) $\mathcal{G}_{\mathbf{i}} \cong \mathcal{G}_{\mathbf{i}'}$ as digraphs.
- (4) $w' = w$ or $w' = w_0 w w_0$.

Remark 4.29. We notice that the isomorphism classes of Schubert varieties are studied in [80].

Remark 4.30. The digraph $\mathcal{G}_{\mathbf{i}}$ in the theorem above is a directed path graph with $n - 1$ vertices. This can be thought of as a directed Dynkin diagram of type A_{n-1} . It turns out that directed Dynkin diagrams appear in the classification of toric Schubert varieties for other Lie types (see [64]).

4.4. Schubert varieties of complexity one. In this subsection, we consider Schubert varieties of complexity one. All toric Schubert varieties are smooth while Schubert varieties of complexity one are not necessarily smooth. We provide similar statements for Schubert varieties of complexity one to Theorem 4.20.

We recall generalized Bott–Samelson varieties from [55]. Let $G = \text{GL}_n(\mathbb{C})$ as before. For a permutation $w \in S_n$, the subvariety P_w of G corresponding to w is defined by

$$P_w = \overline{BwB} \subseteq G.$$




FIGURE 12. Elnitsky 8-gon for the permutation $3241 \in S_4$ and a tiling

Definition 4.31 (cf. [55, §13.4], [40, Definition 2.1], and [48]). Let (w_1, \dots, w_r) be a sequence of elements in S_n . The *generalized Bott–Samelson variety* $Z_{(w_1, \dots, w_r)}$ is defined by the orbit space

$$Z_{(w_1, \dots, w_r)} := (P_{w_1} \times \cdots \times P_{w_r}) / \Theta,$$

where the right action Θ of $B^r := \underbrace{B \times \cdots \times B}_r$ on $\prod_{k=1}^r P_{w_k}$ is defined by

$$\Theta((p_1, \dots, p_r), (b_1, \dots, b_r)) = (p_1 b_1, b_1^{-1} p_2 b_2, \dots, b_{r-1}^{-1} p_r b_r)$$

for $(p_1, \dots, p_r) \in \prod_{k=1}^r P_{w_k}$ and $(b_1, \dots, b_r) \in B^r$. When $\ell(w_1) = \cdots = \ell(w_r) = 1$, $Z_{(w_1, \dots, w_r)}$ is called a *Bott–Samelson variety*.

We notice that there are simpler descriptions of (generalized) Bott–Samelson varieties in [71, 35] in type A . We recall from [35] the description of Bott–Samelson varieties in type A for readers' convenience. Suppose that a sequence (w_1, \dots, w_r) of elements in S_n consists of permutations having length one, that is, $w_k = s_{i_k}$ the simple reflection for some $i_k \in [n - 1]$. Moreover, suppose that it represents a reduced word of a permutation. In other words, $s_{i_1} \cdots s_{i_r}$ is a reduced decomposition of a permutation. In this case, we simply denote the sequence by $\mathbf{i} = (i_1, \dots, i_r)$. For a permutation w , the *Elnitsky 2n-gon* $E(w)$ has sides of length one, and these are labeled, in order, by $1, 2, \dots, n, w(n), w(n - 1), \dots, w(1)$, in which the first n labels form half of a regular $2n$ -gon, and sides with the same label are parallel. In Figure 12(1), we provide the Elnitsky 8-gon for the permutation $3241 \in S_4$. We denote by v_0 the lowest vertex v_0 of $E(w)$.

Elnitsky [34, Theorem 2.2] proved that there is a bijective correspondence between the set of rhombic tilings of $E(w)$ and the commutation classes of reduced words of w . More precisely, for a given rhombic tiling \mathcal{T} , one can associate a commutation class in an inductive way as follows. We first label each tile γ by a simple reflection s_i if

$$i = \min_{v \in V(\gamma)} \{d(v, v_0)\} + 1$$

where $V(\gamma)$ is the set of vertices in γ and d denotes the graph-theoretic distance between two vertices. Let B_1 be the border consisting of edges b_1, \dots, b_{n+1} and \mathcal{T}_1 the set of tiles in \mathcal{T} intersecting B_1 in two edges. We change the border into the new border B_2 by replacing the two edges of each $\gamma \in \mathcal{T}_1$ intersecting B_1 by the other two edges of γ . Then define \mathcal{T}_2 by the set of all tiles $\mathcal{T} \setminus \mathcal{T}_1$ intersecting B_2 with two edges. Continuing this process inductively until B_k consists only of edges lying on the boundary of $E(w)$, we get a partition $\mathcal{T} = \bigsqcup_k \mathcal{T}_k$. Then we assign a reduced word $\mathbf{i}(\mathcal{T}) = (i_1, \dots, i_r)$ to the partition by reading the labeling of each tile in \mathcal{T}_k in ascending order in k . Note that some partition \mathcal{T}_k may have more than one tile, however, the associated commutation class is well-defined. For instance, the tiling \mathcal{T} in Figure 12(2) corresponds to a reduced word $\mathbf{i} = \mathbf{i}(\mathcal{T}) = (2, 1, 2, 3)$.

To define a Bott–Samelson variety, we associate a vector space to each vertex of a tiling \mathcal{T} . Starting with the vertex between the edges labeled 1 and $w(1)$, label the vertices of $E(w)$ in clockwise order by

$$\mathbb{C}^0, \mathbb{C}^1, \dots, \mathbb{C}^n, G_{n-1}, G_{n-2}, \dots, G_1.$$

Moreover, associate vector space V_x to a vertex x in the tiling. The dimension of V_x is $d(x, v_0)$. For a given tiling \mathcal{T} , define

$$Z_{\mathcal{T}} := \{(V_x)_{x \in V(\mathcal{T})} \mid V_y \subseteq V_z \text{ if } \dim(V_y) + 1 = \dim(V_z) \text{ and } y, z \text{ are adjacent}\}.$$

Here, $V(\mathcal{T})$ is the set of vertices in \mathcal{T} . It is proved in [35, Theorem 1.1] that

$$Z_{\mathcal{T}} \cong Z_{\mathbf{i}(\mathcal{T})}.$$

For instance, the Bott–Samelson variety $Z_{(s_2, s_1, s_2, s_3)} = Z_{(2, 1, 2, 3)}$ is isomorphic to $Z_{\mathcal{T}}$

$$\begin{aligned} Z_{\mathcal{T}} = \{(V_1, V_2, V_3, V_4) \mid \dim(V_k) = i_k, \\ \mathbb{C}^1 \subseteq V_1 \subseteq \mathbb{C}^3, \mathbb{C}^0 \subseteq V_2 \subseteq V_1, \\ V_2 \subseteq V_3 \subseteq \mathbb{C}^3, V_3 \subseteq V_4 \subseteq \mathbb{C}^4\}. \end{aligned}$$

For any $w \in S_n$, the generalized Bott–Samelson variety $Z_{(w)}$ is the Schubert variety X_w . Indeed, not every generalized Bott–Samelson varieties are smooth. If all Schubert varieties X_{w_1}, \dots, X_{w_r} corresponding to $w_1, \dots, w_r \in S_n$ are smooth, then the generalized Bott–Samelson variety $Z_{(w_1, \dots, w_r)}$ is smooth (see [55, (6) in §13.4]).

Theorem 4.32 ([30, 47, 62, 65, 85]). *For a permutation w in S_n , the following statements are equivalent:*

- (1') X_w is smooth and of complexity one.
- (2') w contains the pattern 321 exactly once and avoids the pattern 3412.
- (3') There exists a reduced decomposition of w containing $s_i s_{i+1} s_i$ as a factor and no other repetitions.
- (4') X_w is isomorphic to a generalized Bott–Samelson variety $Z_{(w_1, \dots, w_r)}$ such that $r = \ell(w) - 2$, $w_k = s_j s_{j+1} s_j$ for some $1 \leq k \leq r$, $w_i = s_{j_i}$ for $i \neq k$, and $j_1, \dots, j_{k-1}, j_{k+1}, \dots, j_r, j, j+1$ are pairwise distinct.
- (5') The Bruhat interval $[e, w]$ is isomorphic to $S_3 \times \mathfrak{B}_{\ell(w)-3}$ as posets.
- (6') The polytope \mathbf{Q}_w is combinatorially equivalent to the product of the hexagon and the cube of dimension $\ell(w) - 3$.

Theorem 4.33 ([30, 47, 62, 65, 85]). *For a permutation w in S_n , the following statements are equivalent:*

- (1'') X_w is singular and of complexity one.
- (2'') w contains the pattern 3412 exactly once and avoids the pattern 321.
- (3'') There exists a reduced decomposition of w containing $s_{i+1} s_i s_{i+2} s_{i+1}$ as a factor and no other repetitions.
- (4'') X_w is isomorphic to a generalized Bott–Samelson variety $Z_{(w_1, \dots, w_r)}$ such that $r = \ell(w) - 3$, $w_k = s_{j+1} s_j s_{j+2} s_{j+1}$ for some $1 \leq k \leq r$, $w_i = s_{j_i}$ for $i \neq k$, and $j_1, \dots, j_{k-1}, j_{k+1}, \dots, j_r, j, j+1, j+2$ are pairwise distinct.
- (5'') The Bruhat interval $[e, w]$ is isomorphic to $[e, 3412] \times \mathfrak{B}_{\ell(w)-4}$ as posets.
- (6'') The polytope \mathbf{Q}_w is combinatorially equivalent to the product of \mathbf{Q}_{3412} and the cube of dimension $\ell(w) - 4$.

5. GENERIC TORUS ORBIT CLOSURES IN RICHARDSON VARIETIES

In this section, we consider generic torus orbit closures in Richardson varieties. The topology and geometry of generic torus orbit closures in Richardson varieties are related to the combinatorics of Bruhat interval polytopes. Studying the combinatorial properties of Bruhat interval polytopes, we show that every smooth toric Richardson variety is a Bott manifold. We also give a sufficient condition for a pair $(v, w) \in S_n \times S_n$ to give rise to a smooth toric Richardson variety X_w^v . Motivated by this sufficient condition, we define a toric variety of Catalan type and then show that the number of isomorphism classes of smooth toric Richardson varieties of Catalan type is the Wedderburn–Etherington number.

5.1. Richardson varieties and Bruhat interval polytopes. For $v, w \in S_n$ with $v \leq w$ in the Bruhat order, the Richardson variety X_w^v is defined to be the intersection of the Schubert variety X_w and the opposite Schubert variety $X^v = w_0 X_{w_0 v}$. It is known that every Richardson variety X_w^v is a T -invariant irreducible subvariety of $\text{Fl}(n)$ with respect to the T -action on $\text{Fl}(n)$ and

$$\dim_{\mathbb{C}} X_w^v = \ell(w) - \ell(v).$$

See [20, Section 1.3] for more details. Moreover, a T -fixed point uB is contained in X_w^v if and only if $v \leq u \leq w$ in the Bruhat order. Accordingly, the set of T -fixed points of X_w^v is identified with the Bruhat interval

$$[v, w] = \{u \in S_n \mid v \leq u \leq w\}.$$

Recall that the moment map $\mu: G/B \rightarrow \mathbb{R}^n$ in (2.5) sends uB to $\mathbf{u} = (u^{-1}(1), \dots, u^{-1}(n))$. Hence we have

$$\mu(X_w^v) = \text{Conv}\{(u^{-1}(1), \dots, u^{-1}(n)) \mid u \in [v, w]\}.$$

Motivated by this fact, we provide the following definition.

Definition 5.1 ([58]). For elements v and w in S_n with $v \leq w$, we define a polytope \mathbf{Q}_w^v by

$$\mathbf{Q}_w^v := \text{Conv}\{(u^{-1}(1), \dots, u^{-1}(n)) \mid u \in [v, w]\}.$$

This polytope is called a *Bruhat interval polytope*. Note that \mathbf{Q}_w^e is the polytope \mathbf{Q}_w introduced in Section 4.

By the definition of \mathbf{Q}_w^v , we have

$$\mu: X_w^v \rightarrow \mathbf{Q}_w^v \subset \mathbb{R}^n.$$

Remark 5.2. Bruhat interval polytopes were introduced by Kodama–Williams [58] and their combinatorial properties are studied in [86] and [66]. For $v, w \in S_n$ with $v \leq w$, the notation

$$\mathbf{Q}_{v,w} := \text{Conv}\{(u(1), \dots, u(n)) \mid u \in [v, w]\}$$

is used in [58, 86, 66]. Therefore we have $\mathbf{Q}_w^v = \mathbf{Q}_{v^{-1}, w^{-1}}$.

Bruhat interval polytopes \mathbf{Q}_w^v and $\mathbf{Q}_{w^{-1}}^{v^{-1}}$ are not necessarily combinatorially equivalent even though the intervals $[v, w]$ and $[v^{-1}, w^{-1}]$ are isomorphic as posets. For example, one can check that the polytopes $\mathbf{Q}_{35412}^{12345}$ and $\mathbf{Q}_{45132}^{12345}$ have different numbers of edges using a computer program like SageMath. However, they have the same dimension.


Proposition 5.3 ([66, Proposition 3.4]). *Bruhat interval polytopes \mathbf{Q}_w^v and $\mathbf{Q}_{w^{-1}}^{v^{-1}}$ have the same dimension.*

5.2. Generic torus orbit closures in Richardson varieties. A torus orbit $T \cdot x$ for $x \in X_w^v$ is said to be *generic* if $(\overline{T \cdot x})^T = (X_w^v)^T$. Every Richardson variety admits a generic torus orbit. Its proof is given in [63, Proposition 3.8] when $v = e$ and a similar argument works for any Richardson variety X_w^v . We will denote by Y_w^v the closure of a generic T -orbit in X_w^v . Then Y_w^v is the projective toric variety defined by the polytope \mathbf{Q}_w^v since $\mu(Y_w^v) = \mu(X_w^v) = \mathbf{Q}_w^v$.

Note that the family of Bruhat interval polytopes forms a subfamily of the Φ -polytopes. Hence every edge of a Bruhat interval polytope is parallel to a vector of the form $\mathbf{e}_i - \mathbf{e}_j$. This implies that Y_w^v is smooth at a fixed point uB if and only if \mathbf{Q}_w^v is simple at the vertex \mathbf{u} (see [63, Section 8]).

While $Y_w = Y_w^e$ is smooth at eB , not every toric variety Y_w^v is smooth at vB . For instance, the polytope \mathbf{Q}_{3412}^{1324} is not simple at 1324, see Figure 13. A natural generalization of Theorem 4.11 is the following.

Conjecture 5.4. *The generic torus orbit closure Y_w^v in the Richardson variety X_w^v is smooth if it is smooth at the fixed points vB and wB , in other words, \mathbf{Q}_w^v is simple if it is simple at the vertices \mathbf{v} and \mathbf{w} .*

FIGURE 13. Bruhat interval polytope Q_{3412}^{1324}

5.3. Toric Bruhat interval polytopes. Since Q_w^v is the moment map image of a toric variety Y_w^v , we get

$$\dim Q_w^v = \dim_{\mathbb{C}} Y_w^v \leq \dim_{\mathbb{C}} X_w^v = \ell(w) - \ell(v).$$

Motivated by this observation, we call a Bruhat interval polytope Q_w^v *toric* if $\dim Q_w^v = \ell(w) - \ell(v)$. Hence a Richardson variety X_w^v is a toric variety with respect to the T -action, that is, $X_w^v = Y_w^v$, if and only if the Bruhat interval polytope Q_w^v is toric.

Tsukerman and William [86] showed that every face of the Bruhat interval polytope Q_w^v is realizable as a subinterval of $[v^{-1}, w^{-1}]$, but the converse is not true. For instance, the subinterval $[123, 312]$ of $[123, 321] = [123^{-1}, 321^{-1}]$ does not form a face of the polytope Q_{321}^{123} . See Figure 1. However, the converse holds when Q_w^v is toric. More strongly, we have the following.

Theorem 5.5 ([66, Theorem 5.1] and [24, Theorem 1.1]). *For a Bruhat interval polytope Q_w^v , the following statements are equivalent:*

- (1) Q_w^v is toric (i.e., $\dim Q_w^v = \ell(w) - \ell(v)$).
- (2) Q_y^x is a face of Q_w^v for any $[x^{-1}, y^{-1}] \subset [v^{-1}, w^{-1}]$.
- (3) The Bruhat interval $[v, w]$ does not contain any interval that is isomorphic to symmetric group S_3 with respect to Bruhat order and $\ell(w) - \ell(v) \leq n - 1$.
- (4) The interval $[v, w]$ is a lattice and $\ell(w) - \ell(v) \leq n - 1$.

Here, a poset P is a *lattice* if for all $x, y \in P$, the subposet $\{z \in P \mid z \leq x, z \leq y\}$ has a top element, and the subposet $\{z \in P \mid z \geq x, z \geq y\}$ has a bottom element.

The above statement is obtained by combining the results [66, Theorem 5.1] and [24]. Indeed, the equivalence between (1) and (2) is proved in [66, Theorem 5.1]; the equivalences between (1), (3), and (4) are given in [24, Theorem 1.1]. Hence, if Q_w^v is toric, then its combinatorial type is determined by the poset structure of $[v^{-1}, w^{-1}]$, so Q_w^v and $Q_{w^{-1}}^{v^{-1}}$ are combinatorially equivalent. Here, the assumption of “toric” cannot be removed. Indeed, the interval $[1324, 4231]$ is a Boolean algebra of rank 4, but the corresponding Bruhat interval polytope is of dimension 3. See Figure 14. On the other hand, for every positive integer m there is an example of an interval $[v, w]$ of rank m such that $[v, w]$ is a Boolean algebra and Q_w^v is toric. For instance, when $w = s_m s_{m-1} \cdots s_2 s_1$ and $v = e$, the Bruhat interval polytope Q_w^v is combinatorially equivalent to a cube of dimension m (simply, m -cube).




FIGURE 14. The interval $[1324, 4231]$ is a Boolean algebra, but Q_{4231}^{1324} is not a cube.

In the following, for simplicity, when a polytope Q is combinatorially equivalent to a cube (or a d -cube), we say that Q is a *cube* (or a d -*cube*). We also say that an interval $[v, w]$ is *Boolean* if it is a Boolean algebra.

Using the following two facts

- (1) if Q is a simple polytope of dimension ≥ 2 and every 2-face of Q is a 2-cube, then Q is a cube ([94, Problems and Exercises 0.1 in p.23] and [92, Appendix]), and
- (2) every 2-interval is a diamond ([15, Lemma 2.7.3]),

we can prove:

Proposition 5.6 ([66, Proposition 5.6]). *Suppose that Q_w^v is toric. Then Q_w^v is a cube if and only if it is simple. (This is equivalent to saying that a toric Richardson variety is a Bott manifold if and only if it is smooth.)*

The following gives a characterization of when Q_w^v is a cube.

Theorem 5.7 ([66, Theorem 5.7] and [24, Theorem 1.1]). *The following statements are equivalent.*

- (1) A Bruhat interval polytope Q_w^v is a cube.
- (2) Q_w^v is toric and $[v, w]$ is Boolean.
- (3) A Richardson variety X_w^v is a Bott manifold.
- (4) $\ell(w) - \ell(v) \leq n - 1$ and $[v, w]$ is Boolean.

The equivalences between (1), (2), and (3) are given in [66, Theorem 5.7], and the equivalence between (2) and (4) is given by [24, Theorem 1.1]. In the above theorem, we cannot drop either *toric* or *Boolean*.

Example 5.8. (1) For $n = 4$, $v = 1324$, and $w = 4231$ in S_4 , the interval $[v, w]$ is Boolean of length 4 but $\dim Q_w^v = 3$, so Q_w^v is not toric. We notice that $\ell(w) - \ell(v) = 5 - 1 = 4 \not\leq 3 = n - 1$. Since the vertices v and w have degree 4, Q_w^v is not a cube.
 (2) When $v = 1324$ and $w = 3412$, we have $\ell(w) - \ell(v) = 4 - 1 = 3$ and $\dim Q_w^v = 3$. Hence Q_w^v is toric. However, $[v, w]$ is a 4-crown (see [15, p.52]) and not Boolean. The vertices v and w have degree 4 and the others are simple vertices, so Q_w^v is not a cube. See Figure 13.

It is shown in [79, Theorem 3.5.2] that the largest rank of Boolean Bruhat intervals in S_{n+1} is at least $n + \lfloor \frac{n-1}{2} \rfloor$ by finding a sufficient condition on v and w for $[v, w]$ to be Boolean. This implies that there are infinitely many Boolean Bruhat intervals which are not toric like the Boolean

interval $[1324, 4231]$. In fact, for any non-negative integer k , there is a Boolean interval $[v, w]$ such that $\dim \mathbb{Q}_w^v = \ell(w) - \ell(v) - k$. See [66, Proposition 6.4] for details.

Remark 5.9. Can and Saha considered toric Richardson varieties for any connected, simple, simply-connected algebraic group over an algebraically closed field in [24]. They provide characterizations of toric Richardson varieties in terms of the combinatorics of posets.

5.4. Conditions on v and w for \mathbb{Q}_w^v to be a cube. Now we find a sufficient condition for a Bruhat interval polytope \mathbb{Q}_w^v to be a cube.

It is shown in [51, §5 and §6] that \mathbb{Q}_w^v is toric (in fact, a cube) if $v^{-1} = [a_1, \dots, a_{n-1}, n]$ and $w^{-1} = [n, a_1, \dots, a_{n-1}]$ or $v^{-1} = [1, b_2, \dots, b_n]$ and $w^{-1} = [b_2, \dots, b_n, 1]$. In these cases,

$$\begin{aligned} w &= s_1 s_2 \cdots s_{n-1} v & \text{and} & \quad \ell(w) - \ell(v) = n - 1, \text{ or} \\ w &= s_{n-1} s_{n-2} \cdots s_1 v & \text{and} & \quad \ell(w) - \ell(v) = n - 1. \end{aligned}$$

These examples and Theorem 4.20(3) motivate us to study the following case:

$$(5.1) \quad w = s_{j_1} s_{j_2} \cdots s_{j_m} v \text{ or } v s_{j_1} s_{j_2} \cdots s_{j_m} \text{ where } \ell(w) - \ell(v) = m \text{ and } j_1, \dots, j_m \text{ are distinct.}$$

Proposition 5.10 ([66, Proposition 7.1]). *Suppose that $w = s_{j_1} s_{j_2} \cdots s_{j_m} v$ or $w = v s_{j_1} s_{j_2} \cdots s_{j_m}$ with $\ell(w) - \ell(v) = m$. Then j_1, \dots, j_m are distinct if and only if \mathbb{Q}_w^v is toric.*

Note that not every Bruhat interval polytope in the above proposition is a cube. For instance, if $v = 1324$ and $w = 3412$, then $v = s_2$ and $w = s_2 s_3 s_1 s_2$, so $w = s_2 s_3 s_1 v$. However, the polytope \mathbb{Q}_w^v is not a cube as in Figure 13. On the other hand, for $v = 1243$ and $w = 3412$, since $v = s_3$ and $w = s_2 s_3 s_1 s_2 = s_2 s_1 s_3 s_2$, neither wv^{-1} nor $v^{-1}w$ is a product of distinct adjacent transpositions. However, the Bruhat interval polytope \mathbb{Q}_w^v is a 3-cube. These two examples show that it seems difficult to characterize v and w for which \mathbb{Q}_w^v is a cube.

Let us find a sufficient condition on v and w for \mathbb{Q}_w^v to be a cube. For that, we prepare some notations. For $p, q \in [n - 1]$, we set

$$(5.2) \quad s(p, q) = \begin{cases} s_p s_{p+1} \cdots s_q & \text{if } p \leq q, \\ s_p s_{p-1} \cdots s_q & \text{if } p \geq q. \end{cases}$$

For each $s(p, q)$, we also set

$$\bar{p} = \min\{p, q\}, \quad \bar{q} = \max\{p, q\}.$$

We note that if $j_1, \dots, j_m \in [n - 1]$ are distinct, then we have a *minimal* expression

$$(5.3) \quad s_{j_1} s_{j_2} \cdots s_{j_m} = s(p_1, q_1) s(p_2, q_2) \cdots s(p_r, q_r)$$

where the intervals $[\bar{p}_1, \bar{q}_1], \dots, [\bar{p}_r, \bar{q}_r]$ are disjoint and r is the minimum among such expressions.

Example 5.11. Here are examples of minimal expressions.

- (1) $s_1 s_2 \cdots s_{n-1} = s(1, n - 1), \quad s_{n-1} s_{n-2} \cdots s_1 = s(n - 1, 1).$
- (2) $s_1 s_3 s_8 s_2 s_4 s_7 s_6 = s_3 s_4 s_1 s_2 s_8 s_7 s_6 = s(3, 4) s(1, 2) s(8, 6).$
- (3) $s_2 s_8 s_4 s_7 s_1 s_6 = s_2 s_1 s_4 s_8 s_7 s_6 = s(2, 1) s(4, 4) s(8, 6).$

We say that the product $s_{j_1} s_{j_2} \cdots s_{j_m}$ in (5.3) is *proper* if no two intervals among $[\bar{p}_1, \bar{q}_1], \dots, [\bar{p}_r, \bar{q}_r]$ are adjacent, in other words, the cycles defined by $s(p_1, q_1), \dots, s(p_r, q_r)$ are disjoint. In Example 5.11, (1) and (3) are proper, but (2) is not because the intervals $[3, 4]$ and $[1, 2]$ are adjacent.

The following provides a sufficient condition for v and w such that the Richardson variety X_w^v is smooth and toric.

Proposition 5.12 ([66, Proposition 7.3]). *Suppose that $s_{j_1} s_{j_2} \cdots s_{j_m}$ is a proper minimal expression. If $w = s_{j_1} s_{j_2} \cdots s_{j_m} v$ or $w = v s_{j_1} s_{j_2} \cdots s_{j_m}$ with $\ell(w) - \ell(v) = m$, then the Bruhat interval polytope \mathbb{Q}_w^v is a cube.*

In fact, the converse of Proposition 5.12 is also true. That is, a Bruhat interval polytope \mathbb{Q}_w^v is a cube for any v and w in (5.1) if and only if the product $s_{j_1} s_{j_2} \cdots s_{j_m}$ is proper. See [66, Corollary 7.7].

5.5. Toric varieties of Catalan type. Many smooth toric Richardson varieties arise from polygon triangulations. This subsection is a preparation for that. Namely we explain how to associate a compact smooth toric variety with a polygon triangulation.

Let P_{n+2} denote a convex polygon in the plane with $n + 2$ vertices (or *convex* $(n + 2)$ -gon for simplicity). We label the vertices from 0 to $n + 1$ in counterclockwise order. A *triangulation* of P_{n+2} is a decomposition of P_{n+2} into a set of n triangles by adding $n - 1$ diagonals of P_{n+2} which do not intersect in their interiors. Then the *Catalan number*

$$C_n = \frac{1}{n + 1} \binom{2n}{n}$$

is the number of triangulations of P_{n+2} . Figure 15 shows the triangulations of P_5 . Here, we consider all five triangulations to be different because we are triangulating an $(n + 2)$ -gon with *labelled* vertices.




FIGURE 15. Triangulations of P_5

There are many interpretations of Catalan numbers such as binary trees, Dyck paths, and binary operations, see [83]. In the following, we review the correspondence between polygon triangulations and binary trees and then define a toric variety of Catalan type.

A *binary tree* is a rooted plane tree with at most two children at each vertex defined recursively as follows. An empty set \emptyset is a binary tree. Otherwise, a binary tree has a root vertex v , a left subtree, and a right subtree, both of which are binary trees. The set of triangulations \mathcal{T} of a convex polygon P_{n+2} with $n + 2$ vertices and the set of binary trees $\mathcal{B}_{\mathcal{T}}$ with n vertices have the following bijective connection defined recursively. Assume that \mathcal{T} is a triangulation of P_{n+2} . For $n = 0$, the associated binary tree is an empty set. For $n \geq 1$, we put a vertex of $\mathcal{B}_{\mathcal{T}}$ in the interior of each triangle of \mathcal{T} . The root vertex of $\mathcal{B}_{\mathcal{T}}$ corresponds to the vertex in the triangle having the side $\{0, n + 1\}$. Now by deleting the side $\{0, n + 1\}$ of the $(n + 2)$ -gon, we obtain a union of two triangulated polygons which define the left and right subtrees as follows: the polygon having the vertex 0 defines the left subtree; the polygon having the vertex $n + 2$ defines the right subtree. The binary trees associated with the triangulation of P_5 in Figure 15 are shown in Figure 16. The root vertex is the vertex with the additional circle.




FIGURE 16. Binary trees with three vertices

If a binary tree has zero or two children at each vertex, then it is called *full*. Similarly to $\mathcal{B}_{\mathcal{T}}$, a full binary tree $\mathcal{C}_{\mathcal{T}}$, which contains $\mathcal{B}_{\mathcal{T}}$ as a subgraph, is associated with a triangulation \mathcal{T} of P_{n+2} . Indeed, in addition to the vertices of $\mathcal{B}_{\mathcal{T}}$, we place a vertex outside the side $\{i, i + 1\}$ of P_{n+2} for each $i = 0, \dots, n$, and connect it to the vertex in the interior of the triangle having the side $\{i, i + 1\}$ by an edge. This produces the full binary tree $\mathcal{C}_{\mathcal{T}}$. The root vertex of $\mathcal{C}_{\mathcal{T}}$ is the same as that of $\mathcal{B}_{\mathcal{T}}$. Moreover, if we delete all leaves of $\mathcal{C}_{\mathcal{T}}$, then we obtain the binary tree $\mathcal{B}_{\mathcal{T}}$. Here, a *leaf* is a vertex with no children. Indeed, the number of vertices of $\mathcal{C}_{\mathcal{T}}$ is $2n + 1$. The full binary trees associated with the triangulation of P_5 in Figure 15 are shown in Figure 17, where the colored vertices represent the vertices placed outside the polygon.




FIGURE 17. Full binary trees with three vertices

Now we introduce the left tree and the right tree of a triangulation \mathcal{T} of \mathbb{P}_{n+2} using the corresponding full binary tree $\mathcal{C}_{\mathcal{T}}$. The *left tree* \mathcal{T}^{L} (resp., *right tree* \mathcal{T}^{R}) consists of the sides in \mathcal{T} intersecting with an edge of $\mathcal{C}_{\mathcal{T}}$ connecting a vertex and its left child (resp., right child). See Figure 18. The graphs \mathcal{T}^{L} and \mathcal{T}^{R} are indeed *trees* as shown in [68, Lemma 4.4].




FIGURE 18. The left tree (colored in blue) and the right tree (colored in red and dashed) of a triangulation of \mathbb{P}_{10} . Here, we color the vertices $V(\mathcal{C}_{\mathcal{T}}) \setminus V(\mathcal{B}_{\mathcal{T}})$ purple.

From the definition of the left and the right trees, we obtain the following lemma.

Lemma 5.13 ([68, Lemma 4.5]). *Let \mathcal{T} be a triangulation of \mathbb{P}_{n+2} . For each $1 \leq k \leq n$, there is only one edge $\{k_{\text{L}}, k\}$ with $k_{\text{L}} < k$ in the left tree \mathcal{T}^{L} and similarly there is only one edge $\{k, k_{\text{R}}\}$ with $k < k_{\text{R}}$ in the right tree \mathcal{T}^{R} . Moreover, $k_{\text{L}}, k, k_{\text{R}}$ are the vertices of a triangle in \mathcal{T} .*

For the triangulation \mathcal{T} in Figure 18, we display the vertices k_{L} and k_{R} for each $1 \leq k \leq n$ in the following table:

| k | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|----------------|---|---|---|---|---|---|---|---|
| k_{L} | 0 | 0 | 2 | 3 | 3 | 2 | 2 | 7 |
| k_{R} | 2 | 9 | 6 | 5 | 6 | 7 | 9 | 9 |

For each $k = 1, \dots, n$, we define

- (1) $\mathbf{p}_k = \mathbf{e}_{k_{\text{L}+1}} - \mathbf{e}_{k+1}$,
- (2) $\mathbf{q}_k = -\mathbf{e}_k + \mathbf{e}_{k_{\text{R}}}$,

where $\{\mathbf{e}_1, \dots, \mathbf{e}_{n+1}\}$ is the standard basis of \mathbb{Z}^{n+1} . The vectors $\mathbf{p}_1, \dots, \mathbf{p}_n$ (similarly, $\mathbf{q}_1, \dots, \mathbf{q}_n$) form a basis of the n -dimensional lattice

$$M = \{(x_1, \dots, x_{n+1}) \in \mathbb{Z}^{n+1} \mid x_1 + \dots + x_{n+1} = 0\}.$$

Through the dot product on \mathbb{Z}^{n+1} , the dual lattice N of M can be identified with the quotient lattice of \mathbb{Z}^{n+1} by the sublattice generated by $(1, \dots, 1)$, i.e.,

$$N = \mathbb{Z}^{n+1} / \mathbb{Z}(1, \dots, 1).$$

Let ϖ_i ($i = 1, \dots, n$) be the quotient image of $\mathbf{e}_1 + \dots + \mathbf{e}_i$ in N . Then $\{\varpi_1, \dots, \varpi_n\}$ is a basis of N . For convenience, we set $\varpi_0 = \varpi_{n+1} = \mathbf{0}$.

To each side $\{a, b\}$ in the triangulation \mathcal{T} with $a < b$, we assign the vector $\varpi_a - \varpi_b$ and denote it by \mathbf{v}_a when $\{a, b\} \in \mathcal{T}^R$ and \mathbf{w}_b when $\{a, b\} \in \mathcal{T}^L$, in other words,

$$\begin{aligned} \mathbf{v}_k &= \varpi_k - \varpi_{k_R}, \\ \mathbf{w}_k &= \varpi_{k_L} - \varpi_k, \end{aligned}$$

for $k = 1, \dots, n$ by Lemma 5.13. Note that the zero vector $\mathbf{0}$ is assigned to the distinguished side $\{0, n+1\}$ because $\varpi_0 = \varpi_{n+1} = \mathbf{0}$. Then for the pairing $\langle \cdot, \cdot \rangle$ between N and M induced from the dot product on \mathbb{Z}^{n+1} , we have

$$\langle \mathbf{v}_i, \mathbf{p}_j \rangle = \langle \mathbf{w}_i, \mathbf{q}_j \rangle = \delta_{ij},$$

where δ_{ij} denotes the Kronecker delta. See Figure 19 for the assignment of the vectors \mathbf{w}_k 's and \mathbf{v}_k 's to the left and right trees of the triangulation in Figure 18. One can check that $\langle \mathbf{v}_i, \mathbf{p}_j \rangle = \langle \mathbf{w}_i, \mathbf{q}_j \rangle = \delta_{ij}$ using Table 3.

| k | \mathbf{p}_k | \mathbf{q}_k | \mathbf{v}_k | \mathbf{w}_k |
|-----|-------------------------------|--------------------------------|-----------------------|-----------------------|
| 1 | $\mathbf{e}_1 - \mathbf{e}_2$ | $-\mathbf{e}_1 + \mathbf{e}_2$ | $\varpi_1 - \varpi_2$ | $-\varpi_1$ |
| 2 | $\mathbf{e}_1 - \mathbf{e}_3$ | $-\mathbf{e}_2 + \mathbf{e}_3$ | ϖ_2 | $-\varpi_2$ |
| 3 | $\mathbf{e}_3 - \mathbf{e}_4$ | $-\mathbf{e}_3 + \mathbf{e}_6$ | $\varpi_3 - \varpi_6$ | $\varpi_2 - \varpi_3$ |
| 4 | $\mathbf{e}_4 - \mathbf{e}_5$ | $-\mathbf{e}_4 + \mathbf{e}_5$ | $\varpi_4 - \varpi_5$ | $\varpi_3 - \varpi_4$ |
| 5 | $\mathbf{e}_4 - \mathbf{e}_6$ | $-\mathbf{e}_5 + \mathbf{e}_6$ | $\varpi_5 - \varpi_6$ | $\varpi_3 - \varpi_5$ |
| 6 | $\mathbf{e}_3 - \mathbf{e}_7$ | $-\mathbf{e}_6 + \mathbf{e}_7$ | $\varpi_6 - \varpi_7$ | $\varpi_2 - \varpi_6$ |
| 7 | $\mathbf{e}_3 - \mathbf{e}_8$ | $-\mathbf{e}_7 + \mathbf{e}_9$ | ϖ_7 | $\varpi_2 - \varpi_7$ |
| 8 | $\mathbf{e}_8 - \mathbf{e}_9$ | $-\mathbf{e}_8 + \mathbf{e}_9$ | ϖ_8 | $\varpi_7 - \varpi_8$ |

TABLE 3. Vectors $\mathbf{p}_k, \mathbf{q}_k, \mathbf{v}_k, \mathbf{w}_k$ for the triangulation in Figure 18




FIGURE 19. Vectors assigned to the sides of \mathcal{T} in Figure 18

Lemma 5.14 ([68, Lemma 6.1]). *The collection of cones spanned by \mathbf{u}_i ($i \in I$), where \mathbf{u}_i is either \mathbf{v}_i or \mathbf{w}_i and I runs over all subsets of $[n]$, forms a complete non-singular fan $\Sigma_{\mathcal{T}}$ in $N \otimes \mathbb{R}$.*

The above lemma says that the underlying simplicial complex of the fan $\Sigma_{\mathcal{T}}$ is the boundary complex of an n -dimensional cross-polytope. It is known from [72, Corollary 3.5] that such a fan is indeed the normal fan of an n -cube and the smooth compact toric variety $X(\Sigma_{\mathcal{T}})$ associated with the fan $\Sigma_{\mathcal{T}}$ is a Bott manifold (cf. Proposition A.10).

Definition 5.15. Since $\Sigma_{\mathcal{T}}$ is associated with a polygon triangulation \mathcal{T} , we say that the fan $\Sigma_{\mathcal{T}}$ and the corresponding (smooth compact) toric variety $X(\Sigma_{\mathcal{T}})$ are of *Catalan type*.

Since the vertices k_L, k, k_R form a triangle for each $k \in [n]$, there is a unique $k_0 \in [n]$ such that $\mathbf{v}_{k_0} + \mathbf{w}_{k_0} = \mathbf{0}$, and for $k \in [n] \setminus \{k_0\}$ we have

$$\mathbf{v}_k + \mathbf{w}_k = \begin{cases} \mathbf{v}_{k_L} & \text{if } \{k_L, k_R\} \in E(\mathcal{T}^R), \\ \mathbf{w}_{k_R} & \text{if } \{k_L, k_R\} \in E(\mathcal{T}^L). \end{cases}$$

Note that k_0 is the remaining vertex of the triangle containing the distinguished side $\{0, n+1\}$. See Figure 19 and [68, Lemma 5.1] for more details. Hence the set of primitive collections of $\Sigma_{\mathcal{T}}$ is

$$\{\{\mathbf{v}_k, \mathbf{w}_k\} \mid k \in [n]\}.$$

(See Definition A.6 for definition of primitive collections.) Then Batyrev's criterion ([11, Proposition 2.3.6], see also Appendix A.2) implies the following:

Lemma 5.16 ([68, Lemma 6.4]). *The toric variety $X(\Sigma_{\mathcal{T}})$ is Fano.*

The toric varieties of Catalan type are classified as follows.

Theorem 5.17 ([68, Theorem 6.5]). *Let \mathcal{T} and \mathcal{T}' be triangulations of \mathbb{P}_{n+2} . Then the fans $\Sigma_{\mathcal{T}}$ and $\Sigma_{\mathcal{T}'}$ are isomorphic (equivalently, the toric varieties $X(\Sigma_{\mathcal{T}})$ and $X(\Sigma_{\mathcal{T}'})$ are isomorphic) if and only if the binary trees $\mathcal{B}_{\mathcal{T}}$ and $\mathcal{B}_{\mathcal{T}'}$ are isomorphic as unordered rooted trees.*

The above theorem implies that the number of isomorphism classes of n -dimensional toric varieties of Catalan type is the same as that of unordered binary trees with n vertices. It is known that the latter is the Wedderburn–Etherington number b_{n+1} .

Corollary 5.18 ([68, Corollary 6.6]). *The number of isomorphism classes of n -dimensional toric varieties of Catalan type is the Wedderburn–Etherington number b_{n+1} .*

Here, the Wedderburn–Etherington number b_n ($n \geq 1$) is the number of ways to parenthesize a string of n letters subject to a commutative (but nonassociative) binary operation and it appears in counting several different objects (see [Sequence A001190](#) in OEIS [75], [83, A56 in p.133]). The generating function $B(x) = \sum_{n \geq 1} b_n x^n$ of the Wedderburn–Etherington numbers satisfies the functional equation

$$B(x) = x + \frac{1}{2}B(x)^2 + \frac{1}{2}B(x^2),$$

which was the motivation of Wedderburn in his work [88] and was considered by Etherington [36]. This functional equation is equivalent to the recurrence relation

$$b_{2m-1} = \sum_{i=1}^{m-1} b_i b_{2m-i-1} \quad (m \geq 2), \quad b_{2m} = b_m(b_m + 1)/2 + \sum_{i=1}^{m-1} b_i b_{2m-i}$$

with $b_1 = 1$. Using this recurrence relation, one can calculate the Wedderburn–Etherington numbers, see Table 4.

| | | | | | | | | | | | | | | | |
|-------|---|---|---|---|---|---|----|----|----|----|-----|-----|-----|------|------|
| n | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 |
| b_n | 1 | 1 | 1 | 2 | 3 | 6 | 11 | 23 | 46 | 98 | 207 | 451 | 983 | 2179 | 4850 |

TABLE 4. Wedderburn–Etherington numbers b_n for small values of n

5.6. Smooth toric Richardson varieties of Catalan type. We say that a smooth toric Richardson variety X_w^v is of Catalan type if it is of Catalan type as a toric variety, in other words, if the normal fan of \mathbb{Q}_w^v is of Catalan type. In this subsection, we show that every toric variety of Catalan type appears as a smooth toric Richardson variety. However, the converse is not true, i.e., there are smooth toric Richardson varieties which are not of Catalan type. For instance, X_{3142} is a (smooth) toric Schubert variety but not Fano, so it is not of Catalan type by Lemma 5.16.

For a permutation $u \in S_n$, we define permutations u_h and u_t in S_{n+1} by

$$u_h(i) = \begin{cases} 1 & \text{if } i = 1, \\ u(i-1) + 1 & \text{if } 2 \leq i \leq n+1, \end{cases} \quad u_t(i) = \begin{cases} u(i) & \text{if } 1 \leq i \leq n, \\ n+1 & \text{if } i = n+1. \end{cases}$$

For example, if $u = 2314$, then $u_h = 13425$ and $u_t = 23145$. As one may see, the permutation u_h is obtained from u by putting the additional number 1 at the *head* (with the original numbers increased by 1) while u_t is obtained from u by putting the additional number $n+1$ at the *tail*. In the notation, **h** stands for *head* and **t** stands for *tail*.

One notes that if a pair (v, w) of elements in S_n satisfies

$$w = s(n, 1)v \quad (\text{resp. } w = s(1, n)v) \quad \text{and} \quad \ell(w) = \ell(v) + n,$$

then $v(1) = 1$ (resp. $v(n+1) = n+1$), so that

$$(5.4) \quad (v^{-1}, w^{-1}) = (u_h, u_h s(1, n)) \quad (\text{resp. } (v^{-1}, w^{-1}) = (u_t, u_t s(n, 1))) \quad \text{for some } u \in S_n.$$

Here, $s(n, 1) = s_n s_{n-1} \cdots s_1$ and $s(1, n) = s_1 s_2 \cdots s_n$ as in (5.2).

By Proposition 5.12, the Bruhat interval polytope Q_w^v for the pair (v, w) in (5.4) is an n -cube and our concern is the pairs in (5.4). We first consider the former case $(v^{-1}, w^{-1}) = (u_h, u_h s(1, n))$.

We recall a surjection from S_n to the set of binary trees with n vertices (cf. [70, Appendix A]). To a permutation $u \in S_n$, we first associate a binary tree $\tilde{\psi}(u)$ with vertex labels by finding the smallest number in the one-line notation of u inductively. We start with the one-line notation $u(1)u(2) \cdots u(n)$ of u . The smallest integer, say $u(p)$, in the sequence (which is 1 here) becomes the root of the binary tree $\tilde{\psi}(u)$ with n vertices. Then the subsequence $u(1) \cdots u(p-1)$ will provide the left subtree of the root vertex, and the subsequence $u(p+1) \cdots u(n)$ will provide the right subtree of the root vertex. More precisely, the smallest integer in the subsequence $u(1) \cdots u(p-1)$ presents the root of a binary tree with $p-1$ vertices, and it is the left child of the root vertex of $\tilde{\psi}(u)$. On the other hand, the smallest integer in the subsequence $u(p+1) \cdots u(n)$ presents the root of a binary tree with $n-p$ vertices, and it is the right child of the root vertex of $\tilde{\psi}(u)$. Continuing this process, we obtain the binary tree $\tilde{\psi}(u)$. Finally, erasing the vertex labels, we obtain the desired binary tree $\psi(u)$ with n vertices.

For example, if $u = 31687524$, the root of the binary tree $\tilde{\psi}(u)$ is $u(2)$, and its left and right subtrees have the roots $u(1)$ and $u(7)$, respectively. Continuing this process, we first get the binary tree $\tilde{\psi}(u)$, and then by erasing the labels we obtain the binary tree $\psi(u)$, see Figure 20. Note that $v = 21687534$ gives the same binary tree as $u = 31687524$, i.e., $\psi(v) = \psi(u)$.




FIGURE 20. The binary trees $\tilde{\psi}(u)$ and $\psi(u)$ for $u = 31687524$

Through the canonical bijection between the set of binary trees with n vertices and that of triangulations of P_{n+2} , the assignment

$$\psi: S_n \rightarrow \{\text{binary trees with } n \text{ vertices}\} = \{\text{triangulations of } P_{n+2}\}$$

is surjective (cf. [70, Appendix A]).

Proposition 5.19 ([68, Proposition 7.2]). *Let $u \in S_n$ and let $\mathcal{T} = \psi(u)$ be the corresponding triangulation of P_{n+2} . We denote by \mathcal{T}^L and \mathcal{T}^R the left and right trees of \mathcal{T} as before. Then the edges of \mathcal{T}^L correspond to the atoms of the Bruhat interval $[u_h, u_h s(1, n)]$ while the edges of \mathcal{T}^R correspond to the coatoms of the Bruhat interval $[u_h, u_h s(1, n)]$. More precisely,*

$$\begin{aligned} \{(i, j) \mid u_h < u_h t_{i,j} \leq u_h s(1, n)\} &= \{(i, j) \mid \{i-1, j-1\} \in E(\mathcal{T}^L)\}, \\ \{(i, j) \mid u_h \leq u_h s(1, n) t_{i,j} < u_h s(1, n)\} &= \{(i, j) \mid \{i, j\} \in E(\mathcal{T}^R)\}. \end{aligned}$$

Here, $x < y$ means that $x < y$ and there is no z such that $x < z < y$.

Example 5.20. For $u = 31687524$, the triangulation $\mathcal{T} = \psi(u)$ is as shown in Figure 18. Since $v^{-1} = u_h = 142798635$ and $w^{-1} = u_h s(1, 8) = 427986351$, there are eight atoms of the interval $[v^{-1}, w^{-1}]$ given by $v^{-1} t_{i,j}$, where (i, j) is one of the following pairs:

$$(1, 2), (1, 3), (3, 4), (4, 5), (4, 6), (3, 7), (3, 8), (8, 9).$$

These pairs provide the edges of \mathcal{T}^L by subtracting 1 from every component. On the other hand, there are eight coatoms given by $w^{-1}t_{i,j}$, where (i, j) is one of the following pairs:

$$(1, 2), (2, 9), (3, 6), (4, 5), (5, 6), (6, 7), (7, 9), (8, 9).$$

These pairs are the edges of \mathcal{T}^R .

Theorem 5.21 ([68, Theorem 7.4]). *For $u \in S_n$, the normal fan of the Bruhat interval polytope Q_w^v for $(v^{-1}, w^{-1}) = (u_h, u_h s(1, n))$ is the fan $\Sigma_{\psi(u)}$ associated with the triangulation $\psi(u)$ of P_{n+2} .*

Hence, any n -dimensional fan of Catalan type is realized as the normal fan of Q_w^v with $(v^{-1}, w^{-1}) = (u_h, u_h s(1, n))$ for some $u \in S_n$. As for the latter case $(v^{-1}, w^{-1}) = (u_t, u_t s(n, 1))$, we have the following.

Theorem 5.22 ([68, Theorem 7.5]). *For $u \in S_n$, the normal fan of the Bruhat interval polytope Q_w^v for $(v^{-1}, w^{-1}) = (u_t, u_t s(n, 1))$ is isomorphic to the fan $\Sigma_{\psi(w_0 u w_0)}$ associated with the triangulation $\psi(w_0 u w_0)$ of P_{n+2} , where w_0 denotes the longest element of S_n .*

The following is a direct consequence of Corollary 5.18.

Corollary 5.23 ([68, Corollary 8.2]). *The number of isomorphism classes of n -dimensional smooth toric Richardson varieties of Catalan type is the Wedderburn–Etherington number b_{n+1} .*

6. PROBLEMS

The study of torus orbit closures in the flag variety is related to the geometry of Schubert varieties and the combinatorics of Bruhat interval polytopes. In this section, we pose some possible avenues for further exploration related to the discussions in this chapter.

6.1. Poincaré polynomial of Y_w^v . It is known that a Schubert variety X_w is smooth (in type A) if its Poincaré polynomial is palindromic (see [14, Theorem 6.0.4 and p.208]). Similarly, the generic torus orbit closure Y_w in X_w is smooth if its Poincaré polynomial is palindromic (see [44, Proposition 5.6]⁸). When Y_w is singular, its Poincaré polynomial still has some restrictions. For instance, Brion’s inequalities (see [19]) tell us that if we express

$$\text{Poin}(Y_w, t) = \sum_{i=0}^d b_i t^{2i}$$

where $d = \dim_{\mathbb{C}} Y_w$, then $\sum_{i=0}^k b_i \leq \sum_{i=0}^k b_{d-i}$ for $0 \leq \forall k \leq [d/2]$.

Problem 6.1. Is it true that $b_k \leq b_{d-k}$ for $0 \leq \forall k \leq [d/2]$?

Recall that Y_w^v is the generic torus orbit closure in the Richardson variety X_w^v . The Poincaré polynomial of Y_w^v is computable by Theorem 4.17 but such a formula is not known for Y_w^v in general unless Y_w^v is smooth.

Problem 6.2. (1) Does the virtual Poincaré polynomial of Y_w^v agree with the Poincaré polynomial $\text{Poin}(Y_w^v, t)$ of Y_w^v ? (See Remark 4.13 for virtual Poincaré polynomials.)

(2) Is there a polynomial $A_w^v(t)$ defined similarly to $A_w(t)$ such that $\text{Poin}(Y_w^v, t) = A_w^v(t^2)$?

(3) Similarly to [44, Proposition 5.6], is Y_w^v smooth if $\text{Poin}(Y_w^v, t)$ is palindromic?

6.2. Combinatorics of Q_w^v . We pose five problems on the combinatorics of Bruhat interval polytopes.

There are many pairs (v, w) such that the Bruhat interval polytope Q_w^v is a cube. However, those pairs are not completely understood.

Problem 6.3. Find all pairs (v, w) such that Q_w^v is combinatorially equivalent to a cube, equivalently X_w^v is a smooth toric variety.

The following is a restatement of Conjecture 5.4.

⁸We notice that in the previous version of this article, we posed a problem: *Is the generic torus orbit closure Y_w in X_w smooth if its Poincaré polynomial is palindromic?* and it has been solved in [44, Proposition 5.6].

Problem 6.4. Is \mathbb{Q}_w^v simple when the two vertices v and w are simple in \mathbb{Q}_w^v ?

Since \mathbb{Q}_w^v and $\mathbb{Q}_{w^{-1}}^{v^{-1}}$ have the same dimension, \mathbb{Q}_w^v is toric if and only if $\mathbb{Q}_{w^{-1}}^{v^{-1}}$ is toric. Moreover, when \mathbb{Q}_w^v is toric, its combinatorial type is determined by the poset structure of $[v, w]$. Therefore, when \mathbb{Q}_w^v is toric, $\mathbb{Q}_{w^{-1}}^{v^{-1}}$ is simple if \mathbb{Q}_w^v is simple. We ask whether we can drop the condition *toric* in this statement.

Problem 6.5. Is $\mathbb{Q}_{w^{-1}}^{v^{-1}}$ simple if \mathbb{Q}_w^v is simple?

In the proof of Theorems 4.32 and 4.33, we use the fact that if s_r does not appear in a reduced decomposition of w and $v \leq w$, then $\mathbb{Q}_{s_r w}^v$ and $\mathbb{Q}_{w s_r}^v$ are combinatorially equivalent to $\mathbb{Q}_w^v \times \mathbb{Q}_{s_r}$ (see [65, Proposition 5.7]). We wonder whether this fact can be generalized as follows.

Problem 6.6. Consider $u, v, w \in S_n$ satisfying $v \leq w$ and $[e, u] \cap [e, w] = \{e\}$. Are \mathbb{Q}_{wu}^v and \mathbb{Q}_{uv}^w combinatorially equivalent to $\mathbb{Q}_w^v \times \mathbb{Q}_u$?

We set

$$c(v, w) := \ell(w) - \ell(v) - \dim \mathbb{Q}_w^v.$$

It is the complexity of the T -action on the Richardson variety X_w^v . Note that $c(e, w) = c(w)$, see Definition 4.19 for $c(w)$. We note that studying the complexity $c(v, w)$ of a Richardson variety is closely related to studying the complexity of the corresponding *Kazhdan–Lusztig variety* as is shown in [32, Corollary 4.16].

Recall from Theorem 4.32 that a polytope \mathbb{Q}_w is simple and $c(w) = 1$ if and only if \mathbb{Q}_w is combinatorially equivalent to the product of the hexagon and the cube of dimension $\ell(w) - 3$. Moreover, a polytope \mathbb{Q}_w^v is simple and $c(v, w) = 0$ if and only if \mathbb{Q}_w^v is combinatorially equivalent to a cube (see Theorem 4.20 and Theorem 5.7). Therefore it is natural to ask the following:

Problem 6.7. Is a polytope \mathbb{Q}_w^v simple and $c(v, w) = 1$ if and only if \mathbb{Q}_w^v is combinatorially equivalent to the product of the hexagon and the cube of dimension $\ell(w) - \ell(v) - 3$?

APPENDIX A. TORIC VARIETIES

A.1. Toric varieties. We recall the background of the theory of toric varieties from [28]. We first recall the definition of toric varieties.

Definition A.1 ([28, Definition 3.1.1]). A *toric variety* of complex dimension n is a normal algebraic variety containing an algebraic torus $T := (\mathbb{C}^*)^n$ as a Zariski open dense subset such that the action of the torus on itself extends to the whole variety.

Definition A.2 ([28, Definition 3.1.2]). Let N be a lattice. A *fan* Σ in $N_{\mathbb{R}} := N \otimes_{\mathbb{Z}} \mathbb{R}$ is a finite collection of cones $\sigma \subseteq N_{\mathbb{R}}$ such that:

- (1) Every $\sigma \in \Sigma$ is a strongly convex rational polyhedral cone, i.e., for each σ there exists a finite set $S \subset N$ such that

$$\sigma = \text{Conv}(S) := \left\{ \sum_{u \in S} c_u u \mid c_u \geq 0 \right\} \subset N_{\mathbb{R}}$$

and $\sigma \cap (-\sigma) = \{\mathbf{0}\}$.

- (2) For all $\sigma \in \Sigma$, each face of σ is also in Σ .
- (3) For all $\sigma_1, \sigma_2 \in \Sigma$, the intersection $\sigma_1 \cap \sigma_2$ is a face of each.

Let Σ be a fan in $N_{\mathbb{R}}$. Let M be a dual lattice of N and we set $M_{\mathbb{R}} := M \otimes_{\mathbb{Z}} \mathbb{R}$. We denote by $\langle \cdot, \cdot \rangle$ the pairing between $M_{\mathbb{R}} := M \otimes_{\mathbb{Z}} \mathbb{R}$ and $N_{\mathbb{R}}$. Each cone $\sigma \in \Sigma$ gives an affine toric variety

$$U_{\sigma} = \text{Spec}(\mathbb{C}[\sigma^{\vee} \cap M]),$$

where

$$\sigma^{\vee} := \{m \in M_{\mathbb{R}} \mid \langle m, u \rangle \geq 0 \text{ for all } u \in \sigma\}.$$

By gluing these affine toric varieties, we get a variety X_{Σ} . It turns out that this variety is a toric variety and there is a correspondence between normal separated toric varieties and fans.

Theorem A.3 ([28, Theorem 3.1.5 and Corollary 3.1.8]). *For a fan Σ , the variety X_Σ is a normal separated toric variety. Conversely, for a normal separated toric variety X , there exists a fan Σ such that X is isomorphic to X_Σ .*

A *convex polytope* is the convex hull of a finite set of points in the Euclidean space \mathbb{R}^n . It is well known that every convex polytope is a bounded intersection of finitely many half-spaces. Two polytopes are *combinatorially equivalent* if their face posets are isomorphic. A *lattice polytope* is a convex polytope whose vertices are in the lattice $\mathbb{Z}^n \subset \mathbb{R}^n$.

For a full dimensional lattice polytope $P \subset \mathbb{R}^n$, one can associate a fan, called the *normal fan* of P . Consider the presentation of P given by the intersection of half-spaces:

$$P = \{m \in \mathbb{R}^n \mid \langle m, u_F \rangle \geq -a_F \text{ for every facet } F\}.$$

For a face Q of P , we set

$$\sigma_Q := \text{Cone}(u_F \mid F \text{ contains } Q).$$

Thus the cone σ_F is the ray generated by u_F for a facet F and $\sigma_P = \{0\}$. It is known that

$$\Sigma_P := \{\sigma_Q \mid Q \text{ is a face of } P\}$$

becomes a fan (see [28, Theorem 2.3.2]) and we call it the *normal fan* of P . Moreover, we have the following correspondence.

Theorem A.4 ([28, Theorem 6.2.1]).

$$\begin{array}{c} \{P \subset \mathbb{R}^n \mid P \text{ is a full-dimensional lattice polytope}\} \\ \updownarrow_{1-1} \\ \{(X_P, D) \mid X_P = X_{\Sigma_P} \text{ is a projective toric variety, } D \text{ is a } (\mathbb{C}^*)^n\text{-invariant ample divisor on } X_P\}. \end{array}$$

Furthermore, the torus invariant subvarieties in X_P correspond to faces of the polytope P . For example, the vertices of P correspond to the T -fixed points of X_P .

For a vertex v of a polytope P , the *degree* $d(v)$ of v is the number of edges meeting at v . For an n -dimensional polytope P , a vertex v of P is said to be *simple* if $d(v) = n$. When all the vertices of P are simple, we call P a *simple polytope*.

A vertex v of a lattice polytope P is said to be *smooth* if it is simple and the primitive direction vectors of the edges emanating from v form a basis for \mathbb{Z}^n . We call a vertex of P *singular* if it is not smooth. A lattice polytope P is said to be *smooth* if all the vertices of P are smooth. We call a lattice polytope P *singular* if some vertex of P is singular. See Figure 21.




FIGURE 21. Examples of singular or smooth lattice polytopes.

As one may expect, there are geometric interpretations of these terminologies.

Proposition A.5 ([28, Theorem 2.4.3]). *Let P be a lattice polytope and let X_P be the corresponding toric variety. A vertex v of P is smooth if and only if X_P is smooth at the corresponding fixed point. Moreover, X_P is smooth if and only if the polytope P is smooth.*

There is a combinatorial way to determine whether a smooth compact toric variety is Fano.

Definition A.6. For a fan Σ , a subset R of the primitive ray vectors is called a *primitive collection* of Σ if

$$\text{Cone}(R) \notin \Sigma \quad \text{but} \quad \text{Cone}(R \setminus \{\mathbf{u}\}) \in \Sigma \quad \text{for every } \mathbf{u} \in R.$$

Note that primitive collections of Σ_P correspond to the minimal non-faces of P . For a primitive collection $R = \{\mathbf{u}'_1, \dots, \mathbf{u}'_\ell\}$, we get $\mathbf{u}'_1 + \dots + \mathbf{u}'_\ell = \mathbf{0}$ or there exists a unique cone σ such that $\mathbf{u}'_1 + \dots + \mathbf{u}'_\ell$ is in the interior of σ . That is,

$$(A.1) \quad \mathbf{u}'_1 + \dots + \mathbf{u}'_\ell = \begin{cases} \mathbf{0}, & \text{or} \\ a_1 \mathbf{u}_1 + \dots + a_r \mathbf{u}_r, \end{cases}$$

where $\mathbf{u}_1, \dots, \mathbf{u}_r$ are the primitive generators of σ and a_1, \dots, a_r are positive integers. We call (A.1) a primitive relation, and the *degree* $\deg R$ of a primitive collection R is defined to be $\ell - (a_1 + \dots + a_r)$. Batyrev [11] gave a criterion for a projective toric variety to be Fano or weak Fano.

Proposition A.7 ([11, Proposition 2.3.6]). *A smooth compact toric variety X_Σ is Fano (respectively, weak Fano) if and only if $\deg(R) > 0$ (respectively, $\deg(R) \geq 0$) for every primitive collection R of Σ .*

We can also distinguish two smooth Fano toric varieties using the primitive relations.

Proposition A.8 ([11, Proposition 2.1.8 and Theorem 2.2.4]). *Two smooth Fano toric varieties X_Σ and $X_{\Sigma'}$ are isomorphic as toric varieties if and only if there is a bijection between the sets of rays of Σ and Σ' inducing a bijection between maximal cones and preserving the primitive relations.*

A.2. Fano Bott manifolds. One of the interesting families of smooth toric varieties is the family of Bott manifolds. In this subsection, we first recall the definition of a Bott manifold, and then we characterize Fano Bott manifolds and classify them.

Definition A.9 ([48]). *A Bott tower is an iterated $\mathbb{C}P^1$ -bundle starting with a point:*

$$\begin{array}{ccccccc} B_n & \longrightarrow & B_{n-1} & \longrightarrow & \cdots & \longrightarrow & B_1 & \longrightarrow & B_0, \\ \parallel & & \parallel & & & & \parallel & & \parallel \\ P(\underline{\mathbb{C}} \oplus \xi_n) & & P(\underline{\mathbb{C}} \oplus \xi_{n-1}) & & & & \mathbb{C}P^1 & & \{\text{a point}\} \end{array}$$

where each B_i is the complex projectivization $P(\underline{\mathbb{C}} \oplus \xi_i)$ of the Whitney sum of a holomorphic line bundle ξ_i and the trivial line bundle $\underline{\mathbb{C}}$ over B_{i-1} . The total space B_n is called a *Bott manifold*.

Each Bott manifold is a smooth projective toric variety associated with a smooth lattice polytope combinatorially equivalent to a cube, and the converse also holds.

Proposition A.10 ([72, Corollary 3.5]). *If a smooth lattice polytope P is combinatorially equivalent to a cube, then the toric variety X_P is a Bott manifold. Indeed, the family of Bott manifolds is*

$$\{X_P \mid P \text{ is a smooth lattice polytope that is combinatorially equivalent to a cube}\}.$$

Let B_n be a Bott manifold. Then there is a smooth lattice polytope P combinatorially equivalent to a cube of dimension n such that $B_n = X_P$. The polytope P has $2n$ facets and there are n pairs of facets not intersecting with each other. Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n, \mathbf{w}_1, \dots, \mathbf{w}_n\}$ be the ray generators of the normal fan Σ_P of P such that \mathbf{v}_i and \mathbf{w}_i do not form a cone in Σ_P for each $i \in [n]$. That is, each primitive collection of Σ_P corresponds to the set $\{\mathbf{v}_i, \mathbf{w}_i\}$. By Proposition A.7, we obtain

$$(A.2) \quad B_n \text{ is Fano if and only if } \mathbf{v}_i + \mathbf{w}_i \text{ is either } \mathbf{v}_j \text{ or } \mathbf{w}_j \text{ for some } j \text{ unless } \mathbf{v}_i + \mathbf{w}_i = \mathbf{0}.$$

Using (A.2), to a Fano Bott manifold B_n we associate a signed rooted forest with the vertex set $[n]$ as follows:

- vertex i is a root if $\mathbf{v}_i + \mathbf{w}_i = \mathbf{0}$,
- we draw an edge with $+$ sign between i and j if $\mathbf{v}_i + \mathbf{w}_i = \mathbf{v}_j$, and
- we draw an edge with $-$ sign between i and j if $\mathbf{v}_i + \mathbf{w}_i = \mathbf{w}_j$.

We can also construct a Fano Bott manifold from a signed rooted forest up to isomorphism.

Let F be a signed rooted forest with vertex set $[n]$. For each i , by changing the signs of all edges connecting i and its children simultaneously, we get a new signed rooted forest $r_i(F)$. Then the Bott manifold $B_{r_i(F)}$ corresponding to $r_i(F)$ is isomorphic to the Bott manifold B_F corresponding to F by Proposition A.8. Let \mathcal{SF}_n be the set of all signed rooted forests on the vertex set $[n]$. Denote by \sim the equivalence relation on \mathcal{SF}_n generated by r_i 's for all i 's.

Theorem A.11 ([27, Theorem 3.2]). *The isomorphism classes in Fano Bott manifolds of complex dimension n bijectively correspond to \mathcal{SF}_n/\sim .*

We provide all signed rooted forests having three vertices in Figure 22. Among these ten signed rooted forests, there are four equivalence classes in \mathcal{SF}_3/\sim as follows:

$$(1) \sim (2) \sim (3) \sim (4), \quad (5) \sim (7), \quad (6), \quad (8) \sim (9), \quad (10)$$

In the above, the first and the third equivalence classes arise as toric Richardson varieties of Catalan type, and the last two arise as products of Catalan type. However, the second class does not arise as (products of) Catalan type.




FIGURE 22. Signed rooted forests with 3 vertices

Similarly, there are thirteen equivalence classes in \mathcal{SF}_4/\sim as shown in Figure 23, where + signs on edges are omitted. In the figure, (7), (9), (11) arise as toric Richardson varieties of Catalan type, and (1), (2), (3), (4), (6) arise as products of Catalan type. However, the remaining ones do not arise as (products of) Catalan type.




FIGURE 23. Representatives of \mathcal{SF}_4/\sim .

REFERENCES

[1] Hiraku Abe, Megumi Harada, Tatsuya Horiguchi, and Mikiya Masuda. The cohomology rings of regular nilpotent Hessenberg varieties in Lie type A. *Int. Math. Res. Not. IMRN*, (17):5316–5388, 2019.

- [2] Hiraku Abe and Tatsuya Horiguchi. A survey of recent developments on Hessenberg varieties. In *Schubert calculus and its applications in combinatorics and representation theory*, volume 332 of *Springer Proc. Math. Stat.*, pages 251–279. Springer, Singapore, [2020] ©2020.
- [3] Hiraku Abe and Haozhi Zeng. The integral cohomology rings of Peterson varieties in type A . [arXiv:2203.02629](#), 2022.
- [4] Takuro Abe, Tatsuya Horiguchi, Mikiya Masuda, Satoshi Murai, and Takashi Sato. Hessenberg varieties and hyperplane arrangements. *J. Reine Angew. Math.*, 764:241–286, 2020.
- [5] Alfred V. Aho, Michael R. Garey, and Jeffrey David Ullman. The transitive reduction of a directed graph. *SIAM J. Comput.*, 1(2):131–137, 1972.
- [6] Federico Ardila, Carolina Benedetti, and Jeffrey Doker. Matroid polytopes and their volumes. *Discrete Comput. Geom.*, 43(4):841–854, 2010.
- [7] Federico Ardila, Federico Castillo, Christopher Eur, and Alexander Postnikov. Coxeter submodular functions and deformations of Coxeter permutahedra. *Adv. Math.*, 365:107039, 36, 2020.
- [8] Michael Francis Atiyah. Convexity and commuting Hamiltonians. *Bull. London Math. Soc.*, 14(1):1–15, 1982.
- [9] Michèle Audin. *Torus actions on symplectic manifolds*, volume 93 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, revised edition, 2004.
- [10] Anthony Bahri, Soumen Sarkar, and Jongbaek Song. On the integral cohomology ring of toric orbifolds and singular toric varieties. *Algebr. Geom. Topol.*, 17(6):3779–3810, 2017.
- [11] Victor V. Batyrev. On the classification of toric Fano 4-folds. volume 94, pages 1021–1050. 1999. Algebraic geometry, 9.
- [12] Carolina Benedetti and Kolja Knauer. Lattice path matroids and quotients. *Combinatorica*, 2024. Published online at 04 April 2024.
- [13] Andrzej S. Białynicki-Birula. Some theorems on actions of algebraic groups. *Ann. of Math. (2)*, 98:480–497, 1973.
- [14] Sara Billey and V. Lakshmibai. *Singular loci of Schubert varieties*, volume 182 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 2000.
- [15] Anders Björner and Francesco Brenti. *Combinatorics of Coxeter groups*, volume 231 of *Graduate Texts in Mathematics*. Springer, New York, 2005.
- [16] Armand Borel. Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts. *Ann. of Math. (2)*, 57:115–207, 1953.
- [17] Alexandre V. Borovik, I. M. Gelfand, and Neil White. *Coxeter matroids*, volume 216 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 2003.
- [18] Mireille Bousquet-Mélou and Steve Butler. Forest-like permutations. *Ann. Comb.*, 11(3-4):335–354, 2007.
- [19] Michel Brion. Poincaré duality and equivariant (co)homology. *Michigan Math. J.*, 48:77–92, 2000. Dedicated to William Fulton on the occasion of his 60th birthday.
- [20] Michel Brion. Lectures on the geometry of flag varieties. In *Topics in cohomological studies of algebraic varieties*, Trends Math., pages 33–85. Birkhäuser, Basel, 2005.
- [21] Patrick Brosnan and Timothy Y. Chow. Unit interval orders and the dot action on the cohomology of regular semisimple Hessenberg varieties. *Adv. Math.*, 329:955–1001, 2018.
- [22] Victor M. Buchstaber and Taras E. Panov. *Toric topology*, volume 204 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015.
- [23] Amanda Cameron, Rodica Dinu, Mateusz Michałek, and Tim Seynnaeve. Flag matroids: algebra and geometry. In *Interactions with lattice polytopes*, volume 386 of *Springer Proc. Math. Stat.*, pages 73–114. Springer, Cham, [2022] ©2022.
- [24] Mahir Bilen Can and Pinakinath Saha. Toric Richardson varieties. [arXiv:2310.09901](#).
- [25] James B. Carrell and Alexandre Kurth. Normality of torus orbit closures in G/P . *J. Algebra*, 233(1):122–134, 2000.
- [26] James B. Carrell and Jochen Kuttler. Smooth points of T -stable varieties in G/B and the Peterson map. *Invent. Math.*, 151(2):353–379, 2003.
- [27] Yunhyung Cho, Eunjeong Lee, Mikiya Masuda, and Seonjeong Park. On the enumeration of Fano Bott manifolds. [arXiv:2106.12788v1](#), to appear in Fields Institute Communications “Toric Topology and Polyhedral Products”.
- [28] David A. Cox, John B. Little, and Henry K. Schenck. *Toric varieties*, volume 124 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2011.
- [29] Romuald Dabrowski. On normality of the closure of a generic torus orbit in G/P . *Pacific J. Math.*, 172(2):321–330, 1996.
- [30] Daniel Daly. Reduced decompositions with one repetition and permutation pattern avoidance. *Graphs Combin.*, 29(2):173–185, 2013.
- [31] F. De Mari, C. Procesi, and M. A. Shayman. Hessenberg varieties. *Trans. Amer. Math. Soc.*, 332(2):529–534, 1992.
- [32] Maria Donten-Bury, Laura Escobar, and Irem Portakal. Complexity of the usual torus action on Kazhdan-Lusztig varieties. *Algebr. Comb.*, 6(3):835–861, 2023.

- [33] Jack Edmonds. Submodular functions, matroids, and certain polyhedra. In *Combinatorial Structures and their Applications (Proc. Calgary Internat. Conf., Calgary, Alta., 1969)*, pages 69–87. Gordon and Breach, New York, 1970.
- [34] Serge Elnitsky. Rhombic tilings of polygons and classes of reduced words in Coxeter groups. *J. Combin. Theory Ser. A*, 77(2):193–221, 1997.
- [35] Laura Escobar, Oliver Pechenik, Bridget Eileen Tenner, and Alexander Yong. Rhombic tilings and Bott–Samelson varieties. *Proc. Amer. Math. Soc.*, 146(5):1921–1935, 2018.
- [36] Ivor Malcolm Haddon Etherington. Non-associate powers and a functional equation. *The Mathematical Gazette*, 21(242):36–39, 1937.
- [37] C. Kenneth Fan. Schubert varieties and short braidedness. *Transform. Groups*, 3(1):51–56, 1998.
- [38] Hermann Flaschka and Luc Haine. Torus orbits in G/P . *Pacific J. Math.*, 149(2):251–292, 1991.
- [39] Sergey Fomin and Nathan Reading. Root systems and generalized associahedra. In *Geometric combinatorics*, volume 13 of *IAS/Park City Math. Ser.*, pages 63–131. Amer. Math. Soc., Providence, RI, 2007.
- [40] Naoki Fujita, Eunjeong Lee, and Dong Youp Suh. Algebraic and geometric properties of flag Bott–Samelson varieties and applications to representations. *Pacific J. Math.*, 309(1):145–194, 2020.
- [41] William Fulton. *Introduction to toric varieties*, volume 131 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1993. The William H. Roever Lectures in Geometry.
- [42] William Fulton. *Young tableaux*, volume 35 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1997. With applications to representation theory and geometry.
- [43] William Fulton and Joe Harris. *Representation theory*, volume 129 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1991. A first course, Readings in Mathematics.
- [44] Christian Gaetz. One-skeleton posets of Bruhat interval polytopes. *Adv. Math.*, 430:Paper No. 109216, 22, 2023.
- [45] Izrail’ Moiseevich Gel’fand, Mark Goresky, Robert D. MacPherson, and Vera V. Serganova. Combinatorial geometries, convex polyhedra, and Schubert cells. *Adv. in Math.*, 63(3):301–316, 1987.
- [46] Izrail’ Moiseevich Gel’fand and Vera V. Serganova. Combinatorial geometries and the strata of a torus on homogeneous compact manifolds. *Uspekhi Mat. Nauk*, 42(2(254)):107–134, 287, 1987.
- [47] Richard M. Green and Jozsef Losonczy. Freely braided elements of Coxeter groups. *Ann. Comb.*, 6(3-4):337–348, 2002.
- [48] Michael Grossberg and Yael Karshon. Bott towers, complete integrability, and the extended character of representations. *Duke Math. J.*, 76(1):23–58, 1994.
- [49] Victor W. Guillemin, Tara Suzanne Holm, and Cătălin Zara. A GKM description of the equivariant cohomology ring of a homogeneous space. *J. Algebraic Combin.*, 23(1):21–41, 2006.
- [50] Victor W. Guillemin and Shlomo Sternberg. Convexity properties of the moment mapping. *Invent. Math.*, 67(3):491–513, 1982.
- [51] Megumi Harada, Tatsuya Horiguchi, Mikiya Masuda, and Seonjeong Park. The volume polynomial of regular semisimple Hessenberg varieties and the Gelfand–Tsetlin polytope. *Proc. Steklov Inst. Math.*, 305(Algebraicheskaya Topologiya Kombinatorika i Matematicheskaya Fizika):344–373, 2019.
- [52] Christophe Hohlweg, Carsten E. M. C. Lange, and Hugh Thomas. Permutahedra and generalized associahedra. *Adv. Math.*, 226(1):608–640, 2011.
- [53] Tatsuya Horiguchi and Megumi Harada. The cohomology rings of regular nilpotent Hessenberg varieties. Chapter 9 of *Handbook of Combinatorial Algebraic Geometry: Subvarieties of the flag variety*.
- [54] James E. Humphreys. *Reflection groups and Coxeter groups*, volume 29 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1990.
- [55] Jens Carsten Jantzen. *Representations of algebraic groups*, volume 107 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, second edition, 2003.
- [56] Paramasamy Karuppuchamy. On Schubert varieties. *Comm. Algebra*, 41(4):1365–1368, 2013.
- [57] Aleksandr Anatol’evich Klyachko. Orbits of a maximal torus on a flag space. *Functional Analysis and Its Applications*, 19(1):65–66, 1985.
- [58] Yuji Kodama and Lauren Williams. The full Kostant–Toda hierarchy on the positive flag variety. *Comm. Math. Phys.*, 335(1):247–283, 2015.
- [59] János Kollár. *Fano Varieties*, pages 238–284. Springer Berlin Heidelberg, Berlin, Heidelberg, 1996.
- [60] János Kollár, Yoichi Miyaoka, and Shigefumi Mori. Rational connectedness and boundedness of Fano manifolds. *J. Differential Geom.*, 36(3):765–779, 1992.
- [61] Shintarō Kuroki, Eunjeong Lee, Jongbaek Song, and Dong Youp Suh. Flag Bott manifolds and the toric closure of a generic orbit associated to a generalized Bott manifold. *Pacific J. Math.*, 308(2):347–392, 2020.
- [62] Venkatramani Lakshmibai and B. Sandhya. Criterion for smoothness of Schubert varieties in $Sl(n)/B$. *Proc. Indian Acad. Sci. Math. Sci.*, 100(1):45–52, 1990.
- [63] Eunjeong Lee and Mikiya Masuda. Generic torus orbit closures in Schubert varieties. *J. Combin. Theory Ser. A*, 170:105143, 44pp, 2020.
- [64] Eunjeong Lee, Mikiya Masuda, and Seonjeong Park. Toric Schubert varieties and directed Dynkin diagrams. [arXiv:2311.11535v1](https://arxiv.org/abs/2311.11535).
- [65] Eunjeong Lee, Mikiya Masuda, and Seonjeong Park. On Schubert varieties of complexity one. *Pacific J. Math.*, 315(2):419–447, 2021.

- [66] Eunjeong Lee, Mikiya Masuda, and Seonjeong Park. Toric Bruhat interval polytopes. *J. Combin. Theory Ser. A*, 179:105387, 41pp, 2021.
- [67] Eunjeong Lee, Mikiya Masuda, and Seonjeong Park. Torus orbit closures in flag varieties and retractions on Weyl groups. *Internat. J. Math.*, 33(4):Paper No. 2250028, 21, 2022.
- [68] Eunjeong Lee, Mikiya Masuda, and Seonjeong Park. Toric Richardson varieties of Catalan type and Wedderburn–Etherington numbers. *European J. Combin.*, 108:Paper No. 103617, 25, 2023.
- [69] Eunjeong Lee, Mikiya Masuda, Seonjeong Park, and Jongbaek Song. Poincaré polynomials of generic torus orbit closures in Schubert varieties V. A. Rokhlin-Memorial. In *Topology, geometry, and dynamics*, volume 772 of *Contemp. Math.*, pages 189–208. Amer. Math. Soc., [Providence], RI, [2021] ©2021.
- [70] Jean-Louis Loday, Alessandra Frabetti, Frédéric Chapoton, and François Goichot. *Dialgebras and related operads*, volume 1763 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2001.
- [71] Peter Magyar. Schubert polynomials and Bott–Samelson varieties. *Comment. Math. Helv.*, 73(4):603–636, 1998.
- [72] Mikiya Masuda and Taras E. Panov. Semi-free circle actions, Bott towers, and quasitoric manifolds. *Mat. Sb.*, 199(8):95–122, 2008.
- [73] Jacqueline Morand. Closures of torus orbits in adjoint representations of semisimple groups. *C. R. Acad. Sci. Paris Sér. I Math.*, 328(3):197–202, 1999.
- [74] Kazuo Murota. *Discrete convex analysis*. SIAM Monographs on Discrete Mathematics and Applications. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2003.
- [75] N. J. A. Sloane, editor. The On-Line Encyclopedia of Integer Sequences (OEIS). published electronically at <https://oeis.org> (accessed May 30, 2024).
- [76] Seonjeong Park and Jongbaek Song. Conic decomposition of a toric variety and its application to cohomology. *Proc. Amer. Math. Soc.*, 150(7):2777–2792, 2022.
- [77] Alex Postnikov, Victor Reiner, and Lauren Williams. Faces of generalized permutohedra. *Doc. Math.*, 13:207–273, 2008.
- [78] Alexander Postnikov. Permutohedra, associahedra, and beyond. *Int. Math. Res. Not. IMRN*, (6):1026–1106, 2009.
- [79] Nathan Paul Reading. *On the structure of Bruhat order*. ProQuest LLC, Ann Arbor, MI, 2002. Thesis (Ph.D.)–University of Minnesota.
- [80] Edward Richmond and William Slofstra. The isomorphism problem for Schubert varieties. [arXiv:2103.08114v1](https://arxiv.org/abs/2103.08114), 2021.
- [81] Alexander Schrijver. *Combinatorial optimization. Polyhedra and efficiency. Vol. B*, volume 24 of *Algorithms and Combinatorics*. Springer-Verlag, Berlin, 2003. Matroids, trees, stable sets, Chapters 39–69.
- [82] T. A. Springer. *Linear algebraic groups*. Modern Birkhäuser Classics. Birkhäuser Boston, Inc., Boston, MA, second edition, 2009.
- [83] Richard P. Stanley. *Catalan numbers*. Cambridge University Press, New York, 2015.
- [84] Bridget Eileen Tenner. Pattern avoidance and the Bruhat order. *J. Combin. Theory Ser. A*, 114(5):888–905, 2007.
- [85] Bridget Eileen Tenner. Repetition in reduced decompositions. *Adv. in Appl. Math.*, 49(1):1–14, 2012.
- [86] Emmanuel Tsukerman and Lauren K. Williams. Bruhat interval polytopes. *Adv. Math.*, 285:766–810, 2015.
- [87] Julianna S. Tymoczko. Permutation actions on equivariant cohomology of flag varieties. In *Toric topology*, volume 460 of *Contemp. Math.*, pages 365–384. Amer. Math. Soc., Providence, RI, 2008.
- [88] Joseph Henry Maclagan Wedderburn. The functional equation $g(x^2) = 2\alpha x + [g(x)]^2$. *Ann. of Math. (2)*, 24(2):121–140, 1922.
- [89] Neil White, editor. *Theory of matroids*, volume 26 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1986.
- [90] Hassler Whitney. On the abstract properties of linear dependence. *Amer. J. Math.*, 57(3):509–533, 1935.
- [91] Alexander Woo and Alexander Yong. When is a Schubert variety Gorenstein? *Adv. Math.*, 207(1):205–220, 2006.
- [92] Li Yu and Mikiya Masuda. On Descriptions of Products of Simplices. *Chin. Ann. Math. Ser. B*, 42(5):777–790, 2021.
- [93] Andrei Zelevinsky. Nested complexes and their polyhedral realizations. *Pure Appl. Math. Q.*, 2(3, Special Issue: In honor of Robert D. MacPherson. Part 1):655–671, 2006.
- [94] Günter M. Ziegler. *Lectures on polytopes*, volume 152 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.

(E. Lee) DEPARTMENT OF MATHEMATICS, CHUNGBUK NATIONAL UNIVERSITY, CHEONGJU 28644, REPUBLIC OF KOREA

Email address: `eunjeong.lee@chungbuk.ac.kr`

(M. Masuda) OSAKA CITY UNIVERSITY ADVANCED MATHEMATICS INSTITUTE (OCAMI) & DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA CITY UNIVERSITY, SUMIYOSHI-KU, SUGIMOTO, 558-8585, OSAKA, JAPAN

Email address: `mikiyamsd@gmail.com`

(S. Park) DEPARTMENT OF MATHEMATICS EDUCATION, JEONJU UNIVERSITY, JEONJU 55069, REPUBLIC OF KOREA

Email address: `seonjeongpark@jj.ac.kr`