arXiv:2203.16830v1 [math.GT] 31 Mar 2022

THE MINIMAL DIMENSION OF A SPHERE WITH AN EQUIVARIANT EMBEDDING OF THE BOUQUET OF g CIRCLES IS 2g - 1

ZHONGZI WANG

ABSTRACT. To embed the bouquet of g circles B_g into the *n*-sphere S^n so that its full symmetry group action extends to an orthogonal actions on S^n , the minimal n is 2g - 1. This answers a question raised by B. Zimmermann.

In this note graphs are finite and connected and group actions are faithful. Denote the *n*-dimensional Euclidean space and orthogonal group by \mathbb{R}^n and O(n), and the unit sphere in \mathbb{R}^{n+1} by S^n .

Graphs belong to a most familiar geometric subject to us and the symmetries of graphs is a big and active research area.

For a given graph Γ with an action of a finite group G, an embedding $e: \Gamma \to S^n$ is G-equivariant, if there is a subgroup $G \subset O(n+1)$ so that G acts on the pair $(S^n, e(\Gamma))$ satisfying g(e(x)) = e(g(x)) for any $x \in \Gamma$ and $g \in G$. In the definition of G-equivariant embedding, one may replace $(S^n, O(n+1))$ by $(\mathbb{R}^n, O(n))$. See Remark 0.5 for relations between the two definitions.

The existence of *G*-equivariant embedding for any pair (G, Γ) is known, see [Mos] and [Pa]. On the other hand, for given Γ and *G*, usually it is difficult to find the minimal *n* so that there is an *G*-equivariant embedding $\Gamma \to \mathbb{R}^n$.

The most significant finite group associated to a graph Γ is its full symmetry group Sym(Γ). The simplest and also most symmetric graph is the bouquet of g circles (the graph with one vertex and g closed edges, see Figure 1), denoted as B_g . B_g is the simplest in the sense that B_g has only one vertex. B_g is the most symmetric in the sense that: for $g \geq 3$, the maximum order of finite group action on a hyperbolic graph of rank g (defined as its fundamental group rank) is $2^g g!$, which is realized uniquely by the action of Sym(B_q) on B_q [WZ]. A graph Γ is

²⁰¹⁰ Mathematics Subject Classification. Primary 57M25; Secondary 57S17, 57S25, 05C10.

Key words and phrases. Groups acting on finite graphs, equivariant embeddings into spheres.



FIGURE 1. A bouquet of q circles:

hyperbolic if it has rank > 1 and has no free edge. See Example 0.6 for the rank 2 case.

We will simply use equivariant embedding of Γ for Sym(Γ)-equivariant embedding of Γ .

Recently B. Zimmermann asked a very interesting question (as commented by the Math. Review of the paper).

Question 0.1. ([Zi]) What is the minimal n so that there is an equivariant embedding $B_q \to S^n$?

Zimmermann has pointed that $n \ge g - 1$ from a group theoretical reason. He also described an equivariant embedding of B_g into S^{2g} . So he concluded the minimal n is between g - 1 and 2g [Zi]. We will answer this question.

Theorem 0.2. The minimal n so that there is an equivariant embedding $B_g \to S^n$ is 2g - 1.

Theorem 0.2 follows from the next two propositions.

Proposition 0.3. Suppose there is an equivariant embedding of B_g into S^n , then $n \ge 2g - 1$.

Proposition 0.4. There are equivariant embeddings of B_g into S^{2g-1} for each g > 1.

Proof of Proposition 0.3. We need first recall a careful description of the action of $\operatorname{Sym}(B_g)$ on B_g . Denote the vertex of B_g by v and the g circles by C_1, \ldots, C_g , see Figure 2. From now on we consider that each circle C_i is oriented. It is known that $\operatorname{Sym}(B_g)$ is the semi-direct product $(\mathbb{Z}_2)^g \ltimes S_g$, where S_g is the permutation group of g elements, which permutes those g oriented circles, and the normal subgroup $(\mathbb{Z}_2)^g$ acts by orientation reversing involutions of those g oriented circles.

More precisely, for each $\sigma \in S_g$, denote the corresponding action on B_g by τ_{σ} , which sends C_i to $C_{\sigma(i)}$ and preserves the orientations. Denote the element ρ_i to be the element in $(\mathbb{Z}_2)^g \subset \text{Sym}(B_g)$, which



FIGURE 2. Detailed description $\text{Sym}(B_g)$ action on B_g

is the orientation reversing involution on C_i and is the identity on the remaining C_j , for $j \neq i$. Now ρ_i has two fixed points, one is v, another we denote as p_i . It is clear that

$$\tau_{\sigma}\rho_i\tau_{\sigma^{-1}} = \rho_{\sigma(i)}.$$
 (1)

By (1) we have

$$\rho_{\sigma(i)}\tau_{\sigma}(p_i) = \tau_{\sigma}\rho_i\tau_{\sigma}^{-1}\tau_{\sigma}(p_i) = \tau_{\sigma}\rho_i(p_i) = \tau_{\sigma}(p_i).$$

So we have

$$p_{\sigma(i)}\tau_{\sigma}(p_i) = \tau_{\sigma}(p_i).$$

That is, $\tau_{\sigma}(p_i)$ is the fixed point of $\rho_{\sigma(i)}$ on $C_{\sigma(i)} - \{v\}$. Since the fixed point of $\rho_{\sigma(i)}$ on $C_{\sigma(i)} - v$ is unique, we have

$$\tau_{\sigma}(p_i) = p_{\sigma(i)}. \qquad (2)$$

Suppose now we have an equivariant embedding $B_g \subset S^n$. That is to say we have embedded $\operatorname{Sym}(B_g)$ into O(n+1) which acts on the pair (S^n, B_g) and the restriction on B_g is the action of $\operatorname{Sym}(B_g)$ described as above. Since S^n is the unit sphere of \mathbb{R}^{n+1} and O(n+1) is the orthogonal transformation group of \mathbb{R}^{n+1} , the action of $\operatorname{Sym}(B_g)$ on S^n extends to a $\operatorname{Sym}(B_g)$ -action on \mathbb{R}^{n+1} as an orthogonal transformation group.

Consider each point in \mathbb{R}^{n+1} as a vector in \mathbb{R}^{n+1} . Since $v, p_1, p_2, ..., p_g$ are in the unit sphere, each of them is a non-zero vector. Let

$$V = \langle v, p_1, p_2, \dots, p_g \rangle$$

be the subspace of \mathbb{R}^{n+1} spanned by the vectors $v, p_1, p_2, ..., p_g$.

By (2), the S_g -action on B_g permutes those vectors $p_1, p_2, ..., p_g$ and fixes v. Hence S_g acts faithfully on V, that is $\tau_{\sigma}|_V = \mathrm{id}|_V$ if and only if σ is the identity in S_g .

Now we show that the vectors $v, p_1, ..., p_{g-1}$ are linearly independent: Suppose $a, a_1, ..., a_{g-1}$ are real numbers such that

ZHONGZI WANG

$$av + \sum_{i=1}^{g-1} a_i p_i = 0.$$
 (3)

For each $j \in \{1, 2, ..., g - 1\}$, recall that $(j, g) \in S_g$ exchanges j and g, and fixes all the remaining i, and $\tau_{(j,g)}$ corresponds to the element $(j, g) \in S_g$. Since $\tau_{(j,g)}$ is an orthogonal transformation, $\tau_{(j,g)}$ is linear. Applying $\tau_{(j,g)}$ to (3), we have

$$0 = \tau_{(j,g)}(av + \sum_{i=1}^{g-1} a_i p_i) = a\tau_{(j,g)}(v) + \sum_{i=1}^{g-1} a_i \tau_{(j,g)}(p_i).$$
(4)

Since v is the common fixed point of S_g and $\tau_{(j,g)}(p_i) = p_i$ if $j \neq i$ and $\tau_{(j,g)}(p_j) = p_g$, we have further

$$av + \sum_{i=1, i \neq j}^{g-1} a_i p_i + a_j p_g = 0.$$
 (5)

On the other hand by (3) and some elementary transformations we have

$$av + \sum_{i=1, i \neq j}^{g-1} a_i p_i + a_j p_g = av + \sum_{i=1}^{g-1} a_i p_i - a_j p_j + a_j p_g = a_j (p_g - p_j).$$
(6)

Combine (5) and (6), we have

 $a_j(p_g - p_j) = 0.$

Since $B_g \subset \mathbb{R}^{n+1}$ is an embedding, and $j \neq g$, we have

$$p_g - p_j \neq 0$$

Hence $a_j = 0$. Since j can be any element in $\{1, 2, ..., g - 1\}$. Hence $a_j = 0$ for each $j \in \{1, 2, ..., g - 1\}$. By (3) we have

$$av = 0.$$

Since $v \in S^n$, $v \neq 0$, we have a = 0. That is (3) implies that

$$a = a_1 = a_2 = \dots = a_{g-1} = 0, \qquad (7)$$

that is $v, p_1, ..., p_{g-1}$ are linearly independent. As a conclusion we have

$$\dim V \ge g. \tag{8}$$

Let V^{\perp} be orthogonal complement of V in \mathbb{R}^{n+1} . We have

$$\mathbb{R}^{n+1} = V \oplus V^{\perp}. \qquad (9)$$

For any $i \in \{1, 2, ..., g\}$, ρ_i is orthogonal, so is linear. Moreover each point in $\{v, p_1, p_2, ..., p_g\}$ is a fixed point of ρ_i . Therefore ρ_i is the identity on V, the subspace spanned by $\{v, p_1, p_2, ..., p_g\}$. Further more we conclude that the action of the subgroup $(\mathbb{Z}_2)^g$ on V is trivial. In particular V is an invariant subspace of the $(\mathbb{Z}_2)^g$ -action.

Since the $(\mathbb{Z}_2)^{g}$ -action on \mathbb{R}^{n+1} is orthogonal and V is invariant under the $(\mathbb{Z}_2)^{g}$ -action, $(\mathbb{Z}_2)^{g}$ acts orthogonally on the orthogonal complement V^{\perp} of V. Since the $(\mathbb{Z}_2)^{g}$ -action is trivial on V and is faithful on \mathbb{R}^{n+1} , the $(\mathbb{Z}_2)^{g}$ -action must be faithful on V^{\perp} by (9).

Let q be the dimension of V^{\perp} , and consider O(q) as q by q orthogonal matrix group. Then $(\mathbb{Z}_2)^{g}$ -action on V^{\perp} is by a matrix group in O(q). Let Ω_q be the set of all q by q diagonal matrices whose entries on the diagonal of the matrices are either 1 or -1. Then there are exactly 2^{q} matrices in Ω_q .

Each element in $(\mathbb{Z}_2)^g$ is of order 1 or 2, therefore it can be diagonalized, indeed into Ω_q . By linear algebra, a finite number of commuting diagonalizable matrices can be diagonalized simultaneously. Since $(\mathbb{Z}_2)^g$ is a finite abelian group, we may assume that under a basis of V^{\perp} , all matrices in $(\mathbb{Z}_2)^g$ are diagonalized, therefore are elements in Ω_g . Since $(\mathbb{Z}_2)^g$ has 2^g elements, $(\mathbb{Z}_2)^g$ acts faithfully on V^{\perp} , the image of $(\mathbb{Z}_2)^g$ in Ω_g must be also 2^g pairwise different elements. Therefore $2^g < 2^q$, that is

$$g \le q = \dim V^{\perp}. \tag{10}$$

By (8), (9) and (10) we have

$$n+1 = \dim V + \dim V^{\perp} \ge 2g.$$

That is

$$n \ge 2g - 1.$$

We finish the proof.

Proof of Proposition 0.4. We will contruct two different equivariant embeddings. The first one is an equivariant embedding $B_g \to \mathbb{R}^{2g-1}$, then the required G-equivariant embedding $B_g \to S^{2g-1}$ can be obtained by one point compactification via the inverse of stereographic projection $p: S^{2g-1} \to \mathbb{R}^{2g-1}$. The second one is an equivariant embedding of $B_g \to \mathbb{R}^{2g}$, where B_g stays equivariantly in the unit sphere S^{2g-1} .

The first construction. Before we give a general construction, we like to provide a visible equivariant embedding of $B_2 \subset \mathbb{R}^3$ in the familiar dimension in which we live.

Fix a standard xyz-coordinate system of \mathbb{R}^3 . Let C_1 be the unit circle in the lower-half zx-plane which is tangent to the x-axis at the origin, and C_2 be the unit circle in the upper-half yz-plane which is tangent to

ZHONGZI WANG

the y-axis at the origin, as shown in Figure 3. Then $C_1 \bigcup C_2$ provides an embedding $B_2 \subset \mathbb{R}^3$. The action of $\operatorname{Sym}(B_2) = (\mathbb{Z}_2)^2 \ltimes S_2$ is also realized by a subgroup of O(3): where the inversion of C_1 is given by the reflection about the yz-plane, the inversion of C_2 is given by the reflection about the zx-plane, and the permutation of C_1 and C_2 is given by the π -rotation around a diagonal L in the xy-plane. This provides an equivariant embedding.



FIGURE 3. An equivariant embedding of B_2 into \mathbb{R}^3

Then we provide the equivariant embedding of $B_g \subset \mathbb{R}^{2g-1}$ for g > 2. Let $V_1 = \mathbb{R}^{g-1}$ and $v_1, v_2, ..., v_g$ be the vertices of a regular (g-1)dimensional simplex Δ_{g-1} centered at the origin O_1 of V_1 with the length $|O_1v_i| = 1$. Then S_g , the full symmetry group of Δ_{g-1} , is a subgroup of O(g-1). Let $V_2 = \mathbb{R}^g$, and $e_1, e_2, ..., e_g$ be the standard orthogonal basis of V_2 . Let $V = V_1 \times V_2$, then $V = \mathbb{R}^{2g-1}$, and (v_i, e_i) is a standard orthogonal basis of the Euclidean plane E_i spanned by v_i and $e_i, i \in \{1, ..., g\}$.

Define the isometrical embedding $\iota_i: S^1 \to V$ by

$$\iota_i(\cos\theta, \sin\theta) = (1 + \cos\theta)v_i + \sin\theta e_i, \qquad (11)$$

where $0 \le \theta < 2\pi$ is the parameter on the unit circle.

Let $\iota_i(S^1) = C_i$, $i \in \{1, ..., g\}$. Clearly the origin O of V belongs to each C_i . Moreover C_i belongs to E_i , and $E_i \cap E_j = O$ for $i \neq j$, so $C_i \cap C_j = O$ for $i \neq j$. Hence $\bigcup_{i=1}^g C_i$ is a bouquet of g circles embedded in V with the origin O as the vertex, still denoted as B_q .

For each $i \in \{1, 2, ..., g\}$ and each $\sigma \in S_g$, we define $\iota_i, \tau_\sigma \in O(2g-1)$ as follows

$$\iota_i(v_j) = v_j \text{ for all } j, \ \iota_i(e_j) = e_j \text{ for } j \neq i; \ \iota_i(e_i) = -e_i.$$
(12)

$$\tau_{\sigma}(e_i) = e_{\sigma(i)}, \ \tau_{\sigma}(v_i) = v_{\sigma(i)}.$$
(13)

By (11), (12) and (13), one can check directly

(i) ι_i is the identity on C_j for $j \neq i$ and is an orientation reversing involution on C_i .

(ii) $\tau_{\sigma}(C_i) = C_{\sigma(i)}$ and $\tau_{\sigma}^{-1}(C_i) = C_{\sigma^{-1}(i)}$. Let

$$H = \langle \iota_i, \tau_\sigma : 1 \le i \le g, \sigma \in S_g \rangle.$$

Then H is a subgroup of $\text{Sym}(B_q)$, and each $h \in H$ has the form

$$h = \tau_{\sigma} \prod_{i=1}^{g} \iota_i^{\alpha_i},$$

where $\alpha_i = 0$ or 1, since $\langle \iota_i, 1 \leq i \leq g \rangle$ is a normal subgroup of H. Clearly H acts faithfully on B_g . Moreover if $h \in H$ is the identity on B_g , h must fix all v_i and e_i , which implies that h is the identity on V. So H also acts faithfully on V. The restriction of H on B_g is $\text{Sym}(B_g)$, since it exhausts all symmetries of $\text{Sym}(B_g)$.

Therefore we get an equivariant embedding $B_g \to \mathbb{R}^{2g-1}$.

The second construction. View \mathbb{R}^{2g} as the product

$$\mathbb{R}^{2g} = \mathbb{R}_1^2 \times \mathbb{R}_2^2 \times \dots \times \mathbb{R}_g^2,$$

where each \mathbb{R}_i^2 is an Euclidean plane with standard coordinate system $(x_i, y_i), i = 1, 2, ..., g$. Then $(x_1, y_1, x_2, y_2, ..., x_g, y_g)$ provides a standard coordinate system of the Euclidean space \mathbb{R}^{2g} . Let C_i be the circle in $\mathbb{R}_i^2 \subset \mathbb{R}^{2g}$ given by

$$C_i = \{x_i^2 + y_i^2 = \frac{1}{g}, x_j = \frac{1}{\sqrt{g}}, y_j = 0 \text{ for remaining } j \neq i.\}$$
 (13)

with anti-clockwise orientation in the $x_i y_i$ -plane.

Now we make three observations:

(a) Clearly the point

$$v = (\frac{1}{\sqrt{g}}, 0, \frac{1}{\sqrt{g}}, 0, ..., \frac{1}{\sqrt{g}}, 0) \in \mathbb{R}^{2g}$$

belongs to each C_i , and each pair C_i and C_j intersect only at v for $i \neq j$. Therefore the union $\bigcup_{i=1}^{g} C_i$ provides an embedding of $B_g \to \mathbb{R}^{2g}$.

(b) For each $i \in \{1, 2, ..., g\}$ and each $\sigma \in S_g$, we define

$$\iota_i, \tau_\sigma : \mathbb{R}^{2g} \to \mathbb{R}^{2g}$$

as below:

$$\iota_i(x_1, y_1, \dots, x_{i-1}, y_{i-1}, x_i, y_i, x_{i+1}, y_{i+1}, \dots, x_g, y_g)$$

ZHONGZI WANG

$$= (x_1, y_1, \dots, x_{i-1}, y_{i-1}, x_i, -y_i, x_{i+1}, y_{i+1}, \dots, x_g, y_g).$$
$$\tau_{\sigma}(x_1, y_1, \dots, x_i, y_i, \dots, x_g, y_g)$$
$$= (x_{\sigma(1)}, y_{\sigma(1)}, \dots, x_{\sigma(i)}, y_{\sigma(i)}, \dots, x_{\sigma(g)}, y_{\sigma(g)})$$

Clearly $\iota_i, \tau_{\sigma} \in O(2g)$, ι_i is the identity on C_j for $j \neq i$ and is an orientation reversing involution on C_i . and $\tau_{\sigma}(C_i) = C_{\sigma(i)}$. Moreover the group

$$H = <\iota_i, \tau_\sigma: \ 1 \le i \le g, \ \sigma \in S_g >$$

is isomorphic to $(\mathbb{Z}_2)^g \rtimes S_g \cong \text{Sym}(B_g)$ and acts faithfully on the pair (\mathbb{R}^{2g}, B_g) .

(c) For each $w \in B_q$, $w \in C_i$ for some *i*. By (13) we have

$$\begin{split} |w|^2 &= \sum_{j=1}^g (x_j^2 + y_j^2) = \sum_{j=1, j \neq i}^g (x_j^2 + y_j^2) + (x_i^2 + y_i^2) \\ &= \sum_{j=1, j \neq i}^g ((\frac{1}{\sqrt{g}})^2 + 0^2) + \frac{1}{g} = 1. \end{split}$$

That is to say B_g stays in the unit sphere $S^{2g-1} \subset \mathbb{R}^{2g}$.

By the conclusion of (b), the given $B_g \subset S^{2g-1}$ is an equivariant embedding.

Remark 0.5. For given pair (G, Γ) , as we explained in the proof of Proposition 0.4, each *G*-equivariant embedding $\Gamma \to \mathbb{R}^n$ provides a *G*equivariant embedding $\Gamma \to S^n$. However a *G*-equivariant embedding $\Gamma \to S^n$ does not guarantee a *G*-equivariant embedding $\Gamma \to R^n$, see the example below.

Example 0.6. The maximum order of finite group action on a hyperbolic graph of rank 2 is 12, which is realized uniquely by the action of $\text{Sym}(M_3) = S_3 \ltimes \mathbb{Z}_2$ on M_3 [WZ], where M_3 is the graph with two vertices joined by three edges, S_3 permutes three edges and fixes each vertex, and \mathbb{Z}_2 is an orientation-reversing involution on each edge. There is an equivariant embedding $M_3 \to S^2$, which is indicated by the left side of Figure 4.

There is no equivariant embedding $M_3 \subset \mathbb{R}^2$. A quick proof uses Jordan curve theorem: For any embedding $M_3 \to \mathbb{R}^2$, M_3 divides \mathbb{R}^2 into two bounded regions R_1 , R_2 , and one unbounded region R_3 , see the right side of Figure 4. Then C_2 , the unique edge neighboring two bounded regions, must be invariant under any $\tau \in O(2)$, therefore the embedding is not equivariant.

THE MINIMAL n FOR EQUIVARIANT EMBEDDINGS $B_g \to S^n$ IS n=2g-1 9



FIGURE 4. An equivariant embedding $M_3 \rightarrow S^2$

Remark 0.7. A related question is when a finite group action G on a graph Γ can be G-equivariantly embedded into \mathbb{R}^3 (S^3), see [FNPT], [WWZZ2]. Similar question for surfaces is also addressed, see [CC], [WWZZ1].

Acknowledgements. We thank the referees for their suggestions which enhance the paper.

References

- [CC] W. Cavendish, J.H. Conway, Symmetrically bordered surfaces. Amer. Math. Monthly 117 (2010), no. 7, 571-580.
- [FNPT] E. Flapan, R. Naimi, J. Pommersheim, H. Tamvakis, Topological symmetry groups of graphs embedded in the 3-sphere. Comment. Math. Helv. 80 (2005), no. 2, 317-354.
- [Mos] G. D. Mostow, Equivariant embeddings in Euclidean space. Ann. of Math. (2) 65 (1957), 432-446.
- [Pa] R. Palais, Imbedding of compact, differentiable transformation groups in orthogonal representations. J. Math. Mech. 6 (1957), 673-678.
- [WWZZ1] C. Wang, S.C. Wang, Y.M. Zhang, B. Zimmermann, Embedding compact surfaces into the 3-dimensional Euclidean space with maximum symmetry. Sci. China Math. 60 (2017), no. 9, 1599-1614.
- [WWZZ2] C. Wang, S.C. Wang, Y.M. Zhang, B. Zimmermann, Graphs in the 3sphere with maximum symmetry. Discrete Comput. Geom. 59 (2018), no. 2, 331-362.
- [WZ] S.C. Wang, B. Zimmermann, The maximum order of finite groups of outer automorphisms of free groups. Math. Z. 216 (1994), no. 1, 83-87.
- [Zi] B. Zimmermann, On large groups of symmetries of finite graphs embedded in spheres. J. Knot Theory Ramifications 27 (2018), no. 3, 1840011, 8 pp.

DEPARTMENT OF MATHEMATICS SCIENCE, TSINGHUA UNIVERSITY, BEIJING, 100080, CHINA

Email address: wangzz18@mails.tsinghua.edu.cn