# numerical analysis of a nonsmooth quasilinear elliptic control problem: i. explicit second-order optimality conditions

Christian Clason<sup>∗</sup> Vu Huu Nhu† Arnd Rösch‡

Abstract In this paper, we derive explicit second-order necessary and sufficient optimality conditions of a local minimizer to an optimal control problem for a quasilinear second-order partial differential equation with a piecewise smooth but not differentiable nonlinearity in the leading term. The key argument rests on the analysis of level sets of the state. Specifically, we show that if a function vanishes on the boundary and its the gradient is different from zero on a level set, then this set decomposes into finitely many closed simple curves. Moreover, the level sets depend continuously on the functions defining these sets. We also prove the continuity of the integrals on the level sets. In particular, Green's first identity is shown to be applicable on an open set determined by two functions with nonvanishing gradients. In the second part to this paper, the explicit sufficient second-order conditions will be used to derive error estimates for a finite-element discretization of the control problem.

bluestructureKey words Level set, optimal control, nonsmooth optimization, quasilinear elliptic equation, piecewise differentiable function

#### 1 introduction

In this paper, we investigate the nonsmooth quasilinear elliptic optimal control problem

<span id="page-0-0"></span>
$$
\begin{cases}\n\min_{u \in L^{\infty}(\Omega)} j(u) := \int_{\Omega} L(x, y_u(x)) dx + \frac{\nu}{2} ||u||_{L^2(\Omega)}^2 \\
\text{s.t.} \quad -\operatorname{div}[(b + a(y_u)) \nabla y_u] = u \quad \text{in } \Omega, \quad y_u = 0 \text{ on } \partial \Omega, \\
\alpha \le u(x) \le \beta \quad \text{a.e. } x \in \Omega,\n\end{cases}
$$

where Ω is a bounded, convex and polygonal domain  $\Omega \subset \mathbb{R}^2$ ;  $L : \Omega \times \mathbb{R} \to \mathbb{R}$  is a Carathéodory function of class  $C^2$  with respect to (w.r.t.) the second variable;  $b : \overline{\Omega} \to \mathbb{R}$  is a Lipschitz continuous function;  $a : \mathbb{R} \to \mathbb{R}$  is a nonsmooth function; and  $\alpha, \beta, \nu \in \mathbb{R}$  satisfy  $\beta > \alpha$  and  $\nu > 0$ . For the precise hypotheses on the data of [\(P\)](#page-0-0), we refer to [Section 3.1.](#page-18-0)

The control problem [\(P\)](#page-0-0) is interesting since the corresponding state equation arises, for instance, in models of heat conduction where the coefficient in the divergence term of the state equation is the heat conductivity and depends on the temperature  $\gamma$  and on the spatial coordinate x; see, e.g. [\[2,](#page-36-0) [25\]](#page-37-0). When

<sup>∗</sup>Faculty of Mathematics, Universität Duisburg-Essen, 45117 Essen, Germany; current address: Institute of Mathematics and Scientific Computing, University of Graz, Heinrichstrasse 36, 8010 Graz, Austria [\(c.clason@uni-graz.at,](mailto:c.clason@uni-graz.at) orcid: [0000-](https://orcid.org/0000-0002-9948-8426) [0002-9948-8426\)](https://orcid.org/0000-0002-9948-8426)

<sup>†</sup>Faculty of Fundamental Sciences, PHENIKAA University, Yen Nghia, Ha Dong, Hanoi 12116, Vietnam  $(nhu.vuhuu@phenikaa-uniedu.vn, \text{ORCID: } 0000-0003-4279-3937)$  $(nhu.vuhuu@phenikaa-uniedu.vn, \text{ORCID: } 0000-0003-4279-3937)$ 

<sup>‡</sup>Faculty of Mathematics, University of Duisburg-Essen, Thea-Leymann-Strasse 9, 45127 Essen, Germany [\(arnd.roesch@uni](mailto:arnd.roesch@uni-due.de)[due.de\)](mailto:arnd.roesch@uni-due.de)

the data are of class  $C^2$ , the numerical analysis of the discrete approximation of such optimal control problems was investigated by Casas et al. in  $[6, 7]$  $[6, 7]$  $[6, 7]$  for distributed control and in  $[5]$  for Neumann control. Here, we only assume that the coefficient *a* is continuous and piecewise  $C^2$ , which makes the analysis significantly more challenging.

In [\[9\]](#page-36-4), we studied so-called no-gap second-order necessary and sufficient optimality conditions for  $(P)$  in terms of a general nonsmooth curvature functional of *j*. In this paper, we will formulate an explicit formula for the curvature functional; see [Theorem 3.21](#page-31-0) below. This explicit derivation is based only on an assumption on the gradient of the state function  $y_u$  and on the level set of  $y_u$  with respect to points  $\bar{t}$  where the nonlinearity *a* is not differentiable. The sufficient condition will be used to prove a priori numerical error estimates for the discretization of the nonsmooth optimal control problem using the finite element method in the second part of this work.

Let us comment on related works in  $[8]$  and  $[24]$  for explicitly computing curvature terms of nonsmooth functionals. In [\[8\]](#page-36-5), the authors considered bang-bang optimal control problems, where the associated state equation is linear and the controls satisfy the box-constraint  $-1 \le u \le 1$  a.e. in  $\Omega$ . They observed that the set  $\{u \in L^{\infty}(\Omega) \mid -1 \le u \le 1 \text{ a.e. in } \Omega\}$  possesses a curvature in the space  $\mathcal{M}(\Omega)$ , the dual of the space consisting of all continuous functions vanishing on the boundary. Under a structural assumption imposed on the corresponding adjoint state and its level set, they established the curvature of the set  $\{u \in L^{\infty}(\Omega) \mid -1 \le u \le 1$  a.e. in  $\Omega\}$  by using a directional Taylor-like expression in the  $L^1(\Omega)$ -norm. In [\[24\]](#page-37-1), the authors first considered general optimization problems with a nonuniformly convex and nonsmooth integral functional and then applied the obtained results to bang-off-bang optimal control problems with smooth semilinear elliptic partial differential equations. In these papers, the authors first proved explicit formulas for the case  $\Omega \subset \mathbb{R}$  and then lifted the obtained result to the higher-dimensional situation by using partitions of unity.

Here we construct an explicit formula of the nonsmooth curvature functional through a thorough analysis of level sets. In particular, we study the structure and the continuity in terms of determining functions of the level sets, a Green's formula, and the continuity of integrals over level sets. To fix ideas, let  $y:\overline{\Omega}\to\mathbb{R}$  be a  $C^1$  function with an image  $\mathcal{R}(y)$ . For any  $t\in\mathcal{R}(y)$ , the preimage

$$
\{y = t\} := \{x \in \overline{\Omega} \mid y(x) = t\}
$$

is called level set of  $\gamma$  corresponding to the level value t. These level sets are compact as a result of the continuity of y and the boundedness of  $\overline{\Omega}$ . When  $\Omega = \mathbb{R}^2$  and y is of class  $C^2$ , Sard's theorem (see, e.g. [\[22\]](#page-37-2) and Appendix 1 in [\[16\]](#page-37-3)) states that the set of critical values

$$
\{t \in \mathcal{R}(y) \mid \nabla y(x) = 0 \text{ for some } x \in \{y = t\}\}\
$$

is of one-dimensional (Lebesgue) measure zero. The Implicit Function Theorem further yields that for almost all (a.a.)  $t \in \mathcal{R}(y)$ , the level sets  $\{y = t\}$  are simple curves of class  $C^2$ . When y is Lipschitz continuous, it was shown in [\[1\]](#page-36-6) that for a.a.  $t \in \mathcal{R}(y)$ , any connected component of  $\{y = t\}$  is either a point or a simple curve with Lipschitz parametrization; see statement (iv) in Theorem 2.5 in [\[1\]](#page-36-6). If  $\gamma$  is an element of the Sobolev space  $W^{2,1}(\Omega)$ , then for a.a.  $t \in \mathcal{R}(y)$  the level set  $\{y = t\}$  is a union of a finitely many disjoint  $C^1$  simple curves; see Theorem 5.1 and Corollary 5.2 in [\[3\]](#page-36-7). However, here we are interested in the structure of the level set { $y = t$ } for a given concrete value of t and cannot rely on properties that only hold almost everywhere.

The study of level sets at given level values is also relevant to the analysis of level set methods, first proposed by Osher and Sethian [\[20\]](#page-37-4). These methods rely in part on the theory of curve and surface evolution given in [\[23\]](#page-37-5) and stem from the observation that the level set function  $\gamma$  in compressible flows carries information on the stretching of the fluid-structure interface. A key property in the analysis is the continuity of level sets with respect to changes in the function or the level value. For example, [\[10\]](#page-36-8) proved the continuity at  $t = 0$  of a functional, mapping  $t \in \mathbb{R}$  to a corresponding integral

over the level set  $\{y = t\}$ , provided that the function y belongs to class of  $C^2$  and that the gradient  $\nabla y$ is different from zero on  $\{v = 0\}$ .

In this work, under the sole assumption that y is  $C^1$  and vanishes on the boundary  $\partial\Omega$ , we apply the Implicit Function Theorem to show that any connected component of the level set on which the gradient of y does not vanish is indeed a  $C^1$  closed simple curve. Furthermore, if the gradient of y is different from zero on  $\{y = t\}$ , then the level set consists of finitely many closed simple curves; see [Proposition 2.4](#page-7-0) below.

We then study the continuity of integrals over level sets in term of functions that determine the level sets. Specifically, we shall show that if  $y_n \to y$  strongly in  $C^1(\overline{\Omega})$  and if the gradient of y does not vanish on  $\{y = t\}$ , then the integrals over  $\{y_n = t\}$  tend to the one over  $\{y = t\}$ ; see [Proposition 2.13](#page-17-0) below. In order to show this, we first present in [Lemma 2.2](#page-4-0) an extension of the Implicit Function Theorem applying to a point belonging to  $\{y = t\}$ , where the implicit functions can be extended to the boundary of a neighborhood of the mentioned point. We then use partitions of unity to show the continuity of integrals on level sets. In addition, we prove in [Proposition 2.11](#page-14-0) (see also [Corollary 2.12\)](#page-16-0) that in any small  $\varepsilon$ -neighborhood of a connected component of  $\{y = t\}$ , there is one and only one connected component of  $\{y_n = t\}$  for *n* large enough.

We use these properties to establish a version of Green's first identity; see, e.g. identity (2.10) in Chapter II in [\[14\]](#page-37-6) for the classical version. There, the domains of integration,  $\{y_1 < t < y_2\} \cup \{y_1 > t\}$  $\{x > y_2\}$ , are determined by two C<sup>1</sup> functions  $y_1, y_2$  with nonvanishing gradients, and boundaries of these domains are parts of level sets of  $y_1$  and  $y_2$  at the same level value t. As a result, these domains of integration might admit some cusps and then not have Lipschitz boundaries in the sense of Nečas, see, e.g. Definition 1.2.1.1 [\[15\]](#page-37-7). In order to validate Green's first identity on these nonsmooth domains, we show in [Lemma 2.6](#page-8-0) below that the open set  $\{y_1 < t < y_2\}$  can be decomposed into at most countably many disjoint open connected subdomains. Each of such subdomains is approximated by regular domains with boundaries being curvilinear polygons of class  $C^1$ . By passing to the limit in Green's first identity applied over approximating regular domains and summing up the obtained identities, we arrive at the Green formula on  $\{y_1 < t < y_2\} \cup \{y_1 > t > y_2\}$ ; see [Lemma 2.7](#page-10-0) and [Proposition 2.9.](#page-12-0) Together with continuity properties of level sets and of integrals on them, with the aid of this, we derive an explicit formula for the curvature functional in [Theorem 3.21](#page-31-0) and use this to obtain second-order necessary and sufficient optimality conditions in explicit forms obtained in [Theorems 3.22](#page-32-0) and [3.23,](#page-32-1) respectively.

The plan of the paper is as follows. We conclude this section with some general notation. In the next section, we investigate the structure and the continuity properties of the level sets as well as derive Green's first identity on open sets determined by two functions and a level value. [Section 3](#page-18-1) is then devoted to the study of the nonsmooth quasilinear optimal control problem [\(P\)](#page-0-0): In [Section 3.1,](#page-18-0) we rigorously state the assumptions for [\(P\)](#page-0-0) and provide some preliminary results from [\[9\]](#page-36-4). An explicit formula of the curvature functional of the objective functional is derived in [Section 3.2.](#page-22-0) Finally, in [Section 3.3,](#page-32-2) explicit second-order necessary and sufficient optimality conditions are presented.

Notation. We denote by  $B_X(u, \rho)$  and  $\overline{B}_X(u, \rho)$  the open and closed balls in a Banach space X of radius  $\rho > 0$  centered at  $u \in X$ , respectively. For Banach spaces X and Y, the notation  $X \hookrightarrow (\infty)Y$  is understood that X is continuously (compactly) embedded in Y. For a given function  $q : \Omega \to \mathbb{R}$  and a subset  $A \subset \mathbb{R}$ ,  $\{g \in A\}$  denotes the set of all points  $x \in \overline{\Omega}$  for which  $g(x) \in A$ . For functions  $g_1, g_2$ and subsets  $A_1, A_2 \subset \mathbb{R}$ , we set  $\{g_1 \in A_1, g_2 \in \overline{A}_2\} := \{g_1 \in A_1\} \cap \{g_2 \in A_2\}.$  For any set  $\omega \subset \overline{\Omega}$ , we denote by  $\mathbb{1}_{\omega}$  the characteristic function of  $\omega$ , i.e.,  $\mathbb{1}_{\omega}(x) = 1$  if  $x \in \omega$  and  $\mathbb{1}_{\omega}(x) = 0$  otherwise. The symbol  $\mathcal{H}^1$  stands for the one-dimensional Hausdorff measure on  $\mathbb{R}^2$  that is scaled as in [\[12\]](#page-36-9), Def. 2.1. For a measurable two-dimensional subset A, by meas<sub>ℝ2</sub>(A), we denote the two-dimensional Lebesgue measure of  $A$ . Finally, we write the symbol  $C$  for a generic positive constant, which may be different at

different places of occurrence and the notation, e.g.  $C_{\xi}$  for a constant depending only on the parameter .

# <span id="page-3-4"></span>2 analysis of level sets

In this section,  $\Omega$  is assumed to be a convex and bounded domain in  $\mathbb{R}^2$  only.

### 2.1 structure of level sets

We first recall from [\[17\]](#page-37-8) that a connected component of a set  $V \subset \mathbb{R}^d$ ,  $d \geq 1$ , is any element of the class of connected subsets of V that is maximal with respect to inclusion and that a simple closed  $C^1$ curve in  $\R^d$  is a curve that admits a  $C^1$  parametrization  $\gamma:[h,k]\to\R^d$  which is injective on  $[h,k)$  and satisfies  $v(h) = v(k)$ .

We shall investigate the structure of any connected component of the level sets of  $C^1$ -functions that vanish on the boundary and have nonvanishing gradients on these components. We begin with the following  $C^1$  extension of a smooth function defined over convex domains.

<span id="page-3-3"></span>Proposition 2.1. Let G be a convex domain in  $\mathbb{R}^d$  with  $d \geq 1$  and let  $f \in C^1(\overline{G})$  be arbitrary. Then there exists a function  $\bar{f} : \mathbb{R}^d \to \mathbb{R}$  with the following properties:

- <span id="page-3-0"></span>(a)  $\bar{f}$  is of class  $C^1$ ;
- <span id="page-3-1"></span>(b)  $\bar{f} = f$  and  $\nabla \bar{f} = \nabla f$  on  $\bar{G}$ .

Proof. We will show that Whitney's Extension Theorem (see, e.g. Chapter 6 in [\[12\]](#page-36-9) and Theorem 9.43 in [\[17\]](#page-37-8)) can be applied and thus derive the desired extension function. To this end, for any  $x_1, x_2 \in \overline{G}$ with  $x_1 \neq x_2$ , set

$$
R(x_2, x_1) := \frac{f(x_2) - f(x_1) - \nabla f(x_1) \cdot (x_2 - x_1)}{|x_2 - x_1|}.
$$

For any compact subset  $M \subset \overline{G}$ , we define the function  $\rho_M : (0,1) \to (0,\infty)$  by

$$
\rho_M(\tau) := \sup\{|R(x_2, x_1)| \mid 0 < |x_1 - x_2| < \tau, x_1, x_2 \in M\}, \quad \tau \in (0, 1).
$$

In light of Whitney's Extension Theorem, in order to derive assertions [\(a\)](#page-3-0) and [\(b\),](#page-3-1) we need to show that

$$
\rho_M(\tau) \to 0 \quad \text{as} \quad \tau \to 0^+.
$$

Indeed, for any compact subset  $M\subset \overline{G},$  and any  $x_1,x_2\in M,$   $x_1\neq x_2,$  we deduce from the Mean Value Theorem and the convexity of  $G$  that

<span id="page-3-2"></span>
$$
|R(x_2, x_1)| = \frac{|\nabla f(x_1 + \theta(x_2 - x_1)) \cdot (x_2 - x_1) - \nabla f(x_1) \cdot (x_2 - x_1)|}{|x_1 - x_2|}
$$
  
\$\leq | \nabla f(x\_1 + \theta(x\_2 - x\_1)) - \nabla f(x\_1)|\$

with some  $\theta \in (0,1)$ . There then holds

$$
\rho_M(\tau) \leq \sup\{|\nabla f(x_2) - \nabla f(x_1)| \mid |x_1 - x_2| < \tau, x_1, x_2 \in M\}
$$

for all  $\tau \in (0,1)$ . From this and the uniform continuity of  $\nabla f$  on the compact set M, we have [\(2.1\)](#page-3-2). □

It is well-known from differential geometry (see, e.g. [\[18\]](#page-37-9) and [\[16\]](#page-37-3)) that every compact one-dimensional manifold without boundary is a union of finitely many disjoint closed curves. As a direct consequence, we obtain in [Proposition 2.4](#page-7-0) below the decomposition into finitely many disjoint closed curves of level sets of a  $C<sup>1</sup>$ -function that vanishes on the boundary of a two-dimensional convex domain and has a nonvanishing gradient. For the sake of convenience, we provide its proof based on [Proposition 2.1](#page-3-3) and the following result on the extension of an implicit function. Besides, this extension result plays an important role in proving the continuity of integrals on level sets in [Section 2.4](#page-17-1) below.

<span id="page-4-0"></span>Lemma 2.2. Let  $t \in \mathbb{R}$  be arbitrary and let  $y \in C^1(\overline{\Omega})$  be such that  $y_{|\partial\Omega} = 0$  and that there is a point  $x_0 := (x_0^1, x_0^2) \in \{y = t\}$ . Let C be a connected component of  $\{y = t\}$  such that  $x_0 \in C$ . Assume that there exists a constant  $r > 0$  satisfying

<span id="page-4-6"></span>(2.2) { = } ∩ ℝ<sup>2</sup> (0, ) = C ∩ ℝ<sup>2</sup> (0, )

and

(2.3) 
$$
\{y=t\}\cap \partial \overline{B}_{\mathbb{R}^2}(x_0,r)=\{p,q\},\quad p\neq q.
$$

Assume further that

(2.4) 
$$
\left|\frac{\partial y}{\partial x^2}(x)\right| \ge \alpha > 0
$$
 for all  $x \in \overline{B}_{\mathbb{R}^2}(x_0, r) \cap \overline{\Omega}$ 

with some constant  $\alpha > 0$ . Then the following assertions hold.

<span id="page-4-4"></span>(a) There exist constants  $h_0 > 0$ ,  $k_0 > 0$  and a C<sup>1</sup>-function  $g_0 : [x_0^1 - h_0, x_0^1 + k_0] \to \mathbb{R}$  satisfying

<span id="page-4-3"></span><span id="page-4-2"></span>
$$
(2.5) \qquad \begin{cases} g_0(x_0^1) = x_0^2, \\ \overline{B}_{\mathbb{R}^2}(x_0, r) \cap \{y = t\} = \{(\tau, g_0(\tau)) \mid \tau \in [x_0^1 - h_0, x_0^1 + k_0] \}, \\ \{(x_0^1 - h_0, g_0(x_0^1 - h_0)), (x_0^1 + k_0, g_0(x_0^1 + k_0))\} = \partial \overline{B}_{\mathbb{R}^2}(x_0, r) \cap \{y = t\}. \end{cases}
$$

<span id="page-4-5"></span>(b) If, in addition, there exists a sequence  $\{y_n\} \in C^1(\overline{\Omega})$  satisfying that  $y_n|_{\partial \Omega} = 0$  and that

<span id="page-4-1"></span>
$$
(2.6) \quad\n\begin{cases}\n\mathcal{Y}_n \to \mathcal{Y} \quad \text{strongly in } C^1(\overline{\Omega}), \\
\{\mathcal{Y}_n = t\} \cap \overline{B}_{\mathbb{R}^2}(x_0, r_0) \text{ is an arc for all } n \text{ large enough}, \\
\frac{\partial \mathcal{Y}}{\partial x^2}(x) \frac{\partial \mathcal{Y}_n}{\partial x^2}(x) \ge \tilde{\alpha} > 0 \quad \text{for all } x \in \overline{B}_{\mathbb{R}^2}(x_0, r) \cap \overline{\Omega} \text{ and for all } n \text{ sufficiently large},\n\end{cases}
$$

then there is an integer  $n_0 = n_0(x_0, r)$  such that, for any  $n \ge n_0$ , constants  $h_n, k_n > 0$  and a  $C^1$ function  $g_n : [x_0^1 - h_n, x_0^1 + k_n] \to \mathbb{R}$  exist and satisfy the following properties.

<span id="page-4-7"></span>(i) 
$$
\overline{B}_{\mathbb{R}^2}(x_0, r) \cap \{y_n = t\} = \{(\tau, g_n(\tau)) \mid \tau \in [x_0^1 - h_n, x_0^1 + k_n]\};
$$

<span id="page-4-8"></span>
$$
(ii) \ \{ (x_0^1 - h_n, g_n(x_0^1 - h_n)), (x_0^1 + k_n, g_n(x_0^1 + k_n)) \} = \partial \overline{B}_{\mathbb{R}^2}(x_0, r) \cap \{ y_n = t \};
$$

<span id="page-4-9"></span>
$$
(iii) \ (x_0^1 - h_n, g_n(x_0^1 - h_n)) \to (x_0^1 - h_0, g_0(x_0^1 - h_0)) \ and \ (x_0^1 + k_n, g_n(x_0^1 + k_n)) \to (x_0^1 + k_0, g_0(x_0^1 + k_0));
$$

<span id="page-4-10"></span>(iv) For any  $\varepsilon > 0$  and for all  $\tau \in [x_0^1 - h_0 + \varepsilon, x_0^1 + k_0 - \varepsilon]$ , there hold

<span id="page-4-11"></span>(2.7) 
$$
|g_n(\tau) - g_0(\tau)| \leq C_1 ||y_n - y||_{C(\overline{\Omega})} \quad \text{for } n \text{ large enough}
$$

and

<span id="page-4-12"></span>(2.8) 
$$
\begin{cases} g'_n(\tau) \to g'_0(\tau), \\ g'_n(x_0^1 - h_n) \to g'_0(x_0^1 - h_0), \\ g'_n(x_0^1 + k_n) \to g'_0(x_0^1 + k_0). \end{cases}
$$

<span id="page-4-13"></span>Moreover, for all  $\tau \in [x_0^1 - h_n, x_0^1 + k_n]$ , one has

$$
(2.9) \t\t |g'_n(\tau)| \leq C_2.
$$

Here  $\tilde{\alpha}$  and  $C_1, C_2$  are constants independent of n,  $\varepsilon$ , and  $\tau$ .

Remark 2.3. The second condition in [\(2.6\)](#page-4-1) is, in fact, fulfilled if the first one in (2.6) and the second one in [\(2.5\)](#page-4-2) hold. This fact shall be proven in [Proposition 2.11](#page-14-0) below; see also, the proof of that proposition.

*Proof.* Without loss of generality, assume that  $x_0 = (0, 0)$ , i.e.,  $x_0^1 := 0$  and  $x_0^2 := 0$ . We first see from the condition [\(2.4\)](#page-4-3) that the following implication holds

<span id="page-5-0"></span>(2.10) 
$$
\begin{cases} x = (x^1, x^2), \tilde{x} = (\tilde{x}^1, \tilde{x}^2) \in C \cap \overline{B}_{\mathbb{R}^2}(0, r), \\ x \neq \tilde{x} \end{cases} \implies x^1 \neq \tilde{x}^1.
$$

Consequently, one has  $p^1 \neq q^1$ , where  $p^1$  and  $q^1$  stand, respectively, for the first component of the end points  $p$  and  $q$ . We can assume without loss of generality that

<span id="page-5-3"></span><span id="page-5-1"></span>
$$
p^1 < q^1.
$$

Let  $C_{x^1}$  be the projection of the set  $C\cap \overline{B}_{\R^2}(x_0,r)$  onto the  $x^1$ -axis. Since  $C\cap \overline{B}_{\R^2}(x_0,r)$  is connected and compact,  $C_{x^1}$  is also connected and compact; see, e.g. Theorem 4.22 in [\[21\]](#page-37-10). Therefore,  $C_{x^1}$  must be a closed bounded interval of ℝ. Combining this with the implication [\(2.10\)](#page-5-0) yields

$$
(2.11) \tC_{x^1} = [p^1, q^1].
$$

Moreover, since  $0 \in C \cap \overline{B}_{\mathbb{R}^2}(x_0, r)$ , and  $p$  and  $q$  are the end points of the arc  $C \cap \overline{B}_{\mathbb{R}^2}(x_0, r)$ , there holds

$$
(2.12) \t\t p1 < 0 < q1,
$$

due to the implication [\(2.10\)](#page-5-0). We now prove assertions [\(a\)](#page-4-4) and [\(b\).](#page-4-5)

Ad [\(a\):](#page-4-4) In light of [Proposition 2.1,](#page-3-3) y admits a  $C^1$  extension, denoted also by y, on  $\mathbb{R}^2$ . By the continuity of  $\nabla y$ , we can assume that the condition [\(2.4\)](#page-4-3) is valid on an open neighborhood O of  $\overline{B}_{R^2}(0,r) \cap \{y = t\}$ (possibly with a different positive constant, also denoted by  $\alpha$ ), which contains the arc  $C \cap \overline{B_{\mathbb{R}^2}(0,r)}$ . According to the condition [\(2.2\)](#page-4-6), we can assume that the open set  $O$  satisfies

(2.13) O ∩ { = } = O ∩ C.

Therefore, one can apply the Implicit Function Theorem in every point  $x=(x^1,x^2)$  of  $C\cap\overline{B}_{\mathbb{R}^2}(0,r)$ . We now take  $x = (x^1, x^2) \in C \cap \overline{B}_{\mathbb{R}^2}(0, r)$  arbitrarily but fixed. Thanks to the Implicit Function Theorem, there exist constants  $h_x, k_x, \rho_x > 0$  and a  $C^1$ -function  $g_x : (x^1 - h_x, x^1 + k_x) \to \mathbb{R}$  satisfying

<span id="page-5-2"></span>(2.14) 
$$
\begin{cases} B_{\mathbb{R}^2}(x, \rho_x) \subset O, & g_x(x^1) = x^2, \\ y(\tau, g_x(\tau)) = t \quad \text{for all } \tau \in (x^1 - h_x, x^1 + k_x), \\ ((\tau, \xi) \in B_{\mathbb{R}^2}(x, \rho_x) \text{ and } y(\tau, \xi) = t) \quad \Longrightarrow \quad \xi = g_x(\tau). \end{cases}
$$

Since  $C\cap \overline{B}_{\mathbb{R}^2}(0,r)\subset\bigcup\{B_{\mathbb{R}^2}(x,\rho_x)\mid x\in C\cap \overline{B}_{\mathbb{R}^2}(0,r)\}$  and  $C\cap \overline{B}_{\mathbb{R}^2}(0,r)$  is compact, there exist finitely many points  $x_1, x_2, ..., x_s \in C \cap \overline{B}_{\mathbb{R}^2}(0, r)$  with  $s \in \mathbb{N}$  and  $C^1$ -functions  $g_i : (x_i^1 - h_i, x_i^1 + k_i) \to \mathbb{R}$  with  $g_i := g_{x_i}, h_i := h_{x_i}, k_i := k_{x_i}, 1 \le i \le s$ , such that

$$
C\cap \overline{B}_{\mathbb{R}^2}(0,r)\subset \cup_{i=1}^s B_{\mathbb{R}^2}(x_i,\rho_i)
$$

with  $\rho_i := \rho_{x_i}$ . From this and [\(2.11\)](#page-5-1), we have

$$
[p^1, q^1] \subset \bigcup_{i=1}^s (x_i^1 - h_i, x_i^1 + k_i) =: V.
$$

We now construct the function  $q_0$  defined over the open set V by setting

(2.15) 
$$
g_0(\tau) = g_i(\tau) \text{ if } \tau \in (x_i^1 - h_i, x_i^1 + k_i) \text{ for some } i.
$$

Thanks to the implication in [\(2.14\)](#page-5-2), the function  $g_0$  is actually well-defined and is of class  $C^1$  on the open neighborhood V of  $\lceil p^1, q^1 \rceil$ . Moreover, for later use, we obtain by simple computation that

<span id="page-6-3"></span>(2.16) 
$$
g'_0(\tau) = -\frac{\frac{\partial y}{\partial x^1}(\tau, g_0(\tau))}{\frac{\partial y}{\partial x^2}(\tau, g_0(\tau))}, \quad \tau \in V.
$$

In view of [\(2.4\)](#page-4-3) and from the continuity of  $\nabla y$  over  $\overline{\Omega}$ , there holds

<span id="page-6-4"></span>(2.17) 
$$
|g_0'(\tau)| \le \frac{\|\nabla y\|_{C(\overline{\Omega})}}{\alpha} \quad \text{for all } \tau \in V.
$$

Now setting  $h_0 := -p^1$  and  $k_0 := q^1$  and exploiting [\(2.12\)](#page-5-3), one has  $h_0, p_0 > 0$ . On the other hand, we have [\(2.5\)](#page-4-2) from [\(2.14\)](#page-5-2) and the definition of function  $q_0$ . We thus obtain [\(a\).](#page-4-4)

 $Ad (b)$  $Ad (b)$ : The proof of assertion (b) is divided into several steps as follows.

• Step 1: There is an integer  $n_0^1$  such that, for any  $n \ge n_0^1$ , a number  $\xi_n \in (-r_0, r_0)$  uniquely exists and fulfills

$$
(2.18) \t\t yn(0, \xin) = t.
$$

In fact, as a consequence of the last condition in [\(2.6\)](#page-4-1), the functions  $y(0, \cdot)$  and  $y_n(0, \cdot)$  are both either strictly increasing or strictly decreasing. We now only consider the situation where both functions are strictly increasing, since the other is analyzed analogously. It suffices to prove that  $(2.18)$ , for each n large enough, admits at least one solution in  $(-r_0, r_0)$ . To this end, by contradiction assume that there exists a subsequence, denoted in the same way, of  ${n}$  such that [\(2.18\)](#page-6-0) has no solutions in (-r<sub>0</sub>, r<sub>0</sub>) for all  $n \ge 1$ . We then split  ${n}$  into subsequences, also denoted by  ${n}$ , satisfying one of two inequalities

<span id="page-6-1"></span>(I1)  $y_n(0, \xi) > t$  for all  $n \ge 1$  and  $\xi \in (-r_0, r_0);$  (I2)  $y_n(0, \xi) < t$  for all  $n \ge 1$  and  $\xi \in (-r_0, r_0).$ 

In order to show a contradiction, we now only consider  $(I_1)$  since the case  $(I_2)$  is similarly analyzed. We have for all  $n \geq 1$  that

<span id="page-6-2"></span><span id="page-6-0"></span>
$$
y_n(0, -\frac{r_0}{2}) > t = y(0, 0) > y(0, -\frac{r_0}{2}),
$$

which yields that

$$
||y_n - y||_{C(\overline{\Omega})} \ge |y_n(0, -\frac{r_0}{2}) - y(0, -\frac{r_0}{2})| > t - y(0, -\frac{r_0}{2})
$$
 for all  $n \ge 1$ .

Passing to the limit and exploiting the first condition in [\(2.6\)](#page-4-1) give  $0 \ge t - y(0, -\frac{r_0}{2}) > 0$ , a contradiction. • Step 2: For any  $n \ge n_0^1$ , there exist constants  $h_n, k_n > 0$  and a function  $g_n \in C^1[x_0^1 - h_n, x_0^1 + k_n]$ , all of which fulfill assertions [\(i\)](#page-4-7) and [\(ii\),](#page-4-8) where  $n_0^1$  is defined in Step 1. To this end, we first see that  $x_{0,n} := (0, \xi_n) = (x_0^1, \xi_n) \in B_{\mathbb{R}^2}(x_0, r_0)$  with  $\xi_n \in (-r_0, r_0)$  being a unique solution to [\(2.18\)](#page-6-0). Using the argument in the proof of assertion [\(a\)](#page-4-4) for the point  $x_{0,n}$  and the function  $y_n$  in the places of  $x_0$  and  $y$ , respectively, Step 2 is proven. Moreover, similar to [\(2.16\)](#page-6-3) and [\(2.17\)](#page-6-4), there hold for function  $g_n$ 

<span id="page-6-5"></span>(2.19) 
$$
g'_{n}(\tau) = -\frac{\frac{\partial y_{n}}{\partial x^{1}}(\tau, g_{n}(\tau))}{\frac{\partial y_{n}}{\partial x^{2}}(\tau, g_{n}(\tau))} \text{ and } |g'_{n}(\tau)| \leq C ||\nabla y_{n}||_{C(\overline{\Omega})}
$$

Clason, Nhu, Rösch **Numerical analysis of a nonsmooth quasilinear elliptic ...** 

• Step 3: Verification of assertion [\(iii\).](#page-4-9) Indeed, we set the points belonging the boundary  $\partial B_{\mathbb{R}^2}(x_0, r_0)$  of  $\{y = t\}$  and  $\{y_n = t\}$  as follows

$$
p_0 := (-h_0, g_0(-h_0)),
$$
  $q_0 := (k_0, g_0(k_0)),$   $p_n := (-h_n, g_n(-h_n)),$  and  $q_n := (k_n, g_n(k_n)).$ 

Recall that all of constants  $h_0$ ,  $h_n$ ,  $k_0$ ,  $k_n$  are positive. Thanks to the boundedness of  $\{p_n\}$  and the closedness of  $\partial B_{\mathbb{R}^2}(x_0, r_0)$ , we have from the first condition in [\(2.6\)](#page-4-1) that  $p_n \to p_0$ , i.e.,

$$
h_n \to h_0
$$
 and  $g_n(-h_n) \to g_0(-h_0)$ .

Analogously, we have

$$
k_n \to k_0
$$
 and  $g_n(k_n) \to g_0(k_0)$ .

We then obtain [\(iii\).](#page-4-9)

• Step 4: Verification of assertion [\(iv\)](#page-4-10). Indeed, for any  $\varepsilon > 0$ , there exists an integer  $n_0^2$  such that

$$
[-h_0 + \varepsilon, k_0 - \varepsilon] \subset [-h_n, k_n] \cap [-h_0, k_0] \quad \text{for all } n \ge n_0^2.
$$

For any  $\tau \in [-h_0 + \varepsilon, k_0 - \varepsilon]$ , by employing the Mean Value Theorem, we arrive at

$$
t = y_n(\tau, g_n(\tau)) = y(\tau, g_n(\tau)) + w_n(\tau, g_n(\tau))
$$
  
=  $y(\tau, g_0(\tau)) + \frac{\partial y}{\partial x^2}(\tau, g_0(\tau) + \theta_n(g_n(\tau) - g_0(\tau)))(g_n(\tau) - g_0(\tau)) + w_n(\tau, g_n(\tau))$   
=  $t + \frac{\partial y}{\partial x^2}(\tau, g_0(\tau) + \theta_n(g_n(\tau) - g_0(\tau)))(g_n(\tau) - g_0(\tau)) + w_n(\tau, g_n(\tau))$ 

for some  $\theta_n \in (0,1)$  and  $w_n := y_n - y$ . Obviously, one has  $(\tau, g_0(\tau) + \theta_n(g_n(\tau) - g_0(\tau))) \in B_{\mathbb{R}^2}(x_0, r_0) \cap \overline{\Omega}$ since  $(\tau, g_0(\tau))$  and  $(\tau, g_n(\tau))$  belong to  $B_{\mathbb{R}^2}(x_0, r_0) \cap \overline{\Omega}$ . Combing this with the last condition in [\(2.6\)](#page-4-1) yields the estimate in [\(2.7\)](#page-4-11). From this and the derivatives of  $g_0$  and  $g_n$  in [\(2.16\)](#page-6-3) and [\(2.19\)](#page-6-5), all the limits in  $(2.8)$  then follows from the first condition in  $(2.6)$  and from [\(iii\).](#page-4-9) Finally, the limit in  $(2.6)$  and the estimate in  $(2.19)$  imply  $(2.9)$ .

<span id="page-7-0"></span>Proposition 2.4. Let  $t \in \mathbb{R}$  be arbitrary and let  $y \in C^1(\overline{\Omega})$  be such that  $y_{|\partial\Omega} = 0$  and that  $\{y = t\} \neq \emptyset$ . Assume that C is a connected component of  $\{y = t\}$ . Then the following properties hold:

<span id="page-7-1"></span> $(a)$  If

<span id="page-7-4"></span>
$$
|\nabla y(x)| > 0 \quad \text{for all } x \in C,
$$

then C is a closed  $C^1$  simple curve;

<span id="page-7-3"></span> $(b)$  If

<span id="page-7-5"></span>(2.21) 
$$
|\nabla y(x)| > 0 \quad \text{for all } x \in \{y = t\},
$$

then the level set  $\{y = t\}$  decomposes into finitely many disjoint closed  $C^1$  simple curves.

*Proof.* Ad [\(a\):](#page-7-1) We first show that for any  $x \in C$ , there exist positive constants  $r_x > 0$  and a  $C^1$ -function  $\gamma_x : [-1,1] \rightarrow \mathbb{R}^2$  satisfying

<span id="page-7-2"></span>
$$
(2.22) \quad \overline{B}_{\mathbb{R}^2}(x,r_x) \cap \{y = t\} = \{ \gamma_x(\eta) \mid \eta \in [-1,1] \} \quad \text{and} \quad \gamma_x(\eta_x) = x, \gamma_x(\pm 1) \in \partial B_{\mathbb{R}^2}(x,r_x) \cap \{y = t\}
$$

for some  $\eta_x \in (-1, 1)$ . For that purpose, we first take  $x := (x^1, x^2) \in C$  arbitrarily. Without loss of generality, we can assume that  $\frac{\partial y}{\partial x^2}(x) \neq 0$  and thus  $\left| \frac{\partial y}{\partial x^2}(x) \right|$  $\frac{\partial y}{\partial x^2}(\zeta)$  > 0 for all  $\zeta \in \overline{B}_{\mathbb{R}^2}(x,r_x) \cap \overline{\Omega}$  for some constant  $r_x > 0$ . We now can take  $r_x$  small enough (if necessary) such that the arc  $C \cap \overline{B}_{\mathbb{R}^2}(x, r_x)$  has two different two end points.

[Lemma 2.2](#page-4-0) implies that there exist constants  $h_x, s_x > 0$  and a  $C^1$ -function  $g_x : [x^1 - h_x, x^1 + s_x] \to \mathbb{R}$ satisfying

<span id="page-8-1"></span>(2.23) 
$$
\begin{cases} g_x(x^1) = x^2, \\ \overline{B}_{\mathbb{R}^2}(x, r_x) \cap \{y = t\} = \{(\tau, g_x(\tau)) \mid \tau \in [x^1 - h_x, x^1 + s_x]\}, \\ \{(x^1 - h_x, g_x(x^1 - h_x)), (x^1 + s_x, g_x(x^1 + s_x))\} = \partial \overline{B}_{\mathbb{R}^2}(x, r_x) \cap \{y = t\}. \end{cases}
$$

Setting  $\eta := \frac{2}{s+1}$  $\frac{2}{s_x+h_x}\tau+1-\frac{2(x^1+s_x)}{s_x+h_x}$  $\frac{f(x'+s_x)}{s_x+h_x}$  and  $\gamma_x(\eta) := (\tau, g_x(\tau))$  and exploiting [\(2.23\)](#page-8-1), we have [\(2.22\)](#page-7-2).

Combining [\(2.22\)](#page-7-2) with the compactness of C, there exist points  $x_1, \ldots, x_{m_C} \in C$  and positive constants  $r_i > 0$ , and  $C^1$ -functions  $\gamma_i : [-1, 1] \to \mathbb{R}^2$ ,  $1 \le i \le m_C$ , such that

<span id="page-8-2"></span>(2.24) 
$$
\begin{cases} C \subset \cup_{i=1}^{m_C} B_{\mathbb{R}^2}(x_i, r_i) \cap \{y = t\}, \\ \overline{B}_{\mathbb{R}^2}(x_i, r_i) \cap \{y = t\} = \{\gamma_i(\eta) \mid \eta \in [-1, 1]\}, \\ \gamma_i(\pm 1) \in \partial B_{\mathbb{R}^2}(x_i, r_i) \cap \{y = t\}. \end{cases}
$$

From this and the fact that  $C$  is a connected component of  $\{y = t\}$ , we deduce that  $C$  is a closed  $C^1$ simple curve.

Ad [\(b\):](#page-7-3) Since  $\{y = t\}$  is compact, we also have [\(2.24\)](#page-8-2) with C being replaced by  $\{y = t\}$ . Then the level set  $\{v = t\}$  is a union of finitely many closed simple curves.  $\Box$ 

# 2.2 green's first identity

We first state the well-known result on the decomposition of open sets in a finite dimensional Euclidean space into at most countably many disjoint open connected components. Since we could not find its proof in the literature, we provide here the arguments for the sake of completeness.

<span id="page-8-3"></span>Proposition 2.5. Any nonempty open set  $V \subset \mathbb{R}^d$ ,  $d \in \mathbb{N}$ , can be expressed as an union of disjoint, open, connected components of  $V$ , at most countable in number, that is, there exist an index set  $I$ , which is at most countable, and disjoint, open and connected components  $V_i$ ,  $i \in I$ , of V such that

$$
V = \bigcup_{i \in I} V_i.
$$

*Proof.* For any  $x \in V$ , we define the set

 $O_x = \cup \{Z \mid Z \subset V, x \in Z, Z \text{ is open and connected in } \mathbb{R}^d\}.$ 

Obviously,  $O_x$  is nonempty and open since V is open. Moreover,  $O_x$  is connected; see, e.g. Chapter VI in [\[4\]](#page-36-10). On the other hand,  $O_x$  is maximal with respect to containment and thus is an open component of *V*. Therefore, the family  $\{O_x \mid x \in V\}$  is a collection of open connected components of *V*. From this and the fact that two arbitrary connected components of  $V$  are either identical or disjoint, we can now apply Lindelöf's Theorem to deduce the desired decomposition of  $V$ .

The following lemma represents the decomposition of an open set given by level sets of two functions into at most countably many disjoint open connected subdomains.

<span id="page-8-0"></span>Lemma 2.6. Let  $y_1, y_2 \in C^1(\overline{\Omega}),$  let  $t \in \mathbb{R}$  be given, and let  $O$  be an open subset in  $\overline{\Omega}.$  Define the open sets  $S^+$  and  $S^-$  by

 $S^- := \{y_1 < t < y_2\} \cap O$  and  $S^+ := \{y_1 > t > y_2\} \cap O$ .

<span id="page-9-0"></span>

Figure 1: A decomposition into components of the set  $S^+ \cup S^-$  where  $I^+ = \{1, 2\}$ ,  $I^- = I^0 = \{1\}$  when  $C_1 \cap C_2 \neq \emptyset$ . Here  $C_1$  and  $C_2$  are, respectively, plotted by dashed and solid curves

Assume that the level set  $\{y_i = t\}$  has one and only one closed  $C^1$  simple curve  $C_i$  in  $O$  for  $j = 1, 2$ . Assume further that

<span id="page-9-2"></span>
$$
(2.25) \t\t \t\t \partial S^{\pm} \subset C_1 \cup C_2
$$

and that

<span id="page-9-3"></span>(2.26) 
$$
\nabla y_1(x) \cdot \nabla y_2(x) \ge \alpha > 0 \quad \text{for all } x \in S^+ \cup S^-
$$

for some constant  $\alpha$ . Let  $\widehat{My}_j\widehat{N}$  stands for a curve lying in  $\{y_j = t\}$ ,  $j = 1, 2$  with end points M and N; see [Figure 1.](#page-9-0) Then, the following assertions hold:

- <span id="page-9-1"></span>(a) if  $C_1 = C_2$ , then  $S^+ = S^- = \emptyset$ ;
- <span id="page-9-4"></span>(b) if  $C_1 \cap C_2 = \emptyset$ , then one of the sets  $S^{\pm}$  is empty and the boundary of the other consists of  $C_1$  and  $C_2$ ,
- <span id="page-9-5"></span>(c) if  $C_1 \neq C_2$  and  $C_1 \cap C_2 \neq \emptyset$ , then the intersection  $C_1 \cap C_2$  has at most countably many closed  $\sum_{n=1}^{\infty}$  connected components. In other words, there exist index sets  $I^+, I^-, I^0$ , which are at most countable (and might be empty), and points  $M_i^+, N_i^+, M_m^-, N_m^-,$  and  $M_k^0, N_k^0$  with  $i \in I^+, m \in I^-$  ,  $k \in I^0$ , such that

<span id="page-9-6"></span>
$$
(2.27) \tC1 \cap C2 = \bigcup_{i \in I^+, m \in I^-, k \in I^0} \{M_i^+, N_i^+, M_m^-, N_m^-, \widehat{M_k^0 N_k^0}\},\
$$

where  $\widehat{M_k^0 N_k^0}$  $\widehat{F}_k^0, k\in I^0,$  denote the curves lying in  $C_1\cap C_2$  with end points  $M^0_k$  and  $N^0_k$  $U_k^0$ . Furthermore, there hold:

- (i) If  $S^+ = \emptyset$ , then the index set I<sup>+</sup> is empty,
- (ii) If the set  $S^+$  is nonempty, then it has at most countably many open connected components  $S_i^+$ ,  $i \in I^+$ , such that  $\partial S_i^+$  consists of two curves  $\widehat{M}_i^+ y_1 N_i^+$  and  $\widehat{M}_i^+ y_2 N_i^+$ .

Analogous assertions hold for the set  $S^-$ .

*Proof.* Ad [\(a\):](#page-9-1) By contradiction, assume that  $S^- \cup S^+ \neq \emptyset$ . Assume now that  $S^- \neq \emptyset$ . There then exists a point  $x_0 \in \overline{\mathcal{S}^-}$  such that  $y_1(x_0) = \min\{y_1(x) \mid x \in \overline{\mathcal{S}^-}\}.$  Thanks to [\(2.25\)](#page-9-2) and the fact that  $C_1 = C_2$ , one has  $x_0 \notin \partial S^-$  and thus  $x_0 \in S^-$ . Since  $S^-$  is open, we have  $\nabla y_1(x_0) = 0$ , contradicting [\(2.26\)](#page-9-3). Hence  $S^- = \emptyset$ . Similarly, one has  $S^+ = \emptyset$ .

*Ad (b)*: Without loss of generality, we assume now that  $\Omega^1 := \{y_1 < t\} \cap O$  is surrounded by  $C_1$ . We then have

$$
y_1(x) > t
$$
 for all  $x \in O \setminus (\Omega^1 \cup C_1)$ .

Since  $C_1 \cap C_2 = \emptyset$ , there are two possibilities:

- (I) one of  $C_1$  and  $C_2$  is not surrounded by the other; or
- (II) one of  $C_1$  and  $C_2$  is surrounded by the other.

• For case (I): If  $\{y_2 < t\} \cap O$  is surrounded by  $C_2$ , then  $S^- = \Omega^1$ . An argument as in the proof of [\(a\)](#page-9-1) thus gives a contradiction. Otherwise, if  $\{y_2 > t\} \cap O$  is surrounded by  $C_2$ , then  $\mathcal{S}^+ = O\backslash(\overline{\Omega^1} \cup \overline{\{y_2 > t\} \cap O})$ . Therefore, one has  $C_1 \cup C_2 \subsetneq \partial S^+$ , which contradicts [\(2.25\)](#page-9-2). Thus case (I) is impossible.

• For case (II): We only consider the case where  $C_1$  is surrounded by  $C_2$ . The argument for the other case is similar. If  $\{y_2 > t\} \cap O$  is surrounded by  $C_2$ , then  $S^- = \Omega^1$ , and we thus have a contradiction by using arguments analogous to ones in the proof of [\(a\).](#page-9-1) Otherwise, if  $\{y_2 < t\} \cap O$  is surrounded by  $C_2$ , then  $S^- = \emptyset$ , and the boundary of  $S^+$  consists of  $C_1$  and  $C_2$ .

Ad [\(c\):](#page-9-5) We first consider the case where  $S^+ \neq \emptyset$ . Thanks to [Proposition 2.5,](#page-8-3) the set  $S^+$  decomposes into at most countably many disjoint open components. There then exist an at most countable index set  $I^+$  and disjoint open connected components  $S_i^+$ ,  $i \in I^+$ , of  $S^+$  satisfying

$$
\mathcal{S}^+ = \cup \{ \mathcal{S}^+_i \mid i \in I^+ \}.
$$

Moreover, one has  $\partial S^+ = \cup \{ \partial S^+_i \mid i \in I^+ \} \subset C_1 \cup C_2$ . Besides, for any  $i \in I^+$ , the boundary  $\partial S^+_i$  consists of two arcs  $\widehat{M}_i^+ y_1 \widehat{N}_i^+$  and  $\widehat{M}_i^+ y_2 \widehat{N}_i^+$  lying in  $C_1$  and  $C_2$ , respectively, for some end points  $M_i^+, N_i^+ \in C_1 \cap C_2$ . Similarly, if  $S^- \neq 0$ , then it is an union of at most countably many disjoint open connected components  $S_m^-, m \in I^-$ , of  $S^-$  satisfying that the boundary  $\partial S_m^-$  consists of two arcs  $\widehat{M_m y_1 N_m^-}$  and  $\widehat{M_m y_2 N_m^-}$  lying in  $C_1$  and  $C_2$ , respectively, for some end points  $M_m^-, N_m^- \in C_1 \cap C_2$ . Define the set  $\mathcal A$  of closed arcs with end points in  $P := \{M_i^+, N_i^+, M_m^-, N_m^- : i \in I^+, m \in I^-\}$  via

$$
\mathcal{A} := \{ \widehat{MN} \subset C_1 \cap C_2 \mid M, N \in \mathcal{P}, M \neq N, \widehat{MN} \cap (\partial \mathcal{S}^+ \cup \partial \mathcal{S}^-) = \{M, N\} \}.
$$

Obviously,  $A$  consists of at most countable arcs. By renaming the end points of arcs in  $A$ , we obtain an at most countable index set  $I^0$  and points  $M_k^0$ ,  $N_k^0 \in \mathcal{P}$ ,  $k \in I^0$ , that satisfy [\(2.27\)](#page-9-6).

We now prove Green's first identity over nonempty connected components of an open set determined by two functions with nonvanishing gradients.

<span id="page-10-0"></span>Lemma 2.7. Assume that  $y_1, y_2 \in C^1(\overline{\Omega})$  and  $t \in \mathbb{R}$  such that  $\{y_1 > t > y_2\} \neq \emptyset$ . Let  $S$  be a nonempty open connected component of  $\{y_1 > t > y_2\}$  satisfying  $\partial S \cap \partial \Omega = \emptyset$ . Assume further that there is a positive constant  $\alpha$  satisfying

<span id="page-10-2"></span>(2.28) 
$$
\nabla y_1(x) \cdot \nabla y_2(x) \ge \alpha > 0 \quad \text{for all } x \in \overline{S}.
$$

Let  $v \in H^1(\Omega) \cap C(\overline{\Omega})$  and  $\varphi \in W^{2,1}(\Omega) \cap C^1(\overline{\Omega})$  be arbitrary. Then

<span id="page-10-1"></span>
$$
(2.29)\ \int_{S} \nabla v\cdot\nabla \varphi dx=-\int_{S} v \Delta \varphi dx-\int_{\partial S\cap\{y_1=t\}} v \nabla \varphi\cdot \frac{\nabla y_1}{|\nabla y_1|}d\mathcal{H}^1(x)+\int_{\partial S\cap\{y_2=t\}} v \nabla \varphi\cdot \frac{\nabla y_2}{|\nabla y_2|}d\mathcal{H}^1(x).
$$

*Proof.* We first observe that all the integrals in [\(2.29\)](#page-10-1) are well-defined. Note further that  $\partial S \subset \{\gamma_1 =$ *t*}  $\cup$  {*y*<sub>2</sub> = *t*} and that the outward normal vector  $v(x)$  at point  $x \in \partial S$  is defined as

<span id="page-10-3"></span>(2.30) 
$$
v(x) = \begin{cases} -\frac{\nabla y_1(x)}{|\nabla y_1(x)|} & \text{for } x \in \partial S \cap \{y_1 = t\}, \\ \frac{\nabla y_2(x)}{|\nabla y_2(x)|} & \text{for } x \in \partial S \cap \{y_2 = t\}. \end{cases}
$$

Clason, Nhu, Rösch **Numerical analysis of a nonsmooth quasilinear elliptic ...** 

<span id="page-11-0"></span>

Figure 2: A nonempty open component S of the set  $\{y_1 > t > y_2\}$  with the outward normal vectors  $v_1 = -\frac{\nabla y_1}{\nabla x_1}$  $\frac{\nabla y_1}{|\nabla y_1|}$  on  $\{y_1 = t\}$  and  $v_2 = \frac{\nabla y_2}{|\nabla y_2|}$  $\frac{\nabla y_2}{|\nabla y_2|}$  on  $\{y_2 = t\}$  where  $\{y_1 = t\}$  is thin while  $\{y_2 = t\}$  is thick

We now consider the following two cases.

• Case 1: The boundary  $\partial S$  consists of two closed curves having no common points; see, [Figure 2a.](#page-11-0) In this situation, thanks to [\(2.28\)](#page-10-2) and the fact that  $\partial S \cap \partial \Omega = \emptyset$ , we deduce from the Implicit Function Theorem that the domain S has  $C^1$  boundary in the sense of Definition 1.2.1.1 in [\[15\]](#page-37-7). Applying the classical Green formula over the domain  $S$  and employing [\(2.30\)](#page-10-3), we derive [\(2.29\)](#page-10-1).

• Case 2: The boundary  $\partial S$  consists of two arcs  $\widehat{M}\gamma_1\widehat{N}$  and  $\widehat{M}\gamma_2\widehat{N}$  with end points  $M, N \in \{\gamma_1 = t\} \cap \{\gamma_2 = t\}$ t} (here M and N might be identical); see, [Figure 2b](#page-11-0). If  $\partial S$  has no cusps, i.e., there are no points where two branches of  $\partial S$  meet and the tangents at which of each branch are coincidental, then the boundary  $\partial S$ is a curvilinear polygon of class  $C^1$  in the sense of Definition 1.4.5.1 in [\[15\]](#page-37-7), as a result of the combination of the Implicit Function Theorem and [\(2.28\)](#page-10-2). Similar to Case 1, we also have [\(2.29\)](#page-10-1). For the situation where  $\partial S$  has cusps, we shall prove [\(2.29\)](#page-10-1) via approximating S by regular domains having boundaries, which are curvilinear polygons of class  $C^1$ , and via using the fact that  $\varphi \in C^1(\overline{\Omega})$  and  $v \in C(\overline{\Omega})$ . To this end, without loss of generality, we assume that  $\partial S$  has only one cusp at N. For any  $\varepsilon > 0$  small enough, by applying [Lemma](#page-33-0) A.1, there exist  $N_i \in \{y_i = t\} \cap \partial S$ ,  $i = 1, 2$ , such that

$$
l_{\widehat{NN_1}}+l_{\widehat{NN_2}}+N_1N_2
$$

for some positive constant C independent of  $\varepsilon$ , where  $l_{\widehat{NN}}$  denotes the length of the arc  $\widehat{NN}_i$ . Moreover,  $N$  and  $N$  are not guara of  $2S$ , where  $S$  denotes a guarant of  $S$ , where houndary consists of the  $N_1$  and  $N_2$  are not cusps of  $\partial S_{\varepsilon}$ , where  $S_{\varepsilon}$  denotes a subdomain of S, whose boundary consists of the arcs  $\widehat{My_1N_1}, \widehat{My_2N_2}$  and the line segment  $N_1N_2$ . Now applying Green's first formula on the domain  $\mathcal{S}_\varepsilon$ then implies that

$$
\int_{S_{\varepsilon}} \nabla v \cdot \nabla \varphi dx = - \int_{S_{\varepsilon}} v \Delta \varphi dx + \int_{\partial S_{\varepsilon}} v \nabla \varphi \cdot v_{\varepsilon} d\mathcal{H}^{1}(x),
$$

where  $v_{\varepsilon}$  stands for the outward normal vector on  $\partial S_{\varepsilon}$ . Since  $v_{\varepsilon} = v$  on  $\partial S \cap \partial S_{\varepsilon}$  defined in [\(2.30\)](#page-10-3), we can rewrite the above identity as

<span id="page-11-1"></span>
$$
\begin{split} \text{(2.31)}\\ \int_{S_{\varepsilon}}\nabla v\cdot\nabla\varphi dx&=-\int_{S_{\varepsilon}}v\Delta\varphi dx-\int_{\partial S\cap\{y_1=t\}}v\nabla\varphi\cdot\frac{\nabla y_1}{|\nabla y_1|}d\mathcal{H}^1(x)+\int_{\partial S\cap\{y_2=t\}}v\nabla\varphi\cdot\frac{\nabla y_2}{|\nabla y_2|}d\mathcal{H}^1(x)\\ &+\int_{\widehat{N_1y_1N}}v\nabla\varphi\cdot\frac{\nabla y_1}{|\nabla y_1|}d\mathcal{H}^1(x)-\int_{\widehat{N_2y_2N}}v\nabla\varphi\cdot\frac{\nabla y_2}{|\nabla y_2|}d\mathcal{H}^1(x)+\int_{N_1N_2}v\nabla\varphi\cdot v_{\varepsilon}d\mathcal{H}^1(x). \end{split}
$$

Using the continuity over  $\overline{\Omega}$  of  $v$  and  $\nabla \varphi$ , it follows from the choice of  $N_1$  and  $N_2$  that

$$
\left| \int_{\widehat{N_1y_1N}} v \nabla \varphi \cdot \frac{\nabla y_1}{|\nabla y_1|} d\mathcal{H}^1(x) - \int_{\widehat{N_2y_2N}} v \nabla \varphi \cdot \frac{\nabla y_2}{|\nabla y_2|} d\mathcal{H}^1(x) + \int_{N_1N_2} v \nabla \varphi \cdot v_{\varepsilon} d\mathcal{H}^1(x) \right| \leq C_1 \varepsilon
$$

for some positive constant  $C_1$ . By letting  $\varepsilon \to 0^+$  in [\(2.31\)](#page-11-1), we thus deduce [\(2.29\)](#page-10-1) from Lebesgue's Dominated Convergence Theorem. □

<span id="page-12-2"></span>Remark 2.8. Since  $\partial \emptyset = \emptyset$ ,

it is obvious to have the identity [\(2.29\)](#page-10-1) when  $S = \emptyset$ .

The following version of Green's first identity over an open set determined via functions with nonvanishing gradients is a consequence of [Lemmas 2.6](#page-8-0) and [2.7.](#page-10-0)

<span id="page-12-0"></span>Proposition 2.9. Let  $y_1, y_2 \in C^1(\overline{\Omega})$ , let  $t \in \mathbb{R}$  be given, and let  $O$  be an open subset in  $\overline{\Omega}$  such that

(2.32) 
$$
\nabla y_1(x) \cdot \nabla y_2(x) \ge \alpha > 0 \quad \text{for all } x \in O
$$

for some constant  $\alpha$ . Define the open sets  $S^+$  and  $S^-$  by

$$
S^- := \{y_1 < t < y_2\} \cap O \quad \text{and} \quad S^+ := \{y_1 > t > y_2\} \cap O.
$$

Assume that the level set  $\{y_i = t\}$  has one and only one closed  $C^1$  simple curve  $C_i$  in  $O$  for  $j = 1, 2$ . Assume further that  $C_j \cap \partial \Omega = \emptyset$ ,  $j = 1, 2$ , and that  $\partial S^{\pm} \subset C_1 \cup C_2$ . Let  $v \in H^1(\Omega) \cap C(\overline{\Omega})$  and  $\varphi \in W^{2,1}(\Omega) \cap C^1(\overline{\Omega})$ be arbitrary. Then the following identity holds  $(2.33)$ 

<span id="page-12-1"></span>
$$
\int_{\Omega} (\mathbb{1}_{S^+} - \mathbb{1}_{S^-}) \nabla v \cdot \nabla \varphi dx = - \int_{\Omega} (\mathbb{1}_{S^+} - \mathbb{1}_{S^-}) v \Delta \varphi dx - \int_{C_1} v \nabla \varphi \cdot \frac{\nabla y_1}{|\nabla y_1|} d\mathcal{H}^1(x) + \int_{C_2} v \nabla \varphi \cdot \frac{\nabla y_2}{|\nabla y_2|} d\mathcal{H}^1(x).
$$

*Proof.* For the case where  $C_1 = C_2$ , we have  $S^+ = S^- = \emptyset$  and thus  $\mathbb{1}_{S^+} = \mathbb{1}_{S^-} = 0$ . We then obtain [\(2.33\)](#page-12-1). For the case where  $C_1 \cap C_2 = \emptyset$ , one of the sets  $S^+$  and  $S^-$  is empty and the boundary of the other consists of  $C_1$  and  $C_2$ . We then derive [\(2.33\)](#page-12-1) from the classical Green's first formula. It remains to consider the situation where  $C_1 \cap C_2 \neq \emptyset$ ; see [Figure 1.](#page-9-0) As a result of [Lemma 2.6,](#page-8-0) there exist at most countable index sets  $I^+,I^-,I^0,$  points  $M_i^+,N_i^+,M_m^-,N_m^-,M_k^0,N_k^0,$  and open connected domains  ${\cal S}_i^+, {\cal S}_m^$ with  $i \in I^+, m \in I^-$ ,  $k \in I^0$  satisfying assertion [\(c\)](#page-9-5) in [Lemma 2.6.](#page-8-0) (Note that some of the sets  $I^+, I^-, I^0$ might be empty.) For any  $i \in I^+$ , applying [Lemma 2.7](#page-10-0) and [Remark 2.8](#page-12-2) to open connected domain  $\mathcal{S}_i^+$ yields

$$
\int_{S_i^+} \nabla v \cdot \nabla \varphi dx = -\int_{S_i^+} v \Delta \varphi dx - \int_{\partial S_i^+ \cap C_1} v \nabla \varphi \cdot \frac{\nabla y_1}{|\nabla y_1|} d\mathcal{H}^1(x) + \int_{\partial S_i^+ \cap C_2} v \nabla \varphi \cdot \frac{\nabla y_2}{|\nabla y_2|} d\mathcal{H}^1(x).
$$

Summing up the above identities over all  $i \in I^+$  gives

<span id="page-12-4"></span>
$$
(2.34) \quad \int_{S^+} \nabla v \cdot \nabla \varphi dx = -\int_{S^+} v \Delta \varphi dx - \int_{\partial S^+ \cap C_1} v \nabla \varphi \cdot \frac{\nabla y_1}{|\nabla y_1|} d\mathcal{H}^1(x) + \int_{\partial S^+ \cap C_2} v \nabla \varphi \cdot \frac{\nabla y_2}{|\nabla y_2|} d\mathcal{H}^1(x).
$$

Similarly, there holds

<span id="page-12-3"></span>
$$
(2.35)\quad \int_{S^-}\nabla v\cdot\nabla\varphi dx=-\int_{S^-}v\Delta\varphi dx+\int_{\partial S^-\cap C_1}v\nabla\varphi\cdot\frac{\nabla y_1}{|\nabla y_1|}d\mathcal{H}^1(x)-\int_{\partial S^-\cap C_2}v\nabla\varphi\cdot\frac{\nabla y_2}{|\nabla y_2|}d\mathcal{H}^1(x).
$$

By subtracting  $(2.35)$  from  $(2.34)$ , one has

<span id="page-12-5"></span>
$$
(2.36) \quad \int_{\Omega} \left( \mathbb{1}_{S^+} - \mathbb{1}_{S^-} \right) \nabla v \cdot \nabla \varphi dx = -\int_{\Omega} \left( \mathbb{1}_{S^+} - \mathbb{1}_{S^-} \right) v \Delta \varphi dx - \int_{(\partial S^+ \cup \partial S^-) \cap C_1} v \nabla \varphi \cdot \frac{\nabla y_1}{|\nabla y_1|} d\mathcal{H}^1(x) + \int_{(\partial S^+ \cup \partial S^-) \cap C_2} v \nabla \varphi \cdot \frac{\nabla y_2}{|\nabla y_2|} d\mathcal{H}^1(x).
$$

Clason, Nhu, Rösch **Numerical analysis of a nonsmooth quasilinear elliptic ...** 



Figure 3: sets  $C_{\varepsilon}^{+}$  and  $C_{\varepsilon}^{-}$ 

<span id="page-13-1"></span>On the other hand, due to [\(2.27\)](#page-9-6), it follows that

$$
(2.37) \tCj = ((\partial S+ \cup \partial S-) \cap Cj) \cup {\widehat{M_k^0 N_k^0 \mid k \in I^0} \tfor j = 1, 2
$$

with  $\widehat{M^0_k N^0_k}$  $\widetilde{R^0_k}$  denoting the arcs lying on  $C_1 \cap C_2$  with end points  $M^0_k$  and  $N^0_k$  $U_k^0$ . Obviously, one has

<span id="page-13-0"></span>
$$
\frac{\nabla y_1(x)}{|\nabla y_1(x)|} = \frac{\nabla y_2(x)}{|\nabla y_2(x)|} \quad \text{for all } x \in \widehat{M_k^0 N_k^0} \text{ and } k \in I^0.
$$

Combing this with [\(2.36\)](#page-12-5) and [\(2.37\)](#page-13-0) yields [\(2.33\)](#page-12-1).  $\square$ 

#### 2.3 continuity of level sets in term of functions

This subsection is devoted to the continuity of level sets in term of functions determining these level sets. Namely, we will show that when functions  $y_n$  tend to  $y$  in  $C^1(\overline{\Omega})$ , then, in any  $\varepsilon$ -neighborhood (with  $\varepsilon$  small enough) of an arbitrary connected component of the level set  $\{y = t\}$  on which the gradient of y does not vanish, there exists one and only one connected component of  $\{y_n = t\}$  for each  $n$  sufficiently large, see [Proposition 2.11](#page-14-0) and [Corollary 2.12](#page-16-0) below.

From now on, for any  $\varepsilon>0$  and any set  $V\subset\overline{\Omega},$  we denote by  $V^\varepsilon$  the open  $\varepsilon$ -neighborhood in  $\overline{\Omega}$  of  $V$ , that is,

$$
V^{\varepsilon} := \{ x \in \overline{\Omega} \mid \, \text{dist}(x, V) < \varepsilon \},
$$

where dist(x, V) is the distance from x to V. If C is a closed simple curve in  $\overline{\Omega}$ , we define the open sets  $C_{\varepsilon}^{+}$  and  $C_{\varepsilon}^{-}$  (illustrated in [Figure 3\)](#page-13-1) as follows

$$
C_{\varepsilon}^- := \{ x \in \overline{\Omega} \mid 0 < \text{dist}(x, C) < \varepsilon \text{ and } x \text{ is surrounded by } C \} \quad \text{and} \quad C_{\varepsilon}^+ := C^{\varepsilon} \setminus (C_{\varepsilon}^- \cup C).
$$

In the remainder of this section, we shall consider the following general situation.

<span id="page-13-2"></span>(H1) Let  $y_n, y \in C^1(\overline{\Omega})$ ,  $n \ge 1$ , such that  $y_n = y = 0$  on the boundary  $\partial \Omega$  and

<span id="page-13-3"></span>(2.38) 
$$
y_n \to y
$$
 strongly in  $C^1(\overline{\Omega})$ .

We have the following result on strong positivity of the product of gradients of  $y_n$  and  $y$  around an  $\varepsilon$ -neighborhood of any connected component in the level set  $\{y = t\}$ .

<span id="page-13-4"></span>Proposition 2.10. Let functions  $y_n, y, n \geq 1$ , satisfy hypothesis (H1). Assume that C is a connected component of  $\{y = t\}$  for some  $t \in \mathbb{R}$  and fulfills the condition [\(2.20\)](#page-7-4). Then there exist constants  $\varepsilon_0 > 0$ ,  $\alpha_C > 0$ , and  $n_0 \in \mathbb{N}$  such that

<span id="page-13-5"></span>(2.39)  $\nabla y(x) \cdot \nabla y_n(x') \ge \alpha_C \quad \text{for all } n \ge n_0, x, x' \in C^{\varepsilon_0} \text{ with } |x - x'| \le 2\varepsilon_0.$ 

*Proof.* By contradiction, assume that there exist sequences  $\varepsilon_k \to 0^+$ ,  $n_k \to \infty$  satisfying

<span id="page-14-1"></span>
$$
(2.40) \t x_k, x'_k \in C^{\varepsilon_k}, \t |x_k - x'_k| \le 2\varepsilon_k \text{ and } \nabla y(x_k) \cdot \nabla y_{n_k}(x'_k) < \frac{1}{k} \text{ for all } k \ge 1.
$$

By extracting a subsequence if necessary, we conclude from the boundedness of  $\{x_k\}$  and  $\{x_k'\}$  and the first two conditions in [\(2.40\)](#page-14-1) as well as from the continuity of y that  $x_k \to x$  and  $x'_k \to x$  for some  $x \in C$ . Passing to the limit the last condition in [\(2.40\)](#page-14-1) and using [\(2.38\)](#page-13-3) yield

<span id="page-14-2"></span>
$$
|\nabla y(x)|^2 = 0,
$$

which contradicts [\(2.20\)](#page-7-4). The proposition is proven.  $\Box$ 

<span id="page-14-0"></span>Proposition 2.11. Let functions  $y_n, y, n \ge 1$ , satisfy hypothesis (H1). Let C be a connected component of  $\{y = t\}$ ,  $t \in \mathbb{R}$  that satisfies [\(2.20\)](#page-7-4). Then a constant  $\varepsilon_0 > 0$  exists such that for any  $\varepsilon \in (0, \varepsilon_0)$ , there is a positive integer  $n_0 = n_0(\varepsilon)$  such that  $C^{\varepsilon}$  contains one and only one connected component  $C_n$  of  $\{y_n = t\}$ for all  $n \geq n_0$ .

*Proof.* Since C and  $\{\gamma = t\} \backslash C$  are both closed and disjoint, a constant  $\varepsilon_0 > 0$  thus exists and fulfills

$$
(2.41) \tC\varepsilon_0 \cap (\{y=t\} \setminus C)^{\varepsilon_0} = \emptyset.
$$

Obviously, we can assume that the constant  $\varepsilon_0$  in [\(2.41\)](#page-14-2) is identical to the one in [Proposition 2.10.](#page-13-4) Fix  $\varepsilon \in (0, \varepsilon_0)$  and, without loss of generality, assume that

$$
(2.42) \t\t y < t \t on C\varepsilon_0- and y > t \t on C\varepsilon_0+.
$$

We now consider two possible situations:  $C \cap \partial \Omega \neq \emptyset$  and  $C \cap \partial \Omega = \emptyset$ .

• For the case  $C \cap \partial\Omega \neq \emptyset$ , we have  $t = 0$  and  $C = \partial\Omega$ . Moreover, there hold  $C_{\varepsilon_0}^+ = \emptyset$  and  $\partial\Omega \subset \{y_n = 0\}$ . Thanks to [\(2.39\)](#page-13-5) and [Proposition 2.4,](#page-7-0)  $C_n := \partial\Omega$  is a connected component of  $\{y_n = 0\}$ for all  $n \ge 1$ , and is a closed simple curve in  $C^{\varepsilon}$  for all  $\varepsilon \in (0, \varepsilon_0)$ . If there exists another connected component  $C'_n$  of  $\{y_n = 0\}$  such that  $C'_n \cap C^{\varepsilon} \neq \emptyset$  for *n* sufficient large, then by picking a point  $x_2 = (x_2^1, x_2^2) \in C'_n \cap C^{\varepsilon}$  and taking

<span id="page-14-5"></span><span id="page-14-3"></span>
$$
x_1 \in \operatorname{argmin}\{|x_2 - x| : x \in C\},\
$$

one has  $|x_1 - x_2| \leq \varepsilon$  and

(2.43) 
$$
x_2 - x_1 = \kappa_n \nabla y(x_1) \text{ for some } \kappa_n \in \mathbb{R},
$$

as a result of [Lemma](#page-34-0) A.2. Since  $C_n \cap C'_n = \emptyset$  and  $C_n = \partial \Omega = C$ , we obtain  $x_1 \neq x_2$  and thus

$$
k_n \neq 0.
$$

On the other hand, since  $x_1 \in C = C_n$  and  $x_2 \in C'_n$ ,  $y_n(x_1) = t = y_n(x_2)$ . We thus deduce from the Mean Value Theorem that  $\nabla y_n(x_1 + \theta_n(x_2 - x_1)) \cdot (x_2 - x_1) = 0$  with  $\theta_n \in (0, 1)$ . Combining this with [\(2.43\)](#page-14-3) and  $(2.44)$  yields

<span id="page-14-4"></span>
$$
\nabla y_n(x_1 + \theta_n(x_2 - x_1)) \cdot \nabla y(x_1) = 0,
$$

which contradicts  $(2.39)$  for *n* large enough.

• For the case  $C \cap \partial \Omega = \emptyset$ , we have  $C_{\varepsilon}^{\pm} \neq \emptyset$ . We now split the proof into several Claims below. *Claim 1: There exists an integer*  $n_0^1$  such that  $\{y_n = t\} \cap C^{\varepsilon} \neq \emptyset$  for all  $n \geq n_0^1$ . In fact, arguing by contradiction, there exists a subsequence of  $\{n\}$ , denoted in the same way, such that  $\{y_n = t\} \cap C^{\varepsilon} = \emptyset$ 

$$
(2.45) \tC\varepsilon \subset \{y_n > t\} \t or \tC\varepsilon \subset \{y_n < t\}.
$$

For the first case in  $(2.45)$ , we have

<span id="page-15-0"></span>
$$
C_{\varepsilon}^- \subset \{y < t < y_n\} \subset \{0 < |y - t| \leq \tau_n\}
$$

with  $\tau_n := ||y_n - y||_{C(\overline{\Omega})}$ . This implies that  $meas_{\mathbb{R}^2}(C_{\varepsilon}^-) \leq meas_{\mathbb{R}^2}(\{0 < |y - t| \leq \tau_n\}) \to 0$  as  $n \to \infty$ , contradicting the fact that meas $_{\mathbb{R}^2}(C_{\varepsilon}^-) > 0$ . The first case in [\(2.45\)](#page-15-0) is then impossible. Similarly, the second one is also absurd.

*Claim 2: There exists an integer*  $n_0^2$  *such that*  $\{y_n = t\} \cap \partial C^{\epsilon} = \emptyset$  *for all*  $n \ge n_0^2$ . Arguing by contradiction, there are subsequences  $n_k \to \infty$  and  $\{x_k\}$  with  $x_k \in \{y_{n_k} = t\} \cap \partial C^\varepsilon$  for all  $k \geq 1$ . From the boundedness of  $\{x_k\}$  and the closedness of  $\partial C^\varepsilon$ , we can assume that  $x_k \to x \in \partial C^\varepsilon$  as  $k \to \infty$ . Thanks to [\(2.38\)](#page-13-3), there holds  $x \in \{y = t\}$ . We then have  $\{y = t\} \cap \partial C^{\varepsilon} \neq \emptyset$ , which contradicts [\(2.41\)](#page-14-2).

From Claims 1 and 2, there exists an integer  $n_0$  such that  $\{y_n = t\}$  admits a connected component  $C_n$  being in  $C^{\varepsilon}$  for all  $n \geq n_0$ .

For any  $n \geq n_0$ , we define a subset (depending on *n*) of *C* by setting

$$
(2.46) \qquad \qquad \Lambda_n := \{x_1 \in C \mid \exists x_2 \in \{y_n = t\} \cap C^{\varepsilon} \text{ satisfying } (2.47)-(2.48)\},
$$

where

$$
(2.47) \t\t\t |x_1 - x_2| \leq \varepsilon
$$

and

$$
(2.48) \t\t x2 - x1 = \kappa \nabla y(x1), \tfor some  $\kappa \in \mathbb{R}$ .
$$

In view of [Lemma](#page-34-0) A.2, we have  $\Lambda_n \neq \emptyset$ . It remains to prove the following claims.

*Claim 3: For any n*  $\geq n_0$  and any  $x_1 \in C$ , there exists at most one point  $x_2 \in \{y_n = t\} \cap C^{\varepsilon}$  such that  $x_1$ and  $x_2$  satisfy [\(2.47\)](#page-15-1) and [\(2.48\)](#page-15-2). Suppose this is not the case, i.e., suppose that there were a point  $x_1 \in C$ and two different points  $p_2, p'_2 \in \{y_n = t\} \cap C^{\varepsilon}$  satisfying [\(2.47\)](#page-15-1) and [\(2.48\)](#page-15-2) in place of  $x_2$ , respectively, for some  $\kappa := \kappa$  and  $\kappa := \kappa'$ . The relation [\(2.47\)](#page-15-1) for both  $x_2 := p_2$  and  $x_2 := p'_2$  implies that  $|p_2 - p'_2| \leq 2\varepsilon$ . Similarly, [\(2.48\)](#page-15-2) indicates that

<span id="page-15-2"></span><span id="page-15-1"></span>
$$
p_2 - p'_2 = (\kappa - \kappa') \nabla y(x_1),
$$

which, together with the condition that  $p_2 \neq p'_2$ , yields  $(\kappa - \kappa') \neq 0$ . Since  $p_2, p'_2 \in \{y_n = t\} \cap C^{\varepsilon}$ , one has

$$
y_n(p_2) = y_n(p_2') = t.
$$

By the Mean Value Theorem, there holds

$$
\nabla y_n (p'_2 + \theta (p_2 - p'_2)) \cdot (p_2 - p'_2) = 0
$$

for some constant  $\theta \in (0, 1)$ . Combing this with the fact that  $(p_2 - p_2') = (\kappa - \kappa') \nabla y(x_1)$  and that  $(\kappa - \kappa') \neq 0$  yields  $\nabla y_n (p'_2 + \theta(p_2 - p'_2)) \cdot \nabla y(x_1) = 0$ , contradicting [\(2.39\)](#page-13-5).

Claim 4: There exists an integer  $\bar{n} \ge n_0$  such that  $\Lambda_n = C$  for all  $n \ge \bar{n}$ . Arguing by contradiction, assume that there exist a subsequence  $\{k\}$  of  $\{n\}$  and a sequence of points  $\{x_{1,k}\}\subset C$  such that, for

each  $k \ge 1$ , there is no points  $x_{2,k}$  satisfying both [\(2.47\)](#page-15-1) and [\(2.48\)](#page-15-2) corresponding to  $x_1 := x_{1,k}$  and  $x_2 := x_{2,k}$ . This means that the line  $d_k$  through  $x_{1,k}$  with direction vector  $\nabla y(x_{1,k})$  does not intersect  $\{y_{n_k} = t\} \cap B_{\mathbb{R}^2}(x_{1,k}, \varepsilon)$  for all  $k \ge 1$ . By Claim 2, we have  $\{y_{n_k} = t\} \cap \partial C^{\varepsilon} = \emptyset$ , and thus the sign of  $(y_{n_k} - t)$  does not change over the set  $A_k B_k := d_k \cap \overline{B}_{\mathbb{R}^2}(x_{1,k}, \varepsilon)$  with  $A_k \in \partial C_{\varepsilon}^+$  and  $B_k \in \partial C_{\varepsilon}^-$ . We now split the sequence  $\{k\}$  into subsequences, also denoted by  $\{k\}$ , that satisfy one of the following conditions for all  $k \geq 1$ :

(2.49) 
$$
y_{n_k} - t > 0 \quad \text{over } A_k B_k \quad \text{or} \quad y_{n_k} - t < 0 \quad \text{over } A_k B_k.
$$

We now consider the first case in [\(2.49\)](#page-16-1) only, since the other can be treated similarly. For all  $k \geq 1$ , according to [\(2.42\)](#page-14-5), one has

<span id="page-16-1"></span>
$$
y_{n_k}(B_k) - y(B_k) > t - y(B_k) > t - \sup\{y(x) \mid x \in (\partial C_{\varepsilon}^-) \backslash C\} > m_0 > 0
$$

for some constant  $m_0$  independent of  $k.$  This contradicts the limit  $y_{n_k}\to y$  in  $C^1(\overline{\Omega})$  as  $k\to\infty.$ 

From Claims 3 and 4, we can define for all  $n \geq \bar{n}$  the following map (depending on n)

$$
T: C \to \{y_n = t\} \cap C^s
$$

$$
x_1 \mapsto x_2,
$$

with  $x_1$  and  $x_2$  satisfying [\(2.47\)](#page-15-1)–[\(2.48\)](#page-15-2). Moreover, by [Lemma](#page-34-0) A.2, T is surjective.

*Claim 5: For all n*  $\geq \bar{n}$ , *T is continuous.* In fact, assume that  $x_{1,k} \to x_1$  as  $k \to \infty$  with  $x_{1,k}, x_1 \in C$  and  $x_{2,k} = T(x_{1,k}), k \ge 1$ . Then there exists  $\{ \kappa_k \} \subset \mathbb{R}$  such that

$$
|x_{1,k} - x_{2,k}| \leq \varepsilon \quad \text{and} \quad x_{2,k} - x_{1,k} = \kappa_k \nabla y(x_{1,k})
$$

for all  $k \geq 1$ . Therefore  $\{x_{2,k}\}$  is bounded and  $\{\kappa_k\}$  is also bounded due to [\(2.20\)](#page-7-4). Let  $x_2$  and  $\kappa$  be an arbitrary limit point of  $\{x_{2,k}\}\$  and  $\{\kappa_k\}\$ , respectively. We easily have

$$
|x_1 - x_2| \le \varepsilon \quad \text{and} \quad x_2 - x_1 = \kappa \nabla y(x_1).
$$

Moreover, the closedness of  $\{y_n = t\}$  yields  $x_2 \in \{y_n = t\} \cap C^{\varepsilon}$ . Combing this with Claim 3 yields  $x_2 = T(x_1)$ . Consequently, the full sequence  $\{x_{2,k}\}$  converges to  $x_2$  and T is then continuous.

For any  $n \geq \bar{n}$ , we have  $T(C) = \{y_n = t\} \cap C^{\varepsilon}$  since T is surjective. Moreover, because C is connected and compact, and T is continuous,  $T(C)$  is also connected and compact; see, e.g. Theorem 4.22 in [\[21\]](#page-37-10). Therefore  $\{y_n = t\} \cap C^{\varepsilon}$  is connected. From this and Claim 2, we can conclude that  $\{y_n = t\} \cap C^{\varepsilon}$  is the unique connected component of  $\{y_n = t\}$  in  $C^{\varepsilon}$ . The proof is complete.

The following result is a direct consequence of [Propositions 2.4](#page-7-0) and [2.11.](#page-14-0)

<span id="page-16-0"></span>Corollary 2.12. Let functions  $y_n$ ,  $y$ ,  $n \geq 1$ , satisfy hypothesis (H1). Assume that y, together with some value  $t \in \mathbb{R}$ , satisfies [\(2.21\)](#page-7-5). Then

$$
\{y=t\}=\cup_{i=1}^m C_i,
$$

for some integer  $m \geq 1$ , where  $C_i$ ,  $1 \leq i \leq m$ , are  $C^1$  closed simple curves. Moreover, a constant  $\varepsilon_0 > 0$ exists such that for any  $\varepsilon \in (0, \varepsilon_0)$ , there is a positive integer  $n_0 = n_0(\varepsilon)$  with

$$
\{y_n = t\} = \bigcup_{i=1}^m C_{i,n} \quad \text{and} \quad C_{i,n} \subset C_i^{\varepsilon} \quad \text{for all } n \ge n_0, 1 \le i \le m
$$

for  $C^1$  closed simple curves  $C_{i,n}$ .

#### <span id="page-17-1"></span>2.4 CONTINUITY OF INTEGRALS OVER LEVEL SETS

The following result stating the continuity of integrals over connected components of a level set is shown under the nonvanishing gradient condition [\(2.20\)](#page-7-4) by using partitions of unity.

<span id="page-17-0"></span>Proposition 2.13. Let functions  $y_n$ ,  $y$ ,  $n \ge 1$ , satisfy hypothesis (H1). Let C be a connected component of  $\{y = t\}$ ,  $t \in \mathbb{R}$  that satisfies [\(2.20\)](#page-7-4). Assume that  $f_n(x) \to f(x)$  for all  $x \in C^{\bar{\varepsilon}}$  for some constant  $\bar{\varepsilon} > 0$ . Then

<span id="page-17-2"></span>(2.50) 
$$
\int_{\{y_n=t\}\cap C^{\tilde{\varepsilon}}} f_n(x) d\mathcal{H}^1(x) \to \int_C f(x) d\mathcal{H}^1(x) \quad \text{as} \quad n \to \infty.
$$

*Proof.* In light of [Propositions 2.10](#page-13-4) and [2.11,](#page-14-0) there exist positive constants  $\varepsilon_0$ ,  $\alpha_C > 0$  and an integer  $n_0$ such that

 $(2.51)$ 

<span id="page-17-3"></span> $\int \nabla y(x) \cdot \nabla y_n(x') \geq \alpha_C$  for all  $n \geq n_0, x, x' \in C^{\varepsilon_0}$  with  $|x - x'| \leq 2\varepsilon_0$ ,  $C^{\varepsilon}$  contains one and only one a connected component  $C_n$  of  $\{y_n = t\}$  for all  $n \ge n_0$  and  $\varepsilon \in (0, \varepsilon_0)$ .

Setting  $\alpha_{\min} := \min\{|\nabla y(x)| \mid x \in C\}$  yields  $\alpha_{\min} > 0$ . For each  $x \in C$ , there exists a constant  $r_x \in (0, \min\{\varepsilon_0, \bar{\varepsilon}\})$  such that one of the following two estimates is valid:

$$
\left|\frac{\partial y}{\partial x^i}(z)\right| \ge \frac{\alpha_{\min}}{2} > 0 \quad \text{for all } z \in \overline{B}_{\mathbb{R}^2}(x, r_x) \cap \overline{\Omega}, i = 1, 2.
$$

Since C is compact, it is covered by finitely many open balls  $B_{\mathbb{R}^2}(x_1,r_1),\ldots,B_{\mathbb{R}^2}(x_{m_C},r_{m_C})$  with  $r_j:=r_{x_j}$ for all  $1 \le j \le m_C$ . By using partitions of unity, see. e.g. Theorem C.21 and Exercise C.22 in [\[17\]](#page-37-8), there exist nonnegative functions  $\psi_j \in C_c^{\infty}(\Omega)$  such that

$$
\mathrm{supp}(\psi_j) \subset B_{\mathbb{R}^2}(x_j, r_j) \quad \text{and} \quad \sum_{j=1}^{m_C} \psi_j(x) = 1 \quad \text{for all } 1 \le j \le m_C, x \in \Omega.
$$

Therefore, in order to show [\(2.50\)](#page-17-2), it suffices to prove for all  $1 \le j \le m_C$  that

<span id="page-17-4"></span>
$$
(2.52) \qquad \int_{\{y_n=t\}\cap C^{\bar{\varepsilon}}\cap B_{\mathbb{R}^2}(x_j,r_j)} f_n(x)\psi_j(x) d\mathcal{H}^1(x) \to \int_C f(x)\psi_j(x) d\mathcal{H}^1(x) \quad \text{as} \quad n \to \infty.
$$

To this end, fix j, put  $x_0 := x_i$ ,  $r_0 := r_i$ , and without loss of generality assume that  $x_0 = (0, 0)$  and that

$$
\left|\frac{\partial y}{\partial x^2}(z)\right| \ge \frac{\alpha_{\min}}{2} > 0 \quad \text{for all } z \in \overline{B}_{\mathbb{R}^2}(x_0, r_0) \cap \overline{\Omega}.
$$

Applying [Lemma 2.2,](#page-4-0) exploiting [\(2.51\)](#page-17-3) and the fact that  $0 < r_i < \min\{\varepsilon_0, \bar{\varepsilon}\}\)$ , we conclude that there exist positive constants  $h_0, k_0, h_n, k_n$  and  $C^1$  functions  $g_0, g_n$  satisfying assertions [\(a\)](#page-4-4) and [\(i\)–](#page-4-7)[\(iv\)](#page-4-10) in [\(b\)](#page-4-5) of [Lemma 2.2.](#page-4-0) Hence, the limit [\(2.52\)](#page-17-4) can be expressed as

$$
\int_{-h_n}^{k_n} f_n(\tau, g_n(\tau)) \psi_j(\tau, g_n(\tau)) \sqrt{1 + g'_n(\tau)^2} d\tau \to \int_{-h_0}^{k_0} f(\tau, g_0(\tau)) \psi_j(\tau, g_0(\tau)) \sqrt{1 + g'_0(\tau)^2} d\tau \quad \text{as} \quad n \to \infty,
$$

or, equivalently,

(2.53) 
$$
\int_{-h_n}^{k_n} \phi_n(\tau) d\tau - \int_{-h_0}^{k_0} \phi_0(\tau) d\tau \to 0
$$

with

<span id="page-17-5"></span>
$$
\phi_n(\tau) := f_n(\tau, g_n(\tau))\psi_j(\tau, g_n(\tau))\sqrt{1+g'_n(\tau)^2} \quad \text{and} \quad \phi_0(\tau) := f(\tau, g_0(\tau))\psi_j(\tau, g_0(\tau))\sqrt{1+g'_0(\tau)^2}.
$$

Clason, Nhu, Rösch Numerical analysis of a nonsmooth quasilinear elliptic ...

Taking now  $\varepsilon > 0$  small enough, we rewrite the right-hand side term in [\(2.53\)](#page-17-5) as

$$
\int_{-h_n}^{k_n} \phi_n(\tau) d\tau - \int_{-h_0}^{k_0} \phi_0(\tau) d\tau = \int_{-h_0 + \varepsilon}^{k_0 - \varepsilon} [\phi_n(\tau) - \phi_0(\tau)] d\tau \n- \int_{-h_0}^{-h_0 + \varepsilon} \phi_0(\tau) d\tau - \int_{k_0 - \varepsilon}^{k_0} \phi_0(\tau) d\tau + \int_{-h_n}^{-h_0 + \varepsilon} \phi_n(\tau) d\tau + \int_{k_0 - \varepsilon}^{k_n} \phi_n(\tau) d\tau.
$$

In view of assertion [\(iv\)](#page-4-10) in [Lemma 2.2,](#page-4-0) there holds

$$
\phi_n(\tau) \to \phi_0(\tau)
$$
 for all  $\tau \in [-h_0 + \varepsilon, k_0 - \varepsilon]$  and  $|\phi_n(\tau)| \le C$  for all  $\tau \in [-h_n, k_n]$ .

Lebesgue's Dominated Convergence Theorem then implies that

$$
\lim_{n\to\infty}\left|\int_{-h_n}^{k_n}\phi_n(\tau)d\tau-\int_{-h_0}^{k_0}\phi_0(\tau)d\tau\right|\leq \tilde{C}\varepsilon,
$$

where we have used the limits  $h_n \to h_0$  and  $k_n \to k_0$ ; see assertion [\(iii\)](#page-4-9) in [Lemma 2.2.](#page-4-0) We thus obtain  $(2.53)$ .

#### <span id="page-18-1"></span>3 an nonsmooth quasilinear elliptic optimal control problem

#### <span id="page-18-0"></span>3.1 main assumptions and preliminary results

Let  $a$  be a finitely  $PC^2$ -function of the form

<span id="page-18-2"></span>(3.1) 
$$
a(t) := \mathbb{1}_{(-\infty,\bar{t}]}(t)a_0(t) + \mathbb{1}_{(\bar{t},\infty)}a_1(t) \text{ for all } t \in \mathbb{R},
$$

for a given number  $\bar{t} \in \mathbb{R}$  and given functions  $a_0 \in C^2((-\infty, \bar{t}])$  and  $a_1 \in C^2([\bar{t}, \infty))$  with  $a_0(\bar{t}) = a_1(\bar{t})$ . Obviously, the function *a* is of class  $C^2$  over the intervals  $(-\infty, \bar{t}) \cup (\bar{t}, \infty)$ , but not even of class  $C^1$  in general.

Remark 3.1. Let us emphasize that the results and the underlying analysis in this paper can be applied to the situation in which the function  $a$  is continuous and is twice continuously differentiable on finitely many intervals (i.e., a finitely  $PC^2$  function; see [\[9\]](#page-36-4) for a precise definition). However, in order to keep the presentation concise and to be able to focus on the main arguments, we restrict the presentation to the simplest such situation given by  $(3.1)$ .

By  $\{a'\}_{\bar{t}+0}^{\bar{t}-0}$  $\frac{\bar{t}-0}{\bar{t}+0}$ , we denote the difference between the one-sided derivatives of *a* at  $\bar{t}$  from left and right, i.e.,

<span id="page-18-3"></span>
$$
\{a'\}_{\bar{t}+0}^{\bar{t}-0} := \lim_{t \to \bar{t}^-} a'(t) - \lim_{t \to \bar{t}^+} a'(t) = a'_0(\bar{t}) - a'_1(\bar{t}).
$$

By setting

(3.2) 
$$
\sigma_0 := |\{a'\}_{\bar{t}+0}^{\bar{t}-0}| = |a'_0(\bar{t}) - a'_1(\bar{t})|,
$$

we see that this term determines the differentiability of  $a$  and plays a crucial part in the second-order optimality conditions for  $(P)$ ; see [\[9\]](#page-36-4). Moreover, *a* is directionally differentiable and its directional derivative is given by

$$
(3.3) \quad a'(t;s) = \mathbb{1}_{(-\infty,\bar{t})}(t)a'_0(t)s + \mathbb{1}_{(\bar{t},\infty)}(t)a'_1(t)s + \mathbb{1}_{\{\bar{t}\}}(t)[\mathbb{1}_{(0,\infty)}(s)a'_1(\bar{t})s + \mathbb{1}_{(-\infty,0)}(s)a'_0(\bar{t})s], \quad t,s \in \mathbb{R}.
$$

The following assumptions shall hold throughout the whole paper except in [Section 3.2,](#page-22-0) where we will only require the convexity of  $\Omega$  instead of Assumption (A1) below.

- <span id="page-19-0"></span>(A1)  $\Omega \subset \mathbb{R}^2$  is an open bounded convex polygonal.
- <span id="page-19-5"></span>(A2) The Lipschitz continuous function  $b : \overline{\Omega} \to \mathbb{R}$  satisfies  $b(x) \ge b > 0$  for all  $x \in \overline{\Omega}$ .
- <span id="page-19-2"></span>(A3)  $a : \mathbb{R} \to \mathbb{R}$  is nonnegative and given by [\(3.1\)](#page-18-2).
- <span id="page-19-1"></span>(A4)  $L : \Omega \times \mathbb{R} \to \mathbb{R}$  is a Carathéodory function that is of class  $C^2$  w.r.t. the second variable with  $L(·, 0) ∈ L<sup>1</sup>(Ω)$ . Besides, for any  $\overrightarrow{M} > 0$ , there exist  $C_M > 0$  and  $\psi_M ∈ L<sup>\bar{p}</sup>(Ω)$  ( $\bar{p} > 2$ ) such that |  $\left|\frac{\partial L}{\partial y}(x, y)\right| \leq \psi_M(x)$  and  $\left|\frac{\partial^2 L}{\partial y^2}\right|$  $\frac{\partial^2 L}{\partial y^2}(x, y)| \leq C_M$  for all  $y \in \mathbb{R}$  with  $|y| \leq M$ , and a.e.  $x \in \Omega$ .

From [Assumption \(a4\),](#page-19-1) we deduce by explicit computation, a Taylor expansion, and Lebesgue's Dominated Convergence Theorem that the functional  $L^{\infty}(\Omega) \ni y \mapsto \int_{\Omega} L(x, y(x)) dx \in \mathbb{R}$  is of class  $C^2$ .

In the remainder of this subsection, we state some known results for the state equation, the adjoint state equation, and the optimality conditions for [\(P\)](#page-0-0); see, e.g. [\[9\]](#page-36-4). Let us first consider the state equation

<span id="page-19-3"></span>(3.4) 
$$
-\operatorname{div}[(b + a(y))\nabla y] = u \text{ in } \Omega, \quad y = 0 \text{ on } \partial\Omega.
$$

<span id="page-19-7"></span>Theorem 3.2 (cf. [\[9\]](#page-36-4), Thms. 3.1 and 3.5). Let Assumptions (A1) to (A3) hold. Then, the control-to-state operator  $S: W^{-1,p}(\Omega) \ni u \mapsto y_u \in W_0^{1,p}$  $\widetilde{U}_0^{1,p}(\Omega)$  with  $y_u$  being the unique solution to [\(3.4\)](#page-19-3) is of class  $C^1$ . Moreover, for any  $u, v \in W^{-1,p}(\Omega)$  with  $p > 2$  and  $y_u := S(u), z_v := S'(u)v$  is the unique solution to

<span id="page-19-8"></span>(3.5) 
$$
-\operatorname{div}[(b + a(y_u))\nabla z_v + \mathbb{1}_{\{y_u \neq \bar{t}\}} a'(y_u)z_v \nabla y_u] = v \text{ in } \Omega, \quad z_v = 0 \text{ on } \partial\Omega.
$$

Moreover, there exists a number  $p_* > 2$  such that for any  $p \in [2, p_*)$  and for any bounded set  $U \subset L^p(\Omega)$ , there hold  $S(u) \in W_0^{1,p}$  $U_0^{1,p}(\Omega) \cap W^{2,p}(\Omega)$  and  $||S(u)||_{W^{2,p}(\Omega)} \leq C_U$ .

Proof. The wellposedness and the continuous differentiability of  $S$  follows from Theorems 3.1 and 3.5 in [\[9\]](#page-36-4). On the other hand, the  $H^2$ - and  $W^{1,\infty}(\Omega)$ -regularity of solutions to [\(3.4\)](#page-19-3) was also shown in Theorem 3.1 in [\[9\]](#page-36-4) when the right-hand side u belongs to  $L^q(\Omega)$  with  $q > 2$ . Moreover, if U is a bounded subset of  $L^q(\Omega)$  with  $q > 2$ , then there holds  $||S(u)||_{H^2(\Omega)} + ||S(u)||_{W^{1,\infty}(\Omega)} \leq C_U$  for all  $u \in U$ . To show the higher  $W^{2,p}$ -regularity as well as the corresponding a priori estimate, we observe that [\(3.4\)](#page-19-3) can be rewritten as

<span id="page-19-4"></span>(3.6) 
$$
-\Delta y = \frac{1}{b + a(y)} [u + \nabla b \cdot \nabla y + \mathbb{1}_{\{y \neq \bar{t}\}} a'(y) |\nabla y|^2] \text{ in } \Omega, \quad y = 0 \text{ on } \partial \Omega
$$

(see, e.g. equations (A.1) and (A.3) in [\[9\]](#page-36-4), Lem. A.1). Since  $\Omega$  is assumed to be a convex polygon in  $\mathbb{R}^2,$ Theorem 4.4.3.7 in [\[15\]](#page-37-7) shows that there exists a constant  $p_* := 2/(2 - \min\{\pi \omega_{\max}^{-1}, 2\}) > 2$  depending on the maximal interior angle  $\omega_{\text{max}} < \pi$  of the domain  $\Omega$  such that any solution  $\gamma$  to [\(3.6\)](#page-19-4) belongs to  $W^{2,p}(\Omega)$  provided that  $u \in L^p(\Omega)$  for all  $p \in (2, p_*)$ . Of course, we have  $y \in H^2(\Omega)$  when  $u \in L^2(\Omega)$  due to the convexity of  $\Omega$ . Finally, the  $W^{2,p}$ -estimate of solutions y is derived by applying Theorem 4.3.2.4 in [\[15\]](#page-37-7) to [\(3.6\)](#page-19-4) and using the a priori  $W^{1,\infty}(\Omega)$ -estimates of y, Assumptions (A2) and (A3).  $\Box$ 

<span id="page-19-6"></span>We now consider the adjoint state equation

(3.7) 
$$
-\operatorname{div}[(b + a(y_u))\nabla \varphi] + \mathbb{1}_{\{y_u \neq \bar{t}\}} a'(y_u) \nabla y_u \cdot \nabla \varphi = v \text{ in } \Omega, \quad \varphi = 0 \text{ on } \partial \Omega
$$

for  $u \in W^{-1,p}(\Omega)$ ,  $p > 2$ ,  $v \in H^{-1}(\Omega)$ , and  $y_u := S(u)$ .

<span id="page-19-9"></span>Theorem 3.3 (cf. [\[9\]](#page-36-4), Lem. 4.1). Let Assumptions (A1) to (A3) be satisfied and let  $p, q > 2$  be arbitrary. Then, for any  $u \in W^{-1,p}(\Omega)$ ,  $v \in H^{-1}(\Omega)$ , a unique  $\varphi \in H_0^1(\Omega)$  exists and uniquely solves [\(3.7\)](#page-19-6). Furthermore, if U is a bounded subset in  $L^p(\Omega)$ , then for any  $u \in U$  and any  $v \in L^q(\Omega)$ , the solution  $\varphi$  of [\(3.7\)](#page-19-6) belongs to  $H^2(\Omega) \cap W^{1,\infty}(\Omega)$  and there holds  $\|\varphi\|_{H^2(\Omega)} + \|\varphi\|_{W^{1,\infty}(\Omega)} \leq C_U \|v\|_{L^q(\Omega)}$ . Moreover, if  $u \in L^p(\Omega)$  and  $v \in L^r(\Omega)$  with  $r \in (2, p_*)$ , then  $\varphi \in W^{2,r}(\Omega)$ , where  $p_*$  is defined as in [Theorem 3.2.](#page-19-7)

*Proof.* Thanks to Lemma 4.1 in [\[9\]](#page-36-4), all conclusions except the last one of the theorem are verified. To derive the  $W^{2,r}$ -regularity of  $\varphi$ , we argue as in the proof of [Theorem 3.2.](#page-19-7)

Remark 3.4. Despite the  $W^{2,p}$ -regularity of the state and adjoint state, the function  $z_v := S'(u)v$  defined in [\(3.5\)](#page-19-8) is only in  $W^{1,p}(\Omega)$  due to the fact that we cannot apply the chain rule for the divergence acting on the term  $\mathbb{1}_{\{y\neq i\}} a'(y)z\nabla y$  even with  $y \in W^{2,p}(\Omega)$  and  $z \in W^{1,p}(\Omega)$ . Indeed, for any  $y \in W^{2,p}(\Omega)$ , the vector-valued function  $f(x) := \mathbb{1}_{\{y \neq \bar{t}\}}(x) a'(y(x)) \nabla y(x)$  in general does not belong to  $(W^{1,p}(\Omega))^{N}$  with  $N = 2$ . For example, considering  $N := 1, \Omega := (0, 2) \subset \mathbb{R}^1$ , and  $\bar{t} := 1$ , we define functions

$$
a(t) = |t - 1|
$$
 and  $y(x) = x^2$ ,  $x \in (0, 2)$ .

Easily, we have

$$
f(x) = \mathbb{1}_{\{y \neq 1\}}(x)a'(y(x))\nabla y(x) = 2x\mathbb{1}_{\{x \neq 1\}}(x)\operatorname{sign}(x^2 - 1), \quad x \in (0, 2)
$$

and thus  $f \notin W^{1,1}(0, 2)$ .

The optimal control problem [\(P\)](#page-0-0) can be expressed in the form

<span id="page-20-0"></span>(P) 
$$
\min_{u \in \mathcal{U}_{ad}} j(u) = \int_{\Omega} L(x, S(u)(x)) dx + \frac{\nu}{2} ||u||_{L^{2}(\Omega)}^{2}
$$

with

<span id="page-20-1"></span>
$$
\mathcal{U}_{ad} := \{ u \in L^{\infty}(\Omega) \mid \alpha \le u(x) \le \beta \quad \text{for a.e. } x \in \Omega \}.
$$

Under Assumptions (A1) to (A4), the cost functional  $j: L^2(\Omega) \to \mathbb{R}$  is of class  $C^1$ . Moreover, there holds

(3.8) 
$$
j'(u)v = \int_{\Omega} (\varphi_u + vu)v \,dx \quad \text{for } u, v \in L^2(\Omega)
$$

with  $\varphi_u \in H_0^1(\Omega)$  solving [\(3.7\)](#page-19-6) corresponding to the right-hand side term v substituted by  $\frac{\partial L}{\partial y}(\cdot, S(u))$ ; see [\[9\]](#page-36-4), Thm. 4.2. We have the following first-order necessary optimality conditions from Theorem 4.3 in [\[9\]](#page-36-4) and thus derive the regularity of the optimal control as well as the corresponding state and adjoint state from [Theorems 3.2](#page-19-7) and [3.3](#page-19-9) and from Sobolev embeddings.

<span id="page-20-2"></span>Theorem 3.5 ([\[9\]](#page-36-4), Thm. 4.3). Assume that Assumptions (A1) to (A4) are satisfied. Then there exists at least one minimizer  $\bar{u}$  of [\(P\)](#page-20-0). Moreover, there exists an adjoint state  $\bar{\varphi} \in H_0^1(\Omega)$  such that for  $\bar{y} := S(\bar{u})$ ,

(3.9a) 
$$
-\operatorname{div}[(b + a(\bar{y}))\nabla \bar{y}] = \bar{u} \quad \text{in } \Omega, \quad \bar{y} = 0 \text{ on } \partial \Omega,
$$

(3.9b) 
$$
-\operatorname{div}[(b + a(\bar{y}))\nabla \bar{\varphi}] + \mathbb{1}_{\{\bar{y}\neq \bar{t}\}} a'(\bar{y})\nabla \bar{y} \cdot \nabla \bar{\varphi} = \frac{\partial L}{\partial y}(x, \bar{y}) \text{ in } \Omega, \quad \bar{\varphi} = 0 \text{ on } \partial \Omega,
$$

$$
(3.9c) \qquad \int_{\Omega} (\bar{\varphi} + v\bar{u})(u - \bar{u}) dx \ge 0 \quad \text{for all } u \in \mathcal{U}_{ad}.
$$

Furthermore,  $\bar{y} \in W^{2,p}(\Omega)$  and  $\bar{\varphi} \in W^{2,r}(\Omega)$  for any  $p, r \in (2, p_*)$  and  $r \leq \bar{p}$  with  $\bar{p}$  and  $p_*$ , respectively, defined in Assumption (A4) and [Theorem 3.2.](#page-19-7) Therefore,  $\bar{y}$  and  $\bar{\varphi}$  belong to  $C^1(\overline{\Omega})$  and  $\bar{u}$  is Lipschitz continuous on  $\overline{\Omega}$ .

Assume that  $\bar{\varphi} \in H_0^1(\Omega)$  satisfies [\(3.9\)](#page-20-1). The *critical cone* of the problem [\(P\)](#page-20-0) at  $\bar{u}$  is defined as

$$
(3.10) \qquad C(\mathcal{U}_{ad}; \bar{u}) := \{ v \in L^2(\Omega) \mid v \ge 0 \text{ if } \bar{u} = \alpha, v \le 0 \text{ if } \bar{u} = \beta, v = 0 \text{ if } \bar{\varphi} + v\bar{u} \ne 0 \text{ a.e. in } \Omega \}.
$$

In the rest of this subsection, we shall provide second-order necessary and sufficient optimality conditions for  $(P)$ . To this end, the curvature functional of *j* is first introduced and can be separated into three contributions. For any  $(u, y, \varphi) \in L^2(\Omega) \times H^1(\Omega) \times W^{1,\infty}(\Omega)$ , the smooth part and the first-order nonsmooth part of the curvature in direction  $(v_1, v_2) \in L^2(\Omega)^2$  are given by

$$
Q_s(u, y, \varphi; v_1, v_2) := \frac{1}{2} \int_{\Omega} \frac{\partial^2 L}{\partial y^2}(\cdot, y) z_{v_1} z_{v_2} dx + \frac{\nu}{2} \int_{\Omega} v_1 v_2 dx - \frac{1}{2} \int_{\Omega} \mathbb{1}_{\{y \neq \tilde{t}\}} a''(y) z_{v_1} z_{v_2} \nabla y \cdot \nabla \varphi dx,
$$
  

$$
Q_1(u, y, \varphi; v_1, v_2) := -\frac{1}{2} \int_{\Omega} [a'(y; z_{v_1}) \nabla z_{v_2} + a'(y; z_{v_2}) \nabla z_{v_1}] \cdot \nabla \varphi dx,
$$

for  $z_{v_i} := S'(u)v_i$ ,  $i = 1, 2$ , respectively. The critical part for our analysis is of course the second-order nonsmooth part, which requires some additional notation. For ease of exposition, we use the following notation in the remainder. For any  $y \in C(\overline{\Omega})$  and any  $\tau_1, \tau_2 \in \mathbb{R}$ , we define the set  $\Omega_{\mathcal{V}}^{[\tau_1, \tau_2]} := \{y \in \mathbb{R} \mid \mathcal{V} \in \mathbb{R} \}$  $[\bar{t} + \tau_1, \bar{t} + \tau_2]$ ; similar sets such as  $\Omega_{\nu}^{[\tau_1, \tau_2]}$  are defined in the same way. Let  $\delta > 0$  be arbitrary but fixed. For any  $\gamma$ ,  $\hat{\gamma} \in C(\overline{\Omega})$ , we set

<span id="page-21-0"></span>(3.11) 
$$
\begin{cases} \Omega_{y,\hat{y}}^{2} := \Omega_{\hat{y}}^{(0,\delta)} \cap \Omega_{y}^{(-\delta,0]} = {\{\hat{y} \in (\bar{t}, \bar{t} + \delta), y \in (\bar{t} - \delta, \bar{t}]\}}, \\ \Omega_{y,\hat{y}}^{3} := \Omega_{\hat{y}}^{(-\delta,0)} \cap \Omega_{y}^{[0,\delta)} = {\{\hat{y} \in (\bar{t} - \delta, \bar{t}), y \in [\bar{t}, \bar{t} + \delta)\}}. \end{cases}
$$

For any  $s \in \mathbb{R}$ ,  $u, v \in L^2(\Omega)$ ,  $y \in C(\overline{\Omega}) \cap H^1(\Omega)$ , and  $\varphi \in W^{1,\infty}(\Omega)$ , we set  $(3.12)$ 

$$
\begin{cases}\n\zeta_0(u, y; s, v) := -\{a'\}_{\bar{t}+0}^{\bar{t}-0}(\bar{t}-S(u+s v))\mathbb{1}_{\Omega_{S(u+s v), y}^3}, & \zeta_1(u, y; s, v) := \{a'\}_{\bar{t}+0}^{\bar{t}-0}(\bar{t}-S(u+s v))\mathbb{1}_{\Omega_{S(u+s v), y}^2}, \\
\zeta(u, y; s, v) := \zeta_0(u, y; s, v) + \zeta_1(u, y; s, v) = \{a'\}_{\bar{t}+0}^{\bar{t}-0}(\bar{t}-S(u+s v))\left[\mathbb{1}_{\Omega_{S(u+s v), y}^2} - \mathbb{1}_{\Omega_{S(u+s v), y}^3}\right]\n\end{cases}
$$

with  $\{a'\}_{\bar{i}+0}^{\bar{i}-0}$  $\frac{\bar{t}-0}{\bar{t}+0}$  defined as in [\(3.2\)](#page-18-3). We then define for any  $\{s_n\} \in c_0^+ := \{\{s_n\} \subset (0,\infty) \mid s_n \to 0\}$  and  $v \in L^2(\Omega)$ 

<span id="page-21-3"></span>
$$
(3.13) \quad \tilde{Q}(u, y, \varphi; \{s_n\}, v) := \liminf_{n \to \infty} \frac{1}{s_n^2} \int_{\Omega} \sum_{i=0}^1 \zeta_i(u, y; s_n, v) \nabla y \cdot \nabla \varphi \, dx
$$

$$
= \{a'\}_{\tilde{t}+0}^{\tilde{t}-0} \liminf_{n \to \infty} \frac{1}{s_n^2} \int_{\Omega} (\tilde{t} - S(u+s_n v)) \left[ \mathbb{1}_{\Omega^2_{S(u+s_n v), y}} - \mathbb{1}_{\Omega^3_{S(u+s_n v), y}} \right] \nabla y \cdot \nabla \varphi \, dx.
$$

The second-order nonsmooth part of the curvature in direction  $v \in L^2(\Omega)$  is then given by

<span id="page-21-6"></span>(3.14) 
$$
Q_2(u, y, \varphi; v) := \inf \{ \tilde{Q}(u, y, \varphi; \{s_n\}, v) \mid \{s_n\} \in c_0^+\},
$$

and finally the total curvature in direction  $v$  is

<span id="page-21-1"></span>(3.15) 
$$
Q(u, y, \varphi; v) := Q_s(u, y, \varphi; v, v) + Q_1(u, y, \varphi; v, v) + Q_2(u, y, \varphi; v).
$$

Remark 3.6. The definitions of the sets  $\Omega_{\nu,\hat{\nu}}^2$  and  $\Omega_{\nu,\hat{\nu}}^3$  in [\(3.11\)](#page-21-0) are identical to the ones for  $\Omega_{\nu}^{1,2}$  $\frac{1,2}{v,\hat{v}}$  and  $\Omega_{v,\hat{v}}^{0,3}$  $\frac{0,3}{v,\hat{v}}$  in Lemma 3.3 in [\[9\]](#page-36-4) for the case where  $K := 1$ ,  $t_0 := -\infty$ ,  $t_1 := \overline{t}$ , and  $t_2 := \infty$ . Similarly, the definitions of the functionals  $Q_s$ ,  $Q_1$ , and  $Q_2$  in this subsection can be derived from the associated ones in § 5.1 in [\[9\]](#page-36-4).

According to Proposition 5.6 and Lemma 5.7 in [\[9\]](#page-36-4),  $Q_2$  is weakly lower semicontinuous in the last variable and satisfies

<span id="page-21-5"></span>
$$
(3.16) \qquad |Q_2(u, S(u), \varphi; v)| \le \Sigma(S(u)) \|\nabla \varphi\|_{L^{\infty}(\Omega)} \|S'(u)v\|_{L^{\infty}(\Omega)}^2 \quad \text{for all } u, v \in L^2(\Omega) \text{ and } \varphi \in W^{1, \infty}(\Omega),
$$

with the jump functional

<span id="page-21-2"></span>
$$
(3.17) \qquad \qquad \Sigma(y) := \sigma_0 \limsup_{r \to 0^+} \frac{1}{r} \sum_{m=1}^2 \int_{\Omega} \bigl[ \mathbb{1}_{\{0 < |y - \bar{t}| \le r\}} \bigl| \partial_{x_m} y \bigr| \bigr] \, \mathrm{d}x, \quad y \in W^{1,1}(\Omega) \cap C(\overline{\Omega})
$$

for  $\sigma_0$  defined in [\(3.2\)](#page-18-3). Also, from Corollary 5.5 in [\[9\]](#page-36-4), it holds for any  $u \in L^2(\Omega)$ ,  $\{s_n\} \in c_0^+$  and  $v_n \to v$ in  $L^2(\Omega)$  that

<span id="page-21-4"></span>
$$
(3.18) \qquad \liminf_{n\to\infty}\frac{1}{s_n^2}\int_{\Omega}\sum_{i=0}^1\zeta_i(u,S(u);s_n,v_n)\nabla S(u)\cdot\nabla\varphi\,dx=\tilde{Q}(u,S(u),\varphi;\{s_n\},v)\geq Q_2(u,S(u),\varphi;v),
$$

provided that  $\Sigma(S(u)) < \infty$ .

We are now ready to state the second-order necessary and sufficient optimality conditions for  $(P)$  in general, for which the difference between these conditions is only in the strictness of the inequality ("no-gap").

<span id="page-22-3"></span>Theorem 3.7 (second-order necessary optimality conditions, [\[9\]](#page-36-4), Thm. 5.9). Let Assumptions (A1) to (A4) be fulfilled. Assume that  $\bar{u}$  is a local minimizer to [\(P\)](#page-20-0) such that  $\Sigma(\bar{y}) < \infty$  for  $\bar{y} := S(\bar{u})$ . Then, there is a  $\bar{\varphi} \in W_0^{1,\bar{p}}$  $\chi_0^{1,\bar p}(\Omega)\cap W^{1,\infty}(\Omega)$ , with  $\bar p$  defined in Assumption (A4), that together with  $\bar u,$   $\bar y$  satisfies [\(3.9\)](#page-20-1) and

(3.19) 
$$
Q(\bar{u}, \bar{y}, \bar{\varphi}; v) \ge 0
$$
 for all  $v \in C(\mathcal{U}_{ad}; \bar{u})$ 

with  $Q$  defined in  $(3.15)$ .

<span id="page-22-4"></span>Theorem 3.8 (second-order sufficient optimality conditions, [\[9\]](#page-36-4), Thm. 5.10). Let Assumptions (A1) to (A4) be valid. Assume that  $\bar{u}$  is a feasible point of [\(P\)](#page-20-0) such that  $\Sigma(\bar{v}) < \infty$  for  $\bar{v} := S(\bar{u})$ . Assume further that there is  $a \overline{\varphi} \in W_0^{1,\overline{p}}$  $U_0^{1,\bar{p}}(\Omega)\cap W^{1,\infty}(\Omega)$ , with  $\bar{p}$  defined in Assumption (A4), that together with  $\bar{u},\,\bar{y}$  satisfies [\(3.9\)](#page-20-1) and

$$
(3.20) \tQ(\bar{u}, \bar{y}, \bar{\varphi}; v) > 0 \tfor all  $v \in C(\mathcal{U}_{ad}; \bar{u}) \setminus \{0\}$
$$

with Q defined in [\(3.15\)](#page-21-1). Then there exist constants  $c_0$ ,  $\rho_0 > 0$  satisfying

$$
j(\bar{u})+c_0||u-\bar{u}||^2_{L^2(\Omega)}\leq j(u) \qquad \text{for all } u\in \mathcal{U}_{ad}\cap \overline{B}_{L^2(\Omega)}(\bar{u},\rho_0).
$$

It is noted that the term  $2Q(\bar{u}, \bar{y}, \bar{\varphi}; v)$  can be seen as a second-order generalized derivative of *j* at  $\bar{u}$ in the direction v; see, e.g. the proof of Theorem 5.9 in [\[9\]](#page-36-4) and Remark 5.1 in [\[19\]](#page-37-11)).

# <span id="page-22-0"></span>3.2 an explicit formula for the curvature functional

In this subsection, we assume that the domain  $\Omega$  is open, bounded, and convex in ℝ<sup>2</sup> only. We shall establish an explicit formula for the curvature term  $Q(u, y, \varphi; v)$ , defined in [\(3.15\)](#page-21-1), for two situations:

- (i) The gradient of y does not vanish on connected components of  $\{y = \overline{t}\}$ ;
- (ii) The gradient of y vanishes on connected components of  $\{y = \bar{t}\}.$

For the first situation, we will apply the results shown in [Section 2.](#page-3-4) For the latter situation, we need the following notion.

Definition 3.9. A function  $y : \overline{\Omega} \to \mathbb{R}$  is called *uniformly locally convex-concave* on a set  $V \subset \overline{\Omega}$  if an  $\varepsilon > 0$  exists such that for any  $x \in V$ ,  $\gamma$  is either convex or concave on  $B_{\mathbb{R}^2}(x, \varepsilon) \cap \overline{\Omega}$ .

<span id="page-22-1"></span>Proposition 3.10. Let  $t \in \mathbb{R}$  be arbitrary and let  $y \in C^1(\overline{\Omega})$  be uniformly locally convex-concave on the level set  $\{v = t\}$ . Assume that C is a connected component of  $\{y = t\}$ . If  $\nabla y$  vanishes at some point  $x_0 \in C$ , then  $\nabla y(x) = 0$  for all  $x \in C$ .

*Proof.* There is an  $\varepsilon > 0$  such that, for any  $x \in \{y = t\}$ , the restriction  $y \mid_{B_{\mathbb{R}^2}(x,\varepsilon) \cap \overline{\Omega}}$  is either convex or concave. Since  $\nabla y(x_0) = 0$ , then  $x_0$  is a local extremal point of  $y(x)$  and so is every point in  $B_{\mathbb{R}^2}(x, \varepsilon) \cap C$ . From this and the connection property of C, we have  $\nabla y = 0$  on C.

The following result is a direct consequence of [Proposition 3.10.](#page-22-1)

Corollary 3.11. Let  $t \in \mathbb{R}$  be arbitrary and let  $y \in C^1(\overline{\Omega})$  be uniformly locally convex-concave on the level set  $\{y = t\}$ . Assume that C is a connected component of  $\{y = t\}$ . If  $\nabla y(x_0) \neq 0$  for some point  $x_0 \in C$ , then  $\nabla y(x) \neq 0$  for all  $x \in C$ .

<span id="page-22-2"></span>The following result will play an important role in establishing an explicit formula of the curvature functional [\(3.15\)](#page-21-1). Its proof rests on the following several lemmas.

Theorem 3.12. Let  $\{s_n\} \in c_0^+, \bar{\varphi} \in C^1(\overline{\Omega}) \cap W^{2,1}(\Omega)$  and  $\bar{y}, y_n \in C^1(\overline{\Omega})$  such that  $y_n = \bar{y} = 0$  on  $\partial\Omega, y_n \to \bar{y}$ in  $C^1(\overline{\Omega})$  and  $(y_n - \overline{y})/s_n \to w$  in  $W_0^{1,p}$  $\sigma_0^{1,p}(\Omega)$  for some  $p > 2$  and  $w \in W_0^{1,p}$  $\sigma_0^{1,p}(\Omega)$ . Let  $C$  be a closed connected component of  $\{\bar{y} = \bar{t}\}\$ . Assume that one of the following conditions is satisfied:

<span id="page-23-2"></span>(C1) The gradient of  $\bar{v}$  does not vanish on C, i.e.,

<span id="page-23-11"></span>(3.21) 
$$
\min\{|\nabla \bar{y}(x)| : x \in C\} > 0;
$$

<span id="page-23-5"></span>(C2) There holds that

<span id="page-23-0"></span>(3.22) 
$$
\begin{cases} \bar{y} \text{ is uniformly locally convex-concave on } C, \\ \nabla \bar{y} = 0 \text{ on } C, \\ \mathcal{H}^{N-1}(\{\bar{y} = t\} \cap C^{\tilde{\varepsilon}}) \leq C_0, \quad f.a.a. \ t \in (\bar{t} - r_0, \bar{t} + r_0), N = 2, \end{cases}
$$

for some constants  $\tilde{\varepsilon}$ ,  $r_0, C_0 > 0$ ;

<span id="page-23-7"></span> $(C_3)$  There holds that

<span id="page-23-8"></span>(3.23) 
$$
\begin{cases} \nabla \bar{y} = 0 \text{ on } C, \\ \operatorname{meas}_{\mathbb{R}^2} \{ \{ 0 < |\bar{y} - \bar{t}| < r \} \cap C^{\tilde{e}} \} \le c_s r, \quad \text{for all } r \in (0, r_0), \end{cases}
$$

for some constants  $\tilde{\varepsilon}$ ,  $r_0, c_s > 0$ .

Then there exists an  $\varepsilon_0 = \varepsilon_0(C) \in (0, \tilde{\varepsilon})$  such that for any  $\varepsilon \in (0, \varepsilon_0)$ ,

$$
(3.24) \qquad \frac{A_n(C,\varepsilon)}{s_n^2} \to \frac{1}{2} \int_C \mathbb{1}_{\{|\nabla \bar{y}|>0\}} w^2 \frac{\nabla \bar{y} \cdot \nabla \bar{\varphi}}{|\nabla \bar{y}|} d\mathcal{H}^1(x) \quad \text{as} \quad n \to \infty,
$$

where

<span id="page-23-1"></span>(3.25) 
$$
A_n(C,\varepsilon) := \int_{C^{\varepsilon}} (\bar{t} - y_n) [\mathbb{1}_{\Omega^2_{y_n,\bar{y}}} - \mathbb{1}_{\Omega^3_{y_n,\bar{y}}}]\nabla \bar{y} \cdot \nabla \bar{\phi} dx
$$

with  $\Omega^2_{y_n,\bar{y}}$  and  $\Omega^3_{y_n,\bar{y}}$  defined in [\(3.11\)](#page-21-0).

Remark 3.13. Note that [\(3.22\)](#page-23-0) does not require that the level set  $\{\bar{y} = \bar{t}\}\$  has measure zero. When  $N = 1$ , the last condition in [\(3.22\)](#page-23-0) means that for a.e. *t* in a neighborhood of  $\bar{t}$ , the level sets { $\bar{y} = t$ } consist of finitely many points; in other words, the function  $\bar{y}$  oscillates around the values  $\bar{t}$  only finitely many times. In [\[9\]](#page-36-4), Exam. 5.3, this condition was shown to be equivalent to the finiteness of the jump functional  $\Sigma(\bar{y})$  introduced in [\(3.17\)](#page-21-2) for the case  $N = 1$ .

Remark 3.14. According to the definition of the sets  $\Omega^j_{\nu,\hat{\nu}}$  with  $j=2,3$  in [\(3.11\)](#page-21-0), the sets  $\Omega^2_{\gamma_n,\tilde{\nu}}$  and  $\Omega^3_{\gamma_n,\tilde{\nu}}$  can be expressed as

<span id="page-23-9"></span>
$$
(3.26) \qquad \Omega^2_{y_n,\bar{y}} = \tilde{\Omega}^2_{y_n,\bar{y}} \cup \{\bar{y} \in (\bar{t}, \bar{t} + \delta), y_n = \bar{t}\} \quad \text{and} \quad \Omega^3_{y_n,\bar{y}} = \tilde{\Omega}^3_{y_n,\bar{y}} \cup \{\bar{y} \in (\bar{t} - \delta, \bar{t}), y_n = \bar{t}\}
$$

with

<span id="page-23-3"></span>(3.27)  
\n
$$
\begin{cases}\n\tilde{\Omega}_{y,\hat{y}}^2 := \{\hat{y} \in (\bar{t}, \bar{t} + \delta), y \in (\bar{t} - \delta, \bar{t})\}, \\
\tilde{\Omega}_{y,\hat{y}}^3 := \{\hat{y} \in (\bar{t} - \delta, \bar{t}), y \in (\bar{t}, \bar{t} + \delta)\}\n\end{cases}
$$

for  $\gamma$ ,  $\hat{\gamma} \in C(\overline{\Omega})$ . Obviously, one has

<span id="page-23-10"></span>
$$
\tilde{\Omega}^2_{y,\hat{y}} = \tilde{\Omega}^3_{\hat{y},y}.
$$

Moreover, thanks to [\(3.25\)](#page-23-1), the term  $A_n(C, \varepsilon)$  can be rewritten as

<span id="page-23-4"></span>(3.29) 
$$
A_n(C,\varepsilon):=\int_{C^{\varepsilon}}(\overline{t}-y_n)[\mathbb{1}_{\tilde{\Omega}^2_{y_n,\tilde{y}}}-\mathbb{1}_{\tilde{\Omega}^3_{y_n,\tilde{y}}}] \nabla \bar{y}\cdot \nabla \bar{\varphi}\,dx.
$$

<span id="page-23-6"></span>Clason, Nhu, Rösch Numerical analysis of a nonsmooth quasilinear elliptic ...

Lemma 3.15. Under assumption [\(C1\)](#page-23-2) in [Theorem 3.12,](#page-22-2) then there exists an  $\varepsilon_0 = \varepsilon_0(C) > 0$  such that

<span id="page-24-1"></span>
$$
(3.30) \qquad \frac{A_n(C,\varepsilon)}{s_n^2} \to \frac{1}{2} \int_C w^2 \frac{\nabla \bar{y} \cdot \nabla \bar{\varphi}}{|\nabla \bar{y}|} d\mathcal{H}^1(x) \quad \text{as} \quad n \to \infty \quad \text{for all } \varepsilon \in (0,\varepsilon_0).
$$

*Proof.* By [Proposition 2.4,](#page-7-0)  $C$  is a closed  $C^1$  simple curve in  $\overline{\Omega}$ . Thanks to [Proposition 2.10,](#page-13-4) there are constants  $\varepsilon_1$ ,  $\alpha > 0$  such that, for *n* large enough,

(3.31) 
$$
\nabla y_n(x) \cdot \nabla \bar{y}(x') \ge \alpha > 0 \quad \text{for all} \quad x, x' \in C^{\varepsilon_1} \text{ with } |x - x'| \le 2\varepsilon_1.
$$

Since C is a closed component in  $\{\bar{y} = \bar{t}\}$ , there holds  $C^{\epsilon_2} \cap (\{\bar{y} = \bar{t}\} \setminus C)^{\epsilon_2} = \emptyset$  for some constant  $\varepsilon_2 > 0$ . Moreover, the sign of  $(\bar{y} - \bar{t})$  in  $C_{\varepsilon_2}^-$  is opposite to the one in  $C_{\varepsilon_2}^+$ . Without loss of generality, we can thus assume that

<span id="page-24-0"></span>(3.32) 
$$
\bar{y} < \bar{t} \quad \text{on } C_{\varepsilon_2}^- \quad \text{and} \quad \bar{y} > \bar{t} \quad \text{on } C_{\varepsilon_2}^+.
$$

Set  $\varepsilon_0 := \min\{\varepsilon_1, \varepsilon_2\}, \tau_n := ||y_n - \bar{y}||_{C(\overline{\Omega})}$ , fix any  $\varepsilon \in (0, \varepsilon_0)$ , and define the sets (depending on  $\varepsilon$ )

(3.33) 
$$
\Omega_n^2 := \tilde{\Omega}_{y_n, \tilde{y}}^2 \cap C^{\varepsilon} \text{ and } \Omega_n^3 := \tilde{\Omega}_{y_n, \tilde{y}}^3 \cap C^{\varepsilon}.
$$

<span id="page-24-3"></span>Here the sets  $\tilde{\Omega}^2_{\gamma_n,\tilde\gamma}$  and  $\tilde{\Omega}^3_{\gamma_n,\tilde\gamma}$  are defined in [\(3.27\)](#page-23-3). Obviously, for  $n$  large enough such that  $\tau_n<\delta/2,$ we have

(3.34a)  $\Omega_n^2 = \{ \bar{y} \in (\bar{t}, \bar{t} + \delta), y_n \in (\bar{t} - \delta, \bar{t}) \} \cap C^{\varepsilon} = \{ y_n < \bar{t} < \bar{y} \} \cap C^{\varepsilon} \subset C_{\varepsilon}^+$ 

$$
(3.34b) \qquad \Omega_n^3 = \{ \bar{y} \in (\bar{t} - \delta, \bar{t}), y_n \in (\bar{t}, \bar{t} + \delta) \} \cap C^{\varepsilon} = \{ \bar{y} < \bar{t} < y_n \} \cap C^{\varepsilon} \subset C_{\varepsilon}^-.
$$

As a result of [Proposition 2.11,](#page-14-0) for *n* large enough,  $C^{\varepsilon}$  contains one and only one connected component  $C_n$  of  $\{y_n = \overline{t}\}$ . By [Proposition 2.4,](#page-7-0) the set  $C_n = \{y_n = \overline{t}\} \cap C^{\varepsilon}$  is a closed  $C^1$  simple curve in  $\mathbb{R}^2$ . We now consider two cases.

*Case 1: C* =  $\partial\Omega$ . In this case, we have  $\bar{t} = 0$  and thus  $C_n = \partial\Omega$  for sufficiently large *n*. Moreover, one has  $C_{\varepsilon}^+ = \emptyset$ ,  $C^{\varepsilon} = C \cup C_{\varepsilon}^-$  and then  $\Omega_n^2 = \emptyset$  for *n* large enough. On the other hand, since  $C_n = \{y_n = \overline{t}\} \cap C^{\varepsilon}$ is a closed simple curve in  $\mathbb{R}^2$  for *n* large enough and  $y_n \to \bar{y}$  in  $C(\overline{\Omega})$ , we deduce from [\(3.32\)](#page-24-0) that  $y_n \leq \overline{t}$  on  $C^{\varepsilon}$  and therefore  $\Omega_n^3 = \emptyset$  for sufficiently large *n*. From this and the definition of  $A_n(C, \varepsilon)$ , we have [\(3.30\)](#page-24-1) because of the vanishing on  $\partial\Omega$  of w.

Case 2:  $C \neq \partial \Omega$ . In this case, both  $C_{\varepsilon}^+$  and  $C_{\varepsilon}^-$  are nonempty. To estimate  $A_n(C, \varepsilon)$ , we use the expression [\(3.29\)](#page-23-4) of  $A_n(C, \varepsilon)$  and split it into two terms as follows:

<span id="page-24-2"></span>
$$
(3.35) \quad A_n(C, \varepsilon) := \int_{C^{\varepsilon}} (\bar{t} - y_n) [\mathbb{1}_{\tilde{\Omega}^2_{y_n, \tilde{y}}} - \mathbb{1}_{\tilde{\Omega}^3_{y_n, \tilde{y}}} ] \nabla (\bar{y} - y_n) \cdot \nabla \bar{\varphi} \, dx + \int_{C^{\varepsilon}} (\bar{t} - y_n) [\mathbb{1}_{\tilde{\Omega}^{i, 2}_{y_n, \tilde{y}}} - \mathbb{1}_{\tilde{\Omega}^3_{y_n, \tilde{y}}} ] \nabla y_n \cdot \nabla \bar{\varphi} \, dx = B_n + C_n
$$

with

<span id="page-24-4"></span>
$$
(3.36)
$$
  
\n
$$
B_n := \int_{C^{\varepsilon}} (\bar{t} - y_n) [\mathbb{1}_{\tilde{\Omega}^2_{y_n, \tilde{y}}} - \mathbb{1}_{\tilde{\Omega}^3_{y_n, \tilde{y}}}] \nabla (\bar{y} - y_n) \cdot \nabla \bar{\varphi} \, dx \quad \text{and} \quad C_n := \int_{C^{\varepsilon}} (\bar{t} - y_n) [\mathbb{1}_{\tilde{\Omega}^2_{y_n, \tilde{y}}} - \mathbb{1}_{\tilde{\Omega}^3_{y_n, \tilde{y}}}] \nabla y_n \cdot \nabla \bar{\varphi} \, dx.
$$

Moreover, we deduce from the definition of sets  $\Omega_n^2$  and  $\Omega_n^3$  that

$$
B_n = \int_{C^{\varepsilon}} (\bar{t} - y_n) [\mathbb{1}_{\Omega_n^2} - \mathbb{1}_{\Omega_n^3}] \nabla (\bar{y} - y_n) \cdot \nabla \bar{\varphi} \, \mathrm{d}x \quad \text{and} \quad C_n = \int_{C^{\varepsilon}} (\bar{t} - y_n) [\mathbb{1}_{\Omega_n^2} - \mathbb{1}_{\Omega_n^3}] \nabla y_n \cdot \nabla \bar{\varphi} \, \mathrm{d}x.
$$

Clason, Nhu, Rösch Numerical analysis of a nonsmooth quasilinear elliptic ...

From the facts

<span id="page-25-0"></span>(3.37) 
$$
\begin{cases} |\bar{t} - \bar{y}|, |\bar{t} - y_n| \leq |\bar{y} - y_n| & \text{on } \Omega_n^2 \cup \Omega_n^3, \\ \Omega_n^2 \cup \Omega_n^3 \subset \{0 < |\bar{y} - \bar{t}| \leq \tau_n\}, \end{cases}
$$

we deduce from Hölder's inequality for  $p' := \frac{p}{n}$  $\frac{p}{p-1}$  that

<span id="page-25-6"></span>
$$
(3.38) \t |B_n| \leq ||y_n - \bar{y}||_{C(\overline{\Omega})} ||\nabla (y_n - \bar{y})||_{L^p(\Omega)} ||\nabla \bar{\varphi} 1_{\{0 < |\bar{y} - \bar{t}| \leq \tau_n\}}||_{L^{p'}(\Omega)} = o(s_n^2).
$$

From this and [\(3.35\)](#page-24-2), there holds

<span id="page-25-3"></span>(3.39) 
$$
A_n(C, \varepsilon) = o(s_n^2) + C_n.
$$

We now estimate  $C_n$ . To this end, we shall employ [Proposition 2.9.](#page-12-0) We first rewrite  $C_n$  as

<span id="page-25-4"></span>(3.40) 
$$
C_n = -\frac{1}{2} \int_{C^{\varepsilon}} [\mathbb{1}_{\Omega_n^2} - \mathbb{1}_{\Omega_n^3}] \nabla (y_n - \overline{t})^2 \cdot \nabla \overline{\varphi} dx.
$$

From [\(3.34\)](#page-24-3) and the limit  $y_n \to \bar{y}$  in  $C^1(\overline{\Omega})$  that, for  $n$  large enough, there holds

<span id="page-25-7"></span>
$$
\partial\Omega_n^2 \cup \partial\Omega_n^3 \subset C \cup C_n.
$$

Applying [Proposition 2.9](#page-12-0) for functions  $y_1 := \bar{y}$ ,  $y_2 := y_n$  and  $v := (y_n - \bar{t})^2$ ,  $\varphi := \bar{\varphi}$ , and open sets  $S^+ := \Omega_n^2$ ,  $S^- := \Omega_n^3$  (see [\(3.34\)](#page-24-3) for the definition of the sets  $\Omega_n^2$ ,  $\Omega_n^3$ ) yields

$$
C_n = \frac{1}{2} \int_{C^{\varepsilon}} [\mathbb{1}_{\Omega_n^2} - \mathbb{1}_{\Omega_n^3}](y_n - \bar{t})^2 \Delta \bar{\varphi} dx + \frac{1}{2} \int_C (y_n - \bar{t})^2 \nabla \bar{\varphi} \cdot \frac{\nabla \bar{y}}{|\nabla \bar{y}|} d\mathcal{H}^1(x) - \frac{1}{2} \int_{C_n} (y_n - \bar{t})^2 \nabla \bar{\varphi} \cdot \frac{\nabla y_n}{|\nabla y_n|} d\mathcal{H}^1(x),
$$

or equivalently,

<span id="page-25-1"></span>
$$
(3.42) \tC_n = \frac{1}{2} \int_{C^{\varepsilon}} [\mathbb{1}_{\Omega_n^2} - \mathbb{1}_{\Omega_n^3}](y_n - \bar{t})^2 \Delta \bar{\varphi} dx + \frac{1}{2} \int_C (y_n - \bar{y})^2 \nabla \bar{\varphi} \cdot \frac{\nabla \bar{y}}{|\nabla \bar{y}|} d\mathcal{H}^1(x).
$$

In view of  $(3.37)$ , there then holds

<span id="page-25-8"></span>
$$
(3.43) \qquad \left| \int_{C^{\varepsilon}} [\mathbb{1}_{\Omega_n^2} - \mathbb{1}_{\Omega_n^3}](y_n - \overline{t})^2 \Delta \overline{\varphi} dx \right| \leq ||y_n - \overline{y}||_{C(\overline{\Omega})}^2 \int_{\{0 < |\overline{y} - \overline{t}| \leq \tau_n\}} |\Delta \overline{\varphi}| dx = o(s_n^2),
$$

as a result of Lebesgue's Dominated Convergence Theorem. Combining this with [\(3.42\)](#page-25-1) gives

<span id="page-25-2"></span>(3.44) 
$$
C_n = o(s_n^2) + \frac{1}{2} \int_C (y_n - \bar{y})^2 \nabla \bar{\varphi} \cdot \frac{\nabla \bar{y}}{|\nabla \bar{y}|} d\mathcal{H}^1(x).
$$

The combination of [\(3.44\)](#page-25-2) with [\(3.39\)](#page-25-3) and the limits  $(y_n - \bar{y})/s_n \to w$  in  $W_0^{1,p}$  $U_0^{1,p}(\Omega)$  yields [\(3.30\)](#page-24-1). □

Lemma 3.16. Under assumption [\(C2\),](#page-23-5) there is an  $\varepsilon_0 = \varepsilon_0(C) > 0$  such that

<span id="page-25-5"></span>(3.45) 
$$
\frac{A_n(C,\varepsilon)}{s_n^2} \to 0 \quad as \quad n \to \infty \quad \text{for all } \varepsilon \in (0,\varepsilon_0).
$$

 $\overline{C}$ lason, Nhu, Rösch  $\overline{C}$ lason, Nhu, Rösch  $\overline{C}$ lason, Nhu, Rösch  $\overline{C}$ lason, Nhu, Rösch  $\overline{C}$ 



<span id="page-26-0"></span>Figure 4: a closed component C of the level set  $\{\bar{y} = \bar{t}\}$  with positive measure meas<sub>ℝ</sub>2(C)

*Proof.* Without loss of generality, assume that there thus exists an  $\varepsilon_3 > 0$  such that  $\bar{\nu}$  is convex on  $B(x, \varepsilon_3) \cap \overline{\Omega}$  for all  $x \in C$ ; see [Figure 4.](#page-26-0) We first set  $\varepsilon_0 := \min{\{\tilde{\varepsilon}, \varepsilon_2, \varepsilon_3\}}$  with constants  $\tilde{\varepsilon}$  given in [\(3.22\)](#page-23-0) and  $\varepsilon_2$  defined as in the proof of [Lemma 3.15.](#page-23-6) Let us take  $\varepsilon \in (0, \varepsilon_0)$  arbitrarily but fixed and reuse all symbols defined in the proof of [Lemma 3.15.](#page-23-6) Moreover, the relations [\(3.34\)](#page-24-3)–[\(3.40\)](#page-25-4) are still valid. The convexity of  $\bar{y}$  and the fact that  $\nabla \bar{y} = 0$  on C imply that  $\bar{y} > \bar{t}$  on  $C^{\varepsilon} \backslash C$ . This and [\(3.34\)](#page-24-3) guarantee that

<span id="page-26-3"></span> $(3.46)$  $\Omega_n^2 = \{ y_n < \overline{t} \} \cap (C^{\varepsilon} \backslash C)$  and  $\Omega_n^3 = \emptyset$  for all  $n \ge n_0$  for some integer  $n_0 \in \mathbb{N}$ 

and thus

(3.47) 
$$
C_n = \int_{\Omega_n^2} (\bar{t} - y_n) \nabla y_n \cdot \nabla \bar{\varphi} \, dx.
$$

We now split the sequence  ${n}$  into subsequences, also denoted by  ${n}$ , that satisfy one of the following conditions:

<span id="page-26-2"></span><span id="page-26-1"></span>(a)  $\Omega_n^2 \neq \emptyset$  for all  $n \geq n_0$ ; (b)  $\Omega$  $n_n^2 = \emptyset$  for all  $n \geq n_0$ .

For [\(b\),](#page-26-1) we have from [\(3.35\)](#page-24-2) that  $A_n(C, \varepsilon) = 0$  for all  $n \ge n_0$ . The limit [\(3.45\)](#page-25-5) thus follows. It remains to consider [\(a\).](#page-26-2) To this end, by the Morse–Sard Theorem in Sobolev spaces (see; e.g. Theorem 5 in [\[13\]](#page-36-11) and Corollary 5.2 in [\[3\]](#page-36-7)), for a.e.  $t \in \mathbb{R}$ , the level set  $\{y_n = t\}$  is a finite disjoint family of  $C^1$  simple curves in  $\mathbb{R}^2$  and  $\nabla y_n$  does not vanish on  $\{y_n = t\}$ . Therefore, for any  $n \ge n_0$ , there exists a  $\bar{t}_n \in \mathbb{R}$ such that

<span id="page-26-4"></span>(3.48)  
\n
$$
\begin{cases}\n\bar{t} \geq \bar{t}_n \neq 0, \\
\bar{t} - \bar{t}_n = o(s_n^2), \\
\{y_n = \bar{t}_n\} \text{ consists of finite disjoint closed } C^1 \text{ simple curves,} \\
|\nabla y_n(x)| > 0 \quad \text{for all } x \in \{y_n = \bar{t}_n\}.\n\end{cases}
$$

Note that any closed curve in  $\{y_n = \bar{t}_n\}$  does not intersect with the boundary  $\partial\Omega$  since  $\bar{t}_n \neq 0$ . From the expression of  $C_n$  in [\(3.40\)](#page-25-4) and the identities in [\(3.46\)](#page-26-3), one has

$$
C_n = \int_{\Omega_n^2} (\bar{t} - y_n) \nabla y_n \cdot \nabla \bar{\varphi} \, dx = \int_{\{\bar{t}_n \le y_n \le \bar{t}\} \cap C^{\varepsilon}} (\bar{t} - y_n) \nabla y_n \cdot \nabla \bar{\varphi} \, dx + \int_{\{y_n < \bar{t}_n\} \cap C^{\varepsilon}} (\bar{t} - y_n) \nabla y_n \cdot \nabla \bar{\varphi} \, dx =: D_n^1 + D_n^2.
$$

Obviously, one has from the choice of  $\bar{t}_n$  in [\(3.48\)](#page-26-4) and the fact  $||y_n - \bar{y}||_{W_0^{1,p}(\Omega)} = O(s_n)$  that

$$
\begin{split} |D_n^1| &\leq \int_{\mathbb{R}^2} \mathbb{1}_{\{\bar{t}_n \leq y_n \leq \bar{t}\} \cap C^{\varepsilon}| \bar{t} - y_n| |\nabla y_n| |\nabla \bar{\varphi}| \, \mathrm{d}x \\ &\leq \int_{\mathbb{R}^2} \mathbb{1}_{\{\bar{t}_n \leq y_n \leq \bar{t}\} \cap C^{\varepsilon}| \bar{t} - y_n| |\nabla (y_n - \bar{y})| |\nabla \bar{\varphi}| \, \mathrm{d}x + \int_{\mathbb{R}^2} \mathbb{1}_{\{\bar{t}_n \leq y_n \leq \bar{t}\} \cap C^{\varepsilon}| \bar{t} - y_n| |\nabla \bar{y}| |\nabla \bar{\varphi}| \, \mathrm{d}x \\ &\leq |\bar{t} - \bar{t}_n| \int_{\mathbb{R}^2} \mathbb{1}_{\{\bar{t}_n \leq y_n \leq \bar{t}\} \cap C^{\varepsilon}| \nabla (y_n - \bar{y})| |\nabla \bar{\varphi}| \, \mathrm{d}x + \int_{\mathbb{R}^2} \mathbb{1}_{\{\bar{t}_n \leq y_n \leq \bar{t}\} \cap C^{\varepsilon}| \bar{t} - y_n| |\nabla \bar{y}| |\nabla \bar{\varphi}| \, \mathrm{d}x \\ &= o(s_n^2) + \int_{\mathbb{R}^2} \mathbb{1}_{\{\bar{t}_n \leq y_n \leq \bar{t}\} \cap C^{\varepsilon}| \bar{t} - y_n| |\nabla \bar{y}| |\nabla \bar{\varphi}| \, \mathrm{d}x. \end{split}
$$

From the inclusion  $\{\bar{t}_n \leq y_n \leq \bar{t}\} \subset \{\bar{t}_n - \tau_n \leq \bar{y} \leq \bar{t} + \tau_n\}$  and the coarea formula for Lipschitz mappings; see, e.g. [\[12\]](#page-36-9), Thm. 2, p. 117 and [\[1\]](#page-36-6), Sec. 2.7, we have

$$
|D_n^1| \le o(s_n^2) + \int_{\mathbb{R}} \left[ \int_{\{\bar{y}=t\}} \mathbb{1}_{\{\bar{t}_n \le y_n \le \bar{t}\} \cap C^{\varepsilon}} |\bar{t} - y_n| |\nabla \bar{\varphi}| d\mathcal{H}^1(x) \right] dt
$$
  
\n
$$
= o(s_n^2) + \int_{\bar{t}_n - \tau_n}^{\bar{t} + \tau_n} \left[ \int_{\{\bar{y}=t\}} \mathbb{1}_{\{\bar{t}_n \le y_n \le \bar{t}\} \cap C^{\varepsilon}} |\bar{t} - y_n| |\nabla \bar{\varphi}| d\mathcal{H}^1(x) \right] dt
$$
  
\n
$$
\le o(s_n^2) + C_0(\bar{t} - \bar{t}_n + 2\tau_n)(\bar{t} - \bar{t}_n) \|\nabla \bar{\varphi}\|_{L^{\infty}(\Omega)} = o(s_n^2),
$$

due to the choice of  $\bar{t}_n$  in [\(3.48\)](#page-26-4) and conditions in [\(3.22\)](#page-23-0). For  $D_n^2$ , we see from (3.48) that  $\{y_n = \bar{t}_n\} \cap C^{\varepsilon} \neq \emptyset$ for *n* large enough. If  $\{y_n < \bar{t}_n\} \cap C^{\varepsilon} = \emptyset$ , then  $D_n^2 = 0$ . Otherwise, let  $\Gamma_n$  be the boundary of  $\{y_n < \overline{t}_n\} \cap C^{\varepsilon}$ . There are two possibilities in principle: either an infinite subsequence  $\{k\}$  of  $\{n\}$  exists and satisfies  $\Gamma_k \cap \partial C^{\varepsilon} \neq \emptyset$ , or there is no such an subsequence. Let us see that the first possibility is not actually a correct assumption. Indeed, if  $\Gamma_k \cap \partial C^{\varepsilon} \neq \emptyset$ , then  $||y_k - \bar{y}||_{C(\overline{\Omega})} \ge \bar{y}(x) - \bar{t} > 0$  for all  $x \in \partial C^{\varepsilon}$ . This contradicts the limit  $||y_k - \bar{y}||_{C(\overline{\Omega})} \to 0$  as  $k \to \infty$ . Therefore, the second possibility always holds. It then must be true that  $\Gamma_n \cap \partial C^{\varepsilon} = \emptyset$  and so  $\Gamma_n = \{y_n = \overline{t}_n\} \cap C^{\varepsilon}$  for *n* large enough. Combining this with the last two conditions in [\(3.48\)](#page-26-4) and the Implicit Function Theorem, we deduce that the open set  $\{y_n < \bar{t}_n\} \cap C^{\varepsilon}$  decomposes into subdomains with Lipschitz boundaries. By rewriting  $D_n^2$  and then using integration by parts, we have

$$
D_n^2 = \int_{\{y_n < \bar{t}_n\} \cap C^{\epsilon}} (\bar{t} - y_n) \nabla y_n \cdot \nabla \bar{\varphi} \, dx
$$
\n
$$
= \int_{\{y_n < \bar{t}_n\} \cap C^{\epsilon}} (\bar{t} - \bar{t}_n) \nabla y_n \cdot \nabla \bar{\varphi} \, dx + \int_{\{y_n < \bar{t}_n\} \cap C^{\epsilon}} (\bar{t}_n - y_n) \nabla y_n \cdot \nabla \bar{\varphi} \, dx
$$
\n
$$
= \int_{\{y_n < \bar{t}_n\} \cap C^{\epsilon}} (\bar{t} - \bar{t}_n) \nabla y_n \cdot \nabla \bar{\varphi} \, dx - \frac{1}{2} \int_{\{y_n < \bar{t}_n\} \cap C^{\epsilon}} \nabla (y_n - \bar{t}_n)^2 \cdot \nabla \bar{\varphi} \, dx
$$
\n
$$
= \int_{\{y_n < \bar{t}_n\} \cap C^{\epsilon}} (\bar{t} - \bar{t}_n) \nabla y_n \cdot \nabla \bar{\varphi} \, dx + \frac{1}{2} \int_{\{y_n < \bar{t}_n\} \cap C^{\epsilon}} (y_n - \bar{t}_n)^2 \Delta \bar{\varphi} \, dx - \frac{1}{2} \int_{\Gamma_n} (y_n - \bar{t}_n)^2 \nabla \bar{\varphi} \cdot v_n \, d\mathcal{H}^1(x)
$$
\n
$$
= \int_{\{y_n < \bar{t}_n\} \cap C^{\epsilon}} (\bar{t} - \bar{t}_n) \nabla y_n \cdot \nabla \bar{\varphi} \, dx + \frac{1}{2} \int_{\{y_n < \bar{t}_n\} \cap C^{\epsilon}} (y_n - \bar{t}_n)^2 \Delta \bar{\varphi} \, dx,
$$

where  $v_n$  stands for the outward unit normal vector to  $\Gamma_n$ . From this, the first two conditions in [\(3.48\)](#page-26-4), and the fact that  $\bar{y}(x) \geq \bar{t}$  for all  $x \in C^{\varepsilon}$ , we have

$$
\begin{split} |D_n^2| &\leq (\bar{t}-\bar{t}_n)\|\nabla y_n\|_{L^\infty(\Omega)} \|\nabla \bar{\varphi}\|_{L^\infty(\Omega)} \|\mathbb{1}_{\{y_n < \bar{t}_n\}\cap C^\varepsilon}\|_{L^1(\Omega)} + \frac{1}{2} \int_{\{y_n < \bar{t}_n\}\cap C^\varepsilon} (y_n - \bar{y})^2 |\Delta \bar{\varphi}| \, \mathrm{d}x \\ &= o(s_n^2) + \frac{1}{2} \|y_n - \bar{y}\|_{C(\overline{\Omega})}^2 \int_{\{y_n < \bar{t}_n\}\cap C^\varepsilon} |\Delta \bar{\varphi}| \, \mathrm{d}x \\ &= o(s_n^2) + O(s_n^2) \int_{\Omega} \mathbb{1}_{\{y_n < \bar{t}_n\}\cap C^\varepsilon} |\Delta \bar{\varphi}| \, \mathrm{d}x. \end{split}
$$

Since  $\bar{\varphi} \in W^{2,1}(\Omega)$  and  $0 \leq \mathbb{1}_{\{\gamma_n < \bar{t}_n\} \cap C^{\varepsilon}} \leq \mathbb{1}_{\{0 < |\gamma_n - \bar{\gamma}| \leq \tau_n\} \cap C^{\varepsilon}} \to 0$  a.e. in  $\Omega$  as  $n \to \infty$ , we deduce from Lebesgue's Dominated Convergence Theorem that

$$
\int_{\Omega} 1_{\{y_n < \bar{t}_n\} \cap C^{\varepsilon}} |\Delta \bar{\varphi}| \, \mathrm{d} x \to 0 \quad \text{as } n \to \infty.
$$

<span id="page-27-0"></span>We thus have  $D_n^2 = o(s_n^2)$ . In conclusion, we derive  $D_n^1 = o(s_n^2)$ ,  $D_n^2 = o(s_n^2)$  and thus  $C_n = D_n^1 + D_n^2 = o(s_n^2)$ . We then deduce from [\(3.39\)](#page-25-3) that  $A_n(C, \varepsilon) = o(s_n^2)$ . Consequently, the desired conclusion of the lemma follows. □

Lemma 3.17. Under assumption [\(C3\),](#page-23-7) there is an  $\varepsilon_0 = \varepsilon_0(C) > 0$  such that

<span id="page-28-0"></span>(3.49) 
$$
\frac{A_n(C,\varepsilon)}{s_n^2} \to 0 \quad as \quad n \to \infty \quad \text{for all } \varepsilon \in (0,\varepsilon_0).
$$

*Proof.* We first set  $\varepsilon_0 := \min{\{\tilde{\varepsilon}, \varepsilon_2\}}$  with constants  $\tilde{\varepsilon}$  given in [\(3.23\)](#page-23-8) and  $\varepsilon_2$  defined as in the proof of [Lemma 3.15.](#page-23-6) Let us take  $\varepsilon \in (0, \varepsilon_0)$  arbitrarily but fixed and reuse all symbols defined in the proof of [Lemma 3.15.](#page-23-6) Moreover, the relations [\(3.34\)](#page-24-3)–[\(3.39\)](#page-25-3) are still valid. From the definition of  $C_n$  in [\(3.36\)](#page-24-4), we deduce from [\(3.37\)](#page-25-0) that

$$
\frac{1}{s_n^2}|C_n| \leq \|\nabla y_n\|_{L^{\infty}(\{0<|\bar{y}-\bar{t}| \leq \tau_n\}\cap C^{\varepsilon})}\|\nabla \bar{\varphi}\|_{L^{\infty}(\Omega)}\frac{1}{s_n^2}\int_{C^{\varepsilon}}|\bar{y}-\bar{y}_n|\mathbb{1}_{\{0<|\bar{y}-\bar{t}| \leq \tau_n\}}\,dx
$$
  
\n
$$
\leq \|\nabla y_n\|_{L^{\infty}(\{0<|\bar{y}-\bar{t}|\leq \tau_n\}\cap C^{\varepsilon})}\|\nabla \bar{\varphi}\|_{L^{\infty}(\Omega)}\frac{\tau_n}{s_n^2}\operatorname{meas}_{\mathbb{R}^2}(\{0<|\bar{y}-\bar{t}| \leq \tau_n\}\cap C^{\varepsilon})
$$
  
\n
$$
\leq c_s\|\nabla y_n - \nabla \bar{y}\|_{L^{\infty}(\Omega)}\|\nabla \bar{\varphi}\|_{L^{\infty}(\Omega)}\frac{\tau_n^2}{s_n^2} + c_s\|\nabla \bar{y}\|_{L^{\infty}(\{0<|\bar{y}-\bar{t}| \leq \tau_n\}\cap C^{\varepsilon})}\|\nabla \bar{\varphi}\|_{L^{\infty}(\Omega)}\frac{\tau_n^2}{s_n^2}
$$

for all *n* large enough, where we have exploited the last estimate in [\(3.23\)](#page-23-8). Letting  $n \to \infty$  and using the identity in [\(3.23\)](#page-23-8) yields

$$
C_n = o(s_n^2),
$$

which along with  $(3.39)$  gives  $(3.49)$ .

Similar to [Lemmas 3.15](#page-23-6) and [3.17,](#page-27-0) we have the following result.

<span id="page-28-3"></span>Proposition 3.18. Assume that either [\(C1\)](#page-23-2) or [\(C3\)](#page-23-7) is fulfilled. Then there exists an  $\varepsilon_0 = \varepsilon_0(C) > 0$  such that, for any  $\varepsilon \in (0, \varepsilon_0)$ ,

<span id="page-28-2"></span>
$$
(3.50) \qquad \frac{\tilde{A}_n(C,\varepsilon)}{s_n^2} \to -\frac{1}{2} \int_C \mathbb{1}_{\{|\nabla \bar{y}|>0\}} w^2 \frac{\nabla \bar{y} \cdot \nabla \bar{\varphi}}{|\nabla \bar{y}|} d\mathcal{H}^1(x) \quad \text{as} \quad n \to \infty,
$$

where

(3.51) 
$$
\tilde{A}_n(C,\varepsilon) := \int_{C^{\varepsilon}} (\bar{t} - \bar{y}) [\mathbb{1}_{\Omega^2_{y_n,\bar{y}}} - \mathbb{1}_{\Omega^3_{y_n,\bar{y}}}]\nabla \bar{y} \cdot \nabla \bar{\phi} dx.
$$

*Proof.* We first note from  $(3.26)$  and  $(3.28)$  that

<span id="page-28-1"></span>
$$
\label{eq:2} \begin{cases} \Omega^2_{y_n,\bar{y}}=\tilde{\Omega}^2_{y_n,\bar{y}}\cup\{\bar{y}\in(\bar{t},\bar{t}+\delta),\,y_n=\bar{t}\}=\tilde{\Omega}^3_{\bar{y},y_n}\cup\{\bar{y}\in(\bar{t},\bar{t}+\delta),\,y_n=\bar{t}\},\\ \Omega^3_{y_n,\bar{y}}=\tilde{\Omega}^3_{y_n,\bar{y}}\cup\{\bar{y}\in(\bar{t}-\delta,\bar{t}),\,y_n=\bar{t}\}=\tilde{\Omega}^2_{\bar{y},y_n}\cup\{\bar{y}\in(\bar{t}-\delta,\bar{t}),\,y_n=\bar{t}\}, \end{cases}
$$

where the sets  $\tilde{\Omega}^2_{\nu,\hat{\nu}}$  and  $\tilde{\Omega}^3_{\nu,\hat{\nu}}$  are defined as in [\(3.27\)](#page-23-3). Obviously, one has

$$
\tilde{\Omega}^2_{\bar{y},y_n} \cap \{\bar{y} \in (\bar{t}-\delta,\bar{t}), y_n = \bar{t}\} = \emptyset = \tilde{\Omega}^3_{\bar{y},y_n} \cap \{\bar{y} \in (\bar{t},\bar{t}+\delta), y_n = \bar{t}\}.
$$

Combining these with the definition of  $\tilde{A}_n(C, \varepsilon)$  in [\(3.51\)](#page-28-1) yields

$$
\tilde{A}_n(C,\varepsilon) = \int_{C^{\varepsilon}} (\bar{t} - \bar{y}) [1_{\{\bar{y} \in (\bar{t}, \bar{t} + \delta), y_n = \bar{t}\}} - 1_{\{\bar{y} \in (\bar{t} - \delta, \bar{t}), y_n = \bar{t}\}}] \nabla \bar{y} \cdot \nabla \bar{\varphi} \, dx \n- \int_{C^{\varepsilon}} (\bar{t} - \bar{y}) [1_{\tilde{\Omega}^2_{\bar{y}, y_n}} - 1_{\tilde{\Omega}^3_{\bar{y}, y_n}}] \nabla \bar{y} \cdot \nabla \bar{\varphi} \, dx =: \tilde{B}_n - \tilde{C}_n.
$$

Since  $\nabla y_n = 0$  a.e. on  $\{y_n = \overline{t}\}, \tilde{B}_n$  can be rewritten as follows

$$
\tilde{B}_n = \int_{C^{\varepsilon}} (y_n - \bar{y}) [\mathbb{1}_{\{\bar{y} \in (\bar{t}, \bar{t} + \delta), y_n = \bar{t}\}} - \mathbb{1}_{\{\bar{y} \in (\bar{t} - \delta, \bar{t}), y_n = \bar{t}\}}] \nabla (\bar{y} - y_n) \cdot \nabla \bar{\varphi} \, \mathrm{d}x.
$$

Analogous to [\(3.38\)](#page-25-6), one has  $\tilde{B}_n = o(s_n^2)$ . This implies that

$$
\tilde{A}_n(C,\varepsilon) = o(s_n^2) - \tilde{C}_n.
$$

From the definitions of  $C_n$  in [\(3.36\)](#page-24-4) and of  $\tilde{C}_n$ , we can get  $\tilde{C}_n$  by interchanging  $\bar{\gamma}$  and  $\gamma_n$  in the integrand of  $C_n$ . For the situation where  $(C_3)$  holds, by using the argument as in the proof of [Lemma 3.17,](#page-27-0) we have

<span id="page-29-0"></span>
$$
\tilde{C}_n = o(s_n^2),
$$

which together with  $(3.52)$  gives  $(3.50)$ . It remains to consider the situation where  $(C_1)$  is satisfied. We shall estimate  $\tilde{C}_n$  similarly to the estimate of  $C_n$  in the proof of [Lemma 3.15.](#page-23-6) To this end, by [\(3.27\)](#page-23-3), we have for  $n$  large enough that

<span id="page-29-2"></span>
$$
(3.53) \qquad \begin{cases} \tilde{\Omega}_{\bar{y},y_n}^2 \cap C^{\varepsilon} = \{y_n \in (\bar{t},\bar{t}+\delta), \, \bar{y} \in (\bar{t}-\delta,\bar{t})\} \cap C^{\varepsilon} = \{\bar{y} < \bar{t} < y_n\} \cap C^{\varepsilon} =: \tilde{\Omega}_n^2, \\ \tilde{\Omega}_{\bar{y},y_n}^3 \cap C^{\varepsilon} = \{\bar{y} \in (\bar{t},\bar{t}+\delta), \, y_n \in (\bar{t}-\delta,\bar{t})\} \cap C^{\varepsilon} = \{\bar{y} > \bar{t} > y_n\} \cap C^{\varepsilon} =: \tilde{\Omega}_n^3. \end{cases}
$$

In comparison with  $(3.34)$ , there also hold

<span id="page-29-1"></span>
$$
\tilde{\Omega}_n^2 = \Omega_n^3 \quad \text{and} \quad \tilde{\Omega}_n^3 = \Omega_n^2.
$$

We now rewrite  $\tilde{C}_n$  as

(3.54) 
$$
\tilde{C}_n = -\frac{1}{2} \int_{C^{\varepsilon}} [\mathbb{1}_{\tilde{\Omega}_n^2} - \mathbb{1}_{\tilde{\Omega}_n^3}] \nabla (\bar{y} - \bar{t})^2 \cdot \nabla \bar{\varphi} \, dx.
$$

Now we consider the following two cases.

• Case 1:  $C = \partial \Omega$ . Then by using the same argument as in Case 1 in the proof of [Lemma 3.15,](#page-23-6) we have  $\tilde{C}_n = 0.$ 

• Case 2:  $C \neq \partial \Omega$ . For this case, one has  $C \cap \partial \Omega = \emptyset$ . Using [\(3.41\)](#page-25-7) and

applying [Proposition 2.9](#page-12-0) for functions  $y_1 := y_n$ ,  $y_2 := \bar{y}$ ,  $v := (\bar{y} - \bar{t})^2$ ,  $\varphi := \bar{\varphi}$  and the sets  $S^+ := \tilde{\Omega}_n^2$ ,  $S^- := \tilde{\Omega}_n^3$ , we deduce from [\(3.54\)](#page-29-1) and [\(3.53\)](#page-29-2) that

$$
\tilde{C}_n = \frac{1}{2} \int_{C^{\varepsilon}} [\mathbb{1}_{\tilde{\Omega}_n^2} - \mathbb{1}_{\tilde{\Omega}_n^3}](\bar{y} - \bar{t})^2 \Delta \bar{\varphi} \, dx + \frac{1}{2} \int_{C_n} (\bar{y} - \bar{t})^2 \nabla \bar{\varphi} \cdot \frac{\nabla y_n}{|\nabla y_n|} \, d\mathcal{H}^1(x) - \frac{1}{2} \int_C (\bar{y} - \bar{t})^2 \nabla \bar{\varphi} \cdot \frac{\nabla \bar{y}}{|\nabla \bar{y}|} \, d\mathcal{H}^1(x)
$$

with  $C_n := \{y_n = \overline{t}\} \cap C^{\varepsilon}$  being a  $C^1$  closed simple curve for *n* large enough; see [Proposition 2.11.](#page-14-0) Consequently, one has

<span id="page-29-3"></span>
$$
(3.55) \qquad \tilde{C}_n = \frac{1}{2} \int_{C^{\varepsilon}} [\mathbb{1}_{\tilde{\Omega}_n^2} - \mathbb{1}_{\tilde{\Omega}_n^3}](\bar{y} - \bar{t})^2 \Delta \bar{\varphi} \, dx + \frac{1}{2} \int_{C_n} (\bar{y} - y_n)^2 \nabla \bar{\varphi} \cdot \frac{\nabla y_n}{|\nabla y_n|} \, d\mathcal{H}^1(x),
$$

which is similar to that in  $(3.42)$ . Moreover, analogous to  $(3.43)$ , the first term in the right-hand side of  $(3.55)$  is  $o(s_n^2)$ . This implies that

<span id="page-29-4"></span>(3.56) 
$$
\tilde{C}_n = o(s_n^2) + \frac{1}{2} \int_{\{y_n = \bar{t}\} \cap C^{\varepsilon}} (\bar{y} - y_n)^2 \nabla \bar{\varphi} \cdot \frac{\nabla y_n}{|\nabla y_n|} d\mathcal{H}^1(x).
$$

This expression is analogous to [\(3.44\)](#page-25-2) and also valid to the case  $C := \partial \Omega$ . From [\(3.56\)](#page-29-4) and [\(3.52\)](#page-29-0), we derive [\(3.50\)](#page-28-2) by using [Proposition 2.13](#page-17-0) together with the limits  $y_n \to \bar{y}$  in  $C^1(\overline{\Omega})$  and  $(y_n - \bar{y})/s_n \to \infty$  $\lim W_0^{1,p}$  $C(\overline{\Omega}).$ 

<span id="page-29-5"></span>As a consequence of [Theorem 3.12](#page-22-2) and [Proposition 3.18,](#page-28-3) we have an explicit formula for the crucial term  $\tilde{O}$  in [\(3.13\)](#page-21-3), and an important limit that will play a significant role in establishing the error estimates for the numerical approximations of [\(P\)](#page-0-0).

<span id="page-30-1"></span>(a) If, for any connected component C of  $\{\bar{v} = \bar{t}\}\$ ,  $\bar{v}$  fulfills either [\(3.21\)](#page-23-11), [\(3.22\)](#page-23-0), or [\(3.23\)](#page-23-8), then

<span id="page-30-0"></span>
$$
(3.57) \qquad \tilde{Q}(\bar{u},\bar{y},\bar{\varphi};\{s_n\},v)=\frac{1}{2}\{a'\}_{\bar{t}+0}^{\bar{t}-0}\int_{\{\bar{y}=\bar{t}\}}\mathbb{1}_{\{|\nabla\bar{y}|>0\}}w^2\frac{\nabla\bar{y}\cdot\nabla\bar{\varphi}}{|\nabla\bar{y}|}d\mathcal{H}^1(x).
$$

<span id="page-30-2"></span>(b) If, for any connected component C of  $\{\bar{y} = \bar{t}\}\$ ,  $\bar{y}$  fulfills either [\(3.21\)](#page-23-11) or [\(3.23\)](#page-23-8), then

<span id="page-30-3"></span>
$$
\lim_{n \to \infty} \frac{1}{s_n^2} \int_{\Omega} (2\bar{t} - \bar{y} - y_n) [\mathbb{1}_{\Omega^2_{y_n, \bar{y}}} - \mathbb{1}_{\Omega^3_{y_n, \bar{y}}}]\nabla \bar{y} \cdot \nabla \bar{\phi} dx = 0.
$$

*Proof.* Assume that  $\{\bar{y} = \bar{t}\} = \bigcup_{k=1}^{m} C_k$  with  $C_k$  being connected components of  $\{\bar{y} = \bar{t}\}$ . Let  $\varepsilon_0 > 0$  be such that  $(C_{k_1})^{\varepsilon_0} \cap (C_{k_2})^{\varepsilon_0} = \emptyset$  for all  $k_1 \neq k_2$ . By [Theorem 3.12,](#page-22-2) for any  $1 \leq k \leq m$ , there exists  $\varepsilon_k > 0$ satisfying the claims of [Theorem 3.12](#page-22-2) in place of  $\varepsilon_0$ . We now set  $\varepsilon_* := \min\{\varepsilon_0, \varepsilon_k \mid 1 \le k \le m\} > 0$  and fix  $\varepsilon \in (0, \varepsilon_*)$  arbitrarily. We first prove [\(3.57\)](#page-30-0). In fact, all assumptions required in [Theorem 3.12](#page-22-2) are fulfilled. From [\(3.11\)](#page-21-0), we have

$$
\Omega^2_{y_n,\bar{y}} \cup \Omega^3_{y_n,\bar{y}} \subset \{0 < |\bar{y} - \bar{t}| \le \tau_n\} \subset \bigcup_{k=1}^m (C_k)^{\varepsilon}
$$

for  $n$  large enough, where  $\tau_n\coloneqq \|y_n-\bar{y}\|_{C(\overline{\Omega})}.$  From the definition of  $\tilde{Q}\coloneqq \tilde{Q}(\bar{u},\bar{y},\bar{\varphi};\{s_n\},v)$  in [\(3.13\)](#page-21-3) and of  $A_n(C, \varepsilon)$  in [Theorem 3.12,](#page-22-2) we then deduce from [\(3.18\)](#page-21-4) that

$$
\tilde{Q} = \liminf_{n \to \infty} \frac{1}{s_n^2} \{a'\}_{\bar{t}+0}^{\bar{t}-0} \sum_{k=1}^m A_n(C_k, \varepsilon) = \{a'\}_{\bar{t}+0}^{\bar{t}-0} \sum_{k=1}^m \frac{1}{2} \int_{C_k} \mathbb{1}_{\{|\nabla \bar{y}| > 0\}} w^2 \frac{\nabla \bar{y} \cdot \nabla \bar{\varphi}}{|\nabla \bar{y}|} d\mathcal{H}^1(x),
$$

which, together with the disjoint decomposition  $\{\bar{y} = \bar{t}\} = \bigcup_{k=1}^{m} C_k$  yields the identity in [\(3.57\)](#page-30-0). Assertion [\(a\)](#page-30-1) is then proven.

For assertion [\(b\),](#page-30-2) by using the definitions of  $A_n(C, \varepsilon)$  and of  $\tilde{A}_n(C, \varepsilon)$ , respectively, in [Theorem 3.12](#page-22-2) and [Proposition 3.18,](#page-28-3) there holds

$$
\int_{\Omega} (2\overline{t} - \overline{y} - y_n) [\mathbb{1}_{\Omega^2_{y_n, \overline{y}}} - \mathbb{1}_{\Omega^3_{y_n, \overline{y}}}]\nabla \overline{y} \cdot \nabla \overline{\varphi} dx = \sum_{k=1}^m [A_n(C_k, \varepsilon) + \tilde{A}_n(C_k, \varepsilon)].
$$

We thus derive  $(3.58)$  by [Theorem 3.12](#page-22-2) and [Proposition 3.18.](#page-28-3) □

As the last preparatory step, the following result shows the finiteness of the jump functional  $\Sigma(\bar{y})$ determined in  $(3.17)$  under one of three assumptions  $(C_1)$ ,  $(C_2)$ , and  $(C_3)$ .

<span id="page-30-5"></span>Proposition 3.20. Let  $\bar{u} \in L^2(\Omega)$  be such that  $\bar{y} := S(\bar{u})$  is Lipschitz continuous on  $\overline{\Omega}$ . Assume further that the level set  $\{\bar{v} = \bar{t}\}$  decomposes into finitely many connected components and that, on each such connected component C, either [\(C1\),](#page-23-2) [\(C2\),](#page-23-5) or [\(C3\)](#page-23-7) in [Theorem 3.12](#page-22-2) is fulfilled. Then  $\Sigma(\bar{v}) < \infty$ .

*Proof.* Assume that  $\{\bar{y} = \bar{t}\} = \bigcup_{k=1}^{m} C_k$  with  $C_k$  being connected components of  $\{\bar{y} = \bar{t}\}\)$ . Let  $\varepsilon_0 \in (0, \tilde{\varepsilon})$ be such that  $(C_{k_1})^{\varepsilon_0} \cap (C_{k_2})^{\varepsilon_0} = \emptyset$  for all  $k_1 \neq k_2$ , where  $\tilde{\varepsilon}$  is given in [Theorem 3.12.](#page-22-2) For  $r > 0$  small enough, one has

<span id="page-30-4"></span>(3.59) 
$$
\int_{\Omega} 1_{\{0<|\bar{y}-\bar{t}|
$$

Clason, Nhu, Rösch Numerical analysis of a nonsmooth quasilinear elliptic ...

If either assumption (C<sub>1</sub>) or (C<sub>3</sub>) is verified on  $C_k$  for some  $1 \leq k \leq m$ , then we deduce for any *r* small enough that

$$
\int_{C_k^{\varepsilon_0}} 1_{\{0<|\bar{y}-\bar{t}|
$$

where we have used Lemma 3.2 in [\[11\]](#page-36-12) to obtain the last estimate. Here  $C_k := \max\{|\nabla \bar{y}(x)| : x \in \overline{C_k^{\varepsilon_0}}\}$  $\binom{\varepsilon_0}{k}$ for  $1 \le k \le m$ . If assumption [\(C2\)](#page-23-5) holds on  $C_k$  for some  $1 \le k \le m$ , then by applying the coarea formula for Lipschitz mappings (see, e.g. [\[12\]](#page-36-9), Thm. 2, p. 117 and [\[1\]](#page-36-6), Sec. 2.7), we have for any  $r \in (0, r_0)$  that

$$
\int_{C_k^{\varepsilon_0}} \mathbb{1}_{\{0<|\bar{y}-\bar{t}|
$$

From these estimates, [\(3.59\)](#page-30-4), and the definition of  $\Sigma(\bar{y})$  in [\(3.17\)](#page-21-2), we conclude that  $\Sigma(\bar{y}) < \infty$ . □

The following theorem now formulates an explicit formula of the nonsmooth curvature functional  $Q$  defined in  $(3.15)$ .

<span id="page-31-0"></span>Theorem 3.21. Let Assumptions (A1) to (A4) be fulfilled. Assume that  $\bar{u} \in \mathcal{U}_{ad}$  and that  $v \in \text{cl}[\text{cone}(\mathcal{U}_{ad}$  –  $\bar{u}$ ]. Assume further that the level set  $\{\bar{y} = \bar{t}\}$  with  $\bar{y} = S(\bar{u})$  decomposes into finitely many connected components and that, on each such connected component C, either [\(C1\),](#page-23-2) [\(C2\),](#page-23-5) or [\(C3\)](#page-23-7) in [Theorem 3.12](#page-22-2) is fulfilled. Then, for any  $\varphi \in C^1(\overline{\Omega}) \cap W^{2,1}(\Omega)$ , there holds

<span id="page-31-2"></span>
$$
(3.60) \quad Q(\bar{u}, \bar{y}, \varphi; v) = \frac{1}{2} \int_{\Omega} \frac{\partial^2 L}{\partial y^2} (\cdot, \bar{y}) z_v^2 \, dx + \frac{v}{2} \int_{\Omega} v^2 \, dx - \frac{1}{2} \int_{\Omega} \mathbb{1}_{\{y \neq \bar{t}\}} a''(\bar{y}) z_v^2 \nabla \bar{y} \cdot \nabla \varphi \, dx - \int_{\Omega} a'(\bar{y}; z_v) \nabla z_v \cdot \nabla \varphi \, dx + \frac{1}{2} [a'_0(\bar{t}) - a'_1(\bar{t})] \int_{\{\bar{y} = \bar{t}\}} \mathbb{1}_{\{|\nabla \bar{y}| > 0\}} z_v^2 \frac{\nabla \bar{y} \cdot \nabla \varphi}{|\nabla \bar{y}|} d\mathcal{H}^1(x)
$$

with  $z_v := S(\bar{u})v$ . Here cl[cone( $\mathcal{U}_{ad} - \bar{u}$ )] denotes the closure in  $L^2(\Omega)$  of the cone generated by ( $\mathcal{U}_{ad} - \bar{u}$ ).

Proof. We first observe that under the stated assumptions,  $\Sigma(\bar{y})$  is finite by [Proposition 3.20.](#page-30-5) As a result of [\(3.16\)](#page-21-5),  $Q_2(\bar{u},\bar{y},\varphi;\cdot)$  and thus  $Q(\bar{u},\bar{y},\varphi;\cdot)$  (defined, respectively, in [\(3.14\)](#page-21-6) and [\(3.15\)](#page-21-1)) are well-defined on  $L^2(\Omega)$ . Let  $\{s_n\} \in c_0^+$  be arbitrary. Taking  $v \in \text{cl}[\text{cone}(\mathcal{U}_{ad} - \bar{u})]$  arbitrarily, then there are sequences  ${v_m} \subset L^2(\Omega)$ ,  ${u_m} \subset \mathcal{U}_{ad}$ , and  ${\lambda_m} \subset (0,\infty)$  such that

$$
v_m \to v
$$
 in  $L^2(\Omega)$  and  $v_m = \frac{u_m - \bar{u}}{\lambda_m}$  for all  $m \in \mathbb{N}$ .

By the definition of  $\tilde{Q}(\bar{u},\bar{y},\bar{\phi};\{s_n\},v)$  in [\(3.13\)](#page-21-3), there holds

<span id="page-31-1"></span>
$$
(3.61) \qquad \tilde{Q}(\bar{u}, \bar{y}, \bar{\varphi}; \{s_n\}, v) = \lim_{k \to \infty} \frac{1}{s_{n_k}^2} \int_{\Omega} \sum_{i=0}^1 \zeta_i(\bar{u}, \bar{y}; s_{n_k}, v) \nabla \bar{y} \cdot \nabla \bar{\varphi} \, dx = \tilde{Q}(\bar{u}, \bar{y}, \bar{\varphi}; \{s_{n_k}\}, v)
$$

for some subsequence  $\{s_{n_k}\}\$  of  $\{s_n\}$ . Since  $s_{n_k} \to 0^+$  as  $k \to \infty$ , for any  $m \ge 1$ , there exists an integer  $k(m)$  such that  $r_m := s_{n_{k(m)}} \in (0, \lambda_m)$  and thus

$$
\bar{u}+r_mv_m=\left(1-\frac{r_m}{\lambda_m}\right)\bar{u}+\frac{r_m}{\lambda_m}u_m\in\mathcal{U}_{ad}.
$$

Setting  $y_m := S(\bar{u} + r_m v_m)$  yields  $(y_m - \bar{y})/r_m \rightarrow S'(\bar{u})v$  in  $W_0^{1,p}$  $y_0^{1,p}(\Omega)$  and  $y_m \in W^{2,p}(\Omega)$  for some  $p > 2$ , according to [Theorem 3.2.](#page-19-7) From this and the compact embedding  $W^{2,p}(\Omega)\Subset C^1(\overline{\Omega}),$  one has  $y_m\to\bar{y}$ 

in  $C^1(\overline{\Omega})$ . Using now the limit  $v_m \to v$  in  $L^2(\Omega)$ , [\(3.61\)](#page-31-1), as well as [\(3.18\)](#page-21-4), and then applying [Theorem 3.12,](#page-22-2) one has

$$
\tilde{Q}(\bar{u}, \bar{y}, \bar{\varphi}; \{s_n\}, v) = \tilde{Q}(\bar{u}, \bar{y}, \bar{\varphi}; \{r_m\}, v)
$$
\n
$$
= \lim_{m \to \infty} \frac{1}{r_m^2} \int_{\Omega} \sum_{i=0}^1 \zeta_i(\bar{u}, \bar{y}; r_m, v_m) \nabla \bar{y} \cdot \nabla \bar{\varphi} \, dx
$$
\n
$$
= \frac{1}{2} \{a'\}_{\bar{t}+0}^{\bar{t}-0} \int_{\{\bar{y}=\bar{t}\}} \mathbb{1}_{\{|\nabla \bar{y}|>0\}} (S'(\bar{u})v)^2 \frac{\nabla \bar{y} \cdot \nabla \varphi}{|\nabla \bar{y}|} \, d\mathcal{H}^1(x).
$$

Since  $\{s_n\}$  is taken arbitrarily, we then deduce from the definition of  $Q_2$  that

$$
Q_2(\bar{u}, \bar{y}, \bar{\varphi}; v) = \tilde{Q}(\bar{u}, \bar{y}, \bar{\varphi}; \{s_n\}, v) = \frac{1}{2} \{a'\}_{\bar{t}+0}^{\bar{t}-0} \int_{\{\bar{y}=\bar{t}\}} \mathbb{1}_{\{|\nabla \bar{y}|>0\}} (S'(\bar{u})v)^2 \frac{\nabla \bar{y} \cdot \nabla \varphi}{|\nabla \bar{y}|} d\mathcal{H}^1(x).
$$

Combining this with the definition of  $Q$  in [\(3.15\)](#page-21-1) yields [\(3.60\)](#page-31-2). □

<span id="page-32-2"></span>3.3 explicit second-order optimality conditions

The following explicit second-order necessary and sufficient optimality conditions for [\(P\)](#page-0-0) are consequences of [Theorems 3.7,](#page-22-3) [3.8](#page-22-4) and [3.19.](#page-29-5)

<span id="page-32-0"></span>Theorem 3.22 (explicit second-order necessary optimality conditions). Let Assumptions (A1) to (A4) be fulfilled. Assume that  $\bar{u}$  is a local minimizer to [\(P\)](#page-20-0). Assume further that the level set  $\{\bar{y} = \bar{t}\}$  with  $\bar{y} := S(\bar{u})$ decomposes into finitely many connected components and that, on each such connected component C, either (C<sub>1</sub>), (C<sub>2</sub>), or (C<sub>3</sub>) in [Theorem 3.12](#page-22-2) is fulfilled. Then, there is a  $\bar{\varphi} \in W_0^{1,\bar{p}}$  $U_0^{1,\tilde{p}}(\Omega)\cap W^{1,\infty}(\Omega)$ , with  $\tilde{p}$  defined in Assumption (A4), that together with  $\bar{u}$ ,  $\bar{v}$  satisfies [\(3.9\)](#page-20-1) and

<span id="page-32-3"></span>
$$
(3.62) \quad Q(\bar{u}, \bar{y}, \bar{\varphi}; v) = \frac{1}{2} \int_{\Omega} \frac{\partial^2 L}{\partial y^2} (\cdot, \bar{y}) z_v^2 \, dx + \frac{v}{2} \int_{\Omega} v^2 \, dx - \frac{1}{2} \int_{\Omega} \mathbb{1}_{\{y \neq \bar{t}\}} a''(\bar{y}) z_v^2 \nabla \bar{y} \cdot \nabla \bar{\varphi} \, dx - \int_{\Omega} a'(\bar{y}; z_v) \nabla z_v \cdot \nabla \bar{\varphi} \, dx + \frac{1}{2} [a'_0(\bar{t}) - a'_1(\bar{t})] \int_{\{\bar{y} = \bar{t}\}} \mathbb{1}_{\{|\nabla \bar{y}| > 0\}} z_v^2 \frac{\nabla \bar{y} \cdot \nabla \bar{\varphi}}{|\nabla \bar{y}|} d\mathcal{H}^1(x) \ge 0
$$

for all  $v \in C(\mathcal{U}_{ad}; \bar{u})$  with  $z_n := S(\bar{u})v$ .

*Proof.* By [Proposition 3.20,](#page-30-5)  $\Sigma(\bar{y})$  is finite. In view of [Theorem 3.7,](#page-22-3) a function  $\bar{\varphi}$  exists and satisfies [\(3.9\)](#page-20-1). By [Theorem 3.5,](#page-20-2)  $\bar{\varphi} \in W^{2,r}(\Omega) \hookrightarrow C^1(\overline{\Omega})$  for some  $r > 2 = N$ . Moreover, there holds  $C(\mathcal{U}_{ad}, \bar{u}) \subset$ cl[cone( $\mathcal{U}_{ad}$  –  $\bar{u}$ )]. Applying [Theorems 3.7](#page-22-3) and [3.21](#page-31-0) yields [\(3.62\)](#page-32-3).  $\Box$ 

<span id="page-32-1"></span>Theorem 3.23 (explicit second-order sufficient optimality conditions). Let Assumptions (A1) to (A4) be valid. Assume that  $\bar{u}$  is a feasible point of [\(P\)](#page-20-0). Assume that the level set  $\{\bar{v} = \bar{t}\}\$  with  $\bar{v} := S(\bar{u})$  decomposes into finitely many connected components and that, on each such connected component C, either [\(C1\),](#page-23-2) [\(C2\),](#page-23-5) or  $(C_3)$  in [Theorem 3.12](#page-22-2) is fulfilled. Assume further that there is a  $\bar{\varphi} \in W_0^{1,\bar{p}}$  $\overline{O}_0^{1,\bar{p}}(\Omega) \cap W^{1,\infty}(\Omega)$ , with  $\bar{p}$  defined in Assumption (A4), that together with  $\bar{u}$ ,  $\bar{y}$  satisfies [\(3.9\)](#page-20-1) and

$$
Q(\bar{u}, \bar{y}, \bar{\varphi}; v) = \frac{1}{2} \int_{\Omega} \frac{\partial^2 L}{\partial y^2} (\cdot, \bar{y}) z_v^2 \, dx + \frac{v}{2} \int_{\Omega} v^2 \, dx - \frac{1}{2} \int_{\Omega} \mathbb{1}_{\{y \neq \bar{t}\}} a''(\bar{y}) z_v^2 \nabla \bar{y} \cdot \nabla \bar{\varphi} \, dx - \int_{\Omega} a'(\bar{y}; z_v) \nabla z_v \cdot \nabla \bar{\varphi} \, dx + \frac{1}{2} [a_0'(\bar{t}) - a_1'(\bar{t})] \int_{\{\bar{y} = \bar{t}\}} \mathbb{1}_{\{|\nabla \bar{y}| > 0\}} z_v^2 \frac{\nabla \bar{y} \cdot \nabla \bar{\varphi}}{|\nabla \bar{y}|} d\mathcal{H}^1(x) > 0
$$

with  $z_v := S'(\bar{u})v$  for all  $v \in C(\mathcal{U}_{ad}; \bar{u}) \setminus \{0\}$ . Then there exist constants  $c_0, \rho_0 > 0$  satisfying

 $j(\bar{u}) + c_0 \|u - \bar{u}\|_{L}^2$  $L^2(\Omega) \leq j(u)$  for all  $u \in \mathcal{U}_{ad} \cap \overline{B}_{L^2(\Omega)}(\overline{u}, \rho_0)$ .

Proof. Similar to the proof of [Theorem 3.22,](#page-32-0) the desired conclusion follows from [Theorems 3.8](#page-22-4) and [3.21.](#page-31-0)  $\Box$ 

### 4 conclusions

We have derived necessary and sufficient second-order optimality conditions for a nonsmooth quasilinear elliptic optimal control problems in terms of an explicit curvature functional. The main technical tools are a detailed study of the properties of level sets of  $C^1$  functions with nonvanishing gradients, including continuity properties and a Green's identity. These results may be of independent interest in, e.g., the analysis of level set methods or in mathematical imaging. In the second part of this work, the sufficient second-order conditions will be applied to derive error estimates for a finite element discretization of the nonsmooth optimal control problem.

#### appendix a existence of regular points

<span id="page-33-0"></span>Lemma A.1. Let  $y_1, y_2 \in C^1(\overline{\Omega})$  and  $t \in \mathbb{R}$  be such that  $\{y_1 > t > y_2\} \neq \emptyset$ . Let  $\mathcal S$  be a nonempty open connected component of  $\{y_1 > t > y_2\}$  that satisfies  $\partial S \cap \partial \Omega = \emptyset$  and [\(2.28\)](#page-10-2) in [Lemma 2.7.](#page-10-0) Assume that the boundary of S has a cusp at N; see, [Figure 2b.](#page-11-0) Then there is a constant  $\varepsilon_0 > 0$  such that, for each  $0 < \varepsilon < \varepsilon_0$ , points  $N_i \in \partial S \cap \{y_i = t\}$ ,  $i = 1, 2$ , exist that satisfy

<span id="page-33-5"></span>
$$
l_{\widehat{NN}_i} < C\epsilon
$$

and

<span id="page-33-4"></span>
$$
(A.2) \t\t N_1N_2 < C\epsilon
$$

for some constant  $C > 0$ , where  $l_{\widehat{NN}_i}$  denotes the length of the arc  $\widehat{NN}_i$ . Furthermore,  $N_1$  and  $N_2$  are not are not are not all the subset of  $S$  and  $N$  and  $N$  and  $N$  are not are not all the subsets of cusps of  $\partial S_{\varepsilon}$ , where  $S_{\varepsilon}$  is the subdomain of S obtained by splitting S via the line N<sub>1</sub>N<sub>2</sub>, and N  $\notin S_{\varepsilon}$ .

*Proof.* Without loss of generality, assume that  $N = (0, 0) \in \mathbb{R}^2$  and that the positive  $x^2$ -axis ray is the common tangent line of two arcs of  $\partial S$  at N. Then the tangent vector of S at N is  $v_N = (-1, 0)$ . Since  $v_N = \kappa \nabla y_1(N) = \kappa \nabla y_2(N)$  for some  $\kappa \in \mathbb{R}$ , we have  $\frac{\partial y_1}{\partial x^2}(N) = \frac{\partial y_2}{\partial x^2}$  $\frac{\partial y_2}{\partial x^2}(N) = 0$  and

<span id="page-33-2"></span>(A.3) 
$$
\frac{\partial y_1}{\partial x^1}(x) \cdot \frac{\partial y_2}{\partial x^1}(x) \ge \alpha' > 0 \text{ for all } x \in \overline{\mathcal{S}} \cap B_{\mathbb{R}^2}(0,r)
$$

for some constants  $r > 0$  and  $\alpha'$ , due to [\(2.28\)](#page-10-2). Applying the Implicit Function Theorem at the point N yields that there exist constants  $h, k > 0, \rho \in (0, r)$  and  $C^1$ -functions  $\kappa_i : (-h, k) \to \mathbb{R}, i = 1, 2$ , such that

(A.4)  

$$
\begin{cases}\n\kappa_i(0) = 0, \\
y_i(\kappa_i(\xi), \xi) = t \quad \text{for all } \xi \in (-h, k), \\
((\tau, \xi) \in B_{\mathbb{R}^2}(0, \rho) \text{ and } y_i(\kappa_i(\xi), \xi) = t)\n\end{cases} \implies \tau = \kappa_i(\xi).
$$

Moreover, similar to  $(2.17)$ , one has

<span id="page-33-3"></span>
$$
|\kappa_i'(\xi)| \le c \quad \text{for all } \xi \in (-h, k)
$$

and for some constant  $c > 0$ . Since  $S \cap B_{\mathbb{R}^2}(N,r)$  is in the half space  $\{x^2 \ge 0\} := \{(x^1, x^2) \in \mathbb{R}^2 \mid x^2 \ge 0\}$ , we therefore have

$$
\{y_i = t\} \cap \partial S \cap B_{\mathbb{R}^2}(0, \rho_0) = \{(\tau, \xi) \in B_{\mathbb{R}^2}(0, \rho_0) \mid \xi \ge 0, y_i(\tau, \xi) = t\}
$$
  
(A.6)  

$$
= \{(\tau, \xi) \in B_{\mathbb{R}^2}(0, \rho_0) \mid \xi \in [0, k_i), \tau = \kappa_i(\xi)\}
$$

for some constants  $\rho_0 \in (0, \rho]$  and  $k_i \in (0, k]$ ,  $i = 1, 2$ . Moreover, the equation  $\kappa_1(\xi) = \kappa_2(\xi)$  has a unique solution  $\xi = 0$  in the interval  $(-h, k)$  because the arcs  $\{\gamma_1 = t\} \cap B_{\mathbb{R}^2}(0, \rho)$  and  $\{\gamma_2 = t\} \cap B_{\mathbb{R}^2}(0, \rho)$  has only one common point N. Consequently, one and only one of the following cases is valid:

<span id="page-33-1"></span>(A.7) 
$$
\kappa_1(\xi) > \kappa_2(\xi)
$$
 for all  $\xi \in (0, k)$  or  $\kappa_1(\xi) < \kappa_2(\xi)$  for all  $\xi \in (0, k)$ .

We now consider the first situation in  $(A.7)$  only since the other is analyzed analogously. Thanks to (A.3), one has two possibilities: For small number  $x^2$ , in the neighborhood of the origin in  $\mathbb R$ ,

- (I) both  $y_1(\cdot, x^2)$  and  $y_2(\cdot, x^2)$  are strictly increasing; or
- (II) both  $y_1(\cdot, x^2)$  and  $y_2(\cdot, x^2)$  are strictly decreasing.

For the case (I), one has for some constant  $0 < \rho_1 \le \rho_0$  that

$$
\{y_1 > t\} \cap B_{\mathbb{R}^2}(0, \rho_1) \cap \{\xi \ge 0\} \subset \{(\tau, \xi) \in \mathbb{R}^2 \mid \xi \in [0, k_1), \tau > \kappa_1(\xi)\}\
$$

and

$$
\{y_2 < t\} \cap B_{\mathbb{R}^2}(0, \rho_1) \cap \{\xi \geq 0\} \subset \{(\tau, \xi) \in \mathbb{R}^2 \mid \xi \in [0, k_2), \tau < \kappa_2(\xi)\}.
$$

Combining these with the first situation in  $(A,7)$  yields a contradiction. Therefore, the case (II) must be true. Thanks to (II) and [\(a.7\)](#page-33-1), there holds

$$
(A.8) \tS \cap B_{\mathbb{R}^2}(0,\underline{\rho}) \subset \{(\tau,\xi) \in \mathbb{R}^2 \mid \xi \in [0,k_*), \kappa_2(\xi) < \tau < \kappa_1(\xi)\} \subset \mathcal{S} \cap B_{\mathbb{R}^2}(0,\overline{\rho})
$$

for some positive constants  $0 < \rho < \overline{\rho} \le \rho_0$ , and  $0 < k_* \le \min\{k_1, k_2\}$ . Setting  $\varepsilon_0 := \rho$  and taking  $\varepsilon \in (0, \varepsilon_0)$  arbitrarily, we now fix  $\overline{\xi_1} \in (0, \varepsilon)$  and set  $N_1 := (\kappa_1(\xi_1), \xi_1)$  and  $N_2 := (\kappa_2(\xi_1), \xi_1)$ . Obviously, there holds  $N_i \in \{y_i = t\} \cap \partial S$ ,  $i = 1, 2$ . Moreover, using the inequalities (A.7) and (A.5) yields

$$
N_1N_2 = |\kappa_2(\xi_1) - \kappa_1(\xi_1)| = \kappa_1(\xi_1) - \kappa_2(\xi_1)
$$
  
\n
$$
\leq |\kappa_1(\xi_1) - \kappa_1(0)| + |\kappa_2(\xi_1) - \kappa_2(0)|
$$
  
\n
$$
\leq 2c\xi_1.
$$

Thus, (A.2) is verified. On the other hand, since  $\{(\kappa_i(\xi), \xi) : 0 \le \xi \le \xi_1\} = \widehat{NN}_i$ ,  $i = 1, 2$ , we conclude from  $(A.5)$  that

$$
l_{\widehat{NN_i}} = \int_0^{\xi_1} \sqrt{1 + (\kappa_i'(\xi))^2} d\xi \le \sqrt{1 + c^2} \xi_1 \le \varepsilon \sqrt{1 + c^2},
$$

which gives [\(a.1\)](#page-33-5).

It remains to prove that  $N_1$  and  $N_2$  are not cusps of  $\partial S_{\varepsilon}$ . To do this, we first consider the point  $N_1(\kappa_1(\xi_1), \xi_1)$ . Let  $\omega_1 \in [0, \pi/2]$  be the angle between the line  $N_1N_2$  and the tangent vector to  $\{y_1 = t\}$ at  $N_1$ . We then have

$$
\cos \omega_1 = \left| \frac{\nabla y_1}{|\nabla y_1|} (N_1) \cdot (0, 1) \right| = \left| \frac{\frac{\partial y_1}{\partial x^2}}{\sqrt{\left(\frac{\partial y_1}{\partial x^1}\right)^2 + \left(\frac{\partial y_1}{\partial x^2}\right)^2}} (N_1) \right| < 1,
$$

according to (A.3). Hence  $\omega_1 > 0$  and thus  $N_1$  is not a cusp of  $\partial S_{\varepsilon}$ . Similarly,  $N_2$  is also not a cusp of  $\partial S_{\varepsilon}$ . . □

<span id="page-34-0"></span>Lemma A.2. Let  $y \in C^1(\overline{\Omega})$  and  $t \in \mathbb{R}$  be such that  $\{y = t\} \neq \emptyset$  and  $y = 0$  on  $\partial \Omega$ . Let  $C$  be a connected component of  $\{y = t\}$  that satisfies [\(2.20\)](#page-7-4). Then, for any  $\varepsilon > 0$  and  $x_2 \in C^{\varepsilon}$ , there holds

$$
Arg(x_2) := argmin\{|x_2 - x| : x \in C\} \neq \emptyset.
$$

<span id="page-34-2"></span><span id="page-34-1"></span> $|x_1 - x_2| \leq \varepsilon$ 

Moreover, if  $x_1 \in \text{Arg}(x_2)$ , then

$$
(A.9)
$$

and

(A.10)  $x_2 - x_1 = \kappa \nabla y(x_1)$ 

for some constant  $\kappa \in \mathbb{R}$ .

*Proof.* By [Proposition 2.4,](#page-7-0) C is a closed C<sup>1</sup> simple curve. Moreover, for any  $x_2 = (x_2^1, x_2^2) \in C^{\varepsilon}$ , we obviously have Arg( $x_2$ ) ≠ 0 since C is compact. Taking  $x_1 \in \text{Arg}(x_2)$  arbitrarily, the definition of the set  $C^{\varepsilon}$  implies (A.9). It remains to prove (A.10). To do this, we first extend y by a  $C^1$  extension, denoted by the same name, on  $\mathbb{R}^2$  in the case where  $C = \partial\Omega$ , as a result of [Proposition 2.1.](#page-3-3) Due to [\(2.20\)](#page-7-4), without lost of generality, we now assume that

<span id="page-35-0"></span>
$$
\left|\frac{\partial y}{\partial x^2}(x)\right| \ge \alpha > 0 \quad \text{for all } x \in B_{\mathbb{R}^2}(x_1, r)
$$

for some constant  $r > 0$ . Applying now the Implicit Function Theorem in the point  $x_1 = (x_1^1, x_1^2) \in C$ yields that there exist constants  $h, k > 0, \rho \in (0, r)$ , and a  $C^1$ -function  $g: (x_1^1 - h, x_1^1 + k) \to \mathbb{R}$  satisfying

(A.11)

\n
$$
\begin{cases}\n g(x_1^1) = x_1^2, \\
 y(\tau, g(\tau)) = t \quad \text{for all } \tau \in (x_1^1 - h, x_1^1 + k), \\
 ((\tau, \xi) \in B_{\mathbb{R}^2}(x_1, \rho) \text{ and } y(\tau, \xi) = t)\n\end{cases}\n\implies \xi = g(\tau).
$$

Moreover, similar to [\(2.16\)](#page-6-3), there holds

(A.12) 
$$
g'(\tau) = -\frac{\frac{\partial y}{\partial x^{1}}(\tau, g(\tau))}{\frac{\partial y}{\partial x^{2}}(\tau, g(\tau))}, \quad \tau \in (x_{1}^{1} - h, x_{1}^{1} + k).
$$

For any  $s \in (-h, k)$ , one has  $x_s := (x_1^1 + s, g(x_1^1 + s)) \in C$  and the function

<span id="page-35-1"></span>
$$
f(s) := |x_2 - x_s|^2
$$

attains its minimum at  $s = 0$ , according to the definition of  $x_1$ . There thus holds  $f'(0) = 0$ . A simply computation gives

$$
f(s) = (x_2^1 - x_1^1 - s)^2 + (x_2^2 - g(x_1^1 + s))^2
$$

and

$$
f'(s) = -2(x_2^1 - x_1^1 - s) - 2(x_2^2 - g(x_1^1 + s))g'(x_1^1 + s).
$$

This implies that

$$
0 = f'(0) = -2(x_2^1 - x_1^1) - 2(x_2^2 - g(x_1^1))g'(x_1^1)
$$
  
=  $-2[(x_2^1 - x_1^1) + (x_2^2 - g(x_1^1))g'(x_1^1)]$   
=  $-2[(x_2^1 - x_1^1) - (x_2^2 - x_1^2)\frac{\frac{\partial y}{\partial x^1}}{\frac{\partial y}{\partial x^2}}(x_1)],$ 

where we have employed the first equation in  $(A.11)$  and the identity  $(A.12)$  to derive the last equation. We then obtain  $\partial \nu$ 

$$
(x_2^1 - x_1^1) = (x_2^2 - x_1^2) \frac{\frac{\partial y}{\partial x^1}(x_1)}{\frac{\partial y}{\partial x^2}(x_1)},
$$

which yields

$$
x_2 - x_1 = \frac{x_2^2 - x_1^2}{\frac{\partial y}{\partial x^2}(x_1)} \nabla y(x_1).
$$

Setting  $\kappa := \frac{x_2^2 - x_1^2}{\frac{\partial y}{\partial x^2}(x_1)}$  finally yields (A.10).

### acknowledgments

This work was supported by the DFG under the grants CL 487/2-1 and RO 2462/6-1, both within the priority programme SPP 1962 "Non-smooth and Complementarity-based Distributed Parameter Systems: Simulation and Hierarchical Optimization". The second author was funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 101.01–2019.308. The authors would like to thank the anonymous referees for their useful suggestions and comments which improved the manuscript greatly.

# **REFERENCES**

- <span id="page-36-6"></span>[1] G. Alberti, S. Bianchini, and G. Crippa, Structure of level sets and Sard-type properties of Lipschitz maps, Ann. Sc. Norm. Super. Pisa Cl. Sci.(5) XII (2013), 863–902, [doi:10.2422/2036-2145.201107\\_006](https://dx.doi.org/10.2422/2036-2145.201107_006).
- <span id="page-36-0"></span>[2] A. Bejan, Convection Heat Transfer, J. Wiley & Sons, 4 edition, 2013, [doi:10.1002/9781118671627](https://dx.doi.org/10.1002/9781118671627).
- <span id="page-36-7"></span>[3] J. Bourgain, M. V. Korobkov, and J. Kristensen, On the Morse–Sard property and level sets of Sobolev and BV functions, Rev. Mat. Iberoam. 29 (2013), 1–23, [doi:10.4171/rmi/710](https://dx.doi.org/10.4171/rmi/710).
- <span id="page-36-10"></span>[4] N.L. Carothers, Real Analysis, Cambridge University Press, Cambridge, 2000, [doi:10.1017/](https://dx.doi.org/10.1017/CBO9780511814228) [cbo9780511814228](https://dx.doi.org/10.1017/CBO9780511814228).
- <span id="page-36-3"></span>[5] E. Casas and V. Dhamo, Error estimates for the numerical approximation of Neumann control problems governed by a class of quasilinear elliptic equations, Comput. Optim. Appl. 52 (2012), 719–756, [doi:10.1007/s10589-011-9440-0](https://dx.doi.org/10.1007/s10589-011-9440-0).
- <span id="page-36-1"></span>[6] E. Casas and F. Tröltzsch, Numerical analysis of some optimal control problems governed by a class of quasilinear elliptic equations, ESAIM:COCV 17 (2011), 771-800, [doi:10.1051/cocv/2010025](https://dx.doi.org/10.1051/cocv/2010025).
- <span id="page-36-2"></span>[7] E. Casas and F. Tröltzsch, A general theorem on error estimates with application to a quasilinear elliptic optimal control problem, Comput. Optim. Appl. 53 (2012), 173–206, [doi:10.1007/](https://dx.doi.org/10.1007/s10589-011-9453-8) [s10589-011-9453-8](https://dx.doi.org/10.1007/s10589-011-9453-8).
- <span id="page-36-5"></span>[8] C. Christof and G. Wachsmuth, No-gap second-order conditions via a directional curvature functional, SIAM Journal on Optimization 28 (2018), 2097–2130, [doi:10.1137/17m1140418](https://dx.doi.org/10.1137/17M1140418).
- <span id="page-36-4"></span>[9] C. Clason, V. H. Nhu, and A. Rösch, No-gap second-order optimality conditions for optimal control of a non-smooth quasilinear elliptic equation, ESAIM: COCV 27 (2021), 62, [doi:10.1051/cocv/](https://dx.doi.org/10.1051/cocv/2020092) [2020092](https://dx.doi.org/10.1051/cocv/2020092).
- <span id="page-36-8"></span>[10] G. H. Cottet and E. Maitre, A level set method for fluid-structure interactions with immersed surfaces, Mathematical Models and Methods in Applied Sciences 16 (2006), 415–438, [doi:10.1142/](https://dx.doi.org/10.1142/S0218202506001212) [s0218202506001212](https://dx.doi.org/10.1142/S0218202506001212).
- <span id="page-36-12"></span>[11] K. Deckelnick and M. Hinze, A note on the approximation of elliptic control problems with bang-bang controls, Comput. Optim. Appl. 51 (2012), 931–939, [doi:10.1007/s10589-010-9365-z](https://dx.doi.org/10.1007/s10589-010-9365-z).
- <span id="page-36-9"></span>[12] L. C. Evans and R. F. Gariepy, Measure Theory and Fine Properties of Function, CRC Press, New York, 4th edition, 1992, [doi:10.1201/b18333](https://dx.doi.org/10.1201/b18333).
- <span id="page-36-11"></span>[13] A. Figalli, A simple proof of the Morse–Sard theorem in Sobolev spaces, Proceedings of the American Mathematical Society 136 (2008), 3675–3681, [doi:s0002-9939\(08\)09321-0](https://dx.doi.org/S0002-9939(08)09321-0).

- <span id="page-37-6"></span>[14] D. Gilbarg and N.S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin, Heidelberg, 2001, [doi:10.1007/978-3-642-61798-0](https://dx.doi.org/10.1007/978-3-642-61798-0).
- <span id="page-37-7"></span>[15] P. Grisvard, Elliptic Problems in Nonsmooth Domains, Pitman Advanced Pub. Program, 1985, [doi:](https://dx.doi.org/10.1137/1.9781611972030) [10.1137/1.9781611972030](https://dx.doi.org/10.1137/1.9781611972030).
- <span id="page-37-3"></span>[16] V. Guillemin and A. Pollack, Differential Topology, American Mathematical Society, Providence, RI, 1974.
- <span id="page-37-8"></span>[17] G. Leoni, A First Course in Sobolev Spaces, American Mathematical Society, Providence, RI, 2nd edition, 2017.
- <span id="page-37-9"></span>[18] J. Milnor, *Topology from the Differentiable Viewpoint*, University of Virginia Press, 1965.
- <span id="page-37-11"></span>[19] V. H. Nhu, On the no-gap second-order optimality conditions for a non-smooth semilinear elliptic optimal control, Optimization 71 (2022), 4289–4319, [doi:10.1080/02331934.2021.1939698](https://dx.doi.org/10.1080/02331934.2021.1939698).
- <span id="page-37-4"></span>[20] S. Osher and J. A. Sethian, Fronts propagating with curvature-dependent speed: Algorithms based on Hamilton-Jacobi formulations, J. Comput. Phys. 79 (1988), 12-49, [doi:10.1016/0021-9991\(88\)](https://dx.doi.org/10.1016/0021-9991(88)90002-2) [90002-2](https://dx.doi.org/10.1016/0021-9991(88)90002-2).
- <span id="page-37-10"></span>[21] W. Rudin, Principles of Mathematical Analysis, Mc Graw Hill Inc., New York, 3rd edition, 1976.
- <span id="page-37-2"></span>[22] A. Sard, The measure of the critical values of differentiable maps, Bull. Amer. Math. Soc. 48 (1942), 883–890.
- <span id="page-37-5"></span>[23] J. A. Sethian, Curvature and the evolution of fronts, Commun. Math. Phys. 101 (1985), 487–499, [doi:10.1007/bf01210742](https://dx.doi.org/10.1007/BF01210742).
- <span id="page-37-1"></span>[24] D. Wachsmuth and G. Wachsmuth, Second-order conditions for non-uniformly convex integrands: quadratic growth in L<sup>1</sup>, *J. Nonsmooth Anal. Optim.* 3 (2022), 8733, [doi:10.46298/jnsao-2022-8733](https://dx.doi.org/10.46298/jnsao-2022-8733).
- <span id="page-37-0"></span>[25] Y. B. Zel'dovich and Y. P. Raizer, Physics of Shock Waves and High-Temperature Hydrodynamic Phenomena, Academic Press, 1966, [doi:10.1115/1.3607836](https://dx.doi.org/10.1115/1.3607836).