

A New Type Distribution Dependent SDE for Singular Nonlinear PDE*

Feng-Yu Wang

Center for Applied Mathematics, Tianjin University, Tianjin 300072, China

Department of Mathematics, Swansea University, Bay Campus, SA1 8EN, United Kingdom

wangfy@tju.edu.cn

November 14, 2022

Abstract

We propose a new type SDE depending on the future distributions with all initial values, and establish the correspondence between this equation and the associated singular nonlinear PDE. Well-posedness and regularities are investigated.

AMS subject Classification: 60B05, 60B10, 35A01, 35D35.

Keywords: Distribution dependent SDE, nonlinear PDE, well-posedness.

1 Introduction

Probability theory has played a very important role in the study of partial differential equations (PDEs). Starting with the work of A. N. Kolmogorov [8], the probabilistic approach has been successfully applied in deducing scaling laws of turbulence in fluid (see [5] and references within). As a founder of stochastic calculus, K. Itô [7] developed stochastic differential equations (SDEs) to construct Markov processes characterizing linear Kolmogorov-Fokker-Planck equations. Since then several probability models have been introduced to solve nonlinear PDEs, which include the backward SDE initiated by Bismut [2] for the linear case and Pardoux-Peng [11] for the nonlinear case (see [3] and references within), the SDE driven by G -Brownian motion introduced by Peng [12] (see [13] and references within), and the distribution dependent (McKean-Vlasov) SDE introduced by McKean [10] (see [1] and references within). The first two models describe the viscosity solution, while the third one corresponds to the weak solution

*Supported in part by NNSFC (11831014, 11921001) and the National Key R&D Program of China (No. 2020YFA0712900).

(in the sense of integration by parts) of nonlinear PDEs. See also [4, 14] and references within for characterizations on the incompressible Navier-Stokes equation using SDEs.

In this paper, we investigate a class of new type distribution dependent SDEs and establish correspondence to nonlinear PDEs, such that the classical existence and uniqueness of singular nonlinear PDEs are derived.

Let $m, d \in \mathbb{N}$, and let $E = \mathbb{R}^d$ or $\mathbb{T}^d := \mathbb{R}^d/\mathbb{Z}^d$. For fixed $T \in (0, \infty)$, we consider the following equation for $u : [0, T] \times E \rightarrow \mathbb{R}^m$:

$$(1.1) \quad \partial_t u_t = (L_t + V_t)u_t + F_t(\cdot, u_t, \nabla u_t) \cdot \nabla u_t + g_t(\cdot, u_t, \nabla u_t, \nabla^2 u_t), \quad t \in [0, T],$$

where

$$L_t := \text{tr}\{a_t \nabla^2\} + b_t \cdot \nabla = \sum_{i,j=1}^d a_t^{ij} \partial_i \partial_j + \sum_{i=1}^d b_t^i \partial_i$$

is a time dependent second order differentiable operator on E with measurable coefficients

$$a : [0, T] \times E \rightarrow \mathbb{R}^{d \otimes d}, \quad b : [0, T] \times E \rightarrow \mathbb{R}^d,$$

$\nabla u_t = (\partial_i u_t^j)_{1 \leq i \leq d, 1 \leq j \leq m}$ takes values in $\mathbb{R}^{d \otimes m}$, $\nabla^2 u_t = (\partial_i \partial_j u_t^k)_{1 \leq i, j \leq d, 1 \leq k \leq m}$ takes values in $\mathbb{R}^{d \otimes d \otimes m}$, and

$$\begin{aligned} g : [0, T] \times E \times \mathbb{R}^m \times \mathbb{R}^{d \otimes m} \times \mathbb{R}^{d \otimes d \otimes m} &\rightarrow \mathbb{R}^m, \\ F : [0, T] \times E \times \mathbb{R}^m \times \mathbb{R}^{d \otimes m} &\rightarrow \mathbb{R}^d, \quad V : [0, T] \times E \rightarrow \mathbb{R} \end{aligned}$$

are measurable. Note that formally one may put $V_t u_t$ and $F_t \cdot \nabla u_t$ into g_t , but conditions we will use for these two terms are not covered by that for g_t , see Remark 2.1 below for details.

To characterize (1.1) using stochastic differential equation (SDE), we take $E = \mathbb{R}^d$ and let $g : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ only depend on the time-space variables. Denote

$$\sigma_t(x) := \sqrt{2a_{T-t}(x)}, \quad t \in [0, T], x \in \mathbb{R}^d,$$

let W_t be a d -dimensional Brownian motion with respect to a complete filtration probability space $(\Omega, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$, and consider the following SDE on \mathbb{R}^d where differentials are in $s \in [t, T]$:

$$(1.2) \quad \begin{aligned} dX_{t,s}^x &= \sigma_s(X_{t,s}^x) dW_s + \{b_{T-s} + F_{T-s}(\cdot, \psi_s, \nabla \psi_s)\}(X_{t,s}^x) ds, \\ \psi_s(y) &:= \mathbb{E} \left[u_0(X_{s,T}^y) e^{\int_s^T V_{T-r}(X_{s,r}^y) dr} + \int_s^T g_{T-r}(X_{s,r}^y) e^{\int_s^r V_{T-r'}(X_{s,r'}^y) dr'} dr \right], \quad y \in \mathbb{R}^d, \\ s &\in [t, T], X_{t,t}^x = x \in \mathbb{R}^d, t \in [0, T]. \end{aligned}$$

Since ψ_s depends on the distribution of $\{X_{s,r}^x\}_{(r,x) \in [s,T] \times E}$, the SDE (1.2) is distribution dependent for the future and for all initial values. This is essentially different from the usual McKean-Vlasov SDEs. Under certain conditions allowing coefficients to be singular in (t, x) , we prove the well-posedness of (1.2) and make application to (1.1). This extends the study of [16] where $V_t = 0$ and $F_t(\cdot, u_t, \nabla u_t) = -(u_t \cdot \nabla)u_t$ is considered. The present model also includes the nonlinear term $F_t(\cdot, u_t, \nabla u_t) = \beta |\nabla u_t|^2$ in the KPZ equation.

We will state the main results of the paper in Section 2, explain the idea of solving the new type SDE and present some lemmas in Section 3, and finally prove the main results in Sections 4 and 5.

2 Main results

We first state the main result on (1.2) and its link to PDE, then present the local and global well-posedness results on (1.1).

2.1 Distribution dependent SDE for nonlinear PDE

In this part, we let $E = \mathbb{R}^d$, and assume that $g_t(x, r_1, r_2, r_3) = g_t(x)$ depends only on (t, x) . Then the PDE (1.1) reduces to

$$(2.1) \quad \partial_t u_t = (L_t + V_t)u_t + F_t(x, u_t, \nabla u_t) \cdot \nabla u_t + g_t, \quad t \in [0, T].$$

Let $D_T := \{(t, s) : 0 \leq t \leq s \leq T\}$. We define the solution of (1.2) as follows.

Definition 2.1. A family $X := (X_{t,s}^x)_{(t,s,x) \in D_T \times \mathbb{R}^d}$ of random variables on \mathbb{R}^d is called a solution of (1.2), if $X_{t,s}^x$ is \mathcal{F}_s -measurable, $(\psi_s, \nabla \psi_s)$ exists such that

$$\mathbb{E} \int_t^T \left\{ |\sigma_s|^2 + |b_{T-s} + F_{T-s}(\cdot, \psi_s, \nabla \psi_s)| \right\} (X_{t,s}^x) ds < \infty$$

holds for $(t, x) \in [0, T] \times \mathbb{R}^d$, and \mathbb{P} -a.s.

$$X_{t,s}^x = x + \int_t^s \sigma_r(X_{t,r}^x) dW_r + \int_t^s \left\{ b_{T-r} + F_{T-r}(\cdot, \psi_r, \nabla \psi_r) \right\} (X_{t,r}^x) dr, \\ (t, s, x) \in D_T \times \mathbb{R}^d.$$

To prove the well-posedness of (1.2), we introduce some conditions which allow the coefficients to be singular in (t, x) . As we mentioned above that the coefficients of (1.2) may be singular in (t, x) . To measure the singularity, we recall some functional spaces introduced in [17].

For any $p, q > 1$ and $0 \leq t < s$, we write $f \in \tilde{L}_q^p(t, s)$ if f is a (real or vector valued) measurable function on $[t, s] \times E$ such that

$$\|f\|_{\tilde{L}_q^p(t,s)} := \sup_{z \in E} \left(\int_t^s \|f_r 1_{B(z,1)}\|_{L^p}^q dr \right)^{\frac{1}{q}} < \infty,$$

where $B(z, 1)$ is the unit ball at z , and $\|\cdot\|_{L^p}$ is the L^p -norm for the Lebesgue measure. When $f_t(x) = f(x)$ does not depend on t , let

$$\|f\|_{\tilde{L}^p} := \sup_{z \in E} \|f 1_{B(z,1)}\|_{L^p}.$$

When $E = \mathbb{T}^d$ which is compact, we may drop $1_{B(z,1)}$ such that \tilde{L}^p and \tilde{L}_q^p reduce to the usual L^p and L_q^p spaces.

We will take (p, q) from the following class:

$$\mathcal{H} := \left\{ (p, q) : p, q \in (2, \infty), \frac{d}{p} + \frac{2}{q} < 1 \right\}.$$

Let $\|\cdot\|_\infty$ denote the uniform norm, and let \mathcal{C}_b^1 be the class of (real or vector valued) weakly differentiable functions f on E such that

$$\|f\|_{\mathcal{C}_b^1} := \|f\|_\infty + \|\nabla f\|_\infty < \infty,$$

where $\|\nabla f\|_\infty$ coincides with the Lipschitz constant of f . Let

$$k(u_0, g, V) := e^{\int_0^T \|V_t\|_\infty dt} \left(\|u_0\|_\infty + \int_0^T \|g_t\|_\infty dt \right),$$

which might be infinite. When $k(u_0, g, V) < \infty$,

$$\bar{F}_t(x) := \sup_{|r_1| \leq k(u_0, g, V)} \|F_t(x, r_1, \cdot)\|_\infty$$

is finite provided $F_t(x, r_1, r_2)$ is bounded in r_2 but locally bounded in r_1 . We assume that V, u_0, a, b, F and g satisfy the following conditions.

(H_{V, u_0}) $u_0 \in \mathcal{C}_b^1$, and there exists $(p_0, q_0) \in \mathcal{K}$ such that $\|V\|_{\tilde{L}_{q_0}^{p_0}(0, T)} < \infty$.

$(H_{a, b})$ a is invertible and positive definite, b is locally bounded,

$$\sup_{t \in [0, T]} \left\{ \sup_{x \neq y} \frac{|b_t(x) - b_t(y)|}{|x - y|} + \|a_t\|_\infty + \|a_t^{-1}\|_\infty \right\} < \infty,$$

$$\lim_{\varepsilon \rightarrow 0} \sup_{|x - y| \leq \varepsilon, t \in [0, T]} |a_t(x) - a_t(y)| = 0,$$

and there exist $l \in \mathbb{N}$, $(p_i, q_i) \in \mathcal{K}$ and $1 \leq f_i \in \tilde{L}_{q_i}^{p_i}(0, T)$, $0 \leq i \leq l$, such that

$$|\nabla a_t(x)| \leq \sum_{i=1}^l f_i(t, x), \quad (t, x) \in [0, T] \times E.$$

$(H_{F, g}^0)$ $\|F(\cdot, 0, 0)\|_{\tilde{L}_{q_0}^{p_0}(0, T)} + \|g\|_{\tilde{L}_{q_0}^{p_0}(0, T)} < \infty$ for $(p_0, q_0) \in \mathcal{K}$ in (H_{V, u_0}) , and there exists a constant $K > 0$ such that

$$|F_t(x, r_1, r_2) - F_t(x, \tilde{r}_1, \tilde{r}_2)| \leq K \{1 \wedge (|r_1 - \tilde{r}_1| + |r_2 - \tilde{r}_2|)\},$$

$$(t, x) \in [0, T] \times \mathbb{R}^d, \quad |r_1| \vee |\tilde{r}_1| \leq k(u_0, g, V), \quad r_2, \tilde{r}_2 \in \mathbb{R}^{d \otimes m}.$$

Under these conditions, we will prove the well-posedness of (1.2). In this case, let

$$P_{t,s} f(x) := \mathbb{E}[f(X_{t,s}^x)], \quad f \in \mathcal{B}(\mathbb{R}^d),$$

$$P_{t,s}^V f(x) := \mathbb{E} \left[f(X_{t,s}^x) e^{\int_t^s V_{T-r}(X_{t,r}^x) dr} \right], \quad (t, s) \in D_T, \quad x \in \mathbb{R}^d, \quad f \in \mathcal{B}(\mathbb{R}^d),$$

$$u_t^V(f) := \int_t^T P_{t,s}^V f_s ds, \quad f \in \mathcal{B}([0, T] \times \mathbb{R}^d)$$

provided the expectations and integrals exist, where $\mathcal{B}(\cdot)$ is the class of measurable functions for a measurable space.

Theorem 2.1. Assume (H_{V,u_0}) , $(H_{a,b})$ and $(H_{F,g}^0)$. Then the following assertions hold.

- (1) The SDE (1.2) has a unique solution $X := (X_{t,s}^x)_{(t,s,x) \in D_T \times \mathbb{R}^d}$.
(2) For any $p \in [1, \infty)$ and $(t, x) \in [0, T] \times \mathbb{R}^d$, the derivative

$$\nabla_v X_{t,s}^x := \lim_{\varepsilon \downarrow 0} \frac{X_{t,s}^{x+\varepsilon v} - X_{t,s}^x}{\varepsilon}, \quad v \in \mathbb{R}^d, s \in [t, T]$$

exists in $L^p(\Omega \rightarrow C([t, T]; \mathbb{R}^d), \mathbb{P})$, and there exists a constant $c(p) > 0$ such that

$$(2.2) \quad \sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \mathbb{E} \left[\sup_{s \in [t,T]} |\nabla_v X_{t,s}^x|^p \right] \leq c(p) |v|^p, \quad v \in \mathbb{R}^d.$$

- (3) For any $0 \leq t < s \leq T$, $v \in \mathbb{R}^d$ and $f \in \mathcal{B}_b(\mathbb{R}^d)$,

$$(2.3) \quad \begin{aligned} \nabla_v P_{t,s} f(x) &:= \lim_{\varepsilon \downarrow 0} \frac{P_{t,s} f(x + \varepsilon v) - P_{t,s} f(x)}{\varepsilon} \\ &= \frac{1}{s-t} \mathbb{E} \left[f(X_{t,s}^x) \int_t^s \left\langle \sigma_r^{-1}(X_{t,r}^x) \nabla_v X_{t,r}^x, dW_r \right\rangle \right]. \end{aligned}$$

- (4) For any $p \in (1, \infty]$, there exists a constant $k_p > 0$ such that for any $0 \leq t < s \leq T$ and $f \in C_b^1(\mathbb{R}^d)$,

$$(2.4) \quad |\nabla P_{t,s} f| \leq k_p \min \left\{ (t-s)^{-\frac{1}{2}} (P_{t,s} |f|^p)^{\frac{1}{p}}, (P_{t,s} |\nabla f|^p)^{\frac{1}{p}} \right\}.$$

- (5) For any $p \in (1, \infty]$ and $V \in \tilde{L}_{q_0}^{p_0}(0, T)$, there exists a constant $c > 0$ determined by $(H_{a,b})$ and $\|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$, such that

$$(2.5) \quad \begin{aligned} \|P_{t,s}^V f\|_{\mathcal{C}_b^1} &\leq c \|f\|_{\mathcal{C}_b^1}, \quad f \in \mathcal{C}_b^1, (t, s) \in D_T, \\ \|\nabla P_{t,s}^V f\|_{\infty} &\leq c (s-t)^{-\frac{1}{2}} \|f\|_{\infty}, \quad 0 \leq t < s \leq T, f \in \mathcal{B}_b(\mathbb{R}^d), \\ \|u_t^V(f)\|_{\infty} + \|\nabla^2 u^V(f)\|_{\tilde{L}_{q_0}^{p_0}(t,T)} &\leq c \|f\|_{\tilde{L}_{q_0}^{p_0}(t,T)}, \quad t \in [0, T], f \in \tilde{L}_{q_0}^{p_0}(0, T). \end{aligned}$$

The next result provides a correspondence between solutions of (2.1) and (1.2).

Theorem 2.2. Assume (H_{V,u_0}) , $(H_{a,b})$ and $(H_{F,g}^0)$.

- (1) If $(u_t)_{t \in [0,T]}$ solves (2.1) in the class $\mathcal{U}(p_0, q_0)$, then

$$(2.6) \quad \begin{aligned} u_t(x) &= \mathbb{E} \left[u_0(X_{T-t,T}^x) e^{\int_{T-t}^T V_{T-r}(X_{T-t,r}^x) dr} \right] \\ &+ \mathbb{E} \int_{T-t}^T g_{T-s}(X_{T-t,s}^x) e^{\int_{T-t}^s V_{T-r}(X_{T-t,r}^x) dr} ds, \quad (t, x) \in [0, T] \times \mathbb{R}^d. \end{aligned}$$

- (2) On the other hand, u_t given by (2.6) solves (2.1) such that

$$\|u_t\|_{\mathcal{C}_b^1} + \|u\|_{\tilde{L}_{q_0}^{p_0}(t,T)} \leq c (\|u_0\|_{\mathcal{C}_b^1} + \|g\|_{\tilde{L}_{q_0}^{p_0}(0,t)}), \quad t \in [0, T]$$

holds for some constant $c > 0$ determined by $(H_{a,b})$ and $\|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$.

2.2 Well-posedness of (1.1)

In this part, we let $E = \mathbb{R}^d$ or \mathbb{T}^d , and allow $g_t(x, u, \nabla u, \nabla^2 u)$ depending on $(u, \nabla u, \nabla^2 u)$. We will prove the existence and uniqueness of (1.1) in the following sense.

Definition 2.2. We call $(u_t)_{t \in [0, T^*)}$ a solution of (1.1) in the class $\mathcal{U}(p_0, q_0)$, if

(1) $T^* \in (0, T]$,

$$(2.7) \quad \sup_{t \in [0, s]} \|u_t\|_{\mathcal{C}_b^1} + \|\nabla^2 u\|_{\tilde{L}_{q_0}^{p_0}(0, s)} < \infty, \quad s \in (0, T^*),$$

and for any $t \in [0, T^*)$,

$$u_t = u_0 + \int_0^t \left\{ (L_s + V_s)u_s + F_s(\cdot, u_s, \nabla u_s) \cdot \nabla u_s + g_s(\cdot, u_s, \nabla u_s, \nabla^2 u_s) \right\} ds.$$

(2) When $T^* < T$, $\limsup_{t \uparrow T^*} \|u_t\|_{\mathcal{C}_b^1} = \infty$.

Instead of $(H_{F, g}^0)$, we will use the following weaker conditions. Let

$$\bar{g}_t(x) = |g_t(x, 0, 0, 0)|, \quad (t, x) \in [0, T] \times E.$$

$(H_{F, g})$ $\|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0, T)} + \|\bar{g}\|_{\tilde{L}_{q_0}^{p_0}(0, T)} < \infty$ for (p_0, q_0) in (H_{V, u_0}) , and there exist a constant $\alpha > 0$ and a map $K : \mathbb{N} \rightarrow (0, \infty)$ such that

$$\begin{aligned} & |F_t(x, r_1, r_2) - F_t(x, \tilde{r}_1, \tilde{r}_2)| + |g_t(x, r_1, r_2, r_3) - g_t(x, \tilde{r}_1, \tilde{r}_2, \tilde{r}_3)| \\ & \leq K_n(|r_1 - \tilde{r}_1| + |r_2 - \tilde{r}_2|) + \alpha|r_3 - \tilde{r}_3|, \\ & (t, x) \in [0, T] \times E, \quad |r_1| \vee |r_2| \vee |\tilde{r}_1| \vee |\tilde{r}_2| \leq n, \quad r_3, \tilde{r}_3 \in \mathbb{R}^{d \otimes d \otimes m}. \end{aligned}$$

$(H'_{F, g})$ There exist constants $C, \alpha > 0$ and $\theta \in (0, \frac{1}{2})$ such that

$$\begin{aligned} & \sup_{(t, x) \in [0, T] \times E} |g_t(x, r_1, r_2, r_3) - g_t(x, 0, 0, 0)| \\ & \leq C(e + |r_1| + |r_2|) \{\log(e + |r_1| + |r_2|)\}^\theta + \alpha|r_3|, \\ & (r_1, r_2, r_3) \in \mathbb{R}^m \times \mathbb{R}^{d \otimes m} \times \mathbb{R}^{d \otimes d \otimes m}. \end{aligned}$$

We have the following result.

Theorem 2.3. Assume (H_{V, u_0}) , $(H_{a, b})$ and $(H_{F, g})$ with small enough $\alpha > 0$ determined by $(H_{a, b})$ and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0, T)} + \|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0, T)}$.

(1) (1.1) has a unique solution in $\mathcal{U}(p_0, q_0)$.

(2) If $(H'_{F, g})$ holds with small enough $\alpha > 0$ determined by $(H_{a, b})$ and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0, T)} + \|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0, T)}$, then $T^* = T$ and

$$(2.8) \quad \sup_{t \in [0, T]} \|u_t\|_{\mathcal{C}_b^1} + \|\nabla^2 u\|_{\tilde{L}_{q_0}^{p_0}(0, T)} < \infty.$$

Remark 2.1. We do not combine Vu and $F \cdot \nabla u$ with g , since in Theorem 2.3 conditions on V and F are not covered by that on g . To see this, we assume $(H_{a,b})$, let $F_t(x, r_1, r_2)$ and $g_t(x, r_1, r_2, r_3) = g_t(x, r_1, r_2)$ be locally Lipschitz continuous in (r_1, r_2) , and let $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} < \infty$. We consider the following two situations.

- (a) Let for instance F and g be bounded. Then Theorem 2.3 implies the well-posedness and non-explosion of (1.1). However, since $|V_t|$ may be unbounded, $(H'_{F,g})$ does not hold if we regard $V_t r_1$ as a part of $g_t(\cdot, r_1, r_2)$.
- (b) Let for instance $\|g\|_\infty + \int_0^T \|V_t\|_\infty < \infty$ and $F_t(\cdot, r_1, r_2) = \beta(r_1)h_t$ for some locally bounded function β on \mathbb{R}^m and some \mathbb{R}^d -valued $h \in \tilde{L}_{q_0}^{p_0}(0, T)$. Then $k(u_0, V, g) < \infty$, so that

$$\|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)} \leq \|h\|_{\tilde{L}_{q_0}^{p_0}(0,T)} \sup_{|r_1| \leq k(u_0, V, g)} |\beta(r_1)| < \infty.$$

Hence, Theorem 2.3 implies the the well-posedness and non-explosion of (1.1). However, since $|\beta(r_1)|$ may have arbitrary growth in $|r_1|$, $(H'_{F,g})$ does not hold if we regard $\beta_t(r_1)h_t \cdot r_2$ as a part of $g_t(\cdot, r_1, r_2)$.

To illustrate Theorem 2.3, we consider the following equation (2.9). When $F = 0$ it reduces to the KPZ type equation

$$\partial_t u_t = L_t u_t + \beta \langle a_t \nabla u_t, \nabla u_t \rangle + \bar{V}_t,$$

and for $\beta = 0$ it becomes the Navier-Stokes type equation

$$\partial_t u_t = L_t u_t + F_t(\cdot, u_t) \cdot \nabla u_t + \bar{V}_t.$$

Example 2.1. Assume $(H_{a,b})$, let $\beta \in \mathbb{R}$, and let

$$F : [0, \infty) \times E \times \mathbb{R}^m \rightarrow \mathbb{R}^d, \quad V : [0, \infty) \times E \rightarrow \mathbb{R}$$

be measurable such that

$$\tilde{F}_t(x, r) := \sup_{x \in E} |F_t(x, r) - F_t(x, 0)|$$

is locally bounded in $(t, r) \in [0, \infty) \times \mathbb{R}^m$, and for some $(p_0, q_0) \in \mathcal{K}$

$$\|F(\cdot, 0)\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} < \infty, \quad T \in (0, T).$$

Consider the PDE

$$(2.9) \quad \partial_t u_t = L_t u_t + \beta \langle a_t \nabla u_t, \nabla u_t \rangle + F_t(\cdot, \varphi_\beta(u_t)) \cdot \nabla u_t + \bar{V}_t, \quad t \geq 0,$$

where $\langle a_t \nabla u_t, \nabla u_t \rangle := (\langle a_t \nabla u_t^i, \nabla u_t^i \rangle)_{1 \leq i \leq m}$, $\bar{V}_t(x) := (V_t(x), \dots, V_t(x)) \in \mathbb{R}^m$, and

$$\varphi_\beta(u_t) := (\beta^{-1}(1 - e^{-\beta u_t^i}))_{1 \leq i \leq m}$$

which reduces to u_t when $\beta = 0$. If either $\|V_t\|_\infty$ or $\|\tilde{F}_t\|_\infty$ is locally integrable in $t \geq 0$, then for any $u_0 \in \mathcal{C}_b^1$, (2.9) has a unique solution $u : [0, \infty) \times E \rightarrow \mathbb{R}^m$ satisfying

$$\sup_{t \in [0, T]} \|\varphi_\beta(u_t)\|_{\mathcal{C}_b^1} + \|\nabla^2 \varphi_\beta(u)\|_{\tilde{L}_{q_0}^{p_0}(0,T)} < \infty, \quad T \in (0, \infty).$$

Proof. It suffices to prove the assertion up to an arbitrarily fixed time $T \in [0, \infty)$. We only consider $\beta \neq 0$, as when $\beta = 0$ the desired assertion follows from Theorem 2.3 for $V = 0$ and $g = \bar{V}$. Let $v_t = e^{-\beta u t} := (e^{-\beta u_i^i})_{1 \leq i \leq m}$. Then (2.9) up totime T s equivalent to

$$(2.10) \quad \partial_t v_t = (L_t - \beta V_t)v_t + F_t(\cdot, \beta^{-1}(1 - v_t)) \cdot \nabla v_t, \quad t \in [0, T], \quad v_0 = e^{-\beta u_0}.$$

If either $\|V_t\|_\infty$ or $\|\tilde{F}_t\|_\infty$ is locally integrable in $t \geq 0$, we have $\|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0, T)} < \infty$ for

$$\bar{F}_t(x) := \sup_{|r| \leq k_0} |F_t(x, \beta^{-1}(1 - r))|, \quad k_0 = k(u_0, -\beta V, 0) := \|u_0\|_\infty e^{\int_0^T \|\beta V_t\|_\infty dt}.$$

By Theorem 2.3 with $g = 0$ and $-\beta V_t$ replacing V_t , (2.10) has a unique solution satisfying

$$\sup_{t \in [0, T]} \|v_t\|_{\mathcal{C}_b^1} + \|\nabla^2 v_t\|_{\tilde{L}_{q_0}^{p_0}(0, T)} < \infty.$$

Then the proof is finished. □

3 Some lemmas

We first explain the idea of solving (1.2) by fixed point theorem.

Let $\mathcal{C}_T := C(D_T; \mathbb{R}^d)$ be the space of all maps

$$\xi = (\xi_{t,s})_{(t,s) \in D_T} : D_T \rightarrow \mathbb{R}^d; \quad \xi_{t,s} \text{ is continuous in } (t, s) \in D_T.$$

It is a Banach space under the uniform norm $\|\cdot\|_\infty$.

Let \mathcal{P}_T be the space of all probability measures on \mathcal{C}_T equipped with the weak topology. We define Γ as the set of all measurable maps

$$\gamma = (\gamma^x)_{x \in \mathbb{R}^d} : \mathbb{R}^d \rightarrow \mathcal{P}_T$$

such that

$$(3.1) \quad \begin{aligned} \psi_s^\gamma(x) &:= \int_{\mathcal{C}_T} u_0(\xi_{s,T}) e^{\int_s^T V_{T-s}(\xi_{s,r}) dr} \gamma^x(d\xi) \\ &+ \int_s^T dr \int_{\mathcal{C}_T} g_{T-r}(\xi_{s,r}) e^{\int_s^r V_{T-r'}(\xi_{s,r'}) dr'} \gamma^x(d\xi) \end{aligned}$$

is well-defined and the weak gradient $\nabla \psi_s^\gamma$ exists for any $s \in [0, T]$.

We will see that for any $\gamma \in \Gamma$, the classical SDE

$$(3.2) \quad dX_{t,s}^{\gamma,x} = \{b_{T-s} + F_{T-s}(\cdot, \psi_s^\gamma, \nabla \psi_s^\gamma)\}(X_{t,s}^{\gamma,x}) ds + \sigma_s(X_{t,s}^{\gamma,x}) dW_s, \quad s \in [t, T], \quad X_{t,t}^{\gamma,x} = x$$

is well-posed. In this case, we define a map

$$(3.3) \quad \gamma \mapsto \Phi(\gamma); \quad x \mapsto \Phi^x(\gamma) := \mathcal{L}_{X^{\gamma,x}},$$

where $\mathcal{L}_{X^{\gamma,x}}$ is the distribution of the \mathcal{C}_T -valued random variable $X^{\gamma,x} := (X_{t,s}^{\gamma,x})_{(t,s) \in D_T}$. It is clear that $X_{t,s}^{\gamma,x}$ solves (1.2) if and only if γ is a fixed point of Φ , i.e. $\Phi(\gamma) = \gamma$. So, for the well-posedness of (1.2), it suffices to show that Φ has a unique fixed point.

To this end, we construct a non-empty subspace $\tilde{\Gamma}$ of Γ , such that Φ is a contractive map on $\tilde{\Gamma}$ under a complete metric. According to the definition of ψ_t^γ , the space $\tilde{\Gamma}$ should be determined by the following functionals: for any $(t, s, x) \in D_T \times \mathbb{R}^d$ and $\gamma \in \Gamma$, let

$$\begin{aligned} \gamma_{t,s}^V(f)(x) &:= \int_{\mathcal{C}_T} f(\xi_{t,s}) e^{\int_t^s V_{T-r}(\xi_{t,r}) dr} \gamma^x(d\xi), \quad f \in \mathcal{B}(\mathbb{R}^d), \\ u_t^{\gamma,V}(f)(x) &:= \int_t^T \gamma_{t,s}^V(f_s)(x) ds \\ &= \int_t^T ds \int_{\mathcal{C}_T} f_s(\xi_{t,s}) e^{\int_t^s V_{T-r}(\xi_{t,r}) dr} \gamma^x(d\xi), \quad f \in \mathcal{B}([t, T] \times \mathbb{R}^d), \end{aligned}$$

provided the integrals exist. The class $\tilde{\Gamma}$ is defined as follows.

Definition 3.1. Let $\tilde{\Gamma}$ be the set of $\gamma \in \Gamma$ satisfying $\gamma_{t,t}^x = \delta_x$ (the Dirac measure at x) for $(t, x) \in [0, T] \times \mathbb{R}^d$, and

$$\begin{aligned} \sup_{t \in [0, T]} \|\gamma_t\|_V < \infty, \quad \|\gamma_t\|_V &:= \sup_{x \in \mathbb{R}^d} \int_{\mathcal{C}_T} e^{\int_t^T |V_{T-r}(\xi_{t,r})| dr} \gamma^x(d\xi) < \infty, \\ \sup_{t \in [0, T]} \left\{ \sup_{\|f\|_{\mathcal{C}_b^1} \leq 1} \|\nabla \gamma_{t,T}^V(f)\|_\infty + \sup_{\|f\|_{\tilde{L}_{q_0}^{p_0}(0, T)} \leq 1} \|u_t^{\gamma,V}(f)\|_{\mathcal{C}_b^1} \right\} &< \infty. \end{aligned}$$

To apply the fixed point theorem to Φ , we introduce a complete metric on $\tilde{\Gamma}$, which consists of three parts: one is the weighted variation distance induced by

$$\eta_t^V(\xi) := e^{\int_t^T |V_{T-r}(\xi_{t,r})| dr}, \quad \xi \in C([t, T]; \mathbb{R}^d),$$

and the other two are induced by the above functionals $\gamma_{t,T}^V$ and $u_t^{\gamma,V}$. More precisely, for any $\gamma \in \Gamma$ and $t \in [0, T]$, let $\gamma_t = (\gamma_t^x)_{x \in \mathbb{R}^d}$ be the marginal distribution of $\gamma = (\gamma^x)_{x \in \mathbb{R}^d}$ for

$$\xi \mapsto \xi_t := (\xi_{t,s})_{s \in [t, T]} \in C([t, T]; \mathbb{R}^d),$$

i.e. $\gamma_t^x := \gamma^x \circ \xi_t^{-1}$, $x \in \mathbb{R}^d$. Then

$$\gamma_t(f)(x) := \int_{C([t, T]; \mathbb{R}^d)} f(\xi) \gamma_t^x(d\xi) = \int_{\mathcal{C}_T} f(\xi_t) \gamma^x(d\xi), \quad x \in \mathbb{R}^d, \quad f \in \mathcal{B}_b(C([t, T]; \mathbb{R}^d)).$$

For any $\gamma, \tilde{\gamma} \in \tilde{\Gamma}$ and $t \in [0, T]$, let

$$\|\gamma_t - \tilde{\gamma}_t\|_V := \sup_{f \in \mathcal{B}_b(C([t, T]; \mathbb{R}^d)), \|f\|_\infty \leq \eta_t^V} \|\gamma_t(f) - \tilde{\gamma}_t(f)\|_\infty,$$

$$\begin{aligned}\|\gamma_t - \tilde{\gamma}_t\|_{\mathcal{C}_b^1} &:= \sup_{\|f\|_{\mathcal{C}_b^1} \leq 1} \|\nabla\{\gamma_{t,T}^V(f) - \tilde{\gamma}_{t,T}^V(f)\}\|_\infty, \\ \|\gamma_t - \tilde{\gamma}_t\|_{p_0, q_0} &:= \sup_{\|f\|_{\tilde{L}_{q_0}^{p_0}(0,T)} \leq 1} \|u_t^{\gamma, V}(f) - u_t^{\tilde{\gamma}, V}(f)\|_{\mathcal{C}_b^1}.\end{aligned}$$

We will prove that for any $\lambda \geq 0$, $\tilde{\Gamma}$ is complete under the metric

$$\rho_\lambda(\gamma, \tilde{\gamma}) := \sup_{(t,s) \in D_T} e^{-\lambda(T-t)} \{ \|\gamma_t - \tilde{\gamma}_t\|_V + \|\gamma_t - \tilde{\gamma}_t\|_{\mathcal{C}_b^1} + \|\gamma_t - \tilde{\gamma}_t\|_{p_0, q_0} \},$$

and that Φ is contractive in ρ_λ for large $\lambda > 0$, so that it has a unique fixed point in $\tilde{\Gamma}$. To this end, we present below some lemmas.

We first extend [18, Theorem 2.1] with $V = 0$ to the the present setting. For any $p, q \geq 1$, let $\tilde{H}_q^{2,p}(0, T)$ be the space of $f : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ such that

$$\|f\|_{\tilde{L}_q^p(0,T)} + \|\nabla f\|_{\tilde{L}_q^p(0,T)} + \|\nabla^2 f\|_{\tilde{L}_q^p(0,T)} < \infty.$$

Lemma 3.1. *Assume $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} < \infty$ and $(H_{a,b})$ but without the condition on $|\nabla a|$. Let $b^{(0)} : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ with $\|b^{(0)}\|_{\tilde{L}_q^p(0,T)} < \infty$ for some $(p, q) \in \mathcal{K}$, and let*

$$L_t^V := L_{T-t} + b_t^{(0)} \cdot \nabla + V_{T-t}, \quad t \in [0, T].$$

Then for any $\lambda \geq 0$ and $f \in \tilde{L}_{q_0}^{p_0}(0, T)$, the PDE

$$(3.4) \quad (\partial_t + L_t^V - \lambda)u_t^\lambda = f_t, \quad t \in [0, T], u_T^\lambda = 0$$

has a unique solution in $\tilde{H}_{q_0}^{2,p_0}(0, T)$. Moreover, for any $0 < \varepsilon < \frac{1}{2}(1 - \frac{d}{p_0} - \frac{2}{q_0})$, there exists a constant $c > 0$ increasing in $\|b^{(0)}\|_{\tilde{L}_q^p(0,T)}$ such that

$$(3.5) \quad \begin{aligned}(1 + \lambda)^\varepsilon (\|u_t^\lambda\|_\infty + \|\nabla u_t^\lambda\|_\infty) + \|\nabla^2 u^\lambda\|_{\tilde{L}_{q_0}^{p_0}(t,T)} &\leq c \|f\|_{\tilde{L}_{q_0}^{p_0}(t,T)}, \\ \lambda \geq 0, f \in \tilde{L}_{q_0}^{p_0}(0, T), t \in [0, T].\end{aligned}$$

Proof. (a) Let $L_t^0 = L_{T-t} + b_t^{(0)} \cdot \nabla$. By [18, Theorem 2.1], for any $\lambda \geq 0$ and $f \in \tilde{L}_{q_0}^{p_0}(0, T)$, the PDE

$$(3.6) \quad (\partial_t + L_t^0 - \lambda)\hat{u}_t^\lambda = f_t, \quad \hat{u}_T^\lambda = 0$$

has a unique solution in the $\tilde{H}_{q_0}^{2,p_0}(0, T)$, and for any $0 < \varepsilon < \frac{1}{2}(1 - \frac{d}{p_0} - \frac{2}{q_0})$, there exists a constant $c_1 > 0$ increasing in $\|b^{(0)}\|_{\tilde{L}_q^p(0,T)}$ such that

$$(3.7) \quad \begin{aligned}(1 + \lambda)^\varepsilon \{ \|\hat{u}_t^\lambda\|_\infty + \|\nabla \hat{u}_t^\lambda\|_\infty \} + \|\nabla^2 \hat{u}^\lambda\|_{\tilde{L}_{q_0}^{p_0}(t,T)} \\ \leq c_1 \|f\|_{\tilde{L}_{q_0}^{p_0}(t,T)}, \quad \lambda \geq 0, f \in \tilde{L}_{q_0}^{p_0}(0, T), t \in [0, T].\end{aligned}$$

Note that in [18, Theorem 2.1] the class $\tilde{H}_{q_0}^{2,p_0}(0, T)$ is replaced by $\tilde{W}_{1,q_0}^{2,p_0,w}(0, T)$, which consists of $f \in \tilde{H}_{q_0}^{2,p_0}(0, T)$ such that for $w(x) := (1+|x|)^{-1}$ (note that $w(x)$ therein should be $(1+|x|^2)^{-\frac{p_0}{2}}$ according to [18, Lemma 2.3]),

$$(3.8) \quad |wf| + |w\nabla f| + |w\nabla^2 f| + |w\partial_t f| \in \tilde{L}_{q_0}^{p_0}(0, T).$$

Since $w \leq 1$ and $(H_{a,b})$ implies that a is bounded and $|b_t(x)| \leq k(1+|x|)$ for some constant $k > 0$, a solution of (3.6) in $\tilde{H}_{q_0}^{p_0}(0, T)$ must satisfy (3.8), so we may replace $\tilde{W}_{1,q_0}^{2,p_0,w}(0, T)$ by $\tilde{H}_{q_0}^{2,p_0}(t, T)$.

(b) To solve (3.4), for any $\eta \in \mathcal{B}_b([0, T] \times \mathbb{R}^d)$, consider the PDE

$$(3.9) \quad (\partial_t + L_t^0 - \lambda)u_t^{\lambda,\eta} = f_t - V_{T-t}\eta_t, \quad u_T^{\lambda,\eta} = 0.$$

Let $\alpha(V) = \|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$. We have

$$\|f - V_{T-\cdot}\eta\|_{\tilde{L}_{q_0}^{p_0}(0,T)} \leq \|f\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \alpha(V)\|\eta\|_\infty.$$

Then (a) implies that (3.9) has a unique solution in $\tilde{H}_{q_0}^{2,p_0}(0, T) \cap \mathcal{B}_b([0, T] \times \mathbb{R}^d)$. Thus, we obtain a map

$$\mathcal{B}_b([0, T] \times \mathbb{R}^d) \ni \eta \mapsto u^{\lambda,\eta} \in \mathcal{B}_b([0, T] \times \mathbb{R}^d).$$

So, to see that (3.4) has a unique solution in $\tilde{H}_{q_0}^{p_0}(0, T) \cap \mathcal{B}_b([0, T] \times \mathbb{R}^d)$, it suffices to prove that this map has a unique fixed point.

To this end, for any $\eta, \tilde{\eta} \in \mathcal{B}_b([0, T] \times \mathbb{R}^d)$, let $u^{\lambda,\eta,\tilde{\eta}} := u^{\lambda,\eta} - u^{\lambda,\tilde{\eta}}$. Then

$$(\partial_t + L_t^0 - \lambda)u_t^{\lambda,\eta,\tilde{\eta}} = V_{T-t}(\tilde{\eta}_t - \eta_t), \quad u_T^{\lambda,\eta,\tilde{\eta}} = 0.$$

Applying (3.7) to this equation, by (a) we find a constant $c_1 > 0$ such that

$$\|u_t^{\lambda,\eta} - u_t^{\lambda,\tilde{\eta}}\|_\infty = \|u_t^{\lambda,\eta,\tilde{\eta}}\|_\infty \leq c_1 \|V_{T-t}(\tilde{\eta} - \eta)\|_{\tilde{L}_{q_0}^{p_0}(t,T)}.$$

Letting

$$\|\eta - \tilde{\eta}\|_{\infty,n} := \sup_{t \in [0,T]} e^{-n(T-t)} \|\eta_t - \tilde{\eta}_t\|_\infty, \quad n \geq 1,$$

this implies

$$\|u^{\lambda,\eta} - u^{\lambda,\tilde{\eta}}\|_{\infty,n} \leq c_1 \|\eta - \tilde{\eta}\|_{\infty,n} \sup_{t \in [0,T]} \|V_{T-\cdot} e^{-n(\cdot-t)}\|_{\tilde{L}_{q_0}^{p_0}(t,T)}.$$

Hence, when n is large enough, the map $u^{\lambda,\cdot}$ is contractive in the complete metric $\|\eta - \tilde{\eta}\|_{\infty,n}$. Therefore, it has a unique fixed point as desired.

(c) It remains to prove the estimate (3.5). Let u^λ be the unique solution of (3.4) in $\tilde{H}_{q_0}^{p_0}(0, T)$. By (3.7) we obtain

$$(3.10) \quad \begin{aligned} & (1 + \lambda)^\varepsilon (\|u_t^\lambda\|_\infty + \|\nabla u_t^\lambda\|_\infty) + \|\nabla^2 u^\lambda\|_{\tilde{L}_{q_0}^{p_0}(t,T)} \\ & \leq c_1 \|f - V_{T-\cdot} u^\lambda\|_{\tilde{L}_{q_0}^{p_0}(t,T)} \leq c_1 \|f\|_{\tilde{L}_{q_0}^{p_0}(t,T)} + c_1 \|V_{T-\cdot} u^\lambda\|_{\tilde{L}_{q_0}^{p_0}(t,T)}, \quad t \in [0, T]. \end{aligned}$$

Letting

$$\|u^\lambda\|_{\infty,n} := \sup_{s \in [0,T]} e^{-n(T-s)} \|u_s^\lambda\|_\infty, \quad n \geq 1,$$

we derive

$$e^{-n(T-t)} \|u_t^\lambda\|_\infty \leq c_1 \|f\|_{\tilde{L}_{q_0}^{p_0}(t,T)} + c_1 \|u^\lambda\|_{\infty,n} \|V_{T-} e^{-n(\cdot-t)}\|_{\tilde{L}_{q_0}^{p_0}(t,T)}.$$

When $n \geq 1$ is large enough, this implies

$$\|u^\lambda\|_{\infty,n} \leq c_1 \|f\|_{\tilde{L}_{q_0}^{p_0}(t,T)} + \frac{1}{2} \|u^\lambda\|_{\infty,n}, \quad t \in [0, T].$$

Therefore, there exists a constant $c_2 > 0$ such that $\|u_t^\lambda\|_\infty \leq c_2 \|f\|_{\tilde{L}_{q_0}^{p_0}(t,T)}$, $t \in [0, T]$. Combining this with (3.10) and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} < \infty$, we finish the proof. \square

Lemma 3.2. *Assume (H_{V,u_0}) , $(H_{a,b})$ and $(H_{F,g}^0)$. Then for any $\gamma \in \Gamma$, the SDE (3.2) is well-posed, and $\Phi(\gamma)$ defined in (3.3) is a measurable map from \mathbb{R}^d to \mathcal{P}_T . Moreover:*

(1) For any $v \in \mathbb{R}^d$,

$$\nabla_v X_{t,s}^{\gamma,x} := \lim_{\varepsilon \downarrow 0} \frac{X_{t,s}^{\gamma,x+\varepsilon v} - X_{t,s}^{\gamma,x}}{\varepsilon}, \quad s \in [t, T]$$

exists in $L^p(\Omega \rightarrow C([t, T]; \mathbb{R}^d), \mathbb{P})$ for any $p \in (1, \infty)$, there exists a constant $k_p > 0$ determined by p , $(H_{a,b})$ and $\|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$, such that

$$(3.11) \quad \sup_{\gamma \in \Gamma} \sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \mathbb{E} \left[\sup_{s \in [t,T]} |\nabla_v X_{t,s}^{\gamma,x}|^p \right] \leq k_p |v|^p, \quad v \in \mathbb{R}^d,$$

(2) For any $p \in (1, \infty]$ there exists a constant $k_p > 0$ determined by p , $(H_{a,b})$ and $\|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$, such that

$$P_{t,s}^\gamma f(x) := \mathbb{E}[f(X_{t,s}^{\gamma,x})], \quad (t,s) \in D_T, x \in \mathbb{R}^d, f \in \mathcal{B}_b(\mathbb{R}^d)$$

satisfies

$$(3.12) \quad |\nabla P_{t,s}^\gamma f| \leq k_p \min \left\{ (s-t)^{-\frac{1}{2}} (P_{t,s}^\gamma |f|^p)^{\frac{1}{p}}, (P_{t,s}^\gamma |\nabla f|^p)^{\frac{1}{p}} \right\}, \\ \gamma \in \Gamma, 0 \leq t < s \leq T, f \in C_b^1(\mathbb{R}^d),$$

and for any $v, x \in \mathbb{R}^d$, any $0 \leq t < s \leq T$,

$$(3.13) \quad \nabla_v P_{t,s}^\gamma f(x) = \frac{1}{s-t} \mathbb{E} \left[f(X_{t,s}^{\gamma,x}) \int_t^s \left\langle \sigma_r^{-1}(X_{t,r}^{\gamma,x}) \nabla_v X_{t,r}^{\gamma,x}, dW_r \right\rangle \right], \quad f \in \mathcal{B}_b(\mathbb{R}^d).$$

Proof. By $(H_{F,g}^0)$, $b_t^\gamma := F_{T-t}(\cdot, \psi_t^\gamma, \nabla \psi_t^\gamma)$ satisfies

$$(3.14) \quad \sup_{\gamma \in \Gamma} \|b^\gamma\|_{\tilde{L}_{q_0}^{p_0}(0,T)} \leq \|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)} < \infty.$$

According to [15, Theorem 2.1], $(H_{a,b})$ and (3.14) imply that the SDE (3.2) is well-posed, and (3.11), (3.12) and (3.13) hold. Noting that the Borel σ -field on \mathcal{P}_T is induced by the maps

$$\mathcal{C}_T \ni \mu \mapsto \mu_{t,s}(f) := \int_{\mathcal{C}_T} f(\xi_{t,s}) \mu(d\xi), \quad (t,s) \in D_T, f \in C_b(\mathbb{R}^d),$$

by combining (3.11) with the continuity of $X_{t,s}^{\gamma,x}$ in $(t,s) \in D_T$, we see that $\Phi(\gamma)$ is a measurable map from \mathbb{R}^d to \mathcal{P}_T . \square

In the following, we intend to prove $\Phi\Gamma \subset \tilde{\Gamma}$, so that we may apply the fixed point theorem to the map $\Phi : \tilde{\Gamma} \rightarrow \tilde{\Gamma}$. To this end, we introduce the Feynman-Kac semigroup

$$P_{t,s}^{\gamma,V} f(x) := \mathbb{E} \left[f(X_{t,s}^{\gamma,x}) e^{\int_t^s V_{T-r}(X_{t,r}^{\gamma,x}) dr} \right], \quad (t,s,x) \in D_T \times \mathbb{R}^d, f \in \mathcal{B}_b(\mathbb{R}^d),$$

where $X_{t,s}^{\gamma,x}$ solves (3.2) with generator

$$(3.15) \quad \tilde{L}_t^\gamma := L_{T-t} + F_{T-t}(\cdot, \psi_t^\gamma, \nabla \psi_t^\gamma) \cdot \nabla, \quad t \in [0, T].$$

Lemma 3.3. *Assume (H_{V,u_0}) , $(H_{a,b})$ and $(H_{F,g}^0)$. Then the following assertions hold.*

- (1) *There exists a constant $c > 0$ determined by $(H_{a,b})$ and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$, such that for any $\gamma \in \Gamma$, any $f : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ with $\|f\|_{\tilde{L}_{q_0}^{p_0}(0,T)} < \infty$,*

$$u_t^{\Phi(\gamma),V}(f) := \mathbb{E} \int_t^T f_s(X_{t,s}^x) e^{\int_t^s V_{T-r}(X_{t,r}^x) dr} ds = \int_t^T P_{t,s}^{\gamma,V} f_s ds, \quad t \in [0, T]$$

satisfies

$$(3.16) \quad (\partial_t + \tilde{L}_t^\gamma + V_{T-t}) u_t^{\Phi(\gamma),V}(f) = -f_t, \quad t \in [0, T],$$

$$(3.17) \quad \|u_t^{\Phi(\gamma),f}\|_\infty + \|\nabla u_t^{\Phi(\gamma),f}\|_\infty + \|\nabla^2 u_t^{\Phi(\gamma),f}\|_{\tilde{L}_{q_0}^{p_0}(t,T)} \leq c \|f\|_{\tilde{L}_{q_0}^{p_0}(t,T)}, \quad t \in [0, T].$$

- (2) *For any $f \in \mathcal{B}_b(\mathbb{R}^d)$ and $\gamma \in \Gamma$,*

$$(3.18) \quad \partial_t P_{t,s}^{\gamma,V} f = -(\tilde{L}_t^\gamma + V_{T-t}) P_{t,s}^{\gamma,V} f, \quad t \in [0, s], s \in (0, T],$$

and there exists a constant $c > 0$ determined by $(H_{a,b})$ and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$, such that

$$(3.19) \quad \sup_{\gamma \in \Gamma} \sup_{(t,s) \in D_T} \|P_{t,s}^{\gamma,V} f\|_{\mathcal{C}_b^1} \leq c \|f\|_{\mathcal{C}_b^1}, \quad f \in \mathcal{C}_b^1,$$

$$(3.20) \quad \|\nabla P_{t,s}^{\gamma,V} f\|_\infty \leq c(s-t)^{-\frac{1}{2}} \|f\|_\infty, \quad 0 \leq t < s \leq T, f \in \mathcal{B}_b(\mathbb{R}^d).$$

Proof. (1) By Lemma 3.1, $(H_{a,b})$, (3.14) and $\|f\|_{\tilde{L}_{q_0}^{p_0}(0,T)} < \infty$ imply that the PDE

$$(3.21) \quad (\partial_t + \tilde{L}_t^\gamma + V_{T-t})\hat{u}_t^\gamma = -f_t, \quad t \in [0, T], \quad \hat{u}_T^\gamma = 0$$

has a unique solution satisfying

$$(3.22) \quad \|\hat{u}_t^\gamma\|_\infty + \|\nabla \hat{u}_t^\gamma\|_\infty + \|\nabla^2 \hat{u}_t^\gamma\|_{\tilde{L}_{q_0}^{p_0}(t,T)} \leq c_0 \|f\|_{\tilde{L}_q^p(t,T)}$$

for some constant $c_0 > 0$ determined by $(H_{a,b})$ and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$.

Next, by $(H_{a,b})$ and (3.14), the Krylov estimate in [18, Lemma 3.2(2)] holds, so that as shown in the proof of [17, Lemma 4.1(2)], we obtain Khasminskii's estimate: for some function $c : (0, \infty) \rightarrow (0, \infty)$ determined by $(H_{a,b})$ and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$,

$$(3.23) \quad \sup_{\gamma \in \Gamma} \sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \mathbb{E} e^{\lambda \int_0^t |V_{T-s}(X_{t,s}^{\gamma,x})| ds} \leq c(\lambda), \quad \lambda \in (0, \infty).$$

Moreover, by (3.21) and Itô's formula in [18, Lemma 3.3],

$$\begin{aligned} d \left[\hat{u}_s^\gamma(X_{t,s}^{\gamma,x}) e^{\int_t^s V_{T-r}(X_{t,r}^{\gamma,x}) dr} \right] &= e^{\int_t^s V_{T-r}(X_{t,r}^{\gamma,x}) dr} (\partial_s + \tilde{L}_s^\gamma + V_{T-s}) \tilde{u}_s^\gamma(X_{t,s}^{\gamma,x}) ds + dM_s \\ &= -f_s(X_{t,s}^{\gamma,x}) e^{\int_t^s V_{T-r}(X_{t,r}^{\gamma,x}) dr} ds + dM_s, \quad s \in [t, T] \end{aligned}$$

holds for some martingale M_s . This together with $\hat{u}_T^\gamma = 0$, (3.22) and (3.23) yields

$$\begin{aligned} 0 &= \mathbb{E} \left[\hat{u}_T^\gamma(X_{t,T}^{\gamma,x}) e^{\int_t^T V_{T-r}(X_{t,r}^{\gamma,x}) dr} \right] \\ &= \hat{u}_t^\gamma(x) - \mathbb{E} \int_t^T f_s(X_{t,s}^{\gamma,x}) e^{\int_t^s V_{T-r}(X_{t,r}^{\gamma,x}) dr} ds = \hat{u}_t^\gamma(x) - u_t^{\Phi(\gamma), V}(f). \end{aligned}$$

Therefore, (3.21) and (3.22) imply (3.16) and (3.17).

(2) By (3.23), there exists a constant $c_0 > 0$ determined by $(H_{a,b})$ and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$, such that

$$(3.24) \quad \sup_{(t,s) \in D_T, \gamma \in \Gamma} \|P_{t,s}^{\gamma, V} f\|_\infty \leq c_0 \|f\|_\infty, \quad f \in \mathcal{B}_b(\mathbb{R}^d).$$

By [18, Theorem 2.1], or Lemma 3.1 for $V = 0$, $(H_{a,b})$ and (3.14) imply that for any $s \in (0, T]$, the PDE

$$(3.25) \quad (\partial_t + \tilde{L}_t^\gamma) \tilde{u}_t = -P_{t,s}^\gamma f, \quad \tilde{u}_s = 0, \quad t \in [0, s]$$

has a unique solution satisfying

$$(3.26) \quad \sup_{t \in [0, s]} \{ \|\tilde{u}_t\|_\infty + \|\nabla \tilde{u}_t\|_\infty \} + \|\nabla^2 \tilde{u}\|_{\tilde{L}_{q_0}^{p_0}(0,s)} \leq c_1 \|f\|_\infty$$

for some constant $c_1 > 0$ determined by $(H_{a,b})$ and $\|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$. By Itô's formula in [18, Lemma 3.3], we find a martingale $(M_r)_{r \in [t,s]}$ such that (3.25) implies

$$d\tilde{u}_r(X_{t,r}^{\gamma,x}) = dM_r - P_{r,s}^\gamma f(X_{t,r}^{\gamma,x}) dr, \quad r \in [t, s],$$

so that

$$0 = \mathbb{E}[\tilde{u}_T(X_{t,s}^{\gamma,x})] = \tilde{u}_t(x) - \int_t^s \mathbb{E}[P_{r,s}^\gamma f(X_{t,r}^{\gamma,x})] dr = \tilde{u}_t(x) - (s-t)P_{t,s}^\gamma f.$$

Then $P_{t,s}^\gamma f = \frac{\tilde{u}_t}{s-t}$ for $t \in [0, s]$, which together with (3.25) and (3.26) implies

$$(3.27) \quad (\partial_t + \tilde{L}_t^\gamma) P_{t,s}^\gamma f = 0, \quad t \in [0, s],$$

and

$$\sup_{t \in [0,r]} \|P_{t,s}^\gamma f\|_{\mathcal{C}_b^1} + \|\nabla^2 P_{t,s}^\gamma f\|_{\tilde{L}_{q_0}^{p_0}(0,r)} < \infty, \quad r \in [0, s].$$

By Itô's formula in [18, Lemma 3.3], we find a martingale \tilde{M}_r such that (3.27) implies

$$d\left\{e^{\int_t^r V_{T-r'}(X_{t,r'}^{\gamma,x}) dr'} P_{r,s}^\gamma f(X_{t,r}^{\gamma,x})\right\} = d\tilde{M}_r + e^{\int_t^r V_{T-r'}(X_{t,r'}^{\gamma,x}) dr'} \{V_{T-r} P_{r,s}^\gamma f\}(X_{t,r}^{\gamma,x}) dr, \quad r \in [t, s].$$

Thus,

$$\begin{aligned} P_{t,s}^{\gamma,V} f(x) &= \lim_{r \uparrow s} \mathbb{E}\left[e^{\int_t^r V_{T-r'}(X_{t,r'}^{\gamma,x}) dr'} P_{r,s}^\gamma f(X_{t,r}^{\gamma,x})\right] \\ &= \lim_{r \uparrow s} \mathbb{E}\left[P_{t,s}^\gamma f(x) + \int_t^r e^{\int_t^{s'} V_{T-r'}(X_{t,r'}^{\gamma,x}) dr'} \{V_{T-s'} P_{s',s}^\gamma f\}(X_{t,s'}) ds'\right] \\ &= P_{t,s}^\gamma f(x) + \int_t^s P_{t,r}^{\gamma,V} \{V_{T-r} P_{r,s}^\gamma f\}(x) dr, \quad f \in C_b(\mathbb{R}^d). \end{aligned}$$

By a standard approximation argument, this implies

$$(3.28) \quad P_{t,s}^{\gamma,V} f = P_{t,s}^\gamma f + u_t^{\Phi(\gamma),V}(\tilde{f}), \quad t \in [0, s], \quad f \in \mathcal{B}_b(\mathbb{R}^d),$$

where $\tilde{f}_r := 1_{[0,s]}(r) V_{T-r} P_{r,s}^\gamma f$ satisfies

$$\|\tilde{f}\|_{\tilde{L}_{q_0}^{p_0}(0,s)} \leq \|f\|_\infty \|V\|_{\tilde{L}_{q_0}^{p_0}(0,s)}.$$

Combining this with (3.12) for $p = \infty$ and (3.17), we find a constant $c_2 > 0$ determined by $(H_{a,b})$ and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$, such that

$$\sup_{(t,s) \in D_T, \gamma \in \Gamma} \|\nabla P_{t,s}^{\gamma,V} f\|_\infty \leq c_2 \|f\|_{\mathcal{C}_b^1}, \quad f \in \mathcal{C}_b^1,$$

$$\sup_{\gamma \in \Gamma} \|\nabla P_{t,s}^{\gamma,V} f\|_\infty \leq c_2 (s-t)^{-\frac{1}{2}}, \quad f \in \mathcal{B}_b(\mathbb{R}^d), \quad 0 \leq t < s \leq T.$$

These together with (3.24) implies (3.19) and (3.20). Finally, by (3.16), (3.27) and (3.28), we prove (3.18). \square

Lemma 3.4. *Assume (H_{V,u_0}) , $(H_{a,b})$ and $(H_{F,g}^0)$. Then $\tilde{\Gamma}$ is non-empty, ρ_λ -complete and satisfies $\Phi\tilde{\Gamma} \subset \tilde{\Gamma}$.*

Proof. (a) We first prove $\Phi\Gamma \subset \tilde{\Gamma}$. Let $\gamma \in \Gamma$. By the definition of Φ in (3.3), we have $\Phi_{t,t}^x(\gamma) = \delta_x$. Next, (3.19) implies that $\{\Phi(\gamma)\}_{t,s}^V(f) = P_{t,s}^{\gamma,V} f$ satisfies

$$\sup_{\gamma \in \Gamma} \|\{\Phi(\gamma)\}_{t,s}^V(f)\|_{\mathcal{E}_b^1} \leq c \|f\|_{\mathcal{E}_b^1}.$$

Moreover, by (3.17) we obtain

$$\sup_{\gamma \in \Gamma} \sup_{t \in [0,T]} \|u_t^{\Phi(\gamma),V}(f)\|_{\mathcal{E}_b^1} \leq c \|f\|_{\tilde{L}_{q_0}^{p_0}(0,T)}.$$

Thus, $(\Phi\tilde{\Gamma} \subset)\Phi\Gamma \subset \tilde{\Gamma}$.

On the other hand, let γ_X be the distribution of the solution $X = (X_{t,s}^x)_{(t,s) \in D_T, x \in \mathbb{R}^d}$ to (3.2) with $F = 0$. Then the above argument implies $\gamma \in \tilde{\Gamma}$. So, $\tilde{\Gamma} \neq \emptyset$.

(b) The proof of ρ_λ -completeness is more or less standard, which is addressed below for readers' convenience.

Let $\{\gamma^{(n)}\}_{n \geq 1} \subset \tilde{\Gamma}$ such that

$$(3.29) \quad \lim_{n,m \rightarrow \infty} \rho_\lambda(\gamma^{(n)}, \gamma^{(m)}) = 0.$$

Then there exists a constant $c > 0$ such that

$$(3.30) \quad \sup_{n \geq 1, t \in [0,T]} \left\{ \|\gamma_t^{(n)}\|_V + \sup_{\|f\|_{\mathcal{E}_b^1} \leq 1} \|\nabla(\gamma^{(n)})_{t,T}^V(f)\|_\infty + \sup_{\|f\|_{\tilde{L}_{q_0}^{p_0}(0,T)} \leq 1} \|u_t^{\gamma^{(n),V}}(f)\|_{p_0, q_0} \right\} \leq c.$$

Moreover, $\{\gamma^{(n)}\}_{n \geq 1}$ is a Cauchy sequence under the weighted variation distance

$$\|\gamma - \tilde{\gamma}\|_V := \sup_{(t,x) \in [0,T] \times \mathbb{R}^d} \|\gamma_t - \tilde{\gamma}_t\|_V,$$

which is complete. So, there exists a unique measurable map $\gamma : \mathbb{R}^d \rightarrow \mathcal{P}_T$ such that

$$(3.31) \quad \lim_{n \rightarrow \infty} \sup_{t \in [0,T]} \|\gamma_t^{(n)} - \gamma_t\|_V = 0.$$

In particular,

$$(3.32) \quad \gamma_{t,T}^V(f)(x) = \int_{\mathcal{E}_T} f(\xi_{t,T}) \eta_t^V(\xi) \gamma_t^x(d\xi) = \lim_{n \rightarrow \infty} (\gamma^{(n)})_{t,T}^V(f)(x), \quad f \in \mathcal{B}_b(\mathbb{R}^d).$$

We aim to prove that $\gamma \in \tilde{\Gamma}$ and $\rho_\lambda(\gamma^{(n)}, \gamma) \rightarrow 0$ as $n \rightarrow \infty$.

By (3.30) and (3.31), we obtain

$$(3.33) \quad \sup_{t \in [0,T]} \|\gamma_t\|_V = \sup_{t \in [0,T]} \lim_{n \rightarrow \infty} \|(\gamma^{(n)})_t\|_V < \infty.$$

By (3.30), we find a constant $c_1 > 0$ such that

$$\sup_{t \in [0, T]} \sup_{\|f\|_{\mathcal{E}_b^1} \leq 1} |(\gamma^{(n)})_{t, T}^V(f)(x) - (\gamma^{(n)})_{t, T}^V(f)(y)| \leq c_1 |x - y|, \quad x, y \in \mathbb{R}^d, n \geq 1.$$

Combining this with (3.32), we obtain

$$\sup_{t \in [0, T]} \sup_{\|f\|_{\mathcal{E}_b^1} \leq 1} |\gamma_{t, T}^V(f)(x) - \gamma_{t, T}^V(f)(y)| \leq c_1 |x - y|, \quad x, y \in \mathbb{R}^d.$$

Hence,

$$\sup_{t \in [0, T]} \sup_{\|f\|_{\mathcal{E}_b^1} \leq 1} \|\nabla \gamma_{t, T}^V(f)\|_\infty \leq c_1 < \infty.$$

Similarly, we can prove

$$\sup_{t \in [0, T]} \sup_{\|f\|_{\tilde{L}_{q_0}^{p_0}(0, T)} \leq 1} \|u_t^{\gamma, V}(f)\|_{\mathcal{E}_b^1} < \infty.$$

Therefore, by Definition 3.1, we have $\gamma \in \tilde{\Gamma}$.

Next, let $\|f\|_{\mathcal{E}_b^1} \leq 1$. By (3.29) and (3.32), we find positive constants $\{\varepsilon_{m, n}\}_{m, n \geq 1}$ uniformly in f with $\varepsilon_{m, n} \rightarrow 0$ as $m, n \rightarrow \infty$, such that for any $(t, x) \in \mathbb{R}^d$ and $v \in \mathbb{R}^d$ with $|v| = 1$,

$$\begin{aligned} & \frac{1}{\varepsilon} |\gamma_{t, T}^V(f)(x + \varepsilon v) - \gamma_{t, T}^V(f)(x) - (\gamma^{(n)})_{t, T}^V(f)(x + \varepsilon v) + (\gamma^{(n)})_{t, T}^V(f)(x)| \\ &= \lim_{m \rightarrow \infty} \frac{1}{\varepsilon} |(\gamma^{(m)})_{t, T}^V(f)(x + \varepsilon v) - (\gamma^{(m)})_{t, T}^V(f)(x) - (\gamma^{(n)})_{t, T}^V(f)(x + \varepsilon v) + (\gamma^{(n)})_{t, T}^V(f)(x)| \\ &\leq \limsup_{m \rightarrow \infty} \frac{1}{\varepsilon} \int_0^\varepsilon |\nabla_v [(\gamma_{t, T}^{(m)})_{t, T}^V(f) - (\gamma_{t, T}^{(n)})_{t, T}^V(f)](x + rv)| dr \\ &\leq \limsup_{m \rightarrow \infty} \varepsilon_{m, n}, \quad n \geq 1, \varepsilon > 0, t \in [0, T]. \end{aligned}$$

Letting $\varepsilon \downarrow 0$ and taking sup in $x, v \in \mathbb{R}^d$ with $|v| = 1$, we derive

$$\sup_{\|f\|_{\mathcal{E}_b^1} \leq 1} \|\nabla \{ \gamma_{t, T}^V(f) - (\gamma^{(n)})_{t, T}^V(f) \}\|_\infty \leq \limsup_{m \rightarrow \infty} \varepsilon_{m, n}, \quad n \geq 1, t \in [0, T],$$

so that $\varepsilon_{m, n} \rightarrow 0$ as $m, n \rightarrow \infty$ yields

$$\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \|\gamma_t - \gamma_t^{(n)}\|_{\mathcal{E}_b^1} = 0.$$

Similarly, the same holds for $\|\gamma_t - \gamma_t^{(n)}\|_{p_0, p_0}$. Combining these with (3.31), we prove $\rho_\lambda(\gamma^{(n)}, \gamma) \rightarrow 0$ as $n \rightarrow \infty$. \square

4 Proofs of Theorem 2.1 and Theorem 2.2

Proof of Theorem 2.1. We intend to prove that Φ has a unique fixed point γ in $\tilde{\Gamma}$, so that (2.1) has a unique solution given by $X_{t,s}^x = X_{t,s}^\gamma$, and (2.2), (2.3), (2.4) and (2.5) follow from (3.11), (3.13), (3.12), (3.17), (3.19) and (3.20).

Let $\gamma, \tilde{\gamma} \in \tilde{\Gamma}$. By $(H_{F,g}^0)$, we find a constant $K > 0$ such that

$$(4.1) \quad \left\| F_{T-r}(\cdot, \psi_r^{\tilde{\gamma}}, \nabla \psi_r^{\tilde{\gamma}}) - F_{T-r}(\cdot, \psi_r^\gamma, \nabla \psi_r^\gamma) \right\|_\infty \leq K \{1 \wedge \|\psi_r^{\tilde{\gamma}} - \psi_r^\gamma\|_{\mathcal{E}_b^1}\}, \quad \gamma, \tilde{\gamma} \in \tilde{\Gamma}.$$

By $\|u_0\|_{\mathcal{E}_b^1} + \|g\|_{\tilde{L}_{q_0}^{p_0}(0,T)} < \infty$, there exists a constant $c_2 > 0$ such that

$$(4.2) \quad \|\psi_{r,T}^\gamma - \psi_{r,T}^{\tilde{\gamma}}\|_{\mathcal{E}_b^1} \leq c_2 \sup_{s \in [r,T]} \{ \|\gamma_s - \tilde{\gamma}_s\|_{\mathcal{E}_b^1} + \|\gamma_s - \tilde{\gamma}_s\|_{p_0, q_0} \}.$$

Let

$$h_r^x := \sigma_r^{-1}(X_{t,r}^{\gamma,x}) \{ F_{T-r}(\cdot, \psi_r^{\tilde{\gamma}}, \nabla \psi_r^{\tilde{\gamma}}) - F_{T-r}(\cdot, \psi_r^\gamma, \nabla \psi_r^\gamma) \}(X_{t,r}^{\gamma,x}), \quad r \in [t, T].$$

By $(H_{a,b})$, (4.1) and (4.2), we find a constant $c_3 > 0$ such that

$$(4.3) \quad |h_r^x| \leq c_2 \sup_{s \in [r,T]} 1 \wedge \{ \|\gamma_s - \tilde{\gamma}_s\|_{\mathcal{E}_b^1} + \|\gamma_s - \tilde{\gamma}_s\|_{p_0, q_0} \} =: \zeta_r.$$

Let

$$R_t^x := e^{\int_t^T \langle h_r^x, dW_r \rangle - \frac{1}{2} \int_t^T |h_r^x|^2 dr}.$$

By Girsanov's theorem,

$$\tilde{W}_r := W_r - \int_t^r h_s^x ds, \quad r \in [t, T]$$

is a Brownian motion under probability $\mathbb{Q}_t^x := R_t^x \mathbb{P}$. Reformulating (3.2) as

$$dX_{t,s}^{\gamma,x} = \{ b_{T-s} + F_{T-s}(\cdot, \psi_s^{\tilde{\gamma}}, \nabla \psi_s^{\tilde{\gamma}}) \}(X_{t,s}^{\gamma,x}) ds + \sigma_s(X_{t,s}^{\gamma,x}) d\tilde{W}_s, \quad s \in [t, T], X_{t,t}^{\gamma,x} = x,$$

by the weak uniqueness we see that the law of $X_{t,\cdot}^x := (X_{t,s}^{\gamma,x})_{s \in [t,T]}$ under \mathbb{Q}_t^x coincides with that of $(X_{t,s}^{\tilde{\gamma},x})_{s \in [t,T]}$ under \mathbb{P} . So, by (3.23), we find a constant $c_3 > 0$ such that

$$\begin{aligned} \|\Phi_t(\gamma) - \Phi_t(\tilde{\gamma})\|_V &= \sup_{\|f\|_\infty \leq \eta_t^V} \sup_{x \in \mathbb{R}^d} |\mathbb{E}[(R_t^x - 1)f(X_{t,\cdot}^{\gamma,x})]| \\ &\leq \sup_{x \in \mathbb{R}^d} \mathbb{E}[\eta_t^V |R_t^x - 1|] \leq (\mathbb{E}[\eta_t^V]^2)^{\frac{1}{2}} (\mathbb{E}|R_t^x|^2 - 1)^{\frac{1}{2}} \leq c_3 (\mathbb{E}|R_t^x|^2 - 1)^{\frac{1}{2}}. \end{aligned}$$

Noting that (4.3) implies

$$\begin{aligned} \mathbb{E}|R_t^x|^2 &\leq \mathbb{E} \left[e^{2 \int_t^T \langle h_r^x, dW_r \rangle - 2 \int_t^T |h_r^x|^2 dr + \int_t^T |h_r^x|^2 dr} \right] \leq e^{\int_t^T |\zeta_r|^2 dr} \\ &\leq 1 + e^{\int_t^T |\zeta_r|^2 dr} \int_t^T |\zeta_r|^2 dr \leq 1 + c_2^2 e^{T c_2^2} \int_t^T \{ \|\gamma_s - \tilde{\gamma}_s\|_{\mathcal{E}_b^1} + \|\gamma_s - \tilde{\gamma}_s\|_{p_0, q_0} \}^2 ds, \end{aligned}$$

we find a constant $c_4 > 0$ such that

$$\|\Phi_t(\gamma) - \Phi_t(\tilde{\gamma})\|_V \leq c_4 \left(\int_t^T \{ \|\gamma_s - \tilde{\gamma}_s\|_{\mathcal{C}_b^1} + \|\gamma_s - \tilde{\gamma}_s\|_{p_0, q_0} \}^2 ds \right)^{\frac{1}{2}}.$$

Consequently,

$$(4.4) \quad e^{-\lambda(T-t)} \|\Phi_t(\gamma) - \Phi_t(\tilde{\gamma})\|_V \leq c_4 \rho_\lambda(\gamma, \tilde{\gamma}) \left(\int_t^T e^{-2\lambda(r-t)} dr \right)^{\frac{1}{2}}, \quad t \in [0, T].$$

Next, by (3.18) and (3.19), for any $f \in \mathcal{C}_b^1(\mathbb{R}^d)$, we may apply Itô's formula in [18, Lemma 3.3] to deduce

$$\begin{aligned} d \left\{ e^{\int_t^r V_{T-r'}(X_{t,r'}^{\tilde{\gamma},x}) dr'} P_{r,s}^{\gamma,V} f(X_{t,r}^{\tilde{\gamma},x}) \right\} &= dM_r \\ + e^{\int_t^r V_{T-r'}(X_{t,r'}^{\tilde{\gamma},x}) dr'} &\left\{ [F_{T-r}(\cdot, \psi_{r,T}^{\tilde{\gamma}}, \nabla \psi_{r,T}^{\tilde{\gamma}}) - F_{T-r}(\cdot, \psi_{r,T}^\gamma, \nabla \psi_{r,T}^\gamma)] \cdot \nabla P_{r,s}^{\gamma,V} f \right\} (X_{t,r}^{\tilde{\gamma},x}) dr \end{aligned}$$

for some martingale $M_r, r \in [0, s]$. Then

$$(4.5) \quad \begin{aligned} P_{t,s}^{\tilde{\gamma},V} f(x) - P_{t,s}^{\gamma,V} f(x) &= \mathbb{E} \left[e^{\int_t^s V_{T-r}(X_{t,r}^{\tilde{\gamma},x}) dr} P_{s,s}^{\gamma,V} f(X_{t,s}^{\tilde{\gamma},x}) \right] - \mathbb{E} \left[e^{\int_t^s V_{T-r}(X_{t,r}^{\tilde{\gamma},x}) dr} P_{t,s}^{\gamma,V} f(X_{t,t}^{\tilde{\gamma},x}) \right] \\ &= \int_t^s P_{t,r}^{\tilde{\gamma},V} \left\{ [F_{T-r}(\cdot, \psi_r^{\tilde{\gamma}}, \nabla \psi_r^{\tilde{\gamma}}) - F_{T-r}(\cdot, \psi_r^\gamma, \nabla \psi_r^\gamma)] \cdot \nabla P_{r,s}^{\gamma,V} f \right\} dr. \end{aligned}$$

Combining this with (3.19), (3.20), (4.1) and (4.2), we find constants $c_5, c_6 > 0$ such that

$$\begin{aligned} &\sup_{\|f\|_{\mathcal{C}_b^1} \leq 1} \|\nabla \{\Phi(\gamma)\}_{t,T}^V(f) - \nabla \{\Phi(\tilde{\gamma})\}_{t,T}^V(f)\|_\infty \\ &= \sup_{\|f\|_{\mathcal{C}_b^1} \leq 1} \|\nabla P_{t,T}^{\gamma,V} f - \nabla P_{t,T}^{\tilde{\gamma},V} f\|_\infty \leq c_5 \int_t^T (r-t)^{-\frac{1}{2}} \|\psi_r^\gamma - \psi_r^{\tilde{\gamma}}\|_{\mathcal{C}_b^1} dr \\ &\leq c_6 \int_t^T (r-t)^{-\frac{1}{2}} \sup_{\theta \in [r,T]} \{ \|\gamma_r - \tilde{\gamma}_r\|_{\mathcal{C}_b^1} + \|\gamma_r - \tilde{\gamma}_r\|_{p_0, q_0} \} dr. \end{aligned}$$

Then

$$(4.6) \quad e^{-\lambda(T-t)} \|\Phi_t(\gamma) - \Phi_t(\tilde{\gamma})\|_{\mathcal{C}_b^1} \leq c_6 \rho_\lambda(\gamma, \tilde{\gamma}) \int_t^s (r-t)^{-\frac{1}{2}} e^{-\lambda(r-t)} dr, \quad t \in [0, T].$$

Finally, let $\|f\|_{\tilde{L}_{q_0}^{p_0}(0,T)} \leq 1$. By (3.16), for any $r \in [0, T)$, we have

$$\begin{aligned} (\partial_t + \tilde{L}_t^\gamma + V_{T-t}) \{ u_t^{\Phi(\gamma),V}(f) - u_t^{\Phi(\tilde{\gamma}),V}(f) \} &= \tilde{f}_t, \quad t \in [0, T], \quad u_T^{\Phi(\gamma),f} - u_T^{\Phi(\tilde{\gamma}),f} = 0, \\ \tilde{f}_t &:= \{ F_{T-t}(\cdot, \psi_t^{\tilde{\gamma}}, \nabla \psi_t^{\tilde{\gamma}}) - F_{T-t}(\cdot, \psi_t^\gamma, \nabla \psi_t^\gamma) \} \cdot \nabla u_t^{\Phi(\tilde{\gamma}),V}(f). \end{aligned}$$

By $\|f\|_{\tilde{L}_{q_0}^{p_0}(0,T)} \leq 1$, (3.17) and (4.1), there exists a constant $c > 0$ such that

$$\|\tilde{f}_t\|_\infty \leq c \|\psi_t^\gamma - \psi_t^{\tilde{\gamma}}\|_{\mathcal{C}_b^1}, \quad t \in [0, T].$$

So, by Lemma 3.1, there exist constants $c_7, c_8 > 0$ such that

$$\sup_{t \in [r, T]} \|u_t^{\Phi(\gamma), V}(f) - u_t^{\Phi(\tilde{\gamma}), V}(f)\|_{\mathcal{C}_b^1} \leq c_7 \|\tilde{f}\|_{\tilde{L}_{q_0}^{p_0}(r, T)} \leq c_8 \left(\int_r^T \|\psi_t^\gamma - \psi_t^{\tilde{\gamma}}\|_{\mathcal{C}_b^1}^{q_0} dt \right)^{\frac{1}{q_0}}.$$

Combining this with (4.2), we find a constant $c_9 > 0$ such that

$$e^{-\lambda(T-t)} \|\{\Phi(\gamma)\}_t - \{\Phi(\tilde{\gamma})\}_t\|_{p_0, q_0} \leq c_9 \rho_\lambda(\gamma, \tilde{\gamma}) \left(\int_t^T e^{-q_0 \lambda(s-t)} ds \right)^{\frac{1}{q_0}}, \quad t \in [0, T].$$

By taking large enough $\lambda > 0$, this and (4.4)-(4.6) yield that

$$\rho_\lambda(\Phi(\gamma), \Phi(\tilde{\gamma})) \leq \frac{1}{2} \rho_\lambda(\gamma, \tilde{\gamma}), \quad \gamma, \tilde{\gamma} \in \tilde{\Gamma}.$$

Therefore, Φ has a unique fixed point in $\tilde{\Gamma}$ as desired.

Since for any solution $X = (X_{t,s}^x)_{(t,s,x) \in D_T \times \mathbb{R}^d}$ of (2.1), Lemmas 3.2 and 3.3 imply that the law γ_X of X is in $\tilde{\Gamma}$, so that γ_X is a fixed point of Φ in $\tilde{\Gamma}$. By the uniqueness of the fixed point, as well as the well-posedness of (3.2), we prove the uniqueness of (1.2). \square

Proof of Theorem 2.2. (1) Let u solve (2.1) for $t \in [0, T]$ and satisfy (2.7), we intend to prove that u satisfies (2.6).

Note that $\psi_t = \psi_t^\gamma$ for the unique fixed point $\gamma \in \tilde{\Gamma}$ of Φ , (3.19) for $P_{T-t, T}^{\gamma, V} u_0$ and (3.17) for $f = g_{T-}$ imply that

$$(4.7) \quad \sup_{t \in [0, T]} (\|\psi_t\|_\infty + \|\nabla \psi_t\|_\infty) < \infty.$$

Next, let

$$\tilde{f}_s := F_{T-s}(\cdot, \psi_s, \nabla \psi_s) - F_{T-s}(\cdot, u_{T-s}, \nabla u_{T-s}).$$

By $(H_{F,g}^0)$, we find a constant $c_1 > 0$ such that

$$(4.8) \quad |\tilde{f}_s| \leq c_1 \|\psi_s - u_{T-s}\|_{\mathcal{C}_b^1}, \quad s \in [0, T].$$

By (1.2) and Itô's formula in [18, Lemma 3.3], we obtain

$$d\{u_{T-s}(X_{t,s}^x) e^{\int_t^s V_{T-r}(X_{t,r}^x) dr}\} = dM_s + e^{\int_t^s V_{T-r}(X_{t,r}^x) dr} [\tilde{f}_s \cdot \nabla u_{T-s} - g_{T-s}](X_{t,s}^x) ds, \quad s \in [t, T]$$

for some martingale M_s . Combining this with the definition of ψ_t in (1.2), we derive

$$\begin{aligned} & \psi_t(x) - u_{T-t}(x) \\ &= \mathbb{E} \left[u_0(X_{t,T}^x) e^{\int_t^T V_{T-r}(X_{t,r}^x) dr} + \int_t^T g_{T-s}(X_{t,s}^x) e^{\int_t^s V_{T-r}(X_{t,r}^x) dr} ds \right] - u_{T-t}(x) \\ (4.9) \quad &= \mathbb{E} \int_t^T e^{\int_t^s V_{T-r}(X_{t,r}^x) dr} [\tilde{f}_s \cdot \nabla u_{T-s}](X_{t,s}^x) ds \\ &= \int_t^T P_{t,s}^V [\tilde{f}_s \cdot \nabla u_{T-s}](x) ds, \quad t \in [0, T], \end{aligned}$$

where $P_{t,s}^V = P_{t,s}^{\gamma,V}$ for γ being the unique fixed point of Φ . Combining this with (3.17), (4.8) and $\|\nabla u\|_\infty < \infty$, we find a constant $c_2 > 0$ such that

$$\|\psi_t - u_{T-t}\|_{\mathcal{E}_b^1} \leq c_2 \left(\int_t^T \|\psi_s - u_{T-s}\|_{\mathcal{E}_b^1}^{q_0} ds \right)^{\frac{1}{q_0}}, \quad t \in [0, T].$$

Since (2.7) and (4.7) imply that $\|\psi_t - u_{T-t}\|_{\mathcal{E}_b^1}$ is bounded in $t \in [0, T]$, we prove $\psi_t - u_{T-t} = 0$ for all $t \in [0, T]$, so that the proof is finished.

(2) Let u be given in (2.6), and let $P_{t,s}^V = P_{t,s}^{\gamma,V}$ for the unique fixed point $\gamma \in \tilde{\Gamma}$ of Φ . We have

$$(4.10) \quad u_t = \psi_{T-t} = P_{T-t,T}^V u_0 + \int_{T-t}^T P_{T-t,s}^V g_{T-s} ds, \quad t \in [0, T].$$

Let

$$\tilde{L}_t = L_{T-t} + F_{T-t}(\cdot, \psi_t, \nabla \psi_t) \cdot \nabla, \quad t \in [0, T].$$

By Lemma 3.3(2),

$$u_t^{(1)} := P_{T-t,T}^V u_0, \quad t \in [0, T]$$

satisfies

$$(4.11) \quad \begin{aligned} \partial_t u_t^{(1)} &= (\tilde{L}_{T-t} + V_t) u_t^{(1)}, \quad t \in [0, T], \\ \|u^{(1)}\|_\infty + \|\nabla u^{(1)}\|_\infty + \|\nabla^2 u^{(1)}\|_{\tilde{L}_{q_0}^{p_0}(0,T)} &< \infty. \end{aligned}$$

On the other hand, by Lemma 3.3(1),

$$u_t^{(2)} := \int_{T-t}^T P_{T-t,s}^V g_{T-s} ds, \quad t \in [0, T]$$

satisfies

$$(4.12) \quad \begin{aligned} \partial_t u_t^{(2)} &= (\tilde{L}_{T-t} + V_t) u_t^{(2)} + g_t, \quad t \in [0, T], \\ \|u^{(2)}\|_\infty + \|\nabla u^{(2)}\|_\infty + \|\nabla^2 u^{(2)}\|_{\tilde{L}_{q_0}^{p_0}(0,T)} &< \infty, \end{aligned}$$

which together with (4.10) and (4.11) finishes the proof. \square

5 Proof of Theorem 2.3

We first observe that by extending functions from \mathbb{T}^d to \mathbb{R}^d periodically, i.e. letting

$$f(x+k) = f(x), \quad x \in \mathbb{T}^d, \quad k \in \mathbb{Z}^d,$$

we extend the PDE (1.1) from $E = \mathbb{T}^d$ to $E = \mathbb{R}^d$ with assumptions (H_{V,u_0}) , $(H_{a,b})$, $(H_{F,g})$ and $(H'_{F,g})$ invariant. So, the existence of the extended equation implies that of the original

equation. On the other hand, if the original equation has two solutions with same initial value, then their periodical extensions solve the extended equation, so that the uniqueness of the extended equation implies that of the original equation. Therefore, it suffices to prove for $E = \mathbb{R}^d$.

In the following we set $E = \mathbb{R}^d$, and prove Theorem 2.3. We first prove under the following stronger assumption replacing $(H_{F,g})$, then make extension by a truncation argument.

$(\tilde{H}_{F,g})$ $\|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|\bar{g}\|_{\tilde{L}_{q_0}^{p_0}(0,T)} < \infty$, and there exist constants $K, \alpha > 0$ such that

$$\begin{aligned} & |F_t(x, r_1, r_2) - F_t(x, \tilde{r}_1, \tilde{r}_2)| + |g_t(x, r_1, r_2, r_3) - g_t(x, \tilde{r}_1, \tilde{r}_2, \tilde{r}_3)| \\ & \leq K(|r_1 - \tilde{r}_1| + |r_2 - \tilde{r}_2|) + \alpha|r_3 - \tilde{r}_3|, \\ & (t, x) \in [0, T] \times \mathbb{R}^d, \quad r_1, \tilde{r}_1 \in \mathbb{R}^m, \quad r_2, \tilde{r}_2 \in \mathbb{R}^{d \otimes m}, \quad r_3, \tilde{r}_3 \in \mathbb{R}^{d \otimes d \otimes m}. \end{aligned}$$

Lemma 5.1. *Assume $(H_{a,b})$, (H_{V,u_0}) and $(\tilde{F}_{F,g})$ with small enough $\alpha > 0$ determined by $(H_{a,b})$ and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$. The equation (1.1) has a unique solution in $\mathcal{U}(p_0, q_0)$ with $T^* = T$ such that*

$$\sup_{t \in [0, T]} \|u_t\|_{\mathcal{C}_b^1} + \|\nabla^2 u\|_{\tilde{L}_{q_0}^{p_0}(0, T)} < \infty.$$

Proof. We will make use of the fixed point theorem for a map Ψ induced by (1.1). Let \mathbf{H} be the Banach space of maps

$$h : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^m$$

satisfying

$$\|h\|_{\mathbf{H}} := \sup_{t \in [0, T]} \|h_t\|_{\mathcal{C}_b^1} + \|\nabla^2 h\|_{\tilde{L}_{q_0}^{p_0}(0, T)} < \infty.$$

Given $h \in \mathbf{H}$, consider the PDE

$$(5.1) \quad \partial_t u_t^h = (L_t + V_t)u_t^h + F_t(\cdot, u_t^h, \nabla u_t^h) \cdot \nabla u_t^h + g_t(\cdot, h_t, \nabla h_t, \nabla^2 h_t), \quad t \in [0, T].$$

By $(\tilde{H}_{F,g})$, we have $\|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0, T)} < \infty$ and

$$(5.2) \quad \begin{aligned} & \|g(\cdot, h, \nabla h, \nabla^2 h)\|_{\tilde{L}_{q_0}^{p_0}(0, t)} \leq \|\bar{g}\|_{\tilde{L}_{q_0}^{p_0}(0, t)} \\ & + K \left(\int_0^t \|h_s\|_{\mathcal{C}_b^1}^{q_0} ds \right)^{\frac{1}{q_0}} + \alpha \|\nabla^2 h\|_{\tilde{L}_{q_0}^{p_0}(0, t)} < \infty, \quad t \in [0, T], \end{aligned}$$

so that $(H_{F,g}^0)$ holds for $g(\cdot, h, \nabla h, \nabla^2 h)$ replacing g . According to Theorem 2.1 and Theorem 2.2, this together with (H_{V,u_0}) and $(H_{a,b})$ implies that (5.1) has a unique solution in the class $\mathcal{U}(p_0, q_0)$ up to time T , such that

$$(5.3) \quad \|u^h\|_{\infty} \leq k(u_0, V, g),$$

and for some constant $c > 0$ determined by $(H_{a,b})$ and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$,

$$\begin{aligned}
& \|u_t^h\|_{\mathcal{E}_b^1} + \|\nabla^2 u^h\|_{\tilde{L}_{q_0}^{p_0}(0,t)} \leq c(\|u_0\|_{\mathcal{E}_b^1} + \|g(\cdot, h, \nabla h, \nabla^2 h)\|_{\tilde{L}_{q_0}^{p_0}(0,t)}) \\
(5.4) \quad & \leq c(\|u_0\|_{\mathcal{E}_b^1} + \|\bar{g}\|_{\tilde{L}_{q_0}^{p_0}(0,t)}) + cK \left(\int_0^t \|h_s\|_{\mathcal{E}_b^1}^{q_0} ds \right)^{\frac{1}{q_0}} + c\alpha \|\nabla^2 h\|_{\tilde{L}_{q_0}^{p_0}(0,t)}, \\
& h \in \mathbf{H}, t \in [0, T].
\end{aligned}$$

So, it suffices to prove that the map

$$\Psi : \mathbf{H} \ni h \mapsto u^h \in \mathbf{H}$$

has a unique fixed point.

To this end, we let $\alpha \in (0, \frac{1}{2c})$ where $c > 0$ is determined by $(H_{a,b})$ and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$, and denote $c_0 := c(\|u_0\|_{\mathcal{E}_b^1} + \|\bar{g}\|_{\tilde{L}_{q_0}^{p_0}(0,t)})$, and let

$$\mathbf{H}_N := \left\{ h \in \mathbf{H} : \|h\|_N := \sup_{t \in [0, T]} e^{-Nt} (\|h_t\|_{\mathcal{E}_b^1} + \|\nabla^2 h\|_{\tilde{L}_{q_0}^{p_0}(0,t)}) \leq Nc_0 \right\}, \quad N \geq 2.$$

We will find a constant $N_0 \geq 2$ such that for any $N \geq N_0$, $\Psi : \mathbf{H}_N \rightarrow \mathbf{H}_N$ has a unique fixed point.

Firstly, we take $N_0 \geq 2$ such that

$$cK \left(\int_0^T e^{-q_0 N_0 s} ds \right)^{\frac{1}{q_0}} \leq \frac{1}{2}.$$

Combining this with (5.4) and $c\alpha \leq \frac{1}{2}$, we obtain

$$\|\Psi(h)\|_N = \|u^h\|_N \leq c_1 + \frac{1}{2}\|h\|_N \leq c_0 + \frac{N}{2}c_0 \leq Nc_0, \quad h \in \mathbf{H}_N, N \geq N_0.$$

So, $\psi : \mathbf{H}_N \rightarrow \mathbf{H}_N$ for $N \geq N_0$. Thus,

$$(5.5) \quad \sup_{t \in [0, T]} \|u_t^h\|_{\mathcal{E}_b^1} \leq \|\Psi(h)\|_{\mathbf{H}} \leq c(N) := c_0 N e^{NT}, \quad h \in \mathbf{H}_N, N \geq N_0.$$

It remains to prove that Ψ has a unique fixed point in \mathbf{H}_N for any $N \geq N_0$.

For any $h, \tilde{h} \in \mathbf{H}_N$, let

$$\begin{aligned}
f_t & := \{F_{T-t}(\cdot, u_{T-t}^{\tilde{h}}, \nabla u_{T-t}^{\tilde{h}}) - F_{T-t}(\cdot, u_{T-t}^h, \nabla u_{T-t}^h)\} \cdot \nabla u_t^{\tilde{h}} \\
& + g_{T-t}(\cdot, \tilde{h}_{T-t}, \nabla \tilde{h}_{T-t}, \nabla^2 \tilde{h}_{T-t}) - g_{T-t}(\cdot, h_{T-t}, \nabla h_{T-t}, \nabla^2 h_{T-t}), \quad t \in [0, T]
\end{aligned}$$

$$L_t^h := L_{T-t} + F_{T-t}(\cdot, u_{T-t}^h, \nabla u_{T-t}^h) \cdot \nabla, \quad t \in [0, T].$$

Then

$$(5.6) \quad w_t := u_{T-t}^h - u_{T-t}^{\tilde{h}} = \Psi_{T-t}(h) - \Psi_{T-t}(\tilde{h})$$

satisfies $w \in \tilde{H}_{q_0}^{2,p_0}(0, T)$ and solves the PDE

$$(\partial_t + L_t^h + V_{T-t})w_t = f_t, \quad t \in [0, T], \quad w_T = 0.$$

By Lemma 3.1, (5.3), (5.5), (H_{V, u_0}) , $(H_{a,b})$ and $(\tilde{H}_{F,g})$ imply that for some constant $c_1 > 0$ determined by $(H_{a,b})$ and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|\tilde{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$,

$$\begin{aligned} & \|w_t\|_{\mathcal{E}_b^1} + \|\nabla^2 w\|_{\tilde{L}_{q_0}^{p_0}(t,T)} \leq c_1 \|f\|_{\tilde{L}_{q_0}^{p_0}(t,T)} \\ & \leq c_1 K \left(\int_t^T \{c(N)\|w_s\|_{\mathcal{E}_b^1} + \|h_s - \tilde{h}_{T-s}\|_{\mathcal{E}_b^1}\}^{q_0} ds \right)^{\frac{1}{q_0}} \\ & \quad + c_1 \alpha \|\nabla^2(h_{T-\cdot} - \tilde{h}_{T-\cdot})\|_{\tilde{L}_{q_0}^{p_0}(t,T)}, \quad t \in [0, T]. \end{aligned}$$

Then for any $\lambda > 0$, this and (5.6) yield

$$\|\Psi(h) - \Psi(\tilde{h})\|_\lambda \leq \varepsilon(\lambda) [\|h - \tilde{h}\|_\lambda + \|\Psi(h) - \Psi(\tilde{h})\|_\lambda], \quad h, \tilde{h} \in \mathbf{H}_N$$

for $\varepsilon(\lambda) := \max \{c_1 \alpha, (c(N) \vee 1)c_1 K \sup_{t \in [0, T]} (\int_t^T e^{-\lambda q_0(s-t)} ds)^{\frac{1}{q_0}}\}$. Taking $\lambda > 0$ large enough and $\alpha \in (0, \frac{1}{2c_1})$ such that $\varepsilon(\lambda) < \frac{1}{2}$, we see that Ψ is $\|\cdot\|_\lambda$ -contractive in H_N , and hence has a unique fixed point. \square

Proof of Theorem 2.3. (1) For any $n \in \mathbb{N}$, let

$$\varphi_n(r) := r \mathbf{1}_{\{|r| \leq n\}} + \frac{nr}{|r|} \mathbf{1}_{\{|r| > n\}}, \quad r \in \mathbb{R}^m \cup \mathbb{R}^{d \otimes m}.$$

We take the following truncation of F_t :

$$\begin{aligned} F_t^{(n)}(x, r_1, r_2) &= F_t(x, \varphi_n(r_1), \varphi_n(r_2)), \quad g_t^{(n)}(x, r_1, r_2, r_3) = g_t(x, \varphi_n(r_1), \varphi_n(r_2), r_3), \\ t \geq 0, x \in \mathbb{R}^d, (r_1, r_2, r_3) &\in \mathbb{R}^m \times \mathbb{R}^{d \otimes m} \times \mathbb{R}^{d \otimes d \otimes d \otimes m}. \end{aligned}$$

Then $(H_{F,g})$ implies $(\tilde{H}_{F,g})$ for $(F^{(n)}, g^{(n)})$ replacing (F, g) . So, by Lemma 5.1, the equation

$$(5.7) \quad \begin{aligned} \partial_t u_t^{(n)} &= (L_t + V_t)u_t^{(n)} + F_t^{(n)}(\cdot, u_t^{(n)}, \nabla u_t^{(n)}) \cdot \nabla u_t^{(n)} + g_t^{(n)}(\cdot, u_t^{(n)}, \nabla u_t^{(n)}, \nabla^2 u_t^{(n)}), \\ t \in [0, T], u_0^{(n)} &= u_0 \end{aligned}$$

has a unique solution in $\mathcal{U}(p_0, q_0)$ with $T^* = T$, and

$$(5.8) \quad \sup_{t \in [0, T]} \|u_t^{(n)}\|_{\mathcal{E}_b^1} + \|\nabla^2 u_t^{(n)}\|_{\tilde{L}_{q_0}^{p_0}(0, T)} < \infty, \quad n \geq 1.$$

Let

$$(5.9) \quad T_n := T \wedge \inf\{t \geq 0 : \|u_t^{(n)}\|_{\mathcal{E}_b^1} \geq n\}.$$

Then $u_t^{(n)}$ solves (1.1) up to time T_n . Letting $T_0 = 0$ and

$$T^* := \lim_{n \rightarrow \infty} T_n,$$

the uniqueness of (5.7) for every $n \geq 1$ implies

$$u_t^{(n)} = u_t^{(m)}, \quad t \in [0, T_n \wedge T_m], \quad n, m \in \mathbb{N},$$

so that (1.1) has a unique solution

$$u_t := \sum_{n=1}^{\infty} u_t^{(n)} 1_{[T_{n-1}, T_n)}(t), \quad t \in [0, T^*)$$

in the class $\mathcal{U}(p_0, q_0)$, provided $T^* > 0$.

To prove $T^* > 0$, it suffices to show that $T_n > 0$ for large $n > 0$. For fixed $T \in (0, \infty)$, let $P_{t,s}^{(n),V}$ be the Feynman-Kac semigroup generated by

$$L_{T-t} + V_{T-t} + F_{T-t}^{(n)}(\cdot, u_{T-t}^{(n)}, \nabla u_{T-t}^{(n)}) \cdot \nabla, \quad t \in [0, T];$$

that is

$$(5.10) \quad P_{t,s}^{(n),V} f(x) := \mathbb{E}[f(X_{t,s}^{(n),x}) e^{\int_t^s V_{T-r}(X_{t,r}^{(n),x}) dr}], \quad 0 \leq t \leq s \leq T, f \in \mathcal{B}_b(\mathbb{R}^d)$$

for $X_{t,s}^{(n),x}$ solving the SDE

$$\begin{aligned} dX_{t,s}^{(n),x} &= \sigma_s(X_{t,s}^{(n),x}) dW_s + \{b_{T-s} + F_{T-s}^{(n)}(\cdot, u_{T-s}^{(n)}, \nabla u_{T-s}^{(n)})\}(X_{t,s}^{(n),x}) ds, \\ s &\in [t, T], \quad X_{t,t}^{(n),x} = x. \end{aligned}$$

Since $(H_{F,g})$ implies $\|F^{(n)}\|_{\bar{L}_{q_0}^{p_0}(0,T)} < \infty$, by $(H_{a,b})$ and [15, Theorem 2.1], this SDE is well-posed. By (5.7) and Itô's formula, we find a martingale M_s such that

$$\begin{aligned} & d \left\{ u_{T-s}^{(n)}(X_{T-t,s}^{(n),x}) e^{\int_{T-t}^s V_{T-r}(X_{t,r}^{(n),x}) dr} \right\} \\ &= dM_s - g_{T-s}^{(n)}(\cdot, u_{T-s}^{(n)}, \nabla u_{T-s}^{(n)}, \nabla^2 u_{T-s}^{(n)})(X_{T-t,s}^{(n),x}) ds, \quad s \in [T-t, T]. \end{aligned}$$

So,

$$(5.11) \quad u_t^{(n)} = P_{T-t,T}^{(n),V} u_0 + \int_{T-t}^T P_{T-t,s}^{(n),V} g_{T-s}^{(n)}(\cdot, u_{T-s}^{(n)}, \nabla u_{T-s}^{(n)}, \nabla^2 u_{T-s}^{(n)}) ds, \quad t \in [0, T],$$

which implies $\|u^{(n)}\|_{\infty} \leq k(u_0, g, V)$, so that

$$|F_{T-s}^{(n)}(\cdot, u_{T-s}^{(n)}, \nabla u_{T-s}^{(n)})| \leq \bar{F}_{T-s}, \quad s \in [0, T].$$

Combining this with $(H_{a,b})$ and $\|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)} < \infty$, by Lemma 3.3, we find a constant $c > 0$ determined by $(H_{a,b})$ and $\|\bar{F}\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \|V\|_{\tilde{L}_{q_0}^{p_0}(0,T)}$, such that for any $n \geq 1$ and any $t \in [0, T]$,

$$(5.12) \quad \|P_{T-t,T}^{(n),V}u_0\|_{\mathcal{E}_b^1} \leq c, \quad \left\| \int_{T-t}^T P_{T-t,s}^{(n),V}f_s ds \right\|_{\mathcal{E}_b^1} \leq c\|f\|_{\tilde{L}_{q_0}^{p_0}(T-t,T)}, \quad f \in \tilde{L}_{q_0}^{p_0}(0,T),$$

$$(5.13) \quad \|\nabla P_{t,s}^{(n),V}f\|_{\infty} \leq c(s-t)^{-\frac{1}{2}}\|f\|_{\infty}, \quad f \in \mathcal{B}_b(\mathbb{R}^d), \quad s \in (t, T].$$

Combining (5.11) with (5.12), and noting that $(H_{F,g})$ implies

$$\|g^{(n)}(\cdot, u^{(n)}, \nabla u^{(n)}, \nabla^2 u^{(n)})\|_{\tilde{L}_{q_0}^{p_0}(0,T)} \leq 2nK_n + \|\bar{g}\|_{\tilde{L}_{q_0}^{p_0}(0,T)} + \alpha\|\nabla^2 u^{(n)}\|_{\tilde{L}_{q_0}^{p_0}(0,T)} < \infty,$$

we obtain

$$\lim_{t \downarrow 0} \|u_t^{(n)}\|_{\mathcal{E}_b^1} \leq \lim_{t \downarrow 0} \left\{ \|P_{T-t,T}^{(n),V}u_0\|_{\mathcal{E}_b^1} + c\|g^{(n)}(\cdot, u^{(n)}, \nabla u^{(n)}, \nabla^2 u^{(n)})\|_{\tilde{L}_{q_0}^{p_0}(0,t)} \right\} \leq c, \quad n \geq 1.$$

By (5.9), this implies $T_n > 0$ for $n > c$.

(2) Assume $(H'_{F,g})$, we intend to prove $T^* = \infty$. For $n \geq 1$ let $P_{t,s}^{(n),V}$ be in (5.10). Since $u_t = u_t^{(n)}$ for $t \in [0, T_n]$, we have

$$g_s^{(n)}(\cdot, u_s^{(n)}, \nabla u_s^{(n)}, \nabla^2 u_s^{(n)}) = g_s(\cdot, u_s, \nabla u_s, \nabla^2 u_s), \quad s \in [0, T_n].$$

Then with the integral transform $s \mapsto T - s$, (5.11) implies

$$u_t = P_{T-t,T}^{(n),V}u_0 + \int_0^t P_{T-t,T-s}^{(n),V}g_s(\cdot, u_s, \nabla u_s, \nabla^2 u_s)ds, \quad t \in [0, T_n].$$

Combining this with (5.12), (5.13), we find a constant $c_1 > 0$ such that

$$\begin{aligned} \|u_t\|_{\mathcal{E}_b^1} + \|\nabla^2 u\|_{\tilde{L}_{q_0}^{p_0}(0,t)} &\leq c\|u_0\|_{\mathcal{E}_b^1} + c\|g(\cdot, 0, 0, \nabla^2 u)\|_{\tilde{L}_{q_0}^{p_0}(0,t)} \\ &+ c \int_0^t (t-s)^{-\frac{1}{2}} \|g_s(\cdot, u_s, \nabla u_s, \nabla^2 u_s) - g_s(\cdot, 0, 0, \nabla^2 u_s)\|_{\infty} ds. \end{aligned}$$

Since $(H'_{F,g})$ implies

$$\begin{aligned} \|g(\cdot, 0, 0, \nabla^2 u)\|_{\tilde{L}_{q_0}^{p_0}(0,t)} &\leq \|\bar{g}\|_{\tilde{L}_{q_0}^{p_0}(0,t)} + \alpha\|\nabla^2 u\|_{\tilde{L}_{q_0}^{p_0}(0,t)}, \\ \|g_s(\cdot, u_s, \nabla u_s, \nabla^2 u_s) - g_s(\cdot, 0, 0, \nabla^2 u_s)\|_{\infty} &\leq C(e + \|u_s\|_{\mathcal{E}_b^1})\{\log(e + \|u_s\|_{\mathcal{E}_b^1})\}^{\theta}, \end{aligned}$$

we find a constant $c_1 > 0$ such that

$$(5.14) \quad \begin{aligned} \|u_t\|_{\mathcal{E}_b^1} + \|\nabla^2 u\|_{\tilde{L}_{q_0}^{p_0}(0,t)} &\leq c_1 + c\alpha\|\nabla^2 u\|_{\tilde{L}_{q_0}^{p_0}(0,t)} \\ &+ c_1 \int_0^t (t-s)^{-\frac{1}{2}} (e + \|u_s\|_{\mathcal{E}_b^1})\{\log(e + \|u_s\|_{\mathcal{E}_b^1})\}^{\theta} ds, \quad t \in [0, T_n]. \end{aligned}$$

Since Hölder's inequality implies

$$\begin{aligned} & \int_0^t (t-s)^{-\frac{1}{2}} (e + \|u_s\|_{\mathcal{E}_b^1}) \{\log(e + \|u_s\|_{\mathcal{E}_b^1})\}^\theta ds \\ & \leq \left(\int_0^t (t-s)^{-\frac{1}{2(1-\theta)}} ds \right)^{1-\theta} \left(\int_0^t (e + \|u_s\|_{\mathcal{E}_b^1})^{\theta-1} \log(e + \|u_s\|_{\mathcal{E}_b^1}) ds \right)^\theta, \end{aligned}$$

by $\theta \in (0, \frac{1}{2})$, when $\alpha \in (0, \frac{1}{c})$ we find a constant $c_2 > 0$ such that (5.14) implies

$$(5.15) \quad \begin{aligned} & (e + \|u_t\|_{\mathcal{E}_b^1} + \|u\|_{\tilde{L}_{q_0}^{p_0}(0,t)})^{\theta-1} \\ & \leq c_2 + c_2 \int_0^t (e + \|u_s\|_{\mathcal{E}_b^1})^{\theta-1} \log(e + \|u_s\|_{\mathcal{E}_b^1}) dr, \quad t \in [0, T_n], n \geq 1. \end{aligned}$$

By Gronwall's inequality, we obtain

$$(e + \|u_t\|_{\mathcal{E}_b^1})^{\theta-1} \leq c_2 e^{c_2 \int_0^t \log(e + \|u_s\|_{\mathcal{E}_b^1}) dr}, \quad t \in [0, T_n], n \geq 1.$$

Thus, there exists a constant $c_3 > 0$ such that

$$\log(e + \|u_t\|_{\mathcal{E}_b^1}) \leq c_3 + c_3 \int_0^t \log(e + \|u_s\|_{\mathcal{E}_b^1}) dr, \quad t \in [0, T_n], n \geq 1.$$

Using Gronwall's inequality again, we derive

$$\log(e + \|u_t\|_{\mathcal{E}_b^1}) \leq c_3 e^{c_3 T}, \quad t \in [0, T_n], n \geq 1.$$

Therefore, there exists a constant $c_4 > 0$ such that

$$(5.16) \quad \sup_{t \in [0, T_n]} \|u_t\|_{\mathcal{E}_b^1} \leq c_4, \quad n \geq 1.$$

By (5.9), if $T_n < T$ then this implies $n \leq c_4$. Thus, $T_n \geq T$ for $n > c_4$. Hence $T^* = T_n = T$ for large $n \geq 1$, so that (2.8) follows from (5.15) and (5.16). □

References

- [1] V. Barbu, M. Röckner, *From nonlinear Fokker-Planck equations to solutions of distribution dependent SDE*, Ann. Probab. 48(2020), 1902–1920.
- [2] J. M. Bismut, *Conjugate convex functions in optimal stochastic control*, J. Math. Anal. App. 44(1973), 384–404.
- [3] P. Cheridito, H. M. Soner, N. Touzi, N. Victoir, *Second-order backward stochastic differential equations and fully nonlinear parabolic PDEs*, Comm. Pure Appl. Math. 60(2007), 1081–1110.
- [4] P. Constantin, G. Iyer, *A stochastic representation of the three-dimensional incompressible Navier-Stokes equations*, Comm. Pure Appl. Math. 61(2008), 330–345.

- [5] U. Frisch, *Turbulence. The legacy of A. N. Kolmogorov*, Cambridge University Press, Cambridge (1995).
- [6] B. Gilding, M. Guedda, R. Kersner, *The Cauchy problem for $u_t = \Delta u + |\nabla u|^q$* , J. Math. Anal. Appl. 284(2003), 733–755.
- [7] K. Itô, *Differential equations determining a Markoff process*, J. Pan-Jpn. Math. Coll. No. 1077(1942). In Kiyosi Itô: Selected Papers, Springer (1987).
- [8] A. N. Kolmogorov, *Local structure of turbulence in an incompressible fluid at very high Reynolds numbers*, Dokl. Akad. Nauk SSSR 30(1941), 299–303.
- [9] J. Krug, H. Spohn, *Universality classes for deterministic surface growth*, Phys. Rev. A(3) 38(1988), 4271–4283.
- [10] H. P. McKean, *A class of Markov processes associated with nonlinear parabolic equations*, Proc. Nat. Acad. Sci. U.S.A. 56 (1966), 1907–1911.
- [11] E. Pardoux, S. Peng, *Adapted solutions of a backward stochastic differential equation*, Syst. Control Lett. 14(1990), 55–61.
- [12] S. Peng, *G-expectation, G-Brownian motion and related stochastic calculus of Itô's type*, in: Benth et al. (eds.) The Abel Symposia, pp. 541–567. Springer (2007).
- [13] S. Peng, *Nonlinear Expectations and Stochastic Calculus under Uncertainty*, Springer 2019.
- [14] Z. Qian, E. Süli, Y. Zhang, *Random vortex dynamics via functional stochastic differential equations*, arXiv:2201.00448v1.
- [15] F.-Y., Wang, *Regularity estimates and intrinsic-Lions derivative formula for singular McKean-Vlasov SDEs*, arXiv:arXiv:2109.02030.
- [16] F.-Y. Wang, *Distribution dependent SDEs for Navier-Stokes type equations*, Electron. Commun. Probab. 27(2022), 1–12.
- [17] P. Xia, L. Xie, X. Zhang, G. Zhao, *$L^q(L^p)$ -theory of stochastic differential equations*, Stoch. Proc. Appl. 130(2020), 5188–5211.
- [18] C. Yuan, S.-Q. Zhang, *A study on Zvonkin's transformation for stochastic differential equations with singular drift and related applications*, J. Diff. Equat. 297(2021), 277–319.