

GLOBAL WELL-POSEDNESS FOR TWO-DIMENSIONAL FLOWS OF VISCOELASTIC RATE-TYPE FLUIDS WITH STRESS DIFFUSION

MIROSLAV BULÍČEK, JOSEF MÁLEK, CASEY RODRIGUEZ

ABSTRACT. We consider the system of partial differential equations governing two-dimensional flows of a robust class of viscoelastic rate-type fluids with stress diffusion, involving a general objective derivative. The studied system generalizes the incompressible Navier–Stokes equations for the fluid velocity \mathbf{v} and pressure p by the presence of an additional term in the constitutive equation for the Cauchy stress expressed in terms of a positive definite tensor \mathbf{B} . The tensor \mathbf{B} evolves according to a diffusive variant of an equation that can be viewed as a combination of corresponding counterparts of Oldroyd-B and Giesekus models. Considering spatially periodic problem, we prove that for arbitrary initial data and forcing in appropriate L^2 spaces, there exists a unique globally defined weak solution to the equations of motion, and more regular initial data and forcing launch a more regular solution with \mathbf{B} positive definite everywhere.

In memory of Antonín Novotný

1. INTRODUCTION

1.1. Viscoelastic rate-type fluids with stress diffusion. Many real fluids are not Newtonian as they cannot be accurately modeled by the standard Navier–Stokes equations. One family of models frequently encountered in the theory of such *non-Newtonian fluids* is the family of incompressible viscoelastic rate-type models with stress diffusion, as these fluids are capable of describing (non-Newtonian) phenomena such as stress relaxation, nonlinear creep, normal stress differences, shear or vorticity banding. For these models, the governing equations of motion generalize the Navier–Stokes equations for the fluid velocity \mathbf{v} (and the mean normal stress p) in the sense that the constitutive equation for the Cauchy stress contains additional terms expressed in terms of a positive definite second order tensor \mathbf{B} modeling the “elastic” energy storage mechanism of the fluid. The evolution of \mathbf{B} is then described by an additional diffusive equation containing the objective time derivative of \mathbf{B} having the form of a nonlinear operator involving the velocity gradient. We refer the reader to [5] and [14] (and the references cited therein) for an introduction to the mechanics of viscoelastic rate-type fluids with stress diffusion and their applications.

In a recent study [1], the authors presented two results of apparently different nature. First of all, using the principles of continuum thermodynamics, they developed a robust class of viscoelastic-rate type models with stress diffusion that, for small elastic responses, coincide with the combination of Oldroyd-B and Giesekus models with stress diffusion, being in the form of the Laplace operator. Robustness of the developed models consists in considering a very general form of the nonlinear objective tensorial time derivative that includes the upper convected Maxwell, Jaumann–Zaremba and Gordon–Schowalter derivatives as special cases. Second, for such robust class of incompressible fluids, they established long-time and large-data existence of weak solution to three-dimensional flows in a closed bounded domains. Thus, the mathematical foundation established by Leray and his followers for the three-dimensional Navier–Stokes equations (see [2,9,12,22]) is, due to [1], available for large class of complex fluids as well.

In this paper, we focus on two-dimensional flows for the same class of fluids (i.e. those studied in [1]) asking the question of existence, uniqueness and full regularity of weak solution for any regular enough data. We give the affirmative answer to this question. For brevity, we investigate the initial, spatially periodic problem. The whole problem can be then formulated in the following way.

Let $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$ denote a two-dimensional torus. Let $a \in \mathbb{R}$, $\beta \in (0, 1)$ and $\delta_1, \delta_2 \geq 0$. Then, for $(t, x) \in [0, T] \times \mathbb{T}^2$ and for given $\mathbf{v}_0 : \mathbb{T}^2 \rightarrow \mathbb{R}^2$, $\mathbf{B}_0 : \mathbb{T}^2 \rightarrow \mathbb{R}^{2 \times 2}$ and $\mathbf{f} : [0, T] \times \mathbb{T}^2 \rightarrow \mathbb{R}^2$, we look for functions \mathbf{v} , p and \mathbf{B} satisfying

$$\operatorname{div} \mathbf{v} = 0, \quad (1.1)$$

$$\partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} = \operatorname{div} \mathbf{T} + \mathbf{f}, \quad (1.2)$$

$$\text{with } \mathbf{T} = -p\mathbf{I} + 2\mathbf{D}(\mathbf{v}) + 2a[(1 - \beta)(\mathbf{B} - \mathbf{I}) + \beta(\mathbf{B}^2 - \mathbf{B})], \quad (1.3)$$

$$\overset{\diamond}{\mathbf{B}} + \delta_1(\mathbf{B} - \mathbf{I}) + \delta_2(\mathbf{B}^2 - \mathbf{B}) = \Delta \mathbf{B}, \quad (1.4)$$

$$\mathbf{v}(0, x) = \mathbf{v}_0(x), \quad \mathbf{B}(0, x) = \mathbf{B}_0(x), \quad (1.5)$$

where, for any tensor \mathbf{A} , the symbol $\overset{\diamond}{\mathbf{A}}$ denotes an objective time derivative¹ of the symmetric, positive definite second order tensor field \mathbf{A} given by

$$\overset{\diamond}{\mathbf{A}} = \partial_t \mathbf{A} + (\mathbf{v} \cdot \nabla) \mathbf{A} - a(\mathbf{D}(\mathbf{v})\mathbf{A} + \mathbf{A}\mathbf{D}(\mathbf{v})) - (\mathbf{W}(\mathbf{v})\mathbf{A} - \mathbf{A}\mathbf{W}(\mathbf{v})).$$

The symmetric and skew parts $\mathbf{D}(\mathbf{v})$ and $\mathbf{W}(\mathbf{v})$ of the velocity gradient $\nabla \mathbf{v}$ are defined as

$$\mathbf{D}(\mathbf{v}) = \frac{1}{2}[(\nabla \mathbf{v}) + (\nabla \mathbf{v})^T] \quad \text{and} \quad \mathbf{W}(\mathbf{v}) = \frac{1}{2}[(\nabla \mathbf{v}) - (\nabla \mathbf{v})^T].$$

¹The additional terms $-a(\mathbf{D}(\mathbf{v})\mathbf{B} + \mathbf{B}\mathbf{D}(\mathbf{v})) - (\mathbf{W}(\mathbf{v})\mathbf{B} - \mathbf{B}\mathbf{W}(\mathbf{v}))$ in the definition of the objective time derivative guarantee that $\overset{\diamond}{\mathbf{B}}$ satisfies the principle of material frame-indifference (see [23] for a discussion of this principle in mechanics).

Equation (1.1) expresses the incompressibility constraint, equation (1.2) expresses the balance of linear momentum where the constant density is set to be one for simplicity, equation (1.3) is the constitutive equation for the Cauchy stress \mathbf{T} where, again for simplicity, we set the viscosity to be one, and equation (1.4) describes the evolution of the “elastic” part of the overall deformation of the fluid (see [19] and [13] for more details providing clear physical interpretation of \mathbf{B} associated with the concept of evolving natural configuration). The last equation (1.5) characterizes the initial state of the system.

As usual, for spatially periodic problem associated with the equations of the Navier–Stokes type (see [21]), we assume that

$$\mathbf{v}_0, \mathbf{f}(t, \cdot), \mathbf{v}(t, \cdot) \text{ and } p(t, \cdot) \text{ have zero mean value over } \mathbb{T}^2.$$

When $\mathbf{B} = \mathbf{I}$, then the governing equations reduce to the standard incompressible Navier–Stokes equations. Note that the Navier–Stokes equations are decoupled from the equation (1.4) for \mathbf{B} if we set $a = 0$ in the formula for the Cauchy stress (1.3). Although usually the parameter a does not occur in the formula for the Cauchy stress, Larson [11] (see pages 131–133) provided arguments why the parameter a should be there. The analysis presented in [1] and in this study holds for all $a \in \mathbb{R}$. If the reader would prefer to consider the model without the presence of a in the form for the Cauchy stress, then the results proved in [1] and herein hold for $a > 0$. It is also worth mentioning that the limiting cases of the model we consider include diffusive variants of the following:

- Oldroyd-B model [16] ($a = 1, \beta = 0, \delta_1 > 0, \delta_2 = 0$),
- Giesekus model [8] ($a = 1, \beta = 0, \delta_1 = 0, \delta_2 > 0$),
- Johnson-Segalman model [10] ($a \in [-1, 1], \beta = 0$).

The results proved in [1] and herein require that $\beta \in (0, 1)$. This essential assumption is linked with a modification of the constitutive equation for the Helmholtz free energy (see (1.7) below). For small elastic strains (i.e. when $|\mathbf{B} - \mathbf{I}|$ is small) the elastic responses of classical models (with $\beta = 0$) and the elastic responses of the models considered here coincide, see [1] for details.

Note that the diffusive Johnson-Segalman model (and consequently also its generalization) is perceived to be the model of the *shear-banding phenomenon*, which is a phenomenon of eminent interest in the applications; see for example [4, 7, 17, 18].

A basic and essential feature for the specification of the concept of weak solution is the energy balance for the system of governing equations that are being considered. In our case, for sufficiently regular solutions to (1.1)–(1.5), the energy balance takes the form

$$\frac{d}{dt} \int_{\mathbb{T}^2} \frac{1}{2} |\mathbf{v}|^2 dx + \frac{d}{dt} \int_{\mathbb{T}^2} \psi(\mathbf{B}) dx = - \int_{\mathbb{T}^2} \xi dx + \int_{\mathbb{T}^2} \mathbf{f} \cdot \mathbf{v} dx \quad (1.6)$$

where the *Helmholtz free energy* ψ is

$$\psi(\mathbf{B}) = (1 - \beta)(\text{tr}\mathbf{B} - 2 - \log \det \mathbf{B}) + \frac{\beta}{2}|\mathbf{B} - \mathbf{I}|^2 \quad (1.7)$$

and non-negative *dissipation rate* ξ is

$$\begin{aligned} \xi &= 2|\mathbf{D}(\mathbf{v})|^2 + (1 - \beta)|\mathbf{B}^{-1/2}\nabla\mathbf{B}\mathbf{B}^{-1/2}|^2 + \beta|\nabla\mathbf{B}|^2 \\ &\quad + (1 - \beta)\delta_1|\mathbf{B}^{1/2} - \mathbf{B}^{-1/2}|^2 + \beta\delta_2|\mathbf{B}^{3/2} - \mathbf{B}^{1/2}|^2 \\ &\quad + (\beta\delta_1 + (1 - \beta)\delta_2)|\mathbf{B} - \mathbf{I}|^2. \end{aligned} \quad (1.8)$$

This can be seen by summing (1.2) scalarly multiplied by \mathbf{v} and (1.4) scalarly multiplied by $\mathbf{J} = \partial_{\mathbf{B}}\psi(\mathbf{B}) = (1 - \beta)(1 - \mathbf{B}^{-1}) + \beta(\mathbf{B} - \mathbf{I})$ followed by the integration over \mathbb{T}^2 and integration by parts, using (1.1). See [1] for details. Note that for the Navier–Stokes equations, when $\mathbf{B} = \mathbf{I}$, $\psi = 0$ and ξ contains only the first term in (1.8). For $\mathbf{B} \neq \mathbf{I}$, the structure of energy storing and dissipative mechanisms is obviously very complex and nontrivial. Also, by comparing the models with $\beta = 0$ to our model with $\beta \in (0, 1)$, we observe that the models with $\beta \in (0, 1)$ has an attractive mathematical feature that the total energy $\int_{\mathbb{T}^2} \frac{1}{2}|\mathbf{v}|^2 dx + \int_{\mathbb{T}^2} \psi(\mathbf{B}) dx$ and dissipation rate ξ are coercive in norms of \mathbf{v} and \mathbf{B} of *equal* regularity.

In fact, there is a thermodynamical approach that specifies the form of the Cauchy stress tensor \mathbf{T} (i.e a *tensorial* quantity) from the knowledge of the constitutive equations for two *scalars* ψ and ξ (such as (1.7) and (1.8) above). For more details regarding this process of determining the form of \mathbf{T} for the fluids based on positing the mechanisms of energy storage and dissipation (two scalar functions), we refer the reader to [19], [20], [15], [13] and [14].

1.2. Main results and outline. In [1], there was established the long-time existence of weak solutions (for large initial data) to (1.1)–(1.5) considered on three-dimensional Lipschitz domains for Navier slip boundary conditions. A major difficulty in using the energy balance (1.6) in an approximation scheme is justifying that \mathbf{B}^{-1} exists almost everywhere and is in an appropriate test space. The work [1] introduced a novel two-step approximation scheme for (\mathbf{v}, \mathbf{B}) with parameters $(k, \epsilon) \in \mathbb{N} \times (0, 1)$ involving the minimal eigenvalue $\Lambda(\mathbf{B})$ of \mathbf{B} so that:

- in step one, a uniform-in- k energy estimate is obtained by pairing a certain Galerkin system only with the approximates $(\mathbf{v}_\epsilon^k, \mathbf{B}_\epsilon^k)$,
- in step two, the weak limit $(\mathbf{v}_\epsilon, \mathbf{B}_\epsilon)$ is shown to satisfy $\Lambda(\mathbf{B}_\epsilon) \geq \epsilon$ almost everywhere allowing for an approximate form of (1.6) (uniform-in- ϵ) to be obtained.

In this work we push this methodology further and establish the global existence and uniqueness of weak solutions to (1.1)–(1.5) for large initial data in two dimensions. We work on \mathbb{T}^2 for simplicity, but it is likely one can incorporate boundaries and different boundary conditions consistent with the second law of thermodynamics. In Section 2 we give the precise

definition of weak solutions we study (see Definition 2.1) and statements of our main results: global existence and uniqueness of weak solutions (see Theorem 2.2) and higher regularity of weak solutions (see Theorem 2.3). In Section 3 we prove uniqueness of weak solutions using a combination of Hölder's inequality, Sobolev embedding and interpolation. An adaptation of the Galerkin approximation scheme introduced in [1], where the existence of weak solutions was established for large data, is given in the appendix. Finally, in Section 4 we establish simple propagation of regularity results (see Lemma 4.1 and Lemma 4.2) and prove that smooth initial data and forcing launch smooth solutions with \mathbf{B} positive definite everywhere, rather than only almost everywhere (see Proposition 2.3), proving Theorem 2.3.

We comment that the proof of our main result can be contrasted with the work [3] on the standard Oldroyd-B model in two dimensions. There, Constantin and Kliegl establish global well-posedness for strong solutions by using the maximum principle to verify \mathbf{B} is positive definite everywhere and using (1.6) to obtain sufficient a priori bounds. In contrast, for our model we prove that a global weak solution exists, is unique, and a posteriori, we show that more regular data launches a more regular solution with \mathbf{B} positive definite everywhere. Finally, we remark that our main results along with [1] effectively prove that solutions to (1.1)–(1.5) enjoy at least the same level of regularity as solutions to the standard incompressible Navier–Stokes equations in dimensions two and three.

2. WEAK SOLUTIONS

2.1. Notation. In what follows we denote scalar, vector and tensor quantities by a , \mathbf{a} and \mathbf{A} respectively. We use the same notation for function spaces of scalar, vector or tensor-valued functions, but the context should be clear. The Lebesgue and L^2 -based Sobolev spaces on \mathbb{T}^2 are denoted by L^p and H^s , respectively. The L^2 pairing and norm are denoted by (\cdot, \cdot) and $\|\cdot\|$ respectively, and the pairing between dual spaces (clear from the context) is denoted by $\langle \cdot, \cdot \rangle$. We denote

$$L_0^2 = \left\{ \mathbf{v} \in L^2 : \int_{\mathbb{T}^2} \mathbf{v} \, dx = \mathbf{0} \right\},$$

and for $s \geq 0$,

$$\begin{aligned} H_{0,\text{div}}^s &= \left\{ \mathbf{v} \in H^s \cap L_0^2 : \text{div } \mathbf{v} = 0 \right\}, \\ H_{0,\text{div}}^{-s} &= \left(H_{0,\text{div}}^s \right)^*. \end{aligned}$$

The second condition in the definition of $H_{0,\text{div}}^s$ is interpreted in the sense of distributions. In addition, we denote the set of 2×2 real symmetric matrices by Sym and the set of 2×2 real symmetric positive definite matrices by

Sym_+ . Moreover, we define

$$\begin{aligned} H_{\text{Sym}}^s &= \{\mathbf{A} \in H^s : \mathbf{A} \in \text{Sym}\}, \\ L_{\text{Sym}}^p &= \{\mathbf{A} \in L^p : \mathbf{A} \in \text{Sym}\}. \end{aligned}$$

For $\mathbf{A} \in \mathbb{R}^{2 \times 2}$, we denote

$$\begin{aligned} \mathbf{S}(\mathbf{A}) &= (1 - \beta)(\mathbf{A} - \mathbf{I}) + \beta(\mathbf{A}^2 - \mathbf{A}), \\ \mathbf{R}(\mathbf{A}) &= \delta_1(\mathbf{A} - \mathbf{I}) + \delta_2(\mathbf{A}^2 - \mathbf{A}). \end{aligned}$$

We note that the Cauchy stress tensor can then be written as

$$\mathbf{T} = -p\mathbf{I} + 2\mathbf{D}(\mathbf{v}) + 2a\mathbf{S}(\mathbf{B}).$$

2.2. The definition of weak solutions and statement of the main result.

Definition 2.1. Let $T > 0$. Let $\beta \in (0, 1)$, $\delta_1, \delta_2 \geq 0$, $a \in \mathbb{R}$. Let $\mathbf{f} \in L^2(0, T; H_{0,\text{div}}^{-1})$. A pair $(\mathbf{v}, \mathbf{B}) : [0, T] \rightarrow \mathbb{R}^2 \times \text{Sym}_+$ is a weak solution to (1.1)–(1.5) if the following hold:

- (a) $\mathbf{v} \in L^2(0, T; H_{0,\text{div}}^1) \cap C([0, T]; L^2)$ and $\partial_t \mathbf{v} \in L^2(0, T; H_{0,\text{div}}^{-1})$,
- (b) $\mathbf{B} \in L^2(0, T; H_{\text{Sym}}^1) \cap C([0, T]; L^2)$ and $\partial_t \mathbf{B} \in [L^2(0, T; H^1) \cap L^4(0, T; L^4)]^*$,
- (c) the following two identities hold:

$$\begin{aligned} & \int_0^T [\langle \partial_t \mathbf{v}, \boldsymbol{\varphi} \rangle + ((\mathbf{v} \cdot \nabla) \mathbf{v}, \boldsymbol{\varphi})] dt \\ &= - \int_0^T (\nabla \mathbf{v}, \nabla \boldsymbol{\varphi}) - 2a(\mathbf{S}(\mathbf{B}), \nabla \boldsymbol{\varphi}) + \langle \mathbf{f}, \boldsymbol{\varphi} \rangle dt, \end{aligned} \quad (2.1)$$

for all $\boldsymbol{\varphi} \in L^2(0, T; H_{0,\text{div}}^1)$,

$$\begin{aligned} & \int_0^T [\langle \partial_t \mathbf{B}, \mathbf{A} \rangle + ((\mathbf{v} \cdot \nabla) \mathbf{B} + 2\mathbf{B}\mathbf{W}(\mathbf{v}) - 2a\mathbf{B}\mathbf{D}(\mathbf{v}), \mathbf{A})] dt \\ &+ \int_0^T (\mathbf{R}(\mathbf{B}), \mathbf{A}) dt = - \int_0^T (\nabla \mathbf{B}, \nabla \mathbf{A}) dt. \end{aligned} \quad (2.2)$$

for all $\mathbf{A} \in L^2(0, T; H_{\text{Sym}}^1) \cap L^4(0, T; L^4)$

A few remarks in order. The pressure p does not appear in Definition 2.1 since we are essentially considering weak solutions to

$$\begin{aligned} \partial_t \mathbf{v} + P((\mathbf{v} \cdot \nabla) \mathbf{v}) &= \Delta \mathbf{v} + 2aP(\text{div } \mathbf{S}(\mathbf{B})) + \mathbf{f}, \\ \hat{\diamond} \mathbf{B} + \mathbf{R}(\mathbf{B}) &= \Delta \mathbf{B}, \end{aligned}$$

where $P : L_0^2 \rightarrow L_{0,\text{div}}^2$ is the Leray projector. However, if the external forces \mathbf{f} are slightly more regular, then the pressure can be defined a posteriori (see Lemma 4.1 or proof of Theorem 2.2).

We also recall the Ladyzhenskaya inequality valid in two dimensions

$$\|u\|_4^2 \leq C \|u\| \|u\|_{H^1}. \quad (2.3)$$

Then it is also remarkable that by Hölder's inequality, it follows that \mathbf{B} from Definition 2.1 satisfies

$$\int_0^T \|\mathbf{B}\|_{L^4}^4 dt \leq C \|\mathbf{B}\|_{L^\infty L^2}^2 \int_0^T \|\nabla \mathbf{B}\|^2 + \|\mathbf{B}\|^2 dt.$$

Thus, \mathbf{B} can be paired with $\partial_t \mathbf{B}$. By an application of the Hahn-Banach theorem, we have

$$[L^2(0, T; H^1) \cap L^4(0, T; L^4)]^* = L^2(0, T; H^{-1}) + L^{4/3}(0, T; L^{4/3}),$$

and a mollification (in time) argument yields $\mathbf{B} \in C([0, T]; L^2)$, $t \mapsto \|\mathbf{B}(t)\|^2$ is absolutely continuous and

$$\frac{d}{dt} \|\mathbf{B}(t)\|^2 = 2 \langle \partial_t \mathbf{B}(t), \mathbf{B}(t) \rangle$$

a.e. on $[0, T]$; see e.g. [6] for the standard argument involving functions in $L^2(0, T; H^1)$ with time derivatives in $L^2(0, T; H^{-1})$.

The main results of this work are the following existence and uniqueness of global weak solutions, and the existence and uniqueness of global smooth solutions for smooth initial data and forcing.

Theorem 2.2. *Let $T > 0$. Let $\beta \in (0, 1)$, $\delta_1, \delta_2 \geq 0$, $a \in \mathbb{R}$. Let*

$$\begin{aligned} \mathbf{f} &\in L^2(0, T; H_{0, \text{div}}^{-1}), \quad \mathbf{v}_0 \in L_{0, \text{div}}^2, \\ \mathbf{B}_0 : \mathbb{T}^2 &\rightarrow \text{Sym}_+, \mathbf{B}_0 \in L^2 \text{ and } \log \det \mathbf{B}_0 \in L^1. \end{aligned}$$

Then there exists a unique weak solution to (1.1)–(1.5) on $[0, T] \times \mathbb{T}^2$ satisfying $\mathbf{v}(0) = \mathbf{v}_0$, $\mathbf{B}(0) = \mathbf{B}_0$.

Theorem 2.3. *Let $\beta \in (0, 1)$, $\delta_1, \delta_2 \geq 0$, $a \in \mathbb{R}$. Suppose that \mathbf{v}_0 is a smooth vector field on \mathbb{T}^2 with $\int_{\mathbb{T}^2} \mathbf{v}_0 dx = \mathbf{0}$, \mathbf{B}_0 is a smooth, symmetric, positive definite tensor field on \mathbb{T}^2 , and \mathbf{f} is a smooth vector field on $[0, \infty) \times \mathbb{T}^2$ with $\text{div } \mathbf{f} = 0$. Then there exist a unique smooth vector field \mathbf{v} with $\int_{\mathbb{T}^2} \mathbf{v} dx = \mathbf{0}$, a unique smooth scalar function p with $\int_{\mathbb{T}^2} p dx = 0$, and a unique symmetric, positive definite tensor field \mathbf{B} on $[0, \infty) \times \mathbb{T}^2$ solving (1.1)–(1.5).*

3. UNIQUENESS OF WEAK SOLUTIONS

We first prove the uniqueness of weak solutions as defined in Section 2. The proof of existence largely follows as in the three-dimensional case [1] and can be found in the appendix of this work.

3.1. Proof of uniqueness of weak solutions. Let (\mathbf{v}, \mathbf{B}) and (\mathbf{u}, \mathbf{A}) be two weak solutions to (1.1)–(1.5) with initial data $(\mathbf{v}_0, \mathbf{B}_0)$. Define $\mathbf{w} = \mathbf{v} - \mathbf{u}$ and $\mathbf{C} = \mathbf{B} - \mathbf{A}$. Then (\mathbf{w}, \mathbf{C}) satisfy, along with the initial condition $\mathbf{w}(0) = \mathbf{0}$, $\mathbf{C}(0) = \mathbf{0}$ two weak identities:

$$\begin{aligned}
& \int_0^T [(\partial_s \mathbf{w}, \boldsymbol{\varphi}) + (\nabla \mathbf{w}, \nabla \boldsymbol{\varphi})] ds \\
&= - \int_0^T ((\mathbf{v} \cdot \nabla) \mathbf{w} + (\mathbf{w} \cdot \nabla) \mathbf{u}, \boldsymbol{\varphi}) ds \\
&\quad - 2a \int_0^T (\mathbf{S}(\mathbf{B}) - \mathbf{S}(\mathbf{A}), \nabla \boldsymbol{\varphi}) ds \\
&\text{for all } \boldsymbol{\varphi} \in L^2(0, T; H_{0, \text{div}}^1)
\end{aligned} \tag{3.1}$$

and

$$\begin{aligned}
& \int_0^T [(\partial_s \mathbf{C}, \mathbf{E}) + (\nabla \mathbf{C}, \nabla \mathbf{E})] ds \\
&= - \int_0^T ((\mathbf{v} \cdot \nabla) \mathbf{C} + (\mathbf{w} \cdot \nabla) \mathbf{A}, \mathbf{E}) ds \\
&\quad + a \int_0^T (\mathbf{D}(\mathbf{v}) \mathbf{C} + \mathbf{C} \mathbf{D}(\mathbf{v}) + \mathbf{D}(\mathbf{w}) \mathbf{A} + \mathbf{A} \mathbf{D}(\mathbf{w})), \mathbf{E}) ds \\
&\quad + \int_0^T (\mathbf{W}(\mathbf{v}) \mathbf{C} - \mathbf{C} \mathbf{W}(\mathbf{v}) + \mathbf{W}(\mathbf{w}) \mathbf{A} - \mathbf{A} \mathbf{W}(\mathbf{w})), \mathbf{E}) ds \\
&\quad - \int_0^T (\mathbf{R}(\mathbf{B}) - \mathbf{R}(\mathbf{A}), \mathbf{E}) ds \\
&\text{for all } \mathbf{E} \in L^2(0, T; H_{\text{Sym}}^1) \cap L^4(0, T; L^4).
\end{aligned} \tag{3.2}$$

Let $t \in [0, T]$. We use $\boldsymbol{\varphi} = \chi_{[0, t]} \mathbf{w}$ as a test function in (3.1) and $\mathbf{E} = \chi_{[0, t]} \mathbf{C}$ as a test function in (3.2) and estimate the individual terms. Since $\text{div } \mathbf{v} = 0$, we conclude via integration by parts that

$$\begin{aligned}
((\mathbf{v} \cdot \nabla) \mathbf{w}, \mathbf{w}) &= \frac{1}{2} \int_{\mathbb{T}^2} \mathbf{v} \cdot \nabla |\mathbf{w}|^2 dx = 0, \\
((\mathbf{v} \cdot \nabla) \mathbf{C}, \mathbf{C}) &= \frac{1}{2} \int_{\mathbb{T}^2} \mathbf{v} \cdot \nabla |\mathbf{C}|^2 dx = 0.
\end{aligned}$$

Let $\epsilon > 0$, to be specified. We focus on the remaining terms on the right hand side of (3.1). Using the point-wise estimate

$$|\mathbf{S}(\mathbf{B}) - \mathbf{S}(\mathbf{A})| \leq C(1 + |\mathbf{A}| + |\mathbf{B}|)|\mathbf{B} - \mathbf{A}|,$$

we conclude with the help of the Hölder inequality and the Ladyzhenskaya inequality (2.3) that

$$\begin{aligned}
& |-(\mathbf{w} \cdot \nabla)\mathbf{u}, \mathbf{w}) - 2a(\mathbf{S}(\mathbf{B}) - \mathbf{S}(\mathbf{A}), \nabla\mathbf{w})| \\
& \leq C \int_{\mathbb{T}^2} |\mathbf{w}|^2 |\nabla\mathbf{u}| + (1 + |\mathbf{A}| + |\mathbf{B}|) |\mathbf{C}| |\nabla\mathbf{w}| \\
& \leq C(\|\nabla\mathbf{u}\| \|\mathbf{w}\|_4^2 + (1 + \|\mathbf{A}\|_4 + \|\mathbf{B}\|_4) \|\mathbf{C}\|_4 \|\nabla\mathbf{w}\|) \\
& \leq C(\|\nabla\mathbf{u}\| \|\mathbf{w}\| \|\nabla\mathbf{w}\| + (1 + \|\mathbf{A}\|_4 + \|\mathbf{B}\|_4) \|\mathbf{C}\|^{\frac{1}{2}} (\|\mathbf{C}\|^{\frac{1}{2}} + \|\nabla\mathbf{C}\|^{\frac{1}{2}}) \|\nabla\mathbf{w}\|) \\
& \leq \epsilon(\|\nabla\mathbf{w}\|^2 + \|\nabla\mathbf{C}\|^2) + C(\|\mathbf{w}\|^2 + \|\mathbf{C}\|^2)(\|\mathbf{u}\|^2 + 1 + \|\mathbf{B}\|_4^4 + \|\mathbf{A}\|_4^4).
\end{aligned}$$

Thus, as long as ϵ is chosen sufficiently small, we can apply the Young inequality and deduce from (3.1) that for all $t \in [0, T]$

$$\begin{aligned}
& \|\mathbf{w}(t)\|^2 + \frac{199}{100} \int_0^t \|\nabla\mathbf{w}\|^2 ds \\
& \leq C \int_0^t (\|\mathbf{w}\|^2 + \|\mathbf{C}\|^2)(\|\mathbf{u}\|^2 + 1 + \|\mathbf{B}\|_4^4 + \|\mathbf{A}\|_4^4) ds \\
& \quad + \frac{1}{100} \int_0^t \|\nabla\mathbf{C}\|^2 ds. \tag{3.3}
\end{aligned}$$

where C is a constant depending on β and a .

Similarly, as we have

$$|\mathbf{R}(\mathbf{B}) - \mathbf{R}(\mathbf{A})| \leq C(1 + |\mathbf{A}| + |\mathbf{B}|) |\mathbf{B} - \mathbf{A}|$$

we can use the Hölder inequality and the Ladyzhenskaya inequality to get for the terms on the right hand side of (3.2)

$$\begin{aligned}
& -(\mathbf{w} \cdot \nabla)\mathbf{A}, \mathbf{C}) + a(\mathbf{D}(\mathbf{v})\mathbf{C} + \mathbf{C}\mathbf{D}(\mathbf{v}) + \mathbf{D}(\mathbf{w})\mathbf{A} + \mathbf{A}\mathbf{D}(\mathbf{w}), \mathbf{C}) \\
& \quad + (\mathbf{W}(\mathbf{v})\mathbf{C} - \mathbf{C}\mathbf{W}(\mathbf{v}) + \mathbf{W}(\mathbf{w})\mathbf{A} - \mathbf{A}\mathbf{W}(\mathbf{w}), \mathbf{C}) - (\mathbf{R}(\mathbf{B}) - \mathbf{R}(\mathbf{A}), \mathbf{C}) \\
& \leq C \int_{\mathbb{T}^2} |\mathbf{w}| |\nabla\mathbf{A}| |\mathbf{C}| + (1 + |\mathbf{A}| + |\mathbf{B}|) |\nabla\mathbf{v}| |\mathbf{C}|^2 + |\nabla\mathbf{w}| |\mathbf{C}| (1 + |\mathbf{A}| + |\mathbf{B}|) \\
& \leq C\|\mathbf{w}\|_4 \|\nabla\mathbf{A}\| \|\mathbf{C}\|_4 + C(1 + \|\mathbf{A}\|_4 + \|\mathbf{B}\|_4) \|\nabla\mathbf{v}\| \|\mathbf{C}\|_4^2 + \\
& \quad + C\|\nabla\mathbf{w}\| \|\mathbf{C}\|_4 (1 + \|\mathbf{A}\|_4 + \|\mathbf{B}\|_4) \\
& \leq C\|\mathbf{w}\|^{\frac{1}{2}} \|\nabla\mathbf{w}\|^{\frac{1}{2}} \|\nabla\mathbf{A}\| \|\mathbf{C}\|^{\frac{1}{2}} (\|\mathbf{C}\|^{\frac{1}{2}} + \|\nabla\mathbf{C}\|^{\frac{1}{2}}) \\
& \quad + C(1 + \|\mathbf{A}\|_4 + \|\mathbf{B}\|_4) \|\nabla\mathbf{v}\| \|\mathbf{C}\| (\|\mathbf{C}\| + \|\nabla\mathbf{C}\|) + \\
& \quad + C\|\nabla\mathbf{w}\| \|\mathbf{C}\|^{\frac{1}{2}} (\|\mathbf{C}\|^{\frac{1}{2}} + \|\nabla\mathbf{C}\|^{\frac{1}{2}}) (1 + \|\mathbf{A}\|_4 + \|\mathbf{B}\|_4) \\
& \leq \epsilon(\|\nabla\mathbf{w}\|^2 + \|\nabla\mathbf{C}\|^2) \\
& \quad + C(\|\mathbf{w}\|^2 + \|\mathbf{C}\|^2) ((1 + \|\mathbf{A}\|_4^4 + \|\mathbf{B}\|_4^4 + \|\nabla\mathbf{v}\|^2 + \|\nabla\mathbf{A}\|^2)
\end{aligned}$$

Let $\epsilon > 0$, again to be specified. Choosing ϵ sufficiently small and combining the previous estimates we conclude that for all $t \in [0, T]$,

$$\begin{aligned} & \|\mathbf{C}(t)\|^2 + \frac{199}{100} \int_0^t \|\nabla \mathbf{C}\|^2 ds \\ & \leq C \int_0^t (\|\mathbf{w}\|^2 + \|\mathbf{C}\|^2) (1 + \|\mathbf{A}\|_4^4 + \|\mathbf{B}\|_4^4 + \|\nabla \mathbf{v}\|^2 \\ & \quad + \|\nabla \mathbf{A}\|^2) ds \\ & + \frac{1}{100} \int_0^t \|\nabla \mathbf{w}\|^2 ds. \end{aligned} \tag{3.4}$$

Adding (3.3) and (3.4) yields, for all $t \in [0, T]$,

$$\begin{aligned} & \|\mathbf{w}(t)\|^2 + \|\mathbf{C}(t)\|^2 + \int_0^t (\|\nabla \mathbf{w}\|^2 + \|\nabla \mathbf{C}\|^2) ds \\ & \leq C \int_0^t (\|\mathbf{w}(s)\|^2 + \|\mathbf{C}(s)\|^2) g(s) ds, \end{aligned}$$

where

$$\begin{aligned} g &= 1 + \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 + \|\nabla \mathbf{u}\|^2 + \|\nabla \mathbf{v}\|^2 + \|\mathbf{B}\|_4^4 \\ & + \|\mathbf{A}\|_4^4 + \|\nabla \mathbf{B}\|^2 + \|\nabla \mathbf{A}\|^2. \end{aligned}$$

Now, $g \in L^1([0, T])$ by our assumptions for (\mathbf{v}, \mathbf{B}) and (\mathbf{u}, \mathbf{A}) . By Gronwall's inequality we conclude that $(\mathbf{w}, \mathbf{C}) = (\mathbf{0}, \mathbf{0})$ as desired. \square

4. HIGHER REGULARITY

In this section we prove simple propagation of regularity results for weak solutions and conclude that smooth initial data and forcing launch smooth globally defined solutions.

4.1. Propagation of regularity.

Lemma 4.1. *Let $T > 0$. Let $\beta \in (0, 1)$, $\delta_1, \delta_2 \geq 0$, $a \in \mathbb{R}$,*

$$\mathbf{f} \in L^2(0, T; L^2_{0,\text{div}}), \quad \mathbf{v}_0 \in H^1_{0,\text{div}},$$

and $\mathbf{B}_0 : \mathbb{T}^2 \rightarrow \text{Sym}_+$ such that $\log \det \mathbf{B}_0 \in L^1$, $\mathbf{B}_0 \in H^1$. Let (\mathbf{v}, \mathbf{B}) be the unique weak solution to (1.1)–(1.5) on $[0, T] \times \mathbb{T}^2$ such that

$$\mathbf{v}(0) = \mathbf{v}_0, \quad \mathbf{B}(0) = \mathbf{B}_0.$$

Then

$$\begin{aligned} & \mathbf{v} \in L^\infty(0, T; H^1) \cap L^2(0, T; H^2) \\ & \partial_t \mathbf{v} \in L^2(0, T; L^2_{0,\text{div}}), \\ & \mathbf{B} \in L^\infty(0, T; H^1) \cap L^2(0, T; H^2), \\ & \partial_t \mathbf{B} \in L^2(0, T; L^2), \end{aligned} \tag{4.1}$$

there exists a unique scalar function $p \in L^2(0, T; H^1)$ with $\int_{\mathbb{T}^2} p \, dx = 0$ such that

$$\begin{aligned} \partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} &= -\nabla p + \Delta \mathbf{v} + 2a \operatorname{div} \mathbf{S}(\mathbf{B}) + \mathbf{f}, \\ \overset{\diamond}{\mathbf{B}} + \mathbf{R}(\mathbf{B}) &= \Delta \mathbf{B}, \end{aligned} \quad (4.2)$$

a.e. on $[0, T] \times \mathbb{T}^2$, and

$$\begin{aligned} \|\mathbf{v}\|_{L^\infty H^1} + \|\mathbf{B}\|_{L^\infty L^2} + \|\mathbf{v}\|_{L^2 H^2} + \|\mathbf{B}\|_{L^2 H^2} + \|\partial_t \mathbf{v}\|_{L^2 L^2} + \|\partial_t \mathbf{B}\|_{L^2 L^2} \\ \leq C_1 (\|\mathbf{v}_0\|_{H^1}, \|\psi(\mathbf{B}_0)\|_{L^1}, \|\mathbf{B}_0\|_{H^1}, \|\mathbf{f}\|_{L^2 L^2}). \end{aligned} \quad (4.3)$$

If, in addition,

$$\mathbf{v}_0 \in H_{0,\operatorname{div}}^2, \quad \mathbf{B} \in H^2, \quad \partial_t \mathbf{f} \in L^2(0, T; L_{0,\operatorname{div}}^2), \quad (4.4)$$

then

$$\begin{aligned} \mathbf{v} &\in L^\infty(0, T; H^2) \\ \partial_t \mathbf{v} &\in L^\infty(0, T; L_{0,\operatorname{div}}^2) \cap L^2(0, T; H^1), \\ \partial_t^2 \mathbf{v} &\in L^2(0, T; H_{0,\operatorname{div}}^{-1}), \\ \mathbf{B} &\in L^\infty(0, T; H^2), \\ \partial_t \mathbf{B} &\in L^\infty(0, T; L^2) \cap L^2(0, T; H^1), \\ \partial_t^2 \mathbf{B} &\in L^2(0, T; H^{-1}), \end{aligned} \quad (4.5)$$

and

$$\begin{aligned} \|\mathbf{v}\|_{L^\infty H^2} + \|\mathbf{B}\|_{L^\infty H^2} + \|\partial_t \mathbf{v}\|_{L^\infty L^2} + \|\partial_t \mathbf{B}\|_{L^\infty L^2} \\ + \|\partial_t \mathbf{v}\|_{L^2 H^1} + \|\partial_t \mathbf{B}\|_{L^2 H^1} + \|\partial_t^2 \mathbf{v}\|_{L^2 H^{-1}} + \|\partial_t^2 \mathbf{B}\|_{L^2 H^{-1}} \\ \leq C'_1 (\|\mathbf{v}_0\|_{H^2}, \|\psi(\mathbf{B}_0)\|_{L^1}, \|\mathbf{B}_0\|_{H^2}, \|\mathbf{f}\|_{H^1 L^2}). \end{aligned} \quad (4.6)$$

Proof. We will give the arguments for (4.3) and (4.6) assuming the regularity (4.1) and (4.5) (respectively) by taking partial derivatives of the equations of motion. The proof assuming only the stated regularity of the initial data then follows from these arguments by using suitable difference quotients rather than partial derivatives.

Restating (A.18) we have

$$\begin{aligned} \|\mathbf{v}\|_{L^\infty L^2} + \|\mathbf{B}\|_{L^\infty L^2} + \|\nabla \mathbf{v}\|_{L^2 L^2} + \|\nabla \mathbf{B}\|_{L^2 L^2} \\ \leq C_0 (\|\mathbf{v}_0\|, \|\psi(\mathbf{B}_0)\|_{L^1}, \|\mathbf{f}\|_{L^2 H_{0,\operatorname{div}}^{-1}}). \end{aligned} \quad (4.7)$$

Let $i \in \{1, 2\}$, $\mathbf{w} = \partial_{x^i} \mathbf{v}$ and $\mathbf{C} = \partial_{x^i} \mathbf{B}$. Then for all $\varphi \in L^2(0, T; H_{0,\text{div}}^1)$

$$\begin{aligned} & \int_0^T [(\partial_s \mathbf{w}, \varphi) + (\nabla \mathbf{w}, \nabla \varphi)] ds \\ &= - \int_0^T ((\mathbf{v} \cdot \nabla) \mathbf{w} + (\mathbf{w} \cdot \nabla) \mathbf{v}, \varphi) ds - \int_0^T (\mathbf{f}, \partial_{x^i} \varphi) ds \\ & \quad - 2a \int_0^T ((1 - \beta) \mathbf{C} + \beta(\mathbf{C}\mathbf{B} + \mathbf{B}\mathbf{C} - \mathbf{C}), \nabla \varphi) ds, \end{aligned} \quad (4.8)$$

and for all $\mathbf{E} \in [L^2(0, T; H^1) \cap L^4(0, T; L^4)]^*$, $\mathbf{E}^T = \mathbf{E}$, we have

$$\begin{aligned} & \int_0^T [(\partial_s \mathbf{C}, \mathbf{E}) + (\nabla \mathbf{C}, \nabla \mathbf{E})] ds \\ &= - \int_0^T ((\mathbf{v} \cdot \nabla) \mathbf{C} + (\mathbf{w} \cdot \nabla) \mathbf{B}, \mathbf{E}) ds \\ & \quad + a \int_0^T (\mathbf{D}(\mathbf{w})\mathbf{B} + \mathbf{B}\mathbf{D}(\mathbf{w}) + \mathbf{D}(\mathbf{v})\mathbf{C} + \mathbf{C}\mathbf{D}(\mathbf{v})), \mathbf{E}) ds \\ & \quad + \int_0^T (\mathbf{W}(\mathbf{v})\mathbf{C} - \mathbf{C}\mathbf{W}(\mathbf{v}) + \mathbf{W}(\mathbf{w})\mathbf{B} - \mathbf{B}\mathbf{W}(\mathbf{w})), \mathbf{E}) ds \\ & \quad - \int_0^T (\delta_1 \mathbf{C} + \delta_2(\mathbf{C}\mathbf{B} + \mathbf{B}\mathbf{C} - \mathbf{C}), \mathbf{E}) ds. \end{aligned} \quad (4.9)$$

We now argue identically as in Section 3.1 and obtain the estimate, for all $t \in [0, T]$,

$$\begin{aligned} & \|\mathbf{w}(t)\|^2 + \|\mathbf{C}(t)\|^2 + \int_0^t (\|\nabla \mathbf{w}\|^2 + \|\nabla \mathbf{C}\|^2) ds \\ & \leq \|\mathbf{w}(0)\|^2 + \|\mathbf{C}(0)\|^2 + \int_0^t \|\mathbf{f}\| \|\mathbf{w}\| ds \\ & \quad + C \int_0^t (\|\mathbf{w}(s)\|^2 + \|\mathbf{C}(s)\|^2) g(s) ds \\ & \leq \|\mathbf{v}_0\|_{H^1}^2 + \|\mathbf{B}_0\|_{H^1}^2 + \int_0^t \|\mathbf{f}\| \|\mathbf{w}\| ds \\ & \quad + C \int_0^t (\|\mathbf{w}(s)\|^2 + \|\mathbf{C}(s)\|^2) g(s) ds, \end{aligned}$$

where

$$g = 1 + \|\nabla \mathbf{v}\|^2 + \|\nabla \mathbf{B}\|^2 + \|\mathbf{v}\|^2 \|\nabla \mathbf{v}\|^2 + \|\mathbf{B}\|^2 \|\nabla \mathbf{B}\|^2.$$

By Gronwall's inequality and (4.7) we conclude that

$$\begin{aligned} & \|\mathbf{w}\|_{L^\infty L^2} + \|\mathbf{C}\|_{L^\infty L^2} + \|\nabla \mathbf{w}\|_{L^2 L^2} + \|\nabla \mathbf{C}\|_{L^2 L^2} \\ & \leq C_{1/3} (\|\mathbf{v}_0\|_{H^1}, \|\psi(\mathbf{B}_0)\|_{L^1}, \|\mathbf{B}_0\|_{H^1}, \|\mathbf{f}\|_{L^2 L^2}), \end{aligned} \quad (4.10)$$

and thus,

$$\begin{aligned} & \|\mathbf{v}\|_{L^\infty H^1} + \|\mathbf{B}\|_{L^\infty H^1} + \|\mathbf{v}\|_{L^2 H^2} + \|\mathbf{B}\|_{L^2 H^2} \\ & \leq C_{2/3}(\|\mathbf{v}_0\|, \|\psi(\mathbf{B}_0)\|_{L^1}, \|\mathbf{B}_0\|_{H^1}, \|\mathbf{f}\|_{L^2 L^2}). \end{aligned} \quad (4.11)$$

The estimate for $\partial_t \mathbf{v}$ and $\partial_t \mathbf{B}$ then follows from (2.1), (2.2), (4.11) and repeated use of Hölder's inequality and the Sobolev embedding $H^2 \hookrightarrow L^\infty$: for example, we estimate the pairing

$$\begin{aligned} \int_0^T (\mathbf{B} \mathbf{D}(\mathbf{v}), \mathbf{A}) ds & \leq \int_0^T \|\mathbf{B}\|_{L^\infty} \|\mathbf{v}\|_{H^1} \|\mathbf{A}\| ds \\ & \leq \|\mathbf{B}\|_{L^2 H^2} \|\mathbf{v}\|_{L^\infty H^1} \|\mathbf{A}\|_{L^2 L^2}. \end{aligned}$$

Moreover, we can then conclude that

$$\begin{aligned} \partial_t \mathbf{v} + P((\mathbf{v} \cdot \nabla) \mathbf{v}) & = \Delta \mathbf{v} + 2aP(\operatorname{div} \mathbf{S}(\mathbf{B})) + \mathbf{f}, \\ \hat{\mathbf{B}} + \mathbf{R}(\mathbf{B}) & = \Delta \mathbf{B}, \end{aligned} \quad (4.12)$$

a.e. on $[0, T] \times \mathbb{T}^2$ where $P : L_0^2 \rightarrow L_{0, \operatorname{div}}^2$ is the Leray projector. The existence and uniqueness of the pressure p then follows immediately from the Helmholtz decomposition of L_0^2 vector fields on \mathbb{T}^2 .

We now assume the additional regularity (4.4). The argument leading to (4.10) also applies to $\mathbf{w} = \partial_t \mathbf{v}$ and $\mathbf{C} = \partial_t \mathbf{B}$ resulting in the additional estimate

$$\begin{aligned} & \|\partial_t \mathbf{v}\|_{L^\infty L^2} + \|\partial_t \mathbf{B}\|_{L^\infty L^2} + \|\partial_t \mathbf{v}\|_{L^2 H^1} + \|\partial_t \mathbf{B}\|_{L^2 H^1} \\ & \leq C'_{1/3}(\|\mathbf{v}_0\|_{H^2}, \|\psi(\mathbf{B}_0)\|_{L^1}, \|\mathbf{B}_0\|_{H^2}, \|\mathbf{f}\|_{H^1 L^2}), \end{aligned} \quad (4.13)$$

where we used (4.12), (4.4) and Sobolev embedding to bound $\|\partial_t \mathbf{v}(0)\|$ and $\|\partial_t \mathbf{B}(0)\|$. By (4.12), (4.13), Sobolev embedding and interpolation we conclude, for a.e. $t \in [0, T]$,

$$\begin{aligned} \|\Delta \mathbf{v}\| & \leq C(\|\partial_t \mathbf{v}\| + \|\mathbf{v}\|_{L^4} \|\nabla \mathbf{v}\|_{L^4} + \|\nabla \mathbf{B}\| + \|\mathbf{B}\|_{L^4} \|\nabla \mathbf{B}\|_{L^4}) \\ & \leq C(\|\partial_t \mathbf{v}\| + \|\mathbf{v}\|_{H^1}^{3/2} \|\Delta \mathbf{v}\|^{1/2} + \|\nabla \mathbf{B}\| + \|\mathbf{B}\|_{H^1}^{3/2} \|\Delta \mathbf{B}\|^{1/2}) \\ & \leq C'_{2/3} + \frac{1}{4} \|\Delta \mathbf{v}\| + \frac{1}{4} \|\Delta \mathbf{B}\|, \end{aligned} \quad (4.14)$$

and similarly

$$\|\Delta \mathbf{B}\| \leq C'_{2/3} + \frac{1}{4} \|\Delta \mathbf{v}\| + \frac{1}{4} \|\Delta \mathbf{B}\|. \quad (4.15)$$

Summing (4.14) and (4.15) yields, for a.e. $t \in [0, T]$,

$$\|\Delta \mathbf{v}\| + \|\Delta \mathbf{B}\| \leq 2C'_{2/3}, \quad (4.16)$$

proving the $L^\infty H^2$ bound. The bound for $\partial_t^2 \mathbf{v}$ and $\partial_t^2 \mathbf{B}$ then follows from (4.8) with $\mathbf{w} = \partial_t \mathbf{v}$, (4.9) with $\mathbf{C} = \partial_t \mathbf{B}$, (4.13), (4.16) and repeated applications of Hölder's inequality and Sobolev embedding: for example we

estimate the pairing

$$\begin{aligned} \int_0^T (\mathbf{B}\mathbf{D}(\partial_s \mathbf{v}), \mathbf{A}) ds &\leq \|\mathbf{B}\|_{L^\infty L^\infty} \|\partial_t \mathbf{v}\|_{L^2 H^1} \|\mathbf{A}\|_{L^2 L^2} \\ &\leq \|\mathbf{B}\|_{L^\infty H^2} \|\partial_t \mathbf{v}\|_{L^2 H^1} \|\mathbf{A}\|_{L^2 H^1}. \end{aligned}$$

This concludes the proof. \square

Via an induction argument, the Leibniz formula and repeated use of Hölder's inequality, Sobolev embedding and interpolation (as in the proof of Lemma 4.1), we have the follow generalization of Lemma 4.1.

Lemma 4.2. *Let $T > 0$, and let $m \in \mathbb{N} \cup \{0\}$. Let $\beta \in (0, 1)$, $\delta_1, \delta_2 \geq 0$, $a \in \mathbb{R}$,*

$$\begin{aligned} \mathbf{v}_0 &\in H_{0,\text{div}}^{2m+1}, \\ \partial_t^j \mathbf{f} &\in L^2(0, T; H_{0,\text{div}}^{2m-2j}), \quad j = 0, \dots, m, \end{aligned}$$

and $\mathbf{B}_0 : \mathbb{T}^2 \rightarrow \text{Sym}_+$ such that $\log \det \mathbf{B}_0 \in L^1$, $\mathbf{B}_0 \in H^{2m+1}$. Let (\mathbf{v}, \mathbf{B}) be the unique weak solution to (1.1)–(1.5) on $[0, T] \times \mathbb{T}^2$ such that

$$\mathbf{v}(0) = \mathbf{v}_0, \quad \mathbf{B}(0) = \mathbf{B}_0.$$

Then for $k = 0, \dots, m+1$,

$$\begin{aligned} \partial_t^k \mathbf{v} &\in L^2(0, T; H_{0,\text{div}}^{2m+2-2k}), \\ \partial_t^k \mathbf{B} &\in L^2(0, T; H^{2m+2-2k}). \end{aligned}$$

4.2. Proof of Theorem 2.3. Let $T > 0$. By (4.2) and Lemma 4.2 with $m = 0, 1, \dots$, we conclude there exist unique smooth \mathbf{v} , \mathbf{B} and p satisfying (1.1)–(1.5) on $[0, T]$ (the regularity of p following from that of \mathbf{v} and \mathbf{B} and (1.2)). Since $T > 0$ was arbitrary we conclude that there exist unique smooth \mathbf{v} , \mathbf{B} and p satisfying (1.1)–(1.5) on $[0, \infty)$. Moreover, \mathbf{B} is positive definite almost everywhere on $[0, \infty) \times \mathbb{T}^2$.

We now prove that \mathbf{B} is positive definite everywhere by performing a calculation similar to that done for the standard Oldroyd-B model in [3]. We write

$$\mathbf{B} = \begin{pmatrix} \frac{g}{2} + e & f \\ f & \frac{g}{2} - e \end{pmatrix}, \quad a\mathbf{D}(\mathbf{v}) + \mathbf{W}(\mathbf{v}) = \begin{pmatrix} \alpha & \beta \\ \gamma & -\alpha \end{pmatrix}.$$

From (1.4) we conclude, with $D_t = \partial_t + \mathbf{v} \cdot \nabla$, that

$$\begin{aligned} D_t e &= \Delta e - (\delta_1 + (g-1)\delta_2)e + (\beta - \gamma)f + \alpha g, \\ D_t f &= \Delta f - (\delta_1 + (g-1)\delta_2)f - (\beta - \gamma)e + \frac{1}{2}(\beta + \gamma)g, \\ D_t g &= \Delta g - (\delta_1 + (g-1)\delta_2)g + 4\alpha e + 2(\beta + \gamma)f \\ &\quad + 2\delta_1 + 2\delta_2 \left(\frac{g^2}{4} - (e^2 + f^2) \right). \end{aligned} \tag{4.17}$$

The maximum and minimum eigenvalues of \mathbf{B} are given by

$$\lambda_{max} = \frac{g}{2} + (e^2 + f^2)^{1/2}, \quad \lambda_{min} = \frac{g}{2} - (e^2 + f^2)^{1/2}.$$

We note that

$$\begin{aligned} & \Delta(e^2 + f^2)^{1/2} - (e^2 + f^2)^{-1/2}e\Delta e - (e^2 + f^2)^{-1/2}f\Delta f \\ &= (e^2 + f^2)^{-1/2} \sum_{j=1}^2 (e\partial_j f - f\partial_j e)^2 \geq 0. \end{aligned} \quad (4.18)$$

Using (4.17) and (4.18) we conclude

$$\begin{aligned} & D_t \lambda_{min} + \Delta \lambda_{min} \\ &= -[\delta_1 + (g-1)\delta_2 + 2\alpha e(e^2 + f^2)^{-1/2} + (\beta + \gamma)f(e^2 + f^2)^{-1/2}] \lambda_{min} \\ & \quad + \delta_1 + \delta_2 \lambda_{max} \lambda_{min} \\ & \quad + \Delta(e^2 + f^2)^{1/2} - (e^2 + f^2)^{-1/2}e\Delta e - (e^2 + f^2)^{-1/2}f\Delta f \\ & \geq -[\delta_1 + (g-1)\delta_2 + 2\alpha e(e^2 + f^2)^{-1/2} + (\beta + \gamma)f(e^2 + f^2)^{-1/2}] \lambda_{min} \\ & \quad + \delta_1 + \delta_2 \lambda_{max} \lambda_{min}. \end{aligned}$$

Since $\lambda_{min}(0, x) > 0$ by assumption that \mathbf{B}_0 is positive definite, the maximum principle implies that $\lambda_{min}(t, x) > 0$ on $[0, \infty)$. This proves that \mathbf{B} is positive definite on $[0, \infty)$ and concludes the proof of Theorem 2.3. \square

APPENDIX A. EXISTENCE OF WEAK SOLUTIONS

In this appendix we prove the existence of weak solutions via an approximation scheme.

A.1. Galerkin system. An orthonormal basis for $L^2_{0,\text{div}}$ is given by

$$\{n^\perp e^{2\pi i n \cdot x} / |n|\}_{n \in \mathbb{Z}^2 \setminus (0,0)}$$

where $n^\perp = -n_2 \mathbf{i} + n_1 \mathbf{j}$. We denote the projection of $\mathbf{v} \in L^2_{0,\text{div}}$ onto

$$\text{Span}\{n^\perp e^{2\pi i n \cdot x} / |n|\}_{|n| \leq k}$$

by $P_k \mathbf{v}$. We denote the projection of $\mathbf{B} \in L^2$ onto

$$\text{Span}\left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} e^{2\pi i n \cdot x}, \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} e^{2\pi i n \cdot x}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} e^{2\pi i n \cdot x} \right\}_{|n| \leq k}$$

by $Q_k \mathbf{B}$.

For $\epsilon > 0$, $\mathbf{A} \in \mathbb{R}^{2 \times 2}$, $\mathbf{A}^T = \mathbf{A}$, we denote the minimal eigenvalue of \mathbf{A} by $\Lambda(\mathbf{A})$ and define

$$\rho_\epsilon(\mathbf{A}) = \frac{\max\{0, \Lambda(\mathbf{A}) - \epsilon\}}{\Lambda(\mathbf{A})(1 + \epsilon|\mathbf{A}|^3)}.$$

Let $\epsilon > 0$, $k \in \mathbb{N}$. By an application of Carathéodory's theorem for ordinary differential equations, it follows that there exist $T^* \in (0, T]$ and unique

$$\begin{aligned} \mathbf{v}_\epsilon^k &\in \text{Span}\{n^\perp e^{2\pi i n \cdot x} / |n|\}_{|n| \leq k}, \\ \mathbf{B}_\epsilon^k &\in \text{Span}\left\{\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} e^{2\pi i n \cdot x}, \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} e^{2\pi i n \cdot x}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} e^{2\pi i n \cdot x}\right\}_{|n| \leq k}, \end{aligned}$$

with absolutely continuous Fourier coefficients solving the following Galerkin system on $[0, T^*]$:

$$\begin{aligned} \partial_t \mathbf{v}_\epsilon^k + P_k(\mathbf{v}_\epsilon^k \cdot \nabla \mathbf{v}_\epsilon^k) - \Delta \mathbf{v}_\epsilon^k \\ = 2a P_k[\text{div}(\rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{S}(\mathbf{B}_\epsilon^k))] + P_k \mathbf{f}, \end{aligned} \quad (\text{A.1})$$

$$\begin{aligned} \partial_t \mathbf{B}_\epsilon^k + Q_k(\mathbf{v}_\epsilon^k \cdot \nabla \mathbf{B}_\epsilon^k) - \Delta \mathbf{B}_\epsilon^k \\ = a Q_k[\rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{D}(\mathbf{v}_\epsilon^k) \mathbf{B}_\epsilon^k + \rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{B}_\epsilon^k \mathbf{D}(\mathbf{v}_\epsilon^k)] \\ + Q_k[\rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{W}(\mathbf{v}_\epsilon^k) \mathbf{B}_\epsilon^k - \rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{B}_\epsilon^k \mathbf{W}(\mathbf{v}_\epsilon^k)] \\ - Q_k[\rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{R}(\mathbf{B}_\epsilon^k)], \end{aligned} \quad (\text{A.2})$$

with initial conditions

$$\begin{aligned} \mathbf{v}_\epsilon^k(0, x) &= P_k \mathbf{v}_0(x), \quad \mathbf{B}_\epsilon^k(0, x) = Q_k \mathbf{B}_{0, \epsilon}(x), \\ \mathbf{B}_{0, \epsilon}(x) &= \begin{cases} \mathbf{B}_0(x) & \text{if } \Lambda(\mathbf{B}_0(x)) > \epsilon, \\ \mathbf{I} & \text{otherwise.} \end{cases} \end{aligned}$$

Lemma A.1. *For all $\epsilon > 0$, $k \in \mathbb{N}$, there exist constants $C_0(\epsilon)$ and C_1 (absolute) such that the solution $(\mathbf{v}_\epsilon^k, \mathbf{B}_\epsilon^k)$ to the Galerkin system (A.1), (A.2) satisfies*

$$\begin{aligned} \sup_{t \in [0, T^*]} (\|\mathbf{v}_\epsilon^k(t)\|^2 + \|\mathbf{B}_\epsilon^k(t)\|^2) + \int_0^{T^*} (\|\nabla \mathbf{v}_\epsilon^k\|^2 + \|\nabla \mathbf{B}_\epsilon^k\|^2) dt \\ \leq C_0(\epsilon) + \|\mathbf{v}_0\|^2 + \|\mathbf{B}_0\|^2 + C_1 \int_0^{T^*} \|\mathbf{f}\|_{H_{0, \text{div}}^{-1}}^2 dt. \end{aligned} \quad (\text{A.3})$$

Moreover, denoting the right-hand side of (A.3) by D , there exists a constant $C_3(D, \epsilon)$ such that

$$\int_0^{T^*} (\|\partial_t \mathbf{v}_\epsilon^k\|_{H_{0, \text{div}}^{-1}}^2 + \|\partial_t \mathbf{B}_\epsilon^k\|_{H^{-1}}^2) dt \leq C_3(D, \epsilon). \quad (\text{A.4})$$

Proof. Pairing (A.1) with \mathbf{v}_ϵ^k and (A.2) with \mathbf{B}_ϵ^k and summing we obtain the identity

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} (\|\mathbf{v}_\epsilon^k\|^2 + \|\mathbf{B}_\epsilon^k\|^2) + \|\nabla \mathbf{v}_\epsilon^k\|^2 + \|\nabla \mathbf{B}_\epsilon^k\|^2 \\ = -2a(\rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{S}(\mathbf{B}_\epsilon^k), \mathbf{D}(\mathbf{v}_\epsilon^k)) + \langle \mathbf{f}, \mathbf{v}_\epsilon^k \rangle \\ + 2a[(\rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{B}_\epsilon^k \mathbf{D}(\mathbf{v}_\epsilon^k), \mathbf{B}_\epsilon^k) - (\rho_\epsilon(\mathbf{v}_\epsilon^k) \mathbf{R}(\mathbf{B}_\epsilon^k), \mathbf{B}_\epsilon^k)]. \end{aligned} \quad (\text{A.5})$$

The definitions of ρ_ϵ , \mathbf{S} , and \mathbf{R} imply the estimate

$$\rho_\epsilon(\mathbf{B}_\epsilon^k)[|\mathbf{S}(\mathbf{B}_\epsilon^k)| + |\mathbf{B}_\epsilon^k|^2 + |\mathbf{B}_\epsilon^k||\mathbf{R}(\mathbf{B}_\epsilon^k)|] \leq C \frac{1 + |\mathbf{B}_\epsilon^k|^3}{1 + \epsilon|\mathbf{B}_\epsilon^k|^3} \leq C_0(\epsilon). \quad (\text{A.6})$$

Integrating (A.5) from 0 to $t \in [0, T^*]$ and using (A.6) to bound the right-hand side of (A.5) we obtain the estimate (A.3).

We now estimate $\partial_t \mathbf{v}_\epsilon^k$ and $\partial_t \mathbf{B}_\epsilon^k$. Let $\mathbf{w} \in H_{0,\text{div}}^1$. By Hölder's inequality, Sobolev embedding $H^{1/2} \hookrightarrow L^4$ and interpolation we obtain the estimates:

$$\begin{aligned} |(P_k((\mathbf{v}_\epsilon^k \cdot \nabla) \mathbf{v}_\epsilon^k), \mathbf{w})| &= |(\mathbf{v}_\epsilon^k, (\mathbf{v}_\epsilon^k \cdot \nabla) P_k \mathbf{w})| \\ &\leq \|\mathbf{v}_\epsilon^k\|_{L^4}^2 \|\nabla \mathbf{w}\| \\ &\leq C \|\mathbf{v}_\epsilon^k\| \|\nabla \mathbf{v}_\epsilon^k\| \|\nabla \mathbf{w}\|, \\ |(P_k[\text{div}(\rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{S}(\mathbf{B}_\epsilon^k))], \mathbf{w})| &= |(\rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{S}(\mathbf{B}_\epsilon^k), \nabla P_k \mathbf{w})| \\ &\leq C \int_{\mathbb{T}^2} (1 + |\mathbf{B}_\epsilon^k| + |\mathbf{B}_\epsilon^k|^2) |\nabla P_k \mathbf{w}| dx \\ &\leq C(1 + \|\mathbf{B}_\epsilon^k\| + \|\mathbf{B}_\epsilon^k\| \|\nabla \mathbf{B}_\epsilon^k\|) \|\nabla \mathbf{w}\|. \end{aligned}$$

These estimates and (A.1) imply

$$\int_0^{T^*} \|\partial_t \mathbf{v}\|_{H_{0,\text{div}}^{-1}}^2 dt \leq C_{7/3}(D).$$

Similarly, if $\mathbf{A} \in H^{-1}$, we obtain

$$\begin{aligned} |(Q_k((\mathbf{v}_\epsilon^k \cdot \nabla) \mathbf{B}_\epsilon^k), \mathbf{A})| &\leq C \|\mathbf{v}_\epsilon^k\|^{1/2} \|\mathbf{v}_\epsilon^k\|^{1/2} \|\mathbf{B}_\epsilon^k\|^{1/2} \|\mathbf{B}_\epsilon^k\|^{1/2} \|\nabla \mathbf{A}\|, \\ |(Q_k(\rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{R}(\mathbf{B}_\epsilon^k)), \mathbf{A})| &\leq C(1 + \|\mathbf{B}_\epsilon^k\| + \|\mathbf{B}_\epsilon^k\| \|\nabla \mathbf{B}_\epsilon^k\|) \|\nabla \mathbf{A}\|, \end{aligned}$$

and since $|\rho_\epsilon(\mathbf{B}_\epsilon^k)| \|\mathbf{B}_\epsilon^k\| \leq C_{8/3}(\epsilon)$, we obtain

$$\begin{aligned} |(aQ_k[\rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{D}(\mathbf{v}_\epsilon^k) \mathbf{B}_\epsilon^k + \rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{B}_\epsilon^k \mathbf{D}(\mathbf{v}_\epsilon^k)], \mathbf{A})| \\ \leq 2|a| C_{8/3}(\epsilon) \|\nabla \mathbf{v}_\epsilon^k\| \|\mathbf{A}\|, \\ |(aQ_k[\rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{W}(\mathbf{v}_\epsilon^k) \mathbf{B}_\epsilon^k - \rho_\epsilon(\mathbf{B}_\epsilon^k) \mathbf{B}_\epsilon^k \mathbf{W}(\mathbf{v}_\epsilon^k)], \mathbf{A})| \\ \leq 2|a| C_{8/3}(\epsilon) \|\nabla \mathbf{v}_\epsilon^k\| \|\mathbf{A}\|. \end{aligned}$$

These estimates and (A.2) imply that

$$\int_0^{T^*} (\|\partial_t \mathbf{v}\|_{H_{0,\text{div}}^{-1}}^2 + \|\partial_t \mathbf{B}_\epsilon^k\|_{H^{-1}}^2) dt \leq C_3(D, \epsilon),$$

completing the proof. \square

Lemma A.2. *For all $\epsilon > 0$, there exist*

$$\begin{aligned} \mathbf{v}_\epsilon &\in L^\infty(0, T; L_{0,\text{div}}^2) \cap L^2(0, T; H_{0,\text{div}}^1) \cap C([0, T]; H_{0,\text{div}}^{-1}), \\ \mathbf{B}_\epsilon &\in L^\infty(0, T; L^2) \cap L^2(0, T; H^1) \cap C([0, T]; H^{-1}), \end{aligned}$$

such that $\mathbf{v}_\epsilon(0, x) = \mathbf{v}_0(x)$, $\mathbf{B}_\epsilon(0, x) = \mathbf{B}_{0,\epsilon}(x)$,

$$\begin{aligned}\partial_t \mathbf{v}_\epsilon &\in L^2(0, T; H_{0,\text{div}}^{-1}), \\ \partial_t \mathbf{B}_\epsilon &\in L^2(0, T; H^{-1}),\end{aligned}$$

for all $\boldsymbol{\varphi} \in L^2(0, T; H_{0,\text{div}}^1)$,

$$\begin{aligned}&\int_0^T [\langle \partial_t \mathbf{v}_\epsilon, \boldsymbol{\varphi} \rangle + ((\mathbf{v}_\epsilon \cdot \nabla) \mathbf{v}_\epsilon, \boldsymbol{\varphi})] dt \\ &= - \int_0^T (\nabla \mathbf{v}_\epsilon, \nabla \boldsymbol{\varphi}) - 2a((\rho_\epsilon(\mathbf{B}_\epsilon)) \mathbf{S}(\mathbf{B}_\epsilon), \mathbf{D}(\boldsymbol{\varphi})) + \langle \mathbf{f}, \boldsymbol{\varphi} \rangle dt, \quad (\text{A.7})\end{aligned}$$

and for all $\mathbf{A} \in L^2(0, T; H^1)$, we have

$$\begin{aligned}&\int_0^T [\langle \partial_t \mathbf{B}_\epsilon, \mathbf{A} \rangle + ((\mathbf{v}_\epsilon \cdot \nabla) \mathbf{B}_\epsilon + 2\rho_\epsilon(\mathbf{B}_\epsilon) \mathbf{B} \mathbf{W}(\mathbf{v}) - 2a\rho_\epsilon(\mathbf{B}_\epsilon) \mathbf{B} \mathbf{D}(\mathbf{v}), \mathbf{A})] dt \\ &+ \int_0^T (\rho_\epsilon(\mathbf{B}_\epsilon) \mathbf{R}(\mathbf{B}_\epsilon), \mathbf{A}) dt = - \int_0^T (\nabla \mathbf{B}, \nabla \mathbf{A}) dt. \quad (\text{A.8})\end{aligned}$$

Moreover, for almost every $(t, x) \in [0, T] \times \mathbb{T}^2$, the minimal eigenvalue of \mathbf{B}_ϵ satisfies

$$\Lambda(\mathbf{B}_\epsilon) \geq \epsilon. \quad (\text{A.9})$$

Proof. For $\epsilon > 0$, $k \in \mathbb{N}$, let $(\mathbf{v}_\epsilon^k, \mathbf{B}_\epsilon^k)$ be the solution to the Galerkin system (A.1), (A.2). By Lemma A.1 and the Banach-Alaoglu theorem there exist subsequences (which we will not relabel) and $(\mathbf{v}_\epsilon, \mathbf{B}_\epsilon)$ such that:

$$\begin{aligned}\mathbf{v}_\epsilon^k &\overset{*}{\rightharpoonup} \mathbf{v}_\epsilon \quad \text{weakly in } L^\infty(0, T; L^2), \\ \mathbf{v}_\epsilon^k &\rightharpoonup \mathbf{v}_\epsilon \quad \text{weakly in } L^2(0, T; H_{0,\text{div}}^1), \\ \partial_t \mathbf{v}_\epsilon^k &\overset{*}{\rightharpoonup} \partial_t \mathbf{v}_\epsilon \quad \text{weakly in } L^\infty(0, T; H_{0,\text{div}}^{-1}), \\ \mathbf{B}_\epsilon^k &\overset{*}{\rightharpoonup} \mathbf{B}_\epsilon \quad \text{weakly in } L^\infty(0, T; L^2), \\ \mathbf{B}_\epsilon^k &\rightharpoonup \mathbf{B}_\epsilon \quad \text{weakly in } L^2(0, T; H^1), \\ \partial_t \mathbf{B}_\epsilon^k &\overset{*}{\rightharpoonup} \partial_t \mathbf{B}_\epsilon \quad \text{weakly in } L^\infty(0, T; H^{-1}).\end{aligned}$$

By the Aubin–Lions lemma and extracting further subsequences if necessary, we have also

$$\begin{aligned}\mathbf{v}_\epsilon^k &\rightarrow \mathbf{v}_\epsilon \quad \text{strongly in } L^2(0, T; H_{0,\text{div}}^{1/2}) \cap C([0, T]; H_{0,\text{div}}^{-1}), \\ \mathbf{B}_\epsilon^k &\rightarrow \mathbf{B}_\epsilon \quad \text{strongly in } L^2(0, T; H^{1/2}) \cap C([0, T]; H^{-1}), \\ \rho_\epsilon(\mathbf{B}_\epsilon^k) &\rightarrow \rho_\epsilon(\mathbf{B}_\epsilon) \quad \text{a.e. on } [0, T] \times \mathbb{T}^2.\end{aligned}$$

Using (A.1), (A.2), the previous convergence properties and standard arguments we can conclude that $(\mathbf{v}_\epsilon, \mathbf{B}_\epsilon)$ satisfies (A.7), (A.8) and $\mathbf{v}_\epsilon(0, x) = \mathbf{v}_0(x)$, $\mathbf{B}_\epsilon(0, x) = \mathbf{B}_\epsilon(x)$.

The proof of (A.9) is the same as the 3d case in [1], but we will briefly sketch it for completeness. Let $\mathbf{z} \in \mathbb{Q}^2$. In (A.8) we choose

$$\mathbf{A}(t, x) = \chi_{[0, \tau]}(t) (\mathbf{B}_\epsilon(t, x) \mathbf{z} \cdot \mathbf{z} - \epsilon |\mathbf{z}|^2)_- (\mathbf{z} \otimes \mathbf{z}) \in L^2(0, T; H^1).$$

This choice of \mathbf{A} , and the facts that $\operatorname{div} \mathbf{v}_\epsilon = 0$ and $\rho_\epsilon(\mathbf{B}_\epsilon) \mathbf{A} = \mathbf{0}$ imply that the only nonzero terms appearing in (A.8) are

$$\begin{aligned} \int_0^T \langle \partial_t \mathbf{B}_\epsilon, \mathbf{A} \rangle dt &= \int_0^\tau \frac{d}{dt} \frac{1}{2} \|(\mathbf{B}_\epsilon \mathbf{z} \cdot \mathbf{z} - \epsilon |\mathbf{z}|^2)_-\|^2 dt \\ &= \frac{1}{2} \|(\mathbf{B}_\epsilon(\tau) \mathbf{z} \cdot \mathbf{z} - \epsilon |\mathbf{z}|^2)_-\|^2 - \frac{1}{2} \|(\mathbf{B}_{0, \epsilon} \mathbf{z} \cdot \mathbf{z} - \epsilon |\mathbf{z}|^2)_-\|^2 \\ &= \frac{1}{2} \|(\mathbf{B}_\epsilon(\tau) \mathbf{z} \cdot \mathbf{z} - \epsilon |\mathbf{z}|^2)_-\|^2, \end{aligned}$$

and

$$\int_0^T (\nabla \mathbf{B}_\epsilon, \nabla \mathbf{A}) dt = \int_0^\tau \|\nabla (\mathbf{B}_\epsilon \mathbf{z} \cdot \mathbf{z} - \epsilon |\mathbf{z}|^2)_-\|^2 dt.$$

By (A.8)

$$\|(\mathbf{B}_\epsilon(\tau) \mathbf{z} \cdot \mathbf{z} - \epsilon |\mathbf{z}|^2)_-\|^2 = 0, \quad \tau \in [0, T].$$

Thus, $\mathbf{B}(t, x) \mathbf{z} \cdot \mathbf{z} \geq \epsilon |\mathbf{z}|^2$ almost everywhere on $[0, T] \times \mathbb{T}^2$. We conclude that for a.e. $(t, x) \in [0, T] \times \mathbb{T}^2$, for all $\mathbf{z} \in \mathbb{R}^2$, we have $\mathbf{B}(t, x) \mathbf{z} \cdot \mathbf{z} \geq \epsilon |\mathbf{z}|^2$, proving (A.9). \square

A.2. An energy identity. We now derive an energy identity for the ϵ -modified equations reminiscent of the energy balance identity (1.6) for the equations of motion.

Lemma A.3. *Let $\epsilon > 0$ and $(\mathbf{v}_\epsilon, \mathbf{B}_\epsilon)$ be as in Lemma A.2. Then for all $\tau \in [0, T]$,*

$$\begin{aligned} &\frac{1}{2} \|\mathbf{v}_\epsilon(\tau)\|^2 + \int_{\mathbb{T}^2} \psi(\mathbf{B}_\epsilon(\tau)) dx + \int_0^\tau \|\nabla \mathbf{v}_\epsilon\|^2 dt \\ &+ \int_0^\tau [(1 - \beta) \|\mathbf{B}_\epsilon^{-1/2} \nabla \mathbf{B}_\epsilon \mathbf{B}_\epsilon^{-1/2}\|^2 + \beta \|\nabla \mathbf{B}_\epsilon\|^2] dt \\ &+ \int_0^\tau (1 - \beta) \delta_1 \|\rho_\epsilon(\mathbf{B}_\epsilon)^{1/2} (\mathbf{B}_\epsilon^{1/2} - \mathbf{B}_\epsilon^{-1/2})\|^2 dt \\ &+ \int_0^\tau \beta \delta_2 \|\rho_\epsilon(\mathbf{B}_\epsilon)^{1/2} (\mathbf{B}_\epsilon^{3/2} - \mathbf{B}_\epsilon^{1/2})\|^2 dt \\ &+ \int_0^\tau (\beta \delta_1 + (1 - \beta) \delta_2) \|\rho_\epsilon(\mathbf{B}_\epsilon)^{1/2} (\mathbf{B}_\epsilon - \mathbf{I})\|^2 dt \\ &= \frac{1}{2} \|\mathbf{v}_0\|^2 + \int_{\mathbb{T}^2} \psi(\mathbf{B}_{0, \epsilon}) dx + \int_0^\tau \langle \mathbf{f}, \mathbf{v} \rangle dt. \end{aligned} \tag{A.10}$$

Proof. By (A.9) it follows that for a.e. $(t, x) \in [0, T] \times \mathbb{T}^2$, \mathbf{B}_ϵ^{-1} exists and

$$|\mathbf{B}_\epsilon^{-1}(t, x)| \leq \frac{1}{\epsilon}.$$

Then $\nabla \mathbf{B}_\epsilon^{-1} = -\mathbf{B}_\epsilon^{-1} \nabla \mathbf{B}_\epsilon \mathbf{B}_\epsilon^{-1}$ implies

$$\int_0^T \|\nabla \mathbf{B}_\epsilon^{-1}\|^2 dt \leq \frac{1}{\epsilon^4} \int_0^T \|\nabla \mathbf{B}_\epsilon\|^2 dt. \quad (\text{A.11})$$

Let

$$\mathbf{J}_\epsilon = (1 - \beta)(\mathbf{I} - \mathbf{B}_\epsilon^{-1}) + \beta(\mathbf{B}_\epsilon - \mathbf{I}).$$

By (A.11), $\mathbf{J}_\epsilon \in L^2(0, T; H^1)$. Since $\frac{\partial \psi(\mathbf{B}_\epsilon)}{\partial \mathbf{B}_\epsilon} = \mathbf{J}_\epsilon$ and $\operatorname{div} \mathbf{v}_\epsilon = 0$, one readily verifies the identities

$$\begin{aligned} \langle \partial_t \mathbf{B}_\epsilon, \mathbf{J}_\epsilon \rangle &= \frac{d}{dt} \int_{\mathbb{T}^2} \psi(\mathbf{B}_\epsilon) dx, \\ ((\mathbf{v}_\epsilon \cdot \nabla) \mathbf{B}_\epsilon, \mathbf{J}_\epsilon) &= \int_{\mathbb{T}^2} \mathbf{v}_\epsilon \cdot \nabla \psi(\mathbf{B}_\epsilon) dx = 0, \\ (\nabla \mathbf{B}_\epsilon, \nabla \mathbf{J}_\epsilon) &= (1 - \beta) \|\mathbf{B}_\epsilon^{-1/2} \nabla \mathbf{B}_\epsilon \mathbf{B}_\epsilon^{-1/2}\|^2 + \beta \|\nabla \mathbf{B}_\epsilon\|^2, \\ (\rho_\epsilon(\mathbf{B}_\epsilon) \mathbf{R}(\mathbf{B}_\epsilon), \mathbf{J}_\epsilon) &= (1 - \beta) \delta_1 \|\rho_\epsilon(\mathbf{B}_\epsilon)^{1/2} (\mathbf{B}_\epsilon^{1/2} - \mathbf{B}_\epsilon^{-1/2})\|^2 \\ &\quad + \beta \delta_2 \|\rho_\epsilon(\mathbf{B}_\epsilon)^{1/2} (\mathbf{B}_\epsilon^{3/2} - \mathbf{B}_\epsilon^{1/2})\|^2 \\ &\quad + (\beta \delta_1 + (1 - \beta) \delta_2) \|\rho_\epsilon(\mathbf{B}_\epsilon)^{1/2} (\mathbf{B}_\epsilon - \mathbf{I})\|^2. \end{aligned}$$

Since

$$\mathbf{B}_\epsilon \mathbf{J}_\epsilon = \mathbf{S}(\mathbf{B}_\epsilon) = \mathbf{J}_\epsilon \mathbf{B}_\epsilon$$

we also have

$$\begin{aligned} 2(\rho_\epsilon(\mathbf{B}_\epsilon) \mathbf{B}_\epsilon \mathbf{W}(\mathbf{v}_\epsilon), \mathbf{J}_\epsilon) &= 0, \\ 2a(\rho_\epsilon(\mathbf{B}_\epsilon) \mathbf{B}_\epsilon \mathbf{D}(\mathbf{v}_\epsilon), \mathbf{J}_\epsilon) &= 2a(\rho_\epsilon(\mathbf{B}_\epsilon) \mathbf{D}(\mathbf{v}_\epsilon), \mathbf{S}(\mathbf{B}_\epsilon)). \end{aligned}$$

Using the previous identities and (A.8) with $\mathbf{A} = \chi_{[0, \tau]} \mathbf{J}_\epsilon$ we obtain the identity

$$\begin{aligned} &\int_{\mathbb{T}^2} \psi(\mathbf{B}_\epsilon(\tau)) dx + \int_0^\tau [(1 - \beta) \|\mathbf{B}_\epsilon^{-1/2} \nabla \mathbf{B}_\epsilon \mathbf{B}_\epsilon^{-1/2}\|^2 + \beta \|\nabla \mathbf{B}_\epsilon\|^2] dt \\ &\quad + \int_0^\tau (1 - \beta) \delta_1 \|\rho_\epsilon(\mathbf{B}_\epsilon)^{1/2} (\mathbf{B}_\epsilon^{1/2} - \mathbf{B}_\epsilon^{-1/2})\|^2 dt \\ &\quad + \int_0^\tau \beta \delta_2 \|\rho_\epsilon(\mathbf{B}_\epsilon)^{1/2} (\mathbf{B}_\epsilon^{3/2} - \mathbf{B}_\epsilon^{1/2})\|^2 dt \\ &\quad + \int_0^\tau (\beta \delta_1 + (1 - \beta) \delta_2) \|\rho_\epsilon(\mathbf{B}_\epsilon)^{1/2} (\mathbf{B}_\epsilon - \mathbf{I})\|^2 dt \\ &= \int_{\mathbb{T}^2} \psi(\mathbf{B}_{0, \epsilon}) dx + \int_0^\tau 2a(\rho_\epsilon(\mathbf{B}_\epsilon) \mathbf{S}(\mathbf{B}_\epsilon), \mathbf{D}(\mathbf{v}_\epsilon)) dt. \quad (\text{A.12}) \end{aligned}$$

Using (A.7) with $\varphi = \chi_{[0,\tau]} \mathbf{v}_\epsilon$ we obtain

$$\begin{aligned} & \frac{1}{2} \|\mathbf{v}_\epsilon\|^2 + \int_0^\tau \|\nabla \mathbf{v}_\epsilon\|^2 dt \\ &= \frac{1}{2} \|\mathbf{v}_0\|^2 - \int_0^\tau 2a(\rho_\epsilon(\mathbf{B}_\epsilon) \mathcal{S}(\mathbf{B}_\epsilon), \mathbf{D}(\mathbf{v}_\epsilon)). \end{aligned} \quad (\text{A.13})$$

Adding (A.12) to (A.13) yields (A.10). \square

A.3. Conclusion of the proof of Theorem 2.2. We now conclude the proof of Theorem 2.2 by establishing the existence of a weak solution. For $\epsilon > 0$, let $(\mathbf{v}_\epsilon, \mathbf{B}_\epsilon)$ be as in Lemma A.2. By the definition of $\mathbf{B}_{0,\epsilon}$, the fact $\psi(\mathbf{I}) = 0$ and Lemma A.3 we conclude for all $\tau \in [0, T]$

$$\begin{aligned} & \frac{1}{2} \sup_{\tau \in [0, T]} \|\mathbf{v}_\epsilon(\tau)\|^2 + \sup_{\tau \in [0, T]} \int_{\mathbb{T}^2} \psi(\mathbf{B}_\epsilon(\tau)) dx + \frac{1}{2} \int_0^T (\|\nabla \mathbf{v}_\epsilon\|^2 + \|\nabla \mathbf{B}_\epsilon\|^2) dt \\ & \leq \frac{1}{2} \|\mathbf{v}_0\|^2 + \int_{\mathbb{T}^2} \psi(\mathbf{B}_0) dx + C \int_0^T \|\mathbf{f}\|_{H_{0,\text{div}}^{-1}}^2 dt \end{aligned} \quad (\text{A.14})$$

where C is an absolute constant. Note that since $\int_{\mathbb{T}^2} |\log \det \mathbf{B}_0| dx < \infty$, the right-hand side of (A.14) is finite and independent of ϵ . Using (A.7), (A.8), (A.14) and arguing as in the proof of (A.4) we obtain

$$\|\partial_t \mathbf{v}_\epsilon\|_{L^2 H_{0,\text{div}}^{-1}} \leq B(\mathbf{v}_0, \mathbf{B}_0, \mathbf{f}), \quad (\text{A.15})$$

$$\|\partial_t \mathbf{B}_\epsilon\|_{[L^2 H^1 \cap L^4 L^4]^*} \leq B(\mathbf{v}_0, \mathbf{B}_0, \mathbf{f}), \quad (\text{A.16})$$

$$\forall \delta > 0, \quad \|\partial_t \mathbf{B}_\epsilon\|_{L^2 H^{-1-\delta}} \leq B(\mathbf{v}_0, \mathbf{B}_0, \mathbf{f}, \delta). \quad (\text{A.17})$$

For example, to establish (A.16) we estimate the pairing

$$\begin{aligned} & \int_0^T (\mathbf{D}(\mathbf{v}_\epsilon) \mathbf{B}_\epsilon + \mathbf{B}_\epsilon \mathbf{D}(\mathbf{v}_\epsilon), \mathbf{A}) dt \\ & \leq C \|\mathbf{v}_\epsilon\|_{L^2 H^1} \|\mathbf{B}_\epsilon\|_{L^4 L^4} \|\mathbf{A}\|_{L^4 L^4} \\ & \leq C \|\mathbf{v}_\epsilon\|_{L^2 H^1} \|\mathbf{B}_\epsilon\|_{L^\infty L^2}^{1/2} \|\mathbf{B}_\epsilon\|_{L^2 H^1}^{1/2} \|\mathbf{A}\|_{L^4 L^4}, \end{aligned}$$

and to establish (A.17) we use Sobolev embedding $H^{1+\delta} \hookrightarrow L^\infty$ to estimate the pairing

$$\begin{aligned} & \int_0^T (\mathbf{D}(\mathbf{v}_\epsilon) \mathbf{B}_\epsilon + \mathbf{B}_\epsilon \mathbf{D}(\mathbf{v}_\epsilon), \mathbf{A}) dt \\ & \leq C(\delta) \|\mathbf{v}_\epsilon\|_{L^2 H^1} \|\mathbf{B}_\epsilon\|_{L^\infty L^2} \|\mathbf{A}\|_{L^2 H^{1+\delta}}. \end{aligned}$$

By (A.14), (A.15), (A.16), (A.17) and the Banach-Alaoglu theorem there exist $\epsilon_j \rightarrow 0$ and (\mathbf{v}, \mathbf{B}) such that:

$$\begin{aligned} \mathbf{v}_{\epsilon_j} &\overset{*}{\rightharpoonup} \mathbf{v} \quad \text{weakly in } L^\infty(0, T; L^2), \\ \mathbf{v}_{\epsilon_j} &\rightharpoonup \mathbf{v} \quad \text{weakly in } L^2(0, T; H_{0,\text{div}}^1), \\ \partial_t \mathbf{v}_{\epsilon_j} &\overset{*}{\rightharpoonup} \partial_t \mathbf{v} \quad \text{weakly in } L^\infty(0, T; H_{0,\text{div}}^{-1}), \\ \mathbf{B}_{\epsilon_j} &\overset{*}{\rightharpoonup} \mathbf{B} \quad \text{weakly in } L^\infty(0, T; L^2), \\ \mathbf{B}_{\epsilon_j} &\rightharpoonup \mathbf{B} \quad \text{weakly in } L^2(0, T; H^1), \\ \partial_t \mathbf{B}_{\epsilon_j} &\overset{*}{\rightharpoonup} \partial_t \mathbf{B} \quad \text{weakly in } [L^2(0, T; H^1) \cap L^4(0, T; L^4)]^*, \\ \forall \delta > 0, \partial_t \mathbf{B}_{\epsilon_j} &\overset{*}{\rightharpoonup} \partial_t \mathbf{B} \quad \text{weakly in } L^\infty(0, T; H^{-1-\delta}). \end{aligned}$$

We note that (A.14) and these weak convergence properties imply that

$$\begin{aligned} \|\mathbf{v}\|_{L^\infty L^2} + \|\mathbf{B}\|_{L^\infty L^2} + \|\nabla \mathbf{v}\|_{L^2 L^2} + \|\nabla \mathbf{B}\|_{L^2 L^2} \\ \leq C_0(\|\mathbf{v}_0\|, \|\psi(\mathbf{B}_0)\|_{L^1}, \|\mathbf{f}\|_{L^2 H_{0,\text{div}}^{-1}}). \end{aligned} \quad (\text{A.18})$$

By the Aubin–Lions lemma and extracting further subsequences if necessary, we have also

$$\begin{aligned} \mathbf{v}_{\epsilon_j} &\rightarrow \mathbf{v} \quad \text{strongly in } L^2(0, T; H_{0,\text{div}}^{1/2}) \cap C([0, T]; H_{0,\text{div}}^{-1}), \\ \mathbf{B}_{\epsilon_j} &\rightarrow \mathbf{B} \quad \text{strongly in } L^2(0, T; H^{1/2}) \cap C([0, T]; H^{-1-\delta}), \quad \forall \delta > 0, \\ \mathbf{B}_{\epsilon_j} &\rightarrow \mathbf{B} \quad \text{a.e. on } [0, T] \times \mathbb{T}^2. \end{aligned}$$

By (A.9) and the continuity of $\Lambda(\cdot)$, $\Lambda(\mathbf{B}) \geq 0$ a.e. on $[0, T] \times \mathbb{T}^2$. In particular, $\det \mathbf{B}_0 \geq 0$ a.e., and by continuity of $\psi(\cdot)$,

$$\psi(\mathbf{B}_{\epsilon_j}) \rightarrow \psi(\mathbf{B}) \in [0, \infty]$$

a.e. on $[0, T] \times \mathbb{T}^2$. Fatou's lemma and (A.14) then imply that

$$\|\psi(\mathbf{B})\|_{L^\infty L^1} < \infty. \quad (\text{A.19})$$

Since $\Lambda(\mathbf{B}) = 0$ on a subset of $[0, T] \times \mathbb{T}^2$ with positive measure would imply that $\psi(\mathbf{B}) = \infty$ on a subset with positive measure, we conclude from (A.19) that

$$\Lambda(\mathbf{B}) > 0 \quad \text{a.e. on } [0, T] \times \mathbb{T}^2.$$

Thus, by the continuity of $\Lambda(\cdot)$,

$$\rho_{\epsilon_j}(\mathbf{B}_{\epsilon_j}) \rightarrow 1 \quad \text{a.e. on } [0, T] \times \mathbb{T}^2.$$

Using (A.7), (A.8), the previous convergence results and standard arguments we conclude that (\mathbf{v}, \mathbf{B}) verify (2.1) and (2.2). Finally, the additional regularity $\mathbf{v} \in C([0, T]; L_{0,\text{div}}^2)$ and $\mathbf{B} \in C([0, T]; L^2)$ follow immediately from the facts that

$$\mathbf{v} \in L^2(0, T; H_{0,\text{div}}^1), \partial_t \mathbf{v} \in L^2(0, T; H_{0,\text{div}}^{-1})$$

and

$$\begin{aligned} \mathbf{B} &\in L^2(0, T; H^1) \cap L^\infty(0, T; L^2), \\ \partial_t \mathbf{B} &\in [L^2(0, T; H^1) \cap L^4(0, T; L^4)]^*. \end{aligned}$$

□

Acknowledgement. Miroslav Bulíček and Josef Málek acknowledge the support of the project No. 20-11027X financed by Czech science foundation (GACR). Miroslav Bulíček and Josef Málek are members of the Nečas center for mathematical modelling.

REFERENCES

- [1] M. Bathory, M. Bulíček, and J. Málek. Large data existence theory for three-dimensional unsteady flows of rate-type viscoelastic fluids with stress diffusion. *Adv. Nonlinear Anal.*, 10(1):501–521, 2021.
- [2] P. Constantin and C. Foias. *Navier-Stokes equations*. University of Chicago Press, 1988.
- [3] P. Constantin and M. Kliegl. Note on global regularity for two-dimensional Oldroyd-B fluids with diffusive stress. *Arch. Ration. Mech. Anal.*, 206(3):725–740, 2012.
- [4] T. Divoux, M. A. Fardin, S. Manneville, and S. Lerouge. Shear banding of complex fluids. *Annu. Rev. Fluid Mech.*, 48(1):81–103, 2016.
- [5] A. W. El-Kareh and L. G. Leal. Existence of solutions for all Deborah numbers for a non-Newtonian model modified to include diffusion. *J. Non-Newton. Fluid Mech.*, 33(3):257–287, 1989.
- [6] L. C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.
- [7] M.-A. Fardin, O. Radulescu, A. Morozov, O. Cardoso, J. Browaeys, and S. Lerouge. Stress diffusion in shear banding wormlike micelles. *J. Rheol.*, 59(6):1335–1362, 2015.
- [8] H. Giesekus. A simple constitutive equation for polymer fluids based on the concept of deformation-dependent tensorial mobility. *Journal of Non-Newtonian Fluid Mechanics*, 11(1):69–109, 1982.
- [9] E. Hopf. Über die anfangswertaufgabe für die hydrodynamischen grundgleichungen. erhard schmidt zu seinem 75. geburtstag gewidmet. *Mathematische Nachrichten*, 4(1-6):213–231, 1950.
- [10] M. W. Johnson and D. Segalman. A model for viscoelastic fluid behavior which allows non-affine deformation. *Journal of Non-Newtonian Fluid Mechanics*, 2(3):255–270, 1977.
- [11] R. G Larson. *Constitutive Equations for Polymer Melts and Solutions*. Butterworths Series in Chemical Engineering. Butterworths, London, 1988.
- [12] J. Leray. Sur le mouvement d’un liquide visqueux emplissant l’espace. *Acta Math.*, 63:193–248, 1934.
- [13] J. Málek and V. Průša. Derivation of equations for continuum mechanics and thermodynamics of fluids. In *Handbook of mathematical analysis in mechanics of viscous fluids*, pages 3–72. Springer, Cham, 2018.
- [14] J. Málek, V. Průša, T. Skřivan, and E. Süli. Thermodynamics of viscoelastic rate-type fluids with stress diffusion. *Physics of Fluids*, 30(2):023101, 2018.
- [15] J. Málek, K. R. Rajagopal, and K. Tůma. On a variant of the Maxwell and Oldroyd-B models within the context of a thermodynamic basis. *Int. J. Non-Linear Mech.*, 76:42–47, 2015.
- [16] J. G. Oldroyd. On the formulation of rheological equations of state. *Proc. Roy. Soc. London Ser. A*, 200:523–541, 1950.

- [17] P. D. Olmsted. Perspectives on shear banding in complex fluids. *Rheol. Acta*, 47(3):283–300, 2008.
- [18] P. D. Olmsted, O. Radulescu, and C.-Y. D. Lu. Johnson-Segalman model with a diffusion term in cylindrical Couette flow. *Journal of Rheology*, 44(2):257–275, 2000.
- [19] K. R. Rajagopal and A. R. Srinivasa. A thermodynamic frame work for rate type fluid models. *Journal of Non-Newtonian Fluid Mechanics*, 88(3):207–227, 2000.
- [20] K. R. Rajagopal and A. R. Srinivasa. On thermomechanical restrictions of continua. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 460(2042):631–651, 2004.
- [21] R. Temam. *Navier-Stokes equations and nonlinear functional analysis*, volume 66 of *CBMS-NSF Regional Conference Series in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, second edition, 1995.
- [22] R. Temam. *Navier-Stokes equations*. AMS Chelsea Publishing, Providence, RI, 2001. Theory and numerical analysis, Reprint of the 1984 edition.
- [23] C. Truesdell and W. Noll. *The non-linear field theories of mechanics*. Springer-Verlag, Berlin, third edition, 2004. Edited and with a preface by Stuart S. Antman.

MIROSLAV BULÍČEK

Charles University, Faculty of Mathematics and Physics, Mathematical Institute,
Sokolovská 83, 18675 Prague 8, Czech Republic
mbul8060@karlin.mff.cuni.cz

JOSEF MÁLEK

Charles University, Faculty of Mathematics and Physics, Mathematical Institute,
Sokolovská 83, 18675 Prague 8, Czech Republic
mbul8060@karlin.mff.cuni.cz

CASEY RODRIGUEZ

Department of Mathematics, University of North Carolina,
Chapel Hill, NC, USA
crodrig@email.unc.edu