Towards reconciliation of completely positive open system dynamics with the equilibration postulate

Marcin Lobejko, Marek Winczewski, Gerardo Suárez, Robert Alicki, Michał Horodecki

International Centre for Theory of Quantum Technologies, University of Gdańsk, 80-308 Gdańsk, Poland (Dated: April 5, 2022)

Almost every quantum system interacts with a large environment, so the exact quantum mechanical description of its evolution is impossible. One has to resort to approximate description, usually in the form of a master equation. There are at least two basic requirements for such a description: first, it should preserve the positivity of probabilities; second, it should correctly describe the equilibration process for systems coupled to a single thermal bath. Existing two widespread descriptions of evolution fail to satisfy at least one of those conditions. The so-called Davies master equation, while preserving the positivity of probabilities, fails to describe thermalization properly. On the other hand, the Bloch-Redfield master equation violates the first condition, but it correctly describes equilibration, at least for off-diagonal elements for several important scenarios. However, is it possible to have a description of open system dynamics that would share both features? In this paper, we partially resolve this problem in the weak-coupling limit: (i) We provide a general form of the proper thermal equilibrium state (the so-called mean-force state) for an arbitrary open system. (ii) We provide the solution for the steady-state coherences for a whole class of master equations, and in particular, we show that the solution coincides with the mean-force Hamiltonian for the Bloch-Redfield equation. (iii) We consider the cumulant equation, which is explicitly completely positive, and we show that its steady-state coherences are the same as one of the Bloch-Redfield dynamics (and the mean-force state accordingly). (iv) We solve the correction to the diagonal part of the stationary state for a two-level system both for the Bloch-Redfield and cumulant equation, showing that the solution of the cumulant is very close to the mean-force state, whereas the Bloch-Redfield differs significantly.

I. INTRODUCTION

A basic property of the open system interacting with a single thermal reservoir is that it tends to a thermal equilibrium state. Another fundamental property of any system is that evolution should preserve the positivity of the measurement outcomes probabilities. Since it is, except for a few particular cases, impossible to describe open systems dynamics exactly, one applies approximate descriptions in the form of master equations [1, 2]. Clearly, such approximate descriptions should satisfy the above two postulates.

A widespread description of the open systems weakly coupled with the bath stems from Born approximation, followed by Markovian approximation. This leads to Bloch-Redfield equation [3, 4]. If one subsequently applies yet another approximation (called the secular approximation) one obtains the most popular master equation in Gorini-Kossakowski-Linblad-Sudarshan form [5, 6], called Davies equation [7]. The first one is more exact but does not preserve the positivity of the state. The second one, while preserving positivity and working perfectly in typical quantum optical regimes, cannot describe subtleties related to nearly degenerated levels. The Born approximation implies that both equations describe the dynamics up to the second order in the coupling strength λ .

One can now ask to what extent the equilibration postulate is satisfied. Since the equations are up to second order in λ , the equilibrium state should not be the Gibbs state of the free Hamiltonian anymore, but rather it should be a partial trace of a total (system plus bath) thermal equilibrium state, expanded up to second order in λ . Imagining that it is a Gibbs state with respect to some Hamiltonian, one calls the latter "mean-force Hamiltonian", and the state is called mean-force Gibbs state. This issue has been a matter of intense research, see [8] for a review.

It is known, in particular, that for some standard coupling to a bosonic bath, the Bloch-Redfield equation satisfies in part this basic requirement. Namely, the state it equilibrates to has the same off-diagonal elements as the mean-force Gibbs state [9-13]. However, the above is not known for arbitrary coupling, i.e., on the level of generality assumed when deriving these master equations. It should be emphasized that the general form of the second-order correction for either the mean-force Hamiltonian or the steady-state Hamiltonian for the Bloch-Redfield equation has not been calculated so far. Regarding diagonal elements, there is an agreement between the steady state and the mean-force Gibbs state only up to zeroth order, and we cannot expect more with second order master equations as argued in [9, 13]. On the other hand, the Davies equation does not satisfy the equilibration postulate even for off-diagonal elements, since its steady-state is a Gibbs state of the bare Hamiltonian, or employing the renormalization proposed in [14], its second-order correction is equal to the so-called Lamb-Stark shift (known also as *dynamical correction*), in general, different than the mean-force Hamiltonian.

We are thus in quite an unfortunate situation. The Bloch-Redfield equation, that predicts the correct equilibration (at least for off-diagonal elements) is not completely positive. On the other hand, the completely positive master equation - the Davies one, fails to describe thermalization much more severely (cf. [13]). Thus, we are quite far away from achieving the two basic postulates:

- (i) open systems thermalize,
- (ii) probabilities are non-negative,

simultaneously in the description of the evolution of open systems in a weak coupling scenario.

There is yet another candidate for dynamical equation that might join the advantages of both the equations above. This is the "cumulant equation" (also called the refined weak coupling limit) derived in [14] and then independently discovered and further developed in [15, 16] (see [17] for recent development). This is a completely positive evolution that is not based on the secular approximation nor Markovian assumption in any form. Since it is just the secular approximation that makes Davies equation problematic in many respects, one would think that this new equation may indeed be a solution for the above quite fundamental problem. However, the cumulant equation is believed to reduce to Davies one for the long time limit [14, 15], which would imply that the steady-state must be that of Davies equation, hence the postulate would not be satisfied.

In this paper, we make a progress towards reconciliation of the two postulates - positivity and thermalization. To this end we provide several separate and general results, that are of independent interest. Then the main message, which follows from putting them together, is that the cumulant equation is at the moment quite close to satisfy both postulates. We also show where and to what extent it fails to do so.

Let us now present the mentioned results. To this end, let us set up a vocabulary. Namely, in the discussion above, three types of corrections to bare Hamiltonian appear: *mean-force* correction, *steady-state* correction and *dynamical* correction. As explained, the mean-force correction is the following: we imagine that the thermal equilibrium state is the Gibbs state according to perturbed bare Hamiltonian, and the perturbation is the mean-force correction. Regarding steady-state correction, we similarly imagine that the steady-state is the Gibbs state according to a perturbed bare Hamiltonian and that (in general different) perturbation is the steady-state Hamiltonian correction. Finally, the dynamical correction is the well-known Hamiltonian contribution from the environment. The form of the latter is known for the Davies equation and Bloch-Redfield equation.

So far, we have put more emphasis on the problem of relations between the mean-force and the steady-state corrections. Of course, ideally, we would like to have all three corrections equal to each other. However, existing results (see, e.g., [12]) show that, in general, this is not possible, yet it is hard to find a proper physical explanation of such a discrepancy. In our studies, we show how those three corrections are related to each other in a general open systems framework. Let us also stress that all our results concern second-order corrections - we will thus not explicitly repeat this in the following.

Here are the most general results that do not involve any particular master equation: We derive

- general form of mean-force correction for an arbitrary open system, which so far was only known for particular couplings. This form has been also independently derived by G. Timofeev and A. Trushechkin [18];
- *coherences* of the steady-state correction for quite a general form of master equation in terms of its Kossakowski matrix.

In addition, our formulas for the above corrections explicitly show their relations with the dynamical correction. We next concentrate on the Bloch-Redfield equation and the cumulant equation: We show that

- for a general coupling, the Bloch-Redfield equation predicts the *coherences* of a *steady-state correction* that coincide with those of the mean-force (this was known before only for some particular models, see, e.g., [10, 12, 19, 20]);
- the generator of cumulant master equation, to lowest order, agrees with Bloch-Redfield equation.

The above results already achieve the goal partially: the last item implies that the cumulant equation has the same steady-state correction coherences as the Bloch-Redfield one. Since the cumulant equation is completely positive, hence we obtain a reconciliation of complete positivity with thermalization for coherences. This also shows that the cumulant equation does not converge to the Davies equation for long times.

Subsequently, we deal with a much more complicated issue of diagonal elements, we provide:

- general methodology for computing diagonal elements of the steady-state correction.
- formula for those elements in the case of the two-level system, obtaining elegant relation between diagonal elements of the mean-force and those of the steady-state correction.

Next, we specialize to Bloch-Redfield and cumulant equation:

• We show that diagonal elements of the mean-force and the steady-state correction are quite close to one another for cumulant equation, while for Bloch-Redfield, they significantly differ.

Combined with the above results on coherences, these last results show that the cumulant equation is quite close to reconciling the postulates of positivity and thermalization.

The paper is organized as follows. First, in Section II we provide the notation and concepts that will be used in the paper, i.e., we define and derive three different corrections to the bare Hamiltonian, namely the dynamical, the steady-state, and the mean force one. Next, in Section III, we describe the main results regarding the off-diagonal elements of the Hamiltonian. Later in section IV, we introduce the cumulant equation, which is a second-order dynamical map whose differential form can be used to obtain diagonal corrections up to the second-order, which is done in Section V. These results are then applied to a two-level system in Section VI.

II. PRELIMINARIA: DYNAMICAL, MEAN-FORCE AND STEADY STATE CORRECTION

We consider a general Hamiltonian of the interacting system S with the thermal reservoir R of the form:

$$H = H_0 + H_R + \lambda H_I, \tag{1}$$

$$H_I = \sum_{\alpha} A_{\alpha} \otimes R_{\alpha},\tag{2}$$

where H_0 is a bare Hamiltonian of the system, H_R is free Hamiltonian of the bath, A_{α}, R_{α} are interaction operators (acting on the system and bath Hilbert spaces, respectively), and λ is a coupling constant. In the following, we define a Gibbs state of the thermal reservoir $\gamma_R = Z_R^{-1} e^{-\beta H_R}$ at inverse temperature β , where $Z_R = \text{Tr}[e^{-\beta H_R}]$. Additionally, we consider the time-dependent operators $A(t) = e^{i(H_0 + H_R)t} R_{\alpha} e^{-i(H_0 + H_R)t}$, and we use an abbreviation $\langle A \rangle_{\gamma_R} = \text{Tr}[A\gamma_R]$. We assume that that bath operators are centralized, i.e., $\langle R_{\alpha} \rangle_{\gamma_R} = 0$.

The main object of interest of this article are three different corrections to the bare Hamiltonian of the system H_0 in the weak coupling limit (i.e., $\lambda \ll 1$), defined according to the Hamiltonians: dynamical H_{dyn} , steady-state H_{st} and the mean-force H_{mf} . Due to centralization of the bath operators, the leading order of the perturbation calculus is λ^2 . In accordance, we introduce the second-order corrections:

$$H_k - H_0 = \lambda^2 H_k^{(2)} + \dots,$$
 (3)

which we represent in the basis of jump operators:

$$H_k^{(2)}(t) = \sum_{\omega,\omega'} \sum_{\alpha,\beta} \Upsilon_{\alpha\beta}^{(k)}(\omega,\omega',t) A_\alpha^{\dagger}(\omega) A_\beta(\omega'), \tag{4}$$

where k indicates to dynamical (dyn), steady-state (st) or mean-force (mf) correction, and the jump operators are given by:

$$A_{\alpha}(\omega) = \sum_{\epsilon'-\epsilon=\omega} \Pi(\epsilon) A_{\alpha} \Pi(\epsilon').$$
(5)

where $\Pi(\epsilon)$ is the projector on the subspace with energy ϵ , such that $H_0 = \sum_{\epsilon} \epsilon \Pi(\epsilon)$.

A. Dynamical correction

We start with the *dynamical correction* $H_{dyn}^{(2)}$, which is defined according to the generator of the master equation in the Schrödinger picture of the following general form:

$$\mathcal{L}_{t}[\rho] = i[\rho, H_{0} + \lambda^{2} H_{\text{dyn}}^{(2)}(t)] + \lambda^{2} \sum_{\omega, \omega'} \sum_{\alpha \beta} K_{\alpha \beta}(\omega, \omega', t) \left(A_{\beta}(\omega')\rho A_{\alpha}^{\dagger}(\omega) - \frac{1}{2} \{ A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega'), \rho \} \right)$$
(6)

$$= i[\rho, H_0] + \lambda^2 \sum_{\omega, \omega'} \sum_{\alpha\beta} \left[i \Upsilon^{(\text{dyn})}_{\alpha\beta}(\omega, \omega', t) [\rho, A^{\dagger}_{\alpha}(\omega) A_{\beta}(\omega')] + K_{\alpha\beta}(\omega, \omega', t) \left(A_{\beta}(\omega') \rho A^{\dagger}_{\alpha}(\omega) - \frac{1}{2} \{ A^{\dagger}_{\alpha}(\omega) A_{\beta}(\omega'), \rho \} \right) \right]$$
(7)

where K is the so-called Kossakowski matrix and for a while we do not determine it: For particular choice of K and $\Upsilon^{(dyn)}$ we will obtain a given master equation, such as the Bloch-Redfield or Davies-GKSL one. Notice that the master equation Eq. (6) leads to a completely positive dynamics if the matrix $K_{\alpha\beta}(\omega, \omega', t)$ is positive semi-definite. Later, we consider the long-time limit, when $t \to \infty$, for which we use the abbreviations:

$$\Upsilon_{\alpha\beta}^{(\mathrm{dyn})}(\omega,\omega') \equiv \lim_{t \to \infty} \Upsilon_{\alpha\beta}^{(\mathrm{dyn})}(\omega,\omega',t), \quad K_{\alpha\beta}(\omega,\omega') \equiv \lim_{t \to \infty} K_{\alpha\beta}(\omega,\omega',t).$$
(8)

1. Bloch-Redfield and Davies master equations

Let us consider generators of the two most known master equations, i.e., the Bloch-Redfield \mathcal{L}_t^R and the Davis generator \mathcal{L}_t^D . For the Bloch-Redfield master equation we have

$$K_{\alpha\beta}(\omega,\omega',t) = \Gamma_{\alpha\beta}(\omega',t) + \Gamma^*_{\beta\alpha}(\omega,t) \equiv \gamma_{\alpha\beta}(\omega,\omega',t), \qquad (9)$$

$$\Upsilon_{\alpha\beta}^{(\mathrm{dyn})}(\omega,\omega',t) = \frac{1}{2i} [\Gamma_{\alpha\beta}(\omega',t) - \Gamma_{\beta\alpha}^{*}(\omega,t)] \equiv \mathcal{S}_{\alpha\beta}(\omega,\omega',t), \tag{10}$$

where

$$\Gamma_{\alpha\beta}(\omega,t) = \int_0^t ds \ e^{i\omega s} \langle R_\alpha(s)R_\beta(0)\rangle_{\gamma_R}.$$
(11)

It is well-known that the Bloch-Redfield equation, in general, does not preserve the positivity of the state since $\gamma_{\alpha\beta}(\omega,\omega')$ is not a positive semi-definite matrix. Commonly, this issue is solved by applying the so-called secular approximation, which leads to the Davies master equation in the GKSL form. In accordance, applying the secular approximation by putting $\omega = \omega'$, and taking a limit $t \to \infty$, we obtain the coefficients for the Davies dynamics:

$$\gamma_{\alpha\beta}(\omega,\omega',t) \xrightarrow[t \to \infty]{\text{sec. approx.}} \gamma_{\alpha\beta}(\omega)\delta_{\omega,\omega'},$$
(12)

$$\mathcal{S}_{\alpha\beta}(\omega,\omega',t) \xrightarrow[t \to \infty]{\text{sec. approx.}} \mathcal{S}_{\alpha\beta}(\omega)\delta_{\omega,\omega'},$$
 (13)

where

$$\gamma_{\alpha\beta}(\omega) = \lim_{t \to \infty} \gamma_{\alpha\beta}(\omega, \omega, t) = \int_{-\infty}^{+\infty} ds \ e^{i\omega s} \langle R_{\alpha}(s) R_{\beta}(0) \rangle_{\gamma_R}, \tag{14}$$

is the Fourier transform of the auto-correlation function, and

$$\mathcal{S}_{\alpha\beta}(\omega) = \lim_{t \to \infty} \mathcal{S}_{\alpha\beta}(\omega, \omega, t) = \mathcal{P} \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \, \frac{\gamma_{\alpha\beta}(\Omega)}{\omega - \Omega},\tag{15}$$

where \mathcal{P} denotes the principal value integral. We also have the following relation:

$$\Gamma_{\alpha\beta}(\omega) = \lim_{t \to \infty} \Gamma_{\alpha\beta}(\omega, t) = \frac{1}{2} \gamma_{\alpha\beta}(\omega) + i \mathcal{S}_{\alpha\beta}(\omega).$$
(16)

One may believe that the dynamical correction (i.e., the so-called Lamb-Stark shift), together with the bare Hamiltonian, describes the physical Hamiltonian of the system, i.e., the one to which the system should thermalize in contact with a single bath. However, there is no simple physical argument why this should be the case. On the contrary, introduced in the next section, the mean-force Hamiltonian should describe a proper local equilibrium based on the second law of thermodynamics. As we will see later, those two corrections, in general, are entirely different. What is more, they have to be different to ensure the proper steady-state of the master equation.

B. Mean-force correction

Let us then introduce the mean-force Hamiltonian $H_{\rm mf}$, defined according to the marginal Gibbs state of the global equilibrium, i.e.,

$$\frac{e^{-\beta H_{\rm mf}}}{\operatorname{Tr}_S[e^{-\beta H_{\rm mf}}]} = \frac{\operatorname{Tr}_R[e^{-\beta H}]}{\operatorname{Tr}_{SR}[e^{-\beta H}]}.$$
(17)

One should notice that the above equation does not specify uniquely the ground state energy of the mean-force Hamiltonian, i.e., the equation is invariant under the transformation $H_{\rm mf} \to H_{\rm mf} + \Delta$ for arbitrary real constant Δ . Commonly, this constant is fixed by the convention [21]:

$$\operatorname{Tr}_{SR}[e^{-\beta H}] = \operatorname{Tr}_{S}[e^{-\beta H_{\mathrm{mf}}}]\operatorname{Tr}_{R}[e^{-\beta H_{R}}], \qquad (18)$$

such that we have the following relation

$$e^{-\beta \left(H_0 + \lambda^2 H_{\rm mf}^{(2)} + \dots\right)} = \mathcal{Z}_R^{-1} \operatorname{Tr}_R[e^{-\beta H}], \tag{19}$$

where we expanded the mean-force Hamiltonian into a series. Later, we concentrate only on the second-order correction $H_{\rm mf}^{(2)}$. Notice, that the zeroth-order Hamiltonian of the mean-force state (19) is fixed by putting $\lambda = 0$, i.e.,

$$\mathcal{Z}_{R}^{-1} \operatorname{Tr}_{R}[e^{-\beta H}]\Big|_{\lambda=0} = \mathcal{Z}_{R}^{-1} \operatorname{Tr}_{R}[e^{-\beta (H_{0}+H_{R})}] = e^{-\beta H_{0}},$$
(20)

which is given by the bare Hamiltonian H_0 .

We stress that the mean-force is the correct physical Hamiltonian of a weakly coupled system to a single thermal bath, in a sense that the system should thermalize to its corresponding Gibbs state. Otherwise, if the open system dynamics in the long time limit tends to a different steady-state, one might still extract work by thermalizing the whole system to its global equilibrium, which would violate the second law of thermodynamics. In accordance, the mean-force Hamiltonian is the reference point for any open system dynamics: The validation of the particular master equation is that its steady-state should be the Gibbs state with respect to the mean-force Hamiltonian. Let us now then introduce the final, steady-state correction.

C. Steady-state correction

In analogy to the mean-force correction, we define the *steady-state correction* $H_{\rm st}^{(2)}$ as the following Gibbs state:

$$\rho = e^{-\beta (H_0 + \lambda^2 H_{\rm st}^{(2)} + \dots)}.$$
(21)

(We do not introduce the normalization of ρ since it is unnecessary for the following discussion.) Based on the general equilibration postulate, we believe that such a Gibbs state should be the stationary state for the system coupled to a single heat bath (at inverse temperature β). Thus, the master equation with the generator \mathcal{L}_t that describes this process, should satisfy the following steady-state condition:

$$\lim_{t \to \infty} \mathcal{L}_t[\varrho] \equiv \mathcal{L}_\infty[\varrho] = 0.$$
⁽²²⁾

To find the solution for the (second-order) steady-state correction $H_{\rm st}^{(2)}$, we adapt the perturbative method. First, we expand the generator of the master equation:

$$\mathcal{L}_{\infty}[\varrho] = \mathcal{L}_{\infty}^{(0)}[\varrho] + \lambda^2 \mathcal{L}_{\infty}^{(2)}[\varrho] + \lambda^4 \mathcal{L}_{\infty}^{(4)}[\varrho] + \dots,$$
(23)

Note that here we consider a general case, such that for any second-order master equation the generators of the fourth and higher orders identically vanish. In particular, for the master equation given by Eq. (6), we have

$$\mathcal{L}_{\infty}^{(0)}[\varrho] = i[\varrho, H_0], \tag{24}$$

$$\mathcal{L}_{\infty}^{(2)}[\varrho] = \sum_{\omega,\omega'} \sum_{\alpha\beta} \left[i\Upsilon_{\alpha\beta}^{(\text{dyn})}(\omega,\omega')[\rho, A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega')] + K_{\alpha\beta}(\omega,\omega') \left(A_{\beta}(\omega')\rho A_{\alpha}^{\dagger}(\omega) - \frac{1}{2} \{A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega'), \rho\} \right) \right], \quad (25)$$

$$\mathcal{L}_{\infty}^{(4)}[\varrho] = \mathcal{L}_{\infty}^{(6)}[\varrho] = \dots = 0.$$

$$(26)$$

Similarly, we expand the postulated stationary-state, i.e.,

$$\varrho = \varrho_0 + \lambda^2 \varrho_2 + \lambda^4 \varrho_4 \dots, \tag{27}$$

where by using the Dyson series for the lowest orders we get:

$$\varrho_0 = e^{-\beta H_0}, \quad \varrho_2 = -e^{-\beta H_0} \int_0^\beta dt \ e^{tH_0} H_{\rm st}^{(2)} e^{-tH_0}. \tag{28}$$

Finally, from (22), we obtain the set of equations (for each order of λ):

. . .

$$\mathcal{L}_{\infty}^{(0)}[\varrho_0] = 0 \tag{29}$$

$$\mathcal{L}_{\infty}^{(0)}[\varrho_2] + \mathcal{L}_{\infty}^{(2)}[\varrho_0] = 0, \tag{30}$$

$$\mathcal{L}_{\infty}^{(0)}[\varrho_4] + \mathcal{L}_{\infty}^{(2)}[\varrho_2] + \mathcal{L}_{\infty}^{(4)}[\varrho_0] = 0, \tag{31}$$

Our main goal is to solve the following equations to find the solution for the steady-state correction
$$H_{\rm st}^{(2)}$$
. However, as it was highlighted in [10], the second-order equation (30) only provides the solution for off-diagonal terms of the correction, whereas the solution for the diagonal part involves the fourth-order equation (31). In the following (Section III and IV), we provide the general solution for off-diagonal elements for the master equation of the form (6) and the cumulant equation (introduced in Section IV). Moreover, in the Section V, we find the total solution for $H_{\rm st}^{(2)}$ (involving the diagonal part) for the two-dimensional system via solving the fourth-order equation. We also prepare

At the end of this section, we want to make two remarks. First, notice that the choice $\rho_0 = e^{-\beta H_0}$ in the definition of (21) satisfies the zeroth-order equation (29) for the generator (24). However, the solution is not unique and, in particular, it cannot be specified (in contrast to the mean-force correction) in the limit $\lambda \to 0$. Nevertheless, this choice is dictated by the general equilibration postulate, i.e., in the limit of a vanishing coupling constant the system should thermalize to the Gibbs state with respect to its bare Hamiltonian. By reason of this ambiguity, later we provide the necessary condition for a master equation (which is in fact the detailed balance condition) to satisfy the higher order equation for the steady-state ρ with $\rho_0 = e^{-\beta H_0}$. Secondly, similarly to the mean-force correction, Eq. (22) determines the diagonal part of the correction $H_{\rm st}^{(2)}$ only up to an arbitrary constant, i.e., if ρ is the solution, then also is $\rho e^{-\beta\delta}$, where δ is the identity matrix multiplied by an arbitrary real number. This provides us an additional degree of freedom that does not affect the physical meaning (i.e., the only differences of energies matter and not is absolute values).

III. RESULTS: MEAN-FORCE AND (OFF-DIAGONAL) STEADY-STATE CORRECTION

We are ready to state our first main result regarding the mean-force correction.

Theorem 1. The mean-force correction is equal to:

the fundamentals for methods for the general solution.

Ou

as

cor

III

$$\Upsilon_{\alpha\beta}^{(mf)}(\omega,\omega') = \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \ D(\omega,\omega',\Omega) \ \gamma_{\alpha\beta}(\Omega), \quad D(\omega,\omega',\Omega) = \frac{1}{\omega'-\Omega} - \frac{(\omega-\omega')(e^{\beta(\omega-\Omega)}-1)}{(\omega-\Omega)(\omega'-\Omega)(e^{\beta(\omega-\omega')}-1)}, \quad (32)$$

or equivalently, in terms of the $S_{\alpha\beta}(\omega)$ function, it takes the form:

$$\Upsilon_{\alpha\beta}^{(mf)}(\omega,\omega') = \frac{1}{e^{\beta\omega} - e^{\beta\omega'}} \left(e^{\beta\omega} \mathcal{S}_{\alpha\beta}(\omega') - e^{\beta\omega'} \mathcal{S}_{\alpha\beta}(\omega) + e^{\beta(\omega+\omega')} \left(\mathcal{S}_{\beta\alpha}(-\omega') - \mathcal{S}_{\beta\alpha}(-\omega) \right) \right).$$
(33)

Proof. The sketch of the proof is as follows. To solve Eq. (17) we write the exponents from both sides of the equality in terms of the Dyson series, which formally can be expressed via the time-ordering operator \mathcal{T} as:

$$e^{-\beta H_{\rm mf}} = e^{-\beta (H_0 + \delta H_{\rm mf})} = e^{-\beta H_0} \mathcal{T} e^{-\int_0^\beta dt \ \delta \hat{H}_{\rm mf}(t)}.$$
(34)

$$e^{-\beta H} = e^{-\beta (H_0 + H_R + \lambda H_I)} = e^{-\beta H_0} \mathcal{T} e^{-\lambda \int_0^\beta dt \ \hat{H}_I(t)},$$
(35)

where we put $\delta H_{\rm mf} = H_{\rm mf} - H_0$ and we define a time-dependent operators $\hat{A}(t) = e^{t(H_0 + H_R)} A e^{-t(H_0 + H_R)}$. Consequently, the equality (17) can be rewritten in the form:

$$\operatorname{Tr}_{R}\left[\mathcal{T}\left(e^{-\int_{0}^{\beta}dt\ \delta\hat{H}_{\mathrm{mf}}(t)}-e^{-\lambda\int_{0}^{\beta}dt\ \hat{H}_{I}(t)}\right)\gamma_{R}\right]=0.$$
(36)

Note that this is exact for arbitrary coupling strength λ . Then, considering the weak-coupling limit ($\lambda \ll 1$), we expand the above equality and obtain within the second-order the following condition for the mean-force correction:

$$\int_{0}^{\beta} dt \ \hat{H}_{\rm mf}^{(2)}(t) = -\int_{0}^{\beta} dt \int_{0}^{t} ds \left\langle \hat{H}_{I}(t) \hat{H}_{I}(s) \right\rangle_{\gamma_{R}}.$$
(37)

After substituting the representation of the mean-force Hamiltonian given by Eq. (4) and the interaction term (2) with the definition of jump operators (5), we are able to calculate the coefficients $\Upsilon_{\alpha\beta}^{(\mathrm{inf})}(\omega,\omega')$ (see the detailed proof in Appendix **B**).

7

Theorem 1 is the first derivation in the literature of the second-order mean-force Hamiltonian for a general weakcoupling of the form (1) (see also [18]). Remarkably the expression (32) does not exhibit any poles, as it is usual for a dynamical correction. One observes that the coefficients are symmetric, i.e., $\Upsilon_{\alpha\beta}^{(mf)}(\omega, \omega') = \Upsilon_{\alpha\beta}^{(mf)}(\omega', \omega)$, which together with $\mathcal{S}^*_{\alpha\beta}(\omega) = \mathcal{S}_{\beta\alpha}(\omega)$, ensures the hermiticity of the Hamiltonian. According to this expression, let us compare the dynamical Hamiltonian with the mean-force. Using Eq. (16), we write down the dynamical correction in terms of $\gamma_{\alpha\beta}(\omega)$ and $\mathcal{S}_{\alpha\beta}(\omega)$, for the Bloch-Redfield master equation:

$$\Upsilon_{\alpha\beta}^{(\mathrm{dyn})}(\omega,\omega') = \frac{1}{2} (\mathcal{S}_{\alpha\beta}(\omega) + \mathcal{S}_{\alpha\beta}(\omega')) + \frac{i}{4} (\gamma_{\alpha\beta}(\omega) - \gamma_{\alpha\beta}(\omega')), \qquad (38)$$

whereas for the Davies equation the off-diagonal elements vanish (due to the secular approximation). It is seen that the mean-force correction is different than the dynamical one; in particular, the dynamical correction has non-zero imaginary part in contrast to the real coefficients of the mean-force. In Section VIA we compare numerically both of the Hamiltonians for the spin-boson model.

Our second main result is the solution of the Eq. (30). In particular, this provides the formula for off-diagonal terms of the steady-state correction $H_{\rm st}^{(2)}$, which later is compared to the mean-force Hamiltonian.

Theorem 2. If the following conditions are satisfied:

(i) the detailed balance relation:

$$K_{\alpha\beta}(\omega,\omega) = K_{\beta\alpha}(-\omega,-\omega)e^{\beta\omega},\tag{39}$$

(ii) and for $\omega \neq \omega'$:

$$\Upsilon_{\alpha\beta}^{(st)}(\omega,\omega') = \Upsilon_{\alpha\beta}^{(dyn)}(\omega,\omega') + \frac{i}{e^{\beta\omega} - e^{\beta\omega'}} \left(K_{\beta\alpha}(-\omega,-\omega')e^{\beta(\omega+\omega')} - \frac{1}{2}K_{\alpha\beta}(\omega',\omega)(e^{\beta\omega} + e^{\beta\omega'}) \right), \tag{40}$$

then $\varrho = e^{-\beta(H_0 + \lambda^2 H_{st}^{(2)} + \dots)}$ is a solution of the equation: $\mathcal{L}_{\infty}^{(0)}[\varrho_2] + \mathcal{L}_{\infty}^{(2)}[\varrho_0] = 0.$

Proof. In order to construct the solution of the Eq. (30) we transform the operator equation into the algebraic one, by introducing the representation:

$$\mathcal{L}_{\infty}^{(k)}[\rho_l] = \sum_{\alpha,\beta} \sum_{\omega,\omega'} g_{\alpha\beta}^{(kl)}(\omega,\omega') e^{-\beta H_0} A_{\alpha}(\omega) A_{\beta}(\omega'), \tag{41}$$

where l + k = 2. In accordance, the Eq. (30) can be rewritten as:

$$\sum_{\alpha,\beta} \sum_{\omega,\omega'} (g_{\alpha\beta}^{(02)}(\omega,\omega') + g_{\alpha\beta}^{(20)}(\omega,\omega')) e^{-\beta H_0} A_{\alpha}(\omega) A_{\beta}(\omega') = 0.$$
(42)

such that the solution is given by:

$$g_{\alpha\beta}^{(02)}(\omega,\omega') + g_{\alpha\beta}^{(20)}(\omega,\omega') = 0$$

$$\tag{43}$$

for each ω, ω' and α, β . In particular, for the master equation of the form (6), we get

$$-i(\omega'-\omega)\Upsilon_{\alpha\beta}^{(\mathrm{st})}(\omega,\omega')\alpha(\omega'-\omega) + i\Upsilon_{\alpha\beta}^{(\mathrm{dyn})}(\omega,\omega')(1-e^{-\beta(\omega'-\omega)}) + e^{\beta\omega}K_{\beta\alpha}(-\omega',-\omega) - \frac{1}{2}K_{\alpha\beta}(\omega,\omega')(e^{-\beta(\omega'-\omega)}+1) = 0, \quad (44)$$

where

$$\alpha(\omega) = \int_0^\beta dt \ e^{-t\omega} = \begin{cases} \frac{1 - e^{-\beta\omega}}{\omega}, & \omega \neq 0\\ \beta, & \omega = 0 \end{cases}$$
(45)

From which follows the Theorem 2.

The non-zero value of the correction (40) predicts the so-called finite "steady-state coherences" of the density matrix ρ . Those coherences represents the off-diagonal elements with respect to the basis of the bare Hamiltonian H_0 , such that according to the Gibbs state of this Hamiltonian, ρ is out of equilibrium. However, as it was mentioned previously, the correct (local) equilibrium of the system should be identified with the mean-force Hamiltonian $H_{\rm mf}$ rather than the bare one. Indeed, as it follows from the presented below corollary, the Bloch-Redfield master equation predicts the same (second-order) steady-state coherences as for the Gibbs state of the mean-force Hamiltonian, which provides that the off-diagonal part of its stationary state is equal to the proper equilibrium state.

Corollary 2.1. For the Bloch-Redfield equation, i.e., if $K_{\alpha\beta}(\omega, \omega') = \gamma_{\alpha\beta}(\omega, \omega')$ and $\Upsilon^{(dyn)}_{\alpha\beta}(\omega, \omega') = S_{\alpha\beta}(\omega, \omega')$, we have

$$\Upsilon^{(st)}_{\alpha\beta}(\omega,\omega') = \Upsilon^{(mf)}_{\alpha\beta}(\omega,\omega') \quad for \quad \omega \neq \omega',$$
(46)

whereas if $K_{\alpha\beta}(\omega,\omega') = \gamma_{\alpha\beta}(\omega)\delta_{\omega,\omega'}$, then

$$\Upsilon^{(st)}_{\alpha\beta}(\omega,\omega') = \Upsilon^{(dyn)}_{\alpha\beta}(\omega,\omega') \quad for \quad \omega \neq \omega'.$$
(47)

On the contrary to the Bloch-Redfield master equation, if the secular approximation is done for $\gamma_{\alpha\beta}(\omega,\omega')$, then the steady-state coherences are equal to the Gibbs state of the bare Hamiltonian with the dynamical correction $H_{\rm dyn}^{(2)}$. However, commonly the secular approximation is also applied for the dynamical part, as it is for the Davies equation (see Eq. (12)), such that the dynamical correction is diagonal and the steady-state coherences vanish.

IV. CUMULANT EQUATION

Let us now introduce the cumulant equation [14] that describes the open system dynamics. Unlike the previous models of open system, which are derived as the solution of the von Neumann equation, the cumulant equation is initially derived in the form of the dynamical map (which does not involve the Markovian approximation):

$$\tilde{\rho}(t) = e^{\tilde{K}_t^{(2)}} \tilde{\rho}(0), \tag{48}$$

where

$$\tilde{K}_{t}^{(2)}[\rho] = \lambda^{2} \int_{0}^{t} ds \sum_{\omega,\omega'} \sum_{\alpha\beta} \left(i \tilde{\mathcal{S}}_{\alpha\beta}(\omega,\omega',s) [\rho, A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega')] + \tilde{\gamma}_{\alpha\beta}(\omega,\omega',s) \left(A_{\beta}(\omega')\rho A_{\alpha}^{\dagger}(\omega) - \frac{1}{2} \{ A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega'), \rho \} \right) \right)$$

$$\tag{49}$$

is the generator of the map in the interaction picture, such that $\tilde{\rho}(t) = e^{iH_0 t} \rho(t) e^{-iH_0 t}$, and

$$\tilde{\gamma}_{\alpha\beta}(\omega,\omega',t) = e^{i(\omega-\omega')t}\gamma_{\alpha\beta}(\omega,\omega',t),\tag{50}$$

$$\tilde{\mathcal{S}}_{\alpha\beta}(\omega,\omega',t) = e^{i(\omega-\omega')t} \mathcal{S}_{\alpha\beta}(\omega,\omega',t).$$
(51)

We observe that the super-operator $\tilde{K}_t^{(2)}$ is closely related to the generator of the Bloch-Redfield master equation. Indeed, the Bloch-Redfield generator in the interaction picture is given by:

$$\tilde{\mathcal{L}}_{t}^{R}[\rho] = e^{iH_{0}t}\mathcal{L}_{t}^{R}[e^{-iH_{0}t}\rho e^{iH_{0}t}]e^{-iH_{0}t} - i[\rho, H_{0}]$$

$$= \lambda^{2} \sum_{\omega,\omega'} \sum_{\alpha\beta} \left(i[\rho, \tilde{\mathcal{S}}_{\alpha\beta}(\omega, \omega', t)A_{\alpha}^{\dagger}(\omega)\rho A_{\beta}(\omega')] + \tilde{\gamma}_{\alpha\beta}(\omega, \omega', t) \left(A_{\beta}(\omega')\rho A_{\alpha}^{\dagger}(\omega) - \frac{1}{2} \{A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega'), \rho\} \right) \right), (53)$$

which simply implies that

$$\frac{d\tilde{K}_t^{(2)}}{dt} = \tilde{\mathcal{L}}_t^R.$$
(54)

Remarkably, the time-integrated coefficients appearing in the disspative part of the super-operator $\tilde{K}_t^{(2)}$, i.e.,

$$\int_{0}^{t} ds \,\,\tilde{\gamma}_{\alpha\beta}(\omega,\omega',s) \tag{55}$$

are the elements of a positive semi-definite matrix (see, e.g., [15]). This property implies that the $\tilde{K}_t^{(2)}$ super-operator is of the GKSL form. In this way, the cumulant equation defines one parameter family of CPTP dynamical maps. This feature of the cumulant equation is its advantage over the Bloch-Redfield equation, for which the fundamental property of completely positive evolution is not satisfied.

On the other hand, we formulate the cumulant master equation by taking a time derivative of Eq. (48), i.e.,

$$\frac{d}{dt}\tilde{\rho}(t) = \left(\frac{d}{dt}e^{\tilde{K}_t^{(2)}}\right)[\tilde{\rho}(0)] = \left(\int_0^1 ds \ e^{s\tilde{K}_t^{(2)}}\frac{d\tilde{K}_t^{(2)}}{dt}e^{(1-s)\tilde{K}_t^{(2)}}\right)[\tilde{\rho}(0)]$$
(56)

$$= \left(\int_0^1 ds \ e^{s\tilde{K}_t^{(2)}} \frac{d\tilde{K}_t^{(2)}}{dt} e^{-s\tilde{K}_t^{(2)}}\right) [\tilde{\rho}(t)] \equiv \tilde{\mathcal{L}}_t^C[\tilde{\rho}(t)].$$
(57)

Equivalently, the generator of the cumulant master equation can be written as:

$$\tilde{\mathcal{L}}_{t}^{C} = \frac{e^{[\tilde{K}_{t}^{(2)}, \cdot]} - 1}{[\tilde{K}_{t}^{(2)}, \cdot]} \frac{d\tilde{K}_{t}^{(2)}}{dt},$$
(58)

such that

$$\frac{d}{dt}\tilde{\rho}(t) = \left(\frac{d}{dt}\tilde{K}_{t}^{(2)} + \frac{1}{2}[\tilde{K}_{t}^{(2)}, \frac{d}{dt}\tilde{K}_{t}^{(2)}] + \dots\right)\tilde{\rho}(t),\tag{59}$$

where $\frac{d}{dt}\tilde{K}_t^{(2)} = \tilde{\mathcal{L}}_t^R$. According to this, we define the generator of the cumulant master equation in the Schrödinger picture:

$$\mathcal{L}_{t}^{C}[\rho] = i[\rho, H_{0}] + e^{-iH_{0}t} \tilde{\mathcal{L}}_{t}^{C}[e^{iH_{0}t}\rho e^{-iH_{0}t}]e^{iH_{0}t}.$$
(60)

From this we obtain the following theorem:

Theorem 3. The generator of the cumulant master equation is, up to the second-order, equal to the generator of the Bloch-Redfield master equation, *i.e.*,

$$\mathcal{L}_t^C = \mathcal{L}_t^R + O(\lambda^4) \tag{61}$$

Proof. Expanding the generator \mathcal{L}_t^C to the second-order and putting Eq. (54), we get

$$\mathcal{L}_{t}^{C}[\rho] = i[\rho, H_{0}] + e^{-iH_{0}t} \frac{d\dot{K}_{t}^{(2)}}{dt} [e^{iH_{0}t}\rho e^{-iH_{0}t}] e^{iH_{0}t} + O(\lambda^{4}) = \mathcal{L}_{t}^{R}[\rho] + O(\lambda^{4})$$
(62)

Since the generator of the cumulant master equation, up to the second-order, is equal to the Bloch-Redfield generator, we see (based on to the Corollary 2.1) that the off-diagonal part of the mean-force Hamiltonian $H_{\rm mf}^{(2)}$ is the solution for a steady-state of the cumulant, too. In this way, we proved a significant advantage of the cumulant equation over the Bloch-Redfield since both predict the same proper coherences of the stationary state, yet the dynamics governed by the cumulant is contrary to the Bloch-Redfield equation, completely positive.

V. DIAGONAL ELEMENTS FROM FOURTH-ORDER TERMS

In this section, we try to derive the diagonal part of the correction $H_{\rm st}^{(2)}$, which according the representation (4) is given by the coefficients $\Upsilon_{\alpha\beta}^{(\rm st)}(\omega,\omega)$. In principle, this could be done by solving the fourth-order equation (31), i.e.,

$$\mathcal{L}_{\infty}^{(0)}[\varrho_4] + \mathcal{L}_{\infty}^{(2)}[\varrho_2] + \mathcal{L}_{\infty}^{(4)}[\varrho_0] = 0.$$
(63)

However, on the contrary to the solution for off-diagonal terms, it is much more complex problem. In the following, we present a general method how to solve this equation and later we present the solution for a two-level system for the Bloch-Redfield and cumulant master equation.

For simplicity, throughout of this section, we assume that the interaction Hamiltonian is given by

$$H_I = S \otimes R \tag{64}$$

i.e., we replace the sum (2) by a single term (which can be straightforwardly generalized by adding the corresponding indices). According to this, we also simplify notation such that $\Upsilon_{\alpha\beta}^{(k)} \equiv \Upsilon_k$. Then, similarly to the second-order (see Eq. (41)), we start with writing the action of the generators in the basis of jump operators:

$$\mathcal{L}_{\infty}^{(k)}[\rho_l] = \sum_{\omega_1,\omega_2,\omega_3,\omega_4} g_{kl}(\omega_1,\omega_2,\omega_3,\omega_4) e^{-\beta H_0} A(\omega_1) A(\omega_2) A(\omega_3) A(\omega_4), \tag{65}$$

where l + k = 4, such that the fourth-order equation reads then as follows:

$$\sum_{\omega_1,\omega_2,\omega_3,\omega_4} (g_{04}(\omega_1,\omega_2,\omega_3,\omega_4) + g_{22}(\omega_1,\omega_2,\omega_3,\omega_4) + g_{40}(\omega_1,\omega_2,\omega_3,\omega_4))e^{-\beta H_0}A(\omega_1)A(\omega_2)A(\omega_3)A(\omega_4) = 0.$$
(66)

Then, we propose the following general proposition.

Proposition 1. Eq. (66) is satisfied if and only if the set of equation:

$$\sum_{(\omega_1,\omega_2,\omega_2,\omega_4)\in G(|k\rangle\to|k\rangle)} \left(g_{22}(\omega_1,\omega_2,\omega_3,\omega_4) + g_{40}(\omega_1,\omega_2,\omega_3,\omega_4)\right) = 0.$$
(67)

for each k, is satisfied, where $G(|k\rangle \rightarrow |k\rangle)$ denotes the set of all four-tuples $(\omega_1, \omega_2, \omega_2, \omega_4)$ of the form:

$$(\omega_1, \omega_2, \omega_2, \omega_4) = (\epsilon_l - \epsilon_k, \epsilon_m - \epsilon_l, \epsilon_j - \epsilon_m, \epsilon_k - \epsilon_j),$$
(68)

and $|k\rangle$ is an eigenstate of the bare Hamilton with energy ϵ_k .

Proof. We consider diagonal elements of the Eq. (66), such that we obtain the following set of equations

$$\sum_{\omega_1,\omega_2,\omega_3,\omega_4} (g_{04}(\omega_1,\omega_2,\omega_3,\omega_4) + g_{22}(\omega_1,\omega_2,\omega_3,\omega_4) + g_{40}(\omega_1,\omega_2,\omega_3,\omega_4))e^{-\beta\epsilon_k} \langle k | A(\omega_1)A(\omega_2)A(\omega_3)A(\omega_4) | k \rangle = 0.$$
(69)

for $k = 0, 1, 2, \ldots$ We see that since for arbitrary ρ we have:

$$\sum_{\omega_1,\omega_2,\omega_3,\omega_4} g_{04}(\omega_1,\omega_2,\omega_3,\omega_4) \langle k | e^{-\beta H_0} A(\omega_1) A(\omega_2) A(\omega_3) A(\omega_4) | k \rangle = \langle k | \mathcal{L}_{\infty}^{(0)}[\rho] | k \rangle = \langle k | [H_0,\rho] | k \rangle = 0, \tag{70}$$

so our condition is now just

$$\sum_{\omega_1,\omega_2,\omega_3,\omega_4} \left(g_{22}(\omega_1,\omega_2,\omega_3,\omega_4) + g_{40}(\omega_1,\omega_2,\omega_3,\omega_4) \right) e^{-\beta\epsilon_k} \left\langle k \right| A(\omega_1)A(\omega_2)A(\omega_3)A(\omega_4) \left| k \right\rangle = 0.$$

$$(71)$$

Moreover, one observes that $\langle k | A(\omega_1)A(\omega_2)A(\omega_3)A(\omega_4) | k \rangle$ is nonzero only if $\sum_k \omega_k = 0$. Consequently, let us denote by $G(|k\rangle \to |k\rangle)$ the set of all four-tuples $(\omega_1, \omega_2, \omega_2, \omega_4)$ of the form:

$$(\omega_1, \omega_2, \omega_2, \omega_4) = (\epsilon_l - \epsilon_k, \epsilon_m - \epsilon_l, \epsilon_j - \epsilon_m, \epsilon_k - \epsilon_j),$$
(72)

form which follows Eq. (67).

The coefficients Υ_{st} are encoded within the function g_{22} . In particular, for the master equation of the form (6) and ρ_2 given by Eq. (28), we have:

$$g_{22}(\omega_1,\omega_2,\omega_3,\omega_4) = \Upsilon_{\rm st}(-\omega_3,\omega_4)\alpha(\omega_3+\omega_4)\left(i\Upsilon_{\rm dyn}(-\omega_1,\omega_2)+\frac{1}{2}K(-\omega_1,\omega_2)\right) - \Upsilon_{\rm st}(-\omega_1,\omega_2)\alpha(\omega_1+\omega_2)\left(i\Upsilon_{\rm dyn}(-\omega_3,\omega_4)-\frac{1}{2}K(-\omega_3,\omega_4)\right) - e^{-\beta\omega_1}\Upsilon_{\rm st}(-\omega_2,\omega_3)\alpha(\omega_2+\omega_3)K(-\omega_4,\omega_1)$$
(73)

where $\alpha(\omega)$ is given by Eq. (45). Finally, (67) provides the general set of equations for the diagonal elements of the steady-state correction in terms of the $\Upsilon_{st}(\omega, \omega)$ coefficients. In the next section, we provide the solution for the two-level system for the Bloch-Redfield and cumulant master equation.

VI. TWO-LEVEL SYSTEM

Let us consider a two-level system with the bare Hamiltonian $H_0 = -\frac{\omega_0}{2}\sigma_z$ coupled to the thermal bath via a single interaction term $H_I = S \otimes R$, where $S = \vec{r} \cdot \vec{\sigma}$, $\vec{r} = (x, y, z)$, and $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ are the Pauli matrices. Then, the correction $H_k^{(2)}$ can be written in the following form:

$$H_{\mathbf{k}}^{(2)} = \begin{bmatrix} (x^2 + y^2)(\Upsilon_{\mathbf{k}}(\omega_0, \omega_0) - \Upsilon_{\mathbf{k}}(-\omega_0, -\omega_0)) & (x - iy)z(\Upsilon_{\mathbf{k}}(0, -\omega_0) - \Upsilon_{\mathbf{k}}(\omega_0, 0)) \\ (x + iy)z(\Upsilon_{\mathbf{k}}(0, -\omega_0) - \Upsilon_{\mathbf{k}}(\omega_0, 0)) & -(x^2 + y^2)(\Upsilon_{\mathbf{k}}(\omega_0, \omega_0) - \Upsilon_{\mathbf{k}}(-\omega_0, -\omega_0)) \end{bmatrix}$$
(74)

For the general master equation (6), the off-diagonal terms can be calculated from Eq. (40). In particular, for the Bloch-Redfield or cumulant master equation, if $\omega \neq \omega'$ we have generally $\Upsilon_{\rm st}(\omega, \omega') = \Upsilon_{\rm mf}(\omega, \omega')$ (see Corollary 2.1). In the following, we present the formulas for diagonal elements.

Proposition 2. The diagonal elements of the two-level system steady-state correction are given by:

(i) For a master equation of the form (6), obeying the detailed balance condition (i.e., $K(\omega, \omega) = e^{\beta \omega} K(-\omega, -\omega)$):

$$\Upsilon_{st}(\omega,\omega) = 0 \tag{75}$$

(ii) For the cumulant master equation (60):

$$\Upsilon_{st}(\omega,\omega) = \Upsilon_{mf}(\omega,\omega) - \Upsilon_{dyn}(\omega,\omega), \tag{76}$$

where $\Upsilon_{dyn}(\omega, \omega) = \mathcal{S}(\omega)$.

Proof. The proof is solely based on Eq. (67), where for the two-level system $\omega \in \{-\omega_0, 0, \omega_0\}$. Moreover, the first part of the Proposition 2 (i) is for the master equation of the form (6), which does not involve the fourth and higher order generators, such that identically $g_{40} = 0$. Thus, we have

$$\sum_{(\omega_1,\omega_2,\omega_2,\omega_4)\in G(|0\rangle\to|0\rangle)} g_{22}(\omega_1,\omega_2,\omega_3,\omega_4) = \sum_{(\omega_1,\omega_2,\omega_2,\omega_4)\in G(|1\rangle\to|1\rangle)} g_{22}(\omega_1,\omega_2,\omega_3,\omega_4) = 0, \tag{77}$$

from which we obtain:

$$\beta \left(\Upsilon_{\rm st}(-\omega,-\omega)K(-\omega,-\omega) - e^{-\beta\omega}\Upsilon_{\rm st}(\omega,\omega)K(\omega,\omega) \right) = 0, \tag{78}$$

where $\omega = \omega_0$ or $-\omega_0$. After applying the detailed balance condition, finally we get

$$\Upsilon_{\rm st}(\omega_0,\omega_0) = -\Upsilon_{\rm st}(-\omega_0,-\omega_0). \tag{79}$$

Since the solution for the stationary state is invariant under arbitrary shift of the energies (see the comment at the end of the Section II C), we simply put $\Upsilon_{\rm st}(0,0) = \Upsilon_{\rm st}(\omega_0,\omega_0) = \Upsilon_{\rm st}(-\omega_0,-\omega_0) = 0$.

The second part of the Proposition 2 (ii) is much more involving since it contains the fourth-order generator from the Eq. (59) (written in the interaction picture). Based on this, the fourth-order generator in the Schrödinger picture is given by:

$$\mathcal{L}_{t}^{(4)}[\rho] = \frac{1}{2}e^{-iH_{0}t} \left(\left[\tilde{K}_{t}^{(2)}, \frac{d}{dt} \tilde{K}_{t}^{(2)} \right] \left[e^{iH_{0}t} \rho e^{-iH_{0}t} \right] \right) e^{iH_{0}t} = \frac{1}{2} \int_{0}^{t} ds \ e^{-iH_{0}t} \left(\left[\tilde{\mathcal{L}}_{s}^{R}, \tilde{\mathcal{L}}_{t}^{R} \right] \left[e^{iH_{0}t} \rho e^{-iH_{0}t} \right] \right) e^{iH_{0}t}$$
(80)

However, to calculate the coefficient g_{40} we act on the state ρ_0 (that commutes with H_0), and we only consider the diagonal term, such that

$$\langle k | \mathcal{L}_{t}^{(4)}[\varrho_{0}] | k \rangle = \frac{1}{2} \int_{0}^{t} ds \ \langle k | \left[\tilde{\mathcal{L}}_{s}^{R}, \tilde{\mathcal{L}}_{t}^{R} \right] [\varrho_{0}] | k \rangle \,.$$

$$\tag{81}$$

From this one can calculate the expression for a time-dependent coefficient g_{40} . Due to its complicated form, we do not present it explicitly. However, for a two-dimensional system we have a significant simplification since similarly to the coefficient g_{22} , the sum in the long time limit is just given by:

$$\sum_{(\omega_1,\omega_2,\omega_2,\omega_4)\in G(|0\rangle\to|0\rangle)} g_{40}(\omega_1,\omega_2,\omega_2,\omega_4) = \sum_{(\omega_1,\omega_2,\omega_2,\omega_4)\in G(|1\rangle\to|1\rangle)} g_{40}(\omega_1,\omega_2,\omega_2,\omega_4)$$
$$= \frac{1}{2}e^{-\beta\omega}(1+e^{\beta\omega})\gamma(\omega)\int_0^\infty ds \ (e^{-\beta\omega}\gamma(\omega,s)-\gamma(-\omega,s)) \quad (82)$$



FIG. 1: Comparison of the diagonal and off-diagonal elements of the two-level system Hamiltonian corrections (74) for a spin-boson model with respect to $\beta\omega_0$, where β is the inverse temperature of the bath and ω_0 is the bare frequency of the qubit. The orange line represents the mean-force coefficient, the blue one represents the dynamical,

and the dashed lines represent the steady-state coefficients of the dynamical equations: green is for the Bloch-Redfield master equation and black for the cumulant equation. All plots were computed for cut-off frequency

Bioch-Redfield master equation and black for the cumulant equation. All plots were computed for cut-off frequency $\beta\omega_c = 50.$

where, similarly to the previous case, $\omega = \omega_0$ or $-\omega_0$. Finally, we need to solve

$$\beta \left(\Upsilon_{\rm st}(-\omega,-\omega)K(-\omega,-\omega) - e^{-\beta\omega}\Upsilon_{\rm st}(\omega,\omega)K(\omega,\omega) \right) + \frac{1}{2}e^{-\beta\omega}(1+e^{\beta\omega})\gamma(\omega) \int_0^\infty ds \, \left(e^{-\beta\omega}\gamma(\omega,s) - \gamma(-\omega,s) \right) = 0 \tag{83}$$

which leads to the solution (76).

Now, we can combine the results on off-diagonal (Theorem 2) and diagonal (Proposition 2) elements for the two-level system, in the following theorem.

Theorem 4. For the two-level system:

$$H_{et}^{(2)} = H_{mf}^{(2)} - \mathcal{D}[H_{dvm}^{(2)}] \quad for \ the \ cumulant \ equation, \tag{84}$$

$$H_{st}^{(2)} = H_{mf}^{(2)} - \mathcal{D}[H_{mf}^{(2)}] \quad for \ the \ Bloch-Red field \ equation, \tag{85}$$

where $\mathcal{D}[H_k^{(2)}]$ denotes the diagonal part of the representation (74).

A. Spin-boson model

Let us now apply our results to the well-known spin-boson model. We consider a qubit coupled to a bosonic bath with the Hamiltonian:

$$H = -\frac{\omega_0}{2}\sigma_z + \sum_k \Omega_k a_k^{\dagger} a_k + (x\sigma_x + y\sigma_y + z\sigma_z) \sum_{k=1}^{\infty} \lambda_k (a_k + a_k^{\dagger})$$
(86)

where a_k and a_k^{\dagger} are the bosonic annihilation and creation operators. For this model we have the autocorrelation function:

$$\langle R(t)R\rangle_{\gamma_R} = \int_0^\infty d\Omega \ J(\Omega) \left(\coth[\frac{\beta\Omega}{2}]\cos(\Omega t) - i\sin(\Omega t) \right)$$
(87)

where we consider the Ohmic environment, i.e.,

$$J(\Omega) = \gamma \Omega e^{-\frac{|\Omega|}{\omega_c}},\tag{88}$$

where γ is a parameter which modifies the strength of the interaction and ω_c is a cutoff frequency. According to the general representation (74), we then compute the coefficients $\Upsilon_k(\omega_0, \omega_0) - \Upsilon_k(-\omega_0, -\omega_0)$ representing the diagonal part, and $\Upsilon_k(0, -\omega_0) - \Upsilon_k(\omega_0, 0)$ which describes the off-diagonal elements, for dynamical, mean-force and steady-state correction. The results are presented in the Fig. 1.

The most important conclusion coming from the numerical simulation is the very good agreement of the diagonal coefficient for the steady-state of the cumulant equation with the mean-force correction, which even though is shifted, still predicts the same behaviour with respect to growing frequency ω_0 . On the contrary, the Bloch-Redfield equation, which does not involve the fourth-order generator, predicts no change of the diagonal elements at all. As it was pointed out before, the off-diagonal elements predicted by the cumulant and Bloch-Redfield equation coincide with the mean-force. For completeness, we also plot the dynamical correction, which as it is seen differs significantly from other corrections. Especially, on the contrary to others, it predicts the imaginary part of the off-diagonal coefficient.

We see that non-vanishing value of the $\Upsilon_{mf}(0, -\omega_0) - \Upsilon_{mf}(\omega_0, 0)$ predicts the steady-state coherences in the equilibrium state. This was already known in the literature, and in the Appendix E we show that our general formulas very easily predicts the same value reported in [12], if applied for a particular case of a two-level system. Thus, our general formulas are verified to reproduce the particular cases known in the literature, as well as provides a simpler framework for their calculation.

VII. CONCLUSIONS

We have shown the significant advantage of the cumulant equation over the Bloch-Redfield or Davies master equation. Let us point out all of its advantages:

- (i) Its derivation from the exact dynamics does not involve the Markovian approximation;
- (ii) It is completely positive;
- (iii) It predicts the proper steady-state coherences at equilibrium in the long-time limit;
- (iv) For the two-level system, the diagonal part of its stationary state (in the long-time limit) is very close to equilibrium.

In comparison, the Bloch-Redfield master equation only satisfies the condition (iii) and the Davies equation only (ii). We thus conclude that cumulant equation is the best candidate to proper description of the thermalization process of the open systems, which simultaneously is completely positive. We believe that involving the fourth-order of the cumulant super-operator (i.e., the additional $K_t^{(4)}$ term in the exponent (48)) should result in exact solution for diagonal part, too. However, this may break down the positivity of the map, such that the trade-off between long times and the short or intermediate ones can appear.

Moreover, our results were proved on a high level of generality, and thus, contribute to other important areas of open quantum systems. In particular, we derive the second-order formula of the mean-force Hamiltonian for an arbitrary form of the coupling, as well as the solution for steady-state coherences for the whole class of master equations. We also discuss the relationship between different corrections to the bare Hamiltonian. Especially, we compute and compare all of the corrections quantitatively for a particular spin-boson model.

The most important direction for future studies is to analyze the stationary states out of equilibrium (e.g., systems coupled to two baths in different temperatures). As it was pointed out in [13], for systems coupled to a single bath, one can always artificially modify the completely positive GKSL master equation, such that it thermalizes to the proper equilibrium. This, however, results in a wrong description of the non-equilibrium situation. Nevertheless, since the cumulant equation is derived according to the Born approximation from the exact dynamics, we believe that it should also predict the proper stationary state out of equilibrium (in the weak-coupling limit).

Few final remarks are here in order. As usual, the obtained corrections are cut-off dependent and often diverge with the growing cut-off frequency. This is actually ubiquitous in the literature of the topic - the problem of cut-off dependence is at the moment swept under the carpet (see however [17] and [22] for discussion). Secondly, we do not touch on the issue of renormalization: the derivation of the master equation should be based not on bare Hamiltonian but the renormalized one (as advocated in [14] and [22]). We have not followed this in the present manuscript to keep clear the main message.

ACKNOWLEDGEMENT

We thank Antonio Mandarino for drawing our attention to Ref. [12]. We also acknowledge discussions with participants of Open Systems Seminars at ICTQT. This work is supported by Foundation for Polish Science (FNP), IRAP project ICTQT, contract no. 2018/MAB/5, co-financed by EU Smart Growth Operational Programme. MH is also partially supported by Polish National Science Centre grant OPUS-21 (grant No: 2021/41/B/ST2/03207). MW acknowledges grant PRELUDIUM-20 (grant number: 2021/41/N/ST2/01349) from the National Science Center.

- [2] H.-P. Breuer and F. Petruccione, The Theory of Open Quantum Systems (2006).
- [3] A. G. Redfield, IBM Journal of Research and Development 1, 19 (1957).
- [4] F. Bloch, Phys. Rev. **105**, 1206 (1957).
- [5] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, J. Math. Phys. 17, 821 (1976).
- [6] G. Lindblad, Comm. Math. Phys. 48, 119 (1976).
- [7] E. B. Davies, Comm. Math. Phys. **39**, 91 (1974).
- [8] A. S. Trushechkin, M. Merkli, J. D. Cresser, and J. Anders, (2021), arXiv:2110.01671 [quant-ph].
- [9] C. H. Fleming and N. I. Cummings, Phys. Rev. E 83, 031117 (2011).
- [10] J. Thingna, J.-S. Wang, and P. Hänggi, The Journal of Chemical Physics 136, 194110 (2012).
- [11] J. D. Cresser and J. Anders, Phys. Rev. Lett. 127, 250601 (2021).
- [12] A. Purkayastha, G. Guarnieri, M. T. Mitchison, R. Filip, and J. Goold, Npj Quantum Inf. 6 (2020).
- [13] D. Tupkary, A. Dhar, M. Kulkarni, and A. Purkayastha, (2022), arXiv:2105.12091 [quant-ph].
- [14] R. Alicki, Phys. Rev. A 40, 4077 (1989).
- [15] A. Rivas, Physical Review A **95** (2017).
- [16] A. Rivas, Entropy **21**, 725 (2019).
- [17] M. Winczewski, A. Mandarino, M. Horodecki, and R. Alicki, (2021), arXiv:2106.05776 [quant-ph].
- [18] G. Timofeev and A. Trushechkin, (2022), arXiv.
- [19] T. Mori and S. Miyashita, Journal of the Physical Society of Japan 77, 124005 (2008).
- [20] Y. Subaş I, C. H. Fleming, J. M. Taylor, and B. L. Hu, Phys. Rev. E 86, 061132 (2012).
- [21] B. Roux and T. Simonson, Biophysical Chemistry 78, 1 (1999).
- [22] M. Winczewski and R. Alicki, (2021), arXiv:2112.11962 [quant-ph].
- [23] M. Hunacek, The Mathematical Gazette 92, 380–382 (2008).

R. Alicki and K. Lendi, *Quantum Dynamical Semigroups and Applications*, Lecture notes in physics (Springer-Verlag, 1987).

Appendix A: Preliminaries

We consider the system and bath Hamiltonian:

$$H = H_0 + H_R + \lambda H_I \tag{A1}$$

such that H_0 and H_R are the free Hamiltonians of the system and the bath, respectively, and H_I is the interaction Hamiltonian. Except the section B1 and B2, where we do not assume any particular form of the interaction term, throughout the paper we consider the following explicit form:

$$H_I = \sum_{\alpha} A_{\alpha} \otimes R_{\alpha}.$$
 (A2)

We introduce the time-dependent operators:

$$A_{\alpha}(t) = e^{iH_{0}t}A_{\alpha}e^{-iH_{0}t}, \quad R_{\alpha}(t) = e^{iH_{R}t}R_{\alpha}e^{-iH_{R}t}.$$
(A3)

and jump operators (acting on the system Hilbert space):

$$A_{\alpha}(\omega) = \sum_{\epsilon'-\epsilon=\omega} \Pi(\epsilon) A_{\alpha} \Pi(\epsilon') \tag{A4}$$

where $\Pi(\epsilon)$ is a projector on subspace with energy ϵ , such that $H_0 = \sum_{\epsilon} \epsilon \Pi(\epsilon)$. These obey the following commutation relation:

$$[A_{\alpha}(\omega), H_0] = \omega A_{\alpha}(\omega). \tag{A5}$$

as well as the relations:

$$A_{\alpha}^{\dagger}(\omega) = A_{\alpha}(-\omega), \quad \sum_{\omega} A_{\alpha}(\omega) = A_{\alpha}.$$
 (A6)

From this follows also

$$A_{\alpha}(\omega)e^{cH_0} = e^{c\omega}e^{cH_0}A_{\alpha}(\omega) \tag{A7}$$

where c is the complex number, such that, in particular, the time-dependent operator is given by:

$$A_{\alpha}(t) = \sum_{\omega} e^{-i\omega t} A_{\alpha}(\omega).$$
(A8)

Finally, we consider the Bloch-Redfield master equation in the Schrödinger picture:

$$\frac{d}{dt}\rho(t) = \mathcal{L}_t[\rho(t)] = i[\rho(t), H_0 + \sum_{\omega,\omega'} \sum_{\alpha,\beta} \mathcal{S}_{\alpha\beta}(\omega,\omega',t) A^{\dagger}_{\alpha}(\omega) A_{\beta}(\omega')]$$
(A9)

$$+\sum_{\omega,\omega'}\sum_{\alpha,\beta}\gamma_{\alpha\beta}(\omega,\omega',t)\left(A_{\beta}(\omega')\rho(t)A_{\alpha}^{\dagger}(\omega)-\frac{1}{2}\{A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega'),\rho(t)\}\right)$$
(A10)

where ρ is the system density matrix and \mathcal{L}_t is the generator, with

$$\gamma_{\alpha\beta}(\omega,\omega',t) = \Gamma_{\alpha\beta}(\omega',t) + \Gamma^*_{\beta\alpha}(\omega,t), \tag{A11}$$

$$S_{\alpha\beta}(\omega,\omega',t) = \frac{1}{2i} \left[\Gamma_{\alpha\beta}(\omega',t) - \Gamma^*_{\beta\alpha}(\omega,t) \right], \qquad (A12)$$

$$\Gamma_{\alpha\beta}(\omega,t) = \int_0^t ds \ e^{i\omega s} \langle R_\alpha(s)R_\beta(0)\rangle_{\gamma_R}.$$
(A13)

In the interaction picture (with respect to the bare Hamiltonian H_0), the Bloch-Redfield equation takes the following form:

$$\frac{d}{dt}\tilde{\rho}(t) = \tilde{\mathcal{L}}_t[\tilde{\rho}(t)] = i[\tilde{\rho}(t), \sum_{\omega,\omega'} \sum_{\alpha,\beta} \tilde{\mathcal{S}}_{\alpha\beta}(\omega,\omega',t) A^{\dagger}_{\alpha}(\omega) A_{\beta}(\omega')]$$
(A14)

$$+\sum_{\omega,\omega'}\sum_{\alpha,\beta}\tilde{\gamma}_{\alpha\beta}(\omega,\omega',t)\left(A_{\beta}(\omega')\tilde{\rho}(t)A_{\alpha}^{\dagger}(\omega)-\frac{1}{2}\{A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega'),\tilde{\rho}(t)\}\right)$$
(A15)

where $\tilde{\rho}(t) = e^{iH_0t}\rho(t)e^{-iH_0t}$, and

$$\tilde{\gamma}_{\alpha\beta}(\omega,\omega',t) = e^{i(\omega-\omega')t}\gamma_{\alpha\beta}(\omega,\omega',t),\tag{A16}$$

$$\tilde{\mathcal{S}}_{\alpha\beta}(\omega,\omega',t) = e^{i(\omega-\omega')t} \mathcal{S}_{\alpha\beta}(\omega,\omega',t).$$
(A17)

Additionally, we also use the abbreviation

$$\mathcal{S}_{\alpha\beta}(\omega) = \lim_{t \to \infty} \mathcal{S}_{\alpha\beta}(\omega, \omega, t), \quad \gamma_{\alpha\beta}(\omega) = \lim_{t \to \infty} \gamma_{\alpha\beta}(\omega, \omega, t)$$
(A18)

From the definition it is seen that $\gamma_{\alpha\beta}(\omega)$ is the Fourier transform of the autocorrelation function, i.e.,

$$\gamma_{\alpha\beta}(\omega) = \Gamma_{\alpha\beta}(\omega) + \Gamma^*_{\beta\alpha}(\omega) \tag{A19}$$

$$= \int_0^\infty ds \ e^{i\omega s} \langle R_\alpha(s)R_\beta(0)\rangle_{\gamma_R} + \int_0^\infty ds \ e^{-i\omega s} \langle R_\beta(s)R_\alpha(0)\rangle_{\gamma_R}^* \tag{A20}$$

$$= \int_0^\infty ds \ e^{i\omega s} \langle R_\alpha(s) R_\beta(0) \rangle_{\gamma_R} + \int_0^\infty ds \ e^{-i\omega s} \langle R_\alpha(-s) R_\beta(0) \rangle_{\gamma_R}$$
(A21)

$$= \int_{-\infty}^{+\infty} ds \ e^{i\omega s} \langle R_{\alpha}(s) R_{\beta}(0) \rangle_{\gamma_R}$$
(A22)

from which it follows that $\gamma_{\alpha\beta}(\omega)$ obeys the detailed balance condition, i.e.,

$$\gamma_{\alpha\beta}(\omega) = \gamma_{\beta\alpha}(-\omega)e^{\beta\omega}.$$
(A23)

Using the inverse Fourier transform for the autocorrelation function and the Sokhostki-Plemelj identity in the form:

$$\int_0^\infty ds \ e^{i\omega s} = \frac{1}{\pi} \delta(\omega) + i\mathcal{P}\frac{1}{\omega},\tag{A24}$$

we shall represent $\Gamma_{\alpha\beta}(\omega)$ as the principal value integral:

$$\Gamma_{\alpha\beta}(\omega) = \int_0^\infty ds \ e^{i\omega s} \langle R_\alpha(s) R_\beta(0) \rangle_{\gamma_R} = \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \ \gamma_{\alpha\beta}(\Omega) \int_0^\infty ds \ e^{i(\omega - \Omega)s}$$
(A25)

$$=\frac{1}{2}\gamma_{\alpha\beta}(\omega) + \mathcal{P}\frac{1}{2\pi}\int_{-\infty}^{+\infty}d\Omega \;\frac{i\gamma_{\alpha\beta}(\Omega)}{\omega-\Omega} \tag{A26}$$

According to this, and since the $\gamma^*_{\alpha\beta}(\omega) = \gamma_{\beta\alpha}(\omega)$, we have

$$\mathcal{S}_{\alpha\beta}(\omega) = \frac{1}{2i} \left[\Gamma_{\alpha\beta}(\omega) - \Gamma^*_{\beta\alpha}(\omega) \right] = \mathcal{P} \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \, \frac{\gamma_{\alpha\beta}(\Omega)}{\omega - \Omega} \tag{A27}$$

Appendix B: Mean-force Hamiltonian

We search for the solution for the mean-force Hamiltonian $H_{\rm mf}$ from the equation:

$$e^{-\beta H_{\rm mf}} = \frac{1}{\mathcal{Z}_R} \operatorname{Tr}_R[e^{-\beta H}],\tag{B1}$$

where $H_{\rm mf} = H_0 + \delta H_{\rm mf}$ and $Z_R = \text{Tr}[e^{-\beta H_R}]$. Notice, that we used here a standard convention, i.e.,

$$\operatorname{Tr}\left[e^{-\beta H}\right] = \operatorname{Tr}\left[e^{-\beta H_{\mathrm{mf}}}\right] \operatorname{Tr}\left[e^{-\beta H_{R}}\right]$$
(B2)

that fixes the ground state energy of the mean-force Hamiltonian.

1. Dyson series

In the following section, we use an abbreviation $\hat{A}(t) = e^{t(H_0 + H_R)} A e^{-t(H_0 + H_R)}$. We start with the LHS of the Eq. (B1), which we represent by the formal Dyson form:

$$e^{-\beta H_{\rm mf}} = e^{-\beta H_0} e^{\beta H_0} e^{-\beta H_{\rm mf}} = e^{-\beta H_0} \mathcal{T} e^{-\int_0^\beta dt \ \delta \hat{H}_{\rm mf}(t)},\tag{B3}$$

which gives us the series expansion:

$$\mathcal{T}e^{-\int_0^\beta dt \ \delta \hat{H}_{\rm mf}(t)} = \mathbb{1} - \int_0^\beta dt_1 \ \delta \hat{H}_{\rm mf}(t_1) + \int_0^\beta dt_1 \int_0^{t_1} dt_2 \ \delta \hat{H}_{\rm mf}(t_1) \ \delta \hat{H}_{\rm mf}(t_2) + \dots \tag{B4}$$

Similarly, for the RHS, we have

$$e^{-\beta H} = e^{-\beta H_0} \mathcal{T} e^{-\lambda \int_0^\beta dt \ \hat{H}_I(t)},\tag{B5}$$

such that

$$\mathcal{T}e^{-\lambda\int_0^\beta dt\hat{H}_I(t)} = \mathbb{1} - \lambda\int_0^\beta dt_1 \ \hat{H}_I(t_1) + \lambda^2\int_0^\beta dt_1\int_0^{t_1} dt_2 \ \hat{H}_I(t_1) \ \hat{H}_I(t_2) + \dots$$
(B6)

Finally, one can write

$$\frac{1}{\mathcal{Z}_R} \operatorname{Tr}_R[e^{-\beta H}] = \frac{1}{\mathcal{Z}_R} \operatorname{Tr}_R[e^{-\beta H_0} \mathcal{T} e^{-\lambda \int_0^\beta dt \ \hat{H}_I(t)}] = e^{-\beta H_0} \operatorname{Tr}_R[\mathcal{T} e^{-\lambda \int_0^\beta dt \ \hat{H}_I(t)} \gamma_R]$$
(B7)

where $\gamma_R = \frac{e^{-\beta \hat{H}_R}}{Z_R}$ is the Gibbs state of the bath. In analogy, we have:

$$e^{-\beta H_{\rm mf}} = e^{-\beta H_0} \operatorname{Tr}_R[\mathcal{T}e^{-\int_0^\beta dt \ \delta \hat{H}_{\rm mf}(t)} \gamma_B] \tag{B8}$$

such that Eq. (B1) can be written as:

$$\operatorname{Tr}_{R}\left[\mathcal{T}\left(e^{-\int_{0}^{\beta}dt\ \delta\hat{H}_{\mathrm{mf}}(t)}-e^{-\lambda\int_{0}^{\beta}dt\ \hat{H}_{I}(t)}\right)\gamma_{R}\right]=0\tag{B9}$$

or in the series form as:

$$\sum_{k=1}^{\infty} (-1)^k \int_0^{\beta} dt_1 \int_0^{t_1} dt_2 \cdots \int_0^{t_{k-1}} dt_k \left(\delta \hat{H}_{\rm mf}(t_1) \ \delta \hat{H}_{\rm mf}(t_2) \dots \delta \hat{H}_{\rm mf}(t_k) - \lambda^k \left\langle \hat{H}_I(t_1) \ \hat{H}_I(t_2) \dots \hat{H}_I(t_k) \right\rangle_{\gamma_R} \right) = 0 \tag{B10}$$

where $\langle \cdot \rangle_{\gamma_R} = \text{Tr}_R[\cdot \gamma_R].$

2. Weak coupling

Now, let us assume that $\lambda \ll 1$, and we expand:

$$H_{\rm mf} = H_0 + \lambda H_{\rm mf}^{(1)} + \lambda^2 \hat{H}_{\rm mf}^{(2)} + \dots$$
(B11)

such that $\delta H_{\rm mf} = \lambda H_{\rm mf}^{(1)} + \lambda^2 H_{\rm mf}^{(2)} + \dots$ Then, we collect terms in the same order of λ appearing in Eq. (B10), i.e.,

$$\lambda : \int_{0}^{\beta} dt_{1} \hat{H}_{\rm mf}^{(1)}(t_{1}) = \int_{0}^{\beta} dt_{1} \left\langle \hat{H}_{I}(t_{1}) \right\rangle_{\gamma_{R}}$$
(B12)

$$\lambda^{2} : -\int_{0}^{\beta} dt_{1} \hat{H}_{mf}^{(2)}(t_{1}) + \int_{0}^{\beta} dt_{1} \int_{0}^{t_{1}} dt_{2} \hat{H}_{mf}^{(1)}(t_{1}) \hat{H}_{mf}^{(1)}(t_{2}) = \int_{0}^{\beta} dt_{1} \int_{0}^{t_{1}} dt_{2} \left\langle \hat{H}_{I}(t_{1}) \hat{H}_{I}(t_{2}) \right\rangle_{\gamma_{R}}$$
(B13)

$$\lambda^{3} : -\int_{0}^{\beta} dt_{1} \hat{H}_{mf}^{(3)}(t_{1}) + \int_{0}^{\beta} dt_{1} \int_{0}^{t_{1}} dt_{2} \left(\hat{H}_{mf}^{(1)}(t_{1}) \hat{H}_{mf}^{(2)}(t_{2}) + \hat{H}_{mf}^{(2)}(t_{1}) \hat{H}_{mf}^{(1)}(t_{1}) \right) \\ -\int_{0}^{\beta} dt_{1} \int_{0}^{t_{1}} dt_{2} \int_{0}^{t_{2}} dt_{3} \hat{H}_{mf}^{(1)}(t_{1}) \hat{H}_{mf}^{(1)}(t_{2}) \hat{H}_{mf}^{(1)}(t_{3}) = \int_{0}^{\beta} dt_{1} \int_{0}^{t_{1}} dt_{2} \int_{0}^{t_{2}} dt_{3} \left\langle \hat{H}_{I}(t_{1}) \hat{H}_{I}(t_{2}) \hat{H}_{I}(t_{3}) \right\rangle_{\gamma_{R}} (B14)$$

In general for the n-th order we have

$$\sum_{m=1}^{n} (-1)^{m} \sum_{k_{1},\dots,k_{m} \in \mathcal{C}_{n}^{m}} \int_{0}^{\beta} dt_{1} \cdots \int_{0}^{t_{m-1}} dt_{m} \ \hat{H}_{S}^{(k_{1})}(t_{1}) \ \hat{H}_{S}^{(k_{2})}(t_{2}) \dots \hat{H}_{S}^{(k_{m})}(t_{m})$$
$$= (-1)^{n} \int_{0}^{\beta} dt_{1} \cdots \int_{0}^{t_{n-1}} dt_{n} \left\langle \hat{H}_{I}(t_{1}) \dots \hat{H}_{I}(t_{n}) \right\rangle_{\gamma_{R}}$$

where \mathcal{C}_n^k is set of the k-th order composition of the number n, e.g., $\mathcal{C}_4^3 = \{(2,1,1), (1,2,1), (1,1,2)\}$.

3. Derivation of general formulas for corrections to the mean-force Hamiltonian

a. First-order correction

Let us first solve the equation for the first-order correction, i.e.,

$$\int_{0}^{\beta} dt_{1} \hat{H}_{\rm mf}^{(1)}(t_{1}) = \int_{0}^{\beta} dt_{1} \left\langle \hat{H}_{I}(t_{1}) \right\rangle_{\gamma_{R}} = \sum_{\alpha} \int_{0}^{\beta} dt_{1} \hat{A}_{\alpha}(t_{1}) \left\langle \hat{R}_{\alpha}(t_{1}) \right\rangle_{\gamma_{R}}.$$
 (B15)

Furthermore, since $\langle \hat{R}_{\alpha}(t_1) \rangle_{\gamma_R} = \langle R_{\alpha} \rangle_{\gamma_R}$ (due to commutation of the Gibbs state γ_R with the free Hamiltonian H_R), we get the solution:

$$H_{\rm mf}^{(1)} = \sum_{\alpha} \langle R_{\alpha} \rangle_{\gamma_R} \ A_{\alpha}. \tag{B16}$$

From now on, we assume that bath operators are centralized such that $\langle R_{\alpha} \rangle_{\gamma_R} = 0$, which implies $H_{\rm mf}^{(1)} = 0$.

b. Second-order correction

In this section we provide general formula for second order correction for mean-force Hamiltonian. Remarkably the expression does not exhibit any poles, as is usual for dynamical correction. Yet we also decompose it into bricks that are used also to build the dynamical corrections, which do exhibit poles, and require principal value to be well defined.

Theorem 5. The explicit form of second order correction for mean-force Hamiltonian is the following:

$$H_{mf}^{(2)} = \sum_{\omega,\omega'} \Upsilon_{\alpha\beta}^{(mf)}(\omega,\omega') A_{\alpha}^{\dagger}(\omega) A_{\beta}(\omega'), \tag{B17}$$

where

$$\Upsilon_{\alpha\beta}^{(mf)}(\omega,\omega') = \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \ D(\omega,\omega',\Omega) \ \gamma_{\alpha\beta}(\Omega), \quad D(\omega,\omega',\Omega) = \frac{1}{\omega'-\Omega} - \frac{(\omega-\omega')(e^{\beta(\omega-\Omega)}-1)}{(\omega-\Omega)(\omega'-\Omega)(e^{\beta(\omega-\omega')}-1)}$$
(B18)

The coefficients $\Upsilon_{\alpha\beta}(\omega,\omega')$ can be also expressed in terms of the imaginary part of $\Gamma_{\alpha\beta}(\omega)$ (see Eq. (A27)) as follows

$$\Upsilon^{(mf)}_{\alpha\beta}(\omega,\omega') = \frac{1}{e^{\beta\omega} - e^{\beta\omega'}} \left(e^{\beta\omega} \mathcal{S}_{\alpha\beta}(\omega') - e^{\beta\omega'} \mathcal{S}_{\alpha\beta}(\omega) + e^{\beta(\omega+\omega')} \left(\mathcal{S}_{\beta\alpha}(-\omega') - \mathcal{S}_{\beta\alpha}(-\omega) \right) \right).$$
(B19)

Remark. From (B19) one sees that $\Upsilon_{\alpha\beta}^{(mf)}$ is symmetric, i.e., $\Upsilon_{\alpha\beta}^{(mf)}(\omega, \omega') = \Upsilon_{\alpha\beta}^{(mf)}(\omega', \omega)$. This can be also seen by writing $D(\omega, \omega', \Omega)$ in explicitly symmetric form

$$D(\omega, \omega', \Omega) = \frac{1}{2} \frac{(e^{\beta\omega} - e^{\beta\omega'})(\omega + \omega' - 2\Omega) + (\omega - \omega')(e^{\beta\omega} + e^{\beta\omega'} - 2e^{\beta(\omega + \omega' - \Omega)})}{(e^{\beta\omega} - e^{\beta\omega'})(\omega - \Omega)(\omega' - \Omega)}$$
(B20)

Proof. We shall first prove that Eq. (B19) comes from (B18). We shall use Eq. (A27), i.e.,

$$S_{\alpha\beta}(\omega) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \frac{\gamma_{\alpha\beta}(\Omega)}{\omega - \Omega},$$
(B21)

from which we also derive

$$-\mathcal{S}_{\beta\alpha}(-\omega) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \frac{\gamma_{\beta\alpha}(\Omega)}{\omega + \Omega} = \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \frac{\gamma_{\beta\alpha}(-\Omega)}{\omega - \Omega} = \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \frac{\gamma_{\alpha\beta}(\Omega)e^{-\beta\Omega}}{\omega - \Omega},$$
(B22)

where we used the detailed balance condition (A23). Thus, to express $\Upsilon_{\alpha\beta}^{(mf)}$ in terms of $S_{\alpha\beta}$ we have to write D in terms of $1/(\omega - \Omega)$ or $1/(\omega' - \Omega)$. Using

$$\frac{\omega - \omega'}{(\omega' - \Omega)(\omega - \Omega)} = \frac{1}{\omega' - \Omega} - \frac{1}{\omega - \Omega}$$
(B23)

we thus get

$$D(\omega, \omega', \Omega) = \frac{1}{e^{\beta(\omega-\omega')} - 1} \left(\frac{e^{\beta(\omega-\omega')} - 1}{\omega' - \Omega} - e^{\beta(\omega-\Omega)} \left(\frac{1}{\omega' - \Omega} - \frac{1}{\omega - \Omega} \right) + \frac{1}{\omega' - \Omega} - \frac{1}{\omega - \Omega} \right)$$
(B24)

$$=\frac{1}{e^{\beta\omega}-e^{\beta\omega'}}\left(\frac{e^{\beta\omega}}{\omega'-\Omega}-\frac{e^{\beta\omega'}}{\omega-\Omega}-e^{\beta(\omega+\omega'-\Omega)}(\frac{1}{\omega'-\Omega}-\frac{1}{\omega-\Omega})\right)$$
(B25)

Rearranging it a bit, and using (B21) and (B22) we obtain (B19).

Let us now prove the expression (B18). We start with second-order equation with centralized bath operators, i.e.,

$$\int_{0}^{\beta} dt \hat{H}_{\rm mf}^{(2)}(t) = -\int_{0}^{\beta} dt \int_{0}^{t} ds \left\langle \hat{H}_{I}(t) \hat{H}_{I}(s) \right\rangle_{\gamma_{R}} \tag{B26}$$

Next, we put the representation (B17) and according to the relation (A7), we have

$$\hat{H}_{\rm mf}^{(2)}(t) = \sum_{\omega,\omega'} \Upsilon_{\alpha\beta}^{\rm (mf)}(\omega,\omega') e^{tH_0} A_{\alpha}^{\dagger}(\omega) A_{\beta}(\omega') e^{-tH_0} = \sum_{\omega,\omega'} \Upsilon_{\alpha\beta}^{\rm (mf)}(\omega,\omega') e^{t(\omega-\omega')} A_{\alpha}^{\dagger}(\omega) A_{\beta}(\omega'). \tag{B27}$$

Then, the LHS of Eq. (B26) is equal to:

$$\int_{0}^{\beta} dt \ \hat{H}_{\rm mf}^{(2)}(t) = \sum_{\omega,\omega'} A_{\alpha}^{\dagger}(\omega) A_{\beta}(\omega') \left(\Upsilon_{\alpha\beta}^{\rm (mf)}(\omega,\omega') \int_{0}^{\beta} dt \ e^{t(\omega-\omega')} \right)$$
(B28)

whereas the RHS is given by:

$$-\int_{0}^{\beta} dt \int_{0}^{t} ds \left\langle \hat{H}_{I}(t) \hat{H}_{I}(s) \right\rangle_{\gamma_{R}} = -\int_{0}^{\beta} dt \int_{0}^{t} ds \ \hat{A}_{\alpha}(t) \hat{A}_{\beta}(s) \left\langle \hat{R}_{\alpha}(t) \ \hat{R}_{\beta}(s) \right\rangle$$

$$= -\sum_{\omega,\omega'} \int_{0}^{\beta} dt \int_{0}^{t} ds \ e^{t\omega - s\omega'} A_{\alpha}^{\dagger}(\omega) A_{\beta}(\omega') \left\langle \hat{R}_{\alpha}(t-s) \ \hat{R}_{\beta} \right\rangle$$

$$= -\sum_{\omega,\omega'} A_{\alpha}^{\dagger}(\omega) A_{\beta}(\omega') \int_{0}^{\beta} dt \ e^{t(\omega-\omega')} \int_{0}^{t} ds \ e^{s\omega'} \left\langle \hat{R}_{\alpha}(s) \hat{R}_{\beta} \right\rangle$$
(B29)
(B2

where in the last line we change a variables $s \to t - s$. Next, according to Eq. (A19), let us observe that

$$\langle \hat{R}_{\alpha}(it)\hat{R}_{\beta}\rangle = \langle R_{\alpha}(t)R_{\beta}\rangle = \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \ e^{-i\Omega t} \ \gamma_{\alpha\beta}(\Omega), \tag{B31}$$

such that

$$\langle \hat{R}_{\alpha}(t)\hat{R}_{\beta}\rangle = \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \ e^{-\Omega t} \ \gamma_{\alpha\beta}(\Omega). \tag{B32}$$

Finally, the RHS is equal to:

$$-\int_{0}^{\beta} dt \int_{0}^{t} ds \left\langle \hat{H}_{I}(t) \hat{H}_{I}(s) \right\rangle_{\gamma_{R}} = -\sum_{\omega,\omega'} A_{\alpha}^{\dagger}(\omega) A_{\beta}(\omega') \left(\frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \,\gamma_{\alpha\beta}(\Omega) \int_{0}^{\beta} dt \, e^{t(\omega-\omega')} \int_{0}^{t} ds \, e^{s(\omega'-\Omega)} \right)$$
(B33)

Equating LHS=RHS, we get

$$\sum_{\omega,\omega'} \left[\Upsilon_{\alpha\beta}^{(\mathrm{mf})}(\omega,\omega') \int_0^\beta dt \ e^{t(\omega-\omega')} + \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \ \gamma_{\alpha\beta}(\Omega) \int_0^\beta dt \ e^{t(\omega-\omega')} \int_0^t ds \ e^{s(\omega'-\Omega)} \right] A_{\alpha}^{\dagger}(\omega) A_{\beta}(\omega') = 0$$
(B34)

which is solved by

$$\Upsilon_{\alpha\beta}^{(\mathrm{mf})}(\omega,\omega') = -\frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \ \gamma_{\alpha\beta}(\Omega) \ \frac{\int_0^\beta dt \ e^{t(\omega-\omega')} \int_0^t ds \ e^{s(\omega'-\Omega)}}{\int_0^\beta dt \ e^{t(\omega-\omega')}} \tag{B35}$$

We thus obtain

$$D(\omega, \omega', \Omega) = -\frac{\int_0^\beta dt \int_0^t ds \ e^{t(\omega-\omega')} e^{s(\omega'-\Omega)}}{\int_0^\beta dt \ e^{t(\omega-\omega')}}$$
(B36)

Appendix C: Steady-state correction

1. General method

We look for a solution of the equation:

$$\mathcal{L}[\varrho] = 0, \tag{C1}$$

where \mathcal{L} is the generator of the master equation and ϱ is its stationary state. We expand the generator and steady-state in the series, i.e.,

$$\mathcal{L}[\rho] = \mathcal{L}_0[\rho] + \lambda^2 \mathcal{L}_2[\rho] + \lambda^4 \mathcal{L}_4[\rho] + \dots$$
(C2)

$$\varrho = \varrho_0 + \lambda^2 \varrho_2 + \lambda^4 \varrho_4 + \dots \tag{C3}$$

such that we have the following set of equations (for each order in λ):

$$\mathcal{L}_0[\varrho_0] = 0 \tag{C4}$$

$$\mathcal{L}_0[\varrho_2] + \mathcal{L}_2[\varrho_0] = 0 \tag{C5}$$

$$\mathcal{L}_0[\varrho_4] + \mathcal{L}_2[\varrho_2] + \mathcal{L}_4[\varrho_0] = 0 \tag{C6}$$

We postulate the stationary state (in the Gibbs form):

$$\varrho = e^{-\beta(H_0 + \lambda^2 H_{\rm st}^{(2)} + \lambda^4 H_{\rm st}^{(4)}} + \dots) = \varrho_0 + \lambda^2 \varrho_2 + \lambda^2 \varrho_4 + \dots$$
(C8)

such that

$$\varrho_0 = e^{-\beta H_0} \tag{C9}$$

$$\varrho_2 = -e^{-\beta H_0} \int_0^{\cdot} dt \ e^{tH_0} H_{\rm st}^{(2)} e^{-tH_0} \tag{C10}$$

$$\varrho_4 = -e^{-\beta H_0} \int_0^\beta dt \ e^{tH_0} H_{\rm st}^{(4)} e^{-tH_0} + e^{-\beta H_0} \int_0^\beta dt_1 \int_0^{t_1} dt_2 \ e^{t_1 H_0} H_{\rm st}^{(2)} e^{-t_1 H_0} e^{t_2 H_0} H_{\rm st}^{(2)} e^{-t_2 H_0} \tag{C11}$$

In the following, we use the summation convention, i.e., the repeating indices are summed up. We start with representation of the second-order correction in the basis of jump operators:

$$H_{\rm st}^{(2)} = \Upsilon_{\alpha\beta}^{\rm (st)}(\omega,\omega')A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega').$$
(C12)

$$\varrho_2 = -e^{-\beta H_0} \Upsilon_{\alpha\beta}^{(\mathrm{st})}(\omega,\omega') \int_0^\beta dt \ e^{tH_0} A_\alpha^\dagger(\omega) A_\beta(\omega') e^{-tH_0} = -\Upsilon_{\alpha\beta}^{(\mathrm{st})}(\omega,\omega') \alpha(\omega'-\omega) e^{-\beta H_0} A_\alpha^\dagger(\omega) A_\beta(\omega'), \quad (C13)$$

where we define:

$$\alpha(\omega) = \int_0^\beta dt \ e^{-t\omega} = \begin{cases} \frac{1 - e^{-\beta\omega}}{\omega}, & \omega \neq 0\\ \beta, & \omega = 0 \end{cases}$$
(C14)

In general, we are going to transform the operator equations (C5) and (C6) into the algebraic ones. For this we define:

$$\mathcal{L}_k[\varrho_l] = g_{\alpha\beta}^{(kl)}(\omega_1, \omega_2) e^{-\beta H_0} A_\alpha(\omega_1) A_\beta(\omega_2)$$
(C15)

for the second-order (such that k + l = 2), and

$$\mathcal{L}_{k}[\varrho_{l}] = g_{\alpha\beta\gamma\delta}^{(kl)}(\omega_{1},\omega_{2},\omega_{3},\omega_{4})e^{-\beta H_{0}}A_{\alpha}(\omega_{1})A_{\beta}(\omega_{2})A_{\gamma}(\omega_{3})A_{\delta}(\omega_{4})$$
(C16)

for k + l = 4. In accordance, for the second-order equation (C5), we have

$$\left(g_{\alpha\beta}^{(02)}(\omega_1,\omega_2) + g_{\alpha\beta}^{(20)}(\omega_1,\omega_2)\right)e^{-\beta H_0}A_{\alpha}(\omega_1)A_{\beta}(\omega_2) = 0$$
(C17)

whereas for the fourth-order:

$$\left(g^{(04)}_{\alpha\beta\gamma\delta}(\omega_1,\omega_2,\omega_3,\omega_4) + g^{(22)}_{\alpha\beta\gamma\delta}(\omega_1,\omega_2,\omega_3,\omega_4) + g^{(40)}_{\alpha\beta\gamma\delta}(\omega_1,\omega_2,\omega_3,\omega_4)\right)e^{-\beta H_0}A_{\alpha}(\omega_1)A_{\beta}(\omega_2)A_{\gamma}(\omega_3)A_{\delta}(\omega_4) = 0$$
(C18)

In the following, we will also use the commutation relations:

$$[A_{\alpha}(\omega), H_0] = \omega A_{\alpha}(\omega) \tag{C19}$$

from which we get:

$$A_{\alpha}(\omega)e^{-\beta H_0} = e^{-\beta\omega}e^{-\beta H_0}A_{\alpha}(\omega).$$
(C20)

The commutation relation (C19) can be further generalize for the product of jump operators, i.e.,

$$[A_{\alpha_1}(\omega_1)A_{\alpha_2}(\omega_2)\dots A_{\alpha_2}(\omega_2), H_0] = (\omega_1 + \omega_2 + \dots + \omega_n)A_{\alpha_1}(\omega_1)A_{\alpha_2}(\omega_2)\dots A_{\alpha_2}(\omega_2).$$
(C21)

Notice also that $A^{\dagger}_{\alpha}(\omega) = A_{\alpha}(-\omega)$.

2. Second-order equation

In the following, we solve Eq. (C17) for a master equation of the form:

$$\mathcal{L}_{0}[\rho] = i[\rho, H_{0}]$$

$$\mathcal{L}_{2}[\rho] = \sum_{\alpha, \beta} \sum_{\omega, \omega'} \left[\Upsilon^{(\text{dyn})}_{\alpha\beta}(\omega, \omega')[\rho, A^{\dagger}_{\alpha}(\omega)A_{\beta}(\omega')] + K_{\alpha\beta}(\omega, \omega') \left(A_{\beta}(\omega')\rho A^{\dagger}_{\alpha}(\omega) - \frac{1}{2} \{ A^{\dagger}_{\alpha}(\omega)A_{\beta}(\omega'), \rho \} \right) \right]$$
(C22)
(C23)

We observe that the zeroth-order equation, i.e., $[\varrho_0, H_0] = 0$ is obviously satisfied for a choice $\varrho_0 = e^{-\beta H_0}$. Let us then calculate the coefficients $g_{\alpha\beta}^{(02)}$ and $g_{\alpha\beta}^{(20)}$ for the second-order equation. We start with:

$$\mathcal{L}_0[\varrho_2] = i[\varrho_2, H_0] = -i\Upsilon^{(\text{st})}_{\alpha\beta}(\omega_1, \omega_2)\alpha(\omega_1 + \omega_2)[e^{-\beta H_0}A^{\dagger}_{\alpha}(\omega)A_{\beta}(\omega_2), H_0]$$
(C24)

$$= -i(\omega_1 + \omega_2)\Upsilon^{(\text{st})}_{\alpha\beta}(-\omega_1, \omega_2)\alpha(\omega_1 + \omega_2)e^{-\beta H_0}A_\alpha(\omega_1)A_\beta(\omega_2)$$
(C25)

where we used Eq. (C21), such that

$$g_{\alpha\beta}^{(02)} = -i(\omega_1 + \omega_2)\Upsilon_{\alpha\beta}^{(\text{st})}(-\omega_1, \omega_2)\alpha(\omega_1 + \omega_2)$$
(C26)

Next, we shall calculate:

$$\mathcal{L}_{2}[\varrho_{0}] = i\Upsilon_{\alpha\beta}^{(\mathrm{dyn})}(\omega_{1},\omega_{2})[\varrho_{0},A_{\alpha}^{\dagger}(\omega_{1})A_{\beta}(\omega_{2})] + K_{\alpha\beta}(\omega_{1},\omega_{2})\left(A_{\beta}(\omega_{2})\varrho_{0}A_{\alpha}^{\dagger}(\omega_{1}) - \frac{1}{2}\{A_{\alpha}^{\dagger}(\omega_{1})A_{\beta}(\omega_{2}),\varrho_{0}\}\right)$$
(C27)

$$=i\Upsilon^{(\mathrm{dyn})}_{\alpha\beta}(\omega_{1},\omega_{2})[e^{-\beta H_{0}},A^{\dagger}_{\alpha}(\omega_{1})A_{\beta}(\omega_{2})]+K_{\alpha\beta}(\omega_{1},\omega_{2})\left(A_{\beta}(\omega_{2})e^{-\beta H_{0}}A^{\dagger}_{\alpha}(\omega_{1})-\frac{1}{2}\{A^{\dagger}_{\alpha}(\omega_{1})A_{\beta}(\omega_{2}),e^{-\beta H_{0}}\}\right)$$

First, let us rewrite the Hamiltonian part in the form:

$$i\Upsilon_{\alpha\beta}^{(\mathrm{dyn})}(\omega_{1},\omega_{2})[e^{-\beta H_{0}},A_{\alpha}^{\dagger}(\omega_{1})A_{\beta}(\omega_{2})] = i\Upsilon_{\alpha\beta}^{(\mathrm{dyn})}(\omega_{1},\omega_{2})(e^{-\beta H_{0}}A_{\alpha}^{\dagger}(\omega_{1})A_{\beta}(\omega_{2}) - A_{\alpha}^{\dagger}(\omega_{1})A_{\beta}(\omega_{2})e^{-\beta H_{0}}) \quad (C28)$$
$$= i\Upsilon_{\alpha\beta}^{(\mathrm{dyn})}(\omega_{1},\omega_{2})(1 - e^{-\beta(\omega_{2}-\omega_{1})})e^{-\beta H_{0}}A_{\alpha}^{\dagger}(\omega_{1})A_{\beta}(\omega_{2}) \quad (C29)$$

and then the dissipative part as follows

$$K_{\alpha\beta}(\omega_1,\omega_2)\left(A_{\beta}(\omega_2)e^{-\beta H_0}A_{\alpha}^{\dagger}(\omega_1) - \frac{1}{2}\{A_{\alpha}^{\dagger}(\omega_1)A_{\beta}(\omega_2), e^{-\beta H_0}\}\right)$$
(C30)

$$=K_{\alpha\beta}(\omega_1\omega_2)\left(A_{\beta}(\omega_2)e^{-\beta H_0}A^{\dagger}_{\alpha}(\omega_1)-\frac{1}{2}A^{\dagger}_{\alpha}(\omega_1)A_{\beta}(\omega_2)e^{-\beta H_0}-\frac{1}{2}e^{-\beta H_0}A^{\dagger}_{\alpha}(\omega_1)A_{\beta}(\omega_2)\right)\right)$$
(C31)

$$=K_{\alpha\beta}(\omega_{1}\omega_{2})\left(e^{-\beta\omega}e^{-\beta H_{0}}A_{\beta}(\omega_{2})A_{\alpha}^{\dagger}(\omega_{1})-\frac{1}{2}e^{-\beta(\omega-\omega_{2})}e^{-\beta H_{0}}A_{\alpha}^{\dagger}(\omega_{1})A_{\beta}(\omega_{2})-\frac{1}{2}e^{-\beta H_{0}}A_{\alpha}^{\dagger}(\omega_{1})A_{\beta}(\omega_{2})\right)$$
(C32)

$$= \left(e^{\beta\omega_1}K_{\beta\alpha}(-\omega_2, -\omega_1) - \frac{1}{2}K_{\alpha\beta}(\omega_1, \omega_2)(e^{-\beta(\omega_2 - \omega_1)} + 1)\right)e^{-\beta H_0}A^{\dagger}_{\alpha}(\omega_1)A_{\beta}(\omega_2)$$
(C33)

Finally, we get

$$g_{\alpha\beta}^{(20)}(\omega_1,\omega_2) = i\Upsilon_{\alpha\beta}^{(\text{dyn})}(-\omega_1,\omega_2)(1-e^{-\beta(\omega_1+\omega_2)}) + e^{\beta\omega_1}K_{\beta\alpha}(-\omega_2,-\omega_1) - \frac{1}{2}K_{\alpha\beta}(-\omega_1,\omega_2)(e^{-\beta(\omega_1+\omega_2)}+1)(\text{C34})$$

Now, we postulate the solution

$$g_{\alpha\beta}^{(20)}(\omega_1,\omega_2) + g_{\alpha\beta}^{(20)}(\omega_1,\omega_2) = 0$$
(C35)

for each ω_1, ω_2 and α, β . First, for $\omega_1 = \omega_2 \equiv \omega$, we have

$$e^{\beta\omega}K_{\beta\alpha}(-\omega,-\omega) - K_{\alpha\beta}(\omega,\omega) = 0$$
(C36)

such that the coefficient $K_{\alpha\beta}(\omega,\omega)$ has to satisfy the detailed balance condition. Furthermore, for $\omega_1 \neq \omega_2$ we get

$$i\Upsilon_{\alpha\beta}^{(\rm st)}(-\omega_{1},\omega_{2})(e^{-\beta(\omega_{1}+\omega_{2})}-1) - i\Upsilon_{\alpha\beta}^{(\rm dyn)}(-\omega_{1},\omega_{2})(e^{-\beta(\omega_{1}+\omega_{2})}-1) + e^{-\beta\omega_{1}}K_{\beta\alpha}(-\omega_{2},\omega_{1}) - \frac{1}{2}K_{\alpha\beta}(-\omega_{1},\omega_{2})(e^{\beta(\omega_{1}+\omega_{2})}+1) = 0 \quad (C37)$$

This can be further simplified to:

$$\Upsilon_{\alpha\beta}^{(\mathrm{st})}(\omega_1,\omega_2) = \Upsilon_{\alpha\beta}^{(\mathrm{dyn})}(\omega_1,\omega_2) + \frac{i}{e^{\beta\omega_1} - e^{\beta\omega_2}} \left(e^{\beta(\omega_1 + \omega_2)} K_{\beta\alpha}(-\omega_2,-\omega_1) - \frac{1}{2} K_{\alpha\beta}(\omega_1,\omega_2)(e^{\beta\omega_1} + e^{\beta\omega_2}) \right).$$
(C38)

a. Solutions for the Bloch-Redfield master equation and for secular approximation

For the Bloch-Redfield master equation, we have:

$$\Upsilon_{\alpha\beta}^{(\mathrm{dyn})}(\omega,\omega') = \frac{1}{2i} (\Gamma_{\alpha\beta}(\omega') - \Gamma_{\beta\alpha}^{*}(\omega)), \qquad (C39)$$

$$K_{\alpha\beta}(\omega,\omega') = \Gamma_{\alpha\beta}(\omega') + \Gamma^*_{\beta\alpha}(\omega)$$
(C40)

Next, we put:

$$\Gamma_{\alpha\beta}(\omega) = \frac{1}{2}\gamma_{\alpha\beta}(\omega) + i\mathcal{S}_{\alpha\beta}(\omega) \tag{C41}$$

such that

$$\Upsilon^{(\mathrm{dyn})}_{\alpha\beta}(\omega,\omega') = \frac{1}{2i} \left(\frac{1}{2} \gamma_{\alpha\beta}(\omega') + i\mathcal{S}_{\alpha\beta}(\omega') - \frac{1}{2} \gamma_{\alpha\beta}(\omega) + i\mathcal{S}_{\alpha\beta}(\omega)\right) \tag{C42}$$

$$=\frac{1}{4i}(\gamma_{\alpha\beta}(\omega')-\gamma_{\alpha\beta}(\omega))+\frac{1}{2}(\mathcal{S}_{\alpha\beta}(\omega')+\mathcal{S}_{\alpha\beta}(\omega))$$
(C43)

and

$$K_{\alpha\beta}(\omega,\omega') = \frac{1}{2}\gamma_{\alpha\beta}(\omega') + i\mathcal{S}_{\alpha\beta}(\omega') + \frac{1}{2}\gamma_{\alpha\beta}(\omega) - i\mathcal{S}_{\alpha\beta}(\omega)$$
(C44)

$$=\frac{1}{2}(\gamma_{\alpha\beta}(\omega')+\gamma_{\alpha\beta}(\omega))+i(\mathcal{S}_{\alpha\beta}(\omega')-\mathcal{S}_{\alpha\beta}(\omega))$$
(C45)

Let us put above expression into Eq. (C38) and collect all of the terms with $S_{\alpha\beta}$:

$$\frac{1}{2}(\mathcal{S}_{\alpha\beta}(\omega') + \mathcal{S}_{\alpha\beta}(\omega)) - \frac{1}{e^{\beta\omega} - e^{\beta\omega'}} \left(e^{\beta(\omega+\omega')}(\mathcal{S}_{\beta\alpha}(-\omega) - \mathcal{S}_{\beta\alpha}(-\omega')) - \frac{1}{2}(\mathcal{S}_{\alpha\beta}(\omega') - \mathcal{S}_{\alpha\beta}(\omega))(e^{\beta\omega} + e^{\beta\omega'}) \right)$$
(C46)
$$\frac{1}{e^{\beta\omega} - e^{\beta\omega'}} \left(e^{\beta(\omega+\omega')}(\mathcal{S}_{\beta\alpha}(-\omega) - \mathcal{S}_{\beta\alpha}(-\omega')) - \frac{1}{2}(\mathcal{S}_{\alpha\beta}(\omega') - \mathcal{S}_{\alpha\beta}(\omega))(e^{\beta\omega} + e^{\beta\omega'}) \right)$$
(C46)

$$=\frac{1}{e^{\beta\omega}-e^{\beta\omega'}}\left[\frac{e^{\beta\omega}-e^{\beta\omega'}}{2}\left(\mathcal{S}_{\alpha\beta}(\omega')+\mathcal{S}_{\alpha\beta}(\omega)\right)-\frac{e^{\beta\omega}+e^{\beta\omega'}}{2}\left(\mathcal{S}_{\alpha\beta}(\omega)-\mathcal{S}_{\alpha\beta}(\omega')\right)-e^{\beta(\omega+\omega')}\left(\mathcal{S}_{\beta\alpha}(-\omega)-\mathcal{S}_{\beta\alpha}(-\omega')\right)\right]$$
(C47)

$$=\frac{1}{e^{\beta\omega}-e^{\beta\omega'}}\left[e^{\beta\omega}\mathcal{S}_{\alpha\beta}(\omega')-e^{\beta\omega'}\mathcal{S}_{\alpha\beta}(\omega)+e^{\beta(\omega+\omega')}(\mathcal{S}_{\beta\alpha}(-\omega')-\mathcal{S}_{\beta\alpha}(-\omega))\right]$$
(C48)

Next, we collect all of the terms with $\gamma_{\alpha\beta}$, i.e.,

$$\frac{1}{4i}(\gamma_{\alpha\beta}(\omega') - \gamma_{\alpha\beta}(\omega)) + \frac{i}{e^{\beta\omega} - e^{\beta\omega'}} \left(e^{\beta(\omega + \omega')} \frac{1}{2} (\gamma_{\alpha\beta}(-\omega) + \gamma_{\alpha\beta}(-\omega')) - \frac{1}{4} (\gamma_{\alpha\beta}(\omega') + \gamma_{\alpha\beta}(\omega))(e^{\beta\omega} + e^{\beta\omega'}) \right)$$
(C49)

$$=\frac{i}{4}\left(-\gamma_{\alpha\beta}(\omega')+\gamma_{\alpha\beta}(\omega)+\frac{1}{e^{\beta\omega}-e^{\beta\omega'}}\left(2(e^{\beta\omega'}\gamma_{\alpha\beta}(\omega)+e^{\beta\omega}\gamma_{\alpha\beta}(\omega'))-(\gamma_{\alpha\beta}(\omega')+\gamma_{\alpha\beta}(\omega))(e^{\beta\omega}+e^{\beta\omega'})\right)\right) \quad (C50)$$

$$=\frac{i}{4}\left(-\gamma_{\alpha\beta}(\omega')+\gamma_{\alpha\beta}(\omega)+\frac{1}{e^{\beta\omega}-e^{\beta\omega'}}\left(2e^{\beta\omega'}\gamma_{\alpha\beta}(\omega)+2e^{\beta\omega}\gamma_{\alpha\beta}(\omega')-\gamma_{\alpha\beta}(\omega')(e^{\beta\omega}+e^{\beta\omega'})-\gamma_{\alpha\beta}(\omega)(e^{\beta\omega}+e^{\beta\omega'})\right)\right)$$
(C51)

$$=\frac{i}{4}\left(-\gamma_{\alpha\beta}(\omega')+\gamma_{\alpha\beta}(\omega)+\frac{1}{e^{\beta\omega}-e^{\beta\omega'}}\left(e^{\beta\omega'}\gamma_{\alpha\beta}(\omega)+e^{\beta\omega}\gamma_{\alpha\beta}(\omega')-e^{\beta\omega'}\gamma_{\alpha\beta}(\omega')-e^{\beta\omega}\gamma_{\alpha\beta}(\omega)\right)\right)=0.$$
 (C52)

One sees that only terms $S_{\alpha\beta}$ survives. Moreover, these are exactly equal to the expression for a mean-force Hamiltonian given by Eq. (B19), such that for the Bloch-Redfield or cumulant master equation we have simply:

$$\Upsilon^{(\text{st})}_{\alpha\beta}(\omega,\omega') = \Upsilon^{(\text{mf})}_{\alpha\beta}(\omega,\omega'), \tag{C53}$$

for $\omega \neq \omega'$.

Let us observe that if we apply the so-called secular approximation for $\gamma_{\alpha\beta}$ coefficients, i.e.

$$\gamma_{\alpha\beta}(\omega,\omega') \xrightarrow{\text{sec. approx}} \gamma_{\alpha\beta}(\omega)\delta_{\omega,\omega'}$$
 (C54)

then

$$e^{\beta(\omega+\omega')}\gamma_{\beta\alpha}(-\omega',-\omega) - \frac{1}{2}\gamma_{\alpha\beta}(\omega,\omega')(e^{\beta\omega} + e^{\beta\omega'} = \delta_{\omega,\omega'}\left(e^{2\beta\omega}\gamma_{\beta\alpha}(-\omega) - e^{\beta\omega}\gamma_{\alpha\beta}(\omega)\right) = 0$$
(C55)

due to the detailed balance condition. Finally, for such master equation, for $\omega \neq \omega'$, we have

$$\Upsilon_{\alpha\beta}^{(\rm st)}(\omega,\omega') = \Upsilon_{\alpha\beta}^{(\rm dyn)}(\omega,\omega'). \tag{C56}$$

3. Fourth-order equation

Now, we are going to solve the fourth-order equation (C18). For simplicity, we assume that the interaction term is given by $H_I = S \otimes R$, such that we drop the indices, i.e.,

$$(g_{04}(\omega_1,\omega_2,\omega_3,\omega_4) + g_{22}(\omega_1,\omega_2,\omega_3,\omega_4) + g_{40}(\omega_1,\omega_2,\omega_3,\omega_4)) e^{-\beta H_0} A(\omega_1) A(\omega_2) A(\omega_3) A(\omega_4) = 0,$$
(C57)

where $g_{\alpha\beta\gamma\delta}^{(kl)} \equiv g_{kl}$. According to the Proposition 1 in the Section V, the above equation is satisfied if and only if the following set of equations is satisfied:

$$\sum_{(\omega_1,\omega_2,\omega_2,\omega_4)\in G(|k\rangle \to |k\rangle)} \left(g_{22}(\omega_1,\omega_2,\omega_3,\omega_4) + g_{40}(\omega_1,\omega_2,\omega_3,\omega_4) \right) = 0.$$
(C58)

where $G(|k\rangle \rightarrow |k\rangle)$ denotes the set of all four-tuples

$$(\omega_1, \omega_2, \omega_2, \omega_4) = (\epsilon_l - \epsilon_k, \epsilon_m - \epsilon_l, \epsilon_j - \epsilon_m, \epsilon_k - \epsilon_j).$$
(C59)

a. g_{22} function

We consider the term:

$$\mathcal{L}_2[\varrho_2] = \sum_{\omega_1, \omega_2, \omega_3, \omega_4} g_{22}(\omega_1, \omega_2, \omega_3, \omega_4) A(\omega_1) A(\omega_2) A(\omega_3) A(\omega_4)$$
(C60)

where

$$\varrho_2 = \alpha(\omega_3 + \omega_4) \Upsilon_{\rm st}(-\omega_3, \omega_4) \varrho_0 A(\omega_3) A(\omega_4) \tag{C61}$$

$$\mathcal{L}_{2}[\rho] = -i\Upsilon_{\rm dyn}(-\omega_{1},\omega_{2})[A(\omega_{1})A(\omega_{2}),\rho] + K(-\omega_{1},\omega_{2})\left(A(\omega_{2})\rho A(\omega_{1}) - \frac{1}{2}\{A(\omega_{1})A(\omega_{2}),\rho\}\right)$$
(C62)

Then, we have:

$$\mathcal{L}_2[\varrho_2] = i\Upsilon_{\rm dyn}(-\omega_1,\omega_2)\Upsilon_{\rm st}(-\omega_3,\omega_4)\alpha(\omega_3+\omega_4)A(\omega_1)A(\omega_2)\varrho_2A(\omega_3)A(\omega_4) \tag{C63}$$

$$-i\Upsilon_{\rm dyn}(-\omega_1,\omega_2)\Upsilon_{\rm st}(-\omega_3,\omega_4)\alpha(\omega_3+\omega_4)\varrho_2A(\omega_3)A(\omega_4)A(\omega_1)A(\omega_2) \tag{C64}$$

$$+\frac{1}{2}\left(\Upsilon_{\rm st}(-\omega_3,\omega_4)\alpha(\omega_3+\omega_4)K(-\omega_1,\omega_2)A(\omega_1)A(\omega_2)\varrho_2A(\omega_3)A(\omega_4)\right)\tag{C65}$$

$$-\Upsilon_{\rm st}(-\omega_3,\omega_4)\alpha(\omega_3+\omega_4)K(-\omega_1,\omega_2)A(\omega_2)\varrho_2A(\omega_3)A(\omega_4)A(\omega_1) \tag{C66}$$

$$+\frac{1}{2}\left(\Upsilon_{\rm st}(-\omega_3,\omega_4)\alpha(\omega_3+\omega_4)K(-\omega_1,\omega_2)\varrho_2A(\omega_3)A(\omega_4)A(\omega_1)A(\omega_2)\right)\tag{C67}$$

which we may rewrite as:

$$\mathcal{L}_{2}[\varrho_{2}] = \left(-i\Upsilon_{\mathrm{dyn}}(-\omega_{1},\omega_{2})\Upsilon_{\mathrm{st}}(-\omega_{3},\omega_{4})\alpha(\omega_{3}+\omega_{4}) + \frac{1}{2}\left(\Upsilon_{\mathrm{st}}(-\omega_{3},\omega_{4})\alpha(\omega_{3}+\omega_{4})K(-\omega_{1},\omega_{2})\right)\right)e^{-\beta H_{0}}A(\omega_{3})A(\omega_{4})A(\omega_{1})A(\omega_{2})A(\omega_{3})A(\omega_{4})A$$

Since all ω_i 's are mute indices, we change them such that one obtains:

$$g_{22}(\omega_1,\omega_2,\omega_3,\omega_4) = \Upsilon_{\rm st}(-\omega_3,\omega_4)\alpha(\omega_3+\omega_4)\left(i\Upsilon_{\rm dyn}(-\omega_1,\omega_2)+\frac{1}{2}K(-\omega_1,\omega_2)\right) - \Upsilon_{\rm st}(-\omega_1,\omega_2)\alpha(\omega_1+\omega_2)\left(i\Upsilon_{\rm dyn}(-\omega_3,\omega_4)-\frac{1}{2}K(-\omega_3,\omega_4)\right) - e^{-\beta\omega_1}\Upsilon_{\rm st}(-\omega_2,\omega_3)\alpha(\omega_2+\omega_3)K(-\omega_4,\omega_1).$$
 (C69)

b. g_{40} function (cumulant equation)

We consider the fourth-order generator of the cumulant in the Schrödinger picture:

$$\mathcal{L}_{t}^{(4)}[\rho] = \frac{1}{2} \int_{0}^{t} ds \ e^{-iH_{0}t} \left[\tilde{\mathcal{L}}_{s}^{R}, \tilde{\mathcal{L}}_{t}^{R} \right] [e^{iH_{0}t} \rho e^{-iH_{0}t}] e^{iH_{0}t}.$$
(C70)

Acting on ρ_0 that commutes with H_0 , this simplifies to:

$$\mathcal{L}_{t}^{(4)}[\varrho_{0}] = \frac{1}{2} \int_{0}^{t} ds \ e^{-iH_{0}t} \left[\tilde{\mathcal{L}}_{s}^{R}, \tilde{\mathcal{L}}_{t}^{R} \right] [\varrho_{0}] e^{iH_{0}t}$$
(C71)

We then define

$$\mathcal{L}_{t}^{(4)}[\varrho_{0}] = \sum_{\omega_{1},\omega_{2},\omega_{3},\omega_{4}} g_{40}(\omega_{1},\omega_{2},\omega_{3},\omega_{4},t) e^{-\beta H_{0}} A(\omega_{1}) A(\omega_{2}) A(\omega_{3}) A(\omega_{4}).$$
(C72)

To get an expression for g_{40} , we first compute the action of $\tilde{\mathcal{L}}_s^R \tilde{\mathcal{L}}_t^R$ on ϱ_0 , i.e.,

$$\tilde{\mathcal{L}}_{s}^{R}\tilde{\mathcal{L}}_{t}^{R}[\varrho_{0}] = \sum_{\omega_{1},\omega_{2},\omega_{3},\omega_{4}} f(\omega_{1},\omega_{2},\omega_{3},\omega_{4})e^{-\beta H_{0}}A(\omega_{1})A(\omega_{2})A(\omega_{3})A(\omega_{4})$$
(C73)

where

$$f(\omega_1, \omega_2, \omega_3, \omega_4, t, s) = \tilde{\mathcal{S}}(-\omega_1, \omega_2, s)\tilde{\mathcal{S}}(-\omega_3, \omega_4, t)e^{-\beta(\omega_1 + \omega_2)} - \tilde{\mathcal{S}}(-\omega_1, \omega_2, s)\tilde{\mathcal{S}}(-\omega_3, \omega_4, t)e^{-\beta(\omega_1 + \omega_2 + \omega_3 + \omega_4)}$$
(C74)
+
$$i \tilde{\mathcal{S}}(-\omega_1, \omega_2, s)\tilde{\mathcal{S}}(-\omega_2, \omega_4, t)e^{-\beta(\omega_1 + \omega_2)} + i \tilde{\mathcal{S}}(-\omega_2, \omega_2, s)\tilde{\mathcal{S}}(-\omega_3, \omega_4, t)e^{-\beta(\omega_1 + \omega_2 + \omega_3 + \omega_4)}$$
(C75)

$$+\frac{1}{2}\mathcal{S}(-\omega_{1},\omega_{2},s)\gamma(-\omega_{3},\omega_{4},t)e^{-\beta(\omega_{1}+\omega_{2}+\omega_{3})} + \frac{1}{2}\mathcal{S}(-\omega_{1},\omega_{2},s)\gamma(-\omega_{3},\omega_{4},t)e^{-\beta(\omega_{1}+\omega_{2}+\omega_{3})} + \tilde{\mathcal{S}}(-\omega_{1},\omega_{2},t)\tilde{\mathcal{S}}(-\omega_{3},\omega_{4},s)e^{-\beta(\omega_{1}+\omega_{2})}$$
(C75)

$$-\tilde{\mathcal{S}}(-\omega_1,\omega_2,t)\tilde{\mathcal{S}}(-\omega_3,\omega_4,s) + \frac{i}{2}\tilde{\mathcal{S}}(-\omega_1,\omega_2,t)\tilde{\gamma}(-\omega_3,\omega_4,s)e^{-\beta(\omega_1+\omega_2)} - \frac{i}{2}\tilde{\mathcal{S}}(-\omega_1,\omega_2,t)\tilde{\gamma}(-\omega_3,\omega_4,s)$$
(C77)

$$-i\tilde{\mathcal{S}}(-\omega_2,\omega_3,t)\tilde{\gamma}(-\omega_4,\omega_1,s)e^{-\beta(\omega_1+\omega_2+\omega_3)} + i\tilde{\mathcal{S}}(-\omega_2,\omega_3,t)\tilde{\gamma}(-\omega_4,\omega_1,s)e^{-\beta\omega_1}$$

$$(C78)$$

$$-\frac{i}{2}\tilde{\mathcal{S}}(-\omega_3,\omega_4,s)\tilde{\gamma}(-\omega_1,\omega_2,t)e^{-\beta(\omega_1+\omega_2)} - \frac{i}{2}\tilde{\mathcal{S}}(-\omega_3,\omega_4,s)\tilde{\gamma}(-\omega_1,\omega_2,t) + i\tilde{\mathcal{S}}(-\omega_3,\omega_4,s)\tilde{\gamma}(-\omega_2,\omega_1,t)e^{-\beta\omega_1}$$
(C79)

$$-\frac{i}{2}\tilde{\mathcal{S}}(-\omega_3,\omega_4,t)\tilde{\gamma}(-\omega_1,\omega_2,s)e^{-\beta(\omega_1+\omega_2)} + \frac{i}{2}\tilde{\mathcal{S}}(-\omega_3,\omega_4,t)\tilde{\gamma}(-\omega_1,\omega_2,s)e^{-\beta(\omega_1+\omega_2+\omega_3+\omega_4)}$$
(C80)

$$+\frac{1}{4}\tilde{\gamma}(-\omega_{1},\omega_{2},s)\tilde{\gamma}(-\omega_{3},\omega_{4},t)e^{-\beta(\omega_{1}+\omega_{2})}+\frac{1}{4}\tilde{\gamma}(-\omega_{1},\omega_{2},s)\tilde{\gamma}(-\omega_{3},\omega_{4},t)e^{-\beta(\omega_{1}+\omega_{2}+\omega_{3}+\omega_{4})}$$
(C81)

$$-\frac{1}{2}\tilde{\gamma}(-\omega_{1},\omega_{2},s)\tilde{\gamma}(-\omega_{4},\omega_{3},t)e^{-\beta(\omega_{1}+\omega_{2}+\omega_{3})} + \frac{1}{4}\tilde{\gamma}(-\omega_{1},\omega_{2},t)\tilde{\gamma}(-\omega_{3},\omega_{4},s)e^{-\beta(\omega_{1}+\omega_{2})}$$
(C82)

$$+\frac{1}{4}\tilde{\gamma}(-\omega_1,\omega_2,t)\tilde{\gamma}(-\omega_3,\omega_4,s) - \frac{1}{2}\tilde{\gamma}(-\omega_2,\omega_1,t)\tilde{\gamma}(-\omega_3,\omega_4,s)e^{-\beta\omega_1}$$
(C83)

$$-\frac{1}{2}\tilde{\gamma}(-\omega_2,\omega_3,t)\tilde{\gamma}(-\omega_4,\omega_1,s)e^{-\beta(\omega_1+\omega_2+\omega_3)} - \frac{1}{2}\tilde{\gamma}(-\omega_2,\omega_3,t)\tilde{\gamma}(-\omega_4,\omega_1,s)e^{-\beta\omega_1}$$
(C84)

$$+\tilde{\gamma}(-\omega_3,\omega_2,t)\tilde{\gamma}(-\omega_4,\omega_1,s)e^{-\rho(\omega_1+\omega_2)} \tag{C85}$$

Consequently, we have

$$g_{40}(\omega_1, \omega_2, \omega_3, \omega_4, t) = \frac{1}{2}e^{i(\omega_1 + \omega_2 + \omega_3 + \omega_4)t} \int_0^t ds \ (f(\omega_1, \omega_2, \omega_3, \omega_4, t, s) - f(\omega_1, \omega_2, \omega_3, \omega_4, s, t)).$$
(C86)

4. Two-level system

Now, we shall specialize to the case of a two-level system. We then have k = 0, 1 and $\epsilon_1 - \epsilon_0 = \omega_0$, such that

$$G(|0\rangle \to |0\rangle) = \{(0,0,0,0), (\omega_0, -\omega_0, 0, 0), (\omega_0, 0, -\omega_0, 0), (\omega_0, 0, 0, -\omega_0), (C87)\}$$

$$(0, \omega_0, -\omega_0, 0), (0, \omega_0, 0, -\omega_0), (0, 0, \omega_0, -\omega_0), (\omega_0, -\omega_0, \omega_0, -\omega_0\})$$
(C88)

The set $S(|1\rangle \rightarrow |1\rangle)$ is the same but with changed sign of the qubit frequency $\omega_0 \rightarrow -\omega_0$. Then, according to Eq. (C69), one can first observe that the coefficient g_{22} summed over first seven four-tuples vanishes, i.e.,

$$g_{22}(0,0,0,0) + g_{22}(\omega_0,-\omega_0,0,0) + g_{22}(\omega_0,0,-\omega_0,0) + g_{22}(\omega_0,0,0,-\omega_0) + g_{22}(0,\omega_0,-\omega_0,0) + g_{22}(0,\omega_0,0,-\omega_0) + g_{22}(0,0,\omega_0,-\omega_0) = 0$$
(C89)

whereas for the last one we have

$$g_{22}(\omega_0, -\omega_0, \omega_0, -\omega_0) = \beta \left(\Upsilon_{\rm st}(-\omega_0, -\omega_0) K(-\omega_0, -\omega_0) - e^{-\beta\omega_0} \Upsilon_{\rm st}(\omega_0, \omega_0) K(\omega_0, \omega_0) \right).$$
(C90)

If additionally $K(\omega, \omega)$ obeys the detailed balance condition, then

$$g_{22}(\omega_0, -\omega_0, \omega_0, -\omega_0) = \beta e^{-\beta\omega_0} \left(\Upsilon_{\rm st}(-\omega_0, -\omega_0) - \Upsilon_{\rm st}(\omega_0, \omega_0)\right) K(\omega_0, \omega_0). \tag{C91}$$

a. Second-order master equation

Now, since for arbitrary master equation of the form (6), which is up to second order in λ , we also have $g_{40} = 0$. From this we conclude that Eq. (C18) is satisfied if

$$\Upsilon_{\rm st}(\omega_0,\omega_0) = \Upsilon_{\rm st}(-\omega_0,-\omega_0). \tag{C92}$$

Since for a two-level system, in general we have

$$\langle 0 | H_{\rm st}^{(2)} | 0 \rangle = \Upsilon_{\rm st}(0,0) + \Upsilon_{\rm st}(\omega_0,\omega_0), \quad \langle 1 | H_{\rm st}^{(2)} | 1 \rangle = \Upsilon_{\rm st}(0,0) + \Upsilon_{\rm st}(-\omega_0,-\omega_0). \tag{C93}$$

Thus, applying the condition (C92), we finally get:

$$\langle 0 | H_{\rm st}^{(2)} | 0 \rangle = \langle 1 | H_{\rm st}^{(2)} | 1 \rangle = \Upsilon_{\rm st}(\omega_0, \omega_0).$$
 (C94)

b. Cumulant equation

To solve the Eq. (C18) for the cumulant master equation we need to additionally calculate the term involving the coefficient g_{40} . Putting the expression (C86), we observe that, similarly to the summation of g_{22} , the sum over first seven tuples vanishes, such that we obtain a very simple expression

$$\lim_{t \to \infty} \sum_{(\omega_1, \omega_2, \omega_2, \omega_4) \in G(|0\rangle \to |0\rangle)} g_{40}(\omega_1, \omega_2, \omega_2, \omega_4, t) = g_{40}(\omega_0, -\omega_0, \omega_0, -\omega_0),$$
(C95)

where

$$g_{40}(\omega_0, -\omega_0, \omega_0, -\omega_0) = \frac{1}{2}e^{-\beta\omega_0}(1 + e^{\beta\omega_0})\gamma(\omega_0)\int_0^\infty ds \ (e^{-\beta\omega_0}\gamma(\omega_0, s) - \gamma(-\omega_0, s)).$$
(C96)

Since, the leading order of the cumulant master equation is the Bloch-Redfield generator, we also have

$$g_{22}(\omega_0, -\omega_0, \omega_0, -\omega_0) = \beta e^{-\beta\omega_0} \left(\Upsilon_{\rm st}(-\omega_0, -\omega_0) - \Upsilon_{\rm st}(\omega_0, \omega_0)\right) \gamma(\omega_0).$$
(C97)

Finally, we need to solve

$$g_{22}(\omega_0, -\omega_0, \omega_0, -\omega_0) + g_{40}(\omega_0, -\omega_0, \omega_0, -\omega_0) = 0$$
(C98)

which gives us

$$\Upsilon_{\rm st}(\omega_0,\omega_0) - \Upsilon_{\rm st}(-\omega_0,-\omega_0) = \frac{1}{2\beta}(1+e^{\beta\omega_0})\int_0^\infty ds \ (e^{-\beta\omega_0}\gamma(\omega_0,s) - \gamma(-\omega_0,s)) \tag{C99}$$

$$=\frac{1}{2\beta}\int_0^\infty ds \,\left(\gamma(\omega_0,s)+e^{-\beta\omega_0}\gamma(\omega_0,s)-\gamma(-\omega_0,s)-e^{\beta\omega_0}\gamma(-\omega_0,s)\right)\tag{C100}$$

$$= \frac{1}{2\beta} \int_0^\infty ds \, \left(\gamma(\omega_0, s) - e^{\beta\omega_0} \gamma(-\omega_0, s)\right) - \frac{1}{2\beta} \int_0^\infty ds \, \left(\gamma(-\omega_0, s) - e^{-\beta\omega_0} \gamma(\omega_0, s)\right)$$
(C101)

Finally, without loss of generality, we can put

$$\Upsilon_{\rm st}(\omega,\omega) = \frac{1}{2\beta} \int_0^\infty ds \ (\gamma(\omega,s) - e^{\beta\omega}\gamma(-\omega,s)). \tag{C102}$$

At the end, let us compare it with the mean-force coefficient, i.e.,

$$\Upsilon_{\rm mf}(\omega,\omega) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \ D(\omega,\omega,\Omega)\gamma(\Omega)$$
(C103)

where

$$D(\omega,\omega,\Omega) = -\frac{1}{\beta} \int_0^\beta dt \int_0^t ds \ e^{s(\omega-\Omega)} = \frac{1 - e^{\beta(\omega-\Omega)} + \beta(\omega-\Omega)}{\beta(\omega-\Omega)^2}.$$
 (C104)

In order to compare both expression, we propose another representation for the mean-force coefficient. Using the detailed balance condition, we see that

$$\int_{-\infty}^{+\infty} d\Omega \ f(\Omega)\gamma(\Omega) = \int_{-\infty}^{+\infty} d\Omega \ f(\Omega)e^{\beta\Omega}\gamma(-\Omega) = \int_{-\infty}^{+\infty} d\Omega \ f(-\Omega)e^{-\beta\Omega}\gamma(\Omega)$$
(C105)

Consequently, we can write

$$\Upsilon_{\rm mf}(\omega,\omega) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \ D(\omega,\omega,\Omega)\gamma(\Omega) = \int_{-\infty}^{+\infty} d\Omega \ \frac{1+\beta(\omega-\Omega)-e^{\beta(\omega-\Omega)}}{\beta(\omega-\Omega)^2}\gamma(\Omega) \tag{C106}$$

$$= \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \, \left(\frac{1 + \beta(\omega - \Omega)}{\beta(\omega - \Omega)^2} - \frac{e^{\beta(\omega - \Omega)}}{\beta(\omega - \Omega)^2} \right) \gamma(\Omega) \tag{C107}$$

$$= \frac{1}{2\pi\beta} \int_{-\infty}^{+\infty} d\Omega \,\left(\frac{1}{(\omega-\Omega)^2} - \frac{e^{\beta\omega}}{(\omega+\Omega)^2} + \frac{\beta}{\omega-\Omega} \right) \gamma(\Omega) \tag{C108}$$

$$= \frac{1}{2\pi\beta} \int_{-\infty}^{+\infty} d\Omega \,\left(\frac{1}{(\omega-\Omega)^2} - \frac{e^{\beta\omega}}{(\omega+\Omega)^2}\right) \gamma(\Omega) + \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \,\frac{\gamma(\Omega)}{\omega-\Omega} \tag{C109}$$

Finally, let us observe that

$$\int_0^\infty ds \ \gamma(\omega, s) = \frac{1}{\pi} \int_{-\infty}^{+\infty} d\Omega \ \frac{\gamma(\Omega)}{(\omega - \Omega)^2},\tag{C110}$$

and

$$\mathcal{S}(\omega) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \, \frac{\gamma(\Omega)}{\omega - \Omega},\tag{C111}$$

such that according to the previous choice for $\Upsilon_{\rm st}(\omega,\omega)$, we have simply

$$\Upsilon_{\rm mf}(\omega,\omega) = \Upsilon_{\rm st}(\omega,\omega) + \mathcal{S}(\omega). \tag{C112}$$

Appendix D: Cumulant equation

Consider a system interacting with a thermal reservoir whose Hamiltonian is given by:

$$H = H_0 + H_R + \lambda H_I \tag{D1}$$

Let us also consider the Born Approximation such that $\rho(0) = \rho_S(0) \otimes \rho_R$ where ρ_R is a stationary state of the environment. In the interaction picture the reduced state at time t is:

$$\rho_S(t) = \operatorname{Tr}_R\left(U(t,t_0)\rho_S(t_0) \otimes \rho_R(t_0)U^{\dagger}(t,t_0)\right)$$
(D2)

One may expand the evolution operator in the interaction picture $U(t,0) = \mathcal{T}e^{-i\int_0^t H_I(t')dt'}$ and rearrange terms (of the same power of H_I) to obtain:

$$\rho_{S}(t) = \rho_{S}(0) \underbrace{-\lambda^{2} \frac{\mathcal{T}}{2} \int_{0}^{t} dt_{1} \int_{0}^{t} dt_{2} \operatorname{Tr}_{R} \left([H_{I}(t_{1}), [H_{I}(t_{2}), \rho_{S}(0) \otimes \rho_{R}]] \right)}_{\tilde{K}_{t}^{(2)}} + \mathcal{O}(\lambda^{3})$$
(D3)

The terms $\mathcal{O}(H_I^3)$ can be neglected for weak coupling or short times. We already considered the initial state of the bath to be thermal $\rho_B(0) = \rho_\beta = e^{-\beta H_B} / \operatorname{Tr} \{e^{-\beta H_B}\}$ and the bath operators to be centralized. Let us know focus on the second term, let us apply time-ordering explicitly so that:

$$\tilde{K}_{t}^{(2)} = -\frac{\lambda^{2}}{2} \int_{0}^{t} dt_{1} \int_{0}^{t} dt_{2} \theta(t_{1} - t_{2}) \operatorname{Tr}_{R} \left([H_{I}(t_{1}), [H_{I}(t_{2}), \rho_{S}(0) \otimes \rho_{R}]] \right) - \frac{\lambda^{2}}{2} \int_{0}^{t} dt_{1} \int_{0}^{t} dt_{2} \theta(t_{2} - t_{1}) \operatorname{Tr}_{R} \left([H_{I}(t_{2}), [H_{I}(t_{1}), \rho_{S}(0) \otimes \rho_{R}]] \right)$$
(D4)

Let us know expand the double commutators:

$$\begin{split} \tilde{K}_{t}^{(2)} &= -\frac{\lambda^{2}}{2} \int_{0}^{t} dt_{1} \int_{0}^{t} dt_{2} \theta(t_{1} - t_{2}) \operatorname{Tr}_{R} \left[H_{I}(t_{1}) H_{I}(t_{2}) \rho_{S}(0) \rho_{R} - H_{I}(t_{1}) \rho_{S}(0) \rho_{R} H_{I}(t_{2}) - H_{I}(t_{2}) \rho_{S}(0) \rho_{R} H_{I}(t_{1}) \right. \\ &+ \rho_{S}(0) \rho_{R} H_{I}(t_{1}) H_{I}(t_{2}) \Big] - \frac{\lambda^{2}}{2} \int_{0}^{t} dt_{1} \int_{0}^{t} dt_{2} \theta(t_{2} - t_{1}) \operatorname{Tr}_{R} \left[H_{I}(t_{2}) H_{I}(t_{1}) \rho_{S}(0) \rho_{R} - H_{I}(t_{2}) \rho_{S}(0) \rho_{R} H_{I}(t_{1}) \right. \\ &- H_{I}(t_{1}) \rho_{S}(0) \rho_{R} H_{I}(t_{2}) + \rho_{S}(0) \rho_{R} H_{I}(t_{2}) H_{I}(t_{1}) \Big] \end{split}$$

From here it can be seen that we have three kind of terms, namely $H_I^2 \rho$, $H_I \rho H_I$, ρH_I^2 . Let us consider each of those independently

$$H_{I}\rho H_{I} : \frac{\lambda^{2}}{2} \int_{0}^{t} dt_{1} \int_{0}^{t} dt_{2} \left(\theta(t_{1} - t_{2}) + \theta(t_{2} - t_{1}) \right) \operatorname{Tr}_{R} \left[H_{I}(t_{1})\rho_{S}(0)\rho_{R}H_{I}(t_{2}) + H_{I}(t_{2})\rho_{S}(0)\rho_{R}H_{I}(t_{1}) \right]$$
(D5)

$$= \frac{\lambda^2}{2} \int_0^t dt_1 \int_0^t dt_2 \operatorname{Tr}_R \left[H_I(t_1)\rho_S(0)\rho_R H_I(t_2) + H_I(t_2)\rho_S(0)\rho_R H_I(t_1) \right]$$
(D6)

$$=\lambda^{2} \int_{0}^{t} dt_{1} \int_{0}^{t} dt_{2} \operatorname{Tr}_{R} \left[H_{I}(t_{1})\rho_{S}(0)\rho_{R}H_{I}(t_{2}) \right]$$
(D7)

where in the last step we used a change of variables on the second term, such that $t_1 \leftrightarrow t_2$. Next, we consider the other two missing terms

$$H_{I}^{2}\rho : -\frac{\lambda^{2}}{2} \int_{0}^{t} dt_{1} \int_{0}^{t} dt_{2} \Big(\theta(t_{1}-t_{2})\operatorname{Tr}_{R} \Big[H_{I}(t_{1})H_{I}(t_{2})\rho_{S}(0)\rho_{R}\Big] + \theta(t_{2}-t_{1})\operatorname{Tr}_{R} \Big[H_{I}(t_{2})H_{I}(t_{1})\rho_{S}(0)\rho_{R}\Big]\Big)$$
(D8)
$$= -\frac{\lambda^{2}}{2} \int_{0}^{t} dt_{2} \Big(\theta(t_{1}-t_{2})\operatorname{Tr}_{R} \Big[H_{I}(t_{1})H_{I}(t_{2})\rho_{S}(0)\rho_{R}\Big] + \operatorname{Tr}_{R} \Big[H_{I}(t_{2})H_{I}(t_{1})\rho_{S}(0)\rho_{R}\Big]\Big)$$
(D9)

$$= -\frac{\lambda}{2} \int_{0}^{1} dt_{1} \int_{0}^{1} dt_{2} \left(\theta(t_{1} - t_{2}) \operatorname{Tr}_{R} \left[[H_{I}(t_{1}), H_{I}(t_{2})] \rho_{S}(0) \rho_{R} \right] + \operatorname{Tr}_{R} \left[H_{I}(t_{2}) H_{I}(t_{1}) \rho_{S}(0) \rho_{R} \right] \right)$$
(D9)

$$\rho H_I^2 : -\frac{\lambda}{2} \int_0^t dt_1 \int_0^t dt_2 \Big(\theta(t_1 - t_2) \operatorname{Tr}_R \Big[\rho_S(0) \rho_R H_I(t_2) H_I(t_1) \Big] + \theta(t_2 - t_1) \operatorname{Tr}_R \Big[H_I(t_1) H_I(t_2) \rho_S(0) \rho_R \Big] \Big) (D10)$$

$$= -\frac{\lambda^2}{2} \int_0^t dt_1 \int_0^t dt_2 \Big(\theta(t_1 - t_2) \operatorname{Tr}_R \Big[\rho_S(0) \rho_R [H_I(t_2), H_I(t_1)] \Big] + \operatorname{Tr}_R \Big[\rho_S(0) \rho_R H_I(t_1) H_I(t_2) \Big] \Big)$$
(D11)

In both cases the step taken from one line to the other was summing a zero so that the terms could be recast in that form, they were $\pm \frac{1}{2} \int_0^t dt_1 \int_0^t dt_2 \theta(t_1 - t_2) \operatorname{Tr}_R \left[H_I(t_2) H_I(t_1) \rho_S(0) \rho_R \right]$ in the first case and $\pm \frac{1}{2} \int_0^t dt_1 \int_0^t dt_2 \theta(t_1 - t_2) \operatorname{Tr}_R \left[\rho_S(0) \rho_R H_I(t_1) H_I(t_2) \right]$ in the second one. Regrouping all terms we have

$$\tilde{K}_{t}^{(2)} = \lambda^{2} \int_{0}^{t} dt_{1} \int_{0}^{t} dt_{2} \left(\operatorname{Tr}_{R} \left[H_{I}(t_{1})\rho_{S}(0)\rho_{R}H_{I}(t_{2}) \right] - \frac{1}{2} \left(\operatorname{Tr}_{R} \left[\rho_{S}(0)\rho_{R}H_{I}(t_{1})H_{I}(t_{2}) \right] + \operatorname{Tr}_{R} \left[H_{I}(t_{2})H_{I}(t_{1})\rho_{S}(0)\rho_{R} \right] \right) \right) - \frac{\lambda^{2}}{2} \int_{0}^{t} dt_{1} \int_{0}^{t} dt_{2}\theta(t_{1} - t_{2}) \left(\operatorname{Tr}_{R} \left[[H_{I}(t_{1}), H_{I}(t_{2})]\rho_{S}(0)\rho_{R} \right] - \operatorname{Tr}_{R} \left[\rho_{S}(0)\rho_{R}[H_{I}(t_{1}), H_{I}(t_{2})] \right] \right)$$
(D12)

$$= \lambda^{2} \int_{0}^{t} dt_{1} \int_{0}^{t} dt_{2} \left(\operatorname{Tr}_{R} \left[H_{I}(t_{1})\rho_{S}(0)\rho_{R}H_{I}(t_{2}) \right] - \frac{1}{2} \left(\operatorname{Tr}_{R} \left[\rho_{S}(0)\rho_{R}H_{I}(t_{1})H_{I}(t_{2}) \right] + \operatorname{Tr}_{R} \left[H_{I}(t_{2})H_{I}(t_{1})\rho_{S}(0)\rho_{R} \right] \right) \right)$$

$$- i\lambda^{2} [\Lambda(t), \rho_{S}(0)]$$
(D13)

where:

$$\Lambda(t) = \frac{1}{2i} \int_{0}^{t} dt_1 \int_{0}^{t} dt_2 \theta(t_1 - t_2) \operatorname{Tr}_R \left[[H_I(t_1), H_I(t_2)] \rho_R \right]$$
(D14)

$$= \frac{1}{2i} \int_0^t dt_1 \int_0^t dt_2 sgn(t_1 - t_2) \operatorname{Tr}_R \left[H_I(t_1) H_I(t_2) \rho_R \right]$$
(D15)

where we used $\theta(x) = \frac{1+sgn(x)}{2}$. Now, if we expand the interaction Hamiltonian in the interaction picture $H_I = \sum_{w,k} e^{iwt} A_k(w) B_k = \sum_{w,k} e^{-iwt} A_k^{\dagger}(w) B_k$

$$\Lambda(t) = \frac{1}{2i} \sum_{w,w'} \sum_{\alpha\beta} \int_0^t dt_1 \int_0^t dt_2 sgn(t_1 - t_2) e^{i(wt_1 - w't_2)} A^{\dagger}_{\alpha}(w) A_{\beta}(w') \langle B_{\alpha}(t_1) B_{\beta}(t_2) \rangle_R$$
$$= \sum_{w,w'} \sum_{\alpha\beta} \Xi(w,w',t) A^{\dagger}_{\alpha}(w) A_{\beta}(w')$$
(D16)

So we obtain:

$$\tilde{K}_{t}^{(2)}[\rho_{S}(0)] = -i\sum_{w,w'}\sum_{\alpha\beta}\Xi(w,w',t)[A_{\alpha}^{\dagger}(w)A_{\beta}(w'),\rho_{S}(0)] + \xi_{\alpha\beta}(w,w',t)\Big(A_{\beta}(w')\rho_{S}(0)A_{\alpha}^{\dagger}(w) - \frac{1}{2}\{A_{\alpha}^{\dagger}(w)A_{\beta}(w'),\rho_{S}(0)\}\Big)$$

where

$$\xi_{\alpha\beta}(w,w',t) = \int_0^t dt_1 \int_0^t dt_2 e^{i(wt_1 - w't_2)} \langle R_\alpha(t_1) R_\beta(t_2) \rangle.$$
(D17)

We may rewrite this in terms of previously obtained quantities as:

$$\xi_{\alpha\beta}(w,w',t) = \int_0^t ds \int_0^t d\omega e^{i(ws-w'\omega)} \langle R_\alpha(s)R_\beta(w)\rangle$$
(D18)

$$= \int_{0}^{t} ds \int_{s}^{t} d\omega e^{i(ws-w'\omega)} \langle R_{\alpha}(s)R_{\beta}(w)\rangle + \int_{0}^{t} ds \int_{0}^{s} d\omega e^{i(ws-w'\omega)} \langle R_{\alpha}(s)R_{\beta}(w)\rangle$$
(D19)

$$= \int_{0}^{t} d\omega \int_{0}^{\omega} ds e^{i(ws-w'\omega)} \langle R_{\alpha}(s)R_{\beta}(w)\rangle + \int_{0}^{t} ds \int_{0}^{s} d\omega e^{i(ws-w'\omega)} \langle R_{\alpha}(s)R_{\beta}(w)\rangle$$
(D20)

$$= \int_{0}^{t} ds \int_{0}^{s} d\omega e^{i(w\omega - w's)} \langle R_{\alpha}(w)R_{\beta}(s)\rangle + \int_{0}^{t} ds \int_{0}^{s} d\omega e^{i(ws - w'\omega)} \langle R_{\alpha}(s)R_{\beta}(w)\rangle$$
(D21)

$$= \int_{0}^{t} ds \int_{0}^{s} d\omega e^{i(w\omega - w's)} \langle R_{\alpha}(w - s)R_{\beta} \rangle + \int_{0}^{t} ds \int_{0}^{s} d\omega e^{i(ws - w'\omega)} \langle R_{\alpha}(s - w)R_{\beta} \rangle$$
(D22)

$$= \int_{0}^{t} ds \int_{0}^{s} d\xi e^{i((w-w')s-w\xi)} \langle R_{\alpha}(-\xi)R_{\beta}\rangle + \int_{0}^{t} ds \int_{0}^{s} d\xi e^{i((w-w')s+\xi w')} \langle R_{\alpha}(\xi)R_{\beta}\rangle$$
(D23)

$$= \int_0^t ds e^{i(w-w')s} (\Gamma^*_{\beta\alpha}(w,s) + \Gamma_{\alpha\beta}(w',s))$$
(D24)

$$=\int_{0}^{t} ds e^{i(w-w')s} \gamma_{\alpha\beta}(w,w',s) = \int_{0}^{t} ds \tilde{\gamma}_{\alpha\beta}(w,w',s)$$
(D25)

Now, we can notice that the derivative of such coefficient corresponds to:

$$\frac{d}{dt}\xi_{\alpha\beta}(w,w',t) = e^{i(w-w')t}\gamma(w,w',t) = \tilde{\gamma}(w,w',t)$$
(D26)

Furthermore from [22] we know that:

$$\frac{d}{dt}\Xi_{\alpha\beta}(w,w',t) = \frac{e^{i(w'-w)t}}{2i}(\Gamma_{\alpha\beta}(w',t) - \Gamma^*_{\beta\alpha}(w,t)) = e^{i(w'-w)t}S_{\alpha\beta}(w,w',t) = \tilde{S}_{\alpha\beta}(w,w',t)$$
(D27)

One may then rewrite $\tilde{K}_t^{(2)}$ as:

$$\tilde{K}_{t}^{(2)}[\rho] = \lambda^{2} \int_{0}^{t} ds \sum_{\omega,\omega'} \sum_{\alpha\beta} \left(i \tilde{\mathcal{S}}_{\alpha\beta}(\omega,\omega',s) [\rho, A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega')] + \tilde{\gamma}_{\alpha\beta}(\omega,\omega',s) \left(A_{\beta}(\omega')\rho A_{\alpha}^{\dagger}(\omega) - \frac{1}{2} \{ A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega'), \rho \} \right) \right)$$
(D28)

,

Appendix E: Comparison with dynamical Hamiltonian and the steady state - qubit case

In this section we consider the particular case of a qubit coupled to a bosonic bath given by

$$H = \frac{\omega_0}{2}\sigma_z + \sum_k \Omega_k a_k^{\dagger} a_k + S \sum_{k=1}^{\infty} \lambda_k (a_k + a_k^{\dagger})$$
(E1)

where we take S to be a general interaction operator in the pauli basis:

$$S = x\sigma_x + y\sigma_y + z\sigma_z \tag{E2}$$

This form of Hamiltonian with y = 0 has been studied previously in [12]. Where it was reported that such Hamiltonian had steady-state coherences, they used a similar expansion to the general results presented in this article. In this section, we see that the general framework presented agrees with their results. Using equation (B17), and this interaction the second-order correction to the Hamiltonian takes the form:

$$H_{\mathbf{k}}^{(2)} = \begin{bmatrix} z^{2} \Upsilon(0,0) + (x^{2} + y^{2}) \Upsilon(\omega,\omega) & (x - iy)z(\Upsilon(0,-\omega) - \Upsilon(\omega,0)) \\ (x + iy)z(\Upsilon(0,-\omega) - \Upsilon(\omega,0)) & z^{2} \Upsilon(0,0) + (x^{2} + y^{2}) \Upsilon(-\omega,-\omega) \end{bmatrix}$$
(E3)

where k indicates to dynamical (dyn), steady-state (st) or mean-force (mf) correction. We can rewrite this correction as a linear combination of the Pauli Matrices such that:

$$H_{\mathbf{k}}^{(2)} = A\mathbb{1} + B\sigma_x + C\sigma_y + D\sigma_z \tag{E4}$$

$$A = z^{2} \Upsilon(0,0) + \frac{x^{2} + y^{2}}{2} (\Upsilon(w,w) + \Upsilon(-w,-w))$$
(E5)

$$B = xz(\Upsilon(0, -w) - \Upsilon(w, 0))$$
(E6)

$$C = yz(\Upsilon(0, -w) - \Upsilon(w, 0))$$
(E7)

$$D = \frac{x^2 + y^2}{2} (\Upsilon(w, w) - \Upsilon(-w, -w))$$
(E8)

We can see how the different approaches differ qualitatively by looking at the structure of the different $\Upsilon(w, w')$ given by each approach. It is important to remark that any approach that performs the secular approximation will have both B and C equal to zero, meaning the correction will be diagonal and as such won't be able to describe the off-diagonal elements of the steady states accordingly. While nonsecular approaches such as the Bloch-Redfield equation, will have non-diagonal corrections, leading to a more appropriate description of the off-diagonal elements of the correction as well as steady state coherences. Let us for a moment recall the structure of the Bloch-Redfield coefficients which are given by (C43), simply substituting the appropriate frequencies for the qubit leads to:

$$\Upsilon(0, -w) - \Upsilon(w, 0) = \frac{S(-w) - S(w)}{2} + i\frac{\gamma(0) - (\gamma(w) + \gamma(-w))}{4}$$
(E9)

and

$$\Upsilon(w,w) + \Upsilon(-w,-w) = \mathcal{S}(w) + \mathcal{S}(-w) \tag{E10}$$

$$\Upsilon(w,w) - \Upsilon(-w,-w) = \mathcal{S}(w) - \mathcal{S}(-w)$$
(E11)

Let us now compare this coefficient with the one obtained with the mean force approach. We will only be considering the off-diagonal of the correction:

$$K(\omega) = \Upsilon(0, -\omega) - \Upsilon(\omega, 0) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\Omega \gamma(\Omega) C(\omega, \Omega)$$
(E12)

where

$$C(\omega, \Omega) = \left(\frac{\omega^2 (1 - e^{-\beta\Omega}) \coth[\frac{\beta\omega}{2}] + (1 + e^{-\beta\Omega})\Omega\omega}{\Omega(\Omega^2 - \omega^2)}\right)$$
(E13)

Additionally, our coefficients satisfy detailed balance conditions such that:

$$\gamma(-\Omega) = \gamma(\Omega)e^{-\beta\Omega}, \quad C(\omega, -\Omega) = C(\omega, \Omega)e^{\beta\Omega}$$
(E14)

Using those we see that $\gamma(-\Omega)C(\omega, -\Omega) = \gamma(\Omega)C(\omega, \Omega)$ and

$$K(\omega) = \frac{1}{\pi} \int_0^{+\infty} d\Omega \ \gamma(\Omega) C(\omega, \Omega)$$
(E15)

Let us now separate $\gamma(\Omega)$ into its symmetric and anti-symmetric parts:

$$\gamma_s(\Omega) = \frac{1}{2}(\gamma(\Omega) + \gamma(-\Omega)) = \frac{1}{2}(1 + e^{-\beta\Omega})\gamma(\Omega)$$
(E16)

$$\gamma_a(\Omega) = \frac{1}{2}(\gamma(\Omega) - \gamma(-\Omega)) = \frac{1}{2}(1 - e^{-\beta\Omega})\gamma(\Omega)$$
(E17)

Then we may write:

$$K(\omega) = \frac{2}{\pi} \int_0^{+\infty} d\Omega \,\left(\frac{\omega^2 \gamma_a(\Omega) \coth[\frac{\beta\omega}{2}] + \gamma_s(\Omega)\Omega\omega}{\Omega(\Omega^2 - \omega^2)}\right)$$
(E18)

As mentioned before this system with y = 0 had been previously considered in [12], let us now compare our results to those previously available in the literature. Their effective Hamiltonian is given by:

$$H_{S} = \begin{bmatrix} \lambda^{2} f_{1}^{2} \Upsilon(0,0) - \frac{1}{2} (\omega - 2\lambda^{2} f_{2}^{2} \Upsilon(\omega,\omega)) & \lambda^{2} f_{1} f_{2} K(\omega) \\ \lambda^{2} f_{1} f_{2} K(\omega) & \lambda^{2} f_{1}^{2} \Upsilon(0,0) + \frac{1}{2} (\omega - 2\lambda^{2} f_{2}^{2} \Upsilon(\omega,\omega)) \end{bmatrix}$$
(E19)

The couplings in this notation are $x = f_2$, $z = f_1$ and y = 0. They also use $\omega' = \omega - 2\lambda^2 f_2^2 \Upsilon(\omega, \omega)$ and $E_0 = \lambda^2 f_1^2 \Upsilon(0, 0)$ such that:

$$H_S = \begin{bmatrix} E_0 - \frac{\omega'}{2} & \lambda^2 f_1 f_2 K(\omega) \\ \lambda^2 f_1 f_2 K(\omega) & E_0 + \frac{\omega'}{2} \end{bmatrix}$$
(E20)

Then we may find that

$$\langle \sigma_x \rangle = \frac{\text{Tr}[\sigma_x e^{-\beta H_S}]}{\text{Tr}[e^{-\beta H_S}]} = -x \frac{\tanh[\sqrt{x^2 + z^2}\beta]}{\sqrt{x^2 + z^2}}$$
(E21)

where $x = \lambda^2 f_1 f_2 K(\omega)$ and $z = \frac{\omega'}{2}$.

$$\langle \sigma_x \rangle = -\frac{2x}{\omega'} \tanh[\frac{\beta\omega}{2}] + O(\lambda^3)$$
 (E22)

In [12] the authors also put $\omega' = \omega$, such that

$$\langle \sigma_x \rangle = -\frac{4\lambda^2 f_1 f_2}{\pi\omega} \int_0^{+\infty} d\Omega \, \left(\frac{\gamma_s(\Omega)\omega \tanh[\frac{\beta\omega}{2}]}{\Omega^2 - \omega^2} + \frac{\omega^2 \gamma_a(\Omega)}{\Omega(\Omega^2 - \omega^2)} \right)$$
(E23)

(E24)

Now let us compare $\gamma_{a,s}(\Omega)$ with the correlation function for a bosonic bath:

$$f(t) = \int_0^\infty d\Omega \ J(\Omega) \left(\coth\left[\frac{\beta\Omega}{2}\right] \cos(\Omega t) - i\sin(\Omega t) \right)$$
(E25)

$$f(t) = \frac{1}{2} \int_{-\infty}^{+\infty} d\Omega \ e^{-i\Omega t} \gamma(\Omega) = \frac{1}{2} \int_{-\infty}^{+\infty} d\Omega \ \gamma(\Omega) \left(\cos(\Omega t) - i\sin(\Omega t)\right)$$
(E26)

$$=\frac{1}{2}\int_{0}^{+\infty}d\Omega \left(\gamma_{s}(\Omega)+\gamma_{a}(\Omega)\right)\left(\cos(\Omega t)-i\sin(\Omega t)\right)+\frac{1}{2}\int_{0}^{+\infty}d\Omega \left(\gamma_{s}(\Omega)-\gamma_{a}(\Omega)\right)\left(\cos(\Omega t)+i\sin(\Omega t)\right)$$
(E27)

$$= \frac{1}{\pi} \int_{0}^{+\infty} d\Omega \ \left(\gamma_s(\Omega)\cos(\Omega t) - i\gamma_a(\Omega)\sin(\Omega t)\right) = \frac{1}{\pi} \int_{0}^{+\infty} d\Omega \ \gamma_a(\Omega) \left(\frac{\gamma_s(\Omega)}{\gamma_a(\Omega)}\cos(\Omega t) - i\sin(\Omega t)\right)$$
(E28)

$$= \frac{1}{\pi} \int_0^{+\infty} d\Omega \,\gamma_a(\Omega) \left(\frac{1 + e^{-\beta\Omega}}{1 - e^{-\beta\Omega}} \cos(\Omega t) - i\sin(\Omega t) \right) = \frac{1}{\pi} \int_0^{+\infty} d\Omega \gamma_a(\Omega) \left(\coth\left[\frac{\beta\Omega}{2}\right] \cos(\Omega t) - i\sin(\Omega t) \right) E^{29}$$

According to this, we have the following relations:

$$\gamma_a(\Omega) = \pi J(\Omega) = \pi \omega_a(\Omega), \quad \gamma_s(\Omega) = \pi J(\Omega) \coth\left[\frac{\beta \Omega}{2}\right] = \pi \omega_s(\Omega)$$
 (E30)

and the final result becomes:

$$\langle \sigma_x \rangle = -\frac{4\lambda^2 f_1 f_2}{\omega} \int_0^{+\infty} d\Omega \, \left(\frac{\omega_s(\Omega)\omega \tanh[\frac{\beta\omega}{2}]}{\Omega^2 - \omega^2} - \frac{\omega^2 \omega_a(\Omega)}{\Omega(\Omega^2 - \omega^2)} \right)$$
(E31)

on the other hand they have

$$\langle \sigma_x \rangle = \frac{2\lambda^2 f_1 f_2}{\omega} [\Delta_s(\omega) \tanh[\frac{\beta\omega}{2}] + \Delta_a(\omega) - \Delta_a(0)]$$
(E32)

where

$$\Delta_s(\omega) = \int_0^\infty d\Omega \,\,\omega_s(\Omega) \left(\frac{1}{\Omega + \omega} - \frac{1}{\Omega - \omega}\right) = -2 \int_0^\infty d\Omega \frac{\omega_s(\Omega)\omega}{\Omega^2 - \omega^2} \tag{E33}$$

$$\Delta_a(\omega) = \int_0^\infty d\Omega \,\,\omega_a(\Omega) \left(\frac{1}{\Omega + \omega} + \frac{1}{\Omega - \omega} \right) = 2 \int_0^\infty d\Omega \frac{\omega_a(\Omega)\Omega}{\Omega^2 - \omega^2} \tag{E34}$$

$$\Delta_a(\omega) - \Delta_a(0) = 2\int_0^\infty d\Omega \ \omega_a(\Omega) \frac{\Omega^2 - (\Omega^2 - \omega^2)}{\Omega(\Omega^2 - \omega^2)} = 2\int_0^\infty d\Omega \frac{\omega_a(\Omega)\omega^2}{\Omega(\Omega^2 - \omega^2)}$$
(E35)

such that

$$\langle \sigma_x \rangle = -\frac{4\lambda^2 f_1 f_2}{\omega} \int_0^\infty d\Omega \left(\frac{\omega_s(\Omega)\omega \tanh[\frac{\beta\omega}{2}]}{\Omega^2 - \omega^2} - \frac{\omega^2 \omega_a(\Omega)}{\Omega(\Omega^2 - \omega^2)} \right)$$
(E36)

Appendix F: Bloch-Redfield master equation (derivation)

We shall derive the Bloch-Redfield master equation in terms of $\tilde{\gamma}$ (A16) and \tilde{S} (A17) coefficients starting from the von Neumann equation:

$$\tilde{\mathcal{L}}_t^R[\tilde{\rho}(t)] = -\int_0^t ds \ \operatorname{Tr}_R[H_I(t), [H_I(s), \tilde{\rho}(t) \otimes \gamma_R]],$$
(F1)

which is derived according to the Born-Markov approximation. We expand commutators and put an explicit form of the interaction Hamiltonian (A2):

$$\begin{split} \tilde{\mathcal{L}}_{t}^{R}[\tilde{\rho}(t)] &= -\int_{0}^{t} ds \ \operatorname{Tr}_{R}[H_{I}(t), [H_{I}(s), \tilde{\rho}(t) \otimes \gamma_{R}]] = \int_{0}^{t} ds \operatorname{Tr}_{R}\left([H_{I}(s)\tilde{\rho}(t) \otimes \gamma_{R}, H_{I}(t)] - [\tilde{\rho}(t) \otimes \gamma_{R} \ H_{I}(s), H_{I}(t)]\right) \\ &= \int_{0}^{t} ds \operatorname{Tr}_{R}\left[H_{I}(s)\tilde{\rho}(t) \otimes \gamma_{R}H_{I}(t) - \tilde{\rho}(t) \otimes \gamma_{R}H_{I}(s)H_{I}(t) - H_{I}(t)H_{I}(s)\tilde{\rho}(t) \otimes \gamma_{R} + H_{I}(t)\tilde{\rho}(t) \otimes \gamma_{R}H_{I}(s)\right] \\ &= \sum_{\alpha,\beta} \int_{0}^{t} ds \left[A_{\alpha}(s)\tilde{\rho}(t)A_{\beta}(t)\langle R_{\beta}(t)R_{\alpha}(s)\rangle_{\gamma_{R}} + A_{\beta}(t)\tilde{\rho}(t)A_{\alpha}(s)\langle R_{\alpha}(s)R_{\beta}(t)\rangle_{\gamma_{R}} \\ &\quad -\tilde{\rho}(t)A_{\alpha}(s)A_{\beta}(t)\langle R_{\alpha}(s)R_{\beta}(t)\rangle_{\gamma_{R}} - A_{\beta}(t)A_{\alpha}(s)\tilde{\rho}(t)\langle R_{\beta}(t)R_{\alpha}(s)\rangle_{\gamma_{R}}\right] \\ &= \sum_{\alpha,\beta} \int_{0}^{t} ds \left[\langle R_{\alpha}(t)R_{\beta}(s)\rangle_{\gamma_{R}} \left(A_{\beta}(s)\tilde{\rho}(t)A_{\alpha}(t) - A_{\alpha}(t)A_{\beta}(s)\tilde{\rho}(t)\right)\right] + \text{h.c.} \end{split}$$

After introducing the jump operators (A4), we get

$$\tilde{\mathcal{L}}_{t}^{R}[\tilde{\rho}(t)] = \sum_{\omega,\omega'} \sum_{\alpha,\beta} \int_{0}^{t} ds \ e^{-i(\omega's+\omega t)} \langle R_{\alpha}(t)R_{\beta}(s) \rangle_{\gamma_{R}} \left(A_{\beta}(\omega')\tilde{\rho}(t)A_{\alpha}(\omega) - A_{\alpha}(\omega)A_{\beta}(\omega')\tilde{\rho}(t)\right) + \text{h.c.}$$
(F2)

$$=\sum_{\omega,\omega'}\sum_{\alpha,\beta}\int_{0}^{t}ds \ e^{i(\omega t-\omega's)}\langle R_{\alpha}(t)R_{\beta}(s)\rangle_{\gamma_{R}}\left(A_{\beta}(\omega')\tilde{\rho}(t)A_{\alpha}^{\dagger}(\omega)-A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega')\tilde{\rho}(t)\right)+\text{h.c.}$$
(F3)

$$=\sum_{\omega,\omega'}\sum_{\alpha,\beta} \tilde{\Gamma}_{\alpha\beta}(\omega,\omega',t) \left(A_{\beta}(\omega')\tilde{\rho}(t)A_{\alpha}^{\dagger}(\omega) - A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega')\tilde{\rho}(t) \right) + \text{h.c.}$$
(F4)

where we put the definition:

$$\tilde{\Gamma}_{\alpha\beta}(\omega,\omega',t) \equiv \int_0^t ds \ e^{i(\omega t - \omega' s)} \langle R_\alpha(t) R_\beta(s) \rangle_{\gamma_R}.$$
(F5)

This can be further simplified to the form:

$$\tilde{\Gamma}_{\alpha\beta}(\omega,\omega',t) = e^{i(\omega-\omega')t} \int_0^t ds \ e^{i\omega's} \langle R_\alpha(s)R_\beta(0)\rangle_{\gamma_R} \equiv e^{i(\omega-\omega')t} \ \Gamma_{\alpha\beta}(\omega',t)$$
(F6)

where we changed the variables in the integrand $s \to t-s$ and use the property $\langle R_{\alpha}(t)R_{\beta}(s)\rangle_{\gamma_{R}} = \langle R_{\alpha}(t-s)R_{\beta}(0)\rangle_{\gamma_{R}}$. Next, we rewritten the hermitian conjugate part in the form:

$$\sum_{\omega,\omega'} \sum_{\alpha,\beta} \tilde{\Gamma}^*_{\alpha\beta}(\omega,\omega',t) \left(A_\beta(\omega')\tilde{\rho}(t)A^{\dagger}_{\alpha}(\omega) - A^{\dagger}_{\alpha}(\omega)A_\beta(\omega')\tilde{\rho}(t) \right)^{\dagger}$$
(F7)

$$=\sum_{\omega,\omega'}\sum_{\alpha,\beta}\tilde{\Gamma}^*_{\alpha\beta}(\omega,\omega',t)\left(A_{\alpha}(\omega)\tilde{\rho}(t)A^{\dagger}_{\beta}(\omega')-\tilde{\rho}(t)A^{\dagger}_{\beta}(\omega')A_{\alpha}(\omega)\right)$$
(F8)

$$=\sum_{\omega,\omega'}\sum_{\alpha,\beta}\tilde{\Gamma}^*_{\beta\alpha}(\omega',\omega,t)\left(A_{\beta}(\omega')\tilde{\rho}(t)A^{\dagger}_{\alpha}(\omega)-\tilde{\rho}(t)A^{\dagger}_{\alpha}(\omega)A_{\beta}(\omega')\right)$$
(F9)

Finally, we get

$$\tilde{\mathcal{L}}_{t}^{R}[\tilde{\rho}(t)] = \sum_{\omega,\omega'} \sum_{\alpha,\beta} \left(\tilde{\Gamma}_{\alpha\beta}(\omega,\omega',t) + \tilde{\Gamma}_{\beta\alpha}^{*}(\omega',\omega,t) \right) A_{\beta}(\omega') \tilde{\rho}(t) A_{\alpha}^{\dagger}(\omega)$$
(F10)

$$-\frac{1}{2}\sum_{\omega,\omega'}\sum_{\alpha,\beta}\left(\tilde{\Gamma}_{\alpha\beta}(\omega,\omega',t)A^{\dagger}_{\alpha}(\omega)A_{\beta}(\omega')\tilde{\rho}(t)+\tilde{\Gamma}^{*}_{\beta\alpha}(\omega',\omega,t)\tilde{\rho}(t)A^{\dagger}_{\alpha}(\omega)A_{\beta}(\omega')\right)$$
(F11)

$$-\frac{1}{2}\sum_{\omega,\omega'}\sum_{\alpha,\beta} \left(\tilde{\Gamma}_{\alpha\beta}(\omega,\omega',t) A^{\dagger}_{\alpha}(\omega) A_{\beta}(\omega') \tilde{\rho}(t) + \tilde{\Gamma}^{*}_{\beta\alpha}(\omega',\omega,t) \tilde{\rho}(t) A^{\dagger}_{\alpha}(\omega) A_{\beta}(\omega') \right)$$
(F12)

$$-\frac{1}{2}\sum_{\omega,\omega'}\sum_{\alpha,\beta}\left(\tilde{\Gamma}^{*}_{\beta\alpha}(\omega',\omega,t))A^{\dagger}_{\alpha}(\omega)A_{\beta}(\omega')\tilde{\rho}(t)+\tilde{\Gamma}_{\alpha\beta}(\omega,\omega',t)\tilde{\rho}(t)A^{\dagger}_{\alpha}(\omega)A_{\beta}(\omega')\right)$$
(F13)

$$+\frac{1}{2}\sum_{\omega,\omega'}\sum_{\alpha,\beta}\left(\tilde{\Gamma}^{*}_{\beta\alpha}(\omega',\omega,t))A^{\dagger}_{\alpha}(\omega)A_{\beta}(\omega')\tilde{\rho}(t)+\tilde{\Gamma}_{\alpha\beta}(\omega,\omega',t)\tilde{\rho}(t)A^{\dagger}_{\alpha}(\omega)A_{\beta}(\omega')\right)$$
(F14)

where the last two lines sum up to zero. After rearranging terms and putting the definition (A16) and (A17), we finally obtain the master equation in the form:

$$\tilde{\mathcal{L}}_{t}^{R}[\tilde{\rho}(t)] = \sum_{\omega,\omega'} \sum_{\alpha,\beta} \left[i \tilde{\mathcal{S}}_{\alpha\beta}(\omega,\omega',t) [\tilde{\rho}(t), A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega')] + \tilde{\gamma}_{\alpha\beta}(\omega,\omega',t) \left(A_{\beta}(\omega')\tilde{\rho}(t)A_{\alpha}^{\dagger}(\omega) - \frac{1}{2} \{ A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega'), \tilde{\rho}(t) \} \right) \right].$$
(F15)

Appendix G: The cumulant equation in the Schrödinger pircture

The cumulant equation is originally derived in the interaction picture. In order to transform the cumulant equation into the Schrödinger picture we start with a simple observation.

$$\tilde{\rho}(t) = e^{i[H_0,\cdot]t} \rho(t). \tag{G1}$$

The super-operator in the r.h.s. of the equation above has its unique inverse, and $\tilde{\rho}(0) = \rho(0)$, therefore:

$$\rho(t) = e^{-i[H_0, \cdot]t} e^{\tilde{K}_t^{(2)}} \rho(0) = e^{K_t^{(2)}} \rho(0).$$
(G2)

The r.h.s. of the equation above defines the Schrödinger picture cumulant equation super-operator $K_t^{(2)}$:

$$e^{K_t^{(2)}} = e^{-i[H_0,\cdot]t} e^{\tilde{K}_t^{(2)}}.$$
(G3)

The explicit form of $K_t^{(2)}$ can be found with the aid of the Baker–Campbell–Hausdorff (BCH) formula.

$$e^{X}e^{Y} = \exp\left\{X + Y + \frac{1}{2}[X,Y] + \frac{1}{12}[X,[X,Y]] - \frac{1}{12}[Y,[X,Y]] + \cdots\right\}.$$
(G4)

We observe that in a generic case the super-operator $K_t^{(2)}$ in not of the GKSL form. This follows from the presence of multi-commutator terms in the formula (G4). These terms do not vanish, as $[K_t^{(2)}, H_0]$ is not central. Therefore, $e^{K_t^{(2)}}$ is an example of a one-parameter family of CPTP dynamical maps that are not of the GKSL form.

Appendix H: The cumulant equation in the differential form

We start this Section with the following Lemma on the properties of the derivative of an exponential map. **Lemma 1.** The derivative of the exponential map is given by

$$\frac{d}{dt}e^{X(t)} = \left(\frac{e^{[X(t),\cdot]} - \mathbb{1}}{[X(t),\cdot]}\frac{dX(t)}{dt}\right)e^{X(t)}.$$
(H1)

Proof. The proof of the above relation is identical to the proof of Theorem 5 in reference [23] up to small modifications.

Using Lemma 1 we instantly obtain the cumulant equation in the differential form:

$$\frac{d}{dt}\tilde{\rho}(t) = \left(\frac{e^{[\tilde{K}_t^{(2)},\cdot]} - \mathbb{1}}{[\tilde{K}_t^{(2)},\cdot]}\frac{d\tilde{K}_t^{(2)}}{dt}\right)\tilde{\rho}(t) = \tilde{\mathcal{L}}_t^C\tilde{\rho}(t).$$
(H2)

This result can also be obtained with integration of equation (G1). When truncated to the leading order, the above

formula reproduces the Bloch-Redfield master equation since $\frac{d\tilde{K}_t^{(2)}}{dt} = \tilde{\mathcal{L}}_t^R$. Equation (H2) can be readily transformed into the Schrödinger picture. This is done with iterative application of the $e^{\pm iH_0 t}$ operators to the jump operators $A_i(\omega)$ inside $\tilde{K}_t^{(2)}$ super-operator.

$$\frac{d}{dt}\rho(t) = \left(-i[H_0, \cdot] + e^{-i[H_0, \cdot]t} \frac{e^{[\tilde{K}_t^{(2)}, \cdot]} - \mathbb{1}}{[\tilde{K}_t^{(2)}, \cdot]} \tilde{\mathcal{L}}_t^R e^{i[H_0, \cdot]t}\right)\rho(t)$$
(H3)

$$= \left(-i[H_0, \cdot] + \frac{e^{[\bar{K}_t^{(2)}, \cdot]} - \mathbb{1}}{[\bar{K}_t^{(2)}, \cdot]} \bar{\mathcal{L}}_t^R\right) \rho(t) = \mathcal{L}_t^C \rho(t),$$
(H4)

where

$$\bar{K}_{t}^{(2)}[\rho] = \lambda^{2} \int_{0}^{t} ds \sum_{\omega,\omega'} \sum_{\alpha\beta} e^{i(\omega-\omega')(s-t)} \left(i\mathcal{S}_{\alpha\beta}(\omega,\omega',s)[\rho,A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega')] + \gamma_{\alpha\beta}(\omega,\omega',s) \left(A_{\beta}(\omega')\rho A_{\alpha}^{\dagger}(\omega) - \frac{1}{2} \{A_{\alpha}^{\dagger}(\omega)A_{\beta}(\omega'),\rho\} \right) \right),$$
(H5)

$$\bar{\mathcal{L}}_t^R = \mathcal{L}_t^R + i[H_0, \cdot]. \tag{H6}$$

Moreover, we observe that:

$$\mathcal{L}_t^C = \mathcal{L}_t^R + \mathcal{O}(\lambda^4). \tag{H7}$$

Equation (H4) can be compared with the differential form of the Schrödinger picture cumulant equation obtained with Lemma 1 and the super-operator in equation (G3).

$$\frac{d}{dt}\rho(t) = \left(\frac{e^{[K_t^{(2)},\cdot]} - \mathbb{1}}{[K_t^{(2)},\cdot]}\frac{dK_t^{(2)}}{dt}\right)\rho(t) = \mathcal{L}_t^C\rho(t).$$
(H8)

The above formula has only a formal meaning, as the $K_t^{(2)}$ super-operator does not possess a closed form formula. We present it only for the curiosity of the reader.