

An analogue of the Riemann Hypothesis via quantum walks

Norio KONNO

Department of Applied Mathematics, Faculty of Engineering
Yokohama National University
Hodogaya, Yokohama, 240-8501, Japan
e-mail: konno-norio-bt@ynu.ac.jp

arXiv:2204.00765v3 [quant-ph] 20 Jan 2024

Abstract

We consider an analogue of the well-known Riemann Hypothesis based on quantum walks on graphs with the help of the Konno-Sato theorem. Furthermore, we give some examples for complete, cycle, and star graphs.

Keywords: Zeta function, Riemann Hypothesis, Quantum walk, Grover walk, Konno-Sato theorem

1 Introduction

Inspired our recent work for a series of Zeta/Correspondence [6, 7, 8, 9, 10, 11, 12, 13, 16] on the relation between zeta functions and some models such as random walk (RW) and quantum walk (QW), we consider an analogue of the well-known Riemann Hypothesis via QWs on graphs with the help of the Konno-Sato theorem. The QW is a quantum counterpart of the RW. The Konno-Sato theorem [15] treats the Grover walk which is one of the most well-investigated QWs. Concerning QW, see [14, 17, 19, 24], as for RW, see [18, 22], and as for the Riemann Hypothesis, see [3, 4, 23], for examples. More precisely, we introduce a new zeta function $\Lambda_G^{QW}(s) = \det(\mathbf{M} - s(1-s)\mathbf{I})$ for a suitable matrix \mathbf{M} on a simple connected graph G by using the Konno-Sato theorem, where s is a complex number and \mathbf{I} is the identity matrix. Here $\Lambda_G^{QW}(s)$ corresponds the completed zeta function $\Lambda(s)$. As for the detailed definition, see Section 3. Moreover, we prove that $\Lambda_G^{QW}(s)$ satisfies the Functional Equation: $\Lambda_G^{QW}(s) = \Lambda_G^{QW}(1-s)$ and the Riemann Hypothesis: if $\rho \in \text{Zero}(\Lambda_G^{QW})$, then $\Re(\rho) = 1/2$, where $\text{Zero}(f)$ is the set of the zeros of $f(s)$ and $\Re(z)$ is the real part of z . Remark that the original Riemann Hypothesis: if $\rho \in \text{Zero}(\Lambda)$, then $\Re(\rho) = 1/2$ for the completed zeta function $\Lambda(s)$. Equivalently, if $\rho \in \text{Zero}(\zeta)$ with $0 < \Re(\rho) < 1$, then $\Re(\rho) = 1/2$ for Riemann's zeta function $\zeta(s) = \sum_{n=1}^{\infty} 1/n^s$. The present manuscript is the first step of the study on a connection between the Riemann Hypothesis and the QW.

The rest of this paper is organized as follows. Section 2 gives a brief overview of the Konno-Sato theorem. In Section 3, we explain an analogue of the Riemann Hypothesis in our setting. Section 4 presents some examples for complete, cycle, and star graphs. Finally, Section 5 is devoted to conclusion.

2 Konno-Sato Theorem

First we introduce the following notation: \mathbb{Z} is the set of integers, $\mathbb{Z}_> = \{1, 2, 3, \dots\}$, \mathbb{R} is the set of real numbers, and \mathbb{C} is the set of complex numbers.

In this section, we briefly review the Konno-Sato theorem given by [15]. This theorem treats a relation for eigenvalues between QWs and RWs. More specifically, the Grover walk (which is QW determined by the Grover matrix) with flip-flop shift type (called F-type) and simple symmetric RW (whose walker jumps to each of its nearest neighbors with equal probability) on a graph. We assume that all graphs are simple.

Let $G = (V(G), E(G))$ be a connected graph (without multiple edges and loops) with the set $V(G)$ of vertices and the set $E(G)$ of unoriented edges uv joining two vertices u and v . Moreover, let $n = |V(G)|$ and $m = |E(G)|$ be the number of vertices and edges of G , respectively. For $uv \in E(G)$, an arc (u, v) is the oriented edge from u to v . Let $D(G)$ be the symmetric digraph corresponding to G , i.e., $D(G) = \{(u, v), (v, u) \mid uv \in E(G)\}$.

For $e = (u, v) \in D(G)$, set $u = o(e)$ and $v = t(e)$. Furthermore, let $e^{-1} = (v, u)$ be the *inverse* of $e = (u, v)$. For $v \in V(G)$, the *degree* $\deg_G v = \deg v = d_v$ of v is the number of vertices adjacent to v in G . If $\deg_G v = k$ (constant) for each $v \in V(G)$, then G is called *k-regular*. A *path* P of length n in G is a sequence $P = (e_1, \dots, e_n)$ of n arcs such that $e_i \in D(G)$, $t(e_i) = o(e_{i+1})$ ($1 \leq i \leq n-1$). If $e_i = (v_{i-1}, v_i)$ for $i = 1, \dots, n$, then we write $P = (v_0, v_1, \dots, v_{n-1}, v_n)$. Put $|P| = n$, $o(P) = o(e_1)$ and $t(P) = t(e_n)$. Also, P is called an $(o(P), t(P))$ -*path*. We say that a path $P = (e_1, \dots, e_n)$ has a *backtracking* if $e_{i+1}^{-1} = e_i$ for some i ($1 \leq i \leq n-1$). A (v, w) -path is called a *v-cycle* (or *v-closed path*) if $v = w$. Let B^r be the cycle obtained by going r times around a cycle B . Such a cycle is called a *multiple* of B . A cycle C is *reduced* if both C and C^2 have no backtracking. The *Ihara zeta function* of a graph G is a function of a complex variable u with $|u|$ sufficiently small, defined by

$$\mathbf{Z}(G, u) = \exp \left(\sum_{r=1}^{\infty} \frac{N_r}{r} u^r \right),$$

where N_r is the number of reduced cycles of length r in G . Let G be a simple connected graph with n vertices v_1, \dots, v_n . The *adjacency matrix* $\mathbf{A}_n = [a_{ij}]$ is the $n \times n$ matrix such that $a_{ij} = 1$ if v_i and v_j are adjacent, and $a_{ij} = 0$ otherwise. The following result was obtained by Ihara [5] and Bass [2].

Theorem 1 (Ihara [5], Bass [2]) *Let G be a simple connected graph with $V(G) = \{v_1, \dots, v_n\}$ and m edges. Then we have*

$$\mathbf{Z}(G, u)^{-1} = (1 - u^2)^{\gamma-1} \det (\mathbf{I}_n - u\mathbf{A}_n + u^2(\mathbf{D}_n - \mathbf{I}_n)).$$

Here γ is the Betti number of G (i.e., $\gamma = m - n + 1$), \mathbf{I}_n is the $n \times n$ identity matrix, and $\mathbf{D}_n = [d_{ij}]$ is the $n \times n$ diagonal matrix with $d_{ii} = \deg v_i$ and $d_{ij} = 0$ ($i \neq j$).

Let G be a simple connected graph with $V(G) = \{v_1, \dots, v_n\}$ and m edges. Set $d_j = \deg v_j$ ($j = 1, \dots, n$). Then the $2m \times 2m$ Grover matrix $\mathbf{U}_{2m} = [U_{ef}]_{e, f \in D(G)}$ of G is defined by

$$U_{ef} = \begin{cases} 2/d_{t(f)} (= 2/d_{o(e)}) & \text{if } t(f) = o(e) \text{ and } f \neq e^{-1}, \\ 2/d_{t(f)} - 1 & \text{if } f = e^{-1}, \\ 0 & \text{otherwise.} \end{cases}$$

The discrete-time QW with the Grover matrix \mathbf{U}_{2m} as a time evolution matrix is the Grover walk with F-type on G . Then the $n \times n$ matrix $\mathbf{P}_n = [P_{uv}]_{u, v \in V(G)}$ is given by

$$P_{uv} = \begin{cases} 1/(\deg_G u) & \text{if } (u, v) \in D(G), \\ 0 & \text{otherwise.} \end{cases}$$

Note that the matrix \mathbf{P}_n is the transition probability matrix of the simple symmetric RW on G . We introduce the *positive support* $\mathbf{F}^+ = [F_{ij}^+]$ of a real matrix $\mathbf{F} = [F_{ij}]$ as follows:

$$F_{ij}^+ = \begin{cases} 1 & \text{if } F_{ij} > 0, \\ 0 & \text{otherwise.} \end{cases}$$

Ren et al. [20] showed that the Perron-Frobenius operator (or edge matrix) of a graph is the positive support $({}^T\mathbf{U}_{2m})^+$ of the transpose of its Grover matrix \mathbf{U}_{2m} , i.e.,

$$\mathbf{Z}(G, u)^{-1} = \det (\mathbf{I}_{2m} - u({}^T\mathbf{U}_{2m})^+) = \det (\mathbf{I}_{2m} - u\mathbf{U}_{2m}^+).$$

The Ihara zeta function of a graph G is just a zeta function on the positive support of the Grover matrix of G . That is, the Ihara zeta function corresponds to the positive support

version of the Grover walk (defined by the positive support of the Grover matrix \mathbf{U}_{2m}^+) with F-type on G .

Now we propose another zeta function of a graph. Let G be a simple connected graph with m edges. Then we define a zeta function $\overline{\mathbf{Z}}(G, u)$ of G satisfying

$$\overline{\mathbf{Z}}(G, u)^{-1} = \det(\mathbf{I}_{2m} - u\mathbf{U}_{2m}).$$

In other words, this zeta function corresponds to the Grover walk (defined by the Grover matrix \mathbf{U}_{2m}) with F-type on G .

In this setting, Konno and Sato [15] presented the following result which is called the *Konno-Sato theorem*.

Theorem 2 (Konno and Sato [15]) *Let G be a simple connected graph with n vertices and m edges. Then*

$$\overline{\mathbf{Z}}(G, u)^{-1} = \det(\mathbf{I}_{2m} - u\mathbf{U}_{2m}) = (1 - u^2)^{m-n} \det((1 + u^2)\mathbf{I}_n - 2u\mathbf{P}_n). \quad (1)$$

If we take $u = 1/\lambda$, then Eq. (1) implies

$$\det(\lambda\mathbf{I}_{2m} - \mathbf{U}_{2m}) = (\lambda^2 - 1)^{m-n} \det((\lambda^2 + 1)\mathbf{I}_n - 2\lambda\mathbf{P}_n). \quad (2)$$

Furthermore, Eq. (2) can be rewritten as

$$\det(\lambda\mathbf{I}_{2m} - \mathbf{U}_{2m}) = (\lambda^2 - 1)^{m-n} \times \prod_{\lambda_{\mathbf{P}_n} \in \text{Spec}(\mathbf{P}_n)} (\lambda^2 + 1 - 2\lambda_{\mathbf{P}_n}\lambda), \quad (3)$$

where $\text{Spec}(\mathbf{B})$ is the set of eigenvalues of a square matrix \mathbf{B} . More precisely, we also use the following notation:

$$\text{Spec}(\mathbf{B}) = \left\{ [\lambda_1]^{l_1}, [\lambda_2]^{l_2}, \dots, [\lambda_k]^{l_k} \right\},$$

where λ_j is the eigenvalue of \mathbf{B} and $l_j \in \mathbb{Z}_{>}$ is the multiplicity of λ_j for $j = 1, 2, \dots, k$. Set $|\text{Spec}(\mathbf{B})| = l_1 + l_2 + \dots + l_k$. It follows from $\lambda^2 + 1 - 2\lambda_{\mathbf{P}_n}\lambda = 0$ that $\lambda_{\mathbf{U}_{2m}} \in \text{Spec}(\mathbf{U}_{2m})$ is given by

$$\lambda_{\mathbf{U}_{2m}} = \lambda_{\mathbf{P}_n} \pm i\sqrt{1 - \lambda_{\mathbf{P}_n}^2}. \quad (4)$$

Remark that $\lambda_{\mathbf{P}_n} \in [-1, 1]$. Noting Eqs. (3) and (4), we introduce $\text{Spec}(\mathbf{U}_{2m} : \text{RW})$ and $\text{Spec}(\mathbf{U}_{2m} : \text{RW}^c)$ as follows:

$$\begin{aligned} & \text{Spec}(\mathbf{U}_{2m} : \text{RW}) \\ &= \left\{ \left[\lambda_{\mathbf{P}_n} + i\sqrt{1 - \lambda_{\mathbf{P}_n}^2} \right]^1, \left[\lambda_{\mathbf{P}_n} - i\sqrt{1 - \lambda_{\mathbf{P}_n}^2} \right]^1 : \lambda_{\mathbf{P}_n} \in \text{Spec}(\mathbf{P}_n) \right\}, \\ & \text{Spec}(\mathbf{U}_{2m} : \text{RW}^c) = \left\{ [1]^{|m-n|}, [-1]^{|m-n|} \right\}. \end{aligned}$$

When $m = n$, we let $\text{Spec}(\mathbf{U}_{2m} : \text{RW}^c) = \emptyset$. Note that $|\text{Spec}(\mathbf{U}_{2m} : \text{RW})| = 2n$ and $|\text{Spec}(\mathbf{U}_{2m} : \text{RW}^c)| = 2|m - n|$. We should remark that $\text{Spec}(\mathbf{U}_{2m} : \text{RW})$ corresponds to the eigenvalue of \mathbf{P}_n which is the transition probability matrix of the simple symmetric RW on G . On the other hand, $\text{Spec}(\mathbf{U}_{2m} : \text{RW}^c)$ does not corresponds to the RW, so superscript ‘‘c’’ (of RW^c) stands for ‘‘complement’’. Therefore we obtain

Corollary 1 *Let G be a simple connected graph with n vertices and m edges.*

(i) *If $m > n$, then*

$$\text{Spec}(\mathbf{U}_{2m}) = \text{Spec}(\mathbf{U}_{2m} : \text{RW}) \cup \text{Spec}(\mathbf{U}_{2m} : \text{RW}^c).$$

(ii) *If $m = n$, then*

$$\text{Spec}(\mathbf{U}_{2m}) = \text{Spec}(\mathbf{U}_{2m} : \text{RW}).$$

(iii) *If $m < n$, then*

$$\text{Spec}(\mathbf{U}_{2m}) = \text{Spec}(\mathbf{U}_{2m} : \text{RW}) \setminus \text{Spec}(\mathbf{U}_{2m} : \text{RW}^c).$$

In Section 4, we will give some examples for each case. Since \mathbf{U}_{2m} is unitary, $\lambda \in \text{Spec}(\mathbf{U}_{2m})$ satisfies $|\lambda| = 1$, so we put $\lambda = e^{i\theta}$ ($\theta \in [0, 2\pi)$). Thus we get

$$\cos \theta = \frac{\lambda + \bar{\lambda}}{2}, \quad (5)$$

where $\bar{\lambda}$ is the complex conjugate of $\lambda \in \mathbb{C}$. This is called the Joukowski transform. It follows from Eq. (5) and $\lambda^{-1} = \bar{\lambda}$ that Eq. (2) becomes

$$\det(\lambda \mathbf{I}_{2m} - \mathbf{U}_{2m}) = (\lambda^2 - 1)^{m-n} (2\lambda)^n \det(\cos \theta \cdot \mathbf{I}_n - \mathbf{P}_n).$$

Therefore we have a relation “ $\cos \theta \in \text{Spec}(\mathbf{P}_n)$ ” \implies “ $\lambda = e^{i\theta} \in \text{Spec}(\mathbf{U}_{2m})$ ” which is sometimes called the *spectral mapping theorem* in the study of QW (see [21], for example). We will explain in a more detailed fashion. We set

$$\text{Spec}(\mathbf{P}_n) = \left\{ [\cos \theta_1]^{l_1}, [\cos \theta_2]^{l_2}, \dots, [\cos \theta_p]^{l_p} \right\},$$

where $|\text{Spec}(\mathbf{P}_n)| = l_1 + l_2 + \dots + l_p = n$ and $0 = \theta_1 < \theta_2 < \dots < \theta_p < 2\pi$. Then we get

$$\begin{aligned} \text{Spec}(\mathbf{U}_{2m} : \text{RW}) \\ = \left\{ [e^{i\theta_1}]^{l_1}, [e^{-i\theta_1}]^{l_1}, [e^{i\theta_2}]^{l_2}, [e^{-i\theta_2}]^{l_2}, \dots, [e^{i\theta_p}]^{l_p}, [e^{-i\theta_p}]^{l_p} \right\}. \end{aligned} \quad (6)$$

Remark that $|\text{Spec}(\mathbf{U}_{2m} : \text{RW})| = 2(l_1 + l_2 + \dots + l_p) = 2n$.

3 Analogue of the Riemann Hypothesis

The *Riemann zeta function* for $s \in \mathbb{C}$ with $\Re(s) > 1$ is defined by

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_{p:\text{prime}} (1 - p^{-s})^{-1},$$

where $\Re(z)$ is the real part of $z \in \mathbb{C}$. It is known that $\zeta(s)$ has a meromorphic continuation to the entire complex plane. Moreover the *completed zeta function* $\Lambda(s)$ is given by

$$\Lambda(s) = \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \zeta(s),$$

where $\Gamma(z)$ is the gamma function (see [1], for example). Then $\Lambda(s)$ satisfies the **Functional Equation**:

$$\Lambda(s) = \Lambda(1 - s).$$

The **Riemann Hypothesis**, which is stated in terms of Λ , is the following:

$$\Re(\rho) = \frac{1}{2} \quad (\rho \in \text{Zero}(\Lambda)),$$

where

$$\text{Zero}(\Lambda) = \{s \in \mathbb{C} : \Lambda(s) = 0\}.$$

Concerning the Riemann Hypothesis, see [3, 4, 23], for instance. In this background, we want to find the following zeta function $\Lambda_G^{QW}(s)$ via the QW for a graph G as a counterpart of completed zeta function $\Lambda(s)$ such as

(i) **Functional Equation**:

$$\Lambda_G^{QW}(s) = \Lambda_G^{QW}(1 - s).$$

(ii) **Riemann Hypothesis**:

$$\Re(\rho) = \frac{1}{2} \quad (\rho \in \text{Zero}(\Lambda_G^{QW})),$$

where

$$\text{Zero}(\Lambda_G^{QW}) = \{s \in \mathbb{C} : \Lambda_G^{QW}(s) = 0\}.$$

That is,

$$\text{if } \Lambda_G^{QW}(\rho) = 0, \quad \text{then } \Re(\rho) = \frac{1}{2}.$$

To do so, we use the Konno-Sato theorem as described below. We assume that G is a simple connected graph with n vertices and m edges. Put $\lambda \in \text{Spec}(\mathbf{U}_{2m})$, where $\lambda = e^{i\theta}$ with $\theta \in [0, 2\pi)$. In the complex plane, λ corresponds to a point $(\cos \theta, \sin \theta)$. Let ℓ be the line through the points $(\cos \theta, \sin \theta)$ and $(1, 0)$. If we consider the complex plane as xy plane, then the equation for line ℓ becomes

$$y = -\frac{\sin \theta}{1 - \cos \theta} (x - 1) \quad (\theta \in (0, 2\pi)). \quad (7)$$

When $x = 1/2$, Eq. (7) yields the value

$$y = \frac{1}{2} \cot\left(\frac{\theta}{2}\right),$$

for $\theta \in (0, 2\pi)$, so the point $(1/2, \cot(\theta/2)/2)$ lies on ℓ . Thus, $\cot(\theta/2)/2$ is the y coordinate of the intersection of ℓ and $x = 1/2$. Therefore, we want to relate $\lambda = e^{i\theta} \in \text{Spec}(\mathbf{U}_{2m})$ with $\theta \in (0, 2\pi)$ to $\rho = \rho(\theta) \in \text{Zero}(\Lambda_G^{QW})$ in the following way:

$$\rho = \rho(\theta) = \frac{1}{2} + \frac{i}{2} \cot\left(\frac{\theta}{2}\right) \quad (\theta \in (0, 2\pi)). \quad (8)$$

On the other hand, if $\theta = 0$, then we put

$$\rho = \rho(0) = \frac{1}{2} + i \cdot (+\infty). \quad (9)$$

Noting Eqs. (6), (8) and (9), we introduce the following sets $\text{Zero}(\Lambda_G^{QW} : \text{RW})$ and $\text{Zero}(\Lambda_G^{QW} : \text{RW}^c)$ corresponding to $\text{Spec}(\mathbf{U}_{2m} : \text{RW})$ and $\text{Spec}(\mathbf{U}_{2m} : \text{RW}^c)$, respectively.

$$\begin{aligned} & \text{Zero}(\Lambda_G^{QW} : \text{RW}) \\ &= \left\{ [\rho(\theta_1)]^{l_1}, [\rho(-\theta_1)]^{l_1}, [\rho(\theta_2)]^{l_2}, [\rho(-\theta_2)]^{l_2}, \dots, [\rho(\theta_p)]^{l_p}, [\rho(-\theta_p)]^{l_p} \right\} \\ &= \left\{ \left[\frac{1}{2} + \frac{i}{2} \cot\left(\frac{\theta_q}{2}\right) \right]^{l_q}, \left[\frac{1}{2} - \frac{i}{2} \cot\left(\frac{\theta_q}{2}\right) \right]^{l_q} : q = 1, 2, \dots, p \right\}, \\ & \text{Zero}(\Lambda_G^{QW} : \text{RW}^c) \\ &= \left\{ [\rho(0)]^{|m-n|}, [\rho(\pi)]^{|m-n|} \right\} = \left\{ \left[\frac{1}{2} + i \cdot (+\infty) \right]^{|m-n|}, \left[\frac{1}{2} \right]^{|m-n|} \right\}. \end{aligned}$$

Remark that $|\text{Zero}(\Lambda_G^{QW} : \text{RW})| = 2(l_1 + l_2 + \dots + l_p) = 2n$ and $|\text{Zero}(\Lambda_G^{QW} : \text{RW}^c)| = 2|m - n|$. Furthermore, we introduce $\text{Zero}(\Lambda_G^{QW})$ as follows:

(i) If $m > n$, then

$$\begin{aligned} \text{Zero}(\Lambda_G^{QW}) &= \text{Zero}(\Lambda_G^{QW} : \text{RW}) \cup \text{Zero}(\Lambda_G^{QW} : \text{RW}^c) \\ &= \left\{ \left[\frac{1}{2} + \frac{i}{2} \cot\left(\frac{\theta_q}{2}\right) \right]^{l_q}, \left[\frac{1}{2} - \frac{i}{2} \cot\left(\frac{\theta_q}{2}\right) \right]^{l_q} : q = 1, 2, \dots, p \right\} \\ &\cup \left\{ \left[\frac{1}{2} + i \cdot (+\infty) \right]^{m-n}, \left[\frac{1}{2} \right]^{m-n} \right\}. \end{aligned}$$

(ii) If $m = n$, then

$$\begin{aligned} \text{Zero}(\Lambda_G^{QW}) &= \text{Zero}(\Lambda_G^{QW} : \text{RW}) \\ &= \left\{ \left[\frac{1}{2} + \frac{i}{2} \cot\left(\frac{\theta_q}{2}\right) \right]^{l_q}, \left[\frac{1}{2} - \frac{i}{2} \cot\left(\frac{\theta_q}{2}\right) \right]^{l_q} : q = 1, 2, \dots, p \right\}. \end{aligned}$$

(iii) If $m < n$, then

$$\begin{aligned} \text{Zero}(\Lambda_G^{QW}) &= \text{Zero}(\Lambda_G^{QW} : \text{RW}) \setminus \text{Zero}(\Lambda_G^{QW} : \text{RW}^c) \\ &= \left\{ \left[\frac{1}{2} + \frac{i}{2} \cot\left(\frac{\theta_q}{2}\right) \right]^{l_q}, \left[\frac{1}{2} - \frac{i}{2} \cot\left(\frac{\theta_q}{2}\right) \right]^{l_q} : q = 1, 2, \dots, p \right\} \\ &\setminus \left\{ \left[\frac{1}{2} + i \cdot (+\infty) \right]^{n-m}, \left[\frac{1}{2} \right]^{n-m} \right\}. \end{aligned}$$

Noting Eq. (8), we define $\Lambda_G^{QW}(s)$ by

$$\Lambda_G^{QW}(s) = \det(\mathbf{M}_n - s(1-s)\mathbf{I}_n), \quad (10)$$

for a suitable $n \times n$ matrix \mathbf{M}_n . By Eq. (10), we easily confirm that $\Lambda_G^{QW}(s)$ satisfies $\Lambda_G^{QW}(s) = \Lambda_G^{QW}(1-s)$, i.e., Functional Equation. On the other hand, $\Lambda_G^{QW}(s)$ can be rewritten as

$$\Lambda_G^{QW}(s) = \prod_{\lambda_{\mathbf{M}_n} \in \text{Spec}(\mathbf{M}_n)} (\lambda_{\mathbf{M}_n} - s(1-s)). \quad (11)$$

We recall that it follows from Eq. (8) that any $\rho \in \text{Zero}(\Lambda_G^{QW})$ should be expressed as

$$\rho = \frac{1}{2} + \frac{i}{2} \cot\left(\frac{\theta}{2}\right). \quad (12)$$

If so, then $\Lambda_G^{QW}(s)$ satisfies the Riemann Hypothesis, that is, if $\rho \in \text{Zero}(\Lambda_G^{QW})$, then $\Re(\rho) = 1/2$. Next noting Eq. (11), we need to confirm that $\rho \in \text{Zero}(\Lambda_G^{QW})$ with Eq. (12) satisfies

$$s^2 - s + \lambda_{\mathbf{M}_n} = 0,$$

for $\lambda_{\mathbf{M}_n} \in \text{Spec}(\mathbf{M}_n)$. This is,

$$(s - \rho)(s - \bar{\rho}) = s^2 - s + \lambda_{\mathbf{M}_n}.$$

So we have

$$\rho + \bar{\rho} = 1 \quad (13)$$

$$\rho\bar{\rho} = \lambda_{\mathbf{M}_n}. \quad (14)$$

Then Eq. (13) comes from Eq. (12). Concerning Eq. (14), by using Eq. (12), we compute

$$\begin{aligned} \rho\bar{\rho} &= \left(\frac{1}{2} + \frac{i}{2} \cot\left(\frac{\theta}{2}\right)\right) \left(\frac{1}{2} - \frac{i}{2} \cot\left(\frac{\theta}{2}\right)\right) \\ &= \frac{1}{4} \left(1 + \cot^2\left(\frac{\theta}{2}\right)\right) = \frac{1}{2} \frac{1}{1 - \cos\theta} \\ &= \frac{1}{2} \frac{1}{1 - \lambda_{\mathbf{P}_n}}, \end{aligned}$$

where $\theta \in (0, 2\pi)$. Thus we have

$$\rho\bar{\rho} = \frac{1}{2} \frac{1}{1 - \lambda_{\mathbf{P}_n}}. \quad (15)$$

Combining Eq. (14) with Eq. (15) yields

$$\lambda_{\mathbf{M}_n} = \frac{1}{2} (1 - \lambda_{\mathbf{P}_n})^{-1}. \quad (16)$$

Then Eq. (16) suggests that \mathbf{M}_n may be expressed as

$$\mathbf{M}_n = \frac{1}{2} (\mathbf{I}_n - \mathbf{P}_n)^{-1}. \quad (17)$$

Therefore we define $\text{Spec}(\mathbf{M}_n)$ by

$$\text{Spec}(\mathbf{M}_n) = \left\{ \left[\frac{1}{2} \frac{1}{1 - \cos\theta_1} \right]^{l_1}, \left[\frac{1}{2} \frac{1}{1 - \cos\theta_2} \right]^{l_2}, \dots, \left[\frac{1}{2} \frac{1}{1 - \cos\theta_p} \right]^{l_p} \right\},$$

since

$$\text{Spec}(\mathbf{P}_n) = \left\{ [\cos \theta_1]^{l_1}, [\cos \theta_2]^{l_2}, \dots, [\cos \theta_p]^{l_p} \right\}.$$

Remark that $|\text{Spec}(\mathbf{M}_n)| = l_1 + l_2 + \dots + l_p = n$ and $0 = \theta_1 < \theta_2 < \dots < \theta_p < 2\pi$. In particular, when $\cos \theta_1 = 1$ (i.e., $\theta_1 = 0$), we put

$$\frac{1}{2} \frac{1}{1 - \cos \theta_1} = +\infty.$$

This matrix “ \mathbf{M}_n ” is exactly what we wanted to find. However, $(\mathbf{I}_n - \mathbf{P}_n)^{-1}$ does not exist, so we understand Eq. (17) “in the sense of the eigenvalue”, that is, as Eq. (16). In fact, we see that $0 \in \text{Spec}(\mathbf{I}_n - \mathbf{P}_n)$, since all one vector $\mathbf{1}_n$ is the eigenvector corresponding to eigenvalue 0, i.e., $(\mathbf{I}_n - \mathbf{P}_n)\mathbf{1}_n = 0 \cdot \mathbf{1}_n$. Thus $\mathbf{I}_n - \mathbf{P}_n$ is not invertible. Therefore Eq. (17) implies that \mathbf{M}_n does not exist. If G is $(q+1)$ -regular, then we have the Laplacian Δ_n as

$$\Delta_n = \mathbf{D}_n - \mathbf{A}_n = (q+1)(\mathbf{I}_n - \mathbf{P}_n). \quad (18)$$

It follows from Eq. (18) that Eq. (17) becomes

$$\mathbf{M}_n = \frac{q+1}{2} \Delta_n^{-1}$$

in the sense of the eigenvalue. We also remark that Δ_n^{-1} does not exist. Then we obtain the following main result.

Theorem 3 *Let G be a simple connected graph with n vertices and m edges. Put*

$$\Lambda_G^{QW}(s) = \det(\mathbf{M}_n - s(1-s)\mathbf{I}_n),$$

where

$$\mathbf{M}_n = \frac{1}{2} (\mathbf{I}_n - \mathbf{P}_n)^{-1}$$

in the sense of the eigenvalue. Then $\Lambda_G^{QW}(s)$ satisfies

(i) *Functional Equation:*

$$\Lambda_G^{QW}(s) = \Lambda_G^{QW}(1-s).$$

(ii) *Riemann Hypothesis:*

$$\Re(\rho) = \frac{1}{2} \quad \text{for any } \rho \in \text{Zero}(\Lambda_G^{QW}),$$

where $\text{Zero}(\Lambda_G^{QW}) = \{s \in \mathbb{C} : \Lambda_G^{QW}(s) = 0\}$.

4 Example

This section gives examples, i.e., (i) $m > n$, (ii) $m = n$, and (iii) $m < n$ cases. The results here can be obtained by the direct computation and the Konno-Sato theorem.

(i) $m > n$ case. $G = K_n$ (complete graph) with n vertices and $m = n(n-1)/2$ edges for $n \geq 4$. We should remark that if $n = 3$, then (ii) $m = n = 3$ case. If $n = 2$, then (iii) $m = 1 < n = 2$ case. Then we get

$$\begin{aligned} \text{Spec}(\mathbf{P}_n) &= \left\{ [1]^1, \left[-\frac{1}{n-1} \right]^{n-1} \right\}, \\ \text{Spec}(\mathbf{U}_{n(n-1)}) &= \left\{ [1]^{(n(n-3)+4)/2}, [-1]^{n(n-3)/2}, \right. \\ &\quad \left. \left[\frac{-1+i\sqrt{n(n-2)}}{n-1} \right]^{n-1}, \left[\frac{-1-i\sqrt{n(n-2)}}{n-1} \right]^{n-1} \right\}. \end{aligned}$$

Remark that we confirm that spectral mapping theorem holds such as

$$\Re \left[\frac{-1+i\sqrt{n(n-2)}}{n-1} \right] = \Re \left[\frac{-1-i\sqrt{n(n-2)}}{n-1} \right] = -\frac{1}{n-1}.$$

Therefore we have

$$\begin{aligned} \text{Zero}(\Lambda_{K_n}^{QW}) &= \left\{ \left[\frac{1}{2} + i \cdot (+\infty) \right]^2, \left[\frac{1}{2} + \frac{i}{2} \sqrt{1-\frac{2}{n}} \right]^{n-1}, \left[\frac{1}{2} - \frac{i}{2} \sqrt{1-\frac{2}{n}} \right]^{n-1} \right\} \\ &\cup \left\{ \left[\frac{1}{2} + i \cdot (+\infty) \right]^{n(n-3)/2}, \left[\frac{1}{2} \right]^{n(n-3)/2} \right\}. \end{aligned}$$

Note that $m - n = n(n-3)/2$. Furthermore, we obtain

$$\text{Spec}(\mathbf{M}_n) = \left\{ [+\infty]^1, \left[\frac{1}{2} \left(1 - \frac{1}{n} \right) \right]^{n-1} \right\},$$

for $n \geq 4$. Formally, if we take $n \rightarrow \infty$, then

$$\lim_{n \rightarrow \infty} \text{Spec}(\mathbf{M}_n) = \left\{ [+\infty]^1, \left[\frac{1}{2} \right]^{+\infty} \right\}.$$

(ii) $m = n$ case. $G = C_n$ (cycle graph) with n vertices and $m = n$ edges. In this case, $2n \times 2n$ matrix \mathbf{U}_{2n} and $n \times n$ matrix \mathbf{P}_n are expressed as follows (see [7, 8], for example):

$$\mathbf{U}_{2n} = \begin{bmatrix} O & P & O & \dots & \dots & O & Q \\ Q & O & P & O & \dots & \dots & O \\ O & Q & O & P & O & \dots & O \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ O & \dots & O & Q & O & P & O \\ O & \dots & \dots & O & Q & O & P \\ P & O & \dots & \dots & O & Q & O \end{bmatrix}, \quad \mathbf{P}_n = \frac{1}{2} \begin{bmatrix} 0 & 1 & 0 & \dots & \dots & 0 & 1 \\ 1 & 0 & 1 & 0 & \dots & \dots & 0 \\ 0 & 1 & 0 & 1 & 0 & \dots & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & 1 & 0 & 1 & 0 \\ 0 & \dots & \dots & 0 & 1 & 0 & 1 \\ 1 & 0 & \dots & \dots & 0 & 1 & 0 \end{bmatrix},$$

where

$$P = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad Q = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \quad O = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Let $\xi_k = 2\pi k/n$ for $k = 0, 1, \dots, n-1$. Thus we get

$$\begin{aligned}\text{Spec}(\mathbf{P}_n) &= \{[\cos \xi_k]^1 : k = 0, 1, \dots, n-1\}, \\ \text{Spec}(\mathbf{U}_{2n}) &= \{[e^{i\xi_k}]^1, [e^{-i\xi_k}]^1 : k = 0, 1, \dots, n-1\}.\end{aligned}$$

Therefore we obtain

$$\text{Zero}(\Lambda_{C_n}^{QW}) = \left\{ \left[\frac{1}{2} + \frac{i}{2} \cot\left(\frac{\xi_k}{2}\right) \right]^1, \left[\frac{1}{2} - \frac{i}{2} \cot\left(\frac{\xi_k}{2}\right) \right]^1 : k = 0, 1, \dots, n-1 \right\}.$$

Formally, if we take $n \rightarrow \infty$, then

$$\lim_{n \rightarrow \infty} \text{Zero}(\Lambda_{C_n}^{QW}) = \left\{ \left[\frac{1}{2} + i \cdot \gamma \right]^1 : \gamma \in \mathbb{R} \right\}.$$

Moreover we get

$$\text{Spec}(\mathbf{M}_n) = \left\{ [+ \infty]^1, \left[\frac{1}{2} \frac{1}{1 - \cos \xi_1} \right]^1, \dots, \left[\frac{1}{2} \frac{1}{1 - \cos \xi_{n-1}} \right]^1 \right\},$$

since $\xi_0 = 0$ (i.e., $\cos \xi_0 = 1$). From now on, we consider $n = 3$ and $n = 4$ cases. First we deal with $n = 3$. Then we have

$$\begin{aligned}\text{Spec}(\mathbf{P}_3) &= \{[\cos(0 \cdot \pi/3)]^1, [\cos(2 \cdot \pi/3)]^1, [\cos(4 \cdot \pi/3)]^1\} = \left\{ [1]^1, \left[-\frac{1}{2} \right]^2 \right\}, \\ \text{Spec}(\mathbf{U}_6) &= \left\{ [e^{i(0 \cdot \pi/3)}]^2, [e^{i(2 \cdot \pi/3)}]^2, [e^{i(4 \cdot \pi/3)}]^2 \right\} \\ &= \left\{ [1]^2, \left[\frac{-1 + i\sqrt{3}}{2} \right]^2, \left[\frac{-1 - i\sqrt{3}}{2} \right]^2 \right\}.\end{aligned}$$

Thus we obtain

$$\text{Zero}(\Lambda_{C_3}^{QW}) = \left\{ \left[\frac{1}{2} + i \cdot (+\infty) \right]^2, \left[\frac{1}{2} + \frac{i\sqrt{3}}{6} \right]^2, \left[\frac{1}{2} - \frac{i\sqrt{3}}{6} \right]^2 \right\}.$$

Note that

$$\text{Spec}(\mathbf{M}_3) = \left\{ [+ \infty]^1, \left[\frac{1}{3} \right]^2 \right\}, \quad \text{Spec}(\mathbf{\Delta}_3) = \{[0]^1, [3]^2\},$$

where

$$\mathbf{\Delta}_3 = 2(\mathbf{I}_3 - \mathbf{P}_3) = \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix}.$$

So we confirm that $\mathbf{M}_3 = \mathbf{\Delta}_3^{-1}$ does not exist. As in the case of $n = 3$, we treat $n = 4$ case. Then we get

$$\begin{aligned}\text{Spec}(\mathbf{P}_4) &= \{[\cos(0 \cdot \pi/4)]^1, [\cos(2 \cdot \pi/4)]^1, [\cos(4 \cdot \pi/4)]^1, [\cos(6 \cdot \pi/4)]^1\} \\ &= \{[1]^1, [0]^2, [-1]^1\}, \\ \text{Spec}(\mathbf{U}_8) &= \left\{ [e^{i(0 \cdot \pi/4)}]^2, [e^{i(2 \cdot \pi/4)}]^2, [e^{i(4 \cdot \pi/4)}]^2, [e^{i(6 \cdot \pi/4)}]^2 \right\} \\ &= \{[1]^2, [i]^2, [-1]^2, [-i]^2\}.\end{aligned}$$

Therefore we have

$$\text{Zero} \left(\Lambda_{C_4}^{QW} \right) = \left\{ \left[\frac{1}{2} + i \cdot (+\infty) \right]^2, \left[\frac{1}{2} + \frac{i}{2} \right]^2, \left[\frac{1}{2} \right]^2, \left[\frac{1}{2} - \frac{i}{2} \right]^2 \right\}.$$

Remark that

$$\text{Spec}(\mathbf{M}_4) = \left\{ [+ \infty]^1, \left[\frac{1}{2} \right]^2, \left[\frac{1}{4} \right]^1 \right\}, \quad \text{Spec}(\mathbf{\Delta}_4) = \{ [0]^1, [2]^2, [4]^1 \},$$

where

$$\mathbf{\Delta}_4 = 2(\mathbf{I}_4 - \mathbf{P}_4) = \begin{bmatrix} 2 & -1 & 0 & -1 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ -1 & 0 & -1 & 2 \end{bmatrix}.$$

Thus we confirm that $\mathbf{M}_4 = \mathbf{\Delta}_4^{-1}$ does not exist.

(iii) $m < n$ case. $G = S_n$ (star graph) with n vertices and $m = n - 1$ edges. Remark that S_n is isomorphic to the complete bipartite graph $K_{1,n-1}$. Then we obtain

$$\begin{aligned} \text{Spec}(\mathbf{P}_n) &= \{ [1]^1, [0]^{n-2}, [-1]^1 \}, \\ \text{Spec}(\mathbf{U}_{2(n-1)}) &= \{ [1]^2, [i]^{n-2}, [-i]^{n-2}, [-1]^2 \} \setminus \{ [1]^1, [-1]^1 \}. \\ &= \{ [1]^1, [i]^{n-2}, [-i]^{n-2}, [-1]^1 \}. \end{aligned}$$

Thus we have

$$\text{Zero} \left(\Lambda_{S_n}^{QW} \right) = \left\{ \left[\frac{1}{2} + i \cdot (+\infty) \right]^1, \left[\frac{1}{2} + \frac{i}{2} \right]^{n-2}, \left[\frac{1}{2} - \frac{i}{2} \right]^{n-2}, \left[\frac{1}{2} \right]^1 \right\}.$$

Note that

$$\text{Spec}(\mathbf{M}_n) = \left\{ [+ \infty]^1, \left[\frac{1}{2} \right]^{n-2}, \left[\frac{1}{4} \right]^1 \right\}, \quad \text{Spec}(\mathbf{\Delta}_n) = \{ [0]^1, [2]^{n-2}, [4]^1 \}.$$

5 Conclusion

In this paper, we introduced a new zeta function $\Lambda_G^{QW}(s) = \det(\mathbf{M}_n - s(1-s)\mathbf{I}_n)$ for a suitable $n \times n$ matrix \mathbf{M}_n on a simple connected graph G with n vertices via the QW by the help of the Konno-Sato theorem. In addition, we showed that $\Lambda_G^{QW}(s)$ satisfies the Functional Equation: $\Lambda_G^{QW}(s) = \Lambda_G^{QW}(1-s)$ and the Riemann Hypothesis: if $\rho \in \text{Zero}(\Lambda_G^{QW})$, then $\Re(\rho) = 1/2$. The challenging problem, of course, is to prove the original Riemann Hypothesis: if $\rho \in \text{Zero}(\Lambda)$, then $\Re(\rho) = 1/2$, by using our approach based on the QW. Moreover, one of the interesting problems might be to clarify the relation between our zeta function $\Lambda_G^{QW}(s)$ and other zeta functions for a graph G .

References

- [1] Andrews, G. E., Askey, R., Roy, R.: Special Functions. Cambridge University Press (1999)

- [2] Bass, H.: The Ihara-Selberg zeta function of a tree lattice. *Internat. J. Math.* **3**, 717-797 (1992)
- [3] Edwards, H. M.: *Riemann's Zeta Function*. Academic Press (1974)
- [4] Gelbart, S. S., Miller, S. D.: Riemann's zeta function and beyond. *Bull. Amer. Math. Soc.* **41**, 59–112 (2003)
- [5] Ihara, Y.: On discrete subgroups of the two by two projective linear group over p -adic fields. *J. Math. Soc. Japan* **18**, 219–235 (1966)
- [6] Komatsu, T., Konno, N., Sato, I.: A note on the Grover walk and the generalized Ihara zeta function of the one-dimensional integer lattice. *Yokohama Math. J.* **67**, 115–123 (2021).
- [7] Komatsu, T., Konno, N., Sato, I.: Grover/Zeta Correspondence based on the Konno-Sato theorem. *Quantum Inf. Process.* **20**, 268 (2021)
- [8] Komatsu, T., Konno, N., Sato, I.: Walk/Zeta Correspondence. arXiv:2104.10287 (2021)
- [9] Komatsu, T., Konno, N., Sato, I.: IPS/Zeta Correspondence. *Quantum Inf. Comput.* **22**, 251–269 (2022)
- [10] Komatsu, T., Konno, N., Sato, I.: Vertex-Face/Zeta Correspondence. *J. Algebraic Combin.* (in press) arXiv:2107.03300 (2021)
- [11] Komatsu, T., Konno, N., Sato, I.: CTM/Zeta Correspondence. *Quantum Stud.: Math. Found.* **9**, 165–173 (2022)
- [12] Komatsu, T., Konno, N., Sato, I., Tamura, S.: A Generalized Grover/Zeta Correspondence. arXiv:2201.03973 (2022)
- [13] Komatsu, T., Konno, N., Sato, I., Tamura, S.: Mahler/Zeta Correspondence. arXiv:2202.05966 (2022)
- [14] Konno, N.: Quantum Walks. In: *Quantum Potential Theory*, Franz, U., and Schurmann, M., Eds., *Lecture Notes in Mathematics: Vol. 1954*, pp.309–452, Springer-Verlag, Heidelberg (2008)
- [15] Konno, N., Sato, I.: On the relation between quantum walks and zeta functions. *Quantum Inf. Process.* **11**, 341–349 (2012)
- [16] Konno, N., Tamura, S.: Walk/Zeta Correspondence for quantum and correlated random walks. *Yokohama Math. J.* **67**, 125–152 (2021)
- [17] Manouchehri, K., Wang, J.: *Physical Implementation of Quantum Walks*. Springer, New York (2014)
- [18] Norris, J. R.: *Markov Chains*. Cambridge University Press, Cambridge (1997)
- [19] Portugal, R.: *Quantum Walks and Search Algorithms*, 2nd edition. Springer, New York (2018)
- [20] Ren, P., Aleksic, T., Emms, D., Wilson, R. C., Hancock, E. R.: Quantum walks, Ihara zeta functions and cospectrality in regular graphs. *Quantum Inf. Process.* **10**, 405–417 (2011)
- [21] Segawa, E., Suzuki, A.: Spectral mapping theorem of an abstract quantum walk. *Quantum Inf. Process.* **18**, 333 (2019)

- [22] Spitzer, F.: Principles of Random Walk, 2nd edition. Springer, New York (1976)
- [23] Terras, A.: Zeta Functions of Graphs. Cambridge Univ. Press (2011)
- [24] Venegas-Andraca, S. E.: Quantum walks: a comprehensive review. Quantum Inf. Process. **11**, 1015–1106 (2012)