

On characterization of the exponential distribution via hypoexponential distributions

George Yanev^{1,2*}

^{1*}School of Mathematical and Statistical Sciences, University of Texas Rio Grande Valley, 1201 W. University Dr., Edinburg 78539, Texas, USA.

^{2*}Institute of Mathematics and Informatics, Bulgarian Academy of Sciences, 8 Acad. G. Bontchev St., Sofia 1113, Bulgaria.

Corresponding author(s). E-mail(s): george.yanev@utrgv.edu;

Abstract

The sum of independent, but not necessary identically distributed, exponential random variables follows a hypoexponential distribution. We focus on a particular case when all but one rate parameters of the exponential variables are identical. This is known as exponentially modified Erlang distribution in molecular biology. We prove a characterization of the exponential distribution, which complements previous characterizations via hypoexponential distribution with all rates different from each other.

Keywords: characterizations, exponential distribution, hypoexponential distribution, exponentially modified Erlang distribution

MSC Classification: 62G30 , 62E10.

1 Introduction and main results

Sums of exponentially distributed random variables play a central role in many stochastic models of real-world phenomena. The *hypoexponential distribution* arises as a convolution of n independent exponential distributions each with their own rate λ_i , the rate of the i^{th} exponential distribution. It belongs to the class of *phase-type distributions*. Many processes can be divided into sequential

2 On characterization of the exponential distribution

phases. If the time periods spent in different phases of the process are independent but not necessary identically distributed exponential variables, then the overall time is hypoexponential. For example, the absorption time for a finite-state Markov chain follows this distribution.




Fig. 1 Absorption time for a Markov chain with five states.

We will write $X_i \sim \text{Exp}(\lambda_i)$, $\lambda_i > 0$, if X_i for $i = 1, 2, \dots$ has density

$$f_i(x) = \lambda_i e^{-\lambda_i x}, \quad x \geq 0 \quad (\text{exponential distribution}). \quad (1)$$

The distribution of the sum

$$Y_n = X_1 + X_2 + \dots + X_n \sim \text{HypoE}(\lambda_1, \lambda_2, \dots, \lambda_n),$$

where λ_i for $i = 1, \dots, n$ are not all identical, is called (general) *hypoexponential distribution* (e.g., [1] and [2]). Assume that all λ_i 's are distinct, i.e., $\lambda_i \neq \lambda_j$ when $i \neq j$. It is well-known that under this condition, the density of Y_n is given by (see [3], p.309 and [4], p.40, Problem 12) for $x > 0$

$$f_{Y_n}(x) = \sum_{j=1}^n \ell_j f_j(x), \quad \lambda_i \neq \lambda_j, \quad i \neq j.$$

Here the weight ℓ_j is defined as $\ell_j = \prod_{i=1, i \neq j}^n \lambda_i (\lambda_i - \lambda_j)^{-1}$. Thus, the density of the sum of independent exponential random variables with distinct parameters is linear combination of the individual densities. For example, the density of Y_2 is

$$f_{Y_2}(x) = \frac{\lambda_2}{\lambda_2 - \lambda_1} f_1(x) + \frac{\lambda_1}{\lambda_1 - \lambda_2} f_2(x), \quad \lambda_1 \neq \lambda_2.$$

Let X_1 and X_2 be two independent copies of a non-negative random variable X and $\mathbb{E}[X] < \infty$. If $X \sim \text{Exp}(\lambda)$, then $X_1 + X_2/2 \sim \text{HypoE}(\lambda, 2\lambda)$. It was proved in [5] that this property of the exponential distribution is not shared by any other continuous distribution, i.e., for $\lambda > 0$

$$X \sim \text{Exp}(\lambda) \quad \text{iff} \quad X_1 + \frac{1}{2}X_2 \sim \text{HypoE}(\lambda, 2\lambda). \quad (2)$$

The key argument in the proof is that the Laplace transform (LT) of the exponential distribution with $\lambda > 0$

$$\Phi(t) = \frac{\lambda}{\lambda + t}, \quad t \geq 0 \quad (3)$$

is the unique LT solution of the equation

$$\Phi(t)\Phi\left(\frac{t}{2}\right) = 2\Phi(t) - \Phi\left(\frac{t}{2}\right), \quad t \geq 0.$$

Motivated by (2), in [6] we extended it in two directions: (i) for any number $n \geq 2$ of independent copies X_1, X_2, \dots, X_n of X , and (ii) for the linear combination

$$\mu_1 X_1 + \mu_2 X_2 + \dots + \mu_n X_n \quad \mu_i \neq \mu_j, \quad i \neq j, \quad \mu_i > 0,$$

i.e., with arbitrary positive and distinct coefficients $\mu_1, \mu_2, \dots, \mu_n$. Namely, it was proved in [6], under some additional assumptions, that for $\lambda > 0$

$$X \sim \text{Exp}(\lambda) \quad \text{iff} \quad \sum_{k=1}^n \mu_k X_k \sim \text{HypoE}\left(\frac{\lambda}{\mu_1}, \frac{\lambda}{\mu_2}, \dots, \frac{\lambda}{\mu_n}\right). \quad (4)$$

This characterization was obtained by showing that (3) is the unique LT solution of the equation

$$\Phi(\mu_1 t)\Phi(\mu_2 t) \cdots \Phi(\mu_n t) = \sum_{j=1}^n \bar{\ell}_j \bar{\Phi}(\mu_j t), \quad t \geq 0,$$

where $\bar{\ell}_j = \prod_{i=1, i \neq j}^n \mu_j(\mu_j - \mu_i)^{-1}$. Thus, the case of the rate parameters λ_i 's in (1) being all different from each other was settled down. Note that characterization results in the case of distinct but not necessary positive $\mu_1, \mu_2, \dots, \mu_n$ were recently obtained, in [7], under an additional assumption.

The other extreme case of all λ_i 's equal leads to Erlang distribution of the sum. Assume $X_i \sim \text{Exp}(\lambda)$, i.e., $\lambda_1 = \lambda_2 = \dots = \lambda_n = \lambda$ and let

$$Y_n = X_1 + X_2 + \dots + X_n.$$

If Φ is the common LT of X_i , then for $t > 0$

$$\Phi_{Y_n}(t) = \Phi^n(t) = \left(\frac{\lambda}{\lambda + t}\right)^n. \quad (5)$$

If we go in the opposite direction, assuming $Y_n \sim \text{Erl}(n, \lambda)$, then (5) yields $\Phi_i(t) = \lambda(\lambda + t)^{-1}$ for each $i = 1, 2, \dots, n$, which in turn implies $X_i \sim \text{Exp}(\lambda)$.

4 On characterization of the exponential distribution

By words, if X_i are independent and identically distributed r.v.'s and Y_n is Erlang, then the common distribution is exponential.

The question arises whether a similar characterization holds when the rate parameters λ_i 's of $\text{HypoE}(\lambda_1, \lambda_2, \dots, \lambda_n)$ are neither all different nor all equal? It is our goal in this paper to show that, at least in one particular case, the answer to this question is affirmative.

Without the condition that all parameters λ_i 's are different or equal, the hypoexponential density has a quite complex form ([8]). This makes the analysis of the general case difficult. In this paper, we consider the particular case of "all-but-one-equal" rate parameters. More precisely, let X_1, X_2, \dots, X_{n+1} be independent copies of $X \sim \text{Exp}(\lambda)$. Consider the sum

$$X_1 + X_2 + \dots + X_n + wX_{n+1}, \quad w > 0, w \neq 1. \quad (6)$$

This sum has a convoluted Erlang distribution, which is also known as *exponentially modified Erlang (EME) distribution* ([9], [10]).

Recall the well-known (e.g., [11], p.240) Cramér's condition. We say that X satisfies Cramér's condition if there is a number $t_0 > 0$ such that $\mathbb{E}[e^{tX}] < \infty$ for all $t \in (-t_0, t_0)$. The next theorem establishes, under Cramér's condition, a necessary and sufficient condition for $X \sim \text{Exp}(\lambda)$.

Theorem *Suppose that X_1, X_2, \dots, X_{n+1} , $n \geq 1$, are independent copies of a non-negative and absolutely continuous random variable X . Assume further that X satisfies Cramér's condition. Then for some $\lambda > 0$, fixed positive integer n and fixed positive real $w \neq 1$*

$$X \sim \text{Exp}(\lambda) \quad \text{iff} \quad \sum_{k=1}^n X_k + wX_{n+1} \sim \text{HypoE} \left(\lambda, \lambda, \dots, \lambda, \frac{\lambda}{w} \right). \quad (7)$$

The hypoexponential family of distributions has found use in diverse applied fields, including queuing theory ([12]), population genetics ([13]), reliability analysis ([14]), medicine ([15]), and cell biology ([10], [16]). We focus on a particular member of this family, namely the exponentially modified Erlang distribution. The (proper) Erlang distribution is applied in modeling the cell cycle phase progression as a series of sub-phase transitions with the same rate λ . The relevant biological interpretation of the Erlang model is that each cell cycle phase can be viewed as a multi-step biochemical process that needs to be completed sequentially in order to advance to the next cell cycle phase.

Although the identical-stage model, is convenient from a mathematical perspective, it has been shown to be outperformed by a number of other distributions. Particularly, it was shown in [9] that one of the most appropriate distributions for representing cell cycle times is the EME distribution, which models a series of exponentially distributed random variables when one of them has a different rate. Under this assumption the multi-stage cell cycle model is

described as follows

$$X_1 \xrightarrow{\lambda_1} X_2 \xrightarrow{\lambda_1} \dots X_k \xrightarrow{\lambda_1} X_{k+1} \xrightarrow{\lambda_2} 2X_1. \quad (8)$$

Note that, in the system (8), the rate of progression is identical through each of the initial k stages of cell cycle and that we have added an additional exponentially distributed stage at the end whose rate, λ_2 is distinct from the rate, λ_1 , of the previous k stages. Assume that the characteristic time (CT) of one of the transitions is significantly bigger than the CTs of all other transitions. Assume also that the CTs are independent. Then the overall cell cycle time distribution may be approximated by the EME distribution capturing both the relatively rapid transitions and the slow transition (see [9]). Using the density convolution formula, for the density of (6), it is not difficult to obtain

$$f_{EME}(x) = \frac{\lambda}{w} e^{-\lambda x/w} \left(\frac{w}{w-1} \right)^n \left[1 - \frac{\Gamma(n, (w-1)\lambda x/w}{(n-1)!} \right],$$

where $\Gamma(n, t) = \int_t^\infty u^{n-1} e^{-u} du$ is the complementary incomplete gamma function.

In Section 2 we present some auxiliary results. We prove the Theorem in Section 3. The last section includes some concluding remarks.

2 Auxiliary results

Due to the independence assumption, the LT of (6) equals $\Phi(wt)\Phi^n(t)$. If Φ is given by (3), then $\Phi(wt)\Phi^n(t)$ is a product of linear fractions and we can decompose it into sum of the Laplace transforms of wX and X . Denote

$$\Phi_1(t) := (w-1)\Phi(wt) \quad \text{and} \quad \Phi_2(t) := \frac{w-1}{w}\Phi(t). \quad (9)$$

Lemma 1 The following identity holds

$$\Phi_1(t)\Phi_2^n(t) = \Phi_1(t) - \sum_{k=1}^n \Phi_2^k(t). \quad (10)$$

Proof. Without loss of generality assume that $X \sim \text{Exp}(1)$. Note that the following linear fraction decomposition holds

$$\frac{w-1}{(1+wt)(1+t)} = \frac{w}{1+wt} - \frac{1}{1+t}. \quad (11)$$

Recalling that $\Phi(t) = (1+t)^{-1}$, notation (9), and multiplying both sides of (11) by $(w-1)/w$, we obtain

$$\Phi_1(t)\Phi_2(t) = \Phi_1(t) - \Phi_2(t).$$

6 On characterization of the exponential distribution

This proves (10) for $n = 1$. Assuming that (10) holds for n , for the $(n + 1)^{th}$ term we have that

$$\begin{aligned}\Phi_1(t)\Phi_2^n(t)\Phi_2(t) &= \left[\Phi_1(t) - \sum_{k=1}^n \Phi_2^k(t) \right] \Phi_2(t) \\ &= \Phi_1(t)\Phi_2(t) - \sum_{k=1}^n \Phi_2^{k+1}(t) \\ &= \Phi_1(t) - \Phi_2(t) - \sum_{k=2}^{n+1} \Phi_2^k(t) \\ &= \Phi_1(t) - \sum_{k=1}^{n+1} \Phi_2^k(t),\end{aligned}$$

which proves (10) for any n .

Lemma 2 Let n be any positive integer and $v \neq 1$ be a real number.

(i) For any integer $j \geq 1$

$$v \sum_{k=0}^{n-1} \binom{k}{j-1} v^k + (v-1) \sum_{k=0}^{n-1} \binom{k}{j} v^k = \binom{n}{j} v^n. \quad (12)$$

(ii) For any integer $j \geq 2$

$$\left(\frac{v}{v-1} \right)^{j-1} v \sum_{k=0}^{n-1} v^k + (v-1) \sum_{k=0}^{n-1} k v^k \neq n v^n. \quad (13)$$

Proof. (i) The left-hand side of (12) is equivalent to

$$\begin{aligned}& \sum_{k=0}^{n-1} \binom{k}{j-1} v^{k+1} + (v-1) \sum_{k=0}^{n-1} \binom{k}{j} v^k \\ &= \sum_{k=0}^{n-1} \left[\binom{k}{j-1} + \binom{k}{j} \right] v^{k+1} - \sum_{k=0}^{n-1} \binom{k}{j} v^k \\ &= \sum_{k=0}^{n-1} \binom{k+1}{j} v^{k+1} - \sum_{k=0}^{n-1} \binom{k}{j} v^k \\ &= \sum_{k=1}^n \binom{k}{j} v^k - \sum_{k=1}^{n-1} \binom{k}{j} v^k \\ &= \binom{n}{j} v^n.\end{aligned}$$

(ii) Using (12) with $j = 1$, for the left-hand side of (13), we obtain

$$\begin{aligned} & \left(\frac{v}{v-1}\right)^{j-1} v \sum_{k=0}^{n-1} v^k - v \sum_{k=0}^{n-1} v^k + v \sum_{k=0}^{n-1} v^k + (v-1) \sum_{k=0}^{n-1} kv^k \\ &= \left[\left(\frac{v}{v-1}\right)^{j-1} - 1 \right] v \sum_{k=0}^{n-1} v^k + nv^n \\ &= \left[\left(\frac{v}{v-1}\right)^j - \frac{v}{v-1} \right] (v^n - 1) + nv^n \neq nv^n. \end{aligned}$$

Remark. It is not difficult to see that (12) can be generalized to

$$v \sum_{k=0}^{n-1} \binom{k+m}{j-1} v^k + (v-1) \sum_{k=0}^{n-1} \binom{k+m}{j} v^k = \binom{n+m}{j} v^n, \quad (14)$$

where m is any non-negative integer.

3 Proof of the theorem

It follows from Lemma 1 that if $X \sim \text{Exp}(\lambda)$, then (10) holds true. We will proceed with the proof of the opposite direction in the claim. The case where $n = 1$ is a particular case of (4) included in [6]. Let $n \geq 2$. Consider the function Ψ with the following series expansion

$$\Psi(t) := \frac{1}{\Phi(t)} = \sum_{j=0}^{\infty} a_j t^j, \quad t > 0. \quad (15)$$

Note that, as a consequence of Cramér's condition, the above series is uniformly convergent in a proper neighborhood of $t = 0$ (see [11], p.240). To prove the theorem, it is sufficient to show that for some $\lambda > 0$

$$\Psi(t) = 1 + \lambda^{-1}t,$$

i.e., the coefficients of the series in (15) are

$$a_0 = 1, \quad a_1 = \lambda^{-1} > 0, \quad a_j = 0, \quad j \geq 2. \quad (16)$$

Clearly,

$$a_0 = \Psi(0) = 1. \quad (17)$$

It follows from (9) and (10) that

$$\frac{(w-1)^{n+1}}{w^n} \Phi(wt) \Phi^n(t) = (w-1) \Phi(wt) - \sum_{k=1}^n \left(\frac{w-1}{w}\right)^k \Phi^k(t). \quad (18)$$

8 On characterization of the exponential distribution

Dividing both sides of (18) by its left-hand side and changing the summation index, we obtain

$$1 = v^n \Psi^n(t) - (v-1) \Psi(wt) \sum_{k=0}^{n-1} v^k \Psi^k(t), \quad (19)$$

where, for notational simplicity, we set $v = w/(w-1)$. To calculate the coefficients a_j for $j \geq 1$, we differentiate both sides of (19) with respect to t at $t = 0$. After differentiating once at $t = 0$ we have

$$\left[nv^n - v \sum_{k=0}^{n-1} v^k - (v-1) \sum_{k=0}^{n-1} kv^k \right] a_1 = 0.$$

It follows from (12) with $j = 1$, that the coefficient in front of a_1 equals zero and thus there exists a $\lambda > 0$ such that

$$a_1 = \lambda^{-1}. \quad (20)$$

Differentiating (19) twice with respect to t at $t = 0$, we have

$$\begin{aligned} & \left[\binom{n}{2} v^n - v \sum_{k=0}^{n-1} \binom{k}{1} v^k - (v-1) \sum_{k=0}^{n-1} \binom{k}{2} v^k \right] a_1^2 \\ & + \left[nv^n - \left(\frac{v}{v-1} \right) v \sum_{k=0}^{n-1} v^k + (v-1) \sum_{k=0}^{n-1} kv^k \right] a_2 = 0. \end{aligned}$$

Lemma 2 with $j = 2$ yields that the coefficient in front of a_1^2 is zero and the coefficient in front of a_2 is not zero. Therefore,

$$a_2 = 0. \quad (21)$$

It remains to prove that $a_j = 0$ for all $j \geq 3$. We will need the general Leibniz rule for differentiating a product of functions. Denote by $y^{(j)}(x)$ the j^{th} derivative of $y(x)$; $y^{(0)}(x) := y(x)$. Define a multi-index set $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \dots, \alpha_n)$ as a n -tuple of non-negative integers. Denote $\|\boldsymbol{\alpha}\| = \alpha_1 + \alpha_2 + \dots + \alpha_n$ and $\Lambda_j := \{\boldsymbol{\alpha} : \|\boldsymbol{\alpha}\| = j\}$. The j^{th} derivative (when exists) of the product $y_1(t)y_2(t) \cdots y_n(t)$ is given by (e.g. [17])

$$\frac{d^j}{dt^j} \prod_{i=1}^n y_i(t) = \sum_{\Lambda_j} \left(\frac{j!}{\alpha_1! \alpha_2! \cdots \alpha_n!} \prod_{i=1}^n y_i^{(\alpha_i)}(t) \right). \quad (22)$$

Let us write Λ_j as union of three disjoint subsets as follows:

$$\Lambda_j = \Lambda'_j \cup \Lambda''_j \cup \Lambda'''_j,$$

where

$$\begin{aligned}\Lambda'_j &= \{\|\boldsymbol{\alpha}\| = j : \text{only one of } \{\alpha_1, \alpha_2, \dots, \alpha_n\} \text{ equals } j \text{ (others are zeros)}\} \\ \Lambda''_j &= \{\|\boldsymbol{\alpha}\| = j : \text{exactly } j \text{ of } \{\alpha_1, \alpha_2, \dots, \alpha_n\} \text{ equal } 1 \text{ (others are zeros)}\} \\ \Lambda'''_j &= \{\|\boldsymbol{\alpha}\| = j : \text{there is an index } \alpha_i \text{ with } 2 \leq \alpha_i < j\}.\end{aligned}$$

Notice that by definition, Λ''_j is not empty only if $j \leq n$.

We will proceed by induction with respect to the index $j \geq 2$ of a_j . For $j = 2$ we have already proved that $a_2 = 0$. Assuming $a_i = 0$ for $2 \leq i \leq j - 1$, we will show that $a_j = 0$. Since $a_0 = 1$, applying (22), we obtain

$$\begin{aligned}\frac{1}{j!} \frac{d^j}{dt^j} \Psi^n(t)|_{t=0} &= \sum_{\Lambda_j} \left(\prod_{i=1}^n a_{\alpha_i} \right) = \sum_{\Lambda'_j} (\cdot) + \sum_{\Lambda''_j} (\cdot) + \sum_{\Lambda'''_j} (\cdot) \\ &= \binom{n}{1} a_j a_0^{n-1} + \binom{n}{j} a_1^j a_0^{n-j} \\ &= n a_j + \binom{n}{j} a_1^j.\end{aligned}$$

Notice that $\sum_{\Lambda'''_j} (\cdot) = 0$ by the induction assumption. Also,

$$\begin{aligned}\frac{1}{j!} \frac{d^j}{dt^j} \Psi(wt) \Psi^k(t)|_{t=0} &= \sum_{\Lambda_j} \left(w^{\alpha_{k+1}} a_{\alpha_{k+1}} \prod_{i=1}^k a_{\alpha_i} \right) = \sum_{\Lambda'} (\cdot) + \sum_{\Lambda''} (\cdot) + \sum_{\Lambda'''} (\cdot) \\ &= \left[w^j a_0^k a_j + \binom{k}{1} a_j a_0^{k-1} a_0 \right] + \left[\binom{k}{j-1} a_1^{j-1} a_0^{k-j+1} w a_1 + \binom{k}{j} a_1^j a_0^{k-j} a_0 \right] \\ &= \left[\binom{k}{j-1} w + \binom{k}{j} \right] a_1^j + (w^j + k) a_j.\end{aligned}$$

Therefore, differentiating (19) j times at $t = 0$ and grouping the coefficients in front of a_1^j and a_j , we write

$$\begin{aligned}\left[\binom{n}{j} v^n - v \sum_{k=0}^{n-1} \binom{k}{j-1} v^k + (v-1) \sum_{k=0}^{n-1} \binom{k}{j} v^k \right] a_1^j \\ + \left[n v^n - \left(\frac{v}{v-1} \right)^{j-1} v \sum_{k=0}^{n-1} v^k - (v-1) \sum_{k=0}^{n-1} k v^k \right] a_j = 0.\end{aligned}$$

It then follows from Lemma 2 that the coefficient in front of a_1^j is zero and the coefficient in front of a_j is not zero. Therefore, for all $j \geq 2$

$$a_j = 0. \quad (23)$$

Now, (17), (20), (21), and (23) lead to (16), which completes the proof.

4 Concluding remarks

In this paper we continue the study of the relation between the exponential and hypoexponential distributions, initiated in [5] and extended in [6]. The obtained characterization complements those in the above papers. Here we deal with a situation where the rate parameters λ_i 's in a convolution of exponential variables are not all different from each other. The obtained result is of interest itself, however it can also serve as a basis for further investigations of more complex compositions of the rate parameters.

The exponential distribution is well known as a lifetime model in reliability theory. In particular, its use as a probability model for failure times of system's components is well justified. Thus, it is important to assess goodness-of-fit of the exponential distribution for a data set prior to applying the exponential model. Characterization results often serve as a useful device in obtaining goodness-of-fit tests. The presented here characterization can be used for testing the validity of a model based on the exponential distribution.

Acknowledgments. The author acknowledges the valuable suggestions from the referees and the editor.

Statement of Conflict of Interest. The author states that there is no conflict of interest.

References

- [1] Li, K.-H.; Li, C.T. Linear combination of independent exponential random variables. *Methodol. Comput. Appl. Probab.* **2019**, *21*, 253–277.
- [2] Smaili, K.H.; Kadri, T.H.; Kadry, S. Finding the PDF of the hypoexponential random variable using the Kad matrix similar to the general Vandermonde matrix. *Commun. Statist. Theory Methods* **2016**, *45*, 1542–1549.
- [3] Ross, S.M. *Introduction to Probability Models*, 12th ed.; Academic Press: New York, NY, USA, 2019.
- [4] Feller, W. *An Introduction to Probability Theory and Its Applications, Vol. II*, 2nd ed.; Wiley: New York, 1971.
- [5] Arnold, B.C.; Villaseñor, J.A. Exponential characterizations motivated by the structure of order statistics in sample of size two. *Statist. Probab. Lett.* **2013**, *83*, 596–601.
- [6] Yanev, G.P. Exponential and hypoexponential distributions: some characterizations. *Mathematics, MDPI* **2020**, *8*, paper 2207.
- [7] Rastegar, R.; Roitershtein, A. On a characterization of exponential and double exponential distributions *REVSTAT*, **2023**, to appear.

- [8] Jasiulewicz, H.; and Kordecki, W. Convolutions of Erlang and of Pascal distributions with applications to reliability *Demonstratio Math.* **2003**, *36*, 231-238.
- [9] Golubev, A. Applications and implications of the exponentially modified gamma distribution as a model for time variabilities related to cell proliferation and gene expression *J Theor Biol* **2016**, *393*, 203–217.
- [10] Yates, Ch.A.; Ford, M.J.; Mori, R.L. A Multi-stage Representation of Cell Proliferation as a Markov Process *Bull Math Biol* **2017**, *79*, 2905–2928.
- [11] Borovkov, A.A. *Probability Theory*; Springer-Verlag: London, 2013.
- [12] Bekkeer, R.; Koeleman, P.M. Scheduling admissions and reducing variability in bed demand *Health Care Manag Sci* **2011**, *14*, 237–249.
- [13] Strimmer, K.; Pybus, O.G. Exploring the Demographic History of DNA Sequences Using the Generalized Skyline Plot *Mol Biol Evol* **2001**, *18*, 2298–2305.
- [14] Kadry, S.; Kadry, T.; Khaled, S. Markov Modeling for Reliability Analysis Using Hypoexponential Distribution (2015). In S. Kadry, A El Hami (Eds.) *Numerical Methods for Reliability and Safety Assessment: Multiscale and Multiphysics Systems* (1st ed., pp. 599-620). Springer.
- [15] Dowling, C.V.; Cevaal, P.M.; Faria, M.; Johnston, S.T. On predicting heterogeneity in nanoparticle dosage *Math. Biosciences* **2022**, *354*, 108928.
- [16] Gavagnin, E.; Ford, M.J.; Mort, R.L.; Rogers, T.; Yates, Ch. The invasion speed of cell migration models with realistic cell cycle time distributions. *J. Theor Biol* **2019**, *481*, 91–99.
- [17] Thaheem, A.B.; Laradji, A. Classroom note: A generalization of Leibniz rule for higher derivatives. *Intern. J. Math. Educ. Sci. Technol.* **2003**, *34*, 905–907.