

Two iterative formulas of largest and smallest singular value of nonsingular matrices

Shun Xu

School of Mathematical Sciences, Tongji University, Shanghai, 200092, China

e-mail: shunxu1997@163.com

Abstract

We obtain an iterative formula that converges incrementally to the smallest singular value. Similarly, we obtain an iterative formula that converges decreasingly to the largest singular value.

Keywords: Singular values, Frobenius norm, Determinant.

1 Lower bound for the smallest singular value

Let $M_n(\mathbb{C})(n \geq 2)$ be the space of $n \times n$ complex matrices. Let σ_i ($i = 1, \dots, n$) be the singular values of $A \in M_n(\mathbb{C})$ which is nonsingular and suppose that $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_{n-1} \geq \sigma_n > 0$. For $A = [a_{ij}] \in M_n(\mathbb{C})$, the Frobenius norm of A is defined by

$$\|A\|_F = \left(\sum_{i,j=1}^n |a_{ij}|^2 \right)^{1/2} = \text{tr}(A^H A)^{\frac{1}{2}}$$

where A^H is the conjugate transpose of A . The relationship between the Frobenius norm and singular values is

$$\|A\|_F^2 = \sigma_1^2 + \sigma_2^2 + \dots + \sigma_n^2$$

It is well known that lower bounds for the smallest singular value σ_n of a nonsingular matrix $A \in M_n(\mathbb{C})$ have many potential theoretical and practical applications [1, 2]. Yu and Gu [5] obtained a lower bound for σ_n as follows:

$$\sigma_n \geq |\det A| \left(\frac{n-1}{\|A\|_F^2} \right)^{(n-1)/2} = l > 0$$

The above inequality is also shown in [4]. In [6], Zou improved the above inequality by showing that

$$\sigma_n \geq |\det A| \left(\frac{n-1}{\|A\|_F^2 - l^2} \right)^{(n-1)/2} = l_0$$

In [3], Lin and Xie improve a lower bound for smallest singular value of matrices by showing that a is the smallest positive solution to the equation

$$x^2 (\|A\|_F^2 - x^2)^{n-1} = |\det A|^2 (n-1)^{n-1}.$$

and $\sigma_n \geq a > l_0$. Under certain conditions, $\sigma_n = a$ will hold. However, in many cases, $\sigma_n = a$ is not true. We give necessary and sufficient conditions such that $\sigma_n = a$ in Proposition 1.

2 Main results

Let

$$l = |\det A| \left(\frac{n-1}{\|A\|_F^2} \right)^{(n-1)/2},$$

$$l_0 = |\det A| \left(\frac{n-1}{\|A\|_F^2 - l^2} \right)^{(n-1)/2}.$$

And a is the smallest positive solution to the equation

$$x^2 (\|A\|_F^2 - x^2)^{n-1} = |\det A|^2 (n-1)^{n-1}.$$

From [3], we have $\sigma_n \geq a > l_0 > l > 0$. Next, we give necessary and sufficient conditions such that $a = \sigma_n$.

Proposition 1. *Let a be the smallest positive solution to the equation*

$$x^2 (\|A\|_F^2 - x^2)^{n-1} = |\det A|^2 (n-1)^{n-1}.$$

then $a = \sigma_n$ if and only if

$$\sigma_1 = \sigma_2 = \cdots = \sigma_{n-1}$$

Proof. \Rightarrow : If $a = \sigma_n$, then

$$\sigma_n^2 (\|A\|_F^2 - \sigma_n^2)^{n-1} = |\det A|^2 (n-1)^{n-1}.$$

Since $\|A\|_F^2 = \sigma_1^2 + \sigma_2^2 + \cdots + \sigma_n^2$ and $|\det A|^2 = \sigma_1^2 \sigma_2^2 \cdots \sigma_n^2$, we have

$$\sigma_n^2 (\sigma_1^2 + \sigma_2^2 + \cdots + \sigma_{n-1}^2)^{n-1} = \sigma_1^2 \sigma_2^2 \cdots \sigma_n^2 (n-1)^{n-1}.$$

$$\left(\frac{\sigma_1^2 + \sigma_2^2 + \cdots + \sigma_{n-1}^2}{n-1} \right)^{n-1} = \sigma_1^2 \sigma_2^2 \cdots \sigma_{n-1}^2$$

By the arithmetic-geometric mean inequality, the above equation holds if and only if

$$\sigma_1^2 = \sigma_2^2 = \cdots = \sigma_{n-1}^2$$

if and only if

$$\sigma_1 = \sigma_2 = \cdots = \sigma_{n-1}$$

\Leftarrow : Let

$$\sigma_1^2 = \sigma_2^2 = \cdots = \sigma_{n-1}^2 = c^2 (c > 0).$$

Then a is the smallest positive solution to the equation

$$x^2 (c^2(n-1) + \sigma_n^2 - x^2)^{n-1} = c^{2(n-1)} \sigma_n^2 (n-1)^{n-1}$$

if and only if a is the smallest positive zero point of

$$f(x) = x^2 \left(1 + \frac{\sigma_n^2 - x^2}{c^2(n-1)} \right)^{n-1} - \sigma_n^2.$$

Next, we proof $a = \sigma_n$. Obviously, we can see that $f(\sigma_n) = 0$ and $f(0) = -\sigma_n^2 < 0$. Next, we prove that $f(x)$ is an strictly increasing function on $[0, \sigma_n]$. Taking the derivative of $f(x)$, we can get

$$f'(x) = \left(1 + \frac{\sigma_n^2 - x^2}{c^2(n-1)} \right)^{n-2} \frac{2x}{c^2(n-1)} (c^2(n-1) + \sigma_n^2 - nx^2).$$

$\forall x_0 \in (0, \sigma_n)$, we have

$$c^2(n-1) + \sigma_n^2 - nx_0^2 \geq n\sigma_n^2 - nx_0^2 > 0.$$

We have $f'(x_0) > 0$. Therefore $f(x)$ is an strictly increasing function on $[0, \sigma_n]$ and σ_n is the smallest positive zero point of

$$f(x) = x^2 \left(1 + \frac{\sigma_n^2 - x^2}{c^2(n-1)} \right)^{n-1} - \sigma_n^2$$

Therefore, σ_n is the smallest positive solution to the equation

$$x^2 (c^2(n-1) + \sigma_n^2 - x^2)^{n-1} = c^{2(n-1)} \sigma_n^2 (n-1)^{n-1}.$$

We get $a = \sigma_n$. □

In the above special condition, a can be equal to σ_n , which is not general. Next, we give our main theorem. We give an iterative formula for the smallest singular value, which converges incrementally to the smallest singular value.

Theorem 1. *Let $A \in M_n(\mathbb{C})$ be nonsingular and $0 < a_1 \leq \sigma_n$ and*

$$a_{k+1} = \left(a_k^2 + |\det(a_k^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)a_k^2} \right)^{n-1} \right)^{1/2}, k = 1, 2, \dots.$$

Then $\sigma_n \geq a_{k+1} \geq a_k > 0 (k = 1, 2, \dots)$ and $\lim_{k \rightarrow +\infty} a_k = \sigma_n$.

Proof. Let $0 \leq \lambda < \sigma_n^2$, by the arithmetic-geometric mean inequality, we have

$$(\sigma_1^2 - \lambda)(\sigma_2^2 - \lambda) \cdots (\sigma_{n-1}^2 - \lambda) \leq \left(\frac{\sigma_1^2 + \cdots + \sigma_{n-1}^2 - (n-1)\lambda}{n-1} \right)^{n-1}$$

Since

$$\begin{aligned} (\sigma_1^2 - \lambda)(\sigma_2^2 - \lambda) \cdots (\sigma_{n-1}^2 - \lambda) &= \frac{(\sigma_1^2 - \lambda)(\sigma_2^2 - \lambda) \cdots (\sigma_n^2 - \lambda)}{\sigma_n^2 - \lambda} \\ &= \frac{|\det(\lambda I_n - A^H A)|}{\sigma_n^2 - \lambda} \end{aligned}$$

we have

$$\begin{aligned} \frac{|\det(\lambda I_n - A^H A)|}{\sigma_n^2 - \lambda} &\leq \left(\frac{\sigma_1^2 + \cdots + \sigma_{n-1}^2 - (n-1)\lambda}{n-1} \right)^{n-1} \\ \sigma_n^2 &\geq \lambda + |\det(\lambda I_n - A^H A)| \left(\frac{n-1}{\sigma_1^2 + \cdots + \sigma_{n-1}^2 - (n-1)\lambda} \right)^{n-1} \\ \sigma_n &\geq \left(\lambda + |\det(\lambda I_n - A^H A)| \left(\frac{n-1}{\sigma_1^2 + \cdots + \sigma_{n-1}^2 + \sigma_n^2 - (n-1)\lambda} \right)^{n-1} \right)^{1/2} \\ \sigma_n &\geq \left(\lambda + |\det(\lambda I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)\lambda} \right)^{n-1} \right)^{1/2} \end{aligned}$$

Let $\lambda \rightarrow \sigma_n^{2-}$ (λ tends to σ_n^2 from the left). We get that the above inequality is also true for $\lambda = \sigma_n^2$. Therefore, for $0 \leq \lambda \leq \sigma_n^2$, we have

$$\sigma_n \geq \left(\lambda + |\det(\lambda I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)\lambda} \right)^{n-1} \right)^{1/2} \quad (1)$$

We show by induction on k that

$$\sigma_n \geq a_{k+1} \geq a_k > 0$$

We have $\sigma_n \geq a_1 > 0$. In equation 1, let $\lambda = a_1^2$, we have

$$\sigma_n \geq a_2 = \left(a_1^2 + |\det(a_1^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)a_1^2} \right)^{n-1} \right)^{1/2} \geq a_1 > 0$$

Assume that our claim is true for $k = m$, that is $\sigma_n \geq a_{m+1} \geq a_m > 0$. Now we consider the case when $k = m + 1$. In equation 1, let $\lambda = a_{m+1}^2$, we have

$$\sigma_n \geq a_{m+2} = \left(a_{m+1}^2 + |\det(a_{m+1}^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)a_{m+1}^2} \right)^{n-1} \right)^{1/2} \geq a_{m+1} > 0$$

Hence $\sigma_n \geq a_{m+2} \geq a_{m+1} > 0$. This proves $\sigma_n \geq a_{k+1} \geq a_k > 0 (k = 1, 2, \dots)$. By the well known monotone convergence theorem, $\lim_{k \rightarrow \infty} a_k$ exists. Let $\lim_{k \rightarrow \infty} a_k = \sigma$, then

$$\sigma = \left(\sigma^2 + |\det(\sigma^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)\sigma^2} \right)^{n-1} \right)^{1/2}, k = 1, 2, \dots$$

We have

$$|\det(\sigma^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)\sigma^2} \right)^{n-1} = 0$$

Since

$$\frac{n-1}{\|A\|_F^2 - (n-1)\sigma^2} \neq 0$$

we have

$$\det(\sigma^2 I_n - A^H A) = 0.$$

We get that σ^2 is the eigenvalue of $A^H A$. Since σ_n^2 is the smallest eigenvalue of $A^H A$, we have $\sigma^2 \geq \sigma_n^2$. According to the definition of σ , we have $\sigma \leq \sigma_n$. Therefore, $\sigma^2 \leq \sigma_n^2$ and we get $\sigma = \sigma_n$. Hence $\lim_{k \rightarrow +\infty} a_k = \sigma_n$. \square

From Theorem 1, we can see that as long as there is a lower bound of σ_n (we set it to b) and let $a_1 = b$ in Theorem 1, we can get a better lower bound than b . For example, we bring the lower bound of Lin and Xie [3] into our Theorem 1 to obtain the following results.

Corollary 1. *Let a be the smallest positive solution to the equation*

$$x^2 (\|A\|_F^2 - x^2)^{n-1} = |\det A|^2 (n-1)^{n-1}.$$

Let $a_1 = a$ and

$$a_{k+1} = \left(a_k^2 + |\det(a_k^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)a_k^2} \right)^{n-1} \right)^{1/2}, k = 1, 2, \dots$$

Then $\sigma_n \geq a_{k+1} \geq a_k \geq a > 0 (k = 1, 2, \dots)$ and $\lim_{k \rightarrow +\infty} a_k = \sigma_n$.

Proof. From [3], we have $\sigma_n \geq a > 0$. We know Corollary 1 is a special case of Theorem 1. \square

We give the iterative formula for the smallest singular value:

$$a_{k+1} = \left(a_k^2 + |\det(a_k^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)a_k^2} \right)^{n-1} \right)^{1/2}, k = 1, 2, \dots,$$

where $a_1 = a$, which converges incrementally to the smallest singular value. Similarly, we can give an iterative formula that converges decreasingly to the largest singular value.

Theorem 2. *Let $A \in M_n(\mathbb{C})$ be nonsingular, $a_1 \geq \sigma_1$. Assume*

$$a_{k+1} = \left(a_k^2 - |\det(a_k^2 I_n - A^H A)| \left(\frac{n-1}{(n+1)a_k^2 - \|A\|_F^2} \right)^{n-1} \right)^{1/2}, k = 1, 2, \dots.$$

Then $\sigma_1 \leq a_{k+1} \leq a_k$ ($k = 1, 2, \dots$) and $\lim_{k \rightarrow +\infty} a_k = \sigma_1$.

Proof. Set $\lambda > \sigma_1^2$, according to the arithmetic geometric mean inequality, we can get

$$(\lambda - \sigma_2^2) (\lambda - \sigma_3^2) \cdots (\lambda - \sigma_n^2) \leq \left(\frac{(n-1)\lambda - (\sigma_2^2 + \cdots + \sigma_n^2)}{n-1} \right)^{n-1}.$$

Since

$$\begin{aligned} (\lambda - \sigma_2^2) (\lambda - \sigma_3^2) \cdots (\lambda - \sigma_n^2) &= \frac{(\lambda - \sigma_1^2) (\lambda - \sigma_2^2) \cdots (\lambda - \sigma_n^2)}{\lambda - \sigma_1^2}, \\ &= \frac{|\det(\lambda I_n - A^H A)|}{\lambda - \sigma_1^2}, \end{aligned}$$

we can get

$$\begin{aligned} \frac{|\det(\lambda I_n - A^H A)|}{\lambda - \sigma_1^2} &\leq \left(\frac{(n-1)\lambda - (\sigma_2^2 + \cdots + \sigma_n^2)}{n-1} \right)^{n-1} \\ \sigma_1^2 &\leq \lambda - |\det(\lambda I_n - A^H A)| \left(\frac{n-1}{(n-1)\lambda + \sigma_1^2 - \|A\|_F^2} \right)^{n-1} \\ &\leq \lambda - |\det(\lambda I_n - A^H A)| \left(\frac{n-1}{(n+1)\lambda - \|A\|_F^2} \right)^{n-1} \\ \sigma_1 &\leq \left(\lambda - |\det(\lambda I_n - A^H A)| \left(\frac{n-1}{(n+1)\lambda - \|A\|_F^2} \right)^{n-1} \right)^{1/2} \end{aligned}$$

Let $\lambda \rightarrow \sigma_1^{2+}$ (λ tend to σ_1^2 from the right). We get that the above equation is also true for $\lambda = \sigma_1^2$. So, for $\lambda \geq \sigma_1^2$, we have

$$\sigma_1 \leq \left(\lambda - |\det(\lambda I_n - A^H A)| \left(\frac{n-1}{(n+1)\lambda - \|A\|_F^2} \right)^{n-1} \right)^{1/2} \quad (2)$$

We use induction on k to prove that:

$$\sigma_1 \leq a_{k+1} \leq a_k$$

For $k = 1$, $\sigma_1 \leq a_1$ can be obtained from the condition. In the equation (2), taking $\lambda = a_1^2$, we can get

$$\sigma_1 \leq a_2 = \left(a_1^2 - |\det(a_1^2 I_n - A^H A)| \left(\frac{n-1}{(n+1)a_1^2 - \|A\|_F^2} \right)^{n-1} \right)^{1/2} \leq a_1$$

Suppose our conclusion holds for $k = m$, that is, $\sigma_1 \leq a_{m+1} \leq a_m$. Now let's consider the case of $k = m + 1$. In equation (2), let $\lambda = a_{m+1}^2$, we can get

$$\sigma_1 \leq a_{m+2} = \left(a_{m+1}^2 - |\det(a_{m+1}^2 I_n - A^H A)| \left(\frac{n-1}{(n+1)a_{m+1}^2 - \|A\|_F^2} \right)^{n-1} \right)^{1/2} \leq a_{m+1}$$

So $\sigma_1 \leq a_{k+1} \leq a_k$. This proves that $\sigma_1 \leq a_{k+1} \leq a_k (k = 1, 2, \dots)$. From the monotone convergence theorem, $\lim_{k \rightarrow \infty} a_k$ exists. Let $\lim_{k \rightarrow \infty} a_k = \sigma$, then

$$\sigma = \left(\sigma^2 - |\det(\sigma^2 I_n - A^H A)| \left(\frac{n-1}{(n+1)\sigma^2 - \|A\|_F^2} \right)^{n-1} \right)^{1/2}$$

We have

$$|\det(\sigma^2 I_n - A^H A)| \left(\frac{n-1}{(n+1)\sigma^2 - \|A\|_F^2} \right)^{n-1} = 0.$$

because

$$\frac{n-1}{(n+1)\sigma^2 - \|A\|_F^2} \neq 0,$$

we can get

$$\det(\sigma^2 I_n - A^H A) = 0.$$

So σ^2 is the eigenvalue of $A^H A$. Because σ_1^2 is the largest eigenvalue of $A^H A$, there is $\sigma^2 \leq \sigma_1^2$. According to the definition of σ , we have $\sigma \geq \sigma_1$, so $\sigma^2 \geq \sigma_1^2$. We get $\sigma = \sigma_1$, so $\lim_{k \rightarrow +\infty} a_k = \sigma_1$. \square

The largest singular value has an obvious upper bound. Because $\sigma_1^2 \leq \sigma_1^2 + \dots + \sigma_n^2 = \|A\|_F^2$, so $\sigma_1 < \|A\|_F$. We give an iterative formula for the largest singular value:

$$a_{k+1} = \left(a_k^2 - |\det(a_k^2 I_n - A^H A)| \left(\frac{n-1}{(n+1)a_k^2 - \|A\|_F^2} \right)^{n-1} \right)^{1/2}, k = 1, 2, \dots,$$

where $a_1 = \|A\|_F$, which converges decreasingly to σ_1 .

References

- [1] R. A. Horn and C. R. Johnson. *Matrix Analysis*. Cambridge University Press, 1985.
- [2] R. A. Horn and C. R. Johnson. *Topics in matrix analysis*. Cambridge University Press, 1994.
- [3] Minghua Lin and Mengyan Xie. On some lower bounds for smallest singular value of matrices. *Applied Mathematics Letters*, 121:107411, 2021.
- [4] G Piazza and T Politi. An upper bound for the condition number of a matrix in spectral norm. *Journal of Computational and Applied Mathematics*, 143(1):141–144, 2002.
- [5] Yu Yisheng and Gu Dunhe. A note on a lower bound for the smallest singular value. *Linear algebra and its Applications*, 253(1-3):25–38, 1997.
- [6] Limin Zou. A lower bound for the smallest singular value. *Journal of Mathematical Inequalities*, 6(4):625–629, 2012.