

# Strong convergence rates of an explicit scheme for stochastic Cahn–Hilliard equation with additive noise\*

Meng Cai <sup>a,b</sup>, Ruisheng Qi <sup>c</sup>, Xiaojie Wang <sup>a</sup>

<sup>a</sup> School of Mathematics and Statistics, HNP-LAMA, Central South University, Changsha, China

<sup>b</sup> Department of Mathematical and Statistical Sciences, University of Alberta, Edmonton, Canada

<sup>c</sup> School of Mathematics and Statistics, Yancheng Teachers University, Yancheng, China

June 3, 2022

## Abstract

In this paper, we propose and analyze an explicit time–stepping scheme for a spatial discretization of stochastic Cahn–Hilliard equation with additive noise. The fully discrete approximation combines a spectral Galerkin method in space with a tamed exponential Euler method in time. In contrast to implicit schemes in the literature, the explicit scheme here is easily implementable and produces significant improvement in the computational efficiency. It is shown that the fully discrete approximation converges strongly to the exact solution, with strong convergence rates identified. To the best of our knowledge, it is the first result concerning an explicit scheme for the stochastic Cahn–Hilliard equation. Numerical experiments are finally performed to confirm the theoretical results.

**AMS subject classification:** 60H35, 60H15, 65C30

**Key Words:** stochastic Cahn–Hilliard equation, strong convergence rates, spectral Galerkin method, tamed exponential Euler method

## 1 Introduction

Let  $\mathcal{D}$  be a bounded convex set of  $\mathbb{R}^d$ ,  $d \in \{1, 2, 3\}$  with smooth boundary. Let  $H := L^2(\mathcal{D}, \mathbb{R})$  denote a real separable Hilbert space with scalar product  $\langle \cdot, \cdot \rangle$  and norm  $\|\cdot\|$  and  $\dot{H} := \{v \in H : \int_{\mathcal{D}} v dx = 0\}$ . In this paper, we consider the numerical approximation of stochastic Cahn–Hilliard equation (SCHE) in the abstract form

$$\begin{cases} dX(t) + A(AX(t) + F(X(t))) dt = dW(t), & t \in (0, T], \\ X(0) = X_0, \end{cases} \quad (1.1)$$

---

\*This work was supported by NSF of China (Nos. 11971488, 12071488), NSF of Hunan Province (No. 2020JJ2040). M. Cai is also supported by the China Scholarship Council.

where  $0 < T < \infty$ ,  $-A$  is the Neumann Laplacian and  $\{W(t)\}_{t \geq 0}$  is a  $Q$ -Wiener process on a filtered probability space  $(\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0})$ , specified later. The nonlinear term  $F$  is assumed to be a Nemytskii operator, given by  $F(u)(x) = f(u(x)) = u^3(x) - u(x)$ ,  $x \in \mathcal{D}$ . As a phenomenological model from metallurgy and physics, the deterministic version of such equation is used to describe the complicated phase separation and coarsening phenomena in a melted alloy [5, 7] and spinodal decomposition for binary mixture [6]. The stochastic version was studied by many authors, e.g., [1, 11, 12, 14, 16, 26].

Since the true solution of the problem can not be known explicitly, it is therefore natural to look for reliable numerical solutions. To do the approximation error analysis, one often faces difficulties, raised by the presence of the unbounded operator  $A$  in front of the nonlinear term  $F$ . In the past few years, many authors investigated strong and weak approximations of stochastic Cahn-Hilliard equation [10, 23, 25, 18, 8, 13, 20, 21, 27, 17], where some attempts to address the issue were proposed in literature. For the linearized stochastic Cahn-Hilliard equation, the readers are referred to [10, 23, 25]. In [22], strong convergence rates for the spectral Galerkin spatial approximation of the nonlinear problem in dimension one are proved by combining a general perturbation theory with the exponential integrability properties of the numerical approximation. The authors in [18, 24] derive the strong convergence of the finite element spatial approximation and the backward Euler full discretization of the SCHE driven by spatial regular noise, but with no rates obtained. Very recently, the paper [27] fills the gap left by [18, 24] and recover the strong convergence rates of the finite element fully discrete scheme. For space-time white noise, authors in [13] obtained the strong convergence rates of a fully discrete scheme performed by a spatial spectral Galerkin method and a temporal accelerated implicit Euler method. Moreover, the strong convergence rates of an implicit fully discrete mixed finite element method for the SCHE with gradient-type multiplicative noise are derived in [17], where the noise process is a real-valued Wiener process. To our great knowledge, the explicit methods are absent for the SCHE and this paper aims to propose an explicit scheme for the equation and identify its strong convergence rates.

As indicated in [2], the fully discrete exponential Euler and fully discrete linear-implicit Euler approximations diverge strongly and numerically weakly in the case of stochastic Allen-Cahn equations. Later, some explicit modified Euler-type schemes have been proposed in [3, 4, 9, 19, 28] to numerically solve the stochastic Allen-Cahn equations. Based on a spectral Galerkin spatial approximation of (1.1), given by

$$dX^N(t) + A(AX^N(t) + P_N F(X^N(t)))dt = P_N dW(t), \quad t \in (0, T]; \quad X^N(0) = P_N X_0, \quad (1.2)$$

we propose a tamed exponential Euler scheme in time to obtain the explicit fully discrete method

$$X_{t_{m+1}}^{M,N} = E(\tau)X_{t_m}^{M,N} - \frac{A^{-1}(I-E(\tau))P_N F(X_{t_m}^{M,N})}{1+\tau\|P_N F(X_{t_m}^{M,N})\|} + E(\tau)P_N \Delta W_m, \quad m \in \{1, 2, \dots, M-1\}, \quad (1.3)$$

where  $\Delta W_m = W(t_{m+1}) - W(t_m)$ ,  $P_N$  is the projection operator onto  $H_N := \text{span}\{e_1, e_2, \dots, e_N\}$  and  $\tau = \frac{T}{M}$  stands for the time step-size. Compared with existing implicit schemes, the proposed scheme is easy to implement and produces significant improvement in the computational efficiency.

Throughout this article,  $C$  denotes a generic nonnegative constant that is independent of the discretization parameters and may change from line to line. Meanwhile, we use  $\mathbb{N}^+$  to denote

the set of all positive integers and  $\mathbb{N} = \{0\} \cup \mathbb{N}^+$ . In summary, the contribution of this article to the numerical analysis of stochastic Cahn-Hilliard equation is twofold. On the one hand, the uniform a priori moment bounds of the full discretization are derived based on a certain bootstrap argument. To do this, a key ingredient lies on bounding

$$\sup_{M, N \in \mathbb{N}^+} \sup_{m \in \{0, 1, \dots, M\}} \mathbb{E} \left[ \|X_{t_m}^{M, N}\|_{L^6}^p \right] < \infty$$

by virtue of Gagliardo-Nirenberg inequality in  $d = 1$  and energy estimate in  $d = 2, 3$ . On the other hand, as implied by Corollary 3.9, we identify the strong convergence rate of the fully discrete method:

$$\sup_{M, N \in \mathbb{N}^+} \sup_{m \in \{0, 1, \dots, M\}} \|X(t_m) - X_{t_m}^{M, N}\|_{L^p(\Omega, \dot{H})} \leq C (\lambda_N^{-\frac{\gamma}{2}} + \tau^{\frac{\gamma}{4}}), \quad \gamma \in (\frac{d}{2}, 4], \quad (1.4)$$

where  $\lambda_N$  is the  $N$ -th eigenvalue of  $A$  and  $\gamma$  from Assumption 2.3 is a parameter used to measure the spatial regularity of the noise process. The above result reveals that the strong convergence rates are essentially governed by the spatial regularity of the noise term. Comparing (1.4) with the sharp temporal Hölder continuity result in Theorem 2.6, one can easily observe, for  $\gamma \in (\frac{d}{2}, 2]$ , the rate of convergence is in accordance with the temporal Hölder continuity of the mild solution. For  $\gamma \in [2, 4]$ , the rate of the convergence can reach 1 and higher than the Hölder continuity of the mild solution. It must be emphasized that the derivation of (1.4) is not an easy task and requires a variety of delicate error estimates, which are elaborated in subsection 3.3.

The outline of the article is organized as follows. In the next section, we present some assumptions and give the well-posedness and regularity of the mild solution. Section 3 is devoted to the strong convergence analysis, where spectral Galerkin method is introduced in subsection 3.1, uniform a priori moment bounds are deduced in subsection 3.2 and the strong convergence rates are derived in subsection 3.3. Numerical examples are finally included in Section 4 to verify the theoretical findings.

## 2 Main assumptions and the considered problem

Given another separable Hilbert space  $(U, \langle \cdot, \cdot \rangle_U, \|\cdot\|_U)$ ,  $\mathcal{L}(U, H)$  represents the space of all bounded linear operators from  $U$  to  $H$  endowed with the usual operator norm  $\|\cdot\|_{\mathcal{L}(U, H)}$  and by  $\mathcal{L}_2(U, H) \subset \mathcal{L}(U, H)$  we denote the space consisting of all Hilbert-Schmidt operators from  $U$  to  $H$ . To simplify the notation, we often write  $\mathcal{L}(H)$  and  $\mathcal{L}_2(H)$  (or  $\mathcal{L}_2$  for short) instead of  $\mathcal{L}(H, H)$  and  $\mathcal{L}_2(H, H)$ , respectively. It is easy to prove that  $\mathcal{L}_2(U, H)$  is a Hilbert space equipped with the inner product and norm,

$$\langle T_1, T_2 \rangle_{\mathcal{L}_2(U, H)} = \sum_{i \in \mathbb{N}^+} \langle T_1 \phi_i, T_2 \phi_i \rangle, \quad \|T\|_{\mathcal{L}_2(U, H)} = \left( \sum_{i \in \mathbb{N}^+} \|T \phi_i\|^2 \right)^{\frac{1}{2}}, \quad (2.1)$$

independent of the choice of orthonormal basis  $\{\phi_i\}$  of  $U$ . If  $T \in \mathcal{L}_2(U, H)$  and  $L \in \mathcal{L}(H, U)$ , then  $TL \in \mathcal{L}_2(H)$  and  $\|TL\|_{\mathcal{L}_2(H)} \leq \|T\|_{\mathcal{L}_2(U, H)} \|L\|_{\mathcal{L}(H, U)}$ . Also,  $|\langle T_1, T_2 \rangle_{\mathcal{L}_2(U, H)}| \leq \|T_1\|_{\mathcal{L}_2(U, H)} \|T_2\|_{\mathcal{L}_2(U, H)}$  holds for  $T_1, T_2 \in \mathcal{L}_2(U, H)$ . Finally,  $V := C(\mathcal{D}, \mathbb{R})$  represents the Banach space of all continuous

functions from  $\mathcal{D}$  to  $\mathbb{R}$  with supremum norm. Throughout this paper, we define  $P : H \rightarrow \dot{H}$  the generalized orthogonal projector by

$$Pv = v - |\mathcal{D}|^{-1} \int_{\mathcal{D}} v dx \quad (2.2)$$

and then  $(I - P)v = |\mathcal{D}|^{-1} \int_{\mathcal{D}} v dx$  is the average of  $v$ . Here and below, by  $L^r(\mathcal{D}, \mathbb{R})$ ,  $r \geq 1$  ( $L^r(\mathcal{D})$  or  $L^r$  for short) we denote a Banach space consisting of all  $r$ -times integrable functions.

In the sequel, the main assumptions are made for the abstract model (1.1).

**Assumption 2.1.** *Let  $-A$  be the Neumann Laplacian, given by  $-Au = \Delta u$ , with  $u \in \text{dom}(A) := \{v \in H^2(\mathcal{D}) \cap \dot{H} : \frac{\partial v}{\partial n} = 0 \text{ on } \partial\mathcal{D}\}$ .*

For  $v \in H$ , we extend the definition as  $Av = APv$ . Then there exists a family of orthonormal eigenbasis  $\{e_j\}_{j \in \mathbb{N}}$  with corresponding eigenvalues  $\{\lambda_j\}_{j \in \mathbb{N}}$  such that

$$Ae_j = \lambda_j e_j, \quad 0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_j \leq \dots, \quad \lambda_j \rightarrow \infty, \quad (2.3)$$

where  $e_0 = |\mathcal{D}|^{-\frac{1}{2}}$  and  $\{e_j\}_{j \in \mathbb{N}^+}$  forms an orthonormal basis of  $\dot{H}$ . We define the fractional powers of  $A$  on  $\dot{H}$  by the spectral theory, e.g.,  $A^\alpha v = \sum_{j=1}^{\infty} \lambda_j^\alpha \langle v, e_j \rangle e_j$ ,  $\alpha \in \mathbb{R}$ . The space  $\dot{H}^\alpha := \text{dom}(A^{\frac{\alpha}{2}})$  is a Hilbert space with the inner product  $\langle \cdot, \cdot \rangle_\alpha$  and the associated norm  $|\cdot|_\alpha$  defined by

$$\langle v, w \rangle_\alpha = \sum_{j=1}^{\infty} \lambda_j^\alpha \langle v, e_j \rangle \langle w, e_j \rangle, \quad |v|_\alpha = \|A^{\frac{\alpha}{2}} v\| = \left( \sum_{j=1}^{\infty} \lambda_j^\alpha |\langle v, e_j \rangle|^2 \right)^{\frac{1}{2}}, \quad \alpha \in \mathbb{R}. \quad (2.4)$$

Note that for integer  $\alpha \geq 0$ ,  $\dot{H}^\alpha$  is a subspace of  $H^\alpha(D) \cap \dot{H}$  characterized by certain boundary conditions and the norm  $|\cdot|_\alpha$  is equivalent on  $\dot{H}^\alpha$  to the standard Sobolev norm  $\|\cdot\|_{H^\alpha(D)}$  for  $\alpha = 1, 2$ . Since  $H^2(\mathcal{D})$  is an algebra, one can deduce that for any  $f, g \in \dot{H}^2$ ,

$$\|fg\|_{H^2(\mathcal{D})} \leq C \|f\|_{H^2(\mathcal{D})} \|g\|_{H^2(\mathcal{D})} \leq C |f|_2 |g|_2. \quad (2.5)$$

Additionally, the operator  $-A^2$  generates an analytic semigroup  $E(t) = e^{-tA^2}$  on  $H$ , given by

$$\begin{aligned} E(t)v &= e^{-tA^2} v = \sum_{j=0}^{\infty} e^{-t\lambda_j^2} \langle v, e_j \rangle e_j = \sum_{j=1}^{\infty} e^{-t\lambda_j^2} \langle v, e_j \rangle e_j + \langle v, e_0 \rangle e_0 \\ &= P e^{-tA^2} v + (I - P)v, \quad v \in H. \end{aligned} \quad (2.6)$$

At last, the properties of  $E(t)$  are obtained by expansion in terms of the eigenbasis of  $A$  and using Parseval's identity,

$$\|A^\mu E(t)\|_{\mathcal{L}(\dot{H})} \leq C t^{-\frac{\mu}{2}}, \quad t > 0, \quad \mu \geq 0, \quad (2.7)$$

$$\|A^{-\nu}(I - E(t))\|_{\mathcal{L}(\dot{H})} \leq C t^{\frac{\nu}{2}}, \quad t \geq 0, \quad \nu \in [0, 2], \quad (2.8)$$

$$\int_{t_1}^{t_2} \|A^\varrho E(s)v\|^2 ds \leq C |t_2 - t_1|^{1-\varrho} \|v\|^2, \quad \forall v \in \dot{H}, \varrho \in [0, 1], 0 \leq t_1 \leq t_2, \quad (2.9)$$

$$\left\| A^{2\rho} \int_{t_1}^{t_2} E(t_2 - \sigma)v d\sigma \right\| \leq C |t_2 - t_1|^{1-\rho} \|v\|, \quad \forall v \in \dot{H}, \rho \in [0, 1], 0 \leq t_1 \leq t_2. \quad (2.10)$$

**Assumption 2.2.** Let  $F : L^6(\mathcal{D}, \mathbb{R}) \rightarrow H$  be the Nemytskii operator given by

$$F(v)(x) = f(v(x)) = v^3(x) - v(x), \quad x \in \mathcal{D}, v \in L^6(\mathcal{D}, \mathbb{R}). \quad (2.11)$$

Then, for  $v, \zeta, \zeta_1, \zeta_2 \in L^6(\mathcal{D}, \mathbb{R})$ , we have

$$\begin{aligned} (F'(v)(\zeta))(x) &= f'(v(x))\zeta(x) = (3v^2(x) - 1)\zeta(x), \quad x \in \mathcal{D}, \\ (F''(v)(\zeta_1, \zeta_2))(x) &= f''(v(x))\zeta_1(x)\zeta_2(x) = 6v(x)\zeta_1(x)\zeta_2(x), \quad x \in \mathcal{D}. \end{aligned} \quad (2.12)$$

As a result, there exists a constant  $C > 0$  such that

$$-\langle F(u) - F(v), u - v \rangle \leq \|u - v\|^2, \quad u, v \in L^6(\mathcal{D}). \quad (2.13)$$

$$\|F'(v)u\| \leq C(1 + \|v\|_V^2)\|u\|, \quad u, v \in V. \quad (2.14)$$

$$\|F(u) - F(v)\| \leq C(1 + \|u\|_V^2 + \|v\|_V^2)\|u - v\|, \quad u, v \in V. \quad (2.15)$$

To simplify the presentation, we assume the average of the Wiener process to be zero so that the covariance operator  $Q$  of the Q-Wiener process belongs to  $\mathcal{L}(\dot{H})$ .

**Assumption 2.3.** Let  $\{W(t)\}_{t \in [0, T]}$  be a  $\dot{H}$ -valued ((possibly cylindrical)) Q-Wiener process with the covariance operator  $Q \in \mathcal{L}(\dot{H})$  satisfying

$$\left\| A^{\frac{\gamma-2}{2}} Q^{\frac{1}{2}} \right\|_{\mathcal{L}_2} < \infty \quad \text{for some } \gamma \in \left(\frac{d}{2}, 4\right]. \quad (2.16)$$

**Assumption 2.4.** Let  $X_0 : \Omega \rightarrow \dot{H}$  be  $\mathcal{F}_0/\mathcal{B}(\dot{H})$ -measurable and satisfy that for a sufficiently large number  $p_0 \in \mathbb{N}$ ,

$$\mathbb{E}[|X_0|_\gamma^{p_0}] < \infty, \quad (2.17)$$

where  $\gamma$  is the parameter from (2.16).

Before moving on, similar to [13, Equations (2.5), (2.7)], we give the following lemma concerning the spatio-temporal regularity results of stochastic convolution

$$\mathcal{O}_t := \int_0^t E(t-s) dW(s). \quad (2.18)$$

**Lemma 2.5.** Suppose Assumptions 2.1 and 2.3 hold. Then for  $\forall p \geq 1$ , the stochastic convolution  $\mathcal{O}_t$  satisfies

$$\mathbb{E} \left[ \sup_{t \in [0, T]} |\mathcal{O}_t|_V^p \right] + \sup_{t \in [0, T]} \mathbb{E} \left[ |\mathcal{O}_t|_\gamma^p \right] < \infty, \quad (2.19)$$

and for  $\alpha \in [0, \gamma]$ ,

$$\|\mathcal{O}_t - \mathcal{O}_s\|_{L^p(\Omega, \dot{H}^\alpha)} \leq C|t - s|^{\min\{\frac{1}{2}, \frac{\gamma-\alpha}{4}\}}. \quad (2.20)$$

At last, we consider the mild solution of (1.1) by following a semigroup approach proposed in [15]. As already proved in [26, Theorem 3.5] and [13, Proposition 6 & Proposition 7], the above assumptions are sufficient to establish well-posedness of the model (1.1) and spatio-temporal regularity of the mild solution. The relevant results are stated in Theorem 2.6 below.

**Theorem 2.6** (Well-posedness and regularity results). *Under Assumptions 2.1-2.4, there is a unique mild solution  $X : [0, T] \times \Omega \rightarrow \dot{H}$  to (1.1) given by*

$$X(t) = E(t)X_0 - \int_0^t E(t-s)APF(X(s)) ds + \int_0^t E(t-s) dW(s), \quad t \in [0, T]. \quad (2.21)$$

Furthermore, for  $\gamma \in (\frac{d}{2}, 4]$  and  $p \geq 1$ ,

$$\sup_{t \in [0, T]} \|X(t)\|_{L^p(\Omega, \dot{H}^\gamma)} < \infty, \quad (2.22)$$

and for  $\alpha \in [0, \gamma]$ ,

$$\|X(t) - X(s)\|_{L^p(\Omega, \dot{H}^\alpha)} \leq C(t-s)^{\min\{\frac{1}{2}, \frac{\gamma-\alpha}{4}\}}, \quad 0 \leq s < t \leq T. \quad (2.23)$$

### 3 Strong convergence analysis of numerical approximation

This section aims to derive strong convergence rates of the numerical discretization, done by a tamed exponential Euler method based on the spectral Galerkin approximation.

#### 3.1 The spectral Galerkin spatial discretization

We start this part by introducing a finite dimension space spanned by the first  $N$  eigenvectors of the dominant linear operator  $A$ , i.e.,  $H_N = \text{span}\{e_1, \dots, e_N\}$  and the projection operator  $P_N : \dot{H}^\beta \rightarrow H_N$  is defined by  $P_N x = \sum_{i=1}^N \langle x, e_i \rangle e_i$  for  $\forall x \in \dot{H}^\beta$ ,  $\beta \geq -2$ . Given the identity mapping  $I \in \mathcal{L}(\dot{H})$ , one can easily obtain that

$$\|(P_N - I)A^{-\alpha}\|_{\mathcal{L}(\dot{H})} \leq C\lambda_N^{-\alpha}, \quad \forall \alpha \geq 0. \quad (3.1)$$

Then applying the spectral Galerkin method to (1.1) results in the finite dimension stochastic differential equation, given by

$$dX^N(t) + A(AX^N(t) + P_N F(X^N(t)))dt = P_N dW(t), \quad t \in (0, T]; \quad X^N(0) = P_N X_0, \quad (3.2)$$

whose unique mild solution is adapted and satisfies

$$X^N(t) = E(t)P_N X_0 - \int_0^t E(t-s)AP_N F(X^N(s))ds + \int_0^t E(t-s)P_N dW(s). \quad (3.3)$$

The following theorem, concerning the strong convergence rate of the spectral Galerkin method, is an immediate consequence of [26, Theorem 3.5].

**Theorem 3.1.** *Let  $X(t)$  be the mild solution of (1.1) and let  $X^N(t)$  be the solution of (3.2). Suppose Assumptions 2.1-2.4 are valid, then for any  $p \in [1, \infty)$ , it holds that*

$$\sup_{t \in [0, T]} \|X(t) - X^N(t)\|_{L^p(\Omega, \dot{H})} \leq C\lambda_N^{-\frac{\gamma}{2}}. \quad (3.4)$$

### 3.2 An explicit fully discrete scheme and its priori moment bounds

This subsection concerns the a priori moment bounds of a spatio-temporal full discretization based on the spatial spectral Galerkin approximation. In order to introduce the fully discrete scheme, we define the nodes  $t_m = m\tau$  with a uniform time step-size  $\tau = \frac{T}{M}$  for  $m \in \{0, 1, \dots, M\}$ ,  $M \in \mathbb{N}^+$  and introduce a notation  $\lfloor t \rfloor_\tau := t_i$  for  $t \in [t_i, t_{i+1})$ ,  $i \in \{0, 1, \dots, M-1\}$ . It is worthwhile to mention that the fully discrete exponential Euler and fully discrete linear-implicit Euler approximations diverge strongly and numerically weakly in the case of stochastic Allen-Cahn equations [2]. Thus, we apply the tamed exponential Euler scheme to (3.2) and get

$$X_{t_{m+1}}^{M,N} = E(\tau)X_{t_m}^{M,N} - \frac{A^{-1}(I-E(\tau))P_N F(X_{t_m}^{M,N})}{1+\tau\|P_N F(X_{t_m}^{M,N})\|} + \int_{t_m}^{t_{m+1}} E(t_{m+1} - \lfloor s \rfloor_\tau) P_N dW(s). \quad (3.5)$$

Particularly, the following continuous version of (3.5) will be used frequently,

$$X_t^{M,N} = E(t)P_N X_0 - \int_0^t \frac{E(t-s)A P_N F(X_{\lfloor s \rfloor_\tau}^{M,N})}{1+\tau\|P_N F(X_{\lfloor s \rfloor_\tau}^{M,N})\|} ds + \mathcal{O}_t^{M,N}, \quad (3.6)$$

which is  $\mathcal{F}_t$ -adapted. Here for simplicity of presentation we denote

$$\mathcal{O}_t^{M,N} := \int_0^t E(t - \lfloor s \rfloor_\tau) P_N dW(s), \quad (3.7)$$

which satisfies the following regularity result.

**Lemma 3.2.** *Suppose Assumptions 2.1 and 2.3 hold. Then for  $\forall p \geq 1$  and  $\theta \in (\frac{d}{2}, \min\{\gamma, 2\})$ , the discrete stochastic convolution  $\mathcal{O}_t^{M,N}$  satisfies*

$$\mathbb{E} \left[ \sup_{t \in [0, T]} |\mathcal{O}_t^{M,N}|_p^\theta < \infty. \quad (3.8)$$

*Proof.* Following a similar approach used in [15, Theorem 5.10], we can rewrite  $\mathcal{O}_t^{M,N}$  as

$$\mathcal{O}_t^{M,N} = \frac{\sin(\alpha\pi)}{\pi} \int_0^t (t-s)^{\alpha-1} E(t-s) Y_\alpha(s) ds, \text{ for } \alpha \in (0, 1) \quad (3.9)$$

with

$$Y_\alpha(s) := \int_0^s (s-r)^{-\alpha} E(s-\lfloor r \rfloor) P_N dW(r). \quad (3.10)$$

Indeed, by stochastic Fubini theorem, we get

$$\begin{aligned} & \frac{\sin(\alpha\pi)}{\pi} \int_0^t (t-s)^{\alpha-1} E(t-s) Y_\alpha(s) ds \\ &= \frac{\sin(\alpha\pi)}{\pi} \int_0^t (t-s)^{\alpha-1} E(t-s) \left[ \int_0^s (s-r)^{-\alpha} E(s-\lfloor r \rfloor) P_N dW(r) \right] ds \\ &= \frac{\sin(\alpha\pi)}{\pi} \int_0^t \left[ \int_r^t (t-s)^{\alpha-1} (s-r)^{-\alpha} ds \right] E(t-\lfloor r \rfloor) P_N dW(r) \\ &= \int_0^t E(t-\lfloor r \rfloor_\tau) P_N dW(r), \end{aligned} \quad (3.11)$$

where a basic fact

$$\int_r^t (t-s)^{\alpha-1} (s-r)^{-\alpha} ds = \frac{\pi}{\sin(\alpha\pi)}, \quad 0 \leq r \leq t, \quad \alpha \in (0, 1) \quad (3.12)$$

was invoked in the last equality. As a result, using the Burkholder–Davis–Gundy inequality and Hölder’s inequality leads to

$$\begin{aligned} \mathbb{E} \left[ \sup_{t \in [0, T]} |\mathcal{O}_t^{M, N}|_\theta^p \right] &= \mathbb{E} \left[ \sup_{t \in [0, T]} \left| \frac{\sin(\alpha\pi)}{\pi} \int_0^t (t-s)^{\alpha-1} E(t-s) Y_\alpha(s) ds \right|_\theta^p \right] \\ &\leq C \mathbb{E} \left[ \sup_{t \in [0, T]} \left( \int_0^t (t-s)^{\alpha-1-\frac{\theta}{4}} |Y_\alpha(s)| ds \right)^p \right] \\ &\leq C \mathbb{E} \left[ \sup_{t \in [0, T]} \left( \int_0^t (t-s)^{(\alpha-1-\frac{\theta}{4})q} ds \right)^{\frac{p}{q}} \cdot \int_0^t |Y_\alpha(s)|^p ds \right] \\ &\leq C \int_0^T \mathbb{E} [|Y_\alpha(s)|^p] ds \\ &\leq C \int_0^T \left( \int_0^s (s-r)^{-2\alpha} \|E(s-[r]_\tau) Q^{\frac{1}{2}}\|_{\mathcal{L}_2}^2 dr \right)^{\frac{p}{2}} ds \\ &\leq C \int_0^T \left( \int_0^s (s-r)^{-2\alpha+\min\{\frac{\gamma-2}{2}, 0\}} \|A^{\frac{\gamma-2}{2}} Q^{\frac{1}{2}}\|_{\mathcal{L}_2}^2 dr \right)^{\frac{p}{2}} ds \\ &< \infty, \end{aligned} \quad (3.13)$$

where  $\alpha > \frac{1}{p} + \frac{\theta}{4}$  was used in the third inequality and  $\alpha \in (0, \min\{\frac{\gamma}{4}, \frac{1}{2}\})$  was used in the last inequality. Finally, choosing sufficiently large  $p > 1$  and  $\theta < \min\{\gamma, 2\}$  completes the proof.  $\square$

The forthcoming lemma is a direct consequence of [26, Lemma 3.2], which is crucial to the moment bound and convergence analysis.

**Lemma 3.3.** *Let  $F : L^6 \rightarrow H$  be the Nemytskii operator in Assumption 2.2. Then it holds for any  $\iota \in (\frac{1}{2}, 1)$  and  $d = 1$ ,*

$$|F'(u)v|_\iota \leq C(1 + |u|_\iota^2)|v|_1, \quad u \in \dot{H}^\iota, \quad v \in \dot{H}^1, \quad (3.14)$$

and for any  $\iota \in (\frac{d}{2}, 2)$ ,  $d = 2, 3$ ,

$$|F'(u)v|_1 \leq C(1 + |u|_\iota^2)|v|_1, \quad u \in \dot{H}^\iota, \quad v \in \dot{H}^1. \quad (3.15)$$

Next we construct a sequence of decreasing subevents

$$\Omega_{R, t_i} := \left\{ \omega \in \Omega : \sup_{j \in \{0, 1, \dots, i\}} \|X_{t_j}^{M, N}(\omega)\|_{L^6} \leq R \right\}, \quad R \in (0, \infty), \quad i \in \{0, 1, \dots, M\}. \quad (3.16)$$

By  $\Omega^c$  and  $\chi_\Omega$  we denote the complement and indicator function of a set  $\Omega$ , respectively. It is easy to confirm that  $\chi_{\Omega_{R, t_i}}$  is  $\mathcal{F}_{t_i}$ -adapted and  $\chi_{\Omega_{R, t_i}} \leq \chi_{\Omega_{R, t_j}}$  for  $t_i \geq t_j$ .



**Lemma 3.4.** Let  $p \in [1, \infty)$ ,  $R_\tau = \tau^{-\min\{\frac{1}{36}, \frac{\gamma}{36}\}}$  for  $d = 1$  and  $R_\tau = \tau^{-\min\{\frac{1}{36}, \frac{\gamma-1}{36}\}}$  for  $d = 2, 3$ . Based on Assumptions 2.1-2.4, the fully discrete solution  $X_{t_i}^{M,N}$  satisfies

$$\sup_{M,N \in \mathbb{N}^+} \sup_{i \in \{0,1,\dots,M\}} \mathbb{E}[\chi_{\Omega_{R_\tau, t_{i-1}}} \|X_{t_i}^{M,N}\|_{L^6}^p] < \infty, \quad (3.17)$$

where with the convention, we set  $\chi_{\Omega_{R_\tau, t_{-1}}} = 1$ .

*Proof.* Firstly, we introduce a process  $Y_t^{M,N}$  by

$$Y_t^{M,N} := X_t^{M,N} - \mathcal{O}_t^{M,N} = E(t)P_N X_0 - \int_0^t \frac{E(t-s)AP_N F(X_{\lfloor s \rfloor_\tau}^{M,N})}{1+\tau\|P_N F(X_{\lfloor s \rfloor_\tau}^{M,N})\|} ds. \quad (3.18)$$

Then, we can recast it as

$$Y_t^{M,N} = E(t)P_N X_0 - \int_0^t E(t-s)AP_N F(X_s^{M,N}) ds + \int_0^t E(t-s)AP_N Z_s^{M,N} ds, \quad (3.19)$$

with  $Z_t^{M,N} := F(X_t^{M,N}) - \frac{F(X_{\lfloor t \rfloor_\tau}^{M,N})}{1+\tau\|P_N F(X_{\lfloor t \rfloor_\tau}^{M,N})\|}$ . As a result, for  $t \in (0, T]$ ,  $Y_t^{M,N}$  satisfies

$$\frac{d}{dt} Y_t^{M,N} + A^2 Y_t^{M,N} + AP_N F(Y_t^{M,N} + \mathcal{O}_t^{M,N}) = AP_N Z_t^{M,N}. \quad (3.20)$$

Multiplying  $A^{-1}Y_t^{M,N}$  on both sides and integrating from 0 to  $t_i$  yield

$$\begin{aligned} & |Y_{t_i}^{M,N}|_{-1}^2 - |Y_0^{M,N}|_{-1}^2 \\ &= -2 \int_0^{t_i} |Y_s^{M,N}|_1^2 ds - 2 \int_0^{t_i} \langle F(Y_s^{M,N} + \mathcal{O}_s^{M,N}), Y_s^{M,N} \rangle ds + 2 \int_0^{t_i} \langle Z_s^{M,N}, Y_s^{M,N} \rangle ds \\ &= -2 \int_0^{t_i} |Y_s^{M,N}|_1^2 ds - 2 \int_0^{t_i} \|Y_s^{M,N}\|_{L^4}^4 ds + 2 \int_0^{t_i} \|Y_s^{M,N}\|^2 ds + 2 \int_0^{t_i} \langle Z_s^{M,N}, Y_s^{M,N} \rangle ds \\ &\quad - 2 \int_0^{t_i} \langle 3(Y_s^{M,N})^2 \mathcal{O}_s^{M,N} + 3Y_s^{M,N} (\mathcal{O}_s^{M,N})^2 + (\mathcal{O}_s^{M,N})^3 - \mathcal{O}_s^{M,N}, Y_s^{M,N} \rangle ds \\ &\leq - \int_0^{t_i} |Y_s^{M,N}|_1^2 ds - \int_0^{t_i} \|Y_s^{M,N}\|_{L^4}^4 ds + C \int_0^{t_i} |Y_s^{M,N}|_{-1}^2 ds \\ &\quad + C \int_0^{t_i} |Z_s^{M,N}|_{-1}^2 ds + C \int_0^{t_i} (1 + \|\mathcal{O}_s^{M,N}\|_{L^4}^4) ds, \end{aligned} \quad (3.21)$$

where Young's inequality was used. It follows from Gronwall's inequality that

$$|Y_{t_i}^{M,N}|_{-1}^2 \leq C \left( |Y_0^{M,N}|_{-1}^2 + \int_0^{t_i} |Z_s^{M,N}|_{-1}^2 ds + \int_0^{t_i} (1 + \|\mathcal{O}_s^{M,N}\|_{L^4}^4) ds \right), \quad (3.22)$$

which implies that

$$\int_0^{t_i} |Y_s^{M,N}|_1^2 ds + \int_0^{t_i} \|Y_s^{M,N}\|_{L^4}^4 ds \leq C \left( |Y_0^{M,N}|_{-1}^2 + \int_0^{t_i} |Z_s^{M,N}|_{-1}^2 ds + \int_0^{t_i} (1 + \|\mathcal{O}_s^{M,N}\|_{L^4}^4) ds \right). \quad (3.23)$$

Since  $\|Y_0^{M,N}\|_{L^p(\Omega, \dot{H}^\gamma)} < \infty$  and  $\|\mathcal{O}_t^{M,N}\|_{L^p(\Omega, \dot{H}^\gamma)} < \infty$ , we deduce

$$\left\| \int_0^{t_i} |Y_s^{M,N}|_1^2 ds \right\|_{L^p(\Omega, \mathbb{R})} + \left\| \int_0^{t_i} \|Y_s^{M,N}\|_{L^4}^4 ds \right\|_{L^p(\Omega, \mathbb{R})} \leq C \left( 1 + \int_0^{t_i} \|Z_s^{M,N}\|_{L^{2p}(\Omega, \dot{H}^{-1})}^2 ds \right). \quad (3.24)$$

Taking inner product of (3.20) by  $Y_t^{M,N}$  and integrating from 0 to  $t_i$  lead to

$$\begin{aligned} & \|Y_{t_i}^{M,N}\|^2 - \|Y_0^{M,N}\|^2 \\ &= -2 \int_0^{t_i} \|AY_s^{M,N}\|^2 ds - 2 \int_0^{t_i} \langle AF(Y_s^{M,N} + \mathcal{O}_s^{M,N}), Y_s^{M,N} \rangle ds + 2 \int_0^{t_i} \langle Z_s^{M,N}, AY_s^{M,N} \rangle ds \\ &\leq - \int_0^{t_i} \|AY_s^{M,N}\|^2 ds - 6 \int_0^{t_i} \|Y_s^{M,N} \nabla Y_s^{M,N}\|^2 ds + 2 \int_0^{t_i} |Y_s^{M,N}|_1^2 ds + \int_0^{t_i} \|Z_s^{M,N}\|^2 ds \\ &\quad - 2 \int_0^{t_i} \langle 3(Y_s^{M,N})^2 \mathcal{O}_s^{M,N} + 3Y_s^{M,N} (\mathcal{O}_s^{M,N})^2 + (\mathcal{O}_s^{M,N})^3 - \mathcal{O}_s^{M,N}, AY_s^{M,N} \rangle ds \\ &\leq -\frac{1}{2} \int_0^{t_i} \|AY_s^{M,N}\|^2 ds - \frac{3}{2} \int_0^{t_i} \|\nabla[(Y_s^{M,N})^2]\|^2 ds + 2 \int_0^{t_i} |Y_s^{M,N}|_1^2 ds + \int_0^{t_i} \|Z_s^{M,N}\|^2 ds \\ &\quad + C \int_0^{t_i} \left( \|Y_s^{M,N}\|_{L^4}^4 \|\mathcal{O}_s^{M,N}\|_V^2 + \|Y_s^{M,N}\|^2 \|\mathcal{O}_s^{M,N}\|_V^4 + \|\mathcal{O}_s^{M,N}\|_{L^6}^6 + \|\mathcal{O}_s^{M,N}\|^2 \right) ds. \end{aligned} \quad (3.25)$$

Again, the use of (3.23) gives

$$\begin{aligned} & \|Y_{t_i}^{M,N}\|^2 + \int_0^{t_i} \|AY_s^{M,N}\|^2 ds + \int_0^{t_i} \|\nabla[(Y_s^{M,N})^2]\|^2 ds \\ &\leq C \left( 1 + \sup_{s \in [0, T]} \|\mathcal{O}_s^{M,N}\|_V^8 \right) \left( 1 + \|Y_0^{M,N}\|^2 + \int_0^{t_i} \|Z_s^{M,N}\|^2 ds \right). \end{aligned} \quad (3.26)$$

At the moment, we turn attention to the estimate  $\mathbb{E}[\chi_{\Omega_{R\tau, t_i-1}} \|Z_s^{M,N}\|^{2p}]$ . For  $s \in [0, t_i]$ , we have

$$\begin{aligned} \chi_{\Omega_{R\tau, t_i-1}} \|Z_s^{M,N}\| &\leq \chi_{\Omega_{R\tau, t_i-1}} \|F(X_s^{M,N}) - F(X_{[s]_\tau}^{M,N})\| \\ &\quad + \chi_{\Omega_{R\tau, t_i-1}} \left\| F(X_{[s]_\tau}^{M,N}) - \frac{F(X_{[s]_\tau}^{M,N})}{1 + \tau \|P_N F(X_{[s]_\tau}^{M,N})\|} \right\| \\ &\leq C \chi_{\Omega_{R\tau, t_i-1}} \left( 1 + \|X_{[s]_\tau}^{M,N}\|_V^2 + \|X_s^{M,N}\|_V^2 \right) (\|X_s^{M,N} - X_{[s]_\tau}^{M,N}\|) \\ &\quad + C \chi_{\Omega_{R\tau, t_i-1}} \tau \|F(X_{[s]_\tau}^{M,N})\|^2. \end{aligned} \quad (3.27)$$

Before further proof, we claim that

$$\chi_{\Omega_{R\tau, t_i-1}} \|X_s^{M,N}\|_V \leq C \left( 1 + \|X_0\|_V + R_\tau^3 + \|\mathcal{O}_s^{M,N}\|_V \right), \quad \forall s \in [0, t_i]. \quad (3.28)$$

Indeed, by stability of the semigroup  $E(t)$  in  $V$  and Sobolev embedding inequality  $\dot{H}^\delta \subset V, \delta > \frac{d}{2}$ ,

$$\begin{aligned}
& \chi_{\Omega_{R\tau, t_{i-1}}} \|X_s^{M,N}\|_V \\
& \leq \chi_{\Omega_{R\tau, t_{i-1}}} \left( \|E(s)P_N X_0\|_V + \int_0^s \|E(s-r)AF(X_{[r]\tau}^{M,N})\|_V dr + \|\mathcal{O}_s^{M,N}\|_V \right) \\
& \leq \chi_{\Omega_{R\tau, t_{i-1}}} \left( \|X_0\|_V + \int_0^s (s-r)^{-\frac{2+\delta}{4}} \|PF(X_{[r]\tau}^{M,N})\| dr + \|\mathcal{O}_s^{M,N}\|_V \right) \\
& \leq C(1 + \|X_0\|_V + R_\tau^3 + \|\mathcal{O}_s^{M,N}\|_V).
\end{aligned} \tag{3.29}$$

Next, owing to (3.6), one can write

$$\begin{aligned}
X_s^{M,N} - X_{[s]\tau}^{M,N} &= [E(s) - E([s]\tau)]P_N X_0 - \int_0^s \frac{E(s-u)AP_N F(X_{[u]\tau}^{M,N})}{1+\tau\|P_N F(X_{[u]\tau}^{M,N})\|} du \\
& \quad + \int_0^{[s]\tau} \frac{E([s]\tau-u)AP_N F(X_{[u]\tau}^{M,N})}{1+\tau\|P_N F(X_{[u]\tau}^{M,N})\|} du + \mathcal{O}_s^{M,N} - \mathcal{O}_{[s]\tau}^{M,N},
\end{aligned} \tag{3.30}$$

which implies that

$$\begin{aligned}
& \chi_{\Omega_{R\tau, t_{i-1}}} \|X_s^{M,N} - X_{[s]\tau}^{M,N}\| \\
& \leq \tau^{\frac{7}{4}}|X_0|_\gamma + \chi_{\Omega_{R\tau, t_{i-1}}} \left\| \int_0^{[s]\tau} E([s]\tau - u)(E(s - [s]\tau) - I) \frac{AP_N F(X_{[u]\tau}^{M,N})}{1+\tau\|P_N F(X_{[u]\tau}^{M,N})\|} du \right\| \\
& \quad + \chi_{\Omega_{R\tau, t_{i-1}}} \int_{[s]\tau}^s \left\| \frac{E(s-u)AP_N F(X_{[u]\tau}^{M,N})}{1+\tau\|P_N F(X_{[u]\tau}^{M,N})\|} \right\| du + \chi_{\Omega_{R\tau, t_{i-1}}} \|\mathcal{O}_s^{M,N} - \mathcal{O}_{[s]\tau}^{M,N}\| \\
& \leq \tau^{\frac{7}{4}}|X_0|_\gamma + C(1 + R_\tau^3)\tau^{\frac{1}{4}} + \|\mathcal{O}_s^{M,N} - \mathcal{O}_{[s]\tau}^{M,N}\|.
\end{aligned} \tag{3.31}$$

Therefore,

$$\begin{aligned}
\chi_{\Omega_{R\tau, t_{i-1}}} \|Z_s^{M,N}\| &\leq C(1 + \|X_0\|_V^2 + R_\tau^6 + \|\mathcal{O}_s^{M,N}\|_V^2 + \|\mathcal{O}_{[s]\tau}^{M,N}\|_V^2) \\
& \quad \times \left( \tau^{\frac{7}{4}}|X_0|_\gamma + (1 + R_\tau^3)\tau^{\frac{1}{4}} + \|\mathcal{O}_s^{M,N} - \mathcal{O}_{[s]\tau}^{M,N}\| \right) + C\tau(1 + R_\tau^6),
\end{aligned} \tag{3.32}$$

Combining it with  $\mathbb{E}\left(\|\mathcal{O}_s^{M,N} - \mathcal{O}_{[s]\tau}^{M,N}\|^p\right) \leq C\tau^{\min\{\frac{1}{2}, \frac{7}{4}\}p}$  and  $R_\tau = \tau^{-\min\{\frac{1}{36}, \frac{7}{36}\}}$  leads to

$$\mathbb{E}\left[\chi_{\Omega_{R\tau, t_{i-1}}} \|Z_s^{M,N}\|^{2p}\right] < \infty. \tag{3.33}$$

As a result, we can deduce that

$$\begin{aligned}
& \left\| \sup_{s \in [0, t_i]} \chi_{\Omega_{R\tau, t_{i-1}}} \|Y_s^{M,N}\|_{L^p(\Omega, \dot{H})}^2 + \left\| \int_0^{t_i} \chi_{\Omega_{R\tau, t_{i-1}}} \|AY_s^{M,N}\|^2 ds \right\|_{L^p(\Omega, \mathbb{R})} \right. \\
& \quad \left. + \left\| \int_0^{t_i} \chi_{\Omega_{R\tau, t_{i-1}}} \|\nabla[(Y_s^{M,N})^2]\|^2 ds \right\|_{L^p(\Omega, \mathbb{R})} \right. < \infty.
\end{aligned} \tag{3.34}$$

This together with the Sobolev embedding inequality and Gagliardo-Nirenberg inequality implies that for  $d = 1$ ,

$$\begin{aligned}
& \mathbb{E} \left[ \chi_{\Omega_{R\tau, t_{i-1}}} \|Y_{t_i}^{M,N}\|_{L^6}^p \right] \\
& \leq \mathbb{E} \left[ \chi_{\Omega_{R\tau, t_{i-1}}} \|E(t_i)X_0\|_{L^6}^p \right] + \mathbb{E} \left[ \left\| \int_0^{t_i} \chi_{\Omega_{R\tau, t_{i-1}}} E(t_i - s) AP_N(Z_s^{M,N} - F(X_s^{M,N})) ds \right\|_{L^6}^p \right] \\
& \leq C + C \mathbb{E} \left[ \int_0^{t_i} (t_i - s)^{-\frac{7}{12}} (1 + \|\mathcal{O}_s^{M,N}\|_{L^6}^3 + \chi_{\Omega_{R\tau, t_{i-1}}} \|Y_s^{M,N}\|_{L^6}^3 + \chi_{\Omega_{R\tau, t_{i-1}}} \|Z_s^{M,N}\|) ds \right]^p \\
& \leq C + \mathbb{E} \left[ \int_0^{t_i} (t_i - s)^{-\frac{7}{12}} (\chi_{\Omega_{R\tau, t_{i-1}}} \|AY_s^{M,N}\|_{L^6}^{\frac{1}{2}} \|Y_s^{M,N}\|_{L^6}^{\frac{5}{2}}) ds \right]^p \tag{3.35} \\
& \leq C + C \left( \int_0^{t_i} (t_i - s)^{-\frac{7}{9}} ds \right)^p \mathbb{E} \left[ \sup_{s \in [0, t_i]} \chi_{\Omega_{R\tau, t_{i-1}}} \|Y_s^{M,N}\|_{L^6}^{\frac{10p}{3}} \right] \\
& \quad + C \mathbb{E} \left[ \int_0^{t_i} \chi_{\Omega_{R\tau, t_{i-1}}} \|AY_s^{M,N}\|_{L^6}^2 ds \right]^p \\
& < \infty.
\end{aligned}$$

For  $d = 2, 3$ , we define a Lyapunov functional  $J(u)$  by

$$J(u) = \frac{1}{2} \|\nabla u\|^2 + \int_{\mathcal{D}} \Phi(u) dx, \tag{3.36}$$

where  $\Phi$  is the primitive of  $F$ . Multiplying (3.20) by  $A^{-1}\dot{Y}_t^{M,N}$  yields

$$|\dot{Y}_t^{M,N}|_{-1}^2 + \frac{1}{2} \frac{d|Y_t^{M,N}|_1^2}{dt} + \langle F(Y_t^{M,N} + \mathcal{O}_t^{M,N}), \dot{Y}_t^{M,N} \rangle = \langle Z_t^{M,N}, \dot{Y}_t^{M,N} \rangle. \tag{3.37}$$

To proceed further, owing to Hölder's inequality, for  $\theta > \frac{d}{2}$ , we have

$$\|(Y_t^{M,N})^2 \nabla \mathcal{O}_t^{M,N}\| \leq \|(Y_t^{M,N})^2\|_{L^{\frac{2(2+\delta)}{\delta}}} \|\nabla \mathcal{O}_t^{M,N}\|_{L^{2+\delta}} \leq C |(Y_t^{M,N})^2|_1 |\mathcal{O}_t^{M,N}|_{\theta}, \tag{3.38}$$

where the Sobolev embedding inequality  $\dot{H}^{\frac{d}{2} - \frac{d}{p}} \subset L^p$  for  $p \geq 2$  was used in the last inequality and we take sufficiently small  $\delta > 0$  for  $d = 2$  and  $\delta = 1$  for  $d = 3$ . Therefore,

$$\begin{aligned}
& |P(3(Y_t^{M,N})^2 \mathcal{O}_t^{M,N} + 3Y_t^{M,N} (\mathcal{O}_t^{M,N})^2 + (\mathcal{O}_t^{M,N})^3 - \mathcal{O}_t^{M,N})|_1 \\
& \leq C (|(Y_t^{M,N})^2|_1 \|\mathcal{O}_t^{M,N}\|_V + \|(Y_t^{M,N})^2 \nabla \mathcal{O}_t^{M,N}\| + \|\nabla Y_t^{M,N}\| \|\mathcal{O}_t^{M,N}\|_V^2 \\
& \quad + \|Y_t^{M,N}\|_V \|\mathcal{O}_t^{M,N}\|_V \|\nabla \mathcal{O}_t^{M,N}\| + \|\nabla \mathcal{O}_t^{M,N}\| \|\mathcal{O}_t^{M,N}\|_V^2 + \|\nabla \mathcal{O}_t^{M,N}\|) \\
& \leq C (1 + |(Y_t^{M,N})^2|_1 + \|AY_t^{M,N}\|) (1 + |\mathcal{O}_t^{M,N}|_{\theta}^3).
\end{aligned} \tag{3.39}$$

Combining it with the fact  $\Phi'(t) = F(t)$  and Cauchy-Schwartz inequality infers that for  $\theta > \frac{d}{2}$ ,

$$\begin{aligned}
& -\langle F(Y_t^{M,N} + \mathcal{O}_t^{M,N}), \dot{Y}_t^{M,N} \rangle + \langle Z_t^{M,N}, \dot{Y}_t^{M,N} \rangle \\
& = -\langle F(Y_t^{M,N}), \dot{Y}_t^{M,N} \rangle + \langle Z_t^{M,N}, \dot{Y}_t^{M,N} \rangle \\
& \quad - \langle 3(Y_t^{M,N})^2 \mathcal{O}_t^{M,N} + 3Y_t^{M,N} (\mathcal{O}_t^{M,N})^2 + (\mathcal{O}_t^{M,N})^3 - \mathcal{O}_t^{M,N}, \dot{Y}_t^{M,N} \rangle \\
& \leq -\frac{d}{dt} \int_{\mathcal{D}} \Phi(Y_t^{M,N}) dx + |Z_t^{M,N}|_1 |\dot{Y}_t^{M,N}|_{-1} \\
& \quad + |P(3(Y_t^{M,N})^2 \mathcal{O}_t^{M,N} + 3Y_t^{M,N} (\mathcal{O}_t^{M,N})^2 + (\mathcal{O}_t^{M,N})^3 - \mathcal{O}_t^{M,N})|_1 |\dot{Y}_t^{M,N}|_{-1} \\
& \leq -\frac{d}{dt} \int_{\mathcal{D}} \Phi(Y_t^{M,N}) dx + |\dot{Y}_t^{M,N}|_{-1}^2 + \frac{1}{2} |Z_t^{M,N}|_1^2 \\
& \quad + C(1 + |(Y_t^{M,N})^2|_1^2 + \|AY_t^{M,N}\|^2)(1 + |\mathcal{O}_t^{M,N}|_\theta^6).
\end{aligned} \tag{3.40}$$

Therefore,

$$\begin{aligned}
J(Y_t^{M,N}) & \leq J(Y_0^{M,N}) + C \int_0^t |Z_s^{M,N}|_1^2 ds \\
& \quad + C \left( 1 + \int_0^t (|(Y_s^{M,N})^2|_1^2 + \|AY_s^{M,N}\|^2) ds \right) \left( 1 + \sup_{s \in [0, T]} |\mathcal{O}_s^{M,N}|_\theta^6 \right).
\end{aligned} \tag{3.41}$$

Applying (3.34) and Lemma 3.2 infers that

$$\mathbb{E}[(\chi_{\Omega_{R_\tau, t_{i-1}}} J(Y_{t_i}^{M,N}))^p] \leq C \left( 1 + \mathbb{E} \left[ \int_0^{t_i} \chi_{\Omega_{R_\tau, t_{i-1}}} |Z_s^{M,N}|_1^2 ds \right]^p \right). \tag{3.42}$$

In the same manner, by definition of  $X_t^{M,N}$  in (3.6), we can easily check that

$$\begin{aligned}
& \|\chi_{\Omega_{R_\tau, t_{i-1}}} X_s^{M,N}\|_{L^{4p}(\Omega, \dot{H}^\kappa)} \leq C(1 + R_\tau^3) \\
& \|\chi_{\Omega_{R_\tau, t_{i-1}}} (X_s^{M,N} - X_{[s]_\tau}^{M,N})\|_{L^{2p}(\Omega, \dot{H}^1)} \leq C \tau^{\min\{\frac{1}{4}, \frac{\gamma-1}{4}\}} (1 + R_\tau^3) \\
& \|\chi_{\Omega_{R_\tau, t_{i-1}}} F(X_{[s]_\tau}^{M,N})\|_{L^{2p}(\Omega, \dot{H}^1)} \leq C(1 + R_\tau^9).
\end{aligned} \tag{3.43}$$

Finally, it suffices to bound  $\mathbb{E}[\chi_{\Omega_{R_\tau, t_{i-1}}} |Z_s^{M,N}|_1^p]$  for  $s \in [0, t_i]$ . Taking  $R_\tau = \tau^{-\min\{\frac{1}{36}, \frac{\gamma-1}{36}\}}$  yields

$$\begin{aligned}
& \|\chi_{\Omega_{R_\tau, t_{i-1}}} Z_s^{M,N}\|_{L^p(\Omega, \dot{H}^1)} \leq \|\chi_{\Omega_{R_\tau, t_{i-1}}} (F(X_s^{M,N}) - F(X_{[s]_\tau}^{M,N}))\|_{L^p(\Omega, \dot{H}^1)} \\
& \quad + \left\| \chi_{\Omega_{R_\tau, t_{i-1}}} \left( F(X_{[s]_\tau}^{M,N}) - \frac{F(X_{[s]_\tau}^{M,N})}{1 + \tau \|P_N F(X_{[s]_\tau}^{M,N})\|} \right) \right\|_{L^p(\Omega, \dot{H}^1)} \\
& \leq C \|\chi_{\Omega_{R_\tau, t_{i-1}}} (X_s^{M,N} - X_{[s]_\tau}^{M,N})\|_{L^{2p}(\Omega, \dot{H}^1)} \left( 1 + \|\chi_{\Omega_{R_\tau, t_{i-1}}} X_{[s]_\tau}^{M,N}\|_{L^{4p}(\Omega, \dot{H}^\kappa)}^2 \right) \\
& \quad + \|\chi_{\Omega_{R_\tau, t_{i-1}}} X_s^{M,N}\|_{L^{4p}(\Omega, \dot{H}^\kappa)}^2 \\
& \quad + C \tau \|\chi_{\Omega_{R_\tau, t_{i-1}}} F(X_{[s]_\tau}^{M,N})\|_{L^{2p}(\Omega, \dot{H}^1)} \|\chi_{\Omega_{R_\tau, t_{i-1}}} F(X_{[s]_\tau}^{M,N})\|_{L^{2p}(\Omega, \dot{H}^1)} \\
& \leq C \tau^{\min\{\frac{1}{4}, \frac{\gamma-1}{4}\}} (1 + R_\tau^9) + C \tau (1 + R_\tau^{12}) \\
& < \infty.
\end{aligned} \tag{3.44}$$

The above estimate in combination with the fact  $|Y_{t_i}^{M,N}|_1^2 \leq 2J(Y_{t_i}^{M,N})$  yields for  $d = 2, 3$ ,

$$\|\chi_{\Omega_{R\tau, t_{i-1}}} Y_{t_i}^{M,N}\|_{L^p(\Omega, \dot{H}^1)} < \infty. \quad (3.45)$$

Gathering (3.35), (3.45), Sobolev embedding inequality  $\dot{H}^1 \subset L^6$ ,  $d = 2, 3$  and Lemma 3.2 together completes the proof.  $\square$

By adopting the similar argument in [28, Theorem 4.6], we can obtain a priori moment bound of  $\|X_{t_m}^{M,N}\|_{L^6}$  via Markov's inequality.

**Theorem 3.5.** *Under Assumptions 2.1-2.4, it holds that for any  $p \geq 1$ ,*

$$\sup_{M, N \in \mathbb{N}^+} \sup_{m \in \{0, 1, \dots, M\}} \mathbb{E} \left[ \|X_{t_m}^{M,N}\|_{L^6}^p \right] < \infty. \quad (3.46)$$

With Theorem 3.5 at hand, it is trivial to verify the regularity of  $X_t^{M,N}$  in the coming corollaries.

**Corollary 3.6.** *Let Assumptions 2.1-2.4 be fulfilled, then for any  $p \geq 1$ , we have*

$$\sup_{M, N \in \mathbb{N}^+} \sup_{t \in [0, T]} \mathbb{E} [ |X_t^{M,N}|_\gamma^p ] < \infty. \quad (3.47)$$

**Corollary 3.7.** *Let Assumptions 2.1-2.4 be fulfilled, then for any  $p \geq 1$  and  $\beta \in [0, \gamma]$ , there exists a constant  $C > 0$  such that*

$$\sup_{M, N \in \mathbb{N}^+} \|X_t^{M,N} - X_s^{M,N}\|_{L^p(\Omega, \dot{H}^\beta)} \leq C (t - s)^{\min\{\frac{1}{2}, \frac{\gamma - \beta}{4}\}}, \quad 0 \leq s < t \leq T. \quad (3.48)$$

### 3.3 Strong convergence rate of the fully discrete scheme

In this subsection, we are well prepared to analyze the strong convergence rate of the tamed exponential Euler method.

**Theorem 3.8** (Strong convergence rate of temporal semi-discretization). *Suppose Assumptions 2.1-2.4 are valid. Let  $X^N(t)$  and  $X_t^{M,N}$  be given by (3.3) and (3.6), respectively. Then for  $\forall p \geq 1$  we have*

$$\sup_{M, N \in \mathbb{N}^+} \sup_{t \in [0, T]} \|X^N(t) - X_t^{M,N}\|_{L^p(\Omega, \dot{H})} \leq C \tau^{\frac{p}{4}}. \quad (3.49)$$

*Proof.* Firstly, we introduce an auxiliary process,

$$\tilde{X}_t^{M,N} = E(t)P_N X_0 - \int_0^t E(t-s)AP_N F(X_s^{M,N})ds + \int_0^t E(t-s)P_N dW(s). \quad (3.50)$$

According to the uniform moment bounds of  $X_s^{M,N}$ , we can follow a standard approach to obtain  $\|\tilde{X}_t^{M,N}\|_{L^p(\Omega, \dot{H}^\gamma)} < \infty$  for  $\forall t \in [0, T]$ . Then we can separate  $\|X^N(t) - X_t^{M,N}\|_{L^p(\Omega, \dot{H})}$  into two terms:

$$\|X^N(t) - X_t^{M,N}\|_{L^p(\Omega, \dot{H})} \leq \|\tilde{X}_t^{M,N} - X_t^{M,N}\|_{L^p(\Omega, \dot{H})} + \|X^N(t) - \tilde{X}_t^{M,N}\|_{L^p(\Omega, \dot{H})}. \quad (3.51)$$

To treat the first error term  $\|\tilde{X}_t^{M,N} - X_t^{M,N}\|_{L^p(\Omega, \dot{H})}$ , we decompose it into three further parts,

$$\begin{aligned}
& \|\tilde{X}_t^{M,N} - X_t^{M,N}\|_{L^p(\Omega, \dot{H})} \\
&= \left\| \int_0^t \frac{E(t-s)AP_N F(X_{[s]_\tau}^{M,N})}{1+\tau\|P_N F(X_{[s]_\tau}^{M,N})\|} ds - \int_0^t E(t-s)AP_N F(X_s^{M,N}) ds \right\|_{L^p(\Omega, \dot{H})} \\
&+ \left\| \int_0^t E(t-s)P_N dW(s) - \int_0^t E(t-[s]_\tau)P_N dW(s) \right\|_{L^p(\Omega, \dot{H})} \\
&\leq \left\| \int_0^t E(t-s)AP_N (F(X_s^{M,N}) - F(X_{[s]_\tau}^{M,N})) ds \right\|_{L^p(\Omega, \dot{H})} \\
&+ \left\| \int_0^t E(t-s)AP_N F(X_{[s]_\tau}^{M,N}) ds - \int_0^t \frac{E(t-s)AP_N F(X_{[s]_\tau}^{M,N})}{1+\tau\|P_N F(X_{[s]_\tau}^{M,N})\|} ds \right\|_{L^p(\Omega, \dot{H})} \\
&+ \left\| \int_0^t E(t-s)P_N dW(s) - \int_0^t E(t-[s]_\tau)P_N dW(s) \right\|_{L^p(\Omega, \dot{H})} \\
&=: J_1 + J_2 + J_3.
\end{aligned} \tag{3.52}$$

By using the Taylor formula and mild form of  $X_t^{M,N}$ , we divide  $J_1$  into four terms,

$$\begin{aligned}
J_1 &\leq \left\| \int_0^t E(t-s)AP_N F'(X_{[s]_\tau}^{M,N})(E(s-[s]_\tau) - I)X_{[s]_\tau}^{M,N} ds \right\|_{L^p(\Omega, \dot{H})} \\
&+ \left\| \int_0^t E(t-s)AP_N F'(X_{[s]_\tau}^{M,N}) \int_{[s]_\tau}^s E(s-r)AP_N F(X_{[r]_\tau}^{M,N}) dr ds \right\|_{L^p(\Omega, \dot{H})} \\
&+ \left\| \int_0^t E(t-s)AP_N F'(X_{[s]_\tau}^{M,N}) \int_{[s]_\tau}^s E(s-[r]_\tau)P_N dW(r) ds \right\|_{L^p(\Omega, \dot{H})} \\
&+ \left\| \int_0^t E(t-s)AP_N \int_0^1 F''(\lambda(X_s^{M,N}, X_{[s]_\tau}^{M,N})) \right. \\
&\quad \left. \times (X_s^{M,N} - X_{[s]_\tau}^{M,N}, X_s^{M,N} - X_{[s]_\tau}^{M,N})(1-\lambda) d\lambda ds \right\|_{L^p(\Omega, \dot{H})} \\
&=: J_{11} + J_{12} + J_{13} + J_{14},
\end{aligned} \tag{3.53}$$

where  $\lambda(X_s^{M,N}, X_{[s]_\tau}^{M,N}) := X_{[s]_\tau}^{M,N} + \lambda(X_s^{M,N} - X_{[s]_\tau}^{M,N})$ .

Next we treat the above four terms separately. It follows from (2.7), (2.8), (2.14) and Corollary 3.6 that

$$\begin{aligned}
J_{11} &\leq \int_0^t \left\| E(t-s)AP_N F'(X_{[s]_\tau}^{M,N})(E(s-[s]_\tau) - I)X_{[s]_\tau}^{M,N} \right\|_{L^p(\Omega, \dot{H})} ds \\
&\leq C \int_0^t (t-s)^{-\frac{1}{2}} \|F'(X_{[s]_\tau}^{M,N})(E(s-[s]_\tau) - I)X_{[s]_\tau}^{M,N}\|_{L^p(\Omega, \dot{H})} ds \\
&\leq C \int_0^t (t-s)^{-\frac{1}{2}} (1 + \|X_{[s]_\tau}^{M,N}\|_{L^{4p}(\Omega, \dot{H}^\gamma)}^2) \\
&\quad \times \|(E(s-[s]_\tau) - I)A^{-\frac{\gamma}{2}}A^{\frac{\gamma}{2}}X_{[s]_\tau}^{M,N}\|_{L^{2p}(\Omega, \dot{H})} ds \\
&\leq C \tau^{\frac{\gamma}{4}}.
\end{aligned} \tag{3.54}$$

For the error term  $J_{12}$ , we apply (2.7), (2.14) and Corollary 3.6 to deduce that for  $\gamma \in (\frac{d}{2}, 2)$ ,

$$\begin{aligned}
J_{12} &\leq \int_0^t \left\| E(t-s)AP_N F'(X_{[s]_\tau}^{M,N}) \int_{[s]_\tau}^s E(s-r)AP_N F(X_{[r]_\tau}^{M,N})dr \right\|_{L^p(\Omega, \dot{H})} ds \\
&\leq \int_0^t \int_{[s]_\tau}^s (t-s)^{-\frac{1}{2}} \|F'(X_{[s]_\tau}^{M,N})E(s-r)AP_N F(X_{[r]_\tau}^{M,N})\|_{L^p(\Omega, \dot{H})} dr ds \\
&\leq \int_0^t \int_{[s]_\tau}^s (t-s)^{-\frac{1}{2}} (1 + \|X_{[s]_\tau}^{M,N}\|_{L^{4p}(\Omega, \dot{H}^\gamma)}^2) (s-r)^{-\frac{1}{2}} \|PF(X_{[r]_\tau}^{M,N})\|_{L^{2p}(\Omega, \dot{H})} dr ds \quad (3.55) \\
&\leq C \tau^{\frac{1}{2}} \int_0^t (t-s)^{-\frac{1}{2}} ds (1 + \sup_{s \in [0, T]} \|X_{[s]_\tau}^{M,N}\|_{L^{4p}(\Omega, \dot{H}^\gamma)}^2) \sup_{r \in [0, T]} \|PF(X_{[r]_\tau}^{M,N})\|_{L^{2p}(\Omega, \dot{H})} \\
&\leq C \tau^{\frac{1}{2}},
\end{aligned}$$

and for  $\gamma \in [2, 4]$ ,

$$\begin{aligned}
J_{12} &\leq \int_0^t \left\| E(t-s)AP_N F'(X_{[s]_\tau}^{M,N}) \int_{[s]_\tau}^s E(s-r)AP_N F(X_{[r]_\tau}^{M,N})dr \right\|_{L^p(\Omega, \dot{H})} ds \\
&\leq \int_0^t \int_{[s]_\tau}^s (t-s)^{-\frac{1}{2}} \|F'(X_{[s]_\tau}^{M,N})E(s-r)AP_N F(X_{[r]_\tau}^{M,N})\|_{L^p(\Omega, \dot{H})} dr ds \\
&\leq \int_0^t \int_{[s]_\tau}^s (t-s)^{-\frac{1}{2}} (1 + \|X_{[s]_\tau}^{M,N}\|_{L^{4p}(\Omega, \dot{H}^\gamma)}^2) \|PF(X_{[r]_\tau}^{M,N})\|_{L^{2p}(\Omega, \dot{H}^2)} dr ds \quad (3.56) \\
&\leq C \tau \int_0^t (t-s)^{-\frac{1}{2}} ds (1 + \sup_{s \in [0, T]} \|X_{[s]_\tau}^{M,N}\|_{L^{4p}(\Omega, \dot{H}^\gamma)}^2) \sup_{r \in [0, T]} \|PF(X_{[r]_\tau}^{M,N})\|_{L^{2p}(\Omega, \dot{H}^2)} \\
&\leq C \tau,
\end{aligned}$$

where in the last step, (2.5) was used. Concerning  $J_{13}$ , we have

$$\begin{aligned}
J_{13} &= \left\| \int_0^t E(t-s)AP_N F'(X_{[s]_\tau}^{M,N}) \int_{[s]_\tau}^s E(s-[r]_\tau)P_N dW(r) ds \right\|_{L^p(\Omega, \dot{H})} \\
&\leq \left\| \int_0^{[t]_\tau} E(t-s)AP_N F'(X_{[s]_\tau}^{M,N}) \int_{[s]_\tau}^s E(s-[r]_\tau)P_N dW(r) ds \right\|_{L^p(\Omega, \dot{H})} \\
&\quad + \left\| \int_{[t]_\tau}^t E(t-s)AP_N F'(X_{[s]_\tau}^{M,N}) \int_{[s]_\tau}^s E(s-[r]_\tau)P_N dW(r) ds \right\|_{L^p(\Omega, \dot{H})} \\
&=: J_{131} + J_{132}.
\end{aligned} \quad (3.57)$$

Without loss of generality, we assume that there exists an integer  $m \in \mathbb{N}$  such that  $[t]_\tau = t_m$  and then apply the stochastic Fubini theorem, the Burkholder-Davis-Gundy-type inequality and the



Hölder inequality to obtain

$$\begin{aligned}
J_{131} &= \left\| \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} E(t-s) AP_N F'(X_{t_k}^{M,N}) \int_{t_k}^s E(s-[r]_\tau) dW(r) ds \right\|_{L^p(\Omega, \dot{H})} \\
&= \left\| \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} \int_{t_k}^{t_{k+1}} \chi_{[t_k, s)}(r) E(t-s) AP_N F'(X_{t_k}^{M,N}) E(s-[r]_\tau) ds dW(r) \right\|_{L^p(\Omega, \dot{H})} \\
&\leq \left( \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} \left\| \int_{t_k}^{t_{k+1}} \chi_{[t_k, s)}(r) E(t-s) AP_N F'(X_{t_k}^{M,N}) E(s-[r]_\tau) Q^{\frac{1}{2}} ds \right\|_{L^p(\Omega, \mathcal{L}_2)}^2 dr \right)^{\frac{1}{2}} \\
&\leq C \tau^{\frac{1}{2}} \left( \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} \sum_{l=1}^{\infty} \|E(t-s) AF'(X_{t_k}^{M,N}) E(s-[r]_\tau) Q^{\frac{1}{2}} \eta_l\|_{L^p(\Omega, \dot{H})}^2 ds dr \right)^{\frac{1}{2}}.
\end{aligned} \tag{3.58}$$

Further, by using (2.7), Lemma 3.3, Corollary 3.6 and (2.16), one can find that for  $\gamma \in (1, 4]$  and  $\kappa = \frac{3}{4}, d = 1$  and  $\kappa = 1, d = 2, 3$ ,

$$\begin{aligned}
J_{131} &\leq C \tau^{\frac{1}{2}} \left( \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} \int_{t_k}^{t_{k+1}} \sum_{l=1}^{\infty} (t-s)^{-\frac{2-\kappa}{2}} \|A^{\frac{\kappa}{2}} F'(X_{t_k}^{M,N}) E(s-[r]_\tau) Q^{\frac{1}{2}} \eta_l\|_{L^p(\Omega, \dot{H})}^2 ds dr \right)^{\frac{1}{2}} \\
&\leq C \tau^{\frac{1}{2}} \left( \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} (t-s)^{-\frac{2-\kappa}{2}} (1 + \|X_{t_k}^{M,N}\|_{L^{2p}(\Omega, \dot{H}^\gamma)}^4) \right. \\
&\quad \times \left. \int_{t_k}^{t_{k+1}} \sum_{l=1}^{\infty} \|A^{\frac{1}{2}} A^{\frac{2-\gamma}{2}} E(s-[r]_\tau) A^{\frac{\gamma-2}{2}} Q^{\frac{1}{2}} \eta_l\|_{L^p(\Omega, \dot{H})}^2 dr ds \right)^{\frac{1}{2}} \\
&\leq C \tau^{\frac{4-\max\{3-\gamma, 0\}}{4}} \left( \sum_{k=0}^m \int_{t_k}^{t_{k+1}} (t-s)^{-\frac{2-\kappa}{2}} ds \right)^{\frac{1}{2}} \|A^{\frac{\gamma-2}{2}} Q^{\frac{1}{2}}\|_{\mathcal{L}_2} \\
&\leq C \tau^{\frac{4-\max\{3-\gamma, 0\}}{4}}.
\end{aligned} \tag{3.59}$$

For  $\gamma \in (\frac{1}{2}, 1]$ , from  $[t]_\tau = t_m$ , (2.7), (2.14), (2.16), Corollary 3.6 and Itô's isometry, we deduce

$$\begin{aligned}
J_{131} &= \left\| \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} E(t-s) AP_N F'(X_{t_k}^{M,N}) \int_{t_k}^s E(s-[r]_\tau) dW(r) ds \right\|_{L^p(\Omega, \dot{H})} \\
&\leq \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} (t-s)^{-\frac{1}{2}} (1 + \|X_{t_k}^{M,N}\|_{L^{2p}(\Omega, V)}) \left\| \int_{t_k}^s E(s-[r]_\tau) dW(r) \right\|_{L^{2p}(\Omega, \dot{H})} ds \\
&\leq C \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} (t-s)^{-\frac{1}{2}} \left( \int_{t_k}^s \|E(s-[r]_\tau)\|_{\mathcal{L}_2}^2 ds \right)^{\frac{1}{2}} ds \\
&\leq C \tau^{\frac{\gamma}{4}}.
\end{aligned} \tag{3.60}$$

In what follows, we use the same argument to estimate  $J_{132}$ ,

$$\begin{aligned}
J_{132} &\leq \int_{[t]_\tau}^t \left\| E(t-s)AP_N F'(X_{[s]_\tau}^{M,N}) \int_{[s]_\tau}^s E(s-r)P_N dW(r) \right\|_{L^p(\Omega, \dot{H})} ds \\
&\leq C \int_{[t]_\tau}^t (t-s)^{-\frac{1}{2}} ds \left( 1 + \sup_{s \in [0, T]} \|X_s^{M,N}\|_{L^{2p}(\Omega, \dot{H}^\gamma)}^2 \right) \tau^{\frac{\min\{\gamma, 2\}}{4}} \\
&\leq C \tau^{\frac{2+\min\{\gamma, 2\}}{4}}.
\end{aligned} \tag{3.61}$$

Owing to the fact  $L^1 \subset \dot{H}^{-\delta_0}$  with  $\delta_0 \in (\frac{3}{2}, 2)$  and the regularity of  $X_t^{M,N}$ , we arrive at,

$$\begin{aligned}
J_{14} &\leq C \int_0^t (t-s)^{-\frac{2+\delta_0}{4}} \left\| \int_0^1 F''(\lambda(X_s^{M,N}, X_{[s]_\tau}^{M,N})) \right. \\
&\quad \times (X_s^{M,N} - X_{[s]_\tau}^{M,N}, X_s^{M,N} - X_{[s]_\tau}^{M,N})(1-\lambda) d\lambda \left. \right\|_{L^p(\Omega, L^1)} ds \\
&\leq C \int_0^t \int_0^1 (t-s)^{-\frac{2+\delta_0}{4}} \|X_s^{M,N} - X_{[s]_\tau}^{M,N}\|_{L^{4p}(\Omega, \dot{H})}^2 \|\lambda(X_s^{M,N}, X_{[s]_\tau}^{M,N})\|_{L^{2p}(\Omega, V)} d\lambda ds \\
&\leq C \tau^{\min\{1, \frac{\gamma}{2}\}} \left( 1 + \sup_{s \in [0, T]} \|X_s^{M,N}\|_{L^{2p}(\Omega, \dot{H}^\gamma)} \right) \int_0^t (t-s)^{-\frac{2+\delta_0}{4}} ds \\
&\leq C \tau^{\min\{1, \frac{\gamma}{2}\}}.
\end{aligned} \tag{3.62}$$

Combining the above estimates together leads to

$$J_1 \leq C \tau^{\frac{\gamma}{4}}. \tag{3.63}$$

Due to the regularity of  $X_t^{M,N}$  in Corollary 3.6 and properties of nonlinear term  $F$ , we obtain

$$\begin{aligned}
J_2 &= \left\| \int_0^t E(t-s)AP_N F(X_{[s]_\tau}^{M,N}) ds - \int_0^t \frac{E(t-s)AP_N F(X_{[s]_\tau}^{M,N})}{1+\tau\|P_N F(X_{[s]_\tau}^{M,N})\|} ds \right\|_{L^p(\Omega, \dot{H})} \\
&\leq C \tau \int_0^t (t-s)^{-\frac{1}{2}} ds \sup_{s \in [0, T]} \|PF(X_s^{M,N})\|_{L^{2p}(\Omega, \dot{H})}^2 \\
&\leq C \tau.
\end{aligned} \tag{3.64}$$

It remains to estimate  $J_3$  by virtue of (2.8), (2.9) and (2.16),

$$\begin{aligned}
J_3 &= \left\| \int_0^t E(t-s)(I - E(s - [s]_\tau)) dW(s) \right\|_{L^p(\Omega, \dot{H})} \\
&= \left( \int_0^t \|AE(t-s)A^{-\frac{\gamma}{2}}(I - E(s - [s]_\tau))A^{\frac{\gamma-2}{2}}Q^{\frac{1}{2}}\|_{\mathcal{L}_2}^2 ds \right)^{\frac{1}{2}} \\
&\leq C \tau^{\frac{\gamma}{4}}.
\end{aligned} \tag{3.65}$$

Therefore, the estimates of  $J_1$ ,  $J_2$  and  $J_3$  imply

$$\|\tilde{X}_t^{M,N} - X_t^{M,N}\|_{L^p(\Omega, \dot{H})} \leq C \tau^{\frac{\gamma}{4}}. \tag{3.66}$$

At the moment, we start to estimate the error term  $\|X^N(t) - \tilde{X}_t^{M,N}\|_{L^p(\Omega, \dot{H})}$ . For short, by  $e^{M,N}(t)$  we denote  $X^N(t) - \tilde{X}_t^{M,N}$ , which satisfies

$$\frac{d}{dt}e^{M,N}(t) + A^2e^{M,N}(t) = AP_N(F(X_t^{M,N}) - F(X^N(t))). \quad (3.67)$$

Multiplying  $A^{-1}e^{M,N}(t)$  on both sides and using (2.13), (2.15) and Hölder's inequality lead to

$$\begin{aligned} \frac{1}{2}\frac{d}{dt}|e^{M,N}(t)|_{-1}^2 + |e^{M,N}(t)|_1^2 &= \langle e^{M,N}(t), F(X_t^{M,N}) - F(X^N(t)) \rangle \\ &= \langle e^{M,N}(t), F(\tilde{X}_t^{M,N}) - F(X^N(t)) \rangle + \langle e^{M,N}(t), F(X_t^{M,N}) - F(\tilde{X}_t^{M,N}) \rangle \\ &\leq \frac{3}{2}\|e^{M,N}(t)\|^2 + \frac{1}{2}\|F(X_t^{M,N}) - F(\tilde{X}_t^{M,N})\|^2 \\ &\leq \frac{1}{2}|e^{M,N}(t)|_1^2 + \frac{9}{8}|e^{M,N}(t)|_{-1}^2 + C\|X_t^{M,N} - \tilde{X}_t^{M,N}\|^2(1 + \|X_t^{M,N}\|_V^4 + \|\tilde{X}_t^{M,N}\|_V^4). \end{aligned} \quad (3.68)$$

Based on the Gronwall inequality and taking expectation, we achieve

$$\left\| \int_0^t |e^{M,N}(s)|_1^2 ds \right\|_{L^p(\Omega, \mathbb{R})} \leq C\tau^{\frac{7}{2}}, \quad (3.69)$$

where the regularity of  $\tilde{X}_t^{M,N}$  and  $X_t^{M,N}$  and (3.66) were also used. Furthermore, we decompose  $\|e^{M,N}(t)\|_{L^p(\Omega, \dot{H})}$  as follows,

$$\begin{aligned} \|e^{M,N}(t)\|_{L^p(\Omega, \dot{H})} &= \left\| \int_0^t E(t-s)A(P_N F(X_s^{M,N}) - P_N F(X^N(s))) ds \right\|_{L^p(\Omega, \dot{H})} \\ &\leq \int_0^t \|E(t-s)A(F(X_s^{M,N}) - F(\tilde{X}_s^{M,N}))\|_{L^p(\Omega, \dot{H})} ds \\ &\quad + \left\| \int_0^t E(t-s)A(F(\tilde{X}_s^{M,N}) - F(X^N(s))) ds \right\|_{L^p(\Omega, \dot{H})} \\ &=: K_1 + K_2. \end{aligned} \quad (3.70)$$

Thanks to the regularity of  $\tilde{X}_t^{M,N}$  and  $X_t^{M,N}$ , (2.7) and (3.66), one can show

$$\begin{aligned} K_1 &\leq \int_0^t (t-s)^{-\frac{1}{2}} \|F(X_s^{M,N}) - F(\tilde{X}_s^{M,N})\|_{L^p(\Omega, \dot{H})} ds \\ &\leq \int_0^t (t-s)^{-\frac{1}{2}} \|X_s^{M,N} - \tilde{X}_s^{M,N}\|_{L^{2p}(\Omega, \dot{H})} \left(1 + \|\tilde{X}_s^{M,N}\|_{L^{4p}(\Omega, V)}^2 + \|X_s^{M,N}\|_{L^{4p}(\Omega, V)}^2\right) ds \\ &\leq C\tau^{\frac{7}{4}}. \end{aligned} \quad (3.71)$$

Resorting to (2.7), Lemma 3.3, the regularity of  $\tilde{X}_t^{M,N}$  and  $X^N(t)$  and Hölder's inequality, we

acquire for  $\eta = \min\{\gamma, \frac{3}{4}\}$ ,

$$\begin{aligned}
K_2 &\leq C \left\| \int_0^t (t-s)^{-\frac{2-\eta}{4}} \|A^{\frac{\eta}{2}}(F(\tilde{X}_s^{M,N}) - F(X^N(s)))\| ds \right\|_{L^p(\Omega, \mathbb{R})} \\
&\leq C \left\| \int_0^t (t-s)^{-\frac{2-\eta}{4}} |e^{M,N}(s)|_1 \left(1 + |X^N(s)|_\gamma^2 + |\tilde{X}_s^{M,N}|_\gamma^2\right) ds \right\|_{L^p(\Omega, \mathbb{R})} \\
&\leq C \left\| \left( \int_0^t |e^{M,N}(s)|_1^2 ds \right)^{\frac{1}{2}} \left( \int_0^t (t-s)^{-\frac{2-\eta}{2}} (1 + |X^N(s)|_\gamma^4 + |\tilde{X}_s^{M,N}|_\gamma^4) ds \right)^{\frac{1}{2}} \right\|_{L^p(\Omega, \mathbb{R})} \\
&\leq C \left\| \int_0^t |e^{M,N}(s)|_1^2 ds \right\|_{L^p(\Omega, \mathbb{R})}^{\frac{1}{2}} \\
&\leq C \tau^{\frac{7}{4}}.
\end{aligned} \tag{3.72}$$

Collecting all the estimates obtained so far finishes the proof.  $\square$

At last, gathering Theorem 3.8 with Theorem 3.1, we get the strong convergence rates of the fully discrete scheme (3.5).

**Corollary 3.9** (Strong convergence rates of the full discretization). *Let Assumptions 2.1-2.4 be satisfied. Then for  $p \geq 1$  it holds that*

$$\sup_{M,N \in \mathbb{N}^+} \sup_{m \in \{0,1,\dots,M\}} \|X(t_m) - X_{t_m}^{M,N}\|_{L^p(\Omega, \dot{H})} \leq C(\lambda_N^{-\frac{\gamma}{2}} + \tau^{\frac{\gamma}{4}}). \tag{3.73}$$

## 4 Numerical experiments

In this section, we include some numerical results to confirm the above assertions. Let us consider the stochastic Cahn-Hilliard equation in one spatial dimension, given by

$$\begin{cases} \frac{\partial u}{\partial t} = \frac{\partial^2 w}{\partial x^2} + \dot{W}, & (t, x) \in (0, T] \times (0, 1), \\ w = -\frac{\partial^2 u}{\partial x^2} - u + u^3, & x \in (0, 1), \\ u(0, x) = \sqrt{2} \cos(\pi x), & x \in (0, 1), \\ \frac{\partial u}{\partial x} \Big|_{x=0} = \frac{\partial u}{\partial x} \Big|_{x=1} = 0, & t \in (0, T], \\ \frac{\partial w}{\partial x} \Big|_{x=0} = \frac{\partial w}{\partial x} \Big|_{x=1} = 0, & t \in (0, T], \end{cases} \tag{4.1}$$

where  $\{W(t)\}_{t \in [0, T]}$  is a cylindrical  $Q$ -Wiener process. In what follows, the fully discrete method (3.5) is used to approximate the numerical example (4.1) and the error bounds are measured in the mean-square sense at the endpoint  $T = 1$ . Note that the expectations are approximated by computing averages over 1000 samples. Since the exact solutions are not available at hand, fixing  $N = 1000$ , the reference solution is identified with a very small time stepsize  $\tau_{exact} = 2^{-15}$ . Five different time stepsizes  $\tau = 2^{-j}$ ,  $j = 9, 10, 11, 12$  are then used to carry out the numerical simulations. Moreover, we choose three different kinds of noise and present the resulting mean-square errors in Fig. 1–Fig. 3, respectively.

We are now ready to make some explanations on the numerical results. Firstly, for the white noise case (i.e.,  $Q = I$ ), the condition (2.16) in Assumption 2.3 is then fulfilled with  $\gamma$  closing to




Figure 1: strong convergence rate of the tamed Euler method (white noise).




Figure 2: strong convergence rate of the tamed Euler method (trace-class noise).

$\frac{3}{2}$  and the convergence order obtained in Theorem 3.8 is almost  $\frac{3}{8}$ . The mean-square errors are depicted in Fig. 1, against  $\tau$  on a log-log scale, where one can observe that the resulting numerical errors decrease at a slope close to  $\frac{3}{8}$ . This coincides with the theoretical result. Secondly, for the trace-class noise case, we choose  $Q$  such that

$$Qe_1 = 0, Qe_i = \frac{1}{i \log(i)^2} e_i, \forall i \geq 2. \quad (4.2)$$

Obviously, (4.2) guarantees  $\text{Tr}(Q) < \infty$  and thus the condition (2.16) is satisfied with  $\gamma = 2$ . As expected, convergence rates of  $\frac{1}{2}$  is detected in Fig. 2, which is consistent with the finding in Theorem 3.8. Finally, smoother noise are taken with the covariance operator  $Q$  satisfying

$$Qe_1 = 0, Qe_i = \frac{1}{i^5 \log(i)^2} e_i, \forall i \geq 2. \quad (4.3)$$

In this case, condition (2.16) holds with  $\gamma = 4$  and the obtained convergence rate in theory is 1. From Fig. 3, it is obvious to find that the approximation errors decrease with order 1, which agrees with the theoretical result.




Figure 3: strong convergence rate of the tamed Euler method (smoother noise).

## References

- [1] D. C. Antonopoulou, G. Karali, and A. Millet. Existence and regularity of solution for a stochastic Cahn–Hilliard/Allen–Cahn equation with unbounded noise diffusion. *Journal of Differential Equations*, 260(3):2383–2417, 2016.
- [2] M. Beccari, M. Hutzenthaler, A. Jentzen, R. Kurniawan, F. Lindner, and D. Salimova. Strong and weak divergence of exponential and linear-implicit Euler approximations for stochastic partial differential equations with superlinearly growing nonlinearities. *arXiv preprint arXiv:1903.06066*, 2019.
- [3] S. Becker, B. Gess, A. Jentzen, and P. E. Kloeden. Strong convergence rates for explicit space-time discrete numerical approximations of stochastic Allen–Cahn equations. *arXiv preprint arXiv:1711.02423*, 2017.

- [4] C-E. Bréhier. Approximation of the invariant distribution for a class of ergodic SPDEs using an explicit tamed exponential Euler scheme. *ESAIM: Mathematical Modelling and Numerical Analysis*, 56(1):151–175, 2022.
- [5] J. W. Cahn. On spinodal decomposition. *Acta Metallurgica*, 9(9):795–801, 1961.
- [6] J. W. Cahn and J. E. Hilliard. Free energy of a nonuniform system. I. Interfacial free energy. *The Journal of Chemical Physics*, 28(2):258–267, 1958.
- [7] J. W. Cahn and J. E. Hilliard. Spinodal decomposition: A reprise. *Acta Metallurgica*, 19(2):151–161, 1971.
- [8] M. Cai, S. Gan, and Y. Hu. Weak error estimate of a fully discrete method for stochastic Cahn–Hilliard equation with additive noise. *arXiv preprint arXiv:2111.08198*, 2021.
- [9] M. Cai, S. Gan, and X. Wang. Weak convergence rates for an explicit full-discretization of stochastic Allen–Cahn equation with additive noise. *Journal of Scientific Computing*, 86(3), 2021.
- [10] S. Chai, Y. Cao, Y. Zou, and W. Zhao. Conforming finite element methods for the stochastic Cahn–Hilliard–Cook equation. *Applied Numerical Mathematics*, 124:44–56, 2018.
- [11] J. Cui and J. Hong. Wellposedness and regularity estimate for stochastic Cahn–Hilliard equation with unbounded noise diffusion. *arXiv preprint arXiv:1907.02249*, 2019.
- [12] J. Cui and J. Hong. Absolute continuity and numerical approximation of stochastic Cahn–Hilliard equation with unbounded noise diffusion. *Journal of Differential Equations*, 269(11):10143–10180, 2020.
- [13] J. Cui, J. Hong, and L. Sun. Strong convergence of full discretization for stochastic Cahn–Hilliard equation driven by additive noise. *SIAM Journal on Numerical Analysis*, 59(6):2866–2899, 2021.
- [14] G. Da Prato and A. Debussche. Stochastic Cahn–Hilliard equation. *Nonlinear Analysis: Theory, Methods & Applications*, 26(2):241–263, 1996.
- [15] G. Da Prato and J. Zabczyk. *Stochastic Equations in Infinite Dimensions*. Cambridge University Press, 2014.
- [16] N. Elezović and A. Mikelić. On the stochastic Cahn–Hilliard equation. *Nonlinear Analysis: Theory, Methods & Applications*, 16(12):1169–1200, 1991.
- [17] X. Feng, Y. Li, and Y. Zhang. A fully discrete mixed finite element method for the stochastic Cahn–Hilliard equation with gradient-type multiplicative noise. *Journal of Scientific Computing*, 83(1), 2020.

- [18] D. Furihata, M. Kovács, S. Larsson, and F. Lindgren. Strong convergence of a fully discrete finite element approximation of the stochastic Cahn–Hilliard equation. *SIAM Journal on Numerical Analysis*, 56(2):708–731, 2018.
- [19] I. Gyöngy, S. Sabanis, and D. Šiška. Convergence of tamed Euler schemes for a class of stochastic evolution equations. *Stochastics and Partial Differential Equations: Analysis and Computations*, 4(2):225–245, 2016.
- [20] J. Hong, D. Jin, and D. Sheng. Convergence analysis of a finite difference method for stochastic Cahn–Hilliard equation. *arXiv preprint arXiv:2202.09055*, 2022.
- [21] J. Hong, D. Jin, and D. Sheng. Finite difference method for stochastic Cahn–Hilliard equation: Strong convergence rate and density convergence. *arXiv preprint arXiv:2203.00571*, 2022.
- [22] M. Hutzenthaler and A. Jentzen. On a perturbation theory and on strong convergence rates for stochastic ordinary and partial differential equations with nonglobally monotone coefficients. *The Annals of Probability*, 48(1):53–93, 2020.
- [23] G. T. Kossioris and G. E. Zouraris. Finite element approximations for a linear Cahn–Hilliard–Cook equation driven by the space derivative of a space-time white noise. *Discrete and Continuous Dynamical Systems - Series B*, 18(7):1845–1872, 2013.
- [24] M. Kovács, S. Larsson, and A. Mesforush. Finite element approximation of the Cahn–Hilliard–Cook equation. *SIAM Journal on Numerical Analysis*, 49(6):2407–2429, 2011.
- [25] S. Larsson and A. Mesforush. Finite-element approximation of the linearized Cahn–Hilliard–Cook equation. *IMA Journal of Numerical Analysis*, 31(4):1315–1333, 2011.
- [26] R. Qi, M. Cai, and X. Wang. Strong convergence rates of a fully discrete scheme for the Cahn–Hilliard–Cook equation. *arXiv preprint arXiv:2204.01630*, 2022.
- [27] R. Qi and X. Wang. Error estimates of semidiscrete and fully discrete finite element methods for the Cahn–Hilliard–Cook equation. *SIAM Journal on Numerical Analysis*, 58(3):1613–1653, 2020.
- [28] X. Wang. An efficient explicit full-discrete scheme for strong approximation of stochastic Allen–Cahn equation. *Stochastic Processes and their Applications*, 130(10):6271–6299, 2020.