



Mesenchymal stem cells as carrier cells to enable effective intratumoral delivery of oncolytic virus for oncolytic virotherapy: a systematic review

Chottiwatt Jittprasong^{1,*}

¹Department of Biomedical Engineering, College of Engineering, City University of Hong Kong

*Corresponding author: Department of Biomedical Engineering, College of Engineering, City University of Hong Kong, Kowloon, Hong Kong. chottiwatt.j@my.cityu.edu.hk

Abstract

Oncolytic viruses, which may be naturally occurring or genetically engineered, are a type of virus that infects and destroy cancer cells preferentially. Owing to their selectivity, they outperform conventional chemotherapy and radiotherapy, which both have a tendency to impact non-target cells and cause unwanted adverse side effects. Oncolytic virotherapy is a type of cancer treatment in which oncolytic viruses are deliberately introduced into patients affected with cancers in order for them to infect and destroy cancer cells locally or systemically, in a manner analogous to chemotherapy but with a greater degree of selectivity. Multiple studies indicate that oncolytic virotherapy is effective in vitro but in vivo findings remain ambiguous due to the approach's primary limitation: inefficient therapeutic agent delivery to its target, which is heavily influenced by the immune system. Here, we propose overcoming this limitation by exploiting a recent discovery in cancer research: a carrier cell. By exploiting their tumor-promoting activities, mesenchymal stem cells may be employed for cancer therapy by serving as a carrier for the oncolytic viruses toward their target. This approach directly addresses the limitation of conventional oncolytic virotherapy, where oncolytic viruses are often poorly delivered after systemic administration.

Keywords: oncolytic virotherapy, oncolytic virus, mesenchymal stem cell

Introduction

Cancer is a group of diseases that affect a large number of humans and animals. Such diseases are characterized by abnormal cell growth, invasion, metastasis, and organ dysfunction and may ultimately result in death. Humans have over 100 different forms of cancer, which are classified both clinically and histologically (National Cancer Institute 2007). While some cancers respond to conventional therapy or exhibit transient remissions, others have been categorized as aggressive and difficult, if not impossible, to cure with traditional therapies, such as glioblastoma multiforme (GBM), the most aggressive form of brain cancer (Hanif *et al.* 2017). In 2015, cancer affected over 90 million people globally and claimed over 9 million lives (GBD 2015 Disease and Injury Incidence and Prevalence Collaborators 2016; GBD 2015 Mortality and Causes of Death Collaborators 2016). Efforts to cure cancer have been ongoing for decades. Several treatment protocols have been established and proven effective for the treatment of many types of cancer.

Chemotherapy and radiotherapy are two of the most frequently used therapies for a wide variety of tumors. They are undeniably successful therapies that have cured millions of people. Additionally, chemotherapy and radiotherapy have become substantially more efficient as a result of continual research. Despite these accomplishments, both therapies depend on the use of poorly selective, highly toxic substances that are administered locally or systemically and eventually have deleterious effects

on adjacent normal cells. This invariably leads to dose-limiting toxicity, which discourages patients from adopting the therapy. Additionally, both therapies are linked with considerable physical and psychological side effects that may have a negative influence on patients' and families' quality of life.

Oncolytic virotherapy (OVT) is a revolutionary technique in cancer treatment that utilizes an oncolytic virus (OV) that infects and destroys cancer cells preferentially (Russell *et al.* 2012). By using nature's evolutionary process, this method leverages viruses as live therapeutic agents. In fact, OVs are quite naturally occurring in our environment and have an undeniable track record against cancer, such as senecavirus (Roberts *et al.* 2006). OVs are able to infect and replicate in cancer cells before destroying these cells (Figure 1). Genetic engineering techniques have been used to modify OVs for increased tumor selectivity, higher replication rate, and more toxicity toward cancer cells to enhance therapy effectiveness while minimizing off-target effects (Harrington *et al.* 2019). OVs' inherent genetic flexibility enables them to be efficient against a diverse variety of malignancies with varying genetic origins.

OVT provides various advantages over conventional cancer therapy, but it also has substantial drawbacks in terms of safety and efficacy. Since OVT is an intentional inoculation of potentially virulent viruses, its safety is considerable. Utilizing OVT requires close monitoring to ensure safe and effective levels are maintained at the target location. Efficacy of OVT is also affected




Figure 1 OVs work by infecting tumor cells but sparing health cells.

Note: From "Oncolytic virotherapy: the questions and the promise," by Aurelian (2013), *Oncolytic Virother*, 2, 19-29. Copyright 2013 by Dove Medical Press Limited under CC BY-NC 3.0 license. Reprinted under license terms.

Table 1 OVs under clinical trial (Aurelian 2013)

Type of virus	Clinical trial ^a	Reference
Adenovirus	Phase I*	Barton <i>et al.</i> (2008)
Vaccinia virus	Phase II*	Park <i>et al.</i> (2008)
Herpes Simplex Virus	Phase II*	Senzer <i>et al.</i> (2009)

^a completed and ongoing.

by an inability to evade the adaptive immune system, inefficient delivery, and infection of non-target cells. Therefore, OVT requires a development to produce effective mechanisms of action to overcome these limitations.

In this literature review paper, I will discuss the limitations of OVT and the bioengineering approach to overcome these limitations.

Discussion

The primary constraint of contemporary OVT is the OV's ability to reach the targeted cancer cells. Several factors contribute to inefficient delivery, including the following:

- Immune system
- Non-cancer cells infection

In case of successful and sufficient delivery of OVs, the effectiveness of OVT remains contingent on a number of parameters, including but not limited to the following (Goradel *et al.* 2021):

- Tumor penetration
- Hypoxic effect

Immune System

Administration of OVs as therapeutic agents can be done locally or systemically. In a local administration, OVs are delivered intratumorally. This type of administration is restricted to




Figure 2 (A) Intratumoral injection for hepatic cancer. (B) Systemic injection for metastasized cancers

Note: From "Novel insights into the role of *Clostridium novyi*-NT related combination bacteriolytic therapy in solid tumors (Review)," by Feng *et al.* (2021), *Oncology Letters*, 21(2), 110. Copyright 2020 by Feng *et al.* under CC BY-NC-ND 4.0 license. Reprinted under license terms.

easily accessible cancer such as melanoma. At the same time, inaccessible tumors will need systemic administration through intravenous (IV) injection.

As seen in Figure 2, when administered intravenously, the immune system may act as a double-edged sword; the therapeutic agent must avoid being neutralized by neutralizing antibodies (Nabs), which may be from a pre-existing immunity, in the blood circulation before reaching its target (Schmidt 2013). While circulating in the body, it must also avoid the subsequent immune system's natural response to infection and complement system that attempt to eliminate the therapeutic agent prematurely as well (Diaz *et al.* 2007).

Immunosuppressive medications and chemotherapy to inhibit the complement system may be administered to prolong the duration of the oncolytic infection, thus prolonging the therapeutic effect (Melcher *et al.* 2011; Lolkema *et al.* 2011; Kottke *et al.* 2009; Magge *et al.* 2013). However, the use of immunosuppressive drugs may cause toxicity (Wijidicks 2001; Miller 2002). A more viable solution would be to allow the OV to bypass the immune system or even use the immune system to aid in tumor clearance since the inflammation induced by the OV infection within the tumor contribute to immune recruitment and subversion of immune evasion by the tumor cells (W. Tong *et al.* 2012; Naik *et al.* 2011; Workenhe *et al.* 2015).

Infection of tumor cells by OVs and its immunologic cell death (ICD) induces the local and systemic antitumor immunity mediated by the adaptive immune system, resulting in subsequent cancer remission. Danger-associated molecular patterns (DAMPs) released during ICD from cancer cells, including HMGB1, ATP, ecto-CRT, and type I IFNs, attract the antigen-presenting cells (APCs), especially dendritic cells (DCs), whereby they acquire tumor antigens (Ags), matured, and prime the T cell-mediated immune response (Figure 3). Pro-inflammatory cytokines secreted by DCs from the binding of DAMPs and tumor Ags presented to tumor-specific CD8⁺ T cells mediated the antitumor immunity by secretion of effector cytokine and cytolysis (van Vloten *et al.* 2018; Serrano-del Valle *et al.* 2019).

Additionally, OVs may operate as viral vectors for gene expressions suppressing cancer's angiogenesis process, so reducing tumor development by depriving it of oxygen (Frentzen *et al.* 2009; Conner and Braidwood 2012).




Figure 5 Numerous mechanisms contribute to ECs vulnerability to OV infection: (1) increased permeability of neovessels allows for the accumulation of OVs; (2) highly active proliferation of ECs provides OVs with an enzyme required for replication; and (3) VEGF binding to VEGFR2 activates a repressor that inhibits type I interferon-mediated antiviral signaling. Note: From "Infection of non-cancer cells: A barrier or support for oncolytic virotherapy?," by Naumenko *et al.* (2022), *Molecular Therapy - Oncolytics*, 24, 663-682. Copyright 1969 by Elsevier under CC BY 4.0 license. Reprinted under license terms.

Mechanism of infection in cancer cells: Tumor EC is most frequently detected for OV infection by vaccinia virus (VV), herpes simplex virus (HSV), reovirus, vesicular stomatitis virus (VSV), and adenovirus. The infection elicited by a variety of viruses across families indicates that the mechanisms of infection are due to the properties of tumor ECs rather than viral properties. The EC's susceptibility to OV infection is mostly due to vascular endothelial growth factor (VEGF), as believed by various studies. Arulanandam *et al.* (2015) have proved that permissiveness of EC to OV infection is largely contributed by VEGF as seen in VV-treated mice with suppression of VEGF/VEGFR2 by Abs.

Cancer cells also attract leukocytes and reprogram them to generate immunosuppressive cytokines such as tumor growth factor 1 (TGF-1) and interleukin 10 (IL-10), hence promoting viral replication and favoring OVs (Ponomarev and Shubina 2019). Proliferative activity that is greater than usual and its metabolism under the influence of angiogenic factors also establish a vulnerability for OVs to infiltrate (Kottke *et al.* 2011).

Finally, tumor neovessels have a high permeability (Tseng *et al.* 2010), which allows viruses to remain in the microenvironment (Figure 5).

Mechanism of infection in non-cancer cells: Increased concentrations of several kinds of OVs in the spleen have been seen in studies, indicating possible infection. The primary target of OVs infection in non-cancerous cells is CD169+ macrophages of marginal metallophilic macrophages (MMMs) and subcapsular sinus macrophages (SSMs). At least two viruses, VSV (Naumenko *et al.* 2018; Kim *et al.* 2017) and MV (Peng *et al.* 2003), were found to infect CD169+ macrophages.




Figure 6 Properties of MMMs determine permissiveness toward OVs. Anatomical location of both MMMs and SSMs also contributed to its prevalence toward OV infection. Note: From "Infection of non-cancer cells: A barrier or support for oncolytic virotherapy?," by Naumenko *et al.* (2022), *Molecular Therapy - Oncolytics*, 24, 663-682. Copyright 1969 by Elsevier under CC BY 4.0 license. Reprinted under license terms.

SSMs and MMMs are located in the spleen's sinusoid system, where blood slows down and allows them to search for antigens. SSMs are also present in lymph nodes, where they fulfill a similar function to antigen scanning. Despite being exposed to a variety of pathogens as part of their primary function, CD169+ cells are incapable of eliminating viruses on their own (Phan *et al.* 2009). However, the mechanism by which OVs infect CD169+ cells is unknown and is actively being explored. Due to their location and incapacity to eliminate viruses on their own, they have increased viral sequestration in their immediate vicinity (Figure 6).

In vitro infection of DCs by OVs is also reported widely (Ilett *et al.* 2009, 2011; Brown *et al.* 2017; Zhang *et al.* 2013, 2014). Farrell *et al.* (2015) suggested that DCs acquire the infection from infected SSMs. Although the fate of non-cancer cells infected with OV has not been investigated, it is anticipated that non-attenuated viruses may disturb the macrophage layer. Following VV and murine cytomegalovirus (MCMV) infection, a large drop in the number of SSMs is seen, which may be due to viral lysis or IFN-mediated apoptosis.

Wide reports of major absorption of systemically administered OVs by phagocytosis are also documented (Naumenko *et al.* 2018). Monocytes are found to be uniquely susceptible to re-

ovirus, VSV, VV, and influenza A virus (Ilett *et al.* 2014; Berkeley *et al.* 2018; Stichling *et al.* 2018; MacLeod *et al.* 2015, 2013).

It is advisable to do an investigation to establish the clinical significance of these non-cancerous cell infections in order to avoid any serious adverse consequences.

Tumor Penetration

One of the most significant barriers to the entry of therapeutic agents into cancer cells in the presence of intracellular junctions of ECs (Lipinski *et al.* 1997; Lavin *et al.* 2007; Green *et al.* 2002). Metastasis via epithelial-to-mesenchymal transition (EMT) and mesenchymal-to-epithelial transition (MET) causes the junctions to become tighter, making therapy more challenging. (Christiansen and Rajasekaran 2006; Turley *et al.* 2008). Adenoviruses are a good example of this since they are often inhibited by the existence of this barrier (Yumul *et al.* 2016).

There have been a variety of ways developed to get around this issue. Oncolytic adenoviruses expressing hyaluronidase allow for enhanced cell fusion in a melanoma xenograft model as a consequence of increased lysis of the extracellular matrix (Guedan *et al.* 2010). Additionally, the extracellular matrix (ECM) may impede therapeutic agent dispersal inside solid tumors (Netti *et al.* 2000; Vargová *et al.* 2003) while also retarding invasion by OV's (Wojton and Kaur 2010). It has been shown that pretreatment of the tumor with collagenase, which breaks down the ECM layer, improves efficiency (Kuriyama *et al.* 2000).

Hypoxic Effect

Sustained hypoxia is known as a common condition developed in the immediate vicinity of cancer cells as a result of uncontrolled cell proliferation and an excessive amount of metabolic activity. The hypoxic effect has been demonstrated to have contradictory consequences on OV's (Wong *et al.* 2010). Shen *et al.* (2006); Shen and Hermiston (2005) have shown that adenovirus's ability to replicate and lysis has been significantly diminished as a result of the hypoxic effect.

Modifying the E1A gene regulated by a hypoxia-response element-containing promoter expression, Hernandez-Alcoceba *et al.* (2002) has created an oncolytic adenovirus that can replicate effectively in a hypoxic environment.

Limitations

Systemic administration of OV's is considered the most beneficial method of conducting OVT because it influences all types of cancers and disseminated cancers, including brain tumors, the majority of which cannot be treated with radiotherapy or chemotherapy due to the inability of most therapeutic agents to cross the blood-brain barrier (BBB); however, viruses, such as semliki forest virus (SFV), can cross the BBB and infect brain cells (Willems *et al.* 1979), this can be modified to make it preferentially target the abnormal brain cells while sparing normal cells as well. In vivo therapy with modified SFV of mouse neuroblastoma and GBM cell lines, as well as patient-derived human glioblastoma cell cultures (HGCC), demonstrated a promising effect, extending survival and curing half of the animals tested (Ramachandran *et al.* 2017). This demonstrated the potential of OVT to even cure the disease considered incurable, GBM.

As discussed in Discussion the current main limitations of OVT lie in circumvention of the immune system and effective delivery of systemically administered OV's.

Hypotheses

By circulating in the bloodstream as pathogens, OV's tend to be neutralized by various natural responses of the body to clear the infection, thus, failing the OVT.

By integrating OV's into circulating normal body cells without inflicting distress to them, OV's may utilize the normal body cells as a "carrier" to move stealthily to their target and infect it, thereby evading neutralization. Thus, effectively solving the problem of neutralization by the immune system during systemic administration.

Approach

To act as a carrier for the cancer cells, the cells must be somewhat needed by cancer itself. Cells associated with tumors are called tumor-associated cells. Common examples of them are tumor-associated macrophages (TAM) which inhibit anti-tumor immune response, cancer-associated fibroblast (CAF), which promote remodeling of the ECM and stimulation of angiogenesis, and myeloid-derived suppressor cells (MDSC) which suppress the immune system. These cells are commonly recruited by the cancer cells to promote their growth.

Mesenchymal stem cells (MSCs) are a kind of multipotent stem cells found throughout the body. As seen in Figure 7, MSCs play a critical role in tissue healing by undergoing cellular differentiation to orchestrate structural repair, immunological regulation, and growth factor secretion to drive vascularization and epithelialization, as well as recruitment of resident stem cells (Balaji *et al.* 2012; Hu *et al.* 2018; Sylakowski *et al.* 2020).

Tumors may be regarded as "wounds that never heal," and in response to this prolonged wound, MSCs are continuously recruited to the tumor micro-environment to aid its reparation by secreting multiple growth factors, chemokines, and cytokines that promote the proliferation of cancer cells such as angiogenesis while also suppressing the immune response.

This way, MSCs are now tumor-associated MSCs (TA-MSCs) (Shi *et al.* 2017). But many studies also suggest that MSCs might play a role in suppressing cancer (Cousin *et al.* 2009; Klopp *et al.* 2011; Rhee *et al.* 2015).




Figure 7 Graphical summary showing an involvement of the MSC in wound healing process orchestrated by numerous types of cells.

Note: From "TLR4-dependent shaping of the wound site by MSCs accelerates wound healing," by Munir *et al.* (2020), *EMBO reports*, 21(5), e48777. Copyright 2020 by John Wiley and Sons. Reprinted with permission.

Regardless of whether MSCs promote or repress tumor development, we can take advantage of the fact that MSCs are recruited directly to the site of cancer proliferation. By loading MSCs with OV, we can migrate OV directly to the site of cancer. Additionally, MSCs have been shown to protect virus that has been pre-loaded against immune-mediated neutralization (Sonabend *et al.* 2008; Mader *et al.* 2009; Komarova *et al.* 2006). Therefore, OVs are efficiently hidden from the immune system by employing MSCs as a carrier. This way, MSCs are effectively acting as "trojan horses," and we may deliver OVs systemically without concern of the immune system neutralizing them before they reach their target.

By combining OVs with MSCs, OVs may directly target the tumor, circumventing almost all of the constraints associated with contemporary OVT.

Methods

Oncolytic virus

OVs of several types may be used. ICOVIR-5, on the other hand, would be the finest fit. ICOVIR-5 is an oncolytic adenovirus whose replication is regulated by the E2F-responsive promoter. It incorporates the Arg-Gly-Asp (RGD) motif to increase infectivity and deletion of the retinoblastoma protein (pRB)-binding site to prevent it from replicating via the activated RB pathway, a pathway that can be used to differentiate normal cells from cancer cells (Ruano *et al.* 2020; Cascallo *et al.* 2007).

The quantity of ICOVIR-5 replicated inside *ex vivo* proliferating MSCs is found to be 3-log lower than the number of cancer cells (Shi *et al.* 2017).

Carrier mesenchymal stem cell

The protocol is based on a work established by García-Castro *et al.* (2010). Autologous MSCs can be collected from the patient through bone marrow aspirate (BMA). Isolation of mononuclear cells can be done by density gradient centrifugation (400 g, 25 min, 20 °C). After centrifugation, the substance is washed twice with phosphate buffer by sedimentation at room temperature for 5 minutes (300 g), resuspended in MSC medium of Dulbecco's Modified Eagle Medium (DMEM), 10% Fetal Calf Serum (FCS), 2mM glutamine, 10 U/L penicillin, and 10 µg ml⁻¹ streptomycin added to the culture at 1.5 × 10⁵ cells per cm² and left to adhere for 24 hours.

Non-adherent hematopoietic stem cells can be removed by washing twice with phosphate-buffered saline (PBS) at 10ml. MSC can then be cultured at 37 °C at 5% CO₂ in an MSC medium, passage after 85% confluence. 0.5% of trypsin and 0.2% ethylenediaminetetraacetic acid (EDTA) can then be used to treat the cells for 5 minutes, wash, sediment at room temperature for 10 minutes (300 g), and plate in MSC medium at 5 × 10³ cells per cm². In the third passage, homogeneous MSC culture should be obtained.

Specific differentiation of hMSC can be initiated with commercially available hMSC differentiation kit.

OV (ICOVIR-5) can be added to the trypsinized, washed, and resuspended homogenous MSC culture in DMEM and incubated for 2 hours at 37 °C. The cells should then be washed and resuspended in DMEM 10% FBS. EGFP-positive cells can be determined with flow cytometry to verify the transduction efficiency. After the infection, the cells will need to undergo a replication cycle to allow a high load of MSCs and OVs at around 80%-100% infection. Once validated, the cells should

be washed twice with PBS and incubated with fresh medium. The harvesting and cell extracting process of OV carriers can be prepared by freeze-thawing.

Expected outcomes

According to all theories discussed in the part of Discussion, this method should be able to facilitate systemic administration of OVs by using the MSCs as a carrier to protect them from the immune system and directly transfer OVs to the cancer site where OVs are unloaded and directly infect cancer cells and cause oncolysis. If the infection is sustained long enough, continuous distress of cancer cells mediated by oncolysis will trigger the secretion of DAMPs recruiting the adaptive immune system to the site, thus inducing an anti-tumor response by the immune system.

The mechanism of infection is similar to that of a local administration of OVs but uses MSCs as a carrier. This also eliminates the risk of deleterious effects associated with the systemic delivery of OVs (Figure 8).

OVT seems likely to succeed current cancer therapies such as radiation and chemotherapy by nearly eliminating all side effects associated with radiotherapy and chemotherapy while also utilizing all of their advantages, thus paving the way for the next stage in cancer treatment.




Figure 8 MSCs used as carriers for systemically administered OVs. **(A)** The immune system directly attacks OVs, impeding them from reaching their targets. **(B)** MSCs shield OVs from the immune system and migrate them directly to the tumor, triggering direct oncolysis and an anti-tumor reaction mediated by the immune system. Note: From "Patient-derived mesenchymal stem cells as delivery vehicles for oncolytic virotherapy: novel state-of-the-art technology," by M. *et al.* (2015), *Oncolytic Virother*, 4, 149-155. Copyright 2015 by Dove Medical Press Limited under CC BY-NC license. Reprinted under license terms.

Implications

OVT will be one of the next therapy procedures that are both successful and safe while also being capable of curing all sorts

of tumors, including the once incurable GBM. The National Institutes of Health has already begun phase one testing of the treatment, which has been effectively used to treat a variety of other malignant tumors, including liver and breast cancer. Clearly, this is a significant triumph for the millions of individuals who are now battling cancer, as well as for the loved ones of cancer patients worldwide.

Cancer has undoubtedly had a profound effect on society. Due to this exceptional tumor therapy and the tireless efforts of medical experts and scientists, cancer treatment has taken a quantum leap forward. Due to the simplicity and superiority of the technique compared to modern cancer therapies such as immunotherapy, chemotherapy, and radiation, it will be economical yet effective for a large number of patients battling cancer. This is particularly remarkable as it represents a significant breakthrough in the treatment of late-stage cancers for which conventional cancer treatments have been ineffective.

I anticipate widespread adoption of OVT in medical oncology, which has enormous potential for the treatment of cancers in humans. I strongly believe that in the next five to ten years, OVT will be routinely used for central nervous system cancer, esophageal and lung tumors.

Data availability

Data sharing not applicable to this report as no datasets were generated or analysed during the current study.

Funding

This article received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Conflicts of interest

The author reports no conflict of interest.

Literature cited

Alvarez-Breckenridge CA, Yu J Fau Price R, Price R Fau Wojton J, Wojton J Fau Pradarelli J, Pradarelli J Fau Mao H, Mao H Fau Wei M, Wei M Fau Wang Y, Wang Y Fau He S, He S Fau Hardcastle J, Hardcastle J Fau Fernandez SA *et al.* 2012. Nk cells impede glioblastoma virotherapy through nkp30 and nkp46 natural cytotoxicity receptors. .

Arulanandam R, Batenchuk C, Angarita FA, Ottolino-Perry K, Cousineau S, Mottashed A, Burgess E, Falls TJ, Silva ND, Tsang J *et al.* 2015. Vegf-mediated induction of prd1-bf1/blimp1 expression sensitizes tumor vasculature to oncolytic virus infection. *Cancer Cell.* 28:210–224.

Aurelian L. 2013. Oncolytic virotherapy: the questions and the promise. .

Balaji S, Keswani SG, Crombleholme TM. 2012. The role of mesenchymal stem cells in the regenerative wound healing phenotype. (New Rochelle).

Barton KN, Stricker H Fau Brown SL, Brown SI Fau Elshaikh M, Elshaikh M Fau Aref I, Aref I Fau Lu M, Lu M Fau Pegg J, Pegg J Fau Zhang Y, Zhang Y Fau Karvelis KC, Karvelis Kc Fau Siddiqui F, Siddiqui F Fau Kim JH *et al.* 2008. Phase i study of noninvasive imaging of adenovirus-mediated gene expression in the human prostate. .

Berkeley RA, Steele LP, Mulder AA, van den Wollenberg DJ, Kottke TJ, Thompson J, Coffey M, Hoeben RC, Vile RG, Melcher A *et al.* 2018. Antibody-Neutralized Reovirus Is Effective in

Oncolytic Virotherapy. *Cancer Immunology Research.* 6:1161–1173.

Bofill-De Ros X, Rovira-Rigau M, Fillat C. 2017. Implications of micrnas in oncolytic virotherapy. *Frontiers in Oncology.* 7.

Brown MC, Holl EK, Boczkowski D, Dobrikova E, Mosaheb M, Chandramohan V, Bigner DD, Gromeier M, Nair SK. 2017. Cancer immunotherapy with recombinant poliovirus induces ifn-dominant activation of dendritic cells and tumor antigen-specific ctls. *Science Translational Medicine.* 9:eaan4220.

Cascallo M, Alonso MM, Rojas JJ, Perez-Gimenez A, Fueyo J, Alemany R. 2007. Systemic toxicity–efficacy profile of icovir-5, a potent and selective oncolytic adenovirus based on the prb pathway. *Molecular Therapy.* 15:1607–1615.

Christiansen JJ, Rajasekaran AK. 2006. Reassessing Epithelial to Mesenchymal Transition as a Prerequisite for Carcinoma Invasion and Metastasis. *Cancer Research.* 66:8319–8326.

Conner J, Braidwood L. 2012. Expression of inhibitor of growth 4 by hsv1716 improves oncolytic potency and enhances efficacy. *Cancer Gene Therapy.* 19:499–507.

Cook M, Chauhan A. 2020. Clinical application of oncolytic viruses: A systematic review. *International Journal of Molecular Sciences.* 21:7505.

Cousin B, Ravet E, Poglio S, De Toni F, Bertuzzi M, Lulka H, Touil I, André M, Grolleau JL, Péron JM *et al.* 2009. Adult stromal cells derived from human adipose tissue provoke pancreatic cancer cell death both in vitro and in vivo. *PLOS ONE.* 4:1–10.

Diaz RM, Galivo F, Kottke T, Wongthida P, Qiao J, Thompson J, Valdes M, Barber G, Vile RG. 2007. Oncolytic Immunovirotherapy for Melanoma Using Vesicular Stomatitis Virus. *Cancer Research.* 67:2840–2848.

Farrell HE, Davis-Poynter N, Bruce K, Lawler C, Dolken L, Mach M, Stevenson PG, Frueh K. 2015. Lymph node macrophages restrict murine cytomegalovirus dissemination. *Journal of Virology.* 89:7147–7158.

Feng X, He P, Zeng C, Li YH, Das SK, Li B, Yang HF, Du Y. 2021. Novel insights into the role of clostridium novyi-nt related combination bacteriolytic therapy in solid tumors (review). *Oncol Lett.* 21:110.

Frentzen A, Yu Ya Fau Chen N, Chen N Fau Zhang Q, Zhang Q Fau Weibel S, Weibel S Fau Raab V, Raab V Fau Szalay AA, Szalay AA. 2009. Anti-vegf single-chain antibody glaf-1 encoded by oncolytic vaccinia virus significantly enhances antitumor therapy. .

García-Castro J, Alemany R Fau Cascalló M, Cascalló M Fau Martínez-Quintanilla J, Martínez-Quintanilla J Fau Arriero MdM, Arriero Mdel M Fau Lassaletta A, Lassaletta A Fau Madero L, Madero L Fau Ramírez M, Ramírez M. 2010. Treatment of metastatic neuroblastoma with systemic oncolytic virotherapy delivered by autologous mesenchymal stem cells: an exploratory study. .

GBD 2015 Disease and Injury Incidence and Prevalence Collaborators. 2016. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the global burden of disease study 2015. *Lancet (London, England).* 388:1545–1602. 27733282[pmid] PMC5055577[pmcid] S0140-6736(16)31678-6[PII].

GBD 2015 Mortality and Causes of Death Collaborators. 2016. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the global

- burden of disease study 2015. *Lancet* (London, England). 388:1459–1544. 27733281[pmid]PMC5388903[pmcid]S0140-6736(16)31012-1[PII].
- Goradel NH, Baker AT, Arashkia A, Ebrahimi N, Ghorghanlu S, Negahdari B. 2021. Oncolytic virotherapy: Challenges and solutions. *Current Problems in Cancer*. 45:100639.
- Green SK, Karlsson MCI, Ravetch JV, Kerbel RS. 2002. Disruption of Cell-Cell Adhesion Enhances Antibody-dependent Cellular Cytotoxicity: Implications for Antibody-based Therapeutics of Cancer. *Cancer Research*. 62:6891–6900.
- Guedan S, Rojas Jj Fau Gros A, Gros A Fau Mercade E, Mercade E Fau Cascallo M, Cascallo M Fau Alemany R, Alemany R. 2010. Hyaluronidase expression by an oncolytic adenovirus enhances its intratumoral spread and suppresses tumor growth. .
- Hanif F, Muzaffar K, Perveen K, Malhi SM, Simjee SU. 2017. Glioblastoma multiforme: A review of its epidemiology and pathogenesis through clinical presentation and treatment. *Asian Pacific journal of cancer prevention : APJCP*. 18:3–9. 28239999[pmid] PMC5563115[pmcid].
- Harrington K, Freeman DJ, Kelly B, Harper J, Soria JC. 2019. Optimizing oncolytic virotherapy in cancer treatment. *Nature Reviews Drug Discovery*. 18:689–706.
- Hernandez-Alcoceba R, Pihajla M, Qian D, Clarke MF. 2002. New oncolytic adenoviruses with hypoxia- and estrogen receptor-regulated replication. *Human Gene Therapy*. 13:1737–1750. PMID: 12396626.
- Hu MAOX, Borrelli MR, Lorenz HP, Longaker MT, Wan DAO. 2018. Mesenchymal stromal cells and cutaneous wound healing: A comprehensive review of the background, role, and therapeutic potential. .
- Ilett E, Kottke T, Donnelly O, Thompson J, Willmon C, Diaz R, Zaidi S, Coffey M, Selby P, Harrington K *et al.* 2014. Cytokine conditioning enhances systemic delivery and therapy of an oncolytic virus. *Molecular Therapy*. 22:1851–1863.
- Ilett EJ, Bárcena M, Errington-Mais F, Griffin S, Harrington KJ, Pandha HS, Coffey M, Selby PJ, Limpens RW, Mommaas M *et al.* 2011. Internalization of Oncolytic Reovirus by Human Dendritic Cell Carriers Protects the Virus from Neutralization. *Clinical Cancer Research*. 17:2767–2776.
- Ilett EJ, Prestwich RJ, Kottke T, Errington F, Thompson JM, Harrington KJ, Pandha HS, Coffey M, Selby PJ, Vile RG *et al.* 2009. Dendritic cells and t cells deliver oncolytic reovirus for tumour killing despite pre-existing anti-viral immunity. *Gene Therapy*. 16:689–699.
- Kim DS, Dastidar H, Zhang C, Zemp FJ, Lau K, Ernst M, Rakic A, Sikdar S, Rajwani J, Naumenko V *et al.* 2017. Smac mimetics and oncolytic viruses synergize in driving anticancer t-cell responses through complementary mechanisms. *Nature Communications*. 8:344.
- Klopp AH, Gupta A, Spaeth E, Andreeff M, Marini, Frank I. 2011. Concise Review: Dissecting a Discrepancy in the Literature: Do Mesenchymal Stem Cells Support or Suppress Tumor Growth? *Stem Cells*. 29:11–19.
- Komarova S, Kawakami Y, Stoff-Khalili MA, Curiel DT, Pereboeva L. 2006. Mesenchymal progenitor cells as cellular vehicles for delivery of oncolytic adenoviruses. *Molecular Cancer Therapeutics*. 5:755–766.
- Kottke T, Chester J, Ilett E, Thompson J, Diaz R, Coffey M, Selby P, Nuovo G, Pulido J, Mukhopadhyay D *et al.* 2011. Precise scheduling of chemotherapy primes vegf-producing tumors for successful systemic oncolytic virotherapy. *Molecular Therapy*. 19:1802–1812.
- Kottke T, Thompson J, Diaz RM, Pulido J, Willmon C, Coffey M, Selby P, Melcher A, Harrington K, Vile RG. 2009. Improved systemic delivery of oncolytic reovirus to established tumors using preconditioning with cyclophosphamide-mediated treg modulation and interleukin-2. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 15:561–569. 19147761[pmid] PMC3046733[pmcid] 15/2/561[PII].
- Kuriyama N, Kuriyama H, Julin CM, Lamborn K, Israel MA. 2000. Pretreatment with protease is a useful experimental strategy for enhancing adenovirus-mediated cancer gene therapy. *Human Gene Therapy*. 11:2219–2230. PMID: 11084679.
- Lavin SR, McWhorter TJ, Karasov WH. 2007. Mechanistic bases for differences in passive absorption. *Journal of Experimental Biology*. 210:2754–2764.
- Lipinski CA, Lombardo F, Dominy BW, Feeney PJ. 1997. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Advanced Drug Delivery Reviews*. 23:3–25. In Vitro Models for Selection of Development Candidates.
- Lolkema MP, Arkenau HT, Harrington K, Roxburgh P, Morrison R, Roulstone V, Twigger K, Coffey M, Mettinger K, Gill G *et al.* 2011. A Phase I Study of the Combination of Intravenous Reovirus Type 3 Dearing and Gemcitabine in Patients with Advanced Cancer. *Clinical Cancer Research*. 17:581–588.
- M. R, J. GC, J. MG, Á GM, L. FL. 2015. Patient-derived mesenchymal stem cells as delivery vehicles for oncolytic virotherapy: novel state-of-the-art technology. .
- MacLeod DT, Nakatsuji T, Wang Z, di Nardo A, Gallo RL. 2015. Vaccinia virus binds to the scavenger receptor marco on the surface of keratinocytes. *Journal of Investigative Dermatology*. 135:142–150.
- MacLeod DT, Nakatsuji T, Yamasaki K, Kobzik L, Gallo RL. 2013. Hsv-1 exploits the innate immune scavenger receptor marco to enhance epithelial adsorption and infection. *Nature Communications*. 4:1963.
- Mader EK, Maeyama Y, Lin Y, Butler GW, Russell HM, Galanis E, Russell SJ, Dietz AB, Peng KW. 2009. Mesenchymal Stem Cell Carriers Protect Oncolytic Measles Viruses from Antibody Neutralization in an Orthotopic Ovarian Cancer Therapy Model. *Clinical Cancer Research*. 15:7246–7255.
- Magge D, Guo ZS, O'Malley ME, Francis L, Ravindranathan R, Bartlett DL. 2013. Inhibitors of c5 complement enhance vaccinia virus oncolysis. *Cancer gene therapy*. 20:342–350. 23661042[pmid] PMC4060830[pmcid] cgt201326[PII].
- Matzinger P. 2002. The danger model: a renewed sense of self. .
- Melcher A, Parato K, Rooney CM, Bell JC. 2011. Thunder and lightning: immunotherapy and oncolytic viruses collide. *Molecular therapy : the journal of the American Society of Gene Therapy*. 19:1008–1016. 21505424[pmid] PMC3129809[pmcid] S1525-0016(16)31904-9[PII].
- Miller LW. 2002. Cardiovascular toxicities of immunosuppressive agents. *American Journal of Transplantation*. 2:807–818.
- Munir S, Basu A, Maity P, Krug L, Haas P, Jiang D, Strauss G, Wlaschek M, Geiger H, Singh K *et al.* 2020. Tlr4-dependent shaping of the wound site by mscs accelerates wound healing. *EMBO reports*. 21:e48777.
- Naik JD, Twelves CJ, Selby PJ, Vile RG, Chester JD. 2011. Immune recruitment and therapeutic synergy: keys to optimizing oncolytic viral therapy? *Clinical cancer research : an official journal of the American Association for Cancer Re-*

- search. 17:4214–4224. 21576084[pmid] PMC3131422[pmcid] 1078-0432.CCR-10-2848[PII].
- National Cancer Institute. 2007. Cancer types.
- Naumenko V, Van S, Dastidar H, Kim DS, Kim SJ, Zeng Z, Deniset J, Lau A, Zhang C, Macia N *et al.* 2018. Visualizing oncolytic virus-host interactions in live mice using intravital microscopy. *Molecular Therapy - Oncolytics*. 10:14–27.
- Naumenko VA, Stepanenko AA, Lipatova AV, Vishnevskiy DA, Chekhonin VP. 2022. Infection of non-cancer cells: A barrier or support for oncolytic virotherapy? *Molecular Therapy - Oncolytics*. 24:663–682.
- Netti PA, Berk DA, Swartz MA, Grodzinsky AJ, Jain RK. 2000. Role of Extracellular Matrix Assembly in Interstitial Transport in Solid Tumors. *Cancer Research*. 60:2497–2503.
- Park BH, Hwang T, Fau Liu TC, Liu Tc Fau Sze DY, Sze Dy Fau Kim JS, Kim Js Fau Kwon HC, Kwon Hc Fau Oh SY, Oh Sy Fau Han SY, Han Sy Fau Yoon JH, Yoon Jh Fau Hong SH, Hong Sh Fau Moon A *et al.* 2008. Use of a targeted oncolytic poxvirus, jx-594, in patients with refractory primary or metastatic liver cancer: a phase i trial. .
- Peng KW, Frenze M, Myers R, Soeffker D, Harvey M, Greiner S, Galanis E, Cattaneo R, Federspiel MJ, Russell SJ. 2003. Biodistribution of oncolytic measles virus after intraperitoneal administration into ifnarTM-cd46ge transgenic mice. *Human Gene Therapy*. 14:1565–1577. PMID: 14577918.
- Phan TG, Green JA, Gray EE, Xu Y, Cyster JG. 2009. Immune complex relay by subcapsular sinus macrophages and noncognate b cells drives antibody affinity maturation. *Nature Immunology*. 10:786–793.
- Ponomarev AV, Shubina IZ. 2019. Insights into mechanisms of tumor and immune system interaction: Association with wound healing. *Frontiers in Oncology*. 9.
- Ramachandran M, Yu D, Dyczynski M, Baskaran S, Zhang L, Lulla A, Lulla V, Saul S, Nelander S, Dimberg A *et al.* 2017. Safe and Effective Treatment of Experimental Neuroblastoma and Glioblastoma Using Systemically Delivered Triple MicroRNA-Detargeted Oncolytic Semliki Forest Virus. *Clinical Cancer Research*. 23:1519–1530.
- Rhee KJ, Lee JI, Eom YW. 2015. Mesenchymal stem cell-mediated effects of tumor support or suppression. *International Journal of Molecular Sciences*. 16:30015–30033.
- Roberts MS, Lorence RM, Groene WS, Bamat MK. 2006. Naturally oncolytic viruses. *Current Opinion in Molecular Therapeutics*. 8:314–321.
- Ruano D, López-Martín JA, Moreno L, Álvaro Lassaletta, Bautista F, Andión M, Hernández C, África González-Murillo, Melen G, Alemany R *et al.* 2020. First-in-human, first-in-child trial of autologous mscs carrying the oncolytic virus icovir-5 in patients with advanced tumors. *Molecular Therapy*. 28:1033–1042.
- Russell SJ, Peng KW, Bell JC. 2012. Oncolytic virotherapy. *Nature Biotechnology*. 30:658–670.
- Schmidt C. 2013. Awaiting a Moment of Truth for Oncolytic Viruses. *JNCI: Journal of the National Cancer Institute*. 105:675–676.
- Senzer NN, Kaufman HI Fau Amatruda T, Amatruda T Fau Nemunaitis M, Nemunaitis M Fau Reid T, Reid T Fau Daniels G, Daniels G Fau Gonzalez R, Gonzalez R Fau Glaspy J, Glaspy J Fau Whitman E, Whitman E Fau Harrington K, Harrington K Fau Goldsweig H *et al.* 2009. Phase ii clinical trial of a granulocyte-macrophage colony-stimulating factor-encoding, second-generation oncolytic herpesvirus in patients with unresectable metastatic melanoma. .
- Serrano-del Valle A, Anel A, Naval J, Marzo I. 2019. Immunogenic cell death and immunotherapy of multiple myeloma. *Frontiers in Cell and Developmental Biology*. 7.
- Shen BH, Bauzon M, Hermiston TW. 2006. The effect of hypoxia on the uptake, replication and lytic potential of group b adenovirus type 3 (ad3) and type 11p (ad11p). *Gene Therapy*. 13:986–990.
- Shen BH, Hermiston TW. 2005. Effect of hypoxia on ad5 infection, transgene expression and replication. *Gene Therapy*. 12:902–910.
- Shi Y, Du L, Lin L, Wang Y. 2017. Tumour-associated mesenchymal stem/stromal cells: emerging therapeutic targets. *Nature Reviews Drug Discovery*. 16:35–52.
- Sonabend AM, Ulasov IV, Tyler MA, Rivera AA, Mathis JM, Lesniak MS. 2008. Mesenchymal Stem Cells Effectively Deliver an Oncolytic Adenovirus to Intracranial Glioma. *Stem Cells*. 26:831–841.
- Stichling N, Suomalainen M, Flatt JW, Schmid M, Pacesa M, Hemmi S, Jungraithmayr W, Maler MD, Freudenberg MA, Plückthun A *et al.* 2018. Lung macrophage scavenger receptor sr-a6 (marco) is an adenovirus type-specific virus entry receptor. *PLOS Pathogens*. 14:1–33.
- Sylakowski K, Bradshaw A, Wells A. 2020. Mesenchymal stem cell/multipotent stromal cell augmentation of wound healing: Lessons from the physiology of matrix and hypoxia support. *The American Journal of Pathology*. 190:1370–1381.
- Tseng JC, Granot T, DiGiacomo V, Levin B, Meruelo D. 2010. Enhanced specific delivery and targeting of oncolytic sindbis viral vectors by modulating vascular leakiness in tumor. *Cancer Gene Therapy*. 17:244–255.
- Turley EA, Veiseh M, Radisky DC, Bissell MJ. 2008. Mechanisms of disease: epithelial–mesenchymal transition—does cellular plasticity fuel neoplastic progression? *Nature Clinical Practice Oncology*. 5:280–290.
- van Vloten JP, Workenhe ST, Wootton SK, Mossman KL, Bridle BW. 2018. Critical interactions between immunogenic cancer cell death, oncolytic viruses, and the immune system define the rational design of combination immunotherapies. *The Journal of Immunology*. 200:450–458.
- Vargová L, Homola A, Zámečník J, Tichý M, Beneš V, Syková E. 2003. Diffusion parameters of the extracellular space in human gliomas. *Glia*. 42:77–88.
- W. Tong A, Senzer N, Cerullo V, Smyth Templeton N, Hemminki A, Nemunaitis J. 2012. Oncolytic viruses for induction of anti-tumor immunity. *Current Pharmaceutical Biotechnology*. 13:1750–1760.
- Wherry EJ, Ahmed R. 2004. Memory cd8 t-cell differentiation during viral infection. .
- Wijdicks EF. 2001. Neurotoxicity of immunosuppressive drugs. *Liver Transplantation*. 7:937–942.
- Willems WR, Kaluza G, Boschek CB, Bauer H, Hager H, Schütz HJ, Feistner H. 1979. Semliki forest virus: Cause of a fatal case of human encephalitis. *Science*. 203:1127–1129.
- Wojton J, Kaur B. 2010. Impact of tumor microenvironment on oncolytic viral therapy. *Cytokine & Growth Factor Reviews*. 21:127–134. Recent Advances in the Development of Oncolytic Viruses as Cancer Therapeutics.
- Wong HH, Lemoine NR, Wang Y. 2010. Oncolytic viruses for cancer therapy: Overcoming the obstacles. *Viruses*. 2:78–106.
- Workenhe ST, Verschoor ML, Mossman KL. 2015. The role of oncolytic virus immunotherapies to subvert cancer immune

- evasion. *Future Oncology*. 11:675–689. PMID: 25686121.
- Yumul R, Richter M, Lu ZZ, Saydaminova K, Wang H, Wang CHK, Carter D, Lieber A. 2016. Epithelial junction opener improves oncolytic adenovirus therapy in mouse tumor models. *Human Gene Therapy*. 27:325–337. PMID: 26993072.
- Zhang J, Tai LH, Ilkow CS, Alkayyal AA, Ananth AA, de Souza CT, Wang J, Sahi S, Ly L, Lefebvre C *et al.* 2014. Maraba mg1 virus enhances natural killer cell function via conventional dendritic cells to reduce postoperative metastatic disease. *Molecular Therapy*. 22:1320–1332.
- Zhang L, Bridle BW, Chen L, Pol J, Spaner D, Boudreau JE, Rosen A, Bassett JD, Lichty BD, Bramson JL *et al.* 2013. Delivery of viral-vectored vaccines by B cells represents a novel strategy to accelerate CD8+ T-cell recall responses. *Blood*. 121:2432–2439.