

# NON-SELF SIMILAR BLOWUP SOLUTIONS TO THE HIGHER DIMENSIONAL YANG MILLS HEAT FLOWS

A. Bensouilah<sup>(1),(2)</sup>, G. K. Duong<sup>(3),(4)</sup>, and T. E. Ghou<sup>(1)</sup>

<sup>(1)</sup> NYUAD Research Institute, New York University Abu Dhabi, PO Box 129188, Abu Dhabi, UAE

<sup>(2)</sup> School of Mathematics and Data Science, Emirates Aviation University, PO Box 53044, Dubai, UAE

<sup>(3)</sup> International Centre for Research and Postgraduate Training, and

Institute of Mathematics, Vietnam Academy of Science and Technology, Hanoi, Vietnam.

<sup>(4)</sup> Institute of Applied Mathematics, University of Economics Ho Chi Minh City, Vietnam.

January 8, 2024

ABSTRACT. In this paper, we consider the Yang-Mills heat flow on  $\mathbb{R}^d \times SO(d)$  with  $d \geq 11$ . Under a certain symmetry preserved by the flow, the Yang-Mills equation can be reduced to the following nonlinear equation:

$$\partial_t u = \partial_r^2 u + \frac{d+1}{r} \partial_r u - 3(d-2)u^2 - (d-2)r^2 u^3, \text{ and } (r, t) \in \mathbb{R}_+ \times \mathbb{R}_+.$$

We are interested in describing the singularity formation of this parabolic equation. More precisely, we aim to construct non self-similar blowup solutions in higher dimensions  $d \geq 11$ , and prove that the asymptotic of the solution is of the form

$$u(r, t) \sim \frac{1}{\lambda_\ell(t)} \mathcal{Q} \left( \frac{r}{\sqrt{\lambda_\ell(t)}} \right), \text{ as } t \rightarrow T,$$

where  $\mathcal{Q}$  is the steady state corresponding to the boundary conditions  $\mathcal{Q}(0) = -1, \mathcal{Q}'(0) = 0$  and the blowup speed  $\lambda_\ell$  verifies

$$\lambda_\ell(t) = (C(u_0) + o_{t \rightarrow T}(1)) (T - t)^{\frac{2\ell}{\alpha}} \text{ as } t \rightarrow T, \ell \in \mathbb{N}_+^*, \alpha > 1.$$

In particular, the case  $\ell = 1$  corresponds to the stable type II blowup regime, whereas for the cases  $\ell \geq 2$  corresponds to a finite co-dimensional stable regime.

Our approach here is not based on energy estimates but on a careful construction of time dependent eigenvectors and eigenvalues combined with maximum principle and semigroup pointwise estimates.

## 1. Introduction

Recently, geometric heat flows received a lot of attention from both the mathematics and physics communities. Among these geometric flows, the Yang-Mills heat flow is of a great interest. Let us give a brief survey of the physics behind it (more details can be found in [21] and [18]). The Yang-Mills theory is in some sense a *non-commutative* version of Maxwell's electromagnetism where in the latter, the gauge group is the *abelian* group  $U(1)$ . In order to describe the weak nuclear force, governing the nuclear decay of some particles, Yang and Mills proposed to substitute for the Maxwell's gauge group  $U(1)$  the *non-abelian* gauge group  $SU(2)$ . Let us describe the mathematical

---

2010 *Mathematics Subject Classification.* Primary: 35K50, 35B40; Secondary: 35K55, 35K57.

*Key words and phrases.* Blowup solutions, Finite time blowup, Type II blowup, Geometric heat flows, Singularity.

**Acknowledgement:** The work by Giao Ky Duong was supported by the International Center for Research and Postgraduate Training in Mathematics, Institute of Mathematics, Vietnam Academy of Science and Technology, code ICRTM04\_2021.05. The work by Tej-Eddine Ghou is based upon work supported by Tamkeen under the NYU Abu Dhabi Research Institute grant CG002.

January 8, 2024.

setting of the theory. Consider a Riemannian manifold  $M$  of dimension  $d$ , with a structure group  $G$  (i.e., a semi-simple Lie group) and denote by  $\pi$  the canonical projection. Let  $\mathcal{G}$  be the Lie algebra of  $G$  and  $E$  a principal fibre bundle over  $M$ . Let  $D_A$  be a covariant derivative from  $E$  to  $Ad(E) \otimes T^*M$ . On each coordinate chart  $U_\alpha$ ,  $D_A$  can be represented by the  $\mathcal{G}$ -valued 1-form of  $\kappa + A_\alpha$  where  $\kappa$  is some fixed reference connection (e.g. usual exterior derivative), and  $A_{(\alpha)}$  a  $\mathcal{G}$ -valued 1-form

$$A_{(\alpha)} = \sum_{\mu=1}^d A_{\alpha,\mu} dx^\mu.$$

Since the transition functions are smooth, we can set  $A_{(\alpha)} = A$ . Physically, the vector  $A$  represents the electromagnetic potential.

Let the curvature  $F_A$  be the tensor  $D_A D_A$ . By using a local chart  $U_\alpha$ , one can represent  $F_A$  by the  $\mathcal{G}$ -valued 2-form

$$F_A = \sum_{\mu,\nu} F_{\mu,\nu} dx^\mu \wedge dx^\nu,$$

where

$$F_{\mu,\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu].$$

The second rank covariant tensor  $F_{\mu,\nu}$  is the well-known electromagnetic tensor. The Yang-Mills connections are defined as the *critical points* of Yang-Mills functional  $\mathcal{F}_A$  given by

$$\mathcal{F}_A := \int_M |F_A|^2 dvol_M.$$

The Euler Lagrange equations corresponding to these critical points are

$$\sum_{\nu=1}^d D^\nu F_{\mu,\nu} = 0, \forall \mu = 1, \dots, d,$$

where  $D_\nu = \partial_\nu + [A_\nu, \cdot]$ .

The Yang-Mills *heat flow* is defined as the gradient flow associated to the above problem where  $A$  is the Yang-Mills connection. By using a local chart, the time-dependent connection locally satisfies

$$\begin{cases} \partial_t A_\mu(x, t) + \partial^\nu F_{\mu\nu}(x, t) + [A^\nu, F_{\mu,\nu}](x, t) = 0, & t > 0, \\ A_\mu(x, 0) = A_{\mu,0}(x). \end{cases} \quad (1.1)$$

Note that equation (1.1) is invariant under the following scaling

$$A_\lambda(x, t) = \lambda A(\lambda x, \lambda^2 t), \text{ for } \lambda > 0. \quad (1.2)$$

However, the Yang-Mills *functional* is invariant under scaling symmetry for  $d = 4$ , this is why we refer to this dimension as the energy critical one. For  $d \geq 5$ , we say that the equation is supercritical. Results on the long time existence and uniqueness were obtained in [28] for  $d = 2, 3$ , [24, 31] for  $d = 4$  for weak solutions (see also [29] and [30] for the existence of smooth solutions). In particular, in the case  $d = 4$ , the authors in [30] conjectured finite time singularities do not occur on a compact manifold which recently confirmed by [33]. For the energy *supercritical* problem, i.e.  $d \geq 5$ , there is few results on the global existence and this due to the the gauge invariance of the Yang-Mills heat flow.

Let us restrict ourselves to a special case where  $M = \mathbb{R}^d$  and  $E = \mathbb{R}^d \otimes SO(d)$  is the trivial bundle. In this case, the Yang-Mills connection  $A_\mu (\mu \in \{1, \dots, d\})$  is globally given by its  $SO(d)$ -valued coefficient functions  $A_\mu (\mu = 1, \dots, d)$ . In particular, the Lie algebra  $\mathcal{SO}(d)$  is simply the

space of *skew-symmetric*  $d \times d$  matrices endowed with the commutator bracket. Let us denote the coefficient functions by  $A_\mu = A_\mu^{i,j}$  and make (as in [14]) the following  $\mathcal{SO}(d)$ -equivariant ansatz

$$A_\mu^{i,j}(x, t) = u(|x|, t) \sigma_\mu^{i,j}(x), \text{ where } \sigma_\mu^{i,j}(x) = \delta_\mu^i x^j - \delta_\mu^j x^i, i, j \in \{1, \dots, d\}.$$

We emphasize here that the covariant derivative of  $\sigma$  is zero, so that the ansatz amounts to consider the problem in the Lorentz gauge. The equation reduces to (see [21])

$$\partial_t u = \partial_r^2 u + \frac{d+1}{r} \partial_r u - 3(d-2)u^2 - (d-2)r^2 u^3, \text{ and } (r, t) \in \mathbb{R}_+ \times \mathbb{R}_+. \quad (1.3)$$

The solution to this equation is invariant under the scaling

$$u_\lambda(x, t) = \frac{1}{\lambda} u\left(\frac{x}{\sqrt{\lambda}}, \frac{t}{\lambda}\right) \quad (1.4)$$

for  $\lambda > 0$ . Let us remark that (1.3) is locally well posed in the weighted  $L^2$  space

$$L_{1+r^\alpha}^\infty(\mathbb{R}_+) = \{f \text{ measurable such that } \|(1+r^\alpha)f(r)\|_{L^\infty(\mathbb{R}_+)} < +\infty, \alpha \geq \frac{2}{3}\}.$$

The local well-posedness can be easily proved (via a fixed point argument) by extending to a  $\mathbb{R}^{d+2}$ -problem. Thus, with an arbitrary initial data in  $L_{1+r^\alpha}^\infty$  the solution is either global or it develops a singularity in finite time  $T$  in this space. As a matter of fact, we are interested in the blowup phenomenon and a variety of papers were devoted to the study of singularity formation. First, in [18], the author constructed self-similar blowup solutions with  $5 \leq d \leq 9$ . Besides that, the authors in [35] also gave explicit examples (so-called Weinkove solutions)

$$u_W(x, t) = \frac{1}{T-t} W\left(\frac{r}{\sqrt{T-t}}\right),$$

with

$$W(r) = -\frac{1}{a_1(d)r^2 + a_2(d)}.$$

Here  $a_1(d) = \frac{\sqrt{d-2}}{2\sqrt{2}}$ ,  $a_2(d) = \frac{1}{2}(6d - 16 - (d+2)\sqrt{2d-4})$ . Recently, the authors in [13] have constructed non trivial solutions in the range  $5 \leq d \leq 9$  which approach  $u_W$  in  $L^\infty(\mathbb{R}^+)$  and these solutions corresponding to similar blowup setting. The stability of Weinkove solutions was also proved by [13] and [22]. For higher dimension  $d \geq 10$ , the authors in [6] excluded the existence of self similar blowup solutions and then non-self similar solutions are expected.

We have been successful in constructing non-self similar blowup solutions (so-called Type II blowup solutions). Our results are stated in the following.

**Theorem 1.1** (Existence of stable blowup solution). *Let  $d \geq 11$  be an integer. Then, there exist initial data  $u_0 \in C_0^\infty(\mathbb{R}_+, \mathbb{R})$  such that the corresponding solution to (1.3) blows up in finite time  $T(u_0)$ . Moreover, the following decomposition holds true*

$$u(r, t) = \lambda^{-1}(t) Q\left(\frac{r}{\sqrt{\lambda(t)}}\right) + \tilde{u}(r, t), t \in [0, T), \quad (1.5)$$

where  $Q$  is the ground state of (1.3) satisfying  $Q(0) = -1$  and  $Q'(0) = 0$ ; and the error  $\tilde{u}(r, t)$  satisfies

$$\lambda(t) \|\tilde{u}(\cdot, t)\|_{L^\infty(\mathbb{R}^+)} \rightarrow 0 \text{ as } t \rightarrow T, \quad (1.6)$$

and the blowup speed  $\lambda(t)$  exactly behaves as follows

$$\lambda(t) = C(u_0)(1 + o(1))(T - t)^{\frac{2}{\alpha}}. \quad (1.7)$$

as  $t \rightarrow T$  and  $\alpha$  defined in (2.27). In particular, the constructed blowup behavior is stable.

By a suitable expansion the construction technique in Theorem 1.1, we can construct *unstable* blowup solutions with different blowup speeds. More precisely, the result reads.

**Theorem 1.2** (Existence of unstable blowup solutions). *Let us consider integer numbers  $\ell \geq 2$  and  $d \geq 11$ . Then, there exist initial data  $u_{0,\ell} \in C_0^\infty(\mathbb{R}_+, \mathbb{R})$  such that the corresponding solution  $u_\ell$  to (1.3) blows up in finite time  $T(u_{0,\ell})$ . Moreover, the following decomposition holds true*

$$u_\ell(r, t) = \lambda_\ell^{-1}(t)Q\left(\frac{r}{\sqrt{\lambda_\ell(t)}}\right) + \tilde{u}_\ell(r, t), \quad (1.8)$$

where  $Q$  is the ground state satisfying  $Q(0) = -1$  and  $Q'(0) = 0$ ; and the error  $\tilde{u}_\ell(r, t)$  satisfies

$$\lambda_\ell(t)\|\tilde{u}_\ell(\cdot, t)\|_{L^\infty(\mathbb{R}_+)} \rightarrow 0 \text{ as } t \rightarrow T, \quad (1.9)$$

and the blowup speed  $\lambda_\ell(t)$  exactly behaves as follows

$$\lambda_\ell(t) = C(u_0)(1 + o(1))(T - t)^{\frac{2\ell}{\alpha}} \text{ as } t \rightarrow T. \quad (1.10)$$

**Remark 1.3** (Related blowup results for PDE's problem). Note that the Yang-Mills heat flow (1.3) has a lot of similarities with the harmonic map heat flow (under corotational symmetry):

$$\partial_t u = \partial_r^2 u + \frac{d+1}{r} \partial_r u - (d-1) \frac{\sin(2u)}{2r^2}, \text{ and } (r, t) \in \mathbb{R}_+ \times \mathbb{R}_+. \quad (1.11)$$

The harmonic map heat flow forms also singularity in finite time, and the self-similar nature of the singularity appears only when  $3 \leq d \leq 6$ , and for  $d \geq 7$  self-similar blowup solutions don't exist [6]. For  $3 \leq d \leq 6$ , the existence of the self-similar solutions is known [17] and the stability has been proved only in the case  $d = 3$  as in [1]. When  $d = 7$  the blowup is not self-similar and the speed  $\lambda$  has a log correction [19], it turns out that the non-self-similar regime is stable when  $d = 7$ . If  $d \geq 8$ , in [20] the authors proved similar results. The results in [20], also in [4], have been proved with a different method. In [20], the result is based on an energy based method, whereas in [4] is based on the maximum principle which does not allow to obtain the stability. In the present paper, we present a new method that has been introduced previously in [7, 10] but combined with ideas from [4]. We will explain those novelties in the remark below.

**Remark 1.4** (Novelty of the paper). We point out that the approach pursued here is more intuitive than the one in [20] for the heat map as it is based on a spectral approach rather than an energy method. Note that here, the selection by the flow of the blow up speed is linked to the eigenvalue  $\lambda_\ell$  of the time dependent linearized operator  $\mathcal{L}_b$ , after perturbing initially  $Q$  in the direction of the eigenvectors  $\phi_\ell$ . Such an idea was not clear in [20]. The length of the paper is due to the heavy and technical construction of the eigenvectors and eigenvalues of  $\mathcal{L}_b$ . In comparison with [20], the use of maximum principle reduce considerably the difficulty of the control of the infinite dimensional part  $\varepsilon_-$ . We believe that this method can be adapted to a large class of parabolic problems.

**Remark 1.5** (Structure of the paper). To be more convenient for the readers, we aim to give the structure of the paper here: We introduce and explain the importance of the different set of variables: self-similar and blowup variables in the second section. In the third and fourth sections we explain the strategy of the proof, and the time dependent spectral analysis strategy. The fifth section aims to provide a proof of the main theorem without technical details where we show that the infinite dimensional problem can be reduced to a finite dimensional one. In other words, we show that the solution can be split into two parts a finite dimensional part and an infinite dimensional one. In the sixth section we study the dynamic of the finite dimensional part under the assumption that the infinite dimensional part of the solution is decaying in a suitable weighted  $L^2$  norm. The seventh section shows that the assumption made in the section 6 on the infinite dimensional part holds after assuming an  $L^\infty$  bound. In the 8th section we prove this  $L^\infty$  bound assumed in the previous section by using maximum principle and pointwise estimates which is

based on the semigroup associating to the linearised operator. The 9th section is devoted to prove the existence of the ground state  $Q$  which solves an non-autonomous second order ODE. To do so, we prove the existence of an heteroclinic trajectory by finding an appropriate trapping set. In the 10th section we sketch the proof of the existence of the unstable blowup solutions and the last section is devoted to the diagonalisation of the time dependent linearised operator  $\mathcal{L}_b$ .

## 2. Mathematical setting

Let  $u$  be a solution to the following equation on  $[0, T)$  for some  $T > 0$

$$\partial_t u = \partial_r^2 u + \frac{d+1}{r} \partial_r u - 3(d-2)u^2 - (d-2)r^2 u^3, \text{ and } (r, t) \in \mathbb{R}_+ \times \mathbb{R}_+. \quad (2.1)$$

Let  $\lambda$  be an unknown blow-up speed satisfying  $\lambda(t) \rightarrow 0$  as  $t \rightarrow T$  and write

$$u(t, r) = \frac{1}{\lambda(t)} v(\xi, s) \quad (2.2)$$

where the blow-up variables  $s$  and  $\xi$  are such that

$$\frac{ds}{dt} = \frac{1}{\lambda}, \quad \xi = \frac{r}{\sqrt{\lambda}}.$$

Simple computation yields

$$\partial_s v = \partial_\xi^2 v + \frac{d+1}{\xi} \partial_\xi v + \frac{1}{2} \frac{\lambda_s}{\lambda} \Lambda_\xi v - 3(d-2)v^2 - (d-2)\xi^2 v^3. \quad (2.3)$$

We anticipate that  $\frac{\lambda_s}{\lambda} \rightarrow 0$  as  $s \rightarrow \infty$ , since the blow-up mechanism is non-self similar, thus,  $v$  is expected to converge to the ground state  $Q$ , which is a solution to

$$Q''(\xi) + \frac{d+1}{\xi} Q'_\xi - 3(d-2)Q^2 - (d-2)\xi^2 Q^3 = 0 \quad (2.4)$$

with the boundary conditions  $Q(0) = -1$  and  $Q'(0) = 0$ .

In order to establish the convergence of  $v$  to the stationary solution  $Q$ , we linearize around the latter and study the operator

$$H_\xi + \frac{1}{2} \frac{\lambda_s}{\lambda} \Lambda_\xi, \quad (2.5)$$

where

$$\Lambda_\xi = 2 + \xi \partial_\xi, \quad (2.6)$$

and

$$H_\xi = \partial_\xi^2 + \frac{d+1}{\xi} \partial_\xi - 3(d-2)(2Q(\xi) + \xi^2 Q^2(\xi)). \quad (2.7)$$

More precisely, we would like to determine the eigenvectors and eigenvalues of the linearized operator which depend on time. To do so, one has to switch to the so-called self-similar variables, i.e., we write the solution  $u$  as

$$u(r, t) = \frac{1}{T-t} w \left( \frac{r}{\sqrt{T-t}}, \tau \right), \quad \tau = -\log(T-t). \quad (2.8)$$

One then finds that  $w$  satisfies

$$\partial_\tau w = \partial_y^2 w + \frac{d+1}{y} \partial_y w - \frac{1}{2} \Lambda_y w - 3(d-2)w^2 - (d-2)y^2 w^3. \quad (2.9)$$

Now, introduce a function  $b$  of time such that

$$b = \frac{\lambda}{T-t}. \quad (2.10)$$

If the blow-up is self-similar,  $b$  would be a (non-zero) constant. In our case, the blow-up is foreseen to be non-self-similar and  $b$  has then to tend to zero as  $t \rightarrow T$ .

Stepping on the fact that our problem is invariant under time translation, we allow the blow-up time to vary. That is, we replace  $T-t$  by some function  $\mu$  and we prove that it behaves like  $T-t$  for  $t \rightarrow T$ . Hence we relax  $b = \frac{\lambda}{\mu}$  instead of  $\frac{\lambda}{T-t}$ . The parameter  $b$  is measuring the non-self similarity of the solution.

- *Notation:* Based on the above, we write

$$u(r, t) = \frac{1}{\mu(t)} w(y, \tau), \quad y = \frac{r}{\sqrt{\mu(t)}} \quad \text{and} \quad \frac{d\tau}{dt} = \frac{1}{\mu(t)}. \quad (2.11)$$

The function  $w$  now satisfies

$$\partial_\tau w = \partial_y^2 w + \frac{d+1}{y} \partial_y w - \beta(\tau) \Lambda_y w - 3(d-2)w^2 - (d-2)y^2 w^3, \quad (2.12)$$

where

$$\beta(\tau) = -\frac{1}{2} \frac{\mu_\tau}{\mu(\tau)}, \quad (2.13)$$

and

$$\Lambda_y f = y \partial_y f + 2f. \quad (2.14)$$

Note that in the self-similar scale  $\mu$ , one needs to linearise around  $Q_b$  instead of  $Q$ , where

$$Q_{b(\tau)}(y) = \frac{1}{b(\tau)} Q\left(\frac{y}{\sqrt{b(\tau)}}\right). \quad (2.15)$$

In addition,  $w$  is global but blows up in infinite time. Indeed, introduce the error

$$\varepsilon(y, \tau) = w(y, \tau) - Q_{b(\tau)}(y). \quad (2.16)$$

By a simple calculations, it leads to

$$\partial_\tau \varepsilon = \mathcal{L}_b(\varepsilon) + B(\varepsilon) + \Phi(y), \quad (2.17)$$

where

$$\mathcal{L}_b = \partial_y^2 + \frac{d+1}{y} \partial_y - \beta(\tau) \Lambda_y - 3(d-2)(2Q_b + Q_b^2 |y|^2), \quad (2.18)$$

and

$$B(\varepsilon) = -3(d-2)(1 + |y|^2 Q_b) \varepsilon^2 - (d-2) |y|^2 \varepsilon^3, \quad (2.19)$$

and

$$\Phi(\cdot, \tau) = \frac{1}{2} \Lambda_y Q_{b(\tau)} \left[ \frac{b'(\tau)}{b(\tau)} - 2\beta(\tau) \right]. \quad (2.20)$$

From the expression of the operator  $H_\xi$ , we have the relation

$$\mathcal{L}_b w(y, \tau) = \frac{1}{b} (H_\xi - b\beta \Lambda_\xi) v(\xi, \tau). \quad (2.21)$$

From Lemma 9.1, we infer that

$$3(d-2) [2Q_b(y) + y^2 Q_b^2] \rightarrow -\frac{3(d-2)}{y^2} \text{ as } b \rightarrow 0 \text{ with } y \neq 0.$$

We next introduce the limit operator

$$\mathcal{L}_\infty^\beta = \partial_y^2 + \frac{d+1}{y} \partial_y - \beta \Lambda_y + \frac{3(d-2)}{y^2}, \quad (2.22)$$

and we set  $\mathcal{L}_\infty^{\frac{1}{2}} = \mathcal{L}_\infty$

$$\mathcal{L}_\infty = \partial_y^2 + \frac{d+1}{y} \partial_y - \frac{1}{2} \Lambda_y + \frac{3(d-2)}{y^2}. \quad (2.23)$$

Let  $\rho = y^{d+1} e^{-\beta \frac{y^2}{2}}$ . Then a simple computation yields

$$\mathcal{L}_\infty^\beta \phi = \frac{1}{\rho} \frac{d}{dy} (\rho \phi') + \frac{3(d-2)}{y^2} \phi - 2\beta \phi. \quad (2.24)$$

In the present paper, we use weighted Sobolev spaces  $L_{\rho_\beta}^2$  and  $H_{\rho_\beta}^1$  where the weight  $\rho_\beta$  is defined by

$$\rho_\beta(y) = \frac{(2\beta)^{\frac{d+2}{2}}}{(4\pi)^{\frac{d+2}{2}}} y^{d+1} e^{-(2\beta) \frac{y^2}{4}}. \quad (2.25)$$

We also denote  $\rho_{\frac{1}{2}} = \rho$ .

The space  $L_{\rho_\beta}^2$  is equipped with the norm

$$\|f\|_{L_{\rho_\beta}^2(\mathbb{R}^+)}^2 = \int_{\mathbb{R}^+} f^2(y) \rho_\beta(y) dy,$$

and  $H_{\rho_\beta}^1(\mathbb{R}^+)$  has the norm

$$\|f\|_{H_{\rho_\beta}^1(\mathbb{R}^+)}^2 = \|f\|_{L_{\rho_\beta}^2(\mathbb{R}^+)}^2 + \|\partial_y f\|_{L_{\rho_\beta}^2(\mathbb{R}^+)}^2.$$

We also define some special constants in our paper and we assume the dimension  $d \geq 11$ . Let

$$\gamma = \frac{1}{2}(d - \sqrt{d^2 - 12d + 24}), \quad (2.26)$$

$$\alpha = \gamma - 2, \quad (2.27)$$

and

$$a_{i,j} = \frac{(-1)^{i-j}}{(i-j)!} 4^{i-j} \frac{i! (\frac{d}{2} - \gamma)_i!}{j! (\frac{d}{2} - \gamma)_j!} = c_{i,j} C_j, \text{ for all } 0 \leq j \leq i,$$

with  $c_{i,j}$  and  $C_j$  defined as follows

$$c_{i,j} = \frac{(-1)^{i-j} 4^i i! (\frac{d}{2} - \gamma)_i!}{(i-j)!}, \quad (2.28)$$

$$C_j = \frac{1}{4^j j! (\frac{d}{2} - \gamma)_j!}, \quad (2.29)$$

where

$$\left(\frac{d}{2} - \gamma\right)_i! = \left(\frac{d}{2} - \gamma + 1\right) \left(\frac{d}{2} - \gamma + 2\right) \dots \left(\frac{d}{2} - \gamma + i\right) \text{ and } \left(\frac{d}{2} - \gamma\right)_0! = 1.$$

We also use the notation

$$\langle y \rangle = \sqrt{1 + |y|^2}.$$

### 3. Strategy of the proof

We aim to summarize in this paragraph our strategy for the proof of our results. As mentioned above, our goal is to prove that  $v \rightarrow Q$  as  $s \rightarrow \infty$  which is equivalent to the control

$$\|\varepsilon(\cdot, \tau)\|_{L^\infty} \ll \|Q_{b(\tau)}\|_{L^\infty} = b^{-1}(\tau) \text{ as } \tau \rightarrow \infty, \quad (3.1)$$

where  $Q_b$  defined as in (2.15),  $b$  determined as in (2.10), and  $w = Q_b + \varepsilon$  with  $w$  defined in (2.8). Our problem mainly focuses on the perturbative problem (2.17). In addition, the perturbative spectral properties of the linear operator  $\mathcal{L}_b$  is studied in Proposition 4.2 which allows us to expand the error  $\varepsilon$  along its eigenmodes  $\phi_{i,b,\beta}, i \in \{0, 1, \dots, \ell\}$ . More precisely, we arrive at the following decomposition

$$\varepsilon(\tau) = \sum_{j=1}^{\ell-1} \varepsilon_j(\tau) \phi_{j,b(\tau),\beta(\tau)} + \varepsilon_\ell(\tau) \left[ \frac{\phi_{\ell,b(\tau),\beta(\tau)}}{c_{\ell,0}} - \phi_{0,b(\tau),\beta(\tau)} \right] + \varepsilon_-(\tau), \quad (3.2)$$

where  $c_{\ell,0}$  defined in (2.28),  $\varepsilon_-$  is the orthogonal part of  $\varepsilon$  to  $\phi_{i,b,\beta}$  for all  $i \leq \ell$  i.e.

$$\langle \varepsilon_-, \phi_{i,b,\beta} \rangle_{L^2_{\rho_\beta}} = 0, \forall j = 0, \dots, \ell. \quad (3.3)$$

Note that the decomposition in (3.3) is crucial to our approach, as first introduced in [9] (see also [7]). On the one hand, this decomposition provides a good approximation to our solution, including the main perturbative term i.e.

$$\varepsilon_\ell(\tau) \left[ \frac{\phi_{\ell,b(\tau),\beta(\tau)}}{c_{\ell,0}} - \phi_{0,b(\tau),\beta(\tau)} \right]$$

which offers a better approximation compared to the profile when the solution is far from the singular domain. On the other hand, it plays an important role in driving the law of the blowup speed,  $b(\tau)$ . In order to ensure the decomposition (3.2) be unique, we couple the problem (2.17) with

$$c_{\ell,0} \|\phi_{\ell,\phi,\beta}\|_{L^2_{\rho_\beta}}^{-2} \langle \varepsilon, \phi_{\ell,b,\beta} \rangle_{L^2_{\rho_\beta}} = -\|\phi_{0,\phi,\beta}\|_{L^2_{\rho_\beta}}^{-2} \langle \varepsilon, \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} \text{ i.e. } \varepsilon_\ell = -c_{\ell,0} \varepsilon_0, \quad (3.4)$$

and the following compatibility condition (for only the case  $\ell = 1$ )

$$\varepsilon_\ell(\tau) = -\frac{2}{\alpha} m_0 b^{\frac{\alpha}{2}}(\tau). \quad (3.5)$$

Finally, the main issue is to control  $(\varepsilon, b, \beta)$  by a suitable asymptotic behaviors. Specifically, we employ the concept of shrinking set,  $V_\ell[A, \eta, \tilde{\eta}]$  as defined in Definition 5.1 to handle the problem. It's worth noting that the set bears resemblance to recent studies on Type I blowup constructions, such as those found in [3], [25], [27], [11], [16], [15], [12]. More precisely, we control  $\varepsilon_j, j = 0, \dots, \ell, \varepsilon_-$ , the blowup speed  $b$ , the parameter  $\beta(\cdot)$ . Due to the nonlinearity  $y^2 \varepsilon^3$  in equation (2.17), we need to control  $\|\varepsilon_-\|_{L^2_\beta}$  to derive *a priori estimates* on  $\varepsilon_j$  and  $\varepsilon_-$ . Besides that, it is not enough to imply (3.1) from  $\varepsilon_j$  and  $\varepsilon_-$ , since the eigenmodes  $\phi_{i,b,\beta}, i \in \mathbb{N}$  are not bounded as  $y \rightarrow +\infty$ . To address this challenge, we also regulate the outer part  $\varepsilon_e$  introduce in (5.9). Furthermore, we propose a simpler way for constructing Type II blowup solutions for parabolic problems, as an alternative to a direct brute force energy method.

Additionally, we also point out main ideas of the proofs of Theorems 1.1 and 1.2.

- For  $\ell = 1$ . This case involves Theorem 1.1. It is sufficient to control  $(\varepsilon, b, \beta)(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau)$ , for all  $\tau \geq \tau_0$  for some  $\tau_0$  a sufficient large value. The main idea is to construct a suitable initial choice  $(\varepsilon, b, \beta)(\tau_0)$  (see more in subsection 5.2), then we reply in *a priori estimates* provided in Lemmas 6.1, 7.1, 7.2 and 7.3 to improve the bounds in the  $V_1[A, \eta, \tilde{\eta}]$ . Thus, by continuity of the solution in time, we easily conclude that the maximum time trapped in the shrinking set is  $+\infty$ . Finally, using the renormalisation in time given in (5.50), we conclude the proof of Theorem 1.1. We also mention



some interesting points in our proof. First, the control of  $\varepsilon_-$  which we sufficiently do on the interval  $[0, b^{-\tilde{\eta}}(\tau)]$ . On the one hand, on  $[0, b^{\frac{\alpha}{4}}]$  we use the maximum principle, initialled in [4], to control it in avoiding a heavy control from energy method. On the other hand, on  $[b^{\frac{\alpha}{4}}, b^{-\tilde{\eta}}]$  which is far the origin enough. Then, the result follows pointwise estimates based on the Poisson semigroup, see more in section C. Second, the control of the outer part  $\varepsilon_e$ , follows pointwise estimates based on the semigroup  $\mathcal{K}_\beta(\tau, \tau')$ .

- For  $\ell \geq 2$ . This case is related to Theorem 1.2. Similar to the first one. We control the solution to be trapped in the shrinking set  $V_\ell[A, \eta, \tilde{\eta}](\tau)$  by *a priori estimates*. However, this case includes unstable modes that  $\varepsilon_j$  for all  $j \in \{0, \dots, \ell\}$ . Thus, we reduce our problem to a finite dimensional one which is solvent by a classical topological argument.

## 4. Spectral analysis

The aim of this section is to study the linear operator  $\mathcal{L}_b^\beta$ . In order to do so, we begin with the limit operator  $\mathcal{L}_\infty^\beta$ .

**Proposition 4.1** (Diagonalisation of  $\mathcal{L}_\infty^\beta$ , [23], [4], [9]). *Let  $d \geq 11$ ,  $\beta \in (\frac{1}{4}, \frac{3}{4})$  and  $\mathcal{L}_\infty^\beta$  defined as in (2.22). Then,  $\mathcal{L}_\infty^\beta$  admits a unique Friedrichs extension, still denoted by  $\mathcal{L}_\infty^\beta$  with domain  $\mathcal{D}(\mathcal{L}_\infty^\beta) \subset H_{\rho_\beta}^1$  and  $H_{\rho_\beta}^2 \subset \mathcal{D}(\mathcal{L}_\infty^\beta)$ , is self-adjoint with compact resolvent. Moreover, the following hold:*

(i) *Spectrum property:  $\mathcal{L}_\infty^\beta$  consists of countable many eigenvalues. More precisely, the eigenvalues and eigenfunctions are given by*

$$\lambda_{i,\infty,\beta} = 2\beta \left( \frac{\alpha}{2} - i \right), i \in \mathbb{N}, \quad (4.1)$$

$$\begin{aligned} \phi_{i,\infty,\beta}(y) &= \mathcal{N}_i \left( \sqrt{2\beta}y \right)^{-\gamma} L_i^{(\frac{d}{2}-\gamma)} \left( \frac{\beta y^2}{2} \right) = \sum_{j=0}^i a_{i,j} (2\beta)^j y^{2j-\gamma} \\ &= \begin{cases} a_{i,0} y^{-\gamma} (1 + O(y^2)) & \text{as } y \rightarrow 0, \\ a_{i,i} (2\beta)^i y^{2i-\gamma} (1 + O(y^{-2})) & \text{as } y \rightarrow +\infty, \end{cases} \end{aligned} \quad (4.2)$$

where  $L_i^{(\nu)}(z)$  denotes the generalized Laguerre polynomial,  $\mathcal{N}_i$  is a normalization constant and  $\gamma, \alpha, a_{i,j}$  are defined in (2.26), (2.27) and (2.28), respectively.

(ii) *Spectral gap estimate: for all  $u \in H_{\rho_\beta}^1$  satisfying  $\langle \phi_{i,\beta,\infty}, u \rangle_{L_{\rho_\beta}^2} = 0, \forall i \in \{1, \dots, \ell\}$ , then*

$$\langle \mathcal{L}_\infty^\beta u, u \rangle_{L_{\rho_\beta}^2} \leq \lambda_{\ell+1,\infty,\beta} \|u\|_{L_{\rho_\beta}^2}^2.$$

As has been noted above,  $\mathcal{L}_\infty^\beta$  is formally the limit ( $b \rightarrow 0$ ) of  $\mathcal{L}_b$  defined in (2.18), and a priori it is a good approximation of the latter for large values of  $\tau$ . However, such an approximation is good only for  $y$  large enough since  $\phi_{i,\infty}$  is singular when  $y$  approaches 0. Hence, to understand well the operator  $\mathcal{L}_b$  around zero (i.e.,  $y$  small), one has to use the blow-up variables  $(\xi, s)$  and our operator then reads

$$\mathcal{L}_b = \frac{1}{b} (H_\xi - b\beta\Lambda_\xi). \quad (4.3)$$

The strategy is to construct the eigenvalues and eigenvectors of  $\mathcal{L}_b$  in two different regions, namely, for  $y > y_0$  (outer region) using the self-similar scale and and for  $\xi \leq \xi_0$  (inner region) using the variable  $\xi$ . Once such a construction is achieved, we glue at  $y_0$  and in a  $C^1$ -manner the obtained eigenvalues and eigenvectors. The result is summarized in the following proposition.

**Proposition 4.2** (Diagonalisation of  $\mathcal{L}_b$ ). *Let  $d \geq 11, b > 0, \beta \in (\frac{1}{4}, \frac{3}{4}), \ell \in \mathbb{N}^*$  and  $\mathcal{L}_b$  be defined as in (2.18). Then,  $\mathcal{L}_b$  admits a unique Friedrichs extension, still denoted by  $\mathcal{L}_b$  with domain  $\mathcal{D}(\mathcal{L}_b) \subset H_{\rho_\beta}^1$  and  $H_{\rho_\beta}^1 \subset \mathcal{D}(\mathcal{L}_b)$ , is self-adjoint with compact resolvent. Moreover, for all  $\ell \in \mathbb{N}^*$  there exists  $b^*(\ell) \ll 1$  such that for all  $b \in (0, b^*)$  and  $j \leq \ell$ , the following hold:*

(I) *Spectrum: the eigenvalues and eigenfunctions are given by*

$$\lambda_{i,\beta,b} = 2\beta \left( \frac{\alpha}{2} - i \right) + \tilde{\lambda}_{i,\beta,b}, \forall i \in \mathbb{N}, \quad (4.4)$$

$$\phi_{i,\beta,b}(y) = \sum_{j=0}^i c_{i,j} (2\beta)^j (\sqrt{b})^{2j-\gamma} T_j \left( \frac{y}{\sqrt{b}} \right) + \tilde{\phi}_{i,\beta,b}, \quad (4.5)$$

where

$$\|\tilde{\phi}_{i,\beta,b}\|_{H_{\rho_\beta}^1} \lesssim b^{1-\frac{\epsilon}{2}}, \quad \rho_\beta \text{ defined in (2.25)}.$$

In particular, we have

$$|\tilde{\lambda}_{i,\beta,b}| \lesssim b^{1-\frac{\epsilon}{2}} \text{ and } \left| \partial_b \tilde{\lambda}_{i,\beta,b} \right| \lesssim b^{-\frac{\epsilon}{2}}, \text{ and } \left| \partial_\beta \tilde{\lambda}_{i,\beta,b} \right| \lesssim 1. \quad (4.6)$$

(II) *Difference estimate:*

$$\|\phi_{i,\beta,b} - \phi_{i,\beta,\infty}\|_{H_{\rho_\beta}^1} \lesssim b^{1-\frac{\epsilon}{2}}, \quad (4.7)$$

where  $\phi_{i,b,\infty}$  defined as in (4.2)

(III) *Pointwise estimate: for  $k \in \{0, 1\}$  we have*

$$\left| \partial_y^k \phi_{i,\beta,b}(y) \right| \lesssim \frac{\langle y \rangle^{2i+2}}{(\sqrt{b} + y)^{\gamma+k}}, \quad (4.8)$$

$$\left| \partial_y^k b \partial_b \phi_{i,\beta,b}(y) \right| \lesssim \frac{\langle y \rangle^{2i+2}}{(\sqrt{b} + y)^{\gamma+k}}, \quad (4.9)$$

and

$$\left| \partial_y^k \tilde{\phi}_{i,\beta,b}(y) \right| + \left| \partial_y^k b \partial_b \tilde{\phi}_{i,\beta,b}(y) \right| + \left| \partial_y^k \partial_\beta \tilde{\phi}_{i,\beta,b}(y) \right| \lesssim \frac{b^{1-\frac{\epsilon}{2}} \langle y \rangle^{2i+2}}{(\sqrt{b} + y)^{\gamma+k}}. \quad (4.10)$$

(iv) *Spectral gap estimate: assume that  $u \in H_{\rho_\beta}^1(\mathbb{R}^+)$  satisfies*

$$\langle u, \phi_{i,\beta,b} \rangle_{L_{\rho_\beta}^2} = 0, \forall i \in \{0, 1, \dots, \ell\},$$

then, there exists  $c(\ell) > 0$  such that

$$\langle \mathcal{L}_b u, u \rangle_{L_{\rho_\beta}^2} \leq -(\lambda_{\ell,b,\beta} + c(\ell)) \|u\|_{L_{\rho_\beta}^2}^2. \quad (4.11)$$

*Proof.* The spectral analysis is quite the same as in [8] and [9]. We kindly refer the reader to check the details. In addition, we also give the matching ODE approach and the pointwise estimates in Section 10.  $\square$

## 5. Proof in the stable case without technical details

In this section, we aim to give the proof of Theorem 1.1 without technical details.

### 5.1. Shrinking set

We define below the shrinking set that controls the asymptotic behavior of  $(\varepsilon, b, \beta)$  leading to the global existence of the solution, and deriving Theorem 1.1. Let  $b$  and  $\beta$  be positive functions satisfying the hypotheses in Proposition 4.2, then we decompose  $\varepsilon$  as in (3.2) by taking  $\ell = 1$

$$\varepsilon(\tau) = \varepsilon_1(\tau) \left( \frac{\phi_{1,b,\beta}}{c_{1,0}} - \phi_{0,b,\beta} \right) + \varepsilon_-(\tau) = \varepsilon_+(\tau) + \varepsilon_-(\tau). \quad (5.1)$$

**Definition 5.1** (Shrinking set). Let  $A, \eta, \tilde{\eta}$  and  $\tau_0$  be positive constants. For each  $\bar{\tau} > \tau_0$  we introduce  $V_1[A, \eta, \tilde{\eta}](\bar{\tau})$  as the set of all triple time-dependent functions  $(\varepsilon, b, \beta)$  on  $[\tau_0, \bar{\tau}]$  such that  $(\varepsilon, b, \beta)(\tau) \in L^\infty(\mathbb{R}_+) \times \mathbb{R}^2$  for all  $\tau' \in [\tau_0, \bar{\tau}]$  and the following estimates are satisfied:

(i) The dominating mode  $\varepsilon_1$  satisfies

$$\varepsilon_1(\tau) = -\frac{2}{\alpha} m_0 b^{\frac{\alpha}{2}}(\tau), \forall \tau \in [\tau_0, \bar{\tau}], \quad (5.2)$$

and functions  $b$  and  $\beta$  satisfy

$$\frac{1}{2} \leq b(\tau) \exp \left( \left( \frac{2}{\alpha} - 1 \right) \left( \int_{\tau_0}^{\tau} 2\beta(\tilde{\tau}) d\tilde{\tau} + \tau_0 \right) \right) \leq 2, \forall \tau \in [\tau_0, \bar{\tau}], \quad (5.3)$$

and

$$\left| \beta(\tau) - \frac{1}{2} \right| \leq AI^\eta(\tau_0), \forall \tau \in [\tau_0, \bar{\tau}], \quad (5.4)$$

where  $m_0$  is given by (6.15), and  $I(\tau)$  is defined by

$$I(\tau) = e^{(1-\frac{2}{\alpha})\tau}. \quad (5.5)$$

(iii) The part  $\varepsilon_-$  of  $\varepsilon$  defined as in (5.1) satisfies

$$\|\varepsilon_-(\cdot, \tau)\|_{L^2_{\rho_{\beta(\tau)}}} \leq A^2 b^{\frac{\alpha}{2} + \eta}(\tau), \forall \tau \in [\tau_0, \bar{\tau}], \quad (5.6)$$

and

$$\left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{\langle y \rangle^4} \right\|_{L^\infty[0, b^{-\tilde{\eta}}(\tau)]} \leq A^3 b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau), \forall \tau \in [\tau_0, \bar{\tau}]. \quad (5.7)$$

(iv) The part  $\varepsilon_e$  satisfy

$$\|y|\varepsilon_e(\cdot, \tau)\|_{L^\infty} \leq A^4 b^{\frac{\alpha}{2} + (\gamma-4)\tilde{\eta}}(\tau), \forall \tau \in [\tau_0, \bar{\tau}], \quad (5.8)$$

where

$$\varepsilon_e(y, \tau) = (1 - \chi_0(2yb^{\tilde{\eta}}(\tau)))\varepsilon(y, \tau), \text{ and } \text{supp}(\varepsilon_e) \subset \left\{ |y| \geq \frac{1}{2}b^{-\tilde{\eta}} \right\}, \quad (5.9)$$

and  $\chi_0$  defined by

$$\chi_0 \in C^\infty, \chi_0(x) = 1, \forall x \in [0, 1], \text{ and } \chi_0(x) = 0, \forall x \geq 2. \quad (5.10)$$

Consequently, once  $(\varepsilon, b, \beta)$  belongs to  $V_1[A, \eta, \tilde{\eta}]$ , one can easily deduce the following pointwise estimates.

**Lemma 5.2** (Pointwise estimates). *For all  $A \geq 1$  and  $0 < \tilde{\eta} < \eta \ll 1$ , then there exists  $\tau_1(A, \tilde{\eta}, \eta) \geq 1$  such that for all  $\tau_0 \geq \tau_1$  the following holds: Assume  $(\varepsilon, b, \beta)(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau)$  for all  $\tau \in [\tau_0, \bar{\tau}]$  with  $\bar{\tau} > \tau_0$  arbitrarily given, then we have*

$$I^{1+\frac{\tilde{\eta}}{10}}(\tau) \leq b(\tau) \leq I^{1-\frac{\tilde{\eta}}{10}}(\tau), \forall \tau \in [\tau_0, \bar{\tau}]. \quad (5.11)$$

Accordingly (5.1),  $\varepsilon_+$  and  $\varepsilon_-$  satisfy the following pointwise estimates

$$|\varepsilon_+(y, \tau)| \leq \frac{Cb^{\frac{\alpha}{2}}}{y^\gamma} \{y^2 + b^{10\eta}(\tau)\langle y \rangle^4\}, \forall y > 0, \tau \in [\tau_0, \bar{\tau}], \quad (5.12)$$

and

$$|\varepsilon_-(y, \tau)| \leq CA^4 b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau) \frac{\langle y \rangle^4}{y^\gamma}, \forall y > 0, \tau \in [\tau_0, \bar{\tau}]. \quad (5.13)$$

*Proof.* The results immediately follow from the bounds given in Definition 5.1 of the shrinking set  $V_1[A, \eta, \tilde{\eta}](\tau)$ .  $\square$

## 5.2. Preparing initial data

In this part, we aim to construct a suitable family of initial data  $(\varepsilon, b, \beta)(\tau_0)$  such that the solution to the problem (2.17-3.4-3.5) globally exists and satisfies

$$(\varepsilon, b, \beta) \in V_1[A, \eta, \tilde{\eta}](\bar{\tau}), \forall \bar{\tau} > \tau_0.$$

Let us define  $\beta_0 = \beta(\tau_0) = \frac{1}{2}$ ,  $b_0 = b(\tau_0) = I^{\frac{\alpha}{2}}(\tau_0)$  where  $I(\tau)$  introduced as in (5.5),  $\delta \ll 1$  satisfying  $0 < \tilde{\eta} \ll \eta \ll \delta \ll 1$ . In addition, we recall  $\chi_0$  defined as in (5.10) and we introduce then

$$\psi(\tau_0) = \chi_0 \left( y b_0^\delta \right) \left( 1 - \chi_0 \left( \frac{y}{b_0^\delta} \right) \right) \left( -\frac{2}{\alpha} m_0 b_0^{\frac{\alpha}{2}} \right) \left\{ [1 + \hat{\psi}(\tau_0)] \frac{\phi_{1, b_0, \beta_0}}{c_{1,0}} - [1 + \tilde{\psi}(\tau_0)] \phi_{0, b_0, \beta_0} \right\}, \quad (5.14)$$

where the corrections  $\tilde{\psi}(\tau_0)$  and  $\hat{\psi}(\tau_0)$  are uniquely determined such that (3.4-3.5) are satisfied at  $\tau = \tau_0$ . More precisely, via a direct computation, they satisfy

$$\left| \tilde{\psi}(\tau_0) \right| + \left| \hat{\psi}(\tau_0) \right| \lesssim b^\delta(\tau_0).$$

Thus, our initial data is of the form

$$(\varepsilon, b, \beta)(\tau_0) = (\psi(\tau_0), b_0, \beta_0). \quad (5.15)$$

In addition, the initial data for problem (2.12) will be of the form

$$w(y, \tau_0) = Q_{b(\tau_0)}(y) + \varepsilon(\tau_0). \quad (5.16)$$

In the sequel, we prove by using modulation that we can propagate (3.4) and (3.5).

**Lemma 5.3** (Modulation technique). *There exists  $\delta_2 \ll 1$  such that for all  $\delta \leq \delta_2$  there exists  $A_2 \geq 1$  such that for all  $A \geq A_2$  there exists  $\eta_2(A, \delta)$  such that for all  $\eta \leq \eta_2$  there exists  $\tilde{\eta}_2(A, \delta, \eta)$  such that for all  $\tilde{\eta} \leq \tilde{\eta}_2$  there exists  $\tau_2(A, \delta, \eta, \tilde{\eta}) \geq 1$  such that the following property holds: Assume that initial datum is of the form in (5.14), then there exists  $\tau_{loc}^* > \tau_0$  and smooth functions  $(b, \beta) \in (C[\tau_0, \tau_{loc}^*], \mathbb{R}^2) \cap C^1((\tau_0, \tau_{loc}^*), \mathbb{R}^2)$  such that the solution  $w$  (corresponding to initial data in (5.16)) to equation (2.12), locally exists on  $[\tau_0, \tau_{loc}^*]$  and uniquely admits the following decomposition*

$$w(\tau) = Q_{b(\tau)} + \varepsilon(\tau), \quad (5.17)$$

where  $(\varepsilon, b, \beta)$  satisfying (2.17-3.4-3.5) and  $Q_{b(\tau)}$  defined as in (2.15). In addition, it holds that  $(\varepsilon, b, \beta) \in V_1[A, \eta, \tilde{\eta}](\bar{\tau})$ , for all  $\bar{\tau} \in [\tau_0, \tau_{loc}^*]$ . In particular, the existence of  $(\varepsilon, b, \beta)$  can be propagated to the interval  $[\tau_0, \tau_{loc}^* + \tilde{\sigma}]$  for some  $\tilde{\sigma}$  small thank to the bounds in  $V_1$ .

*Proof.* Let us consider initial data  $w(\tau_0)$  defined as in (5.16). Thanks to the local well-posedness in  $L_{1+r}^\infty$ ,  $\alpha \geq \frac{2}{3}$  of the problem (1.3), there exists  $\tilde{\tau} > \tau_0$  such that the solution  $w$  to equation (2.12) uniquely exists on  $[\tau_0, \tilde{\tau}]$ . We mention that the existence of modulations  $b$  and  $\beta$  and decomposition (5.17) is a direct consequence of the implicit function theorem. Let us introduce the following maps

$$\begin{cases} F_1(\tau, b, \beta) & := \langle w(\tau) - Q_b, \|\phi_{1, b, \beta}\|_{L_{\rho_\beta}^2}^{-2} c_{1,0} \phi_{1, b, \beta} + \|\phi_{0, b, \beta}\|_{\rho_\beta}^{-2} \phi_{0, b, \beta} \rangle_{L_{\rho_\beta}^2}, \\ F_2(\tau, b, \beta) & := \langle w(\tau) - Q_b, \phi_{1, b, \beta} \rangle_{L_{\rho_\beta}^2} + \frac{2}{\alpha} m_0 \|\phi_{1, b, \beta}\|_{L_{\rho_\beta}^2}^2 b^{\frac{\alpha}{2}}(\tau). \end{cases} \quad (5.18)$$

Since  $\varepsilon(\tau_0) = \psi(\tau_0)$  defined as in (5.14), it immediately follows

$$F(\tau_0, b_0, \beta_0) = (F_1, F_2)(\tau_0, b_0, \beta_0) = 0.$$

Now we admit the following expansion (the proof will be given below)

$$\begin{aligned} \text{Det}(\mathbf{J})(\tau_0, b_0, \beta_0) &= m_0 \|\phi_{1,b_0,\beta_0}\|_{L^2_{\rho\beta_0}}^2 b_0^{\frac{\alpha}{2}-1} \varepsilon_1(\tau_0) \left( \frac{\|\phi_{1,b_0,\beta_0}\|_{L^2_{\rho\beta_0}}^2 \|\phi_{0,b_0,\beta_0}\|_{L^2_{\rho\beta_0}}^{-2}}{4\beta_0} \right) \\ &+ o(b_0^{\alpha-1}), \end{aligned} \quad (5.19)$$

which implies

$$\text{Det}(\mathbf{J})(\tau_0, b_0, \beta_0) \neq 0,$$

provided that  $b_0$  small enough i.e.  $\tau_0 \geq \tau_{2,1}(\delta)$ . Finally, we apply the implicit function theorem to conclude the unique existence of the functions  $(b, \beta) \in C([\tau_0, \tilde{\tau}_1], \mathbb{R}^2) \cap C^1((\tau_0, \tilde{\tau}_1], \mathbb{R}^2)$  for some  $\tilde{\tau}_1 > \tau_0$  such that

$$F(\tau, b(\tau), \beta(\tau)) = 0, \forall \tau \in [\tau_0, \tilde{\tau}_1].$$

Now, we define  $\tau_{loc}^* = \min(\tilde{\tau}_1, \tilde{\tau})$  and we define  $\varepsilon(\tau) = w(\tau) - Q_{b(\tau)}$ ,  $\tau \in [\tau_0, \tau_1]$ . Thus,  $(\varepsilon, b, \beta)$  reads (2.17-3.4-3.5) for all  $\tau \in [\tau_0, \tau_{loc}^*]$ .

Besides that, from definition (5.14) and the continuity of the solution, there exists  $A_2(\delta)$  such that for all  $A \geq A_2$  there exists  $\eta_2(A, \delta)$  such that for all  $\eta \leq \eta_2$  there exists  $\tilde{\eta}_2(A, \delta, \eta)$  such that for all  $\tilde{\eta} \leq \tilde{\eta}_2$  there exists  $\tau_2(A, \delta, \eta, \tilde{\eta}) \geq 1$  such that for all  $\tau_0 \geq \tau_2$  and  $\tau_{loc}^* > \tau_0$  such that  $(\varepsilon, b, \beta) \in V_1[A, \eta, \tilde{\eta}](\tilde{\tau}), \forall \tilde{\tau} \in (\tau_0, \tau_{loc}^*]$ . To finish the proof we now complete the proof of (5.19) provided that  $\delta \leq \delta_2, \eta \leq \eta_2(\delta), \tilde{\eta} \leq \tilde{\eta}_2(\delta, \eta)$  and  $\tau_0 \geq \tau_2(\delta, \eta, \tilde{\eta})$ .

Let us recall Jacobian matrix  $\mathbf{J}$  defined by

$$\mathbf{J}(\tau, b, \beta) = \begin{pmatrix} \frac{\partial F_1}{\partial b} & \frac{\partial F_1}{\partial \beta} \\ \frac{\partial F_2}{\partial b} & \frac{\partial F_2}{\partial \beta} \end{pmatrix} (\tau, b, \beta).$$

We now explicitly write the partial derivatives:

$$\begin{aligned} \frac{\partial F_1}{\partial b} &= \int \frac{1}{2b} \Lambda_y Q_b \left( \|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^{-2} c_{\ell,0} \phi_{1,b,\beta} + \|\phi_{0,b,\beta}\|_{\rho\beta}^{-2} \phi_{0,b,\beta} \right) \rho_\beta dy \\ &+ \int (w(\tau) - Q_b) \partial_b \left( \|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^{-2} c_{\ell,0} \phi_{1,b,\beta} + \|\phi_{0,b,\beta}\|_{\rho\beta}^{-2} \phi_{0,b,\beta} \right) \rho_\beta dy, \end{aligned} \quad (5.20)$$

and

$$\begin{aligned} \frac{dF_1}{d\beta} &= \partial_\beta (\|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^{-2}) c_{\ell,0} \int (w - Q_b) \phi_{1,b,\beta} \rho_\beta dy + \|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^{-2} c_{1,0} \int (w - Q_b) \partial_\beta \phi_{1,b,\beta} \rho_\beta dy \\ &+ \|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^{-2} c_{1,0} \int (w - Q_b) \phi_{1,b,\beta} \partial_\beta \rho_\beta dy + \partial_\beta (\|\phi_{0,b,\beta}\|_{L^2_{\rho\beta}}^{-2}) \int (w - Q_b) \phi_{0,b,\beta} \rho_\beta dy \\ &+ \|\phi_{0,b,\beta}\|_{L^2_{\rho\beta}}^{-2} \int (w - Q_b) \partial_\beta \phi_{0,b,\beta} \rho_\beta dy + \|\phi_{0,b,\beta}\|_{L^2_{\rho\beta}}^{-2} \int (w - Q_b) \phi_{0,b,\beta} \partial_\beta \rho_\beta dy, \end{aligned} \quad (5.21)$$

and

$$\begin{aligned} \frac{\partial F_2}{\partial b} &= \int \frac{1}{2b} \Lambda Q_b \phi_{1,b,\beta} \rho_\beta dy + \int (w(\tau) - Q_\beta) \partial_b \phi_{1,b,\beta} \rho_\beta dy + \frac{2}{\alpha} m_0 \partial_b \|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^2 b^{\frac{\alpha}{2}} \\ &+ m_0 \|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^2 b^{\frac{\alpha}{2}-1}, \end{aligned} \quad (5.22)$$

and

$$\frac{\partial F_2}{\partial \beta} = \int (w(\tau) - Q_b) \partial_\beta \phi_{1,b,\beta} \rho_\beta dy + \int (w(\tau) - Q_b) \phi_{1,b,\beta} \partial_\beta \rho_\beta dy + \frac{2}{\alpha} m_0 \partial_\beta \|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^2 b^{\frac{\alpha}{2}}. \quad (5.23)$$

We now claim the following (which will be proved later)

$$\frac{\partial F_1}{\partial b}(\tau_0, b(\tau_0), \beta(\tau_0)) = m_0 b^{\frac{\alpha}{2}-1} + o(b^{\frac{\alpha}{2}-1}), \quad (5.24)$$

$$\frac{\partial F_1}{\partial \beta}(\tau_0, b(\tau_0), \beta(\tau_0)) = -\frac{1}{\beta} \varepsilon_1(\tau_0) - \|\phi_{1,b,\beta}\|^2 \|\phi_{0,b,\beta}\|^{-2 \frac{\varepsilon_1(\tau_0)}{4\beta}} + o(b^{\frac{\alpha}{2}}), \quad (5.25)$$

$$\frac{\partial F_2}{\partial b}(\tau_0, b(\tau_0), \beta(\tau_0)) = m_0 \|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^2 b^{\frac{\alpha}{2}-1} + o(b^{\frac{\alpha}{2}-1}), \quad (5.26)$$

$$\frac{\partial F_2}{\partial \beta}(\tau_0, b(\tau_0), \beta(\tau_0)) = -\frac{1}{\beta} \|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^2 \varepsilon_1 + o(b^{\frac{\alpha}{2}}). \quad (5.27)$$

Indeed, using estimates ((5.24)-(5.27)) with  $(b(\tau_0), \beta(\tau_0)) = (b_0, \beta_0)$ , we derive

$$\text{Det}(\mathbf{J})(\tau_0, b_0, \beta_0) = m_0 \|\phi_{1,b_0,\beta_0}\|_{L^2_{\rho\beta_0}}^2 b_0^{\frac{\alpha}{2}-1} \varepsilon_1(\tau_0) \left( \frac{\|\phi_{1,b_0,\beta_0}\|_{L^2_{\rho\beta_0}}^2 \|\phi_{0,b_0,\beta_0}\|_{L^2_{\rho\beta_0}}^{-2}}{4\beta_0} \right) + o(b^{\alpha-1}).$$

Thus, we get

$$\text{Det}(\mathbf{J})(\tau_0, b(\tau_0), \beta(\tau_0)) \neq 0,$$

provided that  $b_0$  is small enough. Finally, we apply the implicit function theorem to get existence of  $(b, \beta) \in C([\tau_0, \tau_1], \mathbb{R}^2) \cap C^1((\tau_0, \tau_1], \mathbb{R}^2)$  for some  $\tau_1 > \tau_0$  and  $\varepsilon = w - Q_b$  satisfying the decomposition (7.6) and the compatibility (3.5) for  $\tau \in [\tau_0, \tau_1]$ . In addition to that, since  $\varepsilon = w - Q_b$ ,  $\varepsilon$  evidently solves (2.17)

Let us now give the details of the computation.

- For (5.24):

we have

$$\int \frac{1}{2b} \Lambda_y Q_b \left( \|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^{-2} c_{1,0} \phi_{1,b,\beta} + \|\phi_{0,b,\beta}\|_{L^2_{\rho\beta}}^{-2} \phi_{0,b,\beta} \right) \rho_\beta dy = m_0 b^{\frac{\alpha}{2}-1} + o(b^{\frac{\alpha}{2}-1}).$$

Next, we estimate

$$\partial_b \|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^{-2} = -\|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^{-4} \left( 2 \int \phi_{\ell,b,\beta} \partial_b \phi_{\ell,b,\beta} \rho_\beta dy \right).$$

From (6.25), we get

$$b \partial_b \|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^{-2} = O(b^{1-\frac{\varepsilon}{2}}). \quad (5.28)$$

Since we choose an initial data  $\varepsilon(\tau_0) = \psi(\tau_0)$  defined as in (5.14) and satisfying  $w(\tau_0) - Q_{b_0} = \varepsilon(\tau_0)$ , we obtain that

$$|\varepsilon(\tau_0)| = |w(\tau_0) - Q_b| \leq C b^{\frac{\alpha}{2}} \frac{(1+y^4)}{y^\gamma}.$$

Hence, from (6.25), we infer that

$$\left| \int (w - Q_b) \partial_b \phi_{\ell,b,\beta} \rho_\beta dy \right| = o(b^{\frac{\alpha}{2}-1}).$$

It follows that,

$$\int (w(\tau_0) - Q_b) \partial_b \left( \|\phi_{1,b,\beta}\|_{L^2_{\rho\beta}}^{-2} c_{1,0} \phi_{1,b,\beta} \right) \rho_\beta dy = o(b^{1-\frac{\varepsilon}{2}}).$$

Similarly,

$$\int (w(\tau_0) - Q_b) \partial_b \left( \|\phi_{0,b,\beta}\|_{L^2_{\rho\beta}}^{-2} \phi_{0,b,\beta} \right) \rho_\beta dy = o(b^{1-\frac{\varepsilon}{2}}).$$

Finally, by adding all integrals in (5.20), (5.24) follows.

- For (5.25): from (5.21), we will establish the following estimates

$$\partial_\beta \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} = \left[ \frac{(\frac{d}{2} - \gamma + 1)}{\beta} - \frac{d+2}{2\beta} \right] \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} + O(b^{1-\frac{\epsilon}{2}}), \quad (5.29)$$

$$\partial_\beta \|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} = \left[ \frac{(\frac{d}{2} - \gamma + 1)}{\beta} - \frac{d+2}{2\beta} \right] \|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} + O(b^{1-\frac{\epsilon}{2}}). \quad (5.30)$$

We remark that these estimates are similar, so we only give the proof of (5.29). Indeed, we write

$$\partial_\beta \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} = -\|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^{-4} \left( 2 \int \phi_{1,b,\beta} \partial_\beta \phi_{1,b,\beta} \rho_\beta dy + \int \phi_{1,b,\beta} \phi_{1,b,\beta} \partial_\beta \rho_\beta dy \right).$$

From the construction of  $\phi_{\ell,b,\beta}$  in Proposition 4.2, we have

$$\begin{aligned} \partial_\beta \phi_{1,b,\beta} &= \frac{1}{\beta} c_{1,1}(2\beta)(\sqrt{b})^{2-\gamma} T_1(\xi) + \partial_\beta \tilde{\phi}_{\ell,b,\beta} \\ &= \frac{1}{\beta} \phi_{1,b,\beta} + \sum_{j=0}^{\ell-1} \tilde{c}_j(\beta) \phi_{j,b,\beta} + \tilde{\Phi}_{1,b,\beta}, \text{ with } \|\tilde{\Phi}_1\|_{L^2_{\rho_\beta}} \leq Cb^{1-\frac{\epsilon}{2}}. \end{aligned} \quad (5.31)$$

Then,

$$\int_0^\infty 2\phi_{1,b,\beta} \partial_\beta \phi_{1,b,\beta} \rho_\beta dy = 2 \left( \frac{1}{\beta} \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 + O(b^{1-\frac{\epsilon}{2}}) \right). \quad (5.32)$$

For the second integral, we use the identity

$$\partial_\beta \rho_\beta = \frac{d+2}{2\beta} \rho_\beta - \frac{y^2}{2} \rho_\beta, \quad (5.33)$$

to derive

$$\int_0^\infty \phi_{1,b,\beta}^2 \partial_\beta \rho_\beta dy = \frac{d+2}{2\beta} \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 - \int_0^\infty \phi_{1,b,\beta} \phi_{1,b,\beta} \frac{y^2}{2} \rho_\beta dy.$$

Besides that, by Proposition 4.2 we have

$$\|\phi_{1,b,\beta} - \phi_{1,\infty,\beta}\|_{L^2_{\rho_\beta}} \leq Cb^{1-\frac{\epsilon}{2}},$$

which yields

$$\int_0^\infty \phi_{1,b,\beta} \phi_{1,b,\beta} \frac{y^2}{2} \rho_\beta dy = \int_0^\infty \phi_{1,\infty,\beta} \phi_{1,\infty,\beta} \frac{y^2}{2} \rho_\beta dy + O(b^{1-\frac{\epsilon}{2}}). \quad (5.34)$$

In addition, we use  $\phi_{\ell,\infty,\beta}$  as in (4.2) to get

$$\begin{aligned} \frac{y^2}{2} \phi_{1,\infty,\beta} &= \frac{y^2}{2} \left\{ a_{1,1}(2\beta)y^{2-\gamma} + a_{1,0}(2\beta)^{\ell-1}y^{-\gamma} \right\} \\ &= \frac{1}{4\beta} (2\beta)^2 y^{4-\gamma} - \frac{1}{\beta} \left( \frac{d}{2} - \gamma + 1 \right) (2\beta) y^{2-\gamma} \\ &= \frac{1}{4\beta} \phi_{2,\infty,\beta} - \frac{1}{4\beta} a_{2,1} \phi_{1,\infty,\beta} - \frac{1}{\beta} \left( \frac{d}{2} - \gamma + 1 \right) \phi_{1,\infty,\beta} \\ &= \frac{1}{4\beta} \phi_{2,\infty,\beta} + \left( \frac{2}{\beta} \left( \frac{d}{2} - \gamma + 2 \right) - \frac{1}{\beta} \left( \frac{d}{2} - \gamma + 1 \right) \right) \phi_{1,\infty,\beta}. \end{aligned}$$

Then

$$\begin{aligned} & \int_0^\infty \phi_{1,\infty,\beta} \phi_{1,\infty,\beta} \frac{y^2}{2} \rho_\beta dy = \left( \frac{2}{\beta} \left( \frac{d}{2} - \gamma + 2 \right) - \frac{1}{\beta} \left( \frac{d}{2} - \gamma + 1 \right) \right) \|\phi_{1,\infty,\beta}\|_{L^2_{\rho_\beta}}^2 + O(b^{1-\frac{\epsilon}{2}}) \\ & = \left( \frac{2}{\beta} \left( \frac{d}{2} - \gamma + 2 \right) - \frac{1}{\beta} \left( \frac{d}{2} - \gamma + 1 \right) \right) \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 + O(b^{1-\frac{\epsilon}{2}}). \end{aligned} \quad (5.35)$$

This concludes the proof of (5.29).

Next, we will prove the following

$$\int_0^\infty \phi_{0,b,\beta} \partial_\beta \phi_{1,b,\beta} \rho_\beta dy = \frac{1}{4\beta} \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 + O(b^{1-\frac{\epsilon}{2}}). \quad (5.36)$$

Indeed, using the orthogonality between  $\phi_{0,b,\beta}$  and  $\phi_{1,b,\beta}$  we get

$$\begin{aligned} 0 & = \partial_\beta \int \phi_{0,b,\beta} \phi_{1,b,\beta} \rho_\beta dy = \int \partial_\beta \phi_{0,b,\beta} \phi_{1,b,\beta} \rho_\beta dy + \int \phi_{0,b,\beta} \partial_\beta \phi_{1,b,\beta} \rho_\beta dy \\ & + \int \phi_{0,b,\beta} \phi_{1,b,\beta} \partial_\beta \rho_\beta dy \\ & = \int \phi_{0,b,\beta} \partial_\beta \phi_{1,b,\beta} \rho_\beta dy + \int \phi_{0,b,\beta} \phi_{1,b,\beta} \partial_\beta \rho_\beta dy + O(b^{1-\frac{\epsilon}{2}}). \end{aligned}$$

From (5.33), we obtain

$$\int \phi_{0,b,\beta} \partial_\beta \phi_{1,b,\beta} \rho_\beta dy = \int \frac{y^2}{2} \phi_{0,b,\beta} \phi_{1,b,\beta} \rho_\beta dy + O(b^{1-\frac{\epsilon}{2}}). \quad (5.37)$$

In addition to that, we have the following identity

$$\frac{y^2}{2} \phi_{0,\infty,\beta} = \frac{1}{4\beta} \phi_{1,\infty,\beta} + \frac{\frac{d}{2} - \gamma + 1}{\beta} \phi_{0,\infty,\beta}, \quad (5.38)$$

and (5.36) follows.

- For (5.26): from (5.22), we have

$$\int \frac{\Lambda Q_b}{b} \phi_{1,b,\beta} \rho_\beta dy = o(b^{\frac{\alpha}{2}-1}), \quad (5.39)$$

from (4.5) and the orthogonality between  $\phi_{0,b,\beta}$  and  $\phi_{1,b,\beta}$ . Moreover, (6.25) ensures that

$$\int (w - Q_b) \partial_b \phi_{1,b,\beta} \rho_\beta = o(b^{\frac{\alpha}{2}-1}),$$

and (5.28) implies

$$\frac{2}{\alpha} m_0 \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 b^{\frac{\alpha}{2}} = o(b^{\frac{\alpha}{2}-1}).$$

Finally, we get

$$\frac{\partial F_2}{\partial b}(\tau_0, b(\tau_0), \beta(\tau_0)) = m_0 \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 b^{\frac{\alpha}{2}-1} + o(b^{\frac{\alpha}{2}-1}), \quad (5.40)$$

which concludes (5.26).

- For (5.27) we use

$$|w(\tau_0) - Q_b| \leq C b^{\frac{\alpha}{2}} \frac{(1+y^4)}{y^\gamma},$$

and

$$\partial_\beta \phi_{1,b,\beta} = \frac{1}{\beta} \phi_{1,b,\beta} + \tilde{\Phi}_{1,b,\beta}, \quad \text{with } \|\tilde{\Phi}_1\|_{L^2_{\rho_\beta}} \leq C b^{1-\frac{\epsilon}{2}}.$$



Using

$$w(\tau_0) - Q_b = \varepsilon(\tau_0) = \sum_{j=0}^1 \varepsilon_j(\tau_0) \phi_{j,b,\beta} + \varepsilon_-(\tau_0),$$

we infer

$$\int (w(\tau_0) - Q_b) \partial_\beta \phi_{1,b,\beta} \rho_\beta dy = \varepsilon_\ell(\tau_0) \|\phi_{\ell,b,\beta}\|^2 + o(b^{\frac{\alpha}{2}}).$$

It follows that

$$\int (w(\tau_0) - Q_b) \phi_{1,b,\beta} \partial_\beta \rho_\beta dy = \frac{d+2}{2\beta} \int (w(\tau_0) - Q_b) \phi_{1,b,\beta} \rho_\beta dy - \frac{1}{2} \int (w(\tau_0) - Q_b) \phi_{1,b,\beta} y^2 \rho_\beta dy.$$

Since  $\varepsilon_j = O(b^{\frac{\alpha}{2} + \eta}(\tau_0))$ ,  $|\varepsilon_-|_{L_\rho^2} \leq C b^{\frac{\alpha}{2} + \tilde{\eta}}$ , we get for the first integral is

$$\int (w(\tau_0) - Q_b) \phi_{1,b,\beta} \rho_\beta dy = \varepsilon_1(\tau_0) \|\phi_{1,b,\beta}\|^2. \quad (5.41)$$

For the second integral, we use the expansion of  $w(\tau_0) - Q_b$  to obtain

$$\int (w(\tau_0) - Q_b) \phi_{1,b,\beta} \frac{y^2}{2} \rho_\beta dy = \varepsilon_0(\tau_0) \int \phi_{0,b,\beta} \phi_{1,b,\beta} \frac{y^2}{2} \rho_\beta dy + \varepsilon_1(\tau_0) \int \phi_{1,b,\beta}^2 \frac{y^2}{2} \rho_\beta dy + o(b^{\frac{\alpha}{2}}). \quad (5.42)$$

In addition, we have that

$$\int_0^\infty \phi_{1,b,\beta} \phi_{1,b,\beta} \frac{y^2}{2} \rho_\beta dy = \int_0^\infty \phi_{1,\infty,\beta} \phi_{1,\infty,\beta} \frac{y^2}{2} \rho_\beta dy + O(b^{1-\frac{\epsilon}{2}})$$

and

$$\int_0^\infty \phi_{1,\infty,\beta} \phi_{1,\infty,\beta} \frac{y^2}{2} \rho_\beta dy = \left( \frac{2}{\beta} \left( \frac{d}{2} - \gamma + 2 \right) - \frac{1}{\beta} \left( \frac{d}{2} - \gamma + 1 \right) \right) \|\phi_{1,b,\beta}\|_{L_{\rho_\beta}^2}^2 + O(b^{1-\frac{\epsilon}{2}}).$$

Hence

$$\int (w(\tau_0) - Q_b) \phi_{1,b,\beta} \frac{y^2}{2} \rho_\beta dy = \left( \frac{2}{\beta} \left( \frac{d}{2} - \gamma + 2 \right) - \frac{1}{\beta} \left( \frac{d}{2} - \gamma + 1 \right) \right) \|\phi_{1,b,\beta}\|_{L_{\rho_\beta}^2}^2 + O(b^{1-\frac{\epsilon}{2}}). \quad (5.43)$$

It remains to estimate the last term in  $\frac{dF_2}{d\beta}$ , namely,  $\frac{2}{\alpha} m_0 \partial_\beta \|\phi_{1,b,\beta}\|_{L_{\rho_\beta}^2}^2 b^{\frac{\alpha}{2}}$ . Indeed, we have

$$\partial_\beta \|\phi_{1,b,\beta}\|_{L_{\rho_\beta}^2}^2 = 2 \int \partial_\beta \phi_{1,b,\beta} \phi_{1,b,\beta} \rho_\beta + \int \phi_{1,b,\beta}^2 \partial_\beta \rho_\beta.$$

Arguing as above, we get

$$\partial_\beta \|\phi_{1,b,\beta}\|_{L_{\rho_\beta}^2}^2 = \left( 2 \frac{1}{\beta} + \frac{d+2}{2\beta} - \left( \frac{2}{\beta} \left( \frac{d}{2} - \gamma + 2 \right) - \frac{1}{\beta} \left( \frac{d}{2} - \gamma + 1 \right) \right) \right) \|\phi_{1,b,\beta}\|_{\rho_\beta}^2 + O(b^{1-\frac{\epsilon}{2}}) \quad (5.44)$$

Putting the different contributions of  $\frac{dF_2}{d\beta}$  together, we arrive at

$$\frac{dF_2}{d\beta} = -\frac{1}{\beta} \|\phi_{1,b,\beta}\|_{L_{\rho_\beta}^2}^2 \varepsilon_1 + o(b^{\frac{\alpha}{2}}).$$

as claimed.  $\square$

### 5.3. The proof of Theorem 1.1

In this part, we focus on the proof of Theorem 1.1 which immediately the following result:

**Proposition 5.4.** *There exist  $A, \eta, \tilde{\tau}$  such that we can find  $\delta \gg \eta, \tilde{\eta}$  and  $\tau_0(A, \eta, \tilde{\eta}, \delta)$  small enough such that with initial data  $\varepsilon(\tau_0)$  defined as in (5.14) and  $(b, \beta)(\tau_0) = \left(e^{(1-\frac{2}{\alpha})\tau_0}, \frac{1}{2}\right)$ , the solution  $(\varepsilon, b, \beta)$  exists for all  $\tau \geq \tau_0$  and satisfies*

$$(\varepsilon, b, \beta)(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau), \forall \tau \geq \tau_0.$$

*Proof.* Let us define  $\tau^*$  by

$$\tau^* = \sup\{\tau_1 \geq \tau_0 \text{ such that } (\varepsilon, b, \beta)(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau), \forall \tau \in [\tau_0, \tau_1]\}. \quad (5.45)$$

By contradiction we suppose that  $\tau^* < +\infty$ . Lemmas 7.1, 7.2 and 7.3, the bounds in  $V_1[A, \eta, \tilde{\eta}](\tau^*)$  involving  $\|\varepsilon\|_{L^2_{\rho\beta(\tau)}}$ ,  $\varepsilon_-$  and  $\varepsilon_e$  are improved by a factor  $\frac{1}{2}$ . In addition to that the improvement for  $b$  and  $\beta$  comes from Lemma 6.1. Indeed, we have

$$|\beta(\tau^*) - \beta(\tau_0)| \lesssim A \int_{\tau_0}^{\tau^*} b^{4\eta}(\tau') d\tau' \leq \frac{A}{2} b^\eta(\tau_0), \quad (5.46)$$

provided that  $\tau_0$  is large enough. For the bound on  $b(\tau)$ , we introduce

$$\Psi(\tau) = b(\tau) \exp\left(\left(\frac{2}{\alpha} - 1\right) \left(\int_{\tau_0}^{\tau} 2\beta(\tau') d\tau' + \tau_0\right)\right) \text{ and } \Psi(\tau_0) = 1, \quad (5.47)$$

and from (6.3), we get

$$|\Psi(\tau^*) - 1| \lesssim \int_{\tau_0}^{\tau^*} |\Psi(\tau') b^{4\eta}(\tau')| d\tau' \leq \frac{1}{10}, \quad (5.48)$$

provided that  $\tau_0$  large enough. Thus, the bound of  $b$  in the shrinking set is improved by the factor  $\frac{1}{2}$ . Besides that, by continuity of the solution, there exists  $\nu > 0$  small such that  $(\varepsilon, b, \beta)(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau), \forall \tau \in [\tau^*, \tau^* + \nu]$  which contradicts to  $\tau^*$ 's definition.  $\square$

Now, we aim to give a proof of Theorem 1.1: Let us consider suitable constants such that Proposition 5.4 holds that  $(\varepsilon, b, \beta) \in V[A, \eta, \tilde{\eta}](\tau)$  for all  $\tau > \tau_0$ . Next, we derive the laws of  $b$  and  $\beta$  as follows:

(i): The law of  $b(\tau)$ : Let us introduce

$$\Psi(\tau) = b(\tau) \exp\left(\left(\frac{2}{\alpha} - 1\right) \left[\int_{\tau_0}^{\tau} 2\beta(\tau') + \tau_0\right]\right), \tau \in [\tau_0, +\infty), \text{ with } \Psi(\tau_0) = 1.$$

From Lemma 6.1, we have

$$|\Psi'(\tau)| \lesssim |\Psi(\tau)| b^{4\eta}(\tau), \forall \tau \geq \tau_0,$$

since  $(\varepsilon, b, \beta) \in V_1[A, \eta, \tilde{\eta}](\tau)$  for all  $\tau > \tau_0$ , we get

$$|\Psi(\tau)| \leq C, \text{ and } b^{4\eta}(\tau) \lesssim I^\eta(\tau) \text{ where } I(\tau) \text{ defined in (5.5),}$$

which yields

$$\Psi(\tau) = \Psi(\tau_0) + \int_{\tau_0}^{\infty} \Psi'(\zeta) d\zeta - \int_{\tau}^{\infty} \Psi'(\zeta) d\zeta = \Psi_\infty + O(I^\eta(\tau)) \text{ as } \tau \rightarrow +\infty,$$

with  $\Psi_\infty = \Psi(\tau_0) + \int_{\tau_0}^{\infty} \Psi'(\zeta) d\zeta = 1 + \int_{\tau_0}^{\infty} \Psi'(\zeta) d\zeta$ . Thus, we get

$$b(\tau) = \Psi_\infty \exp\left(\left(1 - \frac{2}{\alpha}\right) \left(\int_{\tau_0}^{\tau} 2\beta(\zeta) d\zeta + \tau_0\right)\right) [1 + O(I^\eta(\tau))] \text{ as } \tau \rightarrow +\infty. \quad (5.49)$$

(ii) Renormalized flow  $\beta(\tau)$  and derivation the law of  $\mu(t)$  defined in (2.11): Using Lemma 6.1 again, we have

$$|\beta'(\tau)| \lesssim b^{4n}(\tau),$$

then we deduce

$$\beta(\tau) = \beta(\tau_0) + \int_{\tau_0}^{\infty} \beta'(\tau') d\tau' - \int_{\tau}^{\infty} \beta'(\tau') d\tau' = \beta_{\infty} + \int_{\tau}^{\infty} \beta'(\tau') d\tau' = \beta_{\infty} + O(I^n(\tau)), \text{ as } \tau \rightarrow +\infty,$$

with

$$\beta_{\infty} = \frac{1}{2} + \int_0^{\infty} \beta'(\zeta) d\zeta.$$

We introduce the renormalized time  $\tilde{\tau}$  by

$$\tilde{\tau} = 2\beta(\tau)\tau, \quad (5.50)$$

which is an invertible function of  $\tau$ . Indeed,

$$\tau = (2\beta_{\infty})^{-1} \tilde{\tau} (1 + O(I^n(\tilde{\tau}))), \text{ as } \tilde{\tau} \rightarrow +\infty. \quad (5.51)$$

We shall remark that we will make an abuse of notation  $\mu(\tau) = \mu(\tilde{\tau}) = \mu(t)$ . The relation

$$\frac{d\mu}{d\tilde{\tau}} = \frac{d\mu}{d\tau} \frac{d\tau}{d\tilde{\tau}},$$

implies, from the fact that  $\mu_{\tau} = -2\beta\mu$  and (5.50)

$$\frac{d\mu}{d\tilde{\tau}} = -\mu(\tilde{\tau}) [1 + O(I^n(\tilde{\tau}))] \text{ as } \tilde{\tau} \rightarrow +\infty.$$

Thus, we get

$$\mu(\tilde{\tau}) = e^{-\tilde{\tau}} (1 + O(I^n(\tilde{\tau}))), \text{ as } \tilde{\tau} \rightarrow +\infty.$$

In addition, we derive from (2.11) that

$$\frac{d\tilde{\tau}}{dt} = \frac{d\tilde{\tau}}{d\tau} \frac{d\tau}{dt} = 2\beta_{\infty} e^{\tilde{\tau}(t)} (1 + O(I^n(\tilde{\tau}(t)))),$$

which implies

$$\tilde{\tau}(t) = -\ln(2\beta_{\infty}(T-t))(1 + O((T-t)^{\tilde{\eta}})), \text{ as } t \rightarrow T,$$

for some  $T = T(\tau_0) > 0$ . From (5.51), we have

$$\tau(t) = (2\beta_{\infty})^{-1} \tilde{\tau}(t) (1 + I^{\tilde{\eta}}(\tilde{\tau}(t))) = -(2\beta_{\infty})^{-1} \ln(2\beta_{\infty}(T-t))(1 + O((T-t)^{\tilde{\eta}})), \text{ as } t \rightarrow T.$$

Substituting  $\mu(t)$ 's formula, we get

$$\mu(t) = \mu(\tilde{\tau}(t)) = 2\beta_{\infty}(T-t) (1 + O((T-t)^{\tilde{\eta}})) \text{ as } t \rightarrow T.$$

Recall that

$$\int_{\tau_0}^{\tau} 2\beta(\tau') d\tau' = 2\beta_{\infty}\tau(1 + O(\tau^{-1})) \text{ as } \tau \rightarrow +\infty,$$

from which we deduce, with the use of (5.49), that

$$\begin{aligned} b(t) &= \Psi_{\infty} \exp \left( \left( 1 - \frac{2}{\alpha} \right) \left( \int_{\tau_0}^{\tau} 2\beta(\zeta) d\zeta + \tau_0 \right) \right) (1 + O(I^n)(\tau(t))) \\ &= \Psi_{\infty} (2\beta_{\infty})^{\frac{2}{\alpha}-1} (T-t)^{\frac{2}{\alpha}-1} \left( 1 + O(|\ln(T-t)|^{-1}) \right) \text{ as } t \rightarrow T. \end{aligned}$$

Introducing  $\lambda(t) = \mu(t)b(t)$  which satisfies

$$\lambda(t) = C(u_0)(T-t)^{\frac{2}{\alpha}} (1 + O(|\ln(T-t)|^{-1})) \text{ as } t \rightarrow T. \quad (5.52)$$

Finally, the conclusion of the Theorem 1.1 immediately follows (2.11), (2.16), the fact  $(\varepsilon, b, \beta) \in V_1[A, \eta, \tilde{\eta}](\tau)$  for all  $\tau > \tau_0$ , and (5.52).  $\square$

## 6. Finite dimensional system

In this part, we study the dynamics of finite modes  $\varepsilon_j(\tau)$  and the modulation parameters  $b$  and  $\beta$ .

**Lemma 6.1.** *Consider  $A \geq 1, \eta > 0, \tilde{\eta} > 0$ , there exists  $\tau_2(A, \eta, \tilde{\eta})$  such that for all  $\tau_0 \geq \tau_2$ , the following holds: Assume that  $(\varepsilon, b, \beta)(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau), \forall \tau \in [\tau_0, \tau_1]$ , for some  $\tau_1 > \tau_0$ , then, we have*

(i) *The dominating mode  $\varepsilon_1$  satisfies*

$$\begin{cases} \partial_\tau \varepsilon_1 - \left[ 2\beta \left( \frac{\alpha}{2} - 1 \right) \right] \varepsilon_1 = O \left( b^{\frac{\alpha}{2} + 4\eta}(\tau) [|\beta'| + |\frac{b_\tau}{b}| + 1] \right), \\ \partial_\tau \varepsilon_1 - \left[ 2\beta \left( \frac{\alpha}{2} \right) \right] \varepsilon_1 + m_0 \left( \frac{b_\tau}{b} - 2\beta \right) b^{\frac{\alpha}{2}} = O \left( b^{\frac{\alpha}{2} + 4\eta}(\tau) [|\beta'| + |\frac{b_\tau}{b}| + 1] \right), \end{cases} \quad (6.1)$$

for all  $\tau \in [\tau_0, \tau_1]$ .

(iii) *For the  $b$  and  $\beta$ , we obtain*

$$|\beta'(\tau)| \leq CA b^{4\eta}(\tau), \quad (6.2)$$

and

$$\left| \frac{b'(\tau)}{b(\tau)} - 2\beta \left( 1 - \frac{2}{\alpha} \right) \right| \leq CA b^{4\eta}(\tau), \quad (6.3)$$

for all  $\tau \in (\tau_0, \tau_1)$ .

*Proof.* Let us consider  $(\varepsilon, b, \beta)(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau), \forall \tau \in [\tau_0, \tau_1]$  and  $\varepsilon(\tau)$  decomposed as in (5.1), and we also recall that

$$\varepsilon_j = \|\phi_{j,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \langle \varepsilon, \phi_{j,b,\beta} \rangle_{L^2_{\rho_\beta}}. \quad (6.4)$$

Then, we obtain from (3.4) that

$$\begin{cases} \varepsilon_1(\tau) = c_{1,0} \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \langle \varepsilon, \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}}, \\ \varepsilon_1(\tau) = -\|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \langle \varepsilon, \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}}. \end{cases} \quad (6.5)$$

By taking  $\tau$ -derivative of the second equation of the above system, we get

$$\begin{aligned} -\partial_\tau \varepsilon_1 &= \partial_\tau \|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \langle \varepsilon, \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} + \|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \langle \partial_\tau \varepsilon, \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} + \|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \langle \varepsilon, \partial_\tau \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} \\ &+ \|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \left\langle \varepsilon, \frac{\partial_\tau \rho_\beta}{\rho_\beta} \right\rangle_{L^2_{\rho_\beta}}, \end{aligned} \quad (6.6)$$

where  $\rho_\beta$  defined as in (2.25). A direct computation gives

$$\begin{aligned} \partial_\tau \|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} &= \partial_\tau \frac{1}{\int_{\mathbb{R}_+} \phi_{0,b,\beta}^2 \rho_\beta dy} = -\frac{2 \int_{\mathbb{R}_+} \phi_{0,b,\beta} \partial_\tau \phi_{0,b,\beta} \rho_\beta dy + \int_{\mathbb{R}_+} \phi_{0,b,\beta} \phi_{0,b,\beta} \partial_\tau \rho_\beta dy}{\left( \int_{\mathbb{R}_+} \phi_{0,b,\beta}^2 \rho_\beta dy \right)^2} \\ &= -\|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-4} \left( 2 \frac{b'}{b} \langle \phi_{0,b,\beta}, b \partial_b \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} + 2\beta' \langle \phi_{0,b,\beta}, \partial_\beta \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} \right. \\ &+ \left. \beta' \left\langle \phi_{0,b,\beta}, \phi_{0,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}} \right), \end{aligned}$$

and

$$\begin{aligned} \langle \varepsilon, \partial_\tau \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} &= \frac{b'}{b} \langle \varepsilon, b \partial_b \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} + \beta' \langle \varepsilon, \partial_\beta \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}}, \\ \left\langle \varepsilon, \phi_{0,b,\beta} \frac{\partial_\tau \rho_\beta}{\rho_\beta} \right\rangle_{L^2_{\rho_\beta}} &= \beta' \left\langle \varepsilon, \phi_{0,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}}. \end{aligned}$$

We plug these equalities into (6.6) to derive

$$-\partial_\tau \varepsilon_1 = \|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \langle \partial_\tau \varepsilon, \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} + \tilde{K}_0, \quad (6.7)$$

where

$$\begin{aligned} \tilde{K}_0 &= -\|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-4} \langle \varepsilon, \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} \left\{ 2 \frac{b'}{b} \langle \phi_{0,b,\beta}, b \partial_b \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} + 2\beta' \langle \phi_{0,b,\beta}, \partial_\beta \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} \right. \\ &\quad \left. + \beta' \left\langle \phi_{0,b,\beta}, \phi_{0,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}} \right\} + \|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \left\{ \frac{b'}{b} \langle \varepsilon, b \partial_b \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} \right. \\ &\quad \left. + \beta' \langle \varepsilon, \partial_\beta \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} + \beta' \left\langle \varepsilon, \phi_{0,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}} \right\}. \end{aligned} \quad (6.8)$$

Similarly, we derive from the first one in (6.31) that

$$c_{1,0}^{-1} \partial_\tau \varepsilon_1 = \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \langle \partial_\tau \varepsilon, \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} + \tilde{K}_1,$$

where

$$\begin{aligned} \tilde{K}_1 &= -\|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^{-4} \langle \varepsilon, \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} \left\{ 2 \frac{b'}{b} \langle \phi_{1,b,\beta}, b \partial_b \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} + 2\beta' \langle \phi_{1,b,\beta}, \partial_\beta \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} \right. \\ &\quad \left. + \beta' \left\langle \phi_{1,b,\beta}, \phi_{1,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}} \right\} + \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \left\{ \frac{b'}{b} \langle \varepsilon, b \partial_b \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} \right. \\ &\quad \left. + \beta' \langle \varepsilon, \partial_\beta \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} + \beta' \left\langle \varepsilon, \phi_{1,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}} \right\}. \end{aligned} \quad (6.9)$$

We have the following system

$$\begin{cases} c_{1,0}^{-1} \partial_\tau \varepsilon_1(\tau) &= \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \langle \partial_\tau \varepsilon, \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} + \tilde{K}_1, \\ -\partial_\tau \varepsilon_1(\tau) &= \|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \langle \partial_\tau \varepsilon, \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} + \tilde{K}_0. \end{cases} \quad (6.10)$$

Since  $\varepsilon$  solves (2.17), we obtain

$$\langle \partial_\tau \varepsilon, \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} = \langle \mathcal{L}_b \varepsilon, \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} + \langle B(\varepsilon), \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} + \langle \Phi, \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}},$$

which implies

$$\begin{cases} c_{1,0}^{-1} \partial_\tau \varepsilon_1 &= \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \left\{ \langle \mathcal{L}_b \varepsilon, \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} + \langle B(\varepsilon), \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} + \langle \Phi, \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} \right\} + \tilde{K}_1 \\ -\partial_\tau \varepsilon_1 &= \|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^{-2} \left\{ \langle \mathcal{L}_b \varepsilon, \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} + \langle B(\varepsilon), \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} + \langle \Phi, \phi_{0,b,\beta} \rangle_{L^2_{\rho_\beta}} \right\} + \tilde{K}_0 \end{cases}. \quad (6.11)$$

We only estimate all terms of the first equation in (6.11), the rest is left to the reader.

- For  $\langle \mathcal{L}_b \varepsilon, \phi_{j,b,\beta} \rangle_{L^2_{\rho_\beta}}$ : Using the fact that  $\mathcal{L}_b$  is self-adjoint and the special decomposition (7.6), we have

$$\langle \mathcal{L}_b \varepsilon, \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} = \langle \varepsilon, \mathcal{L}_b \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} = \lambda_{1,b,\beta} \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 \frac{\varepsilon_1}{c_{1,0}} \quad (6.12)$$

where

$$\lambda_{1,b,\beta} = 2\beta \left( \frac{\alpha}{2} - 1 \right) + \tilde{\lambda}_{1,b,\beta}, \quad \text{with } |\tilde{\lambda}_{1,b,\beta}| \lesssim b^{1-\frac{\varepsilon}{2}}.$$

- For  $\langle B(\varepsilon), \phi_{j,b,\beta} \rangle_{L^2_{\rho_\beta}}$ : We recall  $B(\varepsilon)$  (2.19) in the below

$$B(\varepsilon) = -3(d-2)(1+|y|^2 Q_b) \varepsilon^2 - (d-2)|y|^2 \varepsilon^3.$$

From (5.1) we have

$$\begin{aligned} |3(d-2)(1+y^2 Q_b) \varepsilon^2| &\lesssim \varepsilon_1^2 \left( \frac{\phi_{1,b,\beta}}{c_{1,0}} - \phi_{0,b,\beta} \right)^2 + \varepsilon_-^2, \\ |(d-2)|y|^2 \varepsilon^3| &\lesssim y^2 \left( |\varepsilon_\ell|^3 \left| \frac{\phi_{1,b,\beta}}{c_{1,0}} - \phi_{0,b,\beta} \right|^3 + |\varepsilon_-|^3 \right). \end{aligned}$$

Since  $(\varepsilon, b, \beta)(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau)$ , for all  $\tau \in [\tau_0, \tau_1]$  which ensures the pointwise estimates given in Proposition (4.2) to deduce that

$$\left| \langle B(\varepsilon), \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} \right| \lesssim |\varepsilon_1|^2 + \int_{\mathbb{R}} \left[ \varepsilon_-^2 + y^2 |\varepsilon_-|^3 \right] |\phi_{j,b,\beta}| \rho_\beta dy.$$

Lemma 5.2 yields

$$|\varepsilon_-(y, \tau)| \leq C A^4 b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau) \frac{\langle y \rangle^{2\ell+2}}{y^\gamma}, \quad \forall y \in \mathbb{R}_+^*, \quad (6.13)$$

which yields

$$\int_{\mathbb{R}_+} \left[ \varepsilon_-^2 + y^2 |\varepsilon_-|^3 \right] |\phi_{j,b,\beta}| \rho_\beta dy \lesssim b^{\frac{\alpha}{2} + 4\eta}(\tau).$$

Hence, we get

$$\langle B(\varepsilon), \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} \lesssim b^{\frac{\alpha}{2} + 4\eta}(\tau). \quad (6.14)$$

- For  $\langle \Phi(\cdot, \tau), \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}}$ : Following  $\Phi$ 's definition in (2.20), we have

$$\Phi(y, \tau) = \left[ \frac{b'(\tau)}{b(\tau)} - 2\beta \right] \frac{1}{2b} \Lambda_y Q \left( \frac{y}{\sqrt{b}} \right) = \left[ \frac{b'(\tau)}{b(\tau)} - 2\beta \right] \frac{1}{2b} \Lambda_\xi Q(\xi), \quad \text{with } \xi = \frac{y}{\sqrt{b}}.$$

Accordingly to  $\phi_{0,b,\beta}$ 's formula in Proposition 4.2, and the construction of  $T_0$  in Lemma 10.1, we write  $\Phi$  as follows

$$\Phi = \left[ \frac{b'}{b} - 2\beta \right] m_0 b^{\frac{\alpha}{2}} \phi_{0,b,\beta} + \tilde{\Phi}, \quad (6.15)$$

for some  $m_0 \neq 0$ , and

$$\|\tilde{\Phi}\|_{L^2_{\rho_\beta}} \lesssim \left| \frac{b_\tau}{b} - 2\beta \right| b^{\frac{\alpha}{2} + 1 - \frac{\varepsilon}{2}}.$$

This immediately implies

$$\langle \Phi(\cdot, \tau), \phi_{j,b,\beta} \rangle_{L^2_{\rho_\beta}} = \begin{cases} m_0 b^{\frac{\alpha}{2}} \left[ \frac{b'(\tau)}{b(\tau)} - 2\beta \right] \|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^2 + O\left( \left| \frac{b_\tau}{b} - 2\beta \right| b^{\frac{\alpha}{2} + 1 - \frac{\varepsilon}{2}} \right) & \text{if } j = 0 \\ O\left( \left| \frac{b_\tau}{b} - 2\beta \right| b^{\frac{\alpha}{2} + 1 - \frac{\varepsilon}{2}} \right) & \text{if } j = 1 \end{cases}. \quad (6.16)$$

- For  $\tilde{K}_1$ : Let us consider  $\delta \ll \min(\frac{1}{2}, 1 - \frac{\epsilon}{2})$  with  $\epsilon$  defined as in Proposition (4.2) and  $\delta \gg \eta \gg \tilde{\eta}$ . From  $\tilde{K}_1$ 's definition given in (6.9), we will prove the following bounds:

$$\langle \phi_{1,b,\beta}, b\partial_b \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} \lesssim b^\delta, \quad (6.17)$$

$$\langle \varepsilon, b\partial_b \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} \lesssim b^\delta \left( |\varepsilon_1| + \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{1+y^4} \right\|_{L^\infty_{[0, b^{-\tilde{\eta}}]}} + \|\varepsilon_-\|_{L^2_{\rho_\beta}} \right), \quad (6.18)$$

$$\langle \phi_{1,b,\beta}, \partial_\beta \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} = \frac{1}{\beta} \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 + O(b^\delta), \quad (6.19)$$

$$\begin{aligned} \left\langle \phi_{1,b,\beta}, \phi_{1,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}} &= \left[ \frac{d+2}{2\beta} - \frac{2}{\beta} \left( \frac{d}{2} - \gamma + 2 \right) + \frac{1}{\beta} \left( \frac{d}{2} - \gamma + 1 \right) \right] \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 \\ &+ O(b^\delta), \end{aligned} \quad (6.20)$$

and with the equality (7.6)

$$\langle \varepsilon, \partial_\beta \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} = \frac{1}{\beta} \frac{\varepsilon_1}{c_{1,0}} \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 - \frac{\varepsilon_1}{4\beta} \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 + O \left( \left\| y^\gamma \frac{\varepsilon_-}{1+y^4} \right\|_{L^\infty} \right) \quad (6.21)$$

$$+ O \left( b^\delta \left( |\varepsilon_\ell| + \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{1+y^4} \right\|_{L^\infty_{[0, b^{-\tilde{\eta}}]}} + \|\varepsilon_-\|_{L^2_{\rho_\beta}} \right) \right), \quad (6.22)$$

and

$$\left\langle \varepsilon, \phi_{1,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}} = \begin{cases} \varepsilon_1 \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 \left[ \frac{1}{c_{1,0}} \left( \frac{d+2}{2\beta} - \frac{2}{\beta} \left( \frac{d}{2} - \gamma + 2 \right) + \frac{1}{\beta} \left( \frac{d}{2} - \gamma + 1 \right) \right) + \frac{1}{4\beta} \right] \\ + O \left( b^\delta \left( |\varepsilon_1| + \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{1+y^4} \right\|_{L^\infty_{[0, b^{-\tilde{\eta}}]}} + \|\varepsilon_-\|_{L^2_{\rho_\beta}} \right) \right), \end{cases} \quad (6.23)$$

and

$$\left\langle \varepsilon, \phi_{0,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}} = \frac{\gamma \varepsilon_1}{\beta} \|\phi_{0,b,\beta}\|_{L^2_{\rho_\beta}}^2 + O(b^{\frac{\alpha}{2} + \delta}). \quad (6.24)$$

+ For (6.17): As we assumed  $\delta \ll \frac{1}{2}$ , then, for all  $y \geq b^\delta$ , we have

$$\xi = \frac{y}{\sqrt{b}} \rightarrow +\infty, \text{ as } b \rightarrow 0.$$

Hence, we have

$$\begin{aligned} \left| \partial_b \sum_{j=0}^1 c_{1,j} (\sqrt{b})^{2j-\gamma} T_j(\xi) \right| &= \left| \frac{1}{2b} \sum_{j=0}^1 c_{1,j} (2j-\gamma) (\sqrt{b})^{2j-\gamma} T_j(\xi) - \frac{1}{2b} \sum_{j=0}^1 c_{1,j} (\sqrt{b})^{2j-\gamma} \xi \partial_\xi T_j(\xi) \right| \\ &\lesssim \sum_{j=0}^1 \sqrt{b}^{2j-2-\gamma} \xi^{2j-\gamma-2} |\ln \xi|, \text{ as } \xi \rightarrow +\infty, \end{aligned}$$

and (10.6) ensures that

$$\xi \partial_\xi T_j(\xi) = (2j-\gamma) T_j(\xi) + O(\xi^{-\gamma+2j-2} \ln \xi), \text{ as } \xi \rightarrow +\infty.$$

Then

$$\left| b\partial_b \sum_{j=0}^i c_{i,j} (\sqrt{b})^{2j-\gamma} T_j(\xi) \right| \lesssim \sum_{j=0}^i \sqrt{b}^{2j-\gamma} \xi^{2j-\gamma-2} |\ln \xi|, \text{ as } \xi \rightarrow +\infty,$$

which implies, from (4.5) and the above inequalities, that

$$|b\partial_b\phi_{j,b,\beta}(y)| \lesssim b^{1-\frac{\epsilon}{2}} \frac{\langle y \rangle^2 |\ln y|}{y^{\gamma+2}}, \forall y \geq b^\delta \text{ and } j \leq 1. \quad (6.25)$$

Hence, we derive on the one hand

$$\left| \int_{y \geq b^\delta} \phi_{j,b,\beta} b\partial_b\phi_{i,b,\beta} \rho_\beta dy \right| \leq Cb^\delta, \forall i, j \leq 1. \quad (6.26)$$

On the other hand, we apply the pointwise estimates given in Proposition 4.2 that yields

$$\left| \int_{y \leq b^\delta} \phi_{j,b,\beta} b\partial_b\phi_{i,b,\beta} \rho_\beta dy \right| \lesssim \int_{y \leq b^\delta} y^{d+1-2\gamma} e^{-\frac{2\beta y^2}{4}} dy \lesssim \int_{y \leq b^\delta} y^{d+1-2\gamma} dy \lesssim b^{\delta(d+2-2\gamma)}. \quad (6.27)$$

Combining (6.26) and (6.27), we obtain

$$\left| \langle \phi_{j,b,\beta}, b\partial_b\phi_{i,b,\beta} \rangle_{L^2_{\rho_\beta}} \right| \lesssim b^{1-\frac{\epsilon}{2}} + b^{\delta(d+2-\gamma)} \lesssim b^\delta, i, j \leq 1, \quad (6.28)$$

thus, (6.17) follows.

+ For (6.18): Using (7.6), we estimate as follows

$$|\langle \varepsilon, b\partial_b\phi_{1,b,\beta} \rangle| \lesssim \sum_{j=0}^1 |\varepsilon_j| \left| \langle \phi_{j,b,\beta}, b\partial_b\phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} \right| + \left| \langle \varepsilon_-, b\partial_b\phi_1 \rangle_{L^2_{\rho_\beta}} \right|.$$

Using (6.28), we get

$$\sum_{j=0}^1 |\varepsilon_j| \left| \langle \phi_{j,b,\beta}, b\partial_b\phi_1 \rangle_{L^2_{\rho_\beta}} \right| \lesssim b^\delta \sum_{j=0}^1 |\varepsilon_j|.$$

Next, we estimate the projection on  $\partial_b\phi_1$  of  $\varepsilon_-$ . Indeed, we split the integral:

$$\begin{aligned} \left| \langle \varepsilon_-(\tau), b\partial_b\phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} \right| &\leq \int_0^{b^\delta} |\varepsilon_-(\tau)| |b\partial_b\phi_{1,b,\beta}| \rho_\beta dy + \int_{b^\delta}^{b^{-\bar{\eta}}} |\varepsilon_-(\tau)| |b\partial_b\phi_{1,b,\beta}| \rho_\beta dy \\ &+ \int_{b^{-\bar{\eta}}}^\infty |\varepsilon_-(\tau)| |b\partial_b\phi_{1,b,\beta}| \rho_\beta dy. \end{aligned}$$

For the integral on  $[0, b^\delta]$ , we estimate as follows

$$\begin{aligned} &\left| \int_0^{b^\delta} \varepsilon_-(\tau) b\partial_b\phi_{1,b,\beta} \rho_\beta dy \right| \leq \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{\langle y \rangle^4} \right\|_{L^\infty[0, b^{-\bar{\eta}}]} \int_0^{b^\delta} y^{d+1-2\gamma} \\ &\leq \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{\langle y \rangle^4} \right\|_{L^\infty[0, b^{-\bar{\eta}}]} b^{\delta(d+2-2\gamma)} \leq \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{\langle y \rangle^4} \right\|_{L^\infty[0, b^{-\bar{\eta}}]} b^\delta. \end{aligned}$$

On the interval  $[b^\delta, b^{-\bar{\eta}}]$ , we estimate

$$\begin{aligned} &\left| \int_{b^\delta}^{b^{-\bar{\eta}}} \varepsilon_-(\tau) b\partial_b\phi_{1,b,\beta} \rho_\beta dy \right| \leq b^{1-\frac{\epsilon}{2}} \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{1+y^4} \right\|_{L^\infty[0, b^{-\bar{\eta}}]} \int_{b^\delta}^{b^{-\bar{\eta}}} \frac{(1+y^4)(1+y^2)|\ln y|}{y^{2\gamma+2}} y^{d+1} e^{-\frac{2\beta y^2}{4}} dy \\ &\lesssim b^{1-\frac{\epsilon}{2}} \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{1+y^4} \right\|_{L^\infty[0, b^{-\bar{\eta}}]} \left\{ \int_{b^\delta}^1 y^{d-2\gamma-1} dy + \int_1^{+\infty} \frac{(1+y^4)(1+y^2)|\ln y|}{y^{2\gamma+2}} y^{d+1} e^{-\frac{2\beta y^2}{4}} dy \right\} \\ &\lesssim b^\delta \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{1+y^4} \right\|_{L^\infty[0, b^{-\bar{\eta}}]}. \end{aligned}$$



For the interval  $[b^{-\bar{\eta}}, +\infty)$ , we use Cauchy-Schwarz inequality and (6.25) to arrive at

$$\left| \int_{b^{-\bar{\eta}}}^{+\infty} |\varepsilon_-| |b\partial_b\phi_{1,b,\beta}| \rho_\beta \right| \lesssim \|\varepsilon_-\|_{L^2_{\rho_\beta}} \left( \int_{b^{-\bar{\eta}}}^{+\infty} |b\partial_b\phi_{1,b,\beta}|^2 \rho_\beta \right)^{\frac{1}{2}} \lesssim b^{1-\frac{\varepsilon}{2}} \|\varepsilon_-\|_{L^2_{\rho_\beta}} \lesssim b^\delta \|\varepsilon_-\|_{L^2_{\rho_\beta}},$$

which concludes (6.18) by adding all related terms.

+ For estimates (6.19) and (6.20): Indeed, from (5.31), we immediately deduce (6.19). In addition

to that, by combining (5.34) and (5.35) one gets (6.20).

+ For (6.22): According to (7.6), we decompose  $\varepsilon$  as follows

$$\langle \varepsilon, \partial_\beta \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} = \varepsilon_1(\tau) \left\langle \frac{\phi_{1,b,\beta}}{c_{1,0}} - \phi_{0,b,\beta}, \partial_\beta \phi_{1,b,\beta} \right\rangle_{L^2_{\rho_\beta}} + \langle \varepsilon_-(\tau), \partial_\beta \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}}.$$

Similarly to (6.18), one can deduce that

$$\langle \varepsilon_-(y, \tau), \partial_\beta \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} \lesssim b^\delta \left( \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{1+y^4} \right\|_{L^\infty_{[0, b^{-\bar{\eta}]}}}} + \|\varepsilon_-\|_{L^2_{\rho_\beta}} \right).$$

In addition to that, we derive from (5.32)

$$\left\langle \frac{\phi_{1,b,\beta}}{c_{1,0}}, \partial_\beta \phi_{1,b,\beta} \right\rangle_{L^2_{\rho_\beta}} = \frac{1}{c_{1,0}} \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 + O(b^{1-\frac{\varepsilon}{2}}),$$

and from (5.36), we have

$$\langle \phi_{0,b,\beta}, \partial_\beta \phi_{1,b,\beta} \rangle_{L^2_{\rho_\beta}} = \frac{1}{4\beta} \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 + O(b^{1-\frac{\varepsilon}{2}}).$$

Finally, we use the above facts to get (6.22).

+ For (6.23): We firstly write as follows

$$\begin{aligned} \left\langle \varepsilon, \phi_{1,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}} &= \varepsilon_1(\tau) \left\langle \frac{\phi_{1,b,\beta}}{c_{1,0}} - \phi_{0,b,\beta}, \phi_{1,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}} \\ &+ \left\langle \varepsilon_-(y, \tau), \phi_{1,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}}. \end{aligned}$$

On one hand, we have

$$\begin{aligned} &\left\langle \varepsilon_-(y, \tau), \phi_{1,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}} \\ &\lesssim \sum_{j=1}^{1-1} |\varepsilon_j| + b^\delta \left( \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{1+y^{21+2}} \right\|_{L^\infty_{[0, b^{-\bar{\eta}]}}}} + \|\varepsilon_-\|_{L^2_{\rho_\beta}} \right). \end{aligned}$$

For the rest, we obtain

$$\begin{aligned} \left\langle \frac{\phi_{1,b,\beta}}{c_{1,0}} - \phi_{0,b,\beta}, \phi_{1,b,\beta} \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}} &= \frac{1}{c_{1,0}} \frac{d+2}{2\beta} \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 - \frac{1}{c_{1,0}} \left\langle \phi_{1,b,\beta}, \phi_{1,b,\beta} \frac{y^2}{2} \right\rangle_{L^2_{\rho_\beta}} \\ &+ \left\langle \phi_{0,b,\beta}, \phi_{1,b,\beta} \frac{y^2}{2} \right\rangle_{L^2_{\rho_\beta}}. \end{aligned}$$

Using (5.34), (5.35), (5.36) and (5.37), we have that

$$\left\langle \phi_{0,b,\beta}, \phi_{1,b,\beta} \frac{y^2}{2} \right\rangle_{L^2_{\rho_\beta}} = \frac{1}{4\beta} \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 + O(b^{1-\frac{\epsilon}{2}})$$

and

$$\left\langle \phi_{1,b,\beta}, \phi_{1,b,\beta} \frac{y^2}{2} \right\rangle_{L^2_{\rho_\beta}} = \left( \frac{2}{\beta} \left( \frac{d}{2} - \gamma + 2 \right) - \frac{1}{\beta} \left( \frac{d}{2} - \gamma + 1 \right) \right) \|\phi_{1,b,\beta}\|_{L^2_{\rho_\beta}}^2 + O(b^{1-\frac{\epsilon}{2}}),$$

which implies (6.23).

- For (6.24): The proof is similar to (6.23) which also follows from (5.34), (5.35), (5.36) and (5.37).

Now, combining (6.17) to (6.24), we derive

$$\begin{aligned} \tilde{K}_1 &= -\frac{\varepsilon_1 \beta'}{c_{1,0} \beta} + O\left(\left|\frac{b'}{b}\right| + 1\right) b^\delta \left( |\varepsilon_1| + \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{1+y^4} \right\|_{L^\infty_{[0, b^{-\tilde{\eta}]}}} + \|\varepsilon_-\|_{L^2_{\rho_\beta}} \right) \\ &+ O\left(\beta' b^\delta \left( |\varepsilon_1| + \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{1+y^4} \right\|_{L^\infty_{[0, b^{-\tilde{\eta}]}}} + \|\varepsilon_-\|_{L^2_{\rho_\beta}} \right)\right). \end{aligned}$$

In a convenient way, we denote

$$\begin{aligned} L &= \left(\left|\frac{b'}{b}\right| + 1\right) b^\delta \left( |\varepsilon_1| + \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{1+y^4} \right\|_{L^\infty_{[0, b^{-\tilde{\eta}]}}} + \|\varepsilon_-\|_{L^2_{\rho_\beta}} \right) \\ &+ \left(\beta' \left( b^\delta |\varepsilon_1| + \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{1+y^4} \right\|_{L^\infty_{[0, b^{-\tilde{\eta}]}}} + \|\varepsilon_-\|_{L^2_{\rho_\beta}} \right)\right). \end{aligned}$$

Then, we have

$$\tilde{K}_1 = -\frac{\beta' \varepsilon_1}{c_{1,0} \beta} + O(L). \quad (6.29)$$

- Applying  $\tilde{K}_1$ 's process to  $\tilde{K}_0$ , we get

$$\tilde{K}_0 = -\frac{\|\phi_{1,b,\beta}\|^2 \|\phi_{0,b,\beta}\|^{-2} \beta' \varepsilon_1}{4\beta c_{1,0}} + O(L). \quad (6.30)$$

Now, we are ready to start to the proof of the Lemma.

- *Proof for (i):* We use system (6.11) combined with all of the previous estimates to derive

$$\begin{cases} \partial_\tau \varepsilon_1 &= 2\beta \left(\frac{\alpha}{2} - 1\right) - \frac{\beta'}{\beta} \varepsilon_1 + O(L) + O\left(\left|\frac{b'}{b}\right| - 2\beta \left|b^{\frac{\alpha}{2} + \delta}\right|\right), \\ \partial_\tau \varepsilon_1 &= 2\beta \frac{\alpha}{2} \varepsilon_1 - m_0 \left[\frac{b'}{b} - 2\beta\right] b^{\frac{\alpha}{2}} + O(L) + O\left(\left|\frac{b'}{b}\right| - 2\beta \left|b^{\frac{\alpha}{2} + \delta}\right|\right). \end{cases} \quad (6.31)$$

In particular, since  $(\varepsilon, b, \beta)(\tau) \in V[A, \eta, \tilde{\eta}](\tau), \forall \tau \in [\tau_0, \tau_1]$ , the pointwise estimates in Lemma 5.2, imply (6.1).

- *Proof for (ii):* The results immediately follows item (i).  $\square$

## 7. Control of the infinite dimensional part

In this part, we aim to give *a priori estimates* involving  $\varepsilon_-$  and  $\varepsilon_e$

### 7.1. Energy estimate

In below, we will prove *a priori estimates* on  $\|\varepsilon_-\|_{L^2_{\rho_\beta}}^2$ .

**Lemma 7.1** (A  $L^2_{\rho_\beta}$ -priori estimates on  $\varepsilon_-$ ). *For all  $A \geq 1, \eta$ , and  $\tilde{\eta}$  satisfying  $1 \ll \eta \ll \tilde{\eta}$ , there exists  $\tau_3(A, \eta, \tilde{\eta})$  and  $\tau^*$  such that for all  $\tau_0 \geq \tau_4$  and the solution  $(\varepsilon, b, \beta)(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau^*), \forall \tau \in [\tau_0, \tau^*]$  and*

$$\|\varepsilon_-(\tau)\|_{L^2_{\rho_\beta}} \leq CA b^{\frac{\alpha}{2} + \eta}(\tau), \forall \tau \in [\tau_0, \tau^*]. \quad (7.1)$$

*Proof.* The result is mainly based on the *spectral gap* property. First, we claim that (7.1) follows from

$$\frac{1}{2} \frac{d}{d\tau} \|\varepsilon_-\|_{L^2_{\rho_\beta}}^2 - \left(\frac{\alpha}{2} - 2\right) \|\varepsilon_-\|_{L^2_{\rho_\beta}}^2 \leq CA b^{\alpha+3\eta}. \quad (7.2)$$

Indeed, let us assume (7.2) holds, we infer that

$$\frac{d}{d\tau} \left( e^{2(2-\frac{\alpha}{2})\tau} \|\varepsilon_-(\tau)\|_{L^2_{\rho_\beta}}^2 \right) \leq CA e^{2(2-\frac{\alpha}{2})\tau} b^{\frac{\alpha}{2} + 3\eta}, \forall \tau \in [\tau_0, \tau^*].$$

From the fact  $(\varepsilon, b, \beta)(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau)$ , for all  $\tau \in [\tau_0, \tau^*]$ , we can apply Lemma 5.2 to deduce

$$\begin{aligned} \|\varepsilon_-\|_{L^2_{\rho_\beta}}^2 &\leq e^{-2(2-\frac{\alpha}{2})(\tau-\tau_0)} \|\varepsilon_-(\tau_0)\|_{L^2_{\rho_\beta}}^2 + CA e^{-2(2-\frac{\alpha}{2})\tau} \int_{\tau_0}^{\tau} e^{2(2-\frac{\alpha}{2})\tau'} b^{\frac{\alpha}{2} + 3\eta}(\tau') d\tau' \\ &\leq CA b^{\alpha+2\eta}(\tau), \forall \tau \in [\tau_0, \tau^*]. \end{aligned}$$

Then, (7.1) follows. Now, it remains to give the proof of (7.2). Indeed, we multiply equation (2.17) by  $\varepsilon_-$  and integrate

$$\frac{1}{2} \frac{d}{d\tau} \|\varepsilon_-\|_{L^2_{\rho_\beta}}^2 = \langle \partial_\tau \varepsilon_-, \varepsilon_- \rangle_{L^2_{\rho_\beta}} + \beta' \left\langle \varepsilon_-, \varepsilon_- \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}}. \quad (7.3)$$

Next, we will prove that for all  $\tau \in [\tau_0, \tau^*]$

$$\langle \partial_\tau \varepsilon_-, \varepsilon_- \rangle_{L^2_{\rho_\beta}} \leq \left(\frac{\alpha}{2} - 2\right) \|\varepsilon_-\|_{L^2_{\rho_\beta}}^2 + O(b^{\alpha+3\eta})(\tau), \quad (7.4)$$

$$\left| \beta' \left\langle \varepsilon_-, \varepsilon_- \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L^2_{\rho_\beta}} \right| \lesssim b^{\alpha+3\eta}(\tau). \quad (7.5)$$

Let us start with (7.4). Indeed, from (2.17), and the decomposition

$$\varepsilon = \varepsilon_\ell \left( \frac{\phi_{\ell, b, \beta}}{c_{\ell, 0}} - \phi_{0, b, \beta} \right) + \sum_{j=1}^{\ell-1} \varepsilon_j \phi_{j, b, \beta} + \varepsilon_- := \varepsilon_+ + \varepsilon_-, \quad (7.6)$$

$\varepsilon_-$  solves

$$\partial_\tau \varepsilon_- = \mathcal{L}_b(\varepsilon_-) + B(\varepsilon_+ + \varepsilon_-) + \Phi - \partial_\tau \varepsilon_+ + \mathcal{L}_b \varepsilon_+.$$

Taking  $L^2_{\rho_\beta}$  scalar product to the both sides of the above equation, we deduce

$$\langle \partial_\tau \varepsilon_-, \varepsilon_- \rangle_{\rho_\beta} = \langle \mathcal{L}_b \varepsilon_-, \varepsilon_- \rangle_{\rho_\beta} + \langle B(\varepsilon_+ + \varepsilon_-), \varepsilon_- \rangle_{\rho_\beta} + \langle \Phi - \partial_\tau \varepsilon_+, \varepsilon_- \rangle_{\rho_\beta},$$

since  $\langle \mathcal{L}_b \varepsilon_+, \varepsilon_- \rangle_{L^2_{\rho_\beta}} = 0$ .

+ Estimate to  $\langle \mathcal{L}_b \varepsilon_-, \varepsilon_- \rangle_{L^2_{\rho_\beta}}$ : Using the orthogonality

$$\langle \phi_{j, b, \beta}, \varepsilon_-(\tau) \rangle_{L^2_{\rho_\beta}} = 0, \text{ for } j = 0 \text{ and } j = 1,$$

the spectral gap in Proposition 4.2 ensures

$$\langle \mathcal{L}_b \varepsilon_-, \varepsilon_- \rangle_{L^2_{\rho_\beta}} \leq \lambda_{2, b, \beta} \|\varepsilon_-\|_{L^2_{\rho_\beta}}^2.$$

In addition, we have

$$\lambda_{2,b,\beta} = \frac{\alpha}{2} - 2 + O(b^{1-\frac{\epsilon}{2}}),$$

which yields

$$\langle \mathcal{L}_b \varepsilon_-, \varepsilon_- \rangle_{L_{\rho\beta}^2} \leq \left( \frac{\alpha}{2} - (\ell + 1) \right) \|\varepsilon_-\|_{L_{\rho}^2}^2 + Cb^{\alpha+3\eta}.$$

+ Estimate for  $\langle \Phi - \partial_\tau \varepsilon_+, \varepsilon_- \rangle_{L_{\rho\beta}^2}$ : Recall that

$$\varepsilon(\tau) = \varepsilon_1(\tau) \left( \frac{\phi_{1,b,\beta}}{c_{1,0}} - \phi_{0,b,\beta} \right) + \varepsilon_-(\tau) = \varepsilon_+ + \varepsilon_-.$$

We decompose

$$\langle \Phi - \partial_\tau \varepsilon_+, \varepsilon_- \rangle = \langle \Phi, \varepsilon_- \rangle - \langle \partial_\tau \varepsilon_+, \varepsilon_- \rangle. \quad (7.7)$$

For  $\langle \Phi, \varepsilon_- \rangle_{L_{\rho\beta}^2}$ , we use (6.15) and by Cauchy-Schwarz inequality to deduce that

$$\left| \langle \Phi, \varepsilon_- \rangle_{L_{\rho\beta}^2} \right| = \left| \langle \tilde{\Phi}, \varepsilon_- \rangle_{L_{\rho\beta}^2} \right| \lesssim \|\tilde{\Phi}\|_{L_{\rho\beta}^2} \|\varepsilon_-\|_{L_{\rho\beta}^2} \leq CA^3 b^{\alpha+\tilde{\eta}+1-\frac{\epsilon}{2}} \leq b^{\alpha+6\eta}(\tau), \forall \tau \in [\tau_0, \tau^*].$$

For the second term, we have

$$\begin{aligned} \partial_\tau \varepsilon_+ &= \varepsilon_1' \left[ \frac{\phi_{1,b,\beta}}{c_{1,0}} - \phi_{0,b,\beta} \right] + \varepsilon_j \left[ \frac{b'}{b} b \partial_b \phi_{j,b,\beta} + \beta' \partial_\beta \phi_{j,b,\beta} \right] \\ &+ \varepsilon_1 \left[ \frac{\frac{b'}{b} b \partial_b \phi_{1,b,\beta} + \beta' \partial_\beta \phi_{1,b,\beta}}{c_{1,0}} - \left( \frac{b'}{b} b \partial_b \phi_{0,b,\beta} + \beta' \partial_\beta \phi_{0,b,\beta} \right) \right]. \end{aligned}$$

Note that

$$\langle \varepsilon_-, \phi_{j,b,\beta} \rangle_{L_{\rho\beta}^2} = 0, \text{ for } j = 0 \text{ and } j = 1, \quad (7.8)$$

combining this with (6.2), (6.3), the necessary bounds in  $V_1[A, \eta, \tilde{\eta}](\tau)$ , and Cauchy-Schwarz inequality, we infer

$$\left| \langle \partial_\tau \varepsilon_+, \varepsilon_- \rangle_{L_{\rho\beta}^2} \right| \leq b^{\alpha+3\eta}(\tau).$$

Finally, we give the following estimate

$$\left| \langle \Phi - \partial_\tau \varepsilon_+, \varepsilon_- \rangle_{L_{\rho\beta}^2} \right| \leq b^{\alpha+3\eta}(\tau), \forall \tau \in [\tau_0, \tau^*].$$

- For  $\langle B(\varepsilon), \varepsilon_- \rangle_{L_{\rho\beta}^2}$  with  $\varepsilon = \varepsilon_+ + \varepsilon_-$ . We explicitly write  $B(\varepsilon)$  in (2.19) as follows

$$B(\varepsilon) = -3(n-2)(1+|y|^2 Q_b)(\varepsilon_+^2 + 2\varepsilon_+ \varepsilon_- + \varepsilon_-^2) - (d-2)|y|^2(\varepsilon_+^3 + 3\varepsilon_+^2 \varepsilon_- + 3\varepsilon_+ \varepsilon_-^2 + \varepsilon_-^3).$$

From  $\gamma$ 's definition, we observe that once  $d \geq 11$ , one has

$$\gamma \leq 3.7 < 4.$$

In addition, from the fact that  $(\varepsilon, b, \beta)(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau)$ ,  $\forall \tau \in [\tau_0, \tau^*]$  and (6.13), we have

$$|\varepsilon_+(y)| \leq \frac{Ab^{\frac{\alpha}{2}+\eta}(\tau)\langle y \rangle^4}{y^\gamma} + \frac{b^{\frac{\alpha}{2}} y^2 \langle y \rangle^2}{y^\gamma}, \text{ and } |\varepsilon_-(y)| \leq \frac{A^4 b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau)\langle y \rangle^4}{y^\gamma},$$

which yields

$$\left| \langle B(\varepsilon), \varepsilon_- \rangle_{L_{\rho\beta}^2} \right| \leq Cb^{\alpha+3\eta}.$$

Thus, we finish the proof of (7.4). In particular, using (6.2) and (6.13), we get

$$\left| \beta' \left\langle \varepsilon_-, \varepsilon_- \left( \frac{d+2}{2\beta} - \frac{y^2}{2} \right) \right\rangle_{L_{\rho\beta}^2} \right| \leq b^{\alpha+3\eta}, \forall \tau \in [\tau_0, \tau^*], \quad (7.9)$$

which implies (7.5). Finally, by combining (7.4) and (7.5) we deduce (7.2) and then the proof of the Lemma follows.  $\square$

## 7.2. $L^\infty$ bounds

In order to handle the nonlinear term in the  $L^2_\rho$ -energy estimate, we used the control of a weighted  $L^\infty$ -norm of  $\varepsilon_-$ . The rest of the section is devoted to it. In the next step, we aim to give *a priori* estimates to the infinite part,  $\varepsilon_-$ . More precisely, we have the following proposition:

**Lemma 7.2** (Control of the infinite dimensional part). *Then, there exists  $A_4 \geq 1$  such that for  $A \geq A_3, \delta \ll 1$ , there exists  $\eta_4(A, \delta) \ll 1$  such that for all  $\eta \leq \eta_4$ , there exists  $\tilde{\eta}_4(A, \eta) \ll \eta$  such that for all  $\tilde{\eta} \leq \tilde{\eta}_4$ , there exists  $\tau_4(A, \eta, \tilde{\eta}) \geq 1$ , such that for all  $\tau_0 \geq \tau_5$ , the following holds: assume that initial data is defined as in (5.14) and the solution  $(\varepsilon, b, \beta)(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau), \forall \tau \in [\tau_0, \tau^*]$ , for some  $\tau^* \geq \tau_0$  then we have the following*

$$\left\| \frac{y^\gamma}{\langle y \rangle^4} \varepsilon_-(\cdot, \tau) \right\|_{L^\infty[0, b^{-\tilde{\eta}}(\tau)]} \leq \frac{A^3}{2} I^{\frac{\alpha}{2} + \tilde{\eta}}(\tau), \forall \tau \in [\tau_0, \tau_1]. \quad (7.10)$$

*Proof.* The proof relies on the maximum principal for the control near the origin i.e.  $[0, b^{\frac{\eta}{4}}]$ , and pointwise estimates on  $[b^{\frac{\eta}{4}}, b^{-\tilde{\eta}}]$ .

a) Let us consider  $y \in [0, b^{\frac{\eta}{4}}]$ . We apply Proposition 11.1 to obtain

$$|\varepsilon(y, \tau)| \leq b^{-1}(\tau) H \left( \frac{y}{\sqrt{b(\tau)}} \right) \leq \frac{C b^{\frac{\alpha}{2} + \frac{\eta}{4}}(\tau) \langle y \rangle^4}{y^\gamma}, \text{ for all } y \in [0, b^{\frac{\eta}{4}}].$$

In addition,  $\varepsilon_+$  can be estimated by

$$|\varepsilon_+(y, \tau)| \leq |\varepsilon_1(\tau)| \left| \frac{\phi_{1,b,\beta}}{c_{1,0}} - \phi_{0,b,\beta} \right|. \quad (7.11)$$

On the one hand, we use the pointwise estimates given in Proposition 4.2

$$\left| \frac{\phi_{1,b,\beta}}{c_{1,0}} - \phi_{0,b,\beta} \right| \leq \left| \frac{c_{1,1}}{c_{1,0}} (\sqrt{b})^{2-\gamma} T_1 \left( \frac{y}{\sqrt{b}} \right) \right| + |\tilde{\phi}_{1,b,\beta}| + |\tilde{\phi}_{0,b,\beta}| \leq \frac{C b^{1-\frac{\epsilon}{2}}(\tau) \langle y \rangle^2}{y^\gamma}.$$

On the other hand, from the compatibility

$$\varepsilon_1(\tau) = -\frac{2}{\alpha} m_0 b^{\frac{\alpha}{2}}(\tau),$$

we deduce that

$$|\varepsilon_+(y, \tau)| \leq \frac{C b^{\frac{\alpha}{2} + \frac{\eta}{2}}(\tau) \langle y \rangle^4}{y^\gamma}.$$

Thus, we obtain

$$\sup_{y \in [0, b^{\frac{\eta}{4}}(\tau)]} \frac{y^\gamma}{\langle y \rangle^{2\ell+2}} |\varepsilon_-(y, \tau)| \leq C A b^{\frac{\alpha}{2} + \frac{\eta}{4}}(\tau) \leq \frac{A^3}{2} b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau), \quad (7.12)$$

provided that  $A \geq A_4$ .

b) Let us consider the control on  $[b^{\frac{\eta}{4}}, b^{-\tilde{\eta}}(\tau)]$ . On this domain, we are far the origin so we can not use the spectrum properties of  $\mathcal{L}_\infty$ . The idea is inspired from [4]. We are going to use pointwise estimates based on the semi-group. As for  $\beta = \frac{1}{2}$ ,  $\mathcal{L}_\infty$  has explicit structure. We introduce the basis of  $L^\infty$

$$\phi_{0,\infty} = \phi_{0,\infty,\frac{1}{2}} \text{ and } \phi_{1,\infty} = \phi_{1,\infty,\frac{1}{2}},$$

and for all  $j \geq 2$  we renormalize as in [4, Lemma 3.4] that

$$\phi_{j,\infty}(y) = \mathcal{N}_j y^{-\gamma} L_j^{(\frac{d}{2}-\gamma)} \left( \frac{y^2}{4} \right),$$

where  $L_j^\nu$  denoted by the generalized Laguerre polynomial, and the renormalisation constant  $\mathcal{N}_j$  ensures that  $\|\phi_{j,\infty}\|_{L^2_\rho} = 1$  and

$$\phi_{j,\infty}(y) = \begin{cases} \alpha_j y^{-\gamma} (1 + o(1)) & \text{as } y \rightarrow 0 \\ \beta_j y^{2j-\gamma} (1 + o(1)) & \text{as } y \rightarrow +\infty, \end{cases}$$

with  $\alpha_j$  and  $\beta_j$  satisfies

$$\alpha_j \sim j^{\frac{\omega}{4}} \text{ and } \beta_j = \frac{j^{-\frac{\omega}{4}}}{4^j j!} \text{ as } j \rightarrow +\infty.$$

The pointwise estimates given in Proposition 4.2 ensures that  $\phi_{j,b,\beta}$  is very close to  $\phi_{j,\infty,\beta}$  on this interval by the following

$$|\phi_{j,b(\tau),\beta}(y) - \phi_{j,\infty,\beta}(y)| \lesssim \frac{b^{\frac{\eta}{2}}(\tau) \langle y \rangle^4}{y^\gamma}, \quad \forall y \geq b^{\frac{\eta}{4}}, j \leq 1. \quad (7.13)$$

In addition, the condition

$$\left| \beta(\tau) - \frac{1}{2} \right| \leq A \Gamma^\eta(\tau_0),$$

defined in the Shrinking set  $V_1[A, \eta, \tilde{\eta}](\tau)$  shows that  $\phi_{j,\infty,\beta}$  is close to  $\phi_{j,\infty,\frac{1}{2}} := \phi_{j,\infty}$  since for all  $j$

$$\left| \phi_{j,\infty,\beta}(y) - \phi_{j,\infty,\frac{1}{2}}(y) \right| \lesssim \left| \beta(\tau) - \frac{1}{2} \right| \frac{\langle y \rangle^4}{y^\gamma}, \quad (7.14)$$

$$\left| e^{-\frac{(2\beta)y^2}{4}} - e^{-\frac{y^2}{4}} \right| \lesssim \left| \beta(\tau) - \frac{1}{2} \right| \frac{y^2}{4} e^{-\frac{y^2}{8}}, \quad (7.15)$$

since for all  $\alpha$ , it holds  $|e^\alpha - 1| \leq C\alpha e^\alpha$ , we have

$$|\hat{\varepsilon}_1(\tau)| \lesssim b^{\frac{\alpha}{2}}(\tau), \quad (7.16)$$

where  $\hat{\varepsilon}_j = \|\phi_{j,\infty}\|_{L^2_\rho}^{-2} \langle \varepsilon, \phi_{j,\infty} \rangle_{L^2_\rho}$  is the projection of  $\varepsilon$  on the basis  $\{\phi_{j,\infty}, j \geq 0\}$ . Hence, we use the semi-group pointwise estimates and we decompose  $\varepsilon$  on the basis  $\phi_{j,0}$

$$\varepsilon = \hat{\varepsilon}_+ + \hat{\varepsilon}_-. \quad (7.17)$$

Thus, we will prove that

$$\sup_{y \in [b^{\frac{\eta}{4}}(\tau), b^{-\tilde{\eta}}(\tau)]} \left| \frac{y^\gamma}{\langle y \rangle^4} \hat{\varepsilon}_-(y, \tau) \right| \leq \frac{A^3}{4} b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau). \quad (7.18)$$

Since  $\varepsilon$  satisfies (2.17),  $\hat{\varepsilon}_-$  solves

$$\partial_\tau \hat{\varepsilon}_- = \mathcal{L}_\infty \hat{\varepsilon}_- + \hat{B}(\hat{\varepsilon}_+ + \hat{\varepsilon}_-) + \left( \frac{1}{2} - \beta(\tau) \right) \Lambda_y(\hat{\varepsilon}_-), \quad (7.19)$$

where  $\mathcal{L}_\infty$  was defined in (2.22) by taking  $\beta = \frac{1}{2}$  and  $\hat{B}$  is defined by

$$\begin{aligned} \hat{B} &= -3(d-2) \left[ 2Q_b + y^2 Q_b^2 + \frac{1}{y^2} \right] (\hat{\varepsilon}_+ + \hat{\varepsilon}_-) + B(\hat{\varepsilon}_+ + \hat{\varepsilon}_-) \\ &+ \Phi(\tau) + \mathcal{L}_\infty^\beta(\hat{\varepsilon}_+) - \partial_\tau \hat{\varepsilon}_+, \end{aligned} \quad (7.20)$$

with  $B$  and  $\Phi$  defined in (2.19) and (2.20), respectively. By using Duhamel's formula, we get

$$\hat{\varepsilon}_-(\tau) = e^{(\tau-\tau_0)\mathcal{L}_\infty} \hat{\varepsilon}_-(\tau_0) + \int_{\tau_0}^{\tau} e^{(\tau-\tau')\mathcal{L}_\infty} \left[ \hat{B}(\hat{\varepsilon}_+ + \hat{\varepsilon}_-) + \left( \frac{1}{2} - \beta(\tau') \right) \Lambda_y \hat{\varepsilon}_- \right] (\tau') d\tau'. \quad (7.21)$$

In addition, we denote  $f_-$  as the part of  $f$  which is orthogonal to  $\phi_{0,\infty}$  and  $\phi_{1,\infty}$ . Then

$$f_-(y) = f - \sum_{j=0}^1 \langle f, \phi_{j,\infty} \rangle_{L_\rho^2} \phi_{j,\infty}.$$

In particular, if the series

$$\sum_{j=0}^{\infty} \langle f, \phi_{j,\infty} \rangle_{L_\rho^2} \phi_{j,\infty}$$

is convergent and well defined, then we can define  $f_-$  pointwisely as

$$f_-(y) = \sum_{j=2}^{\infty} \langle f, \phi_{j,\infty} \rangle_{L_\rho^2} \phi_{j,\infty}.$$

Since  $\varepsilon_-$  is orthogonal to  $\phi_{0,\infty}$  and  $\phi_{1,\infty}$ , we can write

$$\begin{aligned} \varepsilon_- &= \left( e^{(\tau-\tau_0)\mathcal{L}_\infty} (\varepsilon_-(\tau_0)) \right)_- \\ &+ \int_{\tau_0}^{\tau} \left( e^{(\tau-\tau')\mathcal{L}_\infty} (\hat{B}(\tau')) \right)_- d\tau' + \int_{\tau_0}^{\tau} \left( \left[ \frac{1}{2} - \beta(\tau') \right] \Lambda \hat{\varepsilon}_-(\tau') \right)_- d\tau'. \end{aligned} \quad (7.22)$$

We remark that (7.18) immediately follows from

$$\sup_{y \in [b^{\frac{\eta}{4}}(\tau), b^{-\tilde{\eta}}(\tau)]} \left| \frac{y^\gamma}{\langle y \rangle^4} \left( e^{(\tau-\tau_0)\mathcal{L}_\infty} \hat{\varepsilon}_-(\tau_0) \right)_-(y, \tau) \right| \leq \frac{A^3}{16} b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau), \quad (7.23)$$

$$\sup_{y \in [b^{\frac{\eta}{4}}(\tau), b^{-\tilde{\eta}}(\tau)]} \left| \frac{y^\gamma}{\langle y \rangle^4} \int_{\tau_0}^{\tau} \left( e^{(\tau-\tau')\mathcal{L}_\infty} [\hat{B}(\tau')] \right)_- d\tau' \right| \leq \frac{A^3}{16} b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau), \quad (7.24)$$

$$\sup_{y \in [b^{\frac{\eta}{4}}(\tau), b^{-\tilde{\eta}}(\tau)]} \left| \frac{y^\gamma}{\langle y \rangle^4} \int_{\tau_0}^{\tau} \left( e^{(\tau-\tau')\mathcal{L}_\infty} \left[ \frac{1}{2} - \beta(\tau') \right] \Lambda_y (\hat{\varepsilon}_-) \right)_- (\tau') d\tau' \right| \leq \frac{A^3}{16} b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau). \quad (7.25)$$

To prove estimates ((7.23) - (7.25)), we need to consider different cases as

- The first case, we consider  $\tau - \tau_0 \leq \frac{\ln A}{K_0}$

- The second case, we consider  $\tau - \tau_0 > \frac{\ln A}{K_0}$ . In addition, the second will be divided again by two sub-cases that  $\frac{1}{L_0} e^{\frac{\tau-\tau_0}{2}(1-\eta(\frac{2\ell}{\alpha}-1))} \leq b^{-\tilde{\eta}}(\tau)$  and  $\frac{1}{L_0} e^{\frac{\tau-\tau_0}{2}(1-\eta(\frac{2\ell}{\alpha}-1))} > b^{-\tilde{\eta}}(\tau)$  and in these sub-cases also includes some smaller case that there are some large constant  $L_0, K_0, R$  appear which are fixed at the end of the proof. Let us go to the details of the proof.

**First case**  $\tau - \tau_0 \leq \frac{\ln A}{K_0}$

- **Proof of (7.23)** : Note that  $K_0 \gg 1$  will be fixed at the end of the proof. Now, we deduce from (5.14) in accordance with the decomposition (7.17), we arrive at

$$|\hat{\varepsilon}_-(\tau_0) y^\gamma| \leq C A b^{\frac{\alpha}{2} + \eta}(\tau_0) \langle y \rangle^4, \quad (7.26)$$

where  $b(\tau_0) = I(\tau_0)$  defined in (5.5). Since  $\langle y \rangle^4$  is increasing, then, we apply Lemma C.1 and we obtain

$$\begin{aligned} \left| e^{(\tau-\tau_0)\mathcal{L}_\infty} \hat{\varepsilon}_-(\tau_0)(y, \tau) \right| &\leq y^{-\gamma} e^{\frac{\alpha(\tau-\tau_0)}{2}} M(\hat{\varepsilon}_-(\tau_0))(y) \\ &\leq CAe^{\frac{\alpha(\tau-\tau_0)}{2}} b^{\frac{\alpha}{2}+\eta}(\tau_0) y^{-\gamma} \frac{\int_y^\infty \langle y' \rangle^4 (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'}{\int_y^\infty (y')^{1+\omega} e^{-\frac{(y')^2}{2}} dy'} \\ &\leq CAe^{\frac{\alpha}{2}(\tau-\tau_0)} b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau) b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau) b^{\frac{\alpha}{2}+\eta}(\tau_0) y^{-\gamma} \langle y \rangle^4. \end{aligned} \quad (7.27)$$

Using (5.11), we get

$$\begin{aligned} e^{\frac{\alpha}{2}(\tau-\tau_0)} b^{\frac{\alpha}{2}+\eta}(\tau_0) b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau) &\leq Ce^{\frac{\alpha}{2}(\tau-\tau_0)} e^{(1-\frac{2}{\alpha})((\frac{\alpha}{2}+\delta)(1-\frac{\tilde{\eta}}{10})\tau_0 - (\frac{\alpha}{2}+\tilde{\eta})(1+\frac{\tilde{\eta}}{10})\tau)} \\ &\leq Ce^{-c(\eta)\tau_0} e^{\frac{\alpha}{2}(\tau-\tau_0)} e^{(1-\frac{2}{\alpha})((\frac{\alpha}{2}+\tilde{\eta})(1+\frac{\tilde{\eta}}{10})\tau_0 - (\frac{\alpha}{2}+\tilde{\eta})(1+\frac{\tilde{\eta}}{10})\tau)} \\ &\leq Ce^{-c(\eta)\tau_0} e^{(\frac{\alpha}{2}+(\frac{2}{\alpha}-1)(\frac{\alpha}{2}+\tilde{\eta})(1+\frac{\tilde{\eta}}{10}))(\tau-\tau_0)} \\ &\leq Ce^{-c(\eta)\tau_0} A^{(\frac{\alpha}{2}+(\frac{2}{\alpha}-1)(\frac{\alpha}{2}+\tilde{\eta})(1+\tilde{\eta}))\frac{1}{K_0}}, \text{ for some } c(\eta) > 0, \end{aligned} \quad (7.28)$$

which yields

$$\left| \frac{y^\gamma}{\langle y \rangle^4} e^{(\tau-\tau_0)\mathcal{L}_\infty} \hat{\varepsilon}_-(\tau_0)(y, \tau) \right| \leq \frac{A^3}{16} b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau), \quad (7.29)$$

provided that  $K_0 \geq K_4, A \geq A_4$ . Finally, we conclude (7.23).

-Proof of (7.24): for  $\tau' \in [\tau_0, \tau]$ , we apply Lemma C.1 to get

$$\left| e^{(\tau-\tau')\mathcal{L}_\infty} [\hat{B}](\tau') \right| \leq Cy^{-\gamma} e^{\frac{\alpha(\tau-\tau')}{2}} \left\{ M(\mathbb{1}_{(0, b^\delta(\tau'))} \hat{B}) + M(\mathbb{1}_{y \geq b^\delta(\tau')} \hat{B}) \right\}. \quad (7.30)$$

To evaluate  $M(\mathbb{1}_{(0, b^\delta(\tau'))} \hat{B})(\tau')$ , we apply the result in Lemma A.1 to obtain

$$\begin{aligned} M(\mathbb{1}_{[0, b^\delta(\tau')] } \hat{B}(\tau')) &\leq C \left[ b^{\frac{\alpha}{2}}(\tau') M(\mathbb{1}_{(0, b^\delta(\tau'))} y^{-\gamma}) + A^3 b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau') M(\mathbb{1}_{(0, b^\delta(\tau'))} y^{-\gamma-2}) \right. \\ &\quad \left. + A^8 b^{2(\frac{\alpha}{2}+\tilde{\eta})}(\tau') M(\mathbb{1}_{(0, b^\delta(\tau'))} y^{-2\gamma}) + A^{12} b^{3(\frac{\alpha}{2}+\tilde{\eta})}(\tau') M(\mathbb{1}_{[0, b^\delta(\tau')] } y^{-3\gamma+2}) \right]. \end{aligned}$$

For the first term on the right hand side of the above inequality, we rewrite from (C.3)

$$M(\mathbb{1}_{[0, b^\delta(\tau')] } y^{-\gamma}) = \sup_{y \in \mathcal{I}} \frac{\int_{\mathcal{I}} |\mathbb{1}_{[0, b^\delta(\tau'))}| (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'}{\int_{\mathcal{I}} (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'},$$

since  $\mathbb{1}_{[0, b^\delta(\tau'))}$  is non increasing, then, we apply the result in Lemma C.1 to get

$$M(\mathbb{1}_{(0, b^\delta(\tau'))} y^{-\gamma}) \leq \frac{\int_0^{b^\delta(\tau')} |\mathbb{1}_{[0, b^\delta(\tau'))}| (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'}{\int_0^y (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'} \lesssim \frac{(b^\delta(\tau'))^{2+\omega}}{\int_0^y (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'}.$$

Besides that, once  $y \geq 1$ , it follows that

$$\int_0^y (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy' \geq C,$$

which yields

$$\left( \int_0^y (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy' \right)^{-1} \lesssim \frac{\langle y \rangle^{2+\omega}}{y^{2+\omega}}.$$



Otherwise, once  $y \leq 1$ , we have

$$\left( \int_0^y (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy' \right)^{-1} \lesssim \left( \int_0^y (y')^{1+\omega} dy' \right)^{-1} \lesssim \frac{\langle y \rangle^{2+\omega}}{y^{2+\omega}}.$$

Then, we derive

$$M(\mathbb{1}_{(0, b^\delta(\tau'))} y^{-\gamma}) \leq \frac{\int_0^{b^\delta(\tau')} |\mathbb{1}_{[0, b^\delta(\tau'))}| (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'}{\int_0^y (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'} \lesssim \frac{(b^\delta(\tau'))^{2+\omega} \langle y \rangle^{2+\omega}}{y^{2+\omega}}.$$

Similarly, we have

$$M(\mathbb{1}_{[0, b^\delta(\tau')]}) y^{-\gamma-2} \lesssim \frac{(b^\delta(\tau'))^\omega \langle y \rangle^{2+\omega}}{y^{2+\omega}}, \text{ and } M(\mathbb{1}_{(0, b^\delta(\tau'))} y^{-2\gamma}) \lesssim \frac{(I^\delta(\tau'))^{\omega+2-\gamma} \langle y \rangle^{2+\omega}}{y^{2+\omega}}$$

and

$$M(\mathbb{1}_{(0, b^\delta(\tau'))} y^{-3\gamma+2}) \lesssim \frac{(b^\delta(\tau'))^{\omega+4-2\gamma} \langle y \rangle^{2+\omega}}{y^{2+\omega}}.$$

Combining all the related terms with the condition that  $y \in [b^{\frac{\eta}{4}}(\tau), b^{-\tilde{\eta}}]$ , we deduce

$$\begin{aligned} \left| e^{(\tau-\tau')\mathcal{L}_\infty} \left[ \mathbb{1}_{(0, b^\delta(\tau'))} \hat{B} \right] (\tau') \right| &\lesssim y^{-\gamma} e^{\frac{\alpha}{2}(\tau-\tau')} b^{\delta(\omega+4-2\gamma)} (\tau') \frac{\langle y \rangle^{\omega+2}}{y^{\omega+2}} \lesssim y^{-\gamma} e^{\frac{\alpha}{2}(\tau-\tau')} b^{\delta(\omega+4-2\gamma) - \frac{\eta}{4}(\omega+2)} (\tau') \\ &\lesssim \frac{b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau) \langle y \rangle^4}{y^\gamma} \left[ e^{\frac{\alpha}{2}(\tau-\tau')} b^{\delta(\omega+4-2\gamma) - \frac{\eta}{4}(\omega+2)} (\tau') b^{-\frac{\alpha}{2} - \tilde{\eta}}(\tau) \right]. \end{aligned} \quad (7.31)$$

In addition, by the same argument used in (7.28) and (7.29) and that fact that  $\tau - \tau_0 \leq \frac{\ln A}{K_0}$ , we have

$$\int_{\tau_0}^{\tau} \left[ e^{\frac{\alpha}{2}(\tau-\tau')} b^{\delta(\omega+4-2\gamma) - \frac{\eta}{4}(\omega+2)} (\tau') b^{-\frac{\alpha}{2} - \tilde{\eta}}(\tau) \right] d\tau' \lesssim A^{\frac{C(\delta, \eta, \tilde{\eta})}{K_0}}.$$

Finally, we conclude

$$\sup_{y \in [b^{\frac{\eta}{4}}(\tau), b^{-\tilde{\eta}}(\tau)]} \left| \frac{y^\gamma}{\langle y \rangle^{2\ell+2}} \int_{\tau_0}^{\tau} e^{(\tau-\tau')\mathcal{L}_\infty} \left| \mathbb{1}_{(0, b^\delta(\tau'))} \hat{B}(\tau') \right| d\tau' \right| \leq \frac{A^3}{16} b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau),$$

provided that  $K_0 \geq K_4$ ,  $A \geq A_5$ , and  $\tau_0 \geq \tau_5(A, K_0, \delta, \eta, \tilde{\eta})$ .

It remains to evaluate  $M(\mathbb{1}_{y \geq b^\delta(\tau')} \hat{B}(\tau'))$ . Using Lemma A.1 with  $\ell = 1$ , we get

$$\begin{aligned} M(\mathbb{1}_{y \geq b^\delta(\tau')} \hat{B}(\tau')) &\lesssim b^{\frac{\alpha}{2}(\tau') + 4\eta} M\left(\mathbb{1}_{y \geq b^\delta(\tau')} \langle y \rangle^4 y^{-\gamma}\right) + b^{\alpha+\delta(1-\gamma)} M\left(\mathbb{1}_{y \geq b^\delta(\tau')} \langle y \rangle^{12} y^{-\gamma}\right) \\ &\quad + b^{\frac{3\alpha}{2} - 2\delta\gamma}(\tau') M\left(\mathbb{1}_{y \geq b^\delta(\tau')} \langle y \rangle^{20} y^{-\gamma}\right). \end{aligned}$$

First, we observe that the function  $\mathbb{1}_{y \geq b^\delta(\tau')} \langle y \rangle^4$  is non decreasing, we apply Lemma C.1 and we have

$$M\left(\mathbb{1}_{y \geq b^\delta(\tau')} \langle y \rangle^4 y^{-\gamma}\right) \lesssim \frac{\int_y^\infty \langle y' \rangle^4 (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'}{\int_y^\infty (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'}.$$

From a standard result on  $\Gamma$  function, we have

$$\frac{\int_y^\infty \langle y' \rangle^4 (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'}{\int_y^\infty (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'} \lesssim \langle y \rangle^4,$$

which implies

$$M \left( \mathbb{1}_{y \geq b^\delta(\tau')} \langle y \rangle^{2\ell+2} y^{-\gamma} \right) \lesssim \langle y \rangle^{2\ell+2}.$$

Similarly, from Lemma A.1 and the fact  $y \leq b^{-\tilde{\eta}}(\tau)$ , we write

$$\begin{aligned} M \left( \mathbb{1}_{y \geq b^\delta(\tau')} \langle y \rangle^{4+8} y^{-\gamma} \right) &\lesssim \langle y \rangle^{4+8} \lesssim b^{-\tilde{\eta}(2+6)}(\tau) \langle y \rangle^4, \\ M \left( \mathbb{1}_{y \geq b^\delta(\tau')} \langle y \rangle^{6+14} y^{-\gamma} \right) &\lesssim \langle y \rangle^{6+14} \lesssim b^{-\tilde{\eta}(4+12)}(\tau) \langle y \rangle^4. \end{aligned}$$

Thus, we derive for all  $y \in [b^\eta(\tau), b^{-\tilde{\eta}}(\tau)]$

$$\begin{aligned} &\left| e^{(\tau-\tau')\mathcal{L}_\infty} \left[ \mathbb{1}_{y \geq b^\delta(\tau')} \hat{B} \right] (\tau') \right| \\ &\lesssim y^{-\gamma} \langle y \rangle^{2+2} e^{\frac{\alpha}{2}(\tau-\tau')} \left[ b^{\frac{\alpha}{2}+4\eta}(\tau') + b^{\alpha+\delta(1-\gamma)}(\tau') b^{-\tilde{\eta}(2+6)}(\tau) + b^{\frac{3\alpha}{2}-2\delta\gamma}(\tau') b^{-\tilde{\eta}(4+12)}(\tau) \right], \end{aligned} \quad (7.32)$$

which implies

$$\begin{aligned} &\frac{y^\gamma}{\langle y \rangle^{2+2}} \int_{\tau_0}^{\tau} e^{(\tau-\tau')\mathcal{L}_\infty} M(\mathbb{1}_{y \geq b^\delta(\tau')} \hat{B}(\tau')) d\tau' \\ &\lesssim b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau) \int_{\tau_0}^{\tau} e^{\frac{\alpha}{2}(\tau-\tau')} b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau) \left[ b^{\frac{\alpha}{2}+4\eta}(\tau') + b^{\alpha+\delta(1-\gamma)}(\tau') b^{-\tilde{\eta}(2+6)}(\tau) + b^{\frac{3\alpha}{2}-2\delta\gamma}(\tau') b^{-\tilde{\eta}(4+12)}(\tau) \right] d\tau'. \end{aligned} \quad (7.33)$$

From the assumption  $\tau - \tau_0 \leq \frac{\ln A}{K_0}$  with  $K_0$  large enough,  $A \geq A_4$  and  $\alpha \gg \delta \gg \eta \gg \tilde{\eta}$ , and  $\tau_0 \geq \tau_4(A, K_0, \delta, \eta, \tilde{\eta})$ , we proceed similarly as in (7.28) and (7.29) and we obtain

$$\int_{\tau_0}^{\tau} e^{\frac{\alpha}{2}(\tau-\tau')} b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau) \left[ b^{\frac{\alpha}{2}+4\eta}(\tau') + b^{\alpha+\delta(1-\gamma)}(\tau') b^{-\tilde{\eta}(2\ell+6)}(\tau) + b^{\frac{3\alpha}{2}-2\delta\gamma}(\tau') b^{-\tilde{\eta}(4\ell+12)}(\tau) \right] d\tau' \leq \frac{A^3}{32}.$$

Finally, we get

$$\frac{y^\gamma}{\langle y \rangle^4} \int_{\tau_0}^{\tau} e^{(\tau-\tau')\mathcal{L}_\infty} M(\mathbb{1}_{y \geq b^\delta(\tau')} \hat{B}(\tau')) d\tau' \leq \frac{A^3}{32} b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau),$$

and (7.24) immediately follows.

- The proof of (7.25): We first recall the following identity

$$\varepsilon_+ + \varepsilon_- = \varepsilon = \hat{\varepsilon}_+ + \hat{\varepsilon}_-,$$

then, we get

$$\hat{\varepsilon}_-(\tau') = \varepsilon_+(\tau') + \varepsilon_-(\tau') + \hat{\varepsilon}_+(\tau').$$

Since  $(\varepsilon, b, \beta)(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau_1), \forall \tau \in [\tau_0, \tau_1]$ , the pointwise estimates given in Lemma (5.2) and also (7.16) hold, so we get a rough estimate for all  $\tau' \in [\tau_0, \tau], \tau \leq \tau_1$

$$\left| \left( \frac{1}{2} - \beta(\tau') \right) \hat{\varepsilon}_-(\tau') \right| \leq CA^6 b^{\frac{\alpha}{2}}(\tau') I^\eta(\tau_0) \frac{\langle y \rangle^4}{y^\gamma}. \quad (7.34)$$

Now, we apply Lemma C.2 and we obtain

$$\left| e^{(\tau-\tau')\mathcal{L}_\infty} \left( \left( \frac{1}{2} - \beta(\tau') \right) \hat{\varepsilon}_-(\tau') \right) \right| \leq CA^6 b^{\frac{\alpha}{2}}(\tau') I^\eta(\tau_0) \frac{\langle y \rangle^6}{y^\gamma}, \forall \tau' \in [\tau_0, \tau],$$

which yields

$$\left| \int_{\tau_0}^{\tau} e^{(\tau-\tau')\mathcal{L}_\infty} \left( \left( \frac{1}{2} - \beta(\tau') \right) \hat{\varepsilon}_-(\tau') \right) d\tau' \right| \leq CA^6 I^\eta(\tau_0) \frac{\langle y \rangle^6}{y^\gamma} \int_{\tau_0}^{\tau} b^{\frac{\alpha}{2}}(\tau') d\tau'.$$

Using Lemma 5.2 and a similar estimate as in (7.28), we derive

$$\left| A^6 I^\eta(\tau_0) \frac{\langle y \rangle^6}{y^\gamma} \int_{\tau_0}^{\tau} b^{\frac{\alpha}{2}}(\tau') d\tau' \right| \leq \frac{A^3}{16} b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau),$$

provided that  $K_0 \geq K_4, A \geq A_4, \tau_0 \geq \tau_4(K_0, A, \eta, \tilde{\eta})$  and  $\tau - \tau_0 \leq \frac{\ln A}{K_0}$ . Finally, (7.25) follows.

**The second case**  $\tau - \tau_0 \geq \frac{\ln A}{K_0}$

As we mentioned, this case will be divided into two sub-cases

$$\frac{1}{L_0} e^{\frac{\tau-\tau_0}{2}(1-\eta(\frac{2\ell}{\alpha}-1))} \leq b^{-\tilde{\eta}}(\tau) \text{ and } \frac{1}{L_0} e^{\frac{\tau-\tau_0}{2}(1-\eta(\frac{2\ell}{\alpha}-1))} > b^{-\tilde{\eta}}(\tau),$$

where  $L_0$  is large enough.

**The first subcase**  $\frac{1}{L_0} e^{\frac{\tau-\tau_0}{2}(1-\eta(\frac{2\ell}{\alpha}-1))} \leq b^{-\tilde{\eta}}(\tau)$

From (5.11) and  $\frac{1}{L_0} e^{\frac{\tau-\tau_0}{2}(1-\eta(\frac{2\ell}{\alpha}-1))} \leq b^{-\tilde{\eta}}(\tau)$ , we get

$$\tau < \frac{\tau_0 \left(1 - \eta \left(1 - \frac{2}{\alpha}\right)\right) + 2 \ln L_0}{1 - \left(\eta - 2\tilde{\eta} \left(1 + \frac{\tilde{\eta}}{10}\right)\right) \left(1 - \frac{2}{\alpha}\right)}, \quad (7.35)$$

which yields

$$\frac{\ln A}{K_0} \leq \tau - \tau_0 \leq \frac{2\tilde{\eta}\tau_0 \left(\frac{2}{\alpha} - 1\right) \left(1 + \frac{\tilde{\eta}}{10}\right) + 2 \ln L_0}{1 - \left(\eta - 2\tilde{\eta} \left(1 + \frac{\tilde{\eta}}{10}\right)\right) \left(\frac{2}{\alpha} - 1\right)}. \quad (7.36)$$

for  $L_0$  large enough. According to (7.36), we see that this the present sub-case can be handled similarly as the fist one, since  $\tau$  is not too far from  $\tau_0$ .

- The proof of (7.23): From (7.27), we have

$$\left| e^{(\tau-\tau_0)\mathcal{L}_\infty} \hat{\varepsilon}_-(\tau_0)(y, \tau) \right| \leq C A b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau) \frac{\langle y \rangle^4}{y^\gamma} \left[ e^{\frac{\alpha}{2}(\tau-\tau_0)} b^{-\frac{\alpha}{2} - \tilde{\eta}}(\tau_0) b^{\frac{\alpha}{2} + \eta}(\tau_0) \right].$$

The same process used for (7.28) yields

$$e^{\frac{\alpha}{2}(\tau-\tau_0)} b^{\frac{\alpha}{2} + \eta}(\tau_0) b^{-\frac{\alpha}{2} - \tilde{\eta}}(\tau) \leq e^{-c(\eta)\tau_0} e^{X(\tau-\tau_0)}, \text{ with } X = \left( \frac{\alpha}{2} + \left( \frac{2}{\alpha} - 1 \right) \left( \frac{\alpha}{2} + \tilde{\eta} \right) \left( 1 + \frac{\tilde{\eta}}{10} \right) \right).$$

From (7.36), we can prove that there exists  $c(\eta) > 0$  such that

$$\begin{aligned} & e^{-c(\eta)\tau_0} e^{X(\tau-\tau_0)}, \text{ with } X = \left( \frac{\alpha}{2} + \left( \frac{2}{\alpha} - 1 \right) \left( \frac{\alpha}{2} + \tilde{\eta} \right) \left( 1 + \frac{\tilde{\eta}}{10} \right) \right) \\ & \lesssim e^{-c(\eta)\tau_0 + X \left( \frac{2\tilde{\eta}\tau_0 \left( \frac{2}{\alpha} - 1 \right) \left( 1 + \frac{\tilde{\eta}}{10} \right) + 2 \ln L_0}{1 - \left( \eta - 2\tilde{\eta} \left( 1 + \frac{\tilde{\eta}}{10} \right) \right) \left( \frac{2}{\alpha} - 1 \right)} \right)} \leq 1, \end{aligned}$$

provided that  $\tilde{\eta} \leq \tilde{\eta}_4(\eta, L_0)$  and this gives (7.23).

- The proof of (7.24): we use (7.31) and (7.32) to get

$$\begin{aligned} & \left| \frac{y^\gamma}{\langle y \rangle^{2+2}} \int_{\tau_0}^\tau e^{(\tau-\tau')\mathcal{L}_\infty} \left[ \hat{B}(\hat{\varepsilon}_{\beta,+}, \hat{\varepsilon}_{\beta,-}) \right] (\tau') d\tau' \right| \\ & \leq C b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau) \left\{ \int_{\tau_0}^\tau \left[ e^{\frac{\alpha}{2}(\tau-\tau')} b^{\delta(\omega+4-2\gamma) - \frac{\eta}{4}(\omega+2)} (\tau') b^{-\frac{\alpha}{2} - \tilde{\eta}}(\tau) \right] d\tau' \right. \\ & \left. + \int_{\tau_0}^\tau e^{\frac{\alpha}{2}(\tau-\tau')} b^{-\frac{\alpha}{2} - \tilde{\eta}}(\tau) \left[ b^{\frac{\alpha}{2} + 4\eta}(\tau') + b^{\alpha + \delta(1-\gamma)}(\tau') b^{-\tilde{\eta}(2+6)}(\tau) + b^{\frac{3\alpha}{2} - 2\delta\gamma}(\tau') b^{-\tilde{\eta}(16)}(\tau) \right] d\tau' \right\}. \end{aligned}$$

From (7.28) and (7.36), we have

$$\begin{aligned} & \int_{\tau_0}^{\tau} \left[ e^{\frac{\alpha}{2}(\tau-\tau')} b^{\delta(\omega+4-2\gamma)-\frac{\eta}{4}(\omega+2)}(\tau') b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau) \right] d\tau' \\ & + \int_{\tau_0}^{\tau} e^{\frac{\alpha}{2}(\tau-\tau')} b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau) \left[ b^{\frac{\alpha}{2}+4\eta}(\tau') + b^{\alpha+\delta(1-\gamma)}(\tau') b^{-\tilde{\eta}(8)}(\tau) + b^{\frac{3\alpha}{2}-2\delta\gamma}(\tau') b^{-\tilde{\eta}(16)}(\tau) \right] d\tau' \\ & \leq C, \end{aligned}$$

provided that  $\tilde{\eta} \leq \tilde{\eta}_4(\eta, \delta, L_0)$ ,  $A \geq A_4$  and  $\tau_0 \geq \tau_4(A, \eta, \tilde{\eta}, L_0)$ . Then, we infer

$$\left| \frac{y^\gamma}{\langle y \rangle^4} \int_{\tau_0}^{\tau} e^{(\tau-\tau')\mathcal{L}_\infty} \left[ \hat{B}(\hat{\varepsilon}_{\beta,+}, \hat{\varepsilon}_{\beta,-}) \right] (\tau') d\tau' \right| \leq \frac{A^3}{16} b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau), \forall y \in [b^\eta(\tau), b^{-\tilde{\eta}}(\tau)],$$

which implies (7.24).

- Proof of (7.25): it is similar to the that of (7.24) in the first case.

$$\text{The second sub-case } \frac{1}{L_0} e^{\frac{\tau-\tau_0}{2}(1-\eta(\frac{2\ell}{\alpha}-1))} > b^{-\tilde{\eta}}(\tau)$$

We introduce  $R$  large, to be fixed later, and we decompose

$$[b^\eta(\tau), b^{-\tilde{\eta}}(\tau)] = [b^\eta(\tau), R] \cup [R, b^{-\tilde{\eta}}(\tau)].$$

Recall that for each  $f \in L_\rho^2(\mathbb{R}^+)$ , for each  $\nu > 0$ , there exists  $Y(\nu) > 0$  satisfying  $Y(\nu) \rightarrow +\infty$  as  $\nu \rightarrow +\infty$  and such that  $\forall y \in [Y^{-1}(\nu), Y(\nu)]$

$$e^{\nu\mathcal{L}_\infty} f(y) = \sum_{j=0}^{\infty} e^{(\frac{\alpha}{2}-j)\nu} \langle f, \phi_{j,\infty} \rangle_{L_\rho^2} \phi_{j,\infty}(y) \text{ pointwisely.} \quad (7.37)$$

Following the above remarks, the expression (7.21) and the fact that  $\hat{\varepsilon}_-$  is orthogonal to  $\phi_{0,\infty}$  and  $\phi_{1,\infty}$  we are led to

$$\begin{aligned} \hat{\varepsilon}_-(y, \tau) &= \left( e^{(\tau-\tau_0)\mathcal{L}_\infty}(\varepsilon_-(\tau_0)) \right)_- + \int_{\tau_0}^{\tau} \left( e^{(\tau-\tau')\mathcal{L}_\infty}(\hat{B}(\tau')) \right)_- d\tau' + \int_{\tau_0}^{\tau} \left( \left[ \frac{1}{2} - \beta(\tau') \right] \Lambda \hat{\varepsilon}_-(\tau') \right)_- d\tau' \\ &= \sum_{j=2}^{\infty} e^{(\frac{\alpha}{2}-j)(\tau-\tau_0)} \langle \hat{\varepsilon}_-(\tau_0), \phi_{j,\infty} \rangle_{L_\rho^2} \phi_{j,\infty}(y) \\ &+ \int_{\tau_0}^{\tau-L_0} \sum_{j=2}^{\infty} e^{(\frac{\alpha}{2}-j)(\tau-\tau')} \langle \hat{B}(\tau'), \phi_{j,\infty} \rangle_{L_\rho^2} \phi_{j,\infty}(y) d\tau' + \int_{\tau-L_0}^{\tau} \left( e^{(\tau-\tau')\mathcal{L}_\infty}(\hat{B}(\tau')) \right)_- d\tau' \\ &+ \int_{\tau_0}^{\tau-L_0} \sum_{j=2}^{\infty} e^{(\frac{\alpha}{2}-j)(\tau-\tau')} \left\langle \left( \left[ \frac{1}{2} - \beta(\tau') \right] \Lambda \hat{\varepsilon}_-(\tau') \right) d\tau', \phi_{j,\infty} \right\rangle_{L_\rho^2} \phi_{j,\infty}(y) d\tau' \\ &+ \int_{\tau-L_0}^{\tau} \left( \left[ \frac{1}{2} - \beta(\tau') \right] \Lambda \hat{\varepsilon}_-(\tau') \right)_- d\tau' \end{aligned} \quad (7.38)$$

on  $[Y^{-1}(\tau), Y(\tau)]$  ( $Y(\tau) \rightarrow +\infty$  as  $\tau \rightarrow +\infty$ ). Let us consider  $y \in [b^\eta(\tau), R]$ . We consider the initial data  $\varepsilon(\tau_0)$  defined in (5.14), we have

$$\left| \langle \hat{\varepsilon}_-(y, \tau_0), \phi_{j,\infty} \rangle_{L_\rho^2} \right| \leq C b^{\frac{\alpha}{2}+\delta}, \forall j \geq 2.$$

Note that  $\delta \gg \eta \gg \tilde{\eta}$  and we are in the case  $\tau - \tau_0 > \frac{\ln A}{K_0}$ , we have the estimate

$$\begin{aligned} & \left| \sum_{j=2}^{\infty} e^{(\frac{\alpha}{2}-j)(\tau-\tau_0)} \langle \hat{\varepsilon}_-(\tau_0), \phi_{j,\infty} \rangle_{L^2_\rho} \phi_{j,\infty}(y) \right| \\ & \leq C b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau) \frac{\langle y \rangle^4}{y^\gamma} \sum_{j=2}^{\infty} e^{(\frac{\alpha}{2}-j)(\tau-\tau_0)} b^{\frac{\alpha}{2}+\delta}(\tau_0) b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau) \alpha_j (1+y)^{2(j-1)} \\ & \leq C C b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau) \frac{\langle y \rangle^4}{y^\gamma} \sum_{j=2}^{\infty} j^{\frac{\omega}{4}} e^{(\frac{\alpha}{2}-j)(\tau-\tau_0)} b^{\frac{\alpha}{2}+\delta}(\tau_0) b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau) (1+y)^{2(j-1)}. \end{aligned}$$

Similarly to the technique given in (7.28), we have

$$\begin{aligned} e^{(\frac{\alpha}{2}-j)(\tau-\tau_0)} b^{\frac{\alpha}{2}+\delta}(\tau_0) b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau) & \lesssim C e^{(\frac{\alpha}{2}-j)(\tau-\tau_0)} e^{(\frac{2}{\alpha}-1)(\frac{\alpha}{2}+\tilde{\eta})(1+\tilde{\eta})(\tau-\tau_0)} \\ & \lesssim C e^{(\tau-\tau_0)(1-j+c(\tilde{\eta}))} \text{ with } c(\tilde{\eta}) \lesssim \tilde{\eta}, j \geq 2, \end{aligned}$$

which yields to

$$\begin{aligned} & \left| \sum_{j=2}^{\infty} e^{(\frac{\alpha}{2}-j)(\tau-\tau_0)} \langle \hat{\varepsilon}_-(\tau_0), \phi_{j,\infty} \rangle_{L^2_\rho} \phi_{j,\infty}(y) \right| \\ & \leq C b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau) \frac{\langle y \rangle^4}{y^\gamma} \sum_{j=2}^{\infty} j^{\frac{\omega}{4}} R^{2(j-1)} \left( A^{\frac{1}{K_0}} \right)^{1-j+c(\tilde{\eta})} \leq \frac{A^3}{32} b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau) \frac{\langle y \rangle^4}{y^\gamma}, \end{aligned} \quad (7.39)$$

provided that  $K_0 \geq K_4$ ,  $A \geq A_4(R, K_0)$ , and  $\tilde{\eta} \leq \tilde{\eta}_4(R, K_0, A, \eta, \tilde{\eta}, L_0)$ .

Next, observe that  $\tau - (\tau - L_0) = L_0 \leq \frac{\ln A}{K_0}$  if  $A \geq A_4(L_0)$ , so we go back to the first case. Indeed, we argue similarly as in (7.31) and (7.33) to get

$$\begin{aligned} \left| \int_{\tau-L_0}^{\tau} e^{(\tau-\tau')\mathcal{L}_\infty}(\hat{B})(\tau') d\tau' \right| & \leq \frac{\langle y \rangle^4}{y^\gamma} \int_{\tau-L_0}^{\tau} e^{\frac{\alpha}{2}(\tau-\tau')} b^{\frac{\alpha}{2}+4\eta}(\tau') d\tau' \\ & \leq C b^{\frac{\alpha}{2}+3\eta}(\tau) \frac{\langle y \rangle^4}{y^\gamma}, \end{aligned}$$

provided that  $s_0 \geq s_4(\eta, L_0)$ . This yields

$$\left| \int_{\tau-L_0}^{\tau} \left( e^{(\tau-\tau')\mathcal{L}_\infty}(\hat{B})(\tau') \right)_- d\tau' \right| \leq C b^{\frac{\alpha}{2}+3\eta}(\tau) \frac{\langle y \rangle^4}{y^\gamma}.$$

For the integral  $\int_{\tau_0}^{\tau-L_0} \sum_{j=2}^{\infty} e^{(\frac{\alpha}{2}-j)(\tau-\tau')} \langle \hat{B}(\tau'), \phi_{j,\infty} \rangle_{L^2_\rho} \phi_{j,\infty}(y) d\tau'$ , we deduce from Lemma A.1 that

$$\left| \langle \hat{B}(\tau'), \phi_{j,\infty} \rangle_{L^2_\rho} \right| \lesssim b^{\frac{\alpha}{2}+4\eta}(\tau').$$

Then, for all  $y \in [b^\eta(\tau), R]$ , we have

$$\begin{aligned} & \left| \int_{\tau_0}^{\tau-L_0} \left( \sum_{j=2}^{\infty} e^{(\frac{\alpha}{2}-j)(\tau-\tau')} \langle \hat{B}(\tau'), \phi_{j,\infty} \rangle_{L^2_\rho} \phi_{j,\infty} \right) d\tau' \right| \\ & \lesssim \frac{\langle y \rangle^4 b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau)}{y^\gamma} \sum_{j=2}^{\infty} |\alpha_j| R^{2(j-1)} \int_{\tau_0}^{\tau-L_0} e^{(\frac{\alpha}{2}-j)(\tau-\tau')} b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau) b^{\frac{\alpha}{2}+4\eta}(\tau') d\tau'. \end{aligned}$$

We repeat the techniques given in (7.28) and (7.29) to obtain

$$\begin{aligned} b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau)e^{(\frac{\alpha}{2}-j)(\tau-\tau')}b^{\frac{\alpha}{2}+\eta}(\tau') &\lesssim b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau)b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau')e^{(\frac{\alpha}{2}-j)(\tau-\tau')} \\ &\lesssim e^{(1-j+c(\tilde{\eta}))(\tau-\tau')}, \text{ with } c(\tilde{\eta}) \lesssim \tilde{\eta}. \end{aligned}$$

It follows that for all  $j \geq 2$ , we have

$$\int_{\tau_0}^{\tau-L_0} b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau)e^{(\frac{\alpha}{2}-j)(\tau-\tau')}b^{\frac{\alpha}{2}+\eta}(\tau')d\tau' \lesssim \exp([1-j+c(\tilde{\eta})](L_0)), \text{ and } 1-j+c(\tilde{\eta}) < 0.$$

By taking  $L_0 \gg R$ , we get

$$\begin{aligned} &\sum_{j=2}^{\infty} |\alpha_j| R^{2(j-1)} \int_{\tau_0}^{\tau-L_0} e^{(\frac{\alpha}{2}-j)(\tau-\tau')} b^{-\frac{\alpha}{2}-\tilde{\eta}}(\tau) b^{\frac{\alpha}{2}+4\eta}(\tau') d\tau' \\ &\lesssim \sum_{j=2}^{\infty} j^{\frac{\omega}{4}} R^{2(j-1)} \exp([1-j+c(\tilde{\eta})](L_0)) \lesssim 1, \end{aligned}$$

provided that  $L_0 \gg R$  and  $\tau_0 \geq \tau_4(A, L_0, R, \delta, \eta, \tilde{\eta})$ . Then

$$\left| \int_{\tau_0}^{\tau-L_0} \sum_{j=2}^{\infty} e^{(\frac{\alpha}{2}-j)(\tau-\tau')} \left\langle \hat{B}(\tau'), \phi_{j,\infty} \right\rangle_{L_p^2} \phi_{j,\infty}(y) d\tau' \right| \leq C b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau) \frac{\langle y \rangle^4}{y^\gamma}, y \in [b^{\frac{\eta}{4}}, R].$$

Recall that

$$\left| \beta(\tau) - \frac{1}{2} \right| \leq C A I^\eta(\tau_0),$$

so by repeating the above arguments to handle the remaining terms in (7.38), the following result holds

$$\begin{aligned} &\left| \int_{\tau_0}^{\tau-L_0} \sum_{j=2}^{\infty} e^{(\frac{\alpha}{2}-j)(\tau-\tau')} \left\langle \left( \left[ \frac{1}{2} - \beta(\tau') \right] \Lambda \hat{\varepsilon}_-(\tau') \right) d\tau', \phi_{j,\infty} \right\rangle_{L_p^2} \phi_{j,\infty}(y) d\tau' \right| \\ &+ \left| \int_{\tau-L_0}^{\tau} \left( \left[ \frac{1}{2} - \beta(\tau') \right] \Lambda \hat{\varepsilon}_-(\tau') \right)_- d\tau' \right| \leq C b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau) \frac{\langle y \rangle^4}{y^\gamma}. \end{aligned}$$

We conclude that

$$\left| \int_{\tau_0}^{\tau} e^{(\tau-\tau')\mathcal{L}_\infty} \left[ \frac{1}{2} - \beta(\tau') \right] \Lambda_y(\hat{\varepsilon}_{\beta,-})(\tau') d\tau' \right| \leq \frac{A^3}{16} b^{\frac{\alpha}{2}+\tilde{\eta}}(\tau) \langle y \rangle^4 y^{-\gamma}, \forall y \in [b^{\frac{\eta}{4}}, R]. \quad (7.40)$$

- For the case  $y \in [R, b^{-\tilde{\eta}}(\tau)]$ . First, we observe that once  $y \leq b^{-\tilde{\eta}}(\tau) \leq \frac{1}{L_0} e^{\frac{\tau-\tau_0}{2}(1-\eta(\frac{2}{\alpha}-1))}$ , then, there exists  $\bar{\tau} \in [\tau_0, \tau-1]$  such that  $y \in \left[ \frac{1}{2L_0} e^{\frac{\tau-\bar{\tau}}{2}(1-\eta(\frac{2}{\alpha}-1))}, \frac{1}{L_0} e^{\frac{\tau-\bar{\tau}}{2}(1-\eta(\frac{2}{\alpha}-1))} \right]$ . Since  $y \geq R$  we have  $\tau - \bar{\tau} \geq C(R) \rightarrow +\infty$  as  $R \rightarrow +\infty$ . Now, write the integral equation at the initial data  $\bar{\tau}$

$$\hat{\varepsilon}_-(\tau) = e^{(\tau-\bar{\tau})\mathcal{L}_\infty} \hat{\varepsilon}_-(\bar{\tau}) + \int_{\bar{\tau}}^{\tau} e^{(\tau-\tau')\mathcal{L}_\infty} \left[ \hat{B}(\tau') + \left( \frac{1}{2} - \beta(\tau') \right) \Lambda_y \hat{\varepsilon}_- \right](\tau') d\tau'. \quad (7.41)$$

Note that for all  $j \geq 2$ , we have

$$\langle \hat{\varepsilon}_-, \phi_{j,\infty} \rangle_{L_p^2} = \langle \varepsilon_+, \phi_{j,\infty} \rangle_{L_p^2} + \langle \varepsilon_-, \phi_{j,\infty} \rangle.$$

Using the fact that  $\varepsilon_+$  is orthogonal to  $\phi_{j,b,\beta}$ , estimate (4.7),  $\phi_{j,\infty,\beta}$ 's definition given in Proposition 4.1 and  $\phi_{i,\infty} = \phi_{j,\infty,\frac{1}{2}}$ , we derive that

$$\left| \langle \varepsilon_+, \phi_{j,\infty} \rangle_{L_p^2} \right| \leq C b^{\frac{\alpha}{2}+\eta}(\bar{\tau}), \forall j \geq 2.$$

Moreover, using the definition of  $\rho_\beta$  given in (2.25) and  $\rho = \rho_{\frac{1}{2}}$ , we apply Cauchy Schwarz inequality and estimate (7.1) to derive

$$\begin{aligned} \left| \langle \varepsilon_-, \phi_{j,\infty} \rangle_{L_\rho^2} \right| &= \left| \int_{\mathbb{R}_+} \varepsilon_- \phi_{j,\infty} \rho dy \right| \leq C \int_{\mathbb{R}_+} |\varepsilon_-| \sqrt{\rho_\beta} |\phi_{j,\infty}| \frac{\rho}{\sqrt{\rho_\beta}} dy \\ &\leq C \|\varepsilon_-\|_{L_\rho^2} \sqrt{\int_{\mathbb{R}_+} |\phi_{j,\infty}|^2 \frac{\rho^2}{\rho_\beta} dy} \leq C b^{\frac{\alpha}{2} + \eta}. \end{aligned}$$

Hence, we have

$$\left| \langle \hat{\varepsilon}_-(\bar{\tau}), \phi_{j,\infty} \rangle_{L_\rho^2} \right| \leq C b^{\frac{\alpha}{2} + \eta}(\bar{\tau}).$$

Now, we use formula (7.37) to get

$$\begin{aligned} \left| e^{(\tau - \bar{\tau})\mathcal{L}_\infty} \hat{\varepsilon}_-(\bar{\tau})(y, \tau) \right| &\lesssim \sum_{j=2}^{\infty} e^{(\frac{\alpha}{2} - j)(\tau - \bar{\tau})} \left| \langle \hat{\varepsilon}_-(\bar{\tau}), \phi_{j,\infty} \rangle_{L_\rho^2} \right| |\phi_{j,\infty}(y)| \\ &\lesssim \sum_{j=2}^{\infty} 4^j j! e^{(\frac{\alpha}{2} - j)(\tau - \bar{\tau})} b^{\frac{\alpha}{2} + \eta}(\bar{\tau}) |\phi_{j,\infty}(y)| \\ &\lesssim \langle y \rangle^4 y^{-\gamma} b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau) \sum_{j=2}^{\infty} |\beta_j| e^{(\frac{\alpha}{2} - j)(\tau - \bar{\tau})} b^{-\frac{\alpha}{2} - \tilde{\eta}}(\tau) b^{\frac{\alpha}{2} + \eta}(\bar{\tau}) y^{2(j-1)}. \end{aligned}$$

Using (5.11), we have for all  $j \geq 2$

$$e^{(\frac{\alpha}{2} - j)(\tau - \bar{\tau})} b^{\frac{\alpha}{2} + \eta}(\bar{\tau}) b^{-\frac{\alpha}{2} - \tilde{\eta}}(\tau) \lesssim e^{(\frac{\alpha}{2} - j)(\tau - \bar{\tau})} e^{(1 - \frac{2}{\alpha})[(\frac{\alpha}{2} + \eta)(1 - \frac{\tilde{\eta}}{10})\bar{\tau} - (\frac{\alpha}{2} + \tilde{\eta})(1 + \frac{\tilde{\eta}}{10})\tau]}.$$

In addition, since  $\tilde{\eta} \ll \eta$ , and  $(1 - \frac{2}{\alpha}) < 0$ , it follows the following

$$\begin{aligned} &\left(1 - \frac{2}{\alpha}\right) \left[ \left(\frac{\alpha}{2} + \eta\right) \left(1 - \frac{\tilde{\eta}}{10}\right) \bar{\tau} - \left(\frac{\alpha}{2} + \tilde{\eta}\right) \left(1 + \frac{\tilde{\eta}}{10}\right) \tau \right] \\ &\leq \left(1 - \frac{2}{\alpha}\right) \left[ \left(\frac{\alpha}{2} + \frac{\eta}{2}\right) \left(1 + \frac{\tilde{\eta}}{10}\right) \bar{\tau} - \left(\frac{\alpha}{2} + \tilde{\eta}\right) \left(1 + \frac{\tilde{\eta}}{10}\right) \tau \right] \\ &= \left(1 - \frac{2}{\alpha}\right) \left(1 + \frac{\tilde{\eta}}{10}\right) \left[ \frac{\alpha}{2}(\bar{\tau} - \tau) + \frac{\eta}{2}\bar{\tau} - \tilde{\eta}\tau \right] \\ &\leq \left(1 - \frac{2}{\alpha}\right) \left(1 + \frac{\tilde{\eta}}{10}\right) \left[ \frac{\alpha}{2}(\bar{\tau} - \tau) + \frac{\eta}{2}(\bar{\tau} - \tau) \right], \end{aligned}$$

and

$$\begin{aligned} &\left(\frac{\alpha}{2} - j\right) (\tau - \bar{\tau}) + \left(1 - \frac{2}{\alpha}\right) \left(1 + \frac{\tilde{\eta}}{10}\right) \left[ \frac{\alpha}{2}(\bar{\tau} - \tau) + \frac{\eta}{2}(\bar{\tau} - \tau) \right] \\ &\leq \left[ 1 - j + \frac{\tilde{\eta}}{10} \left(\frac{2}{\alpha} - 1\right) \frac{\alpha}{2} + \frac{\eta}{2} \left(\frac{2}{\alpha} - 1\right) \left(1 + \frac{\tilde{\eta}}{10}\right) \right] (\tau - \bar{\tau}) \\ &\leq \left[ 1 - j + \eta \left(\frac{2}{\alpha} - 1\right) \right] (\tau - \bar{\tau}), \end{aligned}$$

and  $|\beta_j| \leq C \frac{j^{-\frac{\omega}{4}}}{4^j j!}$ . Hence, we obtain

$$\begin{aligned} \left| e^{(\tau - \bar{\tau})\mathcal{L}_\infty} \hat{\varepsilon}_-(\bar{\tau})(y, \tau) \right| &\lesssim A b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau) \sum_{j=2}^{\infty} \frac{j^{-\frac{\omega}{4}}}{4^j j!} \left( y e^{-\frac{\tau - \bar{\tau}}{2}(1 - \eta(\frac{2\ell}{\alpha} - 1))} \right)^{2(j-1)} \\ &\lesssim A b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau) \sum_{j=2}^{\infty} \frac{j^{-\frac{\omega}{4}}}{4^j j!} (L_0^{-1})^{2(j-1)} \leq \frac{A^3}{16} b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau), \end{aligned}$$

when  $L_0$  is large and  $A \geq A_4$ .

From the restriction  $\frac{1}{2L_0} e^{\frac{\tau-\bar{\tau}}{2}(1-\eta(\frac{2\ell}{\alpha}-1))} \leq y \leq b^{-\bar{\eta}}(\tau)$ , it follows

$$\tau < \frac{\bar{\tau} \left(1 - \eta \left(1 - \frac{2\ell}{\alpha}\right)\right) + 2 \ln L_0}{1 - \left(\eta - 2\bar{\eta} \left(1 + \frac{\bar{\eta}}{10}\right)\right) \left(1 - \frac{2\ell}{\alpha}\right)},$$

which yields

$$\tau - \bar{\tau} \leq \frac{2\bar{\eta}\bar{\tau} \left(\frac{2\ell}{\alpha} - 1\right) \left(1 + \frac{\bar{\eta}}{10}\right) + 2 \ln L_0}{1 - \left(\eta - 2\bar{\eta} \left(1 + \frac{\bar{\eta}}{10}\right)\right) \left(\frac{2\ell}{\alpha} - 1\right)}. \quad (7.42)$$

Hence, it turns out that the second case is obtained by replacing  $\tau_0$  by  $\bar{\tau}$

$$\left| \int_{\bar{\tau}}^{\tau} e^{(\tau-\tau')\mathcal{L}_\infty} \left[ \hat{B}(\hat{\varepsilon}_{\beta,+} + \hat{\varepsilon}_{\beta,-}) \right] (\tau') d\tau' \right| \leq \frac{A^3}{16} b^{\frac{\alpha}{2} + \bar{\eta}}(\tau) \langle y \rangle^{2\ell+2} y^{-\gamma}, \quad (7.43)$$

$$\left| \int_{\bar{\tau}}^{\tau} e^{(\tau-\tau')\mathcal{L}_\infty} \left( \frac{1}{2} - \beta(\tau') \right) \Lambda_y \hat{\varepsilon}_{\beta,-}(\tau') d\tau' \right| \leq \frac{A^3}{16} b^{\frac{\alpha}{2} + \bar{\eta}}(\tau) \langle y \rangle^{2\ell+2} y^{-\gamma}. \quad (7.44)$$

At final, by adding all the terms, we get

$$|\varepsilon_{\beta,-}(\tau)| \leq \frac{A^3}{2} b^{\frac{\alpha}{2} + \bar{\eta}} \frac{\langle y \rangle^{2\ell+2}}{y^\gamma}, \forall y \in [R, b^{-\bar{\eta}}(\tau)],$$

which concludes the third case.  $\square$

The final step is to focus on the *a priori estimates* on the exterior part  $\varepsilon_e$ . More precisely, we have the following Lemma

**Lemma 7.3** (A priori estimates on the exterior part). *Let us consider  $\varepsilon$  to satisfy equation (2.17) with initial data given in (5.16), and  $\varepsilon(\tau) \in V[A, \eta, \bar{\eta}](\tau)$  for all  $\tau \in [\tau_0, \tau_1]$ , for some  $\tau_1 > \tau_0$ . Then, we have the following estimate*

$$\| |y| \varepsilon_e(\cdot, \tau) \|_{L^\infty} \leq \frac{A^4}{2} b^{\frac{\alpha}{2} + (\gamma-4)\bar{\eta}}(\tau).$$

*Proof.* To get the conclusion of the Lemma, we consider the natural  $(d+2)$ -dimensional extension as follows

$$\varepsilon_{ext}(z, \tau) = \varepsilon(y, \tau), |z| = y \text{ and } z \in \mathbb{R}^{d+2}, \quad (7.45)$$

and we introduce

$$\varepsilon_{ext,e}(z, \tau) = |z| \left( 1 - \chi_0 \left( \frac{8}{3} |z| b^{\bar{\eta}}(\tau) \right) \right) \varepsilon_{ext}(z, \tau),$$

where  $\chi_0$  was defined in (5.10) and  $\gamma$  defined was in (B). Note that  $u$ 's extension defined in (7.45) is  $C^2(\mathbb{R}^{d+2})$ , thanks to the parabolic regularity of the semi-group  $e^{t\Delta_{d+2}}$  and so is  $\varepsilon_{ext}$  and we derive from (2.17) that  $\varepsilon_{ext}$  satisfies

$$\begin{aligned} \partial_\tau \varepsilon_{ext} &= \Delta \varepsilon_{ext} - \beta(\tau) y \cdot \nabla \varepsilon_{ext} - 2\beta(\tau) \varepsilon_{ext} \\ &- 3(d-2) [2Q_b(|z|) + |z|^2 Q_b^2(|z|)] \varepsilon_{ext} + B(\varepsilon_{ext}, |z|) + \Phi(|z|, \tau), \end{aligned}$$

where  $\Lambda_z$  is similarly defined as in (2.14),  $B$  and  $\Phi$  were defined in (2.19) and (2.20), respectively. Now, we introduce

$$\varepsilon_{ext,e}(z, \tau) = |z| (1 - \chi_{\bar{\eta}}) \varepsilon_{ext}, \text{ and } \chi_{\bar{\eta}}(z, \tau) = \chi_0 \left( \frac{8}{3} |z| b^{\bar{\eta}}(\tau) \right), \quad (7.46)$$



where  $\chi_0$  was defined in (5.10) and we have the following facts

$$\text{supp}(1 - \chi_{\tilde{\eta}}) \subset \left\{ z \in \mathbb{R}^n \text{ such that } |z| \geq \frac{3}{8}b^{-\tilde{\eta}}(\tau) \right\}, \quad (7.47)$$

and  $\varepsilon_{ext,e} = \varepsilon_{ext}$  for all  $|z| \geq \frac{3}{4}b^{-\tilde{\eta}}(\tau)$ . We here mention that the conclusion of the Lemma immediately follows from

$$\|\varepsilon_{ext,e}(\cdot, \tau)\|_{L^\infty[b^{-\tilde{\eta}}(\tau), +\infty)} \leq \frac{A^4}{2} b^{\frac{\alpha}{2} + (\gamma-4)\tilde{\eta}}(\tau), \forall \tau \in [\tau_0, \tau_1].$$

By using  $\varepsilon_{ext}$ 's equation above, we that  $\varepsilon_{ext,e}$  exactly solves

$$\partial_\tau \varepsilon_{ext,e} = \mathcal{L}_\beta(\varepsilon_{ext,e}) + \mathcal{C}(\varepsilon_{ext}) + \mathcal{N}(\varepsilon_{ext}), \quad (7.48)$$

where  $\mathcal{L}_\beta$  is defined by

$$\mathcal{L}_\beta = \Delta - \beta(\tau)z \cdot \nabla - \beta(\tau)Id \quad (7.49)$$

and the terms  $\mathcal{C}(\varepsilon_{ext})$  and  $\mathcal{N}(\varepsilon_{ext})$  are respectively defined by

$$\begin{aligned} \mathcal{C}(\varepsilon_{ext}) &= -2\text{div}(\varepsilon_{ext}\nabla(|z|(1 - \chi_{\tilde{\eta}}))) - 3(d-2) [2Q_b(|z|) + |z|^2 Q_b^2(|z|)] \varepsilon_{ext,e} \\ &+ \varepsilon_{ext} [\partial_\tau(1 - \chi_{\tilde{\eta}})|z| + \Delta(|z|(1 - \chi_{\tilde{\eta}})) + \beta(\tau)z \cdot \nabla(1 - \chi_{\tilde{\eta}})|z|], \end{aligned}$$

and

$$\mathcal{N}(\varepsilon_{ext}) = |z|(1 - \chi_{\tilde{\eta}}) (B(\varepsilon_{ext}) + \Phi(\cdot, \tau)).$$

- *The semi-group of  $\mathcal{L}_\beta$* : Let us define  $\mathcal{K}_\beta(\tau', \tau)$ ,  $\tau > \tau' \geq \tau_0$ ; the semi-group associated to  $\mathcal{L}_\beta$  with

$$\mathcal{K}_\beta(\tau, \tau')f = \int_{\mathbb{R}^{d+2}} \mathcal{K}_\beta(\tau, \tau', z, z')f(z')dz'$$

and the Kernel  $\mathcal{K}_\beta(\tau, \tau', z, z')$

$$\mathcal{K}_\beta(\tau, \tau', z, z') = \frac{\zeta(\tau, \tau')}{[4\pi \int_{\tau'}^\tau \zeta^2(\tilde{\tau}, \tau')d\tilde{\tau}]^{\frac{d+2}{2}}} \exp\left(-\frac{(z' - z\zeta(\tau, \tau'))^2}{4 \int_{\tau'}^\tau \zeta^2(\tilde{\tau}, \tau')d\tilde{\tau}}\right),$$

$$\zeta(\tau, \tau') = e^{-\int_{\tau'}^\tau \beta(\tilde{\tau})d\tilde{\tau}}, \tau > \tau'.$$

In particular, when  $\beta \equiv \frac{1}{2}$ , our situation is the same as the operator considered in [26, Lemma A.1] since

$$\zeta(\tau, \tau') = e^{-\frac{\tau-\tau'}{2}}, \text{ and } \int_{\tau'}^\tau \zeta^2(\tilde{\tau}, \tau')d\tilde{\tau} = 1 - e^{-(\tau-\tau')}.$$

Since our operator has the same structure as the one in [26, Lemma A.1], we can apply the arguments there to get

$$\|\mathcal{K}_\beta(\tau', \tau)(\varphi)\|_{L^\infty} \leq \zeta(\tau, \tau')\|\varphi\|_{L^\infty} \leq e^{-\frac{(\tau-\tau')}{4}}\|\varphi\|_{L^\infty}, \quad (7.50)$$

$$\|\mathcal{K}_\beta(\tau, \tau')\text{div}(\varphi)\|_{L^\infty} \leq C \frac{\zeta(\tau, \tau')}{\sqrt{\int_{\tau'}^\tau \zeta^2(\tilde{\tau}, \tau')d\tilde{\tau}}}\|\varphi\|_{L^\infty} \leq C \frac{e^{-\frac{\tau-\tau'}{4}}}{\sqrt{\tau-\tau'}}, \quad (7.51)$$

since (5.4) holds for all  $\tau \in [\tau_0, \tau_1]$ . By Duhamel's formula, we now write equation (7.48) as follows

$$\varepsilon_{ext,e}(\tau) = \mathcal{K}_\beta(\tau, \tau')\varepsilon_{ext,e}(\tau') + \int_{\tau'}^\tau \mathcal{K}_\beta(\tau, \tilde{\tau}) [\mathcal{C}(\varepsilon_{ext}) + \mathcal{N}(\varepsilon_{ext})](\tilde{\tau})d\tilde{\tau}. \quad (7.52)$$

We now aim to estimate the terms involving  $\mathcal{N}(\varepsilon_{ext}(\tau))$  and  $\mathcal{C}(\varepsilon_{ext})$ .

- For  $\mathcal{N}$ , since (7.47) holds, we only consider  $|z| \geq \frac{3}{8}b^{-\tilde{\eta}}(\tau)$  that will be divided into two small cases  $|z| \in [\frac{3}{8}b^{-\tilde{\eta}}(\tau), b^{-\tilde{\eta}}(\tau)]$  and  $|z| \geq b^{-\tilde{\eta}}(\tau)$ . Since  $\varepsilon(\tau) \in V_1[A, \eta, \tilde{\eta}](\tau), \forall \tau \in [\tau_0, \tau_1]$ , we get the following

$$\|z|\varepsilon_{ext}(z, \tau)\| \leq A^3 b^{\frac{\alpha}{2} + \gamma \tilde{\eta}} \langle |z| \rangle^4 \leq CA^3 b^{\frac{\alpha}{2} + (\gamma-4)\tilde{\eta}}(\tau), \forall |z| \in \left[ \frac{3}{8}b^{-\tilde{\eta}}(\tau), b^{-\tilde{\eta}}(\tau) \right],$$

and

$$\|z|\varepsilon_{ext}(z, \tau)\| \leq A^4 b^{\frac{\alpha}{2} + (\gamma-4)\tilde{\eta}}(\tau), \forall |z| \geq b^{-\tilde{\eta}}(\tau)$$

which yields

$$\|\mathcal{N}(\varepsilon_{ext})\|_{L^\infty} \leq CA^3 b^{\frac{\alpha}{2} + 10\tilde{\eta}}(\tau), \forall \tau \in [\tau_0, \tau_1],$$

provided that  $\tilde{\eta} \leq \tilde{\eta}_5(\alpha), \tau_0 \geq \tau_5(A, \tilde{\eta})$ . Using (7.50), we deduce

$$\|\mathcal{K}_\beta(\tau, \tilde{\tau})\mathcal{N}(\tilde{\tau})\|_{L^\infty} \leq CA^3 e^{-\frac{\tau-\tilde{\tau}}{4}} b^{\frac{\alpha}{2} + 10\tilde{\eta}}(\tilde{\tau}).$$

- For  $\mathcal{C}(\varepsilon_{ext})$ : From (7.51), we have

$$\begin{aligned} \|\mathcal{K}_\beta(\tau, \tilde{\tau})\operatorname{div}(\varepsilon_{ext}\nabla[(1-\chi_{\tilde{\eta}})|z])\|_{L^\infty} &\leq C \frac{e^{-\frac{\tau-\tilde{\tau}}{4}}}{\sqrt{\tau-\tilde{\tau}}} \|(\varepsilon_{ext}\nabla[(1-\chi_{\tilde{\eta}})|z])\|_{L^\infty}(\tilde{\tau}) \\ &\leq CA^3 \frac{e^{-\frac{\tau-\tilde{\tau}}{4}}}{\sqrt{\tau-\tilde{\tau}}} b^{\frac{\alpha}{2} + (\gamma-4)\tilde{\eta}}(\tilde{\tau}). \end{aligned}$$

Similarly, we have

$$\|\mathcal{K}_\beta(\tau, \tilde{\tau})[3(d-2)(2Q_b + |z|^2 Q_b^2)\varepsilon_{ext,e}]\|_{L^\infty} \leq CA^3 e^{-\frac{\tau-\tilde{\tau}}{4}} b^{\frac{\alpha}{2} + (\gamma-4)\tilde{\eta}}(\tilde{\tau}).$$

Besides that, we derive from (7.46) that

$$\|\varepsilon_{ext}(\tilde{\tau})\partial_\tau(1-\chi_{\tilde{\eta}})|z|\|_{L^\infty} \leq C\tilde{\eta}A^4 b^{\frac{\alpha}{2} + (\gamma-4)\tilde{\eta}}(\tilde{\eta}) \leq CA^3 b^{\frac{\alpha}{2} + (\gamma-4)\tilde{\eta}}(\tilde{\tau}).$$

Hence, we get

$$\|\mathcal{K}_\beta(\tau, \tilde{\tau})\varepsilon_{ext}(\tilde{\tau})\partial_\tau(1-\chi_{\tilde{\eta}})|z|\|_{L^\infty} \leq CA^3 e^{-\frac{\tau-\tilde{\tau}}{4}} b^{\frac{\alpha}{2} + (\gamma-4)\tilde{\eta}}(\tilde{\tau}). \quad (7.53)$$

By the same technique, we can establish the following

$$\|\mathcal{K}_\beta(\tau, \tilde{\tau})\varepsilon_{ext}(\tilde{\tau})[\Delta(|z|(1-\chi_{\tilde{\eta}})) + \beta(\tau)z \cdot \nabla(1-\chi_{\tilde{\eta}})|z]|\|_{L^\infty} \leq CA^3 e^{-\frac{\tau-\tilde{\tau}}{4}} b^{\frac{\alpha}{2} + (\gamma-4)\tilde{\eta}}(\tilde{\tau}).$$

Taking  $L^\infty$ -estimate on both sides of (7.52), we get

$$\begin{aligned} \|\varepsilon_{ext,e}(\tau)\|_{L^\infty} &\leq e^{-\frac{\tau-\tau'}{4}} \|\varepsilon_{ext,e}(\tau')\|_{L^\infty} + CA^3 \int_{\tau'}^{\tau} e^{-\frac{\tau-\tilde{\tau}}{4}} b^{\frac{\alpha}{2} + (\gamma-4)\tilde{\eta}}(\tilde{\tau}) \left[ \frac{1}{\sqrt{\tau-\tilde{\tau}}} + 1 \right] d\tilde{\tau} \\ &\leq e^{-\frac{\tau-\tau'}{4}} \|\varepsilon_{ext,e}(\tau')\|_{L^\infty} + CA^3 b^{\frac{\alpha}{2} + (\gamma-4)\tilde{\eta}}(\tau') \left( \sqrt{\tau-\tau'} + (\tau-\tau') \right). \end{aligned}$$

We now apply the technique used in [26, Proposition 4.5]. Our the choice of the initial data in (5.14) allows us to improve the bound on  $\varepsilon_{ext,e}$  by

$$\|\varepsilon_{ext,e}(\tau)\|_{L^\infty} \leq \frac{A^4}{2} b^{\frac{\alpha}{2} + (\gamma-4)\tilde{\eta}}(\tau), \forall \tau \in [\tau_0, \tau_1],$$

provided that  $A \geq A_5$  and  $\tilde{\eta} \leq \tilde{\eta}_5(A)$  and  $\tau_0 \geq \tau_5(A, \tilde{\eta})$ . Finally, the conclusion of the lemma follows.  $\square$

## 8. The existence of unstable blowup solutions

In this Section, we aim to give a sketch of the proof to the existence of unstable blowup solutions to equation (1.3) with blowup speeds

$$\lambda_\ell(t) = C(u_0)(T-t)^{\frac{2\ell}{\alpha}} \text{ as } t \rightarrow T.$$

The general strategy of the proof is the same as in the stable setting, except to some technical modifications. For that reason, we only give main changes which lead to the unstable existence. Let us consider the similarity variable (2.11) with  $\mu(\tau) = T-t$  and  $\tau = -\ln(T-t)$  then  $w$  satisfies (2.12) with  $\beta \equiv \frac{1}{2}$ . We linearize around  $Q_{b(\tau)}$  given in (2.15) by  $\varepsilon = w - Q_{b(\tau)}$  and it holds that  $\varepsilon$  satisfies (2.17). Note that all terms in the equation remain the same with  $\beta \equiv \frac{1}{2}$ . In particular, the spectral analysis of  $\mathcal{L}_b$  is valid without the appearance of  $\beta$ . Now, we consider the composition (7.6) with  $\ell \geq 2$ .

### 8.1. Shrinking set for $\ell \geq 2$

In this part, we modify a little bit the set in Definition 5.1 to be compatible with the new setting

**Definition 8.1** (Shrinking set). Let  $A, \eta, \tilde{\eta}$  be positive constants satisfying  $A \gg 1$  and  $1 \gg \eta \gg \tilde{\eta}$ , we define  $V_\ell[A, \eta, \tilde{\eta}](\tau)$  as the set of all  $(\varepsilon, b) \in L^\infty \times \mathbb{R}$  satisfying:

(i) The dominating mode  $\varepsilon_\ell$  and  $b$  satisfy

$$\left| \varepsilon_\ell + \frac{2}{\alpha} m_0 b^{\frac{\alpha}{2}} \right| \leq A b^{\frac{\alpha}{2} + \eta}, \quad (8.1)$$

and

$$\frac{1}{2} \leq b I_\ell^{-1}(\tau) \leq 2, \quad (8.2)$$

where

$$I_\ell(\tau) = e^{(\frac{2\ell}{\alpha}-1)\tau}. \quad (8.3)$$

In addition the others modes  $\varepsilon_j \in j \in \{1, \dots, \ell-1\}$  satisfy

$$|\varepsilon_j| \leq A b^{\frac{\alpha}{2} + \eta}. \quad (8.4)$$

(iii)  $L^2_\rho$ -decay: The part  $\varepsilon_-$  satisfies the following:

$$\|\varepsilon_-(\cdot)\|_{L^2_\rho} \leq A^2 b^{\frac{\alpha}{2} + \eta}(\tau).$$

(iv) The remainders  $\varepsilon_-$  given in (7.6), and  $\varepsilon_e$  satisfy

$$\begin{aligned} \left\| y^\gamma \frac{\varepsilon_-(\cdot, \tau)}{\langle y \rangle^{2\ell+2}} \right\|_{L^\infty[0, b^{-\tilde{\eta}}(\tau)]} &\leq A^3 b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau), \\ \| |y| \varepsilon_e(\cdot, \tau) \|_{L^\infty} &\leq A^4 b^{\frac{\alpha}{2} + (\gamma - (2\ell+2))\tilde{\eta}}(\tau), \end{aligned}$$

where  $\varepsilon_e$  defined as in (5.9).

### 8.2. Preparing initial data

In this part, we aim to construct a class of initial data corresponding to the Shrinking set  $V_\ell[A, \eta, \tilde{\eta}]$ . Let us consider  $A \geq 1$ , and  $0 < \tilde{\eta} \ll \eta \ll \delta \leq 1$ ,  $\alpha, m_0$  and  $c_{\ell,0}$  defined as in (2.27), (6.15) and (2.28), respectively.

$$\begin{aligned} \psi(\ell, \tau_0) &= \chi \left( y b^{\frac{\delta}{2}}(\tau_0) \right) \left( 1 - \chi \left( \frac{y}{b^{\frac{\delta}{2}}(\tau_0)} \right) \right) \\ &\times \left\{ \sum_{j=1}^{\ell-1} A d_j b^{\frac{\alpha}{2} + \eta}(\tau_0) \phi_{j, b(\tau_0), \beta(\tau_0)} - \frac{2}{\alpha} m_0 b^{\frac{\alpha}{2}}(\tau_0) \left[ 1 + d_\ell A b^{\frac{\alpha}{2} + \eta}(\tau_0) \right] \left( \frac{\phi_{\ell, b(\tau_0), \beta(\tau_0)}}{c_{\ell,0}} - \phi_{0, b(\tau_0), \beta(\tau_0)} \right) \right\} \end{aligned} \quad (8.5)$$

In particular, the class of initial data implies the following result

**Lemma 8.2** (Preparing the initial data). *There exists  $A_6 \geq 1$ , such that for all  $A \geq A_6$ , there exist  $\eta_6(A) > 0$  such that for all  $\delta \leq \delta_6$  there exists  $\eta_6(A, \delta) > 0$  such that for all  $\eta \leq \eta_6$  there exists  $\tilde{\eta}_6(A, \delta, \eta)$  such that for all  $\tilde{\eta} \leq \tilde{\eta}_6$  there exists  $\tau_6(A, \delta, \eta, \tilde{\eta}) > 1$  such that for all  $\tau_0 \geq \tau_6$ , there exists  $\mathcal{D}_A \subset [-2, 2]^\ell$  such that the following properties are valid*

(i) *the mapping*

$$\begin{aligned} \Gamma : \mathbb{R}^\ell &\rightarrow \mathbb{R}^\ell \\ (d_1, \dots, d_{\ell-1}) &\mapsto (\psi_1, \dots, \psi_\ell)(\tau_0), \end{aligned}$$

*is affine and one to one from  $\mathcal{D}_A$  to  $\hat{V}[A, \eta](\tau_0)$ , where*

$$\hat{V}[A, \eta](\tau) = \left[ -Ab^{\frac{\alpha}{2}+\eta}(\tau), Ab^{\frac{\alpha}{2}+\eta}(\tau) \right]^\ell. \quad (8.6)$$

*In particular, we have the following property*

$$\Gamma|_{\partial\mathcal{D}_A} \in \partial\hat{V}_A(\tau_0),$$

*and  $\deg(\Gamma|_{\partial\mathcal{D}_A}) \neq 0$ .*

(ii) *for all  $(d_1, \dots, d_\ell) \in \mathcal{D}_A$ , it follows that  $\psi_\ell(\tau_0) \in V_\ell[A, \tau, \tilde{\eta}](\tau_0)$  with strictly improved bounds as follows*

$$\begin{aligned} \left| \psi_\ell(\tau_0) + \frac{2}{\alpha} m_0 b^{\frac{\alpha}{2}}(\tau_0) \right| &\leq b^{\frac{\alpha}{2}+\eta}(\tau_0), \\ \|\psi_-(\tau_0)\|_{L^2} &\leq b^{\frac{\alpha}{2}+\eta}(\tau_0), \\ \left\| \frac{y^\gamma}{\langle y \rangle^{2\ell+2}} \psi_-(\cdot, \tau_0) \right\|_{L^\infty[0, b^{-\frac{\tilde{\eta}}{2}}(\tau_0)]} &\leq b^{\frac{\alpha}{2}+\eta}(\tau_0), \\ \|\psi_e(\cdot, \tau)\|_{L^\infty} &\leq b^{\frac{\alpha}{2}+\eta}(\tau_0), \end{aligned}$$

*and  $b(\tau_0)I_\ell^{-1}(\tau_0) \in [\frac{1}{4}, \frac{3}{2}]$ .*

*Proof.* The bounds in item (ii) immediately follow from the explicit form of  $\psi(\ell, \tau_0)$  in (8.5). In addition, the existence of  $\mathcal{D}_A$  and mapping  $\Gamma$  follows from the concentration of modes  $\psi_j, j \in \{0, 1, \dots, \ell\}$  of  $\psi(\ell, \tau_0)$  and the argument is quite the same as in [32, Proposition 4.5]. We kindly refer the reader to check the details of this result.  $\square$

### 8.3. Finite dimensional reduction

Since the shrinking set  $V_\ell[A, \eta, \tilde{\eta}](\tau)$  has special properties, the conclusion of Theorem 1.2 immediately the following

$$(\varepsilon, b)(\tau) \in V_\ell[A, \eta, \tilde{\eta}](\tau) \forall \tau \in [\tau_0, +\infty), \text{ for some } \tau_0 \text{ large enough.}$$

In particular, we prove in this part that this control is reduced to a finite dimensional problem on  $(\varepsilon_j)_{j \in \{1, 2, \dots, \ell\}}$

**Proposition 8.3** (Finite dimensional reduction). *There exists  $A_7 \geq 1$ , such that for all  $A \geq A_7$ , there exists  $\delta_7(A)$  such that for all  $\delta \leq \delta_7$  there exists  $\eta_7(A, \delta)$  such that for all  $\eta \leq \eta_7$  there exists  $\tilde{\eta}_7(A, \delta, \eta)$  such that for all  $\tilde{\eta} \leq \tilde{\eta}_7$  there exists  $\tau_7(A, \delta, \eta, \tilde{\eta})$  such that for all  $\tau_0 \geq \tau_7$ , the following property is valid: If  $(\varepsilon, b)$  is the solution to the coupled system (2.17-3.4) with initial data (8.5),  $(\varepsilon, b)(\tau) \in V_\ell[A, \eta, \tilde{\eta}](\tau)$  for all  $\tau \in [\tau_0, \tau^*]$  for some  $\tau^* > \tau_0$  and  $(\varepsilon, b)(\tau^*) \in \partial V_\ell[A, \eta, \tilde{\eta}](\tau^*)$ , then we have the following:*

(i) *It holds that  $(\varepsilon_1, \dots, \varepsilon_\ell)(\tau^*) \in \partial\hat{V}[A, \eta](\tau^*)$  defined as in (8.6).*

(ii) *Transversality*: There exists  $\nu_0 > 0$  such that

$$(\varepsilon_1, \dots, \varepsilon_\ell)(\tau^* + \nu) \notin \partial\hat{V}[A, \eta](\tau^* + \nu), \forall \nu \in (0, \nu_0)$$

which implies

$$(\varepsilon, b)(\tau^* + \nu) \notin V[A, \eta, \tilde{\eta}](\tau^* + \nu), \forall \nu \in (0, \nu_0).$$

*Proof.* The proof mainly relies on the *priori estimate* which is the same as in Lemmas [6.1-7.3].

- *Proof of item (i)*: Let us consider  $A \geq A_7$ ,  $\delta \leq \delta_7(A)$ ,  $\eta \leq \eta_7(A, \delta)$ ,  $\tilde{\eta} \leq \tilde{\eta}_7(A, \delta, \eta)$ ,  $\tau_0 \geq \tau_7(A, \delta, \eta, \tilde{\eta})$  and  $(\varepsilon, b)(\tau) \in V_\ell[A, \eta, \tilde{\eta}](\tau) \forall \tau \in [\tau_0, \tau^*]$  such that Lemmas 7.1, 7.2, and 7.3 remain true (the technique of the proof is exactly the same and we kindly refer the reader to check the details). So, it immediately follows that the bounds of  $\varepsilon_-$  and  $\varepsilon_e$  given in Definition 8.1 for  $V_\ell[A, \eta, \tilde{\eta}]$  are always improved by  $\frac{1}{2}$ -factor. In addition, we completely reproduce the argument of Lemma 6.1 to establish the following results: For all  $\tau \in [\tau_0, \tau^*]$ , we have

$$\left| \varepsilon'_j(\tau) - \left( \frac{\alpha}{2} - j \right) \varepsilon_j(\tau) \right| \leq C b^{\frac{\alpha}{2} + 4\eta}(\tau), \forall \tau \in [\tau_0, \tau_1], \quad (8.7)$$

and

$$\begin{cases} \partial_\tau \varepsilon_\ell - \left( \frac{\alpha}{2} - \ell \right) \varepsilon_\ell = O\left(b^{\frac{\alpha}{2} + 4\eta}(\tau)\right), \\ \partial_\tau \varepsilon_\ell - \left( \frac{\alpha}{2} \right) \varepsilon_\ell + m_0 \left( \frac{b_\tau}{b} - 1 \right) b^{\frac{\alpha}{2}} = O(b^{\frac{\alpha}{2} + 4\eta}), \end{cases} \quad (8.8)$$

and

$$\left| \frac{b'(\tau)}{b(\tau)} - \left( 1 - \frac{2\ell}{\alpha} \right) \right| \leq C A b^{4\eta}(\tau). \quad (8.9)$$

From (8.9), we derive that

$$\frac{3}{4} < b(\tau) I_\ell^{-1}(\tau) \leq \frac{3}{2} \forall \tau \in [\tau_0, \tau^*],$$

provided that  $\tau_0 \geq \tau_7(A, \eta, \tilde{\eta})$ . Thus, we derive from the fact that  $(\varepsilon, b)(\tau) \in \partial V_\ell[A, \eta, \tilde{\eta}](\tau^*)$  the following

$$(\varepsilon_1, \varepsilon_2, \dots, \varepsilon_\ell)(\tau^*) \in \partial\hat{V}[A, \eta](\tau^*)$$

which concludes item (i).

- *Proof of item (ii)*: it is sufficient to prove that there exists  $\nu_0$  small enough such that

- Either there exists  $j \in \{1, \dots, \ell - 1\}$ , such that

$$|\varepsilon_j(\tau^* + \nu)| > A b^{\frac{\alpha}{2} + \eta}(\tau_1 + \nu), \forall \nu \in (0, \nu_0), \quad (8.10)$$

- or the following holds

$$\left| \varepsilon_\ell(\tau^* + \nu) + \frac{2}{\alpha} m_0 b^{\frac{\alpha}{2}}(\tau_1 + \nu) \right| > A b^{\frac{\alpha}{2} + \eta}(\tau_1 + \nu), \forall \nu \in (0, \nu_0), \quad (8.11)$$

As we proved in item (i), one of the following two cases holds

- *Case 1*: There exists  $j \in \{1, \dots, \ell - 1\}$  such that

$$\varepsilon_j(\tau^*) = \sigma_j A b^{\frac{\alpha}{2} + \eta}(\tau^*),$$

- *Case 2*:

$$\varepsilon_\ell(\tau_1) + \frac{2}{\alpha} m_0 b^{\frac{\alpha}{2}}(\tau_1) = \sigma_\ell A b^{\frac{\alpha}{2} + \eta}(\tau_1),$$

where  $\sigma_j = \pm 1$ . The goal is to prove that the first case implies (8.10) and the second one concludes (8.11). Indeed, we have

+ Assume that case 1 occurs. Without loss of generality, we can assume  $\sigma_j = 1$  and introduce

$$B_j(\tau) = \varepsilon_j(\tau) - A b^{\frac{\alpha}{2} + \eta}(\tau).$$

It is obvious that  $B(\tau^*) = 0$  and we also get from (8.7) and (8.9),

$$\begin{aligned} B'_j(\tau^*) &= \left(\frac{\alpha}{2} - j\right) \varepsilon_j(\tau^*) - A \left(\frac{\alpha}{2} + \eta\right) \frac{b'(\tau^*)}{b(\tau^*)} b^{\frac{\alpha}{2} + \eta}(\tau^*) + O(b^{\frac{\alpha}{2} + 4\eta}(\tau^*)) \\ &= Ab^{\frac{\alpha}{2} + \eta}(\tau^*) \left( (\ell - j) - \eta \left(1 - \frac{2\ell}{\alpha}\right) \right) + O\left(b^{\frac{\alpha}{2} + 4\eta}(\tau^*)\right) > 0, \end{aligned}$$

provided that  $\ell - j \geq 1$  and  $\eta \leq \eta_\tau(A, \eta, \tilde{\eta})$  and  $\tau^* \geq \tau_0 \geq \tau_\tau(A)$ . Then,  $B_j(\tau^* + \nu) > 0$  for all  $\nu \in (0, \nu_0)$  for some  $\nu_0$  small enough. Thus, (8.10) follows.

+ If the case 2 occurs. We also assume  $\sigma = 1$  (the opposite sign is the same), we then define

$$B_\ell(\tau) = \varepsilon_\ell(\tau) + \frac{2}{\alpha} m_0 b^{\frac{\alpha}{2}}(\tau) - Ab^{\frac{\alpha}{2} + \eta}(\tau),$$

and it holds that  $B_\ell(\tau^*) = 0$ . and we derive from (8.8) and (8.9) that

$$\begin{aligned} B'_\ell(\tau^*) &= \left(\frac{\alpha}{2} - \ell\right) \varepsilon_\ell(\tau^*) + \frac{2}{\alpha} m_0 \frac{\alpha b'}{2b} b^{\frac{\alpha}{2}}(\tau^*) - A \left(\frac{\alpha}{2} + \eta\right) \frac{b'}{b} b^{\frac{\alpha}{2} + \eta}(\tau^*) + O(b^{\frac{\alpha}{2} + 4\eta}(\tau_1)) \\ &= Ab^{\frac{\alpha}{2} + \eta}(\tau^*) \left[ \eta \left(\frac{2\ell}{\alpha} - 1\right) \right] + O(b^{\frac{\alpha}{2} + 4\eta}(\tau^*)) > 0, \end{aligned}$$

since  $\eta \left(\frac{2\ell}{\alpha} - 1\right) > 0$  and  $\tau^* \geq \tau_0 \geq \tau_\tau(A, \eta, \tilde{\eta})$ . Thus, we conclude that  $B_\ell(\tau^* + \nu) > 0$  for all  $\nu \in (0, \nu_0)$ , and (8.11) follows. This concludes the proof of the Proposition.  $\square$

#### 8.4. Topological argument

In this part, we aim to prove the existence of an initial datum  $(\varepsilon, b)(\tau_0)$  that leads to the global existence of  $(\varepsilon, b)(\tau) \in V_\ell[A, \eta, \tilde{\eta}](\tau), \forall \tau \in [\tau_0, +\infty)$ . The following is our result:

**Proposition 8.4.** *There exist  $A, \eta, \tilde{\eta}$  and  $\delta \ll 1$  satisfying  $A \gg 1, 1 \gg \delta \gg \eta \gg \tilde{\eta} > 0, \tilde{\eta}$  and  $\tau_0(A, \eta, \tilde{\eta}, \delta) \gg 1$  such that there exists  $\tilde{d} = (d_1, \dots, d_\ell) \in \mathcal{D}_A$  defined in Lemma 8.2 such that with initial data  $\varepsilon(\ell, \tau_0)$  defined as in (8.5), the solution  $(\varepsilon, b)$  to the coupled problem (2.17-3.4), globally exists and the following holds*

$$(\varepsilon, b)(\tau) \in V_\ell[A, \eta, \tilde{\eta}](\tau), \forall \tau \geq \tau_0.$$

*Proof.* The proof follows from the topological argument which was used in [3] and [25]. Let us assume  $A, \eta, \tilde{\eta}$  and  $\delta$  are suitably chosen such that Lemma 8.2 and Proposition 8.3 hold true. We now proceed to the proof by contradiction and we assume that for all  $\tilde{d} = (d_1, \dots, d_\ell) \in \mathcal{D}_A$ , there exists  $\tau(\tilde{d}) \in [\tau_0, +\infty)$  such that  $(\varepsilon, b)(\tau(\tilde{d})) \notin V_\ell[A, \eta, \tilde{\eta}](\tau(\tilde{d}))$ . Then, we can define for each  $\tilde{d} \in \mathcal{D}_A$  the maximum time

$$\tau^*(\tilde{d}) = \sup \{ \tau_1 \text{ such that } (\varepsilon, b)(\tau) \in V_\ell[A, \eta, \tilde{\eta}](\tau), \forall \tau \in [\tau_0, \tau_1], \} \quad (8.12)$$

Now, let the mapping  $\Pi$  be defined by

$$\Pi : \mathcal{D}_{A, \tau_0} \rightarrow \partial[-1, 1]^\ell \quad (8.13)$$

$$\tilde{d} = (d_1, \dots, d_\ell) \mapsto \Pi(\tilde{d}), \quad (8.14)$$

where

$$\Pi(\tilde{d}) = (Ab^{\frac{\alpha}{2} + \eta}(\tau^*(\tilde{d})))^{-1} \left( \varepsilon_1, \dots, \varepsilon_\ell + \frac{2}{\alpha} m_0 b^{\frac{\alpha}{2}} \right) (\tau^*(\tilde{d})).$$

In particular, the following properties hold:

(i)  $\Pi$  is continuous from  $\mathcal{D}_A$  to  $\partial[-1, 1]^\ell$ . Indeed, since  $\tau^*(\tilde{d})$ 's definition implies

$$(\varepsilon, b)(\tau^*(\tilde{d})) \in \partial V_\ell[A, \eta, \tilde{\eta}](\tau^*(\tilde{d})),$$

and item (ii) of Proposition 8.3 immediately yields the result.

(ii)  $\text{Deg}(\Pi|_{\partial\mathcal{D}_A}) \neq 0$ . The result follows from item (i) of Lemma 8.2.

Thus, such a mapping  $\Pi$  can not exist, since it contradicts the index theory and the conclusion of the Proposition follows.  $\square$

## 9. Existence of ground state

We show in this part the asymptotic behavior of the ground state solution to (2.1). Let us introduce  $Q$  to be the function satisfying

$$Q''(\xi) + \frac{d+1}{\xi}Q'_\xi - 3(d-2)Q^2 - (d-2)\xi^2Q^3 = 0, \quad Q(0) = -1 \text{ and } Q'(0) = 0. \quad (9.1)$$

We have the following result:

**Lemma 9.1.** *Let  $d \geq 10$ , then there exists a unique solution  $Q$  to equation (9.1) satisfying the following:*

(i) *Asymptotic behavior of  $Q$ :*

$$Q(\xi) = -1 + \sum_{i=1}^k a_i \xi^{2i} + O(\xi^{2k+2}) \text{ as } \xi \rightarrow 0, \quad (9.2)$$

$$Q(\xi) = -\frac{1}{\xi^2} + q_0 \xi^{-\gamma} + O(\xi^{-3\gamma-4}) \text{ as } \xi \rightarrow +\infty, \quad (9.3)$$

and  $q_0 > 0$ .

(ii) *In particular, when  $d = 10$ , the ground state is explicitly given by*

$$Q_{10}(\xi) = -\frac{1}{\xi^2 + 1}$$

(iii) *Asymptotic of  $\Lambda Q = 2Q + \xi \cdot \partial_\xi Q$ :*

$$\Lambda Q(\xi) < 0, \text{ and } \Lambda Q = \begin{cases} -2 + 4 \left( \frac{3d-6}{3d+6} \right) \xi^2 + \sum_{i=2}^k a'_i \xi^{2i} + O(\xi^{2k+2}) \text{ as } \xi \rightarrow 0, \\ a_0 \xi^{-\gamma} + O(\xi^{-\gamma-g}) \text{ as } \xi \rightarrow +\infty, \end{cases} \quad (9.4)$$

for some  $a_0 < 0$ .

We note that  $\Lambda Q$ 's asymptotic at infinity is stable under  $\partial_\xi^k$  for all  $k \in \mathbb{N}$ , more precisely

$$\partial_\xi^k \Lambda Q = \partial_\xi^k (a_0 \xi^{-\gamma}) + O(\xi^{-\gamma-g-k}) \text{ as } \xi \rightarrow +\infty. \quad (9.5)$$

*Proof.* - The proof of item (i): Following [18], we reformulate the ground state equation as an autonomous ODE. Indeed, let

$$Z(\xi) = -\xi^2 Q(\xi),$$

then

$$Z'' + \frac{d-3}{\xi} Z' - \frac{(d-2)}{\xi^2} Z(Z-1)(Z-2) = 0. \quad (9.6)$$

Again, apply the change of function

$$Z(\xi) = v(x) \text{ where } \xi = e^x,$$

to get

$$v''(x) + (d-4)v'(x) - (d-2)v(v-1)(v-2) = 0, x \in (-\infty, +\infty). \quad (9.7)$$

To prove global existence and asymptotic behavior of the solution, we employ the phase portrait analysis that used in [4] for the ground-state of the heat flow maps. First observe that (9.7) has three critical points  $v = 0, 1, 2$ . We choose  $v = 1$  to start our analysis (for  $v = 0, 2$ , the linear operator will have complex or positive eigenvalues). According to our initial condition  $Q(0) = -1, Q'(0) = 0$ ,

$Q$  locally exists which in turn implies  $v$ 's existence on  $(-\infty, x_0)$  for some  $x_0 < 0$  and  $|x_0|$  large enough. In particular, we have the boundary condition at  $-\infty$ :

$$v(x) = e^{2x} + \sum_{i=2}^k c_i e^{2ix} + O(e^{(2k+2)x}) \text{ as } x \rightarrow -\infty.$$

This allows us to consider  $v(x)$  as the flow starting at  $(0, 0)$  and ending at  $(1, 0)$ . Following [2], we linearize around 1, i.e.

$$\epsilon = v - 1,$$

then  $\epsilon$  solves

$$\epsilon'' + (d-4)\epsilon' - (d-2)\epsilon(\epsilon+1)(\epsilon-1) = 0. \quad (9.8)$$

The linearisation is given by

$$\epsilon'' + (d-4)\epsilon' + (d-2)\epsilon = (d-2)\epsilon^3,$$

We write the above equation in matrix form

$$\begin{pmatrix} \epsilon' \\ \epsilon \end{pmatrix}' = \begin{pmatrix} -(d-4) & -(d-2) \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \epsilon' \\ \epsilon \end{pmatrix} + \begin{pmatrix} (d-2)\epsilon^3 \\ 0 \end{pmatrix}.$$

The eigenvalues are

$$\begin{aligned} \lambda_1 &= \frac{1}{2}(\sqrt{d^2 - 12d + 24} - d + 4) \\ \lambda_2 &= \frac{1}{2}(-\sqrt{d^2 - 12d + 24} - d + 4), \end{aligned} \quad (9.9)$$

provided that

$$d^2 - 12d + 24 \geq 0,$$

otherwise, the solution is spiral at  $+\infty$ . We see that  $\lambda_{1,2}(d) < 0$  for all  $d \geq 10$ .

- *Construction of no-escape region:* Let us define

$$F(\epsilon, \epsilon') = (\epsilon', -(d-4)\epsilon' + (d-2)(\epsilon^3 - \epsilon)).$$

We introduce a trapping region

$$\mathcal{S} = \{(\epsilon, \epsilon') \mid \epsilon^3 - \epsilon \leq \epsilon' \leq 2(\epsilon^3 - \epsilon), \epsilon \in (-1, 0)\}.$$

- The lower boundary curve  $\epsilon' = (\epsilon^3 - \epsilon)$ . In the phase portrait space  $(\epsilon, \epsilon')$ , we define the normal vector  $\nu_{in}$  which points inward  $\mathcal{S}$

$$\nu_{in} = (-(3\epsilon^2 - 1), 1).$$

We easily check that

$$F(\epsilon, \epsilon') \cdot \nu_{in} = (\epsilon^3 - \epsilon)3(1 - \epsilon^2) > 0, \forall \epsilon \in (-1, 0).$$

- The upper boundary curve  $\epsilon' = 2(\epsilon^3 - \epsilon)$ . In the phase portrait space  $(\epsilon, \epsilon')$ , we define the normal vector  $\nu_{in}$  which points inward  $\mathcal{S}$

$$\nu_{in} = (2(3\epsilon^2 - 1), -1).$$

Then,

$$F(\epsilon, \epsilon') \cdot \nu_{in} = (\epsilon^3 - \epsilon)(12\epsilon^2 + d - 10) > 0, \forall \epsilon \in (-1, 0) \text{ and } m \geq 10.$$

The vector field  $F$  points always inward on the whole boundary of  $\mathcal{S}$  (excluding the stationary points  $(-1, 0)$  and  $(0, 0)$ ). This implies that the integral curve of  $F$  starting in  $\mathcal{S}$  must stay in  $\mathcal{S}$ .



- *The boundary conditions trapped in  $\mathcal{S}$* : Note that with initial data  $Q(0) = -1, Q'(0) = 0$ ,  $Q$  locally exists, this leads that  $\epsilon$  exists locally, i.e., it exists on  $(-\infty, x_0)$ , for some  $x_0$  near  $-\infty$ . In particular, we have  $Q \in C^\infty$ . Using Taylor expansion together with equation (9.1), we see that  $Q(y)$  behaves as follows

$$Q(\xi) = -1 + \left(\frac{3d-6}{2d+6}\right)\xi^2 + \left(\frac{1}{4} \cdot \frac{21d^2 - 74d + 64}{(3d+4)(d+4)}\right)\xi^4 + O(\xi^6) \text{ as } \xi \rightarrow 0.$$

- *Asymptotic of the trapped solution in  $\mathcal{S}$* : Let us discuss the boundary condition at  $-\infty$ : we have

$$\epsilon(x) = -1 + e^{2x} - \frac{3d-6}{3d+6}e^{4x} - \frac{1}{4} \frac{21d^2 - 74d + 64}{(3d+4)(d+4)}e^{6x} + O(e^{8x}), \text{ as } x \rightarrow -\infty.$$

Using this asymptotic, the solution can't end up at  $(-1, 0)$ . In addition, at  $(0, 0)$ , it gives the following general asymptotic of  $\epsilon$ :

$$\epsilon(x) = h_+e^{\lambda_1 x}(1 + O(e^{-2x})) + h_-e^{\lambda_2 x}(1 + O(e^{-2x})),$$

where

$$\lambda_1 = \frac{1}{2}(\sqrt{d^2 - 12d + 24} - d + 4) \text{ and } \lambda_2 = \frac{1}{2}(-\sqrt{d^2 - 12d + 24} - d + 4).$$

Assuming that  $h_+ = 0$ , we derive from the shrinking set  $\mathcal{S}$  that

$$-\epsilon < \epsilon' < -2\epsilon \forall$$

Then

$$h_-(1 + \lambda_2) > 0 \text{ and } h_-(2 + \lambda_2) < 0,$$

this contradicts the formula of  $\lambda_2$ .

So,  $h_+ \neq 0$ .

In addition to that, we require the same condition as  $h_-$

$$h_+(1 + \lambda_1) > 0 \text{ and } h_+(2 + \lambda_1) < 0.$$

since

$$\lambda_1 + 1 < 0 \text{ and } \lambda_1 + 2 > 0$$

we see that  $h_+ < 0$  and we get the conclusion.

- The proof of item (ii): can be done in straightforward way, we omit the details.

- The proof of item (iii): We reformulate  $Q(\xi)$  by

$$Q(\xi) = -\frac{(\epsilon(x) + 1)}{e^{2x}}, \xi = e^x. \quad (9.10)$$

Computation yields

$$\xi Q'_\xi = -\frac{\epsilon'_x(x)}{e^{2x}} - 2Q.$$

Thus,

$$\Lambda Q = 2Q + y\partial_y Q = -\frac{\epsilon'_x(x)}{e^{2x}} < 0 \quad \forall x \in (-\infty, \infty). \quad (9.11)$$

Now, we aim to find the higher derivative of  $\epsilon$ , i.e.,  $\partial_x^k \epsilon$  for all  $k \geq 1$ . In fact,  $\epsilon$  satisfies the following integral equation

$$\epsilon(x) = h_+e^{\lambda_1 x} + h_-e^{\lambda_2 x} - \frac{1}{\lambda_1 - \lambda_2} \int_x^\infty \left( e^{\lambda_1(x-x')} - e^{\lambda_2(x-x')} \right) g(\epsilon(x')) dx', \quad (9.12)$$

where  $g(z) = (d-2)z^3$ . This gives us

$$\epsilon(x) = h_+e^{\lambda_1 x} + O\left(e^{3\lambda_1 x}\right),$$

as  $x \rightarrow +\infty$ . In particular, applying  $\partial_x^k$  to the right hand side of equation (9.12), we derive the following

$$\partial_x^k \epsilon(x) = \partial_x^k (h_+ e^{\lambda_1 x}) + O(e^{3\lambda_1 x}), \text{ as } x \rightarrow +\infty. \quad (9.13)$$

Let us remark that, it remains to prove (9.5). Indeed, we have the following

$$\begin{aligned} \Lambda Q &= -\epsilon'(x)e^{-2x}, \\ \partial_y \Lambda Q &= -\epsilon''(x)e^{-3x} + 2\epsilon'(x)e^{-3x}, \\ \partial_y^2 \Lambda Q &= -\epsilon'''(x)e^{-4x} + 3\epsilon''e^{-4x} + 2\epsilon''(x)e^{-4x} - 6\epsilon'(x)e^{-4x}, \\ &= (-\epsilon''' + 5\epsilon'' - 6\epsilon')e^{-4x}, \\ \partial_y^3 \Lambda Q &= (-\epsilon^{(4)} + 5\epsilon''' - 6\epsilon'')e^{-5x} - 4(-\epsilon''' + 5\epsilon'' - 6\epsilon')e^{-5x}. \end{aligned}$$

By induction, we can prove that

$$\partial_y^k \Lambda Q = -\prod_{j=0}^{k-1} (\partial_x - 2 - j) \partial_x \epsilon(x) e^{(-2-k)x}, \forall k \geq 1.$$

Using (9.13) and the fact that  $\xi = e^x$ , we get

$$\begin{aligned} \Lambda_\xi Q(\xi) &= a_0 \xi^{-\gamma} + O(\xi^{-3\gamma-4}), \\ \partial_\xi \Lambda Q(\xi) &= a_0 (\lambda_1 - 2) \xi^{-\gamma-1} + O(\xi^{-3\gamma-5}), \\ \partial_\xi^2 \Lambda Q(\xi) &= a_0 (\lambda^2 - 5\lambda_1 + 6) \xi^{-\gamma-2} + O(\xi^{-3\gamma-6}). \end{aligned}$$

In particular, we have the general case as follows: for all  $k \geq 1$

$$\begin{aligned} \partial_\xi^k \Lambda Q &= a_0 \prod_{j=0}^{k-1} (\lambda_1 - 2 - j) \xi^{\lambda_1 - 2 - k} + O(\xi^{-\lambda_1 - 2 - k}) \\ &= a_0 (-\gamma) \dots (-\gamma - (k-1)) \xi^{-\gamma - k} + O(\xi^{-3\gamma - 4 - k}), \end{aligned}$$

where  $\gamma = 2 - \lambda_1$ ,  $a_0 = -\lambda_1 h_+$ . Thus, (9.5) directly follows. This finishes the proof of the Lemma.  $\square$

## 10. Diagonalisation of $\mathcal{L}_b$

The goal of this section is to give a detailed proof of Proposition 4.2 which is the same as the route map established in the Section 2 of [9].

### 10.1. Interior problem

In the sequel, we construct eigenfunctions for  $\mathcal{L}_b$  in the region  $0 \leq y \leq y_0 \ll 1$ . First, we introduce

$$w(y, \tau) = v(\xi, \tau), \quad \text{with } \xi = \frac{y}{\sqrt{b}}. \quad (10.1)$$

The interior zone can be written in terms of the blow-up variable  $\xi$  as

$$0 \leq \xi \leq \xi_0 := \frac{y_0}{\sqrt{b}}.$$

Recall the definition of  $\mathcal{L}_b$

$$\mathcal{L}_b w(y) = \frac{1}{b} (H_\xi - \beta b \Lambda_\xi) v.$$

where the Shrödinger type operator  $H_\xi$  defined by

$$H = \partial_\xi^2 + \frac{d+1}{\xi} \partial_\xi - 3(d-2) (2Q(\xi) + \xi^2 Q^2(\xi)). \quad (10.2)$$

**Lemma 10.1** (Generators of the Kernel of  $H$ ). *There exists a family  $\{T_i\}_{i \geq 0}$  with initial element  $T_0 = a_0^{-1} \Lambda_\xi Q$  belonging to the kernel of  $H_\xi$  and such that for all  $i \in \mathbb{N}$*

$$H(T_{i+1}) = T_i. \quad (10.3)$$

Moreover,  $T_i$  admits the expansion

$$T_i(\xi) = \begin{cases} \sum_{l=0}^q t_{i,l} \xi^{2i+2l} + O(\xi^{2i+2q+2}), \forall q \in \mathbb{N}, \text{ as } \xi \rightarrow 0, \\ C_i \xi^{-\gamma+2i} \left(1 + O\left(\frac{\ln \xi}{\xi^2}\right)\right), \text{ as } \xi \rightarrow +\infty, \end{cases} \quad (10.4)$$

and the derivatives, up to order  $k = 3$ , are such that

$$\partial_\xi^k T_i(\xi) = \partial_\xi^k (C_i \xi^{-\gamma+2i}) + O\left(\xi^{-\gamma+2i-2-k} \ln \xi\right), \text{ as } \xi \rightarrow +\infty. \quad (10.5)$$

Here  $\gamma$  and  $C_j$  were defined in (2.26), and (2.29), respectively.

Let

$$\Theta_i = \Lambda T_i - (2i - \alpha) T_i.$$

then, for all  $k \in \{0, 1, 2\}$

$$\partial_\xi^k \Theta_i(\xi) = O\left(\xi^{-\gamma+2i-k-2} \ln \xi\right) \text{ as } \xi \rightarrow +\infty. \quad (10.6)$$

*Proof.* A detailed proof is to be presented in Appendix D.  $\square$

The generators of the kernel of  $H_\xi$  are at hand, we are in position to perform the construction of the eigenvalues and the eigenfunctions in the interior zone. More precisely, our result reads

**Proposition 10.2** (Inner eigenfunctions). *Let  $\ell \in \mathbb{N}, \ell \geq 1, i \in \{0, \dots, \ell\}$  and  $\beta \in [\frac{1}{4}, \frac{3}{4}]$ . Then, there exists  $\epsilon_0(\beta) > 0$  small enough such that for all  $\epsilon \in (0, \epsilon_0)$  such that there exists  $y^*(\epsilon) \ll 1$  such that for all  $0 < y_0 \leq y^*$  there exist  $b^*(y_0)$  and  $\tilde{\lambda}^*(y_0)$  such that for all  $0 < b < b^*(y_0)$  and  $|\tilde{\lambda}| \leq \tilde{\lambda}^*$  there exists  $\phi_{i,int} \in C^\infty\left(\left[0, \frac{y_0}{\sqrt{b}}\right], \mathbb{R}\right)$  such that the following hold:*

$$(H - b\beta\Lambda) \phi_{i,int,\beta} = 2\beta b \left(\frac{\alpha}{2} - i + \tilde{\lambda}\right) \phi_{i,int,\beta}, \quad (10.7)$$

where  $\phi_{i,int,\beta}$  has the following decomposition

$$\phi_{i,int,\beta}(\xi) = \sum_{j=0}^i c_{i,j} (2\beta)^j b^j T_j(\xi) + \tilde{\lambda} \sum_{j=0}^i b^{j+1} (c_{i,j} (2\beta)^{j+1} T_{j+1}(\xi) + S_j(\xi)) + b R_i(\xi), \quad (10.8)$$

where the correction  $R_i$  and  $S_j$  satisfy the following estimates

$$\begin{aligned} \|S_j\|_{X_{\xi_0}^{2j+2-\gamma}} &\leq C y_0^2, \|\partial_b S_j\|_{X_{\xi_0}^{2j+4-\gamma}} \leq C, \|\partial_{\tilde{\lambda}} S_j\|_{X_{\xi_0}^{2j+2-\gamma}} \leq C y_0^2, \text{ and } \|\partial_\beta S_j\|_{X_{\xi_0}^{2j+2-\gamma}} \leq C y_0^2, \\ \|R_i\|_{X_{\xi_0}^{-\gamma+\epsilon}} &\leq C(\epsilon), \|\partial_b R_i\|_{X_{\xi_0}^{2-\gamma+\epsilon}} \leq C(\epsilon), \|\partial_{\tilde{\lambda}} R_i\|_{X_{\xi_0}^{2-\gamma+\epsilon}} \leq C(\epsilon) b, \text{ and } \|\partial_\beta R_i\|_{X_{\xi_0}^{2-\gamma+\epsilon}} \leq C(\epsilon). \end{aligned}$$

*Proof.* Due to the lengthy proof, we aim to put the details in Appendix E.  $\square$

## 10.2. Exterior problem

In this part, we aim to construct the eigenfunctions of  $\mathcal{L}_b$  on  $[y_0, +\infty)$ , for some  $y_0 \ll 1$ . The following is our result

**Proposition 10.3** (Outer eigenfunctions). *Let  $\ell \in \mathbb{N}, \ell \geq 1, i \in \{0, \dots, \ell\}$  and  $\beta \in [\frac{1}{4}, \frac{3}{4}]$ . Then, there exists  $y^*(\beta) \ll 1$  such that for all  $y_0 \leq y^*$ , there exist  $b^*(y_0, \beta)$  and  $\tilde{\lambda}^*(y_0, \beta, b^*)$  such that for all  $b \in (0, b^*)$  and  $\tilde{\lambda} \in (-\tilde{\lambda}^*, \tilde{\lambda}^*)$ , there exists a  $C^\infty[y_0, +\infty)$  function  $\phi_{i,out,\beta}$  satisfying*

$$\mathcal{L}_b \phi_{i,out,\beta} = \left(2\beta \left(\frac{\alpha}{2} - i\right) + \tilde{\lambda}\right) \phi_{i,out,\beta},$$

and having the following decomposition

$$\phi_{i,out,\beta} = \phi_{i,\infty,\beta} + \tilde{\lambda}(\tilde{\phi}_{i,\beta} + \tilde{R}_{i,1}) + \tilde{R}_{i,2},$$

where  $\tilde{\phi}_{i,\beta}$  satisfies

$$\left(\mathcal{L}_\infty^\beta - 2\beta\left(\frac{\alpha}{2} - i\right)\right)\tilde{\phi}_{i,\beta} = \phi_{i,\infty,\beta} \text{ with } \phi_{i,\infty,\beta} \text{ defined as in (4.2),}$$

and  $\tilde{R}_{i,1}$  and  $\tilde{R}_{i,2}$  fulfil the following estimates

$$\|\tilde{R}_{i,1}\|_{X_{y_0}^{\gamma-d,2i-\gamma+2}} \leq C|\tilde{\lambda}|, \quad \partial_b \tilde{R}_{i,1} = 0, \quad \|\partial_{\tilde{\lambda}} \tilde{R}_{i,1}\|_{X_{y_0}^{\gamma-d,2i-\gamma+2}} \leq C, \quad \|\partial_\beta \tilde{R}_{i,1}\|_{X_{y_0}^{\gamma-d,2i-\gamma+2}} \leq C,$$

and

$$\|\tilde{R}_{i,2}\|_{X_{y_0}^{-d,a'}} \leq Cb^{\frac{\alpha}{2}}, \quad \|\partial_b \tilde{R}_{i,2}\|_{X_{y_0}^{-d,a'}} \leq Cb^{\frac{\alpha}{2}-1}, \quad \|\partial_{\tilde{\lambda}} \tilde{R}_{i,2}\|_{X_{y_0}^{-d,a'}} \leq Cb^{\frac{\alpha}{2}}, \quad \|\partial_\beta \tilde{R}_{i,2}\|_{X_{y_0}^{-d,a'}} \leq Cb^{\frac{\alpha}{2}},$$

for  $a' = 2i + 2 - \gamma$  and  $X_{y_0}^{a,a'}$  is a Banach space generated by the norm

$$\|f\|_{X_{y_0}^{a,a'}} = \sup_{y \in [y_0, 1]} y^{-a} \left\{ \sum_{i=0}^2 y^i |\partial_y^i f(y)| \right\} + \sup_{y \in [1, +\infty)} y^{-a'} \left\{ \sum_{i=0}^2 y^i |\partial_y^i f(y)| \right\}. \quad (10.9)$$

*Proof.* See Appendix F. □

### 10.3. Matching asymptotic

This part is devoted to conclude the proof of the diagonalisation on  $\mathcal{L}_b$ .

*Proof of Proposition 4.2:* Let  $i \in \{0, 1, \dots, \ell\}$  where  $\ell \in \mathbb{N}, \ell \geq 2, \beta \in [\frac{1}{4}, \frac{3}{4}]$ ,  $y_0 \leq y_1^*, b \leq b_1^*$  such that Propositions (10.2- 10.3) hold and  $\phi_{i,int}$  and  $\phi_{i,out,\beta}$  are defined in there.

A) *The proof of item (I):* We define

$$\phi_{i,b,\beta}(y) = \begin{cases} b^{-\frac{\gamma}{2}} \phi_{i,int,\beta} \left( \frac{y}{\sqrt{b}} \right) & \text{if } y \in [0, y_0], \\ \frac{b^{-\frac{\gamma}{2}} \phi_{i,int,\beta} \left( \frac{y_0}{\sqrt{b}} \right)}{\phi_{i,out,\beta}(y_0)} \phi_{i,out,\beta}(y) & \text{if } y \in [y_0, +\infty). \end{cases} \quad (10.10)$$

The main goal is to prove that there exists  $y_0 \in (0, 1)$  small enough and  $b^*(y_0) > 0$  such that, for all  $b \in (0, b^*)$ , there exists a unique  $\tilde{\lambda}_i(b, \beta) = \tilde{\lambda}$  satisfying

$$\mathcal{L}_b \phi_{i,b,\beta} = \left( 2\beta \left( \frac{\alpha}{2} - i \right) + \tilde{\lambda} \right) \phi_{i,b,\beta}, \quad (10.11)$$

and  $\tilde{\lambda}$  satisfies (4.6).

First, we observe that  $\phi_{i,int,\beta} \in C^\infty \left( \left[ 0, \frac{y_0}{\sqrt{b}} \right] \right)$  and  $\phi_{i,out,\beta} \in C^\infty[y_0, +\infty)$  and they solve the regular second order differential equations, so  $\phi_{i,b,\beta} \in C^\infty[0, +\infty)$  if and only if

$$b^{-\frac{\gamma}{2}-\frac{1}{2}} \partial_\xi \phi_{i,int,\beta} \left( \frac{y_0}{\sqrt{b}} \right) = b^{-\frac{\gamma}{2}} \frac{\phi_{i,int,\beta} \left( \frac{y_0}{\sqrt{b}} \right)}{\phi_{i,out,\beta}(y_0)} \partial_y \phi_{i,out,\beta}(y_0), \quad (10.12)$$

this condition ensures  $\phi_{i,b,\beta}$ 's differential is continuous at  $y_0$ . In particular, it is equivalent to

$$b^{-\frac{1}{2}} \partial_\xi \phi_{i,int,\beta} \left( \frac{y_0}{\sqrt{b}} \right) \phi_{i,out,\beta}(y_0) - \phi_{i,int,\beta} \left( \frac{y_0}{\sqrt{b}} \right) \partial_y \phi_{i,out,\beta}(y_0) = 0. \quad (10.13)$$

Here we use the implicit function theorem by applying it to the function  $\tilde{F}[y_0](\tilde{\lambda}, b, \beta)$  defined by

$$\tilde{F}[y_0](\tilde{\lambda}, b, \beta) = b^{-\frac{1}{2}} \partial_\xi \phi_{i,int,\beta} \left( \frac{y_0}{\sqrt{b}} \right) \phi_{i,out,\beta}(y_0) - \phi_{i,int,\beta} \left( \frac{y_0}{\sqrt{b}} \right) \partial_y \phi_{i,out,\beta}(y_0).$$

We firstly prove the following expansions:

$$\begin{aligned} \tilde{F}[y_0](\tilde{\lambda}, b, \beta) &= \tilde{\lambda} K_0 a_{i,0} (\tilde{\gamma} - \gamma) y_0^{-\gamma - \tilde{\gamma} - 1} (1 + O(y_0^2) + O(|\tilde{\lambda}|)) \\ &\quad + O(b^{1 - \frac{\epsilon}{2}}), \end{aligned} \quad (10.14)$$

$$\partial_b \tilde{F}[y_0](\tilde{\lambda}, b, \beta) = O(y_0^{-2\gamma - 1 + \epsilon} b^{-\frac{\epsilon}{2}}) + O(|\tilde{\lambda}| b^{-1} y_0^{-2\gamma + 3}), \quad (10.15)$$

$$\partial_{\tilde{\lambda}} \tilde{F}[y_0](\tilde{\lambda}, b, \beta) = (\tilde{\gamma} - \gamma) K_0 a_{i,0} y_0^{-\gamma - \tilde{\gamma} - 1} (1 + O(y_0^2) + O(|\tilde{\lambda}|)), \quad (10.16)$$

and for  $\beta$ -derivative

$$\partial_\beta \tilde{F}[y_0](\tilde{\lambda}, b, \beta) = \begin{cases} \tilde{\lambda} 2a_{i,1} K_0 (2 + \tilde{\gamma} - \gamma) y_0^{-\tilde{\gamma} - \gamma + 1} (1 + O(|y_0|^2) + O(|\tilde{\lambda}|)) & \text{if } i \geq 1, \\ \quad + O(b^{1 - \frac{\epsilon}{2}}) & (10.18) \\ O(|\tilde{\lambda}| y_0^{-\tilde{\gamma} - \gamma + 3}) + O(|\tilde{\lambda}|^2) + O(b^{1 - \frac{\epsilon}{2}}) & \text{if } i = 0. \end{cases}$$

Since the proof of asymptotic expansions (10.14-10.18) are technical and long, we complete them when we finish the proof of Proposition 4.2. Assume that the asymptotic expansions hold for all  $b \in (0, b^*(y_0))$ ,  $y_0 \leq y_0^*$ , and  $\beta \in [\frac{1}{4}, \frac{3}{4}]$ . We mention that the expansions are uniform in  $\beta, \tilde{\lambda}$  and  $\beta$ . So, the argument from the implicit function theorem yields that  $\forall b \in (0, b^*(y_0))$  and  $\beta \in [\frac{1}{4}, \frac{3}{4}]$ , there exists a unique  $\tilde{\lambda} = \tilde{\lambda}(b, \beta)$  such that

$$\tilde{F}[y_0](\tilde{\lambda}, b, \beta) = 0.$$

In particular, (10.14) ensures that  $\tilde{\lambda}(b, \beta) = O(b^{1 - \frac{\epsilon}{2}})$  and expansions (10.15-10.18) imply

$$\left| b \partial_b \tilde{\lambda}(b, \beta) \right| \lesssim_{y_0} b^{1 - \frac{\epsilon}{2}} \quad \text{and} \quad \left| \partial_\beta \tilde{\lambda}(b, \beta) \right| \lesssim_{y_0} 1$$

yielding (4.6). Next, we decompose  $\phi_{i,b,\beta}$  as follows

$$\phi_{i,b,\beta}(y) = \sum_{j=0}^i c_{i,j} (2\beta)^j (\sqrt{b})^{2j - \gamma} T_j \left( \frac{y}{\sqrt{b}} \right) + \tilde{\phi}_{i,b}(y), \quad (10.19)$$

and we aim to prove

$$\|\tilde{\phi}_{i,b}\|_{H_\rho^1} \leq C b^{1 - \frac{\epsilon}{2}}. \quad (10.20)$$

In particular, we can specify it by

$$\tilde{\phi}_{i,b,\beta}(y) = \begin{cases} b^{-\frac{\gamma}{2}} \phi_{i,int,\beta} \left( \frac{y}{\sqrt{b}} \right) - \sum_{j=0}^i c_{i,j} (\sqrt{b})^{2j - \gamma} T_j \left( \frac{y}{\sqrt{b}} \right) & \text{if } y \in [0, y_0], \\ \frac{b^{-\frac{\gamma}{2}} \phi_{i,int,\beta} \left( \frac{y_0}{\sqrt{b}} \right)}{\phi_{i,out,\beta}(y_0)} \phi_{i,out,\beta}(y) - \sum_{j=0}^i c_{i,j} (\sqrt{b})^{2j - \gamma} T_j \left( \frac{y}{\sqrt{b}} \right) & \text{if } y \in [y_0, +\infty). \end{cases}$$

Now, we aim to prove that

$$|\partial_y^k \tilde{\phi}_{i,b,\beta}(y)| \leq C \left( y^{-\gamma + 2 - k} I_{y \in [0, y_0]} + y^{-\gamma + 2i + 2 - k} I_{y \in [y_0, +\infty)} \right) b^{1 - \frac{\epsilon}{2}}, \quad y \in \mathbb{R} \text{ and } k = 0, 1. \quad (10.21)$$

Since the proofs for  $k = 0$  and  $k = 1$  are the same, we only give the proof of (10.21) for the case  $k = 0$  and we kindly refer the reader to check the details. Let us start the proof by considering two cases, namely,  $y \in [0, y_0]$  and  $y \in [y_0, +\infty)$ .

1.  $y \in [0, y_0]$ : write  $\tilde{\phi}_{i,b}$  as

$$\tilde{\phi}_{i,b}(y) = \tilde{\lambda} \sum_{j=0}^i b^{j+1 - \frac{\gamma}{2}} \left[ c_{i,j} T_{j+1} \left( \frac{y}{\sqrt{b}} \right) + S_j \left( \frac{y}{\sqrt{b}} \right) \right] + b^{1 - \frac{\gamma}{2}} R_i \left( \frac{y}{\sqrt{b}} \right).$$

According to Lemma 10.1, we have

$$|T_{j+1}(\xi)| \leq C \xi^{-\gamma + 2j + 2}, \quad \forall \xi \in \mathbb{R}^+,$$

so that

$$\left| \tilde{\lambda} b^{1+j-\frac{\gamma}{2}} c_{i,j} T_{j+1} \left( \frac{y}{\sqrt{b}} \right) \right| \leq C |\tilde{\lambda}| y^{2j+2-\gamma}.$$

Proposition 10.2 yields

$$\left| \tilde{\lambda} b^{1+j-\frac{\gamma}{2}} S_j \left( \frac{y}{\sqrt{b}} \right) \right| \leq C |\tilde{\lambda}| y^{2j+2-\gamma}, \quad \text{and} \quad b^{1-\frac{\gamma}{2}} \left| R_i \left( \frac{y}{\sqrt{b}} \right) \right| \leq C b^{1-\frac{\epsilon}{2}} y^{-\gamma+\epsilon}.$$

The above three estimates allows us to infer that, for  $y \in [0, y_0]$

$$|\tilde{\phi}_{i,b}(y)| \leq C b^{1-\frac{\epsilon}{2}} y^{2-\gamma}.$$

2.  $y \in [y_0, +\infty]$ : write  $\tilde{\phi}_{i,b}$  as follows

$$\begin{aligned} \tilde{\phi}_{i,b}(y) &= \frac{b^{-\frac{\gamma}{2}} \phi_{i,int} \left( \frac{y_0}{\sqrt{b}} \right)}{\phi_{i,ext}(y_0)} \phi_{i,ext}(y) - \sum_{j=0}^i c_{i,j} (\sqrt{b})^{2j-\gamma} T_j \left( \frac{y}{\sqrt{b}} \right) \\ &= \underbrace{\phi_{i,ext}(y) - \sum_{j=0}^i c_{i,j} (\sqrt{b})^{2j-\gamma} T_j \left( \frac{y}{\sqrt{b}} \right)}_{=I} + \underbrace{\left[ \frac{b^{-\frac{\gamma}{2}} \phi_{i,int} \left( \frac{y_0}{\sqrt{b}} \right)}{\phi_{i,ext}(y_0)} - 1 \right] \phi_{i,ext}(y)}_{=II}. \end{aligned}$$

Let

$$\tilde{T}_j(\xi) = T_j(\xi) - C_j \xi^{2j-\gamma}, \quad (10.22)$$

and we decompose  $\phi_{i,ext,\beta}$  as follows

$$\begin{aligned} I &= \phi_{i,ext}(y) - \sum_{j=0}^i c_{i,j} (\sqrt{b})^{2j-\gamma} C_j \left( \frac{y}{\sqrt{b}} \right)^{2j-\gamma} - \sum_{j=0}^i c_{i,j} (\sqrt{b})^{2j-\gamma} \tilde{T}_j \left( \frac{y}{\sqrt{b}} \right) \\ &= - \sum_{j=0}^i c_{i,j} (\sqrt{b})^{2j-\gamma} \tilde{T}_j \left( \frac{y}{\sqrt{b}} \right) + \tilde{\lambda} (\tilde{\phi}_i + \tilde{R}_{i,1}) + \tilde{R}_{i,2} \end{aligned}$$

Lemma 10.1 gives, for  $y \geq y_0$

$$\begin{aligned} \left| (\sqrt{b})^{2j-\gamma} \tilde{T}_j \left( \frac{y}{\sqrt{b}} \right) \right| &\leq C y^{-\gamma+2j-2} |\ln y| b |\ln b| \\ &\leq C y^{-\gamma+2j-2} |\ln y| b^{1-\frac{\epsilon}{2}}. \end{aligned}$$

From Proposition 10.3, we deduce that

$$\begin{aligned} \left| \tilde{\phi}_i(y) \right| &\leq C y^{2i-\gamma} |\ln y|, \\ \left| \tilde{R}_{i,1}(y) \right| &\leq C (y^{-\tilde{\gamma}} + y^{2i-\gamma+2}) |\tilde{\lambda}|, \\ \left| \tilde{R}_{i,2}(y) \right| &\leq C (y^{-\tilde{\gamma}-2-\alpha} + y^{2i+2-\gamma}) b^\alpha, \end{aligned}$$

for all  $y \geq y_0$ . Since  $\alpha \geq 1 - \frac{\epsilon}{2}$ , we obtain for all  $y \geq y_0$

$$\left| \phi_{i,out,\beta}(y) - \sum_{j=0}^i c_{i,j} (\sqrt{b})^{2j-\gamma} T_j \left( \frac{y}{\sqrt{b}} \right) \right| \leq C y^{2i+2-\gamma} |\ln y| b^{1-\frac{\epsilon}{2}}. \quad (10.23)$$

For the term II, we use the estimate obtained for I at  $y = y_0$  to get

$$|II| = \left| \frac{b^{-\frac{\gamma}{2}} \phi_{i,int} \left( \frac{y_0}{\sqrt{b}} \right)}{\phi_{i,ext}(y_0)} - 1 \right| \leq C(y_0) b^{1-\frac{\epsilon}{2}}.$$

Putting together the estimates for I and II yields for all  $y \geq y_0$

$$\left| \tilde{\phi}_{i,b}(y) \right| \leq C(y_0)y^{-\gamma+2i+2}b^{1-\frac{\epsilon}{2}}.$$

Similarly for  $\partial_y \tilde{\phi}_{i,b}$ , we establish

$$\left| \partial_y \tilde{\phi}_{i,b}(y) \right| \leq C(y_0)y^{-\gamma+2i+1}b^{1-\frac{\epsilon}{2}}.$$

- Now, we have

$$\|\phi_{i,b} - \phi_{i,\infty}\|_{H_\rho^1} \leq \left\| \sum_{j=0}^i c_{i,j}(\sqrt{b})^{2j-\gamma} T_j \left( \frac{y}{\sqrt{b}} \right) - \phi_{i,\infty} \right\|_{H_\rho^1} + \|\tilde{\phi}_{i,b}\|_{H_\rho^1}.$$

Taking into account, (10.20), it is sufficient to establish

$$\left\| \sum_{j=0}^i c_{i,j}(\sqrt{b})^{2j-\gamma} T_j \left( \frac{y}{\sqrt{b}} \right) - \phi_{i,\infty} \right\|_{H_\rho^1} \leq Cb^{1-\frac{\epsilon}{2}}.$$

As above, we have

$$\sum_{j=0}^i c_{i,j}(\sqrt{b})^{2j-\gamma} T_j \left( \frac{y}{\sqrt{b}} \right) - \phi_{i,\infty}(y) = \sum_{j=0}^i c_{i,j}(\sqrt{b})^{2j-\gamma} \tilde{T}_j \left( \frac{y}{\sqrt{b}} \right).$$

which yields, after splitting the integral in two regions  $\{y \leq \sqrt{b}\}$  and  $\{y \geq \sqrt{b}\}$  then using Lemma 10.1, to

$$\left\| \sum_{j=0}^i c_{i,j}(\sqrt{b})^{2j-\gamma} \tilde{T}_j \left( \frac{y}{\sqrt{b}} \right) \right\|_{H_\rho^1} \leq Cb^{1-\frac{\epsilon}{2}}.$$

- Now, we move to the proof of item (iii) in Proposition 4.2. We distinguish two regions:

- $y \in [0, y_0]$ : from definition (10.10), we have

$$\begin{aligned} \phi_{i,b}(y) &= b^{-\frac{\gamma}{2}} \phi_{i,int} \left( \frac{y}{\sqrt{b}} \right) = \sum_{j=0}^i c_{i,j}(\sqrt{b})^{2j-\gamma} T_j \left( \frac{y}{\sqrt{b}} \right) \\ &\quad + \tilde{\lambda} \sum_{j=0}^i b^{j+1-\frac{\gamma}{2}} \left[ c_{i,j} T_{j+1} \left( \frac{y}{\sqrt{b}} \right) + S_j \left( \frac{y}{\sqrt{b}} \right) \right] + b^{1-\frac{\gamma}{2}} R_i \left( \frac{y}{\sqrt{b}} \right). \end{aligned}$$

Since, for all  $\xi \in \mathbb{R}$

$$|T_j(\xi)| \leq C \frac{\xi^{2j}}{1 + \xi^\gamma},$$

we have, by Lemma 10.1

$$\left| \sum_{j=0}^i b^{j-\frac{\gamma}{2}} T_j \left( \frac{y}{\sqrt{b}} \right) \right| \leq C(i+1) \frac{\langle y \rangle^{2i}}{(\sqrt{b} + y)^\gamma}.$$

and

$$\left| \sum_{j=0}^i b^{j+1-\frac{\gamma}{2}} T_{j+1} \left( \frac{y}{\sqrt{b}} \right) \right| \leq C(i+1) \frac{\langle y \rangle^{2i+2}}{(\sqrt{b} + y)^\gamma}.$$

For  $S_j$ , we have

$$\left| \tilde{\lambda} \sum_{j=0}^i b^{1+j-\frac{\gamma}{2}} S_j \left( \frac{y}{\sqrt{b}} \right) \right| \leq C \frac{\langle y \rangle^{2i+2}}{(\sqrt{b}+y)^\gamma},$$

while for  $R_i$

$$\left| b^{1-\frac{\gamma}{2}} R_i \left( \frac{y}{\sqrt{b}} \right) \right| \leq C \frac{\langle y \rangle^\epsilon}{(\sqrt{b}+y)^\gamma}.$$

The above allows one to conclude that

$$|\phi_{i,b}(y)| \leq C \frac{\langle y \rangle^{2i+2}}{(\sqrt{b}+y)^\gamma}, \forall y \in [0, y_0].$$

- $y \in [y_0, +\infty)$ : from the definition of  $\phi_b$  on this region and the fact that

$$\left| b^{-\frac{\gamma}{2}} \frac{\phi_{i,int}(\frac{y_0}{\sqrt{b}})}{\phi_{i,ext}(y_0)} \right| \leq C(y_0),$$

it is sufficient to estimate  $\phi_{i,ext}$ . Recall that

$$\phi_{i,ext}(y) = \phi_{i,\infty}(y) + \tilde{\lambda}(\tilde{\phi}_i(y) + \tilde{R}_{i,1}) + \tilde{R}_{i,2}.$$

The asymptotic behavior of  $\tilde{\phi}_i$  yields, for all  $y \geq y_0$

$$|\tilde{\phi}_i(y)| \leq C \frac{y^{2i} |\ln y|}{y^\gamma} \leq C(y_0) \frac{\langle y \rangle^{2i+2}}{(\sqrt{b}+y)^\gamma}.$$

Moreover, we have the following facts: for all  $y \geq y_0$

$$\begin{aligned} |\tilde{R}_{i,1}(y)| &\leq C(y_0) |\tilde{\lambda}| \frac{\langle y \rangle^{2i+2}}{(\sqrt{b}+y)^\gamma}, \\ |\tilde{R}_{i,2}(y)| &\leq C(y_0) b^\alpha \frac{\langle y \rangle^{2i+2}}{(\sqrt{b}+y)^\gamma}. \end{aligned}$$

Putting together the above estimates, one gets

$$|\partial_y \phi_{i,b}(y)| \leq C \frac{\langle y \rangle^{2i+2}}{(\sqrt{b}+y)^\gamma}, \forall y \in [y_0, \infty).$$

A similar reasoning allows us to obtain the rest of the estimates, we omit the details.

- *Proof of (10.14)* :

First, we decompose  $\phi_{i,int,\beta}$  and  $\phi_{i,out,\beta}$  by

$$\begin{aligned} \phi_{i,int,\beta} \left( \frac{y_0}{\sqrt{b}} \right) &= b^{\frac{\gamma}{2}} \left\{ \phi_{i,\infty,\beta}(y_0) + \tilde{\lambda} \sum_{j=0}^i c_{i,j} C_{j+1} (2\beta)^{j+1} y_0^{-\gamma+2j+2} + A_{i,1}(\tilde{\lambda}, y_0, b, \beta) \right\}, \\ b^{-\frac{1}{2}} \partial_\xi \phi_{i,int,\beta} \left( \frac{y_0}{\sqrt{b}} \right) &= b^{\frac{\gamma}{2}} \left\{ \partial_y \phi_{i,\infty,\beta}(y_0) + \tilde{\lambda} \sum_{j=0}^i c_{i,j} C_{j+1} (2\beta)^{j+1} (-\gamma + 2j + 2) y_0^{-\gamma+2j+1} \right. \\ &\quad \left. + A_{i,2}(\tilde{\lambda}, y_0, b, \beta) \right\}, \end{aligned}$$

and

$$\begin{aligned} \phi_{i,out,\beta}(y_0) &= \phi_{i,\infty,\beta}(y_0) + \tilde{\lambda} K_0 y_0^{-\tilde{\gamma}} + B_{i,1}(\tilde{\lambda}, y_0, b, \beta), \\ \partial_y \phi_{i,out,\beta}(y_0) &= \partial_y \phi_{i,\infty,\beta}(y_0) - \tilde{\lambda} K_0 \tilde{\gamma} y_0^{-\tilde{\gamma}-1} + B_{i,2}(\tilde{\lambda}, y_0, b, \beta). \end{aligned}$$



where

$$\begin{aligned}
A_{i,1} &= \sum_{j=0}^i c_{i,j} (2\beta)^j b^{j-\frac{\gamma}{2}} \tilde{T}_j \left( \frac{y_0}{\sqrt{b}} \right) + \tilde{\lambda} \sum_{j=0}^i c_{i,j} (2\beta)^{j+1} b^{j+1-\frac{\gamma}{2}} \tilde{T}_{j+1} \left( \frac{y_0}{\sqrt{b}} \right) \\
&+ \tilde{\lambda} \sum_{j=0}^i b^{j+1-\frac{\gamma}{2}} S_j \left( \frac{y_0}{\sqrt{b}} \right) + b^{1-\frac{\gamma}{2}} R_i \left( \frac{y_0}{\sqrt{b}} \right), \\
A_{i,2} &= \sum_{j=0}^i c_{i,j} (2\beta)^j b^{j-\frac{1}{2}-\frac{\gamma}{2}} \partial_\xi \tilde{T}_j \left( \frac{y_0}{\sqrt{b}} \right) + \tilde{\lambda} \sum_{j=0}^i c_{i,j} (2\beta)^{j+1} b^{j+\frac{1}{2}-\frac{\gamma}{2}} \partial_\xi \tilde{T}_{j+1} \left( \frac{y_0}{\sqrt{b}} \right) \\
&+ \tilde{\lambda} \sum_{j=0}^i b^{j+\frac{1}{2}-\frac{\gamma}{2}} \partial_\xi S_j \left( \frac{y_0}{\sqrt{b}} \right) + b^{\frac{1}{2}-\frac{\gamma}{2}} \partial_\xi R_i \left( \frac{y_0}{\sqrt{b}} \right),
\end{aligned}$$

and

$$\begin{aligned}
B_{i,1} &= \tilde{\lambda} (\tilde{\phi}_{i,\beta} - K_0 y_0^{-\tilde{\gamma}}) + \tilde{\lambda} \tilde{R}_{i,1} + \tilde{R}_{i,2}, \\
B_{i,2} &= \tilde{\lambda} \partial_y (\tilde{\phi}_{i,\beta} - K_0 y_0^{-\tilde{\gamma}}) + \tilde{\lambda} \partial_y \tilde{R}_{i,1} + \partial_y \tilde{R}_{i,2},
\end{aligned}$$

and  $\tilde{T}_j$  and  $C_j$  defined as in (10.22) and (2.29), respectively.

We aim to estimate  $A_i$  and  $B_i$  by using the results of Propositions 10.2 and 10.3.

- estimate on  $A_{i,1}$ : From Lemma 10.1, we use  $T_j$ 's expansion at  $\infty$ , to obtain the following

$$|\tilde{T}_j(\xi)| \leq C \xi^{-\gamma+2j-2} \ln \xi,$$

for all  $\xi$  large enough, i.e  $\xi \geq \xi_0 > 1$ . Applying the above for  $\xi_0 = \frac{y_0}{\sqrt{b}}$ , we derive the following

$$\left| \tilde{T}_j \left( \frac{y_0}{\sqrt{b}} \right) \right| \leq C \left( \frac{y_0}{\sqrt{b}} \right)^{-\gamma+2j-2} (|\ln y_0| + |\ln b|), \forall j \geq 1,$$

and for  $j = 0$ , we have

$$\left| \tilde{T}_0 \left( \frac{y_0}{\sqrt{b}} \right) \right| \leq C \left( \frac{y_0}{\sqrt{b}} \right)^{-\gamma-2}.$$

This yields

$$\left| \sum_{j=0}^i c_{i,j} b^{-\frac{\gamma}{2}+j} \tilde{T}_j \left( \frac{y_0}{\sqrt{b}} \right) \right| \leq C y_0^{-\gamma-2} b |\ln b|.$$

For the second term of  $A_{i,1}$ , the same process as above gives

$$\left| \sum_{j=0}^i c_{i,j} b^{j+1-\frac{\gamma}{2}} \tilde{T}_{j+1} \left( \frac{y_0}{\sqrt{b}} \right) \right| \leq C y_0^{-\gamma-2} b |\ln b|.$$

We now estimate to  $S_j$ . Accordingly to Proposition 10.2, and the definition of  $X_{\xi_0}^{2j+2-\gamma}$ , we have

$$\left| S_j \left( \frac{y_0}{\sqrt{b}} \right) \right| \leq C y_0^2 \left( \frac{y_0}{\sqrt{b}} \right)^{2j+2-\gamma},$$

so that

$$\left| \tilde{\lambda} \sum_{j=0}^i b^{j+1-\frac{\gamma}{2}} S_j \left( \frac{y_0}{\sqrt{b}} \right) \right| \leq C \tilde{\lambda} y_0^{4-\gamma}.$$

The last term in  $A_1$  is to be estimated again via Proposition 10.2, where we have

$$\left| R_i \left( \frac{y_0}{\sqrt{b}} \right) \right| \leq C \left( \frac{y_0}{\sqrt{b}} \right)^{-\gamma+\epsilon}, \text{ with } \epsilon \ll 1.$$

Hence

$$\left| b^{1-\frac{\gamma}{2}} R_i \left( \frac{y_0}{\sqrt{b}} \right) \right| \leq C y_0^{-\gamma+\epsilon} b^{1-\frac{\epsilon}{2}}.$$

Finally, we get

$$A_1 = O(\tilde{\lambda} y_0^{4-\gamma}) + O\left(y_0^{-\gamma-2} b^{1-\frac{\epsilon}{2}}\right).$$

- For  $A_{i,2}$ : First, by Lemma 10.1 we get

$$\left| \partial_\xi \tilde{T}_j(\xi_0) \right| \leq C \xi_0^{-\gamma+2j-3} |\ln \xi_0|, \forall j \geq 1,$$

and

$$\left| \partial_\xi \tilde{T}_0(\xi_0) \right| \leq C |\xi_0|^{-\gamma-g-1}.$$

Then

$$\begin{aligned} \left| \sum_{j=0}^i c_{i,j} b^{-\frac{\gamma}{2}-\frac{1}{2}+j} \partial_\xi \tilde{T}_j(\xi_0) \right| &\leq C \sum_{j=1}^i b^{j-\frac{\gamma}{2}-\frac{1}{2}} |\xi_0|^{-\gamma+2j-3} |\ln \xi_0| + C b^{-\frac{\gamma}{2}-\frac{1}{2}} |\xi_0|^{-\gamma-g-1} \\ &\leq C y_0^{-\gamma-1} |\ln y_0| b |\ln b| + C y_0^{-\gamma-g-1} b^{\frac{g}{2}}. \end{aligned}$$

Next, we estimate the second term in  $A_{i,2}$ :

$$\left| \tilde{\lambda} \sum_{j=0}^i c_{i,j} b^{j+\frac{1}{2}-\frac{\gamma}{2}} \partial_\xi \tilde{T}_{j+1} \left( \frac{y_0}{\sqrt{b}} \right) \right| \leq C |\tilde{\lambda}| y_0^{-\gamma-1} |\ln y_0| b^{\frac{3}{2}} |\ln b|.$$

Using Proposition 10.2 for  $S_j$

$$|\partial_\xi S_j(\xi_0)| \leq C |\xi_0|^{a-1} \leq C |\xi_0|^{2j+1-\gamma},$$

we obtain

$$\left| \tilde{\lambda} \sum_{j=0}^i b^{j+\frac{1}{2}-\frac{\gamma}{2}} \partial_\xi S_j \left( \frac{y_0}{\sqrt{b}} \right) \right| \leq C |\tilde{\lambda}| y_0^{-\gamma+1}.$$

The last term in  $A_{i,2}$  is estimated similarly and we have

$$|b^{1-\frac{\gamma}{2}} \partial_\xi R_i(\xi_0)| \leq C y_0^{-\gamma+\epsilon-1} b^{\frac{3}{2}-\frac{\epsilon}{2}}.$$

Hence, the expansion of  $A_{i,2}$  is

$$A_{i,2} = O(|\tilde{\lambda}| y_0^{-\gamma+1}) + O(y_0^{-\gamma-1} |\ln y_0| b |\ln b|).$$

- For  $B_{i,1}$  Using Lemma F.1 we have

$$\left| \tilde{\lambda} (\tilde{\phi}_i(y_0) - K_0 y_0^{-\tilde{\gamma}}) \right| \leq C |\tilde{\lambda}| y_0^{-\tilde{\gamma}+2}.$$

For the second and third term, we use Proposition 10.3

$$|\tilde{R}_{i,1}(y_0)| \leq C |\tilde{\lambda}| y_0^{-\tilde{\gamma}}$$

and

$$\left| \tilde{R}_{i,2}(y_0) \right| \leq C y_0^{-\tilde{\gamma}-2-\alpha} b^\alpha.$$

Then,  $B_{i,1}$  reads as follows

$$B_{i,1}(y_0) = O\left(|\tilde{\lambda}| y_0^{-\tilde{\gamma}+2}\right) + O(|\tilde{\lambda}|^2 y_0^{-\tilde{\gamma}}) + O(y_0^{-\tilde{\gamma}-2-\alpha} b^\alpha).$$

- For  $B_{i,2}$ : a similar reasoning gives

$$B_{i,2}(y_0) = O\left(|\tilde{\lambda}|y_0^{-\tilde{\gamma}+1}\right) + O(|\tilde{\lambda}|^2y_0^{-\tilde{\gamma}-1}) + O(y_0^{-\tilde{\gamma}-3-\alpha}b^\alpha).$$

Putting the above expansions together, we derive for  $\tilde{F}[y_0](\tilde{\lambda}, b)$

$$\begin{aligned} \tilde{F}[y_0](\tilde{\lambda}, b, \beta) &= (\partial_y \phi_{i,\infty,\beta}(y_0) + \tilde{\lambda} \sum_{j=0}^i c_{i,j} (2\beta)^{j+1} C_{j+1} (-\gamma + 2j + 2) y_0^{-\gamma+2j+1} + A_2(\tilde{\lambda}, y_0, b, \beta)) \\ &\quad \times (\phi_{i,\infty,\beta}(y_0) + \tilde{\lambda} K_0 y_0^{-\tilde{\gamma}} + B_1(\tilde{\lambda}, y_0, b, \beta)) \\ &\quad - (\partial_y \phi_{i,\infty,\beta}(y_0) - \tilde{\lambda} K_0 \tilde{\gamma} y_0^{-\tilde{\gamma}-1} + B_2(\tilde{\lambda}, y_0, b, \beta)) \\ &\quad \times (\phi_{i,\infty,\beta}(y_0) + \tilde{\lambda} \sum_{j=0}^i c_{i,j} C_{j+1} (2\beta)^{j+1} y_0^{-\gamma+2j+2} + A_1(\tilde{\lambda}, y_0, b, \beta)) \\ &= \tilde{\lambda} K_0 a_{i,0} (\tilde{\gamma} - \gamma) y^{-\gamma-\tilde{\gamma}-1} \left(1 + O(y_0^2) + O(|\tilde{\lambda}|)\right) \\ &\quad + O\left(y_0^{-2\gamma-2} b^{1-\frac{\varepsilon}{2}}\right) + O\left(y_0^{-\gamma-\tilde{\gamma}-3-\alpha} b^\alpha\right). \end{aligned}$$

The proofs of (10.15) and (10.16) follow the same outline.  $\square$

## 11. Maximum principal

The main goal in this section is to use Maximum principal to construct the sub solution and the super solution to (2.17) on the interval  $[0, b^{\frac{\eta}{4}}(\tau)]$  that leads to suitable estimates for  $\varepsilon$ .

**Proposition 11.1** (Sub and super solutions). *Let us consider  $\eta, \tilde{\eta}$  be positive constants such that  $1 \gg \eta \gg \tilde{\eta}$ ,  $A \geq 1$ . We assume furthermore that  $\varepsilon$  is the solution to (2.17) on  $[\tau_0, \tau_1]$  with initial data given in (5.16) and the flow  $(b, \beta)(\tau) \in (C^1(\tau_0, \tau_1))^2$  satisfy  $(\varepsilon, b, \beta)(\tau) \in V[A, \eta, \tilde{\eta}](\tau)$  for all  $\tau \in [\tau_0, \tau_1]$ . Then, there exists  $H(\xi)$  satisfying*

$$|H(\xi)| \leq C(\eta) \left[ b(\tau) \frac{\xi^2}{1 + \xi^\gamma} + \frac{b^{\frac{\eta}{4}}(\tau)}{1 + \xi^\gamma} \right], \text{ for all } \xi \in \mathbb{R}_+,$$

such that

$$|\varepsilon(y, \tau)| \leq b^{-1}(\tau) H\left(\frac{y}{\sqrt{b(\tau)}}\right), \forall y \in [0, b^{\frac{\eta}{4}}(\tau)], \quad (11.1)$$

where  $Q_b$  defined as in (2.15). In other words, (11.2) remains true with  $\tau$  larger than  $\tau_1$  as long as  $w$  exists and  $b$  satisfies the hypothesis of the Proposition.

*Proof.* First, we claim that the following

$$|w(y, \tau) - Q_{b(\tau)}(y)| \leq b^{-1}(\tau) H\left(\frac{y}{\sqrt{b(\tau)}}\right), \forall y \in [0, b^{\frac{\eta}{4}}(\tau)] \quad (11.2)$$

implies (11.1). We also mention that the proof is similar to the one in [5] where the authors constructed sub-solution and super-solution to equation (2.12) on the small interval  $[0, b^{\frac{\eta}{4}}(\tau)]$ . Let us consider the blowup variable  $(\xi, \tau)$  and

$$w(y, \tau) = \frac{1}{b(\tau)} \omega\left(\frac{y}{\sqrt{b(\tau)}}, \tau\right) = \frac{1}{b(\tau)} \omega(\xi, \tau).$$

Introducing  $v = \omega - Q(\xi)$  and  $v$  reads

$$b(\tau)\partial_\tau v = \partial_\xi^2 v + \frac{d+1}{\xi}\partial_\xi v - 3(d-2)(2Q + \xi^2 Q^2)v + B(v) + \theta(\tau)\Lambda_\xi Q + \theta(\tau)\Lambda_\xi v, \quad (11.3)$$

where  $B$  defined as in (2.19) and  $\theta(\tau)$  defined by

$$\theta(\tau) = \beta(\tau) (b'(\tau) - b(\tau)). \quad (11.4)$$

We also introduce the operator  $\mathcal{P}$  as follows

$$\mathcal{P}(v) := \partial_\xi^2 v + \frac{d+1}{\xi}\partial_\xi v - 3(d-2)(2Q + \xi^2 Q^2)v + \bar{B}(v) + \theta(\tau)\Lambda_\xi Q + \theta(\tau)\Lambda_\xi v - b(\tau)\partial_\tau v, \quad (11.5)$$

In order to construct the sub-solution and the super-solution, we need to construct two functions as follows: let  $Q$  be the ground state satisfying (9.1),  $Q(0) = -1, Q'(0) = 0$ , and we introduce

$$Q_\sigma = \frac{1}{\sigma} Q\left(\frac{\xi}{\sqrt{\sigma}}\right), \sigma > 0,$$

exactly solves (9.1) thanks to the scaling (1.4). Next, define

$$H_0(\xi) = \Lambda_\xi Q_\sigma(\xi), \quad (11.6)$$

satisfying

$$H_0'' + \frac{d+1}{\xi}H_0' - 3(d-2)(2Q_\sigma + \xi^2 Q_\sigma^2)H_0 = 0, \quad (11.7)$$

and let  $H_1(\xi)$  solve the following

$$H_1'' + \frac{d+1}{\xi}H_1' - 3(d-2)(2Q_\sigma + \xi^2 Q_\sigma^2)H_1 = T(\xi), \quad (11.8)$$

where  $T(\xi) = -\Lambda_\xi Q$ . In particular,  $H_1$  is explicitly given by

$$H_1(\xi) = H_0 \int_0^\xi \frac{\mathcal{L}(T)(\xi')}{H_0(\xi')} d\xi', \quad (11.9)$$

where  $\mathcal{L}$  was defined in (D.3) Using (11.6) and (11.9),  $H_0$  and  $H_1$  have the following asymptotics:

$$H_0(\xi) = \begin{cases} -\frac{2}{\sigma} & \text{as } \xi \rightarrow 0, \\ a_0 \sigma^\alpha \xi^{-\gamma} & \text{as } \xi \rightarrow \infty, \end{cases} \quad (11.10)$$

where  $a_0 < 0$  and  $\alpha = 2 - \gamma$  and

$$H_1(\xi) = \begin{cases} \frac{\xi^2}{d+2} & \text{as } \xi \rightarrow 0, \\ \frac{-a_0}{2(d+2-\gamma)} \xi^{2-\gamma} & \text{as } \xi \rightarrow \infty. \end{cases} \quad (11.11)$$

Inspired by [5], we define

$$v^+(\xi, \tau) = \theta^+(\tau)H_1(\xi) - M(\eta)b^{\frac{\eta}{4}}(\tau)H_0(\xi) \quad \text{and} \quad v^- = \theta^-(\tau)H_1(\xi) + M(\eta)b^{\frac{\eta}{4}}(\tau)H_0(\xi), \quad (11.12)$$

where

$$\theta^+(\tau) = b(\tau) \left[ \beta(2\beta - 1) - 4\beta^2 \frac{\ell}{\alpha} - b^{\frac{\eta}{8}}(\tau) \right] \quad \text{and} \quad \theta^-(\tau) = b(\tau) \left[ \beta(2\beta - 1) - 4\beta^2 \frac{\ell}{\alpha} + b^{\frac{\eta}{8}}(\tau) \right] \quad (11.13)$$

Note that  $H_0(\xi) = \Lambda_\xi Q_\sigma < 0$  see more (9.4). In particular, our aim is to prove

$$w^+(y, \tau) = Q_{b(\tau)}(y) + \frac{1}{b(\tau)}v^+ \left( \frac{y}{\sqrt{b(\tau)}} \right) \quad \text{and} \quad w^-(y, \tau) = Q_{b(\tau)}(y) + \frac{1}{b(\tau)}v^- \left( \frac{y}{\sqrt{b(\tau)}} \right)$$

are respectively the super-solution and the sub-solution to (2.12) which immediately implies (11.2). Following [5], it is sufficient to check that

$$(i) \quad \mathcal{P}(v^+) < 0 \quad (\mathcal{P}(v^-) > 0), \forall \tau \in [\tau_0, \tau_1] \quad \text{and} \quad \xi \leq b^{\frac{\eta}{4} - \frac{1}{2}}(\tau).$$

(ii) Initial estimate:  $\frac{1}{b(\tau_0)}v^- \left( \frac{y}{\sqrt{b(\tau_0)}}, \tau_0 \right) < w(y, \tau_0) - Q_{b(\tau_0)}(\xi) < \frac{1}{b(\tau_0)}v^+ \left( \frac{y}{\sqrt{b(\tau_0)}}, \tau_0 \right), \forall y \leq b^{\frac{\eta}{4}}(\tau_0)$ .

We remark that the proof of the estimates on  $v^-$  are quite the same as for  $v^+$ . Thus, we only handle the latter.

- Proof of (i): plugging  $v^+$  into (11.5), we get

$$\begin{aligned} \mathcal{P}(v^+) &= \theta^+(\tau)\partial_\xi^2 H_1 - Mb^{\frac{\eta}{4}}(\tau)\partial_\xi^2 H_0 + \theta^+(\tau) \left( \frac{d+1}{\xi} \partial_\xi H_1 \right) - M\frac{\eta}{4}(\tau)\frac{d+1}{\xi} \partial_\xi H_0 \\ &\quad - 3(d-2) [2Q + \xi^2 Q^2] \left( \mu^+(\tau)H_1 - Mb^{\frac{\eta}{4}}(\tau)H_0(\xi) \right) + \theta(\tau)\Lambda_\xi Q + \bar{B}(v^+) \\ &\quad + \theta(\tau) \left[ \theta^+ \Lambda_\xi H_1 - Mb^{\frac{\eta}{4}} \Lambda_\xi H_0 \right] - b(\tau) \left[ \partial_\tau \theta^+(\tau)H_1 - M\partial_\tau b^{\frac{\eta}{4}}(\tau)H_0(\xi) \right] \\ &= -3(d-2) [2Q + \xi^2 Q^2 - (2Q_\sigma + \xi^2 Q_\sigma^2)] (\theta^+(\tau)H_1(\xi) - Mb^\eta(\tau)H_0(\xi)) \\ &\quad + \bar{B}(v^+) + [\theta - \theta^+] \Lambda_\xi Q + \theta(\tau) \left[ \theta^+ \Lambda_\xi H_1 - Mb^{\frac{\eta}{4}} \Lambda_\xi H_0 \right] - b(\tau) \left[ \partial_\tau \theta^+ H_1 - M\partial_\tau b^{\frac{\eta}{4}} H_0 \right], \end{aligned}$$

where the simplification comes from the facts that  $H_0$  and  $H_1$  solve (11.7) and (11.8), respectively. Since  $\xi \leq b^{\frac{\eta}{4}-\frac{1}{2}}(\tau)$  with  $\eta \ll 1$  and  $b(\tau) \rightarrow 0$ , the range of  $\xi$  will be large, and we should divide it into two cases  $\xi = O(1)$  and  $\xi \gg 1$ .

+ For the case  $\xi \gg 1$ , we derive on the one hand, from the definitions of  $\theta, \theta^+$  in (11.4) (11.13), and (6.3), that

$$\theta(\tau) - \theta^+(\tau) = b^{1+\frac{\eta}{8}} + O(b^{1+4\eta}).$$

Thus we derive from (9.4) that

$$[\theta(\tau) - \theta^+] \Lambda_\xi Q = a_0 b^{1+\frac{\eta}{8}}(\tau) \xi^{-\gamma} + O(\xi^{-\gamma} b^{1+4\eta}) + O(b^{1+\frac{\eta}{8}} \xi^{-\gamma-2}),$$

where  $a_0 < 0$ . On the other hand, by a similar argument, we have

$$\begin{aligned} -b(\tau) \left[ \partial_\tau \theta^+ H_1 - M\partial_\tau b^{\frac{\eta}{4}} H_0 \right] &= O\left(b^{1+\frac{\eta}{4}} \xi^{-\gamma}\right), \\ \theta(\tau) \left[ \theta^+ H_1 - Mb^{\frac{\eta}{4}} H_0 \right] &= O\left(b^{1+\frac{\eta}{4}} \xi^{-\gamma}\right) \text{ as } \xi \rightarrow \infty. \end{aligned}$$

Thus, we conclude the following inequality

$$\mathcal{P}(z^+) \leq -3(d-2) [2Q + \xi^2 Q^2 - (2Q_\sigma + \xi^2 Q_\sigma^2)] v^+ + \bar{B}(v^+),$$

provided that  $b, \eta$  small enough.

Next, note that  $\xi \leq b^{\frac{\eta}{4}-\frac{1}{2}}(\tau)$  implies the following

$$|b(\tau)\xi^2| \leq b^{\frac{\eta}{2}}(\tau).$$

Hence, by using  $v^+$ 's definition in (11.12), along with (11.10) and (11.11), we obtain

$$v^+(\xi, \tau) = -Ma_0 \sigma^\alpha b^{\frac{\eta}{4}} \xi^{-\gamma} + O(b^\eta \xi^{-\gamma-2}) + O(b^{2\beta} \xi^{-\gamma}), \text{ as } \xi \rightarrow +\infty.$$

In addition, recall that

$$\begin{aligned} 2Q(\xi) + \xi^2 Q^2(\xi) &\sim -\frac{1}{\xi^2} + q_0^2 \xi^{-2\gamma+2} + o(\xi^{-2\gamma+2}), \\ 2Q_\sigma(\xi) + \xi^2 Q_\sigma^2(\xi) &\sim -\frac{1}{\xi^2} + q_0^2 \sigma^{\gamma-2} \xi^{-2\gamma+2} + o(\xi^{-2\gamma+2}), \end{aligned}$$

as  $\xi \rightarrow +\infty$ . Then, fixing  $\sigma$  less than 1, we derive

$$2Q(\xi) + \xi^2 Q^2(\xi) - [2Q_\beta(\xi) + \xi^2 Q_\beta^2(\xi)] = q_0^2 (1 - \sigma^{\gamma-2}) \xi^{-3\gamma+2} + o(\xi^{-2\gamma+2}).$$

Thus,

$$-3(d-2) [2Q + \xi^2 Q^2 - (2Q_\sigma + \xi^2 Q_\sigma^2)] v^+ = m_0 b^{\frac{\eta}{4}}(\tau) \xi^{-3\gamma+2} + o(\xi^{-3\gamma+2}), \text{ as } \xi \rightarrow +\infty. \quad (11.14)$$

where  $m_0 = 3(d-2)q_0^2(1-\sigma^{\gamma-2})Ma_0\sigma^\gamma < 0$ . Next, we study  $\bar{B}(v^+)$  defined by

$$\begin{aligned} \bar{B}(v^+) &= -3(d-2)(1 + \xi^2 Q(\xi))(v^+)^2 - (d-2)\xi^2(v^+)^3 \\ &= \left(m_1 b^{\frac{\eta}{2}} + m_2 b^{\frac{3\eta}{4}}\right) \xi^{-3\gamma+2} + o(\xi^{-3\gamma+2}), \text{ as } \xi \rightarrow +\infty, \end{aligned}$$

where

$$m_1 = -3(d-2)q_0 M^2 a_0^2 \sigma^{2\alpha} \text{ and } m_2 = -(d-2)M^3 a_0^3 \sigma^{3\alpha}.$$

Finally, we derive

$$\mathcal{P}(v^+) < 0,$$

provided that  $\xi \gg 1$ ,  $b \leq b_1 \ll 1$  and  $\eta \ll 1$ .

+ For  $\xi = O(1)$ , i.e.,  $\xi \in [0, K]$  for some  $K > 0$  large enough: thanks to the smallness of  $b$ , the dominating term in  $v^+$  is  $-Mb^{\frac{\eta}{4}}(\tau)H_0(\xi) > 0$ . Besides that, we have

$$\partial_\sigma (2Q_\sigma + \xi^2 Q_\sigma^2) = -\frac{1}{\sigma} \Lambda_\xi Q_\sigma - \xi^2 \frac{Q_\sigma}{\sigma} \Lambda_\xi Q_\sigma = -\frac{\Lambda_\xi Q_\sigma}{\sigma} (1 + \xi^2 Q_\sigma(\xi)).$$

Note that the construction of  $Q$  (see more in (9.10)) ensures  $\xi^2 Q_\sigma(\xi) > -1$  and we derive

$$\partial_\sigma (2Q_\sigma + \xi^2 Q_\sigma^2) > 0,$$

from which we infer the existence of  $m_3(\sigma, K) > 0$  such that

$$2Q + \xi^2 Q^2 - (2Q_\alpha + \xi^2 Q_\alpha^2) \geq m_3 \text{ with } \sigma < 1.$$

Since  $[0, K]$  is compact, we get

$$v^+ = \theta^+ H_1(\xi) - Mb^{\frac{\eta}{4}} H_0 \geq m_4(\sigma, K) b^{\frac{\eta}{4}} \text{ with } m_4 > 0.$$

Thus, we get

$$-3(d-2) [2Q + \xi^2 Q^2 - (2Q_\alpha + \xi^2 Q_\alpha^2)] v^+ \leq -3(d-2)m_3 m_4 b^{\frac{\eta}{4}}.$$

This concludes  $\mathcal{P}(z^+) < 0$  for the case  $\xi = O(1)$ .

- Proof of (ii): Notice that for  $M(\eta)$  large enough and  $\sigma$  small, we have

$$\frac{v^+ \left( \frac{y}{\sqrt{b(\tau_0)}} \right)}{b(\tau_0)} > 0,$$

and from  $\varepsilon(\tau_0)$ 's definition in (5.16), we see that it vanishes when  $y \leq b(\tau_0)$  and it is sufficient to check it for  $y \in [b^{\frac{\delta}{2}}(\tau_0), b^{\frac{\eta}{4}}(\tau_0)]$  giving

$$\frac{y}{\sqrt{b(\tau_0)}} \rightarrow +\infty.$$

Thus, using (11.10) and (11.11), we can find a  $c_0$  such that

$$\frac{v^+ \left( \frac{y}{\sqrt{b(\tau_0)}} \right)}{b(\tau_0)} \geq c_0 b^{\frac{\alpha}{2} + \frac{\eta}{4}}(\tau_0) y^{-\gamma}.$$

On the other hand,  $\varepsilon$ 's formula and the fact that  $y \leq b^{\frac{\eta}{4}}(\tau_0)$  imply

$$\left| \frac{\phi_{\ell, b(\tau_0), \beta(\tau_0)}}{c_{\ell, 0}} - \phi_{0, b(\tau_0), \beta(\tau_0)} \right| \lesssim b^{\frac{\eta}{4}}(\tau_0).$$

This yields

$$|\varepsilon(\tau_0)| \lesssim b^{\frac{\alpha}{2} + \frac{\eta}{2}}(\tau_0)y^{-\gamma}.$$

□

## A. Details on pointwise estimates

In the sequel, we give details to some pointwise estimates used in our paper.

**Lemma A.1.** *Let us consider  $\hat{B}$  defined as in (7.20),  $(\varepsilon, b, \beta)(\tau) \in V[A, \eta, \tilde{\eta}](\tau)$ , for all  $\tau \in [\tau_0, \tau^*]$  for some  $\tau^* \geq \tau_0$ , and  $\delta > \eta > \tilde{\eta}$ . Then, there exists  $\tau_9(A, \delta, \eta, \tilde{\eta}) \geq 1$ , such that for all  $\tau_0 \geq \tau_9$ , the following holds*

$$\begin{aligned} \left| \mathbb{1}_{y \in (0, b^\delta]} \hat{B}(y, \tau) \right| &\leq \frac{Cb^{\frac{\alpha}{2}}}{y^\gamma} + \frac{CA^3 b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau)}{y^{\gamma+2}} + \frac{CA^6 b^{2(\frac{\alpha}{2} + \tilde{\eta})}(\tau)}{y^{2\gamma}} + \frac{CA^9 b^{3(\frac{\alpha}{2} + \tilde{\eta})}(\tau)}{y^{3\gamma-2}}, \\ \left| \mathbb{1}_{y \geq b^\delta} \hat{B}(y, \tau) \right| &\leq C \frac{b^{\frac{\alpha}{2} + 4\eta} \langle y \rangle^{2\ell+2}}{y^\gamma} + \frac{Cb^{\alpha+\delta(1-\gamma)} \langle y \rangle^{4\ell+8}}{y^\gamma} + \frac{Cb^{\frac{3\alpha}{2} - 2\gamma\delta} \langle y \rangle^{6\ell+14}}{y^\gamma}, \forall \tau \in [\tau_0, \tau^*]. \end{aligned}$$

*Proof.* Let us consider  $\delta \gg \eta \gg \tilde{\eta}$ . First, since  $(\varepsilon, b, \beta)(\tau) \in V_\ell[A, \eta, \tilde{\eta}](\tau)$  and by applying Lemma 6.1 we have

$$|b'(\tau)| \lesssim Ab^{4\eta}(\tau) \quad \text{and} \quad \left| \frac{b'}{b} - 2\beta \left( 1 - \frac{2\ell}{\alpha} \right) \right| \lesssim Ab^{4\eta}(\tau).$$

Write

$$\hat{\varepsilon}_{\beta, j} = \|\phi_{j, \infty, \beta}\|_{L_{\rho_\beta}^2}^{-2} \langle \varepsilon, \phi_{j, \infty, \beta} \rangle_{L_{\rho_\beta}^2}. \quad (\text{A.1})$$

We observe that even though  $\phi_{j, b, \beta}$  is not orthogonal to  $\phi_{k, \infty, \beta}$ ,  $k \neq j$ , we have

$$\left| \langle \phi_{j, b, \beta}, \phi_{k, \infty, \beta} \rangle_{L_{\rho_\beta}^2} \right| \leq \int_{y \leq b^\delta} |\phi_{j, b, \beta} \phi_{k, \infty, \beta}| \rho_\beta dy + \int_{y \geq b^\delta} |\phi_{j, b, \beta} \phi_{k, \infty, \beta}| \rho_\beta dy \lesssim b^\delta.$$

Thus, we use pointwise estimates in Lemma 5.2 to obtain

$$|\hat{\varepsilon}_{\beta, j}(\tau)| \leq CA b^{\frac{\alpha}{2} + \tilde{\eta}}, \forall j < \ell, \quad \hat{\varepsilon}_{\beta, \ell} = \frac{\varepsilon_\ell}{c_{\ell, 0}} + O(b^{\frac{\alpha}{2} + 4\eta}), \quad \text{and} \quad \hat{\varepsilon}_{\beta, 0} = -\varepsilon_\ell + O(b^{\frac{\alpha}{2} + 4\eta}). \quad (\text{A.2})$$

In particular, repeating the technique in Lemma 6.1 we get

$$\begin{cases} \partial_\tau \hat{\varepsilon}_{\beta, \ell} &= 2\beta \left( \frac{\alpha}{2} - \ell \right) \hat{\varepsilon}_{\beta, \ell} + O(b^{\frac{\alpha}{2} + 4\eta}), \\ \partial_\tau \hat{\varepsilon}_{\beta, 0} &= 2\beta \frac{\alpha}{2} \hat{\varepsilon}_{\beta, 0} + \left[ \frac{b'}{b} - 2\beta \right] m_0 b^{\frac{\alpha}{2}} + O(b^{\frac{\alpha}{2} + 4\eta}). \end{cases} \quad (\text{A.3})$$

- The first case:  $y \in (0, b^\delta(\tau)]$ . From (7.20), we have

$$\begin{aligned} &\left| \hat{B}(\hat{\varepsilon}_{\beta, +} + \hat{\varepsilon}_{\beta, -}) \right| \\ &\leq \left| 3(d-2) \left( 2Q_b + y^2 Q_b^2 + \frac{1}{y^2} \right) (\hat{\varepsilon}_{\beta, +} + \hat{\varepsilon}_{\beta, -}) \right| + |B(\hat{\varepsilon}_{\beta, +} + \hat{\varepsilon}_{\beta, -})| + \left| \Phi + \mathcal{L}_\infty^\beta \hat{\varepsilon}_{\beta, +} - \partial_\tau \hat{\varepsilon}_{\beta, +} \right|. \end{aligned}$$

From  $Q$ 's asymptotic given in Lemma 9.1 and (2.15), we get

$$\left| 2Q_b + y^2 Q_b + \frac{1}{y^2} \right| \lesssim y^{-2}.$$

Besides that, since  $\hat{\varepsilon}_{\beta, +} + \hat{\varepsilon}_{\beta, -} = \varepsilon = \varepsilon_+ + \varepsilon_-$ , and the pointwise estimates in Lemma 5.2, we obtain

$$|\hat{\varepsilon}_{\beta, +} + \hat{\varepsilon}_{\beta, -}| \leq C \left( A^4 b^{\frac{\alpha}{2} + \tilde{\eta}}(\tau) y^{-\gamma} + b^{\frac{\alpha}{2}} y^{2-\gamma} \right),$$

Thus, the following is valid

$$\left| 3(d-2) \left( 2Q_b + y^2 Q_b^2 + \frac{1}{y^2} \right) (\hat{\varepsilon}_{\beta,+} + \hat{\varepsilon}_{\beta,-}) \right| \leq C \left( A^4 b^{\frac{\alpha}{2} + \tilde{\eta}} y^{-2-\gamma} + b^{\frac{\alpha}{2}}(\tau) y^{-\gamma} \right).$$

Similarly, we have

$$|B(\hat{\varepsilon}_{\beta,+} + \hat{\varepsilon}_{\beta,-})| \leq C (|\varepsilon|^2 + y^2 |\varepsilon|^3) \leq C \left( A^8 b^{\alpha+2\tilde{\eta}} y^{-2\gamma} + A^{12} b^{\frac{3}{2}\alpha+3\tilde{\eta}} y^{2-3\gamma} \right).$$

For the last term, we immediately deduce from (A.2) and (A.3) that

$$\left| \Phi + \mathcal{L}_\infty^\beta \hat{\varepsilon}_{\beta,+} - \partial_\tau \hat{\varepsilon}_{\beta,+} \right| \leq C b^{\frac{\alpha}{2}} y^{-\gamma}.$$

By adding all related terms, we conclude the estimate on  $\mathbb{1}_{y \in (0, b^\delta]} \hat{B}$ .

- The second case:  $y \in [b^\delta(\tau) + \infty)$ . Regarding (6.15), we can improve it as follows

$$\Phi = \left[ \frac{b'}{b} - 2\beta \right] m_0 b^{\frac{\alpha}{2}} \phi_{0,\infty,\beta} + \tilde{\Phi}(y, \tau),$$

where

$$\left| \tilde{\Phi}(y, \tau) \right| \leq C b^{\frac{\alpha}{2} + \delta} y^{-\gamma} \langle y \rangle^{2\ell+2}, \forall y \geq b^\delta(\tau).$$

In addition to that, we have

$$\begin{aligned} \left| \mathcal{L}_\infty^\beta \hat{\varepsilon}_{\beta,+} - \sum_{j=0}^{\ell} \left( \frac{\alpha}{2} - j \right) \hat{\varepsilon}_{\beta,j} \phi_{j,\infty,\beta} \right| &\lesssim b^{\frac{\alpha}{2} + 4\eta} \langle y \rangle^{2\ell+2} y^{-\gamma}, \\ \left| \partial_\tau \hat{\varepsilon}_{\beta,+} - \sum_{j=0}^{\ell} \left( \frac{\alpha}{2} - j \right) \hat{\varepsilon}_{\beta,j} \phi_{j,\infty,\beta} \right| &\lesssim b^{\frac{\alpha}{2} + 4\eta} \langle y \rangle^{2\ell+2} y^{-\gamma}. \end{aligned}$$

Finally, we conclude

$$\left| \Phi + \mathcal{L}_\infty^\beta \hat{\varepsilon}_{\beta,+} - \partial_\tau \hat{\varepsilon}_{\beta,+} \right| \leq \frac{C A b^{\frac{\alpha}{2} + 4\eta}(\tau) \langle y \rangle^{2\ell+2}}{y^\gamma}.$$

Next, we study the estimate involving  $Q_b$ . Using (9.3) and the facts that  $y \geq b^\delta, \delta \ll 1$ , we get

$$\xi = \frac{y}{\sqrt{b}} \gg 1,$$

then, it follows

$$Q_b(y) = -\frac{1}{y^2} \left( 1 + \tilde{Q} \right),$$

where

$$\left| \tilde{Q}(y) \right| \leq C(\delta) b^{\frac{\alpha}{2}} y^{2-\gamma} \leq b^{\frac{\alpha}{2} - \gamma \delta} \leq C b^\delta(\tau), y \geq b^\delta.$$

Hence, we have

$$\left| 2Q_b(y) + y^2 Q_b^2(y) + \frac{1}{y^2} \right| \leq C b^\delta,$$

which implies

$$\left| \left( 2Q_b(y) + y^2 Q_b^2(y) + \frac{1}{y^2} \right) (\hat{\varepsilon}_{\beta,+} + \hat{\varepsilon}_{\beta,-}) \right| \leq C b^{\frac{\alpha}{2} + \delta} \langle y \rangle^{2\ell+2} y^{-\gamma}.$$

Similarly, since

$$|1 + y^2 Q_b(y)| \leq C b^{\frac{\alpha}{2}}(\tau) y^{2-\gamma} \leq C b^\delta, \forall y \geq b^\delta(\tau),$$



we deduce

$$\begin{aligned} |B(\hat{\varepsilon}_{\beta,+} + \hat{\varepsilon}_{\beta,-})| &\lesssim b^\delta \left( \frac{b^\alpha y^4 \langle y \rangle^{4\ell+4}}{y^{2\gamma}} + \frac{A^8 b^{\alpha+2\tilde{\eta}} \langle y \rangle^{4\ell+4}}{y^{2\gamma}} \right) + y^2 \left( \frac{b^{\frac{3\alpha}{2}} y^6 \langle y \rangle^{6\ell+6}}{y^{3\gamma}} + \frac{A^{12} b^{\frac{3\alpha}{2}+3\tilde{\eta}} \langle y \rangle^{6\ell+6}}{y^{3\gamma}} \right) \\ &\lesssim b^{\alpha+\delta} \langle y \rangle^{2\ell+8} y^{-2\gamma} + b^{\frac{3\alpha}{2}} \langle y \rangle^{6\ell+14} y^{-3\gamma}. \end{aligned}$$

In particular, once  $y \geq b^\delta(\tau')$ , it follows  $\frac{1}{y^\gamma} \lesssim b^{-\gamma\delta}$ . By adding the related bounds, we conclude the estimate on  $\mathbb{1}_{y \geq b^\delta} \hat{B}$ . This achieves the proof of the Lemma.  $\square$

## B. Detail on spectral analysis computation of $\mathcal{L}_\infty$

In this part, we aim to give a complete computation to formulate constant in Proposition 4.1. Let us consider the following quadratic equation

$$\gamma^2 - d\gamma + 3(d-2) = 0. \quad (\text{B.1})$$

The equation has two distinct solutions

$$\begin{cases} \gamma_1 &= \frac{1}{2}(d - \sqrt{d^2 - 12d + 24}), \\ \gamma_2 &= \frac{1}{2}(d + \sqrt{d^2 - 12d + 24}). \end{cases}$$

We remark that  $\frac{1}{r^{\gamma_2}}$  does not belong to  $H_\rho^1$ , but  $\frac{1}{r^{\gamma_1}}$  does. In addition, we also define

$$\gamma = \gamma_1 = \frac{1}{2}(d - \sqrt{d^2 - 12d + 24}), \text{ and } \tilde{\gamma} = \gamma_2 = \frac{1}{2}(d + \sqrt{d^2 - 12d + 24}).$$

From  $\gamma$ 's formula above, we can get the first eigenfunction and eigenvalue as follows

$$\phi_{0,\beta,\infty}(r) = \frac{1}{r^\gamma} \text{ and } \lambda_{0,\beta,\infty} = 2\beta \left( \frac{1}{2}(\gamma - 2) \right) := 2\beta \left( \frac{\alpha}{2} \right).$$

Following [9] (also [8], and [7]), we search the eigenfunctions and eigenvalues in the following forms

$$\phi_{i,\beta,\infty}(r) = \sum_{j=0}^i a_{i,j} (2\beta)^j r^{2j-\gamma}, \text{ and } a_{i,i} = 1, \text{ and } \lambda_{i,\beta,\infty} = 2\beta \left( \frac{\alpha}{2} - i \right), \forall i \geq 0. \quad (\text{B.2})$$

Plugging the form (B.2) into the following relation

$$\mathcal{L}_\infty^\beta \phi_{i,\beta,\infty} = 2\beta \left( \frac{\alpha}{2} - i \right) \phi_{i,\beta,\infty}, \quad (\text{B.3})$$

we get

$$\begin{aligned} \mathcal{L}_\infty^\beta \phi_{i,\beta,\infty} &= \sum_{j=0}^i a_{i,j} (2\beta)^j A_j r^{2(j-1)-\gamma} + \sum_{j=0}^i a_{i,j} (2\beta)^j (-2\beta - \beta(2j - \gamma)) r^{2j-\gamma} \\ &= 2\beta \left( \frac{\alpha}{2} - i \right) \sum_{j=0}^i a_{i,j} (2\beta)^j r^{2j-\gamma}, \end{aligned} \quad (\text{B.4})$$

where  $A_j = (2j - \gamma)(2j - \gamma - 1) + (d + 1)(2j - \gamma) + 3(d - 2)$ . Fix  $0 \leq j \leq i - 1$ .

+ For  $j = i$ : We choose  $a_{i,i} = 1$ , then, we get

$$(-2\beta - \beta(2i - \gamma)) = 2\beta \left( \frac{\alpha}{2} - i \right),$$

then, (B.4) is satisfied.

+ For all  $j \leq i - 1$ : (B.4) yields

$$a_{i,j+1} (2\beta)^{j+1} A_{j+1} + a_{i,j} (2\beta)^j (-2\beta - \beta(2j - \gamma)) = 2\beta \left( \frac{\alpha}{2} - i \right) a_{i,j} (2\beta)^j,$$

which yields

$$a_{i,j+1}A_{j+1} = (j-i)a_{i,j}. \quad (\text{B.5})$$

By a simple recurrence, we obtain

$$a_{i,j} = (-1)^{i-j} \prod_{k=j}^{i-1} \frac{A_{k+1}}{i-k} = \frac{(-1)^{i-j}}{(i-j)!} \prod_{k=j+1}^i A_k.$$

Thus

$$a_{i,j} = \frac{(-1)^j}{(i-j)!} \times \prod_{k=j+1}^{i-1} ((2k-\gamma)(2k-\gamma-1) + (d+1)(2k-\gamma) + 3(d-2)).$$

In particular, in (B.4), the order  $r^{-\gamma-2}$  remains. However, its coefficient is equal to 0, since  $\gamma$  solves (B.1). Finally, (B.3) is completely satisfied by the choice of  $a_{i,j}$  above. Next, we aim to decompose  $a_{i,j}$  as follows: First, we deduce from (B.1) that

$$\begin{aligned} & (2k-\gamma)(2k-\gamma-1) + (d+1)(2k-\gamma) + 3(d-2) \\ &= 2k \cdot 2k + 2kd - 4k\gamma = 2k(2k+d-2\gamma) = 4k \left( \frac{d}{2} - \gamma + k \right). \end{aligned}$$

Then,  $a_{i,j}$  is decomposed as

$$a_{i,j} = \frac{(-1)^{i-j}}{(i-j)!} \prod_{k=j+1}^i 4k \left( \frac{d}{2} - \gamma + k \right) = \frac{(-1)^{i-j}}{(i-j)!} 4^{i-j} \frac{i!}{j!} \frac{(\frac{d}{2} - \gamma)_i}{(\frac{d}{2} - \gamma)_j} = c_{i,j} C_j,$$

where  $c_{i,j} = \frac{(-1)^{i-j} 4^{i-j} i! (\frac{d}{2} - \gamma)_i!}{(i-j)!}$ ,  $C_j = \frac{1}{4^j j! (\frac{d}{2} - \gamma)_j!}$ , and

$$\left( \frac{d}{2} - \gamma \right)_i = \left( \frac{d}{2} - \gamma + 1 \right) \left( \frac{d}{2} - \gamma + 2 \right) \dots \left( \frac{d}{2} - \gamma + i \right) \text{ and } \left( \frac{d}{2} - \gamma \right)_0 = 1.$$

## C. Poisson kernel for Laguerre expansions

In this part, we aim to provide some pointwise estimates involving semi-group  $e^{\tau \mathcal{L}_\infty}$  with  $\mathcal{L}_\infty$  defined as in (2.22). Recall that for  $f \in L^1(\mathbb{R}_+, x^\omega e^{-x} dx)$ , we have the following presentation

$$\left[ e^{(\tau-\tau_0)\mathcal{L}_\infty} \right] f(y, \tau) = 2^{\omega+1} y^{-\gamma} e^{\frac{\alpha}{2}(\tau-\tau_0)} \int_0^\infty P_{\frac{\omega}{2}} \left( \frac{y^2}{4}, \frac{x^2}{4}, e^{-(\tau-\tau_0)} \right) [f(x)x^\gamma] x^{\omega+1} e^{-\frac{x^2}{4}} dx. \quad (\text{C.1})$$

where  $P_\zeta$  is defined by

$$\begin{aligned} P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) &= e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \frac{\left( -r \frac{x^2 y^2}{16} \right)^{-\frac{\zeta}{2}}}{1-r} J_\zeta \left( \frac{2 \left( -r \frac{x^2 y^2}{16} \right)^{\frac{1}{2}}}{1-r} \right), \\ &= \frac{\left( \sqrt{r \frac{y^2}{4} \frac{x^2}{4}} \right)^{-\zeta}}{1-r} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} i^\zeta J_\zeta \left( \frac{2r^{\frac{1}{2}} \frac{y}{2} \frac{x}{2} i}{1-r} \right) \\ &= \frac{4^\zeta}{\sqrt{r}^\zeta (1-r)(yx)^\zeta} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \mathbf{I}_\zeta \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right), \end{aligned}$$

and  $\mathbf{I}_\zeta$  is the function of imaginary argument corresponding to

$$\mathbf{I}_\zeta(z) = \frac{\left( \frac{1}{2} z \right)^\zeta}{\Gamma(\zeta + \frac{1}{2}) \Gamma(\frac{1}{2})} \int_0^\pi \cosh(z \cos \theta) \sin^{2\zeta}(\theta) d\theta, \quad (\text{C.2})$$

provided that  $\text{Re}(\zeta + \frac{1}{2}) > 0$ , the reader can check the formula at page 79, formula (2) in [34]. We have the following result

**Lemma C.1** (Maximal estimate, [4]). *Let us consider  $f \in L^2_\rho$  with  $\rho = \rho_{\frac{1}{2}}$  defined in (2.25), then,*

$$\left| e^{(\tau-\tau_0)\mathcal{L}_\infty} f(y) \right| \leq C y^{-\gamma} e^{\frac{\alpha}{2}(\tau-\tau_0)} [Mf](y), \forall y > 0, \tau > \tau_0, \quad (\text{C.3})$$

where  $\alpha$  and  $\gamma$  were defined in (2.27) and (2.26), respectively, and  $Mf$  is given by

$$Mf(y) = \sup_{y \in \mathcal{I}} \frac{\int_{\mathcal{I}} |f(y')(y')^\gamma| (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'}{\int_{\mathcal{I}} (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'}, \omega = \sqrt{d^2 - 12d + 24}, \quad (\text{C.4})$$

here the supremum is taken over all sub-intervals  $\mathcal{I}$  containing  $y$ . In particular, if  $|f(y)y^\gamma|$  is a non decreasing function, then the supremum in (C.4) is attained by  $\mathcal{I} = [y, +\infty)$ . Otherwise, if  $|f(y)y^\gamma|$  is a non increasing, then the supremum is attained by  $\mathcal{I} = [0, y]$ .

*Proof.* The proof is quite the same as for Lemma VI.2 in [4].  $\square$

Next, we will estimate the growth of the action  $e^{(\tau-\tau')\mathcal{L}_\infty}$  to  $\Lambda$ :

**Lemma C.2.** *Let us consider  $f \in L^2_\rho$  and  $\Lambda f \in L^2_\rho$  where  $\Lambda$  was defined in (2.14). Assume further more that*

$$|f(y)| \leq \mathcal{B} \frac{\langle y \rangle^{2\ell+2}}{y^\gamma}, \text{ for some } \mathcal{B} \in \mathbb{R}_+^*. \quad (\text{C.5})$$

Then, it holds that for  $\tau > \tau'$

$$\left| e^{(\tau-\tau')\mathcal{L}_\infty} (\Lambda f) \right| \lesssim e^{\tau-\tau'} \mathcal{B} \langle y \rangle^{2\ell+3} y^{-\gamma}. \quad (\text{C.6})$$

*Proof.* Recall from (2.14) that  $\Lambda f = y\partial_y f + 2f$  and apply (C.1) in deriving

$$\left[ e^{(\tau-\tau')\mathcal{L}_\infty} \right] \Lambda f(y) = 2^{\omega+1} y^{-\gamma} e^{\frac{\alpha}{2}(\tau-\tau')} \int_0^\infty P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) (x\partial_x f + 2f) x^{\omega+1+\gamma} e^{-\frac{x^2}{4}} dx,$$

where  $r = e^{-(\tau-\tau')}$  and  $\zeta = \frac{\omega}{2}$ . First, Lemma C.1 results in

$$\left| 2^{\omega+1} y^{-\gamma} e^{\frac{\alpha}{2}(\tau-\tau')} \int_0^\infty P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) f(x) x^{\omega+1+\gamma} e^{-\frac{x^2}{4}} dx \right| \lesssim \mathcal{B} e^{\frac{\alpha}{2}(\tau-\tau')} \langle y \rangle^{2\ell+2} y^{-\gamma}.$$

Then, it is sufficient to prove that

$$|I(y)| \lesssim \mathcal{B} \langle y \rangle^{2\ell+3} \text{ where } I = \int_0^\infty P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) (x\partial_x f(x)) x^{\omega+1+\gamma} e^{-\frac{x^2}{4}} dx.$$

Using the integration by parts provided that the functions go to 0 at  $+\infty$  and 0, we get

$$\begin{aligned} I &= - \int_0^\infty f(x) \partial_x \left( P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) x^{\omega+2+\gamma} e^{-\frac{x^2}{4}} \right) dx \\ &= - \int_0^\infty f(x) \partial_x \left( P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) \right) x^{\omega+2+\gamma} e^{-\frac{x^2}{4}} dx. \\ &\quad - (\omega + 2 + \gamma) \int_0^\infty f(x) P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) x^{\omega+1+\gamma} e^{-\frac{x^2}{4}} dx \\ &\quad + \frac{1}{2} \int_0^\infty f(x) P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) x^{\omega+3+\gamma} e^{-\frac{x^2}{4}} dx. \end{aligned}$$

We now explicitly compute

$$\begin{aligned}
\partial_x \left( P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) \right) &= \partial_x \left( \frac{4^\zeta}{\sqrt{r}^\zeta (1-r)(yx)^\zeta} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \mathbf{I}_\zeta \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right) \right) \\
&= \partial_x \left( \frac{4^\zeta}{\sqrt{r}^\zeta (1-r)(yx)^\zeta} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \right) \mathbf{I}_\zeta \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right) \\
&\quad + \frac{4^\zeta}{\sqrt{r}^\zeta (1-r)(yx)^\zeta} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \partial_x \left( \mathbf{I}_\zeta \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right) \right) \\
&= \partial_x \left( \frac{4^\zeta}{\sqrt{r}^\zeta (1-r)(yx)^\zeta} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \right) \mathbf{I}_\zeta \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right) \\
&\quad + \frac{4^\zeta}{x \sqrt{r}^\zeta (1-r)(yx)^\zeta} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \frac{r^{\frac{1}{2}} yx}{2(1-r)} \mathbf{I}'_\zeta \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right),
\end{aligned}$$

from equality (3) at page 79 in [34]

$$\frac{r^{\frac{1}{2}} yx}{2(1-r)} \mathbf{I}'_\zeta \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right) = \zeta \mathbf{I}_\zeta \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right) + \frac{r^{\frac{1}{2}} yx}{2(1-r)} \mathbf{I}_{\zeta+1} \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right),$$

we infer

$$\begin{aligned}
\partial_x \left( P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) \right) &= \partial_x \left( \frac{4^\zeta}{\sqrt{r}^\zeta (1-r)(yx)^\zeta} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \right) \mathbf{I}_\zeta \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right) \\
&\quad + \frac{4^\zeta}{x \sqrt{r}^\zeta (1-r)(yx)^\zeta} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \zeta \mathbf{I}_\zeta \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right) \\
&\quad + \frac{4^\zeta}{x \sqrt{r}^\zeta (1-r)(yx)^\zeta} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \frac{r^{\frac{1}{2}} yx}{2(1-r)} \mathbf{I}_{\zeta+1} \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right).
\end{aligned}$$

Besides, we have

$$\begin{aligned}
\partial_x \left( \frac{4^\zeta}{\sqrt{r}^\zeta (1-r)(yx)^\zeta} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \right) &= \frac{4^\zeta (-\zeta)}{\sqrt{r}^\zeta (1-r)(yx)^\zeta x} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \\
&\quad - \frac{r}{1-r} \frac{x}{2} \frac{4^\zeta}{\sqrt{r}^\zeta (1-r)(yx)^\zeta} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)}.
\end{aligned}$$

At final, we arrive to

$$\begin{aligned}
x \partial_x \left( P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) \right) &= -\frac{r}{1-r} \frac{x^2}{2} \frac{4^\zeta}{\sqrt{r}^\zeta (1-r)(yx)^\zeta} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \mathbf{I}_\zeta \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right) \\
&\quad + \frac{4^\zeta}{\sqrt{r}^\zeta (1-r)(yx)^\zeta} \frac{r^{\frac{1}{2}} yx}{2(1-r)} e^{-\frac{r}{1-r} \left( \frac{y^2}{4} + \frac{x^2}{4} \right)} \mathbf{I}_{\zeta+1} \left( \frac{r^{\frac{1}{2}} yx}{2(1-r)} \right) \\
&= -\frac{rx^2}{2(1-r)} P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) + \frac{ry^2 x^2}{8(1-r)} P_{\zeta+1} \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right).
\end{aligned}$$

Plugging into  $I$ 's formula, we get

$$\begin{aligned}
I &= \frac{r}{2(1-r)} \int_0^\infty f(x) P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) x^{\omega+3+\gamma} e^{-\frac{x^2}{4}} \\
&\quad - \frac{ry^2}{8(1-r)} \int_0^\infty f(x) P_{\zeta+1} \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) x^{\omega+3+\gamma} e^{-\frac{x^2}{4}} \\
&\quad - (\omega+2+\gamma) \int_0^\infty f(x) P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) x^{\omega+1+\gamma} e^{-\frac{x^2}{4}} \\
&\quad + \frac{1}{2} \int_0^\infty f(x) P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) x^{\omega+3+\gamma} e^{-\frac{x^2}{4}} \\
&= \int_0^\infty \left[ \frac{P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right)}{2(1-r)} - \frac{ry^2}{8(1-r)} P_{\zeta+1} \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) \right] f(x) x^{\omega+3+\gamma} e^{-\frac{x^2}{4}} dx \\
&\quad - (\omega+2+\gamma) \int_0^\infty f(x) P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) x^{\omega+1+\gamma} e^{-\frac{x^2}{4}} dx.
\end{aligned} \tag{C.7}$$

It also follows from Lemma C.1 that

$$\left| \int_0^\infty f(x) P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) x^{\omega+1+\gamma} e^{-\frac{x^2}{4}} dx \right| \lesssim M(f),$$

and from (C.5), we have

$$M(f)(y) = \sup_{y \in \mathcal{I}} \frac{\int_{\mathcal{I}} |f(y')(y')^\gamma| (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'}{\int_{\mathcal{I}} (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'} \lesssim \mathcal{B} \frac{\int_y^\infty \langle y' \rangle^{2\ell+2} (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'}{\int_y^\infty (y')^{1+\omega} e^{-\frac{(y')^2}{4}} dy'} \tag{C.8}$$

$$\lesssim \mathcal{B} \langle y \rangle^{2\ell+2}. \tag{C.9}$$

Now, it remains to prove the following:

$$|I_1(y)| \lesssim \frac{\mathcal{B}}{\sqrt{1-r}} \langle y \rangle^{2\ell+3}, \tag{C.10}$$

where

$$I_1(y) = \int_0^\infty \left[ \frac{P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right)}{2(1-r)} - \frac{ry^2}{8(1-r)} P_{\zeta+1} \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) \right] f(x) x^{\omega+3+\gamma} e^{-\frac{x^2}{4}} dx$$

We recall formulae (2) at page 77 and formula (2) at page 203 in [34] regarding the function  $I_\zeta$  of imaginary argument

$$\begin{aligned}
cz^\zeta &\leq \mathbf{I}_\zeta(z) \leq Cz^\zeta && \text{if } z \in [0, 1], \\
cz^{-\frac{3}{2}}e^z &\leq \left| \mathbf{I}_\zeta(z) - \frac{e^z}{\sqrt{2\pi z}} \right| \leq Cz^{-\frac{3}{2}}e^z && \text{if } z \in [1, +\infty),
\end{aligned} \tag{C.11}$$

In particular, (C.11) implies

$$C^{-1}H_\zeta(y, x, r) \leq P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) \leq CH_\zeta(y, x, r), \tag{C.12}$$

where

$$H_\zeta(y, x, r) = \begin{cases} (1-r)^{-\zeta-1} e^{-\frac{r(\frac{y^2}{4} + \frac{x^2}{4})}{1-r}} & \text{if } x \in \left[ 0, \frac{2(1-r)}{\sqrt{ry}} \right] \\ \frac{(4r)^{-\frac{\zeta}{2} - \frac{1}{4}} (yx)^{-\zeta - \frac{1}{2}}}{e(\sqrt{1-r})} e^{-\frac{-r\frac{y^2}{4} + \frac{1}{2}(r)\frac{1}{2}(yx) - r\frac{x^2}{4}}{1-r}} & \text{if } z \in \left[ \frac{2(1-r)}{\sqrt{ry}}, +\infty \right) \end{cases}. \tag{C.13}$$

Using (C.5), we estimate

$$\begin{aligned} & |I_1(y, \tau, \tau')| \\ & \lesssim \mathcal{B} \left( \int_0^{\frac{2(1-r)}{\sqrt{ry}}} \left| \frac{P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right)}{2(1-r)} - \frac{ry^2}{8(1-r)} P_{\zeta+1} \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) \right| (1+x^{2\ell+2}) x^{\omega+3+\gamma} e^{-\frac{x^2}{4}} dx \right. \\ & \left. + \int_0^{\frac{2(1-r)}{\sqrt{ry}}} \left| \frac{P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right)}{2(1-r)} - \frac{ry^2}{8(1-r)} P_{\zeta+1} \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) \right| (1+x^{2\ell+2}) x^{\omega+3+\gamma} e^{-\frac{x^2}{4}} dx \right). \end{aligned}$$

+ For the integral on  $\left[0, \frac{2(1-r)}{\sqrt{ry}}\right]$ , we use the first asymptotic in (C.13) to obtain

$$\begin{aligned} & \int_0^{\frac{2(1-r)}{\sqrt{ry}}} \left| \frac{P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right)}{2(1-r)} - \frac{ry^2}{8(1-r)} P_{\zeta+1} \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) \right| (1+x^{2\ell+2}) x^{\omega+3+\gamma} e^{-\frac{x^2}{4}} dx \\ & \leq C(1-r)^{-\zeta-2} \int_0^{\frac{2(1-r)}{\sqrt{ry}}} \left(1 + \frac{ry^2}{(1-r)}\right) e^{-\frac{r}{1-r} \left(\frac{y^2}{4} + \frac{x^2}{4}\right)} (1+x^{2\ell+2}) x^{\omega+3} e^{-\frac{x^2}{4}} dx. \end{aligned}$$

On the one hand, once  $r \in (0, \frac{1}{4})$ , it immediately follows that

$$\begin{aligned} & (1-r)^{-\zeta-2} \int_0^{\frac{2(1-r)}{\sqrt{ry}}} \left(1 + \frac{ry^2}{(1-r)}\right) e^{-\frac{r}{1-r} \left(\frac{y^2}{4} + \frac{x^2}{4}\right)} (1+x^{2\ell+2}) x^{\omega+3} e^{-\frac{x^2}{4}} dx \\ & \leq C \int_0^{\frac{2(1-r)}{\sqrt{ry}}} (1+x^{2\ell+2}) x^{\omega+3} e^{-\frac{x^2}{4} - \frac{rx^2}{4(1-r)}} dx \leq C \int_0^\infty (1+x^{2\ell+2}) x^{\omega+3} e^{-\frac{x^2}{4}} dx \leq C, \end{aligned}$$

where  $C$  is independent of  $r$ . On the other hand, once  $r \geq \frac{1}{4}$  and by a change of variable  $z = \frac{\sqrt{r}}{2\sqrt{1-r}}x$  we obtain

$$\begin{aligned} & \int_0^{\frac{2(1-r)}{\sqrt{ry}}} \left(1 + \frac{ry^2}{(1-r)}\right) e^{-\frac{r}{1-r} \left(\frac{y^2}{4} + \frac{x^2}{4}\right)} (1+x^{2\ell+2}) x^{\omega+3} e^{-\frac{x^2}{4}} dx \\ & \leq C e^{-\frac{ry^2}{4(1-r)}} \int_0^{\frac{\sqrt{1-r}}{y}} \left(1 + \frac{ry^2}{(1-r)}\right) e^{-z^2(1+\frac{1-r}{r})} \left(1 + \frac{(1-r)^{\ell+1}}{r^{\ell+1}} z^{2\ell+2}\right) \frac{(1-r)^{\frac{\omega}{2}+2}}{r^{\frac{\omega}{2}+2}} z^{\omega+3} dz \\ & \leq C(1-r)^{\frac{\omega}{2}+2} e^{-\frac{r}{4(1-r)}y^2} \left(1 + \frac{ry^2}{(1-r)}\right) \int_0^{\frac{\sqrt{1-r}}{y}} e^{-z^2(1+\frac{1-r}{r})} (1+(1-r)^{\ell+1} z^{2\ell+2}) z^{\omega+3} dz \\ & \leq (1-r)^{\zeta+2} e^{-\frac{r}{4(1-r)}y^2} \int_0^\infty e^{-z^2(1+z^{2\ell+2+\omega+3})} dz \leq C(1-r)^{\zeta+2}, \text{ with } \zeta = \frac{\omega}{2}. \end{aligned}$$

Finally, we obtain

$$\left| \int_0^{\frac{2(1-r)}{\sqrt{ry}}} \left| \frac{P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right)}{2(1-r)} - \frac{ry^2}{8(1-r)} P_{\zeta+1} \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) \right| (1+x^{2\ell+2}) x^{\omega+3+\gamma} e^{-\frac{x^2}{4}} dx \right| \lesssim \frac{\langle y \rangle^{2\ell+3}}{\sqrt{1-r}}.$$

- For the integral on  $\left[\frac{2(1-r)}{\sqrt{ry}}, +\infty\right)$ , we apply (C.11) and (C.13) and noticing that  $z = \frac{\sqrt{ryx}}{2(1-r)} \geq 1$

$$\frac{x}{2(1-r)} P_\zeta \left( \frac{y^2}{4}, \frac{x^2}{4}, r \right) = \frac{4^\zeta}{2\sqrt{r}^\zeta (1-r)(yx)^\zeta} \left( \frac{x}{2(1-r)} \right) e^{-\frac{r}{4(1-r)}(y^2+x^2)} \left( \frac{e^z}{\sqrt{2\pi z}} + O(e^z z^{-\frac{3}{2}}) \right),$$

and

$$\frac{ry^2x}{8(1-r)}P_{\zeta+1}\left(\frac{y^2}{4}, \frac{x^2}{4}, r\right) = \frac{ry^2x}{8(1-r)}\frac{4^{\zeta+1}}{\sqrt{r}^{\zeta+1}(1-r)(yx)^{\zeta+1}}e^{-\frac{r}{4(1-r)}(y^2+x^2)}\left(\frac{e^z}{\sqrt{2\pi z}} + O(e^z z^{-\frac{3}{2}})\right).$$

Then

$$\begin{aligned} & \left| x\frac{P_{\zeta}\left(\frac{y^2}{4}, \frac{x^2}{4}, r\right)}{2(1-r)} - x\frac{ry^2}{8(1-r)}P_{\zeta+1}\left(\frac{y^2}{4}, \frac{x^2}{4}, r\right) \right| \\ & \leq C\left|\frac{x-\sqrt{ry}}{(1-r)}\right|\frac{4^{\zeta}}{2\sqrt{r}^{\zeta}(1-r)(yx)^{\zeta}}e^{-\frac{r}{4(1-r)}(y^2+x^2)}\frac{e^z}{\sqrt{2\pi z}} \\ & + C\frac{x}{(1-r)}\frac{4^{\zeta}}{2\sqrt{r}^{\zeta}(1-r)(yx)^{\zeta}}e^{-\frac{r}{4(1-r)}(y^2+x^2)}\frac{e^z}{z^{\frac{3}{2}}} \\ & + C\frac{\sqrt{ry}}{(1-r)}\frac{4^{\zeta}}{2\sqrt{r}^{\zeta}(1-r)(yx)^{\zeta}}e^{-\frac{r}{4(1-r)}(y^2+x^2)}\frac{e^z}{z^{\frac{3}{2}}} \\ & \leq Cr^{-\frac{\zeta}{2}-\frac{1}{4}}(yx)^{-\zeta-\frac{1}{2}}\left|\frac{x-\sqrt{ry}}{1-r}\right|e^{\frac{-\frac{ry^2}{4}+\frac{1}{2}\sqrt{ry}x-\frac{rx^2}{4}}{1-r}} \\ & + Cr^{-\frac{\zeta}{2}-\frac{3}{4}}(yx)^{-\zeta-\frac{3}{2}}\left|\frac{x+\sqrt{ry}}{\sqrt{1-r}}\right|e^{\frac{-\frac{ry^2}{4}+\frac{1}{2}\sqrt{ry}x-\frac{rx^2}{4}}{1-r}} \end{aligned}$$

Notice that we have the following identity

$$e^{\frac{-\frac{ry^2}{4}+\frac{1}{2}\sqrt{ry}x-\frac{rx^2}{4}}{1-r}}e^{-\frac{x^2}{4}} = e^{-\frac{1}{4}\left(\frac{x-\sqrt{ry}}{\sqrt{1-r}}\right)^2},$$

and by a change of variable  $z = \frac{x-\sqrt{ry}}{\sqrt{1-r}}$ , we get

$$\begin{aligned} & \int_{\frac{2(1-r)}{\sqrt{ry}}}^{\infty} \left| x\frac{P_{\zeta}\left(\frac{y^2}{4}, \frac{x^2}{4}, r\right)}{2(1-r)} - x\frac{ry^2}{8(1-r)}P_{\zeta+1}\left(\frac{y^2}{4}, \frac{x^2}{4}, r\right) \right| (1+x^{2\ell+2})x^{\omega+2}e^{-\frac{x^2}{4}} dx \\ & \leq C\frac{y^{-\zeta-\frac{1}{2}}r^{-\frac{\zeta}{4}-\frac{1}{4}}}{\sqrt{1-r}}\int_{\frac{2(1-r)}{\sqrt{ry}}}^{\infty} \left|\frac{x-\sqrt{ry}}{1-r}\right| (1+x^{2\ell+2})x^{\zeta+\frac{3}{2}}e^{-\frac{1}{4}\left(\frac{x-\sqrt{ry}}{\sqrt{1-r}}\right)^2} dx \\ & + C\frac{y^{-\zeta-\frac{3}{2}}r^{-\frac{\zeta}{2}-\frac{3}{4}}}{\sqrt{1-r}}\int_{\frac{2(1-r)}{\sqrt{ry}}}^{\infty} \left|\frac{x+\sqrt{ry}}{\sqrt{1-r}}\right| (1+x^{2\ell+2})x^{\zeta+\frac{1}{2}}e^{-\frac{1}{4}\left(\frac{x-\sqrt{ry}}{\sqrt{1-r}}\right)^2} dx \\ & \leq C\frac{(\sqrt{ry})^{-\zeta-\frac{1}{2}}}{\sqrt{1-r}}\int_{\frac{\frac{2(1-r)}{\sqrt{ry}}-\sqrt{ry}}{\sqrt{1-r}}}^{\infty} |z|(1+(z\sqrt{1-r}+\sqrt{ry})^{2\ell+2})|z\sqrt{1-r}+\sqrt{ry}|^{\zeta+\frac{3}{2}}e^{-\frac{z^2}{4}} dz \\ & + C\frac{(\sqrt{ry})^{-\zeta-\frac{3}{2}}}{\sqrt{1-r}}\int_{\frac{\frac{2(1-r)}{\sqrt{ry}}-\sqrt{ry}}{\sqrt{1-r}}}^{\infty} (1+(z\sqrt{1-r}+\sqrt{ry})^{2\ell+2})|z\sqrt{1-r}+\sqrt{ry}|^{\zeta+\frac{3}{2}}e^{-\frac{z^2}{4}} dz. \end{aligned}$$

Now, we have

$$|z|(1+(z\sqrt{1-r}+\sqrt{ry})^{2\ell+2})|z\sqrt{1-r}+\sqrt{ry}|^{\zeta+\frac{3}{2}} \lesssim \langle y \rangle^{2\ell+2} \langle z \rangle^{2\ell+3} \left[ |z\sqrt{1-r}|^{\zeta+\frac{3}{2}} + |\sqrt{ry}|^{\zeta+\frac{3}{2}} \right]$$

and

$$\begin{aligned}
& (\sqrt{ry})^{-\zeta-\frac{1}{2}} \int_{\frac{2(1-r)}{\sqrt{ry}}-\sqrt{ry}}^{\infty} \frac{\langle z \rangle^{2\ell+3}}{\sqrt{1-r}} |z\sqrt{1-r}|^{\zeta+\frac{3}{2}} e^{-\frac{z^2}{4}} dz \\
& \lesssim (\sqrt{ry})^{-\zeta-\frac{1}{2}} \sqrt{1-r}^{\zeta+\frac{3}{2}} \int_{\frac{2(1-r)}{\sqrt{ry}}-\sqrt{ry}}^{\infty} \frac{\langle z \rangle^{2\ell+3+\zeta+\frac{3}{2}}}{\sqrt{1-r}} e^{-\frac{z^2}{4}} dz \\
& \lesssim X^{\zeta+\frac{1}{2}} \int_{2X-\frac{1}{X}}^{\infty} \langle z \rangle^{2\ell+3+\zeta+\frac{3}{2}} e^{-\frac{z^2}{4}} dz \lesssim 1, \text{ with } X = \frac{\sqrt{1-r}}{\sqrt{ry}} > 0,
\end{aligned}$$

yielding

$$\begin{aligned}
& \frac{(\sqrt{ry})^{-\zeta-\frac{1}{2}}}{\sqrt{1-r}} \int_{\frac{2(1-r)}{\sqrt{ry}}-\sqrt{ry}}^{\infty} \frac{|z|(1+(z\sqrt{1-r}+\sqrt{ry})^{2\ell+2})|z\sqrt{1-r}+\sqrt{ry}|^{\zeta+\frac{3}{2}}}{\sqrt{1-r}} e^{-\frac{z^2}{4}} dz \\
& \lesssim \frac{\langle y \rangle^{2\ell+3}}{\sqrt{1-r}}.
\end{aligned}$$

Similarly we have

$$\frac{(\sqrt{ry})^{-\zeta-\frac{3}{2}}}{\sqrt{1-r}} \int_{\frac{2(1-r)}{\sqrt{ry}}-\sqrt{ry}}^{\infty} \frac{(1+(z\sqrt{1-r}+\sqrt{ry})^{2\ell+2})|z\sqrt{1-r}+\sqrt{ry}|^{\zeta+\frac{3}{2}}}{\sqrt{1-r}} e^{-\frac{z^2}{4}} dz \lesssim \frac{\langle y \rangle^{2\ell+3}}{\sqrt{1-r}}.$$

Thus, we obtain

$$\int_{\frac{2(1-r)}{\sqrt{ry}}}^{\infty} \left| x \frac{P_{\zeta}\left(\frac{y^2}{4}, \frac{x^2}{4}, r\right)}{2(1-r)} - x \frac{ry^2}{8(1-r)} P_{\zeta+1}\left(\frac{y^2}{4}, \frac{x^2}{4}, r\right) \right| (1+x^{2\ell+2}) x^{\omega+2} e^{-\frac{x^2}{4}} dx \lesssim \frac{\langle y \rangle^{2\ell+3}}{\sqrt{1-r}}.$$

By adding all related terms, we conclude (C.10) and with it the proof of the Lemma is accomplished.  $\square$

## D. Generators of the Kernel of $H$

We construct the family  $\{T_{i+1}\}$  via the recursive formula

$$T_{i+1} = H^{-1}T_i, \quad \text{and } T_0 = c\Lambda_{\xi}Q \quad (\text{D.1})$$

here  $c$  is some constant which will be chosen later. In other words, we have for  $i \geq 1$

$$T_i = H^{-i}(T_0), \quad i \geq 1.$$

The operator  $H^{-1}$  is explicitly given by

$$H^{-1}f(\xi) = \Lambda Q(\xi) \int_0^{\xi} \frac{Lf(\xi')}{\Lambda Q(\xi')} d\xi', \quad (\text{D.2})$$

with  $L$

$$Lf(\xi) = \frac{1}{\xi^{d+1}\Lambda Q(\xi)} \int_0^{\xi} f(\xi') \Lambda Q(\xi') (\xi')^{d+1} d\xi'. \quad (\text{D.3})$$

We start now our induction argument. For  $i = 0$ , we have by assumption

$$T_0 = c\Lambda Q(\xi).$$

The asymptotic (9.4) yields

$$T_0(\xi) = \begin{cases} -2c + \sum_{i=1}^k a_i'' \xi^{2i} + O(\xi^{2k+2}) & \text{as } \xi \rightarrow 0, \\ a_0 c \xi^{-\gamma} + O(\xi^{-\gamma-2\alpha}) & \text{as } \xi \rightarrow \infty. \end{cases} \quad (\text{D.4})$$



This proves (10.4) for the case  $i = 0$ . Now, let us suppose that (10.4) is true for some  $k$  and let us prove that it holds true for  $k + 1$ .

- At  $\infty$ : we have

$$\mathcal{L}(T_k) = \frac{C_k}{-2\gamma + d + 2 + 2k} \xi^{-\gamma+1+2k} \left( 1 + O\left(\frac{\ln \xi}{\xi^2}\right) \right).$$

Hence

$$T_{k+1} = H^{-1}(T_k) = \frac{C_k}{4(k+1)\left(\frac{d}{2} - \gamma + k + 1\right)} \xi^{-\gamma+2(k+1)} \left( 1 + O\left(\frac{\ln \xi}{\xi^2}\right) \right)$$

The above expansion yields

$$C_{k+1} = \frac{C_k}{4(k+1)\left(\frac{d}{2} - \gamma + k + 1\right)},$$

so that

$$C_k = \frac{ca_0}{4^k k! \left(\frac{d}{2} - \gamma\right)_k!}.$$

Choosing  $c = \frac{1}{a_0}$  concludes the first part of the proof.

In a similar fashion and upon using that  $\partial_\xi^k T_{i+1} = H^{-1} T_i$ , one can establish the mentioned result regarding the asymptotic behavior of the derivatives of  $T_i$ . We omit the details.

## E. Inner eigenfunctions computation

Now we prove Proposition 10.2. Taking into account (10.8), the eigenvalue problem (10.7) reads

$$\begin{aligned} 0 &= \left\{ H - b\beta\Lambda - 2\beta b \left( \frac{\alpha}{2} - i + \tilde{\lambda} \right) \right\} \phi_{i,\text{int}} = \sum_{j=0}^i c_{i,j} (2\beta)^j b^j \left\{ H - b\beta\Lambda - 2\beta b \left( \frac{\alpha}{2} - i + \tilde{\lambda} \right) \right\} T_j \\ &+ \tilde{\lambda} \sum_{j=0}^i c_{i,j} (2\beta)^{j+1} b^{j+1} \left\{ H - b\beta\Lambda - 2\beta b \left( \frac{\alpha}{2} - i + \tilde{\lambda} \right) \right\} T_{j+1} \\ &+ \tilde{\lambda} \sum_{j=0}^i b^{j+1} \left\{ H - b\beta\Lambda - 2\beta b \left( \frac{\alpha}{2} - i + \tilde{\lambda} \right) \right\} S_j \\ &+ b \left\{ H - \beta b\Lambda - 2\beta b \left( \frac{\alpha}{2} - i + \tilde{\lambda} \right) \right\} R_i. \end{aligned}$$

Since  $T_{i+1} = H^{-1} T_i$  and  $H(T_0) = 0$ , we get

$$\begin{aligned} &\sum_{j=0}^i c_{i,j} (2\beta)^j b^j \left\{ H - \beta b\Lambda - 2\beta b \left( \frac{\alpha}{2} - i + \tilde{\lambda} \right) \right\} T_j \\ &= \sum_{j=0}^i c_{i,j} (2\beta)^j b^j \left\{ HT_j - \beta b((2j - \alpha)T_j + \Theta_j) - 2\beta b \left( \frac{\alpha}{2} - i \right) T_j - 2\beta b \tilde{\lambda} T_j \right\} \\ &= -\frac{1}{2} \sum_{j=0}^i c_{i,j} (2\beta)^{2j} b^{j+1} \Theta_j - \tilde{\lambda} \sum_{j=0}^i c_{i,j} (2\beta)^{j+1} b^{j+1} T_j, \end{aligned}$$

where we used the fact that  $c_{i,j}$  defined in (2.28) satisfies  $c_{i,j+1} + c_{i,j}(i-j) = 0$ . Thus, the construction of  $S_j$  and  $R_i$  reduces the following equations (for all  $j \leq i$ )

$$HS_j = 2\beta b \left\{ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) (S_j + c_{i,j}(2\beta)^{j+1}T_{j+1}) \right\}, \quad (\text{E.1})$$

$$HR_i = 2\beta b \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) R_i + \frac{1}{2} \sum_{j=0}^i c_{i,j}(2\beta)^{j+1}b^j\Theta_j. \quad (\text{E.2})$$

We note that the construction of  $S_j$  and  $R_i$  follows [9] (see also [8]) which relies on the Banach fixed point theorem in the functional space  $X_{\xi_0}^a$  for some  $a \in \mathbb{R}$  and  $\xi_0 > 0$  and where the norm is given by

$$\|f\|_{X_{\xi_0}^a} = \sup_{\xi \in [0, \xi_0]} \sum_{i=0}^2 \frac{|(\xi \partial_\xi)^i f(\xi)|}{\langle \xi \rangle^a}, \quad (\text{E.3})$$

with  $\langle \xi \rangle = \sqrt{1 + \xi^2}$ . For sake of shortness and since the determination of both  $S_j$  and  $R_i$  follows the same reasoning, we only consider  $S_j$  in the sequel.

### Step 1: Construction of $S_j$

Identity (E.1) can be put in the form

$$S_j = 2\beta b H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) (S_j + c_{i,j}(2\beta)^{j+1}T_{j+1}) \right]. \quad (\text{E.4})$$

Now, write  $S_j$  as

$$S_j = L(S_j) = L(0) + DL(S_j), \quad (\text{E.5})$$

where

$$L(0) = bc_{i,j}(2\beta)^{j+2}H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) T_{j+1} \right],$$

$$DL(S_j) = b(2\beta)H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) S_j \right].$$

Our goal is to prove that for all  $j \leq i \leq \ell$

$$\|L(0)\|_{X_{\xi_0}^{2j-\gamma+2}} \leq Cy_0^2, \quad (\text{E.6})$$

$$\|DL(S_j)\|_{X_{\xi_0}^{2j-\gamma+2}} \leq Cy_0^2 \|S_j\|_{X_{\xi_0}^{2j-\gamma+2}}. \quad (\text{E.7})$$

Once these estimates are established, we apply the Banach fixed point theorem to  $L(S_j)$  mapping the ball  $B(0, 2Cy_0^2)$  into itself with  $y_0 \leq \frac{1}{2\sqrt{C}}$ . This yields the existence and uniqueness of  $S_j$  satisfying

$$\|S_j\|_{X_{\xi_0}^{2j-\gamma+2}} \leq 2Cy_0^2. \quad (\text{E.8})$$

We are now in position to prove (E.6) and (E.7):

- *Proof of (E.6)*: take  $a = 2j + 2 - \gamma$ , we get

$$\|L(0)\|_{X_{\xi_0}^a} \leq |c_{i,j}|b(2\beta)^{j+2} \left( \left| \frac{\alpha}{2} - i + \tilde{\lambda} \right| \|H^{-1}(T_{j+1})\|_{X_{\xi_0}^a} + \frac{1}{2} \|H^{-1}(\Lambda T_{j+1})\|_{X_{\xi_0}^a} \right).$$

Lemma E.1 yields

$$\|L(0)\|_{X_{\xi_0}^a} \leq C \|T_{j+1}\|_{X_{\xi_0}^{a-2}} \leq Cb\xi_0^2 \|T_{j+1}\|_{X_{\xi_0}^a}.$$

From lemma 10.1 and  $X_{\xi_0}^a$ 's definition, one infers

$$\|T_{j+1}\|_{X_{\xi_0}^a} \leq C.$$

Plugging  $\xi_0 = \frac{y_0}{\sqrt{b}}$ , we get

$$\|L(0)\|_{X_{\xi_0}^a} \leq C y_0^2,$$

which concludes (E.6).

- *Proof of (E.7)*: we argue as in the proof of (E.6). Indeed, we apply Lemma E.1 so that

$$\begin{aligned} \|DL(S_j)\|_{X_{\xi_0}^a} &\leq Cb(2\beta)(\|H^{-1}(S_j)\|_{X_{\xi_0}^a} + \|H^{-1}(\Lambda S_j)\|_{X_{\xi_0}^a}) \\ &\leq C y_0^2 \|S_j\|_{X_{\xi_0}^a}, \end{aligned}$$

We conclude (E.8) as above.

Our task now is to establish the desired estimates for  $\partial_{\tilde{\lambda}} S_j$ ,  $\partial_b S_j$  and  $\partial_\beta S_j$ . Since the proofs are quite the same we only estimate  $\partial_b S_j$ . Apply  $\partial_b$  to both sides of (E.5) to get

$$\begin{aligned} \partial_b S_j &= c_{i,j}(2\beta)^{j+2} H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) T_{j+1} \right] + (2\beta) H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) S_j \right] \\ &\quad + b(2\beta) H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) \partial_b S_j \right]. \end{aligned}$$

Using Lemma E.1 with  $\tilde{a} = 2j + 4 - \gamma$ , we derive

$$\begin{aligned} \|\partial_b S_j\|_{X_{\xi_0}^{\tilde{a}}} &\leq C \left( \|T_{j+1}\|_{X_{\xi_0}^{\tilde{a}-2}} + \|S_j\|_{X_{\xi_0}^{\tilde{a}-2}} + b \|\partial_b S_j\|_{X_{\xi_0}^{\tilde{a}-2}} \right) \\ &\leq C(1 + y_0^2) + C b \xi_0^2 \|\partial_b S_j\|_{X_{\xi_0}^{\tilde{a}}} \leq C(1 + y_0^2) + C y_0^2 \|\partial_b S_j\|_{X_{\xi_0}^a}, \end{aligned}$$

this implies that  $\|\partial_b S_j\|_{X_{\xi_0}^{2j+4-\gamma}} \leq C$ , provided  $y_0 \leq y_0^*$  is small enough.

+ For  $\partial_\beta S_j$ : Applying  $\partial_\beta$  to (E.5), we get

$$\begin{aligned} \partial_\beta S_j &= b2(j+2)(2\beta)^{j+1} c_{i,j} H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) T_{j+1} \right] + 2b H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) S_j \right] \\ &\quad + b(2\beta) H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) \partial_\beta S_j \right]. \end{aligned}$$

using the boundedness of  $j$ , we get (with  $a = 2j + 2 - \gamma$ )

$$\begin{aligned} \|\partial_\beta S_j\|_{X_{\xi_0}^a} &\leq Cb \left( \|T_{j+1}\|_{X_{\xi_0}^{a-2}} + \|S_j\|_{X_{\xi_0}^{a-2}} + \|\partial_\beta S_j\|_{X_{\xi_0}^{a-2}} \right) \\ &\leq C b \xi_0^2 \left( \|T_{j+1}\|_{X_{\xi_0}^a} + \|S_j\|_{X_{\xi_0}^a} + \|\partial_\beta S_j\|_{X_{\xi_0}^a} \right) \\ &\leq C y_0^2 \left( 1 + \|\partial_\beta S_j\|_{X_{\xi_0}^a} \right) \leq C y_0^2. \end{aligned}$$

provided that  $y_0 \leq y_0^*$  small enough.

### Step 2: construction of $R_i$

Taking  $H^{-1}$  to (E.2), we get

$$\begin{aligned} R_i &= b(2\beta) H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) R_i \right] + \frac{1}{2} \sum_{j=0}^i c_{i,j} (2\beta)^{j+1} b^j H^{-1}(\Theta_j) \\ &= J(R_i). \end{aligned} \tag{E.9}$$

Let us consider  $a = -\gamma + \epsilon > -\gamma$ , then, we apply Lemma [E.1](#)

$$\begin{aligned} \|J(R_i)\|_{X_{\xi_0}^a} &\leq C \left( b(2\beta)\|H^{-1}R_i\|_{X_{\xi_0}^a} + \sum_{j=0}^i (2\beta)^{j+1}b^j\|H^{-1}(\Theta_j)\|_{X_{\xi_0}^a} \right) \\ &\leq C \left( b\|R_i\|_{X_{\xi_0}^{a-2}} + \sum_{j=0}^i b^j\|\Theta_j\|_{X_{\xi_0}^{a-2}} \right). \end{aligned}$$

Now, we derive from [\(10.6\)](#) that

$$\|\Theta_j\|_{X_{\xi_0}^{-2-\gamma+\epsilon}} \leq C(\epsilon)\xi_0^{2j}.$$

Thus, we derive

$$\|J(R_i)\|_{X_{\xi_0}^a} \leq Cy_0^2\|R_i\|_{X_{\xi_0}^a} + C(\epsilon). \quad (\text{E.10})$$

Taking  $y_0$  small enough,  $J$  maps the ball  $B(0, 2C(\epsilon))$  into itself. In addition to that, it is similar to prove  $J$  is a contraction. Hence, by using Banach fixed point theorem, we imply the existence and the uniqueness of  $R_i$  satisfying

$$\|R_i\|_{X_{\xi_0}^{-\gamma+\epsilon}} \leq 2C(\epsilon).$$

Similarly for  $S_j$ , we can respectively take  $\partial_b$ ,  $\partial_{\tilde{\lambda}}$ , and  $\partial_\beta$  to [\(E.9\)](#) by using  $R_i \in B(0, 2C(\epsilon))$ , and we get

$$\begin{aligned} \|\partial_b R_i\|_{X_{\xi_0}^{-\gamma+2+\epsilon}} &\leq C(\epsilon), \\ \|\partial_{\tilde{\lambda}} R_i\|_{X_{\xi_0}^{-\gamma+2+\epsilon}} &\leq C(\epsilon)b, \\ \|\partial_\beta R_i\|_{X_{\xi_0}^{-\gamma+2+\epsilon}} &\leq C(\epsilon), \end{aligned}$$

where the constant  $C(\epsilon)$  is universal. Finally, we conclude the proof of the Proposition [10.2](#).  $\square$

$$\begin{aligned} \partial_b R_i &= (2\beta)H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) R_i \right] + b(2\beta)H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) \partial_b R_i \right] \\ &+ \frac{1}{2} \sum_{j=0}^i c_{i,j} (2\beta)^{j+1} b^{j-1} H^{-1} \Theta_j, \end{aligned}$$

and

$$\begin{aligned} \partial_\beta R_i &= 2bH^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) R_i \right] + b(2\beta)H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) \partial_\beta R_i \right] \\ &+ \sum_{j=0}^i c_{i,j} (j+1) (2\beta)^j b^j H^{-1} \Theta_j, \end{aligned}$$

and

$$\partial_{\tilde{\lambda}} R_i = (2\beta)bH^{-1}R_i + b(2\beta)H^{-1} \left[ \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) \partial_{\tilde{\lambda}} R_i \right].$$

The rest of this part is devoted to the results which are used to complete the proof of Proposition [10.2](#).

**Lemma E.1** (Continuity of  $H^{-1}$  in  $X_{\xi_0}^a$ ). *For all  $a \geq -\gamma$ , we have*

$$\|H^{-1}f\|_{X_{\xi_0}^a} \leq C \sup_{\xi \in [0, \xi_0]} \langle \xi \rangle^{2-a} |f(\xi)|, \quad (\text{E.11})$$

and

$$\|H^{-1}(\Lambda f)\|_{X_{\xi_0}^a} \leq C(a) \sup_{\xi \in [0, \xi_0]} [\langle \xi \rangle^{2-a} |f(\xi)| + \langle \xi \rangle^{3-a} |\partial_\xi f(\xi)|]. \quad (\text{E.12})$$

In particular

$$\|H^{-1}f\|_{X_{\xi_0}^a} \leq C \|f\|_{X_{\xi_0}^{a-2}}, \quad (\text{E.13})$$

and

$$\|H^{-1}(\Lambda f)\|_{X_{\xi_0}^a} \leq C \|f\|_{X_{\xi_0}^{a-2}}. \quad (\text{E.14})$$

Using the fact that, for  $\xi_0 \geq 1$ , we have

$$\|f\|_{X_{\xi_0}^{a-2}} \leq 2\xi_0^2 \|f\|_{X_{\xi_0}^a},$$

Formulae (E.11) and (E.12) read, for  $\xi_0 \geq 1$

$$\|H^{-1}f\|_{X_{\xi_0}^a} \leq C\xi_0^2 \|f\|_{X_{\xi_0}^a}, \quad (\text{E.15})$$

and

$$\|H^{-1}(\Lambda f)\|_{X_{\xi_0}^a} \leq C\xi_0^2 \|f\|_{X_{\xi_0}^a}. \quad (\text{E.16})$$

*Proof.* Regarding (E.3), we have

$$\sum_{j=0}^2 \frac{|(\xi \partial_\xi)^j g(\xi)|}{\langle \xi \rangle^a} \leq \langle \xi \rangle^{-a} |g(\xi)| + 2\langle \xi \rangle^{1-a} |\partial_\xi g| + \langle \xi \rangle^{2-a} |\partial_\xi^2 g|.$$

In addition to that, we derive from (D.2) that

$$\begin{aligned} \partial_\xi (H^{-1}f) &= \partial_\xi \Lambda Q \int_0^\xi \frac{\mathcal{L}f}{\Lambda Q}(\xi') d\xi' + \mathcal{L}(f)(\xi), \\ \partial_\xi^2 (H^{-1}f) &= \partial_\xi^2 \Lambda Q \int_0^\xi \frac{\mathcal{L}f}{\Lambda Q}(\xi') d\xi' + \frac{\partial_\xi \Lambda Q}{\Lambda Q} \mathcal{L}(f) + \partial_\xi (\mathcal{L}(f)), \end{aligned}$$

where  $\mathcal{L}$  defined as in (D.3). We remark that (E.11) follows from: for all  $\xi \in [0, \xi_0]$ :

$$\langle \xi \rangle^{-a} |H^{-1}f(\xi)| \leq \frac{C}{|a|} \sup_{\xi \in [0, \xi_0]} \xi^{2-a} (1 - \xi^\gamma) |f(\xi)|, \quad (\text{E.17})$$

$$\langle \xi \rangle^{1-a} |\partial_\xi (H^{-1}f(\xi))| \leq \frac{C}{|a|} \sup_{\xi \in [0, \xi_0]} \xi^{2-a} (1 - \xi^\gamma) |f(\xi)|, \quad (\text{E.18})$$

$$\langle \xi \rangle^{2-a} |\partial_\xi^2 (H^{-1}f(\xi))| \leq \frac{C}{|a|} \sup_{\xi \in [0, \xi_0]} \xi^{2-a} (1 - \xi^\gamma) |f(\xi)|, \quad (\text{E.19})$$

where  $C$  does not depend on  $\xi_0$ . Let us start with the proof of these estimates:

- *The proof of (E.17):* From  $\mathcal{L}$ 's formula in (D.3), we have

$$\begin{aligned} |\mathcal{L}(f)| &\leq \frac{1}{|\xi|^{d+1} |\Lambda Q|} \int_0^\xi \langle \xi' \rangle^{2-a} |f(\xi')| \langle \xi' \rangle^{a-2} |\Lambda Q(\xi')| (\xi')^{d+1} d\xi' \\ &\leq \sup_{\xi \in [0, \xi_0]} \{ \langle \xi \rangle^{2-a} |f(\xi)| \} \frac{1}{|\xi|^{d+1} |\Lambda Q|} \int_0^\xi \langle \xi' \rangle^{a-2} |\Lambda Q(\xi')| (\xi')^{d+1} d\xi' \\ &= \sup_{\xi \in [0, \xi_0]} \{ \langle \xi \rangle^{2-a} |f(\xi)| \} \tilde{L}(\xi). \end{aligned}$$

Plugging this estimate to  $H^{-1}f$ , we obtain

$$|H^{-1}f(\xi)| \leq \left[ \sup_{\xi \in [0, \xi_0]} \langle \xi \rangle^{2-a} |f(\xi)| \right] |\Lambda Q| \int_0^\xi \frac{\tilde{L}(\xi')}{|\Lambda Q|} d\xi'.$$

Hence, it is sufficient to prove

$$|\Lambda Q| \int_0^\xi \frac{\tilde{L}(\xi')}{|\Lambda Q|} d\xi' \leq C \langle \xi \rangle^a,$$

where  $C$  does not depend on  $\xi_0$ . Indeed, we consider two cases where  $\xi_0 \ll 1$  and  $\xi_0 \gg 1$ :

+ The case  $\xi_0 \ll 1$ : We have the following for all  $\xi \in [0, \xi_0]$

$$\frac{1}{C} \leq |\Lambda Q| \leq C,$$

and

$$\tilde{L}(\xi) = \frac{1}{|\xi|^{d+1} |\Lambda Q|} \int_0^\xi \langle \xi' \rangle^{a-2} |\Lambda Q| (\xi')^{d+1} d\xi' \leq C \xi,$$

this yields

$$|\Lambda Q| \int_0^\xi \frac{\tilde{L}(\xi')}{|\Lambda Q|} d\xi' \leq C \xi^2 \leq 2C.$$

which concludes the case  $\xi_0 \ll 1$ .

+ The case  $\xi_0 \gg 1$ : We observe that there exists  $M > 0$  such that for all  $\xi \in [M, \xi_0]$

$$\frac{1}{C} \xi^{-\gamma} \leq |\Lambda Q| \leq C \xi^{-\gamma}.$$

Then, we have

$$\begin{aligned} \int_0^\xi \frac{\tilde{L}(\xi')}{|\Lambda Q|} d\xi' &= \int_0^M \frac{\tilde{L}(\xi')}{|\Lambda Q|} d\xi' + \int_M^\xi \frac{\tilde{L}(\xi')}{|\Lambda Q|} d\xi' \\ &\leq C(M) + C \int_M^\xi \tilde{L}(\xi') (\xi')^\gamma d\xi'. \end{aligned}$$

Besides that, we estimate  $\tilde{L}(\xi')$ , for all  $\xi' \in [M, \xi_0]$  as follows

$$\begin{aligned} \tilde{L}(\xi) &= \frac{1}{\xi^{d+1} |\Lambda Q|} \left( \int_0^M \langle \xi' \rangle^{a-2} |\Lambda Q| (\xi')^{d+1} d\xi' + \int_M^\xi \langle \xi' \rangle^{a-2} |\Lambda Q| (\xi')^{d+1} d\xi' \right) \\ &\leq C(M) \left( \xi^{-d-1+\gamma} + \xi^{a-1} \right), \end{aligned}$$

it follows that

$$\int_M^\xi \tilde{L}(\xi') (\xi')^\gamma d\xi' \leq C(M) (1 + \xi^{-d+2\gamma} + \xi^{a+\gamma}).$$

Thus, we derive

$$|\Lambda Q| \int_0^\xi \frac{\tilde{L}(\xi')}{|\Lambda Q|} d\xi' \leq C(M, a) (\xi^{-\gamma} + \xi^a).$$

Finally, we have

$$|H^{-1}f(\xi)| \leq C(a) \langle \xi \rangle^a,$$

provided that

$$a > -\gamma.$$

- The proofs of (E.18) and (E.19) are the same.

Now, it remains to prove that if  $f \in X_{\xi_0}^a$ , then we have

$$\|f\|_{X_{\xi_0}^{a-2}} \leq C(a)\xi_0^2 \|f\|_{X_{\xi_0}^a},$$

provided that  $\xi_0 \geq 1$ . Indeed, this comes from

$$\langle \xi \rangle^{2-a} = \langle \xi \rangle^{-a} \langle \xi \rangle^2 \leq 2\xi_0^2 \langle \xi \rangle^{-a},$$

provided that  $\xi_0 \geq 1$ . Thus, we have

$$\|f\|_{X_{\xi_0}^{a-2}} = \sup_{\xi \in [0, \xi_0]} \sum_{j=0}^2 \left| \frac{\langle \xi \rangle \partial_\xi^j f}{\langle \xi \rangle^{a-2}} \right| \leq \sup_{\xi \in [0, \xi_0]} \sum_{j=0}^2 2\xi_0^2 \left| \frac{\langle \xi \rangle \partial_\xi^j f}{\langle \xi \rangle^a} \right| \leq 2\xi_0^2 \|f\|_{X_{\xi_0}^a}.$$

This concludes the proof of the Lemma.  $\square$

In the following Lemma, we aim to estimate  $\partial_\xi^3 S_j$  and  $\partial_\xi^3 R_i$ :

**Lemma E.2** (Higher estimates for  $\partial_\xi^3 S_j$  and  $\partial_\xi^3 R_i$ ). *Let us consider  $S_j$  and  $R_i$  which satisfying (E.4) and (E.9). Furthermore, we assume that the following estimates hold*

$$\|S_j\|_{X_{\xi_0}^{2j+2-\gamma}} \leq C y_0^2 \text{ and } \|R_i\|_{X_{\xi_0}^{\epsilon-\gamma}} \leq C, \text{ with } \xi_0 = \frac{y_0}{\sqrt{b}}.$$

Then, the following holds: for all  $\xi \in [0, \xi_0]$

$$|\partial_\xi^3 S_j(\xi)| \leq C \langle \xi \rangle^{2j-\gamma-1}, \quad (\text{E.20})$$

$$|\partial_\xi^3 R_i(\xi)| \leq C \langle \xi \rangle^{\epsilon-\gamma-3}. \quad (\text{E.21})$$

*Proof.* - The proof of (E.20): We remark that when  $b \rightarrow 0$ ,  $\xi_0 \rightarrow +\infty$ . Then, we will consider two situations, namely,  $\xi \ll 1$  and  $\xi \gg 1$ . Recall the inverse formula

$$S_j = bH^{-1}(f), f = \left( \frac{\alpha}{2} - i + \tilde{\lambda} + \frac{1}{2}\Lambda \right) (S_j + c_{i,j} T_{j+1})$$

and write  $\partial_\xi^3 S_j$  as follows

$$\begin{aligned} b^{-1} \partial_\xi^3 S_j(\xi) &= \partial_\xi^3 \Lambda Q \int_0^\xi \frac{\mathcal{L}(f)}{\Lambda Q} d\xi' + \frac{\partial_\xi^2 \Lambda Q}{\Lambda Q} \mathcal{L}(f) + \partial_\xi f - \frac{(d+1)(d+2)}{\xi^2} \mathcal{L}(f) - \frac{(d+1)\partial_\xi \Lambda Q}{\xi \Lambda Q} \mathcal{L}(f) \\ &- \frac{(d+1)}{\xi} f. \end{aligned}$$

+ The case  $\xi \in [0, 1]$ : We observe that when  $\xi \leq 1$ , it follows that  $|f(\xi)| \leq C$ , then, plugging to (E.4), we obtain

$$|S_j(\xi)| = |H^{-1}f| \leq C\xi^2,$$

Hence, we refine the behavior near 0 as follows

$$\begin{aligned} |f(\xi)| &\leq C\xi^2, \\ |\mathcal{L}(f)(\xi)| &\leq C\xi^3, \end{aligned}$$

continuing this process we enhance the behavior to

$$\begin{aligned} |f(\xi)| &\leq C\xi^{2j+2}, \\ |\mathcal{L}(f)(\xi)| &\leq C\xi^{2j+3}, \end{aligned}$$

(we can get a precise behavior for  $S_j$  at 0 by  $S_j(\xi) = O(\xi^{2j+4})$ ). Then, it is easy to derive

$$b^{-1} \partial_\xi^3 S_j(\xi) = O(\xi^{2j+1}) \text{ as } \xi \rightarrow 0.$$

+ The case  $\xi \geq 1$ : we have the following fact

$$|S_j(\xi)| \leq C \langle \xi \rangle^{2j+2-\gamma} \text{ and } |T_{j+1}(\xi)| \leq C \langle \xi \rangle^{2j+2-\gamma},$$

which yields

$$|f(\xi)| \leq C\langle \xi \rangle^{2j+2-\gamma}.$$

Since

$$|\mathcal{L}(f)(\xi)| \leq C\langle \xi \rangle^{2j+3-\gamma}, \text{ and } |S_j(\xi)| \leq C\langle \xi \rangle^{2j+2-\gamma},$$

we get

$$|\partial_\xi^3 S_j(\xi)| \leq Cb\langle \xi \rangle^{2j+1-\gamma} \leq C\langle \xi \rangle^{2j-\gamma-1},$$

due to the fact that

$$b\xi^2 = y_0 \leq 1.$$

Thus, we conclude the proof of (E.20). By the same technique, we derive (E.21). This finalizes the proof of the Lemma.  $\square$

## F. Outer eigenfunctions construction

This paragraph is devoted to give the complete proof to Proposition 10.3:

*Proof.* Now, let  $\phi_{i,out,\beta}$  be of the form

$$\phi_{i,out,\beta}(y) = \phi_{i,\infty,\beta}(y) + \tilde{\lambda}(\tilde{\phi}_{i,\beta}(y) + R_{i,1}(y)) + R_{i,2}(y), \quad (\text{F.1})$$

where  $R_{i,1}$  and  $R_{i,2}$  are to be constructed. Rewrite (2.18) as

$$\mathcal{L}_b = \mathcal{L}_\infty^\beta - 3(d-2) \left( \frac{1}{y^2} + 2Q_b + Q_b^2 y^2 \right),$$

where  $\mathcal{L}_\infty^\beta$  was defined in (2.22) and  $\mathcal{L}_{i,ext}^\beta = \mathcal{L}_\infty^\beta - 2\beta \left( \frac{\alpha}{2} - i \right)$ . Plugging (F.1) into

$$\left[ \mathcal{L}_b - 2\beta \left( \frac{\alpha}{2} - i \right) - \tilde{\lambda} \right] (\phi_{i,out,\beta}) = 0,$$

which yields

$$\begin{aligned} & \mathcal{L}_{i,ext}^\beta \phi_{i,out,\beta} - \tilde{\lambda} \phi_{i,out,\beta} - 3(d-2) \left( \frac{1}{y^2} + 2Q_b + Q_b^2 y^2 \right) \phi_{i,out,\beta} = \left( \mathcal{L}_{i,ext}^\beta R_{i,1} - \tilde{\lambda}(\tilde{\phi}_{i,\beta} + R_{i,1}) \right) \\ & + \left( \mathcal{L}_{i,ext}^\beta R_{i,2} - 3(d-2) \left( \frac{1}{y^2} + 2Q_b + Q_b^2 y^2 \right) \phi_{i,out} - \tilde{\lambda} R_{i,2} \right) = 0, \end{aligned}$$

where we used  $\mathcal{L}_{i,ext}^\beta \phi_{i,\infty} = 0$  as well as  $\mathcal{L}_{i,ext}^\beta(\tilde{\phi}_{i,\beta}) = \phi_{i,\infty,\beta}$ . Thus, it is sufficient to construct  $R_{i,1}$  and  $R_{i,2}$  satisfying

$$\begin{aligned} \mathcal{L}_{i,ext}^\beta(R_{i,1}) &= \tilde{\lambda} R_{i,1} + \tilde{\lambda} \tilde{\phi}_{i,\beta} \\ \mathcal{L}_{i,ext}^\beta(R_{i,2}) &= \left( \tilde{\lambda} + 3(d-2) \left( \frac{1}{y^2} + 2Q_b + Q_b^2 y^2 \right) \right) R_{i,2} \\ &+ 3(d-2) \left( \frac{1}{y^2} + 2Q_b + Q_b^2 y^2 \right) (\phi_{i,\infty,\beta} + \tilde{\lambda}(\tilde{\phi}_{i,\beta} + R_{i,1})), \end{aligned}$$

or equivalently

$$R_{i,1} = \tilde{\lambda} \mathcal{L}_{i,ext}^{-1}(R_{i,1}) + \tilde{\lambda} \phi_{i,\infty,\beta}, \quad (\text{F.2})$$

$$R_{i,2} = \mathcal{L}_{i,ext}^{-1}(H_1 R_{i,2}) + \mathcal{L}_{i,ext}^{-1}(H_2). \quad (\text{F.3})$$



Here

$$\begin{aligned} H_1(y) &= \left( \tilde{\lambda} + 3(d-2) \left( \frac{1}{y^2} + 2Q_b + Q_b^2 y^2 \right) \right) \\ H_2(y) &= 3(d-2) \left( \frac{1}{y^2} + 2Q_b + Q_b^2 y^2 \right) (\phi_{i,\infty,\beta} + \tilde{\lambda}(\tilde{\phi}_{i,\beta} + R_{i,1})). \end{aligned}$$

*Step 1: Construction of  $R_{i,1}$ :* The construction is based on Banach fixed point theorem on Banach space  $X_{y_0}^{a,a'}$  equipped with the norm introduced in (10.9). Now, denote the right hand side of (F.2) by  $K(R_{i,1})$ , and apply Lemma F.2 with  $a = \gamma - d$  and  $a' = 2i + 2 - \gamma$

$$\|K(R_{i,1})\|_{X_{y_0}^{a,a'}} \leq C_1(a, a')|\tilde{\lambda}| \left( \|R_{i,1}\|_{X_{y_0}^{a,a'}} + \|\phi_{i,\infty,\beta}\|_{X_{y_0}^{a,a'}} \right) \text{ and } \|\phi_{i,\infty,\beta}\|_{X_{y_0}^{a,a'}} \leq C_2.$$

Then,  $K$  maps the ball  $B(0, 2C_1C_2|\tilde{\lambda}|)$  into itself provided that  $|\tilde{\lambda}| \leq \min\left(\frac{1}{2}, \frac{1}{2C_1}\right)$ . In addition to that, for all  $X_1, X_2 \in B(0, 2C_1C_2|\tilde{\lambda}|)$ , we have

$$\|K(X_1) - K(X_2)\|_{X_{y_0}^{a,a'}} \lesssim |\tilde{\lambda}| \|X_1 - X_2\|_{X_{y_0}^{a,a'}}.$$

Hence, for  $\tilde{\lambda}$  small enough,  $K$  is a contraction and the existence of  $R_{i,1}$  follows with the bound

$$\|R_{i,1}\|_{X_{y_0}^{a,a'}} \leq 2C_1C_2|\tilde{\lambda}|.$$

Now, we establish the estimates for  $\partial_b R_{i,1}$ ,  $\partial_b R_{i,1}$  and  $\partial_\beta R_{i,1}$ .

- For  $\partial_b R_{i,1}$ : From (F.2), we see that  $R_{i,1}$  is independent of  $b$  so  $\partial_b R_{i,1} = 0$ .
- For  $\partial_{\tilde{\lambda}} R_{i,1}$ : Applying  $\partial_{\tilde{\lambda}}$  to (F.2), we obtain

$$\partial_{\tilde{\lambda}} R_{i,1} = \left( \mathcal{L}_{i,ext}^\beta \right)^{-1} (R_{i,1}) + \tilde{\lambda} \left( \mathcal{L}_{i,ext}^\beta \right)^{-1} (\partial_{\tilde{\lambda}} R_{i,1}) + \phi_{i,\infty,\beta}.$$

Lemma F.2 implies

$$\|\partial_{\tilde{\lambda}} R_{i,1}\|_{X_{y_0}^{a,a'}} \leq C_1 \|R_{i,1}\|_{X_{y_0}^{a,a'}} + C_1 |\tilde{\lambda}| \|\partial_{\tilde{\lambda}} R_{i,1}\|_{X_{y_0}^{a,a'}} + \|\phi_{i,\infty,\beta}\|_{X_{y_0}^{a,a'}},$$

hence

$$\|\partial_{\tilde{\lambda}} R_{i,1}\|_{X_{y_0}^{a,a'}} \leq C.$$

- For  $\partial_\beta R_{i,1}$ : Applying  $\partial_\beta$  to (F.2), we obtain

$$\partial_\beta R_{i,1} = \tilde{\lambda} \partial_\beta \left( \left( \mathcal{L}_{i,ext}^\beta \right)^{-1} \right) (R_{i,1}) + \tilde{\lambda} \left( \mathcal{L}_{i,ext}^\beta \right)^{-1} (\partial_\beta R_{i,1}) + \tilde{\lambda} \partial_\beta \phi_{i,\infty,\beta}.$$

Applying  $X_{y_0}^{a,a'}$  norm to the above equality and Lemma F.2, we deduce

$$\|\partial_\beta R_{i,1}\|_{X_{y_0}^{a,a'}} \leq C_1 |\tilde{\lambda}| \|R_{i,1}\|_{X_{y_0}^{a,a'}} + C_2 |\tilde{\lambda}| \|\partial_\beta R_{i,1}\|_{X_{y_0}^{a,a'}} + |\tilde{\lambda}| \|\partial_\beta \phi_{i,\infty,\beta}\|_{X_{y_0}^{a,a'}}.$$

On the one hand, we have

$$\|R_{i,1}\|_{X_{y_0}^{a,a'}} + \|\partial_\beta \phi_{i,\infty,\beta}\|_{X_{y_0}^{a,a'}} \leq C.$$

Finally, we get

$$\|\partial_\beta R_{i,1}\|_{X_{y_0}^{a,a'}} \leq C|\tilde{\lambda}|.$$

The construction and estimates on  $R_{i,2}$  are very similar to those established above and are left to the reader.

- *Step 2: Construction of  $R_{i,2}$ :*

The construction is also based on the Banach fixed point theorem on the Banach space  $X_{y_0}^{a,a'}$  with  $a = -\tilde{\gamma} - 2 - \alpha$ , and  $a' = 2i + 2 - \gamma$ . First, we define the right hand side of (F.3) to be  $J(R_{i,2})$ . Using Lemma F.2, we have the following

$$\|J(R_{2,i})\|_{X_{y_0}^{a,a'}} \leq C_1 \left( \|H_1 R_{i,2}\|_{X_{y_0}^{a,a'}} + \|H_2\|_{X_{y_0}^{a,a'}} \right).$$

We now aim to prove the following estimates

$$\|H_1 R_{i,2}\|_{X_{y_0}^{a,a'}} \leq C(y_0)b^\alpha \|R_{i,2}\|_{X_{y_0}^{a,a'}}, \quad (\text{F.4})$$

$$\|H_2\|_{X_{y_0}^{a,a'}} \leq C(y_0)b^\alpha. \quad (\text{F.5})$$

- *The proof of (F.4):* From Lemma 9.1, we have

$$Q_b(y) = \frac{1}{b} Q\left(\frac{y}{\sqrt{b}}\right) = \frac{1}{b} \left( -\frac{1}{\left(\frac{y}{\sqrt{b}}\right)^2} + q_0 \left(\frac{y}{\sqrt{b}}\right)^{-\gamma} + O_{b \rightarrow 0} \left(\frac{y}{\sqrt{b}}\right)^{-\gamma-g} \right),$$

since  $y \geq y_0$ ,  $\frac{y}{\sqrt{b}} \rightarrow +\infty$ . This implies that for all  $y \geq y_0$  and  $b \in (0, b^*(y_0))$

$$\begin{aligned} \left| 2Q_b(y) - \left( -\frac{2}{y^2} + 2q_0 y^{-\gamma} b^{\frac{\alpha}{2}} \right) \right| &\leq C(y_0)b^\alpha, \\ \left| y^2 Q_b^2(y) - \left( \frac{1}{y^2} - 2q_0 y^{-\gamma} b^{\frac{\alpha}{2}} \right) \right| &\leq C(y_0)b^\alpha, \end{aligned}$$

since  $g = -2\lambda_1 = 2(\gamma - 2)$ . Thus, for all  $y \geq y_0$ , we have

$$\left| 2Q_b(y) + Q_b^2(y)y^2 + \frac{1}{y^2} \right| \leq C(y_0)b^\alpha \lesssim b^{\frac{\alpha}{2}}. \quad (\text{F.6})$$

- *Proof of (F.5):* Recall that

$$\|R_{i,1}\|_{X_{y_0}^{-\tilde{\gamma},a'}} \leq C|\tilde{\lambda}|,$$

which implies

$$\|R_{i,1}\|_{X_{y_0}^{a,a'}} \leq C(y_0)|\tilde{\lambda}| \text{ with } a = -\tilde{\gamma} - 2 - \alpha.$$

Similarly, we also have

$$\|\phi_{i,\infty,\beta}\|_{X_{y_0}^{a,a'}} + \|\tilde{\phi}_{i,\beta}\|_{X_{y_0}^{a,a'}} \leq C(y_0).$$

Thus, the above estimates and (F.6) immediately conclude (F.5).

Using estimates (F.4) and (F.5), we get

$$\|J(R_{2,i})\|_{X_{y_0}^{a,a'}} \leq C_1(y_0)b^\alpha \left( \|R_{i,2}\|_{X_{y_0}^{a,a'}} + 1 \right).$$

Consequently, once  $b \ll 1$ ,  $J$  becomes a contraction from the ball  $B(C_1 b^\alpha, 0)$  to itself. Thus, it follows Banach fixed point theorem the existence of  $R_{i,2}$  satisfying

$$\|R_{i,2}\|_{X_{y_0}^{a,a'}} \leq Cb^{\frac{\alpha}{2}}. \quad (\text{F.7})$$

Next, we focus on evaluating  $\partial_{\tilde{\lambda}} R_{i,2}$ ,  $\partial_b R_{i,2}$  and  $\partial_\beta R_{i,2}$ :

+ For  $\partial_{\tilde{\lambda}} R_{i,2}$ : We have

$$\partial_{\tilde{\lambda}} R_{i,2} = \mathcal{L}_{i,ext}^{-1}(R_{i,2}) + \mathcal{L}_{i,ext}^{-1}(H_1 \partial_{\tilde{\lambda}} R_{i,2}) + \mathcal{L}_{i,ext}^{-1}(\partial_{\tilde{\lambda}} H_2)$$

then

$$\begin{aligned} \|\partial_{\bar{\lambda}} R_{i,2}\|_{X_{y_0}^{a,a'}} &\leq C_1 \|R_{i,2}\|_{X_{y_0}^{a,a'}} + C_2 \|H_1 \partial_{\bar{\lambda}} R_{i,2}\|_{X_{y_0}^{a,a'}} + C_3 \|\partial_{\bar{\lambda}} H_2\|_{X_{y_0}^{a,a'}} \\ &\leq C_1 b^{\frac{\alpha}{2}} + C_2 b^{\frac{\alpha}{2}} \|\partial_{\bar{\lambda}} R_{i,2}\|_{X_{y_0}^{a,a'}} + C_3 b^{\frac{\alpha}{2}} \end{aligned}$$

which yields to

$$\|\partial_{\bar{\lambda}} R_{i,2}\|_{X_{y_0}^{a,a'}} \leq C b^{\frac{\alpha}{2}}.$$

- For  $\partial_b R_{i,2}$ : We have

$$\partial_b R_{i,2} = \mathcal{L}_{i,ext}^{-1}(\partial_b H_1 R_{i,2}) + \mathcal{L}_{i,ext}^{-1}(H_1 \partial_b R_{i,2}) + \mathcal{L}_{i,ext}^{-1}(\partial_b H_2)$$

then

$$\begin{aligned} \|\partial_b R_{i,2}\|_{X_{y_0}^{a,a'}} &\leq C_1 \|\partial_b H_1 R_{i,2}\|_{X_{y_0}^{a,a'}} + C_2 \|\Theta_1 \partial_b R_{i,2}\|_{X_{y_0}^{a,a'}} + C_3 \|\partial_b H_2\|_{X_{y_0}^{a,a'}} \\ &\leq C_1 b^{\frac{\alpha}{2}-1} + C_2 b^{\frac{\alpha}{2}} \|\partial_{\bar{\lambda}} R_{i,2}\|_{X_{y_0}^{a,a'}} + C_3 b^{\frac{\alpha}{2}-1} \leq C b^{\frac{\alpha}{2}-1}. \end{aligned}$$

- For  $\partial_{\beta} R_{i,2}$ : We have

$$\partial_{\beta} R_{i,2} = \partial_{\beta}(\mathcal{L}_{i,ext}^{-1}(\Theta_1 R_{i,2})) + \mathcal{L}_{i,ext}^{-1}(\Theta_1 \partial_{\beta} R_{i,2}) + \partial_{\beta}(\mathcal{L}_{i,ext}^{-1}(\Theta_2)),$$

then we use Lemma F.2 to get

$$\|\partial_{\beta} R_{i,2}\|_{X_{y_0}^{a,a'}} \leq C(y_0) \left( b^{\frac{\alpha}{2}} \|R_{i,2}\|_{X_{y_0}^{a,a'}} + b^{\frac{\alpha}{2}} \|\partial_{\beta} R_{i,2}\|_{X_{y_0}^{a,a'}} + b^{\alpha} \right) \leq C b^{\frac{\alpha}{2}}.$$

Finally, we conclude the proof of the Proposition.  $\square$

In the sequel, we aim to complete the results used in the proof of Proposition 10.3. To be begin with, we need the following result on the resonance of  $\mathcal{L}_{i,ext}^{\beta}$ . For sake of shortness we set

$$\mathcal{L}_{i,ext}^{\beta} u = \left( \mathcal{L}_{\infty}^{\beta} - 2\beta \left( \frac{\alpha}{2} - i \right) \right) u. \quad (\text{F.8})$$

We have

**Lemma F.1** (Resonance of  $\mathcal{L}_{i,ext}^{\beta}$ ). *We consider  $i \in \mathbb{N}$ , then, there exists  $\tilde{\psi}_{i,\beta}$  such that it solves  $\mathcal{L}_{i,ext}^{\beta} \tilde{\psi}_{i,\beta} = 0$ . Moreover, we have*

$$\text{Ker}(\mathcal{L}_{i,ext}^{\beta}) = \text{Span}\{\phi_{i,\infty,\beta}, \tilde{\psi}_{i,\beta}\},$$

where  $\phi_{i,\infty,\beta}$  is the  $i$ -th eigenfunction of  $\mathcal{L}_{\infty}^{\beta}$ , given in Proposition 4.1; and  $\tilde{\psi}_{i,\beta}$  has the following asymptotic:

$$\tilde{\psi}_{i,\beta}(y) = \begin{cases} \frac{y^{\gamma-d}}{a_{i,0}(d-2\gamma)} (1 + O(y^2)) \text{ as } y \rightarrow 0, \\ -\frac{2}{a_{i,i}(2\beta)^i} y^{-2i+\gamma-(d+2)} e^{2\beta \frac{y^2}{4}} [1 + O(y^{-2})] \text{ as } y \rightarrow +\infty \end{cases} \quad (\text{F.9})$$

where  $\gamma$  and  $a_{i,j}$  were defined in (2.26) and (2.28). In particular, there exists a solution  $\tilde{\phi}_{i,\beta}$  to  $\mathcal{L}_{i,ext}^{\beta} \tilde{\phi}_i = \phi_{i,\infty,\beta}$ , satisfying the following asymptotic

$$\tilde{\phi}_{i,\beta}(y) = \begin{cases} K_0 y^{-\tilde{\gamma}} (1 + O(y^2)) \text{ as } y \rightarrow 0, \\ K_{\infty} y^{2i-\gamma} (\ln y + O(1)) \text{ as } y \rightarrow +\infty, \end{cases} \quad (\text{F.10})$$

where  $K_0 = \frac{1}{(2\gamma-d)(d+2-2\gamma)}$ ,  $K_{\infty} = 2(2\beta)^i$  and  $\tilde{\gamma}$  is defined by

$$\tilde{\gamma} = \frac{1}{2}(d + \sqrt{d^2 - 12d + 24}). \quad (\text{F.11})$$

*Proof.* Recall that  $\phi_{i,\infty,\beta}$  solves  $\mathcal{L}_{i,ext}^\beta \phi_{i,\infty,\beta} = 0$ , this follows from the fact that  $\phi_{i,\infty}$  is the  $i$ -th eigenfunction of  $\mathcal{L}_\infty$ . According to  $\mathcal{L}_{i,ext}^\beta$ , the Wronskian is given by

$$W(y) = y^{-(d+1)} e^{2\beta \frac{y^2}{4}}. \quad (\text{F.12})$$

Then, we can formulate an independent linear solution  $\tilde{\psi}_{i,\beta}$  to  $\mathcal{L}_{i,ext}^\beta \tilde{\psi}_{i,\beta} = 0$

$$\tilde{\psi}_{i,\beta}(y) = -\phi_{i,\infty,\beta} \int_1^y \frac{(y')^{-(d+1)} e^{2\beta \frac{(y')^2}{4}}}{\phi_{i,\infty,\beta}^2(y')} dy'. \quad (\text{F.13})$$

From (4.2), we derive

$$\phi_{i,\infty,\beta}(y) = \begin{cases} a_{i,0} y^{-\gamma} (1 + O(y^2)) & \text{as } y \rightarrow 0, \\ a_{i,i} (2\beta)^i y^{2i-\gamma} (1 + O(y^{-2})) & \text{as } y \rightarrow +\infty, \end{cases} \quad (\text{F.14})$$

where  $a_{i,j}$ 's general formula given in (2.28), and we plug the above fact into (F.13) to get

$$\tilde{\psi}_i(y) = \begin{cases} \frac{y^{\gamma-d}}{a_{i,0}(2\gamma-d)} (1 + O(y^2)) & \text{as } \xi \rightarrow 0, \\ -\frac{2}{a_{i,i}(2\beta)^i} y^{-2i+\gamma-d-2} e^{2\beta \frac{y^2}{4}} (1 + O(y^{-2})) & \text{as } \xi \rightarrow +\infty. \end{cases} \quad (\text{F.15})$$

In particular, it is easy to see that

$$\text{Ker}(\mathcal{L}_{i,ext}^\beta) = \text{Span}\{\phi_{i,\infty,\beta}, \tilde{\psi}_{i,\beta}\},$$

which leads to

$$\left(\mathcal{L}_{i,ext}^\beta\right)^{-1} f(y) = -\phi_{i,\infty,\beta} \int_1^y f(y') \frac{\tilde{\psi}_{i,\beta}(y')}{W(y')} d\xi' + \tilde{\psi}_{i,\beta} \int_y^{+\infty} f(y') \frac{\phi_{i,\beta,\infty}(y')}{W(y')} dy' + c_1 \phi_{i,\beta,\infty} + c_2 \tilde{\psi}_{i,\beta},$$

and since we need to construct a special solution with explicit asymptotics, we choose  $c_1 = c_2 = 0$ , then, we can omit the generality and write

$$\left(\mathcal{L}_{i,ext}^\beta\right)^{-1} f(y) = -\phi_{i,\infty,\beta} \int_1^y f(y') \frac{\tilde{\psi}_{i,\beta}(y')}{W(y')} dy' + \tilde{\psi}_{i,\beta} \int_y^{+\infty} f(y') \frac{\phi_{i,\beta,\infty}(y')}{W(y')} dy'. \quad (\text{F.16})$$

Thus, the solution  $\tilde{\phi}_{i,\beta}$  to  $\mathcal{L}_{i,ext}^\beta \tilde{\phi}_{i,\beta} = \phi_{i,\infty,\beta}$  can be written

$$\tilde{\phi}_{i,\beta} = \left(\mathcal{L}_{i,ext}^\beta\right)^{-1} (\phi_{i,\infty,\beta}).$$

- *Behavior at 0:* From (F.14) and (F.15), we have

$$-\phi_{i,\infty,\beta} \int_1^y \frac{\phi_{i,\infty,\beta}(\xi') \tilde{\psi}_{i,\infty,\beta}(\xi')}{W(\xi')} d\xi' = \frac{a_{i,0}}{2(2\gamma-d)} y^{-\gamma+2} (1 + O(\xi^2)) \text{ as } y \rightarrow 0$$

and also

$$\tilde{\psi}_{i,\beta} \int_y^\infty \frac{\phi_{i,\infty,\beta}^2(\xi')}{W(\xi')} d\xi' = \frac{a_{i,i}}{(2\gamma-d)(-2\gamma+d+2)} y^{\gamma-d} (1 + O(\xi^2)) \text{ as } y \rightarrow 0,$$

noting that  $\gamma - d = -\frac{1}{2}(d + \sqrt{d^2 - 12d + 24}) = -\tilde{\gamma}$ , we get

$$\tilde{\phi}_{i,\beta}(y) = K_0 y^{-\tilde{\gamma}} (1 + y^2) \text{ as } y \rightarrow 0,$$

where  $K_0 = \frac{1}{(2\gamma-d)(d+2-2\gamma)}$ .

- *Behavior at  $+\infty$ :* Using (F.14) and (F.15) again, we obtain

$$-\phi_{i,\infty,\beta} \int_1^y \frac{\phi_{i,\infty,\beta}(\xi') \tilde{\psi}_{i,\infty,\beta}(\xi')}{W(\xi')} d\xi' = 2a_{i,i} (2\beta)^i y^{2i-\gamma} (\ln y + O(1)) \text{ as } y \rightarrow +\infty,$$

and for the second one

$$\tilde{\psi}_{i,\beta} \int_y^\infty \frac{\phi_{i,\infty,\beta}^2(\xi')}{W(\xi')} d\xi' = -\frac{2}{a_{i,i}(2\beta)^i} y^{-2i+\gamma-d-2} e^{\frac{2\beta y^2}{4}} O(y^{4i-2\gamma+d} e^{-\frac{2\beta y^2}{4}}) \text{ as } \xi \rightarrow +\infty.$$

Thus, we have

$$\tilde{\phi}_{i,\beta}(y) = 2(2\beta)^i y^{2i-\gamma} (\ln \xi + O(1)) \text{ as } \xi \rightarrow +\infty,$$

and this concludes the proof of the Lemma.  $\square$

It remains only to prove the continuity of  $\mathcal{L}_{i,ext}^{-1}$  in the Banach space  $X^{a,a'}$ , a result that we have used above.

**Lemma F.2** (Continuity of  $\mathcal{L}_{i,ext}^{-1}$ ). *For all  $a \leq \gamma - d$ ,  $a \neq -d - 2$  and  $a' > 2i - \gamma$  with  $y_0 \in (0, 1)$  and  $\beta \in (\frac{1}{4}, \frac{3}{4})$ , we have the following estimate*

$$\left\| \left( \mathcal{L}_{i,ext}^\beta \right)^{-1} f \right\|_{X_{y_0}^{a,a'}} \leq C(a, a', \beta) \|f\|_{X_{y_0}^{a,a'}} \text{ and } \left\| \partial_\beta \left( \mathcal{L}_{i,ext}^\beta \right)^{-1} f \right\|_{X_{y_0}^{a,a'}} \leq C(a, a', \beta) \|f\|_{X_{y_0}^{a,a'}}.$$

*Proof.* Define  $g = \left( \mathcal{L}_{i,ext}^\beta \right)^{-1} f$  and recall from (F.16) that

$$g(y) = \left( \mathcal{L}_{i,ext}^\beta \right)^{-1} f(y) = -\phi_{i,\infty,\beta} \int_1^y f(\xi') \frac{\tilde{\psi}_{i,\beta}(\xi')}{W(\xi')} d\xi' + \tilde{\psi}_{i,\beta} \int_y^{+\infty} f(\xi) \frac{\phi_{i,\beta,\infty}(\xi')}{W(\xi')} d\xi'.$$

Then

$$\begin{aligned} \partial_y g(y) &= -\partial_y \phi_{i,\infty,\beta} \int_1^y f(\xi') \frac{\tilde{\psi}_{i,\beta}(\xi')}{W(\xi')} d\xi' + \partial_y \tilde{\psi}_{i,\beta} \int_y^{+\infty} f(\xi) \frac{\phi_{i,\beta,\infty}(\xi')}{W(\xi')} d\xi' \phi_{i,\infty,\beta} \\ &\quad - 2\phi_{i,\infty,\beta} f(y) \frac{\tilde{\psi}_{i,\beta}(y)}{W(y)} \end{aligned}$$

and

$$\begin{aligned} \partial_y^2 g(y) &= -\partial_y^2 \phi_{i,\infty,\beta} \int_1^y f(\xi') \frac{\tilde{\psi}_{i,\beta}(\xi')}{W(\xi')} d\xi' + \partial_y^2 \tilde{\psi}_{i,\beta} \int_y^{+\infty} f(\xi) \frac{\phi_{i,\beta,\infty}(\xi')}{W(\xi')} d\xi' \phi_{i,\infty,\beta} \\ &\quad - 3\partial_y \phi_{i,\infty,\beta} f(y) \frac{\tilde{\psi}_{i,\beta}(y)}{W(y)} - 3\partial_y \tilde{\psi}_{i,\beta} f(y) \frac{\phi_{i,\beta,\infty}(y)}{W(y)} \\ &\quad - 2\phi_{i,\infty,\beta} \partial_y f(y) \frac{\tilde{\psi}_{i,\beta}(y)}{W(y)} + 2\phi_{i,\infty,\beta} f(y) \frac{\tilde{\psi}_{i,\beta}(y) \partial_y W(y)}{W^2(y)}. \end{aligned}$$

In order to establish the desired estimate, we will only need to control higher order derivatives, namely,  $y^{2-a} \partial_y^2 g(y)$  for  $y \in [y_0, 1]$  and  $y^{2-a'} \partial_y^2 g(y)$  for  $y \in [1, \infty)$ .

- $y \in [y_0, 1]$ , we have

$$\begin{aligned}
|y^{2-a} \partial_y^2 g(y)| &\lesssim y^{2-a} |\partial_y^2 \phi_{i,\infty,\beta}| \int_y^1 (\xi')^{-a} |f(\xi')| \frac{|(\xi')^a \|\tilde{\psi}_{i,\beta}(\xi')\|}{W(\xi')} d\xi' \\
&+ y^{2-a} |\partial_y^2 \tilde{\psi}_{i,\beta}| \left( \int_y^1 (\xi')^{-a} |f(\xi')| \frac{|(\xi')^a \|\phi_{i,\beta,\infty}(y)\|}{W(\xi')} d\xi' + \int_1^\infty (\xi')^{-a'} |f(\xi')| \frac{|(\xi')^{a'} \|\phi_{i,\beta,\infty}(y)\|}{W(\xi')} d\xi' \right) \\
&+ y^{2-a} |f(y)| \left( |\partial_y \phi_{i,\infty,\beta}| \frac{|\tilde{\psi}_{i,\beta}(y)|}{W(y)} + |\partial_y \tilde{\psi}_{i,\beta}| \frac{|\phi_{i,\beta,\infty}(y)|}{W(y)} + |\phi_{i,\infty,\beta}| \frac{|\tilde{\psi}_{i,\beta}(y) \partial_y W(y)|}{W^2(y)} \right) \\
&+ y^{1-a} |\partial_y f(y)| \left( y |\phi_{i,\infty,\beta}| \frac{|\tilde{\psi}_{i,\beta}(y)|}{W(y)} \right) \\
&\lesssim \sup_{y \in [y_0, 1]} |y^{-a} f(y)| + \sup_{y \in [y_0, 1]} |y^{1-a} \partial_y f(y)| + \sup_{y \in [1, +\infty)} |y^{-a'} f(y)| \lesssim \|f\|_{X_{y_0}^{a,a'}},
\end{aligned}$$

provided that  $a \leq \gamma - d$  and  $a' \geq 2i - \gamma$ .

- $y \in [1, +\infty)$ , we have

$$\begin{aligned}
|y^{2-a'} \partial_y^2 g(y)| &\lesssim y^{2-a'} |\partial_y^2 \phi_{i,\infty,\beta}| \int_1^\infty (\xi')^{-a'} |f(\xi')| \frac{|(\xi')^{a'} \|\tilde{\psi}_{i,\beta}(\xi')\|}{W(\xi')} d\xi' \\
&+ y^{2-a'} |\partial_y^2 \tilde{\psi}_{i,\beta}| \int_1^\infty (\xi')^{-a'} |f(\xi')| \frac{|(\xi')^{a'} \|\phi_{i,\beta,\infty}(y)\|}{W(\xi')} d\xi' \\
&+ y^{-a'} |f(y)| \left( y^2 |\partial_y \phi_{i,\infty,\beta}| \frac{|\tilde{\psi}_{i,\beta}(y)|}{W(y)} + y^2 |\partial_y \tilde{\psi}_{i,\beta}| \frac{|\phi_{i,\beta,\infty}(y)|}{W(y)} + y^2 |\phi_{i,\infty,\beta}| \frac{|\tilde{\psi}_{i,\beta}(y) \partial_y W(y)|}{W^2(y)} \right) \\
&+ y^{1-a'} |\partial_y f(y)| \left( y |\phi_{i,\infty,\beta}| \frac{|\tilde{\psi}_{i,\beta}(y)|}{W(y)} \right) \\
&\lesssim \sup_{y \in [1, +\infty)} |y^{-a'} f(y)| + \sup_{y \in [1, +\infty)} |y^{1-a'} \partial_y f(y)| \lesssim \|f\|_{X_{y_0}^{a,a'}}
\end{aligned}$$

which yields

$$\|g\|_{X_{y_0}^{a,a'}} = \left( \mathcal{L}_{i,ext}^\beta \right)^{-1} \|f\|_{X_{y_0}^{a,a'}} \leq C(a, a') \|f\|_{X_{y_0}^{a,a'}}$$

as claimed. Finally, we finish the proof of the Lemma.  $\square$

## REFERENCES

- [1] Paweł Biernat, Roland Donninger, and Birgit Schörkhuber. Stable self-similar blowup in the supercritical heat flow of harmonic maps. *Calc. Var. Partial Differential Equations*, 56(6):Paper No. 171, 31, 2017.
- [2] P. Biernat. Non-self-similar blow-up in the heat flow for harmonic maps in higher dimensions. *Nonlinearity*, 28(1):167–185, 2015.
- [3] J. Bricmont and A. Kupiainen. Universality in blow-up for nonlinear heat equations. *Nonlinearity*, 7(2):539–575, 1994.
- [4] P. Biernat and Y. Seki. Type II blow-up mechanism for supercritical harmonic map heat flow. *Int. Math. Res. Not. IMRN*, (2):407–456, 2019.
- [5] Paweł Biernat and Yukihiro Seki. Transition of blow-up mechanisms in k-equivariant harmonic map heat flow. *Nonlinearity*, 33(6):2756–2796, 2020.
- [6] P. Bizoń and A. Wasserman. Nonexistence of shrinkers for the harmonic map flow in higher dimensions. *Int. Math. Res. Not. IMRN*, (17):7757–7762, 2015.
- [7] C. Collot, T. E. Ghoul, N. Masmoudi, and V. T. Nguyen. Refined description and stability for singular solutions of the 2d keller-segel system. *Communications on Pure and Applied Mathematics*, 75(7):1419–1516, 2022.
- [8] C. Collot, T. E. Ghoul, N. Masmoudi, and V. T. Nguyen. Spectral analysis for singularity formation of the two dimensional keller-segel system. *Ann. PDE*, 8(5), 2022.
- [9] C. Collot, F. Merle, and P. Raphaël. Strongly anisotropic type II blow up at an isolated point. *J. Amer. Math. Soc.*, 33(2):527–607, 2020.
- [10] Charles Collot, Frank Merle, and Pierre Raphaël. Strongly anisotropic type II blow up at an isolated point. *J. Amer. Math. Soc.*, 33(2):527–607, 2020.
- [11] G. K. Duong, V. T. Nguyen, and H. Zaag. Construction of a stable blowup solution with a prescribed behavior for a non-scaling invariant semilinear heat equation. *Tunisian J. Math.*, 1(1):13–45, 2019.
- [12] G. K. Duong, N. Nouaïli, and H. Zaag. Construction of blow-up solutions for the complex ginzburg-landau equation with critical parameters. *Mem. Amer. Math. Soc.*, 2020. to appear.
- [13] R Donninger and B Schörkhuber. Stable blowup for the supercritical Yang-Mills heat flow. *J. Differential Geom.*, 113(1):55–94, 2019.
- [14] O Dumitrașcu. Equivariant solutions of the Yang-Mills equations. *Stud. Cerc. Mat.*, 34(4):329–333, 1982.
- [15] G. K. Duong. A blowup solution of a complex semi-linear heat equation with an irrational power. *J. Differential Equations*, 267(9):4975–5048, 2019.
- [16] G. K. Duong. Profile for the imaginary part of a blowup solution for a complex-valued semilinear heat equation. *J. Funct. Anal.*, 277(5):1531–1579, 2019.
- [17] Huijun Fan. Existence of the self-similar solutions in the heat flow of harmonic maps. *Sci. China Ser. A*, 42(2):113–132, 1999.
- [18] A Gastel. Singularities of first kind in the harmonic map and Yang-Mills heat flows. *Math. Z.*, 242(1):47–62, 2002.
- [19] T Ghoul. Stable type ii blowup for the 7 dimensional 1-corotational energy supercritical harmonic map heat flow. *Arxiv*.
- [20] T. E. Ghoul, S. Ibrahim, and V. T. Nguyen. On the stability of type II blowup for the 1-corotational energy-supercritical harmonic heat flow. *Anal. PDE*, 12(1):113–187, 2019.
- [21] J. F. Grotowski. Finite time blow-up for the Yang-Mills heat flow in higher dimensions. *Math. Z.*, 237(2):321–333, 2001.
- [22] I. Glogić and B. Schörkhuber. Nonlinear stability of homothetically shrinking Yang-Mills solitons in the equivariant case. *Comm. Partial Differential Equations*, 45(8):887–912, 2020.
- [23] M. A Herrero and J. J. J Velázquez. A blowup result for the semilinear heat equations in the supercritical case. 1992. Preprint.
- [24] H Kozono, Y Maeda, and H Naito. Global solution for the Yang-Mills gradient flow on 4-manifolds. *Nagoya Math. J.*, 139:93–128, 1995.
- [25] F. Merle and H. Zaag. Stability of the blow-up profile for equations of the type  $u_t = \Delta u + |u|^{p-1}u$ . *Duke Math. J.*, 86(1):143–195, 1997.
- [26] N. Masmoudi and H. Zaag. Blow-up profile for the complex Ginzburg-Landau equation. *J. Funct. Anal.*, 255(7):1613–1666, 2008.
- [27] N. Nouaïli and H. Zaag. Profile for a simultaneously blowing up solution to a complex valued semilinear heat equation. *Comm. Partial Differential Equations*, 40(7):1197–1217, 2015.
- [28] J Rade. On the Yang-Mills heat equation in two and three dimensions. *J. Reine Angew. Math.*, 431:123–163, 1992.

- [29] A Schlatter. Global existence of the Yang-Mills flow in four dimensions. *J. Reine Angew. Math.*, 479:133–148, 1996.
- [30] A. E. Schlatter, M. Struwe, and A. S. Tahvildar-Zadeh. Global existence of the equivariant Yang-Mills heat flow in four space dimensions. *Amer. J. Math.*, 120(1):117–128, 1998.
- [31] M Struwe. The Yang-Mills flow in four dimensions. *Calc. Var. Partial Differential Equations*, 2(2):123–150, 1994.
- [32] S. Tayachi and H. Zaag. Existence of a stable blow-up profile for the nonlinear heat equation with a critical power nonlinear gradient term. *Trans. Amer. Math. Soc.*, 317:5899–5972, 2019.
- [33] A. Waldron. Long-time existence for Yang-Mills flow. *Invent. Math.*, 217(3):1069–1147, 2019.
- [34] G. N Watson. A treatise on the theory of Bessel functions. *Cambridge Uni. Press*, Cambridge, 1922.
- [35] B Weinkove. Singularity formation in the Yang-Mills flow. *Calc. Var. Partial Differential Equations*, 19(2):211–220, 2004.