

LADUMA: Discovery of a luminous OH megamaser at $z > 0.5$

MARCIN GLOWACKI,^{1,2,3} JORDAN D. COLLIER,^{1,4,5,6} AMIR KAZEMI-MORIDANI,⁷ BRADLEY FRANK,^{8,1,4} HAYLEY ROBERTS,⁹ JEREMY DARLING,⁹ HANS-RAINER KLÖCKNER,¹⁰ NATHAN ADAMS,^{11,12} ANDREW J. BAKER,^{7,2} MATTHEW BERSHADY,^{13,14,4} TARIQ BLECHER,¹⁵ SARAH-LOUISE BLYTH,⁴ REBECCA BOWLER,^{11,12} BARBARA CATINELLA,^{16,17} LAURENT CHEMIN,¹⁸ STEVEN M. CRAWFORD,¹⁹ CATHERINE CRESS,² ROMEEEL DAVÉ,^{20,2} ROGER DEANE,^{21,22} ERWIN DE BLOK,^{23,4,24} JACINTA DELHAIZE,⁴ KENNETH DUNCAN,²⁰ ED ELSON,² SEAN FEBRUARY,⁸ ERIC GAWISER,⁷ PETER HATFIELD,¹¹ JULIA HEALY,²³ PATRICIA HENNING,^{25,26} KELLEY M. HESS,^{27,23,24} IAN HEYWOOD,^{11,15,8} BENNE W. HOLWERDA,²⁸ MUNIRA HOOSAIN,⁴ JOHN P. HUGHES,⁷ ZACKARY L. HUTCHENS,²⁹ MATT JARVIS,^{11,2} SHEILA KANNAPPAN,²⁹ NEAL KATZ,³⁰ DUŠAN KEREŠ,³¹ MARIE KORSAGA,^{32,33} RENÉE C. KRAAN-KORTEWEG,⁴ PHILIP LAH,³⁴ MICHELLE LOCHNER,^{2,8} NATASHA MADDOX,³⁵ SPHEHSILE MAKHATHINI,²¹ GERHARDT R. MEURER,¹⁶ MARTIN MEYER,¹⁶ DANAIL OBRESCHKOW,¹⁶ SE-HEON OH,³⁶ TOM OOSTERLOO,^{23,24} JOSHUA OPPOR,¹³ HENGXING PAN,^{2,11} D. J. PISANO,^{37,38} NANDRIANINA RANDRIAMIARINARIVO,² SWARA RAVINDRANATH,³⁹ ANJA C. SCHRÖDER,⁴⁰ ROSALIND SKELTON,¹⁴ OLEG SMIRNOV,^{15,8} MATHEW SMITH,^{41,42} RACHEL S. SOMERVILLE,⁴³ RAGHUNATHAN SRINANAND,⁴⁴ LISTER STAVELEY-SMITH,^{16,17} MASAYUKI TANAKA,^{45,46} MATTIA VACCARI,^{1,2,47} WIM VAN DRIEL,⁴⁸ MARC VERHELJEN,⁴⁹ FABIAN WALTER,⁵⁰ JOHN F. WU,³⁹ AND MARTIN A. ZWAAN⁵¹

¹Inter-University Institute for Data Intensive Astronomy (IDIA)

²Department of Physics and Astronomy, University of the Western Cape, Robert Sobukwe Road, Bellville 7535, South Africa

³International Centre for Radio Astronomy Research (ICRAR), Curtin University, Bentley, WA 6102, Australia

⁴Department of Astronomy, University of Cape Town, Private Bag X3, Rondebosch 7700, South Africa

⁵School of Science, Western Sydney University, Locked Bag 1797, Penrith, NSW 2751, Australia

⁶CSIRO Astronomy and Space Science, PO Box 1130, Bentley, WA, 6102, Australia

⁷Department of Physics and Astronomy, Rutgers, the State University of New Jersey, 136 Frelinghuysen Road, Piscataway, NJ 08854-8019, USA

⁸South African Radio Astronomy Observatory, 2 Fir Street, Black River Park, Observatory, 7925, South Africa

⁹Center for Astrophysics and Space Astronomy, Department of Astrophysical and Planetary Sciences, University of Colorado, 389 UCB, Boulder, CO 80309-0389, USA

¹⁰Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121, Bonn, Germany

¹¹Astrophysics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, UK

¹²Jodrell Bank Centre for Astrophysics, School of Physics and Astronomy, University of Manchester, Manchester M13 9PL, UK

¹³Department of Astronomy, University of Wisconsin-Madison, 475 N. Charter Street, Madison, WI 53706, USA

¹⁴South African Astronomical Observatory, P.O. Box 9, Observatory 7935, South Africa

¹⁵Department of Physics and Electronics, Rhodes University, PO Box 94, Makhanda (Grahamstown), 6140, South Africa

¹⁶International Centre for Radio Astronomy Research (ICRAR), The University of Western Australia, 35 Stirling Highway, Perth, WA 6009, Australia

¹⁷ARC Centre of Excellence for All Sky Astrophysics in 3 Dimensions (ASTRO 3D)

¹⁸Centro de Astronomia – CITEVA, Universidad de Antofagasta, Avenida Angamos 601, Antofagasta 1270300, Chile

¹⁹unaffiliated

²⁰Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh, EH9 3HJ, UK

²¹Wits Centre for Astrophysics, School of Physics, University of the Witwatersrand, 1 Jan Smuts Avenue 2000, South Africa

²²Department of Physics, University of Pretoria, Private Bag X20, Pretoria 0028, South Africa

²³ASTRON, the Netherlands Institute for Radio Astronomy, Oude Hoogeveensedijk 4, 7991 PD, Dwingeloo, The Netherlands

²⁴Kapteyn Astronomical Institute, University of Groningen, P.O. Box 800, 9700 AV Groningen, The Netherlands

²⁵National Radio Astronomy Observatory, Pete V. Domenici Science Operations Center, P.O. Box O, 1003 Lopezville Road, Socorro, NM 87801-0387, USA

²⁶Department of Physics and Astronomy, University of New Mexico, 210 Yale Boulevard NE, Albuquerque, NM 87106, USA

²⁷Instituto de Astrofísica de Andalucía (CSIC), Glorieta de la Astronomía s/n, 18008 Granada, Spain

²⁸Department of Physics and Astronomy, 102 Natural Science Building, University of Louisville, Louisville, KY 40292, USA

²⁹Department of Physics & Astronomy, CB3255, University of North Carolina, Chapel Hill, NC 27516, USA

³⁰Department of Astronomy, University of Massachusetts, Amherst, MA 01003, USA

Corresponding author: Marcin Glowacki

marcin.glowacki@curtin.edu.au

³¹Center for Astrophysics and Space Sciences, University of California San Diego, 9500 Gilman Dr, La Jolla, CA 92093, USA

³²Université de Strasbourg, CNRS, Observatoire astronomique de Strasbourg, UMR 7550, F-67000 Strasbourg, France

³³Laboratoire de Physique et de Chimie de l'Environnement, Observatoire d'Astrophysique de l'Université Joseph Ki-Zerbo (ODAOU), 03 BP 7021, Ouaga 03, Burkina Faso

³⁴Research School of Astronomy and Astrophysics, Australian National University, Canberra, ACT 2611, Australia

³⁵University Observatory, Faculty of Physics, Ludwig-Maximilians-Universität, Scheinerstr. 1, 81679 Munich, Germany

³⁶Department of Physics and Astronomy, Sejong University, 209 Neungdong-ro, Gwangjin-gu, Seoul, Republic of Korea

³⁷Department of Physics & Astronomy, West Virginia University, Morgantown, WV 26506, USA

³⁸Gravitational Wave and Cosmology Center, Chestnut Ridge Research Building, Morgantown, WV 26505, USA

³⁹Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218-2410, USA

⁴⁰Max-Planck-Institut für extraterrestrische Physik, Giessenbachstraße 1, D-85748 Garching bei München, Germany

⁴¹School of Physics and Astronomy, University of Southampton, Southampton, SO17 1BJ, UK

⁴²Univ Lyon, Univ Claude Bernard Lyon 1, CNRS, IP2I Lyon / IN2P3, IMR 5822, F-69622, Villeurbanne, France

⁴³Center for Computational Astrophysics, Flatiron Institute, New York, NY 10010, USA

⁴⁴IUCAA, Postbag 4, Ganeshkhind, Pune 411007, India

⁴⁵Department of Astronomical Science, The Graduate University for Advanced Studies, SOKENDAI, 2-21-1 Osawa, Mitaka, Tokyo, 181-8588, Japan

⁴⁶National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo, 181-8588, Japan

⁴⁷INAF - Istituto di Radioastronomia, via Gobetti 101, 40129 Bologna, Italy

⁴⁸GEPI, Observatoire de Paris, PSL Université, CNRS UMR 8111, 5 place Jules Janssen, 92190 Meudon, France

⁴⁹Kapteyn Astronomical Institute, University of Groningen, Landleven 12, 9747 AD, Groningen, The Netherlands

⁵⁰Max-Planck-Institut für Astronomie (MPIA), Königstuhl 16, 69117 Heidelberg, Germany

⁵¹European Southern Observatory, Karl-Schwarzschild-Strasse 2, D-85748 Garching, Germany

(Received February 11, 2022; Revised March 28, 2022; Accepted March 29, 2022)

ABSTRACT

In the local Universe, OH megamasers (OHMs) are detected almost exclusively in infrared-luminous galaxies, with a prevalence that increases with IR luminosity, suggesting that they trace gas-rich galaxy mergers. Given the proximity of the rest frequencies of OH and the hyperfine transition of neutral atomic hydrogen (HI), radio surveys to probe the cosmic evolution of HI in galaxies also offer exciting prospects for exploiting OHMs to probe the cosmic history of gas-rich mergers. Using observations for the Looking At the Distant Universe with the MeerKAT Array (LADUMA) deep HI survey, we report the first untargeted detection of an OHM at $z > 0.5$, LADUMA J033046.20–275518.1 (nicknamed “Nkalakatha”). The host system, WISEA J033046.26–275518.3, is an infrared-luminous radio galaxy whose optical redshift $z \approx 0.52$ confirms the MeerKAT emission line detection as OH at a redshift $z_{\text{OH}} = 0.5225 \pm 0.0001$ rather than HI at lower redshift. The detected spectral line has 18.4σ peak significance, a width of $459 \pm 59 \text{ km s}^{-1}$, and an integrated luminosity of $(6.31 \pm 0.18 [\text{statistical}] \pm 0.31 [\text{systematic}]) \times 10^3 L_{\odot}$, placing it among the most luminous OHMs known. The galaxy’s far-infrared luminosity $L_{\text{FIR}} = (1.576 \pm 0.013) \times 10^{12} L_{\odot}$ marks it as an ultra-luminous infrared galaxy; its ratio of OH and infrared luminosities is similar to those for lower-redshift OHMs. A comparison between optical and OH redshifts offers a slight indication of an OH outflow. This detection represents the first step towards a systematic exploitation of OHMs as a tracer of galaxy growth at high redshifts.

Keywords: megamasers — hydroxyl masers — radio loud quasars — galaxy mergers — starburst galaxies — ULIRG

1. INTRODUCTION

OH megamasers (OHMs) are luminous 18 cm wavelength masers, produced in the centers of luminous and ultra-luminous infrared galaxies (LIRGs and ULIRGs) that have undergone merger-induced starburst activity (Lo 2005), are rich in dense gas (Darling 2007), and in some cases host luminous active galactic nuclei

(AGN; Klöckner et al. 2003). Four 18 cm lines connect the four hyperfine levels within the ${}^2\Pi_{3/2}(J = 3/2)$ ground state of the OH molecule: two main lines at 1665 and 1667 MHz, and two satellite lines at 1612 and 1720 MHz. In contrast to Galactic OH masers, extragalactic OHMs have large line widths and main line flux ratios $F_{1667}/F_{1665} > 1$. These attributes, along with the

weakness of their satellite lines (McBride et al. 2013), can be naturally explained by a model in which OHMs are powered by radiative pumping through $53\ \mu\text{m}$ lines that overlap in velocity, and in which different projected distributions of masing clumps can account for observations of both diffuse and compact emitting structures (Lockett & Elitzur 2008).

The most extensive OHM survey in the local Universe has been conducted with Arecibo, with a focus on the detection of $z > 0.1$ systems (Darling & Giovanelli 2002a). In total, 53 OHMs were detected, spanning the redshift range $0.10 \leq z \leq 0.27$. Although Combes et al. (2021) report a tentative 3σ detection of the 1720 MHz satellite line in emission in a targeted MeerKAT observation of the $z = 0.89$ quasar PKS 1830–211, the *main* lines of OH have not been detected in emission at $z > 0.27$ up to now.

The local demographics of OHMs have been used to predict their occurrence at higher redshifts, where they are likely to represent a significant source of contamination for HI surveys (Briggs 1998; Darling & Giovanelli 2002b; Suess et al. 2016; Roberts et al. 2021). Such surveys will be conducted by the interferometers that include the Square Kilometre Array (SKA) and its pathfinder facilities, such as MeerKAT (Jonas & MeerKAT Team 2016) and the Australian SKA Pathfinder (ASKAP; DeBoer et al. 2009). Although ULIRGs are more prevalent at higher redshifts (Takeuchi et al. 2005) in part because normal star-forming galaxies at earlier epochs are likely to have $L_{\text{FIR}} > 10^{12} L_{\odot}$ even in the absence of recent merging (e.g., Reddy et al. 2008), OHMs have the potential to provide new constraints on the cosmic history of gas-rich mergers.

We report the first untargeted detection of an OHM at $z > 0.27$ from early observations for the Looking At the Distant Universe with the MeerKAT Array (LADUMA; Blyth et al. 2016) deep HI survey. Section 2 describes the acquisition and processing of the MeerKAT data in which the OH line was detected, and Section 3 presents our measurements and interpretation of the line parameters. In Section 4, we discuss the implications of this OHM in the context of previous knowledge of OHM hosts at lower redshifts and future observations probing to higher redshifts; §5 summarizes our conclusions. The paper assumes a flat Λ CDM cosmology with $\Omega_m = 0.3$, $\Omega_{\Lambda} = 0.7$, and $H_0 = 73.3\ \text{km s}^{-1}\ \text{Mpc}^{-1}$ (Wong et al. 2020).

2. OBSERVATIONS

2.1. MeerKAT

MeerKAT is a fixed-configuration array of 64 antennas equipped with receivers spanning the L (900–1670 MHz) and UHF (580–1015 MHz) bands. These overlapping frequency ranges access redshift ranges of $0 \leq z_{\text{HI}} \leq 0.58$ and $0.40 \leq z_{\text{HI}} \leq 1.45$ for the HI line, and $0 \leq z_{\text{OH}} \leq 0.85$ and $0.64 \leq z_{\text{OH}} \leq 1.87$, adopting the stronger of the two main OH lines (at a rest frequency of 1667.359 MHz) to define z_{OH} . The LADUMA deep HI survey is using both L and UHF bands to probe the evolution of gas in galaxies over cosmic time. LADUMA is targeting a single pointing on the sky (03:32:30.4 –28:07:57 J2000) that encompasses the extended Chandra Deep Field South (ECDFS) and lies roughly at the center of near-IR imaging coverage from the VISTA Deep Extragalactic Observations (VIDEO) survey (Jarvis et al. 2013). Because the solid angle of MeerKAT’s primary beam is inversely proportional to the square of the observing frequency, the cosmic volume LADUMA probes for any single spectral line expands with redshift like a trumpet (e.g., a South African vuvuzela). Across the L band in particular, MeerKAT’s circular field of view increases (at the half-power level) from $0.7\ \text{deg}^2$ at 1667 MHz to $0.9\ \text{deg}^2$ at 1420.4 MHz to $2.3\ \text{deg}^2$ at 900 MHz.

The first official L-band survey observation for LADUMA that used the 32k mode of the MeerKAT correlator (featuring 32,768 channels, each of width 26.1 kHz) was taken on 2019 December 12 with 58 of the 64 antennas in operation. The bright radio galaxy PKS 1934–63 was used as a flux and bandpass calibrator (observed for 10 minutes); the nearby quasar PKS 0237–233 was observed as a gain calibrator for 3.5 minutes after each 20-minute observation of the LADUMA field. The total on-source integration time was 7.3 hours.

We reduced the data across the full L band, excluding regions affected by radio frequency interference (RFI), after splitting into 25 MHz spectral windows (SPWs). For the source discussed in this paper, the relevant SPW spans 1086–1111 MHz. Bandpass, flux, and phase calibration were performed using the PROCESSMEERKAT pipeline¹, which is written in Python, uses a purpose-built CASA (McMullin et al. 2007) Singularity container, and employs MPICASA (a parallelized form of CASA). Flux calibration used the Reynolds (1994) model for the spectrum of PKS 1934–63, which is ultimately tied to northern hemisphere calibrators with flux uncertainties of $\sim 5\%$ at $\sim 1\ \text{GHz}$ (e.g., Perley & Butler 2017). The CASA task *tclean* was used with `robust = 0` to create an initial continuum model as a ba-

¹ <https://idia-pipelines.github.io/docs/processMeerKAT>

sis for phase and amplitude self-calibration, after which model continuum visibility data were subtracted from the corrected visibility data using the CASA task *uvsb*. A third-order polynomial fit to the continuum was then calculated and subtracted using the CASA task *uvcontsub* for all channels in each SPW to remove residual continuum emission from the spectral line data. Finally, spectral line cubes were created using *tclean* with *robust* = 0.5 and no cleaning; all channels in the resulting (dirty) cubes were convolved to a common synthesized beam of $14.2'' \times 12.2''$ at a position angle of -15.8° . The RMS per 26.1 kHz channel was found to increase with frequency across the 1086–1111 MHz SPW, with a value of $0.40 \text{ mJy beam}^{-1}$ near 1095 MHz. All data were reduced on the ilifu cloud computing facility².

Visual inspection of the continuum-subtracted 26.1 kHz channel data cube with the Cube Analysis and Rendering Tool for Astronomy (CARTA; Comrie et al. 2020) revealed a bright spectral line at an observed frequency of 1095 MHz, which we designate as LADUMA J033046.20–275518.1 and describe in detail in §3.1.

2.2. Archival data

LADUMA J033046.20–275518.1 has a clear galaxy counterpart in WISE imaging (WISEA J033046.26–275518.3) and in previous radio continuum mapping by the Australia Telescope Large Area Survey (Mao et al. 2012; Franzen et al. 2015). The galaxy has a bent-tail radio morphology (Dehghan et al. 2014), and has been classified as a narrow-angle (or head-tail) radio galaxy (that is, the bent radio structure lies on one side of the optical host galaxy). The source has also been detected in dust emission by the Multiband Imaging Photometer for *Spitzer* (MIPS; Rieke et al. 2004) on the *Spitzer Space Telescope* (Hanish et al. 2015) and by the Spectral and Photometric Imaging Receiver (SPIRE; Griffin et al. 2010) on the *Herschel Space Observatory* (Oliver et al. 2012; Shirley et al. 2021). Table 1 lists selected photometry for the source in the infrared and radio. Optical spectroscopy has yielded two independent redshift measurements of $z = 0.5245$ (Eales et al. 2009) and $z = 0.5247$ (Mao et al. 2012), neither with a quoted uncertainty; in this paper, we average these two measurements and adopt the $\sim 100 \text{ km s}^{-1}$ uncertainty that is typical for AAT redshifts of similar vintage (see, e.g., §1 of Baldry et al. 2014), yielding $z_{\text{opt}} = 0.5246 \pm 0.0005$.

We can determine the host galaxy’s infrared luminosity using the source redshift and far-infrared photom-

etry. By fitting emissivity-weighted blackbody functions to 1000 realizations of the *Spitzer*/MIPS and *Herschel*/SPIRE flux densities between 70–350 μm (sampling their respective uncertainties), we derive a rest-frame dust temperature $T_d = 41.9 \pm 0.6 \text{ K}$, a dust emissivity index $\beta = 1.67 \pm 0.06$ (for $\tau_\nu \propto \nu^\beta$), and a far-IR (rest-frame 42.5–122.5 μm) flux $(6.30 \pm 0.05) \times 10^{-12} \text{ erg s}^{-1} \text{ cm}^{-2}$. Using the cosmology adopted for this paper and the redshift measured in §3.1 below, we then arrive at $L_{\text{FIR}} = (1.576 \pm 0.013) \times 10^{12} L_\odot$, confirming the source as a ULIRG as previously noted by Moncelsi et al. (2011).

3. RESULTS

3.1. OH emission properties

The spatial centroid of LADUMA J033046.20–275518.1 lies at right ascension 03:30:46.20 and declination $-27:55:18.17$ (J2000) with uncertainties of $\pm 0.7''$ (calculated as beam size divided by S/N) — consistent with the centroids determined for its counterparts at other wavelengths, within their own uncertainties of $\sim 0.5''$ or greater. At this position and the observed line frequency, the primary beam correction was a factor $\sim (0.72)^{-1}$, which we applied using CASA before measuring line parameters.

In the right panel of Figure 1, we show the spectrum integrated over a $15''$ aperture, with frequency Doppler-corrected to the heliocentric reference frame and rebinned by a factor of 2. At the observed frequency, each rebinned (52.2 kHz) channel corresponds to 14.3 km s^{-1} in rest-frame velocity. A single-Gaussian fit to the line spectrum yields (with fit uncertainties noted as errors) a central frequency of $\nu = 1095.07 \pm 0.09 \text{ MHz}$, a rest-frame line width $\Delta\nu = 459 \pm 59 \text{ km s}^{-1}$ (full-width at half-maximum, which corresponds to an observed frequency width of 1.68 MHz), a peak flux density of $1.42 \pm 0.16 [\pm 0.07] \text{ mJy}$,³ and a total spectral line flux $F_{\text{line}} = 0.69 \pm 0.02 [\pm 0.03] \text{ Jy km s}^{-1}$ that agrees well with a direct integral of the spectrum ($0.69 \pm 0.02 [\pm 0.03] \text{ Jy km s}^{-1}$).

We note that the 1665 MHz OH line may contribute to the measured line flux and width, although this fainter feature is not yet detected in this single-track observation, nor currently preferred over a single Gaussian fit. Given the presence of a multiwavelength counterpart with a previously measured optical redshift $z = 0.5246$, it is clear that our detection is hydroxyl emission at $z_{\text{OH}} = 0.5225 \pm 0.0001$ (identified as the 1667.359 MHz transition) rather than H I emission at $z_{\text{HI}} = 0.2970$.

² https://docs.ilifu.ac.za/#/about/what_is

³ Errors in brackets represent 5% flux scale uncertainties.

Table 1. Selected photometry for LADUMA J033046.20–275518.1.

Source	Observed wavelength	Flux density	Reference
WISE	$3.4 \mu\text{m}$	$0.122 \pm 0.005 \text{ mJy}$	Wright et al. (2010)
WISE	$4.6 \mu\text{m}$	$0.084 \pm 0.007 \text{ mJy}$	Wright et al. (2010)
WISE	$12 \mu\text{m}$	$0.928 \pm 0.085 \text{ mJy}$	Wright et al. (2010)
<i>Spitzer</i> /MIPS	$70 \mu\text{m}$	$141.11 \pm 0.66 \text{ mJy}^a$	Hanish et al. (2015)
<i>Spitzer</i> /MIPS	$160 \mu\text{m}$	$208.34 \pm 2.13 \text{ mJy}^a$	Hanish et al. (2015)
<i>Herschel</i> /SPIRE	$250 \mu\text{m}$	$90.93 \pm 0.65 \text{ mJy}$	Shirley et al. (2021)
<i>Herschel</i> /SPIRE	$350 \mu\text{m}$	$42.36 \pm 1.53 \text{ mJy}$	Shirley et al. (2021)
<i>Herschel</i> /SPIRE	$500 \mu\text{m}$	$17.6 \pm 3.7 \text{ mJy}$	Shirley et al. (2021)
ATCA	20 cm	$0.42 \pm 0.029 \text{ mJy}$	Franzen et al. (2015)

NOTE— ^a flux density extracted from a point spread function (PSF) fit




Figure 1. *Left:* RGB composite image made from the grz filters of archival Hyper Suprime-Cam (HSC) data. White contours are from the zeroth moment map of our MeerKAT observation with no masking applied, integrated across $\sim 600 \text{ km s}^{-1}$, at multiples of 5σ ($1\sigma = 0.0295 \text{ Jy beam}^{-1} \text{ km s}^{-1}$). The cyan cross indicates the centroid position and uncertainties from the Franzen et al. (2015) observation with ATCA. The $14.2'' \times 12.2''$ MeerKAT synthesized beam is plotted as a dashed yellow ellipse. *Right:* Spectrum of the detected OHM, with frequency Doppler-corrected to a heliocentric reference frame and rebinned from its native 26.1 kHz resolution by a factor of 2. The rest-frame velocity range included in the moment map is shaded grey; the best Gaussian fit to the OH emission profile is overlaid in blue. The red dashed line indicates the frequency (1093.7 MHz)

where the 1667 MHz line would have appeared for $z_{\text{OH}} = z_{\text{opt}}$; that $z_{\text{OH}} < z_{\text{opt}}$ suggests a possible OH outflow.

From the integrated line flux, we use the general relation for spectral line luminosity

$$\frac{L_{\text{line}}}{L_{\odot}} = 1.0234 \left(\frac{\nu}{\text{MHz}} \right) \left(\frac{F_{\text{line}}}{\text{Jy km s}^{-1}} \right) \left(\frac{D_L}{\text{Gpc}} \right)^2 \quad (1)$$

in terms of observed frequency $\nu = 1095.1 \pm 0.1 \text{ MHz}$ and luminosity distance $D_L = 2.85 \pm 0.02 \text{ Gpc}$ for our adopted cosmology, obtaining an equivalent isotropic luminosity of $L_{\text{OH}} = (6.31 \pm 0.18 [\pm 0.31]) \times 10^3 L_{\odot}$.

The left panel of Figure 1 shows contours from the zeroth moment map for the source, integrated over 600 km s^{-1} and overlaid on a composite image made from the grz filters of archival Hyper Suprime-Cam (HSC; Miyazaki et al. 2018) data. The cyan cross indicates the ATCA position and its uncertainties from Franzen et al. (2015), and the yellow ellipse shows the MeerKAT synthesized beam. The OH emission is spatially unresolved, with $\text{S/N} \sim 18.4$ relative to the RMS in the zeroth moment map away from the source. Fig-




Figure 2. Independent velocity channel maps of a $8 \times$ rebinned 26.1 kHz data cube, with velocities indicated relative to our measured systemic redshift of $z_{\text{OH}} = 0.5225$. Contours are multiples of 2σ ($1\sigma = 0.283 \text{ mJy beam}^{-1}$). The $14.2'' \times 12.2''$ synthesized beam is plotted in the top left panel.

ure 2 shows the velocity channel maps of the source after rebinning by an additional factor of 4 to a rest-frame velocity resolution of 57.2 km s^{-1} .

3.2. Confirmation of the 1667 MHz identification

Notwithstanding the good agreement with a previously measured optical redshift, we have considered whether the detected emission could correspond to a weaker OH line (e.g., the 1612 or 1720 MHz satellite line, which can exhibit conjugate behavior: Darling 2004) rather than the main 1667 MHz transition. In this scenario, we would expect an additional detection of the (brighter) 1667 MHz main line at a different frequency (1132.5 or 1061.2 MHz) in our spectrum. No such emission feature is seen at the radio position. We also looked for emission in the satellite lines assuming our detection is indeed the 1667 MHz main line at 1058.9 or 1130.0 MHz, with no such features seen. This result is not surprising given the weakness of satellite lines in OHMs (McBride et al. 2013), although as the LADUMA survey proceeds and reaches greater depths, we may be able to detect them in some systems. We also found no evidence of H I emission or absorption at this redshift, although such features may become evident from the upcoming deeper LADUMA observations.

In Figure 3, we consider the implications of WISE magnitudes and colors for our identification of the detected emission line. We employ the algorithms presented in Roberts et al. (2021), who use machine learning to determine the redshift evolution in WISE magnitude and color space for an OHM host and a typical H I source. The predicted distributions in WISE properties expected for gas-rich disk galaxies emitting H I at redshift $z \sim 0.30$ (red) and a galaxy merger traced through OH at redshift $z \sim 0.52$ (blue) show that an OH identification for our source is plausible for all diagnostics, while an H I identification is not always supported (see right panel). Therefore, we are confident that the emission detected corresponds to the main OH transition at 1667 MHz.

4. DISCUSSION

4.1. OH and FIR luminosities

LADUMA J033046.20–275518.1 is one of the most luminous OHMs known at any redshift; only three sources within the sample of 53 OHMs presented in Darling & Giovanelli (2002a) and Darling & Giovanelli (2002b) have higher L_{OH} values, and it is only marginally less luminous than the $z = 0.1961$ OHM recently detected by Apertif (Hess et al. 2021), which would have $L_{\text{OH}} = 7.18 \times 10^3 L_{\odot}$ for our cosmology. In recognition of this power and its unprecedentedly high redshift, we have given it the nickname “Nkalakatha,” an isiZulu word that means “big boss.” It is not overly surprising that a detection reached in a single LADUMA track will be among the most luminous OHMs when compared to sources from the local Universe at $z < 0.27$. Using the relationship between OH and far-IR luminosities derived by Darling & Giovanelli (2002a) for an Arecibo survey + literature sample of OHMs, i.e., $\log(L_{\text{OH}}/L_{\odot}) = (1.57 \pm 0.11) \log(L_{\text{FIR}}/L_{\odot}) - (15.76 \pm 1.22)$, we find that the predicted OH luminosity for LADUMA J033046.20–275518.1 is $L_{\text{OHpred}} \approx 2.4 \times 10^3 L_{\odot}$. This is roughly a factor of 2.6 smaller than what we observe, but well within the large scatter observed for the local relation (Fig. 4). The dust temperature recovered from the fit to the system’s FIR photometry ($T_d \approx 42 \text{ K}$) is roughly consistent with theoretical expectations and observational results that $T_d \geq 45 \text{ K}$ is required for OH masing to occur (Lockett & Elitzur 2008; Willett et al. 2011), particularly if cooler dust outside the masing region contributes to L_{FIR} , and is at the lower end of the distribution of global dust temperatures for OHM hosts.




Figure 3. Predicted distributions of gas-rich mergers traced by OH at $z_{\text{OH}} \sim 0.52$ (blue), and HI-rich disk galaxies at $z_{\text{HI}} \sim 0.30$ (red), in WISE color-magnitude and color-color space, generated using algorithms presented in Roberts et al. (2021). Contours enclose the upper 99% (dashed), 90%, 80%,... 20%, and 10% of the Gaussian kernel density estimates for the respective galaxy distributions. In the left panel, both OH and HI identifications are supported for LADUMA J033046.20–275518.1, while in the right panel, OH is clearly preferred.




Figure 4. L_{OH} vs. L_{FIR} for known OHMs (Roberts et al. 2022, in preparation). LADUMA J033046.20–275518.1 (green star) is one of the most luminous OHMs on both axes.

We have also considered the FIR–radio flux ratio parameter q , defined by Helou et al. (1985) as

$$q = \log \left\{ \left(\frac{F_{\text{FIR}}}{\text{W m}^{-2}} \right) \left(\frac{10^{29}}{3.75 \times 10^{12} \text{ Hz}} \right) \left(\frac{S_{1.4 \text{ GHz}}}{\text{mJy}} \right)^{-1} \right\} \quad (2)$$

For LADUMA J033046.20–275518.1, we correct the observed radio continuum flux density⁴ (Table 1) to 1.4 GHz in the rest frame assuming a spectrum $S_\nu \propto \nu^{-0.7}$, and arrive at an estimated $q \approx 2.7$. This value is consistent with the infrared–radio correlation for star-forming galaxies with no evidence of AGN (see, e.g., Fig. 16 of Delhaize et al. 2017), higher than the median $q = 2.37$ recently measured for 89 star-forming galaxies in the COSMOS field (An et al. 2021), and much higher than the threshold $q \approx 1.8$ at which galaxies are three times more radio-loud than the mean for star-forming systems (Condon et al. 2002). *Spitzer*/IRAC photometry for this source (Lonsdale et al. 2003) also disfavor an AGN identification according to well-established criteria (Stern et al. 2005; Donley et al. 2012), as does a classification of its optical spectrum (Mao et al. 2012). As a result, we view the far-IR emission from LADUMA J033046.20–275518.1 as more likely due to starburst activity than to a bolometrically significant AGN. Such a conclusion is also not surprising given evidence from the local Universe that ULIRGs hosting OHMs are less likely to show evidence of AGN at infrared wavelengths than non-masing ULIRGs (Willett et al. 2011).

4.2. A possible molecular outflow

⁴ This value is comparable to the value measured from MeerKAT continuum imaging of the LADUMA field, which will be published in a forthcoming paper analogous to Heywood et al. (2022).

Large-scale outflows emanating from the central regions of galaxies and powered by both starbursts and AGN have been known for decades to play an important role in the evolution of galaxies and the intergalactic medium (e.g., Heckman et al. 1990). Such outflows have been detected through OH observations in a number of systems. Baan et al. (1989) detect three distinct outflows in emission in a sample of five OHMs, with one galaxy showing a maximum outflow velocity of 800 km s^{-1} . González-Alfonso et al. (2014) meanwhile find far-IR OH features blueshifted by over 1000 km s^{-1} in Mrk 231, with the central AGN likely responsible for the high mass outflow rate and outflow velocities detected.

The OH emission in LADUMA J033046.20–275518.1 is blueshifted by $407 \pm 118 \text{ km s}^{-1}$ relative to the redshift measured from optical spectroscopy. The simplest explanation for a one-sided velocity offset is a starburst-driven outflow, given the high value of q noted above. However, the situation could be more complex, involving multiple nuclei or a disk close to an (obscured) AGN, possibly interacting with outflows. As our MeerKAT observation of this galaxy and its OH emission is unresolved, higher spatial resolution observations would be required to further characterize the outflow’s extent and energetics. The study of Gowardhan et al. (2018), investigating molecular gas outflows in two starburst ULIRGs in mid- J CO and 18 cm OH lines, is instructive in this regard. CO outflow velocities are seen to exceed 1600 km s^{-1} , with corresponding mass outflow rates of $300\text{--}700 M_{\odot} \text{ yr}^{-1}$; meanwhile, OH outflow velocities are seen to extend to 1000 km s^{-1} , with velocity wings for one source agreeing “remarkably well” with previous detections of outflowing molecular gas. A higher-resolution study of other molecular species (e.g., CO) in LADUMA J033046.20–275518.1 may thus be a productive strategy for confirming the existence of an outflow in OH.

5. CONCLUSIONS

We present the first detection of an OHM in the LADUMA field, LADUMA J033046.20–275518.1 “Nkalakatha,” which is also the highest redshift detection of such a system to date in the main 1667 MHz OH emission line. OH emission is found to be redshifted to $z_{\text{OH}} = 0.5225$, which agrees well with an optical redshift of $z_{\text{opt}} = 0.5246$ for a host galaxy already known to be a ULIRG. The system’s total OH luminosity of $L_{\text{OH}} = (6.31 \pm 0.18 [\pm 0.31]) \times 10^3 L_{\odot}$ makes it one of the most luminous OHMs known (all other 1667 MHz detections have redshifts $z < 0.27$), and is consistent with its large far-IR luminosity. The $\sim 400 \text{ km s}^{-1}$ off-

set between the OH and optical redshifts is most simply explained by a starburst-driven outflow.

This detection highlights the potential of upcoming spectral line surveys, whose wide frequency coverage will enable further high-redshift measurements. Roberts et al. (2021) predict that 83 ± 20 OHMs will be detected in LADUMA alone, which will nearly double the number of known OHMs. Greater numbers are expected in lower redshift but wider area surveys, such as the Widefield ASKAP L-band Legacy All-sky Blind survey (WALLABY; Koribalski et al. 2020), the Apertif Wide-area Extragalactic Survey (AWES; Hess et al., in prep), and the HI component (MIGHTEE-HI; Maddox et al. 2021) of the MeerKAT International GigaHertz Tiered Extragalactic Exploration (MIGHTEE) survey (Jarvis et al. 2016). As LADUMA reaches greater depths, we expect to set further OHM redshift records, which will enable studies of the cosmic rate of gas-rich galaxy mergers and further constrain galaxy evolution models.

ACKNOWLEDGMENTS

The authors sincerely thank Zolile Tibane for suggesting the nickname “Nkalakatha” for LADUMA J033046.20–275518.1. We also thank the anonymous referee for comments that have helped improve the paper.

The MeerKAT telescope is operated by the South African Radio Astronomy Observatory (SARAO; www.sarao.ac.za), which is a facility of the National Research Foundation (NRF), an agency of the Department of Science and Innovation. The authors thank the members of the SARAO engineering, commissioning, and science data processing teams for building and operating an absolutely superb facility. The MeerKAT data presented in this paper were processed using the ilifu cloud computing facility (www.ilifu.ac.za), which is operated by a consortium that includes the University of Cape Town (UCT), the University of the Western Cape, the University of Stellenbosch, Sol Plaatje University, the Cape Peninsula University of Technology and the South African Radio Astronomy Observatory. The ilifu facility is supported by contributions from the Inter-University Institute for Data Intensive Astronomy (IDIA, which is a partnership between the UCT, the University of Pretoria and the University of the Western Cape), the Computational Biology division at UCT, and the Data Intensive Research Initiative of South Africa (DIRISA). Data processing used pipelines that were developed at IDIA and are available at <https://idia-pipelines.github.io>.

MG acknowledges support from IDIA and was partially supported by the Australian Government through

the Australian Research Council’s Discovery Projects funding scheme (DP210102103). AKM and AJB acknowledge support from NSF grant AST-1814421. AKM also thanks the LSSTC Data Science Fellowship Program, which is funded by the LSST Corporation, NSF grant OAC-1829740, the Brinson Foundation, and the Moore Foundation; his participation in the program has been helpful for this work. HR and JeD acknowledge support from NSF grant AST-1814648; MB and JO acknowledge support from NSF grant AST-1814682; RB acknowledges support from an STFC Ernest Rutherford Fellowship (grant number ST/T003596/1); JaD and HP acknowledge the financial assistance of SARA0; KMH acknowledges funding from the State Agency for Research of the Spanish Ministry of Science, Innovation and Universities through the “Center of Excellence Severo Ochoa” awarded to

the Instituto de Astrofísica de Andalucía (SEV-2017-0709); from grant RTI2018-096228-B-C31 (Ministry of Science, Innovation and Universities / State Agency for Research / European Regional Development Funds, European Union); and from the coordination of the participation in SKA-SPAIN, funded by the Ministry of Science and innovation (MICIN); and ZLH and SK acknowledge support from NSF grant AST-1814486. Parts of this research were supported by the Australian Research Council Centre of Excellence for All Sky Astrophysics in 3 Dimensions (ASTRO 3D), through project number CE170100013, and by the South African Research Chairs Initiative of the Department of Science and Technology and the NRF.

Facilities: MeerKAT

REFERENCES

- An, F., Vaccari, M., Smail, I., et al. 2021, *MNRAS*, 507, 2643, doi: [10.1093/mnras/stab2290](https://doi.org/10.1093/mnras/stab2290)
- Baan, W. A., Haschick, A. D., & Henkel, C. 1989, *ApJ*, 346, 680, doi: [10.1086/168050](https://doi.org/10.1086/168050)
- Baldry, I. K., Alpaslan, M., Bauer, A. E., et al. 2014, *MNRAS*, 441, 2440, doi: [10.1093/mnras/stu727](https://doi.org/10.1093/mnras/stu727)
- Blyth, S., Baker, A. J., Holwerda, B., et al. 2016, in MeerKAT Science: On the Pathway to the SKA, 4
- Briggs, F. H. 1998, *A&A*, 336, 815
- Combes, F., Gupta, N., Muller, S., et al. 2021, *A&A*, 648, A116, doi: [10.1051/0004-6361/202040167](https://doi.org/10.1051/0004-6361/202040167)
- Comrie, A., Wang, K.-S., Ford, P., et al. 2020, CARTA: The Cube Analysis and Rendering Tool for Astronomy, 1.3.0, Zenodo, doi: [10.5281/zenodo.3746095](https://doi.org/10.5281/zenodo.3746095)
- Condon, J. J., Cotton, W. D., & Broderick, J. J. 2002, *AJ*, 124, 675, doi: [10.1086/341650](https://doi.org/10.1086/341650)
- Darling, J. 2004, *ApJ*, 612, 58, doi: [10.1086/422450](https://doi.org/10.1086/422450)
- . 2007, *ApJL*, 669, L9, doi: [10.1086/523756](https://doi.org/10.1086/523756)
- Darling, J., & Giovanelli, R. 2002a, *AJ*, 124, 100, doi: [10.1086/341166](https://doi.org/10.1086/341166)
- . 2002b, *ApJ*, 572, 810, doi: [10.1086/340372](https://doi.org/10.1086/340372)
- DeBoer, D. R., Gough, R. G., Bunton, J. D., et al. 2009, *IEEE Proceedings*, 97, 1507, doi: [10.1109/JPROC.2009.2016516](https://doi.org/10.1109/JPROC.2009.2016516)
- Dehghan, S., Johnston-Hollitt, M., Franzen, T. M. O., Norris, R. P., & Miller, N. A. 2014, *AJ*, 148, 75, doi: [10.1088/0004-6256/148/5/75](https://doi.org/10.1088/0004-6256/148/5/75)
- Delhaize, J., Smolčić, V., Delvecchio, I., et al. 2017, *A&A*, 602, A4, doi: [10.1051/0004-6361/201629430](https://doi.org/10.1051/0004-6361/201629430)
- Donley, J. L., Koekemoer, A. M., Brusa, M., et al. 2012, *ApJ*, 748, 142, doi: [10.1088/0004-637X/748/2/142](https://doi.org/10.1088/0004-637X/748/2/142)
- Eales, S., Chapin, E. L., Devlin, M. J., et al. 2009, *ApJ*, 707, 1779, doi: [10.1088/0004-637X/707/2/1779](https://doi.org/10.1088/0004-637X/707/2/1779)
- Franzen, T. M. O., Banfield, J. K., Hales, C. A., et al. 2015, *MNRAS*, 453, 4020, doi: [10.1093/mnras/stv1866](https://doi.org/10.1093/mnras/stv1866)
- González-Alfonso, E., Fischer, J., Graciá-Carpio, J., et al. 2014, *A&A*, 561, A27, doi: [10.1051/0004-6361/201321709](https://doi.org/10.1051/0004-6361/201321709)
- Gowardhan, A., Spoon, H., Riechers, D. A., et al. 2018, *ApJ*, 859, 35, doi: [10.3847/1538-4357/aabccc](https://doi.org/10.3847/1538-4357/aabccc)
- Griffin, M. J., Abergel, A., Abreu, A., et al. 2010, *A&A*, 518, L3, doi: [10.1051/0004-6361/201014519](https://doi.org/10.1051/0004-6361/201014519)
- Hanish, D. J., Capak, P., Teplitz, H. I., et al. 2015, *ApJS*, 217, 17, doi: [10.1088/0067-0049/217/1/17](https://doi.org/10.1088/0067-0049/217/1/17)
- Heckman, T. M., Armus, L., & Miley, G. K. 1990, *ApJS*, 74, 833, doi: [10.1086/191522](https://doi.org/10.1086/191522)
- Helou, G., Soifer, B. T., & Rowan-Robinson, M. 1985, *ApJL*, 298, L7, doi: [10.1086/184556](https://doi.org/10.1086/184556)
- Hess, K. M., Roberts, H., Dénes, H., et al. 2021, *A&A*, 647, A193, doi: [10.1051/0004-6361/202040019](https://doi.org/10.1051/0004-6361/202040019)
- Heywood, I., Jarvis, M. J., Hale, C. L., et al. 2022, *MNRAS*, 509, 2150, doi: [10.1093/mnras/stab3021](https://doi.org/10.1093/mnras/stab3021)
- Jarvis, M., Taylor, R., Agudo, I., et al. 2016, in MeerKAT Science: On the Pathway to the SKA, 6. <https://arxiv.org/abs/1709.01901>
- Jarvis, M. J., Bonfield, D. G., Bruce, V. A., et al. 2013, *MNRAS*, 428, 1281, doi: [10.1093/mnras/sts118](https://doi.org/10.1093/mnras/sts118)
- Jonas, J., & MeerKAT Team. 2016, in MeerKAT Science: On the Pathway to the SKA, 1
- Klöckner, H.-R., Baan, W. A., & Garrett, M. A. 2003, *Nature*, 421, 821, doi: [10.1038/nature01381](https://doi.org/10.1038/nature01381)
- Koribalski, B. S., Staveley-Smith, L., Westmeier, T., et al. 2020, *Ap&SS*, 365, 118, doi: [10.1007/s10509-020-03831-4](https://doi.org/10.1007/s10509-020-03831-4)

- Lo, K. Y. 2005, *ARA&A*, 43, 625,
doi: [10.1146/annurev.astro.41.011802.094927](https://doi.org/10.1146/annurev.astro.41.011802.094927)
- Lockett, P., & Elitzur, M. 2008, *ApJ*, 677, 985,
doi: [10.1086/533429](https://doi.org/10.1086/533429)
- Lonsdale, C. J., Smith, H. E., Rowan-Robinson, M., et al. 2003, *PASP*, 115, 897, doi: [10.1086/376850](https://doi.org/10.1086/376850)
- Maddox, N., Frank, B. S., Ponomareva, A. A., et al. 2021, *A&A*, 646, A35, doi: [10.1051/0004-6361/202039655](https://doi.org/10.1051/0004-6361/202039655)
- Mao, M. Y., Sharp, R., Norris, R. P., et al. 2012, *MNRAS*, 426, 3334, doi: [10.1111/j.1365-2966.2012.21913.x](https://doi.org/10.1111/j.1365-2966.2012.21913.x)
- McBride, J., Heiles, C., & Elitzur, M. 2013, *ApJ*, 774, 35,
doi: [10.1088/0004-637X/774/1/35](https://doi.org/10.1088/0004-637X/774/1/35)
- McMullin, J. P., Waters, B., Schiebel, D., Young, W., & Golap, K. 2007, in *Astronomical Society of the Pacific Conference Series*, Vol. 376, *Astronomical Data Analysis Software and Systems XVI*, ed. R. A. Shaw, F. Hill, & D. J. Bell, 127
- Miyazaki, S., Komiyama, Y., Kawanomoto, S., et al. 2018, *PASJ*, 70, S1, doi: [10.1093/pasj/psx063](https://doi.org/10.1093/pasj/psx063)
- Moncelsi, L., Ade, P. A. R., Chapin, E. L., et al. 2011, *ApJ*, 727, 83, doi: [10.1088/0004-637X/727/2/83](https://doi.org/10.1088/0004-637X/727/2/83)
- Oliver, S. J., Bock, J., Altieri, B., et al. 2012, *MNRAS*, 424, 1614, doi: [10.1111/j.1365-2966.2012.20912.x](https://doi.org/10.1111/j.1365-2966.2012.20912.x)
- Perley, R. A., & Butler, B. J. 2017, *ApJS*, 230, 7,
doi: [10.3847/1538-4365/aa6df9](https://doi.org/10.3847/1538-4365/aa6df9)
- Reddy, N. A., Steidel, C. C., Pettini, M., et al. 2008, *ApJS*, 175, 48, doi: [10.1086/521105](https://doi.org/10.1086/521105)
- Reynolds, J. 1994, *ATNF Technical Memos*, 39.3/040
- Rieke, G. H., Young, E. T., Engelbracht, C. W., et al. 2004, *ApJS*, 154, 25, doi: [10.1086/422717](https://doi.org/10.1086/422717)
- Roberts, H., Darling, J., & Baker, A. J. 2021, *ApJ*, 911, 38,
doi: [10.3847/1538-4357/abe944](https://doi.org/10.3847/1538-4357/abe944)
- Shirley, R., Duncan, K., Campos Varillas, M. C., et al. 2021, *MNRAS*, 507, 129, doi: [10.1093/mnras/stab1526](https://doi.org/10.1093/mnras/stab1526)
- Stern, D., Eisenhardt, P., Gorjian, V., et al. 2005, *ApJ*, 631, 163, doi: [10.1086/432523](https://doi.org/10.1086/432523)
- Suess, K. A., Darling, J., Haynes, M. P., & Giovanelli, R. 2016, *MNRAS*, 459, 220, doi: [10.1093/mnras/stw666](https://doi.org/10.1093/mnras/stw666)
- Takeuchi, T. T., Buat, V., & Burgarella, D. 2005, *A&A*, 440, L17, doi: [10.1051/0004-6361:200500158](https://doi.org/10.1051/0004-6361:200500158)
- Willett, K. W., Darling, J., Spoon, H. W. W., Charmandaris, V., & Armus, L. 2011, *ApJ*, 730, 56,
doi: [10.1088/0004-637X/730/1/56](https://doi.org/10.1088/0004-637X/730/1/56)
- Wong, K. C., Suyu, S. H., Chen, G. C. F., et al. 2020, *MNRAS*, 498, 1420, doi: [10.1093/mnras/stz3094](https://doi.org/10.1093/mnras/stz3094)
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, 140, 1868, doi: [10.1088/0004-6256/140/6/1868](https://doi.org/10.1088/0004-6256/140/6/1868)