LARGE INTERSECTION PROPERTY FOR LIMSUP SETS IN METRIC **SPACE**

ZHANG-NAN HU, BING LI, AND LINQI YANG*

Abstract. We show that limsup sets generated by a sequence of open sets in compact Ahlfors s-regular space $(X, \mathcal{B}, \mu, \rho)$ belong to the classes of sets with large intersections with index λ , denoted by $\mathcal{G}^{\lambda}(X)$, under some conditions. In particular, this provides a lower bound on Hausdorff dimension of such sets. These results are applied to obtain that limsup random fractals with indices γ_2 and δ belong to $\mathcal{G}^{s-\delta-\gamma_2}(\tilde{X})$ almost surely, and random covering sets with exponentially mixing property belong to $\mathcal{G}^{s_0}(X)$ almost surely, where s_0 equals to the corresponding Hausdorff dimension of covering sets almost surely. We also investigate the large intersection property of limsup sets generated by rectangles in metric space.

1. INTRODUCTION

Sets with large intersection were introduced by Falconer in [\[9\]](#page-18-0). Given $s \in (0, d]$, he defined $\mathcal{G}^s(\mathbb{R}^d)$ to be the class of all G_{δ} sets F in \mathbb{R}^d such that

> $\dim_{\mathrm{H}} \bigcap^{\infty}$ $n=1$ $f_n(F) \geq s$

holds for all sequences of similarity transformations $\{f_n\}_{n\geq 1}$, where dim_H denotes the Hausdorff dimension. It is known that $\mathcal{G}^s(\mathbb{R}^d)$ is the maximal class of G_{δ} sets satisfying (i) Hausdorff dimension larger than s, and (ii) closed under similarity transformations and countable intersections. The packing dimension of sets in $\mathcal{G}^s(\mathbb{R}^d)$ is d from the fact that they are dense G_{δ} sets. In [\[9\]](#page-18-0), Falconer also gave several equivalent definitions and established various properties of $\mathcal{G}^s(\mathbb{R}^d)$. In 2004, Bugeaud [\[3\]](#page-18-1) generalized the class for more general gauge functions. Later, Negreira and Sequeira [\[21\]](#page-18-2) extended the class of sets in Euclidean space with large intersection property to metric space endowed with doubling measure.

There are many applications of large intersection property. Lots of mathematicians applied this property for estimating Hausdorff dimension from below in the study of Diophantine approximations, and refer to [\[3\]](#page-18-1), [\[4\]](#page-18-3), [\[9\]](#page-18-0), [\[21\]](#page-18-2) for more details. In 2007, Durand [\[5\]](#page-18-4) investigated the size and large intersection properties of limsup sets generated by homogeneous ubiquitous systems in \mathbb{R}^d . In 2010, Durand [\[8\]](#page-18-5) showed that random covering sets $\limsup B(\xi_n, r_n)$ are sets with large intersection, where $\{\xi_n\}_{n\geq 1}$ are independent and $n\rightarrow\infty$ uniformly distributed random variables on the circle. In 2019, Persson [\[22\]](#page-18-6) proved that dynamical covering sets lim sup $B(T^n x, n^{-\alpha})$ has large intersection property for dynamical $n\rightarrow\infty$ systems (T, μ) which have summable decay of correlations. Ding [\[4\]](#page-18-3) showed that under a full Hausdorff measure assumption, the limsup sets generated by rectangles with some conditions in compact metric space are sets with large intersection. In 2021, Persson [\[23\]](#page-18-7) considered various sequences of open sets with general shapes, proved the corresponding limsup sets have large intersection properties, and obtained the lower bound on the Hausdorff dimension. Aubry and Jaffard [\[1\]](#page-18-8) noticed that large intersection property also

Key words and phrases. limsup sets; large intersection property; metric space; limsup random fractals; random covering sets.

^{*} Corresponding author.

occurred in probability theory, such as the multifractal analysis of random wavelet series. See also [\[6,](#page-18-9)[7,](#page-18-10)[10\]](#page-18-11) for more study in fractals, dynamical systems and the multifractal analysis of other stochastic processes.

Limsup sets, the upper limits of sequences of sets, play an important role in many areas, such as random covering problem, shrinking target problem, the study of Brownian motion and so on. Motivated by the study on sets with large intersections, we are interested in the large intersection properties of limsup sets generated by open sets in metric spaces $(X, \mathscr{B}, \mu, \rho)$ and those applications.

Definition 1.1. A Borel measure μ on metric space (X, ρ) is Ahlfors s-regular $(0 < s < \infty)$ if there exists a constant $1 \leq C \leq \infty$ such that

$$
C^{-1}r^s \le \mu\big(B(x,r)\big) \le Cr^s \tag{1.1}
$$

holds for all $x \in X$ and $0 \le r \le \text{diam } X$, where $\text{diam } X$ is the diameter of X. Here $B(x,r) = \{y \in X : \rho(x,y) < r\}$ is an open ball with centre x and radius r. A space $(X, \mathscr{B}, \mu, \rho)$ is said to be Ahlfors s-regular if μ satisfies formula [\(1.1\)](#page-1-0).

In this paper, we consider a probability space $(X, \mathscr{B}, \mu, \rho)$ where μ is Ahlfors s-regular $(0 \lt s \lt \infty)$. For $B = B(x, r)$, we adopt the convention that $cB = B(x, cr)$, $B^t =$ $B(x, r^{t/s})$ and $B = \emptyset$ if $r = 0$.

Definition 1.2. For $0 \le t \le s$, the t-potential at $y \in X$ of the measure μ is defined as

$$
\phi_t(\mu, y) = \int_X \rho(x, y)^{-t} d\mu(x).
$$

The t-energy of the measure μ is defined as

$$
I_t(\mu) = \int_X \int_X \rho(x, y)^{-t} d\mu(x) d\mu(y).
$$

Denote

$$
\phi_t(\mu, U, y) = \int_U \rho(x, y)^{-t} d\mu(x),
$$

$$
I_t(\mu, U) = \int_U \int_U \rho(x, y)^{-t} d\mu(x) d\mu(y)
$$

for $U \in \mathcal{B}$.

Theorem 1.1. Let $(X, \mathcal{B}, \mu, \rho)$ be a compact Ahlfors s-regular space $(0 \lt s \lt \infty)$, and ${B_n}_{n>1}$ be a sequence of balls in X with diam B_n decreasing to 0 as $n \to \infty$. For $n \geq 1$, E_n is an open subset of B_n and let

$$
\lambda = \sup \Big\{ t \ge 0 \colon \sup_{n \ge 1} \frac{I_t(\mu, E_n) \mu(B_n)}{\mu(E_n)^2} < \infty \Big\}.
$$

Then μ (lim sup $\limsup_{n\to\infty} B_n$ = 1 *implies that* $\limsup_{n\to\infty}$ $\max_{n\to\infty} E_n \in \mathcal{G}^{\lambda}(X).$

Corollary 1.1. Under the setting in Theorem [1.1,](#page-1-1) if μ (lim sup $\limsup_{n\to\infty} B_n$ = 1, then we have $\dim_{\mathrm{H}} \big(\limsup$ $\limsup_{n\to\infty} E_n$ $\Big) \geq \lambda$ and $\dim_{\rm P} \Big(\limsup_{n\to\infty}$ $\limsup_{n\to\infty} E_n\Big) = s, \hbox{ where } \dim_{\rm P} \hbox{ denotes the packing }$ dimension.

When ${E_n}_{n>1}$ is a sequence of balls, an important tool in determining the Hausdorff dimension of $\limsup E_n$ is the Mass Transference Principle, which was established by Beres $n\rightarrow\infty$ nevich and Velani [\[2\]](#page-18-12).

Theorem 1.2 (Mass Transference Principle [\[2\]](#page-18-12)). Let $(X, \mathcal{B}, \mu, \rho)$ be a locally compact Ahlfors s-regular space $(0 \lt s \lt \infty)$, Let $\{E_n\}_{n\geq 1}$ be a sequence of balls in X with diam $E_n \to 0$ as $n \to \infty$. Let $t > 0$ and suppose that

$$
\mathcal{H}^s(\limsup_{n\to\infty}E_n^t)=\mathcal{H}^s(X).
$$

Then,

$$
\mathcal{H}^t(\limsup_{n\to\infty}E_n)=\mathcal{H}^t(X).
$$

Here $\mathcal{H}^t(F)$ denotes the Hausdorff t-measure of a set $F \subset X$.

For an Ahlfors s-regular space $(X, \mathcal{B}, \mu, \rho)$, Heinonen [\[12,](#page-18-13) Section 8.7] proved that $C^{-1}\mathcal{H}^s \leq \mu \leq C'\mathcal{H}^s$, where $C' \geq 1$ is a constant. Then by Theorem [1.2,](#page-2-0) given $t \in (0, s]$, we can deduce that $\dim_H \left(\limsup E_n \right) \ge t$, if $\mu \left(\limsup E_n^t \right) = 1$. Such estimation can be derived by Corollary [1.1](#page-1-2) when X is compact. In addition, as the following corollary shown, $n\rightarrow\infty$ $\limsup E_n$ also has large intersection property. $n \rightarrow \infty$

Corollary 1.2. Let $(X, \mathcal{B}, \mu, \rho)$ be a compact Ahlfors s-regular space $(0 \lt s \lt \infty)$, and ${B_n}_{n\geq 1}$ be a sequence of balls with diam B_n decreasing to 0 as $n \to \infty$. For $t \in (0, s]$, if $\mu\Big(\limsup$ n→∞ B_n^t = 1, then lim sup $\limsup_{n\to\infty} B_n \in \mathcal{G}^t(X).$

Applying Theorem [1.1](#page-1-1) and Corollary [1.2,](#page-2-1) we prove that limsup random fractals, random covering sets and limsup sets generated by rectangles have large intersection property (Theorems 4.1, 4.7 and 4.8).

The rest of this paper is organized as follows. In Section 2 we give a brief review of the class of sets with large intersection properties. In Sections 3, we give the proofs of Theorem [1.1](#page-1-1) and Corollary [1.2](#page-2-1) which are our main results. In the last section, there are some examples. We apply our results to the study of the large intersection properties of limsup random fractals under some conditions and random covering sets lim sup $B(\xi_n, r_n)$, $n\rightarrow\infty$

where the centers $\{\xi_n\}_{n\geq 1}$ are uniformly distributed and exponentially mixing random variables, are sets with large intersection properties. We also show limsup sets generated by rectangles have large intersection property.

2. Preliminaries

In this section, we refer (X, ρ) to general metric space, and μ is a Borel measure on X. Denote $B(x,r) = \{y \in X : \rho(x,y) < r\}$ and $\overline{B}(x,r) = \{y \in X : \rho(x,y) \leq r\}.$

2.1. Generalized dyadic cubes in metric spaces.

Definition 2.1. A metric space (X, ρ) has the finite doubling property if any closed ball $B(x, 2r) \subset X$ may be covered by finitely many closed balls of radius r. Furthermore, such a space is doubling if there exists $N \in \mathbb{N}$ independent of x and r such that $\overline{B}(x, 2r)$ can be covered by at most N balls of radius r.

Letting (X, ρ) be a metric space with the finite doubling property, Käenmäki, Rajala and Suomala [\[16\]](#page-18-14) showed that there exists a nesting family of cubes which are similar as the dyadic cubes of Euclidean spaces.

Theorem 2.1 ([\[16\]](#page-18-14)). Let (X, ρ) be a metric space with the finite doubling property and let $0 < b < \frac{1}{3}$ be a constant. Then there exists a collection $\{Q_{n,i}: n \in \mathbb{Z}, i \in \mathbb{N}_n \subset \mathbb{N}\}\$ of Borel sets that have the following properties:

- (1) $X = \bigcup_{i \in \mathbb{N}_n} Q_{n,i}$ for every $n \in \mathbb{Z}$.
- (2) $Q_{n,i} \cap Q_{m,j} = \emptyset$ or $Q_{n,i} \subset Q_{m,j}$, where $n, m \in \mathbb{Z}$, $n \geq m$, $i \in \mathbb{N}_n$ and $j \in \mathbb{N}_m$.

(3) For every $n \in \mathbb{Z}$ and $i \in \mathbb{N}_n$, there exists a point $x_{n,i} \in X$ such that

$$
B(x_{n,i}, c_1 b^n) \subset Q_{n,i} \subset \overline{B}(x_{n,i}, c'_1 b^n), \tag{2.1}
$$

where $c_1 = \frac{1}{2} - \frac{b}{1 - b}$ $\frac{b}{1-b}$, $c'_1 = \frac{1}{1-b}$ $\frac{1}{1-b}$.

- (4) There exists a point $x_0 \in X$ so that for every $n \in \mathbb{Z}$, there is an index $i \in \mathbb{N}_n$ with $B(x_0, c_1 b^n) \subset Q_{n,i}.$
- (5) $\{x_{n,i} : i \in \mathbb{N}_n\} \subset \{x_{n+1,i} : i \in \mathbb{N}_{n+1}\}\$ for all $n \in \mathbb{Z}$.

Proposition 2.1. Let (X, ρ) be a metric space with an Ahlfors s-regular measure μ .

(1) Suppose that μ is a probability measure, then for $n \in \mathbb{Z}$, we have

$$
C^{-1}c_1'^{-s}b^{-ns} \leq #\{Q_{n,i}: i \in \mathbb{N}_n\} \leq Cc_1^{-s}b^{-ns}.
$$

(2) The metric space (X, ρ) has the doubling property.

Proof. (1) From the construction of $\{Q_{n,i} : n \in \mathbb{Z}, i \in \mathbb{N}_n \subset \mathbb{N}\}\)$, we see that for any $n \in \mathbb{Z}, Q_{n,i} \cap Q_{n,j} = \emptyset$ for $i \neq j \in \mathbb{N}_n$. Since μ is Ahlfors s-regular, combining (1), (3) in Theorem [2.1](#page-2-2) and the equalities [\(1.1\)](#page-1-3), we have

$$
\mu(X) = \sum_{i \in \mathbb{N}_n} \mu(Q_{n,i}) \ge \sum_{i \in \mathbb{N}_n} \mu(B(x_{n,i}, c_1 b^n)) \ge C^{-1} \# \mathbb{N}_n (c_1 b^n)^s,
$$

and

$$
\mu(X) = \sum_{i \in \mathbb{N}_n} \mu(Q_{n,i}) \le \sum_{i \in \mathbb{N}_n} \mu(\overline{B}(x_{n,i}, c'_1 b^n)) \le C \# \mathbb{N}_n (c'_1 b^n)^s,
$$

which implies that

$$
C^{-1}c_1'^{-s}b^{-ns} \le \#\mathbb{N}_n \le Cc_1^{-s}b^{-ns}.
$$

(2) Given $x \in X$ and $r > 0$, let $n_0 = \min\{n \ge 1 : c'_1 b^n < r\}$. Let

$$
I = \{ i \in \mathbb{N}_{n_0} \colon Q_{n_0, i} \cap \overline{B}(x, 2r) \neq \emptyset \}.
$$

Then $\{B(x_{n_0,i}, r)\}_{i\in I}$ is a cover of $B(x, 2r)$. Note that $\bigcup_{i\in I} Q_{n_0,i} \subset B(x, 4r)$ and $r \leq$ $c'_1 b^{n_0-1}$, then by (2.1) , we have

$$
C4^{s}r^{s} \geq \mu(B(x, 4r)) \geq \mu\left(\bigcup_{i \in I} Q_{n_{0}, i}\right)
$$

=
$$
\sum_{i \in I} \mu(Q_{n_{0}, i}) \geq \sum_{i \in I} \mu(B(x_{n_{0}, i}, c_{1}b^{n_{0}}))
$$

$$
\geq C^{-1}c_{1}^{s}b^{n_{0}s} \# I \geq C^{-1}(bc_{1}/c_{1}')^{s}r^{s} \# I,
$$

$$
\leq C^{2}\left(\frac{4c_{1}'}{bc}\right)^{s}.
$$

which follows that $\#I \leq C^2 \left(\frac{4c'_1}{bc_1} \right)$ \setminus^s

Therefore for a compact Ahlfors s-regular space $(X, \mathscr{B}, \mu, \rho)$, there exists the family ${Q_{n,i}: n \geq 0, i \in \mathbb{N}_n}$ satisfying the properties in Theorem [2.1,](#page-2-2) which are called "generalized dyadic cubes". For convenience, we write $\mathcal{Q}_0 = \{X\}$ and $\mathcal{Q}_n = \{Q_{n,i} : i \in \mathbb{N}_n\}$ for $n \geq 1$, and $\mathcal{Q} = \bigcup_{n \geq 0} \mathcal{Q}_n$.

2.2. Large intersection properties in metric spaces.

Definition 2.2. Let (X, ρ) be a metric space. A Borel measure μ on X is said to be doubling if it is finite and positive in every ball and there exists a constant $1 \leq c_2 < \infty$ such that for all $x \in X$ and $r > 0$,

$$
0 < \mu\left(\overline{B}(x, 2r)\right) \le c_2 \mu\left(\overline{B}(x, r)\right) < \infty.
$$

Let (X, ρ) be a metric space endowed with doubling measure μ . Denote $\tau = \dim_{\mathrm{H}} X$, and τ is finite, see [\[11,](#page-18-15) Section 13]. In [\[21\]](#page-18-2), Negreira and Sequeira gave the definition of the classes of G_{δ} sets with large intersection property. Recall that a G_{δ} set is a countable intersection of open sets.

Definition 2.3. Let $0 < t < \tau$. Given $F \subset X$, define the net content

$$
\mathscr{M}^t_\infty(F) = \inf \Biggl\{ \sum_{i \ge 1} \mu(Q_i)^{t/\tau} : F \subset \bigcup_{i \ge 1} Q_i \text{ where } Q_i \in \mathcal{Q} \Biggr\}.
$$

We denote $\mathcal{G}^t(X)$ the class of all G_{δ} sets $F \subset X$ such that for any $t' < t$,

$$
\mathcal{M}_{\infty}^{t'}(F \cap Q) = \mathcal{M}_{\infty}^{t'}(Q)
$$

holds for all $Q \in \mathcal{Q}$.

Remark 2.1. (1) Negreira and Sequeira [\[21\]](#page-18-2) showed that the net contents given by different dyadic decomposition are equivalent, and pointed out that the net content is defined by $\mu(Q)^{t/\tau}$ instead of $(\text{diam } Q)^t$, since the function $Q \mapsto (\text{diam } Q)^t$ is not sub-additive if $t > 1$ which is important in the study of \mathcal{M}_{∞}^{t} and large intersection property.

(2) Notice that the class $\mathcal{G}^t(X)$ given in Definition [2.3](#page-4-0) depends on μ . Since Ahlfors s-regular measures on X are equivalent, therefore the classes $\mathcal{G}^t(X)$ defined by different Ahlfors s-regular measures are same. In this paper we only consider such measures, hence we use the notation $\mathcal{G}^t(X)$ instead of $\mathcal{G}^t_\mu(X)$.

There are several equivalent definitions of $\mathcal{G}^t(X)$, see [\[21\]](#page-18-2). In this paper, we will use the following equivalent definition.

Theorem 2.2 ([\[21\]](#page-18-2)). Let (X, ρ) be a metric space enowed with doubling measure μ , and $\tau = \dim_{\mathrm{H}} X$. Take $F \subset X$ a G_{δ} subset and $0 < t \leq \tau$. Then the following statements are equivalent:

(1) For all generalized dyadic cubes $Q \in \mathcal{Q}$, we have

$$
\mathcal{M}_{\infty}^{t'}(F \cap Q) = \mathcal{M}_{\infty}^{t'}(Q) \quad (\forall t' < t).
$$

(2) There exists a constant $0 < c \leq 1$ such that for all generalized dyadic cubes $Q \in \mathcal{Q}$, we have

$$
\mathcal{M}^{t'}_{\infty}(F \cap Q) \ge c \mathcal{M}^{t'}_{\infty}(Q) \quad (\forall t' < t). \tag{2.2}
$$

Remark 2.2. Note that, given $0 < t \leq \tau$, $\mathscr{M}^t_\infty(Q) = \mu(Q)^{t/\tau}$ holds for all dyadic cubes $Q \in \mathcal{Q}$. Then [\(2.2\)](#page-4-1) can be rewritten as

$$
\mathcal{M}^{t'}_{\infty}(F \cap Q) \ge c\mu(Q)^{t'/\tau} \quad (\forall t' < t). \tag{2.3}
$$

The following are some properties of the class $\mathcal{G}^t(X)$.

Proposition 2.2 ([21]). Let (X, ρ) be a metric space endowed with doubling measure μ , and $\tau = \dim_{\mathrm{H}} X$ and take $0 < t \leq \tau$.

(1) If $0 < t_1 \leq t$, then $\mathcal{G}^t(X) \subset \mathcal{G}^{t_1}(X)$.

- (2) If $F \subset E \subset X$, where E, F are G_{δ} sets, and $F \in \mathcal{G}^t(X)$, then $E \in \mathcal{G}^t(X)$.
- (3) If (X, ρ) is complete, then $\mathcal{G}^t(X)$ is closed under countable intersections.
- (4) When μ is Ahlfors τ -regular, then $F \in \mathcal{G}^t(X)$ implies that $\dim_\text{H} F \geq t$.

Remark 2.3. Let (X, ρ) be a metric space endowed with doubling measure μ , and $\tau =$ $\dim_{\mathrm{H}} X$. Take $0 < t \leq \tau$, then from the definition of $\mathcal{G}^t(X)$, we have

$$
\mathcal{G}^t(X) = \bigcap_{t' < t} \mathcal{G}^{t'}(X).
$$

3. Proofs of Theorem [1.1](#page-1-1) and Corollary [1.2](#page-2-1)

In 2019, Persson [\[22\]](#page-18-6) used potentials and energies of measures to prove that some sets have large intersection property, which is a new and useful method in the study of large intersection property. Later, Persson [\[23\]](#page-18-7) applied this idea to show that the limsup sets, generated by sequences of open sets, belong to $\mathcal{G}^{\gamma}(\mathbb{T}^d)$ for some $\gamma > 0$. In this paper, we borrowed some ideas from [\[23\]](#page-18-7) to prove Theorem [1.1.](#page-1-1) Before that, we give some related lemmas.

In this section, we always assume that $(X, \mathcal{B}, \mu, \rho)$ is a probability space where the metric space (X, ρ) is compact and μ is an Ahlfors s-regular Borel measure $(0 < s < \infty)$. Let $\mathcal{Q} = \bigcup_{k \geq 0} \mathcal{Q}_k$ be the collection of generalized dyadic cubes in (X, ρ) given in Section 2.1.

Lemma 3.1. For $0 \leq t < s$ and $U \in \mathcal{B}$ with diam $U > 0$, we have

$$
(\text{diam}\, U)^{-t}\mu(U)^2 \le I_t(\mu, U) \le C_1(\text{diam}\, U)^{s-t}\mu(U),\tag{3.1}
$$

where $C_1 > 0$ is an absolute constant.

Proof. Write $l = \text{diam } U$, $l_j = 2^{-j}l$. For $t < s$ and $y \in U$, we have

$$
\phi_t(\mu, U, y) = \int_U \rho(x, y)^{-t} d\mu(x) \le \int_{\overline{B}(y, l)} \rho(x, y)^{-t} d\mu(x) = \sum_{j \ge 0} \int_{\overline{B}(y, l_j) \setminus \overline{B}(y, l_{j+1})} \rho(x, y)^{-t} d\mu(x)
$$

\n
$$
\le \sum_{j \ge 0} l_{j+1}^{-t} \left(\overline{B}(y, l_j) \setminus \overline{B}(y, l_{j+1}) \right) = \sum_{j \ge 0} l_{j+1}^{-t} \left(\mu(\overline{B}(y, l_j)) - \mu(\overline{B}(y, l_{j+1})) \right)
$$

\n
$$
\le \sum_{j \ge 0} l_{j+1}^{-t} \left(Cl_j^s - C^{-1}l_{j+1}^s \right) = l^{s-t} \sum_{j \ge 0} 2^{(1+j)(t-s)} (C2^s - C^{-1})
$$

\n
$$
= C_1 (\text{diam } U)^{s-t},
$$

where $C_1 > 0$ is a constant. Then $I_t(\mu, U) = \int_U \phi_t(\mu, U, y) d\mu(y) \leq C_1(\text{diam } U)^{s-t} \mu(U)$. Also

$$
I_t(\mu, U) = \int_U \int_U \rho(x, y)^{-t} d\mu(x) d\mu(y) \ge \int_U \int_U (\text{diam } U)^{-t} d\mu(x) d\mu(y) = (\text{diam } U)^{-t} \mu(U)^2.
$$

Remark 3.1. For $0 \le t < s$, since (X, ρ) is compact, we have $I_t(\mu) \le C_1 \max\{1, (\text{diam } X)^s\}$ $< \infty$.

Lemma 3.2. Let $\{\mu_n\}_{n>1}$ be a sequence of Borel measures on X which are absolutely continuous with respect to μ . Suppose there exists a constant $M > 0$ such that for any ball $B\subset X$,

$$
M^{-1} \le \liminf_{n \to \infty} \frac{\mu_n(B)}{\mu(B)} \le \limsup_{n \to \infty} \frac{\mu_n(B)}{\mu(B)} \le M,
$$
\n(3.2)

and $\sup_{n\geq 1} I_{\gamma}(\mu_n) < M$ for some $\gamma > 0$. Then for $t < \gamma$, there exists a constant $M_1 > 0$ such that

$$
M_1^{-1} \le \liminf_{n \to \infty} \frac{I_t(\mu_n, Q)}{I_t(\mu, Q)} \le \limsup_{n \to \infty} \frac{I_t(\mu_n, Q)}{I_t(\mu, Q)} \le M_1
$$

holds for all $Q \in \mathcal{Q}$.

Remark 3.2. In Lemma [3.2,](#page-5-0) the condition $\sup_{n\geq 1} I_\gamma(\mu_n) < M$ can be weaken to $\sup_{n\geq N_0} I_\gamma(\mu_n)$ $< M$ for some $N_0 \in \mathbb{N}$.

Proof. Fix any $Q \in \mathcal{Q}$. For any $\alpha > 0$, given $y \in Q$, let $Q_{\alpha}(y) = \{x \in Q : \rho(x, y) < \alpha^{-1/\gamma}\}.$ Then

$$
\phi_t(\mu_n, Q, y) = \int_{Q_{\alpha}(y)} \rho(x, y)^{-t} d\mu_n(x) + \int_{Q \setminus Q_{\alpha}(y)} \rho(x, y)^{-t} d\mu_n(x). \tag{3.3}
$$

Since $I_{\gamma}(\mu_n) < M$, we have

$$
M > \iint \rho(x,y)^{-\gamma} d\mu_n(x) d\mu_n(y) \ge \int_Q \Big(\int_{Q_{\alpha}(y)} \rho(x,y)^{-\gamma} d\mu_n(x) \Big) d\mu_n(y) > \alpha \int_Q \mu_n(Q_{\alpha}(y)) d\mu_n(y),
$$

and it follows that

$$
\int_{Q} d\mu_n(y) \int_{Q_{\alpha}(y)} \rho(x, y)^{-t} d\mu_n(x) = \int_{Q} d\mu_n(y) \int_{Q_{\alpha}(y)} d\mu_n(x) \int_0^{\rho(x, y)^{-t}} du
$$
\n
$$
= \int_{Q} d\mu_n(y) \int_{Q_{\alpha}(y)} d\mu_n(x) \int_0^{\alpha^{t/\gamma}} du + \int_{Q} d\mu_n(y) \int_{Q_{\alpha}(y)} d\mu_n(x) \int_{\alpha^{t/\gamma}}^{\rho(x, y)^{-t}} du
$$
\n
$$
= \alpha^{t/\gamma} \int_{Q} \mu_n(Q_{\alpha}(y)) d\mu_n(y) + \int_{\alpha^{t/\gamma}}^{\infty} du \int_{Q} d\mu_n(y) \int_{Q_{\alpha^{t/\gamma}}} d\mu_n(x)
$$
\n
$$
\leq M \alpha^{t/\gamma - 1} + M \int_{\alpha^{t/\gamma}}^{\infty} u^{-\gamma/t} du
$$
\n
$$
= M \alpha^{t/\gamma - 1} + \frac{Mt}{(\gamma - t)} \alpha^{t/\gamma - 1} = \frac{M\gamma}{(\gamma - t)} \alpha^{t/\gamma - 1},
$$
\n(3.4)

where we apply Fubini's theorem to derive the third equality. For $\epsilon > 0$, for α large enough, $\int_Q d\mu_n(y) \int_{Q_\alpha(y)} \rho(x, y)^{-t} d\mu_n(x) < \epsilon$. Observe that

$$
\int_{Q} d\mu_n(y) \int_{Q \setminus Q_{\alpha}(y)} \rho(x,y)^{-t} d\mu_n(x) \leq \int_{Q} \int_{Q} \min\{\rho(x,y)^{-t}, \alpha^{t/\gamma}\} d\mu_n(x) d\mu_n(y).
$$

Denote $f(x,y) = \min\{\rho(x,y)^{-t}, \alpha^{t/\gamma}\}\.$ Note that $f(x,y)$ is continuous on $Q \times Q$. Let ${Q_{k,i} \in \mathcal{Q}_k : 1 \leq i \leq N(k)}$ be a partition of Q by Borel sets satisfying diam $Q_{k,i} \to 0$ as $k \to \infty$, and take $(x_i, y_j) \in Q_{k,i} \times Q_{k,j}$, then for all $(x, y) \in Q \times Q$ we have

$$
\sum_{i,j}^{N(k)} f(x_i, y_j) \chi_{\{Q_{k,i} \times Q_{k,j}\}}(x, y) \to f(x, y), \text{ as } k \to \infty.
$$

Since $f(x, y) \leq \alpha^{t/\gamma}$, using Dominated Convergence Theorem, we obtain

$$
\int_{Q} \int_{Q} f(x, y) d\mu_n(x) d\mu_n(y) = \lim_{k \to \infty} \int_{Q} \int_{Q} \sum_{i,j}^{N(k)} f(x_i, y_j) \chi_{\{Q_{k,i} \times Q_{k,j}\}}(x, y) d\mu_n(x) d\mu_n(y)
$$

$$
= \lim_{k \to \infty} \sum_{i,j}^{N(k)} f(x_i, y_j) \mu_n(Q_{k,i}) \mu_n(Q_{k,j})
$$

$$
\leq \lim_{k \to \infty} \sum_{i,j}^{N(k)} f(x_i, y_j) \mu_n(\overline{B}_{k,i}) \mu_n(\overline{B}_{k,j}),
$$

where $\overline{B}_{k,i} = \overline{B}(x_{k,i}, c'_1 b^k)$. Recall that $B(x_{k,i}, c_1 b^k) \subset Q_{k,i} \subset \overline{B}_{k,i}$, there exists a constant $C_2 > 0$ such that $\mu(B_{k,i}) \leq C_2 \mu(Q_{k,i})$. We derive from [\(3.2\)](#page-5-1) that

$$
\limsup_{n \to \infty} \frac{\mu_n(\overline{B})}{\mu(\overline{B})} \le C^2 M.
$$

Then for any $\theta > 0$, for *n* large enough, we have

$$
\lim_{k \to \infty} \sum_{i,j}^{N(k)} f(x_i, y_j) \mu_n(\overline{B}_{k,i}) \mu_n(\overline{B}_{k,j}) \le (C^2 M + \theta)^2 \lim_{k \to \infty} \sum_{i,j}^{N(k)} f(x_i, y_j) \mu(\overline{B}_{k,i}) \mu(\overline{B}_{k,j})
$$
\n
$$
\le (C^2 M + \theta)^2 C_2^2 \lim_{k \to \infty} \sum_{i,j}^{N(k)} f(x_i, y_j) \mu(Q_{k,i}) \mu(Q_{k,j})
$$
\n
$$
= (C^2 M + \theta)^2 C_2^2 \int_Q \int_Q f(x, y) d\mu(x) d\mu(y)
$$
\n
$$
\le (C^2 M + \theta)^2 C_2^2 \int_Q \int_Q \int_Q \rho(x, y)^{-t} d\mu(x) d\mu(y).
$$

Since $\epsilon > 0$ is arbitrary, we prove that

$$
\limsup_{n \to \infty} \frac{I_t(\mu_n, Q)}{I_t(\mu, Q)} \le C^4 M^2 C_2^2
$$

.

Notice that $I_t(\mu_n, Q) \geq (\text{diam } Q)^{-t} \mu_n(Q)^2$ and $\mu_n(Q) \geq C^{-1} 2^{-s} (c_1/c_1')^s (\text{diam } Q)^s$, then combining these with Lemma [3.1,](#page-5-2) there exists an absolute constant $C_3 > 0$ such that

$$
\frac{I_t(\mu_n, Q)}{I_t(\mu, Q)} \ge C_3.
$$

Let $M_1 = \max\{C_3^{-1}, C^4M^2C_2^2\}$, then we proved this lemma.

Lemma 3.3. Under the same conditions as Lemma [3.2,](#page-5-0) let $\{A_n\}_{n\geq 1}$ be a sequence of open sets in X satisfying $\mu_n(X \setminus A_n) = 0, n \ge 1$. Then $\limsup_{n \to \infty} A_n \in \mathcal{G}^{\gamma}(X)$.

Proof. Let $Q \in \mathcal{Q}$ and $t < \gamma$. By [\(3.2\)](#page-5-1), $\mu_n(Q) > 0$ for n large enough. Define

$$
\nu_n(F) = \int_{F \cap Q} \phi_t^{-1}(\mu_n, Q, y) d\mu_n(y)
$$

for any $F \in \mathscr{B}$. Since $\mu_n(X \setminus A_n) = 0$, $\nu_n(Q) = \nu_n(Q \cap A_n)$. Given $\epsilon > 0$, combining Jensen's inequality $[20]$ and Lemma [3.2,](#page-5-0) for n large enough, we have

$$
\nu_n(F) = \mu_n(F \cap Q) \int_{F \cap Q} \Big(\int_Q \rho(x, y)^{-t} d\mu_n(x) \Big)^{-1} \frac{1}{\mu_n(F \cap Q)} d\mu_n(y)
$$

\n
$$
\geq \mu_n(F \cap Q) \Big(\int_{F \cap Q} \Big(\int_Q \rho(x, y)^{-t} d\mu_n(x) \Big) \frac{1}{\mu_n(F \cap Q)} d\mu_n(y) \Big)^{-1}
$$

\n
$$
\geq (\mu_n(F \cap Q))^2 \Big(\int_Q \int_Q \rho(x, y)^{-t} d\mu_n(y) d\mu_n(x) \Big)^{-1}
$$

\n
$$
> (\mu_n(F \cap Q))^2 (M_1 + \epsilon)^{-1} \Big(I_t(\mu, Q) \Big)^{-1} .
$$
\n(3.5)

Hence ν_n is a nonzero measure and absolutely continuous with respect to μ_n . By Lemma [3.1](#page-5-2) and (3.5) , for *n* large enough, we obtain

$$
\nu_n(Q) \ge (M_1 + \epsilon)^{-1} \frac{(\mu_n(Q))^2}{I_t(\mu, Q)} \ge C_1^{-1} (M_1 + \epsilon)^{-1} \frac{(\mu_n(Q))^2}{(\text{diam } Q)^{s-t} \mu(Q)} = c_3 \mu(Q)^{t/s}, \quad (3.6)
$$

where $c_3 > 0$ independ of Q and n. By Jensen's inequality,

$$
\phi_t^{-1}(\mu_n, Q, y) = \frac{1}{\mu_n(Q)} \Big(\int_Q \frac{\rho(x, y)^{-t}}{\mu_n(Q)} d\mu_n(x) \Big)^{-1} \leq \frac{1}{(\mu_n(Q))^2} \int_Q \rho(x, y)^t d\mu_n(x).
$$

Then for any $J \in \mathcal{Q}$ and $J \subset Q$, we get that

$$
\nu_n(J) = \int_J \phi_t^{-1}(\mu_n, Q, y) d\mu_n(y) \le \int_J \phi_t^{-1}(\mu_n, J, y) d\mu_n(y)
$$

\n
$$
\le \frac{1}{(\mu_n(J))^2} \int_J \int_J \rho(x, y)^t d\mu_n(x) d\mu_n(y)
$$

\n
$$
\le (\text{diam } J)^t \le c_4 \mu(J)^{t/s}
$$
\n(3.7)

for *n* large enough and $c_4 > 0$ is an absolute constant.

Let $\{Q_k\}_{k\geq 1}$ be a cover of $Q \cap A_n$, where $Q_k \in \mathcal{Q}$. Without loss of generality, we assume $Q_k \cap Q_j = \emptyset$ for $k \neq j$ and $Q_k \subset Q$. For n large enough, we derive from [\(3.6\)](#page-7-1) and [\(3.7\)](#page-8-0) that

$$
\sum_{k} \mu(Q_k)^{t/s} \ge c_4^{-1} \sum_{k} \nu_n(Q_k) \ge c_4^{-1} \nu_n(Q \cap A_n) = c_4^{-1} \nu_n(Q) \ge c_3 c_4^{-1} \mu(Q)^{t/s}.
$$

It follows that

$$
\liminf_{n \to \infty} \mathcal{M}_{\infty}^t(Q \cap A_n) \ge c_3 c_4^{-1} \mu(Q)^{t/s}
$$

for n large enough. Therefore, for any $m \geq 1$ and generalized dyadic cube Q,

$$
\mathcal{M}_{\infty}^{t}\Big(\bigcup_{n\geq m}A_{n}\cap Q\Big)\geq \sup_{n\geq m}\mathcal{M}_{\infty}^{t}(A_{n}\cap Q)\geq c_{3}c_{4}^{-1}\mu(Q)^{t/s}.\quad (\forall t<\gamma)
$$

So $\bigcup_{n\geq m} A_n \in \mathcal{G}^{\gamma}(X)$ by [\(2.3\)](#page-4-2). Thus from (3) in Proposition [2.2,](#page-4-3) we prove that

$$
\limsup_{n \to \infty} A_n = \bigcap_{m \ge 1} \bigcup_{n \ge m} A_n \in \mathcal{G}^{\gamma}(X).
$$

Recall the 5r-covering theorem in metric space.

Lemma 3.4 ([\[19\]](#page-18-17)). Let (X, ρ) be a separable metric space and A be a family either of closed balls or open balls such that

 $\sup{\{\operatorname{diam} B\colon B\in\mathcal{A}\}}<\infty$.

Then there is a finite or countable sequence ${B_i}_{i \in I}$ of pairwise disjoint balls such that

$$
\bigcup_{B \in \mathcal{A}} B \subset \bigcup_{i \in I} 5B_i.
$$

Now we prove Theorem [1.1.](#page-1-1)

Proof of Theorem [1.1](#page-1-1): Since $\mu\left(\, \limsup \right.$ $\limsup_{n\to\infty} B_n$ = 1, for $n \ge 1$, there is some N_n such that

$$
\mu\left(\bigcup_{m=n}^{N_n} B_m\right) > 1 - \frac{1}{2n}.
$$

Denote $A_n = \bigcup_{m=n}^{N_n} B_m$. By Lemma [3.4,](#page-8-1) there is $\mathcal{I}_n \subset \{n, n+1, \ldots N_n\}$ such that

$$
B_i \cap B_j = \emptyset
$$
, for $i \neq j \in \mathcal{I}_n$ and $A_n \subset \bigcup_{m \in \mathcal{I}_n} 5B_m$.

(i) Construct measures

Let $A'_n = \bigcup_{m \in \mathcal{I}_n} B_m$. Since

$$
\mu(A'_n) = \sum_{m \in \mathcal{I}_n} \mu(B_m) \ge C^{-2} 5^{-s} \sum_{m \in \mathcal{I}_n} \mu(5B_m) > C^{-2} 5^{-s} (1 - \frac{1}{2n}) > 0,
$$

and $\mu(A'_n) < \infty$, we define

$$
\eta_n(F) = \frac{1}{\mu(A'_n)} \mu(F \cap A'_n)
$$

for any $F \in \mathcal{B}$. Then η_n is a nonzero probability measure supported on A'_n satisfying $\eta_n(A'_n) = \eta_n(X) = 1.$

Let $F_n = \bigcup_{m \in \mathcal{I}_n} E_m$. Define

$$
\mu_n(F) = \sum_{m \in \mathcal{I}_n} \eta_n(B_m) \frac{\mu(E_m \cap F)}{\mu(E_m)}
$$

for $F \in \mathscr{B}$. Then μ_n is a nonzero probability measure supported on F_n satisfying $\mu_n(F_n) =$ $\mu_n(X) = 1.$

Let $B \subset X$ be a ball. Firstly we will show the following inequalities

$$
C^{-4}5^{-s} \le \liminf_{n \to \infty} \frac{\eta_n(B)}{\mu(B)} \le \limsup_{n \to \infty} \frac{\eta_n(B)}{\mu(B)} \le C^4 5^s. \tag{3.8}
$$

Denote $B = B(x_B, r_B)$. Since r_n decreases to 0 as $n \to \infty$, for any $\theta > 0$, there is some $N > 0$ such that for $n \geq N$, $m \in \mathcal{I}_n$, we have $r_m < r_B \theta/2$. Note that

$$
\bigcup_{m \in \mathcal{I}_n \atop B \cap B_m \neq \varnothing} B_m \subset (1 + \theta)B,
$$

hence

$$
\eta_n(B) = \frac{1}{\mu(A'_n)} \mu(B \cap A'_n) \le \frac{1}{\mu(A'_n)} \mu\left(\bigcup_{m \in \mathcal{I}_n \atop B \cap B_m \ne \varnothing} B_m\right) \le \frac{\mu((1+\theta)B)}{\mu(A'_n)} < \frac{C^2(1+\theta)^s \mu(B)}{C^{-2}5^{-s}(1-1/2n)},
$$

which implies

$$
\limsup_{n \to \infty} \frac{\eta_n(B)}{\mu(B)} \le C^4 5^s.
$$

Let $\mathcal{I}_n(B) = \{m \in \mathcal{I}_n : B_m \subset B\}$, then we have

$$
\frac{\eta_n(B)}{\mu(B)} \ge \frac{1}{\mu(A'_n)\mu(B)}\mu\Big(\bigcup_{m \in \mathcal{I}_n(B)} B_m \cap A'_n\Big) \ge \frac{1}{\mu(B)} \sum_{m \in \mathcal{I}_n(B)} \mu(B_m).
$$

We claim that

$$
\liminf_{n \to \infty} \sum_{m \in \mathcal{I}_n(B)} \mu(B_m) \ge C^{-4} 5^{-s} \mu(B). \tag{3.9}
$$

Then

$$
\liminf_{n \to \infty} \frac{\eta_n(B)}{\mu(B)} \ge \liminf_{n \to \infty} \frac{1}{\mu(B)} \sum_{m \in \mathcal{I}_n(B)} \mu(B_m) \ge C^{-4} 5^{-s}.
$$

Now we prove the claim. For any $0 < \epsilon < 1/2$, take *n* large enough such that $\mu(B)$ $C^2 2^{s-1}/n$, and $r_m < \epsilon r_B/10$, $m \in \mathcal{I}_n$. Then if $m \in \mathcal{I}_n \setminus \mathcal{I}_n(B)$, that is, $B^c \cap B_m \neq \emptyset$, we have $5B_m \cap (1 - \epsilon)B = \emptyset$. Therefore

$$
\mu\Big((1-\epsilon)B\cap A_n\Big) \leq \mu\Big(\bigcup_{m\in\mathcal{I}_n(B)} 5B_m\Big) + \mu\Big(\bigcup_{m\in\mathcal{I}_n\setminus\mathcal{I}_n(B)} 5B_m\cap (1-\epsilon)B\Big)
$$

=
$$
\mu\Big(\bigcup_{m\in\mathcal{I}_n(B)} 5B_m\Big) \leq \sum_{m\in\mathcal{I}_n(B)} \mu(5B_m) \leq C^2 5^s \mu\Big(\bigcup_{m\in\mathcal{I}_n(B)} B_m\Big).
$$

We also note that

$$
\mu\Big((1-\epsilon)B\cap A_n\Big)\geq\mu\Big((1-\epsilon)B\Big)-\frac{1}{2n}\geq C^{-2}(1-\epsilon)^s\mu(B)-\frac{1}{2n}.
$$

Combining the inequalities above, we prove the claim.

Secondly we will prove that

$$
C^{-8}5^{-2s} \le \liminf_{n \to \infty} \frac{\mu_n(B)}{\mu(B)} \le \limsup_{n \to \infty} \frac{\mu_n(B)}{\mu(B)} \le C^6 5^s
$$

hold for any ball B. Given any $\theta > 0$, Since $\bigcup_{B_m \cap B \neq \varnothing} B_m \subset (1 + \theta)B$ holds for n large enough, we obtain

$$
\mu_n(B) = \sum_{m \in \mathcal{I}_n} \eta_n(B_m) \frac{\mu(E_m \cap B)}{\mu(E_m)} = \sum_{\substack{m \in \mathcal{I}_n \\ B_m \cap B \neq \varnothing}} \eta_n(B_m) \frac{\mu(E_m \cap B)}{\mu(E_m)}
$$

$$
\leq \sum_{\substack{m \in \mathcal{I}_n \\ B_m \cap B \neq \varnothing}} \eta_n(B_m) \leq \eta_n((1+\theta)B) \leq (C^4 5^s + \theta)(1+\theta)^s C^2 \mu(B).
$$

The last inequality follows from [\(3.8\)](#page-9-0). So lim sup n→∞ $\frac{\mu_n(B)}{\mu(B)} \leq C^6 5^s.$

We derive from [\(3.8\)](#page-9-0) and [\(3.9\)](#page-9-1) that

$$
\liminf_{n \to \infty} \frac{\mu_n(B)}{\mu(B)} = \liminf_{n \to \infty} \frac{1}{\mu(B)} \sum_{m \in \mathcal{I}_n} \eta_n(B_m) \frac{\mu(E_m \cap B)}{\mu(E_m)}
$$
\n
$$
\geq \liminf_{n \to \infty} \frac{1}{\mu(B)} \sum_{m \in \mathcal{I}_n(B)} \eta_n(B_m)
$$
\n
$$
\geq \liminf_{n \to \infty} \frac{C^{-4} 5^{-s}}{\mu(B)} \sum_{m \in \mathcal{I}_n(B)} \mu(B_m)
$$
\n
$$
\geq C^{-8} 5^{-2s}.
$$

(ii) Show that $\sup_{n\geq N} I_t(\mu_n) < \infty$ for some $N \geq 1$ Note that

$$
I_t(\mu_n) = \int_X \int_X \rho(x, y)^{-t} d\mu_n(x) d\mu_n(y) = \int_{F_n} \int_{F_n} \rho(x, y)^{-t} d\mu_n(x) d\mu_n(y)
$$

=
$$
\sum_{m \in \mathcal{I}_n} \sum_{i \in \mathcal{I}_n} \int_{E_m} \int_{E_i} \rho(x, y)^{-t} d\mu_n(x) d\mu_n(y).
$$

When $i = m$, since $\mu_n|_{E_m} = \frac{\eta_n(B_m)}{\mu(E_m)}$ $\frac{\partial n(B_m)}{\mu(E_m)} \mu|_{E_m}$, we have

$$
\int_{E_m} \int_{E_m} \rho(x, y)^{-t} d\mu_n(x) d\mu_n(y) = \left(\frac{\eta_n(B_m)}{\mu(E_m)}\right)^2 \int_{E_m} \int_{E_m} \rho(x, y)^{-t} d\mu(x) d\mu(y)
$$

= $\left(\frac{\eta_n(B_m)}{\mu(E_m)}\right)^2 I_t(\mu, E_m) \le (C^4 5^s + \theta)^2 \left(\frac{\mu(B_m)}{\mu(E_m)}\right)^2 I_t(\mu, E_m)$

for *n* large enough. Due to $t < \lambda$, there is some absolute constant $M_3 > 0$ such that

$$
\sup_{n\geq 1} \frac{I_t(\mu, E_n)\mu(B_n)}{\mu(E_n)^2} < M_3.
$$

Hence

$$
\int_{E_m} \int_{E_m} \rho(x, y)^{-t} d\mu_n(x) d\mu_n(y) \le (C^4 5^s + \theta)^2 \Big(\frac{\mu(B_m)}{\mu(E_m)}\Big)^2 I_t(\mu, E_m) \le (C^4 5^s + \theta)^2 M_3 \mu(B_m).
$$
\n(3.10)

For $i \neq m$, we suppose that there is a ball cB_m (0 < c < 1) such that $E_m \subset cB_m$ for any $m \geq 1$. Otherwise, we consider $c^{-1}B_m$ instead of B_m in the previous proof. For $x \in E_m$, $y \in E_i$, we get

$$
\rho(x, y) \ge \rho(x_i, x_m) - c(r_m + r_i) \ge (1 - c)\rho(x_i, x_m).
$$

Then

$$
\int_{E_m} \int_{E_i} \rho(x, y)^{-t} d\mu_n(x) d\mu_n(y) \leq \int_{E_m} \int_{E_i} ((1 - c)\rho(x_i, x_m))^{-t} d\mu_n(x) d\mu_n(y)
$$

= $((1 - c)\rho(x_i, x_m))^{-t} \mu_n(E_m) \mu_n(E_i) = ((1 - c)\rho(x_i, x_m))^{-t} \eta_n(B_m) \eta_n(B_i)$
 $\leq (1 - c)^{-t} (C^4 5^s + \theta)^2 \rho(x_i, x_m)^{-t} \mu(B_m) \mu(B_i)$
 $\leq (1 - c)^{-t} (C^4 5^s + \theta)^2 2^t \int_{B_m} \int_{B_i} \rho(x, y)^{-t} d\mu(x) d\mu(y).$

The last inequality follows from $\rho(x, y) \leq \rho(x_i, x_m) + r_i + r_m \leq 2\rho(x_i, x_m)$ for $x \in B_m$, $y \in$ B_i .

Therefore by Lemma [3.1](#page-5-2) and the fact that ${B_m}_{m \in I_n}$ are disjoint balls, for *n* large enough, we obtain

$$
I_t(\mu_n) \le (C^4 5^s + \theta)^2 \sum_{m \in \mathcal{I}_n} \left(\frac{\mu(B_m)}{\mu(E_m)}\right)^2 I_t(\mu, E_m)
$$

+
$$
\frac{(C^4 5^s + \theta)^2 2^t}{(1 - c)^t} \sum_{m \in \mathcal{I}_n} \sum_{\substack{i \in \mathcal{I}_n \\ i \neq m}} \int_{B_m} \int_{B_i} \rho(x, y)^{-t} d\mu(x) d\mu(y)
$$

$$
\le (2C^4 5^s)^2 M_3 \sum_{m \in \mathcal{I}_n} \mu(B_m) + (1 - c)^{-t} (2C^4 5^s)^2 2^t I_t(\mu)
$$

$$
\le (2C^4 5^s)^2 M_3 + (1 - c)^{-t} (2C^4 5^s)^2 2^t C_1 \max\{1, (\text{diam } X)^s\} < \infty.
$$

By Lemma [3.3,](#page-7-2) $\limsup F_n \in \mathcal{G}^t(X)$ for all $t < \lambda$, hence \limsup n→∞ $\limsup_{n\to\infty} F_n \in \mathcal{G}^{\lambda}(X)$ which implies that $\limsup E_n \in \mathcal{G}^{\lambda}$ $(X).$

Proof of Corollary [1.2](#page-2-1): Without loss of generality, we assume that diam $B_n < 1$. Given $t \in (0, s],$ write $F_n = B_n^t$, then $\mu\left(\limsup$ $\limsup_{n\to\infty} F_n$ = 1. For $\beta \geq 0$, by Lemma [3.1,](#page-5-2) we have

$$
C^{-1}2^{-\beta}r_n^{t-\beta} \le \frac{I_\beta(\mu, B_n)\mu(F_n)}{\mu(B_n)^2} \le C^2C_12^{s-\beta}r_n^{t-\beta}.
$$

Notice that if $\beta \leq t$, we have $\sup_{n\geq 1} \frac{I_{\beta}(\mu, B_n)\mu(F_n)}{\mu(B_n)^2}$ $\frac{d_i B_n \mu(r_n)}{\mu(B_n)^2} < \infty$, and if $\beta > t$, then $\limsup_{n \to \infty}$ n→∞ $I_\beta(\mu,B_n)\mu(F_n)$ $\frac{\mu(B_n)\mu(r_n)}{\mu(B_n)^2} =$ ∞ , since r_n decreases to 0 as $n \to \infty$. Hence

$$
t = \sup \Big\{ \beta \ge 0 \colon \sup_{n \ge 1} \frac{I_{\beta}(\mu, B_n) \mu(F_n)}{\mu(B_n)^2} < \infty \Big\},\
$$

and we applying Theorem [1.1](#page-1-1) to have $\limsup B_n \in \mathcal{G}^t(X)$. $∞ √∞$

4. Applications

Let (X, ρ) be a compact metric space endowed with an Ahlfors s-regular probability measure μ . In this section we investigate the large intersection property of limsup random fractals, random covering sets, and limsup sets generated by rectangles.

4.1. Limsup random fractals.

Given $0 < b < 1/3$, there is a family of generalized dyadic cubes of X defined in Subsection 2.1. In this subsection, we consider a model of limsup random fractals which is constructed by using generalized dyadic cubes $Q = \bigcup_{n \geq 0} Q_n$.

For $n \geq 1$, let $\{Z_n(Q), Q \in \mathcal{Q}_n\}$ be a sequence of random variables, each taking values in {0, 1}. Let

$$
A(n) = \bigcup_{\substack{Q \in \mathcal{Q}_n, \\ Z_n(Q) = 1}} Q^o,
$$

where Q^o is the interior of Q . The random set

$$
A = \limsup_{n \to \infty} A(n)
$$

is called a limsup random fractal associated to $\{Z_n(Q), Q \in \mathcal{Q}_n, n \geq 1\}$. For $n \geq 1$, and $Q \in \mathcal{Q}_n$, denote the probability $P_n(Q) := \mathbb{P}(Z_n(Q) = 1)$, and

$$
\gamma_1: = -\limsup_{n \to \infty} \frac{\max_{Q \in \mathcal{Q}_n} \log_{b^{-1}} P_n(Q)}{n},\tag{4.1}
$$

$$
\gamma_2: = -\limsup_{n \to \infty} \frac{\min_{Q \in \mathcal{Q}_n} \log_{b^{-1}} P_n(Q)}{n}.
$$
\n(4.2)

We refer to γ_1 and γ_2 as the indices of the limsup random fratcal A.

We assume the following condition, which is similar as the condition $(H2)$ in [\[15\]](#page-18-18).

Correlation Condition: Suppose that there is a constant $\delta \geq 0$ such that for all $\epsilon > 0$,

$$
\limsup_{n \to \infty} \frac{1}{n} \log_{b^{-1}} f(n, \epsilon) \le \delta,
$$
\n(4.3)

where

$$
f(n,\epsilon) = \max_{Q \in \mathcal{Q}_n} \# \{ Q' \in \mathcal{Q}_n : \text{Cov}(Z_n(Q), Z_n(Q')) \ge \epsilon P_n(Q) P_n(Q') \}.
$$

Theorem 4.1. Let $(X, \mathcal{B}, \mu, \rho)$ be a compact Ahlfors s-regular space $(0 \lt s \lt \infty)$. Let $A = \limsup A(n)$ be a limsup random set with indices γ_1 , γ_2 and satisfy the Correlation $n{\rightarrow}\infty$ Condition. If $\gamma_2 + \delta < s$, then $A \in \mathcal{G}^{s - \gamma_2 - \delta}(X)$ a.s.

Corollary 4.1. Under the setting in Theorem [4.1,](#page-12-0) we have

 $\max\{0, s - \gamma_2 - \delta\} \le \dim_{\mathrm{H}} A \le \max\{0, s - \gamma_1\}$ a.s.

Remark 4.1. We can also use the hitting probability of limsup random fractals to get the estimation on Hausdorff dimension in Corollary [4.1.](#page-12-1) For instance, let A be a limsup random fratcal in T with $\gamma_1 = \gamma_2 < 1$, $\delta = 0$, and if $P_n(Q)$ is independent of Q, Khoshnevisan, Peres and Xiao [\[17\]](#page-18-19) showed that $\dim_{\text{H}} A = 1 - \gamma_1$ a.s. For more related research, see [\[13,](#page-18-20)[15\]](#page-18-18).

Before proving Theorem [4.1,](#page-12-0) we give some lemmas. For $n \geq 1$ and $Q \in \mathcal{Q}_n$, recalling (2.1) , there is some $x_Q \in Q$ such that

$$
B(x_Q, c_1 b^n) \subset Q \subset \overline{B}(x_Q, c'_1 b^n),
$$

and here $c_1 = \frac{1}{2} - \frac{b}{1 - b}$ $\frac{b}{1-b}$ and $c'_1 = \frac{1}{1-b}$ $\frac{1}{1-b}$.

Lemma 4.1. Let C and c'_1 be the constants in [\(1.1\)](#page-1-0) and [\(2.1\)](#page-3-0) respectively. Let $B =$ $B(x_B, r)$ be a ball with $r > 0$ and

$$
\mathcal{Q}_n(B) = \{Q \in \mathcal{Q}_n \colon Q \subset B\}.
$$

Then there exists some $N \geq 1$ such that for all $n \geq N$, we have

$$
\#\mathcal{Q}_n(B) \ge C^{-2} (r c_1'^{-1} b^{-n} - 2)^s.
$$

Proof. For $Q \in \mathcal{Q}_n(B)$, there exists $x_Q \in Q$ such that

$$
Q \subset \overline{B}(x_Q, c'_1 b^n).
$$

For $r > 0$, there exists some $N \ge 1$ such that for all $n \ge N$, we have $r > 2c'_1b^n$. Then for $Q \in \mathcal{Q}_n$, observe that if $Q \notin \mathcal{Q}_n(B)$, then $Q \cap B(x_B, r - 2c'_1b^n) = \varnothing$, and we deduce that

$$
B(x_B, r - 2c'_1 b^n) \subset \bigcup_{Q \in \mathcal{Q}_n(B)} Q \cup \bigcup_{Q \notin \mathcal{Q}_n(B) \atop Q \cap B \neq \varnothing} Q = \bigcup_{Q \in \mathcal{Q}_n(B)} Q.
$$

Hence

$$
C^{-1}(r - 2c'_1b^n)^s \le \mu\big(B(x_B, r - 2c'_1b^n)\big) \le \mu\bigg(\bigcup_{Q \in \mathcal{Q}_n(B)} Q\big) \le \mu\bigg(\bigcup_{Q \in \mathcal{Q}_n(B)} \overline{B}(x_Q, c'_1b^n)\big)
$$

$$
\le \#\mathcal{Q}_n(B)C(c'_1b^n)^s,
$$

which derives that

$$
\#\mathcal{Q}_n(B) \ge \frac{(r - 2c'_1 b^n)^s}{C^2 (c'_1 b^n)^s} \quad (\forall n \ge N).
$$

Recall the notation $B^t(x,r) = B(x, r^{t/s})$.

Lemma 4.2. Suppose that $\gamma_2 + \delta < s$. For $t > 0$ with $t < s - \gamma_2 - \delta$, let

$$
B^{t}(n) = \bigcup_{Q \in \mathcal{Q}_n \atop Z_n(Q)=1} B^t_Q,
$$

where $B_Q = B(x_Q, c_1 b^n) \subset Q$. Given $x \in X$, let $J_{n,x}$ be the unique dyadic cube in \mathcal{Q}_n containing x. Then for $k \geq 1$, we have

$$
\mathbb{P}\Big(\bigcup_{n\geq k} \{J_{n,x} \subset B^t(n)\}\Big) = 1.
$$

Proof. Given $x \in X$, there is a $x' \in J_{n,x}$ such that

$$
B(x', c_1 b^n) \subset J_{n,x}.
$$

Let

$$
\mathcal{B}_n(x) = \left\{ J \in \mathcal{Q}_n : J \subset B(x', (c_1 b^n)^{t/s}) \right\}.
$$

By Lemma [4.1,](#page-12-2) for n large enough, we have

$$
\#\mathcal{B}_n(x) \ge \frac{1}{C^2} \Big(\frac{c_1^{t/s}}{c_1'} b^{n(t/s-1)} - 2\Big)^s. \tag{4.4}
$$

Define $M_n(x) = \sum_{J \in \mathcal{B}_n(x)} Z_n(J)$, and note that

$$
\big\{J_{n,x} \subset B^t(n)\big\} \supset \big\{Z_n(J) = 1 \text{ for some } J \in \mathcal{B}_n(x)\big\} = \big\{M_n(x) > 0\big\}.
$$

From Chebyshev's inequality, we obtain

$$
\mathbb{P}(M_n(x) = 0) \le \frac{\text{Var}(M_n(x))}{(\mathbb{E}(M_n(x)))^2}.
$$

Since $Cov(Z_n(J), Z_n(J')) \leq \mathbb{E}(Z_n(J)Z_n(J')) \leq P_n(J)$, we have

$$
\operatorname{Var}(M_n(x)) = \sum_{J \in \mathcal{B}_n(x)} \sum_{J' \in \mathcal{B}_n(x)} \operatorname{Cov}(Z_n(J), Z_n(J'))
$$

\n
$$
= \sum_{J \in \mathcal{B}_n(x)} \Big(\sum_{J' \in \mathcal{G}_n(J)} \operatorname{Cov}(Z_n(J), Z_n(J')) + \sum_{J' \in \mathcal{T}_n(J)} \operatorname{Cov}(Z_n(J), Z_n(J')) \Big)
$$

\n
$$
\leq \sum_{J \in \mathcal{B}_n(x)} \Big(\sum_{J' \in \mathcal{G}_n(J)} \epsilon P_n(J) P_n(J') + \sum_{J' \in \mathcal{T}_n(J)} P_n(J) \Big)
$$

\n
$$
\leq \epsilon \Big(\sum_{J \in \mathcal{B}_n(x)} P_n(J) \Big) \Big(\sum_{J' \in \mathcal{G}_n(J)} P_n(J') \Big) + \max_{J \in \mathcal{B}_n(x)} \# \mathcal{T}_n(J) \Big(\sum_{J \in \mathcal{B}_n(x)} P_n(J) \Big),
$$
\n(4.5)

where

$$
\mathcal{G}_n(J) = \{ J' \in \mathcal{B}_n(x) \colon \text{Cov}(Z_n(J), Z_n(J')) < \epsilon P_n(J) P_n(J') \},
$$
\n
$$
\mathcal{T}_n(J) = \mathcal{B}_n(x) \setminus \mathcal{G}_n(J).
$$

Recalling the notation of the Correlation Condition, we obtain

$$
\text{Var}(M_n(x)) \le \epsilon \Big(\sum_{J \in \mathcal{B}_n(x)} P_n(J) \Big)^2 + f(n, \epsilon) \Big(\sum_{J \in \mathcal{B}_n(x)} P_n(J) \Big).
$$

Note that $\mathbb{E}(M_n(x)) = \sum_{J \in \mathcal{B}_n(x)} P_n(J) \geq \# \mathcal{B}_n(x) (\min_{Q \in \mathcal{Q}_n} P_n(Q)),$ we derive that $f(\mathbb{F}(M(n)))^2 + f(\mathbb{F}(M(n)))^2$

$$
\mathbb{P}(M_n(x) = 0) \le \frac{\epsilon(\mathbb{E}(M_n(x)))^2 + f(n, \epsilon)(\mathbb{E}(M_n(x)))}{(\mathbb{E}(M_n(x)))^2}
$$

$$
= \epsilon + \frac{f(n, \epsilon)}{\mathbb{E}(M_n(x))}
$$

$$
\le \epsilon + \frac{f(n, \epsilon)}{\#B_n(x)(\min_{Q \in \mathcal{Q}_n} P_n(Q))}.
$$
(4.6)

From [\(4.3\)](#page-12-3) and [\(4.4\)](#page-13-0), for $\theta > 0$ with $2\theta < s - \delta - \gamma_2 - t$, we get

$$
f(n, \epsilon) \le b^{-n(\delta + \theta)}
$$
 and $\#B_n(x) \ge c_5 b^{n(t-s)}$

for *n* large enough, where $c_5 > 0$ is an absolute constant, and from [\(4.2\)](#page-12-4), we have

$$
\min_{Q \in \mathcal{Q}_n} P_n(Q) \ge b^{n(\gamma_2 + \theta)}
$$

for infinitely many n, denoted by $\mathcal N$. Then

$$
\limsup_{\substack{n\in\mathcal{N} \\ n\to\infty}}\frac{f(n,\epsilon)}{\#\mathcal{B}_n(x)\min_{Q\in\mathcal{Q}_n}P_n(Q)} \le \limsup_{\substack{n\in\mathcal{N} \\ n\to\infty}}c_5^{-1}b^{-n(2\theta+\delta+\gamma_2-s+t)} = 0,
$$

which implies $\limsup \mathbb{P}(M_n(x) = 0) = 0$. Therefore n∈N n→∞

$$
\limsup_{n \to \infty} \mathbb{P}(M_n(x) > 0) = 1.
$$

So we show that

$$
\limsup_{n \to \infty} \mathbb{P}(J_{n,x} \subset B^t(n)) = 1.
$$

Hence for $k \geq 1$,

$$
\mathbb{P}\Big(\bigcup_{n=k}^{\infty} \{J_{n,x} \subset B^t(n)\}\Big) \ge \limsup_{n\to\infty} \mathbb{P}\big(J_{n,x} \subset B^t(n)\big) = 1.
$$

Proof of Theorem [4.1](#page-12-0): Let $x \in X$ and $t > 0$ with $t < s - \gamma_2 - \delta$. By Lemma [4.2,](#page-13-1) we have $\mathbb{P}\Big(\bigcup_{n=k}^{\infty} \{J_{n,x} \subset B^t(n)\}\Big) = 1$ for any $k \geq 1$. Note that

$$
\left\{x \in \bigcup_{n=k}^{\infty} B^t(n)\right\} \supset \left\{\exists n \geq k \text{ such that } J_{n,x} \subset B^t(n)\right\}.
$$

This yields that $\mathbb{P}\left\{x \in \bigcup_{n=k}^{\infty} B^t(n)\right\} = 1$. Write the set $F_k = \bigcup_{n=k}^{\infty} B^t(n)$, then

$$
\int d\mu(x) \int \chi_{\{x \in F_k\}} d\mathbb{P}(\omega) = 1.
$$

Using Fubini's Theorem, we obtain

$$
\int d\mathbb{P}(\omega) \int \chi_{F_k}(x) d\mu(x) = 1,
$$

implying that $\mu(F_k) = 1$ a.s. for any $k \geq 1$. Hence $\mu\left(\text{lim sup}\right)$ n→∞ $B^{t}(n)$ = 1 a.s.

By Corollary [1.2,](#page-2-1) for $t > 0$ with $t < s - \gamma_2 - \delta$, we have $\limsup B(n) \in \mathcal{G}^t(X)$ a.s., n→∞ which implies $\limsup B(n) \in \mathcal{G}^{s-\gamma_2-\delta}(X)$ a.s. Hence by Proposition [2.2,](#page-4-3) $\limsup A(n) \in$ n→∞ n→∞ $\mathcal{G}^{s-\gamma_2-\delta}$ (X) a.s.

Proof of Corollary [4.1](#page-12-1): If $s \geq \gamma_1$, let $t > s - \gamma_1$. For $\delta > 0$, there exists $k_0 \geq 1$ such that $2c'_1b^{k_0} \le \delta < 2c'_1b^{k_0-1}$. Since

$$
A\subset \bigcup_{n=k}^{\infty}\bigcup_{Q\in\mathcal{Q}_n\atop Z_n(Q)=1}Q^o
$$

for all $k > 1$, we have

$$
\mathcal{H}_{\delta}^t(A) \leq \sum_{n \geq k_0} \sum_{Q \in \mathcal{Q}_n} (\text{diam } Q)^t Z_n(Q).
$$

Then

$$
\mathbb{E}(\mathcal{H}_{\delta}^{t}(A)) \leq \sum_{n \geq k_0} \sum_{Q \in \mathcal{Q}_n} (\text{diam } Q)^t P_n(Q).
$$

Considering $\theta > 0$ with $\gamma_1 > \theta > s - t$, by [\(4.1\)](#page-12-5), for *n* large enough,

$$
\max_{Q \in \mathcal{Q}_n} P_n(Q) \le b^{n\theta}
$$

which implies that

$$
\mathbb{E}(\mathcal{H}_{\delta}^{t}(A)) \leq \sum_{n \geq k_0} \# \mathcal{Q}_n(2c'_1 b^n)^t b^{n\theta}
$$
\n
$$
\leq C c_1^{-s} (2c'_1)^t \sum_{n \geq k_0} b^{n(t-s+\theta)} < \infty,\tag{4.7}
$$

,

giving $\dim_{\text{H}} A \leq s - \gamma_1$ a.s. If $\gamma_1 > s$, taking $t > 0$ and $\theta = s$ in [\(4.7\)](#page-15-0), we have $\mathbb{E}(\mathcal{H}_{\delta}^t(A)) \leq$ $Cc_1^{-s}(2c'_1)^t\sum_{n\geq k_0}b^{nt}<\infty$ a.s., which implies $\dim_{\rm H} A\leq 0$ a.s.

When $s > \gamma_2 + \delta$, from Theorem [4.1,](#page-12-0) we have $A \in \mathcal{G}^{s-\gamma_2-\delta}(X)$ a.s. Then we obtain from Proposition [2.2](#page-4-3) (4) that $\dim_{\rm H} A \geq s - \gamma_2 - \delta$ a.s.

4.2. Random covering sets.

Durand [\[8\]](#page-18-5) showed that random covering sets in T, defined as limsup sets of a sequence of balls whose centers are independent and uniformly distributed random variables, have large intersection property. In this subsection, we consider random covering sets in $(X, \mathscr{B}, \mu, \rho)$ with weak dependence condition. Let $\{\xi_n\}_{n\geq 1}$ be a stationary process on a probability space $(\Omega, \mathcal{B}, \mathbb{P})$ having μ as probability law.

Definition 4.1. We say that $\{\xi_n\}_{n\geq 1}$ is exponentially mixing if for any $n \geq 1$, there exist two constants $c > 0$ and $0 < \gamma < 1$ such that

$$
\left| \mathbb{P}(\xi_1 \in A | D) - \mathbb{P}(\xi_1 \in A) \right| \le c\gamma^n
$$

holds for any ball $A \subset X$ and $D \in \mathcal{B}_{n+1}$, where \mathcal{B}_{n+1} is the sub- σ -field generated by $\{\xi_{n+i}\}_{i\geq 1}$.

Let ${r_n}_{n\geq1}$ be a sequence of positive real numbers decreasing to zero. The random covering set is defined as

$$
E: = \limsup_{n \to \infty} B(\xi_n, r_n) = \{ y \in X : y \in B(\xi_n, r_n) \text{ for infinitely many } n \ge 1 \}.
$$

The set E consists of the points which are covered by ${B(\xi_n, r_n)}_{n>1}$ infinitely often.

Theorem 4.2 ([\[14\]](#page-18-21)). Let $\{\xi_n\}_{n>1}$ be exponentially mixing and the probability law μ be Ahlfors s-regular. Then we have

$$
\mu(E) = \begin{cases} 0 & \text{if } \sum_{n=1}^{\infty} r_n^s < \infty \\ 1 & \text{if } \sum_{n=1}^{\infty} r_n^s = \infty \end{cases} \quad a.s.
$$

Theorem 4.3. Let $(X, \mathcal{B}, \mu, \rho)$ be a compact Ahlfors s-regular space $(0 \lt s \lt \infty)$. Let ${\xi_n}_{n\geq 1}$ be exponentially mixing and the probability law μ be Ahlfors s-regular, then the random covering set $E \in \mathcal{G}^{s_0}(X)$ a.s., where $s_0 = \inf\{t \geq 0: \sum_{n=1}^{\infty} r_n^t < \infty\}.$

Proof. For $t < s_0$, by Theorem [4.2,](#page-16-0) we have

$$
\mu(\limsup_{n\to\infty} B(\xi_n, r_n^{t/s})) = 1 \quad a.s.,
$$

since $\sum_{n=1}^{\infty} (r_n^{t/s})^s = \infty$. Applying Corollary [1.2,](#page-2-1) we finish the proof.

4.3. Limsup sets generated by rectangles.

The mass transference principle from balls to limsup sets generated by rectangles was established by [\[24\]](#page-18-22) in 2015. In 2021, Ding [\[4\]](#page-18-3) extended the result to product compact metric spaces, and proved that such sets have large intersection property. In this subsection, we use a different method to show the large intersection property of limsup sets generated by rectangles.

For $1 \leq i \leq d$, let (X_i, ρ_i) be a compact metric spaces equipped with a Borel probability measure μ_i which is Ahlfors s_i -regular, that is, there exists a constant $\delta_i > 1$ such that δ_i^{-1} $\sum_{i=1}^{i} r^{s_i} \leq \mu_i(B_i(x,r)) \leq \delta_i r^{s_i}$, here $B_i(x,r)$ is the open ball in X_i . We consider the metric space $(\prod_{i=1}^{d} X_i, \rho)$ with the measure μ , where

$$
\rho(x,y) = \max_{1 \le i \le d} \{\rho_i(x_i, y_i)\}
$$

for $x = (x_1, \ldots, x_d), y = (y_1, \ldots, y_d) \in \prod_{i=1}^d X_i$ and μ satisfies that $\mu(\prod_{i=1}^d A_i) =$ $\prod_{i=1}^d \mu_i(A_i)$ for A_i is a Borel set in X_i . The metric space $(\prod_{i=1}^d X_i, \rho)$ is compact.

Theorem 4.4. For $1 \leq i \leq d$, let $\{B_i(x_{n,i}, r_n)\}_{n \geq 1}$ be a sequence of balls in X_i with $r_n \to 0$ as $n \to \infty$. Let $\{a_i\}_{1 \leq i \leq d}$ be a sequence of positive numbers with $1 \leq a_1 \leq \cdots \leq a_d$. Assume that μ (lim sup n→∞ $\prod_{i=1}^d B_i(x_{n,i}, r_n) = 1$. Then $\limsup_{n \to \infty}$ $\prod_{i=1}^{d} B_i(x_{n,i}, r_n^{a_i}) \in \mathcal{G}^s(\prod_{i=1}^{d} X_i)$, where

$$
s = \min_{1 \le i \le d} \left\{ \frac{\sum_{j=1}^d s_j + a_i \sum_{j=1}^i s_j - \sum_{j=1}^i a_j s_j}{a_i} \right\}.
$$

Proof. We assume that $r_n < 1$, for $n \ge 1$. Denote $B_n = \prod_{i=1}^d B_i(x_{n,i}, r_n)$ and $R_n =$ $\prod_{i=1}^d B_i(x_{n,i}, r_n^{a_i})$. Let $x = (x_1, \ldots, x_d)$ and $y = (y_1, \ldots, y_d)$. For $t < s$ and $y \in R_n$, we have

$$
\phi_t(\mu, R_n, y) = \int_{R_n} \rho(x, y)^{-t} d\mu(x) \le \int_{\prod_{i=1}^d B_i(y_i, 2r_n^{a_i})} \rho(x, y)^{-t} d\mu(x).
$$

Let j be the integer with $\sum_{i=1}^{j-1} s_i < t \leq \sum_{i=1}^{j} s_i$. Denote

$$
F_{1,j} = \left\{ x \in \prod_{i=1}^d B_i(y_i, 2r_n^{a_i}) \colon \max_{1 \le i \le j} \rho_i(x_i, y_i) < 4r_n^{a_j} \right\},
$$
\n
$$
F_{2,j} = \left\{ x \in \prod_{i=1}^d B_i(y_i, 2r_n^{a_i}) \colon \max_{1 \le i \le j-1} \rho_i(x_i, y_i) \ge r_n^{a_j} \right\}.
$$

Note that $R_n \subset F_{1,j} \cup F_{2,j}$. Using Fubini's theorem, we derive that

$$
\phi_t(\mu, R_n, y) \leq \int_{F_{1,j}} \rho(x, y)^{-t} d\mu(x) + \int_{F_{2,j}} \rho(x, y)^{-t} d\mu(x)
$$
\n
$$
\leq \int_{F_{1,j}} (\max_{1 \leq i \leq j} \rho_i(x_i, y_i))^{-t} d\mu_1(x_1) \dots d\mu_d(x_d) + \int_{F_{2,j}} (\max_{1 \leq i \leq j-1} \rho_i(x_i, y_i))^{-t} d\mu_1(x_1) \dots d\mu_d(x_d)
$$
\n
$$
\leq \beta_1 \prod_{i=j+1}^d r_n^{a_i s_i} \int_{\prod_{i=1}^j B_i(y_i, 4r_n^{a_j})} (\max_{1 \leq i \leq j} \rho_i(x_i, y_i))^{-t} d\mu_1(x_1) \dots d\mu_j(x_j)
$$
\n
$$
+ \beta_1 \prod_{i=j}^d r_n^{a_i s_i} \int_{\prod_{i=1}^{j-1} B_i(y_i, 2r_n^{a_i}) \setminus \prod_{i=1}^{j-1} B_i(y_i, r_n^{a_j})} (\max_{1 \leq i \leq j-1} \rho_i(x_i, y_i))^{-t} d\mu_1(x_1) \dots d\mu_{j-1}(x_{j-1})
$$
\n
$$
\leq \beta_1' \Big(\prod_{i=j+1}^d r_n^{a_i s_i} r_n^{-a_j t} \prod_{i=1}^j r_n^{a_i s_i} + \prod_{i=j}^d r_n^{a_i s_i} I_2 \Big), \tag{4.8}
$$

where β_1 , β'_1 are absolute constants. Writting $f = (\max_{1 \le i \le j-1} \rho_i(x_i, y_i))^{-t}$, we have

$$
I_{2} = \int_{\prod_{i=1}^{j-1} B_{i}(y_{i}, 2r_{n}^{a_{i}}) \setminus \prod_{i=1}^{j-1} B_{i}(y_{i}, r_{n}^{a_{j}})} \left(\int_{0}^{f} du \right) d\mu_{1}(x_{1}) \dots d\mu_{j-1}(x_{j-1})
$$

\n
$$
= \int_{\prod_{i=1}^{j-1} B_{i}(y_{i}, 2r_{n}^{a_{i}}) \setminus \prod_{i=1}^{j-1} B_{i}(y_{i}, r_{n}^{a_{j}})} r_{n}^{-a_{j}t} d\mu_{1}(x_{1}) \dots d\mu_{j-1}(x_{j-1})
$$

\n
$$
- \int_{\prod_{i=1}^{j-1} B_{i}(y_{i}, 2r_{n}^{a_{i}}) \setminus \prod_{i=1}^{j-1} B_{i}(y_{i}, r_{n}^{a_{j}})} \left(\int_{f}^{r_{n}^{-a_{j}t}} du \right) d\mu_{1}(x_{1}) \dots d\mu_{j-1}(x_{j-1})
$$

\n
$$
\leq r_{n}^{-a_{j}t} \Biggl(\prod_{i=1}^{j-1} \mu_{i} (B_{i}(y_{i}, 2r_{n}^{a_{i}})) - \prod_{i=1}^{j-1} \mu_{i} (B_{i}(y_{i}, r_{n}^{a_{j}})) \Biggr)
$$

\n
$$
- \int_{0}^{r_{n}^{-a_{j}t}} du \int_{\prod_{i=1}^{j-1} B_{i}(y_{i}, 2r_{n}^{a_{i}}) \setminus \prod_{i=1}^{j-1} B_{i}(y_{i}, u^{-1/t})} d\mu_{1}(x_{1}) \dots d\mu_{j-1}(x_{j-1})
$$

\n
$$
= -r_{n}^{-a_{j}t} \prod_{j=1}^{j-1} \mu_{i} (B_{i}(y_{i}, r_{n}^{a_{j}})) + \int_{0}^{r_{n}^{-a_{j}t} \prod_{j=1}^{j-1} \mu_{i} (B_{i}(y_{i}, u^{-1/t})) du
$$

\n(4.9)

$$
\leq -r_n^{-a_j t} \prod_{i=1}^{j-1} \delta_i^{-1} r_n^{a_j s_i} + \prod_{i=1}^{j-1} \delta_i \int_0^{r_n^{-a_j t}} u^{-\sum_{i=1}^{j-1} s_i/t} du \leq \alpha r_n^{-a_j t} \prod_{i=1}^{j-1} r_n^{a_j s_i},
$$

where $\alpha > 0$ is an absolute constant. Therefore by [\(4.8\)](#page-17-0) and [\(4.9\)](#page-17-1),

$$
\phi_t(\mu, R_n, y) \le \beta_1'(1+\alpha) \prod_{i=j+1}^d r_n^{a_i s_i} r_n^{-a_j t} \prod_{i=1}^j r_n^{a_j s_i}.
$$

Hence there exists an absolute constant $\beta_2 > 0$ such that

$$
\frac{I_t(\mu, R_n)\mu(B_n)}{\mu(R_n)^2} \le \beta_2 r_n^{-a_j t + a_j \sum_{i=1}^j s_i + \sum_{i=1}^d s_i - \sum_{i=1}^j a_i s_i} \le \beta_2.
$$
\n(4.10)

From the proof of Theorem [1.1,](#page-1-1) for any t such that $\sup_{n\geq 1} \frac{I_t(\mu, R_n)\mu(B_n)}{\mu(R_n)^2}$ $\frac{\mu(R_n)\mu(B_n)}{\mu(R_n)^2} < \infty$, we have lim sup $\limsup_{n\to\infty} R_n \in \mathcal{G}^t(\prod_{i=1}^d X_i)$. Note that [\(4.10\)](#page-17-2) holds for any $t < s$, then $\limsup_{n\to\infty} R_n \in$ $\mathcal{G}^s(\prod_{i=1}^d X_i).$

Acknowledgements. The work was supported by NSFC 11671151 and 11871208.

REFERENCES

- [1] J. M. Aubry and S. Jaffard, Random wavelet series. Comm. Math. Phys. 227 (2002), no. 3, 483514.
- [2] V. Beresnevich and S. Velani, A mass transference principle and the Duffin-Schaeffer conjecture for Hausdorff measures. Ann. of Math. (2) 164 (2006), no. 3, 971992.
- [3] Y. Bugeaud, Intersective sets and Diophantine approximation. Michigan Math. J. 52 (2004), no. 3, 667682.
- [4] N. Ding, Large intersection properties for limsup sets generated by rectangles in compact metric spaces. Fractals 29 (2021), no. 6, 2150137, 13 pp.
- [5] A. Durand, Sets with large intersection and ubiquity. Math. Proc. Cambridge Philos. Soc. 144 (2008), no. 1, 119144.
- [6] A. Durand, Large intersection properties in Diophantine approximation and dynamical systems. J. Lond. Math. Soc. (2) 79 (2009), no. 2, 377398.
- [7] A. Durand, *Singularity sets of Lévy processes*. Probab. Theory Related Fields 143 (2009), no. 3-4, 517544.
- [8] A. Durand, *On randomly placed arcs on the circle*. Recent developments in fractals and related fields, 343351, Appl. Numer. Harmon. Anal., Birkh¨auser Boston, Boston, MA, 2010.
- [9] K. Falconer, Sets with large intersection properties. J. London Math. Soc. (2) 49 (1994), no. 2, 267280.
- $[10]$ D. Färm and T. Persson, *Large intersection classes on fractals*. Nonlinearity 24 (2011), no. 4, 12911309.
- [11] J. Fraser, Assouad Dimension and Fractal Geometry (Cambridge Tracts in Mathematics). Cambridge University Press, Cambridge, 2020. doi:10.1017/9781108778459
- [12] J. Heinonen, Lectures on analysis on metric spaces, Universitext. Springer-Verlag, New York, 2001.
- [13] Z.-N. Hu, W.-C. Cheng and B. Li, On the hitting probabilities of limsup random fractals. Fractals (2022) doi: 10.1142/S0218348X22500554.
- [14] Z.-N. Hu and B. Li, Random covering sets in metric space with exponentially mixing property. Statist. Probab. Lett. 168 (2021), Paper No. 108922, 7 pp.
- [15] Z.-N. Hu, B. Li and Y.-M. Xiao, On the intersection of dynamical covering sets with fractals. Math. Z. (2022) doi:10.1007/s00209-021-02924-2.
- [16] A. Käenmäki, T. Rajala and V. Suomala, *Existence of doubling measures via generalised nested cubes.* Proc. Amer. Math. Soc. 140 (2012), no. 9, 32753281.
- [17] D. Khoshnevisan, Y. Peres, Y.-M. Xiao, Limsup random fractals. Electron. J. Probab. 5 (2000), no. 5, 24 pp.
- [18] P. Mattila, Absolute continuity and uniqueness of measures on metric spaces. Trans. Amer. Math. Soc. 266 (1981), no. 1, 233242.
- [19] P. Mattila, Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability. Cambridge Studies in Advanced Mathematics, 44. Cambridge University Press, Cambridge, 1995.
- [20] E. J. McShane, Jensen's inequality. Bull. Amer. Math. Soc. 43 (1937), no. 8, 521527.
- [21] F. Negreira and E. Sequeira, Sets with large intersection properties in metric spaces. J. Math. Anal. Appl. 511 (2022), no. 1, Paper No. 126064, 25 pp.
- [22] T. Persson, Inhomogeneous potentials, Hausdorff dimension and shrinking targets. Ann. H. Lebesgue 2 (2019), 137.
- [23] T. Persson, A mass transference principle and sets with large intersections. arXiv:1911.10399v2
- [24] B.-W. Wang, J. Wu and J. Xu, Mass transference principle for limsup sets generated by rectangles. Math. Proc. Cambridge Philos. Soc. 158 (2015), no. 3, 419437.

Zhang-nan Hu, School of Mathematics, South China University of Technology, Guangzhou, 510641, China

Email address: hnlgdxhzn@163.com

Bing Li, School of Mathematics, South China University of Technology, Guangzhou, 510641, China

Email address: scbingli@scut.edu.cn

Linqi Yang, School of Mathematics, South China University of Technology, Guangzhou, 510641, China

Email address: 2981502134@qq.com