# Robust Model-based Inference for Non-Probability Samples

Ali Rafei<sup>1</sup>, Michael R. Elliott<sup>\*1</sup>, and Carol A. C. Flannagan<sup>2</sup>

<sup>1</sup>Survey Methodology Program, University of Michigan <sup>2</sup>University of Michigan Transportation Research Institute

**Abstract:** With the ubiquitous availability of unstructured data, growing attention is paid as how to adjust for selection bias in such non-probability samples. The majority of the robust estimators proposed by prior literature are either fully or partially design-based, which may lead to inefficient estimates if outlying (pseudo-)weights are present. In addition, correctly reflecting the uncertainty of the adjusted estimator remains a challenge when the available reference survey is complex in the sample design. This article proposes a fully model-based method for inference using non-probability samples where the goal is to predict the outcome variable for the entire population units. We employ a Bayesian bootstrap method with Rubin's combing rules to derive the adjusted point and interval estimates. Using Gaussian process regression, our method allows for kernel matching between the non-probability sample units and population units based on the estimated selection propensities when the outcome model is misspecified. The repeated sampling properties of our method are evaluated through two Monte Carlo simulation studies. Finally, we examine it on a real-world non-probability sample with the aim to estimate crash-attributed injury rates in different body regions in the United States.

Keywords and phrases: Non-probability sample, doubly robust, Bayesian bootstrapping, Kernel matching, Gaussian process regression.

## 1. Introduction

With rapid advances in automated measurement technologies, massive amount of unstructured data become ubiquitously available in various fields. Political views shared on social media, Google searches for particular terms, payment transactions recorded by online stores, electronic health records of patients admitted to a chain of hospitals, videos captured by traffic cameras, and GPS trajectory data by satellite are among the common examples of this revolution. This phenomenon coincides with downward trends in the response rate of traditional probability surveys over recent decades, which not only escalate the survey implementation costs, but imperil the validity of their findings in the absence of relevant auxiliary information for non-response adjustment (Groves, 2011; Miller, 2017). As a result, there is growing demand to leverage these novel data sources, as a cheaper and faster alternative to probability surveys, to inform policy (Johnson and Smith, 2017; Senthilkumar et al., 2018). This has led to a surge of research into methods of inference for non-probability samples to mitigate the inherent selection bias in such data (Kim and Tam, 2021; Meng et al., 2018; Rao, 2021).

Aside from this shift, there are situations where data collection is expensive or must be administered onsite, and therefore conducting a probability survey may not be practical. For instance, the national automotive sampling system (NASS) is responsible for collecting detailed medical and biomechanical information about the occupants of traffic accidents in the U.S., which demands for well-equipped labs. Although the Crash Injury Research Engineering Network (CIREN) fills this gap in NASS, where more succinct data are gathered regularly through larger probability surveys, the additional data items in CIREN are achieved at the expense of a lack of representativity (Elliott et al., 2010). While it is unknown how well the pool of the traumatic patients referring to CIREN's level 1 trauma centers represents the population of all occupants involved in traffic accidents across the U.S., the collaborating centers tend

<sup>\*</sup> Corresponding author; address: 426 Thompson St. Ann Arbor, MI 48109. Rm 4068 ISR, email: mrelliot@umich.edu.

to be located in the urban areas. More importantly, CIREN tends to recruit severely injured occupants, which may result in overestimated risk of injuries (Elliot, 2013; Flannagan and Rupp, 2009).

The prior literature proposed various robust approaches for finite population inference based on a non-probability sample under situations where a parallel "reference survey" is available as the external benchmark (Chen et al., 2019; Rafei et al., 2022, 2021). One common assumption among these approaches is that units of the reference survey have been selected independently with unequal probabilities of selection. This, however, may not hold in practice as probability samples often come with additional complexity in their design. For instance, U.S. federal statistical agencies widely use a stratified multi-stage cluster design in the conduct of their large-scale surveys, such as the American Community Survey (ACS), Current Population Survey (CPS), and National Health Interview Survey (NHIS). These sampling designs are expected to improve sampling efficiency not only statistically but with respect to cost and logistics (Cochran, 1977; Kish, 1965). To maximize statistical efficiency while assuring broad representativeness, a common practice involves partitioning the population into a large number of strata and then selecting a few clusters, ideally two, randomly within each stratum (Zhou, 2014).

To mitigate the potential selection bias in CIREN, which shapes the goal of the application of this article, the literature suggests combining the data with the CDS as a reference survey, because the two samples potentially represent the same population and have a wide range of auxiliary variables in common (Elliott et al., 2010). However, one notable challenge is that units of CDS are selected through a stratified three-stage cluster sampling design (Chen et al., 2015). Elliott et al. (2010) recognize the importance of adjusting for the cluster-level variance of CDS, and suggest using a jackknife repeated replication method to incorporate the uncertainty of the estimated pseudo-weights into the variance estimator. Under a pseudo-likelihood setting, a modified linearization technique is proposed by Wang et al. (2020) to estimate the variance under inverse propensity score weighting (IPSW). When prediction modeling (PM) is incorporated, it is also essential to account for the fixed and random effects of stratification and clustering, respectively, in addition to the sampling weights (Little, 2004; Little and Zheng, 2007). Failure to do so may cause some degrees of model misspecification, which appear as bias in point and variance estimates.

Furthermore, to avoid imposing a pseudo-likelihood structure under a Bayesian framework while adjusting for unequal sampling weights of the reference survey in estimating the PS, Rafei et al. (2022) employed a two-step pseudo-weighting approach proposed by Elliott and Valliant (2017). A noteworthy complication arose from the first step where the sampling weights of the reference survey had to be modeled conditional on the observed auxiliary information explicating the selection mechanism in the non-probability sample. However, the correct structure of such a model may look complicated in the finite population as the weights are deemed to be a deterministic function of the design features of the reference survey. The association between the sampling weights and auxiliary information thus depends on how the latter correlates with those design features. Unless the strength of this association is trivial, misspecifying the working model for sampling weights is likely to induce some degree of residual bias in the final estimates (Rafei et al., 2020).

More importantly, the ultimate form of the adjusted estimators in previous literature mostly involve design-based terms, and therefore, are subject to the general drawbacks of design-based methods. For instance, both the QR and PM terms in the proposed estimator of Rafei et al. (2021) appeared as (pseudo-)weighted sums, and that proposed in Rafei et al. (2022) still contained a similar structure in the PM term. Despite being design-consistent, it is well-understood that any presence of influential (pseudo-)weights may cause inflated variance in such estimators, especially when the sample size is small (Chen et al., 2017; Zangeneh, 2012; Zhang and Little, 2011). From a model-based perspective, Zheng and Little (2003) argue that the HT-estimator is equivalent to a rigid no-intercept regression model with heteroscedasticity. Although the HT-estimator is unbiased, if the true data model is far from the implied model, the resulting estimator will be highly inefficient (Basu, 1971).

Alternatively, one can use a fully model-based approach where the goal is to impute the outcome variable(s) for the entire non-sampled units of the population (Little and Zheng, 2007; Smith, 1983).

Given the observed outcome for the entire population, one can directly quantify the unknown population parameter of interest. This idea clearly eliminates the need for the remaining design-based term in Rafei et al. (2022)'s adjusted estimator, and therefore not only fills the above-mentioned gaps in bias adjustment when using more flexible models but fully satisfies the likelihood principle (Berger and Wolpert, 1988; Little, 2004). As a direct advantage, one can easily expand such a method to a Bayesian setting. Alternative pseudo-likelihood approaches are limited to distributions with an exponential family and have known problems in uncertainty propagation under a Bayesian setting (Gunawan et al., 2020; Savitsky et al., 2016; Williams and Savitsky, 2021).

The present article proposes a fully model-based approach for robust inference based on non-probability samples when the available reference survey is based on a complex probability sample. To this end, a partially linear Gaussian process regression is employed to non-parametrically link the estimated propensity scores (PS) to the response surface (Kammann and Wand, 2003). This approach was termed the Gaussian Process of Propensity Prediction (GPPP) by Rafei et al. (2022). However, because the auxiliary variables are unmeasured for the non-sampled units, we generate synthetic populations beforehand by undoing the sampling mechanism of the reference survey through a Pólya urn scheme (Dong et al., 2014). The use of a synthetic population eliminates the need for a pseudo-likelihood approach and for modeling the sampling weights when estimating the PS. To be able to correctly account for the cluster-level variance of the reference survey, we choose to use the two-step bootstrap-Pólya posterior technique proposed by Zhou et al. (2016c). Mercer (2018) uses a similar idea for inference under a non-probability setting, though with limited theoretical details expressed. Nevertheless, his method utilizes Bayesian Additive Regression Trees (BART) and penalized spline regression for modeling the outcome while our method is based on a GP regression model.

The rest of the article is organized as follows: In Section 2, we start by defining the necessary notation and assumptions required for valid inference under a non-probability sample setting and then describe the proposed method under a rigorous mathematical framework. Section 3 evaluates the repeated sampling properties of the proposed method and contrasts its performance with alternative approaches through a simulation study. Section 4 describes the datasets and auxiliary variables we utilize for the empirical study and reports results of bias adjustment on CIREN/CDS data. Finally, Section 5 reviews the strengths and weaknesses of the study in more detail and suggests some future research directions. Supplemental information about the simulation and empirical studies are provided in Appendix 8.

## 2. Methods

#### 2.1. Notation and Assumptions

Suppose U is a finite population of known size  $N < \infty$ , which is partitioned into H homogeneous strata with  $M_h$  clusters nested within stratum h (h = 1, 2, ..., H) and  $M = \sum_{h=1}^{H} M_h$  the total number of clusters in U. Also, let  $N_{hj}$  be the size of cluster j in stratum h, and thus  $N = \sum_{h=1}^{H} \sum_{j=1}^{N_h} N_{hj}$ . One can identify the stratum and cluster of unit i in U through an (H-1)-dimensional vector of dummies denoted as  $d_i$  and an (M-H)-dimensional vector of dummies denoted as  $c_i$ , respectively. For each i = 1, 2, ..., N, we consider  $y_i$  to be the realized values of a scalar outcome variable, Y, and  $x_i = [x_{i1}, x_{i2}, ..., x_{ip}]^T$  the values of a p-dimensional set of relevant auxiliary variables, X, in U.

Let  $S_A$  be a non-probability sample selected from U with  $[y_i, x_i^T, d_i^T, c_i^T]$  observed and  $n_A$  being the sample size. Descriptive inference aims at learning about an unknown parameter of U that is a function of Y. The goal here is to draw point and interval estimates for the population mean, i.e.  $Q(y) = \overline{y}_U = \sum_{i=1}^N y_i/N$ . Suppose  $\delta_i^A = I(i \in S_A)$  represents the inclusion indicator variable of  $S_A$  for  $i \in U$  whose distribution can be explained by  $x_i$ . To make valid inference for  $S_A$  possible, we assume that conditions **C1-C3** determined in Rafei et al. (2022) hold.

Consider  $S_R$  to be a parallel reference survey of size  $n_R$ , for which the same set of covariates, X, has been measured, though Y has to be unobserved. Also, let  $\delta_i^R = I(i \in S_R)$  denote the inclusion indicator variable

associated with  $S_R$  for  $i \in U$ . As highlighted in the introduction, we concentrate on a setting where units of  $S_R$  are selected via a stratified two-stage clustered sampling design. Note that probability samples often involve more than two stages of clustering, but the majority of existing theories for complex sample survey analysis are built upon the "ultimate clustering principle" where the second and higher stages of clustering are ignored, and the sample is treated as if it follows a single-stage clustering design (Wolter, 2007). From each given stratum h (h = 1, 2, ..., H), we draw  $m_h$  clusters as the primary sampling units (PSU) with first-order inclusion probabilities of  $\pi_{j|h}^R$  ( $j = 1, 2, ..., M_h$ ), and  $n_{hj}$  subjects from the  $j^{\text{th}}$  cluster as the secondary sampling units (SSU) with second-order inclusion probabilities of  $\pi_{i|hj}^R$  ( $i = 1, 2, ..., N_{hj}$ ). Note that  $n_R = \sum_{h=1}^H \sum_{j=1}^{m_h} n_{hj}$ .

While the proposed methods in this article impose no restriction on the number of PSUs per stratum, the reference surveys across both the simulation and application parts involve a two-PSU-per-stratum design, i.e.  $m_h = 2$  for all h = 1, 2, ..., H. As discussed earlier, such a sampling design is widely used among U.S. government statistical agencies. One common practical way to randomly select the PSUs within strata is probability proportional-to-size (PPS). If  $v_{hj}$  denotes the measure of size (MOH) associated with PSU j of stratum h, under a PPS technique,  $\pi_{hj}^R = v_{hj} / \sum_{j=1}^{M_h} v_{hj}$ . In practice, probability survey data are only accompanied by an external set of sampling weights that are supposed to be inversely proportional to the joint selection probabilities, i.e.  $w_i^R \propto 1/\pi_{hji}^R$ , where  $\pi_{hij}^R = \pi_{hj}^R \pi_{i|hj}^R$ , which are only observed for  $i \in S_R$ . It is likely that the sampling weights already involve post-survey adjustments for ineligibility, non-response, and non-coverage errors as well (Valliant et al., 2018). Throughout this article, we assume that the given weights correctly adjust for all these potential sources of bias. Defining  $Z = \{D, C, w^R\}$  as the set of all sampling design variables in  $S_R$ , we also assume Z is fully observed in both  $S_R$  and  $S_A$  with no item missingness.

Despite the possible overlap or correlation between X and Z, in addition to the aforementioned conditions, we also assume:

C4. Independence of Samples— conditional on  $\{X, Z\}$ ,  $S_R$  and  $S_A$  are selected independently, i.e.  $\delta^A \perp \delta^R | X, Z$ .

Then, considering **C1-C4**, the joint density of  $y_i$ ,  $\delta_i^A$  and  $\delta_i^R$  can be factored as follows:

$$p(y_i, \delta_i^A, \delta_i^R | x_i, z_i; \theta, \beta) = p(y_i | x_i, z_i; \theta) p(\delta_i^A | x_i; \beta) p(\delta_i^R | z_i), \quad \forall i \in U$$

$$(2.1)$$

where  $\eta = (\theta, \beta)$  are unknown parameters indexing the conditional distribution of Y|X, Z and  $\delta^A|X$ , respectively. In the following subsection, we describe a fully model-based strategy for inference based on  $S_A$  that models both  $p(y_i|x_i, z_i; \theta)$  and  $p(\delta_i^A|x_i; \beta)$ .

#### 2.2. Model-based inference

Consider  $\hat{\pi}^A$  and  $\hat{y}$  to be consistent estimates of the pseudo-selection probabilities in  $S_A$  and outcome variable in  $S_C = S_A \cup S_R$ , respectively. The proposed robust estimators by Chen et al. (2019) and Rafei et al. (2021) take the following form:

$$\widehat{\overline{y}}_U = \left(\sum_{i \in S_A} \frac{(y_i - \widehat{y}_i)}{\widehat{\pi}_i^A} + \sum_{i \in S_R} \frac{\widehat{y}_i}{\pi_i^R}\right) / N$$
(2.2)

which involves two Horvitz-Thompson (HT) estimators from  $S_A$  and  $S_R$ . While preserving double robustness, Rafei et al. (2022) propose to drop the denominator from the first term as below:

$$\widehat{\overline{y}}_U = \left(\sum_{i \in S_A} \left(y_i - \widehat{y}_i\right) + \sum_{i \in S_R} \frac{\widehat{y}_i}{\pi_i^R}\right) / N$$
(2.3)

The main goal here was to reduce the computational burden under a Bayesian framework by limiting the imputations to the combined sample. However, Eq. 2.3 still relies on a HT term which can yield

inefficient estimates if there are extremely small values of  $\pi_i^R$ . To get rid of this design-based term, we use the estimator proposed by Liu et al. (2021):

$$\widehat{\overline{y}}_U = \left(\sum_{i \in S_A} \left(y_i - \widehat{y}_i\right) + \sum_{i \in U} \widehat{y}_i\right) / N$$
(2.4)

where  $\hat{y}_i$  is the prediction of  $y_i$  for  $i \in U$ . This is a fully model-based estimator in the sense that the unobserved outcome, Y, is imputed for the entire population units.

Analogous to Rafei et al. (2022), the goal is to specify a model for the joint distribution of  $(y_i, \delta_i^A)$  across the units of U, which can be formulated by

$$p(y_i, \delta_i^A | x_i, z_i; \theta, \beta) = p(y_i | x_i, z_i, \delta_i^A; \theta) p(\delta_i^A | x_i; \beta), \quad i \in U$$

$$(2.5)$$

In a Bayesian setting, descriptive inference about  $\overline{y}_U$  is attained by deriving its posterior predictive distribution given the observed data. By integrating out the unknown parameters, we have

$$p(\overline{y}_U|y_A, \delta_U^A, x_U, z_U) = \int \int p(\overline{y}_U|y_A, \delta_U^A, x_U, z_U, \theta, \beta) p(\theta, \beta|y_A, \delta_U^A, x_U, z_U) d\theta d\beta$$
(2.6)

where subscripts U and A denote a vector of the variables defined in U and  $S_A$ , respectively. Thus  $\{X, Z\}$  have to be observed for all of the units of U in order to simulate the posterior predictive distribution of  $\overline{y}_U$ .

In a practical setting, the measurement of  $\{X, Z\}$  is often confined to the pooled sample. To circumvent this issue, we choose to generate a finite set of synthetic populations as the first step by undoing the sampling mechanism in  $S_R$ , which can be performed non-parametrically through the idea of finite population Bayesian bootstrapping (FPBB) (Dong et al., 2014; Little and Zheng, 2007). Theoretical details of such a method adapted to reference surveys with a complex design, are provided in the following subsection. For any given synthetic population, we then predict the pseudo-selection probabilities of  $S_R$ , i.e.  $\pi_i^A$ , as the second step via a QR model. Having  $\pi_i^A$  imputed for the entire units of the synthetic population, the third step involves predicting  $y_i$  for the entire synthetic population by fitting a PM on  $S_A$  with a flexible function of  $\hat{\pi}_i^A$  as a predictor in the model. As demonstrated in Rafei et al. (2022), this leads to double robustness in the adjusted estimator.

Once  $y_i$  is imputed for all units of a synthetic population, one can compute Eq. 2.4 and derive point and interval estimates for the population mean by combining the estimates across synthetic populations based on Rubin's multiple imputation rules (Rubin, 1976). This research implements a frequentist approach joint with resampling techniques to train the model parameters based on maximum likelihood estimation (MLE), and to suitably capture the uncertainties due to modeling and complex sampling. We do this for two reasons: (1) fitting Bayesian models repeatedly on synthetic populations is demanding computationally for the joint estimation of  $(y_i, \pi_i^A)$  especially when N is large, and (2) public-use survey data usually lack detailed information about the selection probabilities across different stages that are essential for reverting the selection mechanism in multi-stage designs. Figure 1 visualizes these steps for the proposed bias adjustment procedure. The shaded blue areas show the observed data while the shaded red areas display the imputed data at each step.

## 2.2.1. STEP 1: Weighted finite population Bayesian bootstrap

The first step involves generating synthetic populations using Finite Population Bayesian Bootstrapping (FPBB). Rafei et al. (2022) discussed in detail how FPBB operates in a probability sample setting, where the goal is to derive the posterior distribution of a given set of variables, such as  $\{X, Z\}$ , for the non-sampled units of U under an exchangeability assumption. Recently, Zhou et al. (2016a) have expanded the weighted variant of FPBB based on Pólya posterior (Cohen, 1997; Dong et al., 2014) to create synthetic populations based on a probability sample with a stratified two-stage clustering design.



Fig 1: Data structure in the population and the combined sample

However, their method requires the first and second-order probabilities of inclusion to be known to the analyst such that sampling stages can be reverted hierarchically. Most often, public-use survey data contain merely a variable representing the sampling weights whose reciprocal can be regarded as marginal selection probabilities, but lack selection probabilities at different stages. Because of this constraint, Zhou et al. (2016c) propose an alternative two-step approach, where design-adjusted bootstrap resamples are initially selected, and a weighed pólya posterior is used to create synthetic populations as the second step.

In this study, we employ a similar method using Rao and Wu (1988)'s bootstrap technique for  $S_R$ , which is complex in design. It basically uses the idea of "with replacement" bootstrap proposed by McCarthy and Snowden (1985), with an additional rescaling factor applied to the original sampling weights. The asymptotic properties of the bootstrap technique, including consistency, have been well-established for both smooth and non-smooth statistics, such as quantiles (Shao and Chen, 1998). Besides, a bootstrap method can be easily adapted to designs with more than two PSUs per stratum, unlike some other resampling techniques. Compared to the conventional bootstrap, the Rao-Wu variant yields more stable variance estimates when the frequency of PSUs within strata is small at the population level. This is often the case in two-PSU-per-stratum designs where a large pool of strata exists in the sample. For  $S_A$ , whose elements are assumed to be selected independently according to C3, we use a regular bootstrap method where with replacement random samples of size  $n_A$  are drawn repeatedly from the original sample (Efron, 1992).

For Rao-Wu's bootstrap technique, the simplest procedure involves drawing random samples of  $m_h^* = m_h - 1$  PSUs with replacement from the stratum h (h = 1, 2, ..., H) of the original sample. These re-samples are then pooled to construct a bootstrap sample of size  $m^* = m - H$ , and the sampling weights are then rescaled to  $w_{hji}^* = w_{hji}^R m_h/(m_h - 1)$ . This procedure is replicated B times for the  $S_R$  to create  $S_R^{(b)}$ 's (b = 1, 2, ..., B) with  $n_R^* = n_R - H$  being the new sample size. Zhou et al. (2016a) note that because the new weights implicitly carry over the information in strata, PSUs, and unequal selection probabilities, one can treat the bootstrapped samples as if units are selected independently with unequal weights, and accordingly, the subscript hj can be dropped, and  $w_i^*$ 's are normalized such that  $N = \sum_{i=1}^{n-H} w_i^*$ . In the following, we show how one can create a synthetic population based on a given  $S_R^{(b)}$  using a Pólya posterior distribution. We also denote the  $b^{\text{th}}$  bootstrap replicate of  $S_A$  by  $S_A^{(b)}$  (b = 1, 2, ..., B).

Let  $t = \{t_1, t_2, ..., t_J\}$  represents a *J*-dimensional vector containing all distinct values of  $(x_i, z_i)$  in  $S_R^{(b)}$ , and  $\xi = \{\xi_1, \xi_2, ..., \xi_J\}$  denotes the vector of probabilities  $p(x_i = t_j | \xi) = \xi_j$  for  $i = 1, 2, ..., n_R^*$ , j = 1, 2, ..., J, where  $\sum_{j=1}^J \xi_j = 1$ . Now, suppose  $n_j^*$  and  $r_j^*$  are the frequencies of units taking the vector  $t_j$  in  $S_R^{(b)}$  and  $\overline{S}_R^{(b)} = U - S_R^{(b)}$ , respectively, for j = 1, 2, ..., J. It is clear that  $\sum_{j=1}^J n_j^* = n_R^*$ , and  $\sum_{j=1}^{J} r_j^* = N - n_R^*.$  Considering a non-informative Haldane prior, i.e.  $\xi \sim Dirichlet(0, ..., 0)$ , with a multinomial likelihood function of  $p(n_1^*, n_2^*, ..., n_J^*|\xi) \propto \prod_{j=1}^{J} \xi_j^{n_j^*}$ , the posterior distribution of  $\xi$  in  $\overline{S}_R^{(b)}$  is given by  $(\xi|n_1^*, n_2^*, ..., n_J^*|) \sim Dirichlet(n_1^* - 1, n_2^* - 1, ..., n_J^* - 1)$ . Using a weighted Pólya Urn Scheme, Dong et al. (2014) prove that the posterior predictive draws of  $r_j^*$  can be directly generated by

$$p(r_1^*, r_2^*, ..., r_J^* | w_1^*, w_2^*, ..., w_J^*) = \frac{\prod_{j=1}^K \Gamma(w_j^* + r_j^*) / \Gamma(w_j^*)}{\Gamma(2N - n_R^*) / \Gamma(N)}$$
(2.7)

where the  $w_j^*$ 's are the normalized rescaled weights of  $S_R^{(b)}$  such that  $\sum_{j=1}^K w_j^* = N$ .

Finally, one can use Cohen (1997)'s computational algorithm to obtain the weighted-Polýa posterior draws of  $\{X, Z\}$  based on  $S_R^{(b)}$ . This involves taking a Polýa sample of size  $N - n_R^*$  denoted by  $(X^{(ls)}, Z^{(ls)})$  from the urn  $(X^{(l)}, Z^{(l)})$  using the following selection probabilities:

$$\zeta_i = \frac{l_{i,k-1} \left(\frac{N - n_R^*}{n_R^*}\right) + w^{(l)^*} - 1}{(k-1) \left(\frac{N - n_R^*}{n_R^*}\right) + N - n_R^*}, \quad k = 1, 2, ..., N - n_R^* + 1$$
(2.8)

where  $w^{(l)^*}$  denotes the bootstrap weights associated with the *i*-th unit in the *l*-th replicate BB sample, and setting  $l_{i,0} = 0$ ,  $l_{i,k-1}$  denotes the frequency of unit *i* selected up to (k-1)-th selection (Zhou et al., 2016c). For the *l*-th posterior predictive draw,  $n_j^* + r_j^*$  represents the frequency of  $(x_i, d_i)$  in the synthetic population whose value is  $t_j$ . Therefore, for any given  $S_R^{(b)}$  (b = 1, 2, ..., B), one can use this approach to generate *L* synthetic populations, denoted by  $\hat{U}^{(b,l)}$  (l = 1, 2, ..., L). Note that the total number of generated synthetic populations will be  $B \times L$ .

## 2.2.2. STEP 2: Imputing the pseudo-selection probabilities in $S_A$

The second step involves modeling  $p(\delta_i^A = 1|x_i)$  with the aim to predict  $\pi_i^A$  for  $i \in \widehat{U}^{(b,l)}$  (b = 1, 2, ..., B; 1, 2, ..., L). Assuming that  $S_A$  has a Poisson sampling design with  $n_A$  being random, one can formulate the likelihood function across the population units by

$$L(\beta|\delta_U^A, x_U) = \prod_{i=1}^N p(\delta_i^A = 1|x_i, \beta)^{\delta_i^A} \left[1 - p(\delta_i^A = 1|x_i, \beta)\right]^{1-\delta_i^A}$$
(2.9)

where  $\beta$  the vector of parameters associated with the QR model. Note that although the population units are autocorrelated due to clustering, L is written as if units of U are independent. Considering a logistic regression model, we have

$$\pi_i^A = p(\delta_i^A = 1 | x_i; \beta) = \frac{exp\{\beta_0 + x_i^T \beta_1\}}{1 + exp\{\beta_0 + x_i^T \beta_1\}}$$
(2.10)

where  $\beta^T = [\beta_0, \beta_1^T]$ . A consistent estimate for  $\beta$  can be attained through an MLE method by solving a set of estimating equations. However, one major obstacle is that  $\delta_i^A$  is unobserved in a generated synthetic population  $\widehat{U}^{(b,l)}$ , and imputing this missing variable would require one to match units of  $S_A$  to units of  $\widehat{U}^{(b,l)}$  (Liu et al., 2021). To avoid this cumbersome task, we utilize a duplicate removal trick proposed by Wang et al. (2020).

For any given (b, l) (b = 1, 2, ..., B; l = 1, 2, ..., L), we initially append  $S_A^{(b)}$  to  $\hat{U}^{(b,l)}$ , and define  $\hat{U}^{(b,l)*} = S_A^{(b)} \cup^* \hat{U}^{(b,l)}$ , where  $\cup^*$  denotes the union of  $S_A$  and  $\hat{U}^{(b,l)}$  while allowing for duplicated units of  $S_A^{(b)}$  in  $\hat{U}^{(b,l)}$ . We then define a new indicator variable  $\delta_i^*$  in  $\hat{U}^{(b,l)*}$  taking the value of 1 for  $i \in S_A^{(b)}$  and 0 for

 $i \in \widehat{U}^{(b,l)}$ . According to Wang et al. (2020), for a given (b, l), one can show that

$$\begin{aligned} \pi_i^A &= p\left(i \in S_A^{(b)} | x_i, i \in \widehat{U}^{(b,l)}\right) \\ &= \frac{p\left(i \in S_A^{(b)} | x_i\right)}{p\left(i \in \widehat{U}^{(b,l)} | x_i\right)} \\ &= \frac{p\left(i \in S_A^{(b)} | x_i\right) / p\left(i \in \widehat{U}^{(b,l)*} | x_i\right)}{p\left(i \in \widehat{U}^{(b,l)} | x_i\right) / p\left(i \in \widehat{U}^{(b,l)*} | x_i\right)} \end{aligned}$$
(2.11)  
$$&= \frac{p\left(i \in S_A^{(b)} | x_i, i \in \widehat{U}^{(b,l)*}\right)}{p\left(i \in \widehat{U}^{(b,l)} | x_i, i \in \widehat{U}^{(b,l)*}\right)} \\ &= \frac{p_i(\beta^*)}{1 - p_i(\beta^*)} \end{aligned}$$

where  $\beta^*$  is a new vector of parameters associated with  $\widehat{U}^{(b,l)*}$  and  $p_i(\beta^*) = p(\delta_i^* = 1 | x_i, \beta^*)$ . Note that

$$p_i(\beta^*) = \frac{p\left(i \in S_A^{(b)} | \widehat{U}^{(b,l)}\right)}{p\left(i \in S_A^{(b)} | \widehat{U}^{(b,l)}\right) + p\left(i \in \widehat{U}^{(b,l)} | \widehat{U}^{(b,l)}\right)} \le \frac{1}{2}$$
(2.12)

and the equality holds only if  $p\left(i \in S_A | \widehat{U}^{(b,l)}\right) = 1$ . This ensures that  $0 < \widehat{\pi}_i^A < 1$ . Again, under the logistic regression model, we have

$$p_i(\beta^*) = p(\delta_i^* = 1 | x_i; \beta_i^*) = \frac{exp\{x_i^T \beta_i^*\}}{1 + exp\{x_i^T \beta_i^*\}}$$
(2.13)

where  $\beta^*$  can be estimated again using the standard MLE method. Considering (2.5), the estimate of  $\pi_i^A$  for  $i \in \hat{U}^{(b,l)}$  is given by

$$\widehat{\pi}_i^A = \frac{\widehat{p}_i(\beta^*)}{1 - \widehat{p}(\beta_i^*)} = \exp\{x_i^T \widehat{\beta}_i^*\}$$
(2.14)

Hence, for a given  $\hat{U}^{(b,l)}$ , to obtain the posterior distribution  $p(\pi_U^A|\delta_U^A, x_U)$ , all one needs is to append  $S_A^{(b)}$  to  $\hat{U}^{(b,l)}$  and fit a Bayesian logistic regression model on the pooled data. Note that the term odds of PS was also seen in the Propensity-adjusted Probability Prediction (PAPP) method in Rafei et al. (2020). However, its first term, which involved modeling the sampling weights, is dropped by expanding the reference survey to the population.

#### 2.2.3. STEP 3: Prediction modeling using a partially linear Gaussian process regression

The goal of PM here is to impute  $y_i$  for  $i \in U^{(b,l)}$  (b = 1, 2, ..., B; l = 1, 2, ..., L) by estimating  $E(y_i|x_i, z_i; \theta)$ , which is essential for the calculation of Eq. 2.4. To this end, one has to know about the true functional form of the response surface that governs the superpopulation structure from which U has been selected. Considering the fact that U is clustered, suppose the true PM is given by

$$g\left(E\left[y_{ij}|x_{ij}, d_{ij}, c_i, w_{ij}^R; \theta\right]\right) = \theta_0 + x_{ij}^T \theta_1 + d_{ij}^T \theta_2 + \log\left(w_{ij}^R\right) \theta_3 + u_i$$
(2.15)

where the subscript ij indicates the observed variable in the  $j^{\text{th}}$  unit of the  $i^{\text{th}}$  cluster,  $\theta$  is a (p + H + 1)-dimensional vector of unknown model parameters, g is an appropriate link function, and  $u_i \sim N(0, \sigma_u^2)$  denotes a random effect term associated associated with the cluster i (i = 1, 2, ..., M). Note that the

notational convention used in Eq. 2.15 suppresses showing the cluster indicator,  $c_i$ , on the right side of the equation.

In a non-probability sample setting, since  $y_i$  is only observed for  $i \in S_A$  the PM has to be trained based on  $S_A^{(b)}$  (b = 1, 2, ..., B). Considering **C2** and by the use of the Bayes rule, one can show that  $p(y_i|x_i, z_i; \theta) = p(y_i|\delta_i^A = 1, x_i, z_i; \theta)$ . This implies that a consistent estimate for  $\theta$  can be derived from  $S_A$ . Therefore, one can use a marginal likelihood function, restricted to  $S_A$ , to obtain a consistent estimate for  $\theta$  as below:

$$L(\theta|y_A, x_A, z_A) = \prod_{i=1}^m \prod_{j=1}^{n_i^A} \int_u p(y_{ij}|x_{ij}, z_{ij}, u_i; \theta) \phi(u_i; 0, \sigma_u^2) du_i$$
(2.16)

where  $\phi(.; \mu, \sigma^2)$  denotes a normal density function with mean  $\mu$  and variance  $\sigma^2$ . To obtain an MLE for  $\theta$  when the outcome is non-normal, one can utilize adaptive Gaussian quadrature to assess the integral in Eq. 2.15 (Pinheiro and Bates, 1995). It is important to note that there must be adequate sample from all H strata observed in  $S_A$  and  $(d_i, w_i^R)$  must be known for all  $i \in S_A$  such that one can fit the model in Eq. 2.15 on  $S_A$ . This introduces a big challenge for inference based on a non-probability sample, which will be further elaborated in the discussion.

In reality, however, the true functional form of the response surface in U is not known to the analyst. It is likely that the auxiliary variables, especially MOS, are non-linearly associated with the conditional expectation of the outcome variable or multi-way interaction effects are present among the covariates. Similar to Rafei et al. (2022), one can add the estimated  $\pi_i^A$  from Step II as a predictor to the PM with the aim to further protect against the misspecification of the PM. This leads to a doubly robust (DR) estimator, which preserves consistency if the underlying model for either QR or PM is correctly specified (Robins et al., 1994). Linear-in-weight (LWP) prediction is the most common type of a model-based DR estimator, which uses the inverse of  $\hat{\pi}_i^A$  as a linear predictor in the PM in addition to  $\{x_i, z_i\}$ , and is equivalent to the earliest class of DR methods, i.e. augmented inverse propensity weighting (AIPW) (Bang and Robins, 2005; Scharfstein et al., 1999). Zheng and Little (2003) realized that the use of a more flexible function of  $\hat{\pi}_i^A$  can yield improved efficiency in the ultimate DR estimator. The improvement is more tangible when extremely low values of  $\hat{\pi}_i^A$  are present (Zhang and Little, 2011). This problem often arises when there is a partial lack of common distributional support in  $x_A$ .

While Zheng and Little (2003) suggest using a penalized spline structure to non-parametrically link  $\hat{\pi}_i^A$  to the response surface, Rafei et al. (2022) proposed to use a partially linear Gaussian process (GP) regression as the PM. By choosing a stationary isotropic covariance function, we showed that GP behaves as an optimal kernel matching technique based on  $\hat{\pi}_i^A$ . The interest in this study is to use GP for matching the units of  $S_A^{(b)}$  to those of  $U^{(b,l)}$  for any given b = 1, 2, ..., B; l = 1, 2, ..., L. Therefore, we update the PM given in Eq. 2.16 as below:

$$g\left(E\left[y_{ij}|x_{ij}, d_{ij}, w_{ij}^R, \widehat{\pi}_{ij}; \theta\right]\right) = \theta_0 + x_{ij}^T \theta_1 + d_{ij}^T \theta_2 + \log(\pi_{ij}^R)\theta_3 + f\left(\log(\widehat{\pi}_{ij}^A)\right)$$
(2.17)

where  $f(x) \sim GP(\mu(x), K(x, x'))$  is a GP with mean  $\mu(x)$  and covariance matrix K(x, x'). As illustrated, we propose to use a log transformation of  $\hat{\pi}_i^A$  as the GP inputs. This is mainly because  $\log(\hat{\pi}_i^A)$  tends to be normally distributed as  $\log(\hat{\pi}_i^A) = x_i^T \hat{\beta}^*$  according to Step II. Note that to minimize the computational burden, we dropped the random intercept term associated with the cluster effects in Eq. 2.13, as the mean of random effects is expected to be zero, and we already accounted for the cluster-level variance through the Rao-Wu's bootstrapping method.

Rafei et al. (2022) dealt with a fully Bayesian framework where we aimed to jointly estimate  $(\pi_i^A, y_i)$ . The proposed GP was characterized by a two-parameter Mátern covariance function added to an inharmonious polunomial kernel. It is well-understood that the joint posterior distribution of the length-scale and marginal standard error parameters are weakly identifiable while their ratio is well-identifiable (Zhang, 2004). Hence, such a method may not be computationally efficient in this study, as the PM in Eq. 2.17 has to be fitted *BL* we employ a frequentist variant of GP with a single smoothing parameter where model parameters are estimated through a maximum a posteriori (MAP) technique. Suppose  $K(x, x_i) =$   $k(||x, x_i||)$  is a non-negative function with ||.|| being the Euclidean norm, such that k(0) = 1 and  $k(d) \to 0$  monotically as  $d \to \infty$ . Now, we define f as

$$f(log(\widehat{\pi}^A)) = log(\widehat{\pi}^A)\theta_4 + \sum_{i=1}^{n_A^{(b)}} v_i K\left(log(\widehat{\pi}^A), log(\widehat{\pi}^A_i)\right)$$
(2.18)

where  $v \sim N(0, (\lambda K)^{-1})$  with  $\lambda$  being a smoothing parameter. Note that the covariance matrix of f is  $K/\lambda$  (?). The MAP estimate for  $\theta$  is then given by

$$\widehat{\theta} = argmin_{\theta}||y - x^{T}\theta_{1} - log(w^{R})\theta_{2} - d^{T}\theta_{3} - \theta_{4}log(\widehat{\pi}^{A}) - Kv||^{2}/\sigma^{2} + \lambda v^{T}Kv$$
(2.19)

For the structure of K, we use a simplified version of Matérn covariance function as below:

$$K(x, x_i) = \left(1 + \frac{||x - x_i||}{\rho}\right) exp\left\{-\frac{||x - x_i||}{\rho}\right\}$$
(2.20)

where  $\rho = \max_{1 \le i,j \le n_A} ||x_i - x_j||$  as recommended by Kammann and Wand (2003). This ensures scale invariance and numerical stability in finding the parameter estimates. This partially linear GP regression model can be implemented in R using the package 'mgcv'.

Having  $y_i$  imputed for all  $i \in U^{(b,l)*}$ , we estimate the population mean of Y for a given (b,l) by

$$\widehat{\overline{y}}_{U}^{(b,l)} = \left(\sum_{i \in S_{A}^{(b)}} (y_{i} - \widehat{y}_{i}) + \widehat{y}_{U}^{(b,l)}\right) / N$$
(2.21)

where  $\hat{y}_U^{(b,l)} = \sum_{i \in U^{(b,l)}} \hat{y}_i$ . Upon calculating the expression 2.21 for all b = 1, 2, ..., B and l = 1, 2, ..., L, one can build a DR estimate for the population mean via Rubin's combining rule

$$\widehat{\overline{y}}_{U} = \frac{1}{BL} \sum_{b=1}^{B} \sum_{l=1}^{L} \widehat{\overline{y}}_{U}^{(b,l)}$$
(2.22)

and the corresponding variance estimator is given by

$$\widehat{Var}(\widehat{\bar{y}}_{U}) = \frac{B+1}{B(B-1)} \sum_{b=1}^{B} \left(\widehat{\bar{y}}_{U}^{(b)} - \widehat{\bar{y}}_{U}\right)^{2}$$
(2.23)

where  $\hat{y}_U^{(b)} = \sum_{l=1}^L \hat{y}_U^{(b,l)}/L$ . It is important noting that the within-imputation variance in Rubin's combining rule, i.e.  $Var(\hat{y}_U^{(b)})$ , is zero as we impute the PS and outcome for the entire population (Zhou et al., 2016c). One can construct the 95% CI for the population mean of Y based on a t-student reference distribution with degrees of freedom equal to  $min\{m-H, B-1\}$  (Zhou et al., 2016b).

Note that the population size N was assumed to be known. Most often, this population parameter is unknown to the analyst. Although an estimate of the population size can always be obtained from the public-use weights of the reference survey by  $\hat{N} = \sum_{h} \sum_{j} \sum_{i} w_{hji}^{R}$ , one has to account for the uncertainty of  $\hat{N}$  as well. For this purpose, we propose to replace N in Eq. 2.21 with  $N_R^{(b)} = \sum_{h} \sum_{j} \sum_{i} w_{hji}^{*}$ where  $w_{hji}^{*}$ 's denote the rescaled sampling weights in  $S_R^{(b)}$  (b = 1, 2, ..., B) based on Rao-Wu's method. Algorithm 1 summarizes the entire procedure of the proposed GPPP method.

Algorithm 1: Fully model-based inference for non-probability samples

 $\begin{array}{l} \textbf{Result: Adjusted point and variance estimates of finite population mean} \\ \textbf{Combine } S_A \text{ with } S_R; \\ \textbf{for } b \ from 1 \ to B \ \textbf{do} \\ \\ \hline \textbf{Draw } S_A^{(b)}, \text{ a regular bootstrap sample of size } n_A, \ from S_A; \\ \textbf{for } h \ from 1 \ to H \ \textbf{do} \\ \\ \\ \hline \textbf{Draw } m_h - 1 \ \text{bootstrap PSUs from } h^{\text{th}} \ \text{stratum of } S_R; \\ \textbf{end} \\ \hline \textbf{Construct } S_R^{(b)} \ \text{by combining bootstrapped PSUs across } H \ \text{strata;} \\ \textbf{Apply Rao and Wu (1988) rescaling method to sampling weights of } S_R^{(b)}; \\ \textbf{for } l \ from 1 \ to L \ \textbf{do} \\ \\ \hline \textbf{Generate } \widehat{U}^{(b,l)}, \ \text{a synthetic population of size } N, \ \text{based on } S_R^{(b)}; \\ \textbf{Attach } S_A^{(b)} \ \text{to } \widehat{U}^{(b,l)} \ \text{to construct } \widehat{U}^{(b,l)*}, \ \text{and define } \delta_i^{A*} = I(i \in S_A^{(b)}); \\ \textbf{Estimate } \widehat{\pi}_i^A \ \text{for } i \in \widehat{U}^{(b,l)*} \ \text{by modeling } p(\delta_i^{A*} = 1|x_i) \ \text{on } \widehat{U}^{(b,l)*}; \\ \textbf{Predict } \widehat{y}_i \ \text{for } i \in \widehat{U}^{(b,l)*} \ \text{by modeling } E[y_i | \widehat{\pi}_i^A, x_i, z_i] \ \text{on } S_A^{(b)}; \\ \textbf{Compute } \widehat{\overline{y}}_U^{(b)} = \left(\sum_{i \in S_A^{(b)}} (y_i - \widehat{y}_i) + \sum_{i \in U^{(b,l)}} \widehat{y}_i\right) / N; \\ \textbf{end} \\ \textbf{Compute } \widehat{\overline{y}}_U^{(b)} = \left(\sum_{l=1}^L \widehat{\overline{y}}_U^{(b,l)}\right) / L; \\ \textbf{end} \\ \textbf{Compute } \widehat{\overline{y}}_U^{(b)} = \sum_{b=1}^L \widehat{\overline{y}}_U^{(b,l)} / B \ \text{and } \widehat{Var}(\widehat{\overline{y}}_U) = (1 + B^{-1}) \sum_{b=1}^R \left(\widehat{\overline{y}}_U^{(b)} - \widehat{\overline{y}}_U\right)^2 / (B-1); \end{aligned}$ 

## 3. Simulation study

We design an extensive simulation study to gauge how the proposed GPPP method operates with respect repeated sampling properties, including unbiasedness and efficiency, and contrast the results with those based on the PSPP, LWP, AIPW and IPSW methods. While all the competing methods are recognized as DR except for the IPSW, the two latter constitute a design-based form unlike the rest that are fully model-based. For any given pair of bootstrap samples,  $S_A^{(b)}$  and  $S_R^{(b)}$  (b = 1, 2, ..., B) and a synthetic population  $\hat{U}^{(b,l)}$  (l = 1, 2, ...L) created based on  $S_R^{(b)}$ , an AIPW estimator for the population mean is given by

$$\widehat{y}_{U}^{(b,l)} = \frac{1}{N} \sum_{i \in S_{A}^{(b)}} \frac{(y_{i} - \widehat{y}_{i})}{\widehat{\pi}_{i}^{A}} + \frac{1}{N} \sum_{i \in U^{(b,l)}} \widehat{y}_{i}$$
(3.1)

where  $\hat{y}_i^{(b,l)}$  is the prediction of  $y_i$  for  $i \in \hat{U}^{(b,l)}$  based the PM presented in Eq. 2.16. Note that the main difference between Eq. 3.1 and Eq. 2.4 is in how  $\hat{\pi}_i^A$  is applied. In the earlier,  $\hat{\pi}_i^A$  is used to inversely weight the PM residuals, whereas the latter utilizes  $\hat{\pi}_i^A$  as a model covariate to predict  $y_i$  in PM.

Regarding the PSPP method, we use a P-spline smoothing technique proposed by Eilers et al. (1996) in the PM, which specifies a discrete penalty to the basis functions. This is equivalent to a situation where knot coefficients are treated as random effects (Ruppert et al., 2003). Here a cubic spline is employed under a ridge penalty with 10 evenly spaced knots. Setting B = 50 and L = 10, we then utilize Rubin's combining rules to obtain point and interval estimates of the population mean for all the competing methods. Our focus here is on a situation where  $S_R$  is complex in design, but units of  $S_A$  are selected independently with unequal probabilities of selection. Once  $S_A$  and  $S_R$  are drawn from U, we assume that  $\pi_i^A$  for  $i \in S_A$  and  $y_j$  for  $j \in S_R$  are unobserved, and the aim is to adjust for the selection bias in  $S_A$  based on the combined sample. The simulation is then iterated K = 1,008 times (which is a multiple of 36, the number of cores we employed for parallel computing), where the bias-adjusted point estimates, SE, and associated 95% confidence intervals (CI) are estimated for  $\overline{y}_U$  in each iteration.

#### 3.1. Simulation design

Initially, we generate a hypothetical population of size N = 50,000 with H = 50 strata and  $M_h = 20$  clusters, each with a fixed size of  $N_{hj} = 50$  (j = 1, 2, ..., 20), within the stratum h (h = 1, 2, ..., 50). Then, two random variables,  $v_1$  and  $v_2$ , are created at the cluster and unit levels, respectively with the following distributions

$$V_1 \sim EXP(\mu = 1),$$
  $V_2 \sim Unif(1,5)$  (3.2)

which are supposed to be the MOS associated with first- and second-order probabilities of inclusion in  $S_R$ . The first-order selection probabilities are produced by  $\pi_{j|h}^R \propto c + v_1$  where c is defined such that  $max\{\pi_{j|h}^R\}/min\{\pi_{j|h}^R\} = 30$ . we then normalize  $\pi_{j|h}^R$  such that  $\sum_{h=1}^H \sum_{j=1}^{M_h} \pi_{j|h}^R = 1$ . Also, for each  $i^{\text{th}}$  unit of the  $j^{\text{th}}$  cluster in the stratum h, the second-order probabilities of inclusion are obtained by  $\pi_{i|hj}^R = v_{2hji}/\sum_{j=1}^{M_h} \sum_{i=1}^{N_{hj}} v_{2hji}$ . As a result, the joint selection probabilities and sampling weights in  $S_R$  are then calculated by  $\pi_{hji}^R = \pi_{j|h}^R \pi_{i|hj}^R$  and  $w_{hji}^R \propto 1/\pi_{hji}^R$ . Finally, we normalize the weights such that  $\sum_{h=1}^H \sum_{j=1}^{M_h} \sum_{i=1}^{N_{hj}} w_{hji}^R = N$ .

Associated with the selection mechanism of  $S_A$ , we also create a random variable X, which is assumed to be correlated with  $w_{hji}^R$  through  $x_{hji} = -7 + \log(w_{hji}^R) + \rho \epsilon_{hji}$  where  $\epsilon_{hji} \sim N(0,1)$  and  $\rho$  is determined such that  $cor(x_{hji}, log(w_{hji}^R)) = 0.5$ . A *logistic* function is then employed to produce the selection probabilities related to  $S_A$  by

$$\pi^{A}(x_{hji}) = p(\delta^{A}_{hji} = 1 | x_{hji}) = \frac{exp\{\gamma_0 + \gamma_1 x_i\}}{1 + exp\{\gamma_0 + \gamma_1 x_i\}}$$
(3.3)

We pick two values for  $\gamma_1$ , 0.3 and 0.6, where the latter produces extremely low selection probabilities. The goal here is to assess how the competing adjustment methods perform in the presence of influential weights. For a given  $\gamma_1$ , the value of  $\gamma_0$  is then found such that  $\sum_{h=1}^{H} \sum_{j=1}^{M_h} \sum_{i=1}^{N_{hj}} \pi_{hji}^A = n_A$ .

Using a non-linear mixed effect model, as the next step, we generate a continuous outcome variable denoted by  $y_{ij}^c$  and a binary outcome variable denoted by  $y_{ij}^b$  by

$$y_{kij}^{c}|x_{ij}, w_{ij}^{R}, d_{ij} \sim N(\mu = 1 + f_{k}(x_{ij}) + \log(w_{ij}^{R}) - 0.1d_{ij} + 0.2x_{ij}\log(w_{ij}^{R}) + u_{i}, \sigma^{2} = 4)$$
  
$$y_{kij}^{b}|x_{ij}, w_{ij}^{R}, d_{ij} \sim Ber\left(\frac{exp\{-7 + f_{k}(x_{ij}) + \log(w_{ij}^{R}) - 0.1d_{ij} + 0.2x_{ij}\log(w_{ij}^{R}) + u_{i}\}}{1 + exp\{-7 + f_{k}(x_{ij}) + \log(w_{ij}^{R}) - 0.1d_{ij} + 0.2x_{ij}\log(w_{ij}^{R}) + u_{i}\}}\right)$$
(3.4)

where  $d_{ij}$  is a categorical variable indicating the strata in U which takes values of 0, 1, ..., 49,  $u_i \sim N(0, \sigma_u^2)$ , and  $\sigma_u^2$  is determined such that the intraclass correlation (ICC) equals 0.2 (??). The function  $f_k(.)$  is assumed to take one of the following forms:

$$LIN: f_1(x) = x CUB: f_2(x) = (x/3)^3 (3.5)$$
  

$$EXP: f_3(x) = exp(x/2)/5 SIN: f_4(x) = 5sin(\pi x/2)$$

Figure 2 depicts the association between  $y_i$  and  $\pi_i^A$  and also between  $y_i$  and  $w_i^A = 1/\pi_i^A$  in the generated hypothetical population under each of these scenarios.

Associated with  $S_R$  and  $S_A$ , independent samples of size  $n_R = 1,000$  and  $n_A = 500$  were selected randomly from U. Regarding  $S_R$ , we consider a stratified two-stage clustering design with two PSU selected per stratum with  $m_h = 2$  and  $n_{hj} = 10$  (h = 1, 2, ..., 50) and (j = 1, 2). At each stage, without replacement random samples are selected with unequal inclusion probabilities from PSUs and SSUs. In addition, a Poisson sampling design is used to select units of  $S_A$  with unequal inclusion probabilities  $\pi_{hji}^A$  directly from U. Finally, to test the DR property of the proposed methods, we investigate different scenarios regarding whether models for QR and PM are correctly specified or not. To misspecify a model, we only include  $x_{ij}^2$  as a predictor in the model.



Fig 2: The proposed relationships between the outcome variable Y and  $log(\pi^A)$  in U for (a) LIN, (b) CUB, (c) EXP and (d) SIN scenarios, and between the outcome Y and sampling weights  $w^A$  for (e) LIN, (f) CUB, (g) EXP and (h) SIN scenarios.

To evaluate the repeated sampling properties of the competing method, relative bias (rBias), relative root mean square error (rMSE), the nominal coverage rate of 95% CIs (crCI), relative length of 95% CIs (rlCI) and SE ratio (rSE) are calculated as below:

$$rbias\left(\widehat{\overline{y}}_{U}\right) = 100 \times \frac{1}{K} \sum_{k=1}^{K} \left(\widehat{\overline{y}}_{U}^{(k)} - \overline{\overline{y}}_{U}\right) / \overline{\overline{y}}_{U}$$
(3.6)

$$rMSE\left(\widehat{\overline{y}}_{U}\right) = 100 \times \sqrt{\frac{1}{K} \sum_{k=1}^{K} \left(\widehat{\overline{y}}_{U}^{(k)} - \overline{y}_{U}\right)^{2} / \overline{y}_{U}}$$
(3.7)

$$crCI\left(\widehat{\overline{y}}_{U}\right) = 100 \times \frac{1}{K} \sum_{k=1}^{K} I\left(\left|\widehat{\overline{y}}_{U}^{(k)} - \overline{\overline{y}}_{U}\right| < z_{0.975} \sqrt{var\left(\widehat{\overline{y}}_{U}^{(k)}\right)}\right)$$
(3.8)

$$rlCI\left(\widehat{\overline{y}}_{U}\right) = 100 \times \frac{2}{K} \sum_{k=1}^{K} z_{0.975} \sqrt{var\left(\widehat{\overline{y}}_{U}^{(k)}\right)} / \overline{y}_{U}$$
(3.9)

$$rSE\left(\widehat{\overline{y}}_{U}\right) = \frac{1}{K} \sum_{k=1}^{K} \sqrt{var(\widehat{\overline{y}}_{U}^{(k)})} / \sqrt{\frac{1}{K-1} \sum_{k=1}^{K} \left(\widehat{\overline{y}}_{U}^{(k)} - \overline{\overline{y}}_{U}\right)^{2}}$$
(3.10)

where  $\hat{\overline{y}}_{U}^{(k)}$  denotes the adjusted sample mean from iteration k,  $\overline{\overline{y}}_{U} = \sum_{k=1}^{K} \hat{\overline{y}}_{U}^{(k)} / K$ ,  $\overline{y}_{U}$  is the finite population true mean, and var(.) represents the variance estimate of the adjusted mean based on the sample.

## 3.2. Simulation results

Figure 3 visualizes the magnitude of bias and efficiency of the competing methods for various types of response surface structures and across different scenarios of model specification when  $\gamma_1 = 0.3$ . Note that the error bars point out the expected rlCI over the simulation iterations with narrower bands indicating

higher efficiency. As illustrated, naive estimates are severely biased while adjustment based on the true sampling weight (if known to the analyst) removes the bias entirely. Generally, we observe that all the competing methods produce unbiased estimates as long as the working model is valid for either QR or PM. This finding holds across all the four outcome variables when there is no evidence of influential pseudo-weights. When both working models are wrong, not surprisingly, estimates based on all the applied methods are biased.



Fig 3: Comparing the performance of the adjusted estimators for the *continuous* outcome variable across different model-specification scenarios under (a) LIN, (b) CUB, (c) EXP, and (d) SIN when  $\gamma_1 = 0.3$ . The error bars point out the relative length of 95% CIs. UW: unweighted; FW: Fully weighted; IPSW: Inverse Propensity Score Weighting; GPPP: Gaussian Processes of Propensity Prediction; PSPP: Penalized Spline of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting

When it comes to efficiency, the adjustment methods perform almost the same, except for the design-based estimators, i.e. IPSW and AIPW, that consistently show lower efficiency across the four plots in situations where it is only the QR model that is correctly specified. In addition, when extreme non-linearity is present, such as in Figure 3d, the LWP estimator is slightly less efficient than those based on the GPPP and PSPP methods when the PM is invalid. While non-parametric terms in GPPP and PSPP improves the efficiency under a wrongly specified PM, it turns out that no gain is reached with respect to efficiency when the PM is correctly specified regardless of how the QR model is specified. Analogous results are depicted in Figure 4 but for  $\gamma_1 = 0.6$ , which leads to generating influential pseudo-weights. In this circumstance, slight residual bias is evident for the PSPP, LWP, and AIPW methods where the underlying model is valid just for the QR approach. Furthermore, the LWP method loses more efficiency when outlying pseudo-weights are present, especially for higher degrees of non-linearity in the outcome variable.

We have summarized the simulation results for rSE and crCI with  $\gamma_0 = 0.3$  in Figure 5. At first glance, it can be seen that the values of rSE are very close to one for all the competing methods and outcome variables, indicating accurate estimation of the variance based on the proposed bootstrap-Polýa posterior



Fig 4: Comparing the performance of the adjusted estimators for the *continuous* outcome variable across different model-specification scenarios under (a) LIN, (b) CUB, (c) EXP, and (d) SIN when  $\gamma_1 = 0.6$ . The error bars point out the relative length of 95% CIs. UW: unweighted; FW: Fully weighted; IPSW: Inverse Propensity Score Weighting; GPPP: Gaussian Processes of Propensity Prediction; PSPP: Penalized Spline of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting

method as long as at least one of the models for QR or PM holds. For higher degrees of non-linearity in the outcome, i.e. in the plot (h), there is evidence of slight overestimation in the variance. In addition, plots (a) to (h) reflect that the percentages of crCI are close to the nominal value, 95%. An exception includes plot (b) CUB, where 95% CIs slightly undercover the true population mean when the PM is valid. In Figure 6, we show how ignoring the effects of strata and clusters in  $S_R$  affects the values of rSE and crCI. To this end, we skip the step of bootstrapping from  $S_R$  and directly generate synthetic populations using  $w_i^R$ 's. The new values of rSE indicate that the variance is substantially underestimated if the complex design of  $S_R$  is ignored. In addition, the estimated 95% CIs tend to significantly undercover the true population mean. Note that ignoring the stratification and clustering effects does not affect the magnitude of bias of point estimates.

In Figure 7, we re-evaluate the simulation results for rSE and crCI when outlying pseudo-weights are present, i.e. under  $\gamma_1 = 0.6$ . According to the values of rSE, the overestimation of variance tends to be higher under this circumstance, especially for the GPPP and PSPP methods. While we observe no consistent outperformance across the competing methods, it seems the worst situation is associated with non-linear PM and where the working PM is invalid. The AIPW method exceptionally underestimates the variance for the outcomes associated with LIN, CUB, and EXP. Furthermore, it turns out that the presence of influential pseudo-weights leads to higher variability in the percentages of crCI, though deviations from the nominal 95% seem to be negligible.

In Appendix 8, we replicate the simulation results for the binary outcome variable with  $\pi_i^A$ 's generated based on  $\gamma_1 = 0.3$  and  $\gamma_1 = 0.6$ , respectively. Regarding the bias magnitude and efficiency, the plots reveal



Fig 5: Comparing the 95% CI coverage rates (crCI) of the adjusted means for the *continuous* outcome variable under (a) LIN, (b) CUB, (c) EXP, and (d) SIN, and SE ratios (rSE) under (e) LIN, (f) CUB, (g) EXP, and (h) SIN, across different DR methods under different model specification scenarios when  $\gamma_1 = 0.3$ . UW: unweighted; FW: Fully weighted; PAPP: Propensity Adjusted Probability Prediction; GPPP: Gaussian Processes of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting



Fig 6: Comparing the 95% CI coverage rates (crCI) of the DR adjusted means for the *continuous* outcome variable under (a) LIN, (b) CUB, (c) EXP, and (d) SIN, and SE ratios (rSE) under (e) LIN, (f) CUB, (g) EXP, and (h) SIN, across different DR methods under different model specification scenarios when  $\gamma_1 = 0.3$  and the complexity in the design of  $S_R$  is ignored. UW: unweighted; FW: Fully weighted; PAPP: Propensity Adjusted Probability Prediction; GPPP: Gaussian Processes of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting

almost no difference between the binary and continuous outcome variables. However, unlike the continuous outcome variable where improvements in efficiency were observed when the PM is incorrectly specified,



Fig 7: Comparing the 95% CI coverage rates (crCI) of the DR adjusted means for the *continuous* outcome variable under (a) LIN, (b) CUB, (c) EXP, and (d) SIN, and SE ratios (rSE) under (e) LIN, (f) CUB, (g) EXP, and (h) SIN, across different DR methods under different model specification scenarios when  $\gamma_1 = 0.6$ . UW: unweighted; FW: Fully weighted; PAPP: Propensity Adjusted Probability Prediction; GPPP: Gaussian Processes of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting

this improvement no longer holds for the binary outcome. In addition, we realized that the estimated 95% CIs consistently undercover the population true mean across all the competing methods. In addition to the simulation results for the binary outcome, Appendix 8 involves the numerical tables associated with all the plots in the current section as well as extensions to simulations with  $(n_A, n_R) = (1,000,500)$  and  $(n_A, n_R) = (500,500)$ .

#### 4. Application

Using Crash Injury Research Engineering Network (CIREN) data from 2005 to 2015, we aim at estimating crash injury rates in various body regions, including head, abdomen, thorax and spine. To correct for the selection bias in CIREN using different methods, we consider pooled data of Crashworthiness Data System (CDS) over the same time period as the reference survey. A brief description of these two datasets is given in the following subsections.

#### 4.1. Crashworthiness Data System

The CDS has served as one of the key components of the National Automotive Sampling System (NASS) for about three decades, investigating severe traffic accidents involving passenger cars in the U.S. The CDS entails a series of stratified three-stage and multi-phase probability samples conducted annually with the primary aim to learn about the crashworthiness of vehicles and its consequences to the affected occupants. A PSU consists of geographical locations with a minimum population size of 50,000 that are grouped into 12 strata. Two PSUs are selected randomly from each stratum in the first stage. A secondary sampling unit (SSU) involves police jurisdictions processing reports of motor vehicle crashes within the selected PSUs. Finally, crashes are sampled randomly on a weekly basis from the police jurisdictions selected in the second stage. As the chief inclusion criteria, a police-reported crash must have caused at least one towed passenger vehicle to be eligible for selection in the final stage of sampling. (Note that the selected PSUs and SSUs are fixed across the years of study (Fleming, 2015).) Thorough information about

vehicle crash profiles, restraint system performance, and injury mechanisms are then collected from the selected crashes via face-to-face interviews, medical records, vehicle inspections, and scene investigations. The 2015 round of the survey involves 2,633 randomly selected police-reported accidents with 5,479 occupants investigated (Zhang et al., 2019).

#### 4.2. Crash Injury Research and Engineering Network

Launched by the National Highway Traffic Safety Administration (NHTSA), CIREN collects prospective clinical and biomechanical data about motor vehicle crashes with severe injury across the U.S. More specifically, CIREN investigates crash causality and consequences with the aim to tailor enhanced strategies for preventing, treating, and rehabilitating injured occupants in traffic accidents. Subjects in CIREN are selected from severely injured occupants due to motor vehicle crashes who are admitted to one of the CIREN-assigned medical centers through a set of specific inclusion criteria. Medical data are gathered by expert trauma/emergency physicians in level one trauma centers, whereas biomechanical data are gathered by trained engineers in academic engineering laboratories. While CIREN collects more detailed information than CDS, the sampling mechanism in CIREN is non-probabilistic. The trauma centers are located in urban areas and there is a tendency to recruit patients with more severe injury and with newer model-year vehicles (Elliott et al., 2010; Flannagan and Rupp, 2009). To adjust for the potential selection bias in CIREN, we analyze the data gathered from 2005 through 2015, which consist of 2,104 subjects.

## 4.3. Data harmonization and auxiliary variables

Before performing any statistical analysis, extensive effort was invested to harmonize the data of the two studies at the occupant level such that both datasets represent potentially the same target population. This involves occupants of any non-derivable motor vehicle crash between 2005 and 2015 in the U.S. who have either been fatally injured or hospitalized. As the first step, we excluded all the individuals participating in CDS who failed to meet CIREN-specific inclusion criteria. Details of these criteria have been reported in the supplemental appendix of Elliott et al. (2010). We also restricted the samples to those who were seated in the first two rows of the crashed vehicle with known crash direction and restraint status. In addition, for both datasets, individuals who survived without an urgent need for hospitalization or whose treatment status was unknown were filtered out. This finally left us a sample of size  $n_R = 7,721$  for CDS and a sample of size  $n_A = 1,738$  for CIREN.

We identified and harmonized 20 common auxiliary variables between the two datasets, mainly describing occupants' demographic, position, and behavior at the incident, vehicle characteristics, and the intensity of crash and resulting injuries. Two variables associated with race/ethnicity were set aside due to a high rate of item-level missingness in CDS. Among the others, we chose the variables whose distribution was significantly imbalanced in CIREN compared to weighted CDS. To this end, we performed stepwise variable selection based on Bayesian Information Criterion (BIC) in a weighted logistic regression whose outcome was the indicator of being in CIREN given the combined sample. Note that the sampling weights were set to 1 for the units of CIREN (Wang et al., 2020). We specifically found eight variables that are significant in the model. Figure 8 depicts these variables, comparing their sample distribution in CIREN with the target population using weighted CDS data.

The bias adjustment procedure in the next step of analysis is restricted to this set of auxiliary variables for modeling the response indicator in CIREN. To be able to evaluate the performance of the adjustment methods on the actual data, in addition to the common auxiliary variables, we recognize a set of injury-level variables that were available in both datasets as the outcome variables. This entailed the indicators of being diagnosed with a 3+ level injury (according to AIS codes) to the following body regions: head, thorax, abdomen, and spine. One would expect that unweighted estimates of the prevalence of such injuries using CIREN-only data are overestimated, as CIREN tends to recruit severely injured occupants. To construct the PM, we conduct the BIC-based stepwise procedure of variable selection for each outcome variable separately. Once the bias-adjusted estimates of the prevalence are obtained for these outcome variables, we compare them with the corresponding weighted estimates using CDS as the benchmark.



Fig 8: Comparing the distributions of common auxiliary variables in CIREN with weighted CDS

## 4.4. Results

As the first step of the analysis, we visually compared the empirical density of the estimated propensity scores (PS) in CIREN data with that of the target population to see if there is any lack of common support in the distribution of estimated PS in CIREN. The estimates of PS were obtained through a weighted logistic regression fitted on the combined sample with the weights equal to 1 for the CIREN units. This comparison has been displayed in Figure 9a. As illustrated there is evidence of a lack of common support in the left tail of the distribution. This may suggest that further filtering should be done on the CDS data such that the undercoverage of the common representing population by CIREN is minimized. We then estimated the pseudo-weights associated with the CIREN units based on both the proposed PAPP method and the PMLE method proposed by Wang et al. (2020).

Figure 9b compares the distribution of the estimated pseudo-weights between the two methods through a boxplot. As illustrated, there is evidence of influential pseudo-weights. However, we believe that the proposed GPPP method is not widely affected by this issue. This mainly originates from the specific structure of the GP covariance, which is a function of  $||\hat{\pi}_i^A - \hat{\pi}_j^A||$ , not  $\hat{\pi}_i^A \hat{\pi}_j^A$  as in the LWP method. In addition, we showed in Rafei et al. (2022) how GP behaves like a matching technique, and it is well-understood that PS-based matching reduces the adverse effects of influential pseudo-weights. We observed a Pearson correlation of  $\rho = 0.69$  in the estimates of pseudo-weights between the two PAPP and PMLE methods. Now, we compare the pseudo-weighted distribution of auxiliary variables in CIREN with their weighted distribution in CDS based on the PAPP method to see how well it mitigates the previously observed after assigning the pseudo-weights to the CIREN units. If the association of these variables is strong enough with the outcome variables of interest, one would expect that pseudo-weighted estimates would improve the potential bias in those variables to a significant extent as well.



**Fig 9:** Comparing the empirical density of (a) estimated propensity scores between CIREN and CDS and (b) estimated pseudo-weights in CIREN across the applied quasi-randomization methods



Fig 10: Comparing the distribution of common auxiliary variables in pseudo-weighted CIREN based on the PAPP method with weighted CDS

For the PM, as was concluded from the joint likelihood factorization in Eq. 2.1, one has to use both X and Z as the predictors in the model. However, one major obstacle is that Z, the design features of CDS including strata indicators and sampling weights are not known or calculable for units of CIREN. Even if the CDS strata was identifiable for units of CIREN, it is most likely that for some strata, there is no observation in CIREN. This may imply a lack of coverage by the CIREN sample, and if the strata indicators are associated with the outcome variable of interest, final estimates may be biased. However, there is still a hope that this association is at least partially explained by the observed set of X. Since the chosen outcome variables in this study are available for both samples, one can check the role of Z in addition to X in the goodness-of-fit of the PM by fitting it on the CDS data.

Table 11 compares the model's goodness-of-fit before and after including the design features Z across different outcome variables. Note that the models under X + Z involve two GP terms, one linking  $\pi_i^R$ and the other linking  $\hat{\pi}_i^A$  to the response surface non-linearly. As the values of adjusted- $R^2$  and explained deviance reflect, the improvement in the PM's goodness-of-fit after including the Z variables, i.e.  $GP(\pi_i^R)$ and  $d_i$ , is small for all the outcome variables. Figure 11 visualizes the predictive power of Z in the PM associated with different outcome variables using the Area Under the Curve (AUC) of receiver operating characteristic (ROC) curve. The values of AUC suggest that Z does not improve the predictive power of X in the PM substantially.

TABLE 1 Assessing goodness-of-fit of the prediction model in CDS when the design features (Z) are excluded

	Adjus	$ted-R^2$	Deviar	nce (%)
Outcome	Х	X+Z	Х	X+Z
CDS				
Head	0.355	0.360	31.00	31.90
Abdomen	0.164	0.172	21.10	22.10
Thorax	0.334	0.336	29.20	29.70
Spine	0.050	0.056	12.70	14.70
CIREN				
Head	0.187		20.40	
Abdomen	0.180		21.10	
Thorax	0.261		24.30	
Spine	0.137		17.10	



Fig 11: Assessing the predictive power of the design features of CDS (Z) in the prediction models using the area under ROC

As the last step, we apply the adjustment methods on the outcome variables of interest and compare the bias-adjusted estimates based on the GPPP approach with those based on the AIPW method and the weighted CDS. The results are shown in Figure 12. The red dashed line and the surrounding shaded area illustrate the true population estimates using the CDS weighted data and associated 95%CI. As expected, naive percentages based on the CDS-only and CIREN-only data are overestimated substantially across the four outcome variable. An exception includes the outcome variable associated with the spinal injury where the CDS-only unweighted estimate is unbiased. At first glance, one can infer that all DR bias

adjustment methods shift the prevalence towards the true population estimates, except for the PAPP method. It turns out that estimates based on the IPSW method are biased, and the magnitude of bias is even larger than that in naive estimates for injuries in the head and thorax. The minimum amount of residual bias seems to be associated with the GPPP and PSPP methods.

Regarding the efficiency, the widest 95% CIs are associated with the IPSW and LWP methods, and the narrowest 95% CI is associated with the PSPP method. This is contrary to the simulation results where we observed better efficiency in GPPP estimator than the PSPP estimator. A notable discrepancy between the application and simulation studies is that, unlike the simulation, all the auxiliary variables used for predicting both PS and outcome variable in the application were categorical. However, in terms of bias, we found no consistent differences between the GPPP and PSPP methods. Further numerical comparisons across different outcome variables and different DR methods by levels of auxiliary variables are provided in tables 14-17 in Appendix 8.2.



Fig 12: Comparing the performance of adjustment methods for estimating the percentage of occupants with severe injury in (a) head, (b) abdomen, (c) thorax, and (d) spine based on CIREN/CDS. The dashed line and surrounding shadowed area represent weighted estimates and 95% CIs in CDS, respectively. UW: unweighted; FW: Fully weighted; IPSW: Inverse Propensity Score Weighting; GPPP: Gaussian Processes of Propensity Prediction; PSPP: Penalized Spline of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting.

## 5. Discussion

This article developed a fully model-based approach for finite population inference using a non-probability sample. Our focus was on a situation where the available reference sample is complex in design. To address this, we particularly used a bootstrap-Polýa posterior technique proposed by Zhou et al. (2016c) to generate synthetic populations based on the reference survey. Generally, the simulation results demonstrate that the proposed GPPP estimator is DR, producing the least degree of residual bias with

improved efficiency among the competing methods. The superiority of the GPPP was more evident when influential pseudo-weights are present. On the contrary, the empirical study showed that adjustments based on PSPP lead to more efficient estimates than those based on the GPPP method. This specific contradiction may have arisen from the fact that all the auxiliary variables used in the application were categorical.

Unlike the proposed estimator in prior literature, the one introduced by this study involved no design-based term, because the outcome variable was predicted for all non-sampled units. This along with the use of a flexible function of estimated propensity based on GP not only yielded double robustness in the consistency of the ultimate estimator but reduced the sensitivity to the presence of influential pseudo-weights. More importantly, the new proposed method eliminated the need for modeling the sampling weights of the reference survey as the outcome in the process of estimating the PS, which can be challenging to the analyst. In fact, one can get rid of the complexity in the design of the reference survey, especially the unequal sampling weights, by generating synthetic populations as the first step. This is advantageous for Bayesian inference and for the use of alternative algorithmic predictive tools such as BART. Liu et al. (2021); Mercer (2018) uses BART for prediction modeling through a similar model-based idea with the estimated PS used as a predictor in the model. In his comparison with the PSPP method, the author shows by simulation that estimates based on the PSPP method are more efficient than those based on BART.

One can easily expand the proposed method to a fully Bayesian setting where the outcome and PS are jointly estimated. However, unlike the proposed method in Rafei et al. (2022), developing a unified Bayesian framework may not be possible under a fully model-based approach given the available Bayesian platforms. This is mainly because one has to generate the synthetic population as the beginning step, while software like winBUGS and Stan does not allow for using the simulated posterior predictive distribution as the input for the following steps. Therefore, one has to use a two-step approach where initially synthetic populations are created repeatedly, and then for any given synthetic population, one can fit the models for QR and PM jointly under a Bayesian setting. Therefore, simulating the posterior predictive distribution of the population's unknown quantity will be prohibitive as one would have to rely on Rubin's combining rules for the variance estimation. In addition, such a method can turn out to be very intensive computationally, as for any given synthetic population, one has to fit the QR model on the entire population and predict the outcome for all units of the population repeatedly.

Another major challenge with the PM, which was argued in the results section of the application, is the fact that design features of the reference sample are often unobserved for units of the non-probability sample. In the application part, because we had picked outcome variables that were available in both samples, we could check the improvements in the PM's goodness-of-fit using the reference survey where both X and Z are available. The good news is that our proposed method is still DR. Even if the PM lacks the reference survey's design features as predictors, one can still rely on the QR model for producing unbiased estimates. As a DR method, if the QR method is correctly specified, one would expect that the final estimates are unbiased.

## 6. Acknowledgement

The present study was part of the doctoral research of the first author of this article at the Michigan Program in Survey and Data Science. Therefore, we would like to thank the respected members of the dissertation committee, Professors Brady T. West, Roderick J. Little, and Philip S. Boonstra at the University of Michigan, who have continuously supported this research with their excellent comments and critical feedback. Our gratitude also goes to Professors Katharine Abraham, Stanley Presser, and Joseph Sedarski at the University of Maryland who have significantly contributed to the development of the main idea of this paper with their valuable comments and feedback over the doctoral seminar course. Last but not least, the authors would like to thank all the researchers and staff who have been involved in collecting the data of CIREN and CDS from 2005 to 2015.

## 7. Conflict of Interest

The authors declare that there was no conflict of interest in the current research.

#### References

- Bang, H. and J. M. Robins (2005). Doubly robust estimation in missing data and causal inference models. Biometrics 61(4), 962–973.
- Basu, D. (1971). An essay on the logical foundations of survey sampling, pp. 203–204. Springer International Publishing.
- Berger, J. O. and R. L. Wolpert (1988). The likelihood principle: A review, generalizations, and statistical implications. Lecture Notes—Monograph Series 6.
- Chen, C.-L., R. Subramanian, F. Zhang, and E. Y. Noh (2015). Nhtsa's data modernization project. In Proceedings of the 2015 Federal Committee on Statistical Methodology (FCSM) Research Conference, Washington, DC, USA, pp. 1–3.
- Chen, Q., M. Elliot, D. Haziza, Y. Yang, M. Gosh, R. Little, J. Sedransk, and M. Thompson (2017). Weights and estimation of a survey population mean: A review. *Statistical Science* 32, 227–248.
- Chen, Y., P. Li, and C. Wu (2019). Doubly robust inference with nonprobability survey samples. *Journal* of the American Statistical Association, 1–11.
- Cochran, W. G. (1977). Sampling Techniques: 3d Ed. Wiley.
- Cohen, M. P. (1997). The bayesian bootstrap and multiple imputation for unequal probability sample designs. In Proceedings of the Survey Research Methods Section, American Statistical Association, pp. 635–638.
- Dong, Q., M. R. Elliott, and T. E. Raghunathan (2014). A nonparametric method to generate synthetic populations to adjust for complex sampling design features. Survey methodology 40(1), 29.
- Efron, B. (1992). Bootstrap methods: another look at the jackknife. In *Breakthroughs in statistics*, pp. 569–593. Springer.
- Eilers, P. H., B. D. Marx, et al. (1996). Flexible smoothing with b-splines and penalties. Statistical science 11(2), 89–121.
- Elliot, M. R. (2013). Combining data from probability and non-probability samples using pseudo-weights. Survey Practice 2(6).
- Elliott, M. R., A. Resler, C. A. Flannagan, and J. D. Rupp (2010). Appropriate analysis of ciren data: using nass-cds to reduce bias in estimation of injury risk factors in passenger vehicle crashes. Accident Analysis & Prevention 42(2), 530–539.
- Elliott, M. R. and R. Valliant (2017). Inference for nonprobability samples. *Statistical science* 32(2), 249–264.
- Flannagan, C. A. and J. D. Rupp (2009). Comparison of distributions of key predictor variables in circn and nass-cds cases meeting circn inclusion criteria. *Traffic injury prevention* 10(5), 451–457.
- Fleming, S. (2015). Auto safety: Status of nhtsa's redesign of its crashworthiness data system. Technical report.
- Groves, R. M. (2011). Three eras of survey research. Public Opinion Quarterly 75(5), 861–871.
- Gunawan, D., A. Panagiotelis, W. Griffiths, and D. Chotikapanich (2020). Bayesian weighted inference from surveys. Australian & New Zealand Journal of Statistics 62(1), 71–94.
- Johnson, T. P. and T. W. Smith (2017). Big data and survey research: Supplement or substitute? In Seeing Cities Through Big Data, pp. 113–125. Springer.
- Kammann, E. and M. P. Wand (2003). Geoadditive models. Journal of the Royal Statistical Society: Series C (Applied Statistics) 52(1), 1–18.
- Kim, J.-K. and S.-M. Tam (2021). Data integration by combining big data and survey sample data for finite population inference. *International Statistical Review* 89(2), 382–401.
- Kish, L. (1965). Survey sampling.
- Little, R. J. (2004). To model or not to model? competing modes of inference for finite population sampling. *Journal of the American Statistical Association* 99(466), 546–556.

- Little, R. J. and H. Zheng (2007). The bayesian approach to the analysis of finite population surveys. Bayesian Statistics  $\mathcal{S}(1)$ , 1–20.
- Liu, Y., A. Gelman, and Q. Chen (2021). Inference from non-random samples using bayesian machine learning. arXiv preprint arXiv:2104.05192.
- McCarthy, P. J. and C. B. Snowden (1985). The bootstrap and finite population sampling.
- Meng, X.-L. et al. (2018). Statistical paradises and paradoxes in big data (i): Law of large populations, big data paradox, and the 2016 us presidential election. *The Annals of Applied Statistics* 12(2), 685–726.
- Mercer, A. W. (2018). Selection Bias in Nonprobability Surveys: A Causal Inference Approach. Ph. D. thesis.
- Miller, P. V. (2017). Is there a future for surveys? Public Opinion Quarterly 81(S1), 205–212.
- Pinheiro, J. C. and D. M. Bates (1995). Approximations to the log-likelihood function in the nonlinear mixed-effects model. Journal of computational and Graphical Statistics 4(1), 12–35.
- Rafei, A., M. R. Elliott, and C. A. Flannagan (2022). Robust and efficient bayesian inference for non-probability samples. arXiv preprint arXiv:2203.14355.
- Rafei, A., C. A. C. Flannagan, and M. R. Elliott (2020). Big data for finite population inference: Applying quasi-random approaches to naturalistic driving data using bayesian additive regression trees. *Journal* of Survey Statistics and Methodology In press.
- Rafei, A., C. A. C. Flannagan, B. T. West, and M. R. Elliott (2021). Robust bayesian inference for big data: Combining sensor-based records with traditional survey data.
- Rao, J. (2021). On making valid inferences by integrating data from surveys and other sources. Sankhya B 83(1), 242–272.
- Rao, J. N. and C. Wu (1988). Resampling inference with complex survey data. *Journal of the american* statistical association 83(401), 231–241.
- Robins, J. M., A. Rotnitzky, and L. P. Zhao (1994). Estimation of regression coefficients when some regressors are not always observed. *Journal of the American statistical Association 89*(427), 846–866. Rubin, D. B. (1976). Inference and missing data. *Biometrika 63*(3), 581–592.
- Ruppert, D., M. P. Wand, and R. J. Carroll (2003). *Semiparametric regression*. Number 12. Cambridge university press.
- Savitsky, T. D., D. Toth, et al. (2016). Bayesian estimation under informative sampling. *Electronic Journal of Statistics* 10(1), 1677–1708.
- Scharfstein, D. O., A. Rotnitzky, and J. M. Robins (1999). Adjusting for nonignorable drop-out using semiparametric nonresponse models. *Journal of the American Statistical Association* 94 (448), 1096–1120.
- Senthilkumar, S., B. K. Rai, A. A. Meshram, A. Gunasekaran, and S. Chandrakumarmangalam (2018). Big data in healthcare management: A review of literature. *American Journal of Theoretical and Applied Business* 4(2), 57–69.
- Shao, J. and Y. Chen (1998). Bootstrapping sample quantiles based on complex survey data under hot deck imputation. *Statistica Sinica*, 1071–1085.
- Smith, T. (1983). On the validity of inferences from non-random sample. Journal of the Royal Statistical Society. Series A (General), 394–403.
- Valliant, R., J. A. Dever, and F. Kreuter (2018). Nonprobability Sampling, pp. 565–603. Springer International Publishing.
- Wang, L., B. I. Graubard, H. A. Katki, and Y. Li (2020). Efficient and robust propensity-score-based methods for population inference using epidemiologic cohorts. arXiv preprint arXiv:2011.14850.
- Wang, L., R. Valliant, and Y. Li (2020). Adjusted logistic propensity weighting methods for population inference using nonprobability volunteer-based epidemiologic cohorts. arXiv preprint arXiv:2007.02476.
- Williams, M. R. and T. D. Savitsky (2021). Uncertainty estimation for pseudo-bayesian inference under complex sampling. *International Statistical Review* 89(1), 72–107.
- Wolter, K. (2007). Introduction to variance estimation. Springer Science & Business Media.
- Zangeneh, S. Z. (2012). Model-based methods for robust finite population inference in the presence of external information.
- Zhang, F., E. Y. Noh, R. Subramanian, and C.-L. Chen (2019). Crash investigation sampling system:

Sample design and weighting. Technical report.

- Zhang, G. and R. Little (2011). A comparative study of doubly robust estimators of the mean with missing data. *Journal of Statistical Computation and Simulation* 81(12), 2039–2058.
- Zhang, H. (2004). Inconsistent estimation and asymptotically equal interpolations in model-based geostatistics. *Journal of the American Statistical Association* 99(465), 250–261.
- Zheng, H. and R. J. Little (2003). Penalized spline model-based estimation of the finite populations total from probability-proportional-to-size samples. *JOURNAL OF OFFICIAL* STATISTICS-STOCKHOLM- 19(2), 99–118.
- Zhou, H. (2014). Accounting for Complex Sample Designs in Multiple Imputation Using the Finite Population Bayesian Bootstrap. Ph. D. thesis.
- Zhou, H., M. R. Elliott, and T. E. Raghunathan (2016a). Synthetic multiple-imputation procedure for multistage complex samples.
- Zhou, H., M. R. Elliott, and T. E. Raghunathan (2016b). Synthetic multiple-imputation procedure for multistage complex samples. *Journal of Official Statistics* 32(1), 231–256.
- Zhou, H., M. R. Elliott, and T. E. Raghunathan (2016c). A two-step semiparametric method to accommodate sampling weights in multiple imputation. *Biometrics* 72(1), 242–252.

## 8. Appendix

## 8.1. Extension of the simulation study

Figures 13-17 show the simulation results for the binary outcome variable with respect to rBias, lCI, rSE and crCI. In addition, tables 2-13 provides numerical results of the simulation extended to sample sizes of  $(n_A, n_R) = (1,000,500)$  and  $(n_A, n_R) = (500,500)$ .



Fig 13: Comparing the performance of the DR estimators under different model-specification scenarios for the *binary* outcome variable under (a) LIN, (b) CUB, (c) EXP, and (d) SIN when  $\gamma_1 = 0.3$ . The error bars point out the relative length of 95% CIs. UW: unweighted; FW: Fully weighted; GPPP: Gaussian Processes of Propensity Prediction; PSPP: Penalized Spline of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting



Fig 14: Comparing the performance of the DR estimators under different model-specification scenarios for the *binary* outcome variable under (a) LIN, (b) CUB, (c) EXP, and (d) SIN when  $\gamma_1 = 0.6$ . The error bars point out the relative length of 95% CIs. UW: unweighted; FW: Fully weighted; GPPP: Gaussian Processes of Propensity Prediction; PSPP: Penalized Spline of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting



Fig 15: Comparing the 95% CI coverage rates (crCI) of the DR adjusted means for the *continuous* outcome variable under (a) LIN, (b) CUB, (c) EXP, and (d) SIN, and SE ratios (rSE) under (e) LIN, (f) CUB, (g) EXP, and (h) SIN, across different DR methods under different model specification scenarios when  $\gamma_1 = 0.3$ . UW: unweighted; FW: Fully weighted; PAPP: Propensity Adjusted Probability Prediction; GPPP: Gaussian Processes of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting



Fig 16: Comparing the 95% CI coverage rates (crCI) of the DR adjusted means for the *continuous* outcome variable under (a) LIN, (b) CUB, (c) EXP, and (d) SIN, and SE ratios (rSE) under (e) LIN, (f) CUB, (g) EXP, and (h) SIN, across different DR methods under different model specification scenarios when  $gamma_1 = 0.3$  and the complexity in the design of  $S_R$  is ignored. UW: unweighted; FW: Fully weighted; PAPP: Propensity Adjusted Probability Prediction; GPPP: Gaussian Processes of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting



Fig 17: Comparing the 95% CI coverage rates (crCI) of the DR adjusted means for the *continuous* outcome variable under (a) LIN, (b) CUB, (c) EXP, and (d) SIN, and SE ratios (rSE) under (e) LIN, (f) CUB, (g) EXP, and (h) SIN, across different DR methods under different model specification scenarios when  $\gamma_1 = 0.6$ . UW: unweighted; FW: Fully weighted; PAPP: Propensity Adjusted Probability Prediction; GPPP: Gaussian Processes of Propensity Prediction; LWP: Linear-in-weight Prediction; AIPW: Augmented Inverse Propensity Weighting

UW: 1 predic	AIPW	LWP	PSPP	GPPP	Model sp	AIPW	LWP	PSPP	GPPP	Model sp	AIPW	LWP	PSPP	GPPP	Model sp	AIPW	LWP	PSPP	GPPP	Model sp	Doubly	IPSW	Model sp	IPSW	Model sp	Non-rok	FW	UW	Non-pro	FW	UW	Probabi	Measure		
Unweight tion; AIF	23.625	23.483	23.569	23.523	ecification	-0.844	0.281	0.877	0.382	ecification	0.690	0.595	0.603	0.580	ecification	0.542	0.531	0.559	0.547	ecification	robust r	23.624	ecification	-0.807	ecification	oust met	-0.002	33.632	bability	0.575	-31.577	lity sam	e rBias		) F
ed; FW: <sup>9</sup> W: Aug	23.902	23.746	23.851	23.802	n: QR–Fa	9.046	5.371	5.321	5.287	n: QR–Fa	6.246	6.261	6.245	6.258	n: QR–Tr	6.246	6.256	6.238	6.244	n: QR–Tr	nethods	23.936	n: QR–Fa	7.220	n: QR–Tr	hod	4.433	33.837	sample	6.592	31.829	ple $(S_R)$	rMSE		
Fully w mented	0.000	0.000	1.091	0.893	ılse, PM	95.536	95.734	95.635	95.238	ılse, PM	95.238	94.940	95.337	95.139	ue, PM	95.833	95.833	96.032	95.734	ue, PM		0.000	lse	95.734	ue.		94.444	0.000	$(S_A)$	94.345	0.000		crCI	LIN	
eighted: Inverse	14.239	14.020	15.829	15.401	–False	36.755	21.534	21.682	21.551	-True	24.808	24.871	24.934	25.016	–False	25.199	25.188	25.210	25.204	-True		15.020		29.561			16.857	14.373		25.873	9.476		lCI		1
; GPPP e Propei	1.001	1.014	1.104	1.079		1.041	1.024	1.053	1.042		1.019	1.018	1.023	1.024		1.033	1.030	1.035	1.033			0.995		1.051			0.970	0.985		1.005	0.604		rSE		1
: Gaussia nsity Wei	17.290	17.250	17.375	17.230		-0.606	0.384	0.915	0.517		0.519	0.515	0.524	0.514		0.511	0.498	0.521	0.503			17.277		-0.495			0.032	26.107		0.400	-25.314		rBias		
m Proce ighting;	17.568	17.523	17.900	17.533		7.323	4.367	4.376	4.321		5.366	5.374	5.380	5.369		5.360	5.358	5.352	5.352			17.587		5.698			3.599	26.300		5.581	25.534		rMSE		1
ss of Pr	0.000	0.000	2.480	1.786		95.933	94.940	95.139	94.742		94.246	94.345	94.444	94.643		94.643	94.940	94.444	94.444			0.000		95.040			94.841	0.000		93.552	0.000		crCI	CUB	1
opensity	12.276	12.217	18.527	14.060		29.641	17.599	17.975	17.664		21.004	21.007	21.232	21.271		21.153	21.168	21.228	21.362			12.880		22.974			13.921	12.565		21.858	7.895		lCI		1
<sup>7</sup> Predic	1.006	1.011	1.097	1.105		1.036	1.032	1.071	1.050		1.003	1.001	1.011	1.015		1.011	1.012	1.016	1.022			1.000		1.032			0.986	1.007		1.001	0.603		rSE		1
tion; PS	15.737	15.655	15.670	15.657		-0.659	0.140	0.665	0.242		0.286	0.322	0.319	0.320		0.322	0.328	0.287	0.302			15.739		-0.608			-0.088	23.527		0.333	-23.741		rBias		1
PP: Pen	16.013	15.921	15.950	15.936		6.581	4.067	4.082	4.027		5.071	5.081	5.072	5.110		5.088	5.082	5.078	5.082			16.043		5.222			3.197	23.712		5.279	23.950		rMSE		1
alized S	0.000	0.000	1.389	2.083		94.345	94.544	95.139	95.238		94.444	94.345	94.345	94.643		93.948	94.147	94.048	94.246			0.099		95.337			95.437	0.000		94.444	0.000		crCI	EXP	1
pline of	11.660	11.549	12.967	12.941		26.312	16.124	16.602	16.317		19.865	19.881	20.012	20.254		19.823	19.853	20.057	20.065			12.204		20.781			12.691	11.612		20.565	7.280		lCI		1
Propens	1.004	1.015	1.112	1.111		1.025	1.012	1.051	1.035		1.000	1.000	1.008	1.013		0.995	0.998	1.009	1.008			1.001		1.022			1.013	1.001		0.995	0.589		rSE		1
sity Pred	28.366	28.059	28.080	28.044		-0.869	-0.486	0.958	0.618		0.578	0.557	0.574	0.558		0.550	0.552	0.557	0.553			28.363		-0.945			-0.033	32.309		0.545	-33.737		rBias		
iction; I	28.651	28.330	28.352	28.317		8.225	6.869	5.647	5.628		6.608	6.615	6.617	6.601		6.554	6.556	6.554	6.573			28.657		7.794			4.947	32.543		6.820	34.038		rMSE		1
MP: Li	0.000	0.000	0.099	0.198		94.940	95.040	95.536	95.536		94.841	94.742	94.841	95.040		94.742	94.643	94.940	95.040			0.000		95.437			95.337	0.000		94.147	0.000		crCI	NIS	1
near-in-	15.857	15.767	15.816	15.841		33.733	27.506	22.702	23.133		26.260	26.288	26.441	26.620		26.397	26.412	26.483	26.622			16.066		32.004			19.326	15.189		26.873	11.272		ICI		
weight	1.004	1.029	1.029	1.031		1.052	1.024	1.040	1.054		1.017	1.017	1.023	1.032		1.031	1.031	1.034	1.036			1.000		1.055			0.996	0.994		1.008	0.636		rSE		I

	Table
	2
$\gamma_1=0.3$	Comparing
	th
	e performance
	of 1
	the
	bias
	adjustment
	methods i
	n tł
	ne simulation
	ı study
	for
	the
	continuous o
	outcome
	with
	$(n_A,$
	$n_R)$
	ت ت
	500, 1
	., 000
	) and

**Table 3:** Comparing the performance of the bias adjustment methods in the simulation study for the *continuous* outcome with  $(n_A, n_R) = (1, 000, 500)$  and  $\gamma_1 = 0.3$ 

			LIN					CUB					EXP					SIN		
Measure	rBias	rMSE	crCI	ICI	rSE	rBias	rMSE	crCI	ICI	rSE	rBias	rMSE	crCI	ICI	rSE	rBias	rMSE	crCI	ICI	rSE
Probabili	ty samp	ole $(S_R)$																		
UW FW	-32.193	32.514 7 362	0.000	13.346	0.747 0.979	-25.780	26.036 6.150	0.000	$\frac{11.118}{24.037}$	0.780	-24.168 -0.306	24.422 5 041	0.000	10.264 22.876	0.746 0.983	-34.387 -0.296	34.793 7 867	0.000 03 452	15.911 30.374	0.766
Non-nrol	hility	sample	(84)	044.04	0.00	71-0-0-	007-00	001.20	100.117	0,000	00000-	TECO	001.40		00000	0.4.0		101-00	-	
	33 130	33 220	0.000	10.150	1 000	25.641	95 734	0.000	8 875	1 035	93 909	23.206	0.000	8 107	1 004	32.060	32-176	0.000	10 744	1 044
FW	-0.143	3.036	95.040	11.958	1.005	-0.114	2.478	94.940	9.882	1.018	-0.093	2.289	94.147	8.969	1.000	0.048	3.346	94.841	13.613	1.037
Non-robi	ist meth	pot																		
Model spe	cification	1: QR-Th	en.																	
IPSW	-2.670	9.146	91.766	33.905	0.988	-2.075	7.056	92.560	26.030	0.984	-1.770	6.239	92.857	23.366	0.996	-2.672	9.423	92.063	35.748	1.009
Model spe	cification	1: QR-F	ulse																	
IPSW	22.769	23.070	0.000	14.501	0.995	16.574	16.873	0.000	12.370	0.998	15.171	15.451	0.000	11.407	0.993	27.891	28.054	0.000	12.288	1.037
Doubly r	obust n	lethods																		
Model spe	cification	1: QR-Th	.ue, PM-	-True																
GPPP	-0.403	7.024	92.956	26.771	0.973	-0.432	5.810	92.460	22.190	0.977	-0.375	5.426	92.361	20.812	0.980	-0.213	7.489	93.353	29.064	0.990
PSPP	-0.391	7.023	92.857	26.763	0.973	-0.429	5.815	92.262	22.168	0.975	-0.386	5.428	92.361	20.817	0.980	-0.213	7.485	93.254	29.036	0.990
LWP	-0.424	7.036	92.857	26.752	0.971	-0.441	5.807	92.063	22.151	0.975	-0.365	5.420	92.163	20.789	0.980	-0.218	7.478	93.254	29.030	0.990
AIPW	-0.434	7.056	92.659	26.828	0.971	-0.435	5.805	92.361	22.179	0.977	-0.356	5.400	92.163	20.781	0.983	-0.229	7.492	93.056	29.064	0.990
Model spe	cificatior	1: QR-Th	ue, PM-	-False																
GPPP	-0.449	7.035	92.560	26.762	0.972	-0.389	5.785	92.361	22.169	0.979	-0.375	5.435	92.262	20.817	0.979	-0.207	7.577	93.651	29.008	0.977
PSPP	-0.423	7.025	92.560	26.754	0.973	-0.392	5.789	92.361	22.146	0.978	-0.387	5.438	92.063	20.817	0.979	-0.198	7.571	93.750	28.994	0.977
LWP	-0.450	7.032	92.659	26.725	0.971	-0.402	5.787	92.361	22.144	0.978	-0.385	5.437	92.163	20.797	0.978	-0.206	7.566	93.651	29.004	0.978
AIPW	-0.335	7.001	92.659	26.656	0.972	-0.400	5.810	92.063	22.126	0.973	-0.428	5.438	91.964	20.802	0.979	-0.183	7.552	93.849	28.981	0.979
Model spe	cificatior	1: QR-F	ulse, PM	-True																
GPPP	-0.370	6.080	91.567	23.011	0.967	-0.220	4.731	92.262	18.194	0.982	-0.238	4.359	92.262	16.587	0.972	0.005	6.556	92.956	25.445	0.990
PSPP	-0.141	6.049	92.063	23.040	0.971	-0.052	4.706	93.056	18.231	0.988	-0.019	4.343	91.865	16.635	0.977	0.140	6.536	92.956	25.376	0.990
LWP	-0.247	5.957	91.766	22.571	0.967	-0.221	4.717	92.063	18.143	0.982	-0.186	4.274	92.262	16.322	0.975	-0.026	6.598	93.948	25.603	0.989
AIPW	-3.276	11.468	92.659	42.552	0.987	-2.575	9.175	94.048	34.192	0.990	-2.202	8.033	92.460	29.757	0.982	-2.820	10.042	92.063	37.905	1.003
Model spe	cificatior.	1: QR-F	ulse, PM	-False																
GPPP	23.085	23.346	0.000	13.851	1.015	16.852	17.118	0.099	12.023	1.020	15.413	15.651	0.099	10.979	1.030	27.846	28.017	0.000	12.644	1.044
PSPP	23.093	23.353	0.397	14.099	1.035	16.899	17.211	0.794	13.589	1.063	15.425	15.663	0.298	11.093	1.038	27.841	28.012	0.000	12.695	1.048
LWP	23.068	23.330	0.000	13.694	1.002	16.837	17.103	0.000	11.764	1.000	15.399	15.637	0.000	10.846	1.017	28.010	28.159	0.000	12.121	1.068
AIPW	23.053	23.323	0.000	13.735	0.990	16.837	17.106	0.000	11.776	0.993	15.393	15.639	0.000	10.861	1.001	27.990	28.146	0.000	12.059	1.039
TTWV-TF	nwei ahte	Nd. FW.	Eully wa	aiahtad.	CPPP.	Gameeia	n Proces	se of Pro	mensity	- Predict	-ion PSP	P. Pana	lized Sr	line of l	Dronens	ity Predi	ction. I	WP. Lin	-ui-rear	weight
			Turry w	rigmen.		· /III ··			forerro de	TICATO	1011, 1 01		In nazire		errodot i	mat t Ann				verguu
predict.	ion; AlF	W: Aug	mented	Inverse	Proper.	isity Wei	ghting;													

UW: U predict	AIPW	LWP	PSPP	GPPP	Model spe	AIPW	LWP	PSPP	GPPP	Model spe	AIPW	LWP	PSPP	GPPP	Model spe	AIPW	LWP	PSPP	GPPP	Model spe	Doubly 1	IPSW	Model spe	IPSW	Model spe	Non-rob	FW	UW	Non-pro	FW	UW	Probabil	Measure		
nweight ion; AIF	23.159	23.169	23.239	23.208	ecification	-3.371	-0.474	0.045	-0.442	ecification	-0.349	-0.478	-0.441	-0.476	ecification	-0.453	-0.459	-0.397	-0.416	ecification	obust n	22.879	ecification	-2.816	ecificatior	ust metl	-0.097	33.538	bability	-0.395	-32.193	ity samı	rBias		
ed; FW: W: Aug	23.512	23.511	23.592	23.555	1: QR–Fa	12.232	6.189	6.185	6.204	1: QR–Fa	7.171	7.200	7.199	7.200	1: QR-T	7.178	7.170	7.145	7.151	1: QR–T	nethods	23.267	1: $QR-F_2$	9.640	1: QR–Ti	pot	4.370	33.739	sample	7.335	32.514	ple $(S_R)$	rMSE		
Fully w gmented	0.000	0.000	0.992	1.091	alse, PM	94.246	93.452	93.056	92.163	alse, PM	91.964	91.964	91.964	92.163	rue, PM-	92.163	92.063	92.361	92.163	rue, PM-		0.000	alse	92.857	rue		94.048	0.000	$(S_A)$	92.956	0.000		crCI	LIN	
eighted; Inverse	15.863	15.785	17.006	16.675	–False	45.300	23.853	24.170	23.930	-True	27.329	27.413	27.484	27.494	-False	27.478	27.418	27.469	27.480	-True		16.832		36.078			16.894	14.374		28.305	13.346		lCI		
GPPP: Propen	0.997	1.007	1.067	1.056		0.982	0.986	0.996	0.986		0.973	0.973	0.975	0.976		0.978	0.977	0.982	0.982			1.015		0.998			0.986	0.996		0.985	0.747		rSE		
Gaussia sity Wei	16.862	16.854	16.895	16.864		-2.445	-0.219	0.231	-0.147		-0.410	-0.420	-0.404	-0.408		-0.415	-0.429	-0.420	-0.422			16.608		-2.150			-0.082	25.970		-0.338	-25.780		rBias		
m Proce ghting;	17.238	17.222	17.282	17.241		9.765	4.974	4.955	4.945		5.931	5.931	5.928	5.939		5.914	5.931	5.922	5.928			17.015		7.463			3.616	26.160		6.109	26.036		rMSE		
ss of Pro	0.000	0.000	2.183	0.893		94.345	93.254	93.651	93.254		91.667	91.865	91.667	91.964		92.063	91.964	91.865	92.063			0.000		93.849			94.742	0.000		92.262	0.000		crCI	CUB	
opensity	13.662	13.679	15.411	14.527		36.292	19.331	19.624	19.356		22.760	22.781	22.886	22.918		22.809	22.791	22.818	22.941			14.371		27.889			13.925	12.566		24.115	11.118		lCI		
Predict	0.974	0.986	1.080	1.033		0.979	0.992	1.011	0.998		0.981	0.982	0.987	0.986		0.986	0.982	0.985	0.989			0.991		0.995			0.982	1.019		1.008	0.780		rSE		
ion; PSI	15.417	15.412	15.470	15.449		-2.309	-0.343	0.159	-0.304		-0.514	-0.446	-0.458	-0.421		-0.431	-0.415	-0.450	-0.433			15.199		-1.945			-0.097	23.492		-0.331	-24.168		rBias		
PP: Pena	15.767	15.751	15.819	15.799		8.690	4.532	4.593	4.567		5.557	5.541	5.557	5.540		5.525	5.543	5.557	5.549			15.579		6.626			3.310	23.681		5.907	24.422		rMSE		
alized Sp	0.099	0.099	1.687	1.091		94.544	93.452	93.750	92.857		92.361	92.163	92.262	92.262		92.560	92.460	92.659	92.560			0.198		93.254			94.246	0.000		92.659	0.000		crCI	EXP	
oline of I	12.785	12.734	13.592	13.411		32.432	17.846	18.225	18.001		21.342	21.331	21.455	21.423		21.510	21.513	21.623	21.670			13.450		25.123			12.694	11.610		22.926	10.264		ICI		
<sup>o</sup> ropensi	0.988	0.998	1.050	1.035		0.987	1.007	1.012	1.007		0.984	0.985	0.988	0.989		0.996	0.992	0.995	0.999			1.002		1.011			0.978	0.992		0.991	0.746		rSE		
ty Predi	28.121	28.074	28.014	27.977		-3.081	-0.890	0.298	-0.021		-0.231	-0.251	-0.237	-0.246		-0.253	-0.252	-0.224	-0.220			28.020		-2.840			0.055	32.315		-0.318	-34.387		rBias		
ction; L	28.420	28.358	28.319	28.284		10.672	7.763	6.739	6.805		7.609	7.606	7.618	7.618		7.612	7.604	7.616	7.633			28.331		9.985			4.904	32.539		7.858	34.793		rMSE		
WP: Lir	0.000	0.000	0.000	0.099		91.865	94.643	93.552	93.651		93.552	93.651	93.452	93.254		93.651	93.750	93.750	93.651			0.000		92.560			95.238	0.000		93.056	0.000		crCI	NIS	
1ear-in-v	16.220	16.187	16.495	16.494		40.608	30.292	26.163	26.639		29.506	29.567	29.639	29.664		29.377	29.361	29.444	29.602			16.501		38.102			19.299	15.176		30.282	15.911		lCI		
veight	1.005	1.030	1.016	1.012		1.013	1.002	0.991	0.998		0.989	0.992	0.992	0.993		0.985	0.985	0.986	0.989			1.006		1.015			1.003	1.016		0.983	0.766		rSE		

	Table
	<u>4</u>
$\gamma_1$	Q
	mp
	arin
	ю́ t
	ne t
	erfo
	orm
	anc
	e of
	th∈
	bia
	as a
	dju
	stm
	ent
	met
	hoc
	ls ir
	1 th
	e sir
	nula
	atio
	n st
	udy
	for
	the
	cor
	ntin
	non
	10 8
	ıtco
	me
	wit]
	h (n
	A, r
	$\iota_R)$
	=
	500;
	- 50C
	)) a:
	nd

hud	
0) a	
1,00	
00,	
= (5	
R =	
A, n	
n (n	
wit]	
ome	
utco	
ns o	
onu	
conti	
he c	
or t	
dy f	
ı stu	
tion	
nule	
e sii	
n th	
ds i	
etho	
t me	
men	
just	
s ad	
bia	
the	
e of	
lanc	
forn	
per	
$_{\mathrm{the}}$	
ring	9
npa	0.
Col	$\gamma_1$ :
۰ ت	
able	
H	

																		1		
Measure	rBias	$_{\rm rMSE}$	LLIN crCI	ICI	rSE	rBias	rMSE	crCI	ICI	rSE	rBias	$\mathbf{r}\mathbf{MSE}$	erCI	ICI	rSE	rBias	$\mathbf{r}\mathbf{MSE}$	s1N crCI	ICI	rSE
Probabili	ty samp	le $(S_R)$																		
UW FW	-31.577 0.544	31.829 6.601	$0.000 \\ 93.552$	9.476 25.637	$0.604 \\ 0.994$	-25.314 0.365	25.534 5.619	$0.000 \\ 93.750$	7.895 21.558	0.603 0.980	-23.741 0.323	23.950 5.261	0.000 93.948	$7.280 \\ 20.656$	$0.589 \\ 1.003$	-33.737 0.606	34.038 6.848	0.000 93.948	11.272 26.869	$0.636 \\ 1.004$
Non-prob	ability	sample	$(S_A)$																	
M	67.162	67.257	0.000	14.469	1.030	54.231	54.329	0.000	13.571	1.058	48.409	48.503	0.000	12.142	1.024	54.984	55.080	0.000	12.872	1.014
FW	0.214	7.308	91.667	26.736	0.933	0.087	5.907	91.171	21.731	0.938	-0.041	5.175	93.056	19.335	0.953	0.127	7.239	93.552	28.085	0.989
Non-robu	ist meth	pou																		
Model spe	cification	CR-Th	ene																	
IPSW	-2.198	12.496	92.659	47.329	0.981	-1.641	9.851	93.651	36.404	0.956	-1.691	8.734	93.155	33.136	0.986	-2.445	12.350	93.552	48.815	1.028
Model spe	cification	1: QR-F	alse																	
IPSW	37.613	37.906	0.000	18.408	0.997	27.228	27.529	0.000	15.746	0.989	25.215	25.501	0.000	14.666	0.984	47.301	47.483	0.000	16.469	1.012
Doubly r	obust n	nethods																		
Model spec	cification	CR-Th	ue, PM-	-True																
GPPP	0.634	6.481	94.940	26.557	1.050	0.365	5.701	94.345	23.324	1.045	0.270	5.380	94.147	21.977	1.043	0.555	6.730	95.635	28.602	1.087
PSPP	0.727	6.448	95.337	26.385	1.050	0.442	5.634	94.147	22.616	1.027	0.154	5.352	94.048	21.614	1.030	0.572	6.690	95.933	27.883	1.067
LWP	0.484	6.623	95.040	26.601	1.027	0.371	5.810	93.750	22.827	1.004	0.318	5.478	93.552	21.715	1.013	0.516	6.792	95.337	27.849	1.049
AIPW	0.588	6.517	94.643	26.259	1.032	0.397	5.764	94.147	22.412	0.994	0.266	5.436	93.254	21.336	1.002	0.537	6.719	95.437	27.381	1.042
Model spe	cification	1: QR-T	rue, PM-	-False																
GPPP	0.574	6.496	95.635	25.962	1.023	0.408	5.807	94.544	23.289	1.025	0.270	5.402	93.254	21.233	1.003	0.457	6.715	95.139	27.287	1.039
PSPP	0.692	6.448	95.635	25.861	1.029	0.467	5.599	94.246	22.261	1.017	0.170	5.336	93.651	20.972	1.003	0.531	6.656	95.040	26.988	1.037
LWP	0.726	6.506	95.437	25.888	1.021	0.471	5.677	94.444	22.094	0.996	0.167	5.384	92.956	21.010	0.996	0.474	6.692	95.437	26.992	1.031
AIPW	0.926	6.432	96.032	25.640	1.027	0.428	5.538	93.750	21.776	1.006	0.059	5.347	93.254	20.739	0.989	0.590	6.629	95.040	26.723	1.032
Model spe	cification	1: QR-F	alse, PM	-True																
GPPP	0.473	5.841	95.238	24.700	1.082	1.173	5.124	94.544	21.555	1.102	0.377	4.722	95.734	20.135	1.091	0.583	6.340	95.734	28.878	1.166
PSPP	2.367	6.515	94.048	26.531	1.114	2.392	5.917	92.262	23.361	1.101	1.968	5.317	93.750	21.951	1.133	1.783	6.260	94.345	26.286	1.117
LWP	-3.023	9.265	96.329	36.224	1.055	-1.289	6.272	95.833	25.461	1.058	-2.038	6.653	96.131	25.586	1.030	-7.996	17.881	91.865	62.917	1.003
AIPW	-2.472	15.606	93.056	57.317	0.948	-1.936	13.033	92.758	46.933	0.928	-1.871	11.081	93.056	40.685	0.950	-2.444	12.657	93.651	49.579	1.018
Model spe	cification	1: QR-F	alse, $PM$	-False																
GPPP	34.240	34.426	0.000	14.190	1.012	24.837	25.044	0.000	12.727	1.010	22.870	23.079	0.000	11.997	0.987	44.075	44.248	0.000	15.383	1.005
PSPP	34.287	34.473	0.000	14.242	1.015	24.940	25.146	0.000	12.766	1.013	22.967	23.175	0.000	12.035	0.991	43.983	44.155	0.000	15.419	1.011
LWP	34.158	34.350	0.000	14.344	1.009	24.704	24.917	0.000	12.785	1.002	22.762	22.976	0.000	12.053	0.981	45.852	46.020	0.000	15.687	1.019
AIPW	35.685	35.890	0.000	15.018	1.001	25.334	25.549	0.000	12.935	0.998	23.645	23.860	0.000	12.313	0.982	46.967	47.144	0.000	16.223	1.013
UW: U	aweighte	d: FW:	Fully w	eighted:	GPPP:	Gaussia	n Proce	ss of Prc	pensity	Predict	tion: PSF	P: Pena	lized Sp	line of I	ropens	itv Predi	ction: I	WP: Li	near-in-v	veight
predicti	on; ĂIP	W: Aug	mented	Inverse	Proper	isity Wei	ghting;		> 	1		1	•		4	6	~	1		p

UW: U predict	AIPW	LWP	PSPP	GPPP	Model sp	AIPW	LWP	PSPP	GPPP	Model sp	AIPW	LWP	PSPP	GPPP	Model sp	AIPW	LWP	PSPP	GPPP	Model sp	Doubly	IPSW	Model sp	IPSW	Model sp	Non-rob	FW	UW	Non-pro	FW	UW	Probabi	Measure		
Jnweight sion; AIF	34.909	33.761	33.720	33.746	ecificatio	-5.685	-2.228	0.739	-0.339	ecificatio	-0.174	-0.439	-0.418	-0.527	ecificatio	-0.542	-0.571	-0.382	-0.446	ecificatio	robust r	36.123	ecificatio	-4.867	ecificatio	ust met	0.189	65.596	bability	-0.525	-32.193	ity sam	rBias		
ed; FW: <sup>9</sup> W: Aug	35.127	33.972	33.925	33.950	n: QR–Fa	17.931	9.094	6.360	6.274	n: QR–Fa	7.144	7.204	7.181	7.224	n: QR–Ti	7.258	7.228	7.106	7.130	a: QR–Ti	nethods	36.423	n: QR–Fa	14.193	n: QR–Ti	hod	5.124	65.646	sample	7.302	32.514	ple $(S_R)$	rMSE		
Fully w mented	0.000	0.000	0.000	0.000	alse, PM-	93.750	94.147	93.155	92.063	alse, PM-	91.468	90.873	91.071	90.972	rue, PM-	92.262	92.361	92.460	92.361	rue, PM-		0.000	alse	91.766	rue		92.460	0.000	$(S_A)$	92.659	0.000	_	crCI	LIN	
eighted; Inverse	14.878	14.581	14.394	14.357	-False	63.563	34.246	24.830	24.082	-True	27.063	27.226	27.221	27.305	-False	27.894	27.723	27.585	27.574	-True		17.563		51.753			19.090	10.167		28.356	13.346		lCI		
GPPP: Propens	0.971	0.983	0.986	0.983		0.953	0.990	1.002	0.980		0.966	0.965	0.968	0.966		0.983	0.981	0.991	0.988			0.960		0.990			0.951	1.013		0.993	0.747		rSE		
Gaussia sity Wei	24.751	24.281	24.398	24.333		-4.543	-1.293	0.789	0.249		-0.327	-0.320	-0.326	-0.362		-0.399	-0.447	-0.382	-0.465			25.933		-3.670			0.156	52.643		-0.341	-25.780		rBias		
n Proces ghting;	24.973	24.504	24.612	24.549		14.755	6.413	5.201	5.047		5.912	5.897	5.892	5.957		5.928	5.924	5.878	5.916			26.232		11.036			4.164	52.693		6.133	26.036		rMSE		
ss of Pro	0.000	0.000	0.000	0.000		94.940	95.536	94.246	93.254		92.262	92.163	92.262	92.262		92.956	92.857	92.857	92.857			0.000		92.560			93.452	0.000		92.560	0.000		crCI	CUB	
pensity	12.705	12.606	12.445	12.472		51.751	24.766	20.681	19.747		22.427	22.576	22.590	22.889		22.827	22.770	22.622	22.849			14.990		39.534			15.456	9.483		23.982	11.118		ICI		
Predict	0.976	0.975	0.979	0.978		0.940	1.005	1.026	0.999		0.969	0.978	0.979	0.982		0.984	0.983	0.983	0.988			0.968		0.969			0.947	1.052		0.999	0.780		rSE		
ion; PSF	23.136	22.464	22.530	22.495		-3.861	-1.467	0.650	-0.101		-0.589	-0.455	-0.462	-0.371		-0.318	-0.307	-0.489	-0.371			24.127		-3.159			0.118	47.162		-0.328	-24.168		rBias		
PP: Penz	23.333	22.656	22.715	22.681		12.612	6.143	4.777	4.573		5.524	5.563	5.526	5.538		5.541	5.570	5.573	5.540			24.396		9.561			3.639	47.209		5.973	24.422		rMSE		
dized Sp	0.000	0.000	0.000	0.000		94.742	94.841	94.048	93.056		91.766	92.361	92.262	92.460		93.353	93.056	92.758	92.659			0.000		91.171			93.254	0.000		92.560	0.000		crCI	EXP	
line of F	11.856	11.697	11.541	11.546		44.501	23.780	19.098	18.084		21.391	21.502	21.489	21.580		21.668	21.565	21.585	21.555			13.830		35.012			13.690	8.512		22.907	10.264		ICI		
<sup>o</sup> ropensi	1.000	1.012	1.016	1.015		0.945	1.016	1.029	1.009		0.993	0.989	0.995	0.996		0.999	0.989	0.991	0.994			0.975		0.989			0.960	1.033		0.979	0.746		rSE		
ty Predi	46.416	45.559	43.725	43.763		-4.997	-4.775	0.707	0.134		-0.145	-0.234	-0.202	-0.242		-0.287	-0.289	-0.233	-0.255			46.633		-4.899			0.295	54.653		-0.308	-34.387		rBias		
ction; L	46.524	45.658	43.855	43.892		14.525	15.831	6.824	6.915		7.516	7.527	7.526	7.576		7.691	7.659	7.589	7.631			46.748		14.133			5.043	54.700		7.900	34.793		rMSE		
WP: Lin	0.000	0.000	0.000	0.000		91.171	93.452	93.353	93.452		93.552	93.651	93.552	93.750		93.155	93.452	94.246	94.048			0.000		91.071			94.345	0.000		93.254	0.000		crCI	SIN	
ıear-in-v	12.466	11.969	13.229	13.207		54.283	55.293	26.604	27.301		29.321	29.419	29.398	29.494		29.800	29.760	29.646	29.894			12.846		52.612			19.700	9.133		30.372	15.911		ICI		
weight	1.004	1.017	1.000	1.002		1.015	0.934	0.999	1.007		0.995	0.997	0.996	0.993		0.989	0.991	0.996	0.999			1.001		1.012			0.998	1.023		0.981	0.766		rSE		

	Table
	е <b>6</b> :
$\gamma_1$	Co
= 0	mp
.6	arin
	g th
	e pe
	rfor
	mar
	ice (
	of th
	ıe bi
	ias a
	ldju
	stm
	ent i
	met]
	hods
	in
	the
	sim
	ılati
	ons
	tud
	y foj
	r the
	00
	ntin
	uou
	no s
	tcor
	ne w
	rith
	$(n_A,$
	$n_R)$
	, O(
	00, 5
	(00
	and

**Table 7:** Comparing the performance of the bias adjustment methods in the simulation study for the *continuous* outcome with  $(n_A, n_B) = (500, 500)$  and  $\gamma_1 = 0.6$ 

;	ļ	1.05 F	LIN	101	ç	Ē		CUB	101	ļ	ļ		EXP	101	Ę	ļ	103	SIN	101	1
Measure	rBias	rMSE	crCI	ICI	rse	rbias	rMSE	crCI	ICI	rse	rBias	rMSE	crCI	ICI	rSE	rbias	rMSE	crCI	ICI	SE E
Probabili	ty samp	ole $(S_R)$																		
MD	-32.193	32.514	0.000	13.346	0.747	-25.780	26.036	0.000	11.118	0.780	-24.168	24.422	0.000	10.264	0.746	-34.387	34.793	0.000	15.911	0.766
F.W	-0.478	7.302	92.560	28.264	0.989	-0.323	6.128	92.262	24.001	1.000	-0.330	5.892	92.063	22.883	0.992	-0.277	7.854	93.452	30.285	0.984
Non-prot	ability	sample	(SA)																	
ΜŪ	66.963	67.069	0.000	14.460	0.978	54.004	54.111	0.000	13.565	1.018	48.290	48.392	0.000	12.131	0.985	54.966	55.063	0.000	12.869	1.008
FW	0.267	7.234	91.964	26.459	0.933	0.138	5.927	93.155	21.376	0.920	0.173	5.243	92.361	19.079	0.928	0.312	7.228	94.841	28.042	0.99
Non-robı	ıst metl	por																		
Model spe	cification	1: QR-Th	en.																	
IPSW	-5.988	16.290	92.460	57.597	0.969	-4.537	12.766	93.155	44.298	0.947	-3.931	10.860	91.766	39.169	0.987	-6.370	15.995	90.079	59.186	1.029
Model spe	cification	1: QR-Fe	ulse																	
IPSW	36.099	36.481	0.000	20.047	0.971	25.809	26.200	0.000	17.184	0.972	24.049	24.419	0.000	15.911	0.958	46.657	46.863	0.000	17.101	0.994
Doubly r	obust n	nethods																		
Model spe	cification	1: QR-Tr	ue, PM-	-True																
GPPP	-0.356	7.303	92.758	28.865	1.009	-0.543	6.284	92.758	25.177	1.025	-0.334	5.741	94.643	23.626	1.051	-0.191	7.809	95.536	31.859	1.041
PSPP	-0.298	7.272	92.758	28.643	1.005	-0.471	6.223	92.758	24.304	0.999	-0.521	5.763	94.147	23.141	1.028	-0.149	7.789	95.139	30.904	1.012
LWP	-0.530	7.521	92.857	29.081	0.988	-0.486	6.319	92.956	24.642	0.997	-0.212	5.845	94.147	23.347	1.019	-0.218	7.917	94.841	31.093	1.002
AIPW	-0.481	7.478	92.361	28.902	0.988	-0.461	6.267	93.056	24.214	0.988	-0.287	5.753	94.048	23.050	1.023	-0.200	7.935	94.345	30.681	0.986
Model spe	cification	1: QR-Th	ue, PM-	-False																
GPPP	-0.470	7.353	92.560	28.513	0.991	-0.434	6.317	93.155	24.923	1.008	-0.352	5.793	93.254	22.718	1.002	-0.341	7.962	94.246	30.438	0.976
PSPP	-0.346	7.296	92.460	28.364	0.992	-0.413	6.145	92.560	23.883	0.993	-0.490	5.751	93.056	22.460	0.999	-0.258	7.857	94.444	30.105	776.0
LWP	-0.374	7.329	92.560	28.405	0.989	-0.380	6.184	92.361	23.722	0.980	-0.500	5.830	92.659	22.456	0.986	-0.303	7.824	94.246	30.134	0.983
AIPW	-0.127	7.252	92.460	28.115	0.989	-0.459	6.153	92.857	23.438	0.974	-0.655	5.755	92.758	22.206	0.990	-0.210	7.837	94.147	29.890	0.973
Model spe	cification	$1: QR-F_{5}$	ulse, PM	-True																
GPPP	-0.352	6.580	94.345	27.253	1.058	0.507	5.550	94.444	22.889	1.056	0.021	5.145	94.940	21.731	1.077	-0.179	7.539	95.337	32.738	1.108
PSPP	1.528	6.825	95.337	28.887	1.107	1.684	6.065	94.643	25.175	1.102	1.533	5.678	94.345	23.655	1.103	1.069	7.255	94.345	29.606	1.052
LWP	-4.549	12.067	94.940	42.565	0.971	-2.484	8.075	96.230	30.197	1.002	-2.878	8.105	94.940	30.271	1.019	-9.622	21.835	91.567	71.398	0.929
AIPW	-6.795	21.022	94.841	71.617	0.918	-5.265	17.481	94.940	57.749	0.883	-4.513	14.430	94.048	49.842	0.927	-6.441	16.546	90.179	60.223	1.008
Model spe	cification	1: QR-Fe	$_{\rm Mse, PM}$	-False																
GPPP	33.879	34.125	0.000	16.200	1.010	24.355	24.629	0.000	14.299	0.995	22.560	22.814	0.000	13.346	1.003	43.989	44.182	0.000	16.532	1.021
PSPP	33.921	34.166	0.000	16.214	1.013	24.450	24.721	0.000	14.288	0.998	22.653	22.905	0.000	13.332	1.003	43.895	44.091	0.000	16.603	1.021
LWP	33.794	34.049	0.000	16.405	1.006	24.210	24.492	0.000	14.405	0.992	22.444	22.706	0.000	13.447	0.996	45.676	45.853	0.000	16.073	1.018
AIPW	34.925	35.195	0.000	16.849	0.987	24.677	24.960	0.000	14.515	0.987	23.106	23.377	0.000	13.619	0.978	46.466	46.662	0.000	16.718	0.998
TIVAZ. II.	amoi whto	.d. EVAI.	Ev.11	oichtod.	CDDD.	Conseries C	Duccord	or of Duc		Duchica	ion. DCD	D. Dono	lizod Cr	lino of l	Ducanon	itor Duodi	otion. I	VA/D. T :.	n ai aoo	+daior
	UIWEIBIIU	du, 1 vv.	r uny w	eiguveu,	5	ereeneb.	1 I I UUG	1 T T T T T T T T T T T T T T T T T T T	GULEILO	LIGUIU	1011, I DI	Г. т сла	Insen of	r to attri	erradou	ing i teur	споп, т	111 · T /	A -TIT - TRAT	/IIRII/
predict.	ion; AIF	W: Aug	mented	Inverse	Proper.	nsity Wei	ghting;													

UW: U predict	AIPW	LWP	PSPP	GPPP	Model spe	AIPW	LWP	PSPP	GPPP	Model spe	AIPW	LWP	PSPP	GPPP	Model spe	AIPW	LWP	PSPP	GPPP	Model spe	Doubly r	IPSW	Model spe	IPSW	Model spe	Non-rob	FW	UW	Non-prol	FW	UW	Probabil	Measure		
nweight ion; AIF	30.827	30.671	30.670	30.673	cification	-1.744	0.161	1.632	0.210	cification	0.341	0.311	0.367	0.498	cification	0.294	0.376	0.308	0.305	cification	obust n	31.422	cification	-1.380	cification	ist met	-0.065	44.519	oability	0.177	-40.760	ity samj	rBias		
ed; FW: W: Aug	31.416	31.262	31.264	31.269	1: QR–Fa	11.622	7.591	7.675	7.589	1: QR–Fa	9.126	9.124	9.127	9.123	1: QR-Ti	9.098	9.066	9.099	9.095	1: QR–Ti	nethods	32.111	1: QR–Fa	9.343	1: QR–Ti	hod	5.792	44.897	sample	9.453	41.085	ple $(S_R)$	rMSE		
Fully w mented	0.099	0.099	0.099	0.099	alse, PM	94.544	93.353	94.246	93.849	alse, PM	94.048	94.246	94.147	94.048	rue, PM-	94.048	94.345	94.345	94.444	rue, PM-		0.298	alse	93.552	ue		95.040	0.000	$(S_A)$	92.460	0.000		crCI	LIN	
eighted; Inverse	24.312	24.389	24.584	24.600	–False	47.373	30.803	30.754	30.850	-True	36.734	36.796	36.845	36.881	-False	36.641	36.601	36.695	36.708	-True		26.210		38.396			23.261	22.957		36.982	13.610		lCI		
GPPP: Propen	1.024	1.029	1.034	1.032		1.051	1.035	1.046	1.037		1.027	1.029	1.030	1.032		1.027	1.030	1.029	1.030			1.010		1.059			1.024	1.007		0.998	0.673		rSE		
Gaussia sity Wei	29.456	29.349	29.369	29.385		-1.869	-0.019	1.403	0.070		0.344	0.283	0.305	0.453		0.170	0.249	0.197	0.209			30.009		-1.417			-0.186	45.983		0.334	-44.127		rBias		
ın Proce ghting;	30.326	30.220	30.243	30.261		12.454	8.269	8.286	8.274		11.075	11.084	11.091	11.105		10.962	10.988	10.978	10.976			30.986		9.703			6.463	46.508		11.357	44.513		rMSE		
ss of Pro	1.290	1.488	1.587	1.587		95.238	94.544	95.238	94.940		93.452	93.552	93.452	93.849		93.651	93.452	93.353	93.155			2.381		93.552			94.940	0.000		93.155	0.000		crCI	CUB	
pensity	28.331	28.355	28.467	28.489		50.703	33.496	33.442	33.506		43.352	43.373	43.491	43.564		43.504	43.597	43.507	43.570			30.531		39.646			25.507	26.997		44.750	14.703		ICI		
Predict	1.001	1.003	1.005	1.005		1.050	1.033	1.044	1.033		0.999	0.998	1.000	1.001		1.012	1.012	1.011	1.012			1.009		1.053			1.007	0.987		1.005	0.642		rSE		
ion; PSF	28.355	28.241	28.260	28.263		-1.878	-0.114	1.161	-0.050		0.023	-0.046	0.042	0.235		-0.041	0.066	0.045	0.052			28.849		-1.530			-0.425	42.958		0.058	-42.643		rBias		
P: Pena	29.135	29.023	29.044	29.047		11.830	8.025	8.017	8.019		10.279	10.291	10.305	10.286		10.296	10.274	10.283	10.292			29.724		9.478			6.084	43.413		10.753	43.014		rMSE		
lized Sp	0.992	1.091	1.091	1.091		94.643	93.948	94.048	93.651		94.544	94.345	94.544	94.444		93.750	94.048	93.552	93.849			1.786		93.155			95.437	0.000		93.353	0.000		crCI	EXP	
line of F	26.651	26.674	26.808	26.805		47.349	31.474	31.547	31.554		40.624	40.694	40.885	40.953		40.663	40.745	40.709	40.779			28.479		38.062			24.581	25.222		41.452	14.229		ICI		
ropensi	1.015	1.017	1.020	1.019		1.034	1.000	1.014	1.003		1.008	1.008	1.012	1.016		1.007	1.011	1.009	1.010			1.014		1.038			1.033	1.026		0.983	0.644		rSE		
ty Predi	29.875	29.585	29.338	29.357		-1.389	0.119	1.202	0.242		0.229	0.535	0.309	0.614		0.132	0.489	0.200	0.206			30.020		-1.462			-0.209	36.027		0.093	-34.513		rBias		
ction; L	30.299	30.003	29.765	29.783		9.396	6.409	6.556	6.494		7.263	7.270	7.277	7.277		7.329	7.337	7.325	7.339			30.469		8.588			5.306	36.340		7.220	34.804		rMSE		
WP: Lin	0.000	0.000	0.000	0.000		93.155	93.849	93.353	93.948		94.048	94.544	94.246	94.544		92.659	92.758	92.956	93.155			0.000		93.254			94.246	0.000		92.659	0.000		crCI	NIS	
1ear-in-v	19.381	19.114 .	19.417	19.439		37.347	25.012	25.177	25.292		28.246	28.260	28.312	28.324		28.293	28.334	28.328	28.380			20.056		34.342			20.774	18.349		28.324	12.322		lCI		
veight	0.978	0.977	0.985	0.988		1.025	0.995	0.996	0.994		0.992	0.994	0.993	0.996		0.985	0.987	0.986	0.986			0.981		1.035			0.999	0.982		1	0.700		rSE		

**Table 8:** Comparing the performance of the bias adjustment methods in the simulation study for the binary outcome with  $(n_A, n_R) = (500, 1, 000)$  and  $\gamma_1 = 0.3$ 

12

**Table 9:** Comparing the performance of the bias adjustment methods in the simulation study for the *bimary* outcome with  $(n_A, n_B) = (1, 000, 500)$  and  $\gamma_1 = 0.3$ 

Measure	rBias	rMSE	LIN crCI	lCI	rSE	rBias	rMSE	CUB crCI	ICI	rSE	rBias	rMSE	EXP crCI	ICI	rSE	rBias	rMSE	SIN crCI	ICI	rSE
Probabili UW FW	ty sam <u>f</u> -41.047 -0.300	$\frac{\text{ole}(S_R)}{41.487}$ 10.798	0.000 93.849	$19.213 \\ 42.547$	$0.813 \\ 1.005$	-44.574 -0.588	45.028 12.476	0.000 93.353	20.730 50.758	0.828 1.039	-42.945 -0.555	43.406 11.617	0.000 93.651	20.084 $47.476$	$0.812 \\ 1.043$	-34.898 -0.358	35.315 8.568	$0.000 \\ 93.651$	17.395 33.456	0.820
Non-prok	44 057	sample 44 254	$(S_A)$	16.239	0.993	45304	45 566	0.000	19 086	0 998	42.865	43 094	0000	17 834	1 026	35,889	36 043	0000	12.980	0.992
FW	0.170	4.160	95.139	16.418	1.007	-0.039	4.564	95.139	18.010	1.006	0.158	4.328	95.635	17.384	1.025 $1.025$	0.142	3.739	95.635	14.658	1.000
Non-rob1	ist meth	pot																		
Model spe	cification	C QR-T	ene																	
IPSW	-2.157	10.471	92.857	42.950	1.069	-2.257	10.919	92.956	43.494	1.038	-1.980	10.273	92.857	41.675	1.054	-2.058	9.476	94.246	38.427	1.059
Model spe	cification	1: QR-F	ulse																	
IPSW	30.837	31.398	0.099	23.746	1.025	29.353	30.150	0.694	27.671	1.024	28.686	29.364	0.496	25.303	1.029	29.823	30.082	0.000	16.020	1.038
Doubly r	obust m	nethods																		
Model spe	cification	R-T	rue, PM-	-True																
GPPP	-0.329	9.828	93.452	39.581	1.027	-0.491	11.391	92.361	45.621	1.022	-0.279	10.520	93.948	42.782	1.037	-0.347	8.042	93.552	31.670	1.005
PSPP	-0.331	9.825	93.552	39.571	1.028	-0.497	11.397	92.460	45.574	1.021	-0.275	10.517	93.849	42.718	1.036	-0.353	8.030	93.452	31.655	1.006
LWP	-0.318	9.819	93.452	39.545	1.027	-0.468	11.408	92.560	45.582	1.020	-0.278	10.516	93.849	42.698	1.036	-0.249	8.029	93.552	31.664	1.006
AIPW	-0.344	9.827	93.254	39.557	1.027	-0.500	11.402	92.560	45.548	1.020	-0.313	10.522	93.651	42.663	1.034	-0.380	8.020	93.651	31.635	1.007
Model spe	cification	CR-T	ue, PM-	-False																
GPPP	-0.243	9.885	93.651	39.576	1.021	-0.446	11.315	92.758	45.663	1.030	-0.181	10.569	93.948	42.585	1.028	-0.248	8.029	93.452	31.783	1.010
PSPP	-0.293	9.889	93.552	39.552	1.020	-0.514	11.319	92.758	45.628	1.029	-0.272	10.575	93.948	42.539	1.026	-0.368	8.036	93.552	31.802	1.010
LWP	-0.306	9.892	93.651	39.546	1.020	-0.523	11.309	92.659	45.595	1.029	-0.305	10.573	93.849	42.463	1.024	-0.290	8.028	93.651	31.771	1.010
AIPW	-0.288	9.902	93.452	39.494	1.017	-0.476	11.304	92.659	45.534	1.028	-0.289	10.563	93.750	42.418	1.024	-0.422	8.042	93.651	31.768	1.009
Model spe	cification	1: QR-Fi	ulse, PM	-True																
GPPP	-0.027	8.255	93.651	33.254	1.027	-0.268	8.722	93.552	34.618	1.012	-0.050	8.168	93.948	32.870	1.026	-0.187	7.194	94.048	28.743	1.019
PSPP	0.664	8.282	93.552	33.140	1.024	0.324	8.716	93.353	34.499	1.010	0.514	8.163	94.345	32.827	1.027	0.297	7.201	93.948	28.654	1.016
LWP	-0.052	8.262	93.750	33.260	1.026	-0.296	8.754	93.552	34.672	1.010	-0.084	8.174	93.948	32.824	1.024	-0.211	7.029	94.841	28.203	1.024
AIPW	-2.397	13.400	94.048	53.981	1.044	-2.748	14.221	93.948	57.030	1.042	-2.459	13.296	94.742	53.741	1.049	-2.081	10.389	94.345	42.436	1.063
Model spe	cification	1: QR-F	Alse, PM	-False																
GPPP	30.818	31.348	0.099	22.751	1.010	29.492	30.201	0.496	26.142	1.025	28.797	29.405	0.298	24.074	1.032	29.590	29.893	0.000	16.838	1.012
PSPP	30.815	31.347	0.099	22.752	1.008	29.486	30.196	0.496	26.129	1.023	28.792	29.401	0.298	24.082	1.032	29.580	29.879	0.000	16.685	1.009
LWP	30.840	31.366	0.099	22.408	0.999	29.484	30.195	0.397	26.043	1.020	28.791	29.397	0.198	23.909	1.027	29.853	30.103	0.000	15.380	1.014
AIPW	30.885	31.393	0.099	22.123	1.003	29.522	30.224	0.298	25.858	1.018	28.836	29.433	0.198	23.739	1.026	29.903	30.143	0.000	15.305	1.027
UW: U	nweight€	∋d; FW:	Fully w	sighted;	GPPP:	Gaussia	n Proces	s of Pro	pensity	Predict	ion; PSP	P: Pena	lized Sp	line of I	ropensi	ity Predi	ction; L	WP: Lii	iear-in-v	veight

2 3 ÷ Ś 5, prediction; AIPW: Augmented Inverse Propensity Weighting;

UW	AIPV	LWP	PSPI	GPP	Model :	AIPV	LWP	PSPI	GPP	Model :	AIP	LWP	PSPI	GPP	Model :	AIPV	LWP	PSPI	GPP	Model :	Doubly	IPSV	Model :	IPSV	Model :	Non-ro	FW	Non-p	FW FW	Probal	Measu	
Unwei	V 31.1	31.0	31.0	P 31.0	specifica	V -2.5.	-0.1	1.29	P -0.0	specifica	V -0.1	-0.1	-0.1	P -0.0	specifica	V -0.3	-0.2	-0.3	P -0.3	specifica	v robus	V = 31.0	specifica	V -2.2.	specifica	bust n	$^{44.0}_{0.18}$	robabil	-41.	oility sa	re rBia	
ghted; I	11 31	60 31	76 31	65 31	tion: Q	59 14	13 8.8	8.8	74 8.8	tion: Q	80 10	89 10	61 10	22 10	tion: Q	34 10	41 10	12 10	04 10	tion: Q	t meth	49 31	tion: Q	43 11	tion: Q	nethod	8 5.9 5.9	ity san	$\begin{array}{ccc} 047 & 41 \\ 35 & 10 \end{array}$	ample	ıs rM	
FW: F	.865 0	.825 0	.846 0	.834 0	R–Fals	.081 9	312 9	384 9	302 9	R-Fals	.278 9	.269 9	.277 9	.274 9	R-Tru	.246 9	.233 9	.254 9	.252 9	R-Tru	ıods	.894 1	R–Fals	.249 9	R-Tru		.904 U	iple (	.487 0 .901 9	$(S_R)$	ISE c	
ully we	.595	.595	.595	.595	e, PM-	5.139	3.056	3.750	3.353	e, PM-	3.155	3.552	3.452	3.552	e, PM-	4.147	4.048	3.849	3.948	e, PM−		.290	е	2.659	(J)		5.238	$S_A)$	3.552		rCI	NI,
eighted;	26.638	26.790	27.108	27.121	-False	57.536	34.873	34.903	34.932	-True	40.695	40.775	40.787	40.822	False	40.796	40.783	40.855	40.870	-True		28.822		45.293			22.900 23.269	200	$\frac{19.213}{42.588}$		lCI	
GPPP	0.986	0.985	0.993	0.994		1.060	1.009	1.013	1.012		1.010	1.013	1.012	1.013		1.016	1.016	1.016	1.017			1.008		1.048			1.004	0000	$0.813 \\ 0.997$		rSE	
: Gaussia	29.743	29.720	29.738	29.748		-2.702	-0.168	1.242	-0.108		-0.327	-0.403	-0.368	-0.232		-0.340	-0.239	-0.309	-0.278			29.561		-2.251			40.010 0.094	20010	-44.574 -0.575		rBias	
n Proce	30.769	30.751	30.764	30.772		15.082	9.391	9.401	9.361		11.655	11.701	11.703	11.714		11.717	11.747	11.731	11.735			30.735		11.567			$\begin{array}{c} 40.004\\ 6.541 \end{array}$		$45.028 \\ 12.433$		rMSE	
ss of Pr	2.877	2.877	2.877	2.679		94.643	93.750	94.742	94.147		93.056	93.254	93.353	93.353		94.147	94.147	94.246	94.246			6.151		92.758			94.742	0000	$0.000 \\ 92.560$		crCI	CUB
opensit	31.144	31.231	31.423	31.444		61.474	37.675	37.718	37.706		47.403	47.504	47.627	47.716		47.183	47.290	47.293	47.397			33.381		46.382			20.997 25.533	2001	20.730 50.551		lCI	
y Predic	1.008	1.009	1.017	1.019		1.056	1.023	1.032	1.027		1.037	1.036	1.038	1.039		1.027	1.027	1.028	1.030			1.011		1.042			0.995	000	$     \begin{array}{r}       0.828 \\       1.038     \end{array}   $		rSE	
tion; PS	28.847	28.806	28.818	28.823		-2.527	-0.169	1.110	-0.099		-0.312	-0.367	-0.276	-0.065		-0.487	-0.373	-0.401	-0.374			28.698		-2.284			43.260 0.038	10 001	-42.945 -0.570		rBias	
PP: Pen	29.804	29.764	29.779	29.783		14.254	8.795	8.819	8.807		10.815	10.855	10.853	10.868		10.837	10.819	10.822	10.829			29.785		11.271			43.783 6.316	10 10 1	$43.406 \\ 11.625$		rMSE	
alized S	2.877	2.480	2.579	2.778		93.849	94.742	94.742	94.444		94.048	93.849	93.849	93.651		94.444	94.643	94.940	95.040			5.060		92.262			95.734	0000	$0.000 \\ 93.254$		crCI	EXP
pline of	28.960	29.036	29.258	29.269		57.570	35.309	35.405	35.413		43.809	43.904	44.023	44.175		44.240	44.304	44.352	44.445			31.006		44.587			23.219 24.616	2020	$20.084 \\ 47.208$		lCI	
Propen	0.985	0.988	0.994	0.995		1.046	1.024	1.032	1.025		1.033	1.032	1.035	1.036		1.042	1.045	1.046	1.047			0.992		1.030			$0.974 \\ 0.994$	201	$     \begin{array}{r}       0.812 \\       1.037     \end{array} $		rSE	
sity Pred	30.078	30.003	29.783	29.821		-2.289	-0.325	0.675	-0.271		-0.372	-0.108	-0.291	-0.030		-0.349	-0.021	-0.297	-0.276			30.014		-2.216			0.156	22	-34.898 -0.326		rBias	
liction; ]	30.482	30.404	30.231	30.265		11.031	7.269	7.426	7.442		8.203	8.174	8.197	8.185		8.213	8.216	8.219	8.244			30.449		10.028			5.027	222	$35.315 \\ 8.522$		rMSE	
LWP: Li	0.000	0.000	0.000	0.000		94.048	93.651	93.948	93.452		93.452	93.948	93.651	93.651		93.353	93.750	93.452	93.254			0.000		93.155			95.337	0000	0.000 94.544		crCI	SIN
inear-in-	20.037	19.992	20.989	20.999		45.025	29.004	29.415	29.490		32.545	32.588	32.673	32.685		32.233	32.302	32.270	32.312			20.837		40.735			10.343 20.785	10010	$17.395 \\ 33.469$		lCI	
-weight	1.033	1.035	1.032	1.037		1.064	1.018	1.014	1.011		1.013	1.017	1.017	1.018		1.002	1.003	1.002	1.000			1.036		1.062			1.037 1.055		$     \begin{array}{r}       0.820 \\       1.002     \end{array} $		rSE	

Table
e 10
С
mpar
ing t
he ]
perfe
orm
ance
of
the
bias
ad
just
mei
at n
ıetk
lods
in
the
$\sin$
ula
tion
stu
ldy i
for 1
the
bin
ary
out
com
le w
ith
$(n_A$
$, n_{R}$
<u> </u>
(500)
), 50
0) a
$\gamma_1$
=
0.3

prediction; AIPW: Augmented Inverse Propensity Weighting;

**Table 11:** Comparing the performance of the bias adjustment methods in the simulation study for the *binary* outcome with  $(n_A, n_R) = (500, 1, 000)$  and  $\gamma_1 = 0.6$ 

Measure	rBias	rMSE	LIN crCI	ICI	rSE	rBias	rMSE	CUB crCI	ICI	rSE	rBias	rMSE	EXP or CI	ICI	rSE	rBias	rMSE	SIN $crCI$	ICI	rSE
Probabili UW	tty sami -40.760	$\frac{\log (S_R)}{41.085}$	0.000	13.610	0.673	-44.127	44.513	0.000	14.703	0.642	-42.643	43.014	0.000	14.229	0.644	-34.513	34.804 7.11	0.000	12.322	0.700
F.W	0.256	9.389	92.758	37.069	1.007	0.371	11.344	93.254	45.111	1.014	0.101	10.820	92.758	41.488	0.978	0.095	7.111	93.849	28.415	1.019
Non-prot	ability	sample	$(S_A)$																	
MU	86.387	86.546	0.000	20.970	1.019	93.225	93.459	0.000	26.157	1.011	86.223	86.433	0.000	24.102	1.021	63.287	63.420	0.000	16.349	1.015
ΕW	0.131	7.764	95.238	30.716	1.009	0.046	8.475	94.841	32.731	0.985	-0.118	8.148	94.147	32.023	1.002	-0.025	7.263	95.734	28.730	1.009
Non-robu	ist meth	por																		
Model spe	cification	: QR-Th	en.																	
IPSW	-3.227	13.982	93.353	56.871	1.066	-3.152	14.463	92.460	57.679	1.042	-3.082	13.955	92.758	55.562	1.041	-3.285	12.805	92.956	52.146	1.074
Model spe	cificatior.	1: QR-F	ulse																	
IPSW	53.562	54.137	0.000	31.653	1.025	50.993	51.855	0.000	37.317	1.010	49.050	49.812	0.000	34.351	1.009	51.063	51.358	0.000	21.820	1.012
Doubly r	obust n	lethods																		
Model spe	cification	1: QR-Th	.ue, PM-	-True																
GPPP	0.337	9.267	94.742	37.662	1.037	0.357	11.473	93.254	44.914	0.999	0.391	10.659	94.444	42.397	1.015	0.313	7.382	93.056	28.939	1.000
PSPP	0.307	9.265	94.742	37.616	1.036	0.312	11.436	93.254	44.827	1.000	0.383	10.639	94.742	42.158	1.011	0.277	7.361	93.056	28.724	0.996
LWP	0.887	9.303	95.139	37.769	1.040	0.679	11.588	93.452	45.211	0.997	0.563	10.697	94.544	42.513	1.015	2.122	7.827	93.552	29.736	1.006
AIPW	0.242	9.280	94.940	37.601	1.033	0.147	11.439	93.254	44.799	0.999	-0.006	10.682	94.444	42.023	1.003	0.133	7.387	92.956	28.793	0.994
Model spe	cification	1: QR-Th	.ue, PM-	-False																
GPPP	1.021	9.354	94.643	38.386	1.053	0.790	11.546	92.857	45.111	0.999	0.739	10.591	94.444	42.842	1.034	1.411	7.633	93.948	29.422	1.000
PSPP	0.469	9.290	94.147	38.002	1.044	0.322	11.365	93.056	44.503	0.999	0.285	10.541	94.147	42.234	1.022	0.298	7.440	93.452	29.029	0.996
LWP	0.282	9.303	94.048	37.791	1.036	0.296	11.384	92.857	44.395	0.995	0.206	10.517	94.048	41.911	1.016	0.650	7.450	94.048	29.049	0.998
AIPW	0.335	9.278	93.948	37.612	1.034	0.472	11.339	93.155	44.319	0.997	0.396	10.522	93.750	41.799	1.014	0.072	7.418	93.651	28.780	0.989
Model spe	cificatior	1: QR-F	ulse, PM	-True																
GPPP	0.279	7.814	95.635	32.201	1.051	0.515	9.122	94.048	35.787	1.002	0.337	8.482	94.742	34.458	1.037	0.259	6.655	93.552	26.511	1.017
PSPP	4.350	9.046	93.948	33.320	1.071	4.859	10.448	92.857	37.065	1.022	4.419	9.750	94.048	35.700	1.047	2.881	7.177	93.155	26.425	1.025
LWP	0.031	7.792	95.238	31.935	1.045	0.010	9.015	93.849	35.327	0.999	-0.070	8.411	94.544	33.866	1.027	0.116	6.618	93.353	26.025	1.003
AIPW	-3.794	17.578	94.940	70.650	1.050	-4.091	19.231	94.246	75.598	1.026	-3.969	17.898	95.536	70.722	1.033	-3.231	13.525	94.345	54.283	1.054
Model spe	cificatior.	1: QR-F	ulse, PM	-False																
GPPP	46.258	46.727	0.000	26.958	1.041	43.406	44.150	0.000	31.503	0.995	42.182	42.813	0.000	29.504	1.027	46.251	46.541	0.000	20.569	1.011
PSPP	46.229	46.695	0.000	26.939	1.043	43.369	44.112	0.000	31.478	0.995	42.150	42.779	0.000	29.486	1.028	46.153	46.445	0.000	20.470	1.004
LWP	46.320	46.787	0.000	26.826	1.038	43.358	44.107	0.000	31.488	0.992	42.145	42.777	0.000	29.429	1.024	48.246	48.528	0.000	20.352	0.994
AIPW	46.840	47.297	0.000	26.698	1.038	43.745	44.481	0.000	31.341	0.992	42.734	43.354	0.000	29.297	1.022	50.152	50.439	0.000	21.149	1.003
TIVAV- II,	minimpt o	N. FW.	Eully w	sichtod.	CPPP.	Cancera	Droces	s of Duc	maneity	Dradio	HON. DCD	D. Done	lized Sr	line of l	Dronone	ity Dradi	ction. I	W/D. L	- ui-reor	moiaht
	ULV	JU, L VV	т. чт. т.	UBUUCU,		Preenport		17 7 70 60	νιστολ	T TCOTT	1011, 1 MT	די ד בייים	In north		erradat :	nna i raan	CUUUII, -	· · · · ·	1_TCOT _ TT - 1	VEISITU
preatct.	10n; Alf	W: Aug	mentea	Inverse	Propen	ISITY WEL	ghung;													

UW: U predict	AIPW	LWP	PSPP	GPPP	Model spe	AIPW	LWP	PSPP	GPPP	Model spe	AIPW	LWP	PSPP	GPPP	Model spe	AIPW	LWP	PSPP	GPPP	Model spe	Doubly r	IPSW	Model spe	IPSW	Model spe	Non-rob	FW	UW.	Non-pro	FW	IIW	Probabil	Measure		
nweight, ion; AIF	46.885	46.391	46.282	46.307	cification	-4.750	-0.093	2.302	0.021	cification	-0.210	-0.258	-0.165	0.112	cification	-0.313	-0.010	-0.276	-0.261	cification	obust n	52.198	cification	-3.783	cification	ist met	0.203	84.864	pability	-0.326	_A1 047	ity sam	rBias		
ed; FW: W: Aug	47.282	46.808	46.703	46.727	1: QR–Fa	18.998	8.411	8.643	8.403	1: QR–Fa	9.956	9.942	9.926	9.911	1: QR–T	9.942	9.904	9.928	9.930	1: QR–T	nethods	52.687	1: QR–Fa	15.498	1: QR–Ti	hod	5.406	84.948	sample	10.847	41 487	ple $(S_R)$	rMSE		
Fully w gmented	0.000	0.000	0.000	0.000	alse, PM	94.742	93.651	94.841	93.849	alse, PM	93.552	94.345	94.444	94.544	rue, PM-	94.544	94.940	94.643	94.643	rue, PM-		0.000	alse	90.179	rue		95.536	0.000	(S )	93.552		-	crCI	LIN	
eighted; Inverse	24.089	24.587	24.788	24.810	–False	77.083	33.554	33.926	33.606	-True	39.995	40.075	40.142	40.299	-False	40.021	39.981	40.018	40.048	-True		28.997		62.509			21.575	14.914		42.776	10 913		lCI		
GPPP: Propens	1.005	1.005	1.009	1.012		1.069	1.017	1.038	1.020		1.025	1.028	1.031	1.037		1.027	1.029	1.028	1.029			1.032		1.061			1.018	1.009		1.006	0 813		rSE		
Gaussia sity Wei	43.817	43.422	43.452	43.480		-5.189	-0.196	2.340	0.028		-0.298	-0.475	-0.429	-0.195		-0.535	-0.272	-0.482	-0.448			49.275		-3.878			0.015	91.029		-0.675	-44 574		rBias		
n Proces ghting;	44.373	43.994	44.019	44.046		20.238	8.926	9.163	8.886		11.469	11.489	11.490	11.524		11.589	11.604	11.562	11.556			50.008		15.620			5.762	91.148		12.615	45 098		rMSE		
ss of Pro	0.000	0.000	0.000	0.000		95.139	93.948	93.948	94.246		93.849	93.849	93.552	94.048		93.452	93.552	93.452	93.552			0.000		89.782			95.933	0.000		92.758	000		crCI	CUB	
pensity	28.079	28.379	28.360	28.364		81.947	35.751	36.205	35.807		46.334	46.395	46.458	46.627		46.411	46.460	46.369	46.462			34.156		62.778			22.988	18.565		51.013	01230		ICI		
Predicti	1.023	1.023	1.026	1.027		1.068	1.021	1.042	1.027		1.030	1.031	1.032	1.032		1.022	1.021	1.024	1.026			1.021		1.058			1.017	1.016		1.033	868 U		rSE		
on; PSP	43.130	42.579	42.547	42.571		-4.784	-0.119	2.233	0.084		-0.230	-0.297	-0.299	-0.101		-0.346	-0.072	-0.172	-0.204			47.956		-3.599			0.090	84.651		-0.509	-49 045		rBias		
P: Pena	43.613	43.081	43.049	43.072		18.933	8.295	8.580	8.297		10.601	10.567	10.611	10.628		10.660	10.646	10.636	10.641			48.604		15.118			5.638	84.761		11.780	43 406		rMSE		
lized Sp	0.000	0.000	0.000	0.000		94.643	94.345	94.643	94.643		94.147	93.948	93.849	93.750		94.444	94.643	94.643	94.444			0.000		90.675			95.437	0.000		93.353	000		crCI	EXP	
line of F	25.882	26.285	26.356	26.363		77.221	33.557	34.188	33.793		42.867	42.907	43.080	43.280		42.987	43.106	43.046	43.177			31.095		60.239			22.508	17.102		47.385	00 084 G		ICI		
ropensit	1.020	1.022	1.026	1.027		1.075 .	1.032 .	1.052	1.039 -		L.031 ·	1.036 .	1.036 .	1.038		1.029 .	1.032	1.032 .	1.035 .			1.002		1.046 .			1.018	1.007		1.027	1 8 1 9		SE		
zy Predi	49.826	48.301	46.336	46.321		-3.902	-0.269	1.294	-0.331		-0.412	-0.094	-0.240	0.391		-0.396	0.617	-0.314	-0.335			50.472		-3.782			0.231	62.735		-0.403	3/ 808		rBias		
ction; L <sup>v</sup>	49.990	48.464	46.552	46.541		14.639	7.155	7.406	7.326		8.136	8.127	8.113	8.149		8.128	8.144	8.105	8.120			50.655		14.133			5.039	62.802		8.536	27 21 7		rMSE		
WP: Lin	0.000	0.000	0.000	0.000		92.560	92.659	93.452 :	93.452		93.651 :	93.849 :	94.147 :	93.651 :		93.353	94.345 :	93.452 :	93.056			0.000		91.964			95.337	0.000		93.849			crCI	SIN	
ear-in-w	16.533 i	16.010	17.919 ;	18.121		50.317 i	28.604	29.067 1	29.197		31.864	31.970	31.929	32.036 i		32.166	32.321	32.015	32.132			17.591		57.700			20.205 i	11.602		33.518 1	17 305 1		lCI		
reight	1.041	1.029	1.018	1.023		090	1.020	1.016	1.017		1.000	1.003	1.004	1.004		1.010	1.015	1.008	1.010			1.042		1.080			1.023	.017		1.002	0081		SE	ſ	

Table 12: Comparing the performance of the bias adjustment methods in the simulation study for the binary outcome with  $(n_A, n_R) = (1, 000, 500)$  and  $\gamma_1 = 0.6$ 

9.
0 =
3
, nc
6
00
Ó.
50
-
$n_{R}$
Α,
<i>u</i> )
$_{\mathrm{th}}$
.W
ne
[]
ut
y o
ar
iin
ie ł
$^{\mathrm{th}}$
$\mathbf{for}$
Jy.
ng
1 ST
ioi
lat
nu
sii
he
n t
s i
lod
eth
Ĥ
'nt
me
ıstı
dju
3 ac
ia£
еþ
$^{\mathrm{th}}$
$\mathbf{of}$
lce
ıar
rn
rfc
$\mathbf{pe}$
he
g t
cin
paı
lm
S
ŝ
÷,
ole
at

$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	LIN ISE crCl ICl 1 (SP)	SE rBia	s rMSE	CUB crCI	ICI	rSE	rBias	rMSE d	EXP STCI	ICI	rSE	rBias	rMSE	SIN crCI	ICI	rSE
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		0.813 -44.5 0.995 -0.62	$\begin{array}{rrr} 74 & 45.028 \\ 0 & 12.474 \end{array}$	$0.000 \\ 92.560$	20.730 50.979	0.828 1.043	-42.945 -0.508	43.406 ( 11.640 9	0.000 93.750	20.084 47.428	$0.812 \\ 1.040$	-34.898 -0.302	$35.315 \\ 8.481$	$0.000\\93.651$	17.395 33.327	$0.820 \\ 1.003$
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$																
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		0.985 92.80 $0.999 0.100$	59 93.119 5 8.183	0.000 96.032	26.171 32.740	$0.979 \\ 1.020$	$86.198 \\ 0.146$	86.428 ( 8.318 3	0.000 93.948	24.101 32.044	$0.977 \\ 0.982$	63.186 0.248	63.318 7.226	0.000 95.139	16.359 28.741	1.019 1.015
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$																
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$																
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		1.082 - 5.19	2 17.348	90.873	69.336	1.068	-5.013	16.767	90.079	66.988	1.068	-5.259	15.947	91.270	63.328	1.073
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$																
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		1.018  49.30	39 50.447	0.397	40.870	1.005	47.969	48.914 (	0.099	37.609	1.002	50.477	50.805	0.000	23.045	1.019
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$																
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	I															
$ \begin{array}{[c]ccccccccccccccccccccccccccccccccccc$		1.021 - 0.19	5 12.013	93.651	48.898	1.038	0.067	11.263	94.048	45.943	1.040	-0.094	8.322	94.048	33.008	1.011
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		1.020 -0.25	3 12.007	92.956	48.684	1.034	0.042	11.238	94.643	45.765	1.038	-0.151	8.278	93.849	32.812	1.011
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		1.028 0.22	12.129	93.849	49.125	1.033	0.310	11.319	94.048	46.165	1.040	1.629	8.595	93.750	33.723	1.019
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		1.022 -0.40	1 12.071	93.254	48.666	1.029	-0.318	11.284	94.345	45.620	1.031	-0.274	8.394	93.452	33.054	1.005
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$																
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		1.038 0.203	3 12.135	93.452	49.055	1.031	0.345	11.208	93.948	46.329	1.054	0.882	8.415	93.353	33.316	1.015
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		1.030 -0.26	6 12.046	93.651	48.487	1.027	-0.131	11.131	93.750	45.665	1.046	-0.187	8.351	92.956	32.960	1.007
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		1.023 -0.30	3 12.042	93.353	48.357	1.024	-0.183	11.103	93.849	45.407	1.043	0.162	8.377	93.155	33.080	1.007
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	-	1.017 -0.07	5 12.003	93.254	48.215	1.024	-0.019	11.115	93.948	45.235	1.038	-0.407	8.372	92.560	32.762	0.999
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$																
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	<u> </u>	1.030 0.26	7 9.903	94.643	39.802	1.025	0.346	9.341	94.246	37.453	1.023	-0.179	7.651	93.750	30.244	1.008
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	-	1.070 4.63	2 11.070	94.048	40.979	1.039	4.346	10.435	93.452	38.781	1.042	2.419	7.961	93.849	30.197	1.015
$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$		1.025 -0.20	2 9.907	94.147	39.346	1.013	-0.022	9.262	93.948	36.989	1.018	-0.209	7.591	93.651	29.763	1.000
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	-	1.092 -6.91	3 23.067	95.933	93.312	1.081	-6.310	21.508	95.536	87.184	1.081	-5.406	16.466	92.262	66.288	1.087
$\begin{array}{cccccccccccccccccccccccccccccccccccc$																
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0	0.991 43.8	3 44.673	0.000	34.430	1.006	42.785	43.578 (	0.099	32.147	0.990	46.636	46.972	0.000	22.280	1.013
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0	0.989  43.7	79 44.640	0.000	34.406	1.006	42.753	43.547 (	0.099	32.117	0.989	46.536	46.875	0.000	22.174	1.005
0.985  44.101  44.957  0.000  34.142  0.997  43.241  44.019  0.099  31.741  0.983  49.864  50.171  0.000  21.994  1.012  0.982  49.864  50.171  0.000  21.994  1.012  0.982  0.982  0.983  0.983  0.984  0.983  0.983  0.984  0.983  0.983  0.984  0.983  0.984  0.983  0.984	~	0.984  43.73	58 44.630	0.000	34.367	0.998	42.752	43.549 (	D.099	32.022	0.985	48.424	48.721	0.000	21.085	1.002
	~	0.985 44.10	11 44.957	0.000	34.142	0.997	43.241	44.019 (	0.099	31.741	0.983	49.864	50.171	0.000	21.994	1.012

TABLE 14

#### 8.2. Supplemental results on CIREN/CDS data

Percentage of severe injury in HEAD and associated 95% CIs by different covariates across DR adjustment methods Covariate  $\mathbf{n}$ Unweighted (95% CI) GPPP (95% CI) LWP (95% CI) AIPW (95% CI) 25.036(23.964,26.108) 10.36(4.559,16.161) 9.658(6.398,12.918) Total 6.2719.552(7.33, 11.774)Gender 28.359(26.805,29.914) Male 3.23011.464(7.982.14.947)12.526(4.69, 20.361)10.81(6.382.15.239) 21.506(20.046,22.966) 8.245(5.697,10.794) 8.882(3.84,13.924) 8.918(4.303,13.533) Female 3.041Age group 659 32.625(29.046, 36.205)9.498(4.117,14.878) 10.874(0.606, 21.143)8.557(0, 18.879)16 - 1910.137(4.987,15.286) 20 - 392.73211.027(5.256.16.799) 26.354(24.702.28.006)10.292(7.258, 13.326)40-641.94620.863(19.058.22.669)8.527(5.688, 11.365)9.442(2.439.16.445)9.222(3.92, 14.525)65 +93424.518(21.759,27.277) 9.999(6.371,13.626) 10.43(5.369, 15.491)10.024(1.537, 18.511)Occupant role Driver 1,547 32.515(30.18,34.849) 9.681(4.677, 14.685)10.958(0.535, 21.382)9.924(0, 20.727)Passenger 4,72422.587(21.394,23.779) 9.594(7.445,11.743) 10.296(5.322, 15.27)9.614(6.267,12.961) Seating Row 24.056(22.961,25.151) 9.4(7.326, 11.474)9.41(6.121,12.699) Front 5.85310.125(4.791, 15.458)38.756(34.086,43.426) 12.632(0.87,24.394) 15.661(0, 36.448)14.995(3.168,26.822) Rear 418Restraint use Restraint 2,275 32.967(31.035,34.899) 15.351(9.796,20.906) 17.135(4.552, 29.718)14.327(7.613, 21.042)Unrestraint 3,996 20.521 (19.268,21.773) 7.01(4.926,9.094)  $7.346(\dot{4}.113, \dot{1}0.579)$ 7.34(3.457,11.222) Air bag Deployed  $2.194 \quad 30.447(28.521.32.372)$ 13.067(0.993.25.14)13.113(6.087.20.138)10.956(5.005.16.907)4.07722.124(20.85,23.398) 8.995(6.466.11.524) 7.804(3.654, 11.955)Not deployed 9.057(6.161, 11.953)Injury scale (AIS) 1.97712.903(8.281,17.525) 3 18.412(16.703,20.12) 13.609(9.765, 17.453)13.61(8.5, 18.719)4 761 55.979(52.452,59.506) 31.675(24.511, 38.839)31.549(23.938, 39.161)33.75(21.739, 45.761) 5+78176.056(73.063,79.049) 59.405(41.668,77.143) 59.683(37.787,81.578) 59.896(39.338,80.453) Days hospitalized 2,25926.737(24.912,28.563) 2.633(0, 11.319)3.842(0, 16.924)4.406(0, 18.412)0 1 - 31,757 17.758(15.971,19.544) 8.564(4.978,12.15) 8.619(4.918,12.32) 9.408(5.25,13.566) 4 - 720.902(18.555,23.249) 10.174(6.976,13.371) 9.981(6.845,13.117) 10.826(6.88,14.771) 1,1538-14 64028.438(24.943,31.932) 16.411(11.916, 20.907)16.313(11.886,20.74) 17.117(11.688,22.547) 15 +46250(45.441, 54.559)27.032(21.157,32.907) 26.964(21.18,32.747) 23.857(17.808, 29.905)Delta-v (Km/h) 404 Minor 14.109(10.714.17.503)9.347(0.209.18.485)11.878(0.26.253) 17.364(1.382,33.346) 19.613(18.259,20.967) Moderate 3,3047.748(5.154,10.343) 8.597(2.33, 14.864)6.326(2.517, 10.135)2,56333.75(31.919,35.58) 15.234(11.167,19.301) 15.395(10.571, 20.219)15.357(8.964, 21.749)Severe Damage distribution 23.148(21.937,24.359) 8.304(6.187,10.421) 9.104(3.819, 14.389)7.276(3.906, 10.647)Wide 4.657Narrow 320 27.812(22.903,32.722) 10.447(1.981, 18.912)11.33(0,23.056) 11.031(4.628,17.434) 58420.205(16.949.23.462)12.282(7.027, 17.537)12.667(6.537.18.798)14.807(3.911.25.702) Corner 17.071(3.194,30.949) Other 710 40.141(36.535,43.746) 15.842(6.849.24.835)21.066(6.792,35.341) Rollover 4.942 21.833(20.681.22.985) 7.526(4.333.10.719) 8.566(6.467.10.665) 9.295(4.417.14.173)Yes  $1,329 \quad 36.945 (34.35, 39.54)$ No 12.579(6.805, 18.354)13.867(2.222,25.512) 17.555(8.204, 26.906)**Deformation** location 7.905(3.188,12.621) Front 3,671 16.78(15.571,17.989) 7.353(4.777, 9.93)6.853(2.596.11.11)Right 1,12332.591(29.85,35.333) 12.15(8.516,15.783) 13.916(3.853,23.979) 10.703(5.523,15.883) 857 39.44(36.168, 42.712) 13.239(1.592,24.887) 14.049(1.539,26.559) 16.421(2.379,30.463) Left Rear 620 40.323(36.461,44.184) 15.445(7.088,23.801) 16.007(4.144,27.869) 17.129(2.342,31.916) Model year <2003 2.52928.035(26.284,29.785) 13.405(8.529,18.282) 14.961(2.984,26.938) 17.958(10.59.25.325)  $\overline{2004}$ -2007 2,37724.443(22.715,26.17) 10.117(7.061,13.174) 10.558(5.971,15.146) 7.776(4.489, 11.063)2008-2011 1.019 20.51(18.031,22.989) 3.33(0.8.253)3.656(0, 9.11)1.094(0, 7.874)2012 +34620.52(16.265,24.776) 8.831(3.093,14.569) 9.234(3.369,15.1) 3.904(0, 11.404)Vehicle make 3,63525.777(24.355,27.199) 10.091(6.99, 13.192)11.175(3.082, 19.268)9.535(4.856,14.213) American 1,927 24.754(22.827,26.68) 9.997(6.843, 13.15)10.479(5.934, 15.025)12.601(7.211,17.991) Japanese 387 21.189(17.117,25.26) 5.661(0, 12.117)6.576(0, 16.798)5.023(0, 12.299)Korean Other 322 22.981(18.386.27.577) 5.801(0.13.635)5.954(0, 13.893)4.832(0, 11.801)Vehicle type 4.06325.966(24.618, 27.314)9.167(6.918, 11.415)9.928(4.629, 15.228)7.384(3.613.11.154)Passenger car 1.213 9.113(5.987,12.239) 9.52(3.705, 15.335)Light truck 22.341(19.997,24.685) 13.447(6.271, 20.623)27.241(22.117,32.365) Van SUV 2904.829(0,10.283)6.556(0, 17.997)7.793(0.18.304)705 16.67(6.355,26.985) 23.404(20.279.26.53)14.224(9.007.19.441)15.617(5.833,25.401)

19

TABLE 15

Percentage of severe injury in ABDOMEN and associated 95% CIs by different covariates across DR adjustment methods

Covariate	n	Unweighted (95% CI)	GPPP (95% CI)	LWP (95% CI)	AIPW (95% CI)
Total	6.271	9.919(9.179,10.658)	6.677(4.408.8.946)	7.621(1.93.13.313)	6.479(2.354, 10.604)
Gender	- , .				
Male	3.230	10.464(9.409.11.52)	7.694(4.134.11.253)	8.837(1.872, 15.802)	3.552(0.11.317)
Female	3.041	9.339(8.305,10.373)	5.931(3.485.8.376)	6.724(1.411.12.036)	8.852(3.96,13.744)
Age group	0,0				
16-19	659	11.836(9.37.14.302)	6.592(2.583.10.601)	7 886(0 16 685)	0(0.22.12)
20-39	2.732	10.505(9.355.11.655)	6.8(4.066.9.535)	7.782(1.873.13.691)	7539(155313524)
40-64	1.946	8 941(7 674 10 209)	7.109(4.012, 10.205)	8 157(1 388 14 926)	7 149(3 55 10 749)
65+	934	8.887(7.062.10.711)	5.75(2.911.8.589)	6.179(2.056.10.301)	8.945(0.561.17.329)
Occupant role					
Driver	1.547	12.864(11.195.14.532)	8.85(4.445.13.255)	10.308(1.336.19.28)	13.363(2.809.23.917)
Passenger	4.724	8.954(8.14.9.768)	6.1(3.73.8.47)	6.921(1.726.12.116)	4.778(0.9.568)
Seating Bow	-,	0.0001(0.11,01100)	011(0110,0111)	0.021(1120,121110)	
Front	5.853	9 465(8 715 10 215)	6414(4287,8541)	7.28(2.052.12.507)	6.264(2.088,10.44)
Bear	418	$16\ 268(12\ 73\ 19\ 806)$	13.822(2.075.25.57)	16.764(0.38.476)	12.916(3.318.22.514)
Restraint use	110	101200(12110;101000)	101022(21010),20101)	101101(0,001110)	121010(01010,221011)
Restraint	2.275	11.648(10.33.12.967)	8 836(4 185 13 488)	10.761(0.22.227)	13,03(5,326,20,733)
Unrestraint	3 996	8 934(8 05 9 818)	5.629(3.583.7.675)	$6\ 103(2\ 64\ 9\ 565)$	2.976(0.8.312)
Air bag	0,000	0.001(0.00,0.010)	0.020(0.000,1.010)	0.100(2.01,0.000)	2.010(0,0.012)
Deployed	2 1 9/1	10 255(8 986 11 525)	5 091(0 10 404)	6.952(0.19.7)	0.928(0.8.901)
Not doployed	4 077	0.738(8.828.10.648)	7718(5486005)	7.007(5.302.10.512)	10.032(5.212.14.852)
Inium coole (A	4,011	9.156(8.828,10.048)	1.118(3.480,3.33)	1.907(3.302,10.312)	10:032(0:212,14:032)
njury scale (A	1.077	7 081 (5 951 8 919)	0 233(5 070 19 487)	0.28(5.232.13.327)	10.554(5.247.15.861)
1	761	21.682(18.754.24.61)	9.233(0.979,12.407) 97.838(18.134.37.549)	28.076(10.007.37.146)	24.053(13.553.36.352)
4 5.1	781	21.002(10.794,24.01) 23.201(20.086.36.506)	27.030(10.134, 37.042) 35.512(15.655.55.37)	27705(13.063.62.348)	24.905(15.005, 50.002) 31.665(5.388.57.042)
Deve hoopiteli	701 rod	33.291(29.980,30.390)	35.512(15.055,55.57)	37.703(13.003,02.348)	31.005(5.588,57.942)
	2 250	11.554(10.226.12.872)	12 206(4 864 21 747)	16 510(2 786 20 252)	6 87(0.26.651)
1 2	1,209	4.008(3.171.5.025)	3808(07787017)	3158(01326183)	3 120(0.835 5 424)
1-3	1,757	4.098(3.171, 3.023) 0.102(7.526.10.861)	7.787(4.405.11.160)	7.487(4.275, 10.508)	0.752(5.241.14.166)
4-7	640	15, 156(12, 278, 17, 024)	1272(847216068)	12107(959217671)	9.755(5.541, 14.100) 12.052(8.400.17.606)
0-14	469	18.615(15.066.22.164)	12.72(0.472,10.908) 17 104(11 228 22 05)	17.920(11.910.92.959)	16.720(0.864.92.615)
	402	18.015(15.000,22.104)	17.194(11.556,25.05)	17.855(11.815,25.858)	10.755(9.804,25.015)
Dena-v (Km/I	1)	4 202 (2 25 6 166)	6.288(1.47.11.206)	8 <del>7</del> 92(0 91 901)	2 851(0 10 225)
Madanata	404	4.208(2.25, 0.100) 5 72(4.028 6 512)	0.388(1.47,11.300)	5.725(0,21.201)	3.851(0,10.325)
Source	0,504	16.921(14.802.17.650)	4.55(2.512, 0.588) 11 707(8 409 15 011)	12.226(7.064.16.508)	(1.001, 7.329)
Demore distril	2,000	10.231(14.803,17.033)	11.707(8.402,15.011)	12.230(7.304,10.308)	11.988(2.975,21.005)
Wide	4.657	10.715(0.827.11.602)	6 078(4 720 0 217)	7 967(9 419 19 917)	6 087(1 455 12 510)
Norrow	320	10.713(9.827,11.003) 10.312(6.08,13.645)	6.750(0.837.12.681)	8 316(0 10 15)	7 17(0 17 036)
Corner	520	8 210(5 002 10 447)	0.112(2.785, 14.441)	0.510(0,19.15) 0.514(2.20,15,727)	0.046(1.052.16.120)
Other	710	5.219(5.992,10.447)	9.113(3.765,14.441)	4.08(0.12.124)	9.040(1.952,10.159)
Dellemen	710	5.915(4.18,7.051)	2.404(0,0.822)	4.08(0,13.134)	0.848(0,3.708)
Voc	4 0 4 9	0.006(0.16.10.822)	7.026(4.712.0.26)	7 864(9 674 19 054)	0.018/5.224.12.702)
Ne	4,942	9.990(9.10,10.032) 0.621(8.045.11.917)	5.026(0.276.0.677)	6522(0.15,004)	9.018(3.334,12.703)
Defermention le	1,329	9.031(8.043,11.217)	5.020(0.370,9.077)	0.332(0,13.904)	0(0,8:029)
Front	2 671	1 8 179(7 986 0 059)	7 344(4 002 0 795)	8 000 (3 232 12 787)	0 683(4 672 14 602)
Picht	1 1 1 9 9	0.112(1.200, 9.000) 14 604(12 539 16 660)	0.325(5.04.12.600)	10.078(1.525.20.491)	1 689(0 14 607)
Loft	1,120 957	14.004(12.006,10.009) 14.009(11.670.16.296)	4507(0.0712)	5.798(0.14.156)	5.494(0.19)
Deen	607	(4.002(11.079,10.320))	4.397(0,9.713)	3.728(0,14.130) 2.512(0.11.282)	5.454(0,12)
Model year	020	0.123(4.241,0.017)	2.200(0,0.002)	3.313(0,11.303)	1.410(0,0.101)
< 2002	9 590	10 874(0 661 19 087)	5 716(1 198 10 909)	6.679(0.16.791)	1.023(0.10.724)
2003	2,029	0.34(8.17.10.500)	8557(590111794)	8 084(3 895 14 149)	19 916(6 554 17 979)
2004-2007	2,377	3.34(0.17,10.309) 10.01(8.167.11.959)	5 080(1 648 10 22)	5.504(3.023,14.143) 7 077(9 546 11 607)	12.210(0.004,11.018) 6 405(0 13 18)
2006-2011	246	6.647(4.023.0.272)	0.009(1.040,10.00) 2.470(0.7.511)	3.11(0.8.306)	3 667(0 12 272)
Vehicle make	040	0.047(4.023, 9.212)	2.479(0,7.511)	3.11(0,8.300)	5.007(0,12.272)
A morizon	2 625	0.684(8.722.10.645)	6 709(2 842 0 56)	7 017(0 565 15 268)	7 191(0 779 19 594)
Innerican	1,000	5.004(0.722,10.040) 10.171(8.899.11.591)	7 5(4 520 10 471)	0.110(2.510.10.200) 0.110(2.510.10.710)	6.051(0.110,13.004)
Japanese	1,921	0.561(6.631.12.40)	1.0(4.029,10.4/1) 5.316(2.03.9.602)	6.110(3.310,12.719) 6.971(0.13.169)	0.951(2.257,11.045) 0.001(0.7.026)
Other	201 201	5.501(0.051, 12.49) 11 $401(8.007.14.074)$	3.667(0.61.6.799)	4 116(0 172 8 06)	1.600(0.4.707)
Vohiole toma	044	11.431(0.001,14.314)	5.007(0.01,0.725)	4.110(0.173,0.00)	1.003(0,4.101)
Pageonger and	4.062	11 140(10 189 19 117)	6 76(1 625 9 995)	7 677(2 215 12 020)	0.094(4.759.19.904)
Fassenger car	4,003	11.149(10.162, 12.117) 7 007(6 47 0 599)	0.70(4.000,0.000) 7 219(2 205 11 411)	2 159(1 200 15 029)	9.024(4.700,10.294)
Von	1,213	1.331(0.41,3.323) 8 976(5 105 11 447)	7.310(3.223,11.411) 2.850(0.7.441)	(1.100(1.209, 10.020))	5(0,13.002)
SUV	290 705	6 800(4 040 8 668)	2.009(0,1.441) 6 500(2 655 10 262)	7 666(0 15 495)	4.766(1.067.9.466)
50 V	705	0.009(4.949,0.000)	0.509(2.055,10.362)	1.000(0,10.400)	4.700(1.007,8.400)

TABLE 16

Percentage of severe injury in THORAX and associated 95% CIs by different covariates across DR adjustment methods

Covariate	n	Unweighted (95% CI)	GPPP (95% CI)	LWP (95% CI)	AIPW (95% CI)
Total	6.271	33.934(32.762.35.106)	21.087(14.925,27.248)	29.62(17.969.41.272)	21.39(18.238.24.541)
Gender	- ) -				
Male	3,230	37.337(35.669, 39.006)	23.605(17.945, 29.264)	31.925(20.364, 43.486)	21.954(18.66, 25.249)
Female	3,041	30.319(28.685,31.953)	19.353(12.307,26.399)	28.049(15.549,40.548)	21.074(16.628, 25.52)
Age group	,				
16-19	659	37.329(33.636,41.022)	17.62(6.253, 28.988)	34.815(9.195,60.436)	21.41(13.6, 29.219)
20-39	2,732	31.442(29.701,33.183)	15.872(8.609, 23.136)	22.863(11.708,34.017)	15.318(9.955, 20.681)
40-64	1,946	32.066(29.992,34.139)	22.877(16.442,29.312)	31.709(18.647,44.771)	21.864(16.993,26.734)
65 +	934	42.719(39.547,45.892)	35.715(29.256,42.175)	41.792(31.461,52.122)	38.108(31.117,45.098)
Occupant role		× · · · · ·			
Driver	1,547	41.047(38.596, 43.498)	25.589(15.135, 36.044)	39.822(19.233,60.411)	22.188(13.747, 30.629)
Passenger	4,724	31.605(30.279,32.93)	20.007(13.585,26.429)	27.096(16.702,37.489)	21.249(17.329,25.17)
Seating Row			· · · · · ·		
Front	$^{5,853}$	33.18(31.973, 34.386)	20.399(14.202, 26.596)	28.514(17.063, 39.964)	20.94(17.786, 24.095)
Rear	418	44.498(39.733, 49.262)	40.118(25.011, 55.225)	59.85(33.897, 85.803)	34.6(20.286, 48.914)
Restraint use					
Restraint	2,275	39.253(37.246, 41.259)	25.078(15.779, 34.377)	40.585(21.1,60.069)	28.13(21.873, 34.387)
Unrestraint	3,996	30.906(29.473, 32.339)	18.765(12.572, 24.958)	23.818(14.563, 33.074)	17.835(14.413, 21.256)
Air bag					
Deployed	2,194	36.372(34.359, 38.385)	22.703(12.088, 33.318)	38.464(17.814, 59.114)	23.947(17.058, 30.836)
Not deployed	4,077	32.622(31.183, 34.061)	20.517(14.113, 26.92)	24.339(14.383, 34.294)	20.319(15.73, 24.908)
Injury scale (A	AIS)				
3	1,977	42.742(40.561, 44.922)	44.551(39.909, 49.194)	45.565(41.221, 49.909)	44.497(38.612, 50.382)
4	761	72.405(69.229, 75.581)	71.412(63.902, 78.923)	71.434(63.938, 78.931)	74.925(64.879, 84.971)
5+	781	70.679(67.486, 73.871)	82.191(74.038,90.344)	83.318(73.489,93.148)	79.427(68.421, 90.433)
Days hospitali	$\mathbf{zed}$				
0	2,259	30.102(28.21, 31.993)	16.397(3.115, 29.679)	33.637(10.187,57.087)	20.958(15.035, 26.88)
1-3	1,757	24.986(22.961, 27.01)	16.58(10.727, 22.433)	16.916(10.974, 22.857)	15.234(10.013, 20.456)
4-7	1,153	35.386(32.626, 38.146)	30.08(23.956,36.203)	30.835(24.826,36.843)	34.479(27.631,41.326)
8-14	640	49.375(45.502,53.248)	41.973(34.31,49.636)	41.683(34.216, 49.15)	39.04(31.172,46.909)
15+	462	61.688(57.255, 66.121)	53.435(46.093, 60.777)	52.866(45.985, 59.748)	48.834(40.281,57.387)
Delta-v (Km/ł	1)				
Minor	404	19.059(15.229,22.889)	11.995(1.156,22.834)	30.193(8.357,52.029)	8.541(0,20.813)
Moderate	3,304	25.938(24.444,27.433)	17.681(10.649,24.712)	27.582(13.704,41.46)	16.779(13.611, 19.947)
Severe	2,363	40.380(44.035,48.317)	32.809(27.293,38.324)	34.293(28.012,40.575)	35.216(29.025,41.407)
Damage distri	bution	22 000(22 540 25 000)	01 069(14 011 07 014)	22 (24/12 042 20 200)	02 114(10 500 00 7)
Naman	4,007	35.900(32.340,35.200)	21.003(14.911,27.214) 27.706(10.020.26.564)	28.024(18.042, 39.200)	23.114(19.526,20.7)
Corpor	520	30.302(31.280,41.839) 36.884(32.388,20.470)	27.790(19.029, 30.304) 17.022(10.176.25.67)	37.404(23.902, 50.900) 24.828(10.202, 20.472)	22.302(10.103, 33.022) 15 681(5 005 25 458)
Other	710	20.864(23.288, 30.479) 28 722(25 140 42 216)	17.923(10.170,23.07) 22 155(10 401 22 82)	24.838(10.202,39.473) 28.604(15.207.62.161)	16.240(4.610.27.870)
Pollovon	710	38.732(33.149,42.310)	22.135(10.491,33.82)	38.094(15.227,02.101)	10.249(4.019,27.879)
Voc	4 0 4 2	22 436(31 131 33 741)	20 100(13 803 26 415)	27 518(16 031 38 105)	20 760(16 031 24 608)
No	1 320	39 503(36 875 42 132)	25.651(17.024.34.278)	39.077(20.706.57.449)	24.238(15.951,24.008)
Deformation		35.505(50.610,42.152)	20.001(11.024,04.210)	55.011(20.100,51.445)	24.200(10.000,02.022)
Front	3 671	26 832(25 399 28 265)	19.037(12.539.25.536)	25 505(14 472 36 537)	19 842(14 653 25 03)
Right	1.123	$45\ 77(42\ 856\ 48\ 684)$	30.62(20.887.40.352)	42.05(25.575.58.525)	30.289(22.563.38.016)
Left	857	$44\ 924(41\ 594\ 48\ 254)$	20.7(2.847.38.554)	30.117(7.802.52.431)	20 237(4 837 35 637)
Bear	620	39.355(35.509.43.2)	21.911(8.05.35.771)	37.303(12.58.62.027)	20.181(7.315.33.046)
Model year			(====,====)		
<2003	2.529	35.389(33.526.37.253)	24.05(16.137.31.962)	37.778(20.449.55.108)	22.382(17.842.26.922)
2004 - 2007	2.377	34.034(32.13.35.939)	22.589(15.129.30.049)	29.434(17.015.41.853)	24.151(17.576.30.727)
2008-2011	1,019	31.403(28.554,34.253)	15.693(6.189,25.198)	20.036(6.962,33.11)	14.735(5.492,23.979)
2012 +	346	30.058(25.227,34.889)	18.322(7.572,29.073)	20.336(7.789,32.883)	23.776(12.118,35.433)
Vehicle make					
American	$^{3,635}$	33.92(32.381, 35.459)	25.321(18.17, 32.471)	37.287(20.562,54.012)	24.194(19.862, 28.526)
Japanese	1,927	34.51(32.387, 36.632)	17.178(8.904, 25.451)	21.274(9.996, 32.551)	17.272(10.022,24.522)
Korean	387	32.558(27.89, 37.227)	21.338(10.317, 32.358)	30.873(11.583, 50.162)	25.404(14.335, 36.473)
Other	322	32.298(27.191, 37.406)	11.894(0.98, 22.807)	13.358(0.14, 26.576)	14.575(5.943, 23.206)
Vehicle type					
Passenger car	4,063	35.565(34.093, 37.037)	20.78(14.442,27.117)	28.72(17.684, 39.755)	21.834(18.109, 25.559)
Light truck	1,213	31.904(29.281, 34.527)	20.445(10.611, 30.28)	28.5(12.712, 44.289)	20.931(14.712,27.149)
Van	290	26.897(21.793, 32)	25.071(10.311, 39.832)	39.315(14.396, 64.234)	19.598(6.469, 32.727)
SUV	705	30.922(27.51, 34.334)	21.916(12.779, 31.053)	32.376(15.318, 49.435)	20.32(12.643, 27.996)

TABLE 17

Percentage of severe injury in SPINE and associated 95% CIs by different covariates across DR adjustment methods

Covariate	n	Unweighted (95% CI)	GPPP (95% CI)	LWP (95% CI)	AIPW (95% CI)
Total	6,271	3.716(3.247, 4.184)	6.735(4.696, 8.774)	6.978(2.777, 11.18)	6.205(3.15, 9.261)
Gender	,				
Male	3,230	4.18(3.489, 4.87)	6.819(4.049, 9.588)	7.154(2.499, 11.809)	8.946(5.07, 12.822)
Female	3,041	3.223(2.595, 3.85)	6.719(3.861,9.577)	6.891(2.261, 11.522)	4.21(0,9.072)
Age group		· · · · · ·			
16-19	659	3.49(2.089, 4.891)	1.823(0, 4.007)	2.358(0, 9.212)	0(0, 3.847)
20-39	2,732	3.807(3.089, 4.524)	5.099(2.982, 7.217)	5.406(1.363, 9.448)	3.964(0.399, 7.529)
40-64	1,946	3.443(2.633, 4.253)	7.68(5.417,9.943)	7.948(3.292,12.603)	9.501(5.524, 13.479)
65 +	934	4.176(2.893, 5.458)	12.795(7.935, 17.655)	12.616(7.457, 17.775)	11.194(1.223, 21.164)
Occupant role					
Driver	1,547	5.624(4.476, 6.772)	7.059(2.952, 11.166)	7.549(0, 16.523)	3.584(0, 10.498)
Passenger	4,724	3.091(2.597, 3.584)	6.683(4.361,9.005)	6.868(3.323, 10.412)	6.93(3.137, 10.722)
Seating Row		,			
Front	5,853	3.434(2.968, 3.901)	6.498(4.559, 8.438)	6.73(2.905, 10.554)	5.999(2.883, 9.116)
Rear	418	7.656(5.107,10.204)	12.752(4.811, 20.692)	13.399(0, 29.785)	9.724(1.107, 18.341)
Restraint use					
Restraint	2,275	4.308(3.473, 5.142)	5.92(1.798, 10.042)	6.805(0, 16.123)	6.475(0, 14.4)
Unrestraint	3,996	3.378(2.818, 3.939)	6.766(4.721, 8.812)	6.797(4.24, 9.353)	6.268(3.289, 9.247)
Air bag		,	· · · · · ·	× · · · ·	
Deployed	2,194	3.874(3.067, 4.682)	8.504(3.145, 13.864)	10.364(0.927, 19.801)	6.919(0, 14.085)
Not deployed	4,077	3.63(3.056, 4.204)	5.876(4.41, 7.343)	5.826(4.01, 7.641)	5.954(3.153, 8.756)
Injury scale (A	IS)				
3	1,977	5.513(4.507, 6.52)	17.854(14.154, 21.553)	17.101(12.953, 21.249)	18.924(12.225, 25.623)
4	761	4.731(3.222,6.239)	16.246(9.105,23.387)	15.862(8.16,23.564)	11.393(0.586,22.2)
5+	781	8.707(6.729,10.684)	22.462(10.034,34.889)	24.098(8.787,39.409)	22.264(8.191,36.336)
Days hospitaliz	zed	X / / /			
0	2,259	2.745(2.071, 3.418)	3.815(0.804, 6.826)	4.92(0, 12.6)	0(0,6.667)
1-3	1,757	2.732(1.97, 3.494)	4.476(2.538, 6.415)	4.054(2.23, 5.877)	5.118(1.926, 8.31)
4-7	1,153	4.51(3.312, 5.708)	13.038(9.061, 17.015)	12.105(8.546, 15.665)	13.629(8.565, 18.692)
8-14	640	5.625(3.84, 7.41)	14.223(10.23, 18.215)	13.837(10.044, 17.63)	13.278(8.82,17.737)
15 +	462	7.576(5.163, 9.989)	25.109(19.802,30.416)	24.763(19.61, 29.915)	25.392(20.03, 30.754)
Delta-v (Km/ł	1)			· · · · · ·	· · · · · · · · · · · · · · · · · · ·
Minor	404	2.228(0.789, 3.667)	11.229(1.136, 21.322)	12.934(0, 26.013)	8.643(0, 30.853)
Moderate	3,304	2.542(2.006, 3.079)	6.269(3.382, 9.156)	6.597(1.454, 11.741)	6.322(3.002, 9.642)
Severe	2,563	5.462(4.583, 6.342)	8.53(5.654, 11.406)	8.535(5.507, 11.562)	7.466(3.136, 11.796)
Damage distril	bution				
Wide	4,657	3.393(2.873, 3.913)	5.28(3.515, 7.045)	5.532(1.939, 9.125)	3.803(0.55, 7.057)
Narrow	320	5(2.612, 7.388)	2.417(0, 5.39)	2.967(0, 8.287)	2.673(0, 5.984)
Corner	584	1.884(0.781, 2.986)	7.107(2.759, 11.454)	7.097(2.231, 11.963)	15.586(0.739, 30.432)
Other	710	6.761(4.914, 8.607)	18.668(10.843, 26.494)	19.005(6.446, 31.564)	17.857(7.203, 28.512)
Rollover					
Yes	4,942	3.157(2.669, 3.644)	5.129(3.248, 7.01)	5.354(1.836, 8.872)	4.668(1.216, 8.119)
No	1,329	5.794(4.538, 7.05)	13.905(9.303, 18.508)	14.245(5.965, 22.524)	13.444(6.794, 20.094)
Deformation lo	ocatior	1			
Front	$^{3,671}$	3.242(2.669, 3.815)	5.676(3.614, 7.738)	5.811(2.379, 9.244)	3.488(0, 7.784)
Right	1,123	3.562(2.478, 4.646)	3.673(0, 7.795)	4.021(0, 10.545)	7.553(1.56, 13.546)
Left	857	3.851(2.562, 5.139)	7.337(0.385, 14.289)	7.933(0.269, 15.598)	7.972(0, 17.665)
Rear	620	6.613(4.657, 8.569)	22.12(12.372, 31.869)	22.254(9.529, 34.98)	20.896(8.037, 33.755)
Model year					
$\leq 2003$	2,529	3.598(2.872, 4.324)	7.792(5.022, 10.561)	8.376(0.769, 15.983)	4.647(0, 10.046)
2004 - 2007	2,377	3.366(2.641, 4.091)	8.083(5.475, 10.691)	8.184(4.72, 11.647)	10.824(6.32, 15.328)
2008-2011	1,019	4.514(3.239, 5.789)	3.349(1.139, 5.558)	3.365(1.098, 5.633)	4.816(0.267, 9.366)
2012+	346	4.624(2.411, 6.837)	5.411(1.644, 9.178)	5.207(1.219, 9.195)	0(0,15.701)
Vehicle make					
American	$^{3,635}$	3.301(2.72, 3.882)	7.388(4.517, 10.258)	7.965(1.09, 14.839)	8.747(4.478, 13.016)
Japanese	1,927	4.307(3.401, 5.214)	6.099(2.908, 9.29)	6.032(2.707, 9.357)	1.562(0, 7.543)
Korean	387	3.359(1.564, 5.154)	8.952(2.792, 15.112)	9.172(2.504, 15.841)	8.469(0.439,16.499)
Other	322	5.28(2.837, 7.722)	3.023(0, 8.831)	2.777(0, 8.478)	3.854(0, 9.39)
Vehicle type					
Passenger car	4,063	3.569(2.998, 4.139)	6.075(3.814, 8.337)	6.245(2.521, 9.969)	4.639(0.478, 8.8)
Light truck	1,213	4.122(3.003, 5.241)	8.549(3.935, 13.164)	8.846(2.517, 15.175)	9.311(2.577, 16.044)
Van	290	1.379(0.037, 2.722)	3.755(0, 11.694)	4.766(0, 15.574)	12.787(0,27.049)
SUV	705	4.823(3.241,6.404)	8.681(1.348,16.013)	9.09(0.315, 17.865)	8.533(0,19.472)