

WEYL'S LAW FOR THE STEKLOV PROBLEM ON SURFACES WITH ROUGH BOUNDARY

MIKHAIL KARPUKHIN, JEAN LAGACÉ, AND IOSIF POLTEROVICH

ABSTRACT. The validity of Weyl's law for the Steklov problem on domains with Lipschitz boundaries is a well-known open question in spectral geometry. We answer this question in two dimensions and show that Weyl's law holds for an even larger class of surfaces with rough boundaries. This class includes domains with interior cusps as well as "slow" exterior cusps. Moreover, the condition on the speed of exterior cusps cannot be improved, which makes our result in a sense optimal. The proof is based on the methods of Suslina and Agranovich combined with some observations about the boundary behaviour of conformal mappings.

1. INTRODUCTION AND MAIN RESULTS

1.1. **Asymptotics of Steklov eigenvalues.** Let Ω be a bounded domain in a smooth complete Riemannian manifold (\mathcal{M}, g) of dimension $d \geq 2$. Consider the Steklov eigenvalue problem

$$\begin{cases} \Delta u = 0 & \text{in } \Omega; \\ \partial_\nu u = \sigma u & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where Δ is the Laplace-Beltrami operator on \mathcal{M} associated with the Riemannian metric g and ∂_ν is the outward normal derivative. Under some regularity conditions on the boundary, for instance, if $\partial\Omega$ is Lipschitz [AM12], the spectrum is discrete and forms a sequence accumulating only at infinity:

$$0 = \sigma_0(\Omega) \leq \sigma_1(\Omega) \leq \sigma_2(\Omega) \leq \dots \nearrow \infty.$$

We will discuss weaker conditions under which the spectrum is discrete later on. To study eigenvalue asymptotics, it is convenient to introduce the eigenvalue counting function

$$N(\sigma) := \#\{j \in \mathbb{N} : \sigma_j(\Omega) < \sigma\}.$$

If $\partial\Omega$ is piecewise C^1 , it is known [Agr06] that the counting function satisfies the Weyl asymptotics

$$N(\sigma) = \frac{\omega_{d-1}}{(2\pi)^{d-1}} \text{Vol}_{d-1}(\partial\Omega) \sigma^{d-1} + o\left(\sigma^{d-1}\right), \quad (1.2)$$

where ω_d is the volume of the d -dimensional unit ball. We refer also to [San55, Sha71, Sus99b, AA96] for earlier results on this topic, as well as to [Roz79, Edw93, GPPS14, GKLP22] for improvements of the error estimate under stronger regularity assumptions. Extending the asymptotic formula (1.2) to domains with Lipschitz boundaries is a well known open problem, see e.g. [GP17, RS21, Sus21, GKLP22], to which we provide an answer in two dimensions.

Theorem 1.1. *Let Ω be a bounded domain with Lipschitz boundary in a smooth complete Riemannian manifold of dimension two. Then its Steklov eigenvalues satisfy the asymptotics (1.2).*

In fact, we prove that (1.2) holds for domains satisfying weaker regularity conditions defined via the boundary behaviour of conformal maps, see Section 1.2. We give examples of domains satisfying those conditions in Section 4, they include the so-called chord-arc domains, as well as domains with inward and “slow” outward cusps, see Proposition 4.2.

The proof of Theorem 1.1 relies on the variational characterisation of Steklov eigenvalues. While this characterisation is standard for Lipschitz domains, certain subtleties arise for domains with less regular boundary which we clarify in Section 1.3. We use a conformal map to obtain an isospectral weighted Steklov problem on a surface with smooth boundary. The isospectrality follows from the equivalence of the corresponding variational characterisations; for Lipschitz domains, it can be deduced almost directly from the analogous results for the Neumann problem [GU16]. Finally, we use the methods developed in [BS80, Sus99b], see also [Agr06], in order to obtain spectral asymptotics for these weighted Steklov problems. For Lipschitz domains we could use the results from [Sus99b, Agr06] in a straightforward way; we extend these techniques to allow for more singular weights corresponding to less regular boundaries.

1.2. Conformal regularity. As was mentioned above, our first goal is to reduce the Steklov problem on a surface with rough boundary to a weighted Steklov problem on a surface with smooth boundary, via a conformal map. Slightly abusing terminology, we refer to domains in two-dimensional Riemannian manifolds whose boundary is a finite collection of disjoint closed simple curves as *surfaces with boundary*. We say that two surfaces Ω_1 and Ω_2 with (potentially empty) boundary are *conformally equivalent*, or *in the same conformal class* if there exists $\varphi : \Omega_1 \rightarrow \Omega_2$ a conformal diffeomorphism of their interior extending to a homeomorphism of their boundary. This defines an equivalence relation on surfaces with boundary, and it is clear that every conformal class consists of surfaces with the same topological type, i.e. same orientability, genus and number of boundary components.

The uniformisation theorems are concerned with finding a canonical representative in every conformal class \mathcal{C} . These canonical representatives are *circle domains*, which are the complement of b geodesic disks in a closed surface M endowed with a metric of constant curvature. It follows from the uniformisation theorems [Haa84, Mas89] that there is a circle domain in every orientable conformal class with finite topology; this result was extended to the non-orientable setting in [KS21, pp. 11–12]. We shall denote this canonical representative $(\Omega_{\mathcal{C}}, g_{\mathcal{C}})$.

Many boundary regularity results in the literature are proven for conformal maps from the disk to simply connected surfaces with boundary. It follows from [BK87, p.24] that any such result for maps from the disk is also valid for maps from an annulus into a doubly connected surface with boundary, by alternately filling the boundary components of the target with a disk and conjugating with inversions. As observed in [KS21, Remark 2.2], this allows one to extend the regularity theory to conformal maps from arbitrary circle domains with finite topology. Indeed, in that situation the restriction of the conformal map to a neighbourhood of one boundary component in the circle domain is a map from an annulus into a doubly connected surface with boundary. In particular, Carathéodory’s Theorem tells us that any conformal diffeomorphism of the interiors $\varphi : \Omega_{\mathcal{C}} \rightarrow \Omega$ extends to a homeomorphism of the boundary [Pom92, Theorem 2.6].

Definition 1.2. Let \mathcal{C} be a conformal class and $\varphi : \Omega_{\mathcal{C}} \rightarrow \Omega$ be a conformal diffeomorphism. We define when they exist,

$$\beta := \left| d\varphi \Big|_{\partial\Omega_{\mathcal{C}}} \right| \quad \text{and} \quad \eta := |d\varphi|^2.$$

We call β the *boundary conformal factor* and η the *interior conformal factor*.

The interior conformal factor $\eta \in L^1(\Omega_{\mathcal{C}})$ and the Riemannian volume measure dv_g on Ω is the pushforward measure $\varphi_*(\eta dv_{g_{\mathcal{C}}})$. If a surface with boundary has finite perimeter the boundary conformal factor $\beta \in L^1(\partial\Omega_{\mathcal{C}})$, and the boundary length measure $d\ell_g$ on $\partial\Omega$ is the pushforward measure $\varphi_*(\beta d\ell_{g_{\mathcal{C}}})$ [Pom92, Theorem 6.8]. Integrability properties of the conformal factors β and η are controlled by the regularity of the boundary $\partial\Omega$. This motivates the following definition of regularity classes.

Definition 1.3. Let \mathcal{C} be a conformal class, $\Omega \in \mathcal{C}$, and $\mathcal{X}(\Omega_{\mathcal{C}})$ and $\mathcal{Y}(\partial\Omega_{\mathcal{C}})$ be function spaces. We say that Ω has *boundary conformal regularity* \mathcal{Y} if for some conformal diffeomorphism $\varphi : \Omega_{\mathcal{C}} \rightarrow \Omega$, the boundary conformal factor $\beta \in \mathcal{Y}$. We say that Ω has *interior conformal regularity* \mathcal{X} if for some conformal diffeomorphism the interior conformal factor $\eta \in \mathcal{X}$.

We note that our definition of interior conformal regularity differs from that in [GU16, GPU18] by a factor of 2 since in those papers it was stated in terms of the integrability of $|d\varphi|$ rather than $|d\varphi|^2$. In other words, interior conformal regularity L^p corresponds to domains which are $2p$ -conformally regular in their definitions.

The integrability class of the interior conformal factor Ω has been used to investigate the properties of the Neumann Laplacian, see [GU16, GPU18]. As expected, the regularity of the boundary conformal factor also appears in the study of the Steklov problem.

Remark 1.4. By the Kellogg–Warschawski Theorem [Pom92, Theorem 3.6], surfaces with boundary of class $C^{n,\alpha}$, $n \geq 1$ and $0 < \alpha < 1$, have boundary conformal regularity $C^{n-1,\alpha}$. It follows furthermore from the arguments in the proof of [BBL16, Lemma 5.1] that any surface with Lipschitz boundary (or, more generally, a chord–arc domain, see Section 4) has boundary conformal regularity L^p for some $p > 1$, and surfaces of finite perimeter have boundary conformal regularity L^1 .

Note that domains with exterior cusps do not have boundary conformal regularity L^p for any $p > 1$. In order to include some of these domains in our analysis we need to recall the following definition [BS88, Sections IV.6, IV.8].

Definition 1.5. Given (Ξ, μ) a measure space of finite measure and $a \geq 0$ the space $L(\log L)^a(\Xi)$ is a space of functions f on Ξ such that

$$\int_{\Xi} |f| (\log(2 + |f|))^a d\mu < \infty$$

endowed with the norm

$$\|f\|_{L(\log L)^a(\Xi)} = \inf \left\{ t > 0 : \int_{\Xi} \left| \frac{f}{t} \right| \left(\log \left(2 + \left| \frac{f}{t} \right| \right) \right)^a d\mu \leq 1 \right\}.$$

One can show that $L(\log L)^a(\Xi)$ is a Banach space for every $a \geq 0$. The dual of $L(\log L)^a(\Xi)$ is the space $\exp L^{1/a}(\Xi)$ of functions f on Ξ such that

$$\|f\|_{\exp L^{1/a}(\Xi)} := \inf \left\{ t > 0 : \int_{\Xi} \exp \left(\left| \frac{f}{t} \right|^{1/a} \right) d\mu \leq 1 \right\} < \infty. \quad (1.3)$$

The norm (1.3) is equivalent to the dual norm on $\exp L^{1/a}$ so that there is C depending only on (Ξ, μ) such that for all $f \in L(\log L)^a(\Xi)$ and $u \in \exp L^{1/a}(\Xi)$ the Hölder-type inequality

$$\|fu\|_{L^1(\Xi)} \leq C \|f\|_{L(\log L)^a(\Xi)} \|u\|_{\exp L^{1/a}(\Xi)} \quad (1.4)$$

holds.

Given a conformal class \mathcal{C} and $\Omega \in \mathcal{C}$ with boundary conformal factor β we consider the weighted Steklov problem

$$\begin{cases} \Delta u = 0 & \text{in } \Omega_{\mathcal{C}}; \\ \partial_{\nu} u = \beta \sigma u & \text{on } \partial\Omega_{\mathcal{C}}, \end{cases} \quad (1.5)$$

The spectrum of this weighted problem is discrete and accumulates at infinity if the trace $W^{1,2}(\Omega_{\mathcal{C}}) \hookrightarrow L^2(\partial\Omega_{\mathcal{C}}, \beta d\ell_{g_{\mathcal{C}}})$ is compact, see [GKL21, Sections 3 and 4]. This is the case if $\beta \in L \log L(\Omega_{\mathcal{C}})$, see Proposition 2.2.

Theorem 1.6. *Let \mathcal{C} be a conformal class and $(\Omega, g) \in \mathcal{C}$ be a surface with boundary conformal regularity $L \log L$. Then, problems (1.1) and (1.5) are isospectral in the sense that $\sigma_k(\Omega) = \sigma_k(\Omega_{\mathcal{C}}, \beta)$ for all $k \in \mathbb{N}$.*

Remark 1.7. A simple computation shows that Theorem 1.6 holds when the boundary is smooth, see e.g. [JS14, Lemma 3.3]. If the boundary is sufficiently rough, isospectrality is not a priori clear, and is resolved through the study of composition operators between Sobolev spaces with appropriately chosen norm. In a similar way, the existence of those bounded composition operators gives rise to isospectrality of the weighted Neumann problems, see [GU16]. This issue was not previously addressed in the literature on the Steklov problem on Lipschitz domains, cf. [GP10, Proposition 2.1.4].

Our main technical theorem is the following.

Theorem 1.8. *Let Ω be a surface with boundary of boundary conformal regularity $L \log L$. Then its Steklov eigenvalues satisfy the asymptotic formula (1.2).*

Equivalently, for any surface Ω with smooth boundary and $\beta \in L \log L(\partial\Omega)$, the eigenvalues of the weighted problem (1.5) satisfy the asymptotic formula (1.2), with $\text{Vol}_{d-1}(\partial\Omega)$ replaced by $\int_{\partial\Omega} \beta d\ell_g$.

The equivalence of the two formulations follows from Theorem 1.6. Note that in view of Remarks 1.4, Theorems 1.6 and 1.8 imply Theorem 1.1.

1.3. Variational characterisation and natural domains for the Steklov problem. On a surface with boundary Ω , consider the Sobolev space

$$W^{1,2}(\Omega) := \{f \in L^2(\Omega) : |\nabla f| \in L^2(\Omega)\},$$

where ∇f is the weak gradient. If Ω is a surface with Lipschitz boundary, there are two equivalent norms on $W^{1,2}(\Omega)$:

$$\|f\|_{W^{1,2}(\Omega)}^2 = \int_{\Omega} |\nabla f|^2 dv_g + \int_{\Omega} f^2 dv_g, \quad (1.6)$$

and

$$\|f\|_{W_{\partial}^{1,2}(\Omega)}^2 = \int_{\Omega} |\nabla f|^2 dv_g + \int_{\partial\Omega} f^2 d\ell_g. \quad (1.7)$$

The norm (1.6) is the standard one and is commonly used in interior problems, for instance the Neumann problem. On the other hand the norm (1.7) is a natural norm of choice for the Steklov problem. When the boundary is only some collection of Jordan curves these norms may not be equivalent, even when the boundary has finite perimeter (one can show that this the case for domains with fast cusps as defined in subsection 4.2). By the Meyers–Serrin theorem, for any surface with boundary Ω the space $W^{1,2}(\Omega)$ is the completion of

$$\mathcal{W}(\Omega) := \left\{ f \in C^{\infty}(\Omega) : \|f\|_{W^{1,2}(\Omega)} < \infty \right\}$$

under the $\|\cdot\|_{W^{1,2}(\Omega)}$ norm, which motivates the following definition.

Definition 1.9. Let Ω be a surface with boundary. The boundary Sobolev space $W_{\partial}^{1,2}(\Omega)$ is defined as the completion of

$$\mathcal{W}_{\partial}(\Omega) := \left\{ f : \bar{\Omega} \rightarrow \mathbb{R} : f \in C^{\infty}(\Omega) \text{ and } \|f\|_{W_{\partial}^{1,2}(\Omega)} < \infty \right\}$$

under the $\|\cdot\|_{W_{\partial}^{1,2}(\Omega)}$ norm.

Again, for surfaces with sufficiently regular boundary, $W^{1,2}(\Omega)$ and $W_{\partial}^{1,2}(\Omega)$ are isomorphic. We give the following condition for their equivalence in terms of interior and boundary conformal regularity.

Proposition 1.10. *Let Ω be a surface with boundary with both interior and boundary conformal regularity $L \log L$. Then, $W^{1,2}(\Omega)$ and $W_{\partial}^{1,2}(\Omega)$ are isomorphic.*

The appropriate space to define the Steklov problem (especially when they are not isomorphic) is $W_{\partial}^{1,2}(\Omega)$, see [NT08]. The Steklov eigenvalues $\sigma_k(\Omega)$ satisfy the variational characterisation

$$\sigma_k(\Omega) = \inf_{E_k} \sup_{u \in E_k \setminus \{0\}} \frac{\int_{\Omega} |\nabla u|^2 dv_g}{\int_{\partial\Omega} u^2 d\ell_g},$$

where E_k is a $k + 1$ dimensional subspace of $W_{\partial}^{1,2}(\Omega)$. For the weighted problem on $\Omega_{\mathcal{C}}$, we have that for $\beta \in L \log L(\Omega_{\mathcal{C}})$ the weighted Steklov eigenvalues satisfy the characterisation

$$\sigma_k(\Omega_{\mathcal{C}}, \beta) = \inf_{E_k} \sup_{u \in E_k \setminus \{0\}} \frac{\int_{\Omega_{\mathcal{C}}} |\nabla u|^2 dv_g}{\int_{\partial\Omega_{\mathcal{C}}} u^2 \beta d\ell_g},$$

where again E_k is a $k + 1$ dimensional subspace of $W_{\partial}^{1,2}(\Omega_{\mathcal{C}})$. The isospectrality Theorem 1.6 is a consequence of the following result on composition operator between Sobolev spaces.

Proposition 1.11. *Let \mathcal{C} be a conformal class and $(\Omega, g) \in \mathcal{C}$ be a surface with boundary. Let $\varphi : \Omega_{\mathcal{C}} \rightarrow \Omega$ be a conformal diffeomorphism with boundary conformal factor $\beta \in L \log L$. Then, the composition operator*

$$\varphi^* : W_{\partial}^{1,2}(\Omega) \rightarrow W_{\partial}^{1,2}(\Omega_{\mathcal{C}}) \quad \varphi^* f := f \circ \varphi$$

induced by φ is an isomorphism.

Plan of the paper. The paper is organised as follows. Section 2 is concerned with the proof of Theorem 1.6. We then use the variational isospectrality to prove that domains of boundary conformal regularity $L \log L$ have discrete Steklov spectrum. Section 3 is dedicated to proving Theorem 1.8, expanding on the theory of spectral asymptotics for variational eigenvalues developed in [BS80, Sus99b]. In Section 4 we give a few examples of domains satisfying the hypotheses of Theorem 1.8. Finally, in Section 5 we discuss some further extensions and applications of our methods, in particular to the Steklov problem with an indefinite weight and to the Neumann eigenvalue problem.

Acknowledgements. The authors would like to thank G. Rozenblum for helpful comments and proposed simplifications of the proofs of Proposition 2.2 and Lemma 3.2. We are also grateful to A. Ukhlov for useful discussions. The research of M.K. was partially supported by NSF grant DMS-2104254. The research of J.L. was partially supported by EPSRC award EP/T030577/1. He also thanks the University of Bristol for its hospitality while this paper was written. The research of I.P. was partially supported by NSERC and FRQNT.

2. ISOSPECTRALITY AND COMPOSITION OPERATORS

We first prove the following lemma about composition operators on some Orlicz spaces, in similar fashion to [GU16, Theorem 4] which is stated for Lebesgue spaces.

Lemma 2.1. *For $j \in \{1, 2\}$, let (Ξ_j, μ_j) be measure spaces with finite measure, $\varphi : \Xi_1 \rightarrow \Xi_2$ be measurable and suppose that the pushforward measure $\varphi_*(\mu_1) = \beta \mu_2$, where $\beta : \Xi_2 \rightarrow (0, \infty)$. Then, φ induces a bounded composition operator*

$$\varphi^* : \exp L^2(\Xi_2) \rightarrow L^2(\Xi_1), \quad \varphi^* f := f \circ \varphi$$

if and only if $\beta \in L \log L(\Xi_2)$.

Proof. To prove that the condition $\beta \in L \log L(\Xi_2)$ is sufficient, assume that $f \in \exp L^2(\Xi_2)$, so that $|f|^2 \in \exp L(\Xi_2)$. Since $L \log L(\Xi_2)$ is a reflexive space with dual $\exp L(\Xi_2)$, we can compute

$$\begin{aligned} \int_{\Xi_1} |\varphi^* f|^2 d\mu_1 &= \int_{\Xi_2} |f|^2 \beta d\mu_2 \\ &\leq \|\beta\|_{L \log L(\Xi_2)} \|f^2\|_{\exp L(\Xi_2)} \\ &= \|\beta\|_{L \log L(\Xi_2)} \|f\|_{\exp L^2(\Xi_2)}^2. \end{aligned} \tag{2.1}$$

Let us now show that the condition is necessary. Indeed, if $\beta \notin L \log L(\Xi_2)$, it is not a bounded linear functional on $\exp L(\Xi_2)$, so we can find $|f|^2 \in \exp L(\Xi_2)$ so that the second term on the first line in (2.1) is not finite. \square

In the next proposition, we show compactness of a weighted boundary trace. The proof is similar in nature to the ideas in [GKL21, Example 3.19 (iii)] where the weight is instead in the interior.

Proposition 2.2. *Let Ω be a surface with smooth boundary and $0 \leq \beta \in L \log L(\partial\Omega)$, $\beta \not\equiv 0$. Then, the trace $T_\beta : W^{1,2}(\Omega) \rightarrow L^2(\partial\Omega, \beta d\ell)$ is compact.*

Proof. Define $\theta : \partial\Omega \rightarrow \mathbb{R}$ as $\theta := \frac{1}{\beta} \mathbf{1}_{\{\beta > 0\}}$. Consider the diagram

$$\begin{array}{ccccc} W^{1,2}(\Omega) & \xrightarrow{T} & \exp L^2(\partial\Omega) & \xrightarrow{M_{\sqrt{\beta}}} & L^2(\partial\Omega) \\ & & & & \downarrow M_{\sqrt{\theta}} \\ & & & & L^2(\partial\Omega, \beta d\ell), \\ & & \searrow T_\beta & & \end{array}$$

where T is the trace and M_h is the operator of multiplication by the function h . The trace operator T is bounded; $\exp L^2(\partial\Omega)$ is in fact the optimal target space on $\partial\Omega$ for bounded traces from $W^{1,2}(\Omega)$, see [CP16, Example 5.3]. By Hölder inequality (1.4) between $\exp L(\partial\Omega)$ and $L \log L(\partial\Omega)$ there exists $C > 0$ such that

$$\left\| M_{\sqrt{\beta}} f \right\|_{L^2(\partial\Omega)}^2 = \int_{\partial\Omega} f^2 \beta \, d\ell \leq C \|f^2\|_{\exp L(\partial\Omega)} \|\beta\|_{L \log L(\partial\Omega)} = C \|f\|_{\exp L^2(\partial\Omega)}^2 \|\beta\|_{L \log L(\partial\Omega)}.$$

In other words, $M_{\sqrt{\beta}}$ is bounded with norm at most $C \|\beta\|_{L \log L(\partial\Omega)}$. As for $M_{\sqrt{\theta}}$, we have that

$$\left\| M_{\sqrt{\theta}} f \right\|_{L^2(\partial\Omega; \beta d\ell)}^2 = \int_{\partial\Omega \cap \{\beta > 0\}} f^2 \, d\ell \leq \|f\|_{L^2(\partial\Omega)}^2$$

Thus, for a probably different constant $C > 0$ independent of β we have

$$\|T_\beta\| \leq C \|\beta\|_{L \log L(\partial\Omega_c)}.$$

To prove compactness, it is sufficient to prove that $M_{\sqrt{\beta}} \circ T$ is compact; if β is a nonnegative smooth function this follows from the usual trace restriction theorem. By density of smooth functions in $L \log L$, for every $\varepsilon > 0$, there is a nonnegative $\beta_\varepsilon \in C^\infty(\partial\Omega)$ such that $\|\beta - \beta_\varepsilon\|_{L \log L(\partial\Omega)} < \varepsilon$ and $\beta_\varepsilon \leq \beta$ almost everywhere, so that $\sqrt{\beta} - \sqrt{\beta_\varepsilon} \leq \sqrt{\beta - \beta_\varepsilon}$. But then,

$$\left\| M_{\sqrt{\beta}} \circ T - M_{\sqrt{\beta_\varepsilon}} \circ T \right\| \leq \left\| M_{\sqrt{\beta - \beta_\varepsilon}} \right\| \|T\| \leq \|\beta - \beta_\varepsilon\|_{L \log L(\partial\Omega)} \|T\| \leq \varepsilon \|T\|.$$

Thus $M_{\sqrt{\beta}} \circ T$ is a norm limit of compact operators hence compact itself and T_β is compact also. \square

We now have the right tools to prove the composition Proposition 1.11, following the structure of the proof of [GU16, Theorem 6].

Proof of Proposition 1.11. Let $f \in \mathcal{W}_\partial(\Omega)$ as in Definition 1.9. Invariance of the Dirichlet energy under conformal diffeomorphisms tells us that

$$\|\nabla f\|_{L^2(\Omega)} = \|\nabla(\varphi^* f)\|_{L^2(\Omega_c)}.$$

Since the boundary conformal factor is in $L \log L$, φ induces the bounded composition operator $(\varphi^{-1})^* : \exp L^2(\partial\Omega_C) \rightarrow L^2(\partial\Omega)$. Therefore, for every $a \in \mathbb{R}$, we have that

$$\begin{aligned} |a| &= \text{Per}(\Omega)^{-1/2} \|c\|_{L^2(\partial\Omega)} \leq \text{Per}(\Omega)^{-1/2} \left(\|f\|_{L^2(\partial\Omega)} + \|f - a\|_{L^2(\partial\Omega)} \right) \\ &\leq \text{Per}(\Omega)^{-1/2} \left(\|f\|_{L^2(\partial\Omega)} + \|\varphi^* f - a\|_{\exp L^2(\partial\Omega_C)} \right). \end{aligned} \quad (2.2)$$

We claim that there exists a constant $C > 0$ such that

$$\begin{aligned} \|\varphi^* f\|_{L^2(\partial\Omega_C)} &\leq \inf_{a \in \mathbb{R}} \left(\|a\|_{L^2(\partial\Omega_C)} + \|\varphi^* f - a\|_{L^2(\partial\Omega_C)} \right) \\ &\leq \inf_{a \in \mathbb{R}} \left(\text{Per}(\Omega)^{-1/2} \text{Per}(\Omega_C)^{1/2} + C \|1\|_{L \log L(\partial\Omega_C)} \right) \left(\|f\|_{L^2(\partial\Omega)} + \|\varphi^* f - a\|_{\exp L^2(\partial\Omega_C)} \right). \end{aligned}$$

Indeed, the first inequality is just the triangle inequality. We then use (2.2) to estimate the first term, and inequality (1.4) together with the relations $\|h\|_{L^2}^2 = \|h^2\|_{L^1}$, $\|h\|_{\exp L^2}^2 = \|h^2\|_{\exp L^1}$ to estimate the second.

The space $\exp L^2(\partial\Omega_C)$ is the optimal target space for traces from $W^{1,2}(\Omega_C)$, and this is equivalent to the validity of a Poincaré trace inequality, see [CP16, Theorems 1.3 and 5.3],

$$\inf_a \|\varphi^* f - a\|_{\exp L^2(\partial\Omega_C)} \leq C \|\nabla \varphi^* f\|_{L^2(\Omega_C)}.$$

Combining the previous display formulas gives us the existence of some constant $C > 0$ such that

$$\|\varphi^* f\|_{W_\partial^{1,2}(\Omega_C)} \leq C \|f\|_{W_\partial^{1,2}(\Omega)}.$$

Since $\mathcal{W}_\partial(\Omega)$ is dense in $W_\partial^{1,2}(\Omega)$, the pullback φ^* extends to the whole space as a bounded operator as well. Proving the analogous result for $(\varphi^{-1})^*$ is simpler. Since Ω_C has smooth boundary, the spaces $W^{1,2}(\Omega_C)$ and $W_\partial^{1,2}(\Omega_C)$ are isomorphic. Compactness (in fact, boundedness is enough here) of the trace $W^{1,2}(\Omega_C) \rightarrow L^2(\partial\Omega_C, \beta d\ell_{g_C})$ obtained in Proposition 2.2 then implies that for every $h \in \mathcal{W}(\Omega_C)$,

$$\|(\varphi^{-1})^* h\|_{W_\partial^{1,2}(\Omega)}^2 = \|\nabla h\|_{L^2(\Omega_C)}^2 + \|h\|_{L^2(\partial\Omega_C, \beta d\ell_{g_C})}^2 \leq C \|h\|_{W_\partial^{1,2}(\Omega_C)}^2.$$

By density we once again have that $(\varphi^{-1})^*$ extends to the whole space as a bounded operator, completing the proof. \square

We can now prove Theorem 1.6.

Proof of Theorem 1.6. Let E_k be a $k + 1$ dimensional subspace of $W_\partial^{1,2}(\Omega)$. Then, by Proposition 1.11, $\varphi^*(E_k)$ is a $k + 1$ dimensional subspace of $W_\partial^{1,2}(\Omega_C)$, and for every $u \in E_k$,

$$\frac{\int_\Omega |\nabla u|^2 dv_g}{\int_{\partial\Omega} u^2 d\ell_g} = \frac{\int_{\Omega_C} |\nabla u|^2 dv_{g_C}}{\int_{\partial\Omega_C} u^2 \beta d\ell_{g_C}}.$$

This implies directly that $\sigma_k(\Omega_C, \beta) \leq \sigma_k(\Omega)$. The analogous reasoning with $(\varphi^{-1})^*$ instead of φ^* gives the reverse inequality. \square

In order to prove Proposition 1.10, we extend the results of [GU16] to a slightly more singular interior conformal factor.

Lemma 2.3. *Let \mathcal{C} be a conformal class and $(\Omega, g) \in \mathcal{C}$ be a surface with boundary, which has interior conformal regularity $L \log L$ through the conformal diffeomorphism $\varphi : \Omega_{\mathcal{C}} \rightarrow \Omega$. Then, the composition operator*

$$\varphi^* : W^{1,2}(\Omega) \rightarrow W^{1,2}(\Omega_{\mathcal{C}}) \quad \varphi^* f := f \circ \varphi$$

induced by φ is an isomorphism.

Proof. The proof is essentially identical to the proof of Proposition 1.11. We replace the result on the traces $W_{\partial}^{1,2}(\Omega_{\mathcal{C}}) \rightarrow \exp L^2(\partial\Omega_{\mathcal{C}})$ and the corresponding Poincaré inequality with the optimal Sobolev embedding $W^{1,2}(\Omega_{\mathcal{C}}) \rightarrow \exp L^2(\Omega_{\mathcal{C}})$, and use the fact that the interior conformal factor $|\mathrm{d}\varphi|^2 \in L \log L$ to get a bounded composition operator $\exp L^2(\Omega_{\mathcal{C}}) \rightarrow L^2(\Omega)$. \square

We can now prove Proposition 1.10.

Proof of Proposition 1.10. Since $\Omega_{\mathcal{C}}$ is a surface with smooth boundary, the spaces $W^{1,2}(\Omega_{\mathcal{C}})$ and $W_{\partial}^{1,2}(\Omega_{\mathcal{C}})$ are isomorphic, via some linear map ι . Interior conformal regularity $L \log L$ provides us with an isomorphism $\varphi^* : W^{1,2}(\Omega) \rightarrow W^{1,2}(\Omega_{\mathcal{C}})$ and boundary conformal regularity $L \log L$ with an isomorphism $\varphi_{\partial}^* : W_{\partial}(\Omega) \rightarrow W^{1,2}(\Omega_{\mathcal{C}})$. The composition $(\varphi_{\partial}^*)^{-1} \circ \iota \circ \varphi^*$ provides the desired isomorphism. \square

3. SPECTRAL ASYMPTOTICS

3.1. Eigenvalue counting functions of compact operators. We first present some known results about spectral asymptotics of compact operators defined via quadratic forms. These results can be found in the works of Suslina [Sus99b], Birman–Solomyak [BS80], and Sukochev–Zanin [SZ20], in a more general form. For the convenience of the reader we state them here in a form which is specific for our purposes.

Let \mathcal{H} be a Hilbert space and $K \in \mathcal{K}(\mathcal{H})$ be a self-adjoint nonnegative compact operator. The non-zero spectrum of K consists of a discrete set of nonincreasing nonnegative eigenvalues $\{\lambda_j(K) : j \in \mathbb{N}\}$ counted with multiplicity and converging to 0. The variational characterisation of the eigenvalues yields

$$\lambda_j(K) = \max_{E_j \subset \mathcal{H}} \min_{u \in E_j \setminus \{0\}} \frac{(Ku, u)}{(u, u)}, \quad (3.1)$$

where E_j ranges over j dimensional subspaces. Note that the operator K can be equivalently defined via the associated bilinear form appearing in the numerator (3.1); we will use this observation further on. For $\lambda > 0$, define the eigenvalue counting function

$$n(\lambda; K) := \#\{j : \lambda_j(K) > \lambda\},$$

and, for a given $\alpha > 0$, the functionals

$$\bar{n}_{\alpha}(K) = \limsup_{\lambda \searrow 0} \lambda^{\alpha} n(\lambda; K) \quad \text{and} \quad \underline{n}_{\alpha}(K) = \liminf_{\lambda \searrow 0} \lambda^{\alpha} n(\lambda; K).$$

Note that if $\bar{n}_{\alpha}(K) = \underline{n}_{\alpha}(K) = C_{\alpha}$, then

$$n(\lambda; K) = C_{\alpha} \lambda^{-\alpha} (1 + o(1)).$$

We make use of the following general properties of this counting function, which are collected in [BS80, Appendix 1], see also the references therein.

Lemma 3.1. *The following properties hold:*

- (1) [BS80, Lemma 1.16] *For any $\alpha > 0$, the functionals $\bar{n}_\alpha(K)$ and $\underline{n}_\alpha(K)$ are invariant under compact perturbations of the inner product on \mathcal{H} , as well as restriction to subspaces of finite codimension.*
- (2) [BS80, Lemma 1.18 and its proof], [Sus99b, Lemma 1.5], *Weyl–Fan Ky lemma.* *Let $K_1 \leq K_2 \in \mathcal{K}(\mathcal{H})$ be nonnegative self-adjoint compact operators. Then, for any $\alpha > 0$,*

$$\left| \bar{n}_\alpha(K_1)^{\frac{1}{1+\alpha}} - \bar{n}_\alpha(K_2)^{\frac{1}{1+\alpha}} \right| \leq \bar{n}_\alpha(K_2 - K_1)^{\frac{1}{1+\alpha}}$$

and

$$\left| \underline{n}_\alpha(K_1)^{\frac{1}{1+\alpha}} - \underline{n}_\alpha(K_2)^{\frac{1}{1+\alpha}} \right| \leq \bar{n}_\alpha(K_2 - K_1)^{\frac{1}{1+\alpha}}.$$

- (3) [BS80, Lemma 1.15] *Let $K_1 \in \mathcal{K}(\mathcal{H}_1)$ and $K_2 \in \mathcal{K}(\mathcal{H}_2)$ be nonnegative self-adjoint compact operators. Let $B : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ be a bounded operator such that $(K_1 u, u)_{\mathcal{H}_1} = 0$ for all $u \in \ker B$. If there is a $a > 0$ such that for all $u \in \mathcal{H}_1 \setminus \ker B$*

$$\frac{(K_1 u, u)_{\mathcal{H}_1}}{(u, u)_{\mathcal{H}_1}} \leq a \frac{(K_2 B u, B u)_{\mathcal{H}_2}}{(B u, B u)_{\mathcal{H}_2}},$$

then for all $\lambda > 0$, $n(\lambda; K_1) \leq n(a^{-1}\lambda; K_2)$ for all $\lambda > 0$.

We will use these abstract results in the concrete situation where $\mathcal{H} = W^{1/2,2}(\Gamma)$, where Γ is a finite collection of smooth curves with length measure $d\ell$. For $\beta : \Gamma \rightarrow [0, \infty)$ let K_β be the operator in $W^{1/2,2}(\Gamma)$ be defined by the bilinear form

$$(K_\beta u, v)_{\mathcal{H}} = \int_{\Gamma} u \bar{v} \beta \, d\ell, \quad u, v \in \text{Dom}(K_\beta). \quad (3.2)$$

The following lemma is a direct reinterpretation of [SZ20, Lemma 4.4] (see also [RS21, Theorem 2.1]) in view of the variational characterisation of the eigenvalue counting function [BS80, Lemma 1.14].

Lemma 3.2. *Let Γ be a finite collection of smooth curves and $0 \leq \beta \in L \log L(\Gamma)$. Let K_β be the self-adjoint operator on $W^{1/2,2}(\Gamma)$ defined via the bilinear form (3.2). Then there exists a constant $C(\Gamma) > 0$ such that*

$$n(\lambda; K_\beta) \leq C(\Gamma) \lambda^{-1} \|\beta\|_{L \log L(\Gamma)}.$$

We now have the required tools to prove Theorem 1.8.

3.2. Proof of Theorem 1.8. We turn to the second, equivalent, statement. Recall that the eigenvalues of the weighted Steklov problem on a surface with smooth boundary are characterised variationally as

$$\sigma_k(\Omega, \beta) = \min_{E_k} \max_{U \in E_k \setminus \{0\}} \frac{\int_{\Omega} |\nabla U|^2 \, dA_g}{\int_{\partial\Omega} u^2 \beta \, d\ell_g}. \quad (3.3)$$

Here $E_k \subset W^{1,2}(\Omega)$ (which is isomorphic to $W_{\partial}^{1,2}(\Omega)$) is a $k+1$ dimensional subspace, and $u := \tau U$, where $\tau : W^{1,2}(\Omega) \rightarrow W^{1/2,2}(\partial\Omega)$ is the trace operator, which is continuous. Here and further

on we adopt the following convention: capital letters denote functions in the interior, and the corresponding lower case letters denote their boundary traces.

Let

$$\mathcal{X} := \left\{ V \in W^{1,2}(\Omega) : \int_{\partial\Omega} v\beta \, d\ell_g = 0 \right\}.$$

be the orthogonal complement in $L^2(\partial\Omega; \beta \, d\ell_g)$ to the kernel of the weighted Dirichlet-to-Neumann map, i.e. to the constant functions. We equip \mathcal{X} with the inner product

$$(U, U)_{\mathcal{X}} = \int_{\Omega} |\nabla U|^2 \, dA_g.$$

Let us define an operator Q_{β} on \mathcal{X} via the bilinear form

$$(Q_{\beta}U, V)_{\mathcal{X}} = \int_{\partial\Omega} uv\beta \, d\ell_g, \quad u, v \in \mathcal{X}. \quad (3.4)$$

Clearly, we have that

$$\lambda_k(Q_{\beta}) = \max_{E_k \subset \mathcal{X}} \min_{u \in E_k \setminus \{0\}} \frac{\int_{\partial\Omega} u^2 \beta \, d\ell_g}{(U, U)_{\mathcal{X}}}. \quad (3.5)$$

In view of (3.5) and (3.3) we have that for $k \geq 1$, $\sigma_k(\Omega, \beta)^{-1} = \lambda_k(Q_{\beta})$, so that

$$N(\sigma; M, \beta) - 1 = n(\sigma^{-1}; Q_{\beta}), \quad (3.6)$$

where we have subtracted one on the left to account for the eigenvalue zero. Let us find the asymptotics of $n(\sigma^{-1}; Q_{\beta})$ as $\sigma^{-1} =: \lambda \searrow 0$. It follows from Lemma 3.1(1) that the asymptotics of $n(\lambda; Q_{\beta})$ does not change if we first replace $(U, U)_{\mathcal{X}}$ with $(U, U)_{\mathcal{X}} + (U, U)_{L^2(\Omega)}$ (this is a compact perturbation), and then lift the orthogonality condition, in order to consider $U \in W^{1,2}(\Omega)$ as in (3.3). By the density of smooth functions in $L \log L$, for every $\varepsilon > 0$ we can find a smooth $\beta_{\varepsilon} \in C^{\infty}(\partial\Omega)$ such that $\|\beta - \beta_{\varepsilon}\|_{L \log L} < \varepsilon$. Without loss of generality, we suppose that $\beta_{\varepsilon} \leq \beta$ almost everywhere so that $Q_{\beta} - Q_{\beta_{\varepsilon}}$ is a positive operator. Since we know, by the general theory of pseudodifferential operators, that as $\lambda \searrow 0$

$$n(\lambda; Q_{\beta_{\varepsilon}}) = \frac{\lambda^{-1}}{\pi} \int_{\partial\Omega} \beta_{\varepsilon} \, d\ell_g + o(\lambda^{-1}),$$

it is sufficient by Lemma 3.1(2) to show that

$$n(\lambda; Q_{\beta} - Q_{\beta_{\varepsilon}}) \leq C\lambda^{-1} \|\beta - \beta_{\varepsilon}\|_{L \log L},$$

with C depending only on Ω . It immediately follows from (3.4) that $\ker \tau \subset \ker(Q_{\beta} - Q_{\beta_{\varepsilon}})$. Defining K_{β} as in Lemma 3.2 with $\Gamma = \partial\Omega$, we have that for all $U \in W^{1,2}(\Omega)$,

$$((K_{\beta} - K_{\beta_{\varepsilon}})u, u)_{W^{1/2,2}(\partial\Omega)} = ((Q_{\beta} - Q_{\beta_{\varepsilon}})U, U)_{W^{1,2}(\Omega)}$$

By the trace theorem, we also have that there exists C_{Ω} such that

$$(\tau U, \tau U)_{W^{1/2,2}(\partial\Omega)} \leq C_{\Omega}(U, U)_{W^{1,2}(\Omega)}.$$

By applying first Lemma 3.1(3) then Lemma 3.2 we deduce that

$$\begin{aligned} n(\lambda; Q_\beta - Q_{\beta_\varepsilon}) &\leq n(C_\Omega \lambda; K_\beta - K_{\beta_\varepsilon}) \\ &\leq C'_\Omega \lambda^{-1} \|\beta - \beta_\varepsilon\|_{L \log L(\partial\Omega)} \\ &\leq C'_\Omega \lambda^{-1} \varepsilon. \end{aligned}$$

Since this holds for arbitrary $\varepsilon > 0$, we deduce that

$$n(\lambda; Q_\beta) = \frac{\lambda^{-1}}{\pi} \int_{\partial\Omega} \beta \, d\ell_g + o(\lambda^{-1})$$

and in view of (3.6) this completes the proof of the theorem.

4. EXAMPLES

In this last section, we present examples of domains having conformal regularity $L \log L$ and explore the sharpness of Theorem 1.8. We give planar domains as example, but they extend in a straightforward manner to domains in a complete Riemannian surface.

4.1. Chord-arc domains. Recall that a Jordan domain $\Omega \subset \mathbb{R}^2$ is called a *chord-arc* (or *Lavrentiev*) *domain* if there exists a constant C such that for any $x, y \in \partial\Omega$

$$\text{dist}_{\partial\Omega}(x, y) \leq C \text{dist}_{\mathbb{R}^2}(x, y),$$

where the left-hand side denotes the length of the shortest arc of the boundary joining x and y , and the right-hand side denotes the distance between x and y in \mathbb{R}^2 . It is clear that any Lipschitz domain is a chord-arc domain. The class of chord-arc domains is larger than Lipschitz and includes, in particular, the domain bounded between two logarithmic spirals. Note that domains with cusps are not chord-arc.

There is a large literature on the conformal regularity of chord-arc domains, see, for instance [Jer83] and references therein. The following result is well-known. We outline its proof below for the convenience of the reader.

Proposition 4.1. *Let $\Omega \subset \mathbb{R}^2$ be a chord-arc domain and let $\varphi : \mathbb{D} \rightarrow \Omega$ be a conformal map. Then $\varphi' \in L^p(\partial\Omega)$ for some $p > 1$.*

Proof. Since Ω is a chord-arc domain, by [Zin82, Theorem 1] we have that $\varphi' \in A_q$ for some $q > 1$, where A_q denotes a Muckenhoupt class of weights (see, for instance [Gar07, Section VI.6] for a definition). By [Gar07, Corollary 6.10], every Muckenhoupt weight on $\partial\Omega$ of class A_q , $q > 1$ is in $L^p(\partial\Omega)$ for some $p > 1$, which is our claim. \square

4.2. Domains with cusps. Let $\Omega \subset \mathbb{R}^2$ be a domain with boundary $\partial\Omega$ which is a finite union of smooth curves. If two curves meet at an interior angle zero we say that they form an *outward cusp*, and if the interior angle is equal to 2π we say that they form an *inward cusp*.

At the tip x_0 of an inward cusp $\varphi'(x_0) = 0$, in fact domains with inward cusps have $\varphi' \in C^{0,1}(\partial\Omega)$ [Pom92, Theorem 3.9]. A typical example is the standard cardioid domain defined in polar coordinates as $\{(r, \theta) : r = 2(1 + \cos \theta)\}$, for which $\varphi(z) = (z + 1)^2$.

Consider now domains with outward cusps. Suppose that in a neighbourhood of the outward cusp at x_0 the boundary of Ω consists of two smooth curves $\gamma_1(t)$, $\gamma_2(t)$, where t is the arc length

parameter and $\gamma_1(0) = \gamma_2(0) = x_0$. We say that Ω has a *slow* cusp at x_0 if there is $\alpha \in (0, 1)$ (the speed of the cusp) such that

$$\lim_{t \searrow 0} \frac{|\gamma_1(t) - \gamma_2(t)|}{t^{1+\alpha}} = s_\alpha > 0$$

In turn if there is C such that $t^{-2} |\gamma_1(t) - \gamma_2(t)| \leq C < \infty$ for all $t > 0$ we say that Ω has a *fast* cusp at x_0 .

It is shown in [NT08] that whenever a domain has a fast cusp, the Dirichlet-to-Neumann map does not have a compact resolvent. Therefore, its spectrum is not discrete and Weyl's law can not hold. However, the Dirichlet-to-Neumann map for a domain whose boundary is Lipschitz except at a finite number of slow cusps has a compact resolvent, and hence a discrete spectrum.

Let $\varphi : \Omega_C \rightarrow \Omega$ be a conformal diffeomorphism and let $z_0 = \varphi^{-1}(x_0)$ be the pre-image of a cusp of speed α . Applying [KL17, Proposition 2.10] to [Pro17, Corollary 1], we see that as $z \rightarrow z_0$ the conformal factor $|\mathrm{d}\varphi(z)|$ behaves asymptotically as

$$|\mathrm{d}\varphi(z)| = O\left(|z - z_0|^{-1} (-\log(|z - z_0|))^{-1 - \frac{1}{\alpha}}\right). \quad (4.1)$$

A direct calculation gives that $|\mathrm{d}\varphi| \in L \log L$ if and only if $0 < \alpha < 1$, in other words precisely when the spectrum is discrete. This shows that Theorem 1.8 gives in a sense an optimal condition for the validity of Weyl's law.

Let us summarize the results of this subsection in the following

Proposition 4.2. *Let $\Omega \subset \mathbb{R}^2$ be a domain with piecewise smooth boundary, possibly with interior and exterior cusps. If all exterior cusps are slow then Weyl's law (1.2) holds for the counting function of the Steklov eigenvalues on Ω . Moreover, if Ω has at least one fast cusp, then the Steklov spectrum of Ω is not discrete.*

Remark 4.3. It would be interesting to understand if there exist domains for which the Steklov spectrum is discrete but the Weyl's law (1.2) does not hold. To construct such an example one needs to find a domain Ω with the boundary conformal factor in $L^1 \setminus L \log L$, and yet for which the resolvent of the Dirichlet-to-Neumann map is still compact. We note that in terms of the weighted problem it seems like this would require going beyond the Orlicz scale: indeed, for every $0 \leq a < 1$ one can find $\beta \in L(\log L)^a(\partial\Omega)$ so that the embedding $W^{1,2}(\partial\Omega) \rightarrow L^2(\partial\Omega, \beta \mathrm{d}\ell)$ is not compact, following the proof found in [GKL21, Example 3.19].

5. FURTHER REMARKS AND EXTENSIONS

5.1. The Steklov problem with indefinite weight. Suppose for now that Ω is a surface with smooth boundary, and given $\beta : \partial\Omega \rightarrow \mathbb{R}$ consider the Steklov problem with an indefinite weight:

$$\begin{cases} \Delta u = 0 & \text{in } \Omega; \\ \partial_\nu u = \beta \sigma u & \text{on } \partial\Omega. \end{cases} \quad (5.1)$$

Indefinite eigenvalue problems of this type have been considered in the literature, see e.g. [San55, Sus99a, Sus99b, Agr06]. If $0 \neq \beta \in L \log L(\partial\Omega)$ changes sign on sets of positive measure in $\partial\Omega$, then the non-zero eigenvalues form two sequences $\{\sigma_k^\pm(\Omega, \beta) : k \in \mathbb{N}\}$ consisting of the positive and

negative eigenvalues, accumulating respectively at $\pm\infty$. To define the variational principle, let us first denote

$$\|f\|_{\mathbb{W}_\partial^{1,2}(\Omega;\beta)}^2 = \int_\Omega |\nabla f|^2 dv_g + \int_{\partial\Omega} f^2 |\beta| d\ell_g$$

and

$$\mathcal{W}_\partial(\Omega; \beta) := \left\{ f : \bar{\Omega} \rightarrow \mathbb{R} : f \in C^\infty(\Omega) \text{ and } \|f\|_{\mathbb{W}_\partial^{1,2}(\Omega;\beta)} < \infty \right\}.$$

We denote by $\mathbb{W}_\partial^{1,2}(\Omega; \beta)$ the closure of $\mathcal{W}_\partial(\Omega; \beta)$ under the $\|\cdot\|_{\mathbb{W}_\partial^{1,2}(\Omega;\beta)}$ norm, and \mathcal{X} to be the subset of $\mathbb{W}_\partial^{1,2}(\Omega; \beta)$ orthogonal to β . Following [BS80, Sus99b], the non-zero eigenvalues of problem (5.1) satisfy the variational principle

$$\frac{\pm 1}{\sigma_k^\pm(\Omega, \beta)} = \min_{F_k} \max_{u \in F_k \setminus \{0\}} \pm \frac{\int_{\partial\Omega} u^2 \beta d\ell_g}{\int_\Omega |\nabla u|^2 dA_g},$$

where F_k is a codimension $k-1$ subspace of \mathcal{X} . Denoting by

$$N^\pm(\sigma; \Omega, \beta) := \# \{k : 0 < \pm\sigma_k^\pm(\Omega, \beta) < \sigma\}$$

the counting functions for each of those sequences, it follows from the work of Birman–Solomyak [BS77, BS79] (see [Pon21, Theorem 6.1] for a modern proof, in English) that if β is smooth, then

$$N^\pm(\sigma; \Omega, \beta) = \frac{\sigma}{\pi} \int_{\partial\Omega} \beta_\pm d\ell + o(\sigma) \quad (5.2)$$

where $\beta_\pm = \max\{0, \pm\beta\}$ are the positive and negative parts of β . This formula is valid whether or not β takes both positive and negative values. Using the same methods as in Section 3 allows us to extend this result to $\beta \in L \log L(\partial\Omega)$. We note that the results in Lemma 3.1 are in fact proven in [BS80, Sus99b] for operators with both positive and negative spectrum, with the obvious redefinition of the functions \bar{n}_α^\pm and n_α^\pm .

When Ω has non-smooth boundary, we consider once again a conformal map $\varphi : \Omega_C \rightarrow \Omega$. If the product $\varphi^* \beta |d\varphi| \in L \log L(\partial\Omega_C)$, then the proof of Proposition 1.10 carries through and φ induces an isomorphism φ^* between $\mathbb{W}_\partial^{1,2}(\Omega; \beta)$ and $\mathbb{W}_\partial^{1,2}(\Omega_C; \varphi^* \beta)$. If both $\varphi^* \beta$ and $\varphi^* \beta |d\varphi|$ are in $L \log L(\partial\Omega_C)$ then

$$\mathbb{W}_\partial^{1,2}(\Omega_C; \varphi^* \beta) \cong \mathbb{W}_\partial^{1,2}(\Omega_C) \cong \mathbb{W}_\partial^{1,2}(\Omega_C; \varphi^* \beta |d\varphi|)$$

and as in Theorem 1.6 problem (5.1) is isospectral to

$$\begin{cases} \Delta u = 0 & \text{in } \Omega_C \\ \partial_\nu u = |d\varphi| \varphi^* \beta \sigma u & \text{on } \partial\Omega_C. \end{cases}$$

We see directly that a sufficient condition for having the Weyl law (5.2) is also that $|d\varphi| \varphi^* \beta \in L \log L(\partial\Omega_C)$. For any surface with Lipschitz boundary, we have that $|d\varphi|$ and $|d\varphi|^{-1}$ are in L^p for some $p > 1$ with Hölder conjugate p' , see [BBL16, proof of Lemma 5.1]. Therefore, if $\varphi^* \beta \in L^q(\partial\Omega_C)$ for $q > p'$, the Weyl law (5.2) holds. Arguing as in [GU16, Theorem 4], one can show that for $q > p'$ the map φ induces a bounded composition operator $\varphi^* : L^{\frac{qp}{p-1}}(\partial\Omega) \rightarrow L^q(\partial\Omega_C)$. Therefore, a sufficient condition for the Weyl law to hold is $\beta \in L^r(\partial\Omega)$, for some $r > p^2/(p-1)^2$. In particular, if Ω is a Lipschitz domain and β is in some Orlicz space contained in $L^q(\partial\Omega)$ for any $q < \infty$, then the weighted Steklov problem, definite or not, satisfies the Weyl law (5.2).

5.2. The Neumann problem. The methods developed in this paper can be also applied to the Neumann problem

$$\begin{cases} -\Delta_g u = \lambda u & \text{in } \Omega \\ \partial_\nu u = 0 & \text{on } \partial\Omega. \end{cases}$$

In this case, the conformal map $\varphi : \Omega_{\mathcal{C}} \rightarrow \Omega$ gives rise to the variationally isospectral weighted Neumann problem

$$\begin{cases} -\Delta_g u = \lambda |\mathrm{d}\varphi|^2 u & \text{in } \Omega_{\mathcal{C}} \\ \partial_\nu u = 0 & \text{on } \partial\Omega_{\mathcal{C}}. \end{cases}$$

If the boundary is regular enough, the Neumann spectrum is discrete and we aim for a Weyl law of the form

$$N_{\mathrm{Neu}}(\lambda) = \frac{\mathrm{Area}(\Omega)}{4\pi} \lambda + o(\lambda). \quad (5.3)$$

This problem is well studied, and, in particular, we already know that the Weyl élaw holds for a large class of domains with rough boundary. For instance, it is shown in [NS05] that (5.3) for every domain whose boundary is of the Hölder class $C^{0,\alpha}$ for $\alpha > 1/2$, see also [Net07] It is also shown in [NS05] that domains with finite straight cusps of any speed satisfy (5.3). Moreover, sharp remainder estimates have been obtained in many cases.

A straightforward adaptation of the methods developed in this paper yields an alternative proof of (5.3) provided $|\mathrm{d}\varphi|^2 \in L \log L(\Omega_{\mathcal{C}})$. This is shown in essentially the same way as Lemma 3.2. While this approach does not give sharp remainder estimates, it is significantly more elementary. Using (4.1), we note that the class of domains for which $|\mathrm{d}\varphi|^2 \in L \log L(\Omega_{\mathcal{C}})$ includes any cusp of polynomial speed.

REFERENCES

- [AA96] M. S. Agranovich and B. A. Amosov. Estimates for s -numbers, and spectral asymptotics for integral operators of potential type on nonsmooth surfaces. *Funktsional. Anal. i Prilozhen.*, 30(2):1–18, 96, 1996.
- [Agr06] M. S. Agranovich. On a mixed Poincaré-Steklov type spectral problem in a Lipschitz domain. *Russ. J. Math. Phys.*, 13(3):239–244, 2006.
- [AM12] W. Arendt and R. Mazzeo. Friedlander’s eigenvalue inequalities and the Dirichlet-to-Neumann semigroup. *Commun. Pure Appl. Anal.*, 11(6):2201–2212, 2012.
- [BBL16] L. Baratchart, L. Bourgeois, and J. Leblond. Uniqueness results for inverse Robin problems with bounded coefficient. *J. Funct. Anal.*, 270(7):2508–2542, 2016.
- [BK87] S. R. Bell and S. G. Krantz. Smoothness to the boundary of conformal maps. *Rocky Mountain J. Math.*, 17(1):23–40, 1987.
- [BS77] M. Š. Birman and M. Z. Solomjak. Asymptotic behavior of the spectrum of pseudodifferential operators with anisotropically homogeneous symbols. *Vestnik Leningrad. Univ.*, (13 Mat. Meh. Astronom. vyp. 3):13–21, 169, 1977.
- [BS79] M. Š. Birman and M. Z. Solomjak. Asymptotic behavior of the spectrum of pseudodifferential operators with anisotropically homogeneous symbols. II. *Vestnik Leningrad. Univ. Mat. Mekh. Astronom.*, (vyp. 3):5–10, 121, 1979.
- [BS80] M. Š. Birman and M. Z. Solomjak. *Quantitative analysis in Sobolev imbedding theorems and applications to spectral theory*, volume 114 of *American Mathematical Society Translations, Series 2*. American Mathematical Society, Providence, R.I., 1980. Translated from the Russian by F. A. Cezus.
- [BS88] C. Bennett and R. Sharpley. *Interpolation of operators*, volume 129 of *Pure and Applied Mathematics*. Academic Press, Inc., Boston, MA, 1988.

- [CP16] A. Cianchi and L. Pick. Optimal Sobolev trace embeddings. *Trans. Amer. Math. Soc.*, 368(12):8349–8382, 2016.
- [Edw93] J. Edward. An inverse spectral result for the Neumann operator on planar domains. *J. Funct. Anal.*, 111(2):312–322, 1993.
- [Gar07] J. B. Garnett. *Bounded analytic functions*, volume 236 of *Graduate Texts in Mathematics*. Springer, New York, first edition, 2007.
- [GKL21] A. Girouard, M. Karpukhin, and J. Lagacé. Continuity of eigenvalues and shape optimisation for Laplace and Steklov problems. *Geom. Funct. Anal.*, 31(3):513–561, 2021.
- [GKLP22] A. Girouard, M. Karpukhin, M. Levitin, and I. Polterovich. The Dirichlet-to-Neumann map, the boundary Laplacian and Hörmander’s rediscovered manuscript. *J. Spectr. Theory*, 2022. to appear.
- [GP10] A. Girouard and I. Polterovich. Shape optimization for low Neumann and Steklov eigenvalues. *Math. Methods Appl. Sci.*, 33(4):501–516, 2010.
- [GP17] A. Girouard and I. Polterovich. Spectral geometry of the Steklov problem (survey article). *J. Spectr. Theory*, 7(2):321–359, 2017.
- [GPPS14] A. Girouard, L. Parnowski, I. Polterovich, and D. Sher. The Steklov spectrum of surfaces: asymptotics and invariants. *Math. Proc. Cambridge Philos. Soc.*, 157(3):379–389, 2014.
- [GPU18] V. Gol’dshstein, V. Pchelintsev, and A. Ukhlov. Integral estimates of conformal derivatives and spectral properties of the Neumann-Laplacian. *J. Math. Anal. Appl.*, 463:19–39, 2018.
- [GU16] V. Gol’dshstein and A. Ukhlov. On the first eigenvalues of free vibrating membranes in conformal regular domains. *Arch. Ration. Mech. Anal.*, 221:893–915, 2016.
- [Haa84] A. Haas. Linearization and mappings onto pseudocircle domains. *Trans. Amer. Math. Soc.*, 282(1):415–429, 1984.
- [Jer83] D. Jerison. The failure of L^p estimates for harmonic measure in chord-arc domains. *Michigan Math. J.*, 30(2):191–198, 1983.
- [JS14] A. Jollivet and V. Sharafutdinov. On an inverse problem for the Steklov spectrum of a Riemannian surface. In *Inverse Problems and Applications*, volume 615 of *Contemporary Mathematics*, pages v+235. American Mathematical Society, 2014.
- [KL17] T. Kaiser and S. Lehner. Asymptotic behaviour of the Riemann mapping function at analytic cusps. *Ann. Acad. Sci. Fenn. Math.*, 42(1):3–15, 2017.
- [KS21] M. Karpukhin and D. Stern. From Steklov to Laplace: Free boundary minimal surfaces with many boundary components. *Preprint arXiv:2109.11029*, 2021.
- [Mas89] B. Maskit. Canonical domains on Riemann surfaces. *Proc. Amer. Math. Soc.*, 106(3):713–721, 1989.
- [Net07] Yu. Netrusov. Sharp remainder estimates in the Weyl formula for the Neumann Laplacian on a class of planar regions. *J. Funct. Anal.*, 250(1):21–41, 2007.
- [NS05] Yu. Netrusov and Yu. Safarov. Weyl asymptotic formula for the Laplacian on domains with rough boundaries. *Comm. Math. Phys.*, 253(2):481–509, 2005.
- [NT08] S. A. Nazarov and J. Taskinen. On the spectrum of the Steklov problem in a domain with a peak. *Vestnik St. Petersburg Univ. Math.*, 41(1):45–52, 2008.
- [Pom92] Ch. Pommerenke. *Boundary behaviour of conformal maps*, volume 299 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1992.
- [Pon21] R. Ponge. Connes’ integration and Weyl’s laws. 2021.
- [Pro17] D. Prokhorov. Conformal mapping asymptotics at a cusp. *Rev. Mat. Complut.*, 30(1):79–89, 2017.
- [Roz79] G. Rozenbljum. Asymptotic behavior of the eigenvalues for some two-dimensional spectral problems. In *Boundary value problems. Spectral theory (Russian)*, volume 7 of *Probl. Mat. Anal.*, pages 188–203, 245. Leningrad. Univ., Leningrad, 1979.
- [RS21] G. Rozenbljum and E. Shargorodsky. Eigenvalue estimates and asymptotics for weighted pseudodifferential operators with singular measures in the critical case. In *Partial Differential Equations, Spectral Theory, and Mathematical Physics: The Ari Laptev Anniversary Volume*, pages 331–354. European Mathematical Society, 2021.
- [San55] L. Sandgren. A vibration problem. *Medd. Lunds Univ. Mat. Sem.*, 13:1–84, 1955.

- [Sha71] S. E. Shamma. Asymptotic behavior of Stekloff eigenvalues and eigenfunctions. *SIAM J. Appl. Math.*, 20:482–490, 1971.
- [Sus99a] T. A. Suslina. Asymptotics of the spectrum for two model problems in the theory of liquid vibrations. In *Proceedings of the St. Petersburg Mathematical Society, Vol. IV*, volume 188 of *Amer. Math. Soc. Transl. Ser. 2*, pages 197–222. Amer. Math. Soc., Providence, RI, 1999.
- [Sus99b] T. A. Suslina. Spectral asymptotics of variational problems with elliptic constraints in domains with piecewise smooth boundary. *Russ. J. Math. Phys.*, 6(2):214–234, 1999.
- [Sus21] T. A. Suslina. Asymptotics of the spectrum of variational problems arising in the theory of fluid oscillations. *CMFD*, 67(2):363–407, 2021. (In Russian).
- [SZ20] F. Sukochev and D. Zanin. Cwikel–Solomyak estimates on tori and Euclidean spaces. *to appear in Mat. Sbornik.*, 2020.
- [Zin82] M. Zinsmeister. Courbes de Jordan vérifiant une condition corde-arc. *Ann. Inst. Fourier (Grenoble)*, 32(2):ix, 13–21, 1982.

MATHEMATICS 253-37, CALTECH, PASADENA, CA 91125, USA

Email address: `mikhailk@caltech.edu`

DEPARTMENT OF MATHEMATICS, KING'S COLLEGE LONDON, THE STRAND, LONDON, WC2R 2LS, UNITED KINGDOM

Email address: `jean.lagace@kcl.ac.uk`

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ DE MONTRÉAL, CP 6128, SUCC. CENTRE-VILLE, MONTRÉAL, QUÉBEC, H3C 3J7, CANADA

Email address: `iossif@dms.umontreal.ca`