

THE EXTREMAL FUNCTION FOR APEX GRAPHS

ELENA PAVELESCU

ABSTRACT. McCarty and Thomas conjectured that a linklessly embeddable graph with $n \geq 7$ vertices and t triangles has at most $3n - 9 + \frac{t}{3}$ edges. Thomas and Yoo proved this to be true for apex graphs. We give a shorter and simpler proof for the apex case.

1. INTRODUCTION

All graphs in this paper are finite and simple. A graph is *linklessly embeddable* if it can be embedded in \mathbb{R}^3 (or, equivalently, S^3) in such a way that no two cycles of the graph are linked. In [1], McCarty and Thomas proved that a bipartite linklessly embeddable graph on $n \geq 5$ vertices has at most $3n - 10$ edges, unless it is isomorphic to the complete bipartite graph $K_{3,n-3}$. They conjectured that a linklessly embeddable graph with $n \geq 7$ vertices and t triangles has at most $3n - 9 + \frac{t}{3}$ edges. Thomas and Yoo proved the conjecture holds for apex graphs [3]. A graph G is apex if it can be made planar by deleting one vertex. Apex graphs are known to be linklessly embeddable [2]. Here we give a simpler and shorter proof for the apex case. We use Euler's identity for connected planar graphs: $v - e + f = 2$, where v , e , and f represent the number of vertices, edges, and faces of a plane embedding.

2. MAIN RESULT

Theorem 1. *Every apex graph with $n \geq 7$ vertices and t triangles has at most $3n - 9 + \frac{t}{3}$ edges.*

Proof. We first show the theorem is true for maximal apex graphs. To construct a maximal apex graph G , start with a maximal planar graph H and add one vertex v together with all the edges from v to the vertices of H . We write $G = H * v$. If G has n vertices, then G has $4n - 10$ edges: $n - 1$ edges incident to v , and $3n - 9$ edges which are edges of H . The graph G has $5n - 15 + L$ triangles: $3n - 9$ triangles containing v and $2n - 6$ triangles which are the faces of a planar embedding of H , by Euler's identity, and L triangles of H which are not faces of a planar embedding of H . Since $4n - 10 \leq 3n - 9 + \frac{5n-15+L}{3}$, for $n \geq 7$, $L \geq 0$, the theorem is true for maximal apex graphs.

Let G be an apex graph with n vertices. Then G is obtained from a maximal apex graph $H * v$ through removal of edges. Assume G is obtained by removing k edges

Date: April 20, 2022.

2010 Mathematics Subject Classification. 05C35, 05C10.

Key words and phrases. extremal function, apex graph.

incident to v and s edges of H . Let $a_1, a_2, \dots, a_k \in V(H)$ and consider the graph G' obtained from $H * v$ by removing edges va_1, va_2, \dots, va_k . Then

$$(1) \quad t(H * v) - t(G') = |E(H)| - |E(H - \{a_1, a_2, \dots, a_k\})|,$$

since the triangles of $H * v$ containing the edge va_i correspond to the edges of H incident to the vertex a_i . We claim that for $k \geq 1$ and an arbitrary choice of vertices $a_1, a_2, \dots, a_k \in V(H)$,

$$|E(H)| - |E(H - \{a_1, a_2, \dots, a_k\})| \leq 2n + 3k - 12,$$

unless $k = 2$, and H has at least $2n - 5$ edges incident to a_1, a_2 , or both. This exceptional case is discussed at the very end of the proof.

To prove the claim, we consider $k = 1$, $k = 2$, and $k \geq 3$ separately.

For $k = 1$,

$$|E(H)| - |E(H - \{a_1\})| = \deg_H(a_1) \leq n - 2 \leq 2n + 3 - 12, \text{ for } n \geq 7.$$

For $k = 2$ in non-exceptional cases,

$$|E(H)| - |E(H - \{a_1, a_2\})| \leq 2n - 6 = 2n + 6 - 12.$$

For $k \geq 3$, the graph H has at most $3k - 6$ edges $a_i a_j$, with $1 \leq i < j \leq k$ (by Euler's identity), and at most $2(n - 1) - 4 = 2n - 6$ edges $a_i b_j$, with $1 \leq i \leq k$ and $1 \leq j \leq n - 1 - k$ (by Euler's identity for bipartite graphs). Then the claim follows, since

$$|E(H - \{a_1, a_2, \dots, a_k\})| \geq |E(H)| - (3k - 6) - (2n - 6) = |E(H)| - (2n + 3k - 12).$$

This last inequality together with (1) gives

$$(2) \quad t(H * v) - t(G') \leq 2n + 3k - 12,$$

meaning that when deleting k edges incident to v from $H * v$, at most $2n + 3k - 12$ triangles are deleted.

To obtain G , delete s edges of G' which are also edges of H . Each such edge belongs to at most one triangle containing the vertex v , at most two triangles which are faces of H , and possibly to some non-face triangles of H . These s edges of H belong to at most L non-face triangles of H . These observations yield

$$(3) \quad t(G') \leq t(G) + 3s + L.$$

Adding the inequalities (2) and (3) gives

$$5n - 15 + L = t(H * v) \leq t(G) + 2n + 3k + 3s + L - 12,$$

and further,

$$3n - 3k - 3s - 3 \leq t(G).$$

We check the inequality holds for the apex graph G :

$$(4) \quad \begin{aligned} |E(G)| &= 4n - 10 - k - s = (3n - 9) + (n - k - s - 1) = \\ &= 3n - 9 + \frac{3n - 3k - 3s - 3}{3} \leq 3n - 9 + \frac{t(G)}{3}. \end{aligned}$$

The exceptional case. Consider $a_1, a_2 \in V(H)$ such that H has at least $2n - 5$ edges incident to a_1, a_2 , or both. This is only possible if $\deg_H(a_1) = \deg_H(a_2) = n - 2$, that is, a_1 and a_2 are both adjacent to all other vertices of H and $a_1 a_2 \in E(H)$. The graph

H is isomorphic to the join $K_2 + P_{n-3}$, where P_{n-3} denotes the path with $n-3$ vertices. Exactly $2n-5$ edges of H are incident to a_1 , a_2 , or both. For $G' = H * v - \{va_1, va_2\}$,

$$(5) \quad t(H * v) - t(G') = 2n - 5.$$

To obtain G , delete s edges of G' which are also edges of H . Edge a_1a_2 belongs to $L+2$ triangles of G' . Edges a_1b_i (and a_2b_i), $i = 1, 2, \dots, n-3$, belong to at most three triangles of G' : two triangles which are faces in H and at most one non-face triangle of H . Edges b_ib_j , $1 \leq i < j \leq n-3$, belong to three triangles of G' : two triangles which are faces in H and one triangle containing the vertex v . These observations yield ($L \geq 1$ for $n \geq 7$)

$$(6) \quad t(G') \leq t(G) + L + 2 + 3(s-1) = t(G) + 3s + L - 1.$$

Adding the identities (5) and (6) gives

$$5n - 15 + L = t(H * v) \leq t(G) + 2n + 3s + L - 6,$$

and further

$$3n - 3s - 9 \leq t(G)$$

Since $k = 2$, this inequality is equivalent to

$$3n - 3k - 3s - 3 \leq t(G),$$

and the same check as in (4) completes the proof. □

Acknowledgements. The author would like to thank Youngho Yoo for noticing that non-face triangles were missing in the first version of the proof, and for suggesting a more detailed explanation for the cases $k = 1$ and $k = 2$.

REFERENCES

1. McCarty, R. and Thomas, R. *The extremal function for bipartite linklessly embeddable graphs*, Combinatorica **39**, no 5 (2019), 1081–1104
2. Sachs, H. *On spatial representations of finite graphs* A. Hajnal, L. Lovasz, V.T. Sós (Eds.), Colloq. Math. Soc. János Bolyai, Vol. **37**, North-Holland, Amsterdam, 1984, 649–662
3. Thomas, R. and Yoo, Y. *The extremal functions for triangle-free graphs with excluded minors*, European J. Combin. **75** (2019), 1–10.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SOUTH ALABAMA, MOBILE, AL, 36688, elenapavelescu@southalabama.edu