Boundary regularity of mixed local-nonlocal operators and its application

ANUP BISWAS, MITESH MODASIYA AND ABHROJYOTI SEN

ABSTRACT. Let Ω be a bounded C^2 domain in \mathbb{R}^n and $u \in C(\mathbb{R}^n)$ solves

 $\Delta u + aIu + C_0 |Du| \ge -K \quad \text{in } \Omega, \quad \Delta u + aIu - C_0 |Du| \le K \quad \text{in } \Omega, \quad u = 0 \quad \text{in } \Omega^c,$

in viscosity sense, where $0 \le a \le A_0$, $C_0, K \ge 0$, and I is a suitable nonlocal operator. We show that u/δ is in $C^{\kappa}(\bar{\Omega})$ for some $\kappa \in (0, 1)$, where $\delta(x) = \operatorname{dist}(x, \Omega^c)$. Using this result we also establish that $u \in C^{1,\gamma}(\bar{\Omega})$. Finally, we apply these results to study an overdetermined problem for mixed local-nonlocal operators.

1. INTRODUCTION AND RESULTS

We are interested in the integro-differential operator L of the form

$$Lu = \Delta u + aIu,$$

where $0 \le a \le A_0$, and I is a nonlocal operator given by

$$Iu(x) = \int_{\mathbb{R}^n} \left(u(x+y) - u(x) - \mathbb{1}_{\{|y| \le 1\}} y \cdot \nabla u(x) \right) k(y) \mathrm{d}y$$
$$= \frac{1}{2} \int_{\mathbb{R}^n} (u(x+y) + u(x-y) - 2u(x)) k(y) \mathrm{d}y$$

for some nonnegative, symmetric kernel k, that is, $k(y) = k(-y) \ge 0$ for all y. Operator L appears as a generator of a Lévy process which is obtained by superimposing a Brownian motion, running twice as fast as standard n-dimensional Brownian motion, and an independent pure-jump Lévy process corresponding to the nonlocal operator aI. Throughout this article we impose the following assumptions on the kernel k.

Assumption 1.1.

(a) For some $\alpha \in (0,2)$ and a kernel \hat{k} we have that, for all $r \in (0,1]$,

$$r^{n+\alpha}k(ry) \le k(y) \quad \text{for } y \in \mathbb{R}^n$$

and $\widehat{k}(y) = \frac{\Lambda}{|y|^{n+\alpha}} \mathbb{1}_{B_1}(y) + J(y)\mathbb{1}_{B_1^c}(y), \text{ where } \Lambda > 0 \text{ and}$
$$\int_{B_1^c} J(y) \mathrm{d}y < \infty.$$

(b) There exists a $\beta > 0$ such that for any $r \in (0,1]$ and $x_0 \in \mathbb{R}^n$ the following holds: for all $x, y \in B_{\frac{r}{2}}(x_0)$ and $z \in B_r^c(x_0)$ we have

$$k(x-z) \le \varrho k(y-z)$$
 for $|y-z| < \beta$,

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RE-SEARCH, DR. HOMI BHABHA ROAD, PUNE 411008, INDIA. EMAIL: ANUP@IISERPUNE.AC.IN; MITESH.MODASIYA@STUDENTS.IISERPUNE.AC.IN; ABHROJYOTI.SEN@ACADS.IISERPUNE.AC.IN

²⁰¹⁰ Mathematics Subject Classification. Primary: 35D40, 47G20, 35J61, 35B65 .

Key words and phrases. Operators of mixed order, semilinear equation, overdetermined problems, gradient estimate.

for some $\rho > 1$.

Assumption 1.1(a) will be used to study certain scaled operators and to find the exact behaviour of $I\delta(x)$ near the boundary where $\delta(x)$ denotes the distance function from the complement Ω^c . Assumption 1.1(b) will be used to apply the Harnack estimate from [24]. It should be noted that [24] uses a stronger hypothesis compared to Assumption 1.1(b). Assumption 1.1 is satisfied by a large class of nonlocal kernels as shown in the examples below.

Example 1.1. The following class of nonlocal kernels satisfy Assumption 1.1.

- (i) k(y) = 1/|y|^{n+α} for some α ∈ (0,2). More generally, we may take k(y) = 1/|y|^{n+α} 𝔅 g) for some ball B centered at the origin.
 (ii) k(y) ≍ Ψ(|y|⁻²)/|y|ⁿ where Ψ is Bernstein function vanishing at 0. In particular, Ψ is strictly
- (ii) $k(y) \simeq \frac{\Psi(|y|^{-1})}{|y|^n}$ where Ψ is Bernstein function vanishing at 0. In particular, Ψ is strictly increasing and concave. These class of nonlocal kernels correspond to a special class of Lévy processes, known as subordinate Brownian motions (see [37]). Assume that Ψ satisfies a global weak scaling property with parameters $\mu_1, \mu_2 \in (0, 1)$, that is,

$$\lambda^{\mu_1} \Psi(s) \lesssim \Psi(\lambda s) \lesssim \lambda^{\mu_2} \Psi(s) \quad \text{for } s > 0, \lambda \ge 1.$$

Then it is easily seen that

$$r^{n+2\mu_2}k(ry) \asymp r^{2\mu_2} \frac{\Psi(|ry|^{-2})}{|y|^n} \lesssim \frac{\Psi(|y|^{-2})}{|y|^n} \lesssim \mathbb{1}_{B_1}(y) \frac{\Psi(1)}{|y|^{n+2\mu_2}} + \mathbb{1}_{B_1^c}(y) \frac{\Psi(1)}{|y|^{n+2\mu_1}}$$

for all $r \in (0, 1]$. Thus Assumption 1.1(a) holds. Using the weak scaling property we can also check that Assumption 1.1(b) holds.

Let Ω be a bounded C^2 domain in \mathbb{R}^n . Let $u \in C(\mathbb{R}^n)$ be a viscosity solution to

$$Lu + C_0 |Du| \ge -K \quad \text{in } \Omega,$$

$$Lu - C_0 |Du| \le K \quad \text{in } \Omega,$$

$$u = 0 \qquad \text{in } \Omega^c,$$
(1.1)

for some nonnegative constants C_0, K . Though the results in this article are obtained for viscosity solutions, the results are also applied for weak solutions, see Remark 1.2 below for more details. Our equations in (1.1) are motived by the operators of the form

$$Lu + H(Du, x) := Lu + \inf_{\mu} \sup_{\nu} \{ b_{\mu,\nu}(x) \cdot Du(x) + f_{\mu,\nu}(x) \} = 0 \quad \text{in } \Omega, \quad u = 0 \quad \text{in } \Omega^c.$$
(1.2)

Such equations arise in the study of stochastic control problems where the control can influence the dynamics only through the drift $b_{\mu,\nu}$.

Integro-differential operators involving both local and nonlocal operators have gained interest very recently. See for instance, [1-6, 21, 25, 26, 32, 33]. For linear equations, one may recover up to the boundary $C^{1,\gamma}$ regularity of u from the $W^{2,p}$ regularity (cf. [25, Theorem 3.1.22]). Recently, inspired by [12, 13], interior regularity of the solutions of (1.2) are studied in [32, 33]. Let us also mention the recent works [17, 27] where interior Hölder regularity of the gradient is established for the weak solutions of degenerate elliptic equations of mixed type. In this article, we are interested in up to the boundary regularity of the solutions. It should also be noted that we are dealing with inequalities.

Our first result deals with the Lipschitz regularity of the solution.

Theorem 1.1. Suppose that Assumption 1.1(a) holds and $u \in C(\mathbb{R}^n)$ is a viscosity solution to (1.1). Then, for some constant C, dependent only on $n, \Omega, C_0, A_0, \hat{k}$, we have

$$\|u\|_{C^{0,1}(\mathbb{R}^n)} \le CK. \tag{1.3}$$

BOUNDARY REGULARITY

Proof of Theorem 1.1 is based on two standard ingredients, interior estimates from [33] and barrier function. It can be easily shown that $\operatorname{dist}(\cdot, \Omega^c)$ gives a barrier function at the boundary (cf. Lemma 2.1). Next we investigate finer regularity property of u near $\partial\Omega$. Let $\delta(x) = \operatorname{dist}(x, \Omega^c)$ be the distance function from the boundary. Modifying $\delta(x)$ inside Ω , if required, we may assume that $\delta \in C^2(\overline{\Omega})$ (cf. [16, Theorem 5.4.3]). Our next result establishes Hölder regularity of u/δ up to the boundary.

Theorem 1.2. Suppose that Assumption 1.1 holds. Let u be a viscosity solution to (1.1). Then there exists $\kappa \in (0, (2 - \alpha) \land 1)$ such that

$$\|u/\delta\|_{C^{\kappa}(\bar{\Omega})} \le C_1 K,\tag{1.4}$$

for some constant C_1 , where κ , C_1 depend on $n, C_0, A_0, \hat{k}, \Omega$.

The regularity of $\partial\Omega$ in the above result can be relaxed to $C^{1,1}$. See Remark 2.3 for more detail. For elliptic operators similar estimate is obtained by Krylov [30]. Boundary estimate for fractional Laplacian operators are studied by Ros-Oton and Serra in [34–36]. Result of [34] has been extended for nonlocal operators with kernel of variable orders by Kim et al. [29] whereas extension to the fractional *p*-Laplacian operator can be found in [31]. For the proof of Theorem 1.2 we follow the approach of [34] which is inspired by a method of Caffarelli [28, p. 39]. A key step in this analysis is the oscillation lemma (see Proposition 2.1) for u/δ which involves computation of $L((u - \kappa\delta)^+)$ for some suitable constant κ . Note that, by Theorem 1.1, Iu is bounded in Ω for $\alpha \in (0, 1)$, and therefore, in this case we can follow standard approach of local operators to get the estimate (1.4). But for $\alpha \in [1, 2)$, $I\delta$ becomes singular near $\partial\Omega$. So we have to do several careful estimates to apply the method of [34].

Using Theorem 1.2 we establish boundary regularity of Du (compare it with Fall-Jarohs [18]). This is the content of our next result.

Theorem 1.3. Let Assumption 1.1 hold. There exist constants γ , C, dependent on Ω , C_0 , A_0 , n, \hat{k} , such that for any solution u of (1.1) we have

$$\|u\|_{C^{1,\gamma}(\bar{\Omega})} \le CK. \tag{1.5}$$

Remark 1.1. By the dependency of the constants in Theorems 1.1 to 1.3 on \hat{k} we mean the dependency on α, Λ and $\int_{|y|>1} J(y) dy$.

To cite a specific application of the above results, let us consider $u, v \in C(\mathbb{R}^n)$ satisfying

$$Lu + H_1(Du, x) = 0, \quad \text{in } \Omega, \quad u = 0 \quad \text{in } \Omega^c,$$

$$Lv + H_2(Dv, x) = 0, \quad \text{in } \Omega, \quad v = 0 \quad \text{in } \Omega^c,$$

respectively. If $|H_1(p, x) - H_2(q, x)| \leq C_0 |p - q| + K$ for all $p, q \in \mathbb{R}^n$ and $x \in \Omega$, then using the interior regularity of u, v from [33] and the coupling result [10, Theorem 5.1] it can be easily seen that w = u - v satisfies (1.1). Our result Theorem 1.3 then gives a $C^{1,\gamma}$ estimate of w up to the boundary. The above results can also be used to establish anti-maximum principle for the generalized principal eigenvalues of nonlinear operators of the form (1.2) (cf. [7,9,15]).

Remark 1.2. Though the above results are mentioned for viscosity solutions, Theorem 1.1-1.3 can also be applied for weak solutions (at least for equations). To see this, let us assume that Ω be a $C^{2,\kappa}$ domain, $\kappa \in (0,1)$. Suppose that for some given Lipschitz function $f : \mathbb{R}^n \to \mathbb{R}$, there exists a unique weak solution $u \in H_0^1(\Omega)$ to

$$Lu + f(Du) = g \quad \text{in } \Omega, \quad u = 0 \quad \text{in } \Omega^c, \tag{1.6}$$

for every $g \in L^{\infty}(\Omega)$. Now consider a sequence of smooth mollifications g_{ε} of g such that $\sup_{\Omega} |g_{\varepsilon} - g| \to 0$, as $\varepsilon \to 0$. Let u_{ε} be the unique weak solution to (1.6) corresponding to g_{ε} . Since, by

Sobolev embedding $u_{x_i} \in L^p(\Omega)$ for $p \in [1, \frac{2n}{n-2}]$, applying [25, Theorem 3.1.22] and a bootstrapping argument we obtain that for some p > n

$$\|u_{\varepsilon}\|_{W^{2,p}(\Omega)} \le C \left(1 + \|Du_{\varepsilon}\|_{L^{2}(\Omega)} + \|g_{\varepsilon}\|_{L^{\infty}(\Omega)}\right), \qquad (1.7)$$

for some constant C independent of u_{ε} . This, of course, implies $u_{\varepsilon} \in C^{1,\gamma}(\bar{\Omega})$. Applying [25, Theorem 3.1.12] we have $u_{\varepsilon} \in C^{2,\gamma}(\bar{\Omega})$ and therefore, u_{ε} is a viscosity solution to (1.6) when g is replaced by g_{ε} . Hence we can apply Theorem 1.1-1.3 on u_{ε} . In particular,

$$\sup_{\varepsilon \in (0,1)} \|Du_{\varepsilon}\|_{L^{\infty}(\Omega)} < \infty.$$

Now, using the stability estimate (1.7), we can pass the limit, as $\varepsilon \to 0$, to show that $u_{\varepsilon} \to u$ where u is the weak solution to (1.6) with data g and u also satisfies the estimates in Theorem 1.1-1.3.

Next we apply Theorems 1.2 and 1.3 to study an overdetermined problem. More precisely, we consider a solution u to the problem

$$Lu + H(|Du|) = f(u) \quad \text{in } \Omega,$$

= 0 in Ω^c , $u > 0$ in Ω , $\frac{\partial u}{\partial n} = c \quad \text{on } \partial\Omega,$ (1.8)

where n is the unit inward normal and $H : \mathbb{R} \to \mathbb{R}, f : \mathbb{R}^n \to \mathbb{R}$ are locally Lipschitz. In Theorem 3.1 we show that Ω must be ball, provided the nonlocal kernel k satisfies certain conditions. Overdetermined problem was first studied in the celebrated work of Serrin [38] where it was shown that if there exists a positive solution to

$$-\Delta u = 1$$
 in Ω , $u = 0$, $\frac{\partial u}{\partial n} = c$ on $\partial \Omega$,

then Ω must be a ball. Serrin's work has been generalized for a vast class of operators, see for instance, [8,9,14,19,20,22,23,39]. In this article, we follow the method of [8,19] to establish our result on overdetermined problem concerning (1.8).

The rest of the article is organized as follows: In the next Section 2 we provide the proofs of Theorems 1.1 to 1.3 and Section 3 discusses the overdetermined problems.

2. Proofs of Theorems 1.1 to 1.3

We begin by showing that δ is a barrier function to u at the boundary.

u

Lemma 2.1. Suppose that Assumption 1.1(a) holds and u be a viscosity solution to (1.1). Then there exists a constant C, dependent only on $n, A_0, C_0, \operatorname{diam}(\Omega)$, radius of exterior sphere and $\int_{\mathbb{R}^n} (|y|^2 \wedge 1) \hat{k}(y) dy$, such that

$$|u(x)| \le CK\delta(x) \quad for \ all \ x \in \Omega, \tag{2.1}$$

where $\delta(x) = \operatorname{dist}(x, \Omega^c)$.

Proof. We first show that

$$|u(x)| \le \kappa K \quad x \in \mathbb{R}^n, \tag{2.2}$$

for some constant κ . From [32, Lemma 5.5] we can find a non-negative function $\chi \in C^2(\bar{\Omega}) \cap C_b(\mathbb{R}^n)$, with $\inf_{\mathbb{R}^n} \chi > 0$, satisfying

 $L\chi + C_0 |D\chi| \le -1$ in Ω .

Note that, since $\chi \in C^2(\overline{\Omega})$, the above equation holds in the classical sense. Defining $\psi = 2(K+\varepsilon)\chi$, $\varepsilon > 0$, we have that $\inf_{\mathbb{R}^n} \psi > 0$ and

$$L\psi + C_0 |D\psi| \le -2(K + \varepsilon) \quad \text{in } \Omega.$$
(2.3)

We claim that $u \leq \psi$ in \mathbb{R}^n . Suppose, on the contrary, that $(u - \psi)(z) > 0$ at some point in $z \in \Omega$. Define

$$\theta = \inf\{t : u \le t + \psi \text{ in } \mathbb{R}^n\}.$$

Since $(u - \psi)(z) > 0$, we must have $\theta \in (0, \infty)$. Again, since u = 0 in Ω^c , there must be a point $x_0 \in \Omega$ such that $u(x_0) = \theta + \psi(x_0)$ and $u \leq \theta + \psi$ in \mathbb{R}^n . Since ψ is C^2 in Ω , we get from the definition of viscosity subsolution that

$$-K \le L(\theta + \psi)(x_0) + C_0 |D\psi(x_0)| = L\psi(x_0) + C_0 |D\psi(x_0)| \le -2(K + \varepsilon),$$

using (2.3). But this is a contradiction. This proves the claim that $u \leq \psi$ in \mathbb{R}^n . Similar calculation using -u will also give us $-u \leq \psi$ in \mathbb{R}^n . Thus

$$|u| \le 2 \sup_{\mathbb{R}^n} |\chi| (K + \varepsilon)$$
 in \mathbb{R}^n .

Since ε is arbitrary, we get (2.2).

Now we can prove (2.1). In view of (2.2), it is enough to consider the case K > 0. Since Ω belongs to the class C^2 , it satisfies a uniform exterior sphere condition from outside. Let r_{\circ} be a radius satisfying uniform exterior condition. From [32, Lemma 5.4] there exists a bounded, Lipschitz continuous function φ , Lipschitz constant being r_{\circ}^{-1} , satisfying

$$\begin{split} \varphi &= 0 \quad \text{in} \quad \bar{B}_{r_{\circ}}, \\ \varphi &> 0 \quad \text{in} \quad \bar{B}^{c}_{r_{\circ}}, \\ \varphi &\geq \varepsilon \quad \text{in} \quad B^{c}_{(1+\delta)r_{\circ}}, \\ L\varphi + C_{0}|D\varphi| &\leq -1 \quad \text{in} \quad B_{(1+\delta)r_{\circ}} \setminus \bar{B}_{r_{\circ}}, \end{split}$$

for some constants ε, δ , dependent on C_0, A_0 and $\int_{\mathbb{R}^n} (|y|^2 \wedge 1) \hat{k}(y) dy$. Furthermore, φ is C^2 in $B_{(1+\delta)r_o} \setminus \overline{B}_{r_o}$. For any point $y \in \partial \Omega$, we can find another point $z \in \Omega^c$ such that $\overline{B}_{r_o}(z)$ touches $\partial \Omega$ at y. Let $w(x) = \varepsilon^{-1} \kappa K \varphi(x-z)$. Also $L(w) + C_0 |Dw| \leq -K$. Then

$$L(u-w) + C_0 |D(u-w)| \ge 0 \quad \text{in } B_{(1+\delta)r_o}(z) \cap \Omega$$

Since, by (2.2), $u - w \leq 0$ in $(B_{(1+\delta)r_{\circ}}(z) \cap \Omega)^c$, from comparison principle [10, Theorem 5.2] it follows that $u(x) \leq w(x)$ in \mathbb{R}^n . Note that the operator in [10, Theorem 5.2] does not have any gradient term, but the same proof (using the supersolution in [32, Lemma 5.5]) also gives a comparison principle for the above operator. Repeating a similar calculation for -u, we can conclude that $-u(x) \leq w(x)$ in \mathbb{R}^n . This relation holds for any $y \in \partial \Omega$. For any point $x \in \Omega$ with $\operatorname{dist}(x, \partial \Omega) < r_{\circ}$ we can find $y \in \partial \Omega$ satisfying $\operatorname{dist}(x, \partial \Omega) = |x - y| < r_{\circ}$. By previous estimate we then obtain

$$|u(x)| \le \varepsilon^{-1} \kappa K \varphi(x-z) \le \varepsilon^{-1} \kappa K (\varphi(x-z) - \varphi(y-z)) \le \varepsilon^{-1} \kappa K r_{\circ}^{-1} \operatorname{dist}(x, \partial \Omega).$$

es us (2.1).

This gives us (2.1).

Now we are ready to prove that $u \in C^{0,1}(\mathbb{R}^n)$.

Proof of Theorem 1.1. Let $x \in \Omega$ and $r \in (0,1)$ be such that $4r = \text{dist}(x, \partial \Omega) \wedge 1$. Without loss of any generality, we assume x = 0. Define v(y) = u(ry) in \mathbb{R}^n . From Lemma 2.1, we then have

$$|v(y)| \le C_1 K \min\{r^{\alpha/2}(1+|y|^{\alpha/2}), r(1+|y|)\} \qquad y \in \mathbb{R}^n,$$
(2.4)

for some constant C_1 . We let

$$I_r f(x) = r^{\alpha} \frac{1}{2} \int_{\mathbb{R}^n} (f(x+y) + f(x-y) - f(x)) k(ry) r^n dy,$$

and $L_r f = \Delta f + r^{2-\alpha} a I_r f$. Let us compute $L_r v(x) + C_0 r |Dv|$ in B_2 . Clearly, we have $\Delta v(x) = r^2 \Delta u(rx)$ and Dv(x) = r Du(rx). Also

$$r^{2-\alpha}I_r v(x) = r^2 \frac{1}{2} \int_{\mathbb{R}^n} (u(rx+ry) + u(rx-ry) - 2u(rx))k(ry)r^n dy$$

= $r^2 Iu(rx).$

Thus, it follows from (1.1) that

$$L_r v + C_0 r |Dv| \ge -Kr^2 \quad \text{in} \quad B_2,$$

$$L_r v - C_0 r |Dv| \le Kr^2 \quad \text{in} \quad B_2.$$
(2.5)

Now consider a smooth cut-off function $\varphi, 0 \leq \varphi \leq 1,$ satisfying

$$\varphi = \begin{cases} 1 & \text{in } B_{3/2}, \\ 0 & \text{in } B_2^c. \end{cases}$$

Let $w = \varphi v$. Clearly, $((\varphi - 1)v)(y) = 0$ for all $y \in B_{3/2}$, which gives $D((\varphi - 1)v) = 0$ and $\Delta((\varphi - 1)v) = 0$ in $x \in B_{3/2}$. Since $w = v + (\varphi - 1)v$, we obtain

$$L_r w + C_0 r |Dw| \ge -Kr^2 - A_0 r^{2-\alpha} |I_r((\varphi - 1)v)| \quad \text{in} \quad B_1,$$

$$L_r w - C_0 r |Dw| \le Kr^2 + A_0 r^{2-\alpha} |I_r((\varphi - 1)v)| \quad \text{in} \quad B_1,$$
(2.6)

from (2.5). Again, since $(\varphi - 1)v = 0$ in $B_{3/2}$, we have in B_1 that

$$\begin{aligned} |r^{2-\alpha}I_r((\varphi-1)v)(x)| &= r^{n+2}\frac{1}{2} \Big| \int_{|y|\ge 1/2} ((\varphi-1)v)(x+y) + ((\varphi-1)v)(x-y)k(ry)dy \Big| \\ &\leq r^{n+2}\int_{|y|\ge 1/2} |v(x+y)|k(ry)dy \\ &\leq r^{n+2}\int_{\frac{1}{2}\le |y|\le \frac{1}{r}} |v(x+y)|k(ry)dy + r^{n+2}\int_{r|y|>1} |v(x+y)|k(ry)dy, \\ &:= I_{r,1} + I_{r,2}. \end{aligned}$$

By Assumption 1.1(a)

$$\begin{split} I_{r,1} &= r^{n+2} \int_{\frac{1}{2} \le |y| \le \frac{1}{r}} |v(x+y)| k(ry) \mathrm{d}y \le r^{2-\alpha} \int_{\frac{1}{2} \le |y| \le \frac{1}{r}} |v(x+y)| \widehat{k}(y) \mathrm{d}y \\ &= \Lambda r^{2-\alpha} \int_{\frac{1}{2} \le |y| \le \frac{1}{r}} |v(x+y)| \frac{1}{|y|^{n+\alpha}} \mathrm{d}y \\ &\le r^{2-\alpha} \Lambda 3^{n+\alpha} \int_{|y| \ge 1/2} \frac{|v(x+y)|}{1+|x+y|^{\alpha/2}} \frac{1+|x+y|^{\alpha/2}}{1+|y|^{n+\alpha}} \mathrm{d}y \\ &\le \kappa_2 K r^{2-\alpha/2} \int_{|y| \ge 1/2} \frac{1+|x+y|^{\alpha/2}}{1+|x+y|^{n+\alpha}} \mathrm{d}y \\ &\le \kappa_3 K r^{2-\alpha/2}, \end{split}$$

for some constants κ_2, κ_3 , and in the fifth line we use (2.4). Again, by (2.2), we have

$$I_{r,2} \leq \kappa r^2 K \int_{r|y|>1} r^n k(ry) dy = \kappa r^2 K \int_{|y|>1} k(y) dy$$
$$\leq \kappa r^2 K \int_{|y|>1} \widehat{k}(y) dy \leq \kappa r^2 K \int_{|y|>1} J(y) dy \leq \kappa_4 r^2 K,$$

for some constant κ_4 . Therefore, putting the estimates of $I_{1,r}$ and $I_{2,r}$ in (2.6) we obtain

$$L_r v + C_0 r |Dv| \ge -\kappa_5 K r^{2-\alpha/2}$$
 in B_1 , (2.7)

$$L_r v - C_0 r |Dv| \ge \kappa_5 K r^{2-\alpha/2} \quad \text{in} \quad B_1,$$

for some constant κ_5 . Applying [33, Theorem 4.1] we obtain from (2.7)

$$\|v\|_{C^{1}(B_{\frac{1}{2}})} \le \kappa_{6} \left(\|v\|_{L^{\infty}(B_{2})} + r^{2-\alpha/2}K \right)$$
(2.8)

for some constant κ_6 . The proof in [33, Theorem 4.1] is stated for equations but it is easily seen that the same proof also works for a system of inequalities as in (2.7). From (2.4) and (2.8) we then obtain

$$\sup_{y \in B_{r/2}(x), y \neq x} \frac{|u(x) - u(y)|}{|x - y|} \le \kappa_7 K,$$
(2.9)

for some constant κ_7 .

Now we can complete the proof. Not that if $|x - y| \ge \frac{1}{8}$, then

$$\frac{|u(x) - u(y)|}{|x - y|} \le 2\kappa K,$$

by (2.2). So we consider $|x - y| < \frac{1}{8}$. If $|x - y| \ge 8^{-1}(\delta(x) \lor \delta(y))$, then using Lemma 2.1 we get

$$\frac{u(x) - u(y)|}{|x - y|} \le 4CK(\delta(x) + \delta(y))(\delta(x) \lor \delta(y))^{-1} \le 8CK.$$

Now let $|x - y| < 8^{-1} \min\{\delta(x) \lor \delta(y), 1\}$. Then either $y \in B_{\frac{\delta(x) \land 1}{8}}(x)$ or $x \in B_{\frac{\delta(y) \land 1}{8}}(y)$. Without loss of generality, we suppose $y \in B_{\frac{\delta(x) \land 1}{8}}(x)$. From (2.9) we get

$$\frac{|u(x) - u(y)|}{|x - y|} \le \kappa_7 K.$$

This completes the proof.

With the help of Theorem 1.1 we may choose $\beta = \infty$ in Assumption 1.1(b).

Remark 2.1. Since u is globally Lipschitz, choosing $\mathcal{J}(y) = |y|^{-n-\zeta}, \zeta \in (0, 1 \land \alpha)$, we see from Theorem 1.1 that

$$\left|\int_{\mathbb{R}^{n}} (u(x+y) + u(x-y) - 2u(x))\mathcal{J}(y)dy\right| \le \kappa \|u\|_{C^{0,1}(\mathbb{R}^{n})} \le \kappa CK$$
(2.10)

for some constant κ . Let $\tilde{k}(y) = k(y) \mathbb{1}_{\{|y| \le \beta'\}} + \mathcal{J}(y)$, where $\beta' < \frac{2}{3}\beta$. It is easy to see that \tilde{k} satisfies Assumption 1.1(a).

We now show that Assumption 1.1(b) also holds for this kernel with $\beta = \infty$. Fix $r \in (0, 1]$ and $x_0 \in \mathbb{R}^n$ and choose $x, y \in B_{\frac{r}{2}}(x_0)$ and $z \in B_r^c(x_0)$. Without any loss of generality we may assume that given β is in (0, 1/2). If $\max\{|x - z|, |y - z|\} \leq \beta'$, we have from Assumption 1.1(b) that

$$k(x-z) = k(x-z) + \mathcal{J}(x-z) \le \varrho k(y-z) + 3^{n+\zeta} \mathcal{J}(y-z) \le (\varrho \lor 3^{n+\zeta}) k(y-z),$$

using the fact

$$\frac{1}{3}|x-z| \le |y-z| \le 3|x-z|.$$

Also, if $|x - z| > \beta'$, then

$$\tilde{k}(x-z) = \mathcal{J}(x-z) \le 3^{n+\zeta} \tilde{k}(y-z).$$

Suppose $|x - z| \leq \beta'$ and $|y - z| > \beta'$. Note that $|y - z| \leq \frac{3}{2}\beta' < \beta$. Using Assumption 1.1(a) we find that

$$\tilde{k}(x-z) \le \varrho k(y-z) + 3^{n+\zeta} \mathcal{J}(y-z) \le \varrho \Lambda |y-z|^{-n-\alpha} + 3^{n+\zeta} \mathcal{J}(y-z)$$

$$\leq \left(\varrho\Lambda(\beta')^{-\alpha+\zeta} + 3^{n+\zeta}\right)\mathcal{J}(y-z)$$
$$= \left(\varrho\Lambda(\beta')^{-\alpha+\zeta} + 3^{n+\zeta}\right)\tilde{k}(y-z).$$

Thus, the kernel \tilde{k} satisfies Assumption 1.1(b) for $\beta = \infty$.

On the other hand, replacing the kernel k by \tilde{k} and using Theorem 1.1, (2.10), we obtain from (1.1) that

$$\begin{aligned} \Delta u + aI_{\tilde{k}}u + C_0|Du| &\geq -C_1 K \quad \text{in } \Omega, \\ \Delta u + aI_{\tilde{k}}u - C_0|Du| &\leq C_1 K \quad \text{in } \Omega, \\ u &= 0 \quad \text{in } \Omega^c, \end{aligned}$$
(2.11)

for some constant C_1 , dependent on \hat{k}, ζ, A_0 . This modification of nonlocal kernel would be useful apply the Harnack inequality from [24].

2.1. Fine boundary regularity. In this section we study the regularity of u/δ in Ω . Since u is Lipschitz, using the estimate (1.3) we may write (1.1) as follows

$$|Lu| = |\Delta u + aIu| \le CK \quad \text{in } \Omega, \quad \text{and} \quad u = 0 \quad \text{in } \Omega^c, \tag{2.12}$$

where C is a constant depending on \hat{k} , A_0 , C_0 . Also, in view of Remark 2.1, we can assume that k satisfies Assumption 1.1(b) for $\beta = \infty$. The rest of the section is devoted to the proofs of Theorems 1.2 and 1.3. Towards the proof of Theorem 1.2, our first goal is to get the oscillation estimate Proposition 2.1. To obtain this result we need Lemmas 2.2 to 2.6.

Lemma 2.2. There exists a constant $\tilde{\kappa}$, dependent on n, A_0, \hat{k} , such that for any $r \in (0, 1]$, we have a bounded radial function ϕ_r satisfying

$$\begin{cases} L\phi_r \ge 0 & \text{in } B_{4r} \setminus B_r, \\ 0 \le \phi_r \le \tilde{\kappa}r & \text{in } B_r, \\ \phi_r \ge \frac{1}{\tilde{\kappa}}(4r - |x|) & \text{in } B_{4r} \setminus B_r, \\ \phi_r \le 0 & \text{in } \mathbb{R}^n \setminus B_{4r}. \end{cases}$$

Moreover, $\phi_r \in C^2(B_{4r} \setminus \overline{B}_r).$

Proof. Fix $r \in (0,1]$ and define $v_r(x) = e^{-\eta q(x)} - e^{-\eta (4r)^2}$, where $q(x) = |x|^2 \wedge 2(4r)^2$ and $\eta > 0$. Clearly, $1 \ge v_r(0) \ge v_r(x)$ for all $x \in \mathbb{R}^n$. Thus

$$v_r(x) \le 1 - e^{-\eta(4r)^2} \le \eta(4r)^2,$$
(2.13)

using the fact that $1 - e^{-s} \leq s$ for all $s \geq 0$. Again, for $x \in B_{4r} \setminus B_r$, we have

$$v_r(x) = e^{-\eta(4r)^2} (e^{\eta((4r)^2 - q(x))} - 1) \ge \eta e^{-\eta(4r)^2} ((4r)^2 - |x|^2)$$

= $\eta e^{-\eta(4r)^2} (4r + |x|)(4r - |x|) \ge 5\eta r e^{-\eta(4r)^2} (4r - |x|).$ (2.14)

Now we estimate Lv_r in $B_{4r} \setminus \overline{B}_r$. Fix $x \in B_{4r} \setminus \overline{B}_r$. Then

$$\Delta v_r = \eta e^{-\eta |x|^2} \left(4\eta |x|^2 - 2n \right),$$

and, since $Iv_r = I(v_r + e^{-\eta(4r)^2})$, using the convexity of exponential map we obtain

$$I(e^{-\eta q(\cdot)})(x) \ge -\eta e^{-\eta |x|^2} \int_{\mathbb{R}^n} \left(q(x+y) + q(x-y) - 2q(x) \right) k(y) dy$$

$$\ge -\eta e^{-\eta |x|^2} \left[\int_{|y| \le 1} \left(q(x+y) + q(x-y) - 2q(x) \right) k(y) dy + \int_{|y| > 1} (8r)^2 k(y) dy \right]$$

BOUNDARY REGULARITY

$$\geq -\eta e^{-\eta |x|^2} \left[\int_{|y| < r} \frac{2|y|^2}{|y|^{n+\alpha}} \mathrm{d}y + \int_{r < |y| < 1} \frac{(8r)^2}{|y|^{n+\alpha}} \mathrm{d}y + (8r)^2 \int_{|y| > 1} J(y) \mathrm{d}y \right]$$

$$\geq -\eta e^{-\eta |x|^2} \kappa r^{2-\alpha},$$

for some constant κ , independent of η . Combining the above estimates we see that, for $x \in B_{4r} \setminus \overline{B}_r$,

$$Lv_r(x) \ge \eta e^{-\eta |x|^2} \Big[4\eta |x|^2 - 2n - A_0 \kappa r^{2-\alpha} \Big] \ge \eta e^{-\eta |x|^2} \Big[4\eta r^2 - 2n - A_0 \kappa r^{2-\alpha} \Big].$$

Thus, letting $\eta = \frac{1}{r^2}(n + A_0\kappa)$, we obtain

$$Lv_r > 0$$
 in $B_{4r} \setminus B_r$

We set $\phi_r = \frac{v_r}{r}$ and the result follows from (2.13)-(2.14).

Let us now define the sets that we use for our oscillation estimates. We borrow the notations of [34].

Definition 2.1. Let $\kappa \in (0, \frac{1}{16})$ be a fixed small constant and let $\kappa' = 1/2 + 2\kappa$. Given a point $x_0 \in \partial\Omega$ and R > 0, we define

$$D_R = D_R(x_0) = B_R(x_0) \cap \Omega,$$

and

$$D_{\kappa'R}^{+} = D_{\kappa'R}^{+}(x_0) = B_{\kappa'R}^{+}(x_0) \cap \{x \in \Omega : (x - x_0) \cdot \mathbf{n}(x_0) \ge 2\kappa R\},\$$

where $n(x_0)$ is the unit inward normal at x_0 . Using the C^2 regularity of the domain, there exists $\rho > 0$, depending on Ω , such that the following inclusions hold for each $x_0 \in \partial \Omega$ and $R \leq \rho$:

$$B_{\kappa R}(y) \subset D_R(x_0) \qquad \text{for all } y \in D^+_{\kappa' R}(x_0), \tag{2.15}$$

and

 $B_{4\kappa R}(y^* + 4\kappa Rn(y^*)) \subset D_R(x_0)$, and $B_{\kappa R}(y^* + 4\kappa Rn(y^*)) \subset D^+_{\kappa'R}(x_0)$ (2.16) for all $y \in D_{R/2}$, where $y^* \in \partial\Omega$ is the unique boundary point satisfying $|y - y^*| = \operatorname{dist}(y, \partial\Omega)$. Note that, since $R \leq \rho$, $y \in D_{R/2}$ is close enough to $\partial\Omega$ and hence the point $y^* + 4\kappa Rn(y^*)$ belongs to the line joining y and y^* .

Remark 2.2. In the remaining part of this section, we fix $\rho > 0$ to be a small constant depending only on Ω , so that (2.15)-(2.16) hold whenever $R \leq \rho$ and $x_0 \in \partial \Omega$. Also, every point on $\partial \Omega$ can be touched from both inside and outside Ω by balls of radius ρ . We also fix $\gamma > 0$ small enough so that for $0 < r \leq \rho$ and $x_0 \in \partial \Omega$ we have

$$B_{\eta r}(x_0) \cap \Omega \subset B_{(1+\sigma)r}(z) \setminus \overline{B}_r(z) \quad \text{for} \quad \eta = \sigma/8, \ \sigma \in (0,\gamma),$$

for any $x' \in \partial \Omega \cap B_{\eta r}(x_0)$, where $B_r(z)$ is a ball contained in $\mathbb{R}^n \setminus \Omega$ that touches $\partial \Omega$ at point x'.

We first treat the case $\alpha \in (0, 1)$. Note that in this situation Iu can be defined in the classical sense and is bounded in Ω , by Theorem 1.1.

Lemma 2.3. Let $\alpha \in (0,1)$ and Ω be a bounded C^2 domain. Let u be such that $u \ge 0$ in \mathbb{R}^n , and $|Lu| \le C_2$ in D_R , for some constant C_2 . Then, there exists a positive constant C, depending only on n, Ω, A_0, \hat{k} , such that

$$\inf_{D_{\kappa'R}^+} \left(\frac{u}{\delta}\right) \le C \left(\inf_{D_{\frac{R}{2}}} \frac{u}{\delta} + C_2 R\right)$$
(2.17)

for all $R \leq \rho_0$, where the constant ρ_0 depends only on n, Ω, A_0 and $\int_{\mathbb{R}^n} (|y|^2 \wedge 1) \widehat{k}(y) dy$.

Proof. We split the proof in two steps.

Step 1. Suppose $C_2 = 0$ and $R \le \rho$, where ρ is given by Remark 2.2. Define $m = \inf_{\substack{D_{\kappa'R} \\ \kappa'R}} u/\delta \ge 0$. By (2.15),

$$u \ge m\delta \ge m(\kappa R)$$
 in $D^+_{\kappa' R}$. (2.18)

Again, by (2.16), for any $y \in D_{R/2}$, we have either $y \in D_{\kappa'R}^+$ or $\delta(y) < 4\kappa R$. If $y \in D_{\kappa'R}^+$ it follows from the definition of m that $m \le u(y)/\delta(y)$. Now let $\delta(y) < 4\kappa R$. Let y^* be the nearest point to y on $\partial\Omega$ and $\tilde{y} = y^* + 4\kappa R n(y^*)$. Again by (2.16), we have $B_{4\kappa R}(\tilde{y}) \subset D_R$ and $B_{\kappa R}(\tilde{y}) \subset D_{\kappa'R}^+$. Recall that Lu = 0 in D_R and $u \ge 0$ in \mathbb{R}^n .

Now take $r = \kappa R$ and let ϕ_r be the subsolution in Lemma 2.2. Define $\tilde{\phi}_r(x) = \frac{1}{\tilde{\kappa}}\phi_r(x-\tilde{y})$. Using (2.18) and the comparison principle [10, Theorem 5.2] in $B_{4r}(\tilde{y}) \setminus \bar{B}_r(\tilde{y})$ it follows that $u(x) \ge m\tilde{\phi}_r(x)$ in all of \mathbb{R}^n . In particular, we have $u/\delta \ge \frac{1}{(\tilde{\kappa})^2}m$ on the segment joining y^* and \tilde{y} , that contains y. Hence

$$\inf_{D_{\kappa'R}^+} \left(\frac{u}{\delta}\right) \le C \inf_{D_{\frac{R}{2}}} \frac{u}{\delta}.$$

Step 2. Suppose $C_2 > 0$. Define $r' = \eta r$ for $r \leq \rho$ and $\eta \leq 1$ to be chosen later. Let \tilde{u} to be the solution of (cf. [10, Theorem 1.1])

$$\begin{cases} L\tilde{u} = 0 & \text{ in } D_{r'} ,\\ \tilde{u} = u & \text{ in } \mathbb{R}^n \setminus D_{r'} , \end{cases}$$

From step 1, we see that \tilde{u} satisfies (2.17). Define $w = \tilde{u} - u$. Applying [10, Theorem 5.1], we obtain that $|Lw| \leq C_2$ in $D_{r'}$ and w = 0 in $\mathbb{R}^n \setminus D_{r'}$. Since $r \leq \rho$, points of $\partial\Omega$ can be touched by exterior ball of radius r. Thus for any point $y \in \partial\Omega$ we can find another point $z \in \Omega^c$ such that $\bar{B}_r(z)$ touches $\partial\Omega$ at y. From the proof of [32, Lemma 5.4] there exists a bounded, Lipschitz continuous function φ_r , with Lipschitz constant r^{-1} , that satisfies

$$\begin{cases} \varphi_r = 0, & \text{in } \bar{B}_r, \\ \varphi_r > 0, & \text{in } \bar{B}_r^c, \\ L\varphi_r \le -\frac{1}{r^2}, & \text{in } B_{(1+\sigma)r} \setminus \bar{B}_r, \end{cases}$$

for some constant σ , independent of r. Without any loss of generality we may assume $\sigma \leq \gamma$ (see Remark 2.2). We set $\eta = \frac{\sigma}{8}$. Then $D_{r'} \subset B_{(1+\sigma)r}(z) \setminus \overline{B}_r(z)$. Letting $v(x) = C_2 r^2 \varphi_r(x-z)$ will give us a desired supersolution and therefore, by comparison principle we get $|w| \leq v$ in \mathbb{R}^n . For any point $x \in D_{r'}$ we can find $y \in \partial\Omega$ satisfying dist $(x, \partial\Omega) = |x-y|$. By above estimate we obtain

$$|w(x)| \le C_2 r^2 \varphi_r(x-z) \le C_2 r^2 (\varphi_r(x-z) - \varphi_r(y-z)) \le C_2 r \operatorname{dist}(x, \partial \Omega) = C_2 r \delta(x).$$

Thus we obtain

$$|w(x)| \le C_2 \frac{r'}{\eta} \delta(x)$$
 in $D_{r'}$.

Combining with step 1 we have

$$\inf_{D_{\kappa'r'}^+} \left(\frac{u}{\delta}\right) \le \frac{C}{\eta} \left(\inf_{D_{\frac{r'}{2}}} \frac{u}{\delta} + C_2 r'\right).$$

Setting $\rho_0 = \eta \rho$ and R = r' we have the desired result.

Next we obtain a similar estimate when $\alpha \in [1, 2)$.

Lemma 2.4. Let Ω be a bounded C^2 domain and u be such that $u \ge 0$ in all of \mathbb{R}^n and $|Lu| \le C_2 g$ in D_R for some positive constant C_2 and g is given by

$$g(x) = \begin{cases} (\delta(x))^{1-\alpha} & \text{if } \alpha > 1, \\ -\log(\delta(x)) + C_3 & \text{if } \alpha = 1, \end{cases}$$

for some constant C_3 . Set $\hat{\alpha} = 2 - \alpha$ for $\alpha \in (1, 2)$ and for $\alpha = 1$, $\hat{\alpha}$ is any number in (0, 1). Then there exists a positive constant C, depending on Ω , n and \hat{k} , such that

$$\inf_{D_{k'R}^+} \frac{u}{\delta} \le C \left(\inf_{D_{R/2}} \frac{u}{\delta} + C_2 R^{\hat{\alpha}} \right)$$
(2.19)

for all $R < \rho_0$, where ρ_0 is a positive constant depending only on Ω , n, $\hat{\alpha}$, A_0 and $\int_{\mathbb{R}^n} (|y|^2 \wedge 1) \hat{k}(y) dy$. *Proof.* When $C_2 = 0$, the proof follows from Step 1 of Lemma 2.3. So we let $C_2 > 0$. As before, we consider \tilde{u} to be the solution of

$$\begin{split} L\tilde{u} &= 0 \quad \text{in } D_R, \\ \tilde{u} &= u \quad \text{in } \mathbb{R}^n \setminus D_R. \end{split}$$

Then

$$\inf_{D_{k'R}^+} \frac{\tilde{u}}{\delta} \le C \inf_{D_{R/2}} \frac{\tilde{u}}{\delta}$$

holds, by step 1 of Lemma 2.3. Defining $w = \tilde{u} - u$, we get $|Lw| \leq C_2 g$ in D_R by using [10, Theorem 5.1] and w = 0 in D_R^c . As before, we would consider an appropriate supersolution and then apply comparison principle to establish (2.19).

For this construction of supersolution we take inspiration from [32, Lemma 5.8]. We set

$$\tilde{\psi}(s) = \int_0^s 2\mathrm{e}^{-ql-q\int_0^l \Theta(\tau)\mathrm{d}\tau}\mathrm{d}l - s,$$

where q > 0 is to be chosen later and Θ is given by

$$\Theta(s) = \int_{|z|>s} \min\{1, |z|\} \widehat{k}(z) \mathrm{d}z.$$

Since Θ is integrable in a neighbourhood of 0, there exists a sufficiently small constant s(q) > 0such that, for 0 < s < s(q), $\tilde{\psi}'(s) = 2e^{-qs-q\int_0^s \Theta(\tau)d\tau} - 1 \ge \frac{1}{2}$. Set $\sigma_1 = \min\{\frac{s(q)}{8}, 1, \gamma\}$. For any $r \in (0, 1)$, we define

$$\psi_{r,z}(x) = \begin{cases} \tilde{\psi}\left(\frac{d_{B_r(z)}(x)}{r}\right) & \text{if } d_{B_r(z)}(x) < r\sigma_1, \\ \tilde{\psi}(\sigma_1) & \text{if } d_{B_r(z)}(x) \ge r\sigma_1, \end{cases}$$

$$(2.20)$$

where $d_{B_r(z)}(x) = \text{dist}(x, B_r(z))$. Let $\eta = \frac{\sigma_1}{8}, 0 < r \le \rho$ and $B_{\eta r}(x_0) \cap \Omega = D_{\eta r}$. We define

$$\Phi_r(x) = \begin{cases} \tilde{\psi}\left(\frac{\delta(x)}{r}\right), & \text{if } \delta(x) < r\sigma_1, \\ \tilde{\psi}(\sigma_1) & \text{if } \delta(x) \ge r\sigma_1. \end{cases}$$

For $x \in D_{\eta r}$ then we have $x^* \in \partial \Omega$ such that $\delta(x) = |x - x^*|$. Let $z_x^* = z$ be a point in Ω^c such that $B_r(z)$ touches $\partial \Omega$ at x^* . From Remark 2.2 we have that

$$D_{\eta r} \subset B_{(1+\sigma_1)r}(z) \setminus B_r(z)$$

Since $\tilde{\psi}'' < 0$ and $|D\delta(x)| \ge \kappa > 0$ for $\delta(x) \in (0, \rho_1)$, ρ_1 sufficiently small, it follows that

$$\Delta \Phi_r(x) \le \frac{C}{r} + \tilde{\psi}''(\frac{\delta(x)}{r})\frac{\kappa^2}{r^2}.$$
(2.21)

Consider $\psi_{r,z}$ from (2.20) and notice that $\psi_{r,z}(x) = \Phi_r(x)$ and $\delta(x+y) \leq d_{B_r(z)}(x+y)$ for all $y \in \mathbb{R}^n$. Hence

$$\psi_{r,z}(x+y) + \psi_{r,z}(x-y) - 2\psi_{r,z}(x) \ge \Phi_r(x+y) + \Phi_r(x-y) - 2\Phi_r(x).$$

This readily gives (see [32, Lemma 5.8])

$$I\Phi_r(x) \le I\psi_{r,z}(x) \le \frac{C}{r} \left(1 + \Theta\left(\frac{d_{B_r(z)}(x)}{r}\right)\right) = \frac{C}{r} \left(1 + \Theta\left(\frac{\delta(x)}{r}\right)\right), \tag{2.22}$$

using the fact $\delta(x) = |x - x^*| = d_{B_r(z)}(x)$. Combining (2.21) and (2.22) we have

$$L\Phi_r \le \frac{CA_0}{r} \left(1 + \Theta\left(\frac{\delta(x)}{r}\right) \right) - \frac{2\kappa^2 q}{r^2} \left(1 + \Theta\left(\frac{\delta(x)}{r}\right) \right)$$

for all $x \in D_{\eta r}$. Now choose $q = \frac{1}{2\kappa^2}(CA_0 + 1)$ in the expression of ψ we obtain

$$L\Phi_r \le -\frac{1}{r^2} \left(1 + \Theta\left(\frac{\delta(x)}{r}\right) \right) \le -\frac{1}{r^2} \Theta\left(\frac{\delta(x)}{r}\right), \tag{2.23}$$

for all $x \in D_{\eta r}$.

Next we estimate the function Θ in $D_{\eta r}$. For $\xi \in (0, 1]$, we see that

$$\Theta(\xi) = \int_{|z|>\xi} \min\{1, |z|\} \widehat{k}(z) dz = \Lambda \int_{\xi < |z| \le 1} \frac{|z|}{|z|^{n+\alpha}} dz + \int_{|z|\ge 1} J(z) dz$$
$$= \Lambda \omega_n \int_{\xi}^1 \frac{r^n}{r^{n+\alpha}} dr + \kappa_1$$
$$= \begin{cases} \frac{\omega_n \Lambda}{\alpha - 1} \Big[\xi^{1-\alpha} - 1 \Big] + \kappa_1 & \text{for } \alpha \in (1, 2), \\ \omega_n \Lambda(-\log \xi) + \kappa_1 & \text{for } \alpha = 1, \end{cases}$$
(2.24)

for some positive constant κ_1 . Here ω_n denotes the surface area of the unit sphere in \mathbb{R}^n . Since $\frac{\delta(x)}{r} < \frac{1}{2}$ in $D_{\eta r}$, we get from the above estimate that

$$\Theta\left(\frac{\delta(x)}{r}\right) = \frac{\omega_n \Lambda}{\alpha - 1} \left[\left(\frac{\delta(x)}{r}\right)^{1 - \alpha} - 1 \right] + \kappa_1 \ge \Lambda \omega_n \left(\frac{\delta(x)}{r}\right)^{1 - \alpha} \left(\frac{2^{\alpha - 1} - 1}{2^{\alpha - 1}(\alpha - 1)}\right) \ge \kappa_2 \left[\frac{\delta(x)}{r}\right]^{1 - \alpha}.$$

for $\alpha \in (1,2)$, where the constant κ_2 is independent of α . Again, for $\alpha = 1$, we have

$$\Theta\left(\frac{\delta(x)}{r}\right) = \Lambda \omega_n \log(\frac{r}{\delta(x)}) + \kappa_1 \tag{2.25}$$

for $x \in D_{\eta r}$. We claim that for any $0 < \zeta < 1$, there exists a $r_{\theta} < 1$ such that for all $r < r_{\theta}$

$$\log(rz) \ge r^{\zeta} \log(z) \tag{2.26}$$

for all $z \ge \frac{1}{\theta r}$, where $0 < \theta < 1$ is a fixed positive constant. To prove the claim, we let

$$h(z) = \frac{\log(rz)}{\log(z)}$$

By our choice of parameters z, r, θ , we have h(z) > 0. Since $\log z \ge \log(rz)$, we have

$$h'(z) = \frac{(\log z - \log(rz))}{z(\log z)^2} > 0$$

for $z > \frac{1}{\theta r}$. Thus *h* is strictly increasing in $[(\theta r)^{-1}, \infty)$, and therefore,

$$h(z) \ge h((\theta r)^{-1}) = \frac{\log\left(\frac{1}{\theta}\right)}{\log\left(\frac{1}{\theta r}\right)} = \frac{\log\theta}{\log(r\theta)} \ge r^{\zeta},$$

for all $r \in (0, r_{\theta})$, where r_{θ} depends only on θ and ζ . The gives us (2.26). Putting (2.26) in (2.25) we have

$$\Theta\left(\frac{\delta(x)}{r}\right) \ge \Lambda \omega_n r^{\zeta} \log(\frac{1}{\delta(x)}) + \kappa_1 \ge C r^{\zeta} \left(\log\left(\frac{1}{\delta(x)}\right) + \kappa_1\right),$$

in $D_{\eta r}$, for all $r \leq r_{\theta}$. Using the above estimate and (2.23), we define the supersolutions as

$$v(x) = \mu r^{1+\hat{\alpha}} \Phi_r(x),$$

where the constant μ is chosen suitably so that $Lv \leq -g$ in $D_{\eta r}$, for all $r \leq r_{\theta}$. Thus, in $x \in D_{\eta r}$, we have $L(C_2v)(x) \leq -C_2g(x)$ and $C_2v(x) \geq 0$ in \mathbb{R}^n . Using comparison principle [10, Theorem 5.2] we then obtain $C_2v(x) \geq w(x)$ in \mathbb{R}^n . Repeating the same argument with -w, we get $|w(x)| \leq C_2v(x)$ in $D_{\eta r}$. Now we can complete the proof repeating the same argument as in Lemma 2.3.

Lemma 2.5. Let Ω be a bounded C^2 domain, and u be a bounded continuous function such that $u \geq 0$ in all of \mathbb{R}^n , and $|Lu| \leq C_2(1 + \mathbb{1}_{[1,2)}(\alpha)g)$ in D_R , for some constant C_2 . Let $\hat{\alpha} = 1 \wedge (2 - \alpha)$ for $\alpha \neq 1$, and for $\alpha = 1$, $\hat{\alpha}$ be any number in (0,1). Then, there exists a positive constant C, depending only on n, Ω, A_0 and \hat{k} , such that

$$\sup_{D^+_{\kappa'R}} \left(\frac{u}{\delta}\right) \le C \left(\inf_{D^+_{\kappa'R}} \frac{u}{\delta} + C_2 R^{\hat{\alpha}}\right)$$
(2.27)

for all $R \leq \rho_0$, where constant ρ_0 depends only on $\Omega, n, A_0, \hat{\alpha}$ and $\int_{\mathbb{R}^n} (|y|^2 \wedge 1) \hat{k}(y) dy$.

Proof. Recall from Remark 2.1 that we may take $\beta = \infty$ in Assumption 1.1(b). This property will be useful to apply the Harnack inequality from [24]. We split the proof in two steps.

Step 1. Let $C_2 = 0$. In this case (2.27) follows from the Harnack inequality for L. Let $R \leq \rho$. Then for each $y \in D^+_{\kappa'R}$ we have $B_{\kappa R}(y) \subset D_R$. Hence we have Lu = 0 in $B_{\kappa R}(y)$. Without loss of generality, we may assume y = 0. Let $r = \kappa R$ and define v(x) = u(rx) for all $x \in \mathbb{R}^n$. Then, it can be easily seen that

$$r^{2}Lu(rx) = L_{r}v(x) := \Delta v(x) + r^{2}\frac{1}{2}\int_{\mathbb{R}^{n}} (v(x+y) + v(x-y) - 2v(x))k(ry)r^{n}dy \quad \text{for all } x \in B_{1}.$$

This gives $L_r v(x) = 0$ in B_1 and $v \ge 0$ in whole \mathbb{R}^n . From the stochastic representation of v [10, Theorem 1.1], it follows that v is also a harmonic function in the probabilistic sense as considered in [24]. Hence by the Harnack inequality [24, Theorem 2.4] we obtain

$$\sup_{B_{\frac{1}{2}}} v \le C \inf_{B_{\frac{1}{2}}} v$$

where constant C does not depend on r. This of course, implies

$$\sup_{B_{\frac{\kappa R}{2}}} u \le C \inf_{B_{\frac{\kappa R}{2}}} u$$

Now cover $D_{\kappa'R}^+$ by a finite number of balls $B_{\kappa R/2}(y_i)$, independent of R, to obtain

$$\sup_{D^+_{\kappa'R}} u \le C \inf_{D^+_{\kappa'R}} u.$$

(2.27) follows since $\kappa R/2 \leq \delta \leq 3\kappa R/2$ in $D^+_{\kappa'R}$.

Step 2. Let $C_2 > 0$. The proof follows by combining Step 1 above and Step 2 of Lemmas 2.3 and 2.4.

Next we compute $L\delta$ in Ω .

Lemma 2.6. We have $|L\delta(x)| \leq Cg(x)$, where g is given by

$$g(x) = \begin{cases} (\delta(x) \wedge 1)^{1-\alpha} & \text{for } \alpha > 1, \\ \log(\frac{1}{\delta(x) \wedge 1}) + 1 & \text{for } \alpha = 1, \\ 1 & \text{for } \alpha \in (0, 1). \end{cases}$$
(2.28)

Proof. Since $\delta \in C^{0,1}(\mathbb{R}^n) \cap C^2(\overline{\Omega})$ [16, Theorem 5.4.3], (2.28) easily follows for the case $\alpha \in (0,1)$. Let $\Omega_{\rho_0} = \Omega \cap \{\delta < \rho_0\}$ where $\rho_0 < 1$. It is enough to show that

$$|L\delta(x)| \le C\Theta(\delta(x)) \quad \text{for } x \in \Omega_{\rho_0}, \tag{2.29}$$

where Θ is defined as before

$$\Theta(\xi) = \int_{|z| > \xi} \min\{1, |z|\} \widehat{k}(z) \mathrm{d}z$$

First of all

$$|L\delta(x)| \le |\Delta\delta(x)| + A_0|I\delta(x)| \le \kappa + A_0|I\delta(x)|, \tag{2.30}$$

for some constant κ , depending on Ω . Again,

$$I\delta(x) = \int_{\mathbb{R}^n} \left(\delta(x+z) + \delta(x-z) - 2\delta(x) \right) k(z) dz$$
$$= \int_{|z| \le \delta(x)/2} + \int_{|z| > \delta(x)/2}.$$

Since $\delta(x+z) + \delta(x-z) - 2\delta(x) \le \kappa_2 |z|^2$ for $|z| \le \delta(x)/2$, we have

$$\int_{|z| \le \delta(x)/2} \left(\delta(x+z) + \delta(x-z) - 2\delta(x) \right) k(z) \mathrm{d}z \le \kappa_2 \int_{|z| \le \delta(x)/2} |z|^2 \widehat{k}(z) \mathrm{d}z \le \kappa_3,$$

for some constant κ_3 . Since δ is Lipschitz, it follows that

$$\delta(x+z) + \delta(x-z) - 2\delta(x) \le 2(\operatorname{diam}(\Omega) \lor 1) \min\{|z|, 1\}.$$

Thus

$$\int_{|z|>\delta(x)/2} \left(\delta(x+z) + \delta(x-z) - 2\delta(x)\right) k(z) \mathrm{d}z \le \kappa_4 \int_{|z|>\delta(x)/2} \min\{|z|,1\} \widehat{k}(z) \mathrm{d}z = \kappa_4 \Theta(\delta(x)/2),$$

for some constant κ_4 . Inserting these estimates in (2.30) we obtain

$$|L\delta(x)| \le \kappa_5(1 + \Theta(\delta(x)/2))$$
 for all $x \in \Omega_{\rho_0}$

for some constant κ_5 . Choosing ρ_0 sufficiently small, (2.29) follows from (2.24).

Now we are ready to prove our key estimate towards the regularity of u/δ .

Proposition 2.1. Let u be a bounded continuous function such that $|Lu| \leq K$ in Ω , for some constant K, and u = 0 in Ω^c . Given any $x_0 \in \partial\Omega$, let D_R be as in the Definition 2.1. Then for some $\tau \in (0, 1 \land (2 - \alpha))$ there exists C, dependent on Ω, n, A_0, α and \hat{k} but not on x_0 , such that

$$\sup_{D_R} \frac{u}{\delta} - \inf_{D_R} \frac{u}{\delta} \le CKR^{\tau}$$
(2.31)

for all $R \leq \rho_0$, where $\rho_0 > 0$ is a constant depending only on $\Omega, n, \hat{\alpha}, A_0$ and $\int_{\mathbb{R}^n} (|y|^2 \wedge 1) \widehat{k}(y) dy$.

BOUNDARY REGULARITY

Proof. For the proof we follow a standard method, similar to [34], with the help of Lemmas 2.4 to 2.6. Fix $x_0 \in \partial\Omega$ and consider $\rho_0 > 0$ to be chosen later. With no loss of generality, we assume $x_0 = 0$. In view of (2.2), we only consider the case K > 0. By considering u/K instead of u, we may assume that K = 1, that is, $|Lu| \leq 1$ in Ω . From Lemma 2.1 we note that $|u|_{C^{0,1}(\mathbb{R}^n)} \leq C_1$. For $\alpha \in (0, 1)$, we can calculate Iu classically and $|Iu| \leq \tilde{C}$ in Ω , we can combine the nonlocal term on the rhs and only deal with Δu . In this case the proof is simpler and can be done following the same method as for the local case (the proof below also works with minor modifications). Therefore, we only deal with $\alpha \in [1, 2)$.

We show that there exists G > 0, $\rho_1 \in (0, \rho_0)$ and $\tau \in (0, 1)$, dependent only on Ω, n, A_0 and k, and monotone sequences $\{M_k\}$ and $\{m_k\}$ such that, for all $k \ge 0$,

$$M_k - m_k = \frac{1}{4^{k\tau}}, \quad -1 \le m_k \le m_{k+1} < M_{k+1} \le M_k \le 1,$$
 (2.32)

and

$$m_k \le G^{-1} \frac{u}{\delta} \le M_k$$
 in $D_{R_k} = D_{R_k}(x_0)$, where $R_k = \frac{\rho_1}{4^k}$. (2.33)

Note that (2.33) is equivalent to the following

$$m_k \delta \le G^{-1} u \le M_k \delta$$
, in $B_{R_k} = B_{R_k}(x_0)$, where $R_k = \frac{\rho_1}{4^k}$. (2.34)

Next we construct monotone sequences $\{M_k\}$ and $\{m_k\}$ by induction.

The existence of M_0 and m_0 such that (2.32) and (2.34) hold for k = 0 is guaranteed by Lemma 2.1. Assume that we have the sequences up to M_k and m_k . We want to show the existence of M_{k+1} and m_{k+1} such that (2.32)-(2.34) hold. We set

$$u_k = \frac{1}{G}u - m_k\delta. \tag{2.35}$$

Note that to apply Lemma 2.5 we need u_k to be nonnegative in \mathbb{R}^n . Therefore we work with u_k^+ , the positive part of u_k . Let $u_k = u_k^+ - u_k^-$ and by the induction hypothesis,

$$u_k^+ = u_k \text{ and } u_k^- = 0 \text{ in } B_{R_k}.$$
 (2.36)

We need a lower bound on u_k . Since $u_k \ge 0$ in B_{R_k} , we get for $x \in B_{R_k}^c$ that

$$u_k(x) = u_k(R_k x_u) + u_k(x) - u_k(R_k x_u) \ge -C_L |x - R_k x_u|,$$
(2.37)

where $z_{\rm u} = \frac{1}{|z|} z$ for $z \neq 0$ and C_L denotes a Lipschitz constant of u_k which can be chosen independent of k. Using Lemma 2.1 we also have $|u_k| \leq G^{-1} + \operatorname{diam}(\Omega) = C_1$ for all $x \in \mathbb{R}^n$. Thus using (2.36) and (2.37) we calculate Lu_k^- in $D_{\frac{R_k}{2}}$. Let $x \in D_{R_k/2}(x_0)$. By (2.36), $\Delta u_k^-(x) = 0$. Denote by

$$\tilde{g}(r) = \begin{cases} |\log(r)| & \text{for } r > 0, \ \alpha = 1, \\ r^{1-\alpha} & \text{for } r > 0, \ \alpha \in (1,2). \end{cases}$$

Then

$$\begin{split} 0 &\leq Iu_{k}^{-}(x) = \int_{x+y \notin B_{R_{k}}} u_{k}^{-}(x+y)k(y)\mathrm{d}y \\ &\leq \int_{\left\{|y| \geq \frac{R_{k}}{2}, x+y \neq 0\right\}} u_{k}^{-}(x+y)k(y)\mathrm{d}y \\ &\leq C_{L} \int_{\left\{\frac{R_{k}}{2} \leq |y| \leq 1, \ x+y \neq 0\right\}} \left|(x+y) - R_{k}(x+y)_{u}\right| \hat{k}(y)\mathrm{d}y + C_{1} \int_{|y| \geq 1} J(y)\mathrm{d}y \\ &\leq C_{L} \int_{\frac{R_{k}}{2} \leq |y| \leq 1} (|x| + R_{k}) \frac{1}{|y|^{n+\alpha}} \,\mathrm{d}y + C_{L} \int_{\frac{R_{k}}{2} \leq |y| \leq 1} \frac{1}{|y|^{n+\alpha-1}} \,\mathrm{d}y + \kappa_{1}C_{1} \end{split}$$

$$\leq \kappa_3((R_k)^{1-\alpha} + \tilde{g}(R_k/2) + 1)$$

$$\leq \kappa_4 \tilde{g}(R_k)$$

for some constants $\kappa_1, \kappa_3, \kappa_4$, independent of k. Now we write $u_k^+ = G^{-1}u - m_k\delta + u_k^-$ and applying the operator L, we get

$$|Lu_{k}^{+}| \leq G^{-1}|Lu| + m_{k}|L\delta| + |Lu_{k}^{-}| \leq G^{-1} + m_{k}Cg(x) + \kappa_{4}\tilde{g}(R_{k}), \qquad (2.38)$$

using Lemma 2.6. Since $\rho_1 \ge R_k \ge \delta$ in D_{R_k} , for $\alpha \ge 1$, we have $R_k^{1-\alpha} \le \delta^{1-\alpha}$, and hence, from (2.38), we have

$$|Lu_k^+| \le \left[G^{-1}[\tilde{g}(\rho_1)]^{-1} + C + \kappa_4\right]g(x) := \kappa_5 g(x) \quad \text{in} \quad D_{R_k/2}$$

Now we are ready to apply Lemma 2.5. Recalling that

$$u_k^+ = u_k = G^{-1}u - m_k\delta \quad \text{in} \quad D_{R_k}$$

we get from Lemmas 2.4 and 2.5 that

$$\sup_{\substack{D_{\kappa'R_k/2}^+}} \left(G^{-1} \frac{u}{\delta} - m_k \right) \le C \left(\inf_{\substack{D_{\kappa'R_k/2}^+}} \left(G^{-1} \frac{u}{\delta} - m_k \right) + \kappa_5 R_k^{\hat{\alpha}} \right)$$

$$\le C \left(\inf_{\substack{D_{R_k/4}^+}} \left(G^{-1} \frac{u}{\delta} - m_k \right) + \kappa_5 R_k^{\hat{\alpha}} \right).$$
(2.39)

Repeating a similar argument for the function $\tilde{u}_k = M_k \delta - G^{-1} u$, we obtain

$$\sup_{\substack{D_{\kappa'R_k/2}^+}} \left(M_k - G^{-1} \frac{u}{\delta} \right) \le C \left(\inf_{\substack{D_{R_k/4}}} \left(M_k - G^{-1} \frac{u}{\delta} \right) + \kappa_5 R_k^{\hat{\alpha}} \right)$$
(2.40)

Combining (2.39) and (2.40) we obtain

$$M_{k} - m_{k} \leq C \Big(\inf_{\substack{D_{R_{k}/4}^{+}}} \Big(M_{k} - G^{-1} \frac{u}{\delta} \Big) + \inf_{\substack{D_{R_{k}/4}^{+}}} \Big(G^{-1} \frac{u}{\delta} - m_{k} \Big) + \kappa_{5} R_{k}^{\hat{\alpha}} \Big)$$

= $C \Big(\inf_{\substack{D_{R_{k+1}}}} G^{-1} \frac{u}{\delta} - \sup_{\substack{D_{R_{k+1}}}} G^{-1} \frac{u}{\delta} + M_{k} - m_{k} + \kappa_{5} R_{k}^{\hat{\alpha}} \Big).$ (2.41)

Putting $M_k - m_k = \frac{1}{4^{\tau k}}$ in (2.41), we have

$$\sup_{D_{R_{k+1}}} G^{-1} \frac{u}{\delta} - \inf_{D_{R_{k+1}}} G^{-1} \frac{u}{\delta} \le \left(\frac{C-1}{C} \frac{1}{4^{\tau k}} + \kappa_5 R_k^{\hat{\alpha}}\right) \\ = \frac{1}{4^{\tau k}} \left(\frac{C-1}{C} + \kappa_5 R_k^{\hat{\alpha}} 4^{\tau k}\right).$$
(2.42)

Since $R_k = \frac{\rho_1}{4^k}$ for $\rho_1 \in (0, \rho_0)$, we can choose ρ_0 and τ small so that

$$\left(\frac{C-1}{C} + \kappa_5 R_k^{\hat{\alpha}} 4^{\tau k}\right) \le \frac{1}{4^{\tau}}.$$

Putting in (2.42) we obtain

$$\sup_{D_{R_{k+1}}} G^{-1} \frac{u}{\delta} - \inf_{D_{R_{k+1}}} G^{-1} \frac{u}{\delta} \le \frac{1}{4^{\tau(k+1)}}.$$

Thus we find m_{k+1} and M_{k+1} such that (2.32) and (2.33) hold. It is easy to prove (2.31) from (2.32)-(2.33).

Now we are ready to complete the proof of Theorem 1.2.

Proof of Theorem 1.2. As mentioned before, it is enough to consider (2.12). Replacing u by $\frac{u}{CK}$ we may assume that $|Lu| \leq 1$ in Ω . Let $v = u/\delta$. From Lemma 2.1 we then have

$$\|v\|_{L^{\infty}(\Omega)} \le C,\tag{2.43}$$

for some constant C. Also, from Theorem 1.1 we have

$$\|u\|_{C^{0,1}(\mathbb{R}^n)} \le C. \tag{2.44}$$

It is also easily seen that for any $x \in \Omega$ with $R = \delta(x)$ we have

z

$$\sup_{1,z_2 \in B_{R/2}(x)} \frac{|\delta^{-1}(z_1) - \delta^{-1}(z_2)|}{|z_1 - z_2|} \le CR^{-2}.$$

Combining it with (2.44) gives

$$\sup_{z_1, z_2 \in B_{R/2}(x)} \frac{|v(z_1) - v(z_2)|}{|z_1 - z_2|} \le C(1 + R^{-2}).$$
(2.45)

Again, by Proposition 2.1, for each $x_0 \in \partial \Omega$ and for all r > 0 we have

$$\sup_{D_r(x_0)} v - \inf_{D_r(x_0)} v \le C r^{\tau}.$$
(2.46)

where $D_r(x_0) = B_r(x_0) \cap \Omega$ as before. To complete the proof it is enough to show that

$$\sup_{x,y\in\Omega, x\neq y} \frac{|v(x) - v(y)|}{|x - y|^{\kappa}} \le C,$$
(2.47)

for some $\eta > 0$. Consider $x, y \in \Omega$ and let r = |x - y|. We also suppose that $\delta(x) \ge \delta(y)$. If $r \ge 1/2$, then

$$\frac{|v(x) - v(y)|}{|x - y|^{\kappa}} \le C2^{1+\kappa},$$

by (2.43). So we suppose |x-y| = r < 1/2. Let $R = \delta(x)$ and $x_0, y_0 \in \partial\Omega$ satisfying $\delta(x) = |x-x_0|$ and $\delta(y) = |y-y_0|$. Fix p > 2. Set $\kappa = [2 + \operatorname{diam} \Omega]^{-p}$. If $r \leq \kappa R^p$, then $r < \frac{1}{2}R$. In this case, it follows from (2.45) that

$$|v(x) - v(y)| \le C(1 + R^{-2})r \le C(r + \kappa^{2/p}r^{1-2/p}) \le C_1r^{1-2/p}.$$

Again, if $r \ge \kappa R^p$, we have $R \le [r/\kappa]^{\frac{1}{p}}$. Thus, $y \in B_{\kappa_1 r^{\frac{1}{p}}}(x_0)$ for $\kappa_1 = 1 + \kappa^{-1/p}$. From (2.46) we then have

$$|v(x) - v(y)| \le C_2 r^{r_F}$$
.
Thus (2.47) follows by fixing $\kappa = \min\{\frac{\tau}{n}, 1 - \frac{2}{n}\}$. This completes the proof.

Remark 2.3. The regularity of $\partial\Omega$ in Theorem 1.2 can be relaxed to $C^{1,1}$. In this case, δ will be a $C^{1,1}$ function. Therefore, $I\delta$ is defined classically and $L\delta$ can be interpreted in the viscosity sense (see [11]). The proof of Theorem 1.2 goes through due to the coupling result in [10, Lemma 5.2].

2.2. Boundary regularity of Du. Using Theorem 1.2 we show that $Du \in C^{1,\gamma}(\Omega)$ for some $\gamma > 0$. Recall (1.1)

$$Lu + C_0 |Du| \ge -K \quad \text{in } \Omega,$$

$$Lu - C_0 |Du| \le K \quad \text{in } \Omega,$$

$$u = 0 \quad \text{in } \Omega^c.$$
(2.48)

Let $v = u/\delta$. From Theorem 1.2 we know that $v \in C^{\kappa}(\Omega)$. We extend v in all of \mathbb{R}^n as a C^{κ} function without altering its C^{κ} norm (cf. [34, Lemma 3.8]). Below we find the equations satisfied by v.

Lemma 2.7. If $|Lu| \leq C$ in Ω and u = 0 in Ω^c , then we have

$$\frac{1}{\delta}[-C - vL\delta - Z[v,\delta]] \le Lv + 2\frac{D\delta}{\delta} \cdot Dv \le \frac{1}{\delta}[C - vL\delta - Z[v,\delta]],$$
(2.49)

in Ω , where

$$Z[v,\delta](x) = \int_{\mathbb{R}^n} (v(y) - v(x))(\delta(y) - \delta(x))k(y - x)dy$$

Proof. First of all, since $u \in C^1(\Omega)$ [33, Theorem 4.1], we have $v \in C^1(\Omega)$. Therefore, $Z[v, \delta]$ is continuous in Ω . Consider a test function $\psi \in C^2(\Omega)$ that touches v from above at $x \in \Omega$. Define

$$\psi_r(z) = \begin{cases} \psi(z) & \text{in } B_r(x), \\ v(z) & \text{in } B_r^c(x). \end{cases}$$

By our assertion, we must have $\psi_r \geq v$ for all r small. To verify (2.49) we must show that

$$L\psi_r(x) + 2\frac{D\delta}{\delta} \cdot D\psi_r(x) \ge \frac{1}{\delta(x)} [-C + v(x)L\delta(x) - Z[v,\delta](x)], \qquad (2.50)$$

for some r small. We define

$$\tilde{\psi}_r(z) = \begin{cases} \delta(z)\psi(z) & \text{ in } B_r(x), \\ u(z) & \text{ in } B_r^c(x). \end{cases}$$

Then, $\tilde{\psi}_r \geq u$ for all r small. Since $|Lu| \leq C$ and $\delta \psi_r = \tilde{\psi}_r$, we obtain

$$-C \le L\tilde{\psi}_r(x) = \delta(x)L\psi_r(x) + v(x)L\delta(x) + 2D\delta(x) \cdot D\psi_r(x) + Z[\psi_r,\delta](x)$$

for all r small. Rearranging the terms we have

$$-C - v(x)L\delta(x) - Z[\psi_r, \delta](x) \le \delta(x)L\psi_r(x) + 2D\delta(x) \cdot D\psi_r(x).$$
(2.51)

Let $r_1 \leq r$. Since ψ_r is decreasing with r, we get from (2.51) that

$$\delta(x)L\psi_r(x) + 2D\delta(x) \cdot D\psi_r(x) \ge \delta(x)L\psi_{r_1}(x) + 2D\delta(x) \cdot D\psi_r(x)$$
$$\ge \lim_{r_1 \to 0} [-C - v(x)L\delta(x) - Z[\psi_{r_1}, \delta](x)$$
$$= [-C - v(x)L\delta(x) - Z[v, \delta](x)],$$

by dominated convergence theorem. This gives (2.50). Similarly we can verify the other side of (2.49).

In order to prove Theorem 1.3, we also need the following estimate on v. Define $\Omega_{\sigma} = \{x \in \Omega : \text{dist}(x, \Omega^c) \geq \sigma\}$. Then we have

Lemma 2.8. For some constant C it holds that

$$\|Dv\|_{L^{\infty}(\Omega_{\sigma})} \le CK\sigma^{\kappa-1} \quad for \ all \ \sigma \in (0,1).$$

$$(2.52)$$

Furthermore, there exists $\eta \in (0,1)$ such that for any $x \in \Omega_{\sigma}$ and $0 < |x-y| \le \sigma/8$ we have

$$\frac{|Dv(y) - Dv(x)|}{|x - y|^{\eta}} \le CK\sigma^{\kappa - 1 - \eta}$$

for all $\sigma \in (0,1)$.

Proof. As earlier, we suppose K > 0. Diving u by K in (2.48) we may assume K = 1. Using Theorem 1.1 we can write $|Lu| \leq C_1$ in Ω , for some constant C_1 . By Lemma 2.7 we then have

$$\frac{1}{\delta} [-C_1 - vL\delta - Z[v,\delta]] \le Lv + 2\frac{D\delta}{\delta} \cdot Dv \le \frac{1}{\delta} [C_1 - vL\delta - Z[v,\delta]],$$
(2.53)

in Ω . Fix $x_0 \in \Omega_{\sigma}$ and define

$$w(x) = v(x) - v(x_0).$$

From (2.53) we then obtain

$$-\frac{1}{\delta}C_1 - \ell \le Lw + 2\frac{D\delta}{\delta} \cdot Dw \le \frac{1}{\delta}C_1 - \ell, \qquad (2.54)$$

in Ω , where

$$\ell(x) = \frac{1}{\delta(x)} [w(x)L\delta(x) + Z[v,\delta](x) + v(x_0)L\delta(x)].$$

Set $r = \frac{\sigma}{2}$. We claim that

$$\|\ell\|_{L^{\infty}(B_r(x_0)} \le \kappa_1 \sigma^{\kappa-2}, \quad \text{for all } \sigma \in (0,1),$$

$$(2.55)$$

for some constant κ_1 . Let us denote by

$$\xi_1 = \frac{wL\delta}{\delta}, \quad \xi_2 = \frac{1}{\delta}Z[v,\delta] \quad \text{and} \quad \xi_3 = \frac{v(x_0)}{\delta}L\delta$$

Recall that $\kappa \in (0, (2 - \alpha) \land 1)$ from Theorem 1.2. Since

$$\|\Delta\delta\|_{L^{\infty}(\Omega)} < \infty \quad \text{and} \quad \|I\delta\|_{L^{\infty}(\Omega_{\sigma})} \lesssim \begin{cases} (\delta(x) \wedge 1)^{1-\alpha} & \text{for} \quad \alpha > 1, \\ \log(\frac{1}{\delta(x) \wedge 1}) + 1 & \text{for} \quad \alpha = 1, \\ 1 & \text{for} \quad \alpha \in (0, 1), \end{cases}$$

(cf. Lemma 2.6), and

$$\|v\|_{L^{\infty}}(\mathbb{R}^n) < \infty, \quad \|w\|_{L^{\infty}(B_r(x_0))} \lesssim r^{\kappa},$$

it follows that

$$\|\xi_3\|_{L^{\infty}(B_r(x_0))} \lesssim \begin{cases} \sigma^{-(\alpha \vee 1)} & \text{for } \alpha \neq 1, \\ \sigma^{-1}|\log \sigma| & \text{for } \alpha = 1, \end{cases} \lesssim \sigma^{-2+\kappa},$$

and

$$\|\xi_1\|_{L^{\infty}(B_r(x_0))} \lesssim \begin{cases} \sigma^{\kappa-1+1-(1\vee\alpha)} & \text{for } \alpha \neq 1, \\ \sigma^{\kappa-1}|\log\sigma| & \text{for } \alpha = 1, \end{cases} \lesssim \sigma^{-2+\kappa}.$$

So we are left to compute the bound for ξ_2 . Let $x \in B_r(x_0)$. Denote by $\hat{r} = \delta(x)/4$. Note that

$$\delta(x) \ge \delta(x_0) - |x - x_0| \ge 2r - r = r \Rightarrow \hat{r} \ge r/4.$$

Thus, since $u \in C^1(\Omega)$ by [33, Theorem 4.1] (as mentioned before, the proof of [33] works for inequations),

$$|Dv| \le |\frac{Du}{\delta}| + |\frac{uD\delta}{\delta^2}| \lesssim [\delta(x)]^{-1} \quad \text{in } B_{\hat{r}}(x),$$

using (2.1). Since δ is Lipscitz and bounded in \mathbb{R}^n , we obtain

$$\begin{split} |Z[v,\delta](x)| &\leq \Lambda \int_{y \in B_{\hat{r}}(x)} \frac{|\delta(x) - \delta(y)| |v(x) - v(y)|}{|x - y|^{n + \alpha}} \mathrm{d}y + \int_{y \in B_{\hat{r}}^c(x)} |\delta(x) - \delta(y)| |v(x) - v(y)| \hat{k}(y - x) \mathrm{d}y \\ &\lesssim [\delta(x)]^{-1} \int_{y \in B_{\hat{r}}(x)} |x - y|^{2 - n - \alpha} \mathrm{d}y + \int_{y \in B_1(x) \setminus B_{\hat{r}}(x)} \frac{(\delta(x) - \delta(y))(v(x) - v(y))}{|x - y|^{n + \alpha}} \mathrm{d}y \\ &+ \int_{|y| > 1} |\delta(x) - \delta(y + x)| |v(x) - v(y + x)| J(y) \mathrm{d}y \\ &\lesssim [\delta(x)]^{1 - \alpha} + \int_{y \in B_1(x) \setminus B_{\hat{r}}^c(x)} \frac{|\delta(x) - \delta(y)| |v(x) - v(y)|}{|x - y|^{n + \alpha}} \mathrm{d}y + \kappa_2, \end{split}$$

for some constant κ_2 . The second integration on the right hand side can be computed as follows: for $\alpha \leq 1$ we write

$$\int_{y\in B_1(x)\setminus B_{\hat{r}}^c(x)} \frac{|\delta(x)-\delta(y)||v(x)-v(y)|}{|x-y|^{n+\alpha}} \mathrm{d}y \lesssim \int_{y\in B_1(x)\setminus B_{\hat{r}}^c(x)} |x-y|^{-n-\alpha+1+\kappa} \mathrm{d}y \lesssim (1-\hat{r}^{1-\alpha+\kappa}) \lesssim \sigma^{-1+\kappa},$$

whereas for $\alpha \in (1,2)$ we can compute it as

$$\int_{y\in B_1(x)\setminus B_{\hat{r}}^c(x)} \frac{|\delta(x)-\delta(y)||v(x)-v(y)|}{|x-y|^{n+\alpha}} \mathrm{d}y \lesssim \int_{y\in B_1(x)\setminus B_{\hat{r}}^c(x)} |x-y|^{-n-\alpha+1} \lesssim \hat{r}^{-\alpha+1} \lesssim \sigma^{-1+\kappa}.$$

Combining the above estimates we obtain

$$\|\xi_2\|_{L^\infty B_r(x_0)} \lesssim \sigma^{-2+\kappa}$$

Thus we have established the claim (2.55).

Let us now define $\zeta(z) = w(\frac{r}{2}z + x_0)$. Letting $b(z) = 2\frac{D\delta(\frac{r}{2}z + x_0)}{\delta(\frac{r}{2}z + x_0)}$ and $r_1 = \frac{r}{2}$ it follows from (2.54) that

$$r_1^2 \left(-\frac{C}{\delta} - \ell \right) (r_1 z + x_0) \le \Delta \zeta + r_1^{2-\alpha} I_{r_1} \zeta + r_1 b(z) \cdot D\zeta \le r_1^2 \left(\frac{C}{\delta} - \ell \right) (r_1 z + x_0)$$
(2.56)

in $B_2(0)$, where

$$I_{r_1}f(x) = r_1^{\alpha} \frac{1}{2} \int_{\mathbb{R}^n} (f(x+y) + f(x-y) - f(x))k(r_1y)r_1^n \mathrm{d}y.$$

Consider a cut-off function φ satisfying $\varphi = 1$ in $B_{3/2}(0)$ and $\varphi = 0$ in $B_2^c(0)$. Defining $\tilde{\zeta} = \zeta \varphi$ we get from (2.56) that

$$|\Delta \tilde{\zeta}(z) + r_1^{2-\alpha} I_{r_1} \tilde{\zeta}(z) + r_1 b(z) \cdot D\tilde{\zeta}| \le r_1^2 (\frac{C}{\delta} + |\ell|) (r_1 z + x_0) + r_1^{2-\alpha} |I_{r_1}((\varphi - 1)\zeta)|$$

in $B_1(0)$. Since

 $||r_1b||_{L^{\infty}(B_1(0))} \le \kappa_3 \text{ for all } \rho \in (0,1),$

applying [33, Theorem 4.1] (this result works for inequality) we obtain, for some $\eta \in (0, 1)$,

$$\|D\zeta\|_{C^{\eta}(B_{1/2}(0))} \le \kappa_6 \left(\|\tilde{\zeta}\|_{L^{\infty}(\mathbb{R}^n)} + Cr_1 + r_1^2 \|\ell(r_1 \cdot + x_0)\|_{L^{\infty}(B_1)} + r_1^{2-\alpha} \|I_{r_1}((\varphi - 1)\zeta)\|_{L^{\infty}(B_1)} \right),$$
(2.57)

for some constant κ_6 independent of $\rho \in (0, 1)$. Since v is in $C^{\kappa}(\mathbb{R}^n)$, it follows that

$$\|\tilde{\zeta}\|_{L^{\infty}(\mathbb{R}^n)} = \|\tilde{\zeta}\|_{L^{\infty}(B_2)} \le \|\zeta\|_{L^{\infty}(B_2)} \lesssim r^{\kappa}.$$

Also, by (2.55),

$$r_1^2 \|\ell(r_1 \cdot + x_0)\|_{L^\infty(B_1)} \lesssim \sigma^{\kappa}.$$

Note that, for $z \in B_1(0)$,

$$|I_{r_1}(\varphi - 1)\zeta| \leq \int_{|y| \geq 1/2} |(\varphi(x+y) - 1)\zeta(x+y)|\widehat{k}(y)dy$$
$$\leq 2||v||_{L^{\infty}(\mathbb{R}^n)} \int_{|y| \geq 1/2} \widehat{k}(y)dy$$
$$\leq \kappa_3$$

for some constant κ_3 . Putting these estimates in (2.57) and calculating the gradient at z = 0 we obtain

$$|Dv(x_0)| \le \kappa_4 \sigma^{-1+\kappa_4}$$

for all $\sigma \in (0, 1)$. This proves the first part.

For the second part, compute the Hölder ratio with $D\zeta(0) - D\zeta(z)$ where $z = \frac{2}{r}(y - x_0)$ for $|x_0 - y| \le \sigma/8$. This completes the proof.

Now we can establish the Hölder regularity of the gradient up to the boundary.

Proof of Theorem 1.3. Since $u = v\delta$ it follows that

$$Du = vD\delta + \delta Dv.$$

Since $\delta \in C^2(\bar{\Omega})$, it follows from Theorem 1.2 that $vD\delta \in C^{\kappa}(\bar{\Omega})$. Thus, we only need to concentrate on $\vartheta = \delta D v$. Consider η from Lemma 2.8 and with no loss of generality, we may fix $\eta \in (0, \kappa)$. For $|x - y| \ge \frac{1}{8}(\delta(x) \lor \delta(y))$ it follows from (2.52) that

$$\frac{|\vartheta(x) - \vartheta(y)|}{|x - y|^{\eta}} \le C(\delta^{\kappa}(x) + \delta^{\kappa}(y))(\delta(x) \vee \delta(y))^{-\eta} \le 2C$$

So consider the case $|x - y| < \frac{1}{8}(\delta(x) \lor \delta(y))$. Without loss of generality, we may assume that $|x - y| < \frac{1}{8}\delta(x)$. Then

$$\frac{9}{8}\delta(x) \ge |x-y| + \delta(x) \ge \delta(y) \ge \delta(x) - |x-y| \ge \frac{7}{8}\delta(x).$$

By Lemma 2.8, it follows

$$\frac{\vartheta(x) - \vartheta(y)|}{|x - y|^{\eta}} \le |Dv(x)| \frac{|\delta(x) - \delta(y)|}{|x - y|^{\eta}} + \delta(y) \frac{|Dv(x) - Dv(y)|}{|x - y|^{\eta}}$$
$$\lesssim \delta(x)^{-1+\kappa} (\delta(x))^{1-\eta} + \delta(y) [\delta(x)]^{\kappa-1-\eta}$$
$$\le C.$$

This completes the proof by setting $\gamma = \eta$.

3. Overdetermined problems

In this section we solve an overdetermined problem. Let $H : \mathbb{R} \to \mathbb{R}$ be a given locally Lipschitz function. We also assume that $a \in (0, A_0)$. Our main result of this section is the following.

Theorem 3.1. Let $\Omega \subset \mathbb{R}^n$, $n \geq 2$, be an open bounded set with C^2 boundary. Suppose that Assumption 1.1 holds, k = k(|y|) and $k : (0, \infty) \to (0, \infty)$ is strictly decreasing. Let $f : \mathbb{R} \to \mathbb{R}$ be locally Lipschitz and u be a viscosity solution to

$$Lu + H(|Du|) = f(u) \quad in \Omega,$$

$$u = 0 \quad in \Omega^{c}, \quad u > 0 \quad in \Omega,$$

$$\frac{\partial u}{\partial n} = c \quad on \partial\Omega,$$
(3.1)

for some fixed c > 0, where n is the unit inward normal on $\partial \Omega$. Then Ω must be a ball. Furthermore, u is radially symmetric and strictly decreasing in the radial direction.

Proof of Theorem 3.1 follows from the boundary estimates in Theorem 1.2 combined with the approach of [8,19]. Also, note that we have taken k to be positive valued. This is just for convenience and the proofs below can be easily modified to include kernel k that is non-increasing but strictly decreasing in a neighbourhood of 0, provided we assume Ω to be connected. We provide a sketch for the proof of Theorem 3.1 and the finer details can be found in [8,19]. From Theorems 1.1 and 1.3 we see that $u \in C^{0,1}(\mathbb{R}^n) \cap C^{1,\gamma}(\overline{\Omega})$, and therefore, $u \in C^{2,\eta}(\Omega)$ by [33]. Therefore, we can assume that u is a classical solution to (3.1).

Given a unit vector e, let us define the half space

$$\mathcal{H} = \mathcal{H}_{\lambda, e} = \{ x \in \mathbb{R}^n : x \cdot e > \lambda \},\$$

and let $\bar{x} = \Re_{\lambda,e}(x) = x - 2(x \cdot e)e + 2\lambda e$ be the reflection of x along $\partial \mathcal{H} = \{x \cdot e = \lambda\}$. We say $v : \mathbb{R}^n \to \mathbb{R}$ is anti-symmetric if $v(x) = -v(\bar{x})$ for all $x \in \mathbb{R}^n$. Now let $D \subset \mathcal{H}$ be a bounded open

set and u be a bounded anti-symmetric solution to

$$\begin{aligned} Lu - \beta |Du| &\leq g \quad \text{in } D\\ u &\geq 0 \quad \text{in } \mathcal{H} \setminus D, \end{aligned}$$

where $\beta > 0$ is a fixed constant. Let

$$v = \begin{cases} -u & \text{in } \{u < 0\} \cap D\\ 0 & \text{otherwise.} \end{cases}$$

Then it can be easily seen that v solves

$$Lv - \beta |Dv| \ge -g \quad \text{in} \quad \Sigma := \{u < 0\} \cap D, \tag{3.2}$$

in the viscosity sense. To check (3.2), consider $x \in \Sigma$ and test function ϕ such that $\phi(x) = v(x)$ and $\phi(y) > v(y)$ for $y \in \mathbb{R}^n \setminus \{x\}$. Define $\psi := \phi + (-u - v)$. Then $\psi(x) = -u(x)$ and $\psi(y) > -u(y)$ for $y \in \mathbb{R}^d \setminus \{x\}$. Furthermore, $\psi = \phi$ in Σ . Thus, we get $L\psi(x) + \beta |D\phi(x)| \ge -g(x)$. This implies $L\phi + \beta |D\phi| + L(-u - v)(x) + |D\phi(x)| \ge -g(x)$. Since k is radially decreasing and u is anti-symmetric, it follows that $L(-u - v)(x) \le 0$ (cf. [8, p. 11]). This gives us (3.2).

The following narrow domain maximum principle is a consequence of the ABP estimate in [32, Theorem A.4].

Lemma 3.1. Let \mathcal{H} be the half space and $D \subset \mathcal{H}$ be open and bounded. Also, assume c to be bounded. Then there exists a positive constant C, depending on diam D, n, k, such that if $u \in C_b(\mathbb{R}^n)$ is an anti-symmetric supersolution of

$$Lu - \beta |Du| - c(x)u = 0 \quad in D,$$
$$u \ge 0 \quad in \mathcal{H} \setminus D,$$

then we have

$$\sup_{\Omega} u^{-} \le C \|c^{+}\|_{L^{\infty}(D)} \|u^{-}\|_{L^{n}(D)}.$$

In particular, given $\kappa_{\infty} > 0$ and $c^+ \leq \kappa_{\infty}$ on D, there is a $\delta > 0$ such that if $|D| < \delta$, we must have $u \geq 0$ in \mathcal{H} .

Proof. Set $\Sigma = \{u < 0\} \cap D$ and define v as above. From (3.2) we see that

 $Lv + \beta |Dv| + c(x)v \ge 0$ in Σ .

Since $v \ge 0$ in Σ and $c \le c^+$, we get

$$Lv + \beta |Dv| + c^+ v \ge 0$$
 in Σ

Taking $f = -c^+ v$ and using [32, Theorem A.4] we obtain, for some constant C_1 , that

$$\sup_{\Omega} u^{-} = \sup_{\Sigma} v \leq \sup_{\Sigma^{c}} |v| + C_{1} ||f||_{L^{n}(D)} \leq C_{1} ||c^{+}||_{L^{\infty}(D)} ||v||_{L^{n}(\Sigma)} = C_{1} ||c^{+}||_{L^{\infty}(D)} ||u^{-}||_{L^{n}(D)}.$$

This completes the proof of the lemma.

Next result is a Hopf's lemma for anti-symmetric functions.

Lemma 3.2. Let \mathcal{H} be a half space, $D \subset \mathcal{H}$, and $c \in L^{\infty}(D)$. If $u \in C_b(\mathbb{R}^n)$ is an anti-symmetric supersolution of $Lu - \beta |Du| - c(x)u = 0$ in D with $u \ge 0$ in \mathcal{H} , then either $u \equiv 0$ in \mathbb{R}^n or u > 0 in D. Furthermore, if $u \ne 0$ in D and there exists a $x_0 \in \partial D \setminus \partial \mathcal{H}$ with $u(x_0) = 0$ such that there is a ball $B \subset D$ with $x_0 \in \partial B$, then there exists a C > 0 such that

$$\lim\inf_{t\to 0}\frac{u(x_0-t\mathbf{n})}{t} \ge C,$$

where n is the inward normal at x_0 .

Proof. With any loss of generality, we may assume that $c \ge 0$. Suppose that $u \ne 0$ and $u \ne 0$ in D. Then there exists a compact set $K \subset D$ such that $\inf_K u = \delta > 0$ and a point $x_1 \in D$ such that $u(x_1) = 0$. For ε small enough we can choose the test function

$$\phi(y) = \begin{cases} 0 & \text{for } y \in B_{\varepsilon}(x_1), \\ u & \text{for } y \in B_{\varepsilon}^c(x_1). \end{cases}$$

Note that $\Delta \phi(x_1) = 0$, $D\phi(x_1) = 0$. Since k is radially non-increasing and positive, from the proof of [8, Theorem 3.2] it follows that $u \equiv 0$ in D. This contradicts our assertion. Thus either $u \equiv 0$ in \mathbb{R}^n or u > 0 in D.

Now we prove the second part of the lemma. Assume that u > 0 in D. Let B be a ball in D that touches ∂D at x_0 and $B \subseteq \mathcal{H}$. This is possible since $x_0 \in \partial D \setminus \partial \mathcal{H}$. Let ϑ be the positive solution to

$$L\vartheta - \beta |D\vartheta| = -1$$
 in B , and $\vartheta = 0$ in B^c

Existence of ϑ follows from Theorem 1.3 and Leray-Schauder fixed point theorem theorem. Define $w = \kappa(\vartheta - \vartheta \circ \Re)$. Then we have $Lw \ge -\kappa C$ in *B* for some positive constant *C*. Now repeating the arguments of [8, Theorem 3.2] it follows that for some $\kappa > 0$ we have $u \ge w$ in *B*. To complete the proof we need to apply Hopf's lemma on ϑ at the point x_0 (cf. [10, Theorem 2.2]).

Given $\lambda \in \mathbb{R}$, $e \in \partial B_1(0)$, define

$$v(x) = v_{\lambda,e}(x) = u(x) - u(\bar{x}) \quad x \in \mathbb{R}^n,$$
(3.3)

where $\bar{x} = \Re_{\lambda,e}(x)$ denotes the reflection of x by $T_{\lambda,e} := \partial \mathcal{H}_{\lambda,e}$ and $\mathcal{H}_{\lambda,e} = \{x \in \mathbb{R}^n : x \cdot e > \lambda\}$. We note that $\mathbb{R}^n \setminus \overline{\mathcal{H}}_{\lambda,e} = \mathcal{H}_{-\lambda,-e}$. Moreover, let $\lambda < l := \sup_{x \in D} x \cdot e$. Then $\mathcal{H} \cap \Omega$ is nonempty for all $\lambda < l$ and we put $D_{\lambda} := \Re_{\lambda,e}(\Omega \cap \mathcal{H})$. Then for all $\lambda < l$ the function v satisfies

$$Lv + \beta |Dv(x)| - c(x)v \ge 0 \quad \text{in } D_{\lambda},$$

$$Lv - \beta |Dv(x)| - c(x)v \le 0 \quad \text{in } D_{\lambda},$$

$$v \ge 0 \quad \text{in } \mathcal{H}_{-\lambda,-e} \setminus D_{\lambda},$$

$$v(x) = -v(\bar{x}) \quad \text{for all } x \in \mathbb{R}^{n},$$
(3.4)

where β is the Lipschitz constant of H on the interval $[0, \sup |Du|]$ and

$$c(x) = \frac{f(u(x)) - f(u(\bar{x}))}{u(x) - u(\bar{x})}$$

In view of Lemmas 3.1-3.2 and (3.4), we see that $v = v_{\lambda,e}$ is either 0 in \mathbb{R}^n or positive in D_{λ} for λ close to l. Now as we decrease λ one of following two situation may occur.

Situation A: there is a point $p_0 \in \partial \Omega \cap \partial D_{\lambda} \setminus T_{\lambda,e}$,

Situation B: $T_{\lambda,e}$ is orthogonal to $\partial\Omega$ at some point $p_0 \in \partial\Omega \cap T_{\lambda,e}$.

 λ_0 be the maximal value in $(-\infty, l)$ such that one of these situations occur. We show that Ω is symmetric with respect to $T_{\lambda_0,e}$. This would complete the proof of Theorem 3.1 since e is arbitrary. Also, note that, since u > 0 in Ω , to establish the symmetry of Ω with respect to $T_{\lambda_0,e}$, it is enough to show that v = 0 in \mathbb{R}^n . Suppose, to the contrary, that v > 0 in D_{λ_0} .

Situation A: In this case we have $v(p_0) = 0$ and therefore, by Theorem 1.3 and Lemma 3.2, we get $\frac{\partial v}{\partial n}(p_0) > 0$. But, by (3.1), we have

$$\frac{\partial v}{\partial \mathbf{n}}(p_0) = \frac{\partial u}{\partial \mathbf{n}}(p_0) - \frac{\partial u}{\partial \mathbf{n}}(\mathcal{R}(p_0)) = 0.$$

This is a contradiction.

Situation B: This situation is a bit more complicated than the previous one. Set $T = T_{\lambda_0,e}$, $\mathcal{H} = \mathcal{H}_{\lambda_0,e}$ and $\mathcal{R} = \mathcal{R}_{\lambda_0,e}$. By rotation and translation, we may set $\lambda_0 = 0$, $p_0 = 0$, $e = e_1$ and $e_2 \in T$ is the interior normal at ∂D .

Next two lemmas are crucial to get a contradiction in Situation B.

Lemma 3.3. We have

$$v(t\bar{\eta}) = o(t^2), \quad as \ t \to 0^+,$$

where $\bar{\eta} = e_2 - e_1 = (-1, 1, 0, ..., 0) \in \mathbb{R}^n$.

Proof of Lemma 3.3 follows from Theorem 1.2 and [8, Lemma 3.2].

Lemma 3.4. Let $D \subset \mathbb{R}^n$, $n \geq 2$, be an open bounded domain such that $0 \in \partial D$ and $\{x_1 = 0\}$ is orthogonal and there is a ball $B \subset D$ with $\overline{B} \cap \partial \overline{D} = \{0\}$. Denote

$$D^* := D \cap \{x_1 < 0\}$$

and assume that $w \in C_b(\mathbb{R}^n)$ is an anti-symmetric supersolution of

$$Lw - \beta |Dw| - c(x)w = 0 \quad in D^*$$
$$w \ge 0 \quad in \{x_1 < 0\}$$
$$w > 0 \quad in D^*.$$

Let $\bar{\eta} = e_2 - e_1 = (-1, 1, \dots, 0) \in \mathbb{R}^n$, then there exist positive C, t_0 , dependent on D^*, n , such that $w(t\bar{\eta}) > Ct^2$

for all $t \in (0, t_0)$.

Clearly, Lemmas 3.3 and 3.4 give a contradiction to the Situation B.

Proof of Theorem 3.1. Proof follows from the above discussion and the arguments in [19, p. 11]. \Box

In the remaining part of this section we provide a proof of Lemma 3.4.

Proof of Lemma 3.4. We follow the arguments of [8, Lemma 3.3]. Fix a ball $B = B_R(Re_2) \subset D$ for some R > 0 small enough with $\partial B \cap \partial D = \{0\}$. Denote

$$K = \{x_1 < 0\} \cap B.$$

Let $M_1 \in D^*$ such that $\theta := \inf_{M_1} w > 0$ and $M_2 = \Re(M_1)$, that is, reflection of M_1 with respect to $\{x_1 = 0\}$. Furthermore, we may assume that M_1 to be an open ball and taking R smaller, we also assume that $\operatorname{dist}(K, M_1) > 0$ and $|K| < \varepsilon$ for some small $\varepsilon > 0$. Now let g be the unique positive viscosity solution to

$$Lg - \beta |Dg| = -1 \quad \text{in } B$$
$$g = 0 \quad \text{in } B^c.$$

From Theorem 1.1, we know that $||g||_{C^{0,1}(\mathbb{R}^n)} \leq C$. Let $\phi \in C_c^{\infty}(\mathbb{R}^d)$, $\operatorname{support}(\phi) \subset M_1$, $0 \leq \phi \leq 1$ and there exists a $U \subset M_1$ such that $\phi = 1$ in U, |U| > 0. Construct a barrier function h of the following form:

$$h(x) = -\kappa x_1 g(x) + \theta \phi(x) - \theta \phi(\mathcal{R}(x))$$

Choosing $\kappa > 0$ small enough, it can be easily checked that (see [8, Lemma 3.3])

$$Lh - \beta |Dh| + c(x)h \ge 0$$

in K. It is also standard to see that g is radial about the point Re_2 (cf. [10, Theorem 4.1]). Thus we have : (i) w - h is anti symmetric, (ii) $w - h \ge 0$ in $\{x_1 < 0\} \setminus K$, since because $\theta > 0$, and (iii) $L(w-h) - \beta |D(w-h)| - c(x)(w-h) \le 0$. Applying Lemma 3.1, we obtain $w - h \ge 0$ in $\{x_1 < 0\}$. Hence

$$w(t\bar{\eta}) \ge h(t\bar{\eta}) \ge Ct^2$$

for $t \in (0, t_0)$, where we used Hopf's lemma on g. This completes the proof.

Acknowledgement. We thank the referee for his/her careful reading of the manuscript and suggestions. This research of Anup Biswas was supported in part by a Swarnajayanti fellowship (DST/SJF/MSA-01/2019-20). Mitesh Modasiya is partially supported by CSIR PhD fellowship (File no. 09/936(0200)/2018-EMR-I).

References

- R. Arora and V. D Rădulescu: Combined effects in mixed local-nonlocal stationary problems, Preprint, 2022, arXiv: 2111.06701
- [2] G. Barles, E. Chasseigne and C. Imbert: On the Dirichlet problem for second-order elliptic integro-differential equations, *Indiana Univ. Math. J.* 57 (2008), no. 1, 213–246.
- [3] G. Barles and C. Imbert: Second-order elliptic integro-differential equations: viscosity solutions' theory revisited, Ann. Inst. H. Poincaré Anal. Non Linéaire 25 (2008), 567–585.
- [4] S. Biagi, S. Dipierro, E. Valdinoci and E. Vecchi: A Faber-Krahn inequality for mixed local and nonlocal operators, Preprint, 2021. arXiv:2104.00830
- [5] S. Biagi, S. Dipierro, E. Valdinoci and E. Vecchi: Mixed local and nonlocal elliptic operators: regularity and maximum principles, *Communications in Partial Differential Equations* 47 (2022), no. 3, 585–629
- [6] S. Biagi, E. Vecchi, S. Dipierro and E. Valdinoci: Semilinear elliptic equations involving mixed local and nonlocal operators, Proceedings of the Royal Society of Edinburgh Section A: Mathematics, DOI:10.1017/prm.2020.75
- [7] A. Biswas: Principal eigenvalues of a class of nonlinear integro-differential operators, J. Differential Equations 268 (2020), no. 9, 5257–5282.
- [8] A. Biswas and S. Jarohs: On overdetermined problems for a general class of nonlocal operators. Journal of Differential Equations 268, no. 5, 2368–2393, (2020)
- [9] A. Biswas and J. Lőrinczi: Hopf's Lemma for viscosity solutions to a class of non-local equations with applications, *Nonlinear Analysis* 204 (2021), 112194
- [10] A. Biswas and M. Modasiya: Mixed local-nonlocal operators: maximum principles, eigenvalue problems and their applications, preprint, 2021. arXiv: 2110.06746
- [11] L. Caffarelli, M. G. Crandall, M. Kocan and A. Święch: On viscosity solutions of fully nonlinear equations with measurable ingredients. Comm. Pure Appl. Math. 49 (1996), no. 4, 365–397
- [12] L. A. Caffarelli and L. Silvestre: Regularity theory for fully nonlinear integro-differential equations, Comm. Pure Appl. Math. 62 (2009), 597–638.
- [13] L. A. Caffarelli and L. Silvestre: Regularity results for nonlocal equations by approximation, Arch. Ration. Mech. Anal. 200 (2011), no. 1, 59–88
- [14] A. Cianchi and P. Salani: Overdetermined anisotropic elliptic problems, Math. Ann.345.4 (2009), 859–881
- [15] P. Clément and L. A. Peletier: An anti-maximum principle for second-order elliptic operators. J. Differential Equations, 34(2) (1979, 218–229
- [16] M. C. Delfour and J.-P. Zolésio: Shapes and geometries. Metrics, analysis, differential calculus, and optimization. Second edition. Advances in Design and Control, 22. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2011. xxiv+622 pp.
- [17] C De Filippis and G. Mingione: Gradient regularity in mixed local and nonlocal problems, Preprint. Arxiv:2204.06590
- [18] M. M. Fall and S. Jarohs: Gradient estimates in fractional Dirichlet problems, Potential Anal. 54.4 (2021), 627–636
- [19] M. M. Fall and S. Jarohs: Overdetermined problems with fractional Laplacian, ESAIM Control Optim. Calc. Var. 21 (2015), no. 4, 924–938
- [20] M. M. Fall, I. A. Milend and T. Weth: Unbounded periodic solutions to Serrin's overdetermined boundary value problem, Arch. Ration. Mech. Anal. 223 (2017), no. 2, 737–759.
- [21] Y. Fang, B. Shang and C. Zhang: Regularity Theory for Mixed Local and Nonlocal Parabolic p-Laplace Equations, Journal of Geometric Analysis 32 (2022), 22
- [22] A. Farina and B. Kawohl: Remarks on an overdetermined boundary value problem, Calc. Var. Partial Differential Equations 31 (2008), no. 3, 351–357
- [23] A. Farina and E. Valdinoci: Flattening results for elliptic PDEs in unbounded domains with applications to overdetermined problems, Arch. Rational Mech. Anal. 195 (2010), no. 3, 1025–1058.
- [24] M. Foondun: Harmonic functions for a class of integro-differential operators, Potential Anal. 31(1) (2009), 21-44
- [25] M.G. Garroni and J.L. Menaldi: Second order elliptic integro-differential problems, Chapman & Hall/CRC Research Notes in Mathematics, 430. Chapman & Hall/CRC, Boca Raton, FL, 2002. xvi+221 pp.

BOUNDARY REGULARITY

- [26] P. Garain and J. Kinnunen: On the regularity theory for mixed local and nonlocal quasilinear elliptic equations, to appear in Transactions of the AMS, 2022
- [27] P. Garain and E. Lindgren: Higher Hölder regularity for mixed local and nonlocal degenerate elliptic equations. Preprint. Arxiv: 2204.13196
- [28] J. L. Kazdan: Prescribing The Curvature Of A Riemannian Manifold, CBMS Reg. Conf. Ser. Math.57, Amer. Math. Soc., Providence, 1985.
- [29] M. Kim, P. Kim, J. Lee and K-A Lee: Boundary regularity for nonlocal operators with kernel of variable orders, J. Funct. Anal. 277 (2019), no. 1, 279–332.
- [30] N. Krylov: Boundedly inhomogeneous elliptic and parabolic equations in a domain, Izv. Akad. Nauk SSSR Ser. Mat. 47 (1983), 75–108.
- [31] A. Iannizzotto, S. Mosconi and M. Squassina: Fine boundary regularity for the degenerate fractional p-Laplacian. J. Funct. Anal. 279 (2020), no. 8, 108659, 54 pp.
- [32] C. Mou: Existence of C^{α} solutions to integro-PDEs, Calc. Var. Partial Diff. Equ. 58(4) (2019), 1–28
- [33] C. Mou and Y.P. Zhang: Regularity Theory for Second Order Integro-PDEs, Potential Anal 54 (2021), 387–407
- [34] X. Ros-Oton and J. Serra: The Dirichlet problem for the fractional Laplacian: regularity up to the boundary, J. Math. Pures Appl. 101 (2014), 275–302.
- [35] X. Ros-Oton and J. Serra: Boundary regularity for fully nonlinear integro-differential equations, Duke Math. J. 165 (2016), 2079–2154.
- [36] X. Ros-Oton and J. Serra: Boundary regularity estimates for nonlocal elliptic equations in C^1 and $C^{1,\alpha}$ domains, Annali di Matematica Pura ed Applicata 196 (2017), 1637–1668.
- [37] R. Schilling, R. Song, and Z. Vondraček: Bernstein Functions, Walter de Gruyter, 2010.
- [38] J. Serrin: A symmetry problem in potential theory, Arch. Rational Mech. Anal. 43 (1971), 304–318.
- [39] N. Soave and E. Valdinoci: Overdetermined problems for the fractional Laplacian in exterior and annular sets, Journal d'analysis Mathematique 137 (2019), 101–134