

# HEXAGONAL GEOMETRIC TRIANGULATIONS

XIAOPING ZHU

ABSTRACT. It is well-known that the Euclidean plane has a standard 6-regular geodesic triangulation, and the unit sphere has a 5-regular geodesic triangulation, which is induced from the regular Dodecahedron, and the hyperbolic plane has an  $n$ -regular geodesic triangulation for any  $n > 6$ . Here we constructed a 6-regular geodesic triangulation of the hyperbolic plane.

## CONTENTS

1. Introduction	1
2. Preliminary case: closed surfaces	2
3. General facts about regular geometric triangulations	3
4. $k$ -Regular geometric triangulations on The Euclidean plane	4
5. 6-regular geometric triangulations on the Hyperbolic Plane	5
6. Related work	8
References	9

## 1. INTRODUCTION

Here we consider the problem on the existence of special classes of geometric triangulations on the Euclidean plane and the hyperbolic plane.

A triangulation  $T$  on a surface  $S$  is  $k$ -regular for some integer  $k > 0$  if the degrees of all its vertices are  $k$ , and it is *geometric* if all the edges are embedded as geodesic arcs with respect to the metric. The most familiar geometric 6-regular triangulation is the standard hexagonal triangulation on the plane, and the 7-regular equilateral triangulation of the hyperbolic plane. The triangular groups [3]

$$\Delta(3, k) = \{a, b, c | a^3 = b^k = c^2 = 1\}$$

produce  $k$ -regular equilateral triangulations for any  $k > 6$ .

We ask whether we can construct 6-regular geometric triangulations on the hyperbolic plane. Moreover, we can consider the general question as follows.

**Question:** Does there exist  $k$ -regular geometric triangulations on the Euclidean plane  $\mathbb{E}^2$ , the round 2-sphere  $\mathbb{S}^2$  and the hyperbolic plane  $\mathbb{H}^2$  for any  $k \geq 3$ ? Furthermore, can we construct such a geometric triangulation  $T$  with a uniform bound on the edge lengths of  $T$ ?

If  $k = 6$ , the number of vertices grows linearly with respect to the combinatorial distance from one point. If  $k > 6$ , the number of vertices grows exponentially. Intuitively, they correspond to the Euclidean and hyperbolic geometry. We show that we can construct 6-regular geometric triangulation on  $\mathbb{H}^2$ .




FIGURE 1. Degree-regular geometric triangulations.

The semi-regular tiling of the Euclidean and hyperbolic plane by polygons has been studied by Datta and Gupta [1].

## 2. PRELIMINARY CASE: CLOSED SURFACES

The basic combinatorial formulas below of Euler characteristics of closed surfaces show that there is no  $k$ -regular triangulation on  $\mathbb{E}^2$  and  $\mathbb{H}^2$  if  $k = 3, 4, 5$ , and there is no  $k$ -regular triangulation on  $\mathbb{S}^2$  if  $k \geq 6$ .

From the combinatorial formula

$$12(1 - g) = 6\chi(S) = \sum_{i \in V} (6 - d_i) = |V|(6 - k),$$

we have a necessary condition for a closed orientable surface  $S$  to admit a  $k$ -regular triangulation  $T$ , namely

$$(k - 6) | 12(g - 1).$$

It implies immediately that there is no  $k$ -regular triangulation on  $\mathbb{S}^2$  if  $k \geq 6$ , and the only  $k$ -regular triangulation on tori is when  $k = 6$ . This condition turns out to be also sufficient for the existence of  $k$ -regular triangulation on  $S$  with genus  $g$ .

**Lemma 2.0.1.** *Let  $S$  be a surface with genus  $g$ . Then there exists a  $k$ -regular triangulation on  $S$  if and only if*

$$(k - 6) | 12(g - 1).$$

*Proof.* The necessity is proved above. The sufficiency follows from the result by Jucovic and Trenkler [2]. First notice that if a  $k$ -regular triangulation  $T$  exists on  $S$ , then set  $V_0$ ,  $E_0$ , and  $F_0$  be the number of vertices, edges, and faces of  $T$ . Then  $V_0 = 6\chi(S)/(6 - k)$ ,  $E_0 = 3k\chi(S)/(6 - k)$ , and  $F_0 = 2k\chi(S)/(6 - k)$ .

By the Main Theorem in [2], if three positive numbers  $V$ ,  $E$ , and  $F$  on  $S$  satisfies two conditions

- (1)  $F - (4 - k)V = 8(1 - g)$ ;
- (2)  $kV = 3F = 2E$ ,

then a  $k$ -regular triangulation with  $V$  vertices,  $E$  edges and  $F$  faces exists on  $S$ . It is straightforward to check that  $V_0$ ,  $E_0$ , and  $F_0$  satisfies the two conditions.  $\square$

**Lemma 2.0.2.** *There is no  $k$ -regular triangulation on  $\mathbb{E}^2$  and  $\mathbb{H}^2$  if  $k = 3, 4, 5$ .*

*Proof.* If such a triangulation  $T$  exists, take an arbitrary subcomplex  $T'$  which is homeomorphic disk. A combinatorial formula for a triangulated disk is given by

$$\sum_{i \in V_I} (6 - d_i) = 6 + \sum_{i \in V_B} (d_i - 4)$$

where  $V_I$  and  $V_B$  are the index sets for interior vertices and boundary vertices. Since  $T$  is  $k$ -regular for  $k < 6$ , if  $v_i$  is a boundary vertex of  $T'$ , then  $d_i < 5$  since  $T'$  is part of  $T$ . Then the left side is no larger than 6. Hence the disk  $T'$  can only contains at most 6 vertices. This leads to a contradiction because we can take  $T'$  as large as we want. □

For  $k = 3, 4, 5$ , we have  $k$ -regular geometric triangulations on  $\mathbb{S}^2$ , given by projecting regular tetrahedron, octahedron, and icosahedron to the unit sphere from the center. It is straightforward to construct a 6-regular triangulation on a torus using the standard hexagonal triangulation on  $\mathbb{E}^2$ .




FIGURE 2. Degree-regular geometric triangulations on the unit 2-sphere.

### 3. GENERAL FACTS ABOUT REGULAR GEOMETRIC TRIANGULATIONS

The goal is to construct  $k$ -regular geometric triangulations on  $\mathbb{E}^2$  with  $k > 6$ , and 6-regular geometric triangulations on  $\mathbb{H}^2$ . We first point out that such triangulations contain skinny triangles.

**Lemma 3.0.1.** *Let  $T$  be a  $k$ -regular geometric triangulations on  $\mathbb{E}^2$  with  $k > 6$ , or a 6-regular geometric triangulations on  $\mathbb{H}^2$ . Then the infimum of the angles in  $T$  is zero.*

*Proof.* If not, in the first case for  $\mathbb{E}^2$  we can construct a quasiconformal homeomorphism from  $\mathbb{E}^2$  to  $\mathbb{H}^2$  by sending a triangle in  $T$  to the corresponding triangle in the standard geometric triangulation of  $\mathbb{H}^2$  generated by reflecting equilateral triangles with inner angle  $2\pi/k$ . Similarly, in the second case we can construct a quasiconformal homeomorphism from  $\mathbb{H}^2$  to  $\mathbb{E}^2$  sending  $T$  to the standard equilateral hexagonal triangulation. This contradicts to the fact that there is no quasiconformal map from  $\mathbb{E}^2$  to  $\mathbb{H}^2$ . □

In this sense, the “best”  $k$ -regular geometric triangulation we can expect is a geometric triangulation with a uniform low bound  $c > 0$  and a uniform upper  $C$  on the lengths of edges.

**Conjecture 3.0.2.** *There is no  $k$ -regular geometric triangulation on  $\mathbb{E}^2$  with a uniform low bound  $c > 0$  and a uniform upper  $C$  on the lengths of edges for  $k \geq 7$ . Similarly, there is no 6-regular geometric triangulation on  $\mathbb{H}^2$  with a uniform low bound  $c > 0$  and a uniform upper  $C$  on the lengths of edges.*

4.  $k$ -REGULAR GEOMETRIC TRIANGULATIONS ON THE EUCLIDEAN PLANE

It is known that one can construct  $k$ -regular geometric triangulation on  $\mathbb{E}^2$  for  $k > 6$ . The idea is to construct chains of circles with radius  $n$  and distribute the points on the circle evenly. The similar computation is given in [1]. We can show that these triangulations have the desired uniform bounds.

**Lemma 4.0.1.** *For any  $k \geq 6$ , there exist  $k$ -regular geometric triangulations on  $\mathbb{E}^2$  with uniform upper bounds on the lengths of edges.*

*Proof.* The construction of a  $k$ -regular geometric triangulation is given above. Let  $C_n$  be circles with radius  $r_n = n$  centered at the origin of  $\mathbb{E}^2$ . Let  $a_n$  be the number of vertices on  $C_n$  on this construction. These vertices have combinatorial distance  $n$  from the vertex at the origin.

We have the following recursive relation:

$$a_{n+1} = (k - 4)a_n - a_{n-1}$$

with  $a_0 = 0$  and  $a_1 = k$ . Solve this series and we have the formula for  $a_n$  with  $n \geq 1$ :

$$a_n = k \frac{\alpha^n - \beta^n}{\alpha - \beta}$$

where  $\alpha = \cosh^{-1}((k - 4)/2)$  and  $\beta = 1/\alpha$ . Thus, asymptotically  $a_n$  grows as  $\alpha^{n-1}$  to infinity. But the length of  $C_n$  is given by  $l_n = 2\pi n$ . So the arclength of each small arc goes to zero as  $n$  goes to infinity, hence the length between two consecutive vertices on  $C_n$  goes to zero.

By the construction above, if  $v$  is a vertex in  $C_n$ , then the  $k - 4$  or  $k - 3$  vertices  $v_i$  in  $C_{n-1}$  connecting to  $v$  forms an arc  $\gamma$  in  $C_{n+1}$ , and the ray starting from the origin passing through  $v$  intersects with  $\gamma$ . As the length of  $\gamma$  goes to zero, all the lengths of the edges connecting  $v$  to  $v_i$  goes to  $r_{n+1} - r_n = 1$ .

So as  $n \rightarrow \infty$ , the lengths of edges approaches either 1 or 0. Hence we can take the maximal length of a compact part of  $T$  bounded by  $C_n$  with  $n$  large enough so that its complement contains edges no longer than 2. Then we can find a uniform bound. □

Using the same idea, we can construct a  $k$ -regular geometric triangulation of  $\mathbb{E}^2$  whose edges has a uniform lower bound. We can pick  $r_n = 2^n$  and follow the construction above.




FIGURE 3. A 7-regular geometric triangulations on the Euclidean plane.




FIGURE 4. Tianqi's Picture.

### 5. 6-REGULAR GEOMETRIC TRIANGULATIONS ON THE HYPERBOLIC PLANE

We now give a construction of the desired 6-regular geometric triangulations with uniform bound in the hyperbolic plane using the Klein disk model.

- (1) Put one vertex  $v_0$  in the origin, and shoot out two rays  $R_1$  and  $R_2$  along the positive  $x$ -axis and along the direction  $e^{i\pi/3}$ .
- (2) Let  $r_n = \alpha \log n$  with  $0 < \alpha < 1/2$ . Construct a sequence of points  $a_n$  and  $b_n$  on  $R_1$  and  $R_2$  with distance  $r_n$  from  $v_0$  in hyperbolic metric. Connect  $a_n$  with  $a_{n+1}$  and  $b_n$  with  $b_{n+1}$  using geodesics. Call these edges of *type 0*.
- (3) Connect  $a_n$  with  $b_n$  by circle arcs  $C_n$ , called the  $n$ -th layer in this construction. Equally distribute  $n - 1$  points on  $C_n$ . Then each  $C_n$  are divided to  $n$  arcs. Connect the consecutive vertices on  $C_n$  to generate  $n$  edges. Call these edges of *type 1*. By symmetry, these edge have the same length.

- (4) Connect points on  $C_n$  and  $C_{n+1}$  based on the combinatorics of the hexagonal triangulation to generate  $2(n+1)$  edges. Call these edges of *type 2*.
- (5) Rotate the configuration by  $\pi/3$  to generate the full hexagon geometric triangulation.

We can give explicit coordinates of the vertices on  $C_n$  in this triangulation in the plane

$$v_n^k = \tanh r_n e^{i \frac{k\pi}{3n}}, \quad k = 0, 1, \dots, n.$$

And the vertex  $v_n^k$  is connected with vertices  $v_{n+1}^k$  and  $v_{n+1}^{k+1}$  in this triangulation.

**Proposition 5.0.1.** *The construction above generates a hexagonal geometric triangulation of  $\mathbb{H}^2$  with uniform bound  $K > 0$  on the lengths of edges.*

The proof of this proposition consists of two lemmas.

**Lemma 5.0.2.** *The construction above generates a valid geometric triangulations with no intersection of edges and degenerate triangles.*

*Proof.* The proof is based on induction on  $n$ . The base case  $n = 1$  is trivial. Assume it is true for the triangulation generated up to the  $n$ -th layer. We will show that the triangles between  $C_n$  and  $C_{n+1}$  lie in the ring  $R_n$  bounded  $C_n$  and  $C_{n+1}$ . There are two types of triangles:

- (1) Type 1: two vertices on  $C_n$  and one vertex on  $C_{n+1}$ . The three vertices are

$$v_n^k = \tanh r_n e^{i \frac{k\pi}{3n}}, v_n^{k+1} = \tanh r_n e^{i \frac{(k+1)\pi}{3n}}, \text{ and } v_{n+1}^{k+1} = \tanh r_{n+1} e^{i \frac{(k+1)\pi}{3(n+1)}}.$$

Notice that

$$\frac{k}{n} < \frac{k+1}{n+1} < \frac{k+1}{n}, \text{ and } r_n \leq r_{n+1},$$

- (2) Type 2: one vertex on  $C_n$  and two vertex on  $C_{n+1}$ . In this case, the three vertices are

$$v_n^k = \tanh r_n e^{i \frac{k\pi}{3n}}, v_{n+1}^{k+1} = \tanh r_{n+1} e^{i \frac{(k+1)\pi}{3n}}, \text{ and } v_{n+1}^k = \tanh r_{n+1} e^{i \frac{k\pi}{3(n+1)}}.$$

Notice that

$$\frac{k}{n+1} \leq \frac{k}{n} \leq \frac{k+1}{n+1}, \text{ and } r_n \leq r_{n+1}.$$

We want to show that the edges of type 2 lie in the ring bounded by two consecutive  $C_n$  and  $C_{n+1}$ , so it does not intersect with previous layers. If  $r_n$  increase too slowly, then it is possible that the edge determined by  $v_{n+1}^k$  and  $v_{n+1}^{k+1}$  might intersect with  $v_n^k$ . We need to show that if  $r_n = \alpha \log n$  with  $0 < \alpha < 1/2$ , this will not occur.

The critical figures for triangles of type 1 and type 2 are given below, where the angles at  $v_n^k$  or  $v_n^{k+1}$  are right angles.




FIGURE 5. Critical Cases for Two Types of Triangles

In this case, we can show that for sufficiently large  $n > N$ ,

$$\tanh r_n < \tanh r_{n+1} \cos \frac{\pi}{6(n+1)},$$

and

$$\tanh r_n < \tanh r_{n+1} \cos \frac{\pi}{6n},$$

which is equivalent to

$$\sinh(r_{n+1} - r_n) > \cosh r_n \sinh r_{n+1} \left(1 - \cos \frac{\pi}{6n}\right).$$

One can show that the left side has order  $O(1/n)$  and right side has order  $O(1/n^{2-2\alpha})$ , so the inequality holds for large  $n$  if  $0 < \alpha < 1/2$ .

This means that for large  $n > N$ , the triangles of type 2 will not intersect with the previous layers. For  $0 \leq n \leq N$ , we can treat it as a polygon in the Euclidean plane, and adjust the radius to avoid the intersections.  $\square$

**Lemma 5.0.3.** *The lengths of edges of the geometric triangulation above are uniformly bounded.*

*Proof.* The length of the edge of type 1 is bounded by the length of the arc of the circle connecting  $v_n^k$  and  $v_n^{k+1}$  given by

$$\sinh |v_n^k v_n^{k+1}| < \frac{\pi}{3n} \sinh r_n \sim O\left(\frac{1}{n^{1-\alpha}}\right) \rightarrow 0.$$

From the previous lemma, all the edges of type 0 and type 2 in the  $n$ -th layer lie in the ring  $R_n$ . The length of the edges in  $R_n$  with vertices on  $C_n$  is bounded by the following situation




FIGURE 6. Estimate for Edges of Type 2.

The half length  $L$  of this edge is given by hyperbolic Pythagorean theorem

$$L = 2 \cosh^{-1}\left(\frac{\cosh r_{n+1}}{\cosh r_n}\right) \leq 2 \cosh^{-1}\left(2\left(1 + \frac{1}{n}\right)^\alpha\right) < \infty.$$

Hence we can bound the edges of all the three types for large  $n > 0$ . The remaining part is naturally bounded by compactness.  $\square$

## 6. RELATED WORK

The existence of certain types of geodesic triangulations is a fundamental question. Recently various theory upon geodesic triangle meshes were developed, where people discuss problems about rigidity, convergence, variational principles, discrete maps and discrete geometric structures. See [3-21] for example.

## REFERENCES

- [1] Basudeb Datta and Subhojoy Gupta, *Semi-regular tilings of the hyperbolic plane*, *Discrete & Computational Geometry* **65** (2021), no. 2, 531–553.
- [2] Ernest Jucovič and Marián Trenkler, *A theorem on the structure of cell-decompositions of orientable 2-manifolds*, *Mathematika* **20** (1973), no. 1, 63–82.
- [3] Bruno Martelli, *An introduction to geometric topology*, arXiv preprint arXiv:1610.02592 (2016).

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ, 08854

*Email address:* `zx349@rutgers.edu`