# The Lack of Convexity of the Relevance-Compression Function

Albert E. Parker Center for Biofilm Engineering Department of Mathematical Sciences Montana State University Bozeman, MT USA parker@math.montana.edu Tomáš Gedeon Department of Mathematical Sciences Montana State University Bozeman, MT USA gedeon@math.montana.edu

Alexander G. Dimitrov Department of Mathematics Department of Neuroscience Washington State University Vancouver, WA alex.dimitrov@wsu.edu

## Abstract

In this paper we investigate the convexity of the relevance-compression function for the Information Bottleneck and the Information Distortion problems. This curve is an analog of the rate-distortion curve, which is convex. In the problems we discuss in this paper, the distortion function is not a linear function of the quantizer, and the relevance-compression function is not necessarily convex (concave), but can change its convexity. We relate this phenomena with existence of first order phase transitions in the corresponding Lagrangian as a function of the annealing parameter.

## 1 Introduction

In previous work [1, 2, 3] we have described the bifurcation structure for solutions to problems of the form

$$\max_{q \in \Delta} G(q)$$

$$D(q) \ge D_0$$
(1)

where  $\Delta$  is a constraint space of valid conditional probabilities, G and D are continuous, real valued functions of q, smooth in the interior of  $\Delta$ , and the functions G(q) and D(q)are invariant under the group of symmetries  $S_N$ . This type of problem, which arises in Rate Distortion Theory [4, 5] and Deterministic Annealing [6], is NP complete [7] when D(q) is the mutual information  $I(X; Y_N)$  as in the Information Bottleneck [8, 9, 10] and the Information Distortion [11, 12, 13] methods. In this paper, we address the relationship between the bifurcation structure of solutions to (1) and the relevance compression function [10].

# 2 Preliminaries

## 2.1 Rate Distortion Theory

We assume that the random variable  $X \in \{x_1, x_2, ..., x_K\}$  is an input source, and that  $Y \in \{y_1, y_2, ..., y_K\}$  is an output source. In rate distortion theory [4], the random variable Y is represented by using N symbols or *classes*, which we call  $Y_N$ , where we assume without loss of generality that  $Y_N \in \{1, 2, ..., N\}$ . We denote a stochastic clustering or *quantization*, of the realizations of Y to the classes of  $Y_N$ , by  $q(Y_N|Y)$ . To find the quantization that yields the minimum information rate  $I(Y; Y_N)$  at a given distortion, one can find points on the rate distortion curve for each value of  $D_0 \in [0, D_{\text{max}}]$ . The rate distortion curve is defined as [4, 5]

$$R(D_0) := \begin{array}{c} \min_{q \in \Delta} I(Y; Y_N) \\ D(Y, Y_N) \le D_0 \end{array},$$

$$(2)$$

where D(q) is a distortion function. A quantization  $q(Y_N|Y)$  that satisfies (2) yields an approximation of the probabilistic relationship, p(X, Y), between X and Y [11, 8, 9]. The constraint space  $\Delta$  is the space of valid finite conditional probabilities  $q(Y_N|Y)$ , where we will write  $q(Y_N = \nu|Y = y_k) = q_{\nu k}$ .

The Information Bottleneck method [8, 9, 10] uses the *information distortion function*  $D(q) := I(X;Y) - I(X;Y_N)$ . Since the spaces X and Y are fixed, then I(X;Y) is fixed, and so the rate distortion problem (2) in the case of the Information Bottleneck problem can rewritten as

$$R_I(I_0) := \begin{array}{c} \max_{q \in \Delta} -I(Y; Y_N) \\ I(X; Y_N) \ge I_0 \end{array}$$
(3)

where  $I_0 > 0$  is some information rate. The function  $R_I(I_0)$  is referred to as the *relevance-compression function* in [10]. Observe that there is a one to one correspondence between  $I_0$  and  $D_0$  via  $I_0 = I(X;Y) - D_0$ . To solve the neural coding problem, the Information Distortion method [11, 12, 13] considers a problem of the form

$$R_H(I_0) := \begin{array}{c} \max_{q \in \Delta} H(Y_N|Y) \\ I(X;Y_N) \ge I_0 \end{array}$$

$$\tag{4}$$

where  $H(Y_N|Y)$  is the conditional entropy.

## 2.2 Annealing

Using the method of Lagrange multipliers, an arbitrary problem of the form (1) is rewritten as

$$\max_{q \in \Delta} \left( G(q) \right) + \beta (D(q) - D_0) \right).$$
(5)

As we will see, solutions of (1) are not always solutions of (5). Similarly, the problem (3) can be rewritten [8, 9, 10] as

$$\max_{q \in \Delta} \left( -I(Y_N, Y) + \beta I(X; Y_N) \right), \tag{6}$$

and problem (4) can be rewritten [11, 12, 13], in analogy with Deterministic Annealing [6], as

$$\max_{q \in \Delta} \left( H(Y_N | Y) + \beta I(X; Y_N) \right). \tag{7}$$

In (5), (6) and (7), the Lagrange multiplier  $\beta$  can be viewed as an annealing parameter.



Figure 1: (A) The bifurcation structure of stationary points of the Information Distortion problem (4), a problem of the form (1). We found these points by annealing in  $\beta$  and finding stationary points for the problem (7) using the algorithm which we presented in [14]. A square indicates where a bifurcation occurs. (B) A close up of the subcritical bifurcation at  $\beta \approx 1.038706$ , indicated by an square. Observe the subcritical bifurcating branch, and the subsequent saddle-node bifurcation at  $\beta \approx 1.037479$ , indicated by another square. We applied Theorem 1 to show that the subcritical bifurcating branch is composed of quantizers which are solutions of (4) but not of (7).

### 2.3 Bifurcation Structure of solutions

In [14], we presented an algorithm which can be used to determine the bifurcation structure of stationary points of (5) for each value of  $\beta \in [0, \beta_{\max})$  for some  $\beta_{\max} > 0$ . These stationary points are quantizers  $q^*(\beta) \in \Re^{NK}$  where there exists a vector of Lagrange multipliers  $\lambda^* \in \Re^K$  such that the gradient of the Lagrangian of (1) is a vector of 0's,

$$\nabla_{q,\lambda}(G(q^*) + \beta D(q^*) + \sum_{k=1}^{K} \lambda_k^* (\sum_{\nu} q^*(\nu | y_k) - 1) = \mathbf{0}.$$

This condition is also known as the Karush-Kuhn-Tucker necessary condition for constrained optimality [15]. It is well known in optimization theory that a stationary point, i.e. the point satisfying (8), is a solution of (1) if the matrix of second derivatives, the Hessian  $\Delta_q(G(q) + \beta D(q))$ , is negative definite on the kernel of the Jacobian of the constraints [15]. We have the following results.

**Theorem 1** [1] A stationary point  $q^*$ , is a solution of (5) if  $\Delta_q(G(q^*) + \beta D(q^*))$  is negative definite on ker  $(I_K \ I_K \ \dots \ I_K)$ . A stationary point  $q^*$  is a solution of (1) if  $\Delta_q(G(q^*) + \beta D(q^*))$  is negative definite on ker  $\begin{pmatrix} \nabla_q D(q^*)^T \\ I_K \ I_K \ \dots \ I_K \end{pmatrix}$ .

From Theorem 1 we see that there may be solutions of (1) which are not solutions of (5). We illustrate this fact numerically. For the Information Distortion problem (4) [11, 12, 13], and the synthetic data set composed of a mixture of four Gaussians which the authors used in [11], we determined the bifurcation structure of solutions to (4) by annealing in  $\beta$  and finding the corresponding stationary points to (7) (see Figure 1).

Similar to the results which we presented in [14], the close up of the bifurcation at  $\beta \approx 1.038706$  in Figure 1(B) shows a subcritical bifurcating branch (a first order phase transition) which consists of stationary points of the problem (7). By projecting the Hessian  $\Delta_q(G(q^*)+\beta D(q^*))$  onto each of the kernels referenced in Theorem 1, we determined



Figure 2: (A) The distortion curve  $R_H(I_0)$  defined in (9). For each value of  $I_0$ , we solved the problem (4) to ascertain  $H(Y_N|Y)$ . Observe that the quantizers which yield  $I(X;Y_N) \ge I_0$  for the values  $I_0 \in [0,.03]$  correspond to solutions on the subcritical bifurcating branch shown in Figure fig:bifstructure(B). (B) For each value of  $I_0$ , we solved (4), and found the corresponding Lagrange multiplier. This plot shows  $I_0$  as function of this Lagrange multiplier. This plot is identical to the subcritical bifurcation shown in Figure 1(B), which shows  $I(X;Y_N)$  as a function of the annealing parameter  $\beta$ .

that the points on this subcritical branch are **not** solutions of (7), and yet they **are** solutions of (1).

Furthermore, observe that Figure 1(B) indicates that a saddle-node bifurcation occurs at  $\beta \approx 1.037479$ . That this is indeed the case was proved in [1]. In fact, for any problem of the form (5), there are only two types of bifurcations to be expected.

**Theorem 2** [1] Generically, for problems of the form (1), only symmetry breaking pitchfork-like and saddle-node bifurcations occur.

Clearly, the existence of saddle-node bifurcation at  $\beta \approx 1.037479$  is tied to the existence of subcritical bifurcation (first order phase transition) at  $\beta \approx 1.038706$ . We now investigate the connection between existence of subcritical bifurcations and the convexity of the relevance-compression function.

# 3 The Relevance-Compression Function

Given the generic existence of subcritical pitchfork-like and saddle-node bifurcations of solutions to problems of the form (1), a natural question arises: What are the implications for the rate distortion curve (2)? We examine this question for the information distortion  $D(q) = I(X;Y) - I(X;Y_N)$ , used by the Information Bottleneck and the Information Distortion methods. Recall that the relevance-distortion function is

$$R_I(I_0) := \max_{\Delta \cap \mathcal{Q}_{I_0}} -I(Y_N, Y).$$
(8)

where

$$\mathcal{Q}_{I_0} := \{ q \in \Delta \mid I(X, Y_N) \ge I_0 \}$$

For the Information Distortion problem the relevance-distortion function is

$$R_H(I_0) := \max_{\Delta \cap \mathcal{Q}_{I_0}} H(Y_N | Y).$$
(9)

In Figure 2(A), we present a plot of  $R_H(I_0)$ , which was computed using the same data set of a mixture of four Gaussians which we used in Figures 1(A) and (B). The plot was obtained by solving the problem (4) for each value of  $I_0$ .

To make explicit the relationship between the bifurcation structure shown in Figure 1, which was obtained by annealing in  $\beta$ , and the distortion curve shown in Figure 2(A), which was obtained by annealing in  $I_0$ , we present Figure 2(B). When solving (4) for each  $I_0$ , we computed the corresponding Lagrange multiplier  $\beta$ . Thus,  $\beta = \beta(I_0)$ , which is the curve we show in Figure 2(B). This plot is matches precisely the subcritical bifurcating branch from Figure 1(B), which we obtained by annealing in  $\beta$ .

**Lemma 3** For a fixed value of  $I_0 > 0$  the solution  $q^*$  of (3) and (4) satisfies the equality constraint  $I = I_0$ .

*Proof.* Assume  $q^*$  is a maximizer of (4) and  $I(q^*) > I_0$ . Then the constraint is not active and we must have  $\nabla H(q^*) = 0$ . Since  $\nabla H(q^*) = 0$  implies  $q^* = 1/N$ , we get  $I(q^*) = 0$ . This is a contradiction and thus  $I(q^*) = I_0$ .

Now assume  $q^*$  is a maximizer of (3) and  $I(q^*) > I_0$ . Then again the constraint is not active and we must have  $\nabla I(Y, Y_N) = 0$ . Short computation shows that the condition  $\nabla I(Y, Y_N) = 0$  implies  $q^* = (q_{\nu k})$  is of the form  $q_{\nu k} = a_{\nu}$ , i.e.  $q_{\nu k}$  does not depend on k. However, at such value of  $q^*$  we get  $I(q^*) = 0$ . This is again a contradiction and thus  $I(q^*) = I_0$ .

As a consequence of the Lemma, for each  $I_0 > 0$  there exists a Lagrange multiplier  $\beta(I_0)$ . The existence of subcritical bifurcation branch implies that along this branch  $\beta(I_0)$  is not a one-to-one function of  $I_0$ , and therefore not invertible.

#### 3.1 Properties of relevance-compression function

It is well known that if the distortion function D(q) is linear in q, that  $R(D_0)$  is a nonincreasing and convex [4, 5]. The proof of this result first establishes that the rate-distortion curve is monotone and that it is convex. These two properties together imply continuity and strict monotonicity of the rate distortion curve. Since the information distortion  $D = I(X;Y) - I(X;Y_N)$  is not a linear function of q, the convexity proof given in [4, 5] does not generalize to prove that either (8) or (9) is convex. Therefore we need to prove continuity of the relevance-compression function using other means.

**Lemma 4** The curves  $R_H$  and  $R_I$  are non-increasing curves on  $I_0 \in [0, I_{\max}]$  and are continuous for  $I_0 \in (0, I_{\max})$ .

*Proof.* Observe that since  $I(X, Y_N)$  is convex [11] in quantizer q, we have that

 $\mathcal{Q}_{I_1} \subset \mathcal{Q}_{I_2}$  whenever  $I_1 \geq I_2$ .

Therefore, if  $I_1 \ge I_2$ , then the maximization at  $I_1$  happens over a smaller set than in  $\mathcal{Q}_{\mathcal{I}_{\epsilon}}$ , and so  $R(I_1) \le R(I_2)$ .

Now we prove continuity. Take an arbitrary  $I_0 \in (0, I_{\text{max}})$ . Let

$$M_{I_0} := \{ y \mid y = H(q) \text{ where } q \in \Delta \cap \mathcal{Q}_{D_0} \}$$

be the range (in IR) of the function H(q) with the domain  $\Delta \cap Q_{I_0}$ . Given an arbitrary  $\epsilon > 0$ , let  $M_{I_0}^{\epsilon}$  be an  $\epsilon$  neighborhood of  $M_{I_0}$  in IR. A direct computation shows that  $\nabla_q H(q) = \mathbf{0}$  if and only if q is homogeneous, i.e. qnuk = 1/N, where N is the number of classes of  $Y_N$ . Since H(q) is continuous on  $\Delta$ , then the set  $H^{-1}(M_{I_0}^{\epsilon})$  is a relatively open set in  $\Delta$ . Because by definition  $H(\Delta \cap Q_{I_0}) = M_{I_0}$ , we see that

$$\mathcal{Q}_{I_0} \cap \Delta \subset H^{-1}(M_{I_0}^{\epsilon}). \tag{10}$$

Furthermore, since  $\nabla H(q) \neq 0$  for  $q \in Q_{I_0}$ , then, by the Inverse Mapping Theorem,  $H^{-1}(M_{I_0}^{\epsilon})$  is an open neighborhood of  $Q_{I_0}$ .

The function  $I(X; Y_N)$  is also continuous in the interior of  $\Delta$ . Observe that

$$Q_{I_0} = I^{-1}([I_0, I_{\max}])$$

is closed, and thus  $Q_{I_0} \cap \Delta$  is closed and hence compact. Thus, by (10)  $H^{-1}(M_{I_0}^{\epsilon})$  is an relatively open neighborhood of a compact set  $Q_{I_0} \cap \Delta$ . Therefore, since  $I(X; Y_N)$  is continuous, there exists a  $\delta > 0$  such that the set

$$\operatorname{Int} \mathcal{Q}_{I_0+\delta} \cap \Delta = I^{-1}((I_0+\delta, D_{\max}]) \cap \Delta$$

is a relatively open set in  $\Delta$  such that

$$\mathcal{Q}_{I_0} \cap \Delta \subset \operatorname{Int} \mathcal{Q}_{I_0+\delta} \subset H^{-1}(M_{I_0}^{\epsilon}).$$

It then follows that

$$|\max_{\Delta \cap \mathcal{Q}_{I_0+\delta}} H - \max_{\Delta \cap \mathcal{Q}_{I_0}} H| < \epsilon.$$

By definition of the rate distortion function, this means that

$$|R(I) - R(I_0)| < \epsilon$$
 whenever  $I - I_0 < \delta$ .

Since  $\epsilon$  was arbitrary, this implies continuity of R(I) at  $I = I_0$ .

# **3.2** The Derivative $\frac{\partial R}{\partial I_0}$

In [8, 10], using variational notation, it is shown that

$$\frac{\delta R_I}{\delta D} = -\beta$$

For the sake of completeness, we will reprove this, acknowledging explicitly the fact that the problems (3) and (4) are constrained problems.

**Theorem 5** If relevance-compression functions  $R_I(I_0)$  and  $R_H(I_0)$  are differentiable, then

$$\frac{dR}{dI_0} = -\beta(I_0) \text{ and } \frac{d^2R}{dI_0^2} = -\frac{d\beta(I_0)}{dI_0}$$
(11)

**Corollary 6** Since  $\frac{d\beta(I_0)}{dI_0}$  changes sign at saddle-node bifurcation, then the relevancecompression functions  $R_I(I_0)$  and  $R_H(I_0)$  are neither concave, nor convex.

## Proof of Theorem 5: We start with

$$\max_{q \in \Delta} R(q) + \beta D(q) + \sum_{k} \lambda_k (\sum_{\nu} q_{\nu k} - 1),$$
(12)

where R(q) is one of  $R_I(q) := H(Y_N|Y), R_H(q) := -I(Y_N, Y)$ . We parameterize the solution  $q^*$  locally by  $\beta$ . This can be done everywhere except if  $q^*$  is at a saddle-node bifurcation. At  $q^*(\beta)$ ,

$$\nabla_q R + \beta \nabla_q D + \lambda = 0. \tag{13}$$

**Lemma 7** For  $q \in \Delta$ ,

$$q \cdot \nabla_q R_H = R_H + 1, \ q \cdot \nabla_q R_I = R_I, \ \text{and} \ q \cdot \nabla_q I = I.$$
(14)

*Proof.* Direct calculation.

Hence (13) implies

$$R(q^*(\beta)) + c + \beta I(q^*(\beta)) + q \cdot \vec{\lambda} = 0$$
(15)

Here c = 1 for  $R_H$  and c = 0 for  $R_I$  is a constant. For  $q \in \Delta$ , we set

$$\Lambda(\beta) := q \cdot \vec{\lambda} = \sum_k \lambda_k.$$

The equation (15) defines a relation between R and I. Recall, that we can always express  $\beta = \beta(I_0)$ . Then the term  $I(q^*(\beta(I_0))) = I_0$  and we have a relationship

$$R(I_0) + c + \beta(I_0)I_0 + \Lambda(I_0) = 0.$$
(16)

We differentiate (16):

$$\frac{dR}{dI_0} + \frac{d\beta}{dI_0}I_0 + \beta(I_0) + \frac{d\Lambda}{dI_0} = 0 \Rightarrow$$
(17)

which shows that  $\frac{dR}{dI_0} = -\frac{d\beta}{dI_0}(I_0 + \frac{d\Lambda}{d\beta}) - \beta(I_0)$  since  $\frac{d\Lambda}{dI_0} = \frac{d\Lambda}{d\beta}\frac{d\beta}{dI_0}$ .

In [11](equation (10)) we have an explicit expression for  $\lambda_k$  as a function of  $\beta$ :

$$\lambda_k = p_k (1 - \ln \sum_{\nu} e^{\beta(\nabla I)_{\nu k}/p_k}).$$
(18)

Differentiating this with respect to  $\beta$  yields

$$\frac{d\lambda_k}{d\beta} = -p_k \frac{\sum_{\nu} e^{\beta(\nabla I)_{\nu k}/p_k} (\nabla I)_{\nu k}/p_k}{\sum_{\nu} e^{\beta(\nabla I)_{\nu k}/p_k}}.$$

Since  $\Lambda = \sum_k \lambda_k$ , this implies that

$$\frac{d\Lambda}{d\beta} = \sum_{k} \frac{d\lambda_k}{d\beta} = -\sum_{\nu k} \frac{e^{\beta(\nabla I)_{\nu k}/p_k}}{\sum_{\mu} e^{\beta(\nabla I)_{\mu k}/p_k}} (\nabla I)_{\nu k}$$

For a solution  $q^*$ ,  $\frac{e^{\beta(\nabla I)_{\nu k}/p_k}}{\sum_{\mu} e^{\beta(\nabla I)_{\mu k}/p_k}} = q^*_{\nu k}$  ([11], (12)), hence

$$\frac{d\Lambda}{d\beta}|_{q^*} = -\sum_{\nu k} q^*_{\nu k} (\nabla I)_{\nu k} = -q^* \cdot \nabla I = -I$$

This shows that the term  $I_0 + \frac{d\Lambda}{d\beta} = 0$  at  $q^*$ , hence from (17) we get the first part of (11). The second part follows immediately.

## Acknowledgments

This research is partially supported by NSF grants DGE 9972824, MRI 9871191, and EIA-0129895; and NIH Grant R01 MH57179.

## References

- [1] Albert E. Parker and Tomas Gedeon. Bifurcations of a class of  $S_n$ -invariant constrained optimization problems. *Journal of Dynamics and Differential Equations*, 16(3):629–678, July 2004. Second special issue dedicated to Shui-Nee Chow.
- [2] A Parker, A Dimitrov, and T Gedeon. Symmetry breaking clusters in soft clustering decoding of neural codes. *IEEE Trans. Inform. Theor.*, 56:901–927, 2010.

- [3] T. Gedeon, A. Parker, and A. Dimitrov. The mathematical structure of information bottleneck methods. *Entropy: Special issue for the Information Bottleneck Method*, 14:456–479, 2012.
- [4] Thomas Cover and Jay Thomas. *Elements of Information Theory*. Wiley Series in Communication, New York, 1991.
- [5] Robert M. Gray. Entropy and Information Theory. Springer-Verlag, 1990.
- [6] Kenneth Rose. Deterministic annealing for clustering, compression, classification, regression, and related optimization problems. *Proc. IEEE*, 86(11):2210–2239, 1998.
- [7] Brendan Mumey and Tomas Gedeon. Optimal mutual information quantization is np-complete. Neural Information Coding (NIC) workshop, Snowbird UT, 2003.
- [8] Noam Slonim and Naftali Tishby. Agglomerative information bottleneck. In S. A. Solla, T. K. Leen, and K.-R. Muller, editors, *Advances in Neural Information Processing Systems*, volume 12, pages 617–623. MIT Press, 2000.
- [9] Naftali Tishby, Fernando C. Pereira, and William Bialek. The information bottleneck method. The 37th annual Allerton Conference on Communication, Control, and Computing, 1999.
- [10] Noam Slonim. The information bottleneck: Theory and applications. Doctoral Thesis, Hebrew University, 2002.
- [11] Alexander G. Dimitrov and John P. Miller. Neural coding and decoding: communication channels and quantization. *Network: Computation in Neural Systems*, 12(4):441– 472, 2001.
- [12] Alexander G. Dimitrov and John P. Miller. Analyzing sensory systems with the information distortion function. In Russ B Altman, editor, *Pacific Symposium on Biocomputing 2001*. World Scientific Publishing Co., 2000.
- [13] Tomas Gedeon, Albert E. Parker, and Alexander G. Dimitrov. Information distortion and neural coding. *Canadian Applied Mathematics Quarterly*, 10(1):33–70, 2003.
- [14] Albert Parker, Tomas Gedeon, and Alexander Dimitrov. Annealing and the rate distortion problem. In S. Thrun S. Becker and K. Obermayer, editors, *Advances in Neural Information Processing Systems* 15, volume 15, pages 969–976. MIT Press, Cambridge, MA, 2003.
- [15] J. Nocedal and S. J. Wright. Numerical Optimization. Springer, New York, 2000.