The non-GRS properties for the twisted generalized Reed-Solomon code and its extended code

Canze Zhu, Qunying Liao

Abstract-In 2017, Beelen et al. firstly introduced twisted generalized Reed-Solomon (in short, TGRS) codes, and constructed a large subclass of MDS TGRS codes. Later, they proved that TGRS code is non-GRS when the code rate is less than one half. In this letter, basing on the dual code of the TGRS code or the extended TGRS code, by using the Schur product, we prove that almost all of TGRS codes and extended TGRS codes are non-GRS when the code rate more than one half.

Index Terms-Twisted generalized Reed-Solomon codes, Extended twisted generalized Reed-Solomon codes, Generalized **Reed-Solomon codes, Non-GRS properties**

I. INTRODUCTION

T HROUGHOUT this paper, let \mathbb{F}_q be the finite field with q elements, where q is the power of a prime. An [n, k, d]linear code C over \mathbb{F}_q is a k-dimensional subspace of \mathbb{F}_q^n with minimum (Hamming) distance d and length n. If the parameters reach the Singleton bound, namely, d = n - k + 1, then C is maximum distance separable (in short, MDS). Especially, generalized Reed-Solomon (in short, GRS) codes are a well known class of MDS codes, it is very important in coding theory and applications [12], [8], [18], [20], [25], [6], [17], [24], [26], [10], [1], [7], [13], [5]. Other known MDS codes have been constructed from n-arcs in projective geometry [9], circulant matrices [22], Hankel matrices [22], or twisted Reed-Solomon (in short, TGRS) codes [2], [3].

In 2017, inspired by the construction for twisted Gabidulin codes [23], Beelen et al. firstly introduced TGRS codes, which is a generalization for GRS codes, they also showed that TGRS codes could be well decoded. Different from GRS codes, they showed that a TGRS code is not necessarily MDS and presented a sufficient and necessary condition for a TGRS code to be MDS [2], [4], [3]. Especially, the authors showed that TGRS codes are not GRS when the code rate is less than one half [3]. Later, by TGRS codes, Lavauzelle et al. presented an efficient key-recovery attack used in the McEliece cryptosystem [19]. TGRS codes are also used to construct LCD MDS codes by their applications in cryptography [15], [21]. Recently, the authors gave the parity check matrix for the TGRS code and obtained some self-dual TGRS codes with small Singleton defect [14], [27]. More relative results about self-orthogonal TGRS codes can be seen in [28], [29], [11].

Qunying Liao is with the College of Mathematical Science, Sichuan Normal University, Chengdu Sichuan, 610066 (qunyingliao@sicnu.edu.cn).

The second author is supported by National Natural Science Foundation of China (Grant No. 12071321).

In this letter, we focus on the non-GRS properties for the TGRS code and its extended code. Let C be the TGRS code or extended TGRS (in short, ETGRS) code with code rate more than one half, by calculating the dimension and the minimun Hamming distance of the Schur square for \mathcal{C}^{\perp} under some assumptions, we show that \mathcal{C}^{\perp} is non-GRS, and then \mathcal{C} is non-GRS. The rest of this letter is organized as follows. In section 2, some basic notations and results about linear codes are given. In section 3, we show that almost all of the TGRS code and ETGRS code are non-GRS. In section 4, we conclude the whole paper.

1

II. PRELIMINARIES

In this section, we review some basic knowledge.

A. The dual codes, the Schur product and equivalence for linear codes

The nonation of the dual code is given in the following.

For $\mathbf{a} = (a_1, \ldots, a_n)$, $\mathbf{b} = (b_1, \ldots, b_n) \in \mathbb{F}_q^n$, the inner product is defined as $\langle \mathbf{a}, \mathbf{b} \rangle = \sum_{i=1}^{n} a_i b_i$. And then the dual code of C is defined as

$$\mathcal{C}^{\perp} = \{ \mathbf{c}^{'} \in \mathbb{F}_{q}^{n} \mid \langle \mathbf{c}^{'}, \mathbf{c} \rangle = 0 \text{ for any } \mathbf{c} \in \mathcal{C} \}.$$

The Schur product is defined as follows.

Definition 1 ([5]): For $\mathbf{x} = (x_1, \dots, x_n), \mathbf{y}$ = $(y_1,\ldots,y_n) \in \mathbb{F}_q^n$, the Schur product between ${f x}$ and ${f y}$ is defined as $\mathbf{x} \star \mathbf{y} := (x_1 y_1, \dots, x_n y_n)$. The Schur product of two q-ary codes C_1 and C_2 with length n is defined as

$$\mathcal{C}_1 \star \mathcal{C}_2 = \langle \mathbf{c}_1 \star \mathbf{c}_2 \mid \mathbf{c}_1 \in \mathcal{C}_1, \mathbf{c}_2 \in \mathcal{C}_2 \rangle.$$

Especially, for a code C, we call $C^2 := C \star C$ the Schur square of \mathcal{C} .

Remark 1: For any linear code $C = \langle v_1, \ldots, v_k \rangle$ with $v_i \in$ \mathbb{F}_{q}^{n} $(i = 1, \ldots, k)$, we have

$$\mathcal{C}^2 = \langle \boldsymbol{v}_i \star \boldsymbol{v}_j \ (i, j \in \{1, \dots, k\}) \rangle. \tag{1}$$

The definition of the equivalence for linear codes is given in the following.

Definition 2: ([16]) Let C_1 and C_2 be q-ary linear codes with length n. We say that C_1 and C_2 are equivalent if there is a permutation π in the permutation group with order n and $\boldsymbol{v} = (v_1, \dots, v_n) \in (\mathbb{F}_q^*)^n$ such that $\mathcal{C}_2 = \Phi_{\pi, \boldsymbol{v}}(\mathcal{C}_1)$, where

$$\Phi_{\pi,\boldsymbol{v}}: \mathbb{F}_q^n \to \mathbb{F}_q^n, \quad (c_1,\ldots,c_n) \mapsto (v_1c_{\pi(1)},\ldots,v_nc_{\pi(n)}).$$

Remark 2: By Definition 2, if C_1 and C_2 are equivalent, then • C_1^2 and C_2^2 are equivalent; • C_1^{\perp} and C_2^{\perp} are equivalent.

Canze Zhu is with the College of Mathematical Science, Sichuan Normal University, Chengdu Sichuan, 610066 (canzezhu@163.com).

B. The GRS, TGRS and ETGRS code

The definition of the GRS code is given in the following. Definition 3 ([16]): Let $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{F}_q^n$ with $\alpha_i \neq \alpha_j$ $(i \neq j)$ and $v = (v_1, \ldots, v_n) \in (\mathbb{F}_q^*)^n$. Then the GRS code is defined as

$$\mathcal{GRS}_{k,n}(\boldsymbol{\alpha}, \boldsymbol{v}) = \{ (v_1 f(\alpha_1), \dots, v_n f(\alpha_n)) | f(x) \in \mathbb{F}_q[x], \deg f(x) \le k-1 \}.$$

The dual code of the GRS code is given in the following. Lemma 1 ([18]): Let $\boldsymbol{u} = (u_1, \ldots, u_n)$ with $u_j = \prod_{i=1, i \neq j}^n (\alpha_j - \alpha_i)^{-1}$ $(j = 1, \ldots, n)$, then

$$\left(\mathcal{GRS}_{k,n}(oldsymbollpha,oldsymbol 1)
ight)^{oldsymbol -}=\mathcal{GRS}_{n-k,n}(oldsymbollpha,oldsymbol u)$$

By the definition of the GRS code, Lemma 1 and Remark 1, we have

Proposition 1: For $\frac{n}{2} \leq k < n$, let $\boldsymbol{u} = (u_1, \ldots, u_n)$ with $u_j = -\prod_{i=1, i\neq j}^n (\alpha_j - \alpha_i)$ $(j = 1, \ldots, n)$, then $\left(\mathcal{GRS}_{k,n}^{\perp}(\boldsymbol{\alpha}, \mathbf{1})\right)^2 = \mathcal{GRS}_{2(n-k)-2,n}(\boldsymbol{\alpha}, \boldsymbol{u}^2)$ with dimension 2(n-k)-1 and minimum Hamming distance 2k-n+2.

The definition of the twisted polynomials linear space $V_{k,t,h,\eta}$ is given in the following.

Definition 4 ([2]): Let $\eta \in \mathbb{F}_q^*$, and $t, h, k \in \mathbb{N}$ with $0 \le h < k \le q$. Then the set of (k, t, h, η) -twisted polynomials is defined as

$$\mathcal{V}_{k,t,h,\eta} = \{f(x) = \sum_{i=0}^{k-1} a_i x^i + \eta a_h x^{k-1+t} | a_i \in \mathbb{F}_q \ (i = 0, \dots, k-1) \},\$$

which is a k-dimensional \mathbb{F}_q -linear subspace. h and t are the hook and the twist, respectively.

From the twisted polynomials linear space $V_{k,t,h,\eta}$, the definitions of the TGRS code and the ETGRS code are given in the following, respectively.

Definition 5 ([2]): For any t, h, k, $n \in \mathbb{N}$ with $0 \le h \le k-1 < k-1+t < n \le q$, let $\eta \in \mathbb{F}_q^*$, $\boldsymbol{\alpha} = (\alpha_1, \ldots, \alpha_n) \in \mathbb{F}_q^n$ with $\alpha_i \ne \alpha_j$ $(i \ne j)$ and $\boldsymbol{v} = (v_1, \ldots, v_n) \in (\mathbb{F}_q^*)^n$. Then the TGRS code is defined as

$$\mathcal{C}_{t,h,k,n}(\boldsymbol{\alpha},\boldsymbol{v},\eta) = \{ (v_1 f(\alpha_1), \dots, v_n f(\alpha_n)) \mid f(x) \in \mathcal{V}_{k,t,h,\eta} \}.$$

For h > 0, the ETGRS code is defined as

$$\mathcal{C}_{t,h,k,n}(\boldsymbol{\alpha}, \boldsymbol{v}, \eta, \infty) = \{ (v_1 f(\alpha_1), \dots, v_n f(\alpha_n), f_h) \mid f(x) \in \mathcal{V}_{k,t,h,\eta} \}$$

where f_h is the coefficient of x^h in f(x). Especially, $C_{t,h,k,n}(\alpha, 1, \eta)$ and $C_{t,h,k,n}(\alpha, 1, \eta, \infty)$ are called the TRS code and the ETRS code, respectively.

Remark 3: • Note that $f(0) = f_0$, thus it is not necessary to define the ETGRS code when hook h = 0.

• In [2], $C_{t,h,k,n}(\alpha, v, \eta)$ and $C_{t,h,k,n}(\alpha, v, \eta, \infty)$ are called TGRS codes without difference. But it is not different from Definition 5.

Beelen et al. gave the non-GRS property for the TGRS code as the following.

Lemma 2 (Theorem 30 [3]): If $3 < k < \frac{n}{2}$, then $C_{t,h,k,n}(\boldsymbol{\alpha}, \boldsymbol{v}, \eta)$ is non-GRS.

Note that any punctured code of the GRS code is a GRS code and $C_{t,h,k,n}(\alpha, v, \eta)$ is a punctured code of $C_{t,h,k,n}(\alpha, v, \eta, \infty)$. By Lemma 2, we have

Lemma 3: If $3 < k < \frac{n+1}{2}$, then $C_{t,h,k,n}(\alpha, v, \eta, \infty)$ is non-GRS.

The authors gave the parity check matrix for $C_{t,h,k,n}(\alpha, v, \eta)$, which is useful to show that $C_{t,h,k,n}(\alpha, v, \eta)$ is non-GRS when $2k \ge n$.

Lemma 4 (Theorem 3.1 [27]): For any $m \in \mathbb{N}$, let

$$u_{j} = \prod_{i=1, i \neq j}^{n} (\alpha_{j} - \alpha_{i})^{-1} \ (i = 1, \dots, n), \quad L_{m} = \sum_{l=1}^{n} u_{l} \alpha_{l}^{n-1+m}$$

and $\tilde{L} = \sum_{m=1}^{k-h-1} L_{m} L_{k+t-h-1-m} - \eta^{-1} (1 + \eta L_{k+t-h-1}).$

Then $\mathcal{C}_{t,h,k,n}(\boldsymbol{\alpha},\boldsymbol{v},\eta,\infty)$ has the parity check matrix

$$\widetilde{H}_{n-k}(\boldsymbol{v}) = \left(\widetilde{\boldsymbol{\beta}}_1, \widetilde{\boldsymbol{\beta}}_2, \dots, \widetilde{\boldsymbol{\beta}}_n\right),$$

where

$$\tilde{\boldsymbol{\beta}}_{j} = \begin{pmatrix} \frac{u_{j}}{v_{j}} \\ \vdots \\ \frac{u_{j}}{v_{j}} \alpha_{j}^{n-(k+t+1)} \\ \frac{u_{j}}{v_{j}} (\tilde{L}\alpha_{j}^{n-k-t} + \alpha_{j}^{n-(h+1)} - \sum_{m=1}^{k-h-1} L_{m}\alpha_{j}^{n-(h+1)-m}) \\ \frac{u_{j}}{v_{j}} (\alpha_{j}^{n-(k+t-1)} - L_{1}\alpha_{j}^{n-k-t}) \\ \vdots \\ \frac{u_{j}}{v_{j}} (\alpha_{j}^{n-(k+1)} - L_{t-1}\alpha_{j}^{n-k-t}) \end{pmatrix}_{(n-k) \times n}$$

Remark 4: Under the assumption in Lemma 4, let

$$\mathcal{V}_{k,t,h,\eta}^{\perp} = \left\{ f(x) = \sum_{i=0}^{n-(k+t+1)} a_i x^i + \sum_{j=0}^{t-1} a_{n-k-t+j} h_{n-k-t+j}(x) \right|$$
$$a_i \in \mathbb{F}_q \ (i = 0, \dots, n-k-1) \right\}$$

with

$$h_{n-k-t+j}(x) = \begin{cases} x^{n-(h+1)} + \tilde{L}x^{n-k-t} - \sum_{m=1}^{k-h-1} L_m x^{n-(h+1)-m}, & \text{if } j = 0; \\ x^{n-k-t+j} - L_j x^{n-k-t}, & \text{if } 1 \le j \le t-1. \end{cases}$$

Then

$$\mathcal{C}_{t,h,k,n}^{\perp}(\boldsymbol{\alpha},\boldsymbol{1},\eta) = \Big\{ \big(u_1 g(\alpha_1), \dots, u_n g(\alpha_n) \big) | g(x) \in \mathcal{V}_{k,t,h,\eta}^{\perp} \Big\}.$$

III. NON-GRS PROPERTIES FOR THE TGRS CODE AND ETGRS CODE

In this section, by using the Schur product, we prove that almost all of TGRS codes and ETGRS codes are non-GRS when the code rate more than one half. and $k+t \leq n \leq q$. If $\frac{n}{2} \leq k \leq n-3$ and one of the $n-2k+h+2 \leq n-k-1$. Now by $2k \geq n+2$, one has following conditions (1.1)-(1.6) holds, then $C_{t,h,k,n}(\boldsymbol{\alpha}, \boldsymbol{v}, \eta)$ is non-GRS.

(1.1) n = k + t, $2k \ge n + 2$, and $3 \le h \le k - 3$; (1.2) n = k + t + 1, $2k \ge n + 2$, and $2 \le h \le k - 3$; (1.3) $n \ge k + t + 2$, $2k \ge n$, t = 1 and $2 \le h \le k - 2$; (1.4) $n \ge k + t + 2$, $2k \ge n + 1$, t = 2 and $1 \le h \le k - 3$; (1.5) $n \ge k + t + 2, 2k \ge n, t \ge 3$ and $h \in \{0, \dots, k - 1\}$ $1\} \setminus \{k-t\};$ (1.6) $n \ge k + t + 2$, $2k \ge n + 1$, $t \ge 3$ and h = k - t.

Proof. We show that dim $\left(\left(\mathcal{C}_{t,h,k,n}^{\perp}(\boldsymbol{\alpha},\mathbf{1},\eta)\right)^{2}\right) \geq 2(n-k)$ as follows.

By Remark 4, we have

 $\mathcal{C}_{t,h,k,n}^{\perp}(\boldsymbol{\alpha},\boldsymbol{1},\eta) = \left\{ \left(u_1 f(\alpha_1), \dots, u_n f(\alpha_n) \right) \middle| f(x) \in \mathcal{V}_{k,t,h,n}^{\perp} \right\},\$ where

$$\mathcal{V}_{k,t,h,\eta}^{\perp} = \left\{ f(x) = \sum_{i=0}^{n-(k+t+1)} a_i x^i + \sum_{j=0}^{t-1} a_{n-k-t+j} h_{n-k-t+j}(x) \right|$$
$$a_i \in \mathbb{F}_q \ (i = 0, \dots, n-k-1) \right\}$$

with

$$h_{n-k-t+i}(x) = \begin{cases} x^{n-(h+1)} + \tilde{L}x^{n-k-t} - \sum_{m=1}^{k-h-1} L_m x^{n-(h+1)-m}, \text{ if } i = 0; \\ x^{n-k-t+i} - L_i x^{n-k-t}, & \text{ if } 1 \le i \le t-1. \end{cases}$$
(2)

Thus

$$\left(\mathcal{C}_{t,h,k,n}^{\perp}(\boldsymbol{\alpha},\boldsymbol{1},\eta)\right)^{2} = \left\{ \left(g(\alpha_{1}),\ldots,g(\alpha_{n})\right) | g(x) \in (\mathcal{V}_{k,t,h,\eta}^{\perp})^{2} \right\}$$

with

$$(\mathcal{V}_{k,t,h,\eta}^{\perp})^2 = \langle g(x) = f_1(x)f_2(x) | f_1(x), f_2(x) \in \mathcal{V}_{k,t,h,\eta}^{\perp} \rangle.$$

Now we show that if one of the conditions (1.1)-(1.6) holds, then there exists $g_i(x) \in (\mathcal{V}_{k,t,h,\eta}^{\perp})^2$ $(i = 0, \dots, 2(n-k)-1)$ such that

$$\deg g_i(x) \le n-1 \text{ and } \deg g_i(x) \ne \deg g_j(x) \ (i \ne j).$$
 (3)

(1.1) For n = k + t, $2k \ge n + 2$, and $3 \le h \le k - 3$, we have

$$\begin{aligned} (\mathcal{V}_{k,t,h,\eta}^{\perp})^2 \\ = \langle h_l(x)h_s(x), \ h_0(x)h_i(x), \ h_0^2(x) \ (l,s,i \in \{1,\ldots,n-k-1\}) \rangle. \end{aligned}$$

By (2), one has

$$\deg (h_l(x)h_s(x)) = l + s \le 2(n-k) - 2 \le n-1$$

(l, s \equiv \{1, \dots, n-k-1\}\). (4)

Furthermore, for $h_0(x)h_i(x)$, we have the following two cases. If $3 \le h \le 2k - n + 1$, then $2(n-k) - 2 \le n - (h+1) \le n - 4$. Thus

$$2(n-k) - 1 \le \deg (h_0(x)h_i(x))$$

= $n - (h+1) + i$ (5
 $\le n - 1 \ (i \in \{1, 2, 3\}).$

Theorem 1: Let t, h, k, $n \in \mathbb{N}$ with $0 \le h \le k-1$ If $2k-n+2 \le h \le k-3$, then $1 \le n-2k+h < n-2k+h+1 <$

$$2(n-k) - 1 \le \deg \left(h_0(x)h_{n-2k+h-1+i}(x) \right)$$

=2(n-k) - 2 + i \le n - 1 (i \in \{1,2,3\}).
(6)

So far, by (4)-(6), (3) holds.

(1.2) For n = k + t + 1, $2k \ge n + 2$, and $2 \le h \le k - 3$, we have

$$\begin{aligned} (\mathcal{V}_{k,t,h,\eta}^{\perp})^2 = & \langle 1, h_0(x), h_m(x), h_l(x)h_s(x), h_0(x)h_i(x), h_0^2(x) \\ & (m,l,s,i \in \{2,\ldots,n-k-1\}) \rangle. \end{aligned}$$

By (2), one has

$$\deg h_m(x) = m \ (m = 2, 3),$$

$$4 \le \deg \left(h_l(x) h_s(x) \right) = l + s \le n - 1 \ (l, s \in \{2, \dots, n - k - 1\}).$$
(7)

Furthermore, for $h_0(x)h_i(x)$, we have the following two cases.

If $3 \le h \le 2k - n + 1$, then $2(n - k) - 2 \le n - (h + 1) \le n - 4$. Thus

$$2(n-k) - 1 \le \deg(h_0(x)h_i(x)) = n - (h+1) + i \le n - 1 \ (i \in \{2,3\}).$$
(8)

If $2k - n + 2 \le h \le k - 3$, then $2 \le n - 2k + h + 1 < k - 3$ $n - 2k + h + 2 \le n - k - 1$. Thus

$$2(n-k) \le \deg \left(h_0(x)h_{n-2k+h+i}(x) \right) = 2(n-k) - 1 + i \le n-1 \ (i \in \{1,2\}).$$
(9)

Now by $1 \in (\mathcal{V}_{k,t,h,n}^{\perp})^2$ and (7)-(9), (3) holds.

(1.3) For $n \ge k + t + 2$, $2k \ge n$, t = 1 and $2 \le h \le k - 2$, we have

$$(\mathcal{V}_{k,t,h,\eta}^{\perp})^2 = \langle x^{l+s}, x^i h_0(x), h_0^2(x) \quad (l,s,i \in \{0,1,\ldots,n-k-2\}) \rangle.$$

Obviously,

$$\deg (x^{l+s}) = l + s \le n - 1 \ (l, s \in \{0, 1, \dots, n - k - 2\}).$$
(10)

Furthermore, for $x^{j}h_{i}(x)$, we have the following two cases. If $2 \le h \le 2k - n + 2$, then $2(n-k) - 3 \le n - (h+1) \le n - 3$. Thus

$$2(n-k) - 3 \le \deg \left(x^i h_0(x)\right)$$

= $n - (h+1) + i \le n - 1 \ (i \in \{0, 1, 2\}).$
(11)

n - 2k + h - 1 < n - 2k + h < n - k - 2, thus

$$2(n-k) - 4 \le \deg \left(x^{n-2k+h-2+i}h_0(x)\right)$$

=2(n-k) - 3 + i \le n - 1 (i \in \{0,1,2\}).
(12)

Now by (10)-(12), (3) holds.

(1.4) For $n \geq k+t+2, \, 2k \geq n+1, \, t=2$ and $1 \leq h \leq k-3,$ we have

$$\begin{aligned} (\mathcal{V}_{k,t,h,\eta}^{\perp})^2 = & \langle x^{l+s}, \ x^i h_0(x), \ x^i h_1(x), \ h_0^2(x), \ h_0(x) h_1(x), \ h_1^2(x) \\ & (l,s,i \in \{0,1,\ldots,n-k-3\}) \rangle. \end{aligned}$$

By (2), one has

$$deg(x^{l+s}) = l + s \le 2n - 2k - 6 \le n - 1$$

(l, s \in {0, 1, ..., n - k - 3}),
$$deg(x^{n-k-5+m}h_1(x)) = 2(n-k) - 6 + m \ (m = 1, 2),$$

(13)
$$deg(h_1^2(x)) = 2(n-k) - 2.$$

Furthermore, for $x^l h_0(x)$, we have the following two cases. If $1 \le h \le 2k - n$, then $2(n-k) - 1 \le n - (h+1) \le n - 2$, thus

$$2(n-k) - 1 \le \deg \left(x^i h_0(x)\right)$$

= n - (h + 1) + i \le n - 1 (i \in \{0, 1\}). (14)

If $2k - n + 1 \le h \le k - 3$, then $h - 1 \le n - 2k + h - 2 < n - 2k + h \le n - k - 3$, thus

$$\deg \left(x^{n-2k+h-2}h_0(x)\right) = 2(n-k) - 3,$$

$$\deg \left(x^{n-2k+h}h_0(x)\right) = 2(n-k) - 1 \le n-1.$$
 (15)

Now by (13)-(15), (3) holds.

(1.5) For
$$n \ge k + t + 2$$
, $2k \ge n$ and $h \in \{0, \dots, k-1\} \setminus \{k-t\}$, we have
 $(\mathcal{V}_{k,t,h,\eta}^{\perp})^2$
 $= \left\langle x^{s_1+s_2}, x^s h_{n-k-t+l}(x), h_{n-k-t+l_1}(x) h_{n-k-t+l_2}(x) \right.$
 $\left(s, s_1, s_2 \in \{0, \dots, n-k-t-1\}, l, l_1, l_2 \in \{0, 1, \dots, t-1\}\right) \right\rangle$

and

$$\deg \left(x^{l+s}\right) = l+s \le n-1 \ (l,s \in \{0,1,\ldots,n-k-t-1\}), \\ \deg \left(x^{n-k-t-2}h_{n-k-t+l}(x)\right) = 2(n-k) - 2t - 2 + l \\ (l = 1,2,\ldots,t-1), \\ \deg \left(h_{n-k-3}(x)h_{n-k-t+l_1}(x)\right) = 2(n-k) - t - 3 + l_1 \\ (l_1 = 1,\ldots,t-1), \\ \deg \left(h_{n-k-2}(x)h_{n-k-1}(x)\right) = 2(n-k) - 3, \\ \deg \left(h_{n-k-1}^2(x)\right) = 2(n-k) - 2.$$

$$(16)$$

Furthermore, for $x^{l}h_{0}(x)$ and $h_{i}(x)h_{0}(x)$, we have the following three cases.

If $0 \le h \le 2k - n$, then $2(n - k) - 1 \le n - (h + 1) \le n - 1$, thus

$$\deg(h_0(x)) = n - (h+1) \le n - 1.$$
(17)

If $2k - n + 1 \le h \le k - t - 1$, then $h + 1 \le n - 2k + h \le n - k - t - 1$, thus

$$\deg\left(x^{n-2k+h}h_0(x)\right) = 2(n-k) - 1 \le n-1.$$
(18)

If $k - t + 1 \le h \le k - 1$, then $n - k - t + 1 \le n - 2k + h \le n - k - 1$, thus

$$\deg(h_{n-2k+h}(x)h_0(x)) = 2(n-k) - 1 \le n-1.$$
(19)

Now by (16)-(19), (3) holds.

(1.6) For $n \ge k + t + 2$, $2k \ge n + 1$ and h = k - t, in the similar proof as that for (1.5), we can get (3).

Let
$$\mathcal{V} = \left\{ \sum_{i=0}^{2(n-k)-1} a_i g_i(x) \mid a_i \in \mathbb{F}_q \right\}$$
, where $g_i(x)$ $(i = 0, \ldots, 2(n-k)-1)$ is given in (3), then $\dim(\mathcal{V}) = 2(n-k)$ and $\mathcal{V} \subseteq (\mathcal{V}_{k,t,h,\eta}^{\perp})^2$. Thus

$$\mathcal{C}_{\mathcal{V}} = \left\{ \left(g(\alpha_1), \dots, g(\alpha_n) \right) \mid g(x) \in \mathcal{V} \right\} \subseteq \left(\mathcal{C}_{t,h,k,n}^{\perp}(\boldsymbol{\alpha}, \mathbf{1}, \eta) \right)^2.$$

Now by $\deg(g(x)) \leq n-1$ ($\forall g(x) \in \mathcal{V}$), we have $(g(\alpha_1), \ldots, g(\alpha_n)) \neq 0$ ($\forall g(x) \in \mathcal{V} \setminus \{0\}$). It implies that $\dim \left(\left(\mathcal{C}_{t,h,k,n}^{\perp}(\boldsymbol{\alpha}, \boldsymbol{v}, \eta) \right)^2 \right) \geq \dim(\mathcal{C}_{\mathcal{V}}) = \dim \mathcal{V} = 2(n-k).$ By Propositon 1, for any [n,k] GRS code \mathcal{C} ,

By Proposition 1, for any $[n, \kappa]$ GKS code \mathcal{C} , $(\mathcal{C}^{\perp})^2 = 2(n-k) - 1$. Thus if one of conditions (1.1)-(1.6) holds, then $\mathcal{C}^{\perp}_{t,h,k,n}(\boldsymbol{\alpha}, \boldsymbol{v}, \eta)$ is non-GRS, and so $\mathcal{C}_{t,h,k,n}(\boldsymbol{\alpha}, \boldsymbol{v}, \eta)$ is non-GRS.

The following lemma is necessary to determine some codewords in $C_{t,h,k,n}^{\perp}(\boldsymbol{\alpha}, \boldsymbol{v}, \eta, \infty)$.

Lemma 5: For any $m \in \mathbb{N}$ and $A \subseteq \mathbb{F}_q$ with |A| > 2, let $L_A(m) = \sum_{\alpha \in A} \alpha^m \prod_{\beta \in \mathbb{F}_q \setminus A} (\alpha - \beta)$, then

$$L_A(m) = \begin{cases} 0, & \text{if } m \le |A| - 2; \\ -1, & \text{if } m = |A| - 1; \\ -\sum_{\alpha \in A} \alpha, & \text{if } m = |A|. \end{cases}$$
(20)

Proof. For any $l \in \mathbb{Z}^+$, it is well-known that

$$\sum_{\beta \in \mathbb{F}_q} \beta^l = \begin{cases} -1, & \text{if } q-1 \mid l; \\ 0, & \text{otherwise.} \end{cases}$$
(21)

Note that

$$\prod_{\beta \in \mathbb{F}_q \setminus A} (\alpha - \beta)$$

= $\alpha^{q - |A|} - \sum_{\beta \in \mathbb{F}_q \setminus A} \beta \alpha^{q - |A| - 1} + \dots + (-1)^{q - |A|} \prod_{\beta \in \mathbb{F}_q \setminus A} \beta$
= $\alpha^{q - |A|} + \sum_{\gamma \in A} \gamma \alpha^{q - |A| - 1} + \dots + (-1)^{q - |A|} \prod_{\beta \in \mathbb{F}_q \setminus A} \beta,$

thus we have

$$L_{A}(m) = \sum_{\alpha \in A} \alpha^{m} \prod_{\beta \in \mathbb{F}_{q} \setminus A} (\alpha - \beta)$$

=
$$\sum_{\alpha \in \mathbb{F}_{q}} \left(\alpha^{q-|A|+m} + \sum_{\gamma \in A} \gamma \alpha^{q-|A|-1+m} + \cdots + (-1)^{q-|A|+m} \prod_{\beta \in \mathbb{F}_{q} \setminus A} \beta \alpha^{m} \right).$$
 (22)

Now by (21)-(22), we obtain (20).

Theorem 2: For $t \ge 2$, $3 \le k \le n-2$ and $n \ge k+t+1$, $\mathcal{C}_{t,h,k,n}(\boldsymbol{\alpha}, \boldsymbol{v}, \eta, \infty)$ is non-GRS.

Proof. By Lemma 3, if $3 \le k < \frac{n+1}{2}$, then $C_{t,h,k,n}(\alpha, v, \eta, \infty)$ is non-GRS. Thus it is enought to prove that the theorem is true for $2k \ge n+1$.

In the following, we prove that $c_i \in C_{t,h,k,n}^{\perp}(\alpha, v, \eta, \infty)$ (i = 1, 2, 3), where

$$c_{1} = \left(\begin{array}{c} \frac{u_{1}}{v_{1}} \alpha_{1}^{n-k-t-1}, \dots, \frac{u_{n}}{v_{n}} \alpha_{n}^{n-k-t-1}, 0 \end{array} \right),$$

$$c_{2} = \left(\begin{array}{c} \frac{u_{1}}{v_{1}} \alpha_{1}^{n-k-t}, \dots, \frac{u_{n}}{v_{n}} \alpha_{n}^{n-k-t}, -\eta \end{array} \right),$$

$$c_{3} = \left(\begin{array}{c} \frac{u_{1}}{v_{1}} \alpha_{1}^{n-k-t+1}, \dots, \frac{u_{n}}{v_{n}} \alpha_{n}^{n-k-t+1}, -\eta \sum_{i=1}^{n} \alpha_{i} \end{array} \right).$$

Denote $A_{\alpha} = \{\alpha_1, \dots, \alpha_n\}$, for $s \in \{n - k - t - 1, n - k - t, n - k - t + 1\}$ and $l \in \{0, \dots, k - 1, k - 1 + t\}$, by the assumption $t \ge 2$ and Lemma 5, we have

$$\begin{split} &\sum_{i=0}^{n} u_i \alpha_i^s \alpha_i^l \\ &= -\sum_{\alpha \in A_{\alpha}} \alpha^{s+l} \prod_{\beta \in \mathbb{F}_q \setminus A_{\alpha}} (\alpha - \beta) \\ &= \begin{cases} 1, & \text{if } s = n - k - t \text{ and } l = k - 1 + t; \\ \sum_{i=1}^{n} \alpha_i, & \text{if } s = n - k - t + 1 \text{ and } l = k - 1 + t; \\ 0, & \text{otherwise.} \end{cases} \end{split}$$

By above equation, we can verify that

$$\boldsymbol{c}_i \in \mathcal{C}_{t,h,k,n}^{\perp}(\boldsymbol{\alpha}, \boldsymbol{v}, \eta, \infty) \ (i = 1, 2, 3)$$

directly. Thus

$$c = c_1 \star c_3 - c_2 \star c_2 = (0, 0, \dots, 0, \eta^2) \in (\mathcal{C}_{k,n}^{\perp}(\boldsymbol{\alpha}, \boldsymbol{v}, \eta, \infty))^2.$$

By Proposition 1, for an [n+1,k] GRS code C, $(C^{\perp})^2$ is with minimun Hamming distance

$$d = 2k - (n+1) + 2 \ge 2.$$

Thus $c \notin (\mathcal{C}^{\perp})^2$, and then $\mathcal{C}_{t,h,k,n}^{\perp}(\alpha, v, \eta, \infty)$ is non-GRS, it implies that $\mathcal{C}_{t,h,k,n}(\alpha, v, \eta, \infty)$ is non-GRS. \Box

Remark 5: • Note that any punctured code of the GRS code is a GRS code and $C_{t,h,k,n}(\alpha, v, \eta)$ is a punctured code of $C_{t,h,k,n}(\alpha, v, \eta, \infty)$. Thus, if one of conditions (1.1)-(1.6) in Theorem 1 holds, then $C_{t,h,k,n}(\alpha, v, \eta, \infty)$ is non-GRS.

• The assumption in Theorem 2 is weaker than that in Theorem 1 when $t \ge 2$ and $n \ge k + t + 1$.

IV. CONCLUSION

In this letter, by using the Schur product, we prove that almost all of TGRS codes and ETGRS codes are non-GRS when the code rate more than one half. However, if n, k, t, h do not satisfy one of the conditions in Theorem 1, we can not obtain the non-GRS property for the TGRS code by the method given in the proof for Theorem 1. The main reason is that we can not obtain dim $\left(\left(\mathcal{C}_{t,h,k,n}^{\perp}(\boldsymbol{\alpha},\mathbf{1},\eta)\right)^2\right) \geq 2k$ without the conditions in Theorem 1.

REFERENCES

- Bartz H., Sven P., Decoding of Interleaved Linearized Reed-Solomon Codes with Applications to Network Coding, IEEE ISIT, 160-165 (2021).
- [2] Beelen P., Puchinger S., Nielsen J., Twisted Reed-Solomon codes, IEEE ISIT, 336-340 (2017).
- [3] Beelen P., Puchinger S., Nielsen J., Twisted Reed-Solomon codes, IEEE Trans. Inf. Theory, doi: 10.1109/TIT.2022.3146254.
- [4] Beelen P., Bossert M., Puchinger S., Rosenkilde J., Structural properties of twisted Reed-Solomon codes with applications to code-based cryptography, IEEE ISIT, 946-950 (2018).
- [5] Couvreur A., Gaborit P., Gauthier-Umaña V., Otmani A., Tillich J. P., Distinguisher-based attacks on public-key cryptosystems using Reed–Solomon codes, Des. Codes Cryptogr., 73(2), 641-666 (2014).
- [6] Chen B., Liu H., New constructions of MDS codes with complementary duals, IEEE Trans. Inf. Theory, 64(8), 5776-5782 (2018).
- [7] Couvreur A., Lequesne M., On the security of subspace subcodes of Reed–Solomon codes for public key encryption, IEEE Trans. Inf. Theory, 68 (1), 632-648 (2021).
- [8] Fang X., Lebed K., Liu H., Luo J., New self-dual MDS codes over finite fields of odd characteristic, Des. Codes Cryptogr., 88(6), 1127-1138 (2020).
- [9] F. Macwilliams, N. Sloane, The Theory of Error-Correcting Codes, North-Holland, Amsterdam, (1977).
- [10] Gao, Y., Yue, Q., Huang, X., Zhang, J., Hulls of generalized Reed-Solomon codes via Goppa codes and their applications to quantum codes, IEEE Trans. Inf. Theory, 67 (10), 6619-6626 (2021).
- [11] Guo G., Li R., Liu Y., Song h., Duality of generalized twisted Reed-Solomon codes and Hermitian self-dual MDS or NMDS codes, https://doi.org/10.48550/arXiv.2202.11457.
- [12] Huang Z., Fang W., Fu W., New constructions of MDS self-dual and selforthogonal codes via GRS codes, https://arxiv.53yu.com/abs/2103.11665.
- [13] Holzbaur L., Polyanskaya R., Polyanskii N., Vorobyev I., Lifted Reed-Solomon Codes with Application to Batch Codes, IEEE ISIT, 634-639 (2020).
- [14] Huang, D., Yue Q., Niu Y., Li X., MDS or NMDS self-dual codes from twisted generalized Reed–Solomon codes, Des, Codes Cryptogr., 89(9), 2195-2209 (2021).
- [15] Huang D., Yue Q., Niu Y., MDS or NMDS LCD codes from twisted Reed-Solomon codes, https://seta-2020.etu.ru/assets/files/program/papers/paper-43.pdf.
- [16] Huffman W., Pless V., Fundamentals of Error Correcting Codes, Cambridge University Press, Cambridge (2003).
- [17] Jin L., Construction of MDS codes with complementary duals, IEEE Trans. Inf. Theory, 63(5), 2843-2847 (2017).
- [18] Jin L., Xing C., New self-dual MDS codes from generalized Reed-Solomon codes, IEEE Trans. Inf. Theory, 63(3), 1434-1438 (2017).
- [19] Lavauzelle J., Renner J., Crypt analysis of a system based on twisted Reed-Solomon codes, Des. Codes Cryptogr., 88(7), 1285-1300 (2020).
- [20] Lebed K., Liu H., Luo J., Construction of self-dual MDS codes over finite fields, Finite Fields Appl., 59, 199-207 (2019).
- [21] Liu H., Liu S., Construction of MDS twisted Reed-Solomon codes and LCD MDS codes, Des, Codes Cryptogr., 89(9), 2051-2065 (2021).
- [22] Roth R., Lempel A., On MDS Codes via Cauchy Matrices, IEEE Trans. Inf. Theory 35(6), 1314–1319 (1989)
- [23] Sheekey J., A new family of linear maximum rank distance codes, Adv. Math. Commun., 10(3), 475-488 (2016).
- [24] Shi X., Yue Q., Yang S., New LCD MDS codes constructed from generalized Reed-Solomon codes, J. Alg. Appl. 1950150 (2018).
- [25] Zhang A., Feng K., A unified approach to construct self-dual MDS codes via Reed-Solomon codes, IEEE Trans. Inf. Theory, 66(6), 3650-3656 (2020).
- [26] Wicker, S. B., Vijay K. B., Reed-Solomon codes and their applications, John Wiley, Sons (1999).
- [27] Zhu C., Liao Q., Self-dual twisted generalized Reed-Solomon codes, https://arxiv.org/abs/2111.11901.
- [28] Zhu C., Liao Q., Self-orthogonal generalized twisted Reed-Solomon codes, https://doi.org/10.48550/arXiv.2201.02758.
- [29] Zhang J., Zhou Z., Tang C., A class of twisted generalized Reed-Solomon codes, https://doi.org/10.48550/arXiv.2202.09011.