

# LIUVILLE TYPE THEOREMS FOR THE STATIONARY HALL-MHD EQUATIONS IN LOCAL MORREY SPACES

ZHOUYU LI<sup>1</sup> AND YIFAN SU<sup>2</sup>

**ABSTRACT.** This paper is concerned with the Liouville type theorems for the 3D stationary incompressible Hall-MHD equations. We establish that under some sufficient conditions in local Morrey spaces, solutions of the stationary Hall-MHD equations are identically zero. In particular, we also prove Liouville type results for the stationary incompressible MHD equations on  $\mathbb{R}^3$ . Our theorems extend and generalize the classical results for the stationary incompressible Navier-Stokes equations.

*Keywords:* Liouville type theorem; Hall-MHD equations; local Morrey spaces

*AMS Subject Classification (2000):* 35Q30, 76D03.

## 1. INTRODUCTION

We are concerned in this paper with the following stationary incompressible Hall-MHD equations on the whole space  $\mathbb{R}^3$ :

$$(1.1) \quad \begin{cases} -\Delta \mathbf{u} + \operatorname{div}(\mathbf{u} \otimes \mathbf{u}) + \nabla \Pi - \operatorname{curl} \mathbf{B} \times \mathbf{B} = 0, \\ -\Delta \mathbf{B} + \operatorname{curl}(\mathbf{B} \times \mathbf{u}) + \operatorname{curl}(\operatorname{curl} \mathbf{B} \times \mathbf{B}) = 0, \\ \operatorname{div} \mathbf{u} = \operatorname{div} \mathbf{B} = 0. \end{cases}$$

Here  $x = (x_1, x_2, x_3) \in \mathbb{R}^3$  is the space variable. The unknown functions  $\mathbf{u}$ ,  $\mathbf{B}$  and  $\Pi$  denote the velocity field, the magnetic field and the pressure, respectively. The Hall term  $\operatorname{curl}(\operatorname{curl} \mathbf{B} \times \mathbf{B})$  is derived from the Ohm's law and describes deviation from charge neutrality between the electrons and the ions. Hall term plays an important role in understanding the magnetic reconnection phenomena in plasmas, neutron stars, star formation, etc. For more physical explanations, see [1, 10, 12, 21].

The main purpose of this paper is to study Liouville type problem for the system (1.1), which is inspired by the development of Navier-Stokes equations. If  $\mathbf{B} = 0$ , the system (1.1) reduces to the stationary incompressible Navier-Stokes equations. A very challenging open question is when the Dirichlet integral

$$\int_{\mathbb{R}^3} |\nabla \mathbf{u}|^2 dx < +\infty,$$

whether there exist a nontrivial solution. This uniqueness problem, or equivalently the Liouville type problem originated from Leray's paper [16], and has been studied extensively recently. The most well-known one is due to Galdi (see [11], Remark X. 9.4, pp.729), which proved that if the smooth solution  $\mathbf{u}$  satisfies

$$\mathbf{u} \in L^{\frac{9}{2}}(\mathbb{R}^3),$$

<sup>1</sup> School of Sciences, Xi'an University of Technology, Xi'an 710054, China.

<sup>2</sup> School of Mathematics, Northwest University, Xi'an 710127, China.

E-mail address: zyli@163.com (Z. Li); mayifansu@163.com (Y. Su).

then  $\mathbf{u} = 0$ . The key to the proof is essentially based on the following local estimate (also called the Caccioppoli type inequality):

$$\begin{aligned} \int_{|x| \leq R} |\nabla \mathbf{u}|^2 dx &\leq C \|\mathbf{u}\|_{L^{\frac{9}{2}}(R \leq |x| \leq 2R)}^3 + CR^{-\frac{1}{3}} \|\mathbf{u}\|_{L^{\frac{9}{2}}(R \leq |x| \leq 2R)}^2 \\ &\quad + C \|\mathbf{u}\|_{L^{\frac{9}{2}}(R \leq |x| \leq 2R)} \|\Pi\|_{L^{\frac{9}{4}}(R \leq |x| \leq 2R)} \end{aligned}$$

for all  $R > 1$  and a constant  $C$  independent of  $R$ . Later, Chae and Wolf in [5] obtained a logarithmic improvement to Galdi's result under the assumption

$$\int_{\mathbb{R}^3} |\mathbf{u}|^{\frac{9}{2}} \left\{ \log \left( 2 + \frac{1}{|\mathbf{u}|} \right) \right\}^{-1} dx < \infty.$$

Seregin in [23] established that  $\mathbf{u} \in L^6(\mathbb{R}^3) \cap BMO^{-1}(\mathbb{R}^3)$  leads to the Liouville type theorem. The result of Galdi was also extended to the setting of Lorentz space  $L^{\frac{9}{2}, \infty}(\mathbb{R}^3)$  by Kozono et al. in [15] and a certain kind of local Lorentz type space by Seregin and Wang in [24]. Moreover, the authors in [6, 13, 14] proved the Liouville type results in a more general setting of Morrey spaces and some kind of local Morrey spaces.

However, the Liouville type problem for MHD type system is quite different due to the lack of maximum principle. For the case of MHD equations, Chae and Weng in [3] proved that the smooth solution  $(\mathbf{u}, \mathbf{B}) = 0$  under the condition

$$(\mathbf{u}, \mathbf{B}) \in L^3(\mathbb{R}^3) \quad \text{and} \quad (\nabla \mathbf{u}, \nabla \mathbf{B}) \in L^2(\mathbb{R}^3).$$

Schulz in [22] showed if  $(\mathbf{u}, \mathbf{B}) \in L^6(\mathbb{R}^3) \cap BMO^{-1}(\mathbb{R}^3)$ , then  $(\mathbf{u}, \mathbf{B}) = 0$ . For the case of Hall-MHD equations, in the famous paper [2], Chae, et al. proved that if  $(\mathbf{u}, \mathbf{B})$  is a smooth solution of (1.1) such that

$$\int_{\mathbb{R}^3} |\nabla \mathbf{u}|^2 + |\nabla \mathbf{B}|^2 dx < \infty \quad \text{and} \quad (\mathbf{u}, \mathbf{B}) \in L^\infty(\mathbb{R}^3) \cap L^{\frac{9}{2}}(\mathbb{R}^3),$$

then  $(\mathbf{u}, \mathbf{B}) = 0$ . Recently, the first author and Niu in [20] extend the result of Chae, et al. in [2] to the Lorentz space  $L^{\frac{9}{2}, \infty}(\mathbb{R}^3)$ . For more interesting Liouville type results about MHD and Hall-MHD equations, we refer to [4, 7, 18, 19, 25] and the references therein.

Here we consider the Liouville type theorems for the stationary incompressible Hall-MHD equations (1.1). Motivated by [6, 14, 15], we investigate some Liouville type theorems provided the velocity and magnetic field belongs to a fairly general functional setting, that is, some local Morrey spaces. It is a natural way to extend the space widely and improve the previous results.

An outline of this paper is as follows: we state our main results in Section 2. In Section 3, we collect some elementary facts, which will be used later. In Section 4, we obtain a characterization of the pressure term for the Hall-MHD equations. Section 5 is devoted to obtaining the Caccioppoli type inequalities, which are essential for achieving our results. In Section 6, the proofs of Theorems 1 and 2 are given.

## 2. STATEMENT OF THE RESULTS

Before stating the main theorems, we first recall the definitions of Morrey spaces and local Morrey spaces. For  $1 < p < r < +\infty$ , the homogeneous Morrey space  $\dot{M}^{p,r}(\mathbb{R}^3)$  is the set of all  $f \in L^p_{loc}(\mathbb{R}^3)$  such that

$$(2.1) \quad \|f\|_{\dot{M}^{p,r}} = \sup_{R>0, x_0 \in \mathbb{R}^3} R^{\frac{3}{r}} \left( R^{-3} \int_{B(x_0, R)} |f(x)|^p dx \right)^{\frac{1}{p}} < +\infty,$$

where  $B(x_0, R)$  is a ball of radius  $R$  centered at  $x_0$ . Here the term  $R^{\frac{3}{r}}$  characterizes the decaying of the averaged quantity  $\left( R^{-3} \int_{B(x_0, R)} |f(x)|^p dx \right)^{\frac{1}{p}}$  when  $R$  is large. The spaces

satisfy the following embedding relation

$$L^r(\mathbb{R}^3) \hookrightarrow L^{r,q}(\mathbb{R}^3) \hookrightarrow \dot{M}^{p,r}, \quad 1 < p < r \leq q \leq +\infty,$$

where the space  $L^{r,q}(\mathbb{R}^3)$  is the Lorentz space.

The local Morrey spaces describe the averaged decaying of functions in a more general setting. Let  $\gamma \geq 0$  and  $1 < p < +\infty$ , the local Morrey space  $M_\gamma^p(\mathbb{R}^3)$  is the set of functions  $f \in L_{loc}^p(\mathbb{R}^3)$  such that

$$(2.2) \quad \|f\|_{M_\gamma^p} = \sup_{R \geq 1} \left( R^{-\gamma} \int_{B(0,R)} |f(x)|^p dx \right)^{\frac{1}{p}} < +\infty.$$

It is easy to verify that  $M_\gamma^p(\mathbb{R}^3)$  is a Banach space under the norm  $\|\cdot\|_{M_\gamma^p}$ . Remark that the parameter  $\gamma$  describes the behavior of the quantity  $\left( \int_{B(0,R)} |f(x)|^p dx \right)^{\frac{1}{p}}$  when  $R$  is large. Furthermore, the spaces satisfy the continuous embedding

$$M_{\gamma_1}^p(\mathbb{R}^3) \hookrightarrow M_{\gamma_2}^p(\mathbb{R}^3), \quad \gamma_1 \leq \gamma_2.$$

We emphasize that for  $1 < p < r < +\infty$ , taking the parameter  $\gamma$  such that  $3(1 - p/r) < \gamma$ , then

$$\dot{M}^{p,r}(\mathbb{R}^3) = M_{3(1-p/r)}^p(\mathbb{R}^3) \hookrightarrow M_\gamma^p(\mathbb{R}^3).$$

Hence, in this sense local Morrey space  $M_\gamma^p(\mathbb{R}^3)$  can be regarded as a generalization of homogeneous Morrey space  $\dot{M}^{p,r}(\mathbb{R}^3)$ . For more references about the Morrey spaces and the local Morrey spaces, see [8, 17].

Finally, we define the space  $M_{\gamma,0}^p(\mathbb{R}^3)$  as the subspace of all functions  $f \in M_\gamma^p(\mathbb{R}^3)$  such that

$$\lim_{R \rightarrow +\infty} \left( R^{-\gamma} \int_{\frac{R}{2} \leq |x| \leq R} |f(x)|^p dx \right)^{\frac{1}{p}} = 0.$$

To simplify the notations, we denote

$$\eta = \eta(p, \gamma) = \frac{\gamma}{p} + \frac{2}{3} - \frac{3}{p},$$

$B_R = \{x \in \mathbb{R}^3 : |x| \leq R\}$  and  $C_R = \{x \in \mathbb{R}^3 : R \leq |x| \leq 2R\}$ . We state the first result of this paper as follows:

**Theorem 1.** Suppose that  $(\mathbf{u}, \mathbf{B})$  is a smooth solution to (1.1) with  $\nabla \mathbf{B} \in L^2(\mathbb{R}^3)$  and  $(\mathbf{u}, \mathbf{B}) \in M_\gamma^p(\mathbb{R}^3)$  for  $0 < \gamma < 3 \leq p < +\infty$ .

(i) If  $\eta(p, \gamma) < 0$ , then  $\mathbf{u} = \mathbf{B} = 0$  on  $\mathbb{R}^3$ .

(ii) If  $\eta(p, \gamma) = 0$ , and if moreover

$$(2.3) \quad \|\mathbf{u}\|_{M_\gamma^p(\mathbb{R}^3)}^3 + \|\mathbf{u}\|_{M_\gamma^p(\mathbb{R}^3)} \|\mathbf{B}\|_{M_\gamma^p(\mathbb{R}^3)}^2 \leq \delta \|\nabla \mathbf{u}, \nabla \mathbf{B}\|_{L^2(\mathbb{R}^3)}^2$$

for some  $0 < \delta < 1/C_0$ , then  $\mathbf{u} = \mathbf{B} = 0$  on  $\mathbb{R}^3$ .

(iii) If  $\eta(p, \gamma) > 0$ , and if moreover

$$(2.4) \quad \lim_{R \rightarrow +\infty} R^{3\eta(p,\gamma)} \left( (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} + (R^{-\gamma} \int_{C_R} |\mathbf{B}|^p dx)^{\frac{1}{p}} \right) = 0,$$

then  $\mathbf{u} = \mathbf{B} = 0$  on  $\mathbb{R}^3$ .

On the other hand, in the framework of a slightly smaller space  $M_{\gamma,0}^p(\mathbb{R}^3)$ , we do not need any additional assumption similar to (2.3). Specifically, this idea can be stated as follows.

**Theorem 2.** Suppose that  $(\mathbf{u}, \mathbf{B})$  is a smooth solution to (1.1) with  $\nabla \mathbf{B} \in L^2(\mathbb{R}^3)$ ,  $\mathbf{u} \in M_{\gamma,0}^p(\mathbb{R}^3)$  and  $\mathbf{B} \in M_\gamma^p(\mathbb{R}^3)$  for  $0 < \gamma < 3 \leq p < +\infty$ . If  $(p, \gamma)$  is such that  $\eta(p, \gamma) = 0$ , then  $\mathbf{u} = \mathbf{B} = 0$  on  $\mathbb{R}^3$ .

It should be noted that in Theorems 1 and 2 we always set the parameters  $0 < \gamma < 3 \leq p < +\infty$ . Since our methods deeply rely on the Cacciopoli type inequalities, for more details see Propositions 5.1 and 5.2 below, we need the condition  $3 \leq p < +\infty$ . In addition, in order to obtain the estimate for the pressure term, we need to use the Calderón-Zygmund inequality in the spaces  $M_\gamma^p(\mathbb{R}^3)$ , that is, point (i) of Lemma 3.2, so the condition  $0 < \gamma < 3$  is required.

The main interest of our results is based on the fact that we use a fairly general framework to solve the Liouville type problem. Indeed, we know that if  $\eta(p, \gamma) \leq 0$ , then for  $3 < r < \frac{9}{2}$  and  $3 \leq p < r$ , we have the embeddings

$$L^r(\mathbb{R}^3) \hookrightarrow L^{r,\infty}(\mathbb{R}^3) \hookrightarrow \dot{M}^{p,r}(\mathbb{R}^3) \hookrightarrow M_\delta^p(\mathbb{R}^3) \hookrightarrow M_{\gamma,0}^p(\mathbb{R}^3),$$

involving the Lebesgue, Lorentz, Morrey and local Morrey spaces. Here, the parameter  $\delta$  satisfies  $3(1 - p/r) < \delta < \gamma$ . In particular, for the values  $\gamma = 1$  and  $p = 3$ , and for  $r = \frac{9}{2}$  and  $\frac{9}{2} < q < +\infty$ , we also have the embeddings

$$L^{\frac{9}{2}}(\mathbb{R}^3) \hookrightarrow L^{\frac{9}{2},q}(\mathbb{R}^3) \hookrightarrow M_{1,0}^3(\mathbb{R}^3).$$

Notice that when  $\gamma = 1$  and  $p = 3$ , we have  $\eta(3, 1) = 0$ . For more references about the inclusion relationship of local Morrey spaces, we recommend [9, 14]. With the above facts in hand, we now give several remarks.

(i). Our results improve the result of Chae, et al. in [2], and the result of [20] is extended to the local Morrey spaces.

(ii). When removing the Hall term  $\text{curl}(\text{curl} \mathbf{B} \times \mathbf{B})$ , the system (1.1) reduces to the stationary incompressible MHD system. In this case, we do not need the assumption condition  $\nabla \mathbf{B} \in L^2(\mathbb{R}^3)$ , because of the fact that this condition is only used in dealing with the Hall term, for more details see (5.5) below. Hence we also obtain two new results for incompressible MHD system, which generalize the related result given in [19] to the local Morrey spaces.

(iii). When  $\mathbf{B} = 0$ , the system (1.1) reduces to the stationary incompressible Navier-Stokes system. Clearly, our results generalize the corresponding results of Galdi [11] and Jarrín [14].

### 3. PRELIMINARIES

For the convenience of readers, we will summarize some elementary facts about the local Morrey spaces and the weighted Lebesgue spaces.

Let us recall the definition of the weighted Lebesgue spaces. For  $\gamma \geq 0$  and  $1 < p < +\infty$ , we consider the weight  $\omega_\gamma(x) = \frac{1}{(1+|x|)^\gamma}$ , then the weighted Lebesgue spaces  $L_{\omega_\gamma}^p(\mathbb{R}^3)$  is defined as  $L_{\omega_\gamma}^p(\mathbb{R}^3) = L^p(\omega_\gamma dx)$ . It should be stressed that by using the Cauchy-Schwarz inequality we have the following continuous embedding:

$$(3.1) \quad L_{\omega_\gamma}^p(\mathbb{R}^3) \hookrightarrow L_{\omega_\gamma}^{\frac{p}{2}}(\mathbb{R}^3)$$

for  $\frac{9}{4} < \gamma < 3$ .

In order to study the characterization of the pressure term for the general Hall-MHD equations, we introduce a time-space version of the local Morrey spaces. Let  $\gamma > 0$  and  $1 < p < +\infty$ . We define the space  $M_\gamma^p L^p(0, T)$  as the set of functions  $f \in L_{loc}^p([0, T] \times \mathbb{R}^3)$  such that

$$\|f\|_{M_\gamma^p L^p(0, T)} = \sup_{R \geq 1} \left( R^{-\gamma} \int_0^T \int_{B(0, R)} |f(x, t)|^p dx dt \right)^{\frac{1}{p}} < +\infty.$$

Furthermore, we define the space  $M_{\gamma,0}^p L^p(0,T)$  as the set of functions  $f \in M_\gamma^p L^p(0,T)$  such that

$$\lim_{R \rightarrow +\infty} \left( R^{-\gamma} \int_0^T \int_{\frac{R}{2} \leq |x| \leq R} |f(x)|^p dx dt \right)^{\frac{1}{p}} = 0.$$

For details see Section 7 of [8]. Finally, we give two useful lemmas.

**Lemma 3.1** (Lemma 2.1 of [9]). *Assume  $0 \leq \gamma < \delta$ ,  $1 < p < +\infty$  and  $0 < T < +\infty$ . Then the following inclusion relationships are true:*

- (i)  $L_{\omega_\gamma}^p(\mathbb{R}^3) \hookrightarrow M_{\gamma,0}^p(\mathbb{R}^3) \hookrightarrow M_\gamma^p(\mathbb{R}^3) \hookrightarrow L_{\omega_\delta}^p(\mathbb{R}^3)$ ;
- (ii)  $L^p([0,T]; L_{\omega_\gamma}^p(\mathbb{R}^3)) \hookrightarrow M_{\gamma,0}^p L^p(0,T) \hookrightarrow M_\gamma^p L^p(0,T) \hookrightarrow L^p([0,T]; L_{\omega_\delta}^p(\mathbb{R}^3))$ .

**Lemma 3.2** (Lemma 2.2 and Corollary 2.1 of [9]). *Assume  $0 < \gamma < 3$  and  $1 < p < +\infty$ . Then there hold*

- (i)  $\|\nabla^2(-\Delta)^{-1}f\|_{M_\gamma^p(\mathbb{R}^3)} \leq C(p,\gamma)\|f\|_{M_\gamma^p(\mathbb{R}^3)}$ ;
- (ii)  $\|\mathcal{M}_f\|_{M_\gamma^p(\mathbb{R}^3)} \leq C(p,\gamma)\|f\|_{M_\gamma^p(\mathbb{R}^3)}$ , where  $\mathcal{M}$  is the Hardy-Littlewood maximal function operator;
- (iii) The points (i) and (ii) also hold for the weighted Lebesgue spaces  $L_{\omega_\gamma}^p(\mathbb{R}^3)$ .

#### 4. CHARACTERIZATION OF THE PRESSURE TERM

In this section, we aim at establishing a characterization of the pressure term for the following time-dependent Hall-MHD equations:

$$(4.1) \quad \begin{cases} \partial_t \mathbf{u} - \Delta \mathbf{u} + \operatorname{div}(\mathbf{u} \otimes \mathbf{u}) + \nabla \Pi - \operatorname{curl} \mathbf{B} \times \mathbf{B} = 0, & (t, x) \in \mathbb{R}^+ \times \mathbb{R}^3, \\ \partial_t \mathbf{B} - \Delta \mathbf{B} + \operatorname{curl}(\mathbf{B} \times \mathbf{u}) + \operatorname{curl}(\operatorname{curl} \mathbf{B} \times \mathbf{B}) = 0, & (t, x) \in \mathbb{R}^+ \times \mathbb{R}^3, \\ \operatorname{div} \mathbf{u} = \operatorname{div} \mathbf{B} = 0, & (t, x) \in \mathbb{R}^+ \times \mathbb{R}^3, \\ (u, B)|_{t=0} = (u_0, B_0), & x \in \mathbb{R}^3. \end{cases}$$

More precisely, we show that the pressure  $\Pi$  is always related to the velocity field  $\mathbf{u}$  and the magnetic field  $\mathbf{B}$  in the general setting of the time-space local Morrey spaces. It should be noted that the following proposition has an independent interest when seeking for very general frameworks in which the pressure is related to the other unknowns in the system (4.1).

**Proposition 4.1.** *Assume that  $(\mathbf{u}, \mathbf{B}, \Pi)$  is a solution of the system (4.1) with  $\Pi \in \mathcal{D}'([0,T] \times \mathbb{R}^3)$  and  $(\mathbf{u}, \mathbf{B}) \in M_\gamma^p L^p(0,T)$  for  $0 < \gamma < 3$ ,  $2 < p < +\infty$ . Then the term  $\nabla \Pi$  is given by the formula*

$$\nabla \Pi = \nabla \left( \sum_{i,j=1}^3 (-\Delta)^{-1} \partial_i \partial_j (u_i u_j - B_i B_j) - \frac{|\mathbf{B}|^2}{2} \right).$$

*Proof.* Motivated by the method in [14], we first define  $P$  given by the expression

$$(4.2) \quad P = \sum_{i,j=1}^3 (-\Delta)^{-1} \partial_i \partial_j (u_i u_j - B_i B_j) - \frac{|\mathbf{B}|^2}{2}.$$

Noting  $(\mathbf{u}, \mathbf{B}) \in M_\gamma^p L^p(0,T)$  for  $0 < \gamma < \frac{3}{2}$ , we apply point (ii) of Lemma 3.1 to get  $(\mathbf{u}, \mathbf{B}) \in L^p([0,T]; L_{\omega_\delta}^p(\mathbb{R}^3))$ , which gives  $(\mathbf{u} \otimes \mathbf{u}, \mathbf{B} \otimes \mathbf{B}, \frac{|\mathbf{B}|^2}{2}) \in L^{\frac{p}{2}}([0,T]; L_{\omega_\delta}^{\frac{p}{2}}(\mathbb{R}^3))$ . Thus, by point (iii) of Lemma 3.2, one has  $P \in L^{\frac{p}{2}}([0,T]; L_{\omega_\delta}^{\frac{p}{2}}(\mathbb{R}^3))$  for  $\frac{9}{4} < \delta < 3$ .

Next, we will show that  $\nabla \Pi - \nabla P = 0$ , where  $\Pi$  is the pressure term in (4.1). Let  $\varepsilon > 0$  (small enough) and  $\psi(x) \in C_0^\infty(\mathbb{R}^3)$ . Let  $\alpha(t) \in C_0^\infty(\mathbb{R})$  be a function such that for  $|t| > \varepsilon$ ,  $\alpha(t) = 0$ . We see that for all  $t \in (\varepsilon, T - \varepsilon)$ ,  $(\alpha\psi) * \nabla \Pi \in \mathcal{D}'((\varepsilon, T - \varepsilon) \times \mathbb{R}^3)$  and  $(\alpha\psi) * \nabla P \in \mathcal{D}'((\varepsilon, T - \varepsilon) \times \mathbb{R}^3)$ . Then, we define  $A_\varepsilon(t)$  given by

$$A_\varepsilon(t) = (\alpha\psi) * \nabla \Pi(\cdot, t) - (\alpha\psi) * \nabla P(\cdot, t).$$

We claim that  $A_\varepsilon(t) \in \mathcal{S}'(\mathbb{R}^3)$ . Indeed, we get by using the vector identity  $\operatorname{curl} \mathbf{B} \times \mathbf{B} = -\nabla \frac{|\mathbf{B}|^2}{2} + (\mathbf{B} \cdot \nabla) \mathbf{B}$  and the system (4.1) that

$$\nabla \Pi = -\partial_t \mathbf{u} + \Delta \mathbf{u} - \operatorname{div}(\mathbf{u} \otimes \mathbf{u}) - \operatorname{div}(\mathbf{B} \otimes \mathbf{B} - \frac{1}{2} |\mathbf{B}|^2 I).$$

Here  $I$  is the identical matrix. Thus,

$$(4.3) \quad \begin{aligned} A_\varepsilon(t) = & [(-(\partial_t \alpha) \psi + \alpha \Delta \psi) * \mathbf{u}](\cdot, t) - [(\alpha * \nabla \psi) * (\mathbf{u} \otimes \mathbf{u})](\cdot, t) \\ & - [(\alpha * \nabla \psi) * (\mathbf{B} \otimes \mathbf{B} - \frac{1}{2} |\mathbf{B}|^2 I)](\cdot, t) - [(\alpha \nabla \psi) * P](\cdot, t). \end{aligned}$$

We successively prove that each term on the right hand side of (4.3) belongs to the space  $L_{\omega_\delta}^{\frac{p}{2}}(\mathbb{R}^3)$  for  $\frac{9}{4} < \delta < 3$ . For the first term of the right in (4.3), since for  $\psi \in C_0^\infty(\mathbb{R}^3)$  and a test function  $f$ , we have the pointwise estimate  $|(\psi * f)(x)| \leq C(\varphi) \mathcal{M}_f(x)$ , where  $\mathcal{M}$  denotes the Hardy-Littlewood maximal function operator. Then, using point (iii) of Lemma 3.2 implies convolution with test functions is a bounded operator on  $L_{\omega_\delta}^p(\mathbb{R}^3)$ . Noting that  $\mathbf{u} \in L^p([0, T]; L_{\omega_\delta}^p(\mathbb{R}^3))$ , we get  $[(-(\partial_t \alpha) \psi + \alpha \Delta \psi) * \mathbf{u}](\cdot, t) \in L_{\omega_\delta}^p(\mathbb{R}^3)$ , which together with the embedding (3.1) gives  $[(-(\partial_t \alpha) \psi + \alpha \Delta \psi) * \mathbf{u}](\cdot, t) \in L_{\omega_\delta}^{\frac{p}{2}}(\mathbb{R}^3)$  for  $\frac{9}{4} < \delta < 3$ .

For the second and third terms of the right in (4.3), since  $\mathbf{u} \otimes \mathbf{u} \in L^{\frac{p}{2}}([0, T]; L_{\omega_\delta}^{\frac{p}{2}}(\mathbb{R}^3))$  and  $(\mathbf{B} \otimes \mathbf{B}, \frac{|\mathbf{B}|^2}{2}) \in L^{\frac{p}{2}}([0, T]; L_{\omega_\delta}^{\frac{p}{2}}(\mathbb{R}^3))$ , we have  $[(\alpha * \nabla \psi) * (\mathbf{u} \otimes \mathbf{u})](\cdot, t) \in L_{\omega_\delta}^{\frac{p}{2}}(\mathbb{R}^3)$  and  $[(\alpha * \nabla \psi) * (\mathbf{B} \otimes \mathbf{B} - \frac{1}{2} |\mathbf{B}|^2 I)](\cdot, t) \in L_{\omega_\delta}^{\frac{p}{2}}(\mathbb{R}^3)$ . Here we have used the fact convolution with test functions is a bounded operator on the space  $L_{\omega_\delta}^{\frac{p}{2}}(\mathbb{R}^3)$ .

For the fourth term of the right in (4.3), noting  $P \in L^{\frac{p}{2}}([0, T]; L_{\omega_\delta}^{\frac{p}{2}}(\mathbb{R}^3))$ , we have  $[(\alpha \nabla \psi) * P](\cdot, t) \in L^{\frac{p}{2}}([0, T]; L_{\omega_\delta}^{\frac{p}{2}}(\mathbb{R}^3))$ .

Hence, we have  $A_\varepsilon(t) \in L_{\omega_\delta}^{\frac{p}{2}}(\mathbb{R}^3)$ , and then  $A_\varepsilon(t) \in \mathcal{S}'(\mathbb{R}^3)$ . In addition, taking the divergence of the first equation in (4.1), we obtain from  $\operatorname{div} \partial_t \mathbf{u} = \operatorname{div} \Delta \mathbf{u} = 0$  that

$$\Delta \Pi = - \sum_{i,j=1}^3 \partial_i \partial_j (u_i u_j - B_i B_j) - \Delta \frac{|\mathbf{B}|^2}{2},$$

which implies  $\Delta(\Pi - P) = 0$ . Then we have  $\Delta A_\varepsilon(t) = 0$  and  $A_\varepsilon(t)$  is a polynomial from  $A_\varepsilon(t) \in \mathcal{S}'(\mathbb{R}^3)$ . Noticing  $A_\varepsilon(t) \in L_{\omega_\delta}^{\frac{p}{2}}(\mathbb{R}^3)$ , one has  $A_\varepsilon(t) = 0$ . Moreover, we use the approximation of the identity  $\frac{1}{\varepsilon^4} \alpha(\frac{t}{\varepsilon}) \psi(\frac{x}{\varepsilon})$  to write

$$\nabla(\Pi - P)(\cdot, t) = \lim_{\varepsilon \rightarrow 0} A_\varepsilon(t) = 0.$$

The proof is completed.  $\square$

## 5. CACCIOPPOLI TYPE INEQUALITIES

In this section, we derive the Caccioppoli type inequalities, which are the key to our proof. Let us first give the estimate in the framework of Lebesgue spaces.

**Proposition 5.1.** *Let  $(\mathbf{u}, \mathbf{B})$  be a smooth solution to (1.1). For  $3 \leq p < +\infty$ , if  $(\mathbf{u}, \mathbf{B}) \in L_{loc}^p(\mathbb{R}^3)$  and  $\nabla \mathbf{B} \in L^2(\mathbb{R}^3)$ , then we have*

$$(5.1) \quad \begin{aligned} & \int_{B_R} |\nabla \mathbf{u}|^2 dx + \int_{B_R} |\operatorname{curl} \mathbf{B}|^2 dx \\ & \leq CR^{1-\frac{6}{p}} \|\mathbf{u}\|_{L^p(C_R)}^2 + CR^{1-\frac{6}{p}} \|\mathbf{B}\|_{L^p(C_R)}^2 + CR^{2-\frac{9}{p}} \|\mathbf{u}\|_{L^p(C_R)}^3 + CR^{-\frac{1}{2}} \|\nabla \mathbf{B}\|_{L^2(C_R)}^3 \\ & \quad + CR^{2-\frac{9}{p}} \|\mathbf{u}\|_{L^p(C_R)} \|\mathbf{B}\|_{L^p(C_R)}^2 + CR^{2-\frac{9}{p}} \|\mathbf{u}\|_{L^p(C_R)} \|\Pi\|_{L^{\frac{p}{2}}(C_R)}^2, \end{aligned}$$

where the constant  $C$  is independent of  $R \geq 1$ .

*Proof.* First, we introduce the following cut-off function: let  $\varphi \in C_0^\infty(\mathbb{R}^3)$  be a nonincreasing radial function such that

$$\varphi(|x|) = \begin{cases} 1, & \text{if } |x| < 1, \\ [0, 1], & \text{if } 1 \leq |x| \leq 2, \\ 0, & \text{if } |x| > 2. \end{cases}$$

Define  $\varphi_R(x) = \varphi(\frac{|x|}{R})$  for each given  $R$ . Then the function  $\varphi_R(x)$  satisfies

$$\|\nabla^k \varphi_R\|_{L^\infty} \leq CR^{-k}$$

for  $k = 0, 1, 2$  with some positive constant  $C$ .

Taking the  $L^2$ - inner product of the  $u$  equation of (1.1) with  $\mathbf{u}\varphi_R^2$  and integrating by parts over  $\mathbb{R}^3$ , we obtain

$$(5.2) \quad \begin{aligned} \int_{\mathbb{R}^3} |\nabla \mathbf{u}|^2 \varphi_R^2 dx &= -2 \int_{\mathbb{R}^3} \varphi_R \nabla \mathbf{u} : (\mathbf{u} \otimes \nabla \varphi_R) dx - \int_{\mathbb{R}^3} (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot \mathbf{u} \varphi_R^2 dx \\ &\quad - \int_{\mathbb{R}^3} \nabla \Pi \cdot \mathbf{u} \varphi_R^2 dx + \int_{\mathbb{R}^3} \text{curl } \mathbf{B} \times \mathbf{B} \cdot \mathbf{u} \varphi_R^2 dx. \end{aligned}$$

Similarly, taking the  $L^2$ - inner product of the  $B$  equation of (1.1) with  $\mathbf{B}\varphi_R^2$  and integrating by parts over  $\mathbb{R}^3$ , we have

$$(5.3) \quad \begin{aligned} \int_{\mathbb{R}^3} |\text{curl } \mathbf{B}|^2 \varphi_R^2 dx &= -2 \int_{\mathbb{R}^3} \varphi_R \text{curl } \mathbf{B} \cdot (\nabla \varphi_R \times \mathbf{B}) dx - \int_{\mathbb{R}^3} (\mathbf{B} \times \mathbf{u}) \cdot (\varphi_R^2 \text{curl } \mathbf{B}) dx \\ &\quad - 2 \int_{\mathbb{R}^3} \varphi_R (\mathbf{B} \times \mathbf{u}) \cdot (\nabla \varphi_R \times \mathbf{B}) dx - 2 \int_{\mathbb{R}^3} \varphi_R (\text{curl } \mathbf{B} \times \mathbf{B}) \cdot (\nabla \varphi_R \times \mathbf{B}) dx, \end{aligned}$$

where we have used the fact  $\Delta \mathbf{B} = \nabla \text{div } \mathbf{B} - \text{curl}(\text{curl } \mathbf{B})$ .

Combining (5.2) and (5.3), it gives

$$(5.4) \quad \begin{aligned} &\int_{\mathbb{R}^3} |\nabla \mathbf{u}|^2 \varphi_R^2 dx + \int_{\mathbb{R}^3} |\text{curl } \mathbf{B}|^2 \varphi_R^2 dx \\ &= -2 \int_{\mathbb{R}^3} \varphi_R \nabla \mathbf{u} : (\mathbf{u} \otimes \nabla \varphi_R) dx - \int_{\mathbb{R}^3} (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot \mathbf{u} \varphi_R^2 dx \\ &\quad - 2 \int_{\mathbb{R}^3} \varphi_R \text{curl } \mathbf{B} \cdot (\nabla \varphi_R \times \mathbf{B}) dx - 2 \int_{\mathbb{R}^3} \varphi_R (\mathbf{B} \times \mathbf{u}) \cdot (\nabla \varphi_R \times \mathbf{B}) dx \\ &\quad - 2 \int_{\mathbb{R}^3} \varphi_R (\text{curl } \mathbf{B} \times \mathbf{B}) \cdot (\nabla \varphi_R \times \mathbf{B}) dx - \int_{\mathbb{R}^3} \nabla \Pi \cdot \mathbf{u} \varphi_R^2 dx \\ &= \sum_{i=1}^6 I_i. \end{aligned}$$

In what follows, we estimate each term on the right-hand side of (5.4) separately. For  $I_1$ , it can be deduced from the integration by parts that

$$\begin{aligned} I_1 &= \int_{\mathbb{R}^3} \mathbf{u} \cdot \text{div}(\mathbf{u} \otimes \nabla(\varphi_R^2)) dx \\ &= \int_{\mathbb{R}^3} \mathbf{u} \cdot (\nabla \mathbf{u} \cdot \nabla(\varphi_R^2) + \mathbf{u} \Delta(\varphi_R^2)) dx \\ &= \frac{1}{2} \int_{\mathbb{R}^3} |\mathbf{u}|^2 \Delta(\varphi_R^2) dx \\ &= \int_{\mathbb{R}^3} |\mathbf{u}|^2 (\varphi_R \Delta \varphi_R + |\nabla \varphi_R|^2) dx. \end{aligned}$$

Applying the Hölder inequality implies

$$\begin{aligned} |I_1| &\leq \int_{C_R} |\mathbf{u}|^2 (|\varphi_R \Delta \varphi_R| + |\nabla \varphi_R|^2) dx \\ &\leq CR^{-2} \|\mathbf{u}\|_{L^{\frac{p}{2}}(C_R)}^2 \|1\|_{L^{\frac{p}{p-2}}(C_R)} \\ &\leq CR^{1-\frac{6}{p}} \|\mathbf{u}\|_{L^p(C_R)}^2. \end{aligned}$$

For  $I_2$ , using integration by parts, we obtain from  $\operatorname{div} \mathbf{u} = 0$  that

$$\begin{aligned} I_2 &= - \int_{\mathbb{R}^3} (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot (\mathbf{u} \varphi_R^2) dx = -\frac{1}{2} \int_{\mathbb{R}^3} \mathbf{u} \cdot \nabla |\mathbf{u}|^2 \varphi_R^2 dx \\ &= \frac{1}{2} \int_{\mathbb{R}^3} |\mathbf{u}|^2 \operatorname{div}(\mathbf{u} \varphi_R^2) dx = \int_{\mathbb{R}^3} |\mathbf{u}|^2 \mathbf{u} \cdot (\varphi_R \nabla \varphi_R) dx. \end{aligned}$$

Thus,

$$\begin{aligned} |I_2| &\leq \int_{C_R} |\mathbf{u}|^2 |\mathbf{u}| |\varphi_R \nabla \varphi_R| dx \\ &\leq CR^{-1} \|\mathbf{u}\|_{L^{\frac{p}{2}}(C_R)}^2 \|\mathbf{u}\|_{L^p(C_R)} \|1\|_{L^{\frac{p}{p-3}}(C_R)} \\ &\leq CR^{2-\frac{9}{p}} \|\mathbf{u}\|_{L^p(C_R)}^3. \end{aligned}$$

For  $I_3$ , using Young's inequality, we get

$$\begin{aligned} |I_3| &\leq \int_{C_R} |\varphi_R \operatorname{curl} \mathbf{B}| |\mathbf{B}| |\nabla \varphi_R| dx \\ &\leq CR^{-1} \|\varphi_R \operatorname{curl} \mathbf{B}\|_{L^2(C_R)} \|\mathbf{B}\|_{L^p(C_R)} \|1\|_{L^{\frac{2p}{p-2}}(C_R)} \\ &\leq CR^{\frac{1}{2}-\frac{3}{p}} \|\varphi_R \operatorname{curl} \mathbf{B}\|_{L^2(C_R)} \|\mathbf{B}\|_{L^p(C_R)} \\ &\leq \frac{1}{2} \|\varphi_R \operatorname{curl} \mathbf{B}\|_{L^2(C_R)}^2 + CR^{1-\frac{6}{p}} \|\mathbf{B}\|_{L^p(C_R)}^2. \end{aligned}$$

Again, applying the Hölder inequality gives

$$\begin{aligned} |I_4| &\leq CR^{-1} \int_{C_R} |\mathbf{B}|^2 |\mathbf{u}| dx \\ &\leq CR^{-1} \|\mathbf{B}\|_{L^{\frac{p}{2}}(C_R)}^2 \|\mathbf{u}\|_{L^p(C_R)} \|1\|_{L^{\frac{p}{p-3}}(C_R)} \\ &\leq CR^{2-\frac{9}{p}} \|\mathbf{B}\|_{L^p(C_R)}^2 \|\mathbf{u}\|_{L^p(C_R)}. \end{aligned}$$

For  $I_5$ , using the embedding  $\dot{H}^1(\mathbb{R}^3) \hookrightarrow L^6(\mathbb{R}^3)$  implies

$$\begin{aligned} |I_5| &\leq CR^{-1} \int_{C_R} |\varphi_R \operatorname{curl} \mathbf{B}| |\mathbf{B}|^2 dx \\ (5.5) \quad &\leq CR^{-1} \|\operatorname{curl} \mathbf{B}\|_{L^2(C_R)} \|\mathbf{B}\|_{L^3(C_R)}^2 \|1\|_{L^6(C_R)} \\ &\leq CR^{-\frac{1}{2}} \|\nabla \mathbf{B}\|_{L^2(C_R)} \|\mathbf{B}\|_{L^6(C_R)}^2 \\ &\leq CR^{-\frac{1}{2}} \|\nabla \mathbf{B}\|_{L^2}^3. \end{aligned}$$

Finally, for  $I_6$ , using integration by parts and the fact  $\operatorname{div} \mathbf{u} = 0$ , we get

$$\begin{aligned} I_6 &= \int_{\mathbb{R}^3} \Pi \operatorname{div}(\mathbf{u} \varphi_R^2) dx \\ &= \int_{\mathbb{R}^3} \Pi \operatorname{div} \mathbf{u} \varphi_R^2 dx + 2 \int_{\mathbb{R}^3} \varphi_R \Pi \mathbf{u} \cdot \nabla \varphi_R dx \\ &= 2 \int_{\mathbb{R}^3} \varphi_R \Pi \mathbf{u} \cdot \nabla \varphi_R dx. \end{aligned}$$



Then,

$$\begin{aligned}
 |I_6| &\leq CR^{-1} \int_{C_R} |\Pi| |\mathbf{u}| dx \\
 &\leq CR^{-1} \|\Pi\|_{L^{\frac{p}{2}}(C_R)} \|\mathbf{u}\|_{L^p(C_R)} \|1\|_{L^{\frac{p}{p-3}}(C_R)} \\
 &\leq CR^{2-\frac{9}{p}} \|\Pi\|_{L^{\frac{p}{2}}(C_R)} \|\mathbf{u}\|_{L^p(C_R)}.
 \end{aligned}$$

Consequently, substituting the estimates  $I_i$  for  $i = 1, 2, \dots, 6$  into (5.4), we deduce that

$$\begin{aligned}
 &\int_{B_R} |\nabla \mathbf{u}|^2 dx + \frac{1}{2} \int_{B_R} |\operatorname{curl} \mathbf{B}|^2 dx \\
 &\leq CR^{1-\frac{6}{p}} \|\mathbf{u}\|_{L^p(C_R)}^2 + CR^{1-\frac{6}{p}} \|\mathbf{B}\|_{L^p(C_R)}^2 + CR^{2-\frac{9}{p}} \|\mathbf{u}\|_{L^p(C_R)}^3 + CR^{2-\frac{9}{p}} \|\mathbf{u}\|_{L^p(C_R)} \|\mathbf{B}\|_{L^p(C_R)}^2 \\
 &\quad + CR^{-\frac{1}{2}} \|\nabla \mathbf{B}\|_{L^2(C_R)}^3 + CR^{2-\frac{9}{p}} \|\mathbf{u}\|_{L^p(C_R)} \|\Pi\|_{L^{\frac{p}{2}}(C_R)}
 \end{aligned}$$

for all  $R \geq 1$  and the constant  $C$  independent of  $R$ . This completes the proof of Proposition 5.1.  $\square$

Next, in the framework of local Morrey spaces, the Caccioppoli type estimate is stated as follows:

**Proposition 5.2.** *Let  $(\mathbf{u}, \mathbf{B})$  be a smooth solution to (1.1). For  $0 < \gamma < 3 \leq p < +\infty$ , if  $(\mathbf{u}, \mathbf{B}) \in M_\gamma^p(\mathbb{R}^3)$  and  $\nabla \mathbf{B} \in L^2(\mathbb{R}^3)$ , then we have*

$$\begin{aligned}
 &\int_{B_R} |\nabla \mathbf{u}|^2 dx + \int_{B_R} |\operatorname{curl} \mathbf{B}|^2 dx \\
 (5.6) \quad &\leq CR^{3\eta} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{3}{p}} + CR^{-\frac{1}{3}} R^{2\eta} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{2}{p}} + CR^{-\frac{1}{2}} \|\nabla \mathbf{B}\|_{L^2(C_R)}^3 \\
 &\quad + CR^{3\eta} (R^{-\gamma} \int_{C_R} |\mathbf{B}|^p dx)^{\frac{2}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} + CR^{-\frac{1}{3}} R^{2\eta} (R^{-\gamma} \int_{C_R} |\mathbf{B}|^p dx)^{\frac{2}{p}} \\
 &\quad + CR^{3\eta} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} (\|\mathbf{u}\|_{M_\gamma^p(\mathbb{R}^3)}^2 + \|\mathbf{B}\|_{M_\gamma^p(\mathbb{R}^3)}^2),
 \end{aligned}$$

where the constant  $C$  is independent of  $R \geq 1$ .

*Proof.* Using Proposition 4.1, we know that  $\nabla \Pi = \nabla P$ , where  $P$  is given in (4.2). Thus, we consider the system (1.1) with the term  $\nabla P$  instead of the term  $\nabla \Pi$ . Since  $(\mathbf{u}, \mathbf{B}) \in M_\gamma^p(\mathbb{R}^3)$ , one has  $(\mathbf{u}, \mathbf{B}) \in L_{loc}^p(\mathbb{R}^3)$ . Then we obtain from Proposition 5.1 that for  $3 \leq p < +\infty$ ,

$$\begin{aligned}
 &\int_{B_R} |\nabla \mathbf{u}|^2 dx + \int_{B_R} |\operatorname{curl} \mathbf{B}|^2 dx \\
 (5.7) \quad &\leq CR^{1-\frac{6}{p}} \|\mathbf{u}\|_{L^p(C_R)}^2 + CR^{1-\frac{6}{p}} \|\mathbf{B}\|_{L^p(C_R)}^2 + CR^{2-\frac{9}{p}} \|\mathbf{u}\|_{L^p(C_R)}^3 + CR^{-\frac{1}{2}} \|\nabla \mathbf{B}\|_{L^2(C_R)}^3 \\
 &\quad + CR^{2-\frac{9}{p}} \|\mathbf{u}\|_{L^p(C_R)} \|\mathbf{B}\|_{L^p(C_R)}^2 + CR^{2-\frac{9}{p}} \|\mathbf{u}\|_{L^p(C_R)} \|P\|_{L^{\frac{p}{2}}(C_R)}^2.
 \end{aligned}$$

A direct computation gives

$$\begin{aligned}
 R^{2-\frac{9}{p}} \|\mathbf{u}\|_{L^p(C_R)}^3 &= R^{-\frac{2}{p}\gamma} R^{\frac{2}{p}\gamma+2-\frac{9}{p}} \left( \int_{C_R} |\mathbf{u}|^p dx \right)^{\frac{2}{p}} \left( \int_{C_R} |\mathbf{u}|^p dx \right)^{\frac{1}{p}} \\
 &= R^{\frac{3}{p}\gamma+2-\frac{9}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{2}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}}.
 \end{aligned}$$

Similarly, we obtain

$$\begin{aligned}
& R^{1-\frac{6}{p}} \|\mathbf{u}\|_{L^p(C_R)}^2 + R^{1-\frac{6}{p}} \|\mathbf{B}\|_{L^p(C_R)}^2 \\
&= R^{-\frac{1}{3}-\frac{2}{p}\gamma} R^{\frac{2}{p}\gamma+\frac{4}{3}-\frac{6}{p}} \left( \int_{C_R} |\mathbf{u}|^p dx \right)^{\frac{2}{p}} + R^{-\frac{1}{3}-\frac{2}{p}\gamma} R^{\frac{2}{p}\gamma+\frac{4}{3}-\frac{6}{p}} \left( \int_{C_R} |\mathbf{B}|^p dx \right)^{\frac{2}{p}} \\
&= R^{-\frac{1}{3}} R^{\frac{2}{p}\gamma+\frac{4}{3}-\frac{6}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{2}{p}} + R^{-\frac{1}{3}} R^{\frac{2}{p}\gamma+\frac{4}{3}-\frac{6}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{B}|^p dx)^{\frac{2}{p}}
\end{aligned}$$

and

$$\begin{aligned}
R^{2-\frac{9}{p}} \|\mathbf{B}\|_{L^p(C_R)}^2 \|\mathbf{u}\|_{L^p(C_R)} &= R^{\frac{2}{p}\gamma+2-\frac{9}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{B}|^p dx)^{\frac{2}{p}} \left( \int_{C_R} |\mathbf{u}|^p dx \right)^{\frac{1}{p}} \\
&= R^{\frac{3}{p}\gamma+2-\frac{9}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{B}|^p dx)^{\frac{2}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}}.
\end{aligned}$$

Finally, we are in a position to estimate  $R^{2-\frac{9}{p}} \|\mathbf{u}\|_{L^p(C_R)} \|P\|_{L^{\frac{p}{2}}(C_R)}^2$ . Noticing that the term  $P$  is defined through  $\mathbf{u}$  and  $\mathbf{B}$  in (4.2), we deduce from point (i) of Lemma 3.2 that for all  $0 < \gamma < 3 \leq p < +\infty$ ,

$$\|P\|_{M_\gamma^{\frac{p}{2}}(\mathbb{R}^3)} \leq C(p, \gamma) (\|\mathbf{u}\|_{M_\gamma^p(\mathbb{R}^3)}^2 + \|\mathbf{B}\|_{M_\gamma^p(\mathbb{R}^3)}^2).$$

Moreover, recalling the definition of  $\|\cdot\|_{M_\gamma^p(\mathbb{R}^3)}$ , see (2.2), we get

$$\begin{aligned}
R^{2-\frac{9}{p}} \|P\|_{L^{\frac{p}{2}}(C_R)} \|\mathbf{u}\|_{L^p(C_R)} &= R^{\frac{3}{p}\gamma+2-\frac{9}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} (R^{-\gamma} \int_{C_R} |P|^{\frac{p}{2}} dx)^{\frac{2}{p}} \\
&\leq R^{\frac{3}{p}\gamma+2-\frac{9}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} \|P\|_{M_\gamma^{\frac{p}{2}}(\mathbb{R}^3)} \\
&\leq C R^{\frac{3}{p}\gamma+2-\frac{9}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} (\|\mathbf{u}\|_{M_\gamma^p(\mathbb{R}^3)}^2 + \|\mathbf{B}\|_{M_\gamma^p(\mathbb{R}^3)}^2).
\end{aligned}$$

Inserting all the above estimates into (5.7), we complete the proof of Proposition 5.2.  $\square$

## 6. THE PROOFS OF THEOREMS

With the help of Proposition 5.2, we can now complete the proofs of Theorems 1 and 2. For simplicity, we denote

$$\begin{aligned}
D_1 &:= R^{3\eta} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{3}{p}} + R^{-\frac{1}{3}} R^{2\eta} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{2}{p}} + R^{-\frac{1}{2}} \|\nabla \mathbf{B}\|_{L^2(C_R)}^3 \\
&\quad + R^{3\eta} (R^{-\gamma} \int_{C_R} |\mathbf{B}|^p dx)^{\frac{2}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} + R^{-\frac{1}{3}} R^{2\eta} (R^{-\gamma} \int_{C_R} |\mathbf{B}|^p dx)^{\frac{2}{p}} \\
&\quad + R^{3\eta} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} (\|\mathbf{u}\|_{M_\gamma^p(\mathbb{R}^3)}^2 + \|\mathbf{B}\|_{M_\gamma^p(\mathbb{R}^3)}^2).
\end{aligned}$$

Then the inequality (5.6) writes the following

$$(6.1) \quad \int_{B_R} |\nabla \mathbf{u}|^2 dx + \int_{B_R} |\operatorname{curl} \mathbf{B}|^2 dx \leq C D_1.$$

*Proof of Theorem 1.* Assume that  $\nabla \mathbf{B} \in L^2(\mathbb{R}^3)$  and  $(\mathbf{u}, \mathbf{B}) \in M_\gamma^p(\mathbb{R}^3)$  with  $0 < \gamma < 3 \leq p < +\infty$ . We divide our proof into three cases.

(i). **The case when  $\eta(p, \gamma) < 0$ .** According to the definition of  $\|\cdot\|_{M_\gamma^p(\mathbb{R}^3)}$ , we obtain

$$(6.2) \quad D_1 \leq R^{3\eta} \left( \|\mathbf{u}\|_{M_\gamma^p}^2 + \|\mathbf{B}\|_{M_\gamma^p}^2 \right) \|\mathbf{u}\|_{M_\gamma^p} + R^{-\frac{1}{3}} R^{2\eta} \left( \|\mathbf{u}\|_{M_\gamma^p}^2 + \|\mathbf{B}\|_{M_\gamma^p}^2 \right) + R^{-\frac{1}{2}} \|\nabla \mathbf{B}\|_{L^2}^2.$$

Passing  $R \rightarrow +\infty$  in (6.1), we see

$$\lim_{R \rightarrow +\infty} D_1 = 0.$$

Thus, it leads to

$$\lim_{R \rightarrow +\infty} \left( \int_{B_R} |\nabla \mathbf{u}|^2 dx + \int_{B_R} |\operatorname{curl} \mathbf{B}|^2 dx \right) = 0,$$

which together with Levi's monotone convergence theorem yields

$$\int_{\mathbb{R}^3} |\nabla \mathbf{u}|^2 dx + \int_{\mathbb{R}^3} |\operatorname{curl} \mathbf{B}|^2 dx = 0.$$

Using the fact that

$$\|\nabla \mathbf{B}\|_{L^2(\mathbb{R}^3)} \leq C \left( \|\operatorname{div} \mathbf{B}\|_{L^2(\mathbb{R}^3)} + \|\operatorname{curl} \mathbf{B}\|_{L^2(\mathbb{R}^3)} \right),$$

we conclude that  $\|(\nabla \mathbf{u}, \nabla \mathbf{B})\|_{L^2(\mathbb{R}^3)} = 0$  and  $\mathbf{u} = \mathbf{B} = 0$  on  $\mathbb{R}^3$  from  $(\mathbf{u}, \mathbf{B}) \in M_\gamma^p(\mathbb{R}^3)$ .

**(ii). The case when  $\eta(p, \gamma) = 0$ .** Passing  $R \rightarrow +\infty$  in (6.1) and by virtue of (6.2), we get

$$\lim_{R \rightarrow +\infty} D_1 \leq \|\mathbf{u}\|_{M_\gamma^p}^3 + \|\mathbf{B}\|_{M_\gamma^p}^2 \|\mathbf{u}\|_{M_\gamma^p},$$

and thus

$$\int_{\mathbb{R}^3} |\nabla \mathbf{u}|^2 dx + \int_{\mathbb{R}^3} |\operatorname{curl} \mathbf{B}|^2 dx \leq C_0 \left( \|\mathbf{u}\|_{M_\gamma^p}^3 + \|\mathbf{B}\|_{M_\gamma^p}^2 \|\mathbf{u}\|_{M_\gamma^p} \right).$$

It follows from the assumption (2.3) that

$$\|(\nabla \mathbf{u}, \nabla \mathbf{B})\|_{L^2(\mathbb{R}^3)}^2 \leq C_0 \left( \|\mathbf{u}\|_{M_\gamma^p}^3 + \|\mathbf{B}\|_{M_\gamma^p}^2 \|\mathbf{u}\|_{M_\gamma^p} \right) \leq C_0 \delta \|(\nabla \mathbf{u}, \nabla \mathbf{B})\|_{L^2(\mathbb{R}^3)}^2.$$

Since  $0 < C_0 \delta < 1$ , we conclude that  $\|(\nabla \mathbf{u}, \nabla \mathbf{B})\|_{L^2(\mathbb{R}^3)} = 0$  and  $\mathbf{u} = \mathbf{B} = 0$  on  $\mathbb{R}^3$ .

**(iii). The case when  $\eta(p, \gamma) > 0$ .** Here, we see  $2\eta(p, \gamma) \leq 6\eta(p, \gamma)$  and  $3\eta(p, \gamma) \leq 6\eta(p, \gamma)$ . By direct calculation, we have

$$R^{3\eta} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{2}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} \leq \left[ R^{3\eta} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} \right]^2 (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}}$$

and

$$R^{-\frac{1}{3}} R^{2\eta} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{2}{p}} \leq R^{-\frac{1}{3}} \left[ R^{3\eta} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} \right]^2.$$

Similarly, we get

$$R^{3\eta} (R^{-\gamma} \int_{C_R} |\mathbf{B}|^p dx)^{\frac{2}{p}} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} \leq \left[ R^{3\eta} (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} \right] (R^{-\gamma} \int_{C_R} |\mathbf{B}|^p dx)^{\frac{2}{p}}$$

and

$$R^{-\frac{1}{3}} R^{2\eta} (R^{-\gamma} \int_{C_R} |\mathbf{B}|^p dx)^{\frac{2}{p}} \leq R^{-\frac{1}{3}} \left[ R^{3\eta} (R^{-\gamma} \int_{C_R} |\mathbf{B}|^p dx)^{\frac{1}{p}} \right]^2.$$

Noting the condition (2.4), passing  $R \rightarrow +\infty$  in (6.1), we deduce

$$\int_{\mathbb{R}^3} |\nabla \mathbf{u}|^2 dx + \int_{\mathbb{R}^3} |\operatorname{curl} \mathbf{B}|^2 dx = 0.$$

Hence, we conclude that  $\mathbf{u} = \mathbf{B} = 0$  on  $\mathbb{R}^3$ .  $\square$

*Proof of Theorem 2.* Assume that  $\nabla \mathbf{B} \in L^2(\mathbb{R}^3)$ ,  $\mathbf{u} \in M_{\gamma,0}^p(\mathbb{R}^3)$  and  $\mathbf{B} \in M_\gamma^p(\mathbb{R}^3)$  for  $0 < \gamma < 3 \leq p < +\infty$ . By the definition of  $\|\cdot\|_{M_\gamma^p(\mathbb{R}^3)}$ , we have

$$(6.3) \quad \begin{aligned} D_1 &\leq R^{3\eta} (\|\mathbf{u}\|_{M_\gamma^p}^2 + \|\mathbf{B}\|_{M_\gamma^p}^2) (R^{-\gamma} \int_{C_R} |\mathbf{u}|^p dx)^{\frac{1}{p}} + R^{-\frac{1}{2}} \|\nabla \mathbf{B}\|_{L^2}^3 \\ &\quad + R^{-\frac{1}{3}} R^{2\eta} (\|\mathbf{u}\|_{M_\gamma^p}^2 + \|\mathbf{B}\|_{M_\gamma^p}^2). \end{aligned}$$

Considering the assumption  $\eta(p, \gamma) = 0$ , letting  $R \rightarrow +\infty$  in (6.1), and applying Levi's monotone convergence theorem, we have

$$\int_{\mathbb{R}^3} |\nabla \mathbf{u}|^2 dx + \int_{\mathbb{R}^3} |\operatorname{curl} \mathbf{B}|^2 dx = 0.$$

Hence, we get the desired identity  $\mathbf{u} = \mathbf{B} = 0$  on  $\mathbb{R}^3$ .  $\square$

**Acknowledgments.** The work is partially supported by the National Natural Science Foundation of China under the grants 11571279 and 11601423.

#### REFERENCES

- [1] M. Acheritogaray, P. Degond, A. Frouvelle, J. G. Liu, *Kinet. Relat. Models*, **4** (2011) no. 4, 901-918.
- [2] D. Chae, P. Degond, J. G. Liu, Well-posedness for Hall-magnetohydrodynamics, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **31** (2014) no. 3, 555-565.
- [3] D. Chae, S. Weng, Liouville type theorems for the steady axially symmetric Navier-Stokes and magneto-hydrodynamic equations. *Discrete Contin. Dyn. Syst.*, **36** (2016) no. 10, 5267-5285.
- [4] D. Chae, J. Wolf, On Liouville type theorems for the stationary MHD and Hall-MHD systems, *J. Differential Equations*, **295** (2021) 233-248.
- [5] D. Chae, J. Wolf, On Liouville type theorems for the steady Navier-Stokes equations in  $\mathbb{R}^3$ , *J. Differential Equations*, **261** (2016) no. 10, 5541-5560.
- [6] D. Chamorro, O. Jarrín, P. G. Lemarié-Rieusset, Some Liouville theorems for stationary Navier-Stokes equations in Lebesgue and Morrey spaces, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **38** (2021) no. 3, 689-710.
- [7] H. Fan, M. Wang, The Liouville type theorem for the stationary magnetohydrodynamic equations, *J. Math. Phys.*, **62** (2021) no. 5, Paper No. 031503, 12 pp.
- [8] P. G. Fernández-Dalgo, P. G. Lemarié-Rieusset. Characterisation of the pressure term in the incompressible Navier-Stokes equations on the whole space, *Discrete Contin. Dyn. Syst. Ser. S*, **14** (2021) no. 8, 2917-2931.
- [9] P. G. Fernández-Dalgo, O. Jarrín, Weak-strong uniqueness in weighted  $L^2$  spaces and weak suitable solutions in local Morrey spaces for the MHD equations, *J. Differential Equations*, **271** (2021) 864-915.
- [10] T. G. Forbes, Magnetic reconnection in solar flares, *Geophys. Astrophys. Fluid Dyn.*, **62** (1991) 15-36.
- [11] G. P. Galdi, An Introduction to the Mathematical Theory of the Navier-Stokes Equations, Steady-State Problems, second edition. Springer Monographs in Mathematics. Berlin-Heidelberg-New York: Springer-Verlag, 2011.
- [12] H. Homann, R. Grauer, Bifurcation analysis of magnetic reconnection in Hall-MHD systems, *Phys. D*, **208** (2005) no. 1-2, 59-72.
- [13] O. Jarrín, A remark on the Liouville problem for stationary Navier-Stokes equations in Lorentz and Morrey spaces, *J. Math. Anal. Appl.*, **486** (2020) no. 1, 123871, 16 pp.
- [14] O. Jarrín, Liouville theorems for a stationary and non-stationary coupled system of liquid crystal flows, arXiv.org: 2006.14502.
- [15] H. Kozono, Y. Terasawa, Y. Wakasugi, A remark on Liouville-type theorems for the stationary Navier-Stokes equations in three space dimensions. *J. Funct. Anal.*, **272** (2017) no. 2, 804-818.
- [16] J. Leray, Étude de diverses équations intégrales non linéaire et de quelques problèmes que pose l'hydrodynamique, *J. Math. Pures Appl.*, **12** (1933) 1-82.
- [17] P. G. Lemarié-Rieusset, The Navier-Stokes Problem in the 21st Century, CRC Press, Boca Raton, FL, (2016).
- [18] Z. Li, X. Pan, Liouville theorem of the 3D stationary MHD system: for D solutions converging to non-zero constant vectors, *NoDEA Nonlinear Differential Equations Appl.*, **28** (2021) no. 2, Paper No. 12, 14 pp.
- [19] Z. Li, P. Liu, P. Niu, Remarks on Liouville type theorems for the 3D stationary MHD equations, *Bull. Korean Math. Soc.*, **57** (2020) no. 5, 1151-1164.
- [20] Z. Li, P. Niu, Liouville type theorems for the 3D stationary Hall-MHD equations, *ZAMM Z. Angew. Math. Mech.*, **100** (2020) no. 5, e201900200, 9 pp.
- [21] M. J. Lighthill, Studies on magnetohydrodynamic waves and other anisotropic wave motions, *Philos. Trans. Roy. Soc. London Ser. A*, **252** (1960) 397-430.
- [22] S. Schulz, Liouville type theorem for the stationary equations of magneto-hydrodynamics, *Acta Math. Sci. Ser. B (Engl. Ed.)*, **39** (2019) no. 2, 491-497.
- [23] G. Seregin, Liouville type theorem for stationary Navier-Stokes equations, *Nonlinearity*, **29** (2016) no. 8, 2191-2195.
- [24] G. Seregin, W. Wang, Sufficient conditions on Liouville type theorems for the 3D steady Navier-Stokes equations, *Algebra i Analiz*, **31** (2019) no. 2, 269-278; reprinted in *St. Petersburg Math. J.*, **31** (2020) no. 2, 387-393.

- [25] B. Yuan, Y. Xiao, Liouville-type theorems for the 3D stationary Navier-Stokes, MHD and Hall-MHD equations, *J. Math. Anal. Appl.*, **491** (2020) no. 2, 124343, 10 pp.