

Multi species asymmetric simple exclusion process with impurity activated flips

Amit Kumar Chatterjee^{1*}, Hisao Hayakawa^{1,2†},

1 Yukawa Institute for Theoretical Physics, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan

2 Center for Gravitational Physics and Quantum Information, Yukawa Institute for Theoretical Physics, Kyoto University Sakyo-ku, Kyoto 606-8502, Japan

* ak.chatterjee@yukawa.kyoto-u.ac.jp, † hisao@yukawa.kyoto-u.ac.jp

May 17, 2022

Abstract

We obtain an exact matrix product steady state for a class of multi species asymmetric simple exclusion process with impurities, under periodic boundary condition. Alongside the usual hopping dynamics, an additional flip dynamics is activated only in the presence of impurities. Although the microscopic dynamics renders the system to be non-ergodic, exact analytical results for observables are obtained in steady states for a specific class of initial configurations. Interesting physical features including negative differential mobility and transition of correlations from negative to positive with changing vacancy density, have been observed. We discuss plausible connections of this exactly solvable model with multi lane asymmetric simple exclusion processes as well as enzymatic chemical reactions.

Contents

1	Introduction	2
2	Model: μ-TASEP-IAF	5
2.1	Microscopic dynamics	5
2.2	Steady state: matrix product ansatz	6
3	Partition function for special initial configuration	9
3.1	$\mu = 2$	10
3.2	$\mu = 3$	11
3.3	general μ	11
4	Observables: comparisons of analytical results with Monte Carlo simulations	12
4.1	Species densities	12
4.2	Drift current	15

4.3	Two-point correlations	17
4.4	Flip current	18
4.5	Non-ergodicity: dependence on initial configuration	19
5	Partially asymmetric generalization: μ-ASEP-IAF	21
5.1	Matrix algebra, auxiliaries and matrix representations	21
5.2	Partition function for special initial configuration	23
5.3	Species densities, drift current and flip current	23
5.4	Negative differential mobility	24
6	Summary and future directions	26
A	Connection between μ-TASEP-IAF and multi lane TASEP	28
B	A variation of μ-TASEP-IAF, connection to multi-lane traffic flow	30
C	Connection between μ-TASEP-IAF and enzymatic chemical reactions	31
	References	33

1 Introduction

Non-equilibrium stochastic processes are ubiquitous in nature, with wide range of applicability in physics [1–3], chemistry [1], biology [4] and interdisciplinary areas [5–7]. In fact, even in one dimension, several models of non-equilibrium statistical mechanics exhibit surprisingly rich physical phenomena including phase transitions [8] along with the important feature of analytical tractability [9]. The asymmetric simple exclusion process (ASEP) [2, 10–12] is broadly regarded as a paradigmatic model for non-equilibrium transport processes as diverse as traffic and pedestrian flow [13], mRNA translation by ribosomes [14] and motor protein transport through single filaments [15, 16] etc.

Apart from its extensive success in modeling numerous real-world phenomena, ASEP and its variations have been instrumental in understanding the mathematical structures and physical characteristics of generic non-equilibrium steady states and dynamics [2, 12, 17–20]. In particular, the exact steady states of the totally asymmetric simple exclusion process (TASEP) [12, 21] and the general ASEP [22] with open boundary conditions, have been obtained using matrix product ansatz. The matrix product ansatz has been extremely effective in deriving the non-equilibrium steady states of several generalizations of ASEP including multi-species processes [23], see Ref. [24] for a detailed review. In fact, the stationary state for the multi species TASEP has been solved remarkably by a different method of multiline queuing process [25], which is explored further in terms of combinatorial R in crystal base theory [26]. Several two point and three point correlations have been studied analytically in the multi species TASEP [27] and inhomogeneous multi species TASEP with species dependent rates have been analyzed [28, 29]. The multi species ASEP has also been investigated with integrable open boundary conditions [30] and matrix product solutions are found [31]. Due to the connection of

TASEP to integrable spin chains [32], the algebraic Bethe Ansatz has been applied to study the dynamics of TASEP [33] and ASEP [34] with open boundaries. Interestingly, ASEP belongs to Kardar-Parisi-Zhang universality class [35] with dynamic exponent $\frac{3}{2}$ [36,37]. With the aid of Monte Carlo simulations and several improved versions of mean-field theories, TASEP has also been generalized to non-conserved dynamics [38], two-lane [39–41] and multi-lane [42–45] models relating to traffic flow and complex networks [46].

It is quite natural to expect the presence of multiple species of particles with variety of microscopic dynamics in a system in general. Often due to the distinction between the dynamics of different species, some species are referred to as *impurities* and give rise to fascinating physical and mathematical structures. For example, the presence of a single impurity which hops with a different rate and allows overtaking of ordinary particles in the TASEP on a periodic lattice, leads to a matrix product steady state with six distinct phases including the creation of a shock in one of the phases [47]. This impurity model has been generalized by considering bidirectional asymmetric hopping of the ordinary particles [48] which allows for finite dimensional matrix representations in certain regions of the parameter space, in comparison to the infinite dimensional representation in Ref. [47]. A phase transition arising from the motion of the single impurity in the direction opposite to the ordinary particles has also been observed [49]. The long time limit behavior of the TASEP with a single impurity has been solved using the Bethe Ansatz [50] and the diffusion constant of the impurity has been calculated from both the Bethe Ansatz [50] and the matrix product ansatz [51]. Remarkably, a disordered ASEP with species dependent hop rates, has been shown to exhibit Bose-Einstein condensation [52]. A variation of the ASEP with ordinary particles and many impurities has been considered in Ref. [53], where the impurities are not allowed to hop to the vacant neighbors but they can exchange positions with ordinary particles. Interestingly, such an impurity model in [53] possesses a different scaling exponent $\frac{5}{2}$ in comparison to the usual KPZ exponent $\frac{3}{2}$ for the ASEP without impurities [36,37]. Other than the disorders or impurities associated with the particles themselves, there are many exciting studies with position dependent or site-wise disorders for ASEP [54–60].

In this article, we study a class of the multi-species ($I = 1, 2, \dots, \mu$) ASEP in the presence of impurities, under periodic boundary conditions. In addition to the usual hopping of particles to vacant sites in ASEP, we consider flips of different species among each other (e.g. species I transforming to species J and vice versa). Importantly, these flip processes are initiated *only in the presence of a special type of particles* (as nearest neighbors) that we denote *impurities*. These impurities activate the flip processes. Therefore, we name this non-equilibrium stochastic process to be *multi species asymmetric simple exclusion process with impurity activated flips* (μ -ASEP-IAF). Note that the total number of impurities, along with that of the vacancies, remain conserved in the μ -ASEP-IAF. Specifically, we emphasize that the flip processes between two non-conserved species (I, J) do not occur through the interaction with any non-conserved species K ($= 1, 2, \dots, \mu$) at the nearest neighbors. Thus, the microscopic dynamics considered here is different from previously studied models like TASEP with internal degrees of freedom [61] and multi-species reaction-diffusion processes [62,63]. Notably, the distinction in the microscopic dynamics also makes μ -ASEP-IAF *non-ergodic* in nature in contrast to the ergodic models [61–63]. We should mention that the non-ergodicity of exactly solvable models is related to undecidability of thermalization in integrable models [64].

The motivations for studying the μ -ASEP-IAF are as follows. (i) We aim to obtain an exact non-equilibrium steady state of the μ -ASEP-IAF under periodic boundary condition, so that it

would be an important addition to the category of exact solvable models in disordered systems.

- (ii) The μ -ASEP-IAF being non-ergodic, it would be interesting to derive exact analytical expressions for partition function and observables for suitable choice of initial configurations and compare the corresponding steady state results with that of a random initial configuration.
- (iii) The μ -TASEP-IAF can be mapped to multi-lane TASEP which is a basic model for multi-lane traffic flow. Different species of particles in μ -ASEP-IAF play the roles of particles in different lanes of multi-lane TASEP and the impurities in μ -ASEP-IAF act as bridges between lanes that allow particles to exchange lanes in multi lane TASEP. See Appendix A for details.
- (iv) Considering the conserved impurities as enzymes (E) and different non-conserved species as substrates (S) and products (P), the flip process of μ -ASEP-IAF can be thought as an enzymatic chemical reaction like $S + E \rightarrow P + E$, which is a crude approximation of the Michaelis-Menten reaction scheme $S + E \rightleftharpoons SE \rightarrow P + E$ [65–67]. See Appendix C for details. Notably, both the mappings in (iii) and (iv) would not be possible if the impurities could also flip to other species.

Below we briefly summarize our main results.

- (i) We find that the steady states of μ -TASEP-IAF (totally asymmetric hopping) and μ -ASEP-IAF (bidirectional hopping) under periodic boundary conditions, can be obtained exactly as matrix product states, where distinct matrices represent different components (each species, impurity, vacancy) of the system. We provide explicit finite dimensional matrix representations for the totally asymmetric case, whereas the matrices for the general asymmetric case are found to be infinite dimensional .
- (ii) For specific choice of initial configuration, we could analytically calculate the partition function in the sector of allowed configurations in the steady state and consequently the observables of interests (average densities of non-conserved species, currents and spatial correlations). The analytical results are in agreement with Monte Carlo simulations. For a fixed set of input parameters, we show considerable quantitative deviations between steady state observable values for different initial configurations, establishing the initial configuration dependence or non-ergodicity of the dynamics.
- (iii) Two-point nearest neighbor correlations exhibit interesting non-trivial behaviors. Particularly, with the variation of the vacancy density, we observe characteristics like certain correlations changing signs i.e. varying from negative to positive with some intermediate zero correlation point, and, non-monotonic behavior with both local maximum and local minimum.
- (iv) We find *negative differential mobility* in μ -ASEP-IAF. For special choices of hopping rates, both the drift current and flip current decrease with increasing bias giving rise to negative differential mobility.

The article is organized as follows. In Sec. 2 we describe the μ -TASEP-IAF in details and show that the steady state can be achieved using matrix product ansatz. The analytical calculation of partition function starting from a suitably chosen initial configuration is presented in Sec. 3. We discuss the behaviors of observables like species densities, drift current, flip current and spatial correlations from both analytical calculation and Monte Carlo simulations with variation of input parameters in Sec. 4. The μ -TASEP-IAF is generalized to μ -ASEP-IAF with bidirectional motions of the species in Sec. 5, where we show the corresponding matrix product states and discuss the negative differential mobility of particles. In Sec. 6, we summarize the results with future directions. We discuss the mapping between μ -TASEP-IAF and multi lane TASEP in Appendix A. A variation of μ -TASEP-IAF that comes up with better features in connection to traffic in multi-lane problems is discussed in Appendix B. The connections between μ -TASEP-IAF and enzymatic chemical reactions are briefly presented in




Figure 1: The figure illustrates all possible microscopic dynamical processes for the 2-TASEP-IAF with $\mu = 2$. The species 1 and species 2 particles can hop to right (if vacant) with rates p_1 and p_2 whereas the corresponding hopping rate for the impurity (+) is ϵ . The flip process between the species 1 and 2 can occur only in the presence of an impurity at the right neighbor. The corresponding flip rates are w_{12} (for 1 transforming to 2) and w_{21} (for 2 transforming to 1).

Appendix C.

2 Model: μ -TASEP-IAF

2.1 Microscopic dynamics

Let us consider a system of μ different species of particles and impurities on a one dimensional periodic lattice with L sites $i = 1, 2, \dots, L$. Each site can either be vacant or it can be occupied by only one particle of any of the species $I = 1, 2, \dots, \mu$ or by an impurity. All the particles obey hardcore exclusion. The impurity and the vacancy are denoted by + and 0, respectively. The system evolves according to the microscopic dynamics given below,

$$\begin{aligned}
 \text{drift (species)} : \quad & I0 \xrightarrow{p_I} 0I, \quad I = 1, 2, \dots, \mu, \\
 \text{drift (impurity)} : \quad & +0, \xrightarrow{\epsilon} 0+ \\
 \text{flip} : \quad & I+ \xrightleftharpoons[w_{KI}]{w_{IK}} K+, \quad I, K = 1, \dots, \mu.
 \end{aligned} \tag{1}$$

According to the dynamics of the 2-TASEP-IAF in Eq. (1), a particle of species I can hop to its right nearest neighbor with rate p_I if the target site is vacant. The impurity (+) hopping rate is ϵ . If the right neighbor of a particle of species I is occupied by an impurity, then the species I can transform to species K with rate w_{IK} and the reverse transformation from species K to I occurs with rate w_{KI} . Clearly, this flip dynamics is activated by the presence of the impurities (+). The total number of impurities N_+ along with the total number of vacancies N_0 are conserved quantities, which can be readily seen from Eq. (1). The complete set of input parameters for the μ -TASEP-IAF is $(p_I, \epsilon, w_{IK}, \rho_+, \rho_0)$, where $\rho_+ = N_+/L$ and $\rho_0 = N_0/L$ are the conserved densities for the impurities and the vacancies respectively. To illustrate the dynamics, we present a schematic figure of the allowed dynamical processes for the $\mu = 2$ case in Fig. 1.

From the microscopic dynamics in Eq. (1), it is clear that starting from a specific initial configuration, the different species and the impurities cannot overtake each other. Indeed the flip dynamics changes the number of accessible configurations by transforming one species to another, but does not allow the dynamics to be ergodic. To discuss the non-ergodicity with an example, let us consider an initial configuration (for 2-TASEP-IAF) of the form $\{\dots 0 + 102011 + 2 \dots\}$. In the rest of this section, we will denote the particle under consideration by italics e.g. *1* for the chosen particle 1, *2* for the chosen particle 2 etc. If we

consider the 1 in $\{\dots 0 + 102011 + 2\dots\}$, it can transform into 2 by the $+$ at its right neighbor, thereby changing the configuration to $\{\dots 0 + 102012 + 2\dots\}$. However, another 1 in the initial configuration $\{\dots 0 + 102011 + 2\dots\}$ can never transform to 2 at any stage of the evolution because it can never come in contact with any $+$ (due to the non-overtaking nature of the dynamics), so that the configuration $\{\dots 0 + 102021 + 2\dots\}$ is never accessible.

2.2 Steady state: matrix product ansatz

Any configuration of the μ -TASEP-IAF can be represented by $\{s_i\} \equiv \{s_1, s_2, \dots, s_L\}$, where s_i denotes the occupation at site i . Clearly, s_i can be one of the species $K = 1, 2, \dots, \mu$ or it can be an impurity ($+$) or it can be a vacancy (0). We find that the steady state of the present model can be written in the following matrix product form

$$P(\{s_i\}) \propto \text{Tr} \left[\prod_{i=1}^L X_i \right],$$

$$X_i = E \delta_{s_i,0} + A \delta_{s_i,+} + \sum_{K=1}^{\mu} D_K \delta_{s_i,K}. \quad (2)$$

In Eq. (2), any configuration $\{s_i\}$ is represented by a string of matrices $\{X_i\}$ where the matrices D_K, A and E corresponds to a particle of species K , impurity and vacancy respectively. The time evolution of any configuration of the μ -TASEP-IAF is dictated by the Master equation

$$\frac{d}{dt}|P(t)\rangle = M|P(t)\rangle, \quad (3)$$

which in steady state becomes $M|P\rangle = 0$. Here $|P\rangle$ is a column vector containing all possible configurations and M is the rate matrix made up of the transition rates between configurations. Since the dynamics in Eq. (1) is a two-site microscopic dynamics, the rate matrix, under the periodic boundary condition, can be expressed as $M = \sum_{i=1}^L \mathcal{M}_{i,i+1}$, where $\mathcal{M}_{i,i+1}$ is a $(\mu+2)^2 \times (\mu+2)^2$ dimensional matrix. Then the steady state $M|P\rangle = 0$ of the μ -TASEP-IAF can be achieved through the following two-site flux (probability current) cancellation condition

$$\mathcal{M}_{i,i+1} \mathbf{X}_i \otimes \mathbf{X}_{i+1} = \tilde{\mathbf{X}}_i \otimes \mathbf{X}_{i+1} - \mathbf{X}_i \otimes \tilde{\mathbf{X}}_{i+1}, \quad (4)$$

where

$$\mathbf{X} = (E, A, D_1, D_2, \dots, D_\mu)^T, \quad (5)$$

and

$$\tilde{\mathbf{X}} = (\tilde{E}, \tilde{A}, \tilde{D}_1, \tilde{D}_2, \dots, \tilde{D}_\mu)^T, \quad (6)$$

where $(.)^T$ denotes the transpose of the row vector $(.)$ and $\tilde{E}, \tilde{A}, \tilde{D}_K$ are auxiliary matrices that are introduced to satisfy the steady state equation and these have to be found out consistently along with the matrix representations for E, D, D_K ($K = 1, 2, \dots, \mu$). We find that suitable choices for the auxiliary matrices for the μ -TASEP-IAF are

$$\tilde{E} = 1, \quad \tilde{A} = 0, \quad \tilde{D}_K = 0 \quad K = 1, 2, \dots, \mu. \quad (7)$$

Correspondingly, the matrices E , A and D_K have to obey the matrix algebra consisting of the equations given below

$$\begin{aligned} p_K D_K E &= D_K, & K &= 1, \dots, \mu \\ \epsilon A E &= A, \\ \sum_{\substack{I=1 \\ I \neq K}}^{\mu} w_{IK} D_I A &= D_K A \sum_{\substack{I=1 \\ I \neq K}}^{\mu} w_{KI}, & K &= 1, \dots, \mu. \end{aligned} \quad (8)$$

The last relation in Eq. (8) is reminiscent of the Kirchhoff's current law for each species K , in the sense that the total flip current from all other species to species K is equal to the total flip current from K to all other species. Note that the matrix algebra in Eq. (8) allows scalar solutions when the hopping rates for every species and the impurity become equal i.e. $p_K = \epsilon$ for all K . Naturally for this special set of rates, since the matrices reduce to scalars, no spatial correlations exist between the constituents of the system. For any other choice of rates, we expect matrix solutions to the Eq. (8). Below we discuss the cases $\mu = 2$, $\mu = 3$ extensively with explicit matrix representations and then generalize them to get the matrix representations for general μ .

$\mu = 2$: For the 2-TASEP-IAF ($K = 1, 2$), the matrix algebra [Eq. (8)] simplifies to

$$\begin{aligned} p_1 D_1 E &= D_1, & p_2 D_2 E &= D_2, \\ \epsilon A E &= A, \\ w_{12} D_1 A &= w_{21} D_2 A. \end{aligned} \quad (9)$$

Clearly, the matrix relation for the flip process becomes trivial for the two-species case implying the absence of net flip current between the two species. More precisely, the flip process satisfies detailed balance condition for $\mu = 2$. However, there are non-zero drift currents in the system. We find the following 3×3 matrix representations that satisfy the matrix algebra in Eq. (9),

$$\begin{aligned} D_1 &= w_{21} \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & D_2 &= w_{12} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\ E &= \begin{pmatrix} \frac{1}{\epsilon} & 0 & 0 \\ \frac{1}{p_1} - \frac{1}{\epsilon} & \frac{1}{p_1} & 0 \\ \frac{1}{p_2} - \frac{1}{\epsilon} & 0 & \frac{1}{p_2} \end{pmatrix}, & A &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \end{aligned} \quad (10)$$

The matrix representation of A , the impurity, resembles to that of the defect or second class particle in case of TASEP with first and second class particles [68] except the fact that the matrices are infinite dimensional in Ref. [68].

$\mu = 3$: The matrix algebra in Eq. (8) for the 3-TASEP-IAF process reads as

$$\begin{aligned} p_1 D_1 E &= D_1, & p_2 D_2 E &= D_2, & p_3 D_3 E &= D_3, \\ \epsilon A E &= A, \\ w_{21} D_2 A + w_{31} D_3 A &= (w_{12} + w_{13}) D_1 A, \\ w_{12} D_1 A + w_{32} D_3 A &= (w_{21} + w_{23}) D_2 A, \end{aligned}$$

$$w_{13}D_1A + w_{23}D_2A = (w_{31} + w_{32})D_3A. \quad (11)$$

In comparison to the last relation in Eq. (9), clearly the flip processes for the three-species case given by the last three relations in Eq. (11) do not require the detailed balance as a necessary condition. Rather, the general condition (without putting any constraint on the set of flip rates) that satisfies the flip processes in Eq. (11) is

$$w_{12}D_1A - w_{21}D_2A = w_{23}D_2A - w_{32}D_3A = w_{31}D_3A - w_{13}D_1A. \quad (12)$$

We obtain the following 4×4 representations of the matrices that satisfy Eq. (11) along with Eq. (12),

$$\begin{aligned} D_1 &= d_1 \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, D_2 = d_2 \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ D_3 &= d_3 \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ E &= \begin{pmatrix} \frac{1}{\epsilon} & 0 & 0 & 0 \\ \frac{1}{p_1} - \frac{1}{\epsilon} & \frac{1}{p_1} & 0 & 0 \\ \frac{1}{p_2} - \frac{1}{\epsilon} & 0 & \frac{1}{p_2} & 0 \\ \frac{1}{p_3} - \frac{1}{\epsilon} & 0 & 0 & \frac{1}{p_3} \end{pmatrix}, A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \end{aligned} \quad (13)$$

where

$$\begin{aligned} d_1 &= w_{21}w_{31} + w_{23}w_{31} + w_{32}w_{21}, \\ d_2 &= w_{12}w_{32} + w_{13}w_{32} + w_{31}w_{12}, \\ d_3 &= w_{13}w_{23} + w_{12}w_{23} + w_{21}w_{13}. \end{aligned} \quad (14)$$

We note that the condition Eq. (12), with the explicit matrix representations from Eq. (13), becomes

$$w_{KI}D_KA - w_{IK}D_IA = \alpha A, \quad \alpha = w_{12}w_{23}w_{31} - w_{21}w_{13}w_{32}. \quad (15)$$

The parameter α in Eq. (15) quantifies the deviations of the flip processes from the detailed balance condition between any pair of species. As we would see later, the net flip current between any two species is proportional to α . In fact $\alpha = 0$, which puts some constraints on the flip-rates, correspond to a straightforward generalization of the two-species process to three-species process with similar flip process matrix relations $w_{KI}D_KA = w_{IK}D_IA$.

general μ : For the general case of μ -TASEP-IAF ($K = 1, 2, \dots, \mu$), the structures of the matrices D_K, E, A would be similar to that of Eq. (10) and Eq. (13). More precisely, the matrices are $(\mu + 1) \times (\mu + 1)$ dimensional and the corresponding explicit representations of the matrices are given by

$$D_K = d_K|1\rangle(\langle 1| + \langle K|), \quad K = 1, \dots, \mu,$$

$$\begin{aligned}
 E &= \frac{1}{\epsilon} |1\rangle\langle 1| + \sum_{K=2}^{\mu+1} \frac{1}{p_{K-1}} |K\rangle\langle K| + \left(\frac{1}{p_{K-1}} - \frac{1}{\epsilon} \right) |K\rangle\langle 1|, \\
 A &= |1\rangle\langle 1|.
 \end{aligned}
 \tag{16}$$

In Eq. (16), the vector $\langle I| = (0, \dots, 0, 1, 0, \dots, 0)$ where 1 is placed at the I -th element with all other elements being zero and $|I\rangle$ is the transpose of $\langle I|$. Notably, the values of the coefficients d_K associated with matrices D_K in Eq. (16), can be calculated by solving the set of μ homogeneous linear equations of the form

$$d_K \sum_{I \neq K} w_{KI} + \sum_{I \neq K} w_{IK} d_I = 0, \quad K = 1, 2, \dots, \mu.
 \tag{17}$$

The summations over the index I in Eq. (17) generally includes all $I = 1, \dots, \mu$ except K . This corresponds to the general dynamics in Eq. (1) where any two species can transform into one another in the presence of the impurity (at right neighbor). However, one might be interested in special cases where the flip processes are restricted between certain pairs of species only. For example, one particular situation can be where any species K can only transform to species numbers $(K + 1)$ and $(K - 1)$. In that case, the parameter α [Eq. (15)] which dictates the flip-current between any two species, will be given by $\alpha = \left(\prod_{K=1}^{\mu} w_{KK+1} - \prod_{K=1}^{\mu} w_{K+1K} \right)$.

Notably, this reduces to the general solution of $\mu = 3$ as given in Eq. (15). However, we should mention that, for $\mu > 3$, this is not the general expression for α , rather α would depend on I, K and the flip current between different pairs of species would be different in general.

3 Partition function for special initial configuration

The non-ergodic nature of the microscopic dynamics in Eq. (1) ensures that we cannot express the partition function of the μ -TASEP-IAF in the usual form of $\text{Tr}[T^L]$, even under periodic boundary conditions. Where the transfer matrix T refers to

$$T = z_0 E + z_+ A + \sum_K D_K,
 \tag{18}$$

with z_0 and z_+ being the fugacities corresponding to the vacancies and impurities respectively, in the grand canonical ensemble. This is because T^L generates all configurations in the configuration space irrespective of the initial ordering of the species, but the dynamics in Eq. (1) allows only those configurations which preserve certain orderings from the initial configuration. To illustrate this with an example, let us consider a string of $(n + 1)$ number of species 1 particles followed by an impurity in the initial configuration i.e. $\{..11..11 + ..\} \equiv \{..1^{n+1} + ..\}$. At any step in the μ -TASEP-IAF, the ordering of the first n number of species 1 particles in this string cannot be broken, i.e. no species 2 particle or impurity can appear inside this string, except for the last 1 (which may transform to 2). On the other hand, T^L generates configurations which do not preserve such orderings. Naturally, to calculate partition function for μ -TASEP-IAF analytically, it becomes essential to choose suitable initial configurations for which we can correctly identify the accessible set of configurations in the steady state. In the rest of this section, we discuss such special initial configurations, corresponding steady states and partition functions.

3.1 $\mu = 2$

One special initial configuration $C(t=0) \equiv C(0)$ (represented by matrices) for the 2-TASEP-IAF is

$$C(0) = \underbrace{D_1 A \dots D_1 A}_{N_+} \underbrace{D_2 A \dots D_2 A}_{\bar{N}} \underbrace{D_2 \dots D_2}_{\bar{N}} \underbrace{D_1 \dots D_1}_{\bar{N}} \underbrace{E \dots E}_{\bar{N}}, \quad (19)$$

where $\underbrace{X \dots X}$ represents an uninterrupted sequence of the matrix X . We consider the densities of the uninterrupted sequences of D_2 -s and D_1 -s to be equal, which is $\bar{\rho} = \bar{N}/L$. Further, we have taken the two sequences of $D_1 A$ and $D_2 A$ to be of equal density ρ_+ so that the density of impurity (A) in each of these sequences is $\rho_+/2$. This ensures that the total density of impurities is ρ_+ . The initial configuration in Eq. (19) satisfies the relation $\bar{\rho} = \frac{1}{2}(1 - \rho_0 - 2\rho_+)$, where ρ_0 and ρ_+ are densities of the vacancies and impurities respectively. In the steady state we have $\rho_0 + \rho_+ + \rho_1 + \rho_2 = 1$, with ρ_1 and ρ_2 being the average densities of species 1 and species 2 particles in the steady state. We emphasize that ρ_0 and ρ_+ are input parameters while ρ_1 and ρ_2 are derived quantities. Starting from Eq. (19), any accessible configuration C_{ss} in the steady state is of the following generic form

$$C_{ss} = \prod_{k=1}^{N_+} (\tau D_1 + (1 - \tau) D_2) E^{m_k} A E^{n_k} \prod_{i=1}^{\bar{N}} D_2 E^{r_i} \prod_{j=1}^{\bar{N}} D_1 E^{s_j}, \quad (20)$$

subjected to the constraint $\sum_{i=1}^{N_+} (m_i + n_i) + \sum_{j=1}^{\bar{N}} (r_j + s_j) = N_0$. The parameter τ in Eq. (20) can take value either 1 or 0. The partition function, obeying the above mentioned constraint, is given by

$$\begin{aligned} Q_{N_0, N_+} = & \sum_{\{n_i\}} \sum_{\{m_i\}} \sum_{\{r_j\}} \sum_{\{s_j\}} \text{Tr} \left[\prod_{k=1}^{N_+} (D_1 + D_2) E^{m_k} A E^{n_k} \prod_{i=1}^{\bar{N}} D_2 E^{r_i} \prod_{j=1}^{\bar{N}} D_1 E^{s_j} \right] \\ & \times \delta \left(\sum_{i=1}^{N_+} (n_i + m_i) + \sum_{j=1}^{\bar{N}} (r_j + s_j) - N_0 \right). \end{aligned} \quad (21)$$

It would be useful to get rid of the $\delta(\cdot)$ constraint by associating a fugacity z_0 to the vacancy (represented by E) and considering the system in a grand canonical ensemble. The matrix strings $(D_1 + D_2) E^m A E^n$, $D_2 E^r$ and $D_1 E^s$ can be evaluated by incorporating the matrix algebra in Eq. (9) along with the explicit representations from Eq. (10). Consequently, the partition function in the grand canonical ensemble under the periodic boundary condition, finally becomes

$$Q_{N_+}(z_0) = \left(\left[\frac{w_{21}}{1 - \frac{z_0}{p_1}} + \frac{w_{12}}{1 - \frac{z_0}{p_2}} \right] \left(\frac{1}{1 - \frac{z_0}{\epsilon}} \right) \right)^{N_+} \left(\frac{w_{21} w_{12}}{(1 - \frac{z_0}{p_1})(1 - \frac{z_0}{p_2})} \right)^{\bar{N}}. \quad (22)$$

For the special initial configuration in Eq. (19), we have derived the partition function in Eq. (22). The fugacity z_0 can be obtained as a function of the vacancy density and other input parameters by inverting the density-fugacity relation $\rho_0 = z_0 \frac{d}{dz_0} \ln(Q_{N_+}(z_0))$. The other conserved quantity, the impurity density is already fixed at $\rho_+ = N_+/L$. The expression Eq. (22) would be used for evaluating the observables of interest in the next Sec. 4.

3.2 $\mu = 3$

Similar to the initial configuration $C(t=0)$ for $\mu = 2$, a suitable initial configuration for the three species case that enables us to perform analytical calculation of partition function and observables, is

$$C(0) \equiv \underbrace{D_1 A \dots D_1 A}_{N_+} \underbrace{D_2 A \dots D_2 A}_{N_+} \underbrace{D_3 A \dots D_3 A}_{N_+} \underbrace{D_3 \dots D_3}_{N_+} \underbrace{D_2 \dots D_2}_{N_+} \underbrace{D_1 \dots D_1}_{N_+} \underbrace{E \dots E}_{N_0}. \quad (23)$$

The density of the uninterrupted sequence of each species 1, 2, 3 in Eq. (23) is taken to be equal to $\bar{\rho} = \bar{N}/L$. Moreover, we have chosen the density of each of the sequences $D_1 A$, $D_2 A$ and $D_3 A$ to be $2\rho_+/3$ where the density of impurities in each of these sequences is $\rho_+/3$. Consequently, the choice of initial configuration in Eq. (23) ensures that the total density of impurities remain ρ_+ . In the steady state, $\rho_0 + \rho_+ + \sum_{I=1}^3 \rho_I = 1$, where ρ_I is the average density of the non-conserved species I . Starting from Eq. (23), the form of any accessible configuration C_{ss} in steady state would be

$$C_{ss} \equiv \prod_{k=1}^{N_+} (D_1 \delta_{\tau,1} + D_2 \delta_{\tau,2} + D_3 \delta_{\tau,3}) E^{m_k} A E^{n_k} \prod_{i=1}^{\bar{N}} D_3 E^{l_i} \prod_{i=1}^{\bar{N}} D_2 E^{r_i} \prod_{i=1}^{\bar{N}} D_1 E^{s_i}, \quad (24)$$

with the conservation of the total number of vacancies $N_0 = \sum_{k=1}^{N_+} (m_k + n_k) + \sum_{i=1}^{\bar{N}} (l_i + r_i + s_i)$, where $\delta_{\tau,K}$ is the Kronecker delta symbol with $K = 1, 2, 3$. As done in case of $\mu = 2$, here also we associate a fugacity z_0 with the vacancy. Using the matrix algebra from Eq. (11) alongside the matrix representations in Eqs.(13) and (14), the partition function in grand canonical ensemble becomes

$$Q_{N_+}(z_0) = \left(\left[\sum_{I=1}^3 \frac{d_I}{1 - \frac{z_0}{p_I}} \right] \left(\frac{1}{1 - \frac{z_0}{\epsilon}} \right) \right)^{N_+} \left(\prod_{I=1}^3 \frac{d_I}{1 - \frac{z_0}{p_I}} \right)^{\bar{N}}, \quad (25)$$

where the explicit expressions for d_I -s have been presented earlier in Eq. (14).

3.3 general μ

For the μ -TASEP-IAF ($K = 1, \dots, \mu$), the generalization of the initial configurations in Eq. (19) ($\mu = 2$) and Eq. (23) ($\mu = 3$) would be

$$C(0) \equiv \left(\prod_{i=1}^{N_+/\mu} D_1 A \prod_{i=1}^{N_+/\mu} D_2 A \dots \prod_{i=1}^{N_+/\mu} D_\mu A \right) \left(\prod_{i=1}^{\bar{N}} D_1 \prod_{i=1}^{\bar{N}} D_2 \dots \prod_{i=1}^{\bar{N}} D_\mu \right) \prod_{i=1}^{N_0} E. \quad (26)$$

The above initial configuration is chosen in a way that the density of each sequence $D_I A$ ($I = 1, \dots, \mu$) is $2\rho_+/\mu$ in which the density of impurities is equal to ρ_+/μ , so that the total impurity density adds up to ρ_+ . We have $\rho_0 + \rho_+ + \sum_{I=1}^{\mu} \rho_I = 1$ in the steady state, with ρ_I being the average density of the non-conserved species I . Proceeding in the same way as shown in

cases of $\mu = 2$ and $\mu = 3$, we obtain the partition function for the general μ -TASEP-IAF to be

$$Q_{N_+}(z_0) = \left(\left[\sum_{I=1}^{\mu} \frac{d_I}{1 - \frac{z_0}{p_I}} \right] \left(\frac{1}{1 - \frac{z_0}{\epsilon}} \right) \right)^{N_+} \left(\prod_{I=1}^{\mu} \frac{d_I}{1 - \frac{z_0}{p_I}} \right)^{\bar{N}}, \quad (27)$$

where d_I is the solution of Eq. (17) and z_0 is the fugacity associated with the vacancy in the grand canonical ensemble.

In this section, we have derived the partition functions of the μ -TASEP-IAF with $\mu = 2$, $\mu = 3$ and general μ in Eq. (22), Eq. (25) and Eq. (27), respectively, for specific initial configuration Eq. (19), Eq. (23) and Eq. (26), respectively, under periodic boundary conditions. These results would be useful to calculate the average values of observables in the next section for the same initial configurations discussed here.

4 Observables: comparisons of analytical results with Monte Carlo simulations

In this section, we analytically calculate the following observables in the steady state, (i) average density ρ_I of the non-conserved species I , (ii) average drift currents J_{I0} and J_{+0} , for species I and impurities respectively, (iii) average flip current $J_{I \leftrightarrow K}$ between species pair (I, K) and (iv) two-point correlations C_{0I} between vacancies (0) and species I . Mostly we will restrict the calculations to the number of species $\mu = 2$, except using $\mu = 3$ for the case of average flip current (since there is no net flip current between pair of species for $\mu = 2$). In particular, starting from the special initial configuration Eq. (19) for $\mu = 2$ (or, Eq. (23) for $\mu = 3$), we will show agreements between the analytical calculations and the Monte Carlo simulation results.

4.1 Species densities

First we consider the average densities (ρ_I) of the non-conserved species $I = 1, 2$. The formal expression for ρ_I in the steady state under the periodic boundary condition, can be written as

$$\rho_1 = \frac{1}{2}(1 - \rho_0 - 2\rho_+) + \frac{\rho_+}{Q_{N_+}} \sum_{n_1=0}^{\infty} \dots \sum_{n_{N_+}=0}^{\infty} \sum_{m_1=0}^{\infty} \dots \sum_{m_{N_+}=0}^{\infty} \sum_{r_1=0}^{\infty} \dots \sum_{r_{\bar{N}}=0}^{\infty} \sum_{s_1=0}^{\infty} \dots \sum_{s_{\bar{N}}=0}^{\infty} \text{Tr} \left[D_1(z_0 E)^{m_1} A(z_0 E)^{n_1} \prod_{k=2}^{N_+} (D_1 + D_2)(z_0 E)^{m_k} A(z_0 E)^{n_k} \prod_{k=1}^{\bar{N}} D_2(z_0 E)^{r_k} \prod_{k=1}^{\bar{N}} D_1(z_0 E)^{s_k} \right]. \quad (28)$$

To elaborate Eq. (28), the main point is to note the expression inside the trace ($\text{Tr}[\cdot]$) that denotes configurations with at least one D_1 . This can be understood more clearly by comparing it with the expression for any possible configuration in Eq. (20). From Eq. (20), to arrive at the matrix string inside the trace in Eq. (28), one has to put $\tau = 1$ for one k value to ensure the presence of at least one species 1 particle (D_1) in the configuration. Since this D_1 could




Figure 2: The figures (a) and (b) show the variations of the average species densities ρ_I ($I = 1, 2$) against vacancy density ρ_0 and flip rate w_{12} respectively in the steady state. The species densities decrease linearly with increasing ρ_0 in (a). In (b), with increasing w_{12} , ρ_1 and ρ_2 decrease and increase respectively, both in nonlinear manners. Notably, in the parameter range $w_{21} < w_{12} < w_{12}^*$, we observe $\rho_2 < \rho_1$ in spite of the higher flip rate of transformation from species 1 to species 2. The common parameters for both figures (a) and (b) are $L = 10^3$, $p_1 = 0.3$, $p_2 = 1.0$, $\epsilon = 0.1$, $\rho_+ = 0.2$. For (a), $w_{12} = 0.4$ and $w_{21} = 1.0$. For (b), $\rho_0 = 0.2$ and $w_{21} = 0.6$. The ensemble average is done over 10^5 samples.

have been placed for any $k = 1, 2, \dots, N_+$, we have a combinatorial pre-factor $\rho_+ = N_+/L$ in Eq. (28). The summations over all the variables $\{m, n, r, s\}$ from zero to infinity are performed as the system is considered in the grand canonical ensemble. The first factor $(1 - \rho_0 - 2\rho_+)/2$ in Eq. (28) is to take care of the \bar{N} number of D_1 -s present in the initial configuration Eq. (19) that cannot flip. Using the matrix algebra and matrix representations from Eqs. (9) and (10) respectively, we finally arrive at the following expressions for the average densities of the non-conserved species,

$$\begin{aligned}
 \rho_1 &= \rho_+ \frac{w_{21}}{\left(1 - \frac{z_0}{p_1}\right)} \frac{1}{\left[\frac{w_{21}}{1 - \frac{z_0}{p_1}} + \frac{w_{12}}{1 - \frac{z_0}{p_2}}\right]} + \frac{1}{2}(1 - 2\rho_+ - \rho_0) \\
 \rho_2 &= \rho_+ \frac{w_{12}}{\left(1 - \frac{z_0}{p_2}\right)} \frac{1}{\left[\frac{w_{21}}{1 - \frac{z_0}{p_1}} + \frac{w_{12}}{1 - \frac{z_0}{p_2}}\right]} + \frac{1}{2}(1 - 2\rho_+ - \rho_0).
 \end{aligned}
 \tag{29}$$

The fugacity z_0 is obtained in terms of the input parameters by solving the density-fugacity relation $\rho_0 = z_0 \frac{d}{dz_0} \ln(Q_{N_+}(z_0))$ using Mathematica. Replacing the solution of z_0 (which is too lengthy to provide here) in Eq. (29), we finally get the average densities as functions of input parameters ($p_{1,2}, w_{12,21}, \epsilon, \rho_0, \rho_+$) only.

We compare the analytical results with those of Monte Carlo simulations, starting from the same initial configuration Eq. (19). In the simulation, we vary the vacancy density in




Figure 3: The figure illustrates the non-monotonic behaviors of the drift currents J_{I0} ($I = 1, 2$) with increasing vacancy density ρ_0 , unlike the drift current for the impurities J_{+0} which increases monotonically with increasing ρ_0 . The analytical results (solid lines) are in agreement with the Monte Carlo simulation results (points). The parameters used are $L = 10^3$, $p_1 = 0.3$, $p_2 = 1.0$, $\epsilon = 0.1$, $w_{12} = 0.4$, $w_{21} = 1.0$, $\rho_+ = 0.2$. The ensemble average is done over 10^5 samples.

the initial configuration by changing the lengths of the uninterrupted strings of D_2 and D_1 in Eq. (19) i.e. simply by tuning $\bar{\rho}$ which is related to vacancy density ρ_0 as $\rho_0 + 2\bar{\rho} = 1 - 2\rho_+$. In Figs. 2(a) and (b), we observe that the analytical and simulation results are in agreement with each other, where ρ_I is plotted against ρ_0 and w_{12} (flip rate of species 1 to species 2), respectively. The species densities decrease linearly with increasing ρ_0 [Fig. 2(a)] whereas they decrease in non-linear fashion with increasing w_{12} [Fig. 2(b)]. Notably, in the absence of any drift, we would have $\rho_2 = \rho_1$ exactly at $w_{12}^* = w_{21}$. However, due to the hopping process, this point shifts to

$$w_{12}^* = w_{21} \frac{(1 - \frac{z_0}{p_2})}{(1 - \frac{z_0}{p_1})}. \quad (30)$$

Consequently in Fig. 2(b), for a particular set of chosen parameters, we observe that for $w_{21} < w_{12} < w_{12}^*$, one still has $\rho_2 < \rho_1$. In other words, when $w_{12} \in (w_{21}, w_{12}^*)$, although the species 1 particles more often transform to species 2 particles, still the average density of species 2 particles is less than that of species 1 particles. For any value of μ , the general expression for the average density ρ_I for the non-conserved species I ($I = 1, 2, \dots, \mu$) is

$$\rho_I = \rho_+ \frac{d_I}{(1 - \frac{z_0}{p_I})} \frac{1}{\sum_{K=1}^{\mu} \frac{d_K}{(1 - \frac{z_0}{p_K})}} + \frac{1}{\mu} (1 - 2\rho_+ - \rho_0), \quad (31)$$

where d_I is the solution of Eq. (17) (e.g. the solution for $\mu = 3$ is explicitly given in Eq. (14)).




Figure 4: The figure shows comparison of drift currents J_{I0} between theory (solid lines) and Monte Carlo simulations (points) by changing flip rate w_{12} . The nonlinear monotonic decrease and increase in J_{10} and J_{20} respectively are similar in nature to the effect of w_{12} on the species densities (Fig. 2). The impurity drift current increases slowly with increasing w_{12} . The parameters used are $L = 10^3$, $p_1 = 0.3$, $p_2 = 1.0$, $\epsilon = 0.1$, $w_{21} = 0.6$, $\rho_+ = 0.2$, $\rho_0 = 0.2$. The ensemble average is done over 10^5 samples.

4.2 Drift current

Next we consider the average drift currents J_{I0} and J_{+0} for the non-conserved species I ($I = 1, 2$) and the impurity respectively. We focus on $I = 1$ to explain the procedure for calculating the current J_{10} , because the parallel procedure applies for any other species. The average drift current J_{10} is equal to $p_1 \langle 10 \rangle$, where $\langle 10 \rangle$ is the ensemble average of the pair 10. In terms of matrices the expression $\langle 10 \rangle$ simply translates to $\langle D_1 E \rangle$. The current J_{10} can be calculated in two parts,

$$J_{10} = p_1 \langle 10 \rangle = p_1 \langle D_1 E \rangle = J_{10}^{(1)} + J_{10}^{(2)} = p_1 \langle D_1 E \rangle^{(1)} + p_1 \langle D_1 E \rangle^{(2)}, \quad (32)$$

where $J_{10}^{(1)} = p_1 \langle D_1 E \rangle^{(1)}$ is the contribution from the drift of species 1 particles that can flip and $J_{10}^{(2)} = p_1 \langle D_1 E \rangle^{(2)}$ is the corresponding contribution from species 1 particles that cannot flip (as they cannot have any impurity as right neighbor) according to the initial configuration in Eq. (19). Correspondingly, the term $D_1 E$ in the averages $\langle D_1 E \rangle^{(1)}$ and $\langle D_1 E \rangle^{(2)}$ would come from the product sequences $(\tau D_1 + (1 - \tau) D_2) E^m A$ and $D_1 E^s$ respectively Eq. (20). The expression for $J_{10}^{(1)}$ is given by

$$J_{10}^{(1)} = p_1 \langle D_1 E \rangle^{(1)} = \frac{\rho_+}{Q_{N_+}} \sum_{n_1=0}^{\infty} \dots \sum_{n_{N_+}=0}^{\infty} \sum_{m_1=1}^{\infty} \dots \sum_{m_{N_+}=0}^{\infty} \sum_{r_1=0}^{\infty} \dots \sum_{r_{\bar{N}}=0}^{\infty} \dots \sum_{s_{\bar{N}}=0}^{\infty} \text{Tr} \left[z_0 p_1 D_1 E (z_0 E)^{m_1-1} A (z_0 E)^{n_1} \prod_{k=2}^{N_+} (D_1 + D_2) (z_0 E)^{m_k} A (z_0 E)^{n_k} \prod_{k=1}^{\bar{N}} D_2 (z_0 E)^{r_k} \prod_{k=1}^{\bar{N}} D_1 (z_0 E)^{s_k} \right]. \quad (33)$$

The construction of Eq. (33) follows similar arguments as of Eq. (28), except now we have to place $D_1 E$ instead of D_1 . This also reflects in the summations, note that the lower limit of

the index m_1 has been changed to 1 instead of 0 to ensure the presence of one $D_1 E$. Similarly, the formal expression for $J_{10}^{(2)}$ is

$$J_{10}^{(2)} = p_1 \langle D_1 E \rangle^{(2)} = \frac{1}{Q_{N_+}} \sum_{n_1=0}^{\infty} \cdots \sum_{r_{\bar{N}}=0}^{\infty} \sum_{s_1=1}^{\infty} \cdots \sum_{s_{\bar{N}}=0}^{\infty} \text{Tr} \left[\prod_{k=1}^{N_+} (D_1 + D_2)(z_0 E)^{m_k} A \prod_{k=1}^{\bar{N}} D_2(z_0 E)^{r_k} z_0 p_1 D_1 E (z_0 E)^{s_1-1} \prod_{k=2}^{\bar{N}} D_1(z_0 E)^{s_k} \right]. \quad (34)$$

Obviously in Eq. (34), the lower limit of the index s_1 is shifted to 1 from 0. Using the matrix algebra and matrix representations from Eqs. (9) and (10), we get from Eqs. (33) and (34):

$$J_{10}^{(1)} = z_0 \rho_+ \frac{w_{21}}{\left(1 - \frac{z_0}{p_1}\right)} \frac{1}{\left[\frac{w_{21}}{1 - \frac{z_0}{p_1}} + \frac{w_{12}}{1 - \frac{z_0}{p_2}}\right]}, \quad J_{10}^{(2)} = \frac{z_0}{2} (1 - 2\rho_+ - \rho_0). \quad (35)$$

Substituting Eq. (35) into Eq. (32), we obtain J_{10} . Following the same procedures, one can calculate J_{20} and J_{+0} . We finally arrive at the analytical expressions for the drift currents under the periodic boundary condition, given below:

$$\begin{aligned} J_{10} &= z_0 \rho_+ \frac{w_{21}}{\left(1 - \frac{z_0}{p_1}\right)} \frac{1}{\left[\frac{w_{21}}{1 - \frac{z_0}{p_1}} + \frac{w_{12}}{1 - \frac{z_0}{p_2}}\right]} + \frac{z_0}{2} (1 - 2\rho_+ - \rho_0), \\ J_{20} &= z_0 \rho_+ \frac{w_{12}}{\left(1 - \frac{z_0}{p_2}\right)} \frac{1}{\left[\frac{w_{21}}{1 - \frac{z_0}{p_1}} + \frac{w_{12}}{1 - \frac{z_0}{p_2}}\right]} + \frac{z_0}{2} (1 - 2\rho_+ - \rho_0), \\ J_{+0} &= \rho_+ z_0. \end{aligned} \quad (36)$$

In Figs. 3 and 4 we present the variation of the drift currents as functions of vacancy density ρ_0 and flip rate w_{12} , respectively. For both cases, the analytical results match with the Monte Carlo simulation results. The drift currents for the species 1 and 2 exhibit non-monotonic behaviors with increasing vacancy density whereas the impurity drift current increases monotonically (Fig. 3). At lower vacancy densities, as we increase ρ_0 , the chances for hopping increase, thereby increasing the drift current in Fig. 3. However, after a particular value of ρ_0 , if we increase it further, the densities of the non-conserved species fall considerably so that drift current ultimately decreases, although there are many vacancies in the system. Since the impurities do not flip, the density of impurities is fixed and the corresponding impurity drift current can only increase with increasing ρ_0 (Fig. 3). The maximum of the drift current for different species generally occur at distinct values of ρ_0 . With variation of the flip rate w_{12} (Fig. 4), the drift currents of the non-conserved species show similar non-linear behaviors like their corresponding densities [Fig. 2(b)]. Notably, although the flip dynamics does not affect the drift of the impurities explicitly, still we observe that the impurity drift current increases with increasing w_{12} , albeit weakly.

For any positive integer value of μ , the general expression for the drift current of any non-conserved species I ($I = 1, \dots, \mu$) is obtained to be

$$J_{I0} = z_0 \rho_+ \frac{d_I}{\left(1 - \frac{z_0}{p_I}\right)} \frac{1}{\sum_{K=1}^{\mu} \frac{d_K}{\left(1 - \frac{z_0}{p_K}\right)}} + \frac{z_0}{\mu} (1 - 2\rho_+ - \rho_0), \quad (37)$$

where the d_I is the solution of Eq. (17).




Figure 5: The figure illustrates the variation of two-point correlation C_{01} between the vacancy and species 1 by tuning the vacancy density ρ_0 . We observe that this correlation initially starts increasing with increasing ρ_0 , reaches to a local maximum, then decreases. However, as ρ_0 is increased further, C_{01} reaches to a local minimum and then starts increasing again. The parameters used are $L = 10^3$, $p_1 = 0.3$, $p_2 = 1.0$, $\epsilon = 0.1$, $w_{12} = 0.8$, $w_{21} = 1.0$, $\rho_+ = 0.2$. The ensemble average is done over 10^5 samples.

4.3 Two-point correlations

Besides the currents, we would like to calculate some other two point functions which have interesting features. We have basically calculated the two-point functions $\langle 10 \rangle$ and $\langle 20 \rangle$ in the process of determining the drift currents J_{10} and J_{20} respectively. Now we focus on the nearest neighbors two-point correlations involving $\langle 01 \rangle$ and $\langle 02 \rangle$. We find the exact expressions for the corresponding two-point correlations under the periodic boundary condition as

$$\begin{aligned}
 C_{01} = \langle 01 \rangle - \langle 0 \rangle \langle 1 \rangle &= \left(\frac{z_0}{\epsilon} - \rho_0 \right) \frac{w_{21}}{\left(1 - \frac{z_0}{p_1} \right)} \frac{\rho_+}{\left[\frac{w_{21}}{1 - \frac{z_0}{p_1}} + \frac{w_{12}}{1 - \frac{z_0}{p_2}} \right]} \\
 &\quad + \left(\frac{z_0}{2p_1} - \frac{\rho_0}{2} \right) (1 - 2\rho_+ - \rho_0), \\
 C_{02} = \langle 02 \rangle - \langle 0 \rangle \langle 2 \rangle &= \left(\frac{z_0}{\epsilon} - \rho_0 \right) \frac{w_{12}}{\left(1 - \frac{z_0}{p_2} \right)} \frac{\rho_+}{\left[\frac{w_{21}}{1 - \frac{z_0}{p_1}} + \frac{w_{12}}{1 - \frac{z_0}{p_2}} \right]} \\
 &\quad + \left(\frac{z_0}{2p_2} - \frac{\rho_0}{2} \right) (1 - 2\rho_+ - \rho_0). \tag{38}
 \end{aligned}$$

The correlation C_{01} is plotted against the vacancy density ρ_0 in Fig. 5. As ρ_0 is increased starting from zero, the correlation also increases and reaches a local maximum, followed by a decrease and reaching a local minimum. After this point, if the vacancy density is increased further, C_{01} increases again. So, instead of a single maximum or single minimum, the correlation C_{01} interestingly exhibits both local maximum and local minimum with the variation of ρ_0 . In other words, C_{01} increases with increasing vacancy density for both sufficiently high and sufficiently low values of ρ_0 , with intermediate non-monotonic character. In Fig. 6, it is




Figure 6: The figure shows a crossover from negative correlation to positive correlation for C_{02} (between vacancy and species 2) with increasing vacancy density ρ_0 . There is some special intermediate density for which C_{02} becomes zero. The parameters used are $L = 10^3$, $p_1 = 0.3$, $p_2 = 1.0$, $\epsilon = 0.1$, $w_{12} = 0.8$, $w_{21} = 1.0$, $\rho_+ = 0.2$. The ensemble average is done over 10^5 samples.

interesting to see that the non-monotonic behavior of C_{02} is such that it goes from negative correlation values to positive correlation values. Naturally, there exists some intermediate value of ρ_0 for which the special arrangements of the accessible steady state configurations makes the average correlation C_{02} to be zero.

4.4 Flip current

All the observables we have discussed up to now (average species densities, drift currents, correlations), correspond to 2-TASEP-IAF. However, the net flip current is zero for $\mu = 2$. Therefore, in order to have a net non-zero flip current between pairs of species, here we consider the case $\mu = 3$ ($I = 1, 2, 3$). We denote the net flip current between species I and K as $J_{I \leftrightarrow K}$. For $\mu = 3$, we find that the net flip current between any two species $[(1, 2), (2, 3), (3, 1)]$ are equal to each other (i.e. independent of the indices I and K) and its exact form is given by

$$J_{I \leftrightarrow K} = w_{IK} \langle I+ \rangle - w_{KI} \langle K+ \rangle = \rho_+ \frac{(w_{12}w_{23}w_{31} - w_{21}w_{13}w_{32})}{\left[\frac{d_1}{1-\frac{z_0}{p_1}} + \frac{d_2}{1-\frac{z_0}{p_2}} + \frac{d_3}{1-\frac{z_0}{p_3}} \right]}. \quad (39)$$

The initial configuration that we have used to arrive at Eq. (40) is the one in Eq. (23). In Eq. (40), we have calculated the current in the cyclic ordering i.e. ($I = 1, K = 2$), ($I = 2, K = 3$), ($I = 3, K = 1$). We have presented the behavior of the flip current as functions of the vacancy density ρ_0 and flip rate w_{12} in Figs. 7(a) and 7(b) respectively. We observe that the analytical calculation are in agreement with the Monte Carlo simulation results. In Fig. 7(a), the flip-current decreases monotonically in a nonlinear manner with increasing vacancy density. The reason behind this, is the decrease in species densities with increasing ρ_0 [Fig. 2(a)]. On the other hand, flip current between any pair of species increases monotonically with increasing flip rate w_{12} (Fig. 7(b)).




Figure 7: The figures exhibit the variation of the flip current $J_{I \leftrightarrow K}$ with vacancy density ρ_0 and flip rate w_{12} in (a) and (b), respectively. The analytical and Monte Carlo simulation results show good agreement. The flip current monotonically decreases with increasing ρ_0 whereas it increases monotonically with increasing w_{12} . The common parameters for both figures (a) and (b) are $L = 10^3$, $p_1 = 0.3$, $p_2 = 1.0$, $p_3 = 1.0$, $\epsilon = 0.1$, $w_{21} = 0.5$, $w_{23} = 0.5$, $w_{32} = 0.2$, $w_{31} = 0.8$, $w_{13} = 0.2$, $\rho_+ = 0.15$. For (a), $w_{12} = 0.4$ and for (b), $\rho_0 = 0.22$. The ensemble average is done over 10^5 samples.

The generalization of the formula Eq. (40) for any μ , under the periodic boundary condition, is obtained as

$$J_{I \leftrightarrow K} = \rho_+ \frac{(d_I w_{IK} - d_K w_{KI})}{\sum_{K=1}^{\mu} \frac{d_K}{(1 - \frac{z_0}{p_K})}}, \quad (40)$$

where d_I, d_K are the solutions of Eq. (17). We should mention that, for $\mu > 3$, the flip currents between different pairs of species would be in general distinct from one another.

4.5 Non-ergodicity: dependence on initial configuration

Here we establish the non-ergodicity of the μ -TASEP-IAF by showing explicitly the dependence of the average steady state values of observables on the choice of the initial configuration. For simplicity, we restrict ourselves to the case of $\mu = 2$. We choose two different initial configurations as follows. (i) The initial configuration given in Eq. (19) for which we know exactly which constituent (any species or impurity or vacancy) is placed at a given lattice site. Since the initial arrangement is specified completely, we call this configuration *specified initial configuration* (SIC). (ii) A random initial configuration where the constituent at each site is selected randomly such that the densities of impurities and vacancies, and the initial densities of species I ($I = 1, 2$), are exactly the same as that of the SIC described in (i). Since the initial arrangement for this configuration is randomly carried out, we call it *random initial configuration* (RIC). To clarify, both SIC and RIC are characterized by the same set of rates ($p_1, p_2, \epsilon, w_{12}, w_{21}$) and the densities ($\rho_+, \rho_0, \rho_1(0), \rho_2(0)$), where $\rho_1(0), \rho_2(0)$ represent the initial ($t = 0$) densities of the non-conserved species 1 and 2 respectively. Although the analysis of the steady state for the SIC can be performed exactly as discussed already, the





Figure 8: The set of figures exhibits the dependence of steady state values of observables on the choice of initial configuration. We observe clear deviations between SIC (denoted by circles) and RIC (denoted by rectangles) in (a)-(d) and the amount of deviation in each figure changes with the variation of input parameter. The common set of parameters used for (a)-(d) are $L = 10^3$, $p_1 = 0.3$, $\epsilon = 0.1$, $\rho_+ = 0.216$, $\rho_0 = 0.203$, $\rho_1(0) = 0.321$, $\rho_2(0) = 0.260$. The other parameters for (a), (b) are $p_2 = 1.0$, $w_{21} = 0.6$ and for (c), (d) are $w_{12} = 0.5$, $w_{21} = 0.4$. The ensemble averages are done over 10^5 samples.

same could not be done for the RIC. Therefore, in this section we use Monte Carlo simulations to compare the steady state observable values for SIC and RIC.

We compare the steady state observable values for SIC and RIC with the same set of input parameters $(p_1, p_2, \epsilon, w_{12}, w_{21}, \rho_+, \rho_0)$ and same initial densities of the species $(\rho_1(0), \rho_2(0))$ in Fig. 8. We denote the data points for SIC and RIC with different symbols, circles and rectangles respectively, in Fig. 8. The variations of the non-conserved species densities ρ_I are presented in Figs. 8(a) and 8(c) as functions of flip rate w_{12} and hop rate p_2 , respectively. Both figures exhibit clear quantitative differences between the density values for SIC and RIC. We observe that the deviations between SIC and RIC decreases (increases) with increasing w_{12} (p_2). The initial configuration dependence of the steady state values of nearest neighbor two-point correlations are shown in Figs. 8(b) and (d). In Fig. 8(b), we observe that the correlation between species 1 particles $C_{11} = \langle 11 \rangle - \rho_1^2$ has distinct numerical values for SIC and RIC when the parameter w_{12} is tuned. Interestingly, the correlation corresponding to RIC changes from negative to positive whereas the same for SIC remains positive with increasing w_{12} . This implies the existence of some intermediate w_{12} which corresponds to uncorrelated species 1 particles for RIC, whereas they are correlated for the SIC. Similar kind of interesting behavior is observed for the correlation between species 2 particles $C_{22} = \langle 22 \rangle - \rho_2^2$ when plotted against p_2 in Fig. 8(d). Thus we have illustrated the dependence of steady state values of species densities and correlations on the choice of the initial configuration. The same can also be investigated in other two point and higher point functions.

5 Partially asymmetric generalization: μ -ASEP-IAF

In this section, we consider the μ -ASEP-IAF under periodic boundary conditions, a generalization of the μ -TASEP-IAF in Eq. (1), by including partially asymmetric motions of different species of particles. A particle of species I can hop towards right with rate p_I and it can hop towards left with rate q_I ($I = 1, \dots, \mu$), if the target site is empty. Notably, the impurities are not allowed to hop to left. This naturally adds another way to distinguish the conserved impurities from all non-conserved species. The microscopic dynamics is given by,



The μ -ASEP-IAF remains non-ergodic in nature. Since we could obtain the steady state of the μ -TASEP-IAF using matrix product ansatz [Eq. (2)], we assume the same can be done for the partially asymmetric motion also.

5.1 Matrix algebra, auxiliaries and matrix representations

The matrix algebra for the dynamics in Eq. (41) under the periodic boundary condition, is

$$\begin{aligned}
 p_K D_K E - q_K E D_K &= D_K, & K = 1, \dots, \mu \\
 \epsilon A E &= A,
 \end{aligned}$$

$$\sum_{\substack{I=1 \\ I \neq K}}^{\mu} w_{IK} D_I A = D_K A \sum_{\substack{I=1 \\ I \neq K}}^{\mu} w_{KI}, \quad K = 1, \dots, \mu. \quad (42)$$

In comparison to the matrix algebra [Eq. (8)] for the μ -TASEP-IAF, the only changes occurring in Eq. (42) correspond to the drifts of the non-conserving species. At this point, we should mention that the matrix equation $p_K D_K E - q_K E D_K = D_K$ has been studied in Ref. [52], in context of a conserved disordered ASEP model. Due to the presence of the impurities and the flip processes activated by them, the matrix algebra for μ -ASEP-IAF in Eq. (42), can be considered as a generalization of the matrix algebra in Ref. [52]. To arrive at the matrix algebra in Eq. (42) from the dynamics (41), ansatz (2) and flux cancellation condition (4), the choice of the auxiliary matrices are the same as the totally asymmetric case, i.e.

$$\tilde{E} = 1, \quad \tilde{A} = 0, \quad \tilde{D}_K = 0 \quad K = 1, 2, \dots, \mu. \quad (43)$$

However, unlike the totally asymmetric case, we find the matrix representations for the μ -ASEP-IAF to be infinite dimensional. Notably, this does not necessarily eliminate the possibility of getting alternate finite dimensional representations of the matrices. Below we present the matrix representations for $\mu = 3$ case explicitly (as we will stick to $\mu = 3$ for the discussion of observable in this section) and mention the changes required to construct the matrices for any $\mu > 0$.

$\mu = 3$: A possible set of representations of the matrices for the 3-ASEP-IAF ($I = 1, 2, 3$) is the following

$$E = \begin{pmatrix} 0 & 0 & 0 & 0 & \cdot & \cdot \\ 1 & 0 & 0 & 0 & \cdot & \cdot \\ 0 & 1 & 0 & 0 & \cdot & \cdot \\ 0 & 0 & 1 & 0 & \cdot & \cdot \\ 0 & 0 & 0 & 1 & & \\ \cdot & \cdot & & & \cdot & \cdot \\ \cdot & \cdot & & & & \cdot \end{pmatrix}, \quad A = \begin{pmatrix} 1 & \frac{1}{\epsilon} & \frac{1}{\epsilon^2} & \frac{1}{\epsilon^3} & \cdot & \cdot \\ 0 & 0 & 0 & 0 & \cdot & \cdot \\ 0 & 0 & 0 & 0 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}$$

$$D_I = \begin{pmatrix} d_I^{1,1} & d_I^{1,2} & d_I^{1,3} & d_I^{1,4} & \cdot & \cdot \\ 0 & d_I^{2,2} & d_I^{2,3} & d_I^{2,4} & \cdot & \cdot \\ 0 & 0 & d_I^{3,3} & d_I^{3,4} & \cdot & \cdot \\ 0 & 0 & 0 & d_I^{4,4} & \cdot & \cdot \\ \cdot & \cdot & & & \cdot & \cdot \\ \cdot & \cdot & & & & \cdot \end{pmatrix}, \quad I = 1, 2, 3$$

$$d_I^{m,m+r} = \frac{\binom{m}{r}}{r! p_I^r} \left(\frac{q_I}{p_I} \right)^{m-1} d_I^{1,1}, \quad \forall r \geq 0$$

$$d_1^{1,1} = w_{21} w_{31} + w_{23} w_{31} + w_{32} w_{21},$$

$$d_2^{1,1} = w_{12} w_{32} + w_{13} w_{32} + w_{31} w_{12},$$

$$d_3^{1,1} = w_{13} w_{23} + w_{12} w_{23} + w_{21} w_{13}. \quad (44)$$

In the absence of the impurities (A) and the flip processes, the term $d_\mu^{1,1}$ for every μ becomes unity, and corresponding the matrix representations for D_I and E in Eq. (44) (and their generalizations for general μ) are the same as that of the conserved disordered ASEP [24, 52].

In Eq. (44), we observe that the matrices (D_I) corresponding to the species (I) are upper triangular. The subscript I in matrix element $d_I^{i,j}$ denotes the species I whereas the superscript (i, j) refers to the i -th row and j -th column of the matrix. The notation $(m)_r$ used in the expression of $d_I^{m,m+r}$ corresponds to the Pochhammer symbol for rising factorials, $(m)_r = m(m+1)(m+2)\dots(m+r-1)$ with $(m)_0 = 1$. The matrix E representing vacancy is a lower shift matrix and the matrix A representing impurity has non-zero terms in a single row only. For the simpler case $\mu = 2$ the only changes in comparison to Eq. (44) will be in the values of $d_I^{1,1}$ ($I = 1, 2$), which would be simply $d_1^{1,1} = w_{21}$ and $d_2^{1,1} = w_{12}$. In fact, the matrix representations [Eq. (44)] for the 3-ASEP-IAF can be generalized for any $\mu > 0$ in a straightforward manner. The representations will remain the same, only the values of $d_I^{1,1}$ ($I = 1, 2, \dots, \mu$) would change where $d_I^{1,1}$ is the solution of Eq. (17).

5.2 Partition function for special initial configuration

In the totally asymmetric case, for general μ , we have considered the specific initial configuration Eq. (26) which leads us to acquire analytical expressions for observables of interest. We choose a particular case of Eq. (26), namely the $\bar{\rho} = 0$ case, as the special initial configuration for analytical calculation in μ -ASEP-IAF. More precisely, the choice of our special initial configuration for μ -ASEP-IAF is,

$$C(0) \equiv \prod_{i=1}^{N_+/ \mu} D_1 A \prod_{i=1}^{N_+/ \mu} D_2 A \cdots \prod_{i=1}^{N_+/ \mu} D_\mu A \prod_{i=1}^{N_0} E. \quad (45)$$

The initial configuration Eq. (45) is chosen in a way that fixes the total impurity density to ρ_+ . In comparison to Eq. (26), the initial configuration in Eq. (45) is simpler and does not contain consecutive D_I -s. We would see shortly that this specific choice is sufficient to show negative differential mobility in μ -ASEP-IAF. We find the partition function in the steady state under the periodic boundary condition corresponding to the initial configuration Eq. (45), to be

$$Q_{N_+}(z_0) = \left(\left[\sum_{I=1}^{\mu} \frac{d_I}{1 - \frac{z_0}{p_I} - \frac{q_I z_0}{p_I \epsilon}} \right] \left(\frac{1}{1 - \frac{z_0}{\epsilon}} \right) \right)^{N_+}. \quad (46)$$

We have used short hand notations $d_I \equiv d_I^{1,1}$ which are essentially the solutions of Eq. (17). When $q_I = 0$ for all species, it is straightforward to check that the partition function in Eq. (46) reduces to the partition function Eq. (27) of the totally asymmetric case, under the condition $\bar{\rho} = 0$.

5.3 Species densities, drift current and flip current

Just like we did in the μ -TASEP-IAF, we can analytically calculate several observables of interest in the partially asymmetric case also, using the matrix algebra (42) and matrix representations (44) following the same procedures as before. Starting from the initial configuration stated in Eq. (45), the average density ρ_I of any non-conserved species I ($I = 1, \dots, \mu$) for the μ -ASEP-IAF is obtained as

$$\rho_I = \rho_+ \frac{d_I}{\left(1 - \frac{z_0}{p_I} - \frac{q_I z_0}{p_I \epsilon}\right)} \frac{1}{\sum_{K=1}^{\mu} \frac{d_K}{\left(1 - \frac{z_0}{p_K} - \frac{q_K z_0}{p_K \epsilon}\right)}}, \quad (47)$$

where ρ_0 and ρ_+ are the conserved densities for the vacancies and the impurities, respectively. If we put $q_I = 0$ for all I in Eq. (47), the expression of ρ_I for the totally asymmetric case [Eq. (31)] is correctly recovered. The drift currents J_{I0} and the flip currents $J_{I\leftrightarrow K}$ for the μ -ASEP-IAF ($I, K = 1, \dots, \mu$) are given by

$$\begin{aligned} J_{I0} &= z_0 \rho_+ \frac{d_I}{\left(1 - \frac{z_0}{p_I} - \frac{q_I z_0}{p_I \epsilon}\right)} \frac{1}{\sum_{K=1}^{\mu} \frac{d_K}{\left(1 - \frac{z_0}{p_K} - \frac{q_K z_0}{p_K \epsilon}\right)}}, \\ J_{I\leftrightarrow K} &= \rho_+ \frac{(d_I w_{IK} - d_K w_{KI})}{\sum_{K=1}^{\mu} \frac{d_K}{\left(1 - \frac{z_0}{p_K} - \frac{q_K z_0}{p_K \epsilon}\right)}}. \end{aligned} \quad (48)$$

5.4 Negative differential mobility

In what follows, we will show that the species in the μ -ASEP-IAF under the periodic boundary condition, exhibit negative differential mobility [69, 70]. More precisely, we would see that both the drift currents and the flip current can decrease with increasing bias (which we define later), giving rise to the phenomena of negative differential mobility (NDM). NDM has been observed for driven tracer particles in the presence of static obstacles [72, 73] or in crowded medium [71] and for many particle systems in presence of kinetic constraints [74] or obstacles [75]. There have been many studies to understand the mechanism of NDM in driven systems and it appears that some kind of trapping that leads to decrease in dynamical activity, acts as a main cause of NDM [73, 76, 77]. Recently in Ref. [78], the authors have proposed that slowing down of non-driven degrees of freedom (modes) through the biasing of the driven mode, can give rise to negative differential mobility for both the driven and non-driven degrees of freedom in an interacting many particle system. Here, we apply this mechanism to show that indeed the μ -ASEP-IAF can exhibit NDM for particular choices of the rates in the microscopic dynamics.

To illustrate NDM in μ -ASEP-IAF, we will focus on the $\mu = 3$ case. We have three species of particles ($I = 1, 2, 3$), impurities (+) and vacancies in the system following the microscopic dynamics Eq. (41). We choose the drift rates p_I and q_I of the species $I = 1, 2, 3$ to be

$$p_1 = 1, q_1 = e^{-b}, \quad p_2 = \frac{1}{1+b^2} = q_2, \quad p_3 = 1 = q_3. \quad (49)$$

The choices of the hopping rates in Eq. (49) are inspired by similar choices in Ref. [78] in context of NDM for different models. The special choices of the hopping rates in Eq. (49) allow us to identify the parameter b as the hopping bias in the system. This is because $\ln(p_1/q_1) = b$ and the unbiased case $p_1 = q_1 = 1$ corresponds to $b = 0$. Then, the species 1 particle is a driven mode for any $b > 0$. Even when $b > 0$, Eq. (49) clearly states that species 2 and species 3 particles are non-driven modes in the system because the right and left hopping rates are equal for both of them. However, there is a key difference between the hopping rates of species 2 and species 3 particles. The hopping rates of species 2 depend explicitly on the bias b of the driven mode (species 1). More precisely, the hopping rates p_2 and q_2 decrease with increasing bias b . This corresponds to the slowing down of non-driven mode and turns out to be the key for negative differential mobility. The hopping rates p_3 and q_3 of the other non-driven mode species 3, does not depend on b . Here we should mention the presence of another driven mode in the system, which is the impurity. Since, the impurity motion is only unidirectional, it is a driven mode by construction. Consequently, even at $b = 0$ the system is




Figure 9: The figure illustrates the negative differential mobility of the currents in the 3-ASEP-IAF. In (a), the drift current of the species 1 (driven by bias b) monotonically decreases with increasing bias. The current of the other driven mode (driven due to unidirectional motion with rate ϵ) impurity (+) shows non-monotonic behavior, it increases initially but ultimately decreases with increasing bias for large values of b . In (b), the drift current of both non-driven modes (species 2 and species 3) decrease with increasing bias b for large values of the bias. The figure (c) shows that the flip current between any pair of species also exhibits negative differential mobility with increasing b . Although the flip current is not related to the drift bias b directly, still it decreases with increasing bias for large values of b . The parameters used here are $L = 10^3$, $\epsilon = 0.1$, $p_3 = q_3 = 1.0$, $w_{31} = w_{12} = 0.8$, $w_{13} = w_{32} = 0.2$, $w_{21} = w_{23} = 0.5$, $\rho_+ = 0.3$, $\rho_0 = 0.4$. The ensemble average is done over 10^6 samples.

in a non-equilibrium steady state for $\epsilon > 0$ where the impurity acts as the lone driven mode. In the present context, we choose ϵ to be a constant independent of the bias b . We focus on the behavior of the currents with the variation of b . To summarize, the driven modes in the 3-ASEP-IAF are (i) species 1 (driven by b) and (ii) impurity (driven due to unidirectional motion with constant rate ϵ , independent of b), whereas the non-driven modes are (i) species 2 (hopping rates are decreasing function of b) and (ii) species 3 (hopping rates independent of b). All the flip rates w_{IK} ($I, K = 1, 2, 3$) are kept constants independent of the drift bias b . With this set up, we now investigate the variation of the drift currents and flip current as functions of the bias b , both from analytical formulae Eq. (48) and Monte Carlo simulations.

In Fig. 9(a), we present the behaviors of the drift currents of the driven modes with variation of the bias b , under the periodic boundary condition. Interestingly, although the bias b is directly applied to species 1 to increase its current, the drift current for species 1 decreases monotonically with increasing bias giving rise to the phenomena of negative differential mobility. The current of the other driven mode, the impurities, initially increase with increasing bias, reaches to a maximum, but then decreases as the bias is further increased, thereby leading to NDM. The drift currents of both the non-driven modes exhibit non-monotonic behaviors with increasing bias as shown in Fig. 9(b). Both of them decrease with increasing bias for sufficiently large values of b , showing negative differential mobility. Notably, the flip dynamics is not directly affected by the drift bias since all the flip rates are kept to be constants independent of b . Therefore, it is intriguing to observe that the net flip current still decreases with increasing bias (for large b) and therefore exhibits NDM, as presented in Fig. 9(c). The mechanism behind the negative differential mobility in drift current is related to the decreasing dynamical activity (number of hops per unit time) of the species 2 particles (one of the non-driven modes) with increasing forward bias b for the species 1 particles. Since with increasing b , the hop rate of species 2 [Eq. (49)] decreases, its waiting time at the residing site increases i.e. it becomes more and more prone to stay at the residing site rather to leave the site as b increases. That is why, although the increasing bias tries to push particles forward, their ways are blocked by the slowed down species 2 particles. The exclusion interaction and the non-overtaking dynamics facilitates the NDM even better by not allowing other species or impurities to overtake the slowed down species 2 particles. The reason behind the negative differential mobility occurring in the flip current requires further investigation.

We end this section with mentioning the possibility of further nontrivial transport properties in the steady state of the μ -ASEP-IAF when one considers the counter flow scenario. The counter flow in the system arises when the net bias of some species of particles are opposite to that of the others. For example, in the 2-ASEP-IAF, species 1 can have net bias to right i.e. $p_1 > q_1$ whereas the species 2 can have net bias in the opposite direction i.e. $q_2 > p_2$. Counter flow can give rise to interesting physical features e.g. phase transitions [45, 49, 79]. This urges for detailed investigation of the counter flow situation in μ -ASEP-IAF in future works.

6 Summary and future directions

In this article, we have obtained an exact steady state probability distribution of the μ -ASEP-IAF on a one dimensional lattice under periodic boundary conditions, using the ma-

trix product ansatz. The μ -ASEP-IAF consists of (i) drift of the species ($I = 1, 2, \dots, \mu$) and impurities, and (ii) flip between different species initiated by the impurities. In steady state, we provide the explicit finite dimensional $[(\mu + 1) \times (\mu + 1)]$ matrix representations for any $\mu > 0$ for the totally asymmetric case i.e. μ -TASEP-IAF. For the partially asymmetric scenario i.e. μ -ASEP-IAF, we obtain the corresponding matrices with infinite dimensional representations. Importantly, due to the non-ergodicity of the μ -ASEP-IAF dynamics, the partition function and observables in the steady state depend on the specific choice of the initial configuration. However, for a special class of initial configurations, we could indeed analytically calculate the partition function for both the totally asymmetric and partially asymmetric cases with any $\mu > 0$, under periodic boundary conditions. We present exact analytical expressions for steady state observables like the average densities of the non-conserved species, drift current, flip current and some other two-point correlations. We show that our analytical calculations are in agreement with the Monte Carlo simulations for the analytically tractable specific initial configuration. In this connection, the non-ergodicity of the model has been established extensively (Monte Carlo simulations) by showing the deviations of the steady state observable values for a random initial configuration from that of the specific initial configuration mentioned above. Along with the important exactly solvable analytical structure, the μ -ASEP-IAF also has interesting physical features. Notably, with the variation of vacancy density, several two-point correlations exhibit interesting behaviors e.g. transiting between negative and positive correlations, showing both local maximum and local minimum etc. The effect of the drift on the flip processes are evident from the functional dependence of the species densities on the flip rates. Interestingly, both the drift current and flip current in the μ -ASEP-IAF are shown (analytically and numerically) to display negative differential mobility (decreasing current with increasing bias) for certain choices of the drift rates. The mechanism behind the negative differential mobility relies on slowing down a non-driven mode in the system through the biasing of a driven mode, which eventually leads to decreased dynamical activity of all the modes in the steady state.

Apart from its own intriguing mathematical and physical characteristics, the μ -ASEP-IAF studied here is relevant in two other important contexts. The μ -ASEP-IAF has interesting connections to (i) multi lane asymmetric simple exclusion process (m -ASEP) which serves as a simple yet remarkable model for multi lane traffic flow, and (ii) enzymatic chemical reactions. For the totally asymmetric model μ -TASEP-IAF, these connections are discussed in details in Appendix A and Appendix C respectively. Importantly, the exact solution of μ -ASEP-IAF suggests possible exact solutions in corresponding multi lane ASEP and traffic models with correlations between particles in different lanes and non-zero net current between lanes. The detailed and rigorous analysis to develop these connections and incorporate them for studying multi-lane traffic flow, constitutes one of the main future directions. We also propose a variation of the μ -ASEP-IAF that exhibits better prospects for being a model for multi lane traffic flow (see Appendix B). In future we plan to investigate the exact steady state and observables of this varied μ -ASEP-IAF model using matrix product ansatz. Just like the multi lane traffic flow, the connections between μ -ASEP-IAF and enzymatic chemical reactions both in steady state as well as dynamics, should be analyzed in more details by considering observables relevant for the chemical reactions. In the present article, we have considered impurities with fixed finite density and constant hopping rate. It would be useful to explore the effects of the variations of impurity density and impurity hopping rate on the observables in the μ -ASEP-IAF. Another important future direction would be to analyze the exact steady state of the μ -ASEP-IAF with open boundary conditions which is more pertinent

in context of transport processes, and might also lead to rich phase transitions. It would be interesting to look into the effect of counter flow (i.e. some species having net bias in opposite direction relative to the other species) on the transport properties and possibility of phase separations in the μ -ASEP-IAF. We would also like to investigate the dynamics of the μ -ASEP-IAF in detail, in particular if the product form of the steady state also prevails in the dynamics (using dynamical matrix product ansatz), the dynamical activity in terms of large deviations and possibility of dynamical phase transitions in related models.

Acknowledgements

We thank Kazuaki Takasan for fruitful discussions. We gratefully acknowledge Arvind Ayyer for pointing out important references. This work is partially supported by the Grants-in-Aid for Scientific Research (Grant No. 21H01006).

A Connection between μ -TASEP-IAF and multi lane TASEP

Here we explore the connections between the multi lane totally asymmetric simple exclusion process (m -TASEP) and the one dimensional μ -ASEP-IAF. For simplicity, we consider $\mu = 2$ i.e. the 2-TASEP-IAF (and correspondingly 2-TASEP or two lane TASEP).

In Fig. 10, we present a two lane TASEP where particles can hop in forward directions in lane 1 and lane 2 with rates p_1 and p_2 respectively. The particles in lane 1 and lane 2 can be interpreted as two types i.e. species 1 and species 2 particles in the 2-TASEP-IAF process. Except for the hopping of particles in the two lanes in the 2-TASEP, the




Figure 10: The figure illustrates a two lane totally asymmetric simple exclusion process and the identification of its components to the equivalent constituents of the 2-TASEP-IAF.

particles can change lanes through bridges connecting the lanes. There are two types of bridges, *active* (green vertical lines in Fig. 10) that allows vertical hopping i.e. lane change of

particles and *inactive* (red vertical lines in Fig. 10) that does not allow lane change of particles. The active bridges in the two lane TASEP mimic the impurities (+) in the 2-TASEP-IAF. However, the inactive bridges are not counted in the equivalent 2-TASEP-IAF. Notably, a neighboring pair of (active, inactive) bridges can change to (inactive, active). This inactive-active transformation of neighboring lanes can be interpreted as a resultant drift of the active bridges through the system. Consequently, this accounts for the forward hopping of impurity in the 2-TASEP-IAF.




Figure 11: The figure shows the connection between each microscopic dynamics of the two lane TASEP with the equivalent microscopic dynamics in the 2-TASEP-IAF.

The equivalence of the microscopic dynamics of the two lane TASEP and the 2-TASEP-IAF is shown in Fig. 11. The last (bottom) panel in Fig. 11 exhibits the connection between the inactive-active lane transformations in 2-TASEP and the impurity hopping in the 2-TASEP-IAF. The two panels above the bottom panel in Fig. 11 describe the equivalence of the lane change of particles in 2-TASEP with the impurity activated flip in the corresponding 2-TASEP-IAF. To elaborate, when the lane 1 (2) particle in 2-TASEP comes in contact with an active bridge, it can go to lane 2 (1) with rate w_{12} (w_{21}). Similarly, when a species 1 (2) particle in the 2-TASEP-IAF encounters an impurity as a right neighbor, it can flip to a species 2 (1) particle with rate w_{12} (w_{21}). The first two panels in Fig. 11 present the relations between usual hopping dynamics in the two processes.

We can generalize the approaches described above to establish connections between the multi-lane TASEP and the μ -TASEP-IAF for any $\mu \geq 3$. It is noteworthy that, for the multi-species case ($\mu \geq 3$), we have shown the existence of non-zero net flip current in the μ -TASEP-IAF. It implies the existence of net non-zero lane change current between neighboring lanes in the multi lane TASEP. Also, the correlations between different species of particles and vacancies in the μ -TASEP-IAF suggest non-zero correlations between particles in the different lanes in the m -TASEP. The mapping can also be extended for the partially asymmetric motion of particles. We must mention that the connections between the multi-lane TASEP and μ -TASEP-IAF described here, are approximate. To establish more accurate relations between the two processes, one has to perform rigorous calculations for observables in the multi-lane TASEP and compare the corresponding results with that of the μ -TASEP-IAF.




Figure 12: The figure shows the variation of the total drift current of species 1 and 2 with the flip rate w_{12} . While the flip rate (equivalent lane change rate in two lane TASEP) is varied over a large range $w_{12} \in (0, 1)$, the corresponding increase in the total drift current (total flow along the two lanes in TASEP) is reasonably small. The parameters used are $L = 10^4$, $p_1 = 0.3$, $p_2 = 1.0$, $\epsilon = 0.1$, $w_{21} = 1.0$, $\rho_+ = 0.2$, $\rho_0 = 0.2$. The ensemble average is done over 10^7 samples.

B A variation of μ -TASEP-IAF, connection to multi-lane traffic flow

The multi lane TASEP has been widely regarded as a simplistic yet important model for multi lane traffic flow [2]. Due to the connections between the μ -TASEP-IAF and the multi-lane TASEP discussed in Appendix A, it is natural to ask about the applicability of μ -TASEP-IAF [Eq. (1)] as a suitable model for multi lane traffic flow. Before addressing this question, we should mention that the lane change dynamics in realistic traffic flow must facilitate the traffic as a whole. More precisely, the change of lanes should increase the total flow or total current along the lanes. To investigate this for the μ -TASEP-IAF with $\mu = 2$, we plot the total drift current of species 1 and species 2, $J_{\text{total}} = J_{10} + J_{20}$ as a function of the flip rate w_{12} in Fig. 12. Of course, for the two lane TASEP, this amounts to investigating the variation of the total drift current of lane 1 and lane 2 by changing the lane change rate w_{12} .

In Fig. 12 we observe that the total current, although increases with the flip rate, the amplitude of the increment is quite small keeping in mind the wide range of variation in the tuning parameter $w_{12} \in (0, 1)$. The reason behind this, as revealed by a careful observation, is the *approximate* mapping between the μ -TASEP-IAF and multi lane TASEP described in Fig. 11. In the μ -TASEP-IAF dynamics, when a species encounters an impurity, it flips but does not change its position. On the other hand, in the multi lane TASEP, when a particle in any lane comes in contact with an active bridge, it actually changes the lane i.e. not only changes its characteristics (lane 1 particle to lane 2 particle or vice versa) but also changes its position. To incorporate this in our present model, we propose a variation of the μ -TASEP-IAF dynamics in Eq. (1). Specifically, the change is made only in the flip dynamics. Earlier in Eq. (1), the flip dynamics has been

$$I+ \xrightarrow{w_{IK}} K+, \quad (50)$$





Figure 13: The figure illustrates the comparison of the total drift current J_{total} between the flip process in Eq. (50) and the proposed variation in Eq. (51). The current J_{total} for the variation Eq. (51) increases considerably with increasing flip rate (equivalent lane change rate in two lane TASEP) w_{12} , whereas the same for the original flip process Eq. (50) increase much slowly (Fig. 12). The parameters used are $L = 10^4$, $p_1 = 0.3$, $p_2 = 1.0$, $\epsilon = 0.1$, $w_{21} = 1.0$, $\rho_+ = 0.2$, $\rho_0 = 0.2$. The ensemble average is done over 10^7 samples.

with $I, K = 1, \dots, \mu$ and $I \neq K$. Whereas we propose the new flip dynamics to be



Note that, in comparison to Eq. (50), the flip process in Eq. (51) accompanies the flip of the species with a hop towards right. The other hopping processes in Eq. (1) remain the same for this varied μ -TASEP-IAF. The effect of the dynamics can be immediately observed in Fig. 13 where we present the variation of the total drift current as a function of the flip rate (lane change rate) for both the dynamics in Eq. (50) and Eq. (51) (using Monte Carlo simulations). Indeed, the Fig. 13 shows that J_{total} increases considerably with increasing lane change rate for Eq. (51) whereas it grows weakly for Eq. (50) (see Fig. 12). This observation implies that the proposed variation of the μ -TASEP-IAF acts as a better model for multi lane traffic flow in comparison to the original model. It would be interesting to study this variation of the μ -TASEP-IAF both analytically and numerically and to build connections with the multi lane traffic flow.

C Connection between μ -TASEP-IAF and enzymatic chemical reactions


In this appendix, we briefly discuss some connections between the μ -TASEP-IAF and enzymatic chemical reactions. One of the simplest form of the enzymatic chemical reaction is,



<i>Enzymatic chemical reaction:</i>	\equiv	<i>2-TASEP-IAF</i>
Substrate: S	\equiv	Species: 1
Product: P	\equiv	Species: 2
Enzyme: E	\equiv	Impurity: +
Substrate diffusion: $S \xrightarrow{p_1}$	\equiv	$10 \xrightarrow{p_1} 01$
Product diffusion: $P \xrightarrow{p_2}$	\equiv	$20 \xrightarrow{p_2} 02$
Enzyme diffusion: $E \xrightarrow{\epsilon}$	\equiv	$+0 \xrightarrow{\epsilon} 0+$
Forward reaction: $S + E \xrightarrow{k_S} P + E$	\equiv	$1+ \xrightarrow{w_{12}} 2+$
Backward reaction: $P + E \xrightarrow{k_P} S + E$	\equiv	$2+ \xrightarrow{w_{21}} 1+$

Figure 14: The figure illustrates the connection between the simplified form of the enzymatic chemical reaction Eq. (53) (in a narrow channel with many units of drifting enzymes, substrates and products) and the 2-TASEP-IAF. The enzyme, substrate and product in the chemical system can be identified as impurity, species 1 and species 2 particle in the 2-TASEP-IAF.

where E, S, P denotes enzyme, substrate, product respectively and ES corresponds to the intermediate complex. The parameters k_f and k_b are the rate constants for forward and backward reaction for the intermediate complex formation, while \bar{k}_f and \bar{k}_b are the rate constants for forward and backward reactions between the intermediate complex and the product (along with enzyme). Clearly, the initially present enzyme in the reaction remains intact after the reaction is completed. Here, to discuss some connections to the μ -ASEP-IAF, we would rather consider a much simplified version of the chemical reaction (52) as



where we ignore the intermediate complex formation. k_S (k_P) is the rate constant for S transforming to P (P transforming to S).

Let us consider a spatially extended narrow channel with many units of substrates, enzymes and products all of which drift through the channel at different rates. This system of chemical reagents can be approximately mapped to an equivalent μ -TASEP-IAF. As explained in Fig. 14 for $\mu = 2$, the impurity in 2-ASEP-IAF plays the role of the enzyme, as it can transform one species of particle to another species. One species, e.g. species 1 can be considered as S for chemical reaction Eq. (53), whereas the species 2 particle acts as P . The flip rates w_{12} and w_{21} mimic the rate constants k_S and k_P . With this set up, our study of the 2-TASEP-IAF reveals the effect of drift on the resultant concentrations of substrates and products in the steady state. Notably, the multi-species ($\mu > 2$) case of the μ -TASEP-IAF can be mapped to generalized version of the chemical reaction Eq. (53) as



where $i = 1, \dots, \mu_1$ are identified as μ_1 number of substrates and $j = 1, \dots, \mu_2$ are identified as μ_2 number of products in the enzymatic chemical reaction system (with E acting as the enzyme for each chemical reaction) where $\mu_1 + \mu_2 = \mu$ (μ being the total number of species in the μ -TASEP-IAF). To study the time evolution of the enzymatic chemical reaction, one has to study the dynamics of the μ -TASEP-IAF. The connection between the μ -TASEP-IAF and the enzymatic chemical reactions should be studied more thoroughly with proper attention to the observables of interest for the chemical reactions.

References

- [1] N. G. van Kampen, 1992, *Stochastic Processes in Physics and Chemistry* (Elsevier, North-Holland).
- [2] A. Schadschneider, D. Chowdury and K. Nishinari, 2011, *Stochastic transport in complex system: From molecules to vehicles* (Elsevier, Amstrdam).
- [3] P. Krapivsky, S. Redner and E. Ben-Naim , 2010, *A Kinetic View of Statistical Physics* (Cambridge: Cambridge University Press).
- [4] E. S. Allman and J. A. Rhodes , 2004, *Mathematical models in biology : an introduction* (Cambridge: Cambridge University Press).
- [5] R. A. Blythe and A. J. McKane, *Stochastic models of evolution in genetics, ecology and linguistics*, J. Stat. Mech. P07018 (2007), doi:10.1088/1742-5468/2007/07/P07018.
- [6] C. Castellano, S. Fortunato and V. Loreto, *Statistical physics of social dynamics*, Rev. Mod. Phys. **81**, 591 (2009), doi:10.1103/RevModPhys.81.591.
- [7] H. Qian and L. M. Bishop, *The Chemical Master Equation Approach to Nonequilibrium Steady-State of Open Biochemical Systems: Linear Single-Molecule Enzyme Kinetics and Nonlinear Biochemical Reaction Networks*, Int. J. Mol. Sci. **11**, 3472 (2010), doi:10.3390/ijms11093472.
- [8] B. Schmittmann and R. K. P. Zia, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. Lebowitz (Academic, London, 1994), Vol. 17.
- [9] V. Privman, 1997, *Nonequilibrium Statistical Mechanics in One Dimension* (Cambridge University Press).
- [10] F. Spitzer, *Interaction of Markov processes*, Adv. in Math. **5**, 246 (1970), doi:10.1016/0001-8708(70)90034-4.
- [11] T. M. Liggett, 1999, *Stochastic Models of Interacting Systems: Contact, Voter and Exclusion Processes*, (Springer-Verlag, New-York).
- [12] G.M. Schütz, in C.Domb and J.Lebowitz (eds.) *Phase Transitions and Critical Phenomena*, Vol.19 (Academic, London, 2001).
- [13] D. Helbing, *Traffic and related self-driven many-particle systems*, Rev. Mod. Phys. **73**, 1067 (2001), doi:10.1103/RevModPhys.73.1067.

- [14] C. T. MacDonald and J. H. Gibbs, *Concerning the kinetics of polypeptide synthesis on polyribosomes*, Biopolymers **7**, 707 (1969), doi:10.1002/bip.1969.360070508.
- [15] K. Nishinari, Y. Okada, A. Schadschneider and D. Chowdury, *Intracellular Transport of Single-Headed Molecular Motors KIF1A*, Phys. Rev. Lett. **95**, 118101 (2005), doi:10.1103/PhysRevLett.95.118101.
- [16] C. Leduc, K. Padberg-Gehle, V. Varga, D. Helbing, S. Diez and J. Howard, *Molecular crowding creates traffic jams of kinesin motors on microtubules*, Proc. Natl Acad. Sci. USA **109**, 6100 (2012), doi:10.1073/pnas.1107281109.
- [17] B. Derrida, *An exactly soluble non-equilibrium system: The asymmetric simple exclusion process*, Phys. Rep. **301**, 65 (1998), doi:10.1016/S0370-1573(98)00006-4.
- [18] T. Sasamoto, *One-dimensional partially asymmetric simple exclusion process with open boundaries: orthogonal polynomials approach*, J. Phys. A: Math. Gen. **32**, 7109 (1999), doi:10.1088/0305-4470/32/41/306.
- [19] O. Golinelli and K. Mallick, *The asymmetric simple exclusion process: an integrable model for non-equilibrium statistical mechanics*, J. Phys. A: Math. Gen. **39**, 12679 (2006), doi:10.1088/0305-4470/39/41/S03.
- [20] K. Mallick, *The exclusion process: A paradigm for non-equilibrium behaviour*, Physica A **418**, 17 (2015), doi:10.1016/j.physa.2014.07.046.
- [21] B. Derrida, M. R. Evans, V. Hakim and V. Pasquier, *Exact solution of a 1D asymmetric exclusion model using a matrix formulation*, J. Phys. A: Math. Gen. **26**, 1493 (1993), doi:10.1088/0305-4470/26/7/011.
- [22] M. Uchiyama, T. Sasamoto and M. Wadati, *Asymmetric simple exclusion process with open boundaries and Askey–Wilson polynomials*, J. Phys. A: Math. Gen. **37**, 4985 (2004), doi:10.1088/0305-4470/37/18/006.
- [23] M. R. Evans, P. A. Ferrari and K. Mallick, *Matrix Representation of the Stationary Measure for the Multispecies TASEP*, J. Stat. Phys. **135**, 217 (2009), doi:10.1007/s10955-009-9696-2.
- [24] R. A. Blythe and M. R. Evans, *Nonequilibrium steady states of matrix-product form: a solver's guide*, J. Phys. A: Math. Theor. **40**, R333 (2007), doi:10.1088/1751-8113/40/46/R01.
- [25] P. A. Ferrari and J. B. Martin, *Stationary distributions of multi-type totally asymmetric exclusion processes*, Ann. Probab. **35**(3), 807 (2007), doi:10.1214/009117906000000944.
- [26] A. Kuniba, S. Maruyama and M. Okado, *Multispecies TASEP and combinatorial R^** , J. Phys. A: Math. Theor. **48**, 34FT02 (2015), doi:10.1088/1751-8113/48/34/34FT02.
- [27] A. Ayyer and S. Linusson, *Correlations in the Multispecies TASEP and a Conjecture by Lam*, Trans. Amer. Math. Soc. **369**, 1097 (2017), doi:10.1090/tran/6806.
- [28] A. Ayyer and S. Linusson, *An Inhomogeneous Multispecies TASEP on a Ring*, Advances in Applied Mathematics **57**, 21 (2014), doi:https://doi.org/10.1016/j.aam.2014.02.001.

- [29] E. Lee, *Integrability of the Multi-Species TASEP with Species-Dependent Rates*, *Symmetry* **13**(9), 1578 (2021), doi:10.3390/sym13091578.
- [30] N. Crampe, C. Finn, E. Ragoucy and M. Vanicat, *Integrable boundary conditions for multi-species ASEP*, *Phys. A: Math. Theor.* **49**, 375201 (2016), doi:10.1088/1751-8113/49/37/375201.
- [31] C. Finn, E. Ragoucy and M. Vanicat, *Matrix product solution to multi-species ASEP with open boundaries*, *J. Stat. Mech.* 043201 (2018), doi:10.1088/1742-5468/aab1b5.
- [32] S. Sandow, *Partially asymmetric exclusion process with open boundaries*, *Phys. Rev. E* **50**, 2660 (1994), doi:10.1103/PhysRevE.50.2660.
- [33] N. Crampé, *Algebraic Bethe ansatz for the totally asymmetric simple exclusion process with boundaries*, *J. Phys. A: Math. Theor.* **48**, 08FT01 (2015), doi:10.1088/1751-8113/48/8/08FT01.
- [34] J. de Gier and F. Essler, *Bethe Ansatz Solution of the Asymmetric Exclusion Process with Open Boundaries*, *Phys. Rev. Lett.* **95**, 240601 (2005), doi:10.1103/PhysRevLett.95.240601.
- [35] M. Kardar, G. Parisi and Y-C. Zhang, *Dynamic Scaling of Growing Interfaces*, *Phys. Rev. Lett.* **56**, 889 (1986), doi:10.1103/PhysRevLett.56.889.
- [36] L. H. Gwa and H. Spohn, *Bethe solution for the dynamical-scaling exponent of the noisy Burgers equation*, *Phys. Rev. A* **46**, 844 (1992), doi:10.1103/PhysRevA.46.844.
- [37] T. Sasamoto and H. Spohn, *One-Dimensional Kardar-Parisi-Zhang Equation: An Exact Solution and its Universality*, *Phys. Rev. Lett.* **104**, 230602 (2010), doi:10.1103/PhysRevLett.104.230602.
- [38] A. Parmeggiani, T. Franosch and E. Frey, *Phase Coexistence in Driven One-Dimensional Transport*, *Phys. Rev. Lett.* **90**, 086601 (2003), doi:10.1103/PhysRevLett.90.086601.
- [39] E. Pronina and A. B. Kolomeisky, *Two-channel totally asymmetric simple exclusion processes*, *J. Phys. A: Math. Gen.* **37**, 9907 (2004), doi:10.1088/0305-4470/37/42/005.
- [40] T. Mitsudo and H. Hayakawa, *Synchronization of kinks in the two-lane totally asymmetric simple exclusion process with open boundary conditions*, *J. Phys. A: Math. Gen.* **38**, 3087 (2005), doi:10.1088/0305-4470/38/14/002.
- [41] R. Jiang, M. Hu, Y. Wu and Q. Wu, *Weak and strong coupling in a two-lane asymmetric exclusion process*, *Phys. Rev. E* **77**, 041128 (2008), doi:10.1103/PhysRevE.77.041128.
- [42] H. J. Hilhorst and C. Appert-Rolland, *A multi-lane TASEP model for crossing pedestrian traffic flows*, *J. Stat. Mech.* P06009 (2012), doi:10.1088/1742-5468/2012/06/P06009.
- [43] Y. Wang, R. Jiang and Q. Wu, *Dynamics in phase transitions of TASEP coupled with multi-lane SEPs*, *Nonlinear Dyn* **88**, 1631 (2017), doi:10.1007/s11071-017-3335-2.
- [44] P. Helms, U. Ray and G. K.-L. Chan, *Dynamical phase behavior of the single- and multi-lane asymmetric simple exclusion process via matrix product states*, *Phys. Rev. E* **100**, 022101 (2019), doi:10.1103/PhysRevE.100.022101.

- [45] Q.-Y. Hao, R. Jiang, M.-B. Hu, Y. Zhang, C.-Y. Wu and N. Guo, *Theoretical analysis and simulation of phase separation in a driven bidirectional two-lane system*, Phys. Rev. E **100**, 032133 (2019), doi:10.1103/PhysRevE.100.032133.
- [46] I. Neri, N. Kern and A. Permezziani, *Totally Asymmetric Simple Exclusion Process on Networks*, Phys. Rev. Lett. **107**, 068702 (2011), doi:10.1103/PhysRevLett.107.068702.
- [47] K. Mallick, *Shocks in the asymmetry exclusion model with an impurity*, J. Phys. A: Math. Gen. **29**, 5375 (1996), doi:10.1088/0305-4470/29/17/013.
- [48] F. H. Jafarpour, *Partially Asymmetric Simple Exclusion Model in the Presence of an Impurity on a Ring*, J. Phys. A: Math. Gen. **33**, 1797 (2000), doi:10.1088/0305-4470/33/9/306.
- [49] F. H. Jafarpour, *Exact solution of an exclusion model in the presence of a moving impurity on a ring*, J. Phys. A: Math. Gen. **33**, 8673 (2000), doi:10.1088/0305-4470/33/48/307.
- [50] B. Derrida and M. R. Evans, *Bethe ansatz solution for a defect particle in the asymmetric exclusion process*, J. Phys. A: Math. Gen. **32**, 4833 (1999), doi:10.1088/0305-4470/32/26/303.
- [51] C. Boutillier, P. Francois, K. Mallick and S. Mallick, *Amatrixansatz for the diffusion of an impurity in the asymmetric exclusion process*, J. Phys. A: Math. Gen. **35**, 9703 (2002), doi:10.1088/0305-4470/35/46/301.
- [52] M. R. Evans, *Bose-Einstein condensation in disordered exclusion models and relation to traffic flow*, Europhys. Lett., **36**, 13 (1996), doi:10.1209/epl/i1996-00180-y.
- [53] M. J. Lazo and A. J. Ferreira, *Asymmetric exclusion model with impurities*, Phys. Rev. E **81**, 050104(R) (2010), doi:10.1103/PhysRevE.81.050104.
- [54] A. B. Kolomeisky, J. Phys. A: Math. Gen. **31**, *Asymmetric simple exclusion model with local inhomogeneity*, 1153 (1998), doi:10.1088/0305-4470/31/4/006.
- [55] S. A. Janowsky and J. L. Lebowitz, *Finite-size effects and shock fluctuations in the asymmetric simple-exclusion process*, Phys. Rev. A **45**, 618 (1992), doi:10.1103/PhysRevA.45.618.
- [56] G. Tripathy and M. Barma, *Driven lattice gases with quenched disorder: Exact results and different macroscopic regimes*, Phys. Rev. E **58**, 1911 (1998), doi:10.1103/PhysRevE.58.1911.
- [57] T. Chou and G. Lakatos, *Clustered Bottlenecks in mRNA Translation and Protein Synthesis*, Phys. Rev. Lett. **93**, 198101 (2004), doi:10.1103/PhysRevLett.93.198101.
- [58] M. Barma, *Driven diffusive systems with disorder*, Physica A **372**, 22 (2006), doi:10.1016/j.physa.2006.05.002.
- [59] J. S. Nossan, *Disordered exclusion process revisited: some exact results in the low-current regime*, J. Phys. A: Math. Theor. **46**, 315001 (2013), doi:10.1088/1751-8113/46/31/315001.

- [60] B. Waclaw, J. Cholewa-Waclaw and P. Greulich, *Totally asymmetric exclusion process with site-wise dynamic disorder*, J. Phys. A: Math. Theor. **52**, 065002 (2019), doi:10.1088/1751-8121/aafb8a.
- [61] U. Basu and P. K. Mohanty, *TASEP on a ring with internal degrees of freedom*, Phys. Rev. E. **82**, 041117 (2010), doi:10.1103/PhysRevE.82.041117.
- [62] S. Zeraati, F.H. Jafarpour and H. Hinrichsen, *Phase transition in an exactly solvable reaction-diffusion process*, Phys. Rev. E. **87**, 062120 (2013), doi:10.1103/PhysRevE.87.062120.
- [63] M. Ghadermazi and F. H. Jafarpour, *A new family of exactly solvable disordered reaction-diffusion systems*, J. Stat. Mech. P09023 (2013), doi:10.1088/1742-5468/2013/09/P09023.
- [64] N. Shiraishi and K. Matsumoto, *Undecidability in quantum thermalization*, Nature Communications **12**, 5084 (2021), doi:10.1038/s41467-021-25053-0.
- [65] K. A. Johnson and R. S. Goody, *The Original Michaelis Constant: Translation of the 1913 Michaelis-Menten Paper*, Biochemistry **50**, 8264 (2011), doi:10.1021/bi201284u.
- [66] M. J. Schnitzer, K. Visscher and S. M. Block, *Force production by single kinesin motors*, Nat. Cell Biol. **2**, 718 (2000), doi:10.1038/35036345.
- [67] R. Grima, *Noise-Induced Breakdown of the Michaelis-Menten Equation in Steady-State Conditions*, Phys. Rev. Lett. **102**, 218103 (2009), doi:10.1103/PhysRevLett.102.218103.
- [68] K. Mallick, S. Mallick and N. Rajewsky, *Exact solution of an exclusion process with three classes of particles and vacancies*, J. Phys. A: Math. Gen. **32**, 8399 (1999), doi:10.1088/0305-4470/32/48/303.
- [69] R. K. P. Zia, E. L. Præstgaard and O. G. Mouritsen, *Getting more from pushing less: Negative specific heat and conductivity in nonequilibrium steady states*, Am. J. Phys. **70**, 384 (2002), doi:10.1119/1.1427088.
- [70] R. Eichhorn, P. Reimann, B. Cleuren and C. Van den Broeck, *Moving backward noisily*, Chaos **15**, 026113 (2005), doi:10.1063/1.1869932.
- [71] O. Bénichou, P. Illien, G. Oshanin, A. Sarracino and R. Voituriez, , Phys. Rev. Lett. **113**, 268002 (2014), doi:10.1103/PhysRevLett.113.268002.
- [72] D. Dhar, *Diffusion and drift on percolation networks in an external field*, J. Phys. A **17**, L257 (1984), doi:10.1088/0305-4470/17/5/007.
- [73] P. Baerts, U. Basu, C. Maes and S. Safaverdi, *Frenetic origin of negative differential response*, Phys. Rev. E **88**, 052109 (2013), doi:10.1103/PhysRevE.88.052109.
- [74] M. Sellitto, *Asymmetric Exclusion Processes with Constrained Dynamics*, Phys. Rev. Lett. **101**, 048301 (2008), doi:10.1103/PhysRevLett.101.048301.
- [75] R. Eichhorn, J. Regtmeier, D. Anselmettib and P. Reimann, *Negative mobility and sorting of colloidal particles*, Soft Matter **6**, 1858 (2010), doi:10.1039/b918716m.

- [76] M. Baiesi, C. Maes and B. Wynants, *Fluctuations and Response of Nonequilibrium States*, Phys. Rev. Lett. **103**, 010602 (2009), doi:10.1103/PhysRevLett.103.010602.
- [77] U. Basu and C. Maes, *Mobility transition in a dynamic environment*, J. Phys. A: Math. Theor. **47**, 255003 (2014), doi:10.1088/1751-8113/47/25/255003.
- [78] A. K. Chatterjee, U. Basu and P. K. Mohanty, *Negative differential mobility in interacting particle systems*, Phys. Rev. E **97**, 052137 (2018), doi:10.1103/PhysRevE.97.052137.
- [79] P. F. Arndt, T. Heinzel and V. Rittenberg, *Spontaneous breaking of translational invariance in one-dimensional stationary states on a ring*, J. Phys. A: Math. Gen. **31**, L45 (1998), doi:10.1088/0305-4470/31/2/001.