dispersive AB system with weighted Sobolev initial data, which allows soliton solutions via the Dbar steepest descent method. Based on the spectral analysis of Lax pair, the Cauchy problem of the coupled dispersive AB system is transformed into a Riemann-Hilbert problem, and its existence and uniqueness of the solution is proved by the vanishing lemma. The stationary phase points play an important role in the long-time asymptotic behavior. We demonstrate that in any fixed time cone $\mathcal{C}(x_1, x_2, v_1, v_2) = \{(x, t) \in \mathbb{R}^2 \mid x = x_0 + vt, x_0 \in [x_1, x_2], v \in [v_1, v_2]\}$, the long-time asymptotic behavior of the solution for the coupled dispersive AB system can be expressed by $N(\mathcal{I})$ solitons on the discrete spectrum, the leading order term $\mathcal{O}(t^{-1/2})$ on the continuous spectrum and the allowable residual $\mathcal{O}(t^{-3/4})$.

arXiv:2205.03179v1 [math.AP] 6 May 2022

1. INTRODUCTION In the past few decades, many solutions of integrable equations have been studied, including soliton solutions, breather solutions, rogue wave solutions and so on [1–5]. These solutions are obtained by classical inverse scattering method, Riemann-Hilbert method under the condition of no reflection [6–8]. However, for continuous spectrum or reflection, it is necessary to analyze the asymptotic behavior of the solution. In fact, as early as 1973, Manakov et al. first studied the long-time asymptotic behavior of the solution of the fast decay initial value problem for nonlinear integrable systems by using the inverse scattering method [9,10]. In 1976, Zakharov and Manakov gave the first term asymptotic expression that the solution of the initial value problem for nonlinear Schrödinger (NLS) equation which depends explicitly on the initial value [11]. In 1981, Its transformed the long-time behavior of the solution for the initial value problem of the NLS equation into the local Riemann-Hilbert problem (RHP) in the small neighborhood of the steady-state phase point by using the single value deformation theory, which provided a set of feasible and

LONG-TIME ASYMPTOTIC BEHAVIOR OF THE COUPLED DISPERSIVE AB SYSTEM IN LOW REGULARITY SPACES

JINYAN ZHU AND YONG CHEN*

ABSTRACT. In this paper, we mainly investigate the long-time asymptotic behavior of the solution for the coupled

strict ways to analyze the long-term behavior of the integrable equation [12]. In 1993, Deift and Zhou developed the nonlinear descent method for solving oscillatory RHP based on the classical descent method, and studied the asymptotic properties of the solution for mKdV equation with decaying initial value [13]. Since then, more and more scholars have paid attention to the nonlinear descent method to study the long-time asymptotic behavior of the solution for the initial value problem of integrable systems, and many equations have been studied by this method [14–21]. Later, McLaughlin and Miller extended the classical Deift-Zhou steepest descent method to Dbar steepest descent method, which was successfully used to study the asymptotic stability of NLS multiple soliton solutions [23–25], and the long-time proximity of KdV equation and NLS equation [26–30]. TheDbar steepest descent method is to transform the discontinuous part on the jump line into the form of Dbar problem, rather than the analysis of the asymptotic properties of orthogonal polynomials in singular integrals on the jump line. Then it became a more powerful tool in the study of long-time asymptotic for integrable equations, such as NLS equation [31], the Hirota equation [32], the Fokas-Lenells equation [33], the modified Camassa-Holm equation [34], the short-pulse equation [35] and so on.

The coupled dispersion AB system [36]

$$A_{xt} - \alpha A - \beta AB = 0,$$

$$B_x + \frac{\gamma}{2} \left(|A|^2 \right)_t = 0,$$
(1.1)

describes the evolution of micro stable or unstable wave packets in baroclinic shear flow through a quasi geostrophic two-layer model on a beta-plane, where $A \equiv A(x,t)$ is the complex function representing the amplitude of the data packet, the real function $B \equiv B(x, t)$ is the measure of the average flow change caused by the baroclinic wave packet, and α represents the critical situation of shear and the real parameter β, γ represent the nonlinear coefficient. And system (1.1) meets the following compatibility condition:

$$\frac{\gamma}{\beta} \left| A_t \right|^2 + B^2 + \frac{2\alpha}{\beta} B = f(t),$$

where f(t) is an integral function of time. If $A(x,t) = 1/\sqrt{\beta\gamma}\psi_x(x,t)$ and $B(x,t) = \pm [\cos(\psi(x,t)) - \alpha]/\beta$ with $\beta\gamma > 0$, system (1.1) can reduce to $\psi_{xt} = \pm \sin \psi$. If $A(x,t) = 1/\sqrt{-\beta\gamma}\psi_x(x,t)$ and $B(x,t) = \pm [\cos(\psi(x,t)) - \alpha]/\beta$ with $\beta\gamma < 0$, system (1.1) can reduce to $\psi_{xt} = \pm \sinh \psi$.

In recent years, many studies have been done for system (1.1). For example, the soliton solutions of system (1.1) are obtained by the inverse scattering method [40]. Kamchatnov and Pavlov found the periodic wave solution of system (1.1) [41]. Guo et al. have studied its classical Darboux transformation(DT) and N-flod DT and obtained breather solutions and multi-soliton solutions [42, 43]. The rogue wave solution of system (1.1) was also obtained by generalized

DT under the condition $\alpha = 0, \beta = \gamma = 1, f(t) = 1$ [44]. In addition, the high-order strange wave and modulation instability of system (1.1) were further given via generalized DT [45]. Recently, Yan et al. studied the long-time asymptotic behavior of solutions whose initial values belong to Schwarz space through the classical Deift-Zhou descent method [47].

In this paper, we mainly study the long-time asymptotic behavior of the solution of system (1) satisfying the following initial value problem

$$A(x,0) = A_0(x) \in H^{1,1}(\mathbb{R}), B(x,0) = B_0(x) \in H^{1,1}(\mathbb{R}),$$

where $H^{1,1}(\mathbb{R})$ is the weighted Sobelev space, its expression is

$$H^{1,1}(\mathbb{R}) = \left\{ f(x) \in L^2(\mathbb{R}) : f'(x), xf(x) \in L^2(\mathbb{R}) \right\}.$$

The main tool used here is the Dbar steepest descent method. Compared with Deift-Zhou steepest descent method, Dbar method avoids estimating Cauchy integral operator in L^P space and rewrites the discontinuous part of RHP into Dbar problem which can be solved by integral equation. Compared with previous studies [47], we consider the initial value in a wider space instead of in Schwarz space and allow the existence of soliton solutions.

The context of this paper is as follows:

In section 2, we analyze the spectrum of the system (1.1) based on Lax pair. The analyticity, symmetry and asymptotic behavior of characteristic function and scattering matrix are studied. In terms of asymptotic behavior, two singular points k = 0 and $k = \infty$ need to be considered. We also prove that when the initial value belongs to weighted Sobelev space, r(k) belongs to $H^{1,1}(\mathbb{R})$.

In section 3, the RHP of system (1.1) is constructed by the piecewise smooth property of matrix M(x,t).

In section 4, we prove the existence and uniqueness of the solution of RHP by using vanishing lemma.

In section 5, by introducing T(k) function, we change M(x,k) into $M^{(1)}(x,k)$ and get a new RHP about $M^{(1)}(x,k)$. This is mainly to allow the jump matrix near the phase points to have two triangular decomposition.

In section 6, the contour is deformed near the phase points, and a mixed $\bar{\partial}$ -RHP is obtained by defining $\mathcal{R}^{(2)}$.

In section 7, we decompose the mixed $\bar{\partial}$ -RHP into a model RHP of $M_{rhp}(k)$ and the pure $\bar{\partial}$ problem of $M^{(3)}(k)$.

In section 8, the RHP about $M_{rhp}(k)$ is studied and the long-time behavior of the soliton solution is analyzed. It is obtained that the soliton solution can be represented by a ring region, and the applicability near the phase points is also explained. Secondly, the error function is calculated with RHP of a small norm.

In section 9, the pure $\bar{\partial}$ problem for $M^{(3)}(k)$ is analyzed.

In section 10, using the above deformation and results, Eq.(10.1) is obtained, and then the long-time behavior of the solution of the coupled dispersion AB system (1.1) in the case of weighted Sobelev initial value is obtained, which is given in the form of theorem.

2. The spectral analysis

System (1.1) has Lax pair is [46]

$$\phi_x = \hat{U}\phi = \begin{pmatrix} -ik & \frac{\sqrt{\beta\gamma}}{2}A \\ -\frac{\sqrt{\beta\gamma}}{2}A^* & ik \end{pmatrix}\phi,$$

$$\phi_t = \hat{V}\phi = \begin{pmatrix} \frac{i(\alpha+\beta B)}{4k} & -\frac{i\sqrt{\beta\gamma}}{4k}A_t \\ -\frac{i\sqrt{\beta\gamma}}{4k}A_t^* & -\frac{i(\alpha+\beta B)}{4k} \end{pmatrix}\phi.$$
(2.1)

In order to better facilitate the presentation of the results, we mainly consider the case of $\beta \gamma = 1$. At this time, the Lax pair of Eq.(1.1) can be simplified to

$$\Phi_x = X\Phi, \ \Phi_t = T\Phi, \tag{2.2}$$

where

$$X = -ik\sigma_3 + U, \ T = \frac{i\alpha}{4k}\sigma_3 + T_1,$$
(2.3)

with

$$\sigma_{3} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad U(x,t) = \frac{1}{2} \begin{pmatrix} 0 & A \\ -A^{*} & 0 \end{pmatrix}, \quad T_{1}(x,t,k) = \frac{i}{4k} \begin{pmatrix} \beta B & -A_{t} \\ -A_{t}^{*} & -\beta B \end{pmatrix}.$$
 (2.4)

The spectral parameter $k \in \mathbb{C}$, under the rapidly decaying initial condition

$$A_0(x), B_0(x) \to 0 \quad \text{as} \quad x \to \infty,$$

make the following correction transformation

$$\Phi = \Psi e^{-ikx\sigma_3 + \frac{i\alpha}{4k}t\sigma_3},\tag{2.5}$$

the modified Lax pair can be obtained

$$\Psi_x + ik [\sigma_3, \Psi] = U\Psi,$$

$$\Psi_t - \frac{i\alpha}{4k} [\sigma_3, \Psi] = T_1 \Psi,$$
(2.6)

3




FIGURE 1. Definition of the $\mathbb{C}^+ = \{k \mid \text{Im } k > 0\}, \ \mathbb{C}^- = \{k \mid \text{Im } k < 0\}$

which can be written in full derivative form

$$d(e^{i\left(kx-\frac{\alpha}{4k}t\right)\widehat{\sigma}_{3}}\Psi(x,t,k)) = e^{i\left(kx-\frac{\alpha}{4k}t\right)\widehat{\sigma}_{3}}[Udx+T_{1}dt)\Psi].$$

From the above Lax pair (2.6), we can see that it has two singular points k = 0 and $k = \infty$. Therefore, we need to investigate the different expansion forms of the characteristic function at two singular points. After simple calculation, the following form is obtained

$$\begin{split} \Psi(x,t,k) &= I + \frac{\Psi^1}{k} + \mathcal{O}\left(k^{-2}\right), \quad k \to \infty, \\ \Psi(x,t,k) &= \Psi^0 + \Psi^1 k + \mathcal{O}\left(k^2\right), \quad k \to 0, \end{split}$$

where

$$\Psi^{0}(x,t) = \begin{pmatrix} e^{\frac{1}{2} \int_{-\infty}^{x} \frac{\beta A B}{A_{t}} dx} & e^{\frac{1}{2} \int_{x}^{\infty} \frac{A A_{t}^{*}}{\beta B} dx} \\ e^{\frac{1}{2} \int_{x}^{\infty} \frac{A^{*} A_{t}}{\beta B} dx} & e^{\frac{1}{2} \int_{-\infty}^{x} \frac{\beta A^{*} B}{A_{t}^{*}} dx} \end{pmatrix}, \Psi^{1}_{12} = -\frac{i}{4} A, \Psi^{1}_{21} = -\frac{i}{4} A^{*}.$$

$$(2.7)$$

The asymptotic solution $\Psi(x, t, k)$ satisfies

$$\Psi_{\pm}(x,t,k) \to I, \quad x \to \pm \infty,$$

the modified solution satisfies the following integral equation

$$\Psi_{\pm}(x,t,k) = I + \int_{\pm\infty}^{x} e^{-ik(x-y)\sigma_3} U(y,t) \Psi_{\pm}(y,t,k) e^{ik(x-y)\sigma_3} \, \mathrm{d}y.$$
(2.8)

Dividing Ψ_{\pm} into columns as $\Psi_{\pm} = \left(\Psi_{\pm}^{(1)}, \Psi_{\pm}^{(2)}\right)$, due to the structure of the potential U, and Volterra integral equation (2.8), we have

Proposition 1. For $A(x), B(x) \in H^{1,1}(\mathbb{R})$ and $t \in \mathbb{R}^+$, there exist unique eigenfunctions Ψ_{\pm} which satisfy Eq.(2.8), respectively, and have the following properties:

- $det\Psi_{\pm}(x,t,k) = 1, x, t \in \mathbb{R};$
- $[\Psi_{-}]_{1}^{-}, [\Psi_{+}]_{2}^{-}$ are analytic in \mathbb{C}^{+} and continuous in $\mathbb{C}^{+} \cup \mathbb{R}$;
- $[\Psi_+]_1^1, [\Psi_-]_2^2$ are analytic in \mathbb{C}^- , and continuous in $\mathbb{C}^- \cup \mathbb{R}$.

The partition of \mathbb{C}^{\pm} is shown in Fig.1. It can be seen from the equation that $\Psi_{+}e^{-ikx\sigma_{3}+\frac{i\alpha}{4k}t\sigma_{3}}$ and $\Psi_{-}e^{-ikx\sigma_{3}+\frac{i\alpha}{4k}t\sigma_{3}}$ are the solutions of Lax equation (2.8), so they are linearly related

$$\Psi_{-}e^{-ikx\sigma_{3}+\frac{i\alpha}{4k}t\sigma_{3}} = \Psi_{+}e^{-ikx\sigma_{3}+\frac{i\alpha}{4k}t\sigma_{3}}S(t,k), \quad k \in \mathbb{R},$$
(2.9)

where

$$S(t,k) = \left[\begin{array}{cc} s_{11} & s_{12} \\ s_{21} & s_{22} \end{array}\right]$$

Proposition 2. $\Psi(x,t,k)$ and S(t,k) satisfy the symmetry relation

$$\Psi(x,t,k) = \sigma_0 \Psi^*(x,t,k^*) \sigma_0^{-1}, \ S(t,k) = \sigma_0 S^*(t,k^*) \sigma_0^{-1},$$
(2.10)

where

$$\sigma_0 = \left[\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array} \right].$$

Proof. The symmetry of Φ can be easily obtained from Lax equation

$$\Phi(x,t,k) = \sigma_0 \Phi^* (x,t,k^*) \, \sigma_0^{-1},$$

then contact transformation (2.5), which is easy to verify

$$\Psi(x,t,k) = \sigma_0 \Psi^*(x,t,k^*) \, \sigma_0^{-1}.$$

Then according to the relationship (2.9), there are

$$S(t,k) = \sigma_0 S^*(t,k^*) \,\sigma_0^{-1}.$$

Based on the symmetry condition of S(k), we can get

$$s_{11}(k) = s_{22}^*(k^*), s_{12}(k) = -s_{21}^*(k^*)$$

S(t,k) is called scattering matrix and $s_{i,j}(i,j=1,2)$ are usually called scattering data. Moreover, Eq.(2.9) leads to

$$s_{11}(k) = \det\left(\left[\Psi_{-}\right]_{1}, \left[\Psi_{+}\right]_{2}\right), \ s_{12}(k) = e^{2ikx - \frac{i\alpha}{2k}t} \det\left(\left[\Psi_{-}\right]_{2}, \left[\Psi_{+}\right]_{2}\right).$$
(2.11)

On the basis of the properties of Ψ in Proposition 1, we can get

Proposition 3. The scattering matrix S(t,k) and scattering data $s_{11}(k), s_{12}(k)$ satisfies:

- $detS(k) = 1, \quad x, t \in \mathbb{R};$
- $|s_{11}(k)|^2 + |s_{12}(k)|^2 = 1$ for $k \in \mathbb{R}$;
- $s_{11}(k)$ is analytic in \mathbb{C}^+ and continuous in $\mathbb{C}^+ \cup \mathbb{R}$;
- $s_{12}(k)$ is continuous on the real k;
- $s_{11}(k) = 1 + \mathcal{O}(k^{-1}) \text{ as } k \to \infty; \ s_{12}(k) = \mathcal{O}(k^{-1}) \text{ as } k \to \infty;$
- $s_{11}(k) = \varphi_0 + \mathcal{O}(k^2)$ as $k \to 0$; $s_{12}(k) = \mathcal{O}(k^2)$ as $k \to 0$,
- where $\varphi_0 = e^{\frac{1}{2}\int_{-\infty}^{\infty} \frac{\beta A^* B}{A_t^*} dx} e^{\frac{1}{2}\int_{-\infty}^{\infty} \frac{A^* A_t}{\beta B} dx}$.

The reflection coefficient is described below, which is usually defined as

$$r(k) = \frac{s_{12}(k)}{s_{11}(k)},\tag{2.12}$$

using the first two properties in proposition 3, we have

$$1 + |r(k)|^2 = \frac{1}{|s_{11}(k)|^2}$$

In order to deal with our future work, we assume that the initial data meets the following assumptions.

Assume 4. The initial data $A, B \in H^{1,1}(\mathbb{R})$ and it generates generic scattering data which satisfy that:

- $s_{11}(k)$ has no zeros on \mathbb{R} ;
- $s_{11}(k)$ only has finite number of simple zeros.

Based on the above analysis, we can get the following theorem about r(k).

Theorem 5. For any given initial value $A, B \in H^{1,1}(\mathbb{R})$, we have $r(k) \in H^{1,1}(\mathbb{R})$.

Proof. According to the definition of r(k) we have and the asymptotic behavior of s_{11} and s_{21} at singular points in Proposition 3, we can obtain

$$r(k) = \mathcal{O}(k^{-1}), k \to \infty; \ r(k) = \mathcal{O}(k^2), k \to 0.$$

From the proposition estimation in Appendix A that

$$s_{11}(k), s_{12}(k) \in L^2(\mathbb{R}),$$

which lead to

$$r(k) \in L^2(\mathbb{R}).$$

In addition, it can also be obtained $s'_{11}(k), s'_{12}(k) \in L^2(\mathbb{R})$. Therefore

$$r'(k) = \frac{s'_{12}(k)s_{11}(k) - s'_{11}(k)s_{12}(k)}{s^2_{11}(k)} \in L^2(\mathbb{R}).$$

Finally come to the map $A_0, B_0 \to r(k)$ is Lipschitz continuous from $H^{1,1}(\mathbb{R})$ into $H^{1,1}$.

3. The construction of a RHP

Suppose $s_{11}(k)$ has N simple zeros $\mathcal{K} = \{k_j\}_{j=1}^N \subset \mathbb{C}^+$, reviewing the symmetry of S, we can see that there are also N simple zeros $\mathcal{K}^* = \{k_j^*\}_{j=1}^N \subset \mathbb{C}^-$. Then define a meromorphic function M(x, t, k) as

$$M(x,t,k) = \begin{cases} \left(\frac{[\Psi_{-}]_{1}}{s_{11}(k)}, [\Psi_{+}]_{2}\right), & k \in \mathbb{C}^{+}, \\ \left([\Psi_{+}]_{1}, \frac{[\Psi_{-}]_{2}}{s_{22}(k)}\right), & k \in \mathbb{C}^{-}, \end{cases}$$
(3.1)

which solves the following RHP.

Riemann-Hilbert Problem 6. Find a matrix-valued function M(x, t, k) which satisfies:

- Analyticity: M(x,t,k) is meromorphic in $\mathbb{C}\backslash\mathbb{R}$ and has single poles;
- Symmetry: $M(x,t,k) = \sigma_0 M^*(x,t,k^*) \sigma_0^{-1};$
- Jump condition: M(x,t,k) has continuous boundary values $M_{\pm}(x,t,k)$ on \mathbb{R} and

$$M_{+}(x,t,k) = M_{-}(x,t,k)V(k), \quad k \in \mathbb{R},$$
(3.2)

where

$$V(k) = \begin{pmatrix} 1 + |r(k)|^2 & r^*(k^*)e^{-2it\theta(k)} \\ r(k)e^{2it\theta(k)} & 1 \end{pmatrix};$$
(3.3)

• Asymptotic behaviors:

$$M(x,t,k) = I + \mathcal{O}(k^{-1}), \quad k \to \infty;$$

• Residue conditions: M has simple poles at each point in $k_j \in \mathbb{C}^+$ and $k_j^* \in \mathbb{C}^-$ with:

$$\operatorname{Res}_{k=k_{j}} M(z) = \lim_{k \to k_{j}} M(k) \begin{pmatrix} 0 & 0 \\ c_{j}e^{2it\theta(k_{j})} & 0 \end{pmatrix},$$

$$\operatorname{Res}_{k=k_{j}^{*}} M(k) = \lim_{k \to k_{j}^{*}} M(k) \begin{pmatrix} 0 & -c_{j}^{*}e^{-2it\theta(k_{j}^{*})} \\ 0 & 0 \end{pmatrix},$$

(3.4)

where $\theta = k \frac{x}{t} - \frac{\alpha}{4k}, c_j = \frac{s_{12}(k_j)}{s'_{11}(k_j)}.$

The solution of the coupled dispersion AB system (1.1) can be expressed by

$$A = 4i \lim_{k \to \infty} (kM)_{12}(x, t, k), \ B = -\frac{4i}{\beta} \lim_{k \to \infty} \frac{d}{dt} (kM)_{11}.$$
 (3.5)

4. EXISTENCE AND UNIQUENESS OF SOLUTION OF RH PROBLEM

In this chapter, we will mainly focus on the existence and uniqueness of the solution of the RH problem constructed above. Our idea here is to use the vanishing lemma to show that the integral equation has only zero solution to the homogeneous equation, and then show that the equation has a unique solution.

In order to facilitate the application of the subsequent lemma, we focus all the residue conditions in RHP 6 on the circle $\{\kappa_j, \kappa_j^*\}$ centered on the pole $\{k_j \in \mathbb{C}^+, k_j^* \in \mathbb{C}^-\}$. As long as these circles are small enough, they will not intersect. See Fig. 2 for details.




FIGURE 2. Distribution of $\Sigma^{(1)}$.

$$\Sigma^{(1)} = \mathbb{R} \cup \left\{ \kappa_j \right\}_{j=1}^N \cup \left\{ \kappa_j^* \right\}_{j=1}^N$$

and rewrite RHP 6 as follows:

Riemann-Hilbert Problem 7. Find a matrix-valued function M(x, t, k) which satisfies:

- Analyticity: M(x,t,k) is analytic in $\mathbb{C}\setminus\Sigma^{(1)}$;
- Symmetry: $M(x, t, k) = \sigma_0 M^*(x, t, k^*) \sigma_0^{-1}$;
- Jump condition: M(x,t,k) has continuous boundary values $M_{\pm}(x,t,k)$ on $\Sigma^{(1)}$ and

$$M_{+}(x,t,k) = M_{-}(x,t,k)V'(k), \quad k \in \mathbb{R},$$
(4.1)

where

$$V'(k) = \begin{cases} \begin{pmatrix} 1+|r(k)|^2 & r^*(k^*)e^{-2ikx} \\ r(k)e^{2ikx} & 1 \end{pmatrix}, & k \in \mathbb{R}, \\ \begin{pmatrix} 1 & 0 \\ \frac{c_je^{2ikx}}{k-k_j} & 1 \end{pmatrix}, & k \in \kappa_j, \\ \begin{pmatrix} 1 & \frac{c_j^*e^{-2ikx}}{k-k_j^*} \\ 0 & 1 \end{pmatrix}, & k \in \kappa_j^*; \end{cases}$$
(4.2)

• Normalization:

(a)
$$M(k,x) = I + \mathcal{O}(k^{-1}), \quad k \to \infty,$$

(b) $M(k,x) = \mathcal{O}(k^{-1}), \quad k \to \infty.$

Here, we deliberately turn t into a gray variable, mainly to better show the importance of x in the proof process. In normalization (a) is mainly to reconstruct the potential A, B. The role of (b) is to illustrate the existence and uniqueness of solutions to the RHP of type (a).

As we all know, V'(k) has the following decomposition

$$V'(k) = \left(I - \Lambda^{-}\right)^{-1} \left(I + \Lambda^{+}\right).$$

Bring it into (4.1) and define it as ω

$$\omega = M_+ \left(I + \Lambda^+ \right)^{-1} = M_- \left(I - \Lambda^- \right)^{-1}, \tag{4.3}$$

the above formula can be rewritten as

$$M_+ - M_- = \omega(\Lambda^+ + \Lambda^-).$$

Using Plemelj formula, it is easy to give the solution of solvable RHP(7) equation with regularized boundary (a) in the form of

$$M(x,k) = I + \frac{1}{2\pi i} \int_{\Sigma^{(1)}} \frac{\omega(\Lambda^+ + \Lambda^-)}{s-k} ds,$$
(4.4)

the solution of the equation with boundary condition (b),

$$M(x,k) = \frac{1}{2\pi i} \int_{\Sigma^{(1)}} \frac{\omega(\Lambda^+ + \Lambda^-)}{s-k} ds.$$
 (4.5)

The following main task is to prove that the solution with boundary condition (b) tends to 0, that is, Eq.(4.5) tends to 0, which is also obvious. Write M(x, k) expansion as

$$M(x,k) = \frac{1}{2\pi i} \left(\int_{\mathbb{R}} \frac{\omega(\Lambda^+ + \Lambda^-)}{s-k} ds + \int_{\Sigma^{(1)} \setminus \mathbb{R}} \frac{\omega(\Lambda^+ + \Lambda^-)}{s-k} ds \right).$$
(4.6)

Here we define

$$\mathcal{N}(k) = M(k)M(k^*)^H,$$

where 'H' represents conjugate transpose of the matrix, and we only need to prove

$$\int_{\mathbb{R}} \mathcal{N}_{\pm}(k) dk = 0 \tag{4.7}$$

below. Obviously, according to Schwartz's reflection principle, \mathcal{N} is analytical in $\mathbb{C} \setminus \Sigma^{(1)}$.

For $k \in \mathbb{R}$, we have

$$\mathcal{N}_{+}(k) = M_{+}(k)M_{-}(k)^{H} = M_{-}(k)V'(k)M_{-}(k)^{H},$$

$$\mathcal{N}_{-}(k) = M_{-}(k)M_{+}(k)^{H} = M_{-}(k)V'(k)^{H}M_{-}(k)^{H}.$$

Therefore, if Eq.(4.7) is satisfied, there is

$$\int_{\mathbb{R}} M_{-}(k) \left(V'(k) + V'(k)^{H} \right) M_{-}(k)^{H} = 0,$$

since $k \in \mathbb{R}$, so $V'(k) = V'(k)^H$. Then we know that $V'(k) + V'(k)^H$ is also an Hermite and is positive definite. Therefore, if the above formula holds, there must be $M_-(k) = 0$ on \mathbb{R} . So

$$M_{+}(k) = M_{-}(k)V'(k) = 0, \ k \in \mathbb{R}$$

According to Morera's theorem, we know that M(k) is analytic in the neighborhood of each point on \mathbb{R} . And M(k) tends to disappear.

For $k \in \Sigma^{(1)} \setminus \mathbb{R}$. then we have

$$\mathcal{N}_{+}(k) = M_{+}(k)M_{-}(k^{*})^{H} = M_{-}(k)V'(k)\left(\mathcal{N}_{-}(k^{*})^{-1}\right)^{H}M_{+}(k^{*})^{H}$$
$$= M_{-}(k)M_{+}(k^{*})^{H} = \mathcal{N}_{-}(k).$$

Similarly, it can be seen from the Morera's theorem that $\mathcal{N}(k)$ is analytic in $\Sigma^{(1)} \setminus \mathbb{R}$. And $M_{\pm}(x, \cdot) \in L^2(\mathbb{R})$, we know that $\mathcal{N}(k)$ is integrable. According to the integral form of Eq.(4.6),

$$\begin{split} M(x,k) &= \frac{1}{2\pi i} \int_{\Sigma^{(1)} \setminus \mathbb{R}} \frac{\omega(\Lambda^+ + \Lambda^-)}{s-k} ds \\ &= \frac{1}{2k\pi i} \int_{\Sigma^{(1)} \setminus \mathbb{R}} \frac{s}{s-k} \omega(x,s) \left(\Lambda^+ + \Lambda^-\right) ds - \frac{1}{2k\pi i} \int_{\Sigma^{(1)} \setminus \mathbb{R}} \omega(x,s) \left(\Lambda^+ + \Lambda^-\right) ds, \end{split}$$

the Taylor expansion of k shows that $M(x,k) \sim \mathcal{O}(k^{-1})$, $k \in \mathbb{C} \setminus \mathbb{R}$. And then we know that $\mathcal{N}(k) \sim \mathcal{O}(k^{-2})$, $k \in \mathbb{C} \setminus \mathbb{R}$. Then the Eq.(4.7) can be obtained by Cauchy's integral theorem and Jordon's theorem. We can apply the same argument $k \in \mathbb{R}$ to $k \in \Sigma^{(1)} \setminus \mathbb{R}$, and we can also get M(x,k) = 0. Therefore, it is concluded that M(x,k) = 0 is on the whole complex plane.

So using the vanishing lemma and Fredholm alternative theorem can get the following proposition.

Proposition 8. Assuming that the initial date $A, B \in H^{1,1}(\mathbb{R})$, then the RHP(7) with normalization boundary condition (a) has a unique solution M(x, k).

5. Conjugation

Through the form of the jump matrix and the residual condition, it can be found that the long-term asymptotic of RHP1 is affected by the growth and attenuation of the exponential function $e^{\pm 2it\theta(k)}$. Therefore, it is necessary to deal with the jump matrix in Eq. (3.3) and the oscillation term in the residual condition, and decompose the jump matrix according to the sign change graph of $Re(i\theta)$ to ensure that any jump matrix is bounded in a given area. We further rewrite θ in the form

$$\theta = -\frac{\alpha k}{4k_0^2} - \frac{\alpha}{4k} = -\frac{\alpha}{4} \left(\frac{k}{k_0^2} + \frac{1}{k}\right),$$

$$Re(i\theta) = \frac{Im(k)\alpha}{4} \left(\frac{1}{k_0^2} - \frac{1}{|k|^2}\right).$$
(5.1)

which leads to

For convenience, we mainly consider
$$\alpha x < 0, t > 0$$
, and can be treated similarly for $t < 0$. Without losing generality, it can be assumed that $\alpha < 0$ and $x > 0$. Based on this, we find two phase points of function $\theta(k)$ are $\pm k_0$, where $k_0 = \sqrt{-\frac{\alpha t}{4x}}$.

According to the sign change of $Re(i\theta)$, the attenuation region of oscillation term $e^{2\pm it\theta}$ can be obtained when $t \to \infty$. which is shown in Fig. 3.




FIGURE 3. Decay domains of exponential oscillatory terms.

7

The jump matrix V(k) in RHP1 has the following decomposition

$$V(k) = \begin{pmatrix} 1 & r^*(k^*)e^{-2it\theta} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ r(k)e^{2it\theta} & 1 \end{pmatrix}, \ |k| > k_0,$$

and

$$V(k) = \begin{pmatrix} 1 & 0\\ \frac{r(k)}{1+|r(k)|^2}e^{2it\theta} & 1 \end{pmatrix} \begin{pmatrix} 1+|r(k)|^2 & 0\\ 0 & \frac{1}{1+|r(k)|^2} \end{pmatrix} \begin{pmatrix} 1 & \frac{r^*(k^*)}{1+|r(k)|^2}e^{-2it\theta}\\ 0 & 1 \end{pmatrix}, |k| < k_0.$$

In order to remove the diagonal matrix in the second decomposition, we introduce a scalar RHP about $\delta(k)$ by using the method in Ref. [33].

Riemann-Hilbert Problem 9. Find a scalar function $\delta(k)$ which satisfies:

- $\delta(k)$ is analytic in $\mathbb{C}\backslash\mathbb{R}$;
- $\delta_+(k) = \delta_-(k) \left(1 + |r(k)|^2 \right), \quad |k| < k_0;$
- $\delta_+(k) = \delta_-(k), \quad |k| > k_0;$
- $\delta(k) \to 1$, as $k \to \infty$.

Using the Plemelij formula, it is easy to write the unique solution of the above RHP 9 as

$$\delta(k) = \exp\left[i\int_{-k_0}^{k_0} \frac{\nu(s)}{s-k}ds\right],\tag{5.2}$$

(5.3)

where $\nu(s) = -\frac{1}{2\pi} \log (1 + |r(s)|^2)$. The trace formula can be given directly according to the calculation **Proposition 10.**

$s_{11}(k) = \prod_{j=1}^{N} \frac{k - k_j}{k - k_j^*} \exp\left[-i \int_{-\infty}^{+\infty} \frac{\nu(z)}{z - k} dz\right],$

$$s_{22}(k) = \prod_{j=1}^{N} \frac{k - k_j^*}{k - k_j} \exp\left[i \int_{-\infty}^{+\infty} \frac{\nu(z)}{z - k} dz\right].$$
(5.4)

Proof. From the above, we know that $s_{11}(k)$ and $s_{22}(k)$ are analytic on \mathbb{C}^+ and \mathbb{C}^- respectively, and according to the assumption, the discrete zeros of $s_{11}(k)$ and $s_{12}(k)$ are k_j and k_j^* respectively, so let

$$\beta^{+}(k) = s_{11}(k) \prod_{j=1}^{N} \frac{k - k_{j}^{*}}{k - k_{j}}, \quad \beta^{-}(k) = s_{22}(k) \prod_{j=1}^{N} \frac{k - k_{j}}{k - k_{j}^{*}}, \tag{5.5}$$

they are still analytic on \mathbb{C}^+ , \mathbb{C}^- , but they no longer have zeros. And have $\beta^+(k)\beta^-(k) = s_{11}(k)s_{22}(k), k \in \mathbb{R}$. It can be obtained that

$$\beta^+(k)\beta^-(k) = s_{11}s_{22} = \frac{1}{1+|r(k)|^2}, \quad k \in \mathbb{R}$$

through

$$\det S(k) = s_{11}(k)s_{22}(k) - s_{21}(k)s_{12}(k) = 1$$

which can leads to that

$$\log \beta^+(k) - \left(-\log \beta^-(k)\right) = -\log\left(1 + |r(k)|^2\right), \quad k \in \mathbb{R}$$

using Plemelj formula, we can get

$$\beta^{\pm}(k) = \exp\left(\pm i \int_{-\infty}^{\infty} \frac{\nu(z)}{z-k} dz\right), \quad k \in \mathbb{C}^{\pm}.$$
(5.6)

Therefore, the trace formula can be obtained by bringing Eq.(5.6) into (5.5).

Next, for the convenience of follow-up research, we first introduce some notations

$$\Delta_{k_0}^+ = \{ j \in \{1, \dots, N\} \mid |k_j| > k_0 \}, \Delta_{k_0}^- = \{ j \in \{1, \dots, N\} \mid |k_j| < k_0 \}.$$
(5.7)

In addition, the function is introduced

$$T(k) = \prod_{j \in \Delta_{k_0^-}} \frac{k - k_j^*}{k - k_j} \delta(k).$$

$$(5.8)$$

Proposition 11. The function T(k) defined by Eq.(5.8) has the following properties:

• T(k) is meromorphic function in $\mathbb{C}\setminus(-k_0,k_0)$. For $j \in \Delta_{k_0}^-$, it is a simple pole at k_j and a simple zero at k_j^* . Other places are non-zero and analytic;

- $T^*(k^*)T(k) = 1$, for $\mathbb{C} \setminus (-k_0, k_0)$;
- For $k \in (-k_0, k_0)$, the boundary values $T_{\pm}(k)$ satisfy

$$T_{+}(k) = T_{-}(k) \left(1 + |r(k)|^{2} \right), k \in (-k_{0}, k_{0})$$

• When $|k| \to \infty$, $|\arg(k)| \le c < \pi$

$$T(k) = 1 + \frac{i}{k} \left[2 \sum_{j \in \Delta_{k_0}^-} \operatorname{Im} k_j - \int_{-k_0}^{k_0} \nu(s) ds \right] + O(k^{-2});$$

• $k \to \pm k_0$, along any ray $\pm k_0 + e^{i\phi} \mathbb{R}_+$ with $|\phi| \le c < \pi$

$$\left|T(k) - T_0(\pm k_0)(k \mp k_0)^{i\nu(\pm k_0)}\right| \le c \left|k \mp k_0\right|^{\frac{1}{2}}$$

where

$$\begin{split} T_{0}\left(\pm k_{0}\right) &= \prod_{j\in\Delta_{k_{0}}^{-}} \left(\frac{\pm k_{0}-k_{j}^{*}}{\pm k_{0}-k_{j}}\right) e^{i\beta(k,\pm k_{0})},\\ \beta(k,-k_{0}) &= \left(i\int_{-k_{0}}^{-k_{0}+1} \frac{\nu\left(-k_{0}\right)}{z-k}dz + i\int_{-k_{0}}^{k_{0}} \frac{\nu(z)-\chi_{1}(z)\nu\left(-k_{0}\right)}{z-k}dz\right),\\ \beta(k,k_{0}) &= \left(i\int_{k_{0}-1}^{k_{0}} \frac{\nu\left(k_{0}\right)}{z-k}dz + i\int_{-k_{0}}^{k_{0}} \frac{\nu(z)-\chi_{2}(z)\nu\left(k_{0}\right)}{z-k}dz\right),\\ \chi_{1}(k) &= \begin{cases} 1, & -k_{0} < k < -k_{0}+1, \\ 0, & elsewhere \end{cases}, \quad \chi_{2}(k) = \begin{cases} 1, & k_{0}-1 < k < k_{0}, \\ 0, & elsewhere \end{cases}. \end{split}$$

Through the defined T(k), the following transformation can be done to eliminate the diagonal matrix in the decomposition of the jump matrix in the interval $[-k_0, k_0]$.

$$M^{(1)}(k) = M(k)T(k)^{-\sigma_3},$$
(5.9)

which satisfies the following RHP:

Riemann-Hilbert Problem 12. Find a matrix $M^{(1)}(k)$ that satisfies:

- $M^{(1)}$ is analytic within $\mathbb{C} \setminus \mathbb{R} \cup \mathcal{K} \cup \mathcal{K}^*$;
- $M^{(1)}(k) = \sigma_0 (M^{(1)})^* (k^*) \sigma_0^{-1};$
- $\begin{array}{l} \bullet M^{(1)}(k) = I + \mathcal{O}\left(k^{-1}\right) \sigma_{0}^{-1}; \\ \bullet M^{(1)}(k) = I + \mathcal{O}\left(k^{-1}\right), \quad k \to \infty; \\ \bullet M^{(1)}_{+}(k) = M^{(1)}_{-}(k) V^{(1)}(k), \quad k \in \mathbb{R}, \\ where \end{array}$

$$V^{(1)}(k) = \begin{cases} \begin{pmatrix} 1 & 0 \\ \frac{r}{1+|r|^2}T_{-}^{-2}e^{2it\theta(k)} & 1 \end{pmatrix} \begin{pmatrix} 1 & \frac{r^*}{1+|r|^2}T_{+}^{2}e^{-2it\theta(k)} \\ 0 & 1 \end{pmatrix}, & |k| < k_0 \\ \begin{pmatrix} 1 & r^*T^2e^{-2it\theta(k)} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ rT^{-2}e^{2it\theta(k)} & 1 \end{pmatrix}, & |k| > k_0; \end{cases}$$

• $M^{(1)}(k)$ has simple poles at each point in $k_j \in \mathbb{C}^+$ and $k_j^* \in \mathbb{C}^-$ with:

$$\operatorname{Res}_{k=k_{j}} M^{(1)}(k) = \begin{cases}
\lim_{k \to k_{j}} M^{(1)}(k) \begin{pmatrix} 0 & c_{j}^{-1} \left(\frac{1}{T}\right)'(k_{j})^{-2} e^{-2it\theta(k_{j})} \\
0 & 0 \end{pmatrix}, & j \in \Delta_{k_{0}}^{-}, \\
\lim_{k \to k_{j}} M^{(1)}(k) \begin{pmatrix} 0 & 0 \\ c_{j}T(k_{j})^{-2} e^{2it\theta(k_{j})} & 0 \end{pmatrix}, & j \in \Delta_{k_{0}}^{+}. \\
\\
\operatorname{Res}_{k=k_{j}^{*}} M^{(1)}(k) = \begin{cases}
\lim_{k \to k_{j}^{*}} M^{(1)}(k) \begin{pmatrix} 0 & -c_{j}^{*}T^{2}(k_{j}^{*})e^{-2it\theta(k_{j}^{*})} \\
0 & 0 \end{pmatrix}, & j \in \Delta_{k_{0}}^{-}, \\
\\
\\
\lim_{k \to k_{j}^{*}} M^{(1)}(k) \begin{pmatrix} 0 & -c_{j}^{*}T^{2}(k_{j}^{*})e^{-2it\theta(k_{j}^{*})} \\
0 & 0 \end{pmatrix}, & j \in \Delta_{k_{0}}^{+}.
\end{cases}$$
(5.10)

In addition, due to $T(k)^{-\sigma^3} \to I$, as $k \to \infty$, so the solution of the coupled dispersion AB system can be expressed as

$$A = 4i \lim_{k \to \infty} (kM^{(1)}(k))_{12}, \ B = -\frac{4i}{\beta} \lim_{k \to \infty} \frac{d}{dt} (kM^{(1)}(k))_{11}.$$
(5.11)

6. CONTOUR DEFORMATION

The next idea is to eliminate the jump on the real axis and open it at a small angle at the steady-state phase point to deform the contour of RHP 12. According to the number of steady-state phase points and spectral singularity points, the following contours can be considered

$$\Sigma^{(2)} = \Sigma_1 \cup \Sigma_2^{\pm} \cup \Sigma_3^{\pm} \cup \Sigma_4 \cup \Sigma_5 \cup \Sigma_6^{\pm} \cup \Sigma_7^{\pm} \cup \Sigma_8, \tag{6.1}$$

shown in Fig. 4. In order to facilitate later applications, new tokens are introduced




FIGURE 4. Deformation from \mathbb{R} to $\Sigma^{(2)}$.

$$\varrho = \frac{1}{2} \min_{k,\mu \in \mathcal{K} \cup \mathcal{K}^*} |k - \mu|$$

Because of the symmetry, and the pole is not on the real axis, then for any $k_j = u_j + iv_j \in \mathcal{K}$, obviously there is $k_j^* = u_j - iv_j \in \mathcal{K}^*$. according to the above representation, there must be $|k_j - k_j^*| = 2|v_j| > \rho$. So there must be dis $(k, \mathbb{R}) = |v| \ge \rho > 0$, here is due to the arbitrariness of j.

In order to keep the residual condition unchanged during contour deformation, an eigenfunction is defined near the discrete spectrum

$$\Upsilon_{\mathcal{K}}(k) = \begin{cases} 1, \operatorname{dist}\left(k, \mathcal{K} \cup \mathcal{K}^*\right) < \varrho/3, \\ 0, \operatorname{dist}\left(k, \mathcal{K} \cup \mathcal{K}^*\right) > 2\varrho/3. \end{cases}$$
(6.2)

Now we find a matrix $R_j \to \mathbb{C}$ with the following boundary conditions,

$$R_{1}(k) = \begin{cases} r(k)T^{-2}(k), & k \in (-\infty, -k_{0}), \\ r(-k_{0})T_{0}^{-2}(-k_{0})(k+k_{0})^{-2i\nu(-k_{0})}(1-\Upsilon_{\mathcal{K}}(k)), & k \in \Sigma_{1}, \end{cases}$$
(6.3)

$$R_{3}(k) = \begin{cases} \frac{r^{*}(k^{*})T_{+}^{2}(k)}{1+|r(k)|^{2}}, & k \in (-k_{0}, k_{0}), \\ \frac{r^{*}(-k_{0})T_{0}^{2}(-k_{0})}{1+|r(-k_{0})|^{2}} \left(k+k_{0}\right)^{2iv(-k_{0})} \left(1-\Upsilon_{\mathcal{K}}(k)\right), & k \in \Sigma_{2}^{+}, \\ 0, & k \in \Sigma_{2}^{-}, \end{cases}$$

$$(6.4)$$

$$R_{4}(k) = \begin{cases} \frac{r^{*}(k^{*})T_{+}^{2}(k)}{1+|r(k)|^{2}}, & k \in (-k_{0}, k_{0}), \\ \frac{r^{*}(k_{0})T_{0}^{2}(k_{0})}{1+|r(k_{0})|^{2}} (k-k_{0})^{2iv(k_{0})} (1-\Upsilon_{\mathcal{K}}(k)), & k \in \Sigma_{3}^{+}, \\ 0, & k \in \Sigma_{3}^{-}, \end{cases}$$
(6.5)

$$R_{5}(k) = \begin{cases} r(k)T^{-2}(k), & k \in (k_{0}, \infty), \\ r(k_{0})T_{0}^{-2}(k_{0})(k-k_{0})^{-2iv(k_{0})}(1-\Upsilon_{\mathcal{K}}(k)), & k \in \Sigma_{4}, \end{cases}$$
(6.6)

$$R_{6}(k) = \begin{cases} r^{*}(k^{*})T^{2}(k), & k \in (k_{0}, \infty), \\ r^{*}(k_{0})T^{2}_{0}(k_{0})(k-k_{0})^{2iv(k_{0})}(1-\Upsilon_{\mathcal{K}}(k)), & k \in \Sigma_{5}, \end{cases}$$
(6.7)

$$R_{7}(k) = \begin{cases} \frac{r(k)T_{-}^{-2}(k)}{1+|r(k_{0})|^{2}}, & k \in (-k_{0}, k_{0}), \\ \frac{r(k_{0})T_{0}^{-2}(k_{0})}{1+|r(k_{0})|^{2}} (k-k_{0})^{-2iv(k_{0})} (1-\Upsilon_{\mathcal{K}}(k)), & k \in \Sigma_{6}^{+}, \\ 0, & k \in \Sigma_{6}^{-}, \end{cases}$$
(6.8)

$$R_{8}(k) = \begin{cases} \frac{r(k)T_{-}^{-2}(k)}{1+|r(k)|^{2}}, & k \in (-k_{0}, k_{0}), \\ \frac{r(-k_{0})T_{0}^{-2}(-k_{0})}{1+|r(-k_{0})|^{2}} (k+k_{0})^{-2i\nu(-k_{0})} (1-\Upsilon_{\mathcal{K}}(k)), & k \in \Sigma_{7}^{+}, \\ 0, & k \in \Sigma_{7}^{-}, \end{cases}$$
(6.9)

$$R_{9}(k) = \begin{cases} r^{*}(k)T^{2}(k), & k \in (-\infty, \xi_{1}), \\ r^{*}(-k_{0})T_{0}^{2}(-k_{0})(k+k_{0})^{2iv(-k_{0})}(1-\Upsilon_{\mathcal{K}}(k)), & k \in \Sigma_{8}. \end{cases}$$
(6.10)

And meet the following estimates

$$R_j \lesssim \sin^2 \left(\arg \left(k \pm k_0 \right) \right) + \langle \operatorname{Re}(k) \rangle^{-\frac{1}{2}}, \ j = 1, 3, ..., 9$$
 (6.11)

In this way, the original jump in $\pm k_0$ will become as shown in Fig. 5 and Fig. 6.




FIGURE 5. Matrix $\mathcal{R}^{(2)}$ around stationary phase point $-k_0$.




FIGURE 6. Matrix $\mathcal{R}^{(2)}$ around stationary phase point k_0 .

In order to facilitate future calculation and estimation, the region needs to be divided appropriately, as shown in the Fig.7 below.




FIGURE 7. Region segmentation.

And making some marks

$$p_1(k) = p_5(k) = r(k), \qquad p_6(k) = p_9(k) = r^*(k) p_3(k) = p_4(k) = \frac{r^*(k)}{1 + |r(k)|^2}, \quad p_7(k) = p_8(k) = \frac{r(k)}{1 + |r(k)|^2},$$

further estimates of $\bar{\partial}R_j$ in different regions can be obtained

Lemma 13. Suppose $r \in H^{1,1}(\mathbb{R})$, $\bar{\partial}R_i$ defined by (6.3)-(6.10) satisfies

$$\left|\bar{\partial}R_j(k)\right| \lesssim \left|\bar{\partial}\Upsilon_{\mathcal{K}}(k)\right| + \left|p'_j(\operatorname{Re}(k))\right| + \left|k \pm k_0\right|^{-\frac{1}{2}},\tag{6.12}$$

on D_1, D_5, D_6, D_9 and $D_3^+, D_4^+, D_7^+, D_8^+$. $\bar{\partial}R_j$ satisfies

$$\left|\bar{\partial}R_j(k)\right| \lesssim \left|\bar{\partial}\Upsilon_{\mathcal{K}}(k)\right| + \left|p_j'(\operatorname{Re}(k))\right| + |k|^{-\frac{1}{2}}$$
(6.13)

on $D_3^-, D_4^-, D_7^-, D_8^-$.

Correspondingly, the jump line is transformed into $\Sigma^{(3)}$. Here, two jump lines are added, which are in the following form . 1

$$\tilde{v} = \begin{pmatrix} I, & k \in \left(\frac{-ik_0}{2} \tan \frac{\pi}{24}, \frac{ik_0}{2} \tan \frac{\pi}{24}\right), \\ \begin{pmatrix} 1 & \left(R_2^+ - R_2^-\right) e^{-2i\theta} \\ 0 & 1 \end{pmatrix}, & k \in \left(\frac{ik_0}{2} \tan \frac{\pi}{24} - \frac{k_0}{2}, \frac{ik_0}{2\sqrt{3}} - \frac{k_0}{2}\right), \\ \begin{pmatrix} 1 & 0 \\ \left(R_7^+ - R_7^-\right) e^{2i\theta} & 1 \\ 1 & \left(R_3^- - R_3^+\right) e^{-2i\theta} \\ 0 & 1 \end{pmatrix}, & k \in \left(-\frac{ik_0}{2} \tan \frac{\pi}{24} - \frac{k_0}{2}, -\frac{ik_0}{2\sqrt{3}} - \frac{k_0}{2}\right), \\ \begin{pmatrix} \left(R_6^- - R_6^+\right) e^{2i\theta} & 1 \right), & k \in \left(-\frac{ik_0}{2} \tan \frac{\pi}{24} + \frac{k_0}{2}, \frac{ik_0}{2\sqrt{3}} + \frac{k_0}{2}\right), \\ \begin{pmatrix} \left(R_6^- - R_6^+\right) e^{2i\theta} & 1 \right), & k \in \left(-\frac{ik_0}{2} \tan \frac{\pi}{24} + \frac{k_0}{2}, -\frac{ik_0}{2\sqrt{3}} + \frac{k_0}{2}\right). \end{cases}$$
(6.14)

The jump line $\Sigma^{(3)}$ is shown in Fig. 8.




FIGURE 8. The contour of $\Sigma^{(3)}$.

At the same time, similar to Ref. [49], we can get \tilde{v} subjects to the following estimate

$$|\tilde{v} - I| \lesssim e^{-t}.$$

Therefore, the following changes can be made

$$M^{(2)}(k) = M^{(1)}(k)\mathcal{R}^{(2)}, \tag{6.15}$$

where

$$\mathcal{R}^{(2)}(k) = \begin{cases} \begin{pmatrix} 1 & 0 \\ (-1)^{m_j} R_j(k) e^{2it\theta} & 1 \end{pmatrix}, & k \in D_j, \quad j = 1, 5, 7, 8 \\ \begin{pmatrix} 1 & (-1)^{m_j} R_j(k) e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & k \in D_j, \quad j = 3, 4, 6, 9 \\ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & k \in D_2 \cup D_{10}, \end{cases}$$

where $m_1 = m_3 = m_4 = m_5 = 1, m_6 = m_7 = m_8 = m_9 = 0$. Matrix $M^{(2)}(k)$ satisfies the following RHP:

Riemann-Hilbert Problem 14. The $M^{(2)}(k)$ obtained from $M^{(1)}(k)$ and $\mathcal{R}^{(2)}(k)$ above is expected to satisfy the following properties:

- $M^{(2)}$ is analytic within $\mathbb{C} \setminus \Sigma^{(2)} \cup \mathcal{K} \cup \mathcal{K}^*$; $M^{(2)}(k) = \sigma_0 (M^{(2)}(k^*))^* \sigma_0^{-1}$; $M^{(2)}(k) = I + \mathcal{O}(k^{-1}), \quad k \to \infty$;

• $M^{(2)}_{\perp}(k) = M^{(2)}_{\perp}(k)V^{(2)}(k), \quad k \in \Sigma^{(2)}, \text{ where }$

$$V^{(2)}(k) = \begin{cases} \begin{pmatrix} 1 & 0 \\ R_1(k)e^{2it\theta} & 1 \end{pmatrix}, & k \in \Sigma_1, \\ \begin{pmatrix} 1 & R_3(k)e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_2, \\ \begin{pmatrix} 1 & R_4(k)e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_3, \\ \begin{pmatrix} 1 & R_5(k)e^{2it\theta} & 1 \end{pmatrix}, & k \in \Sigma_4, \\ \begin{pmatrix} 1 & R_6(k)e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_5, \\ \begin{pmatrix} 1 & 0 \\ R_7(k)e^{2it\theta} & 1 \\ 1 & R_9(k)e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_7, \\ \begin{pmatrix} 1 & R_9(k)e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_8. \end{cases}$$

• $M^{(2)}(k)$ has simple poles at each point in $\mathcal{K} \cup \mathcal{K}^*$ with:

$$\operatorname{Res}_{k=k_{j}} M^{(2)}(k) = \begin{cases} \lim_{k \to k_{j}} M^{(2)}(k) \begin{pmatrix} 0 & c_{j}^{-1} \left(\frac{1}{T}\right)' (k_{j})^{-2} e^{-2it\theta(k_{j})} \\ 0 & 0 \end{pmatrix}, & j \in \Delta_{k_{0}}^{-}, \\ \lim_{k \to k_{j}} M^{(2)}(k) \begin{pmatrix} 0 & 0 \\ c_{j}T(k_{j})^{-2} e^{2it\theta(k_{j})} & 0 \end{pmatrix}, & j \in \Delta_{k_{0}}^{+}. \end{cases}$$

$$\operatorname{Res}_{k=k_{j}^{*}} M^{(2)}(k) = \begin{cases} \lim_{k \to k_{j}^{*}} M^{(2)}(k) \begin{pmatrix} 0 & -c_{j}^{*}T(k_{j}^{*})^{-2} e^{2it\theta(k_{j}^{*})} & 0 \end{pmatrix}, & j \in \Delta_{k_{0}}^{-}, \\ \lim_{k \to k_{j}^{*}} M^{(2)}(k) \begin{pmatrix} 0 & -c_{j}^{*}T(k_{j}^{*})^{2} e^{-2it\theta(k_{j}^{*})} & 0 \end{pmatrix}, & j \in \Delta_{k_{0}}^{+}. \end{cases}$$

$$(6.16)$$

• For $\mathbb{C} \setminus \Sigma^{(2)} \cup \mathcal{K} \cup \mathcal{K}^*$, we have the $\bar{\partial}$ derivative

$$\bar{\partial}M^{(2)}(k) = M^{(2)}(k)\bar{\partial}\mathcal{R}^{(2)}(k),$$
(6.17)

where

$$\mathcal{R}^{(2)}(k) = \begin{cases} \begin{pmatrix} 1 & 0 \\ (-1)^{m_j} \bar{\partial} R_j(k) e^{2it\theta} & 1 \\ 1 & (-1)^{m_j} \bar{\partial} R_j(k) e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & k \in D_j, \quad j = 3, 4, 6, 9 \\ \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, & k \in D_2 \cup D_{10}, \end{cases}$$

where $m_1 = m_3 = m_4 = m_5 = 1, m_6 = m_7 = m_8 = m_9 = 0.$

The relationship between the solution of the coupled dispersion AB system and $M^{(2)}(k)$ is

$$A(x,t) = 4i \lim_{k \to \infty} (kM^{(2)}(k))_{12}, \ B(x,t) = -\frac{4i}{\beta} \lim_{k \to \infty} \frac{d}{dt} (kM^{(2)}(k))_{11}.$$
(6.18)

The $\bar{\partial}$ derivative appears in the above RHP, so it is also called mixed $\bar{\partial}$ -RHP.

7. Decomposition of the RHP 14

The following is mainly about the classification of RHP 14. For the case of $\bar{\partial}\mathcal{R}^{(2)}(k) = 0$, it is called a pure RH problem, and for the case of $\bar{\partial}\mathcal{R}^{(2)}(k) \neq 0$, it is called a pure $\bar{\partial}$ problem. In the process of classification, consider the transformation $M^{(2)}(k) = M^{(3)}(k)M^{(2)}_{rhp}(k)$, if $\bar{\partial}\mathcal{R}^{(2)}(k) = 0$, it corresponds to $M^{(2)}_{rhp}(k)$, if $\bar{\partial}\mathcal{R}^{(2)}(k) \neq 0$, it corresponds to $M^{(3)}_{rhp}(k)$, if $\bar{\partial}\mathcal{R}^{(2)}(k) = 0$, it spectral to $M^{(3)}_{rhp}(k)$, if $\bar{\partial}\mathcal{R}^{(2)}(k) = 0$, it corresponds to $M^{(2)}_{rhp}(k)$, if $\bar{\partial}\mathcal{R}^{(2)}(k) \neq 0$, it corresponds to $M^{(3)}_{rhp}(k)$.

Riemann-Hilbert Problem 15. Find a matrix-valued function $M_{rhp}^{(2)}(k)$ with following properties:

- $M_{rhp}^{(2)}(k)$ is analytic within $\mathbb{C}\setminus\Sigma^{(2)} \cup \mathcal{K} \cup \mathcal{K}^*$; Symmetry: $M_{rhp}^{(2)}(k) = \sigma_0(M_{rhp}^{(2)})^*(k^*)\sigma_0^{-1}$;

• Analytic behavior:

$$M_{rhp}^{(2)}(k) = I + \mathcal{O}\left(k^{-1}\right), \quad k \to \infty,$$

$$(7.1)$$

• Jump condition:

$$M_{rhp+}^{(2)}(k) = M_{rhp-}^{(2)}(k)V^{(2)}(k), \quad k \in \Sigma^{(2)};$$
(7.2)

• Residue conditions: $M_{rhp}^{(2)}(k)$ has simple poles at each point in $\mathcal{K} \cup \mathcal{K}^*$ with

$$\operatorname{Res}_{k=k_{j}} M_{rhp}^{(2)}(k) = \begin{cases}
\operatorname{lim}_{k \to k_{j}} M_{rhp}^{(2)}(k) \begin{pmatrix} 0 & c_{j}^{-1} \left(\frac{1}{T}\right)'(k_{j})^{-2} e^{-2it\theta(k_{j})} \\
0 & 0 \end{pmatrix}, \quad j \in \Delta_{k_{0}}^{-}, \\
\operatorname{lim}_{k \to k_{j}} M_{rhp}^{(2)}(k) \begin{pmatrix} 0 & 0 \\ c_{j}T(k_{j})^{-2} e^{2it\theta(k_{j})} & 0 \end{pmatrix}, \quad j \in \Delta_{k_{0}}^{+}. \\
\operatorname{lim}_{k \to k_{j}^{*}} M_{rhp}^{(2)}(k) \begin{pmatrix} 0 & -(c_{j}^{*})^{-1}T'(k_{j}^{*})^{-2} e^{2it\theta(k_{j}^{*})} & 0 \end{pmatrix}, \quad j \in \Delta_{k_{0}}^{-}, \\
\operatorname{lim}_{k \to k_{j}^{*}} M_{rhp}^{(2)}(k) \begin{pmatrix} 0 & -c_{j}^{*}T(k_{j}^{*})^{2} e^{-2it\theta(k_{j}^{*})} & 0 \end{pmatrix}, \quad j \in \Delta_{k_{0}}^{+}. \end{cases}$$
(7.3)

• $\bar{\partial}$ -Derivative: $\bar{\partial}R^{(2)}(k) = 0$ for $k \in \mathbb{C}$.

When $\bar{\partial}\mathcal{R}^{(2)}(k) \neq 0$, we use the above $M_{rhp}^{(2)}(k)$ to construct a transformation: $M^{(3)}(k) = M^{(2)}(k)(M_{rhp}^{(2)}(k))^{-1}$, which is a pure $\bar{\partial}$ problem. For $M^{(3)}(k)$, we have the following properties:

Riemann-Hilbert Problem 16. Find a matrix-valued function $M^{(3)}(k)$ with following properties:

- $M^{(3)}(k)$ is continuous in \mathbb{C} , continuous first partial derivatives in $\mathbb{C}\setminus\Sigma^{(2)}\cup\mathcal{K}\cup\mathcal{K}^*$; Symmetry: $M^{(3)}(k) = \sigma_0(M^{(3)}(k^*))^*\sigma_0^{-1}$; Jump condition: $M^{(3)}_+(k) = M^{(3)}_-(k)$, $k \in \Sigma^{(2)}$; Analytic behavior: $M^{(3)}(k) = I + \mathcal{O}(k^{-1})$, $k \to \infty$; $\bar{\partial}$ -Derivative: $\bar{\partial}M^{(3)}(k) = M^{(3)}(k)M^{(2)}_{rhp}(k)\bar{\partial}R^{(2)}(k)M^{(2)}_{rhp}(k)^{-1}$.

For the above proof, we can refer to Ref. [48]. Here we avoid repetition and do not present it again. The next focus is to find $M_{rhp}^{(2)}(k)$. First, two open intervals are defined

$$\mathcal{A}_{1} = \left\{ k : |k + k_{0}| \leq \min\left\{\frac{k_{0}}{2}, \rho/3\right\} \triangleq \varepsilon \right\},$$

$$\mathcal{A}_{2} = \left\{ k : |k - k_{0}| \leq \min\left\{\frac{k_{0}}{2}, \rho/3\right\} \triangleq \varepsilon \right\}.$$

(7.4)

Then $M_{rhp}^{(2)}(k)$ is divided into three parts

$$M_{rhp}^{(2)}(k) = \begin{cases} M^{err}(k)M^{out}(k), & k \notin \{\mathcal{A}_1 \cup \mathcal{A}_2\}, \\ M^{err}(k)M^{(-k_0)}(k), & k \in \mathcal{A}_1, \\ M^{err}(k)M^{(k_0)}(k), & k \in \mathcal{A}_2. \end{cases}$$
(7.5)

Due to dist $(\mathcal{K} \cup \mathcal{K}^*, \mathbb{R}) > \varrho$, $M_{rhp}^{(2)}(k)$, $M^{(-k_0)}(k)$ and $M^{(k_0)}(k)$ have no poles in \mathcal{A}_1 and \mathcal{A}_2 . The matrix $M_{rhp}^{(2)}(k)$ is divided into three parts by decomposition: one part can be called the external model RH problem, represented by $M^{out}(k)$, which can be solved directly by considering the standard RH problem without reflection potential. The other two parts are near the phase points $M^{(-k_0)}(k)$ and $M^{(k_0)}(k)$, which can be matched to the known PC model, namely the parabolic cylinder model in \mathcal{A}_1 and \mathcal{A}_2 , to be solved in Section 8.4. In addition, matrix $M^{err}(k)$ is an error function, which can be solved by the small norm RH problem in Section 8.5.

Let's define L_{ϵ} for a fixed ϵ

$$L_{\epsilon} = \left\{ k : k = k_0 + Ak_0 e^{\frac{3i\pi}{4}}, \epsilon \le A \le \frac{1}{\sqrt{2}} \right\}$$
$$\cup \left\{ k : k = k_0 + Ak_0 e^{\frac{i\pi}{4}}, \epsilon \le A \le \infty \right\}$$
$$\cup \left\{ k : k = -k_0 + Ak_0 e^{\frac{i\pi}{4}}, \epsilon \le A \le \frac{1}{\sqrt{2}} \right\}$$
$$\cup \left\{ k : k = -k_0 + Ak_0 e^{\frac{3i\pi}{4}}, \epsilon \le A \le \infty \right\}.$$

Through the above definition, we can write the estimation of $V^{(2)}(k)$ as follows

$$\begin{split} \left\| V^{(2)}(k) - I \right\|_{L^{\infty} \left(\Sigma^{(2)}_{+} \cap \mathcal{A}_{2} \right)} &= \mathcal{O} \left(e^{\frac{\sqrt{2}t}{4} \alpha |k - k_{0}|} \left(k_{0}^{-2} - |k|^{-2} \right) \right), \\ \left\| V^{(2)}(k) - I \right\|_{L^{\infty} \left(\Sigma^{(2)}_{-} \cap \mathcal{A}_{1} \right)} &= \mathcal{O} \left(e^{\frac{\sqrt{2}t}{4} \alpha |k + k_{0}|} \left(k_{0}^{-2} - |k|^{-2} \right) \right), \\ \left\| V^{(2)}(k) - I \right\|_{L^{\infty} \left(\Sigma^{(2)} \setminus \left(\mathcal{A}_{1} \cup \mathcal{A}_{2} \right) \right)} &= O \left(e^{-2t\epsilon} \right), \end{split}$$
(7.6)

where the contours are defined by

$$\Sigma_{+}^{(2)} = \Sigma_3 \cup \Sigma_4 \cup \Sigma_5 \cup \Sigma_6, \quad \Sigma_{-}^{(2)} = \Sigma_1 \cup \Sigma_2 \cup \Sigma_7 \cup \Sigma_8.$$

This implies that the jump matrix $V^{(2)}(k)$ goes to I on both $\Sigma^{(2)} \setminus (\mathcal{A}_1 \cup \mathcal{A}_2)$.

 $M^{out}(k)$ is the solution on the soliton region of $M^{(2)}(k)$, which is defined as no jump on \mathbb{C} , only discrete spectral points, and analytical in $\mathcal{A}_1 \cup \mathcal{A}_2$ and outside discrete spectral points, that is, the following RHP

Riemann-Hilbert Problem 17. Find a matrix-valued function $M^{out}(k)$ with following properties:

- Analyticity: $M^{out}(k)$ is is analytical in $\mathbb{C} \setminus \Sigma^{(2)} \cup \mathcal{K} \cup \mathcal{K}^*$;
- Symmetry: $M^{out}(k) = \sigma_0(M^{out}(k^*))^* \sigma_0^{-1};$ Analytic behavior: $M^{out}(k) = I + \mathcal{O}(k^{-1}), \quad k \to \infty,$

• Residue conditions: $M^{out}(k)$ has simple poles at each point in $\mathcal{K} \cup \mathcal{K}^*$ satisfying the same residue relations with $(7.3) of M_{rhp}^{(2)}(k).$

8. Pure RHP and its asymptotic behavior

In this segment, we will probe into the asymptotic behavior of the external soliton region and the internal non soliton region.

8.1. External soliton solution region. As we all know, solitons are generated when the reflection data is equal to 0, that is, r(k) = 0. At this time, the trace function is simplified to

$$s_{11}(k) = \prod_{j=1}^{N} \frac{k - k_j}{k - k_j^*}, \ s_{22}(k) = \prod_{j=1}^{N} \frac{k - k_j^*}{k - k_j}.$$
(8.1)

And the jump matrix V(k) = I, then RHP 6 can be simplified to

Riemann-Hilbert Problem 18. For a given scattering data $\mathcal{P} = \{(k_j, c_j)\}_{j=1}^N$, a matrix $M(k|\mathcal{P})$ can be found to satisfy:

- Analyticity: $M(k|\mathcal{P})$ is analytical in $\mathbb{C}\setminus (\Sigma^{(2)}\cup\mathcal{K}\cup\mathcal{K}^*)$;
- Symmetry: $M(k|\mathcal{P}) = \sigma_0 M^*(k^*|\mathcal{P}) \sigma_0^{-1}$
- Asymptotic behaviors: $M(k|\mathcal{P}) = I + \mathcal{O}(k^{-1}), \quad k \to \infty,$
- Residue conditions: $M(k|\mathcal{P})$ has simple poles at each point in $k_j \in \mathbb{C}^+$ and $k_i^* \in \mathbb{C}^-$ with:

$$\operatorname{Res}_{k=k_{j}} M(k|\mathcal{P}) = \lim_{k \to k_{j}} M(k|\mathcal{P}) \begin{pmatrix} 0 & 0 \\ c_{j}e^{2it\theta(k_{j})} & 0 \end{pmatrix},$$

$$\operatorname{Res}_{k=k_{j}^{*}} M(k|\mathcal{P}) = \lim_{k \to k_{j}^{*}} M(k|\mathcal{P}) \begin{pmatrix} 0 & -c_{j}^{*}e^{-2it\theta(k_{j}^{*})} \\ 0 & 0 \end{pmatrix}.$$
(8.2)

The uniqueness of RHP 18 solution can be easily proved by using Liouville theorem.

In order to facilitate future research, we divide the scattering data into two parts. Note $\nabla \subseteq \{1, ..., N\}$, and define

$$s_{11\nabla}(k) = \prod_{j \in \nabla} \frac{k - k_j}{k - k_j^*}$$

Next, make a modified transformation of $M(k|\mathcal{P})$ defined above as follows

$$M_{\nabla}(k \mid \mathcal{D}) = M(k \mid \mathcal{P})s_{11,\nabla}(k)^{\sigma_3}, \qquad (8.3)$$

where the scattering data are given by

$$\mathcal{D} = \left\{ \left(k_j, c'_j\right) \right\}_{j=1}^N, \quad c'_j = c_j s_{11,\nabla}(k)^2.$$

Therefore, $M_{\nabla}(k \mid \mathcal{D})$ satisfies the following modified discrete RHP:

Riemann-Hilbert Problem 19. For a given scattering data $\mathcal{D} = \{(k_j, c'_j)\}_{j=1}^N$, a matrix $M_{\nabla}(k \mid \mathcal{D})$ can be found to satisfy:

- Analyticity: $M^{\nabla}(k \mid \mathcal{D})$ is analytical in $\mathbb{C} \setminus (\Sigma^{(2)} \cup \mathcal{K} \cup \mathcal{K}^*)$;
- Symmetry: $M^{\nabla}(k \mid \mathcal{D}) = \sigma_0 M^*_{\nabla}(k^* \mid \mathcal{D}) \sigma_0^-$
- Asymptotic behaviors: $M_{\nabla}(k \mid \mathcal{D}) = I + \mathcal{O}(k^{-1}), \quad k \to \infty;$
- Residue conditions: $M_{\nabla}(k \mid D)$ has simple poles at each point in $k_j \in \mathbb{C}^+$ and $k_j^* \in \mathbb{C}^-$ with:

$$\operatorname{Res}_{k=k_{j}} M_{\nabla}(k \mid \mathcal{D}) = \begin{cases} \lim_{k \to k_{j}} M_{\nabla}(k \mid \mathcal{D}) \begin{pmatrix} 0 & \varpi_{j}^{-1}(s_{11,\nabla}')^{-2}(k_{j}) \\ 0 & 0 \end{pmatrix}, j \in \nabla, \\ \lim_{k \to k_{j}} M_{\nabla}(k \mid \mathcal{D}) \begin{pmatrix} 0 & 0 \\ \varpi_{j}s_{11,\nabla}^{2}(k_{j}) & 0 \end{pmatrix}, j \notin \nabla. \end{cases}$$

$$\operatorname{Res}_{k=k_{j}^{*}} M_{\nabla}(k \mid \mathcal{D}) = \begin{cases} \lim_{k \to k_{j}^{*}} M_{\nabla}(k \mid \mathcal{D}) \begin{pmatrix} 0 & 0 \\ -\varpi_{j}^{*-1}(s_{11,\nabla}^{*+})^{-2}(k_{j}^{*}) & 0 \end{pmatrix}, j \in \nabla, \\ \lim_{k \to k_{j}^{*}} M_{\nabla}(k \mid \mathcal{D}) \begin{pmatrix} 0 & -\varpi_{j}^{*}(s_{11,\nabla}^{*})^{2}(k_{j}^{*}) \\ 0 & 0 \end{pmatrix}, j \notin \nabla. \end{cases}$$

$$(8.4)$$

where $\varpi_j = c_j e^{2it\theta(k_j)}$.

Proposition 20. If $A_{sol}(x,t) = A_{sol}(x,t|\mathcal{D})$, $B_{sol}(x,t) = B_{sol}(x,t|\mathcal{D})$ denote the N-soliton solution of the system (1.1), for the scattering data without reflection $\mathcal{D} = \{(k_j, c'_j)\}_{j=1}^N$, RHP 19 has a unique solution and

$$A_{sol}(x,t|\mathcal{D}) = 4i \lim_{k \to \infty} [kM_{\nabla}(k \mid \mathcal{D})]_{12}$$

= $4i \lim_{k \to \infty} [kM(k \mid \mathcal{P})]_{12} = A_{sol}(x,t|\mathcal{P}),$
$$B_{sol}(x,t|\mathcal{D}) = -\frac{4i}{\beta} \lim_{k \to \infty} \frac{d}{dt} [kM_{\nabla}(k \mid \mathcal{D})]_{11}$$

= $-\frac{4i}{\beta} \lim_{k \to \infty} \frac{d}{dt} [kM(k \mid \mathcal{P})]_{11} = B_{sol}(x,t|\mathcal{P}).$

8.2. Existence and uniqueness of solutions for external RH problems. We see that $M^{out}(k)$ is a reflection soliton solution, and the reflection mainly comes from T(k). In order to connect $M^{out}(x,t,k)$ with the case of non reflection scattering $\mathcal{D} = \{(k_j, c'_j)\}_{j=1}^N$, an idea is to take $\nabla = \triangle_{k_0}^-$ in Eq.(8.3). So there are

$$T(k) = \prod_{j \in \Delta_{k_0}^-} \frac{k - k_j^*}{k - k_j} \exp\left(i \int_{-k_0}^{k_0} \frac{\nu(s)}{s - k} dk\right) = s_{11, \Delta_{k_0}^-} \delta(k),$$

$$T(k_j)^{-2} = s_{11, \Delta_{k_0}^-} (k_j)^2 \delta(k_j)^{-2}, \quad (\frac{1}{T})'(k_j)^{-2} = s'_{11, \Delta_{k_0}^-} (k_j)^{-2} \delta(k_j)^2.$$

The scattering data can be written as

$$\widetilde{\mathcal{D}} = \{(k_j, \widetilde{c}_j)\}_{j=1}^N, \quad \widetilde{c}_j = \begin{cases} c_j^{-1} s'_{11, \Delta_{k_0^-}}(k_j)^{-2} \,\delta(k_j)^2, & j \in \Delta_{k_0}^-\\ c_j s_{11, \Delta_{k_0^-}}(k_j)^2 \,\delta(k_j)^{-2}, & j \notin \Delta_{k_0}^- \end{cases}.$$

$$(8.5)$$

Therefore, RHP 19 is rewritten as

Riemann-Hilbert Problem 21. For a given scattering data $\tilde{\mathcal{D}} = \{(k_j, \tilde{c}_j)\}_{j=1}^N$, a matrix $M_{\Delta_{k_0}^-}(k \mid \tilde{\mathcal{D}})$ can be found to satisfy:

- $M_{\Delta_{k_0}^-}(k \mid \tilde{\mathcal{D}})$ is analytical in $\mathbb{C} \setminus (\mathcal{K} \cup \mathcal{K}^*);$
- $$\begin{split} \bullet \ & M_{\Delta_{k_0}^-}(k \mid \tilde{\mathcal{D}}) = \sigma_0 M_{\Delta_{k_0}^-}^*(k^* \mid \tilde{\mathcal{D}}) \sigma_0^{-1}; \\ \bullet \ & M_{\Delta_{k_0}^-}(k \mid \tilde{\mathcal{D}}) = I + \mathcal{O}\left(k^{-1}\right), \quad k \to \infty; \end{split}$$
- $M_{\Delta_{k_0}^-}(k \mid \tilde{\mathcal{D}})$ has simple poles at each point in $k_j \in \mathbb{C}^+$ and $k_j^* \in \mathbb{C}^-$ with:

$$\operatorname{Res}_{k=k_{j}}^{k}M_{\Delta_{k_{0}}^{-}}(k \mid \tilde{\mathcal{D}}) = \begin{cases}
\operatorname{lim}_{k \to k_{j}}^{k}M_{\Delta_{k_{0}}^{-}}(k \mid \tilde{\mathcal{D}}) \begin{pmatrix} 0 & \Lambda_{j}^{-1}s_{11,\Delta_{k_{0}}^{-2}}^{\prime - 2}(k_{j}) \\ 0 & 0 \end{pmatrix}, j \in \Delta_{k_{0}}^{-}, \\
\operatorname{lim}_{k \to k_{j}}^{k}M_{\Delta_{k_{0}}^{-}}(k \mid \tilde{\mathcal{D}}) \begin{pmatrix} 0 & \Lambda_{j}^{-1}s_{11,\Delta_{k_{0}}^{-2}}^{\prime - 2}(k_{j}) \\ 0 & 0 \end{pmatrix}, j \notin \Delta_{k_{0}}^{-}, \\
\operatorname{lim}_{k \to k_{j}^{*}}^{k}M_{\Delta_{k_{0}}^{-}}(k \mid \tilde{\mathcal{D}}) \begin{pmatrix} 0 & \Lambda_{j}s_{11,\Delta_{k_{0}}^{-}}^{2}(k_{j}) & 0 \\ -\Lambda_{j}s_{11,\Delta_{k_{0}}^{-2}}^{\prime - 2}(k_{j}^{*}) & 0 \end{pmatrix}, j \in \Delta_{k_{0}}^{-}, \\
\operatorname{lim}_{k \to k_{j}^{*}}^{k}M_{\Delta_{k_{0}}^{-}}(k \mid \tilde{\mathcal{D}}) \begin{pmatrix} 0 & -\Lambda_{j}s_{11,\Delta_{k_{0}}^{*-2}}^{\ast - 2}(k_{j}^{*}) \\ 0 & 0 \end{pmatrix}, j \notin \Delta_{k_{0}}^{-}.
\end{cases}$$
(8.6)




FIGURE 9. The cone $\mathcal{C}(x_1, x_2, v_1, v_2)$.

where $\Lambda_j = c_j \delta^{-2} e^{2it\theta(k_j)}$.

So we have the following corollary

Corollary 22. There exists and unique solution for the RHP 17, moreover

 $M^{out}(x,t,k) = M_{\Delta_{k_0}^-}(x,t,k \mid \tilde{\mathcal{D}}),$

where scattering data $\tilde{\mathcal{D}}$ is given by (8.5) and

$$\begin{aligned} A_{sol}\left(x,t|\mathcal{D}\right) &= 4i \lim_{k \to \infty} \left[kM^{out}(x,t,k \mid \mathcal{D})\right]_{12} \\ &= 4i \lim_{k \to \infty} \left[kM_{\Delta_{k_0}^-}(x,t,k \mid \tilde{\mathcal{D}})\right]_{12} = A_{sol}\left(x,t|\tilde{\mathcal{D}}\right), \\ B_{sol}\left(x,t|\mathcal{D}\right) &= -\frac{4i}{\beta} \lim_{k \to \infty} \frac{d}{dt} \left[kM^{out}(x,t,k \mid \mathcal{D})\right]_{11} \\ &= -\frac{4i}{\beta} \lim_{k \to \infty} \frac{d}{dt} \left[kM_{\Delta_{k_0}^-}(x,t,k \mid \tilde{\mathcal{D}})\right]_{11} = B_{sol}\left(x,t|\tilde{\mathcal{D}}\right) \end{aligned}$$

8.3. The long-time behavior of soliton solution. This section mainly considers the asymptotic behavior of soliton solutions. On the basis of the residue condition (8.6), let $N = 1, k = \zeta + i\eta$, under the condition of no scattering, the soliton solution of the coupled dispersion AB system is

$$A(x,t) = 4\eta sech(2\eta(x + \frac{\alpha}{4|k|^2}t))e^{i\zeta(-2x + \frac{\alpha}{2|k|^2}t)},$$

$$B(x,t) = -\frac{2\alpha\eta^2}{\beta|k|^2}sech(\eta(2x + \frac{\alpha}{2|k|^2}t))^2.$$
(8.7)

It can be seen that the velocity of soliton solution is $v = -\frac{\alpha}{4|k|^2}$. Suppose $x_1 < x_2$ with $x_1, x_2 \in \mathbb{R}$ and $v_1 < v_2$ with $v_1, v_2 \in \mathbb{R}^-$, a conical region can be defined as

$$\mathcal{C}(x_1, x_2, v_1, v_2) = \{(x, t) \in \mathbb{R}^2 \mid x = x_0 + vt, x_0 \in [x_1, x_2], v \in [v_1, v_2]\}$$

Then make the following marks

$$\mathcal{I} = \{k : f(v_2) < |k| < f(v_1)\}, \quad f(v) \doteq \left(-\frac{\alpha}{4v}\right)^{\frac{1}{2}}, \\ \mathcal{K}(\mathcal{I}) = \{k_j \in \mathcal{K} : k_j \in \mathcal{I}\}, \quad N(\mathcal{I}) = |\mathcal{K}(\mathcal{I})|, \\ \mathcal{K}^+(\mathcal{I}) = \{k_j \in \mathcal{K} : |k_j| > f(v_1)\}, \\ \mathcal{K}^-(\mathcal{I}) = \{k_j \in \mathcal{K} : |k_j| < f(v_2)\}, \\ c_j^{\pm}(\mathcal{I}) = c_j \prod_{\operatorname{Re} k_n \in I_{\pm} \setminus \mathcal{I}} \left(\frac{k_j - k_n}{k_j - k_n^*}\right)^2 \exp\left[\pm \frac{1}{\pi i} \int_{I_{\pm}} \frac{\log[1 + r(\varsigma)r^*(\varsigma^*)]}{\varsigma - k_j} d\varsigma\right].$$

$$(8.8)$$

Proposition 23. Let $\nabla = \Delta_{k_0}^{\mp}$ in $\mathcal{C}(x_1, x_2, v_1, v_2)$ on RHP 19, and ensure the following estimation when $t \to \infty$:

$$\left\|\Theta_{j}^{\Delta_{k_{0}}^{\mp}}\right\| = \begin{cases} \mathcal{O}(1), & k_{j} \in \mathcal{K}(\mathcal{I}), \\ \mathcal{O}\left(e^{-2\mu t}\right), & k_{j} \in \mathcal{K} \setminus \mathcal{K}(\mathcal{I}), \end{cases} \quad t \to \infty$$

$$(8.9)$$

where

$$\mu = \min_{k_j \in \mathcal{K} \setminus \mathcal{K}(\mathcal{I})} \left\{ \operatorname{Im} k_j \operatorname{dist} \left(v_{k_j} - v \right) \right\}, \ v_j = -\frac{\alpha}{4|k_j|^2}$$

and in the formula, $v_j - v$ represents the difference between the speed of a soliton and the speed corresponding to k.




FIGURE 10. Here, the original data has eight pairs zero points of discrete spectrum, but insider the cone C only two pairs points with $\mathcal{K}(\mathcal{I}) = \{k_j \in \mathcal{K} : k_j \in \mathcal{I}\}, j = 2, 3.$

Proof. Here we take $\nabla = \triangle_{k_0}^-$ as an example to prove the above estimation. When $k = k_j \in \mathcal{K}^-(\mathcal{I})$ and $(x,t) \in \mathcal{C}(x_1, x_2, t_1, t_2)$, then the remainder condition for equation (8.4)

$$|\varpi_j(x_0 + vt, k_j)| \le c|e^{-2i\theta(k_j)t}|.$$

Next, focus on the index part

$$-2i\theta(k_j)t = -2i(k_j\frac{x}{t} - \frac{\alpha}{4k_j})t = -2i(k_j\frac{x_0 + vt}{t} - \frac{\alpha}{4k_j})t = -2ik_jx_0 - 2i(k_jv - \frac{\alpha}{4k_j})t$$

Its real part is

so there

$$Re(-2i\theta(k_j)t) = 2Imk_j x_0 - 2Imk_j t(v_j - v),$$
$$e^{-2i\theta(k_j)t}| = e^{2Imk_j x_0} e^{-2Imk_j t(v_j - v)} \le ce^{-2\mu t}.$$

For the discrete spectrum $k_j \in \mathcal{K} \setminus \mathcal{K}(\mathcal{I})$, make a disk D_k with a sufficiently small radius at each point and do not intersect each other, and define the function

$$\Gamma(k) = \begin{cases} I - \frac{1}{k - k_j} \Theta_{\Delta_{\mathcal{K}}^{\pm}}^{j}, & k \in D_k, \\ I - \frac{1}{k - k_j^*} \sigma_2 \Theta_{\Delta_{\mathcal{K}}^{\pm}}^{*j} \sigma_2, & k \in \bar{D}_k, \\ I, & elsewhere \end{cases}$$
(8.10)

where

$$\Theta_{\Delta_{\mathcal{K}}^{\pm}}^{j} = \begin{cases} \begin{pmatrix} 0 & \varpi_{j}^{-1}(s_{11,\nabla}')^{-2}(k_{j}) \\ 0 & 0 \end{pmatrix}, j \in \nabla, \\ \begin{pmatrix} 0 & 0 \\ \\ \varpi_{j}s_{11,\nabla}^{2}(k_{j}) & 0 \end{pmatrix}, & j \notin \nabla. \end{cases}$$

Make transformation

$$\widetilde{M}_{\Delta_{k_0}^{\pm}}\left(k \mid \widetilde{D}_{\Delta_{k_0}^{\pm}}\right) = M_{\Delta_{k_0}^{\pm}}\left(k \mid \widetilde{D}_{\Delta_{k_0}^{\pm}}\right) \Gamma(k),$$

and $\widetilde{M}_{\Delta_{k_0}^{\pm}}\left(k \mid \widetilde{D}_{\Delta_{k_0}^{\pm}}\right)$ satisfies the following relationship

$$\widetilde{M}_{\Delta_{k_0}^{\pm}}\left(k \mid \widetilde{D}_{\Delta_{k_0}^{\pm}}\right) = M_{\Delta_{k_0}^{\pm}}\left(k \mid \widetilde{D}_{\Delta_{k_0}^{\pm}}\right)\widetilde{V}(k), k \in \widetilde{\Sigma} = \bigcup_{k_j \in \mathcal{K} \setminus \mathcal{K}(\mathcal{I})} \partial D_k \cup \partial \overline{D}_k.$$
(8.11)

The jump matrix $\widetilde{V}(k) = \Gamma(k)$ is known based on the above proposition 8

$$\|\widetilde{V}(k) - I\|_{L^{\infty}(\widetilde{\Sigma})} = \mathcal{O}\left(e^{-2\mu t}\right).$$
(8.12)

In addition, let

$$M_{0}(k) = \widetilde{M}_{\Delta_{k_{0}}^{-}}\left(k \mid \widetilde{\mathcal{D}}_{\Delta_{k_{0}}^{-}}\right) \left[M_{\Delta_{k_{0}}^{-}}\left(k \mid \widetilde{\mathcal{D}}(\mathcal{I})\right)\right]^{-1},$$

$$(8.13)$$

where $\widetilde{\mathcal{D}}(\mathcal{I}) = \{k_j, c_j(\mathcal{I})\}, \ c_j(\mathcal{I}) = c_j \prod_{\operatorname{Re} k_n \in I_{\pm} \setminus \mathcal{I}} \left(\frac{k_j - k_n}{k_j - k_n^*}\right)^2$. it is obvious that $\widetilde{M}_{\Delta_{k_0}^-}(k|\widetilde{\mathcal{D}})$ and $M_{\Delta_{k_0}^-}(k|\widetilde{\mathcal{D}})$ have the same residue condition, so $M_0(k)$ has no poles, its jump is

$$M_0^+(k) = M_0^-(k) V_{M_0}(k), \ k \in \widetilde{\Sigma}.$$
(8.14)

Further, the form of jump matrix $V_{M_0}(k)$ is

$$V_{M_0}(k) = M_{\Delta_{k_0}^-}\left(k \mid \widetilde{\mathcal{D}}(\mathcal{I})\right) \widetilde{V}(k) M_{\Delta_{k_0}^-}\left(k \mid \widetilde{\mathcal{D}}(\mathcal{I})\right)^{-1},\tag{8.15}$$

according to equation (8.12)

$$\|V_{M_0}(k) - I\|_{L^{\infty}(\widetilde{\Sigma})} = \|\tilde{V}(k) - I\|_{L^{\infty}(\widetilde{\Sigma})} = \mathcal{O}\left(e^{-2\mu t}\right), \quad t \to \infty.$$
(8.16)

Due to the small norm RH property, it is known that $M_0(k)$ exists and

$$M_0(k) = I + \mathcal{O}\left(e^{-2\mu t}\right), \quad t \to \infty.$$

So there are

$$M_{\Delta_{k_0}^{\pm}}\left(k \mid \widetilde{\mathcal{D}}\right) = \left(I + \mathcal{O}\left(e^{-2\mu t}\right)\right) M_{\Delta_{k_0}^{\pm}}\left(k \mid \widetilde{\mathcal{D}}(\mathcal{I})\right).$$

$$(8.17)$$

Corollary 24. $\widetilde{\mathcal{D}}$ represents the scattering data of the equation (1.1), corresponding to the N-soliton solution of the coupled dispersion AB system, and $\mathcal{D}(\mathcal{I})$ represents the scattering data on $\mathcal{K}(\mathcal{I})$, when $x, t \in \mathcal{C}(x_1, x_2, v_1, v_2)$

$$\begin{split} A_{sol}\left(x,t|\widetilde{\mathcal{D}}\right) &= 4i \lim_{k \to \infty} \left[kM_{\Delta_{k_0}^{\pm}}\left(x,t,k \mid \widetilde{\mathcal{D}}\right)\right]_{12} \\ &= 4i \lim_{k \to \infty} \left[kM_{\Delta_{k_0}^{\pm}(\mathcal{I})}(x,t,k \mid \widetilde{\mathcal{D}}(\mathcal{I})\right]_{12} + \mathcal{O}\left(e^{-2\mu t}\right), \quad t \to \infty, \\ B_{sol}\left(x,t|\widetilde{\mathcal{D}}\right) &= -\frac{4i}{\beta} \lim_{k \to \infty} \frac{d}{dt} \left[kM_{\Delta_{k_0}^{\pm}}\left(x,t,k \mid \widetilde{\mathcal{D}}\right)\right]_{11} \\ &= -\frac{4i}{\beta} \lim_{k \to \infty} \frac{d}{dt} \left[kM_{\Delta_{k_0}^{\pm}(\mathcal{I})}(x,t,k \mid \widetilde{\mathcal{D}}(\mathcal{I})\right]_{11} + \mathcal{O}\left(e^{-2\mu t}\right), \quad t \to \infty. \end{split}$$

8.4. Solvable RHP near phase point. From Eq.(7.6), we can see that ||V(k) - I|| has no consistent small jump in the neighborhood \mathcal{A}_1 and \mathcal{A}_2 when $t \to \infty$, so we need to establish a local model $M^{in}(k)$, which completely matches the jump of $M_{rhp}(k)$ on $(\Sigma^{(2)} \cap \mathcal{A}_1) \cup (\Sigma^{(2)} \cap \mathcal{A}_2)$ of the error function $M^{err}(k)$, so as to achieve a consistent estimation of the attenuation of the transition.

Riemann-Hilbert Problem 25. For a matrix $M^{AB}(k)$, the following properties are satisfied: • $M^{AB}(k)$ is analyticity in $\mathbb{C} \setminus \Sigma^{AB}$, $\Sigma^{AB} = \bigcup_j \Sigma^j$, $j = 1, 2^+, 3^+, 4, 5, 6^+, 7^+, 8$;

- - $M^{AB}(k) = I + \mathcal{O}(k^{-1}), \quad k \to \infty;$
 - $M^{AB}(k)$ has continuous boundary values $M^{AB}_{\pm}(k)$ on Σ^{AB} and

$$M_+^{AB}(k) = M_-^{AB}(k)V^{AB}|_{\mathcal{A}_1 \cup \mathcal{A}_2}(k), \quad k \in \Sigma^{AB},$$

where

$$V^{AB}|_{\mathcal{A}_{1}\cup\mathcal{A}_{2}}(k) = \begin{cases} \begin{pmatrix} 1 & 1 & 0 \\ r\left(-k_{0}\right)\delta^{-2}\left(-k_{0}\right)\left(k+k_{0}\right)^{-2iv\left(-k_{0}\right)}e^{2it\theta} & 1 \end{pmatrix}, & k \in \Sigma_{1}, \\ \begin{pmatrix} 1 & \frac{r^{*}\left(-k_{0}\right)\delta^{2}\left(-k_{0}\right)}{1+|r\left(-k_{0}\right)|^{2}}\left(k+k_{0}\right)^{2iv\left(-k_{0}\right)}e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_{2}^{+}, \\ \begin{pmatrix} 1 & \frac{r^{*}\left(k_{0}\right)\delta^{2}\left(k_{0}\right)}{1+|r\left(k_{0}\right)|^{2}}\left(k-k_{0}\right)^{2iv\left(k_{0}\right)}e^{-2it\theta} \\ 1 & r^{*}\left(-k_{0}\right)\delta^{-2}\left(k_{0}\right)\left(k+k_{0}\right)^{-2iv\left(-k_{0}\right)}e^{-2it\theta} \\ 1 & r^{*}\left(-k_{0}\right)\delta^{2}\left(-k_{0}\right)\left(k+k_{0}\right)^{2iv\left(-k_{0}\right)}e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_{8}, \\ \begin{pmatrix} 1 & r^{*}\left(-k_{0}\right)\delta^{-2}\left(-k_{0}\right)\left(k+k_{0}\right)^{-2iv\left(-k_{0}\right)}e^{-2it\theta} \\ 1+|r\left(-k_{0}\right)|^{2}}\left(k+k_{0}\right)^{-2iv\left(-k_{0}\right)}e^{2it\theta} & 1 \end{pmatrix}, & k \in \Sigma_{7}^{+}, \\ \begin{pmatrix} \frac{r\left(-k_{0}\right)\delta^{-2}\left(-k_{0}\right)}{1+|r\left(-k_{0}\right)|^{2}}\left(k-k_{0}\right)^{-2iv\left(k_{0}\right)}e^{-2it\theta} \\ 1-r^{*}\left(k_{0}\right)\delta^{2}\left(\xi_{2}\right)\left(k-k_{0}\right)^{2iv\left(k_{0}\right)}e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_{5}. \end{cases}$$

This is mainly because when there is no discrete spectrum, $T_0(\pm k_0)$ can be reduced to $\delta(\pm k_0)$ and $1 - \Upsilon_{\mathcal{K}}(k) = 1$.

Here, we need to consider the well-known PC model near the two phase points, as shown in Fig.11. First, expand $\theta(k)$ at k_0 point

$$\theta(k) = -\frac{1}{2}\frac{\alpha}{k_0} - \frac{1}{4}\frac{\alpha}{k_0^3}(k-k_0)^2 + \frac{1}{4}\frac{\alpha}{k_0^4}(k-k_0)^3,$$
(8.18)



FIGURE 11. The jump contour Jump profile of local RHP near phase points $\pm k_0$

then define the following transformation

$$\mathbb{T}: f(k) \to (\mathbb{T}f)(k) = f\left(\sqrt{\frac{-k_0^3}{\alpha t}}\zeta + k_0\right),\tag{8.19}$$

so there is on Σ_4

$$r(k_{0}) \delta^{-2}(k_{0}) (k-k_{0})^{-2i\nu(k_{0})} e^{2it\theta}$$

= $r(k_{0}) \delta^{-2}(k_{0}) \left(\sqrt{\frac{-k_{0}^{3}}{\alpha t}} \zeta_{1}\right)^{-2i\nu(k_{0})} e^{2it(-\frac{1}{2}\frac{\alpha}{k_{0}} - \frac{1}{4}\frac{\alpha}{k_{0}^{3}}(k-k_{0})^{2} + \frac{1}{4}\frac{\alpha}{k_{0}^{4}}(k-k_{0})^{3})$
= $r(k_{0}) \delta^{-2}(k_{0}) e^{i\nu(k_{0})(\ln(k_{0}^{3}) - \ln(-\alpha t))} \zeta_{1}^{-2i\nu(k_{0})} e^{\frac{i\zeta_{1}^{2}}{2}} e^{-i\alpha t[\frac{1}{k_{0}} - \frac{1}{2}\frac{1}{k_{0}^{4}}(k-k_{0})^{3}]},$

we set

$$r_{k_0} = r(k_0) \,\delta^{-2}(k_0) \,e^{i\nu(k_0)\left(\ln(k_0^3) - \ln(-\alpha t)\right)} e^{-i\alpha t\left[\frac{1}{k_0} - \frac{1}{2}\frac{1}{k_0^4}(k - k_0)^3\right]},\tag{8.20}$$

in the same way, we can calculate the values of $\Sigma_2^+, \Sigma_3^+, \Sigma_4, \Sigma_6^+, \Sigma_7^+$ and Σ_8 . Therefore, the jump matrix of V^{AB} in $\mathcal{A}_1 \cup \mathcal{A}_2$ can be rewritten as

$$V^{AB}|_{\mathcal{A}_{1}\cup\mathcal{A}_{2}}(k) = \begin{cases} \begin{pmatrix} 1 & 0 \\ r_{-k_{0}}\zeta^{-2i\nu(-k_{0})}e^{i\zeta^{2}/2} & 1 \end{pmatrix}, & k \in \Sigma_{1}, \\ \begin{pmatrix} 1 & \frac{r^{*}_{-k_{0}}}{1+|r_{-k_{0}}|^{2}}\zeta^{2i\nu(-k_{0})}e^{-i\zeta^{2}/2} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_{2}^{+}, \\ \begin{pmatrix} 1 & \frac{r^{*}_{k_{0}}}{1+|r_{k_{0}}|^{2}}\zeta^{2i\nu(k_{0})}e^{-i\zeta^{2}/2} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_{4}, \\ \begin{pmatrix} 1 & r^{*}_{-k_{0}}\zeta^{2i\nu(-k_{0})}e^{-i\zeta^{2}/2} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_{8}, \\ \begin{pmatrix} 1 & r^{*}_{-k_{0}}\zeta^{2i\nu(-k_{0})}e^{i\zeta^{2}/2} & 1 \\ \frac{1}{1+|r_{-k_{0}}|^{2}}\zeta^{-2i\nu(-k_{0})}e^{i\zeta^{2}/2} & 1 \end{pmatrix}, & k \in \Sigma_{7}^{+}, \\ \begin{pmatrix} 1 & 0 \\ \frac{r_{-k_{0}}}{1+|r_{-k_{0}}|^{2}}\zeta^{-2i\nu(k_{0})}e^{i\zeta^{2}/2} & 1 \\ 1 & r^{*}_{k_{0}}\zeta^{2i\nu(k_{0})}e^{-i\zeta^{2}/2} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_{6}^{+}, \\ \begin{pmatrix} 1 & r^{*}_{k_{0}}\zeta^{2i\nu(k_{0})}e^{-i\zeta^{2}/2} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_{5}. \end{cases}$$

If the PC model considered in k_0 is expanded in the following form

$$M_{k_0}^{PC}(\zeta) = I + \frac{M_1^{pc}(k_0)}{i\zeta} + \mathcal{O}(\zeta^{-2}),$$

where

$$M_1^{PC}(k_0) = \begin{pmatrix} 0 & \Xi_{12}(k_0) \\ -\Xi_{21}(k_0) & 0 \end{pmatrix},$$

$$\Xi_{12}(k_0) = \frac{\sqrt{2\pi}e^{i\pi/4}e^{-\pi\nu(k_0)/2}}{r_1\Gamma(-i\nu(k_0))},$$

$$\Xi_{21}(k_0) = -\frac{\sqrt{2\pi}e^{-i\pi/4}e^{-\pi\nu(k_0)/2}}{r_1^*\Gamma(i\nu(k_0))}.$$




FIGURE 12. The jump contour Σ^{err} for the $M^{err}(k)$.

The same is true for $-k_0$. According to the transformation equation (8.19), the relationship between M^{AB} and M^{PC} is

$$M^{AB} = I + \frac{1}{i\zeta} (M_1^{PC}(k_0) + M_1^{PC}(-k_0)) + \mathcal{O}(\zeta^{-2}),$$

further can be written as

$$M^{AB}(x,t,k) = I + \sqrt{\frac{-k_0^3}{\alpha t}} \frac{M_1^{PC}(k_0)}{k-k_0} + \sqrt{\frac{-k_0^3}{\alpha t}} \frac{M_1^{PC}(-k_0)}{k+k_0} + \mathcal{O}(\zeta^{-2}).$$
(8.21)

From this, we get a consistent estimate

$$\left|M^{AB} - I\right| \lesssim O\left(t^{-\frac{1}{2}}\right). \tag{8.22}$$

Near the local circle of $\pm k_0$ have

$$\left|\frac{1}{k \mp k_0}\right| < c,$$

where c is a constant. Finally, we use M^{AB} to define a local model in $\pm k_0$

$$M^{(in)}(k) = M^{out}(k)M^{AB}(k),$$

it has the same jump matrix as M_{rhp} and is a bounded function in \mathcal{A}_1 and \mathcal{A}_2 .

8.5. Small norm RH problem of $M^{err}(k)$. In the disk $\mathcal{A}_1 \cup \mathcal{A}_2$, $M^{PC}(k)$ is consistent with the jump matrix of $M_{rhp}^{(2)}(k)$, but $M^{out}(k)$ does not have a jump. Therefore, the matrix $M^{err}(k)$ defined by (7.5) erases the jump of $M_{rhp}^{(2)}(k)$ inside the disk $\mathcal{A}_1 \cup \mathcal{A}_2$, and there is still a jump from $M_{rhp}^{(2)}(k)$ outside the disk $\mathbb{C} \setminus \mathcal{A}_1 \cup \mathcal{A}_2$. Therefore, the jump path of $M^{err}(k)$ is

$$\Sigma^{err} = \bigcup_{n=1}^{2} \partial \mathcal{A}_{n} \cup \left(\Sigma^{(2)} \setminus \bigcup_{n=1}^{2} \mathcal{A}_{n} \right) +$$

where ∂A_n is clockwise, see Fig.12. $M^{err}(k)$ can be directly verified to meet the following RHP:

Riemann-Hilbert Problem 26. For a matrix $M^{err}(k)$, the following properties are satisfied:

- $M^{err}(k)$ is analytical in $\mathbb{C}\backslash\Sigma^{err}$; $M^{err}(k) = \sigma_0(M^{err}(k^*))^*\sigma_0^{-1}$; $M^{err}(k) = I + \mathcal{O}(k^{-1}), \quad k \to \infty$;
- $M^{err}(k)$ has continuous boundary values $M^{err}_{\pm}(k)$ on Σ^{err} and

$$M_{+}^{err}(k) = M_{+}^{err}(k)V^{err}(k), \quad k \in \Sigma^{err},$$

where

$$V^{err}(k) = \begin{cases} M^{out}(k)V^{(2)}(k)M^{out}(k)^{-1}, & k \in \Sigma^{(2)} \setminus \bigcup_{n=1}^{2} \mathcal{A}_{n}, \\ M^{out}(k)M^{AB}(k)M^{out}(k)^{-1}, & k \in \bigcup_{n=1}^{2} \partial \mathcal{A}_{n}. \end{cases}$$

Proof. Here we mainly prove the form of $V^{err}(k)$. When $k \in \Sigma^{(2)} \setminus \bigcup_{n=1}^{2} \mathcal{A}_n$, $M^{err}(k) = M_{rhp}^{(2)}(k)(M^{out}(k))^{-1}$ and $M^{out}(k)$ does not jump in (7.5), then

$$\begin{aligned} M^{err}_{+}(k) &= M^{(2)}_{rhp+}(k)(M^{out}(k))^{-1} = M^{(2)}_{rhp-}(k)V^{(2)}(k)(M^{out}(k))^{-1} \\ &= M^{err}_{-}(k)V^{err}(k) = M^{2}_{rhp-}(k)M^{out}(k)^{-1}V^{err}(k), \end{aligned}$$

obviously,

$$V^{err} = M^{out}(k)V^{(2)}(M^{out}(k))^{-1}, \ k \in \Sigma^{(2)} \setminus \bigcup_{n=1}^{2} \mathcal{A}_{n}.$$

When $k \in \bigcup_{n=1}^{2} \mathcal{A}_n$, we have

$$M^{err} = M^{(2)}_{rhp}(k)(M^{(in)}(k))^{-1} = M^{(2)}_{rhp}(k)(M^{AB}(k))^{-1}(M^{out}(k))^{-1},$$

and $M_{rhp}^{(2)}(k)$ does not jump in $\cup_{n=1}^2 \partial \mathcal{A}_n$, then

$$M_{-}^{err}(k) = M_{rhp}^{(2)}(k)(M^{AB}(k))^{-1}(M^{out}(k))^{-1}.$$

 $M_{+}^{err}(k) = M_{rhp}^{(2)}(k)(M^{out}(k))^{-1},$

For $k \in \bigcup_{n=1}^{2} \partial \mathcal{A}_n$,

so from

we get

$$M_+^{err}(k) = M_-^{err}(k)V^{err}(k), \ k \in \bigcup_{n=1}^2 \partial \mathcal{A}_n,$$

$$V^{err}(k) = M^{out}(k)M^{AB}(k)(M^{out}(k))^{-1}, \ k \in \bigcup_{n=1}^{2} \partial \mathcal{A}_{n}.$$

Next, we will show that the small norm RH problem can be well solved by the error function $M^{err}(k)$ for large time.

It is known from equations (5.1) and (7.6) that the jump matrix satisfies the following estimation

$$\|V^{err} - I\| \lesssim O\left(e^{-2t\varepsilon}\right), \ k \in \Sigma^{(2)} \setminus \left(\mathcal{A}_1 \cup \mathcal{A}_2\right).$$
(8.23)

In addition, it is known from equations (8.22) that when $k \in \bigcup_{n=1}^{2} \partial A_n$, $V^{err}(k)$ satisfies the following estimation

$$\|V^{err}(k) - I\| = \left| M^{out}(k)^{-1} \left(M^{AB}(k) - I \right) M^{out}(k) \right| = \mathcal{O}\left(t^{-\frac{1}{2}} \right).$$
(8.24)

Proposition 27. RHP26 has a unique solution in the form of

$$M^{err}(k) = I + \frac{1}{2\pi i} \int_{\Sigma^{err}} \frac{\mu^{err}(s) \left(V^{err}(s) - I\right)}{s - k} ds,$$
(8.25)

where $\mu^{err} \in L^2(\Sigma^{err})$, meet $(1 - C_{err}) \mu^{err} = I$, C_{err} is Cauchy projection operator, defined as

$$C_{err}f(z) = \lim_{k' \to k \in \Sigma^{err}} \frac{1}{2\pi i} \int_{\Sigma^{err}} \frac{f(s)}{s - k'} ds.$$

Proof. It can be seen from Eq.(8.24) that the operator is bounded. It can be seen from Ref. [33]

$$\|\mu\|_{L^{2}(\Sigma^{err})} \lesssim \frac{\|C_{err}\|}{1 - \|C_{err}\|} \lesssim t^{-\frac{1}{2}},\tag{8.26}$$

obviously, $1 - C_{err}$ is reversible, so M^{err} is unique.

Proposition 28. Consider the asymptotic expansion of M^{err} when $k \to \infty$

$$M^{err}(k) = I + \frac{M_1^{err}}{k} + \mathcal{O}\left(\frac{1}{k^2}\right),\tag{8.27}$$

where

$$M_{1}^{err} = \frac{1}{i} \sqrt{\frac{-k_{0}^{3}}{\alpha t}} M^{out}(k_{0}) M_{1}^{PC}(k_{0}) (M^{out})^{-1}(k_{0}) - \frac{1}{i} \sqrt{\frac{-k_{0}^{3}}{\alpha t}} M^{out}(-k_{0}) M_{1}^{PC}(-k_{0}) (M^{out})^{-1}(-k_{0}) + \mathcal{O}(t^{-1}).$$

Proof. From Proposition 10,

$$M_1^{err} = -\frac{1}{2\pi i} \int_{\Sigma^{err}} \mu^{err}(s) \left(V^{err}(s) - I \right) ds,$$

then there are

$$M_{1}^{err} = -\frac{1}{2\pi i} \int_{\Sigma^{err}} (V^{err} - I) \, ds - \frac{1}{2\pi i} \int_{\Sigma^{err}} (\mu^{err}(s) - I) \, (V^{err} - I) \, ds$$

$$= -\frac{1}{2\pi i} \oint_{\bigcup_{n=1}^{2} \partial \mathcal{A}_{n}} (V^{err} - I) \, ds - \frac{1}{2\pi i} \int_{\Sigma^{err} \setminus \bigcup_{n=1}^{2} \partial \mathcal{A}_{n}} (V^{err} - I) \, ds$$

$$- \frac{1}{2\pi i} \int_{\Sigma^{err}} (\mu^{err}(s) - I) \, (V^{err} - I) \, ds.$$

(8.26) M_{1}^{err} can be written as

Using equations (8.23), (8.26), M_1^{err} can be written as

$$M_1^{err} = -\frac{1}{2\pi i} \oint_{\bigcup_{n=1}^2 \partial \mathcal{A}_n} (V^{err}(s) - I) \, ds + O\left(t^{-1}\right).$$

And using equations (8.21) and (8.24), and residue theorem there are

$$M_1^{err} = \frac{1}{i} \sqrt{\frac{-k_0^3}{\alpha t}} M^{out}(k_0) M_1^{PC}(k_0) (M^{out})^{-1}(k_0) - \frac{1}{i} \sqrt{\frac{-k_0^3}{\alpha t}} M^{out}(-k_0) M_1^{PC}(-k_0) (M^{out})^{-1}(-k_0) + \mathcal{O}(t^{-1}).$$

9. Asymptotic analysis on the pure $\bar{\partial}$ -problem

In this section, we mainly solve the part of $\bar{\partial}\mathcal{R}^{(2)}(k) \neq 0$ in $M^{(3)}(k)$, that is, RHP 16. Its solution can be expressed in the following form

$$M^{(3)}(k) = I - \frac{1}{\pi} \iint_{\mathbb{C}} \frac{M^{(3)}W^{(3)}}{s-k} \, \mathrm{d}A(s), \tag{9.1}$$

where $W^{(3)} = M_{rhp}^{(2)}(k)\bar{\partial}R^{(2)}M_{rhp}^{(2)}(k)^{-1}$, and dA(s) is the Lebesgue measure on the real plane. In fact, Eq.(9.1) can also be written in the form of an operator

$$(I - S)M^{(3)}(k) = I, (9.2)$$

where $\mathbb S$ is Cauchy-Green operator,

$$\mathbb{S}[f](k) = -\frac{1}{\pi} \iint_{\mathbb{C}} \frac{f(s)W(s)}{s-k} \, \mathrm{d}A(s).$$

If the operator $(I - S)^{-1}$ exists, then the above equation has a solution. Next, we will prove the existence of operator $(I - S)^{-1}$ in regions D_5 and D_4^{\pm} , which can be proved by similar methods in other regions. Before proving the existence of operator $(I - S)^{-1}$, we give the following lemma

Lemma 29.

$$\left| \bar{\partial} R_{5}^{(2)} e^{2i\theta t} \right| \lesssim \left(\left| \bar{\partial} \Upsilon_{\mathcal{K}}(s) \right| + \left| p_{5}'(\operatorname{Re}(k)) \right| + \left| k - k_{0} \right|^{-\frac{1}{2}} \right) e^{-c_{1}|u||v|t}, \ k \in D_{5}$$

$$\bar{\partial} R_{7}^{(3)} e^{-2i\theta t} \left| \lesssim \left\{ \begin{array}{l} \left(\left| \bar{\partial} \Upsilon_{\mathcal{K}}(s) \right| + \left| p_{4}'(\operatorname{Re}(k)) \right| + \frac{1}{|k|^{1/2}} \right) e^{-c_{2}|v|t}, \ k \in D_{4}^{-1} \\ \left(\left| \bar{\partial} \Upsilon_{\mathcal{K}}(s) \right| + \left| p_{4}'(\operatorname{Re}(k)) \right| + \left| k - k_{0} \right|^{-\frac{1}{2}} \right) e^{-c_{1}|u||v|t}. \ k \in D_{4}^{+} \end{array} \right.$$

$$(9.3)$$

Proof. Recalling the above condition $\alpha < 0$, for the convenience of analysis, the exponential part can be written as $-\frac{\alpha vt}{2}(\frac{1}{u^2+v^2}-\frac{1}{k_0^2})$, which is a decreasing function of u. In the D_5 region, the index part has

$$-\frac{\alpha vt}{2} \left(\frac{1}{(u+k_0)^2 + v^2} - \frac{1}{k_0^2} \right) = -\frac{\alpha vt}{2} \left(\frac{1}{(u+k_0)^2 + v^2} - \frac{1}{k_0^2} \right)$$
$$= -\frac{\alpha vt}{2} \left(\frac{-u^2 - 2k_0 u - v^2}{((u+k_0)^2 + v^2)(k_0^2)} \right)$$
$$\leq -c_1 |u| |v| t.$$

There is a similar analysis in area D_4^+ . In region D_4^- , k = u + iv and $0 \le v < u$ the index part has

$$\frac{\alpha vt}{2} \left(\frac{1}{u^2 + v^2} - \frac{1}{k_0^2} \right) = \frac{\alpha vt}{2} \left(\frac{1}{u^2 + v^2} - \frac{1}{k_0^2} \right)$$

$$\leq -c_2 |v|t.$$
as equation (9.3).

The overall conclusion is summarized as equation (9.3).

Therefore, our next goal is to prove the existence of $(I - S)^{-1}$.

Proposition 30. For sufficiently large t, operator S is a small norm and has

$$\|\mathbb{S}\|_{L^{\infty} \to L^{\infty}} \leqslant ct^{-\frac{1}{4}},\tag{9.4}$$

therefore, $(I - \mathbb{S})^{-1}$ exists.

Proof. This is discussed in detail in the D_5 region, and other regions can be obtained similarly. Let $s = u + k_0 + iv$, $k = \zeta + i\eta$, for any $f \in L^{\infty}$, we have

$$\begin{aligned} |\mathbb{S}(f)| &\leq \frac{1}{\pi} \iint_{D_5} \frac{\left| f M_{rhp}^{(2)} \bar{\partial} R^{(2)} M_{rhp}^{(2)-1} \right|}{|s-k|} dA(s) \\ &\leq \frac{1}{\pi} \|f\|_{L^{\infty}} \left\| M_{rhp}^{(2)} \right\|_{L^{\infty}} \left\| M_{rhp}^{(2)-1} \right\|_{L^{\infty}} \int_{0}^{\infty} \int_{k_0+v}^{\infty} \frac{\left| \bar{\partial} R_5 \right| e^{\frac{\alpha v t}{2} \left(\frac{1}{k_0^2} - \frac{1}{(u+k_0)^2 + v^2}\right)}}{|s-k|} du dv \\ &\leq c \left(\Omega_1 + \Omega_2 + \Omega_3\right), \end{aligned}$$
(9.5)

where

$$\Omega_{1} = \int_{0}^{\infty} \int_{k_{0}+v}^{\infty} \frac{\left|\bar{\partial}\Upsilon_{\mathcal{K}}(s)\right| e^{\frac{\alpha v t}{2} \left(\frac{1}{k_{0}^{2}} - \frac{1}{(u+k_{0})^{2}+v^{2}}\right)}}{|s-k|} du dv,$$

$$\Omega_{2} = \int_{0}^{\infty} \int_{k_{0}+v}^{\infty} \frac{|r'(\operatorname{Re}(s))| e^{\frac{\alpha v t}{2} \left(\frac{1}{k_{0}^{2}} - \frac{1}{(u+k_{0})^{2}+v^{2}}\right)}}{|s-k|} du dv,$$

$$\Omega_{3} = \int_{0}^{\infty} \int_{k_{0}+v}^{\infty} \frac{|s-k_{0}|^{-\frac{1}{2}} e^{\frac{\alpha v t}{2} \left(\frac{1}{k_{0}^{2}} - \frac{1}{(u+k_{0})^{2}+v^{2}}\right)}}{|s-k|} du dv.$$

Next, the main idea is to estimate Ω_j , j = 1, 2, 3. An important inequality to know in advance is

$$\left\|\frac{1}{s-k}\right\|_{L^{2}(v+k_{0},\infty)}^{2} = \int_{v+k_{0}}^{\infty} \frac{1}{|s-k|^{2}} du \le \int_{-\infty}^{\infty} \frac{1}{|s-k|^{2}} du$$
$$= \int_{-\infty}^{\infty} \frac{1}{(u-\zeta)^{2} + (v-\eta)^{2}} du = \frac{1}{|v-\eta|} \int_{-\infty}^{\infty} \frac{1}{1+y^{2}} dy = \frac{\pi}{|v-\eta|},$$
(9.6)

where $y = \frac{u-\zeta}{v-\eta}$. For Ω_1 , it can be obtained by direct calculation using Eq.(9.6)

$$\Omega_{1} \leq \int_{0}^{\infty} \left\| \bar{\partial} \Upsilon_{\mathcal{K}}(s) \right\|_{L^{2}} \left\| \frac{1}{s-k} \right\|_{L^{2}} dv \leq c_{3} \int_{0}^{\infty} \frac{e^{-\frac{\alpha vt}{2} \left(\frac{1}{(u+k_{0})^{2}+v^{2}} - \frac{1}{k_{0}^{2}} \right)}}{\sqrt{|\eta-v|}} dv \\ \leq c_{3} \int_{0}^{\infty} \frac{e^{-c_{1}|u||v|t}}{\sqrt{|\eta-v|}} dv \leq c_{3} \int_{0}^{\infty} \frac{e^{-c_{1}|v|^{2}t}}{\sqrt{|\eta-v|}} dv \\ \leq ct^{-\frac{1}{4}}.$$
(9.7)

Use the same method to estimate Ω_2 . For the estimation of Ω_3 , Here we need the help of Hölder inequality with p > 2and $\frac{1}{p} + \frac{1}{q} = 1$.

$$\left\| |s - k_0|^{-\frac{1}{2}} \right\|_{L^p(k_0 + v, \infty)} = \left(\int_{k_0 + v}^{\infty} |u - k_0 + iv|^{-\frac{p}{2}} du \right)^{\frac{1}{p}} = \left(\int_{v}^{\infty} (u^2 + v^2)^{-\frac{p}{4}} du \right)^{\frac{1}{p}} = \left(\int_{v}^{\infty} (u^2 + v^2)^{-\frac{p}{4}} du \right)^{\frac{1}{p}} = v^{\frac{1}{p} - \frac{1}{2}} \left(\int_{1}^{\infty} (1 + x^2)^{-\frac{p}{4}} dx \right)^{\frac{1}{p}} \le cv^{\frac{1}{p} - \frac{1}{2}}.$$
(9.8)

Similar estimates can be proved

$$\left\|\frac{1}{s-k}\right\|_{L^{q}(v,\infty)} \le c|v-\eta|^{\frac{1}{q}-1}, \quad \frac{1}{q}+\frac{1}{p}=1.$$

Then we have

$$\Omega_{3} \leq c \int_{0}^{+\infty} \left\| |s - k_{0}|^{-\frac{1}{2}} \right\|_{L^{p}} \left\| \frac{1}{s - k} \right\|_{L^{q}} e^{-\frac{\alpha v t}{2} \left(\frac{1}{(u + k_{0})^{2} + v^{2}} \right) - \frac{1}{k_{0}^{2}} dv} \\
\leq c \int_{0}^{+\infty} \left\| |s - k_{0}|^{-\frac{1}{2}} \right\|_{L^{p}} \left\| \frac{1}{s - k} \right\|_{L^{q}} e^{-c_{1}|u||v|t} dv \\
\leq c \int_{0}^{+\infty} \left\| |s - k_{0}|^{-\frac{1}{2}} \right\|_{L^{p}} \left\| \frac{1}{s - k} \right\|_{L^{q}} e^{-c_{1}v^{2}t} dv \\
\leq c \left[\int_{0}^{\eta} v^{\frac{1}{p} - \frac{1}{2}} |v - \eta|^{\frac{1}{q} - 1} e^{-c_{1}v^{2}t} dv + \int_{\eta}^{\infty} v^{\frac{1}{p} - \frac{1}{2}} |v - \eta|^{\frac{1}{q} - 1} e^{-c_{1}v^{2}t} dv \right] \\
\leq H_{1} + H_{2},$$
(9.9)

the following is the estimation of the two integrals. For H_1 , since $0 \le v < \eta$, the following estimates are obtained by replacing $v = w\eta$ with variables

$$H_{1} = \int_{0}^{\eta} v^{\frac{1}{p} - \frac{1}{2}} (\eta - v)^{\frac{1}{q} - 1} e^{-c_{1}v^{2}t} dv$$
$$= \int_{0}^{1} \eta^{\frac{1}{2}} w^{\frac{1}{p} - \frac{1}{2}} (1 - w)^{\frac{1}{q} - 1} e^{-c_{1}w^{2}\eta^{2}t} dw$$
$$\leq ct^{-\frac{1}{4}}.$$

For H_2 , we let $w = v - \eta$, then

$$H_{2} = \int_{\eta}^{\infty} v^{\frac{1}{p} - \frac{1}{2}} (v - \eta)^{\frac{1}{q} - 1} e^{-c_{1}v^{2}t} dv$$

$$= \int_{0}^{\infty} (w + \eta)^{\frac{1}{p} - \frac{1}{2}} w^{\frac{1}{q} - 1} e^{-c_{1}(w + \eta)^{2}t} dw$$

$$\leq ct^{-\frac{1}{4}} \int_{0}^{\infty} w^{-\frac{1}{2}} e^{-c_{1}(w)^{2}t} dw$$

$$\leq ct^{-\frac{1}{4}}.$$

So combining the estimates of H_1 and H_2 , we get $\Omega_3 \leq ct^{-\frac{1}{4}}$.

In the D_4^- region, the estimation is still divided into three parts, at this time, there is s = u + iv.

$$\begin{split} |\mathbb{S}(f)| &\leq \frac{1}{\pi} \iint_{D_{4}^{-}} \frac{\left| f M_{rhp}^{(2)} \bar{\partial} R^{(2)} M_{rhp}^{(2)-1} \right|}{|s-k|} dA(s) \\ &\leq \frac{1}{\pi} \| f \|_{L^{\infty}} \left\| M_{rhp}^{(2)} \right\|_{L^{\infty}} \left\| M_{rhp}^{(2)-1} \right\|_{L^{\infty}} \iint_{D_{4}^{-}} \frac{\left| \bar{\partial} R_{3} \right| e^{\frac{\alpha v t}{2} \left(\frac{1}{u^{2}+v^{2}} - \frac{1}{k_{0}^{2}} \right)}}{|s-k|} dA(s) \\ &\leq c \left(\Omega_{1}' + \Omega_{2}' + \Omega_{3}' \right), \end{split}$$
(9.10)

where

$$\begin{split} \Omega_1' &= \int_0^{\frac{k_0}{2}} \int_v^{\frac{k_0}{2}} \frac{\left| \bar{\partial} \Upsilon_{\mathcal{K}}(s) \right| e^{\frac{\alpha v t}{2} \left(\frac{1}{u^2 + v^2} - \frac{1}{k_0^2}\right)}}{|s - k|} du dv, \\ \Omega_2' &= \int_0^{\frac{k_0}{2}} \int_v^{\frac{k_0}{2}} \frac{|r'(\operatorname{Re}(s))| e^{\frac{\alpha v t}{2} \left(\frac{1}{u^2 + v^2} - \frac{1}{k_0^2}\right)}}{|s - k|} du dv, \\ \Omega_3' &= \int_0^{\frac{k_0}{2}} \int_v^{\frac{k_0}{2}} \frac{|s|^{-\frac{1}{2}} e^{\frac{\alpha v t}{2} \left(\frac{1}{u^2 + v^2} - \frac{1}{k_0^2}\right)}}{|s - k|} du dv. \end{split}$$

Combined with lemma (9.3) and Eq.(9.6), we can estimate each integral. For Ω'_1 ,

$$\Omega_{1}' \lesssim \int_{0}^{\frac{k_{0}}{2}} e^{-c_{2}|v|t} \int_{v}^{\frac{k_{0}}{2}} \frac{r'(u)}{|s-k|} du dv \leq c \int_{0}^{\frac{k_{0}}{2}} \left\| \frac{1}{s-k} \right\|_{L^{2}} e^{-c_{2}|v|t} dv \leq c \int_{0}^{\infty} \frac{e^{-c_{2}|v|t}}{\sqrt{|\eta-v|}} dv \leq ct^{-\frac{1}{2}}.$$
(9.11)

The estimation of Ω'_2 is similar to that of Ω'_1 , the estimation method of Ω'_3 is similar to that of Ω_3 , and can get

$$\Omega_j' \le ct^{-\frac{1}{2}}, \ j = 2, 3. \tag{9.12}$$

Therefore, this proves that all regions meet (9.4).

Next, consider the expansion of $M^{(3)}$,

$$M^{(3)}(k) = I + \frac{M_1^{(3)}(x,t)}{k} + \mathcal{O}(k^{-2}), \quad k \to \infty,$$

from the previous equation (9.1), it is easy to know

$$M_1^{(3)}(x,t) = \frac{1}{\pi} \int_{\mathbb{C}} M^{(3)}(s) W^{(3)}(s) dA(s).$$

Further, we can prove that

Proposition 31. For a large t, we have

$$M_1^{(3)} \le ct^{-\frac{3}{4}}.$$
(9.13)

Proof. Owing to $M_{rhp}^{(2)}$ is bounded outside the pole, there is

$$\begin{split} \left| M_{1}^{(3)} \right| &\leq \frac{1}{\pi} \iint_{D_{5}} \left| M^{(3)} M_{rhp}^{(2)} \bar{\partial} R^{(2)} M_{rhp}^{(2)-1} \right| dA(s) \\ &\leq \frac{1}{\pi} \left\| M^{(3)} \right\|_{L^{\infty}} \left\| M_{rhp}^{(2)} \right\|_{L^{\infty}} \left\| \left(M_{rhp}^{(2)} \right)^{-1} \right\|_{L^{\infty}} \iint_{D_{5}} \left| \bar{\partial} R_{5} e^{2it\theta} \right| dA(s) \\ &\leq c (\iint_{D_{5}} \left| \bar{\partial} \Upsilon_{\mathcal{K}}(s) \right| e^{-\frac{\alpha v t}{2} \left(\frac{1}{(u+k_{0})^{2}+v^{2}} - \frac{1}{k_{0}^{2}} \right)} dA(s) + \iint_{D_{5}} \left| p_{5}'(u) \right| e^{-\frac{\alpha v t}{2} \left(\frac{1}{(u+k_{0})^{2}+v^{2}} - \frac{1}{k_{0}^{2}} \right)} dA(s) \\ &+ \iint_{D_{5}} \frac{1}{|s-k_{0}|^{1/2}} e^{-\frac{\alpha v t}{2} \left(\frac{1}{(u+k_{0})^{2}+v^{2}} - \frac{1}{k_{0}^{2}} \right)} dA(s)) \\ &\leq c \left(\Omega_{4} + \Omega_{5} + \Omega_{6} \right). \end{split}$$
(9.14)

We constrain Ω_4 by using the Cauchy-Schwarz inequality

$$\begin{aligned} |\Omega_4| &\leq \int_0^\infty \left\| \bar{\partial} \Upsilon_{\mathcal{K}} \right\|_{L^2_u(v+k_0,\infty)} \left(\int_v^\infty e^{-c_1 t |u| |v|} du \right)^{\frac{1}{2}} dv \\ &\leq ct^{-\frac{1}{2}} \int_0^\infty \frac{e^{-4tv^2}}{\sqrt{v}} dv \\ &\leq ct^{-\frac{1}{4}} \int_0^\infty \frac{e^{-4w^2}}{\sqrt{w}} dw \leq ct^{-\frac{3}{4}}. \end{aligned}$$

Similar constraints can be done for Ω_5 . For the constraint of Ω_6 , we follow the method of Ω_3 and use Hölder inequality and Eq.(9.8) to obtain

$$\begin{split} I_{3} &= \int_{0}^{+\infty} \int_{k_{0}+v}^{+\infty} \left((u-k_{0})^{2} + v^{2} \right)^{-\frac{1}{4}} e^{-\frac{\alpha v t}{2} \left(\frac{1}{(u+k_{0})^{2}+v^{2}} - \frac{1}{k_{0}^{2}} \right)} du dv \\ &\leq \int_{0}^{+\infty} \left\| \left((u-k_{0})^{2} + v^{2} \right)^{-1/4} \right\|_{L^{p}} \left(\int_{k_{0}+v}^{+\infty} e^{-c_{1}q|u||v|t} du \right)^{\frac{1}{q}} dv \\ &\leq \int_{0}^{+\infty} v^{\frac{1}{p}-\frac{1}{2}} \left(\int_{k_{0}+v}^{+\infty} e^{-c_{1}q|u||v|t} du \right)^{\frac{1}{q}} dv \\ &\leq ct^{-\frac{1}{q}} \int_{0}^{+\infty} v^{\frac{2}{p}-\frac{3}{2}} e^{-c_{1}q|v+k_{0}||v|t} dv, \\ &\leq ct^{-\frac{3}{4}} \int_{0}^{\infty} w^{\frac{2}{p}-\frac{3}{2}} e^{-c_{1}qw^{2}t} dw \leq ct^{-\frac{3}{4}} \end{split}$$

Here we replace the variable $v = wt^{-\frac{1}{2}}$. Notice that here $2 and <math>-1 < \frac{2}{p} - \frac{3}{2} < -\frac{1}{2}$.

10. Long time asymptotic behavior of solution solution region for the coupled dispersive AB $$_{\rm SYSTEM}$$

After many deformations above, we now begin to construct the long-term asymptotic properties of the coupled dispersive AB system (1.1). Looking back at the previous transformation, we have

$$M(k) = M^{(3)}(k)M^{err}(k)M^{out}(k)\mathcal{R}^{(2)}(k)^{-1}T(k)^{\sigma_3}, \quad k \in \mathbb{C}\backslash \mathcal{A}_1 \cup \mathcal{A}_2.$$
(10.1)

In particular, in the vertical direction $k \in D_2, D_{10}$, there is $\mathcal{R}^{(2)} = I$, so we consider $k \to \infty$ in this region, so

$$M = \left(I + \frac{M_1^{(3)}}{k} + \dots\right) \left(I + \frac{M_1^{err}}{k} + \dots\right) \left(I + \frac{M_1^{out}}{k} + \dots\right) \left(I + \frac{T_1^{\sigma_3}}{k} + \dots\right).$$

In order to recover the potential, the coefficient of k^{-1} needs to be collected

$$M_1 = M_1^{out} + M_1^{err} + M_1^{(3)} + T_1^{\sigma_3}$$

so there

$$A = 4i(M_1^{out} + M_1^{err})_{12} + \mathcal{O}(t^{-\frac{3}{4}}),$$

$$B = -\frac{4i}{\beta}\frac{d}{dt}(M_1^{out} + M_1^{err})_{11} + \mathcal{O}(t^{-\frac{3}{4}}).$$
(10.2)

Based on the above formula, the specific asymptotic state can be written in the form of the following theorem

Theorem 32. Suppose $A_0, B_0 \in H^{1,1}(\mathbb{R})$ have general scattering data, and A(x,t) and B(x,t) are the solutions of equation (1.1). For fixed $x_1 < x_2$ with $x_1, x_2 \in \mathbb{R}$ and $v_1 < v_2$ with $v_1, v_2 \in \mathbb{R}^-$, a conical region can be defined as

$$\mathcal{C}(x_1, x_2, v_1, v_2) = \{(x, t) \in \mathbb{R}^2 \mid x = x_0 + vt, x_0 \in [x_1, x_2], v \in [v_1, v_2]\}$$

Then we define two regions for the spectral parameters

$$\mathcal{I} = \{k : f(v_2) < |k| < f(v_1)\}, \quad f(v) \doteq \left(-\frac{\alpha}{4v}\right)^{1/2}$$

as is shown in Fig. 10. Using $A_{sol}(x,t,k|\widetilde{\mathcal{D}})$ and $B_{sol}(x,t,k|\widetilde{\mathcal{D}})$ for the coupled dispersive AB system corresponding to $N(\mathcal{I}) \leq N$ modulated non reflection scattering data

$$\widetilde{D}(\mathcal{I}) = \{k_j, c_j(\mathcal{I})\}, \ c_j(\mathcal{I}) = c_j \prod_{k_n \in \mathcal{K}(\mathcal{I})} \left(\frac{k_j - k_n}{k_j - k_n^*}\right)^2$$

Then when $t \to \infty$ and $(x,t) \in \mathcal{C}(x_1, x_2, v_1, v_2)$, there is

$$\begin{split} A(x,t) &= A_{sol}(x,t | \widetilde{D}(\mathcal{I})) + t^{-\frac{1}{2}}(g_1 + g_2) + \mathcal{O}(t^{-\frac{3}{4}}), \\ B(x,t) &= B_{sol}(x,t | \widetilde{D}(\mathcal{I})) + t^{-\frac{1}{2}}(h_1 + h_2 + f_1 + f_2) + \mathcal{O}(t^{-\frac{3}{4}}), \end{split}$$

where

$$\begin{split} M^{out}\left(\pm k_{0}\right) &= \left(\begin{array}{cc} m_{11}(\pm k_{0}) & m_{12}(\pm k_{0}) \\ m_{21}(\pm k_{0}) & m_{22}(\pm k_{0}) \end{array}\right), \ j = 1,2 \\ g_{1} &= 4\sqrt{\frac{-k_{0}^{3}}{\alpha}}(m_{12}^{2}(-k_{0})\Xi_{21}(-k_{0}) + m_{11}^{2}(-k_{0})\Xi_{12}(-k_{0})), \\ g_{2} &= 4\sqrt{\frac{-k_{0}^{3}}{\alpha}}(m_{12}^{2}(k_{0})\Xi_{21}(k_{0}) + m_{11}^{2}(k_{0})\Xi_{12}(k_{0})), \\ h_{1} &= \frac{4}{\beta}\sqrt{\frac{-k_{0}^{3}}{\alpha}}\frac{d}{dt}(m_{12}(-k_{0})m_{22}(-k_{0})\Xi_{21}(-k_{0}) + m_{11}(-k_{0})m_{21}(-k_{0})\Xi_{12}(-k_{0})), \\ h_{2} &= \frac{4}{\beta}\sqrt{\frac{-k_{0}^{3}}{\alpha}}\frac{d}{dt}(m_{12}(k_{0})m_{22}(k_{0})\Xi_{21}(k_{0}) + m_{11}(k_{0})m_{21}(k_{0})\Xi_{12}(k_{0})), \\ f_{1} &= \frac{2}{\beta t}\sqrt{\frac{-k_{0}^{3}}{\alpha}}m_{12}(-k_{0})m_{22}(-k_{0})\Xi_{21}(-k_{0}) + m_{11}(-k_{0})m_{21}(-k_{0})\Xi_{12}(-k_{0}), \\ f_{2} &= \frac{2}{\beta t}\sqrt{\frac{-k_{0}^{3}}{\alpha}}m_{12}(k_{0})m_{22}(k_{0})\Xi_{21}(k_{0}) + m_{11}(k_{0})m_{21}(k_{0})\Xi_{12}(k_{0}). \end{split}$$

Appendix A: Estimation of characteristic function

In the previous assumption 2.1, there is no zero point of s_{11} on \mathbb{R} , so there is no pole of r(k) on \mathbb{R} . From Volterra integral formula (2.8)

$$\psi_{11}^{\pm}(x,0,k) = 1 + \int_{\pm\infty}^{x} \frac{1}{2}A(y,k)\psi_{21}^{\pm}dy,$$

$$\psi_{21}^{\pm}(x,0,k) = \int_{\pm\infty}^{x} -\frac{1}{2}A^{*}(y,k)\psi_{11}^{\pm}e^{2ik(x-y)}dy.$$

Let

$$\Psi_1^- = (\psi_{11}^-, \ \psi_{21}^-)^T.$$

For $k \in \mathbb{R}$, we introduce an operator mapping

$$\Re_0 f = \int_{-\infty}^x T_0 f(y) dy,$$

and

$$T_0 = \left(\begin{array}{cc} 0 & \frac{1}{2}A \\ -\frac{1}{2}A^* e^{2ik(x-y)} & 0 \end{array} \right).$$

It can be seen from this that

$$|\Re_0 f(x)| \le \int_{-\infty}^x \frac{1}{2} |A(y)| \, dy ||f||_{(L^{\infty}(-\infty,0])},$$

which means that \Re_0 is a bounded linear operator on $L^{\infty}(-\infty, 0]$. Further, it can be obtained by mathematical induction that

$$|\Re_0^n f(x)| \leq \frac{1}{n!} \left(\int_{-\infty}^x \frac{1}{2} |A(y)| \, dy \right)^n \|f\|_{(L^{\infty}(-\infty,0])}.$$
 Therefore, the following series is uniformly convergent on $x \in (-\infty, 0]$,

$$\Psi_1^-(x,k) = \sum_{n=0}^\infty \Re_0^n \begin{pmatrix} 1\\ 0 \end{pmatrix}.$$

(10.3)

So for $k \in \mathbb{R}$ and $x \in (-\infty, 0]$,

$$|\Psi_1(x,k)_1^-| \le e^{\frac{1}{2} ||A||_{L^1(\mathbb{R})}}.$$

Similarly, we can get

$$\begin{split} \left| \Psi_2^-(x,k) \right| &\leq e^{\frac{1}{2} \|A\|_{L^1(\mathbb{R})}}, x \in (-\infty,0];\\ \left| \Psi_1^+(x,k) \right| &\leq e^{\frac{1}{2} \|A\|_{L^1(\mathbb{R})}}, x \in [0,\infty);\\ \left| \Psi_2^+(x,k) \right| &\leq e^{\frac{1}{2} \|A\|_{L^1(\mathbb{R})}}, x \in [0,\infty). \end{split}$$

Let's look at the derivative of the characteristic function with respect to k. According to the uniform convergence property of series (10.3), we have

$$\Psi_{1,k}^{-}(x,k) = \sum_{n=0}^{\infty} \partial_k \Re_0^n \left(\begin{array}{c} 1\\ 0 \end{array}\right) = \sum_{n=1}^{\infty} \sum_{m=0}^n \Re_0^m \Re_0' \Re_0^{n-m-1} \left(\begin{array}{c} 1\\ 0 \end{array}\right) \quad ,$$

where the form of \Re'_0 is

$$\Re'_0 f(x) = \int_{-\infty}^x \left(\begin{array}{cc} 1 & 0 \\ 0 & e^{2ik(x-y)} \end{array} \right) \left(\begin{array}{c} \frac{1}{2}A(y)f_2 \\ -i(x-y)A^*f_1 \end{array} \right) dy.$$

Similarly, \Re'_0 is a bounded liner operator on $x \in (-\infty, 0]$ with

$$\|\Re_0'\|_{L^{\infty}(-\infty,0]} \le \|A\|_{L^{1,1}(\mathbb{R})}.$$

So there is

$$\Psi_{1,k}^{-}(x,k) \leq \sum_{n=1}^{\infty} \left| \partial_k \Re_0^n \begin{pmatrix} 1\\0 \end{pmatrix} \right| \leq \sum_{n=0}^{\infty} \frac{1}{2} \frac{\|A\|_{L^1(\mathbb{R})^{n-1}}}{(n-1)!} \|A\|_{L^{1,1}(\mathbb{R})}$$
$$\leq \|A\|_{L^{1,1}(\mathbb{R})} e^{\frac{1}{2} \|A\|_{L^1(\mathbb{R})}}$$

for $k \in \mathbb{R}$ and $x \in (-\infty, 0]$.

In similar steps, we can also get

$$\begin{split} \Psi_{2,k}^{-}(x,k) &\leq \|A\|_{L^{1,1}(\mathbb{R})} e^{\frac{1}{2}\|A\|_{L^{1}(\mathbb{R})}};\\ \Psi_{1,k}^{+}(x,k) &\leq \|A\|_{L^{1,1}(\mathbb{R})} e^{\frac{1}{2}\|A\|_{L^{1}(\mathbb{R})}};\\ \Psi_{2,k}^{+}(x,k) &\leq \|A\|_{L^{1,1}(\mathbb{R})} e^{\frac{1}{2}\|A\|_{L^{1}(\mathbb{R})}}. \end{split}$$

Next, we will estimate the characteristic function $(\Psi^{\pm})_{ij}(x,k), (i, j = 1, 2)$ respectively. In fact, there are the following lemma

Lemma 33. For $k \in \mathbb{R}$, we have the following estimates

$$\begin{split} \left\| \psi_{21}^{-}(x,0,k) \right\|_{C^{0}(\mathbb{R}^{-},L^{2}(I_{0}))} \lesssim \|A\|_{H^{1,1}}; \\ \left\| \psi_{21}^{-}(x,0,k) \right\|_{L^{2}(\mathbb{R}^{-}\times\mathbb{R})} \lesssim \|A\|_{H^{1,1}}; \\ \left\| \psi_{12}^{-}(x,0,k) \right\|_{C^{0}(\mathbb{R}^{-},L^{2}(\mathbb{R}))} \lesssim \|A\|_{H^{1,1}}; \\ \left\| \psi_{12}^{-}(x,0,k) \right\|_{L^{2}(\mathbb{R}^{-}\times\mathbb{R})} \lesssim \|A\|_{H^{1,1}}; \\ \left\| \psi_{12}^{+}(x,0,k) \right\|_{C^{0}(\mathbb{R}^{+},L^{2}(\mathbb{R}))} \lesssim \|A\|_{H^{1,1}}; \\ \left\| \psi_{12}^{+}(x,0,k) \right\|_{L^{2}(\mathbb{R}^{+}\times\mathbb{R})} \lesssim \|A\|_{H^{1,1}}; \\ \left\| \psi_{21}^{+}(x,0,k) \right\|_{C^{0}(\mathbb{R}^{+},L^{2}(\mathbb{R}))} \lesssim \|A\|_{H^{1,1}}; \\ \left\| \psi_{21}^{+}(x,0,k) \right\|_{L^{2}(\mathbb{R}^{+}\times\mathbb{R})} \lesssim \|A\|_{H^{1,1}}. \end{split}$$

Proof. From the above integral form, we can know that

$$\psi_{21}^{-}(x,0,k) = \int_{-\infty}^{x} -\frac{1}{2}A^{*}(y,k)\psi_{11}^{-}e^{2ik(x-y)}dy,$$

then for $k \in \mathbb{R}, \forall \sigma \in C_0(\mathbb{R})$, we compute

$$\begin{split} \left\| \int_{-\infty}^{x} -\frac{1}{2} A^{*} \psi_{11}^{-} e^{2ik(x-y)} dy \right\|_{L^{2}(I_{0})}^{2} &= \sup_{\|\phi\|=1} \int_{I_{0}} \sigma(k) \int_{-\infty}^{x} -\frac{1}{2} A^{*} \psi_{11}^{-} e^{2ik(x-y)} dy \\ &\lesssim c e^{\frac{1}{2} \|A\|_{L^{1}(\mathbb{R})}} \|A\|_{L^{1}} \,, \end{split}$$

this estimate is direct. In addition,

$$\int_{\mathbb{R}^{+}} \int_{\mathbb{R}} \left| \int_{-\infty}^{x} -\frac{1}{2} A^{*} \psi_{11}^{-} e^{2ik(x-y)} dy \right|^{2} dk dx \lesssim \int_{\mathbb{R}^{+}} \left| \int_{-\infty}^{x} -\frac{1}{2} A^{*} \psi_{11}^{-} e^{2ik(x-y)} dy \right|^{2} dx \lesssim \|A\|_{L^{1,1}},$$

which implies that for $k\in\mathbb{R}$

$$\left\| \psi_{21}^{-}(x,0,k) \right\|_{C^{0}(\mathbb{R}^{-},L^{2}(\mathbb{R}))} \lesssim \|A\|_{H^{1,1}}, \\ \left\| \psi_{21}^{-}(x,0,k) \right\|_{L^{2}(\mathbb{R}^{-}\times\mathbb{R})} \lesssim \|A\|_{H^{1,1}}.$$

Similarly, others can also be obtained.

Its derivative form for k has

$$\psi_{21,k}^{-}(x,0,k) = \int_{-\infty}^{x} -\frac{1}{2}A^{*}2i(x-y)\psi_{11}^{-}e^{2ik(x-y)}dy + \int_{-\infty}^{x} -\frac{1}{2}A^{*}\psi_{11,k}^{-}e^{2ik(x-y)}dy,$$

Similarly, the following estimates can be obtained

$$\left\| \psi_{21,k}^{-}(x,0,k) \right\|_{C^{0}(\mathbb{R}^{-},L^{2}(\mathbb{R}))} \lesssim \|A\|_{H^{1,1}} \\ \left\| \psi_{21,k}^{-}(x,0,k) \right\|_{L^{2}(\mathbb{R}^{-}\times\mathbb{R})} \lesssim \|A\|_{H^{1,1}}$$

According to Eq.(2.11)

$$s_{11}(k) - 1 = (\psi_{11}^- - 1)(\psi_{22}^+ - 1) + (\psi_{11}^- - 1) + (\psi_{22}^+ - 1) - \psi_{21}^- \psi_{12}^+$$

$$s_{12}(k) = e^{2ikx}(\psi_{12}^- \psi_{22}^+ - \psi_{22}^- \psi_{12}^+).$$

Let's examine $\psi_{11}^- - 1$ and $\psi_{22}^+ - 1$.

$$\psi_{11}^{-} - 1 = \int_{-\infty}^{x} \frac{1}{2} A(y) \psi_{21}^{-} dy,$$

$$\psi_{22}^{+} - 1 = \int_{-\infty}^{x} -\frac{1}{2} A^{*}(y) \psi_{12}^{+} dy,$$

it's easy to know that ψ_{11}^--1 is bounded and can be obtained by using Lemma 33 that

$$\begin{aligned} \left\| \psi_{11}^{-}(x,0,k) - 1 \right\|_{C^{0}(\mathbb{R}^{-},L^{2}(\mathbb{R}))} &\lesssim \|A\|_{H^{1,1}} \\ \left\| \psi_{11}^{-}(x,0,k) - 1 \right\|_{L^{2}(\mathbb{R}^{-} \times \mathbb{R})} &\lesssim \|A\|_{H^{1,1}}. \end{aligned}$$

The estimation of $\psi_{22}^+ - 1$ can be obtained similarly. For

$$s_{11,k} = \psi_{11,k}^- \psi_{22}^+ + \psi_{11}^- \psi_{22,k}^+ - \psi_{21,k}^- \psi_{12}^+ - \psi_{21}^- \psi_{12,k}^+$$

where

$$\psi_{11,k}^{-} = \int_{-\infty}^{x} \frac{1}{2} A(y,k) \psi_{21,k}^{-} dy, \ \psi_{22,k}^{+} = \int_{\infty}^{x} \frac{1}{2} A(y,k) \psi_{12,k}^{+} dy.$$

A similar operation is performed on $s_{12}(k)$. Tracing back to the above estimates, when $k \in \mathbb{R}$ and initial value $A \in H^{1,1}$, there is

$$s_{11}(k) \in L^2(\mathbb{R}), \ s_{11,k}(k) \in L^2(\mathbb{R}),$$

 $s_{12}(k) \in L^2(\mathbb{R}), \ s_{12,k}(k) \in L^2(\mathbb{R}).$

APPENDIX B: SOLVABLE PARABOLIC CYLINDER MODEL

Here we mainly describe the solution of the parabolic cylinder model introduced above. For the cAB equation studied in this paper, since there are two steady-state phase points, we need two parabolic cylinders to describe it. but their expansion forms are the same, using the following model.

For
$$r_0 \in \mathbb{R}$$
, if $\nu = -\frac{1}{2\pi} \log \left(1 + |r_0|^2 \right)$, define the contour $\Sigma^{PC} = \bigcup_{j=1}^4 \Sigma_j$,

$$\Sigma_j = \{\xi \in \mathbb{C} : \arg \xi = (2j-1)\pi/4\}, \quad j = 1, 2, 3, 4$$

These four contours divide the plane into six areas D_j , j = 1..6, as shown in Fig. 13.




FIGURE 13. The jump contour V^{PC} for the $M^{PC}(k)$.

Therefore, we can consider the RHP corresponding to the following parabolic cylinder model.

Riemann-Hilbert Problem 34. Find a matrix-valued function $M^{PC}(x,t,k)$ which satisfies:

- Analyticity: M(x,t,k) is analytic in $\mathbb{C} \setminus \Sigma_{PC}^{PC}$;
- Asymptotic behaviors: $M^{PC}(k) = I + \frac{M_1^{PC}}{ik} + O(k^{-2}), \quad k \to \infty;$ Jump condition: $M^{PC}(x,t,k)$ has continuous boundary values $M_{\pm}^{PC}(x,t,k)$ on \mathbb{R} and

$$M_{+}^{PC}(x,t,k) = M_{-}^{PC}(x,t,k)V^{PC}(k), \quad k \in \mathbb{R},$$
(10.4)

where

$$V^{PC}(k) = \begin{cases} \begin{pmatrix} 1 & 0 \\ r_0 k^{-2i\nu} e^{\frac{ik^2}{2}} & 1 \end{pmatrix}, & k \in \Sigma_1, \\ \begin{pmatrix} 1 & \frac{\bar{r}_0}{1+|r_0|^2} k^{2i\nu} e^{-\frac{ik^2}{2}} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_2, \\ \begin{pmatrix} 1 & 0 \\ \frac{r_0}{1+|r_0|^2} k^{-2i\nu} e^{\frac{ik^2}{2}} & 1 \end{pmatrix}, & k \in \Sigma_3, \\ \begin{pmatrix} 1 & \bar{r}_0 k^{2i\nu} e^{-\frac{ik^2}{2}} \\ 0 & 1 \end{pmatrix}, & k \in \Sigma_4. \end{cases}$$
(10.5)

Make changes

$$M^{PC} = \vartheta \mathcal{P} k^{-i\nu\sigma_3} e^{\frac{ik^2}{4}\sigma_3},\tag{10.6}$$

where

$$\mathcal{P}(k) = \begin{cases} \begin{pmatrix} 1 & 0 \\ -r_0 & 1 \end{pmatrix}, & k \in D_1, \\ \begin{pmatrix} 1 & \frac{-\bar{r}_0}{1+|r_0|^2} \\ 0 & 1 \end{pmatrix}, & k \in D_3, \\ \begin{pmatrix} 1 & 0 \\ \frac{r_0}{1+|r_0|^2} & 1 \end{pmatrix}, & k \in D_4, \\ \begin{pmatrix} 1 & \bar{r}_0 \\ 0 & 1 \end{pmatrix}, & k \in D_6, \\ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & k \in D_2 \cup D_5 \end{cases}$$

Then we can get a standard RHP with jump only in k = 0.

Riemann-Hilbert Problem 35. Find a matrix-valued function $\vartheta(x, t, k)$ which satisfies:

- Analyticity: ϑ(x,t,k) is analytic in C\R;
 Asymptotic behaviors: ϑ(k)e^{ik²}/₄σ₃k^{-iνσ₃} = I + O(k⁻¹), k→∞;
 Jump condition: ϑ₊(x,t,k) = ϑ₋(x,t,k)v(0).

The above RH problem can be reduced to Weber equation,

$$\partial_z^2 D_a(z) + \left[\frac{1}{2} - \frac{z^2}{4} + a\right] D_a(z) = 0,$$

where

$$z = e^{-\frac{3i\pi}{4}}k, \ a = i\Xi_{12}\Xi_{21} = i\nu,$$

$$\Xi_{12} = \frac{\sqrt{2\pi}e^{i\pi/4e^{-\pi\nu/2}}}{r_0\Gamma(-a)}, \quad \Xi_{21} = \frac{-\sqrt{2\pi}e^{-i\pi/4e^{-\pi\nu/2}}}{\bar{r}_0\Gamma(a)} = \frac{\nu}{\Xi_{12}}.$$

The solution of parabolic cylindrical explicit solution $\vartheta(k)$ can be given by parabolic cylindrical function. Obtained from Eq.(10.6)

$$M^{PC}(k) = I + \frac{M_1^{PC}}{ik} + O(k^{-2}),$$

where

$$M_1^{PC} = \left(\begin{array}{cc} 0 & \Xi_{12} \\ -\Xi_{21} & 0 \end{array} \right).$$

References

- [1] Belinskii V A, Zakharov V E. Integration of the Einstein equations by means of the inverse scattering problem technique and construction of exact soliton solutions. Zh. Eksp. Teor. Fiz. 1978;75(6): 1955-1971.
- [2] Kaup D J, Newell A C. An exact solution for a derivative nonlinear Schrödinger equation. J. Math. Phys. 1978;19(4): 798-801.
- [3] Arkadiev V A, Pogrebkov A K, Polivanov M C. Inverse scattering transform method and soliton solutions for Davey-Stewartson II equation. Physica D: Nonlinear Phenomena. 1989;36(1-2): 189-197.
- [4] Liu N, Guo B. Solitons and rogue waves of the quartic nonlinear Schrödinger equation by Riemann-Hilbert approach. Nonlinear Dynamics. 2020;100(1): 629-646.
- [5] Zhao L C, Liu J. Rogue-wave solutions of a three-component coupled nonlinear Schrödinger equation. Physical Review E. 2013;87(1): 013201.
- Hu J, Xu J, Yu G F. Riemann-Hilbert approach and N-soliton formula for a higher-order Chen-Lee-Liu equation. Journal of Nonlinear Mathematical Physics. 2018; 25(4): 633-649.
- [7] Yang B, Chen Y. High-order soliton matrices for Sasa-Satsuma equation via local Riemann-Hilbert problem. Nonlinear Anal-Real. 2019;45:918-941.
- [8] Li Y, Tian S F. Inverse scattering transform and soliton solutions of an integrable nonlocal Hirota equation. Communications on Pure, Applied Analysis. 2022, 21(1): 293.
- [9] Manakov S V. Nonlinear Fraunhofer diffraction. Zh. Eksp. Teor. Fiz, 1973, 65(4): 10.
- [10] Ablowitz M J, Newell A C. The decay of the continuous spectrum for solutions of the Korteweg-deVries equation. Journal of Mathematical Physics. 1973; 14(9): 1277-1284.
- [11] Zakharov V E, Manakov S V. Asymptotic behavior of non-linear wave systems integrated by the inverse scattering method. Soviet Journal of Experimental and Theoretical Physics. 1976; 44: 106.
- [12] Its A R. Asymptotics of solutions of the nonlinear schrödinger equation and isomonodromic deformations of systems of linear differential equations. Doklady Akademii Nauk. Russian Academy of Sciences. 1981; 261(1): 14-18.
- [13] Deift P, Zhou X. A steepest descent method for oscillatory Riemann-Hilbert problems. Bulletin of the American Mathematical Society. 1992;26(1): 119-123.
- [14] Zhou X, Deift P. Long-Time Behavior of the Non-focusing Nonlinear Schrödinger Equation-A Case Study, Lectures in Mathematical Sciences, Graduate School of Mathematical Sciences, University of Tokyo, 1994.
- [15] Grunert K, Teschl G. Long-time asymptotics for the Korteweg-de Vries equation via nonlinear steepest descent. Mathematical Physics, Analysis and Geometry. 2009; 12(3): 287-324.
- [16] de Monvel A B, Shepelsky D. A Riemann-Hilbert approach for the Degasperis-Procesi equation. Nonlinearity. 2013;26(7): 2081–2107.

[17] Cheng P J, Venakides S, Zhou X. Long-time asymptotics for the pure radiation solution of the sine-gordon equation. History and philosophy of logic. 1999; 24(7-8): 1195-1262.

- [18] Huang L, Lenells J. Nonlinear Fourier transforms for the sine-Gordon equation in the quarter plane. Journal of Differential Equations. 2018; 264(5): 3445-3499.
- [19] Xu J, Fan E. Long-time asymptotics for the Fokas-Lenells equation with decaying initial value problem: without solitons. Journal of Differential Equations. 2015; 259(3): 1098-1148.
- [20] Xu J, Fan E, Chen Y. Long-time asymptotic for the derivative nonlinear Schrödinger equation with step-like initial value. Mathematical Physics, Analysis and Geometry. 2013; 16(3): 253-288.
- [21] De Monvel A B, Kostenko A, Shepelsky D, Teschl G. Long-time asymptotics for the Camassa-Holm equation. SIAM journal on mathematical analysis. 2009; 41(4): 1559-1588.
- [22] de Monvel A B, Its A, Shepelsky D. Painlevé-type asymptotics for the Camassa-Holm equation. SIAM Journal on Mathematical Analysis. 2010;42(4): 1854-1873.
- [23] McLaughlin K T R, Miller P D. The $\bar{\partial}$ steepest descent method and the asymptotic behavior of polynomials orthogonal on the unit circle with fixed and exponentially varying nonanalytic weights. Int. Math. Res. Not. 2006;48673.
- [24] McLaughlin K T R, Miller P D. The $\bar{\partial}$ steepest descent method for orthogonal polynomials on the real line with varying weights. International Mathematics Research Notice. 2008; 075.
- [25] Dieng M, McLaughlin K D T. Dispersive asymptotics for linear and integrable equations by the Dbar steepest descent method, Nonlinear dispersive partial differential equations and inverse scattering. Fields Inst. Commun. 2019; 83:253-291.
- [26] Cuccagna S, Jenkins R. On asymptotic stability of N-solitons of the defocusing nonlinear Schrödinger equation. Commun. Math. Phys. 2016; 343: 921-969.
- [27] Borghese M, Jenkins R, McLaughlin K D T R. Long time asymptotic behavior of the focusing nonlinear Schrödinger equation. Ann. Inst. Henri Poincaré, Anal. Non Linéaire. 2018; 35(4): 887-920.
- [28] Jenkins R, Liu J, Perry P, Sulem C. Soliton resolution for the derivative nonlinear Schrödinger equation. Communications in Mathematical Physics. 2018; 363(3): 1003-1049.
- [29] Liu J, Perry P, Sulem C. Long-time behavior of solutions to the derivative nonlinear Schrödinger equation for soliton-free initial data, Ann.I. H. Poincaré-AN. 2018; 35: 217-265.
- [30] Giavedoni P. Long-time asymptotic analysis of the Korteweg-de Vries equation via the dbar steepest descent method: the soliton region. Nonlinearity. 2017; 30(3): 1165-1181.
- [31] Dieng M, McLaughlin K D T R, Miller P D. Dispersive Asymptotics for Linear and Integrable Equations by the Dbar Steepest Descent Method, in Nonlinear Dispersive Partial Differential Equations and Inverse Scattering. Springer, New York, NY, 2019; 253-291.
- [32] Yang J J, Tian S F, Li Z Q. Soliton resolution for the Hirota equation with weighted Sobolev initial data[J]. arXiv preprint arXiv:2101.05942, 2021.
- [33] Cheng Q, Fan E. Long-time asymptotics for the focusing Fokas-Lenells equation in the solitonic region of space-time. Journal of Differential Equations. 2022; 309: 883-948.
- [34] Yang Y, Fan E. On the long-time asymptotics of the modified Camassa-Holm equation in space-time solitonic regions. Advances in Mathematics, 2022; 402: 108340.
- [35] Yang Y, Fan E. Soliton resolution for the short-pulse equation. Journal of Differential Equations, 2021, 280: 644-689.
- [36] Gibbon J D, James I N, Moroz I M. An example of soliton behaviour in a rotating baroclinic fluid. Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences. 1979; 367(1729): 219-237.
- [37] Fowler A C, Gibbon J D, McGuinness M J. The real and complex Lorenz equations and their relevance to physical systems. 1983.
- [38] Gibbon J D, McGuinness M J. Amplitude equations at the critical points of unstable dispersive physical systems. Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences, 1981, 377(1769): 185-219.

- [39] Pedlosky J. Finite-amplitude baroclinic wave packets. Journal of Atmospheric Sciences. 1972; 29(4): 680-686.
- [40] Gibbon J D, James I N, Moroz I M. The sine-Gordon equation as a model for a rapidly rotating baroclinic fluid. Physica Scripta. 1979; 20(3-4): 402.
- [41] Kamchatnov A M, Pavlov M V. Periodic solutions and Whitham equations for the AB system. Journal of Physics A: Mathematical and General. 1995; 28(11): 3279.
- [42] Guo R, Hao H Q, Zhang L L. Dynamic behaviors of the breather solutions for the AB system in fluid mechanics. Nonlinear Dynamics, 2013;74(3): 701-709.
- [43] Guo R, Liu Y F. The canonical AB system: conservation laws and soliton solutions. Applied Mathematics and Computation, 2015; 259: 153-163.
- [44] Wang X, Li Y, Huang F, Chen Y. Rogue wave solutions of AB system. Communications in Nonlinear Science and Numerical Simulation, 2015; 20(2): 434-442.
- [45] Wen X Y, Yan Z. Modulational instability and higher-order rogue waves with parameters modulation in a coupled integrable AB system via the generalized Darboux transformation. Chaos: An Interdisciplinary Journal of Nonlinear Science. 2015; 25(12): 123115.
- [46] Gibbon J D, McGuinness M J. Amplitude equations at the critical points of unstable dispersive physical systems. Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences. 1981; 377(1769): 185-219.
- [47] Chen S, Yan Z. Long-time asymptotics of solutions for the coupled dispersive AB system with initial value problems. Journal of Mathematical Analysis and Applications. 2021; 498(2): 124966.
- [48] Yang J J, Tian S F, Li Z Q. Soliton resolution for the Hirota equation with weighted Sobolev initial data. arXiv preprint. arXiv:2101.05942, 2021.
- [49] Chen G, Liu J, Lu B. Long-time asymptotics and stability for the sine-Gordon equation. arXiv preprint. arXiv:2009.04260, 2020.

(JY) SCHOOL OF MATHEMATICAL SCIENCES, SHANGHAI KEY LABORATORY OF PURE MATHEMATICS AND MATHEMATICAL PRACTICE, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200241, PEOPLE'S REPUBLIC OF CHINA

(YC) School of Mathematical Sciences, Shanghai Key Laboratory of Pure Mathematics and Mathematical Practice, East China Normal University, Shanghai 200241, People's Republic of China

(YC) COLLEGE OF MATHEMATICS AND SYSTEMS SCIENCE, SHANDONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, QINGDAO 266590, PEOPLE'S REPUBLIC OF CHINA

Email address: ychen@sei.ecnu.edu.cn