

EQUATORIALLY BALANCED C_4 -FACE-MAGIC LABELINGS ON KLEIN BOTTLE GRID GRAPHS

STEPHEN J. CURRAN, RICHARD M. LOW, AND STEPHEN C. LOCKE

ABSTRACT. For a graph $G = (V, E)$ embedded in the Klein bottle, let $\mathcal{F}(G)$ denote the set of faces of G . Then, G is called a C_k -face-magic Klein bottle graph if there exists a bijection $f : V(G) \rightarrow \{1, 2, \dots, |V(G)|\}$ such that for any $F \in \mathcal{F}(G)$ with $F \cong C_k$, the sum of all the vertex labelings along C_k is a constant S . Let $x_v = f(v)$ for all $v \in V(G)$. We call $\{x_v : v \in V(G)\}$ a C_k -face-magic Klein bottle labeling on G . We consider the $m \times n$ grid graph, denoted by $\mathcal{K}_{m,n}$, embedded in the Klein bottle in the natural way. We show that for $m, n \geq 2$, $\mathcal{K}_{m,n}$ admits a C_4 -face-magic Klein bottle labeling if and only if n is even. We say that a C_4 -face-magic Klein bottle labeling $\{x_{i,j} : (i, j) \in V(\mathcal{K}_{m,n})\}$ on $\mathcal{K}_{m,n}$ is equatorially balanced if $x_{i,j} + x_{i,n+1-j} = \frac{1}{2}S$ for all $(i, j) \in V(\mathcal{K}_{m,n})$. We show that when m is odd, a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$ must be equatorially balanced. Also when m is odd, we show that (up to symmetries on the Klein bottle) the number of C_4 -face-magic Klein bottle labelings on the $m \times 4$ Klein bottle grid graph is $2^m (m-1)! \tau(m)$, where $\tau(m)$ is the number of positive divisors of m .

Furthermore, let $m \geq 3$ be an odd integer and $n \geq 6$ be an even integer. Then, the minimum number of distinct C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,n}$ (up to symmetries on a Klein bottle) is either $(5 \cdot 2^m)(m-1)!$ if $n \equiv 0 \pmod{4}$, or $(6 \cdot 2^m)(m-1)!$ if $n \equiv 2 \pmod{4}$.

1. INTRODUCTION

Graph labelings were formally introduced in the 1970s by Kotzig and Rosa [12]. Graph labelings have been applied to graph decomposition problems, radar pulse code designs, X-ray crystallography and communication network models. The interested reader should consult J.A. Gallian's comprehensive dynamic survey on graph labelings [8] for further investigation.

We refer the reader to Bondy and Murty [4] for concepts and notation not explicitly defined in this paper. All graphs in this paper are simple and connected. The concept of a C_k -face-magic labeling was first applied to planar graphs. For a planar (toroidal, cylindrical, or Klein bottle) graph $G = (V, E)$ embedded in the plane (torus, cylinder, or Klein bottle), let $\mathcal{F}(G)$ denote the set of faces of G . Then, G is called a C_k -face-magic planar (toroidal, cylindrical, or Klein bottle) graph if there exists a bijection $f : V(G) \rightarrow \{1, 2, \dots, |V(G)|\}$ such that for any $F \in \mathcal{F}(G)$ with $F \cong C_k$, the sum of all the vertex labelings along C_k is a constant S . Here, the constant S is called a C_k -face-magic value of G . More generally, C_k -face-magic planar graph labelings are a special case of (a, b, c) -magic labelings introduced by Lih [13]. For assorted values of a, b and c , Baca and others [1, 2, 3, 9, 10, 11, 13] have analyzed the problem for various classes of graphs. Wang [14] showed that the toroidal grid graphs $C_m \times C_n$ are antimagic for all integers $m, n \geq 3$. Butt et al. [5] investigated face antimagic labelings on toroidal and Klein bottle grid graphs. Curran, Low and Locke [6, 7] investigated C_4 -face-magic toroidal labelings on $C_m \times C_n$ and C_4 -face-magic cylindrical labelings on $P_m \times C_n$. Curran and Low [6] showed that (up to symmetries on a torus) there are three C_4 -face-magic toroidal labelings on $C_4 \times C_4$ that are row-sum balanced, column-sum balanced, and diagonal-sum balanced. Curran, Low and Locke [7] showed that for all positive integers n , $C_{4n} \times C_{4n}$ has a C_4 -face-magic toroidal labeling that is row-sum balanced, column-sum balanced, and diagonal-sum balanced. In this paper, we investigate C_4 -face-magic Klein bottle labelings on an $m \times n$ Klein bottle grid graph. We show that for $m, n \geq 2$, $\mathcal{K}_{m,n}$ has a C_4 -face-magic Klein bottle labeling if and only if n is even. When m is odd, we show that (up to symmetries on the Klein bottle) the number of C_4 -face-magic Klein bottle labelings on the $m \times 4$ Klein bottle grid graph is $2^m (m-1)! \tau(m)$, where $\tau(m)$ is the number of positive divisors of m . Furthermore, let $m \geq 3$ be an odd integer and $n \geq 6$ be an even integer. Then, the minimum number of distinct C_4 -face-magic Klein bottle

Date: June 3, 2022.

2010 Mathematics Subject Classification. 05C78.

Key words and phrases. C_4 -face-magic graphs, polyomino, Cartesian products of paths, Klein bottle grid graphs.

labelings X on $\mathcal{K}_{m,n}$ (up to symmetries on a Klein bottle) is either $(5 \cdot 2^m)(m-1)!$ if $n \equiv 0 \pmod{4}$, or $(6 \cdot 2^m)(m-1)!$ if $n \equiv 2 \pmod{4}$.

2. PRELIMINARIES

Notation 1. Let m be a positive integer. For convenience, we let $\lceil m/2 \rceil = \lfloor (m+1)/2 \rfloor$, and $\lceil m \rceil = \lfloor (m-1)/2 \rfloor$.

Definition 2. For a graph $G = (V, E)$ embedded on the Klein bottle (plane or torus or cylinder), let $\mathcal{F}(G)$ denote the set of faces of G . Then, G is called a C_n -face-magic Klein bottle (planar or toroidal or cylindrical) graph if there exists a bijection $f : V(G) \rightarrow \{1, 2, \dots, |V(G)|\}$ such that for any $F \in \mathcal{F}(G)$ with $F \cong C_n$, the sum of all the vertex labelings along C_n is a constant S . We call S the C_4 -face-magic value.

Theorem 3 ([7], Theorem 1). *Let $m, n \geq 2$. Then, $P_m \times P_n$ is C_4 -face-magic planar.*

Lemma 4 ([7], Lemma 2). *Let m and n be integers such that $m \geq 3$ and $n \geq 2$. Suppose $P_m \times C_n$ is a C_4 -face-magic cylindrical graph with the natural embedding of $P_m \times C_n$ on the cylinder. Then, n is even.*

Definition 5. Let m and n be integers such that $m, n \geq 2$. The $m \times n$ Klein bottle grid graph, denoted by $\mathcal{K}_{m,n}$, is the graph whose vertex set is

$$V(\mathcal{K}_{m,n}) = \{(i, j) : 1 \leq i \leq m, 1 \leq j \leq n\},$$

and whose edge set consists of the following edges:

- There is an edge from (i, j) to $(i, j+1)$, for $1 \leq i \leq m$ and $1 \leq j \leq n-1$.
- There is an edge from (i, n) to $(i, 1)$, for $1 \leq i \leq m$.
- There is an edge from (i, j) to $(i+1, j)$, for $1 \leq i \leq m-1$ and $1 \leq j \leq n$.
- There is an edge from (m, j) to $(1, n+1-j)$, for $1 \leq j \leq n$.

The graph $\mathcal{K}_{m,n}$ has a natural embedding on the Klein bottle.

Proposition 6. *Let m and n be integers such that $m, n \geq 2$. Suppose that $\mathcal{K}_{m,n}$ is a C_4 -face-magic Klein bottle graph. Then, n is even.*

Proof. Let $\{x_{i,j} : (i, j) \in V(\mathcal{K}_{m,n})\}$ be a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$. Suppose $m \geq 3$. This labeling yields a C_4 -face-magic cylindrical labeling on $P_m \times C_n$. By Lemma 4, n is even.

For the purposes of contradiction, assume that $m = 2$ and n is odd. Let $n = 2n_1 + 1$ for some positive integer n_1 . When we set the two C_4 -face sums given below equal to each other

$$x_{1,n_1} + x_{1,n_1+1} + x_{2,n_1} + x_{2,n_1+1} = S = x_{2,n_1} + x_{2,n_1+1} + x_{1,n_1+1} + x_{1,n_1+2},$$

we obtain $x_{1,n_1} = x_{1,n_1+2}$. This is a contradiction. Hence, n is even. This completes the proof. \square

Lemma 7. *Let m be an integer such that $m \geq 2$ and let n be an even positive integer. Let $\{x_{i,j} : (i, j) \in V(\mathcal{K}_{m,n})\}$ be a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$. Then, the C_4 -face-magic value of this labeling is $S = 2(mn + 1)$.*

Proof. Consider the sum

$$\begin{aligned} mnS &= \sum_{i=1}^{m-1} \sum_{j=1}^n (x_{i,j} + x_{i+1,j} + x_{i,j+1} + x_{i+1,j+1}) \\ &\quad + \sum_{i=1}^m (x_{m,j} + x_{m,j+1} + x_{1,n-j} + x_{1,n-j+1}) \\ &= 4 \left(\sum_{k=1}^{mn} k \right) = (2mn)(mn + 1), \end{aligned}$$

where the indices i and j are taken modulo m and n , respectively. Thus, $S = 2(mn + 1)$. \square

3. C_4 -FACE-MAGIC LABELINGS ON AN $m \times n$ KLEIN BOTTLE GRID GRAPH

Definition 8. Let $m, n \geq 2$ be integers such that n is even. Let $X = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,n})\}$ be a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$. Let $S = 2(mn + 1)$ be the C_4 -face-magic value of this labeling. We say that X is *equatorially balanced* if, for all $1 \leq i \leq m$ and $1 \leq j \leq \frac{n}{2}$, we have

$$x_{i,j} + x_{i,n+1-j} = \frac{1}{2}S = mn + 1.$$

Lemma 9. Let m be an odd integer such that $m \geq 3$ and let n be an even positive integer. Let $X = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,n})\}$ be a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$. Then, X is equatorially balanced.

Proof. Consider the C_4 -face sums

$$\begin{aligned} S &= x_{i,1} + x_{i+1,1} + x_{i,n} + x_{i+1,n} \text{ and} \\ S &= x_{i+1,1} + x_{i+2,1} + x_{i+1,n} + x_{i+2,n}, \text{ where } i = 1, 2, \dots, m. \end{aligned}$$

When we equate these sums and simplify, we obtain

$$x_{i,1} + x_{i,n} = x_{i+2,1} + x_{i+2,n}, \text{ where } i = 1, 2, \dots, m.$$

Since the indices i and $i + 2$ are taken modulo m , and m is odd, we have $x_{i_1,1} + x_{i_1,n} = x_{i_2,1} + x_{i_2,n}$, for all $1 \leq i_1, i_2 \leq m$. Thus, $S = x_{i,1} + x_{i,n} + x_{i+1,1} + x_{i+1,n} = 2(x_{i,1} + x_{i,n})$, which implies $x_{i,1} + x_{i,n} = \frac{1}{2}S$, for all $i = 1, 2, \dots, m$.

Assume that for some positive integer k such that $k < \frac{n}{2}$, we have for all $1 \leq i \leq m$ and $1 \leq j \leq k$ that $x_{i,j} + x_{i,n+1-j} = \frac{1}{2}S$. When we equate the C_4 -face sums given below together

$$x_{i,k} + x_{i,k+1} + x_{i+1,k} + x_{i+1,k+1} = S = x_{i+1,k} + x_{i+1,k+1} + x_{i+2,k} + x_{i+2,k+1},$$

we obtain

$$x_{i,k} + x_{i,k+1} = x_{i+2,k} + x_{i+2,k+1}, \text{ for all } 1 \leq i \leq m - 2.$$

A similar argument shows that

$$x_{i,n-k} + x_{i,n-k+1} = x_{i+2,n-k} + x_{i+2,n-k+1}, \text{ for all } 1 \leq i \leq m - 2.$$

When we equate the C_4 -face sums given below together

$$x_{m-1,k} + x_{m-1,k+1} + x_{m,k} + x_{m,k+1} = S = x_{m,k} + x_{m,k+1} + x_{1,n-k} + x_{1,n-k+1},$$

we obtain

$$x_{m-1,k} + x_{m-1,k+1} = x_{1,n-k} + x_{1,n-k+1}.$$

Similarly, when we equate the C_4 -face sums given below together

$$x_{m,k} + x_{m,k+1} + x_{1,n-k} + x_{1,n-k+1} = S = x_{1,n-k} + x_{1,n-k+1} + x_{2,n-k} + x_{2,n-k+1},$$

we obtain

$$x_{m,k} + x_{m,k+1} = x_{2,n-k} + x_{2,n-k+1}.$$

Let $S_i = x_{i,k} + x_{i,k+1}$, for $i = 1, 2$. Then,

$$\begin{aligned} S_1 &= x_{2i+1,k} + x_{2i+1,k+1}, \text{ for } i = 0, 1, 2, \dots, \frac{m}{2}, \text{ and} \\ S_1 &= x_{2i,n-k} + x_{2i,n-k+1}, \text{ for } i = 1, 2, \dots, \frac{m}{2}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} S_2 &= x_{2i,k} + x_{2i,k+1}, \text{ for } i = 1, 2, \dots, \frac{m}{2}, \text{ and} \\ S_2 &= x_{2i+1,n-k} + x_{2i+1,n-k+1}, \text{ for } i = 0, 1, 2, \dots, \frac{m}{2}. \end{aligned}$$

Since i and $i + 1$ have opposite parity, we have $x_{i+1,k} + x_{i+1,k+1} = x_{i,n-k} + x_{i,n+1-k}$. From the C_4 -face sum $x_{i,k} + x_{i,k+1} + x_{i+1,k} + x_{i+1,k+1} = S$, we have $x_{i,k} + x_{i,k+1} + x_{i,n-k} + x_{i,n+1-k} = S$. From the inductive hypothesis, we have $x_{i,k} + x_{i,n+1-k} = \frac{1}{2}S$. Thus, it follows that

$$x_{i,k+1} + x_{i,n-k} = S - (x_{i,k} + x_{i,n+1-k}) = \frac{1}{2}S.$$

This completes the proof. \square

Proposition 10. Let m be an odd integer such that $m \geq 3$ and let n be an even positive integer such that $n = 2n_1$, for some integer n_1 . Define a labeling $X = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,n})\}$ on $\mathcal{K}_{m,n}$ as follows:

- $x_{2i-1,2j-1} = (j-1)m + i$, for $i = 1, 2, \dots, \frac{m}{2}^+$ and $j = 1, 2, \dots, \frac{n_1}{2}^+$.

- $x_{2i,2j-1} = (n+1-j)m+1 - \mathfrak{m}^+ - i$, for $i = 1, 2, \dots, \mathfrak{m}$ and $j = 1, 2, \dots, \mathfrak{n}_1^+$.
- $x_{2i-1,2j} = (n_1-j+1)m+1 - i$, for $i = 1, 2, \dots, \mathfrak{m}^+$ and $j = 1, 2, \dots, \mathfrak{n}_1$.
- $x_{2i,2j} = (n_1+j-1)m + \mathfrak{m}^+ + i$, for $i = 1, 2, \dots, \mathfrak{m}$ and $j = 1, 2, \dots, \mathfrak{n}_1$.

Furthermore, for $1 \leq i \leq m$ and $\mathfrak{n} + 1 \leq j \leq n$, let $x_{i,j} = mn + 1 - x_{i,n+1-j}$. Then, X is a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$.

Proof. Let $m = 2m_1 + 1$ for some positive integer m_1 . Then, $\mathfrak{m}^+ = m_1 + 1$ and $\mathfrak{m} = m_1$. We observe that $\mathfrak{n}_1^+ + \mathfrak{n}_1 = n_1$. Let S be the C_4 -face-magic value of X . By Lemma 7, $S = 2(mn + 1)$. We verify the C_4 -face sum

$$(1) \quad x_{i,j} + x_{i+1,j} + x_{i+1,j+1} + x_{i,j+1} = S, \quad \text{for all } 1 \leq i \leq m \text{ and } 1 \leq j \leq n_1.$$

We first verify equation (1) for $j = n_1$. By Lemma 9, we have

$$x_{i,n_1} + x_{i,n_1+1} = \frac{1}{2}S, \quad \text{for all } 1 \leq i \leq m.$$

Thus, for all $1 \leq i \leq m$, we have

$$x_{i,n_1} + x_{i,n_1+1} + x_{i+1,n_1} + x_{i+1,n_1+1} = \frac{1}{2}S + \frac{1}{2}S = S.$$

Case 1. Assume i is odd and j is odd in equation (1). We verify the C_4 -face sum $x_{2i-1,2j-1} + x_{2i-1,2j} + x_{2i,2j-1} + x_{2i,2j} = S$, for $1 \leq i \leq m_1 + 1$ and $1 \leq j \leq \mathfrak{n}_1$. Assume that $1 \leq i \leq m_1$ and $1 \leq j \leq \mathfrak{n}_1$. Then, $x_{2i-1,2j-1} = (j-1)m + i$, $x_{2i,2j-1} = (n+1-j)m + 1 - \mathfrak{m}^+ - i$, $x_{2i-1,2j} = (n_1-j+1)m + 1 - i$ and $x_{2i,2j} = (n_1+j-1)m + \mathfrak{m}^+ + i$. Thus,

$$\begin{aligned} & x_{2i-1,2j-1} + x_{2i-1,2j} + x_{2i,2j-1} + x_{2i,2j} \\ &= ((j-1)m + i) + ((n_1-j+1)m + 1 - i) \\ &+ ((n+1-j)m + 1 - \mathfrak{m}^+ - i) + ((n_1+j-1)m + \mathfrak{m}^+ + i) \\ &= 2(mn + 1) = S. \end{aligned}$$

Assume $i = m_1 + 1$ and $1 \leq j \leq \mathfrak{n}_1$. We have $x_{m,2j-1} = (j-1)m + m_1 + 1$, $x_{m,2j} = (n_1-j+1)m - m_1$, $x_{1,2j-1} = (j-1)m + 1$, and $x_{1,2j} = (n_1-j+1)m$. Thus, $x_{m,2j-1} + x_{m,2j} = n_1m + 1 = x_{1,2j-1} + x_{1,2j}$. Since $x_{1,(n+1)-(2j-1)} = \frac{1}{2}S - x_{1,2j-1}$ and $x_{1,(n+1)-2j} = \frac{1}{2}S - x_{1,2j}$, we have

$$\begin{aligned} & x_{m,2j-1} + x_{m,2j} + x_{1,(n+1)-(2j-1)} + x_{1,(n+1)-2j} \\ &= x_{m,2j-1} + x_{m,2j} + \left(\frac{1}{2}S - x_{1,2j-1}\right) + \left(\frac{1}{2}S - x_{1,2j}\right) \\ &= (n_1m + 1) + S - (n_1m + 1) = S. \end{aligned}$$

Similarly, since $x_{m,(n+1)-(2j-1)} = \frac{1}{2}S - x_{m,2j-1}$ and $x_{m,(n+1)-2j} = \frac{1}{2}S - x_{m,2j}$, we have

$$\begin{aligned} & x_{1,2j-1} + x_{1,2j} + x_{m,(n+1)-(2j-1)} + x_{m,(n+1)-2j} \\ &= x_{1,2j-1} + x_{1,2j} + \left(\frac{1}{2}S - x_{m,2j-1}\right) + \left(\frac{1}{2}S - x_{m,2j}\right) \\ &= (n_1m + 1) + S - (n_1m + 1) = S. \end{aligned}$$

Case 2. Assume i is even and j is odd in equation (1). We verify the C_4 -face sum $x_{2i,2j-1} + x_{2i,2j} + x_{2i+1,2j-1} + x_{2i+1,2j} = S$, for $1 \leq i \leq m_1$ and $1 \leq j \leq \mathfrak{n}_1$. We have $x_{2i,2j-1} = (n+1-j)m + 1 - \mathfrak{m} - i$, $x_{2i,2j} = (n_1+j-1)m + \mathfrak{m} + i$, $x_{2(i+1)-1,2j-1} = (j-1)m + i + 1$ and $x_{2(i+1)-1,2j} = (n_1-j+1)m - i$. Thus,

$$\begin{aligned} & x_{2i,2j-1} + x_{2i,2j} + x_{2(i+1)-1,2j-1} + x_{2(i+1)-1,2j} \\ &= ((n+1-j)m + 1 - \mathfrak{m} - i) + ((n_1+j-1)m + \mathfrak{m} + i) \\ &+ ((j-1)m + i + 1) + ((n_1-j+1)m - i) \\ &= 2(mn + 1) = S. \end{aligned}$$

Case 3. Assume i is odd and j is even in equation (1). We verify the C_4 -face sum $x_{2i-1,2j} + x_{2i-1,2j+1} + x_{2i,2j} + x_{2i,2j+1} = S$, for $1 \leq i \leq m_1 + 1$ and $1 \leq j \leq \mathfrak{n}_1^-$. Assume $1 \leq i \leq m_1$ and $1 \leq j \leq \mathfrak{n}_1^-$. Then,

$x_{2i-1,2j} = (n_1 - j + 1)m + 1 - i$, $x_{2i-1,2j+1} = x_{2i-1,2(j+1)-1} = jm + i$, $x_{2i,2j} = (n_1 + j - 1)m + \mathfrak{m}^+ + i$ and $x_{2i,2j+1} = x_{2i,2(j+1)-1} = (n - j)m + 1 - \mathfrak{m}^+ - i$. Thus,

$$\begin{aligned} & x_{2i-1,2j} + x_{2i-1,2j+1} + x_{2i,2j} + x_{2i,2j+1} \\ &= ((n_1 - j + 1)m + 1 - i) + (jm + i) \\ &+ ((n_1 + j - 1)m + \mathfrak{m}^+ + i) + ((n - j)m + 1 - \mathfrak{m}^+ - i) \\ &= 2(mn + 1) = S. \end{aligned}$$

Next, assume $i = m_1 + 1$ and $1 \leq j \leq \mathfrak{n}_1^-$. We have $x_{m,2j} = x_{2(m_1+1)-1,2j} = (n_1 - j + 1)m - m_1$, $x_{m,2j+1} = x_{2(m_1+1)-1,2(j+1)-1} = jm + m_1 + 1$, $x_{1,2j} = (n_1 - j + 1)m$, and $x_{1,2j+1} = x_{1,2(j+1)-1} = jm + 1$. Then, $x_{m,2j} + x_{m,2j+1} = (n_1 + 1)m + 1 = x_{1,2j} + x_{1,2j+1}$. Since $x_{1,(n+1)-(2j)} = \frac{1}{2}S - x_{1,2j}$ and $x_{1,(n+1)-(2j+1)} = \frac{1}{2}S - x_{1,2j+1}$, we have

$$\begin{aligned} & x_{m,2j} + x_{m,2j+1} + x_{1,(n+1)-2j} + x_{1,(n+1)-(2j+1)} \\ &= x_{m,2j} + x_{m,2j+1} + \left(\frac{1}{2}S - x_{1,2j}\right) + \left(\frac{1}{2}S - x_{1,2j+1}\right) \\ &= ((n_1 + 1)m + 1) + S - ((n_1 + 1)m + 1) = S. \end{aligned}$$

Similarly, since $x_{m,(n+1)-2j} = \frac{1}{2}S - x_{m,2j}$ and $x_{m,(n+1)-(2j+1)} = \frac{1}{2}S - x_{m,2j+1}$, we have

$$\begin{aligned} & x_{1,2j} + x_{1,2j+1} + x_{m,(n+1)-2j} + x_{m,(n+1)-(2j+1)} \\ &= x_{1,2j-1} + x_{1,2j} + \left(\frac{1}{2}S - x_{m,2j}\right) + \left(\frac{1}{2}S - x_{m,2j+1}\right) \\ &= ((n_1 + 1)m + 1) + S - ((n_1 + 1)m + 1) = S. \end{aligned}$$

Case 4. Assume i is even and j is even in equation (1). We verify the C_4 -face sum $x_{2i,2j} + x_{2i,2j+1} + x_{2i+1,2j} + x_{2i+1,2j+1} = S$, for $1 \leq i \leq m_1$ and $1 \leq j \leq \mathfrak{n}_1^-$. We have $x_{2i,2j} = (n_1 + j - 1)m + \mathfrak{m}^+ + i$, $x_{2i,2j+1} = x_{2i,2(j+1)-1} = (n - j)m + 1 - \mathfrak{m}^+ - i$, $x_{2i+1,2j} = x_{2(i+1)-1,2j} = (n_1 - j + 1)m - i$ and $x_{2i+1,2j+1} = x_{2(i+1)-1,2(j+1)-1} = jm + i + 1$. Thus,

$$\begin{aligned} & x_{2i,2j} + x_{2i,2j+1} + x_{2i+1,2j} + x_{2i+1,2j+1} \\ &= ((n_1 + j - 1)m + \mathfrak{m}^+ + i) + ((n - j)m + 1 - \mathfrak{m}^+ - i) \\ &+ ((n_1 - j + 1)m - i) + (jm + i + 1) \\ &= 2(mn + 1) = S. \end{aligned}$$

Next, for $1 \leq i < m$ and $n_1 + 1 \leq j < n$, we have

$$\begin{aligned} & x_{i,j} + x_{i+1,j} + x_{i,j+1} + x_{i+1,j+1} \\ &= \left(\frac{1}{2}S - x_{i,n+1-j}\right) + \left(\frac{1}{2}S - x_{i+1,n+1-j}\right) + \left(\frac{1}{2}S - x_{i,n-j}\right) + \left(\frac{1}{2}S - x_{i+1,n-j}\right) \\ &= 2S - (x_{i,n+1-j} + x_{i+1,n+1-j} + x_{i,n-j} + x_{i+1,n-j}) = S. \end{aligned}$$

Finally, we need to show that every integer k , for $1 \leq k \leq mn$, that k is used exactly once in the labeling X . From conditions (1) and (2) of Proposition 10 and the identity $x_{i,n+1-j} = \frac{1}{2}S - x_{i,j}$ of Lemma 9, we have

$$\begin{aligned} & x_{2i-1,2j-1} = (j - 1)m + i, \quad \text{for } 1 \leq i \leq m_1 + 1 \text{ and } 1 \leq j \leq \mathfrak{n}_1^+, \text{ and} \\ & x_{2i,n+2-2j} = (j - 1)m + m_1 + 1 + i, \quad \text{for } 1 \leq i \leq m_1 \text{ and } 1 \leq j \leq \mathfrak{n}_1^+. \end{aligned}$$

This produces the list of integers k , for $1 \leq k \leq m\mathfrak{n}_1^+$, where each integer k is used exactly once in the labeling X .

From conditions (3) and (4) of Proposition 10 and the identity $x_{i,n+1-j} = \frac{1}{2}S - x_{i,j}$ of Lemma 9, we have

$$\begin{aligned} & x_{2i-1,2j} = (n_1 - j + 1)m + 1 - i, \quad \text{for } 1 \leq i \leq m_1 + 1 \text{ and } 1 \leq j \leq \mathfrak{n}_1, \text{ and} \\ & x_{2i,n+1-2j} = (n_1 - j + 1)m - m_1 - i, \quad \text{for } 1 \leq i \leq m_1 \text{ and } 1 \leq j \leq \mathfrak{n}_1. \end{aligned}$$

This produces the list of integers k , for $m\mathfrak{n}_1^+ + 1 \leq k \leq mn_1$, where each integer k is used exactly once in the labeling X .

When we include the other values of the labeling X that result from using the identity $x_{i,n+1-j} = \frac{1}{2}S - x_{i,j}$ of Lemma 9, we produce the list of integers k , for $mn_1 + 1 \leq k \leq mn$, where each integer k is used exactly once in the labeling X . This completes the proof. \square

Example 11. Table 1 illustrates Proposition 10, when $m = 7$ and $n = 10$.

1	35	8	28	15	56	43	63	36	70
66	40	59	47	52	19	24	12	31	5
2	34	9	27	16	55	44	62	37	69
65	41	58	48	51	20	23	13	30	6
3	33	10	26	17	54	45	61	38	68
64	42	57	49	50	21	22	14	29	7
4	32	11	25	18	53	46	60	39	67

TABLE 1. A C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{7,10}$.

Proposition 12. Let m and n be even positive integers. Let $n = 2n_1$ for some positive integer n_1 . Define a labeling $X = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,n})\}$ on $\mathcal{K}_{m,n}$ as follows:

- $x_{2i-1,2j-1} = 2m(j-1) + i$, for $i = 1, 2, \dots, \frac{n}{2}$ and $j = 1, 2, \dots, n_1^+$.
- $x_{2i-1,2j} = 2mj + 1 - i$, for $i = 1, 2, \dots, \frac{n}{2}$ and $j = 1, 2, \dots, n_1$.
- $x_{2i-1,2j-1} = m(n+1-2j) + i$, for $i = 1, 2, \dots, \frac{n}{2}$ and $j = n_1^+ + 1, n_1^+ + 2, \dots, n_1$.
- $x_{2i-1,2j} = m(n+1-2j) + 1 - i$, for $i = 1, 2, \dots, \frac{n}{2}$ and $j = n_1 + 1, n_1 + 2, \dots, n_1$.

Furthermore for $1 \leq i \leq \frac{n}{2}$ and $1 \leq j \leq n$, let $x_{2i,j} = mn + 1 - x_{2i-1,j}$. Then, X is a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$.

Proof. Since $x_{2i,j} = mn + 1 - x_{2i-1,j}$, for $1 \leq i \leq \frac{n}{2}$ and $1 \leq j \leq n$, we know that $x_{2i,j} + x_{2i-1,j} + x_{2i,j+1} + x_{2i-1,j+1} = 2mn + 2$, for $1 \leq i \leq \frac{n}{2}$ and $1 \leq j \leq n-1$. Also, $x_{2i-1,n} + x_{2i,n} + x_{2i-1,1} + x_{2i,1} = 2mn + 2$.

Let $1 \leq i \leq \frac{n}{2} - 1$ and $1 \leq j \leq n$. Note that for j odd, $x_{2i+1,j} = x_{2i-1,j} + 1$, and for j even, $x_{2i+1,j} = x_{2i-1,j} - 1$. Hence, for $j < n$, $x_{2i,j} + x_{2i+1,j} + x_{2i,j+1} + x_{2i+1,j+1} = x_{2i,j} + x_{2i-1,j} + x_{2i,j+1} + x_{2i-1,j+1} = 2mn + 2$. This also holds for $j = n$, with the understanding that $j + 1 = 1$ in this case.

It is now necessary to calculate the sums involving $x_{1,j}$ and $x_{m,j}$, for $1 \leq j < n$. If j is odd, $x_{1,j} - x_{1,n+1-j} = 1 - m$. If j is even, $x_{1,j} - x_{1,n+1-j} = m - 1$. Thus, for all integers $1 \leq j < n$, we have

$$(x_{1,j} - x_{1,n+1-j}) + (x_{1,j+1} - x_{1,n-j}) = 0.$$

If j is odd, $x_{m,j} = mn + 1 - x_{m-1,j} = mn + 1 - (x_{1,j} + \frac{n}{2} - 1) = mn - \frac{n}{2} - x_{1,j} + 2$. If j is even, $x_{m,j} = mn + 1 - x_{m-1,j} = mn + 1 - (x_{1,j} - \frac{n}{2} + 1) = mn + \frac{n}{2} - x_{1,j}$. Thus, for all integers $1 \leq j < n$, we have

$$(x_{m,n+1-j} + x_{m,n-j}) = 2mn + 2 - x_{1,n+1-j} - x_{1,n-j}.$$

Hence,

$$\begin{aligned} & x_{1,j} + x_{1,j+1} + x_{m,n+1-j} + x_{m,n-j} \\ &= x_{1,j} + x_{1,j+1} + (2mn + 2 - x_{1,n+1-j} - x_{1,n-j}) \\ &= 2mn + 2 + (x_{1,j} - x_{1,n+1-j}) + (x_{1,j+1} - x_{1,n-j}) = 2mn + 2. \end{aligned}$$

For the case when $j = n$, we have

$$x_{1,1} + x_{1,n} + x_{m,1} + x_{m,n} = 1 + m + (\frac{1}{2}S - \frac{n}{2}) + (\frac{1}{2}S - \frac{n}{2} - 1) = S = 2mn + 2.$$

Finally, we need to show that every integer k , for $1 \leq k \leq mn$, that k is used exactly once in the labeling X . A relabeling of the indices i and j in the expressions for $x_{2i-1,j}$ yields the following formulas:

$$(2) \quad x_{2i-1,2j-1} = (2j-2)m + i, \quad \text{for } 1 \leq i \leq \lceil \frac{n}{2} \rceil \text{ and } 1 \leq j \leq \lceil \frac{n}{2} \rceil^+,$$

$$(3) \quad x_{2(\lceil \frac{n}{2} \rceil + 1 - i) - 1, 2j} = (2j)m - \lceil \frac{n}{2} \rceil + i, \quad \text{for } 1 \leq i \leq \lceil \frac{n}{2} \rceil \text{ and } 1 \leq j \leq \lceil \frac{n}{2} \rceil,$$

$$(4) \quad x_{2i-1, 2(n_1+1-j)-1} = (2j-1)m + i, \quad \text{for } 1 \leq i \leq \lceil \frac{n}{2} \rceil \text{ and } 1 \leq j \leq \lceil \frac{n}{2} \rceil, \text{ and}$$

$$(5) \quad x_{2(\lceil \frac{n}{2} \rceil + 1 - i) - 1, 2(n_1+1-j)} = (2j-1)m - \lceil \frac{n}{2} \rceil + i, \quad \text{for } 1 \leq i \leq \lceil \frac{n}{2} \rceil \text{ and } 1 \leq j \leq \lceil \frac{n}{2} \rceil^+.$$

Let $1 \leq j \leq \lceil \frac{n}{2} \rceil$. The sequence of labels

$$\begin{aligned} (x_{2i-1, 2j-1} : 1 \leq i \leq \lceil \frac{n}{2} \rceil) &= ((2j-2)m + i : 1 \leq i \leq \lceil \frac{n}{2} \rceil) \\ &= ((2j-2)m + 1, (2j-2)m + 2, \dots, (2j-2)m + \lceil \frac{n}{2} \rceil), \end{aligned}$$

followed by the sequence of labels

$$\begin{aligned} (x_{2(\lceil \frac{n}{2} \rceil + 1 - i) - 1, 2(n_1+1-j)} : 1 \leq i \leq \lceil \frac{n}{2} \rceil) &= ((2j-1)m - \lceil \frac{n}{2} \rceil + i : 1 \leq i \leq \lceil \frac{n}{2} \rceil) \\ &= ((2j-2)m + \lceil \frac{n}{2} \rceil + 1, (2j-2)m + \lceil \frac{n}{2} \rceil + 2, \dots, (2j-1)m), \end{aligned}$$

followed by the sequence of labels

$$\begin{aligned} (x_{2i-1, 2(n_1+1-j)-1} : 1 \leq i \leq \lceil \frac{n}{2} \rceil) &= ((2j-1)m + i : 1 \leq i \leq \lceil \frac{n}{2} \rceil) \\ &= ((2j-1)m + 1, (2j-1)m + 2, \dots, (2j-1)m + \lceil \frac{n}{2} \rceil), \end{aligned}$$

and then followed by the sequence of labels

$$\begin{aligned} (x_{2(\lceil \frac{n}{2} \rceil + 1 - i) - 1, 2j} : 1 \leq i \leq \lceil \frac{n}{2} \rceil) &= ((2j)m - \lceil \frac{n}{2} \rceil + i : 1 \leq i \leq \lceil \frac{n}{2} \rceil) \\ &= ((2j-1)m + \lceil \frac{n}{2} \rceil + 1, (2j-1)m + \lceil \frac{n}{2} \rceil + 2, \dots, (2j)m), \end{aligned}$$

yields the sequence of integers $(k : (2j-1)m + 1 \leq k \leq (2j)m)$. When we concatenate these sequences of integers together for $1 \leq j \leq \lceil \frac{n}{2} \rceil$, we obtain the sequence $(k : 1 \leq k \leq 2m\lceil \frac{n}{2} \rceil)$. When n_1 is even, this is the sequence $(k : 1 \leq k \leq mn_1)$. However, when n_1 is odd, this is the sequence $(k : 1 \leq k \leq m(n_1 - 1))$. When we let $j = \lceil \frac{n}{2} \rceil^+ = \frac{1}{2}(n_1 + 1)$ in equations (2) and (5), the sequence of labels

$$\begin{aligned} (x_{2i-1, n_1} : 1 \leq i \leq \lceil \frac{n}{2} \rceil) &= ((n_1 - 1)m + i : 1 \leq i \leq \lceil \frac{n}{2} \rceil) \\ &= ((n_1 - 1)m + 1, (n_1 - 1)m + 2, \dots, (n_1 - 1)m + \lceil \frac{n}{2} \rceil), \end{aligned}$$

followed by the sequence of labels

$$\begin{aligned} (x_{2(\lceil \frac{n}{2} \rceil + 1 - i) - 1, n_1+1} : 1 \leq i \leq \lceil \frac{n}{2} \rceil) &= ((n_1 - 1)m - \lceil \frac{n}{2} \rceil + i : 1 \leq i \leq \lceil \frac{n}{2} \rceil) \\ &= ((n_1 - 1)m + \lceil \frac{n}{2} \rceil + 1, (n_1 - 1)m + \lceil \frac{n}{2} \rceil + 2, \dots, (n_1)m), \end{aligned}$$

produces the sequence of integers $(k : m(n_1 - 1) + 1 \leq k \leq mn_1)$. Therefore, the set of labels $\{x_{2i-1,j} : 1 \leq i \leq \lceil \frac{n}{2} \rceil \text{ and } 1 \leq j \leq n\}$ is the set of integers $\{k : 1 \leq k \leq mn_1\}$. From the identity $x_{2i,j} = mn + 1 - x_{2i-1,j}$, the set of labels $\{x_{2i,j} : 1 \leq i \leq \lceil \frac{n}{2} \rceil \text{ and } 1 \leq j \leq n\}$ is the set of integers $\{k : mn_1 + 1 \leq k \leq mn\}$. This completes the proof. \square

Example 13. Table 2 illustrates Proposition 12, when $m = 6$ and $n = 10$.

1	12	13	24	25	30	19	18	7	6
60	49	48	37	36	31	42	43	54	55
2	11	14	23	26	29	20	17	8	5
59	50	47	38	35	32	41	44	53	56
3	10	15	22	27	28	21	16	9	4
58	51	46	39	34	33	40	45	52	57

TABLE 2. A C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{6,10}$.

Theorem 14. *Let m and n be positive integers such that $m, n \geq 2$. Then, $\mathcal{K}_{m,n}$ has a C_4 -face-magic Klein bottle labeling if and only if n is even.*

Proof. (\Rightarrow) Suppose that $\mathcal{K}_{m,n}$ is a C_4 -face-magic Klein bottle graph. Then by Proposition 6, n is even.

(\Leftarrow) Assume n is even. On the one hand, if m is odd, by Proposition 10, $\mathcal{K}_{m,n}$ has a C_4 -face-magic Klein bottle labeling. On the other hand, if m is even, by Proposition 12, $\mathcal{K}_{m,n}$ has a C_4 -face-magic Klein bottle labeling. \square

4. EQUIVALENT C_4 -FACE-MAGIC KLEIN BOTTLE LABELINGS

Definition 15. Consider the natural embedding of $\mathcal{K}_{m,n}$ on the Klein bottle. We say that two C_4 -face-magic Klein bottle labelings on $\mathcal{K}_{m,n}$ are *Klein bottle equivalent* if there is a homeomorphism of the Klein bottle that maps $\mathcal{K}_{m,n}$ onto itself such that the first C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$ is mapped to the second C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$.

Definition 16. We consider the following symmetries of the Klein bottle that result in a graph automorphism on $\mathcal{K}_{m,n}$. Let U be the translation on $\mathcal{K}_{m,n}$ given by

$$U(i, j) = \begin{cases} (i + 1, j), & \text{if } 1 \leq i \leq m - 1 \text{ and } 1 \leq j \leq n, \\ (1, n + 1 - j), & \text{if } i = m \text{ and } 1 \leq j \leq n. \end{cases}$$

Let H be the translation on $\mathcal{K}_{m,n}$ given by $H(i, j) = (i, j + \#)$. Let F be the reflection on $\mathcal{K}_{m,n}$ given by $F(i, j) = (i, n + 1 - j)$. We observe that $U^m = F$. Also, the symmetries of the Klein bottle that result in a graph automorphism of $\mathcal{K}_{m,n}$ is the group of graph automorphisms $KBLS(m, n) = \langle U, H, F \rangle = \{U^i, FU^i, HU^i, FHU^i : 0 \leq i \leq m - 1\} = \{U^i, HU^i : 0 \leq i \leq 2m - 1\}$. Given a C_4 -face-magic Klein bottle labeling $X = \{x_{i,j} : (i, j) \in V(\mathcal{K}_{m,n})\}$ and $A \in KBLS(m, n)$, we let $A(X) = \{x_{A(i,j)} : (i, j) \in V(\mathcal{K}_{m,n})\}$. We call $KBLS(m, n)$ the *Klein bottle labeling symmetry group on $\mathcal{K}_{m,n}$* .

Definition 17. Let m be an odd positive integer such that $m \geq 3$, n be an even positive integer such that $n \geq 4$, $\# = n/2$, and $n_0 = \# - 1$. Let a_1, a_2, \dots, a_{n_0} be positive integers. We define the *labeling search graph corresponding to $(a_1, a_2, \dots, a_{n_0})$* , denoted by $G(a_1, a_2, \dots, a_{n_0})$, to be the graph whose vertex set is

$$V(G(a_1, a_2, \dots, a_{n_0})) = \{\{q, mn + 1 - q\} : 1 \leq q \leq m\#\}$$

and there is an edge from $\{x_1, x_2\}$ to $\{y_1, y_2\}$, where $\{x_1, x_2\} \neq \{y_1, y_2\}$, if and only if there exist $z_1 \in \{x_1, x_2\}$, $z_2 \in \{y_1, y_2\}$, and $1 \leq j \leq n_0$ such that $z_1 + z_2 = a_j$.

Let $P = (\{x_{j,1}, x_{j,2}\} : 1 \leq j \leq n_1)$ be a path on n_1 distinct vertices in $G(a_1, a_2, \dots, a_{n_0})$. We say that P is an $(a_1, a_2, \dots, a_{n_0})$ -*admissible path* provided that, for all $1 \leq j \leq n_1$, there exist $z_j \in \{x_{j,1}, x_{j,2}\}$, such that, for all $1 \leq j \leq n_0$, $z_j + z_{j+1} = a_j$. We call z_1 and z_{n_1} *end vertex labels* of P . For all $1 < j < n_1$, we call z_j an *interior vertex label* of P . We say that z_1 is an *end vertex label of P incident to the edge labeled a_1* . We say that z_{n_1} is an *end vertex label of P incident to the edge labeled a_{n_0}* . For all $1 < j < n_1$, we say that z_j is an *interior vertex label of P incident to the edges labeled a_{j-1} and a_j* . We call the sequence $(z_j : 1 \leq j \leq n_1)$ the *vertex label sequence of P* .

Let H be a spanning subgraph of $G(a_1, a_2, \dots, a_{n_0})$. We say that H is an $(a_1, a_2, \dots, a_{n_0})$ -*admissible path partition* of $G(a_1, a_2, \dots, a_{n_0})$ if H is the disjoint union of m distinct $(a_1, a_2, \dots, a_{n_0})$ -admissible paths in $G(a_1, a_2, \dots, a_{n_0})$.

Remark 18. The condition $\{x_1, x_2\} \neq \{y_1, y_2\}$ in Definition 17 on vertices $\{x_1, x_2\}$ and $\{y_1, y_2\}$ ensures that the graph $G(a_1, a_2, \dots, a_{n_0})$ does not have any loops. Suppose integer a_j is even, for some $1 \leq j \leq n_0$. Let $a_j = 2k$ for some positive integer k . Then, vertex $\{k, mn + 1 - k\}$ does not have a loop labeled a_j incident to it in $G(a_1, a_2, \dots, a_{n_0})$.

Lemma 19. *Let $m \geq 3$ be an odd integer and $n \geq 4$ be an even integer, $\# = n/2$, and $n_0 = \# - 1$. Let $S = 2(mn + 1)$. Let $X = \{x_{i,j} : (i, j) \in V(\mathcal{K}_{m,n})\}$ be a labeling on $\mathcal{K}_{m,n}$ such that each integer k with $1 \leq k \leq mn$ is used exactly once. For all $1 \leq j \leq n_0$, let $a_j = x_{1,j} + x_{1,j+1}$. Then, $X = \{x_{i,j} : (i, j) \in V(\mathcal{K}_{m,n})\}$ is a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$ if and only if*

- for all $1 \leq i \leq \# + 1$ and $1 \leq j \leq n_0$, we have $x_{2i-1,j} + x_{2i-1,j+1} = a_j$,
- for all $1 \leq i \leq \# + 1$ and $1 \leq j \leq \#$, we have $x_{2i-1,n+1-j} = \frac{1}{2}S - x_{2i-1,j}$,

- for all $1 \leq i \leq \overline{m}$ and $1 \leq j \leq n_0$, we have $x_{2i,n-j} + x_{2i,n-j+1} = a_j$, and
- for all $1 \leq i \leq \overline{m}$ and $1 \leq j \leq \overline{n}$, we have $x_{2i,j} = \frac{1}{2}S - x_{2i,n+1-j}$.

Proof. (\Rightarrow) Assume that $X = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,n})\}$ is a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$. When we equate the two C_4 -face sums given below equal to each other

$$\begin{aligned} x_{2i-1,j} + x_{2i-1,j+1} + x_{2i,j} + x_{2i,j+1} &= S \\ &= x_{2i,j} + x_{2i,j+1} + x_{2i+1,j} + x_{2i+1,j+1}, \end{aligned}$$

we obtain

$$x_{2i-1,j} + x_{2i-1,j+1} = x_{2i+1,j} + x_{2i+1,j+1}, \quad \text{for } 1 \leq i \leq \overline{m}.$$

Thus, $x_{2i-1,j} + x_{2i-1,j+1} = x_{1,j} + x_{1,j+1} = a_j$, for $1 \leq i \leq \overline{m}^+$. By Lemma 9, for all $1 \leq i \leq \overline{m}^+$ and $1 \leq j \leq \overline{n}$, we have $x_{2i-1,n+1-j} = \frac{1}{2}S - x_{2i-1,j}$.

When we equate the two C_4 -face sums given below equal to each other

$$x_{m,j} + x_{m,j+1} + x_{1,n-j} + x_{1,n-j+1} = S = x_{1,n-j} + x_{1,n-j+1} + x_{2,n-j} + x_{2,n-j+1},$$

we obtain

$$x_{2,n-j} + x_{2,n-j+1} = x_{m,j} + x_{m,j+1} = a_j.$$

When we equate the two C_4 -face sums given below equal to each other

$$\begin{aligned} x_{2i,n-j} + x_{2i,n-j+1} + x_{2i+1,n-j} + x_{2i+1,n-j+1} &= S \\ &= x_{2i+1,n-j} + x_{2i+1,n-j+1} + x_{2i+2,n-j} + x_{2i+2,n-j+1}, \end{aligned}$$

we obtain

$$x_{2i,n-j} + x_{2i,n-j+1} = x_{2i+2,n-j} + x_{2i+2,n-j+1}, \quad \text{for } 1 \leq i \leq \overline{m} - 1.$$

Thus, $x_{2i,n-j} + x_{2i,n-j+1} = x_{2,n-j} + x_{2,n-j+1} = a_j$, for $1 \leq i \leq \overline{m}$. By Lemma 9, for all $1 \leq i \leq \overline{m}$ and $1 \leq j \leq \overline{n}$, we have $x_{2i,j} = \frac{1}{2}S - x_{2i,n+1-j}$.

(\Leftarrow) Let $X = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,n})\}$ be a labeling on $\mathcal{K}_{m,n}$ such that each integer k with $1 \leq k \leq mn$ is used exactly once; for all $1 \leq i \leq \overline{m}^+$ and $1 \leq j \leq n_0$, we have

$$(6) \quad x_{2i-1,j} + x_{2i-1,j+1} = a_j;$$

for all $1 \leq i \leq \overline{m}^+$ and $1 \leq j \leq n_1$, we have

$$(7) \quad x_{2i-1,n+1-j} = \frac{1}{2}S - x_{2i-1,j};$$

for all $1 \leq i \leq \overline{m}$ and $1 \leq j \leq n_0$, we have

$$(8) \quad x_{2i,n-j} + x_{2i,n-j+1} = a_j, \text{ and}$$

for all $1 \leq i \leq \overline{m}$ and $1 \leq j \leq n_1$, we have

$$(9) \quad x_{2i,j} = \frac{1}{2}S - x_{2i,n+1-j}.$$

We show that $X = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,n})\}$ is a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$.

Case 1. Let $1 \leq i \leq \overline{m}$ and $1 \leq j \leq n_0$. We show that

$$x_{2i-1,j} + x_{2i-1,j+1} + x_{2i,j} + x_{2i,j+1} = S.$$

From equation (6), we have $x_{2i-1,j} + x_{2i-1,j+1} = a_j$. From equation (9), we have $x_{2i,j} = \frac{1}{2}S - x_{2i,n+1-j}$ and $x_{2i,j+1} = \frac{1}{2}S - x_{2i,n-j}$. Also, from equation (8), we have $x_{2i,n-j} + x_{2i,n-j+1} = a_j$. Thus,


$$\begin{aligned} x_{2i-1,j} + x_{2i-1,j+1} + x_{2i,j} + x_{2i,j+1} &= x_{2i-1,j} + x_{2i-1,j+1} \\ &\quad + \left(\frac{1}{2}S - x_{2i,n+1-j}\right) + \left(\frac{1}{2}S - x_{2i,n-j}\right) \\ &= a_j + S - a_j = S. \end{aligned}$$

A similar argument shows that

$$x_{2i,j} + x_{2i,j+1} + x_{2i+1,j} + x_{2i+1,j+1} = S.$$

Case 2. Let $1 \leq i \leq m$. We show that

$$\begin{aligned} x_{i,1} + x_{i+1,1} + x_{i,n} + x_{i+1,n} &= S \text{ and} \\ x_{i,\overline{m}} + x_{i+1,\overline{m}} + x_{i,\overline{m}+1} + x_{i+1,\overline{m}+1} &= S, \end{aligned}$$

FIGURE 1. $(22, 28, 34, 40)$ -admissible path partition K in $G(22, 28, 34, 40)$.

where we take the index i modulo m . From equations (7) and (9), we have $x_{i,1} = \frac{1}{2}S - x_{i,n}$ and $x_{i+1,1} = \frac{1}{2}S - x_{i+1,n}$. Thus,

$$x_{i,1} + x_{i+1,1} + x_{i,n} + x_{i+1,n} = (\frac{1}{2}S - x_{i,n}) + (\frac{1}{2}S - x_{i+1,n}) + x_{i,n} + x_{i+1,n} = S.$$

A similar argument shows that

$$x_{i,\#} + x_{i+1,\#} + x_{i,\#+1} + x_{i+1,\#+1} = S.$$

Case 3. Let $1 \leq j \leq n_0$. From equation (6), we have $x_{1,j} + x_{1,j+1} = a_j$ and $x_{m,j} + x_{m,j+1} = a_j$. From equation (7), we have $x_{1,n+1-j} = \frac{1}{2}S - x_{1,j}$ and $x_{1,n-j} = \frac{1}{2}S - x_{1,j+1}$. Thus,

$$\begin{aligned} x_{m,j} + x_{m,j+1} + x_{1,n+1-j} + x_{1,n-j} &= x_{m,j} + x_{m,j+1} \\ &\quad + (\frac{1}{2}S - x_{1,j}) + (\frac{1}{2}S - x_{1,j+1}) \\ &= a_j + S - a_j = S. \end{aligned}$$

A similar argument shows that, for $\# + 1 \leq j < n$, we have

$$x_{m,j} + x_{m,j+1} + x_{1,n+1-j} + x_{1,n-j} = S.$$

Since every integer k with $1 \leq k \leq mn$ is used in the labeling X , X is a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$. \square

Definition 20. Let $m \geq 3$ be an odd integer and $n \geq 4$ be an even integer, $\# = n/2$, and $n_0 = \# - 1$. Let $X = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,n})\}$ be a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$. We define the *labeling graph* of X , denoted by $\mathcal{L}(X)$, to be the graph with vertex set

$$V(\mathcal{L}(X)) = \{\{q, mn + 1 - q\} : 1 \leq q \leq m\#\}$$

and consists of the following edges:

- For all $1 \leq i \leq \# + 1$ and $1 \leq j \leq n_0$, there is an edge labeled $a_j = x_{2i-1,j} + x_{2i-1,j+1}$ from $\{x_{2i-1,j}, mn + 1 - x_{2i-1,j}\}$ to $\{x_{2i-1,j+1}, mn + 1 - x_{2i-1,j+1}\}$. The path $(\{x_{2i-1,j}, mn + 1 - x_{2i-1,j}\} : 1 \leq j \leq \#)$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in $G(a_1, a_2, \dots, a_{n_0})$ with vertex label sequence $(x_{2i-1,j} : 1 \leq j \leq \#)$.
- For all $1 \leq i \leq \#$ and $1 \leq j \leq n_0$, there is an edge labeled $a_j = x_{2i,n+1-j} + x_{2i,n-j}$ from $\{x_{2i,n+1-j}, mn + 1 - x_{2i,n+1-j}\}$ to $\{x_{2i,n-j}, mn + 1 - x_{2i,n-j}\}$. The path $(\{x_{2i,n+1-j}, mn + 1 - x_{2i,n+1-j}\} : 1 \leq j \leq \#)$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in $G(a_1, a_2, \dots, a_{n_0})$ with vertex label sequence $(x_{2i,n+1-j} : 1 \leq j \leq \#)$.

By Lemma 19, the graph $\mathcal{L}(X)$ is well-defined. Also, $\mathcal{L}(X)$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition of $G(a_1, a_2, \dots, a_{n_0})$.

Example 21. In Table 3 we have a C_4 -face-magic Klein bottle labeling X on the 3×10 Klein bottle grid graph $\mathcal{K}_{3,10}$ such that $\mathcal{L}(X) = K$ is the $(22, 28, 34, 40)$ -admissible path partition K in the labeling search graph $G(22, 28, 34, 40)$ given in Figure 1.

1	21	7	27	13	18	4	24	10	30
29	11	23	5	17	14	26	8	20	2
3	19	9	25	15	16	6	22	12	28

TABLE 3. C_4 -face-magic Klein bottle labeling X on $\mathcal{K}_{3,10}$ such that $\mathcal{L}(X) = K$ is the (22, 28, 34, 40)-admissible path partition K in $G(22, 28, 34, 40)$ given in Figure 1.

Lemma 22. *Let $m \geq 3$ be an odd integer, $n \geq 4$ be an even integer, $n_1 = \# = n/2$, and $n_0 = \# - 1$. Let a_1, a_2, \dots, a_{n_0} be positive integers. Suppose K is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition of $G(a_1, a_2, \dots, a_{n_0})$. Then, $\mathcal{K}_{m,n}$ has a C_4 -face-magic Klein bottle labeling $X = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,n})\}$ on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X) = K$. In addition, we have the following two possibilities:*

- Suppose $(a_1, a_2, \dots, a_{n_0})$ is not a palindrome. Then, there are $(m-1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings $X = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,n})\}$ on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X) = K$.
- Suppose $(a_1, a_2, \dots, a_{n_0})$ is a palindrome. Then, there are $2^{m-1}(m-1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings $X = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,n})\}$ on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X) = K$.

Proof. Suppose $(a_1, a_2, \dots, a_{n_0})$ is not a palindrome. Let T be the set of end vertex labels of paths in K incident to the edge labeled a_1 . In other words, we let

$$T = \left\{ k \in \mathbb{Z} : 1 \leq k \leq mn, \{k, mn+1-k\} \text{ is an end vertex of a path } P \text{ in } K, \right. \\ \left. \begin{aligned} &\{k, mn+1-k\} \text{ is adjacent to } \{k, mn+1-k\} \text{ in } P, k+\ell = a_1, \text{ and the} \\ &\text{sequence of labels on edges in the path } P \text{ from vertex } \{k, mn+1-k\} \text{ to} \\ &\text{the other end vertex of } P \text{ is } (a_1, a_2, \dots, a_{n_0}). \end{aligned} \right\}.$$

Since there are m paths in K and the sequence $(a_1, a_2, \dots, a_{n_0})$ is not a palindrome, there are exactly m positive integers in the set T . For convenience, we label the elements of T in ascending order. Let

$$(10) \quad T = \{k_1, k_2, \dots, k_m\} \quad \text{such that } k_i < k_{i+1}, \text{ for } 1 \leq i < m.$$

Let α be a permutation on the set $\{1, 2, 3, \dots, m\}$. We define a labeling $L(K, \alpha) = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,n})\}$ on $\mathcal{K}_{m,n}$ in the following manner:

- For $1 \leq i \leq \#^+$ and $1 \leq j \leq \#$, let

$$(11) \quad x_{2i-1,j} = (-1)^{j+1} k_{\alpha(i)} + \sum_{\ell=1}^{j-1} (-1)^{\ell+1} a_{j-\ell}.$$

- For $1 \leq i \leq \#^+$ and $1 \leq j \leq \#$, let

$$(12) \quad x_{2i-1, n+1-j} = \frac{1}{2}S - x_{2i-1,j}.$$

- For $\#^+ + 1 \leq i \leq m$ and $1 \leq j \leq \#$, let

$$(13) \quad x_{2(i-\#^+), n+1-j} = (-1)^{j+1} k_{\alpha(i)} + \sum_{\ell=1}^{j-1} (-1)^{\ell+1} a_{j-\ell}.$$

- For $\#^+ + 1 \leq i \leq m$ and $1 \leq j \leq \#$, let

$$(14) \quad x_{2(i-\#^+), j} = \frac{1}{2}S - x_{2(i-\#^+), n+1-j}.$$

We show that for $1 \leq i \leq \#^+$, the sequence $(x_{2i-1,j} : 1 \leq j \leq \#)$ is the vertex label sequence of some $(a_1, a_2, \dots, a_{n_0})$ -admissible path in K . Let $1 \leq i \leq \#^+$. Then, $x_{2i-1,1} = k_{\alpha(i)}$. For all $1 \leq j \leq n_0$, we have $x_{2i-1,j} + x_{2i-1,j+1} = a_j$. Hence for all $1 \leq i \leq \#^+$, the sequence of vertices $(\{x_{2i-1,j}, \frac{1}{2}S - x_{2i-1,j}\} : 1 \leq j \leq \#)$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in K with vertex label sequence $(x_{2i-1,j} : 1 \leq j \leq \#)$.

Similarly, we show that for $\#^+ + 1 \leq i \leq m$, the sequence $(x_{2(i-\#^+), n+1-j} : 1 \leq j \leq \#)$ is the vertex label sequence of some $(a_1, a_2, \dots, a_{n_0})$ -admissible path in K . Let $\#^+ + 1 \leq i \leq m$. Then, $x_{2(i-\#^+), n} = k_{\alpha(i)}$.

For all $1 \leq j \leq n_0$, we have $x_{2(i-\mathfrak{m}^+),n+1-j} + x_{2(i-\mathfrak{m}^+),n-j} = a_j$. Hence, for all $\mathfrak{m}^+ + 1 \leq i \leq m$, the sequence of vertices $(\{x_{2(i-\mathfrak{m}^+),n+1-j}, \frac{1}{2}S - x_{2(i-\mathfrak{m}^+),n+1-j}\} : 1 \leq j \leq \mathfrak{m})$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in K with vertex label sequence $(x_{2(i-\mathfrak{m}^+),n+1-j} : 1 \leq j \leq \mathfrak{m})$.

Next we show that the labeling $L(K, \alpha)$ uses each integer in the set $\{k \in \mathbb{Z} : 1 \leq k \leq mn\}$ exactly once. For $1 \leq i \leq \mathfrak{m}^+$, we have $x_{2i-1,1} = k_{\alpha(i)}$, and for $\mathfrak{m}^+ + 1 \leq i \leq m$, we have $x_{2(i-\mathfrak{m}^+),n} = k_{\alpha(i)}$. Thus, each of the values $x_{2i-1,1}$, $1 \leq i \leq \mathfrak{m}^+$, and $x_{2(i-\mathfrak{m}^+),n}$ for $\mathfrak{m}^+ + 1 \leq i \leq m$, are distinct end vertex labels incident to an edge labeled a_1 of an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in K . For $1 \leq i \leq \mathfrak{m}^+$, the sequence $(x_{2i-1,j} : 1 \leq j \leq n)$ is the sequence of labels $(x_{2i-1,j} : 1 \leq j \leq \mathfrak{m})$ followed by the sequence of labels $(\frac{1}{2}S - x_{2i-1,\mathfrak{m}+1-j} : 1 \leq j \leq \mathfrak{m})$. Furthermore, each label in the sequence $(x_{2i-1,j} : 1 \leq j \leq n)$ appears exactly once as a label for a vertex in the path $P_i = (\{x_{2i-1,j}, \frac{1}{2}S - x_{2i-1,j} : 1 \leq j \leq \mathfrak{m}\})$ in K . Similarly, for $1 \leq i \leq \mathfrak{m}$, the sequence $(x_{2i,n+1-j} : 1 \leq j \leq n)$ is the sequence of labels $(x_{2i,n+1-j} : 1 \leq j \leq \mathfrak{m})$ followed by the sequence of labels $(\frac{1}{2}S - x_{2i,\mathfrak{m}+j} : 1 \leq j \leq \mathfrak{m})$. Furthermore, each label in the sequence $(x_{2i,n+1-j} : 1 \leq j \leq n)$ appears exactly once as a label for a vertex in the path $P_{i+\mathfrak{m}^+} = (\{x_{2i,n+1-j}, \frac{1}{2}S - x_{2i,n+1-j}\} : 1 \leq j \leq \mathfrak{m})$ in K . Since the set $\{V(P_i) : 1 \leq i \leq m\}$ is a partition on the vertices of $G(a_1, a_2, \dots, a_{n_0})$, each integer k in the set $\{k \in \mathbb{Z} : 1 \leq k \leq mn\}$ is used exactly once. By Lemma 19, $L(K, \alpha)$ is a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$.

We consider the graph automorphisms of $\mathcal{K}_{m,n}$ that are induced by a symmetry on the Klein bottle. From Notation 16, these graph automorphisms of $\mathcal{K}_{m,n}$ are in the group $KBLS(m, n)$. We consider the result of each graph automorphism in $KBLS(m, n)$ on $L(K, \alpha)$. We let i be an integer and apply the graph automorphism U^{2i+1} from $KBLS(m, n)$ on $L(K, \alpha)$. Observe that $\mathcal{L}(U^{2i+1}(L(K, \alpha)))$ is an $(S - a_1, S - a_2, \dots, S - a_{n_0-1}, S - a_{n_0})$ -admissible path partition of $G(S - a_1, S - a_2, \dots, S - a_{n_0-1}, S - a_{n_0})$. Thus, $\mathcal{L}(U^{2i+1}(L(K, \alpha))) \neq K$. Next, we apply the graph automorphism HU^{2i} to $L(K, \alpha)$. Then, $\mathcal{L}(HU^{2i}(L(K, \alpha)))$ is an $(S - a_{n_0}, S - a_{n_0-1}, \dots, S - a_2, S - a_1)$ -admissible path partition of $G(S - a_{n_0}, S - a_{n_0-1}, \dots, S - a_2, S - a_1)$. Thus, $\mathcal{L}(HU^{2i}(L(K, \alpha))) \neq K$. Similarly, we apply the graph automorphism HU^{2i+1} to $L(K, \alpha)$. Then, $\mathcal{L}(HU^{2i+1}(L(K, \alpha)))$ is an $(a_{n_0}, a_{n_0-1}, \dots, a_2, a_1)$ -admissible path partition of $G(a_{n_0}, a_{n_0-1}, \dots, a_2, a_1)$. Thus, $\mathcal{L}(HU^{2i+1}(L(K, \alpha))) \neq K$.

On the other hand, when we apply the graph automorphism U^{2i} to $L(K, \gamma)$, where γ is a permutation on the set $\{1, 2, \dots, n\}$, we obtain $\mathcal{L}(U^{2i}(L(K, \gamma))) = K$. Let β be the cyclic permutation on the set $\{1, 2, \dots, m\}$ defined by $\beta(i) = i - 1 \pmod{m}$. Let $X = \{x_{i,j} : (i, j) \in V(\mathcal{K}_{m,n})\}$ be any C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X) = K$. Let γ be the permutation on the set $\{1, 2, \dots, m\}$ such that $x_{2i-1,1} = k_{\gamma(i)}$, for $1 \leq i \leq \mathfrak{m}^+$, and $x_{2(i-\mathfrak{m}^+),n} = k_{\gamma(i)}$, for $\mathfrak{m}^+ + 1 \leq i \leq m$. Then $L(K, \gamma) = X$. Let t be the unique integer such that $0 \leq t < m$ and $\gamma(t) = 1$. Let $\alpha = \beta^{t-1}\gamma$. Then, $\alpha(1) = 1$. Thus, there is a unique permutation α on the set $\{1, 2, \dots, m\}$ with $\alpha(1) = 1$ and $L(K, \alpha) = U^{-2(t-1)}(X)$. Hence, for every C_4 -face-magic labeling X such that $\mathcal{L}(X) = K$, there is a unique permutation α on the set $\{1, 2, \dots, m\}$ with $\alpha(1) = 1$ such that X is Klein bottle equivalent to $L(K, \alpha)$. Furthermore, each distinct permutation α on $\{1, 2, \dots, m\}$ with $\alpha(1) = 1$ results in a distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labeling $L(K, \alpha)$ on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(L(K, \alpha)) = K$. Since there are $(m-1)!$ permutation α on the set $\{1, 2, \dots, m\}$ with $\alpha(1) = 1$, there are $(m-1)!$ Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X) = K$.

Suppose the sequence $(a_1, a_2, \dots, a_{n_0})$ is a palindrome. We define the two-element sets of integers $\{k, \ell\}$ such that k and ℓ are end vertex labels of an $(a_1, a_2, \dots, a_{n_0})$ -admissible path P in K in the following way: Let

$$\begin{aligned} \tilde{T} = \left\{ \{k, \ell\} : k, \ell \in \mathbb{Z}, 1 \leq k, \ell \leq mn, \{k, mn+1-k\} \text{ and } \{\ell, mn+1-\ell\} \text{ are end} \right. \\ \left. \text{vertices of an } (a_1, a_2, \dots, a_{n_0})\text{-admissible path } P \text{ in } K, \text{ and there exist vertices} \right. \\ \left. \{p, mn+1-p\} \text{ and } \{q, mn+1-q\} \text{ on path } P \text{ such that } k+p = a_1 = \ell+q \right\}. \end{aligned}$$

Since there are m paths in K and the sequence $(a_1, a_2, \dots, a_{n_0})$ is a palindrome, there are exactly m two-element sets in \tilde{T} . For convenience, we label the elements of \tilde{T} in the following way: Let

$$(15) \quad \begin{aligned} \tilde{T} = \{ \{k_{i,1}, k_{i,2}\} : 1 \leq i \leq m \} \text{ such that } k_{i,1} < k_{i,2}, \text{ for all } 1 \leq i \leq m, \\ \text{and } k_{i,1} < k_{i+1,1}, \text{ for all } 1 \leq i < m. \end{aligned}$$

Let α be a permutation on the set $\{1, 2, 3, \dots, m\}$. Let ρ be a function from $\{1, 2, \dots, m\}$ to $\{1, 2\}$. We define a labeling $L(K, \alpha, \rho) = \{x_{i,j} : (i, j) \in V(\mathcal{K}_{m,n})\}$ on $\mathcal{K}_{m,n}$ in the following manner:

- For $1 \leq i \leq \frac{m}{2}^+$ and $1 \leq j \leq \frac{n}{2}$, let

$$(16) \quad x_{2i-1,j} = (-1)^{j+1} k_{\alpha(i), \rho(i)} + \sum_{\ell=1}^{j-1} (-1)^{\ell+1} a_{j-\ell}.$$

- For $1 \leq i \leq \frac{m}{2}^+$ and $1 \leq j \leq \frac{n}{2}$, let

$$(17) \quad x_{2i-1, n+1-j} = \frac{1}{2}S - x_{2i-1,j}.$$

- For $\frac{m}{2}^+ + 1 \leq i \leq m$ and $1 \leq j \leq \frac{n}{2}$, let

$$(18) \quad x_{2(i - \frac{m}{2}^+), n+1-j} = (-1)^{j+1} k_{\alpha(i), \rho(i)} + \sum_{\ell=1}^{j-1} (-1)^{\ell+1} a_{j-\ell}.$$

- For $\frac{m}{2}^+ + 1 \leq i \leq m$ and $1 \leq j \leq \frac{n}{2}$, let

$$(19) \quad x_{2(i - \frac{m}{2}^+), j} = \frac{1}{2}S - x_{2(i - \frac{m}{2}^+), n+1-j}.$$

An argument similar to that in the case when $(a_1, a_2, \dots, a_{n_0})$ is not a palindrome establishes each of the following statements. We leave the details of the proof to the reader. Given a C_4 -face-magic labeling $X = \{x_{i,j} : (i, j) \in V(\mathcal{K}_{m,4})\}$ such that $\mathcal{L}(X) = K$, we can find a unique graph automorphism $A \in KBLS(m, 4)$, a unique permutation α on $\{1, 2, \dots, m\}$ such that $\alpha(1) = 1$, and a unique function $\rho : \{1, 2, \dots, m\} \rightarrow \{1, 2\}$ such that $\rho(1) = 1$ for which $A(X) = L(K, \alpha, \rho)$. Furthermore, each distinct choice of a permutation α on $\{1, 2, \dots, m\}$ such that $\alpha(1) = 1$, and each distinct choice of a function $\rho : \{1, 2, \dots, m\} \rightarrow \{1, 2\}$ such that $\rho(1) = 1$ yields a Klein bottle nonequivalent C_4 -face-magic Klein bottle labeling $L(K, \alpha, \rho)$ on $\mathcal{K}_{m,4}$ such that $\mathcal{L}(L(K, \alpha, \rho)) = K$. Thus, there are $2^{m-1}(m-1)!$ Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X) = K$. \square

Definition 23. Let $m \geq 3$ be an odd integer, $n \geq 4$ be an even integer, $n_1 = \frac{n}{2} = n/2$, and $n_0 = \frac{n}{2} - 1$. Let a_1, a_2, \dots, a_{n_0} be positive integers.

- Suppose $(a_1, a_2, \dots, a_{n_0})$ is not a palindrome. Let K be the $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition of $G(a_1, a_2, \dots, a_{n_0})$ such that for each $(a_1, a_2, \dots, a_{n_0})$ -admissible path P of K , there exists $k_i \in T$ such that $\{k_i, mn + 1 - k_i\}$ is an end vertex of P , where k_i is defined in equation (10). Let ϵ be the identity function on $\{1, 2, \dots, m\}$. The labeling $L(K, \epsilon)$ as defined by equations (11) through (14) in the proof of Lemma 22 is called the *standard C_4 -face-magic Klein bottle labeling* on $\mathcal{K}_{m,n}$ associated with K .
- Suppose $(a_1, a_2, \dots, a_{n_0})$ is a palindrome. Let K be the $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition of $G(a_1, a_2, \dots, a_{n_0})$ such that for each $(a_1, a_2, \dots, a_{n_0})$ -admissible path P of K , there exists $\{k_{i,1}, k_{i,2}\} \in \tilde{T}$ such that $\{k_{i,1}, mn + 1 - k_{i,1}\}$ is an end vertex of P , where $\{k_{i,1}, k_{i,2}\}$ is defined in equation (15). Let ϵ be the identity function on $\{1, 2, \dots, m\}$ and let $\mu : \{1, 2, \dots, m\} \rightarrow \{1, 2\}$ be defined by $\mu(i) = 1$, for all $1 \leq i \leq m$. The labeling $L(K, \epsilon, \mu)$ as defined by equations (16) through (19) in the proof of Lemma 22 is called the *standard C_4 -face-magic Klein bottle labeling* on $\mathcal{K}_{m,n}$ associated with K .

Definition 24. Let $m \geq 3$ be an odd integer, $n \geq 4$ be an even integer, and $n_0 = \frac{n}{2} - 1$.

- (1) We let $NAPP(m, n)$ be the set of all non-palindromic sequences $(a_1, a_2, \dots, a_{n_0})$ such that there exists an $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition of $G(a_1, a_2, \dots, a_{n_0})$.
- (2) We let $PAPP(m, n)$ be the set of all palindromic sequences $(a_1, a_2, \dots, a_{n_0})$ such that there exists an $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition of $G(a_1, a_2, \dots, a_{n_0})$.

Given two non-palindromic sequences $(a_1, a_2, \dots, a_{n_0})$ and $(a'_1, a'_2, \dots, a'_{n_0})$ in $NAPP(m, n)$, we say that $(a_1, a_2, \dots, a_{n_0})$ is *path partition equivalent* to $(a'_1, a'_2, \dots, a'_{n_0})$ if there exist an $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition K of $G(a_1, a_2, \dots, a_{n_0})$, an $(a'_1, a'_2, \dots, a'_{n_0})$ -admissible path partition K' of $G(a'_1, a'_2, \dots, a'_{n_0})$, $A \in KBLS(m, n)$, and $\alpha \in S_n$ for which $\alpha(1) = 1$ such that $\mathcal{L}(A(L(K, \alpha))) = K'$. Here, S_n is the symmetry group of degree n .

Given two palindromic sequences $(a_1, a_2, \dots, a_{n_0})$ and $(a'_1, a'_2, \dots, a'_{n_0})$ in

$PAPP(m, n)$, we say that $(a_1, a_2, \dots, a_{n_0})$ is *path partition equivalent* to $(a'_1, a'_2, \dots, a'_{n_0})$ if there exist an $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition K of $G(a_1, a_2, \dots, a_{n_0})$, an $(a'_1, a'_2, \dots, a'_{n_0})$ -admissible path partition K' of $G(a'_1, a'_2, \dots, a'_{n_0})$, $A \in KBLS(m, n)$, $\alpha \in S_n$ for which $\alpha(1) = 1$, and a function $\rho : \{1, 2, \dots, n\} \rightarrow \{1, 2\}$ for which $\rho(1) = 1$ such that $\mathcal{L}(A(L(K, \alpha, \rho))) = K'$.

Remark 25. The path partition equivalence relation is an equivalence relation on both $NAPP(m, n)$ and $PAPP(m, n)$.

Lemma 26. *Let $m \geq 3$ be an odd integer, $n \geq 4$ be an even integer, and $n_0 = \# - 1$.*

- (1) *Let $(a_1, a_2, \dots, a_{n_0})$ and $(a'_1, a'_2, \dots, a'_{n_0})$ be non-palindromic sequences in $NAPP(m, n)$. Then, $(a_1, a_2, \dots, a_{n_0})$ is path partition equivalent to $(a'_1, a'_2, \dots, a'_{n_0})$ if and only if either $(a'_1, a'_2, \dots, a'_{n_0}) = (a_1, a_2, \dots, a_{n_0})$, $(a'_1, a'_2, \dots, a'_{n_0}) = (a_{n_0}, a_{n_0-1}, \dots, a_1)$, $(a'_1, a'_2, \dots, a'_{n_0}) = (S - a_1, S - a_2, \dots, S - a_{n_0})$, or $(a'_1, a'_2, \dots, a'_{n_0}) = (S - a_{n_0}, S - a_{n_0-1}, \dots, S - a_1)$.*
- (2) *Let $(a_1, a_2, \dots, a_{n_0})$ and $(a'_1, a'_2, \dots, a'_{n_0})$ be palindromic sequences in $PAPP(m, n)$. Then, $(a_1, a_2, \dots, a_{n_0})$ is path partition equivalent to $(a'_1, a'_2, \dots, a'_{n_0})$ if and only if either $(a'_1, a'_2, \dots, a'_{n_0}) = (a_1, a_2, \dots, a_{n_0})$ or $(a'_1, a'_2, \dots, a'_{n_0}) = (S - a_1, S - a_2, \dots, S - a_{n_0})$.*

Proof. We prove the lemma only for the case when $(a_1, a_2, \dots, a_{n_0})$ and $(a'_1, a'_2, \dots, a'_{n_0})$ are non-palindromic sequences in $NAPP(m, n)$. The case when $(a_1, a_2, \dots, a_{n_0})$ and $(a'_1, a'_2, \dots, a'_{n_0})$ are palindromic sequences in $PAPP(m, n)$ is similar, and we leave the details of the proof for that case to the reader.

(\Rightarrow) Let $(a_1, a_2, \dots, a_{n_0})$ and $(a'_1, a'_2, \dots, a'_{n_0})$ be non-palindromic sequences in $NAPP(m, n)$ such that $(a_1, a_2, \dots, a_{n_0})$ is path partition equivalent to $(a'_1, a'_2, \dots, a'_{n_0})$. Then, there exist an $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition K of $G(a_1, a_2, \dots, a_{n_0})$, an $(a'_1, a'_2, \dots, a'_{n_0})$ -admissible path partition K' of $G(a'_1, a'_2, \dots, a'_{n_0})$, $A \in KBLS(m, n)$, and $\alpha \in S_n$ for which $\alpha(1) = 1$ such that $\mathcal{L}(A(L(K, \alpha))) = K'$. Since $A \in KBLS(m, n)$, we have for some integer i , either $A = U^{2i}$, $A = U^{2i+1}$, $A = HU^{2i}$, $A = HU^{2i+1}$.

First, suppose $A = U^{2i}$ for some integer i . Then, $\mathcal{L}(A(L(K, \alpha)))$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition of $G(a_1, a_2, \dots, a_{n_0})$. However, $K' = \mathcal{L}(A(L(K, \alpha)))$ is an $(a'_1, a'_2, \dots, a'_{n_0})$ -admissible path partition of $G(a'_1, a'_2, \dots, a'_{n_0})$. Thus, $(a'_1, a'_2, \dots, a'_{n_0}) = (a_1, a_2, \dots, a_{n_0})$.

Second, suppose $A = U^{2i+1}$ for some integer i . Then, $\mathcal{L}(A(L(K, \alpha)))$ is an $(S - a_1, S - a_2, \dots, S - a_{n_0})$ -admissible path partition of $G(S - a_1, S - a_2, \dots, S - a_{n_0})$. However, $K' = \mathcal{L}(A(L(K, \alpha)))$ is an $(a'_1, a'_2, \dots, a'_{n_0})$ -admissible path partition of $G(a'_1, a'_2, \dots, a'_{n_0})$. Thus, $(a'_1, a'_2, \dots, a'_{n_0}) = (S - a_1, S - a_2, \dots, S - a_{n_0})$.

Third, suppose $A = HU^{2i}$ for some integer i . Then, $\mathcal{L}(A(L(K, \alpha)))$ is an $(S - a_{n_0}, S - a_{n_0-1}, \dots, S - a_1)$ -admissible path partition of $G(S - a_{n_0}, S - a_{n_0-1}, \dots, S - a_1)$. However, $K' = \mathcal{L}(A(L(K, \alpha)))$ is an $(a'_1, a'_2, \dots, a'_{n_0})$ -admissible path partition of $G(a'_1, a'_2, \dots, a'_{n_0})$. Thus, $(a'_1, a'_2, \dots, a'_{n_0}) = (S - a_{n_0}, S - a_{n_0-1}, \dots, S - a_1)$.

Fourth, suppose $A = HU^{2i+1}$ for some integer i . Then, $\mathcal{L}(A(L(K, \alpha)))$ is an $(a_{n_0}, a_{n_0-1}, \dots, a_1)$ -admissible path partition of $G(a_{n_0}, a_{n_0-1}, \dots, a_1)$. However, $K' = \mathcal{L}(A(L(K, \alpha)))$ is an $(a'_1, a'_2, \dots, a'_{n_0})$ -admissible path partition of $G(a'_1, a'_2, \dots, a'_{n_0})$. Thus, $(a'_1, a'_2, \dots, a'_{n_0}) = (a_{n_0}, a_{n_0-1}, \dots, a_1)$.

(\Leftarrow) Suppose either $(a'_1, a'_2, \dots, a'_{n_0}) = (a_1, a_2, \dots, a_{n_0})$, $(a'_1, a'_2, \dots, a'_{n_0}) = (a_{n_0}, a_{n_0-1}, \dots, a_1)$, $(a'_1, a'_2, \dots, a'_{n_0}) = (S - a_1, S - a_2, \dots, S - a_{n_0})$, or $(a'_1, a'_2, \dots, a'_{n_0}) = (S - a_{n_0}, S - a_{n_0-1}, \dots, S - a_1)$. We want to show that $(a_1, a_2, \dots, a_{n_0})$ is path partition equivalent to $(a'_1, a'_2, \dots, a'_{n_0})$. Since $(a_1, a_2, \dots, a_{n_0})$ is in $NAPP(m, n)$, there is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition K of $G(a_1, a_2, \dots, a_{n_0})$.

First, suppose $(a'_1, a'_2, \dots, a'_{n_0}) = (a_1, a_2, \dots, a_{n_0})$. Let $A = U^0 \in KBLS(m, n)$. Then, K is an $(a'_1, a'_2, \dots, a'_{n_0})$ -admissible path partition of $G(a'_1, a'_2, \dots, a'_{n_0})$ such that $K = \mathcal{L}(A(L(K, \epsilon)))$. Thus, $(a_1, a_2, \dots, a_{n_0})$ is path partition equivalent to $(a'_1, a'_2, \dots, a'_{n_0})$.

Second, suppose $(a'_1, a'_2, \dots, a'_{n_0}) = (a_{n_0}, a_{n_0-1}, \dots, a_1)$. Let $A = HU \in KBLS(m, n)$ and let $K' = \mathcal{L}(A(L(K, \epsilon)))$. Then, K' is an $(a'_1, a'_2, \dots, a'_{n_0})$ -admissible path partition of $G(a'_1, a'_2, \dots, a'_{n_0})$ such that $K' = \mathcal{L}(A(L(K, \epsilon)))$. Thus, $(a_1, a_2, \dots, a_{n_0})$ is path partition equivalent to $(a'_1, a'_2, \dots, a'_{n_0})$.

The proofs of the remaining two cases when $(a'_1, a'_2, \dots, a'_{n_0}) = (S - a_1, S - a_2, \dots, S - a_{n_0})$ or $(a'_1, a'_2, \dots, a'_{n_0}) = (S - a_{n_0}, S - a_{n_0-1}, \dots, S - a_1)$ are similar to the proof of the case when $(a'_1, a'_2, \dots, a'_{n_0}) = (a_{n_0}, a_{n_0-1}, \dots, a_1)$. If $(a'_1, a'_2, \dots, a'_{n_0}) = (S - a_1, S - a_2, \dots, S - a_{n_0})$, we let $A = U \in KBLS(m, n)$. If $(a'_1, a'_2, \dots, a'_{n_0}) = (S - a_{n_0}, S - a_{n_0-1}, \dots, S - a_1)$, we let $A = H \in KBLS(m, n)$. \square

5. RESULTS ON AN $m \times 4$ KLEIN BOTTLE GRID GRAPH

Corollary 27. *Let $m \geq 3$ be an odd positive integer. Let a be a positive integer. Suppose $G(a)$ has a perfect matching K . Then, $\mathcal{K}_{m,4}$ has $2^{m-1}(m-1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings $X = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,4})\}$ on $\mathcal{K}_{m,4}$ such that $\mathcal{L}(X) = K$.*

Proof. Let K be a perfect matching on $G(a)$. Then, K is an (a) -admissible path partition of $G(a)$. Since the sequence (a) is a palindrome, by Lemma 22, there are $2^{m-1}(m-1)!$ Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,4}$ such that $\mathcal{L}(X) = K$. \square

Notation 28. Let a and b be integers such that $a \leq b$. The *discrete closed interval from a to b* is the set denoted by $[a, b] = \{i \in \mathbb{Z} : a \leq i \leq b\}$.

Theorem 29. *Let m be an odd positive integer such that $m \geq 3$. Then, the number of Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings on $\mathcal{K}_{m,4}$ is $2^m(m-1)!\tau(m)$, where $\tau(m)$ is the number of positive divisors of m .*

Proof. Let $X = \{x_{i,j} : (i,j) \in V(\mathcal{K}_{m,n})\}$ be a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,4}$ and let $a = x_{1,1} + x_{1,2}$. Then, $\mathcal{L}(X)$ is an (a) -admissible path partition of $G(a)$. Since $a = z_1 + z_2$ for integers z_1 and z_2 such that $1 \leq z_1, z_2 \leq 4m$, we have $2 \leq a \leq 8m$. For convenience, let S be the C_4 -face-magic value of X . By Lemma 7, $S = 2(4m + 1)$. If $a \geq \frac{1}{2}S = 4m + 1$, then $U(X)$ is a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,4}$ that is Klein bottle equivalent to X . Furthermore, $\mathcal{L}(U(X))$ is an $(S - a)$ -admissible path partition of $G(S - a)$, where $S - a \leq \frac{1}{2}S = 4m + 1$. By replacing a with $S - a$, we may assume that $2 \leq a \leq 4m + 1$.

If $G(a)$ has a perfect matching, then vertex $\{2m, 2m + 1\}$ is adjacent to some vertex $\{i, 4m + 1 - i\}$. Thus, there is an integer $z \in \{i, 4m + 1 - i\}$ such that $a \geq 2m + z \geq 2m + 1$. Also, all pairs of integers z_1 and z_2 such that $z_1 + z_2 = 4m + 1$ are paired in the set $\{z_1, z_2\}$. Thus, $a \neq 4m + 1$. Hence, $2m + 1 \leq a \leq 4m$.

Let $\{z_1, z_2\} \in V(G(a))$. If there are integers y_1 and y_2 such that $1 \leq y_1, y_2 \leq 4m$, $z_1 + y_1 = a$ and $z_2 + y_2 = a$, then vertex $\{z_1, z_2\}$ in $G(a)$ has degree two. However, the only integers y_1 and y_2 that satisfy this criterion are $y_1 = a - z_1$ and $y_2 = a - z_2$. Thus, the degree of vertex $\{z_1, z_2\}$ is at most two. Hence, $G(a)$ is a disjoint union of paths and cycles. For convenience, let $b = 4m + 1$. We show that there are no cycles in $G(a)$. For the purpose of contradiction, suppose $G(a)$ has a cycle on k vertices for some positive integer k . Let $(\{y_{i,1}, y_{i,2}\} : 1 \leq i \leq k)$ be a walk in $G(a)$ such that $y_{i,2} + y_{i+1,1} = a$, for $1 \leq i \leq k - 1$ and $y_{k,2} + y_{1,1} = a$. For convenience, for $j = 1, 2$, let $y_{k+1,j} = y_{1,j}$. Since $y_{i,1} + y_{i,2} = b$ for $1 \leq i \leq k + 1$, and $y_{i,2} + y_{i+1,1} = a$ for $1 \leq i \leq k$, we have $y_{i,1} = y_{1,1} + (i - 1)(a - b)$, for $1 \leq i \leq k + 1$. Then, $y_{1,1} = y_{k+1,1} = y_{1,1} + k(a - b)$ implies that $a = b$. However, $a < b$. This is a contradiction. Therefore, $G(a)$ is a disjoint union of paths. Since $G(a)$ is a disjoint union of paths, $G(a)$ has a perfect matching if and only if each path in $G(a)$ is a path on an even number of vertices. We next establish that for all integers k such that $1 \leq k \leq 2m$, $G(4m + 1 - k)$ has a perfect matching if and only if either $k = d$ or $k = 2d$ for some positive divisor d of m . We consider five cases to verify this statement.

Case 1. Let $k = d$ for some positive divisor d of m . Let $m = m_2d$ for some positive integer m_2 . Since d is odd, we can write $d = 2d_1 + 1$ for some nonnegative integer d_1 . Let $\ell = \frac{1}{2}(4m + 1 - d) = 2m - d_1$. The sequence of integers $(di + \ell : i \in [-(2m_2 - 1), 2m_2])$ satisfies $di + \ell \equiv d_1 + 1 \pmod{2d_1 + 1}$, and the corresponding descending sequence of integers $(4m + 1 - (di + \ell) : i \in [-(2m_2 - 1), 2m_2])$ satisfies $4m + 1 - (di + \ell) \equiv d_1 + 1 \pmod{2d_1 + 1}$. Thus, the sequence $(4m + 1 - (di + \ell) : i \in [-(2m_2 - 1), 2m_2])$ is the sequence $(di + \ell : i \in [-(2m_2 - 1), 2m_2])$ in reverse order. Hence, the sequence of vertices $(\{di + \ell, 4m + 1 - (di + \ell)\} : i \in [-(2m_2 - 1), 0])$ is a path in $G(4m + 1 - d)$ on $2m_2 = (2m)/d$ vertices.

Let $i_0 \in [1, d_1]$. The sequence of integers $(di + i_0 : i \in [0, 4m_2 - 1])$ satisfies $di + i_0 \equiv 1, 2, \dots, d_1 \pmod{2d_1 + 1}$ for $i_0 = 1, 2, \dots, d_1$, and the corresponding descending sequence of integers $(4m + 1 - (di + i_0) : i \in [0, 4m_2 - 1])$ satisfies $4m + 1 - (di + i_0) \equiv 2d_1 + 1, 2d_1, \dots, d_1 + 2 \pmod{2d_1 + 1}$ for $i_0 = 1, 2, \dots, d_1$. Thus, the sequence of vertices $(\{di + i_0, 4m + 1 - (di + i_0)\} : i \in [0, 4m_2 - 1])$ is a path in $G(4m + 1 - d)$ on $4m_2 = (4m)/d$ vertices, for all $i_0 \in [1, d_1]$. Hence, $G(4m + 1 - d)$ is a disjoint union of d_1 paths where each path has $(4m)/d$ vertices and one path on $(2m)/d$ vertices. Since $G(4m + 1 - d)$ is a disjoint union of paths where each path has an even number of vertices, $G(4m + 1 - d)$ has a unique perfect matching.

Case 2. Let $k = 2d$ for some positive divisor d of m . Let $m = m_2d$ for some positive integer m_2 . Let $i_0 \in [1, d]$. The sequence of integers $((2d)i + i_0 : i \in [0, 2m_2 - 1])$ satisfies $(2d)i + i_0 \equiv 1, 2, \dots, d \pmod{2d}$ for $i_0 = 1, 2, \dots, d$, and the corresponding descending sequence of integers $(4m + 1 - ((2d)i + i_0) : i \in [0, 2m_2 - 1])$

satisfies $4m + 1 - ((2d)i + i_0) \equiv 2d, 2d - 1, \dots, d + 1 \pmod{2d}$ for $i_0 = 1, 2, \dots, d$. Thus, the sequence of vertices $(\{(2d)i + i_0, 4m + 1 - ((2d)i + i_0)\} : i \in [0, 2m_2 - 1])$ is a path in $G(4m + 1 - 2d)$ on $2m_2 = (2m)/d$ vertices, for all $i_0 \in [1, d]$. Hence, $G(4m + 1 - 2d)$ is a disjoint union of d paths where each path has $(2m)/d$ vertices. Since $G(4m + 1 - 2d)$ is a disjoint union of paths where each path has an even number of vertices, $G(4m + 1 - 2d)$ has a unique perfect matching.

Case 3. Let $k = 4d$ for some positive divisor d of m . Let $m = m_2d$ for some positive integer m_2 . Let $i_0 \in [1, 2d]$. The sequence of integers $((4d)i + i_0 : i \in [0, m_2 - 1])$ satisfies $(4d)i + i_0 \equiv 1, 2, \dots, 2d \pmod{4d}$ for $i_0 = 1, 2, \dots, 2d$, and the corresponding descending sequence of integers $(4m + 1 - ((4d)i + i_0) : i \in [0, m_2 - 1])$ satisfies $4m + 1 - ((4d)i + i_0) \equiv 4d, 4d - 1, \dots, 2d + 1 \pmod{4d}$ for $i_0 = 1, 2, \dots, 2d$. Thus, the sequence of vertices $(\{(4d)i + i_0, 4m + 1 - ((4d)i + i_0)\} : i \in [0, m_2 - 1])$ is a path in $G(4m + 1 - 4d)$ on $m_2 = m/d$ vertices, for all $i_0 \in [1, 2d]$. Hence, $G(4m + 1 - 4d)$ is a disjoint union of $2d$ paths where each path has m/d vertices. Since $G(4m + 1 - 4d)$ has at least one path on an odd number of vertices, $G(4m + 1 - 4d)$ has no perfect matching.

Case 4. Let k be an even positive integer such that $k \neq 2d$ for all positive divisors d of m . Let $k = 2k_1$ for some positive integer k_1 . By the division algorithm, there exist integers q and r such that $4m = kq + r$ and $0 \leq r < k$. Let $r_1 = 2m - k_1q$. Then, $r = 2r_1$, $2m = k_1q + r_1$ and $0 \leq r_1 < k_1$. Since k_1 is not a divisor of m , $r_1 \neq 0$. Thus, $0 < r_1 < k_1$.

Let $i_0 \in [1, r_1]$. The sequence of integers $(ki + i_0 : i \in [0, q])$ satisfies $ki + i_0 \equiv 1, 2, \dots, r_1 \pmod{2k_1}$, for $i_0 = 1, 2, \dots, r_1$, and the corresponding descending sequence of integers $(4m + 1 - (ki + i_0) : i \in [0, q])$ satisfies $4m + 1 - (ki + i_0) \equiv 2r_1, 2r_1 - 1, \dots, r_1 + 1 \pmod{2k_1}$, for $i_0 = 1, 2, \dots, r_1$. Thus, the sequence of vertices $(\{ki + i_0, 4m + 1 - (ki + i_0)\} : i \in [0, q])$ is a path in $G(4m + 1 - k)$ on $q + 1$ vertices, for all $i_0 \in [1, r_1]$.

Let $i_0 \in [2r_1 + 1, k_1 + r_1]$. The sequence of integers $(ki + i_0 : i \in [0, q - 1])$ satisfies $ki + i_0 \equiv 2r_1 + 1, 2r_1 + 2, \dots, k_1 + r_1 \pmod{2k_1}$ for $i_0 = 2r_1 + 1, 2r_1 + 2, \dots, k_1 + r_1$, and the corresponding descending sequence of integers $(4m + 1 - (ki + i_0) : i \in [0, q - 1])$ satisfies $4m + 1 - (ki + i_0) \equiv 2k_1, 2k_1 - 1, \dots, k_1 + r_1 + 1 \pmod{2k_1}$, for $i_0 = 2r_1 + 1, 2r_1 + 2, \dots, k_1 + r_1$. Thus, the sequence of vertices $(\{ki + i_0, 4m + 1 - (ki + i_0)\} : i \in [0, q - 1])$ is a path in $G(4m + 1 - k)$ on q vertices, for all $i_0 \in [2r_1 + 1, k_1 + r_1]$.

Let v be the number of vertices in the r_1 paths, where each path has $q + 1$ vertices, and $k_1 - r_1$ paths, where each path has q vertices. Then, $v = r_1(q + 1) + (k_1 - r_1)q = k_1q + r_1 = 2m$. Hence, $G(4m + 1 - k)$ is a disjoint union of r_1 paths, where each path has $q + 1$ vertices, and $k_1 - r_1$ paths, where each path has q vertices. Since $r_1 > 0$, $k_1 - r_1 > 0$, and q and $q + 1$ have opposite parity, $G(4m + 1 - k)$ has at least one path on an odd number of vertices. Therefore, $G(4m + 1 - k)$ has no perfect matching.

Case 5. Let k be an odd positive integer such that $k \neq d$ for all positive divisors d of m . Let $k = 2k_1 + 1$ for some nonnegative integer k_1 . Since k is not a divisor of m , we have $k \neq 1$. Thus, $k_1 > 0$. Let $\ell = \frac{1}{2}(4m + 1 - k) = 2m - k_1$. By the division algorithm, there exist integers q and r such that $4m = kq + r$ and $0 \leq r < k$. Since k is not a divisor of m , $r \neq 0$. Thus, $0 < r < k$. Since $1 \leq k < 2m$, we have $q \geq 2$. We consider the subcases where either q is even or q is odd.

Subcase 5(i). Assume q is even. Let $q = 2q_1$ for some positive integer q_1 . Let $r_1 = 2m - kq_1$. Then, $r = 2r_1$. The sequence of integers $(ki + \ell : i \in [-(q_1 - 1), q_1])$ satisfies $ki + \ell \equiv r_1 + k_1 + 1 \pmod{2k_1 + 1}$, and the corresponding descending sequence of integers $(4m + 1 - (ki + \ell) : i \in [-(q_1 - 1), q_1])$ satisfies $4m + 1 - (ki + \ell) \equiv r_1 + k_1 + 1 \pmod{2k_1 + 1}$. Thus, the sequence $(4m + 1 - (ki + \ell) : i \in [-(q_1 - 1), q_1])$ is the sequence $(ki + \ell : i \in [-(q_1 - 1), q_1])$ in reverse order. Hence, the sequence of vertices $(\{ki + \ell, 4m + 1 - (ki + \ell)\} : i \in [-(q_1 - 1), 0])$ is a path in $G(4m + 1 - k)$ on q_1 vertices.

Let $i_0 \in [1, r_1]$. The sequence of integers $(ki + i_0 : i \in [0, q])$ satisfies $ki + i_0 \equiv 1, 2, \dots, r_1 \pmod{2k_1 + 1}$ for $i_0 = 1, 2, \dots, r_1$, and the corresponding descending sequence of integers $(4m + 1 - (ki + i_0) : i \in [0, q])$ satisfies $4m + 1 - (ki + i_0) \equiv 2r_1, 2r_1 - 1, \dots, r_1 + 1 \pmod{2k_1 + 1}$, for $i_0 = 1, 2, \dots, r_1$. Thus, the sequence of vertices $(\{ki + i_0, 4m + 1 - (ki + i_0)\} : i \in [0, q])$ is a path in $G(4m + 1 - k)$ on $q + 1$ vertices, for all $i_0 \in [1, r_1]$.

Since $2r_1 = r < k = 2k_1 + 1$, we have $r_1 \leq k_1$. Let $i_0 \in [2r_1 + 1, k_1 + r_1]$. The sequence of integers $(ki + i_0 : i \in [0, q - 1])$ satisfies $ki + i_0 \equiv 2r_1 + 1, 2r_1 + 2, \dots, k_1 + r_1 \pmod{2k_1 + 1}$ for $i_0 = 2r_1 + 1, 2r_1 + 2, \dots, k_1 + r_1$, and the corresponding descending sequence of integers $(4m + 1 - (ki + i_0) : i \in [0, q - 1])$ satisfies $4m + 1 - (ki + i_0) \equiv 2k_1 + 1, 2k_1, \dots, k_1 + r_1 + 2 \pmod{2k_1 + 1}$, for $i_0 = 2r_1 + 1, 2r_1 + 2, \dots, k_1 + r_1$.

Thus, the sequence of vertices $(\{ki + i_0, 4m + 1 - (ki + i_0)\} : i \in [0, q - 1])$ is a path in $G(4m + 1 - k)$ on q vertices, for all $i_0 \in [2r_1 + 1, k_1 + r_1]$.

Let v be the number of vertices in the r_1 paths, where each path has $q + 1$ vertices, $k_1 - r_1$ paths, where each path has q vertices, and one path on q_1 vertices. Then, $v = r_1(q + 1) + (k_1 - r_1)q + q_1 = kq_1 + r_1 = 2m$. Hence, $G(4m + 1 - k)$ is a disjoint union of r_1 paths, where each path has $q + 1$ vertices, $k_1 - r_1$ paths, where each path has q vertices, and one path on q_1 vertices. Since $r_1 > 0$ and $q + 1$ is odd, $G(4m + 1 - k)$ has at least one path on an odd number of vertices. Therefore, $G(4m + 1 - k)$ has no perfect matching.

Subcase 5(ii). Assume q is odd. Let $q + 1 = 2q_1$ for some positive integer q_1 . Then, $r = 4m - kq$ is odd. Let $r = 2r_1 + 1$ for some nonnegative integer r_1 . Since $r < k$, we have $r_1 < k_1$. Recall that $\ell = \frac{1}{2}(4m + 1 - k) = 2m - k_1$.

The sequence of integers $(ki + \ell : i \in [-(q_1 - 1), q_1])$ satisfies $ki + \ell \equiv r_1 + 1 \pmod{2k_1 + 1}$ and the corresponding descending sequence of integers $(4m + 1 - (ki + \ell) : i \in [-(q_1 - 1), q_1])$ satisfies $4m + 1 - (ki + \ell) \equiv r_1 + 1 \pmod{2k_1 + 1}$. Thus, the sequence $(4m + 1 - (ki + \ell) : i \in [-(q_1 - 1), q_1])$ is the sequence $(ki + \ell : i \in [-(q_1 - 1), q_1])$ in reverse order. Hence, the sequence of vertices $(\{ki + \ell, 4m + 1 - (ki + \ell)\} : i \in [-(q_1 - 1), 0])$ is a path in $G(4m + 1 - k)$ on q_1 vertices.

Let $i_0 \in [1, r_1]$. The sequence of integers $(ki + i_0 : i \in [0, q])$ satisfies $ki + i_0 \equiv 1, 2, \dots, r_1 \pmod{2k_1 + 1}$ for $i_0 = 1, 2, \dots, r_1$, and the corresponding descending sequence of integers $(4m + 1 - (ki + i_0) : i \in [0, q])$ satisfies $4m + 1 - (ki + i_0) \equiv 2r_1 + 1, 2r_1, \dots, r_1 + 2 \pmod{2k_1 + 1}$, for $i_0 = 1, 2, \dots, r_1$. Thus, the sequence of vertices $(\{ki + i_0, 4m + 1 - (ki + i_0)\} : i \in [0, q])$ is a path in $G(4m + 1 - k)$ on $q + 1$ vertices, for all $i_0 \in [1, r_1]$.

Let $i_0 \in [2r_1 + 2, k_1 + r_1 + 1]$. The sequence of integers $(ki + i_0 : i \in [0, q - 1])$ satisfies $ki + i_0 \equiv 2r_1 + 2, 2r_1 + 3, \dots, k_1 + r_1 + 1 \pmod{2k_1 + 1}$ for $i_0 = 2r_1 + 2, 2r_1 + 3, \dots, k_1 + r_1 + 1$, and the corresponding descending sequence of integers $(4m + 1 - (ki + i_0) : i \in [0, q - 1])$ satisfies $4m + 1 - (ki + i_0) \equiv 2k_1 + 1, 2k_1, \dots, k_1 + r_1 + 2 \pmod{2k_1 + 1}$, for $i_0 = 2r_1 + 2, 2r_1 + 3, \dots, k_1 + r_1 + 1$. Thus, the sequence of vertices $(\{ki + i_0, 4m + 1 - (ki + i_0)\} : i \in [0, q - 1])$ is a path in $G(4m + 1 - k)$ on q vertices, for all $i_0 \in [2r_1 + 2, k_1 + r_1 + 1]$.



Let v be the number of vertices in the r_1 paths, where each path has $q + 1$ vertices, $k_1 - r_1$ paths, where each path has q vertices, and one path on q_1 vertices. Then, $v = r_1(q + 1) + (k_1 - r_1)q + q_1 = kq_1 - k_1 + r_1 = 2m$. Hence, $G(4m + 1 - k)$ is a disjoint union of r_1 paths, where each path has $q + 1$ vertices, $k_1 - r_1$ paths, where each path has q vertices, and one path on q_1 vertices. Since $k_1 - r_1 > 0$ and q is odd, $G(4m + 1 - k)$ has at least one path on an odd number of vertices. Therefore, $G(4m + 1 - k)$ has no perfect matching.

We define the *perfect matching set* of m , denoted by $PMS(m)$, by

$$PMS(m) = \{a = 4m + 1 - k : k = d \text{ or } k = 2d, \text{ for some positive divisor } d \text{ of } m\}.$$

Let $a \in PMS(m)$. Because $G(a)$ is a disjoint union of paths where each path has an even number of vertices, $G(a)$ has exactly one perfect matching K_a . Furthermore, suppose we are given two integers $a, a' \in PMS(m)$ such that $2m + 1 \leq a, a' \leq 4m$, and there are Klein bottle equivalent C_4 -face-magic Klein bottle labeling X and X' for which $\mathcal{L}(X) = K_a$ and $\mathcal{L}(X') = K_{a'}$. Then, there exists $A \in KBLS(m, 4)$ such that $X' = A(X)$. We have $X = B(L(K_a, \alpha))$ for some $\alpha \in S_n$ for which $\alpha(1) = 1$ and $B \in KBLS(m, 4)$. Hence, $K_{a'} = \mathcal{L}((AB)(L(K_a, \alpha)))$. Thus, (a) is path partition equivalent to (a') . By Lemma 26, we have either $a' = a$ or $a' = S - a$. Since $a' \leq 4m$ and $S - a \geq 4m + 2$, we have $a' \neq S - a$. Thus, $a' = a$. For each value $a \in PMS(m)$, by Corollary 27, there are $2^{m-1}(m-1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,4}$ such that $\mathcal{L}(X) = K_a$. We have $|PMS(m)| = 2\tau(m)$, where $\tau(m)$ is the number of positive divisors of m . Therefore, there are $2\tau(m) \cdot 2^{m-1}(m-1)! = 2^m(m-1)!\tau(m)$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings on $\mathcal{K}_{m,4}$. This completes the proof. \square

Example 30. In Figure 2 we have the labeling search graph $G(16)$ for the 5×4 Klein bottle grid graph $\mathcal{K}_{5,4}$. In Figure 3 we have a (16)-admissible path partition K of $G(16)$ that is a perfect matching on $G(16)$. The standard C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{5,4}$ associated with K denoted by $L(K, \epsilon, \mu)$ is given in Table 4. See Definition 23.

FIGURE 2. Labeling search graph $G(16)$ for the 5×4 Klein bottle grid graph.FIGURE 3. (16) -admissible path partition K of $G(16)$ is a perfect matching on $G(16)$.

1	15	6	20
17	9	12	4
2	14	7	19
16	10	11	5
3	13	8	18

TABLE 4. The standard C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{5,4}$ associated with K where K is the (16) -admissible path partition of $G(16)$ given in Figure 3.

6. RESULTS ON AN $m \times 6$ KLEIN BOTTLE GRID GRAPH

Notation 31. For convenience, we let S be the C_4 -face-magic value of a C_4 -face-magic Klein bottle labeling X of $\mathcal{K}_{m,6}$. By Lemma 7, we have $S = 12m + 2$.

Proposition 32. *Let $m \geq 3$ be an odd integer and let b be an integer such that either $3m + 1 \leq b \leq 4m + 1$ or $8m \leq b \leq 9m + 1$. Then, $G(6m, b)$ has a $(6m, b)$ -admissible path partition K_b and thus, there exist $(m - 1)!$ Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,6}$ such that $\mathcal{L}(X) = K_b$.*

Proof. Case 1. Suppose $3m + 1 \leq b \leq 4m + 1$. Let $b = 3m + k$, where k is an integer such that $1 \leq k \leq m + 1$. We determine a $(6m, b)$ -admissible path partition of $G(6m, b)$. For all integers i such that $1 \leq i \leq k - 1$, we

let

$$(20) \quad x_{i,1,k} = 6m - 2i + 1,$$

$$(21) \quad \frac{1}{2}S - x_{i,1,k} = 2i,$$

$$(22) \quad x_{i,2,k} = 2i - 1,$$

$$(23) \quad \frac{1}{2}S - x_{i,2,k} = 6m - 2i + 2,$$

$$(24) \quad x_{i,3,k} = 3m - 2i + k + 1, \text{ and}$$

$$(25) \quad \frac{1}{2}S - x_{i,3,k} = 3m + 2i - k.$$

For all integers i such that $k \leq i \leq m$, we let

$$(26) \quad x_{i,1,k} = 6m - 3i + k + 1,$$

$$(27) \quad \frac{1}{2}S - x_{i,1,k} = 3i - k,$$

$$(28) \quad x_{i,2,k} = 3i - k - 1,$$

$$(29) \quad \frac{1}{2}S - x_{i,2,k} = 6m - 3i + k + 2,$$

$$(30) \quad x_{i,3,k} = 3m - 3i + 2k + 1, \text{ and}$$

$$(31) \quad \frac{1}{2}S - x_{i,3,k} = 3m + 3i - 2k.$$

From equations (21) and (22), for $1 \leq i \leq k - 1$, we obtain the labels $\{q \in \mathbb{Z} : 1 \leq q \leq 2k - 2\}$. From equations (27), (28), and (30), for $k \leq i \leq m$, we obtain the labels $\{q \in \mathbb{Z} : 2k - 1 \leq q \leq 3m - k + 1\}$. From equation (24), for $k + 1 \leq i \leq k - 1$, and equation (25), for $1 \leq i \leq k$, we obtain the labels $\{q \in \mathbb{Z} : 3m - k + 2 \leq q \leq 3m\}$. However, it should be noted that the details of the argument in this last case depends on whether k is even or odd. Thus for $1 \leq i \leq m$ and $1 \leq j \leq 3$, the vertices $\{x_{i,j,k}, \frac{1}{2}S - x_{i,j,k}\}$ in the graph $G(6m, b)$ are distinct.

We observe that for all $1 \leq i \leq m$ and $1 \leq k \leq m + 1$, we have $x_{i,1,k} + x_{i,2,k} = 6m$ and $x_{i,2,k} + x_{i,3,k} = 3m + k = b$. Hence for all $1 \leq i \leq m$ and $1 \leq k \leq m + 1$, the path $P_{i,k} = (\{x_{i,j,k}, \frac{1}{2}S - x_{i,j,k}\} : 1 \leq j \leq 3)$ is a $(6m, b)$ -admissible path in $G(6m, b)$ with vertex label sequence $(x_{i,j,k} : 1 \leq j \leq 3)$. Therefore for all $1 \leq k \leq m + 1$, $K_{3m+k} = \{P_{i,k} : 1 \leq i \leq m\}$ is a $(6m, b)$ -admissible path partition of $G(6m, b)$. Since $(6m, b)$ is not a palindrome, by Lemma 22, for each $3m + 1 \leq b \leq 4m + 1$, there exist $(m - 1)!$ Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X of $\mathcal{K}_{m,6}$ such that $\mathcal{L}(X) = K_b$.

Case 2. Suppose $8m \leq b \leq 9m + 1$. Let $b = 8m + k$, where k is an integer such that $0 \leq k \leq m + 1$. We determine a $(6m, b)$ -admissible path partition of $G(6m, b)$. For all integers i such that $1 \leq i \leq k - 1$, we let

$$(32) \quad x_{i,1,k} = 2m - 3i + k + 1,$$

$$(33) \quad \frac{1}{2}S - x_{i,1,k} = 4m + 3i - k,$$

$$(34) \quad x_{i,2,k} = 4m + 3i - k - 1,$$

$$(35) \quad \frac{1}{2}S - x_{i,2,k} = 2m - 3i + k + 2,$$

$$(36) \quad x_{i,3,k} = 4m - 3i + 2k + 1, \text{ and}$$

$$(37) \quad \frac{1}{2}S - x_{i,3,k} = 2m + 3i - 2k.$$

For all integers i such that $k \leq i \leq m$, we let

$$(38) \quad x_{i,1,k} = 2m - 2i + 1,$$

$$(39) \quad \frac{1}{2}S - x_{i,1,k} = 4m + 2i,$$

$$(40) \quad x_{i,2,k} = 4m + 2i - 1,$$

$$(41) \quad \frac{1}{2}S - x_{i,2,k} = 2m - 2i + 2,$$

$$(42) \quad x_{i,3,k} = 4m - 2i + k + 1, \text{ and}$$

$$(43) \quad \frac{1}{2}S - x_{i,3,k} = 2m + 2i - k.$$

From equations (42) and (43), for $k \leq i \leq m$, we obtain the labels $\{q \in \mathbb{Z} : 1 \leq q \leq 2m - 2k + 2\}$. From equations (32), (35), and (37), for $1 \leq i \leq k - 1$, we obtain the labels $\{q \in \mathbb{Z} : 2m - 2k + 3 \leq q \leq 2m + k - 1\}$. From equation (42), for $m^+ + k^+ \leq i \leq m$, and equation (43), for $k \leq i \leq m^+ + k^+ - 1$, we obtain the labels

$\{q \in \mathbb{Z} : 2m + k \leq q \leq 3m\}$. However, it should be noted that the details of the argument in this last case depends on whether k is even or odd. Thus for $1 \leq i \leq m$ and $1 \leq j \leq 3$, the vertices $\{x_{i,j,k}, \frac{1}{2}S - x_{i,j,k}\}$ in the graph $G(6m, b)$ are distinct.

We observe that, for all $1 \leq i \leq m$ and $0 \leq k \leq m + 1$, we have $x_{i,1,k} + x_{i,2,k} = 6m$ and $x_{i,2,k} + x_{i,3,k} = 8m + k = b$. Hence for all $1 \leq i \leq m$ and $0 \leq k \leq m + 1$, the path $P_{i,k} = (\{x_{i,j,k}, \frac{1}{2}S - x_{i,j,k}\} : 1 \leq j \leq 3)$ is a $(6m, b)$ -admissible path in $G(6m, b)$ with vertex label sequence $(x_{i,j,k} : 1 \leq j \leq 3)$. Therefore for each $0 \leq k \leq m + 1$, $K_{8m+k} = \{P_{i,k} : 1 \leq i \leq m\}$ is a $(6m, b)$ -admissible path partition of $G(6m, b)$. Since $(6m, b)$ is not a palindrome, by Lemma 22, for each $8m \leq b \leq 9m + 1$, there exist $(m - 1)!$ Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X of $\mathcal{K}_{m,6}$ such that $\mathcal{L}(X) = K_b$. \square

Proposition 33. *Let $m \geq 3$ be an odd integer and let k be an integer such that $1 \leq k \leq m + 1$. Then, $G(2m + k, 3m + k)$ has a $(2m + k, 3m + k)$ -admissible path partition K_k and thus, there are $(m - 1)!$ Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,6}$ such that $\mathcal{L}(X) = K_k$.*

Proof. We determine a $(2m + k, 3m + k)$ -admissible path partition of $G(2m + k, 3m + k)$. For all integers i such that $1 \leq i \leq k - 1$, we let

$$(44) \quad x_{i,1,k} = k - i,$$

$$(45) \quad \frac{1}{2}S - x_{i,1,k} = 6m + i - k + 1,$$

$$(46) \quad x_{i,2,k} = 2m + i,$$

$$(47) \quad \frac{1}{2}S - x_{i,2,k} = 4m - i + 1,$$

$$(48) \quad x_{i,3,k} = m - i + k, \text{ and}$$

$$(49) \quad \frac{1}{2}S - x_{i,3,k} = 5m + i - k + 1.$$

For all integers i such that $k \leq i \leq m$, we let

$$(50) \quad x_{i,1,k} = 2m - i + k,$$

$$(51) \quad \frac{1}{2}S - x_{i,1,k} = 4m + i - k + 1,$$

$$(52) \quad x_{i,2,k} = i,$$

$$(53) \quad \frac{1}{2}S - x_{i,2,k} = 6m - i + 1,$$

$$(54) \quad x_{i,3,k} = 3m - i + k, \text{ and}$$

$$(55) \quad \frac{1}{2}S - x_{i,3,k} = 3m + i - k + 1.$$

From equation (44), for $1 \leq i \leq k - 1$, and equation (52), for $k \leq i \leq m$, we obtain the labels $\{q \in \mathbb{Z} : 1 \leq q \leq m\}$. From equation (48), for $1 \leq i \leq k - 1$, and equation (50), for $k \leq i \leq m$, we obtain the labels $\{q \in \mathbb{Z} : m + 1 \leq q \leq 2m\}$. From equation (46), for $1 \leq i \leq k - 1$, and equation (54), for $k \leq i \leq m$, we obtain the labels $\{q \in \mathbb{Z} : 2m + 1 \leq q \leq 3m\}$. Thus for $1 \leq i \leq m$ and $1 \leq j \leq 3$, the vertices $\{x_{i,j}, \frac{1}{2}S - x_{i,j}\}$ in the graph $G(2m + k, 3m + k)$ are distinct.

We observe that, for all $1 \leq i \leq m$ and $1 \leq k \leq m + 1$, we have $x_{i,1,k} + x_{i,2,k} = 2m + k$ and $x_{i,2,k} + x_{i,3,k} = 3m + k$. Hence for all $1 \leq i \leq m$ and $1 \leq k \leq m + 1$, the path $P_{i,k} = (\{x_{i,j,k}, \frac{1}{2}S - x_{i,j,k}\} : 1 \leq j \leq 3)$ is a $(2m + k, 3m + k)$ -admissible path in $G(2m + k, 3m + k)$ with vertex label sequence $(x_{i,j,k} : 1 \leq j \leq 3)$. Therefore, $K_k = \{P_{i,k} : 1 \leq i \leq m\}$ is a $(2m + k, 3m + k)$ -admissible path partition of $G(2m + k, 3m + k)$. Since $(2m + k, 3m + k)$ is not a palindrome, by Lemma 22, for each $1 \leq k \leq m + 1$, there exist $(m - 1)!$ Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X of $\mathcal{K}_{m,6}$ such that $\mathcal{L}(X) = K_k$. \square


7. RESULTS ON AN $m \times n$ KLEIN BOTTLE GRID GRAPH

We first introduce functions ρ_1 and ρ_2 .

Notation 34. Let $i \in \{1, 2\}$. Suppose n_1 is a positive integer. We define $\rho_i : \{1, 2, \dots, n_1\} \rightarrow \{1, 2\}$ by $\rho_i(k) = i$ for all $k \in \{1, 2, \dots, n_1\}$.

Proposition 35. *Let $m \geq 3$ be an odd integer. Let $n \geq 6$ be an even integer, $n_1 = n/2$ and $n_0 = n_1 - 1$. Let $S = 2(mn + 1)$. Let $(a_1, a_2, \dots, a_{n_0})$ be one of the following three sequences:*

- (1) $a_k = mn - n_0 + k$, for all $1 \leq k \leq n_0$,
- (2) $a_k = m(n_1 + k) + 1$, for all $1 \leq k \leq n_0$, or

FIGURE 4. $(27, 28, 29, 30)$ -admissible path partition K_{ρ_1} in $G(27, 28, 29, 30)$.

(3) $a_k = 2mk + 1$, for all $1 \leq k \leq n_0$.

Then, $G(a_1, a_2, \dots, a_{n_0})$ has at least 2^m distinct $(a_1, a_2, \dots, a_{n_0})$ -admissible path partitions and thus, for each distinct $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition K of $G(a_1, a_2, \dots, a_{n_0})$, there exist $(m-1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X) = K$.

Proof. We consider each sequence $(a_1, a_2, \dots, a_{n_0})$ individually. The proof of the proposition for each sequence is similar.

Case 1. Suppose $a_k = mn - n_0 + k$, for all $1 \leq k \leq n_0$. We define the labels of $(a_1, a_2, \dots, a_{n_0})$ -admissible paths in $G(a_1, a_2, \dots, a_{n_0})$ as follows: for all $1 \leq i \leq m$, we let

$$\begin{aligned}
 x_{1,i,2j-1} &= n_1(i-1) + j, & \text{for } 1 \leq j \leq \mathfrak{n}_1^+, \\
 \frac{1}{2}S - x_{1,i,2j-1} &= mn - n_1(i-1) - j + 1, & \text{for } 1 \leq j \leq \mathfrak{n}_1^+, \\
 x_{1,i,2j} &= mn - n_1i + j, & \text{for } 1 \leq j \leq \mathfrak{n}_1, \\
 \frac{1}{2}S - x_{1,i,2j} &= n_1i - j + 1, & \text{for } 1 \leq j \leq \mathfrak{n}_1, \\
 x_{2,i,2j-1} &= mn - n_1i + j, & \text{for } 1 \leq j \leq \mathfrak{n}_1^+, \\
 \frac{1}{2}S - x_{2,i,2j-1} &= n_1i - j + 1, & \text{for } 1 \leq j \leq \mathfrak{n}_1^+, \\
 x_{2,i,2j} &= n_1(i-1) + j, & \text{for } 1 \leq j \leq \mathfrak{n}_1, \text{ and} \\
 \frac{1}{2}S - x_{2,i,2j} &= mn - n_1(i-1) - j + 1, & \text{for } 1 \leq j \leq \mathfrak{n}_1.
 \end{aligned}$$

Let $\rho : \{1, 2, \dots, m\} \rightarrow \{1, 2\}$. Let $P_{\rho,i}$ be the path on the sequence of vertices $(\{x_{\rho(i),i,j}, \frac{1}{2}S - x_{\rho(i),i,j}\} : 1 \leq j \leq n_1)$ with vertex label sequence $(x_{\rho(i),i,j} : 1 \leq j \leq n_1)$. The sets of labels

$$\begin{aligned}
 &\{x_{1,i,2j-1} : 1 \leq j \leq \mathfrak{n}_1^+\} \cup \{\frac{1}{2}S - x_{1,i,2j} : 1 \leq j \leq \mathfrak{n}_1\} \text{ and} \\
 &\{x_{2,i,2j} : 1 \leq j \leq \mathfrak{n}_1\} \cup \{\frac{1}{2}S - x_{2,i,2j-1} : 1 \leq j \leq \mathfrak{n}_1^+\}
 \end{aligned}$$


are the set $\{n_1(i-1) + k : 1 \leq k \leq n_1\}$. Hence for $1 \leq i \leq m$ and $1 \leq j \leq n_1$, the vertices $\{x_{\rho(i),i,j}, \frac{1}{2}S - x_{\rho(i),i,j}\}$ are distinct in the graph $G(a_1, a_2, \dots, a_{n_0})$.

Furthermore for $1 \leq i \leq m$, we have

$$\begin{aligned}
 x_{\rho(i),i,2j-1} + x_{\rho(i),i,2j} &= (n_1(i-1) + j) + (mn - n_1i + j) \\
 &= mn - n_0 + (2j - 1) = a_{2j-1}, & \text{for } 1 \leq j \leq \mathfrak{n}_1, \text{ and} \\
 x_{\rho(i),i,2j} + x_{\rho(i),i,2j+1} &= x_{\rho(i),i,2j} + x_{\rho(i),i,2j-1} + 1 \\
 &= mn - n_0 + 2j = a_{2j}, & \text{for } 1 \leq j \leq \mathfrak{n}_1^-.
 \end{aligned}$$

Thus, $P_{\rho,i}$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in $G(a_1, a_2, \dots, a_{n_0})$.

We consider the example where $m = 3$, $n_1 = 5$ and $n = 10$. Then, Figures 4 and 5 provide the $(27, 28, 29, 30)$ -admissible path partitions K_{ρ_1} and K_{ρ_2} in the graph $G(27, 28, 29, 30)$. See Notation 34 for the definitions of ρ_1 and ρ_2 .

FIGURE 5. $(27, 28, 29, 30)$ -admissible path partition K_{ρ_2} in $G(27, 28, 29, 30)$.

Case 2. Suppose $a_k = m(n_1 + k) + 1$, for all $1 \leq k \leq n_0$. We define the labels of $(a_1, a_2, \dots, a_{n_0})$ -admissible paths in $G(a_1, a_2, \dots, a_{n_0})$ as follows: for all $1 \leq i \leq m$, we let

$$\begin{aligned}
 x_{1,i,2j-1} &= m(j-1) + i, & \text{for } 1 \leq j \leq \pi_1^+, \\
 \frac{1}{2}S - x_{1,i,2j-1} &= m(n-j+1) - i + 1, & \text{for } 1 \leq j \leq \pi_1^+, \\
 x_{1,i,2j} &= m(n_1 + j) - i + 1, & \text{for } 1 \leq j \leq \pi_1, \\
 \frac{1}{2}S - x_{1,i,2j} &= m(n_1 - j) + i, & \text{for } 1 \leq j \leq \pi_1, \\
 x_{2,i,2j-1} &= m(n_1 + j) - i + 1, & \text{for } 1 \leq j \leq \pi_1^+, \\
 \frac{1}{2}S - x_{2,i,2j-1} &= m(n_1 - j) + i, & \text{for } 1 \leq j \leq \pi_1^+, \\
 x_{2,i,2j} &= m(j-1) + i, & \text{for } 1 \leq j \leq \pi_1, \text{ and} \\
 \frac{1}{2}S - x_{2,i,2j} &= m(n-j+1) - i + 1, & \text{for } 1 \leq j \leq \pi_1.
 \end{aligned}$$

Let $\rho : \{1, 2, \dots, m\} \rightarrow \{1, 2\}$. Let $P_{\rho,i}$ be the path on the sequence of vertices $(\{x_{\rho(i),i,j}, \frac{1}{2}S - x_{\rho(i),i,j}\} : 1 \leq j \leq n_1)$ with vertex label sequence $(x_{\rho(i),i,j} : 1 \leq j \leq n_1)$. The sets of labels

$$\begin{aligned}
 &\{x_{1,i,2j-1} : 1 \leq j \leq \pi_1^+\} \cup \{\frac{1}{2}S - x_{1,i,2j} : 1 \leq j \leq \pi_1\} \text{ and} \\
 &\{x_{2,i,2j} : 1 \leq j \leq \pi_1\} \cup \{\frac{1}{2}S - x_{2,i,2j-1} : 1 \leq j \leq \pi_1^+\}
 \end{aligned}$$



are the set $\{mk + i : 0 \leq k \leq n_1 - 1\}$. Hence for $1 \leq i \leq m$ and $1 \leq j \leq n_1$, the vertices $\{x_{\rho(i),i,j}, \frac{1}{2}S - x_{\rho(i),i,j}\}$ are distinct in the graph $G(a_1, a_2, \dots, a_{n_0})$.

Furthermore, for $1 \leq i \leq m$, we have

$$\begin{aligned}
 x_{\rho(i),i,2j-1} + x_{\rho(i),i,2j} &= (m(j-1) + i) + (m(n_1 + j) - i + 1) \\
 &= m(n_1 + 2j - 1) + 1 = a_{2j-1}, & \text{for } 1 \leq j \leq \pi_1, \text{ and} \\
 x_{\rho(i),i,2j} + x_{\rho(i),i,2j+1} &= x_{\rho(i),i,2j} + x_{\rho(i),i,2j-1} + m \\
 &= m(n_1 + 2j) + 1 = a_{2j}, & \text{for } 1 \leq j \leq \pi_1^-.
 \end{aligned}$$

Thus, $P_{\rho,i}$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in $G(a_1, a_2, \dots, a_{n_0})$.

We consider the example where $m = 3$, $n_1 = 5$ and $n = 10$. Then, Figures 6 and 7 provide the $(19, 22, 25, 28)$ -admissible path partitions K_{ρ_1} and K_{ρ_2} in the graph $G(19, 22, 25, 28)$. See Notation 34 for the definitions of ρ_1 and ρ_2 .



FIGURE 6. $(19, 22, 25, 28)$ -admissible path partition K_{ρ_1} in $G(19, 22, 25, 28)$.FIGURE 7. $(19, 22, 25, 28)$ -admissible path partition K_{ρ_2} in $G(19, 22, 25, 28)$.

Case 3. Suppose $a_k = 2mk + 1$, for all $1 \leq k \leq n_0$. We define the labels of $(a_1, a_2, \dots, a_{n_0})$ -admissible paths in $G(a_1, a_2, \dots, a_{n_0})$ as follows: for all $1 \leq i \leq m$, we let

$$\begin{aligned}
 x_{1,i,2j-1} &= 2m(j-1) + i, & \text{for } 1 \leq j \leq \mathfrak{n}_1^+, \\
 \frac{1}{2}S - x_{1,i,2j-1} &= m(n-2j+2) - i + 1, & \text{for } 1 \leq j \leq \mathfrak{n}_1^+, \\
 x_{1,i,2j} &= 2mj - i + 1, & \text{for } 1 \leq j \leq \mathfrak{n}_1, \\
 \frac{1}{2}S - x_{1,i,2j} &= m(n-2j) + i, & \text{for } 1 \leq j \leq \mathfrak{n}_1, \\
 x_{2,i,2j-1} &= 2mj - i + 1, & \text{for } 1 \leq j \leq \mathfrak{n}_1^+, \\
 \frac{1}{2}S - x_{2,i,2j-1} &= m(n-2j) + i, & \text{for } 1 \leq j \leq \mathfrak{n}_1^+, \\
 x_{2,i,2j} &= 2m(j-1) + i, & \text{for } 1 \leq j \leq \mathfrak{n}_1, \text{ and} \\
 \frac{1}{2}S - x_{2,i,2j} &= m(n-2j+2) - i + 1, & \text{for } 1 \leq j \leq \mathfrak{n}_1.
 \end{aligned}$$

Let $\rho : \{1, 2, \dots, m\} \rightarrow \{1, 2\}$. Let $P_{\rho,i}$ be the path on the sequence of vertices $(\{x_{\rho(i),i,j}, \frac{1}{2}S - x_{\rho(i),i,j}\} : 1 \leq j \leq n_1)$ with vertex label sequence $(x_{\rho(i),i,j} : 1 \leq j \leq n_1)$. The sets of labels

$$\begin{aligned}
 &\{x_{1,i,2j-1} : 1 \leq j \leq \mathfrak{n}_1^+\} \cup \{\frac{1}{2}S - x_{1,i,2j} : 1 \leq j \leq \mathfrak{n}_1\} \text{ and} \\
 &\{x_{2,i,2j} : 1 \leq j \leq \mathfrak{n}_1\} \cup \{\frac{1}{2}S - x_{2,i,2j-1} : 1 \leq j \leq \mathfrak{n}_1^+\}
 \end{aligned}$$

FIGURE 8. $(7, 13, 19, 25)$ -admissible path partition K_{ρ_1} in $G(7, 13, 19, 25)$.FIGURE 9. $(7, 13, 19, 25)$ -admissible path partition K_{ρ_2} in $G(7, 13, 19, 25)$.

are the set $\{2mk + i : 0 \leq k \leq n_1 - 1\}$. Also, the sets of labels

$$\begin{aligned} & \left\{ \frac{1}{2}S - x_{1,i,2j-1} : 1 \leq j \leq \#_1^+ \right\} \cup \{x_{1,i,2j} : 1 \leq j \leq \#_1\} \text{ and} \\ & \left\{ \frac{1}{2}S - x_{2,i,2j} : 1 \leq j \leq \#_1\right\} \cup \{x_{2,i,2j-1} : 1 \leq j \leq \#_1^+\} \end{aligned}$$

are the set $\{2mk - i + 1 : 1 \leq k \leq n_1\}$. Hence for $1 \leq i \leq m$ and $1 \leq j \leq n_1$, the vertices $\{x_{\rho(i),i,j}, \frac{1}{2}S - x_{\rho(i),i,j}\}$ are distinct in the graph $G(a_1, a_2, \dots, a_{n_0})$.

Furthermore, for $1 \leq i \leq m$, we have

$$\begin{aligned} x_{\rho(i),i,2j-1} + x_{\rho(i),i,2j} &= (2m(j-1) + i) + (2mj - i + 1) \\ &= 2m(2j - 1) + 1 = a_{2j-1}, & \text{for } 1 \leq j \leq \#_1, \text{ and} \\ x_{\rho(i),i,2j} + x_{\rho(i),i,2j+1} &= x_{\rho(i),i,2j} + x_{\rho(i),i,2j-1} + 2m \\ &= 2m(2j) + 1 = a_{2j}, & \text{for } 1 \leq j \leq \#_1^-. \end{aligned}$$

Thus, $P_{\rho,i}$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in $G(a_1, a_2, \dots, a_{n_0})$.

We consider the example where $m = 3$, $n_1 = 5$ and $n = 10$. Then, Figures 8 and 9 provide the $(7, 13, 19, 25)$ -admissible path partitions K_{ρ_1} and K_{ρ_2} in the graph $G(7, 13, 19, 25)$. See Notation 34 for the definitions of ρ_1 and ρ_2 .

Therefore in each of Cases 1, 2 and 3, $K_\rho = \{P_{\rho,i} : 1 \leq i \leq m\}$ is a $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition of $G(a_1, a_2, \dots, a_{n_0})$. Since $(a_1, a_2, \dots, a_{n_0})$ is not a palindrome, by Lemma 22, for each ρ , there exist $(m-1)!$ Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X of $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X) = K_\rho$. Hence, there exist at least $2^m(m-1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X of $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X)$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition of $G(a_1, a_2, \dots, a_{n_0})$. \square

Proposition 36. *Let $m \geq 3$ be an odd integer. Let $n \geq 6$ be an even integer such that $n_1 = n/2$ is odd. Let $n_0 = n_1 - 1$ and $S = 2(mn + 1)$. Let $(a_1, a_2, \dots, a_{n_0})$ be one of the following three sequences:*

- (1) $a_k = (2m - 1)n_1 + 2k + 1$, for all $1 \leq k \leq n_0$,
- (2) $a_k = mn_1 + 2mk + 1$, for all $1 \leq k \leq n_0$, or
- (3) $a_k = 4mk + 1$, for all $1 \leq k \leq n_0$.

Then, $G(a_1, a_2, \dots, a_{n_0})$ has at least 2^m distinct $(a_1, a_2, \dots, a_{n_0})$ -admissible path partitions and thus, for each distinct $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition K of $G(a_1, a_2, \dots, a_{n_0})$, there exist $(m - 1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X) = K$.

Proof. We consider each sequence $(a_1, a_2, \dots, a_{n_0})$ individually. The proof of the proposition for each sequence is similar.

Case 1. Suppose $a_k = (2m - 1)n_1 + 2k + 1$, for all $1 \leq k \leq n_0$. We define the labels of $(a_1, a_2, \dots, a_{n_0})$ -admissible paths in $G(a_1, a_2, \dots, a_{n_0})$ as follows: for all $1 \leq i \leq m$, we let

$$\begin{aligned}
 x_{1,i,2j-1} &= n_1(i-1) + 2j - 1, & \text{for } 1 \leq j \leq \#\mathfrak{r}_1^+, \\
 \frac{1}{2}S - x_{1,i,2j-1} &= mn - n_1(i-1) - 2j + 2, & \text{for } 1 \leq j \leq \#\mathfrak{r}_1^+, \\
 x_{1,i,2j} &= mn - n_1i + 2j, & \text{for } 1 \leq j \leq \#\mathfrak{r}_1, \\
 \frac{1}{2}S - x_{1,i,2j} &= n_1i - 2j + 1, & \text{for } 1 \leq j \leq \#\mathfrak{r}_1, \\
 x_{2,i,2j-1} &= mn - n_1i + 2j - 1, & \text{for } 1 \leq j \leq \#\mathfrak{r}_1^+, \\
 \frac{1}{2}S - x_{2,i,2j-1} &= n_1i - 2j + 2, & \text{for } 1 \leq j \leq \#\mathfrak{r}_1^+, \\
 x_{2,i,2j} &= n_1(i-1) + 2j, & \text{for } 1 \leq j \leq \#\mathfrak{r}_1, \text{ and} \\
 \frac{1}{2}S - x_{2,i,2j} &= mn - n_1(i-1) - 2j + 1, & \text{for } 1 \leq j \leq \#\mathfrak{r}_1.
 \end{aligned}$$

Let $\rho : \{1, 2, \dots, m\} \rightarrow \{1, 2\}$. Let $P_{\rho,i}$ be the path on the sequence of vertices $(\{x_{\rho(i),i,j}, \frac{1}{2}S - x_{\rho(i),i,j}\} : 1 \leq j \leq n_1)$ with vertex label sequence $(x_{\rho(i),i,j} : 1 \leq j \leq n_1)$. The sets of labels

$$\begin{aligned}
 &\{x_{1,i,2j-1} : 1 \leq j \leq \#\mathfrak{r}_1^+\} \cup \{\frac{1}{2}S - x_{1,i,2j} : 1 \leq j \leq \#\mathfrak{r}_1\} \text{ and} \\
 &\{x_{2,i,2j} : 1 \leq j \leq \#\mathfrak{r}_1\} \cup \{\frac{1}{2}S - x_{2,i,2j-1} : 1 \leq j \leq \#\mathfrak{r}_1^+\}
 \end{aligned}$$



are the set $\{n_1(i-1) + k : 1 \leq k \leq n_1\}$. Hence for $1 \leq i \leq m$ and $1 \leq j \leq n_1$, the vertices $\{x_{\rho(i),i,j}, \frac{1}{2}S - x_{\rho(i),i,j}\}$ are distinct in the graph $G(a_1, a_2, \dots, a_{n_0})$.

Furthermore, for $1 \leq i \leq m$, we have

$$\begin{aligned}
 x_{\rho(i),i,2j-1} + x_{\rho(i),i,2j} &= (n_1(i-1) + 2j - 1) + (mn - n_1i + 2j) \\
 &= (2m - 1)n_1 + 2(2j - 1) + 1 = a_{2j-1}, \quad \text{for } 1 \leq j \leq \#\mathfrak{r}_1, \text{ and} \\
 x_{\rho(i),i,2j} + x_{\rho(i),i,2j+1} &= x_{\rho(i),i,2j} + x_{\rho(i),i,2j-1} + 2 \\
 &= (2m - 1)n_1 + 2(2j) + 1 = a_{2j}, \quad \text{for } 1 \leq j \leq \#\mathfrak{r}_1^-.
 \end{aligned}$$

Thus, $P_{\rho,i}$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in $G(a_1, a_2, \dots, a_{n_0})$.

We consider the example where $m = 3$, $n_1 = 5$ and $n = 10$. Then, Figures 10 and 11 provide the $(28, 30, 32, 34)$ -admissible path partitions K_{ρ_1} and K_{ρ_2} in the graph $G(28, 30, 32, 34)$. See Notation 34 for the definitions of ρ_1 and ρ_2 .

FIGURE 10. $(28, 30, 32, 34)$ -admissible path partition K_{ρ_1} in $G(28, 30, 32, 34)$.FIGURE 11. $(28, 30, 32, 34)$ -admissible path partition K_{ρ_2} in $G(28, 30, 32, 34)$.



Case 2. Suppose $a_k = mn_1 + 2mk + 1$, for all $1 \leq k \leq n_0$. We define the labels of $(a_1, a_2, \dots, a_{n_0})$ -admissible paths in $G(a_1, a_2, \dots, a_{n_0})$ as follows: for all $1 \leq i \leq m$, we let

$$\begin{aligned}
 x_{1,i,2j-1} &= m(2j-2) + i, & \text{for } 1 \leq j \leq \mathfrak{n}_1^+, \\
 \frac{1}{2}S - x_{1,i,2j-1} &= m(n-2j+2) - i + 1, & \text{for } 1 \leq j \leq \mathfrak{n}_1^+, \\
 x_{1,i,2j} &= m(n_1+2j) - i + 1, & \text{for } 1 \leq j \leq \mathfrak{n}_1, \\
 \frac{1}{2}S - x_{1,i,2j} &= m(n_1-2j) + i, & \text{for } 1 \leq j \leq \mathfrak{n}_1, \\
 x_{2,i,2j-1} &= m(n_1+2j-1) - i + 1, & \text{for } 1 \leq j \leq \mathfrak{n}_1^+, \\
 \frac{1}{2}S - x_{2,i,2j-1} &= m(n_1-2j+1) + i, & \text{for } 1 \leq j \leq \mathfrak{n}_1^+, \\
 x_{2,i,2j} &= m(2j-1) + i, & \text{for } 1 \leq j \leq \mathfrak{n}_1, \text{ and} \\
 \frac{1}{2}S - x_{2,i,2j} &= m(n-2j+1) - i + 1, & \text{for } 1 \leq j \leq \mathfrak{n}_1.
 \end{aligned}$$

Let $\rho : \{1, 2, \dots, m\} \rightarrow \{1, 2\}$. Let $P_{\rho,i}$ be the path on the sequence of vertices $(\{x_{\rho(i),i,j}, \frac{1}{2}S - x_{\rho(i),i,j}\} : 1 \leq j \leq n_1)$ with vertex label sequence $(x_{\rho(i),i,j} : 1 \leq j \leq n_1)$. The sets of labels

$$\begin{aligned}
 &\{x_{1,i,2j-1} : 1 \leq j \leq \mathfrak{n}_1^+\} \cup \{\frac{1}{2}S - x_{1,i,2j} : 1 \leq j \leq \mathfrak{n}_1\} \text{ and} \\
 &\{x_{2,i,2j} : 1 \leq j \leq \mathfrak{n}_1\} \cup \{\frac{1}{2}S - x_{2,i,2j-1} : 1 \leq j \leq \mathfrak{n}_1^+\}
 \end{aligned}$$

are the set $\{mk+i : 0 \leq k \leq n_1-1\}$. Hence for $1 \leq i \leq m$ and $1 \leq j \leq n_1$, the vertices $\{x_{\rho(i),i,j}, \frac{1}{2}S - x_{\rho(i),i,j}\}$ are distinct in the graph $G(a_1, a_2, \dots, a_{n_0})$.

FIGURE 12. $(22, 28, 34, 40)$ -admissible path partition K_{ρ_1} in $G(22, 28, 34, 40)$.FIGURE 13. $(22, 28, 34, 40)$ -admissible path partition K_{ρ_2} in $G(22, 28, 34, 40)$.

Furthermore, for $1 \leq i \leq m$, we have



$$\begin{aligned}
 x_{\rho(i), i, 2j-1} + x_{\rho(i), i, 2j} &= mn_1 + 2m(2j-1) + 1 = a_{2j-1}, & \text{for } 1 \leq j \leq \#_1, \text{ and} \\
 x_{\rho(i), i, 2j} + x_{\rho(i), i, 2j+1} &= x_{\rho(i), i, 2j} + x_{\rho(i), i, 2j-1} + 2m = a_{2j}, & \text{for } 1 \leq j \leq \#_1^-.
 \end{aligned}$$

Thus, $P_{\rho, i}$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in $G(a_1, a_2, \dots, a_{n_0})$.

We consider the example where $m = 3$, $n_1 = 5$ and $n = 10$. Then, Figures 12 and 13 provide the $(22, 28, 34, 40)$ -admissible path partitions K_{ρ_1} and K_{ρ_2} in the graph $G(22, 28, 34, 40)$. See Notation 34 for the definitions of ρ_1 and ρ_2 .

Case 3. Suppose $a_k = 4mk + 1$, for all $1 \leq k \leq n_0$. We define the labels of $(a_1, a_2, \dots, a_{n_0})$ -admissible paths in $G(a_1, a_2, \dots, a_{n_0})$ as follows: for all $1 \leq i \leq m$, we let

$$\begin{aligned}
 x_{1, i, 2j-1} &= 4m(j-1) + i, & \text{for } 1 \leq j \leq \#_1^+, \\
 \frac{1}{2}S - x_{1, i, 2j-1} &= m(n-4j+4) - i + 1, & \text{for } 1 \leq j \leq \#_1^+, \\
 x_{1, i, 2j} &= 4mj - i + 1, & \text{for } 1 \leq j \leq \#_1, \\
 \frac{1}{2}S - x_{1, i, 2j} &= m(n-4j) + i, & \text{for } 1 \leq j \leq \#_1, \\
 x_{2, i, 2j-1} &= m(4j-2) - i + 1, & \text{for } 1 \leq j \leq \#_1^+, \\
 \frac{1}{2}S - x_{2, i, 2j-1} &= m(n-4j+2) + i, & \text{for } 1 \leq j \leq \#_1^+, \\
 x_{2, i, 2j} &= m(4j-2) + i, & \text{for } 1 \leq j \leq \#_1, \text{ and} \\
 \frac{1}{2}S - x_{2, i, 2j} &= m(n-4j+2) - i + 1, & \text{for } 1 \leq j \leq \#_1.
 \end{aligned}$$

FIGURE 14. $(13, 25, 37, 49)$ -admissible path partition K_{ρ_1} in $G(13, 25, 37, 49)$.FIGURE 15. $(13, 25, 37, 49)$ -admissible path partition K_{ρ_2} in $G(13, 25, 37, 49)$.

Let $\rho : \{1, 2, \dots, m\} \rightarrow \{1, 2\}$. Let $P_{\rho, i}$ be the path on the sequence of vertices $(\{x_{\rho(i), i, j}, \frac{1}{2}S - x_{\rho(i), i, j}\} : 1 \leq j \leq n_1)$ with vertex label sequence $(x_{\rho(i), i, j} : 1 \leq j \leq n_1)$. The sets of labels

$$\{x_{1, i, 2j-1} : 1 \leq j \leq \#\mathfrak{r}_1^+\} \cup \{\frac{1}{2}S - x_{1, i, 2j} : 1 \leq j \leq \#\mathfrak{r}_1\} \text{ and}$$

$$\{x_{2, i, 2j} : 1 \leq j \leq \#\mathfrak{r}_1\} \cup \{\frac{1}{2}S - x_{2, i, 2j-1} : 1 \leq j \leq \#\mathfrak{r}_1^+\}$$

are the set $\{2mk + i : 0 \leq k \leq n_1 - 1\}$. Also, the sets of labels

$$\{\frac{1}{2}S - x_{1, i, 2j-1} : 1 \leq j \leq \#\mathfrak{r}_1^+\} \cup \{x_{1, i, 2j} : 1 \leq j \leq \#\mathfrak{r}_1\} \text{ and}$$

$$\{x_{2, i, 2j} : 1 \leq j \leq \#\mathfrak{r}_1\} \cup \{\frac{1}{2}S - x_{2, i, 2j-1} : 1 \leq j \leq \#\mathfrak{r}_1^+\}$$

are the set $\{2mk - i + 1 : 1 \leq k \leq n_1\}$. Hence for $1 \leq i \leq m$ and $1 \leq j \leq n_1$, the vertices $\{x_{\rho(i), i, j}, \frac{1}{2}S - x_{\rho(i), i, j}\}$ are distinct in the graph $G(a_1, a_2, \dots, a_{n_0})$.

Furthermore, for $1 \leq i \leq m$, we have

$$x_{\rho(i), i, 2j-1} + x_{\rho(i), i, 2j} = 4m(2j-1) + 1 = a_{2j-1}, \quad \text{for } 1 \leq j \leq \#\mathfrak{r}_1, \text{ and}$$

$$x_{\rho(i), i, 2j} + x_{\rho(i), i, 2j+1} = x_{\rho(i), i, 2j} + x_{\rho(i), i, 2j-1} + 4m = a_{2j}, \quad \text{for } 1 \leq j \leq \#\mathfrak{r}_1^-.$$

Thus, $P_{\rho, i}$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in $G(a_1, a_2, \dots, a_{n_0})$.

We consider the example where $m = 3$, $n_1 = 5$ and $n = 10$. Then, Figures 14 and 15 provide the $(13, 25, 37, 49)$ -admissible path partitions K_{ρ_1} and K_{ρ_2} in the graph $G(13, 25, 37, 49)$. See Notation 34 for the definitions of ρ_1 and ρ_2 .

Therefore in each of Cases 1, 2 and 3, $K_\rho = \{P_{\rho, i} : 1 \leq i \leq m\}$ is a $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition of $G(a_1, a_2, \dots, a_{n_0})$. Since $(a_1, a_2, \dots, a_{n_0})$ is not a palindrome, by Lemma 22, for each ρ , there

exist $(m-1)!$ Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X of $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X) = K_\rho$. Hence, there exist at least $2^m(m-1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X of $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X)$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition of $G(a_1, a_2, \dots, a_{n_0})$. \square

Proposition 37. *Let $m \geq 3$ be an odd integer. Let $n \geq 6$ be an even integer such that $n_1 = n/2$ is even. Let $n_0 = n_1 - 1$, $\#_1 = \lfloor n_1/2 \rfloor$, and $S = 2(mn + 1)$. Let $(a_1, a_2, \dots, a_{n_0})$ be one of the following four sequences:*

- (1) $a_{2k-1} = mn$, for all $1 \leq k \leq \#_1$, and $a_{2k} = mn + 2$, for all $1 \leq k \leq \#_1 - 1$,
- (2) $a_{2k-1} = mn - 1$, for all $1 \leq k \leq \#_1$, and $a_{2k} = mn + 3$, for all $1 \leq k \leq \#_1 - 1$,
- (3) $a_{2k-1} = m(n-1) + 1$, for all $1 \leq k \leq \#_1$, and $a_{2k} = m(n+1) + 1$, for all $1 \leq k \leq \#_1 - 1$, or
- (4) $a_{2k-1} = m(n-2) + 1$, for all $1 \leq k \leq \#_1$, and $a_{2k} = m(n+2) + 1$, for all $1 \leq k \leq \#_1 - 1$.

Then, $(a_1, a_2, \dots, a_{n_0})$ is a palindrome and there exists an $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition K of $G(a_1, a_2, \dots, a_{n_0})$. Furthermore, there exist $2^{m-1}(m-1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X) = K$.

Proof. We consider each sequence $(a_1, a_2, \dots, a_{n_0})$ individually. The proof of the proposition for each sequence is similar.

Case 1. Suppose $a_{2k-1} = mn$, for all $1 \leq k \leq \#_1$. and $a_{2k} = mn + 2$, for all $1 \leq k \leq \#_1^-$. We define the labels of $(a_1, a_2, \dots, a_{n_0})$ -admissible paths in $G(a_1, a_2, \dots, a_{n_0})$ as follows: for all $1 \leq i \leq m$, we let

$$\begin{aligned} x_{i,2j-1} &= n_1(i-1) + 2j - 1, & \text{for } 1 \leq j \leq \#_1, \\ \frac{1}{2}S - x_{i,2j-1} &= n_1(2m-i+1) - 2j + 2, & \text{for } 1 \leq j \leq \#_1, \\ x_{i,2j} &= n_1(2m-i+1) - 2j + 1, & \text{for } 1 \leq j \leq \#_1, \text{ and} \\ \frac{1}{2}S - x_{i,2j} &= n_1(i-1) + 2j, & \text{for } 1 \leq j \leq \#_1. \end{aligned}$$

Let P_i be the path on the sequence of vertices $(\{x_{i,j}, \frac{1}{2}S - x_{i,j}\} : 1 \leq j \leq n_1)$ with vertex label sequence $(x_{i,j} : 1 \leq j \leq n_1)$. The set of labels

$$\{x_{i,2j-1} : 1 \leq j \leq \#_1\} \cup \{\frac{1}{2}S - x_{i,2j} : 1 \leq j \leq \#_1\}$$

is the set $\{n_1(i-1) + k : 1 \leq k \leq n_1\}$. Hence for $1 \leq i \leq m$ and $1 \leq j \leq n_1$, the vertices $\{x_{i,j}, \frac{1}{2}S - x_{i,j}\}$ are distinct in the graph $G(a_1, a_2, \dots, a_{n_0})$.


Furthermore, for $1 \leq i \leq m$, we have

$$\begin{aligned} x_{i,2j-1} + x_{i,2j} &= (n_1(i-1) + 2j - 1) + (n_1(2m-i+1) - 2j + 1) \\ &= mn = a_{2j-1}, \quad \text{for } 1 \leq j \leq \#_1, \text{ and} \\ x_{i,2j} + x_{i,2j+1} &= x_{i,2j} + x_{i,2j-1} + 2 \\ &= mn + 2 = a_{2j}, \quad \text{for } 1 \leq j \leq \#_1. \end{aligned}$$

Thus, P_i is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in $G(a_1, a_2, \dots, a_{n_0})$. Also, we observe that $a_{2k-1} = a_{n_1-2k+1} = mn$ for all $1 \leq k \leq \#_1$, and $a_{2k} = a_{n_1-2k} = mn + 2$ for all $1 \leq k \leq \#_1^-$. Hence, $(a_1, a_2, \dots, a_{n_0})$ is a palindrome.

We consider the example where $m = 3$, $n_1 = 6$ and $n = 12$. Then, Figure 16 provides the $(36, 38, 36, 38, 36)$ -admissible path partition K in the graph $G(36, 38, 36, 38, 36)$.

Case 2. Suppose $a_{2k-1} = mn - 1$ for all $1 \leq k \leq \#_1$, and $a_{2k} = mn + 3$ for all $1 \leq k \leq \#_1^-$. Since $m \geq 3$ is odd, we let $m = 2m_1 + 1$ for some positive integer m_1 . We define the labels of $(a_1, a_2, \dots, a_{n_0})$ -admissible

FIGURE 16. $(36, 38, 36, 38, 36)$ -admissible path partition K in $G(36, 38, 36, 38, 36)$.

paths in $G(a_1, a_2, \dots, a_{n_0})$ as follows: for all $1 \leq i \leq m_1$, we let

$$\begin{aligned}
 x_{2i-1,2j-1} &= n(i-1) + 4j - 3, & \text{for } 1 \leq j \leq \#_1, \\
 \frac{1}{2}S - x_{2i-1,2j-1} &= n(m-i+1) - 4j + 4, & \text{for } 1 \leq j \leq \#_1, \\
 x_{2i-1,2j} &= n(m-i+1) - 4j + 2, & \text{for } 1 \leq j \leq \#_1, \\
 \frac{1}{2}S - x_{2i-1,2j} &= n(i-1) + 4j - 1, & \text{for } 1 \leq j \leq \#_1, \\
 x_{2i,2j-1} &= n(i-1) + 4j - 2, & \text{for } 1 \leq j \leq \#_1, \\
 \frac{1}{2}S - x_{2i,2j-1} &= n(m-i+1) - 4j + 3, & \text{for } 1 \leq j \leq \#_1, \\
 x_{2i,2j} &= n(m-i+1) - 4j + 1, & \text{for } 1 \leq j \leq \#_1, \text{ and} \\
 \frac{1}{2}S - x_{2i,2j} &= n(i-1) + 4j, & \text{for } 1 \leq j \leq \#_1.
 \end{aligned}$$

Also, we let

$$\begin{aligned}
 x_{m,2j-1} &= m_1n + 4j - 3, & \text{for } 1 \leq j \leq \#_1, \\
 \frac{1}{2}S - x_{m,2j-1} &= (m_1+1)n - 4j + 4, & \text{for } 1 \leq j \leq \#_1, \\
 x_{m,2j} &= (m_1+1)n - 4j + 2, & \text{for } 1 \leq j \leq \#_1, \text{ and} \\
 \frac{1}{2}S - x_{m,2j} &= m_1n + 4j - 1, & \text{for } 1 \leq j \leq \#_1.
 \end{aligned}$$


Let P_i be the path on the sequence of vertices $(\{x_{i,j}, \frac{1}{2}S - x_{i,j}\} : 1 \leq j \leq n_1)$ with vertex label sequence $(x_{i,j} : 1 \leq j \leq n_1)$. The set of labels

$$\begin{aligned}
 &\{x_{2i-1,2j-1} : 1 \leq j \leq \#_1\} \cup \{\frac{1}{2}S - x_{2i-1,2j} : 1 \leq j \leq \#_1\} \\
 &\quad \cup \{\frac{1}{2}S - x_{2i,2j-1} : 1 \leq j \leq \#_1\} \cup \{x_{2i,2j} : 1 \leq j \leq \#_1\}
 \end{aligned}$$

is the set $\{n(i-1) + k : 1 \leq k \leq n\}$. Let $n_2 = \#_1$. Also, the set of labels

$$\begin{aligned}
 &\{x_{m,2j-1} : 1 \leq j \leq \#_2^+\} \cup \{\frac{1}{2}S - x_{m,2j} : 1 \leq j \leq \#_2\} \\
 &\quad \cup \{\frac{1}{2}S - x_{m,2j-1} : \#_2^+ < j \leq n_2\} \cup \{x_{m,2j} : \#_2 \leq j \leq n_2\}
 \end{aligned}$$

is the set $\{m_1n + k : 1 \leq k \leq n_1\}$. Hence for $1 \leq i \leq m$ and $1 \leq j \leq n_1$, the vertices $\{x_{i,j}, \frac{1}{2}S - x_{i,j}\}$ are distinct in the graph $G(a_1, a_2, \dots, a_{n_0})$.

FIGURE 17. $(35, 39, 35, 39, 35)$ -admissible path partition K in $G(35, 39, 35, 39, 35)$.

Furthermore for $1 \leq i \leq m_1$, we have

$$\begin{aligned}
 x_{2i-1,2j-1} + x_{2i-1,2j} &= (n(i-1) + 4j - 3) + (n(m-i+1) - 4j + 2) \\
 &= mn - 1 = a_{2j-1}, \quad \text{for } 1 \leq j \leq \#_1, \\
 x_{2i-1,2j} + x_{2i-1,2j+1} &= x_{2i-1,2j} + x_{2i-1,2j-1} + 4 \\
 &= mn + 3 = a_{2j}, \quad \text{for } 1 \leq j \leq \#_1, \\
 x_{2i,2j-1} + x_{2i,2j} &= (n(i-1) + 4j - 2) + (n(m-i+1) - 4j + 1) \\
 &= mn - 1 = a_{2j-1}, \quad \text{for } 1 \leq j \leq \#_1, \text{ and} \\
 x_{2i,2j} + x_{2i,2j+1} &= x_{2i,2j} + x_{2i,2j-1} + 4 \\
 &= mn + 3 = a_{2j}, \quad \text{for } 1 \leq j \leq \#_1.
 \end{aligned}$$

In addition, we have

$$\begin{aligned}
 x_{m,2j-1} + x_{m,2j} &= (m_1n + 4j - 3) + ((m_1 + 1)n - 4j + 2) \\
 &= mn - 1 = a_{2j-1}, \quad \text{for } 1 \leq j \leq \#_1, \text{ and} \\
 x_{m,2j} + x_{m,2j+1} &= x_{m,2j} + x_{m,2j-1} + 4 \\
 &= mn + 3 = a_{2j}, \quad \text{for } 1 \leq j \leq \#_1.
 \end{aligned}$$

Thus, P_i is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in $G(a_1, a_2, \dots, a_{n_0})$. Also, we observe that $a_{2k-1} = a_{n_1-2k+1} = mn$ for all $1 \leq k \leq \#_1$, and $a_{2k} = a_{n_1-2k} = mn + 2$ for all $1 \leq k \leq \#_1^-$. Hence, $(a_1, a_2, \dots, a_{n_0})$ is a palindrome.


We consider the example where $m = 3$, $n_1 = 6$ and $n = 12$. Then, Figure 17 provides the $(35, 39, 35, 39, 35)$ -admissible path partition K in the graph $G(35, 39, 35, 39, 35)$.

Case 3. Suppose $a_{2k-1} = m(n-1) + 1$ for all $1 \leq k \leq \#_1$, and $a_{2k} = m(n+1) + 1$ for all $1 \leq k \leq \#_1^-$. We define the labels of $(a_1, a_2, \dots, a_{n_0})$ -admissible paths in $G(a_1, a_2, \dots, a_{n_0})$ as follows: for all $1 \leq i \leq m$, we let

$$\begin{aligned}
 x_{i,2j-1} &= m(2j-2) + i, & \text{for } 1 \leq j \leq \#_1, \\
 \frac{1}{2}S - x_{i,2j-1} &= m(n-2j+2) - i + 1, & \text{for } 1 \leq j \leq \#_1, \\
 x_{i,2j} &= m(n-2j+1) - i + 1, & \text{for } 1 \leq j \leq \#_1, \text{ and} \\
 \frac{1}{2}S - x_{i,2j} &= m(2j-1) + i, & \text{for } 1 \leq j \leq \#_1.
 \end{aligned}$$

Let P_i be the path on the sequence of vertices $(\{x_{i,j}, \frac{1}{2}S - x_{i,j}\} : 1 \leq j \leq n_1)$ with vertex label sequence $(x_{i,j} : 1 \leq j \leq n_1)$. The set of labels

$$\{x_{i,2j-1} : 1 \leq j \leq \#_1\} \cup \{\frac{1}{2}S - x_{i,2j} : 1 \leq j \leq \#_1\}$$

FIGURE 18. $(34, 40, 34, 40, 34)$ -admissible path partition K in $G(34, 40, 34, 40, 34)$.

is the set $\{mk + i : 0 \leq k \leq n_1 - 1\}$. Hence for $1 \leq i \leq m$ and $1 \leq j \leq n_1$, the vertices $\{x_{i,j}, \frac{1}{2}S - x_{i,j}\}$ are distinct in the graph $G(a_1, a_2, \dots, a_{n_0})$.

Furthermore for $1 \leq i \leq m$, we have

$$\begin{aligned} x_{i,2j-1} + x_{i,2j} &= (m(2j-2) + i) + (m(n-2j+1) - i + 1) \\ &= m(n-1) + 1 = a_{2j-1}, && \text{for } 1 \leq j \leq \#_1, \text{ and} \\ x_{i,2j} + x_{i,2j+1} &= x_{i,2j} + x_{i,2j-1} + 2m \\ &= m(n+1) + 1 = a_{2j}, && \text{for } 1 \leq j \leq \#_1. \end{aligned}$$

Thus, P_i is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in $G(a_1, a_2, \dots, a_{n_0})$. Also, we observe that $a_{2k-1} = a_{n_1-2k+1} = m(n-1) + 1$, for all $1 \leq k \leq \#_1$, and $a_{2k} = a_{n_1-2k} = m(n+1) + 1$, for all $1 \leq k \leq \#_1^-$. Hence, $(a_1, a_2, \dots, a_{n_0})$ is a palindrome.

We consider the example where $m = 3$, $n_1 = 6$ and $n = 12$. Then, Figure 18 provides the $(34, 40, 34, 40, 34)$ -admissible path partition K in the graph $G(34, 40, 34, 40, 34)$.

Case 4. Suppose $a_{2k-1} = m(n-2) + 1$ for all $1 \leq k \leq \#_1$, and $a_{2k} = m(n+2) + 1$ for all $1 \leq k \leq \#_1^-$. We define the labels of $(a_1, a_2, \dots, a_{n_0})$ -admissible paths in $G(a_1, a_2, \dots, a_{n_0})$ as follows: for all $1 \leq i \leq m$, we let

$$\begin{aligned} x_{i,2j-1} &= m(4j-4) + i, && \text{for } 1 \leq j \leq \#_1, \\ \frac{1}{2}S - x_{i,2j-1} &= m(n-4j+4) - i + 1, && \text{for } 1 \leq j \leq \#_1, \\ x_{i,2j} &= m(n-4j+2) - i + 1, && \text{for } 1 \leq j \leq \#_1, \text{ and} \\ \frac{1}{2}S - x_{i,2j} &= m(4j-2) + i, && \text{for } 1 \leq j \leq \#_1. \end{aligned}$$


Let P_i be the path on the sequence of vertices $(\{x_{i,j}, \frac{1}{2}S - x_{i,j}\} : 1 \leq j \leq n_1)$ with vertex label sequence $(x_{i,j} : 1 \leq j \leq n_1)$. The set of labels

$$\{x_{i,2j-1} : 1 \leq j \leq \#_1\} \cup \{\frac{1}{2}S - x_{i,2j} : 1 \leq j \leq \#_1\}$$

is the set $\{2mk + i : 0 \leq k \leq n_1 - 1\}$. Also, the set of labels

$$\{\frac{1}{2}S - x_{i,2j-1} : 1 \leq j \leq \#_1^+\} \cup \{x_{i,2j} : 1 \leq j \leq \#_1\}$$

is the set $\{2mk - i + 1 : 1 \leq k \leq n_1\}$. Hence for $1 \leq i \leq m$ and $1 \leq j \leq n_1$, the vertices $\{x_{i,j}, \frac{1}{2}S - x_{i,j}\}$ are distinct in the graph $G(a_1, a_2, \dots, a_{n_0})$.

FIGURE 19. $(31, 43, 31, 43, 31)$ -admissible path partition K in $G(31, 43, 31, 43, 31)$.

Furthermore for $1 \leq i \leq m$, we have

$$\begin{aligned} x_{i,2j-1} + x_{i,2j} &= (m(4j-4) + i) + (m(n-4j+2) - i + 1) \\ &= m(n-2) + 1 = a_{2j-1}, & \text{for } 1 \leq j \leq \#_1, \text{ and} \\ x_{i,2j} + x_{i,2j+1} &= x_{i,2j} + x_{i,2j-1} + 4m \\ &= m(n+2) + 1 = a_{2j}, & \text{for } 1 \leq j \leq \#_1. \end{aligned}$$

Thus, P_i is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path in $G(a_1, a_2, \dots, a_{n_0})$. Also, we observe that $a_{2k-1} = a_{n_1-2k+1} = m(n-2) + 1$ for all $1 \leq k \leq \#_1$, and $a_{2k} = a_{n_1-2k} = m(n+2) + 1$ for all $1 \leq k \leq \#_1$. Hence, $(a_1, a_2, \dots, a_{n_0})$ is a palindrome.

We consider the example where $m = 3$, $n_1 = 6$ and $n = 12$. Then, Figure 19 provides the $(31, 43, 31, 43, 31)$ -admissible path partition K in the graph $G(31, 43, 31, 43, 31)$.

Therefore in each of Cases 1, 2, 3, and 4, $K = \{P_i : 1 \leq i \leq m\}$ is a $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition of $G(a_1, a_2, \dots, a_{n_0})$. Since $(a_1, a_2, \dots, a_{n_0})$ is a palindrome, by Lemma 22, there exist $2^{m-1}(m-1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X of $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X) = K$. Hence, there exist at least $2^{m-1}(m-1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X of $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X)$ is an $(a_1, a_2, \dots, a_{n_0})$ -admissible path partition of $G(a_1, a_2, \dots, a_{n_0})$. \square

Theorem 38. *Let $m \geq 3$ be an odd integer and let $n \geq 6$ be an even integer.*

- (1) *If $n \equiv 2 \pmod{4}$, then the minimum number of distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,n}$ is $(6 \cdot 2^m)(m-1)!$.*
- (2) *If $n \equiv 0 \pmod{4}$, then the minimum number of distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,n}$ is $(5 \cdot 2^m)(m-1)!$.*

Proof. Let $A_i = (a_{i,1}, a_{i,2}, \dots, a_{i,n_0})$ be the sequence where $a_{1,k} = mn - n_0 + k$, $a_{2,k} = m(n_1 + k) + 1$, and $a_{3,k} = 2mk + 1$, for all $1 \leq k \leq n_0$. Also, let $A_i = (a_{i,1}, a_{i,2}, \dots, a_{i,n_0})$ be the sequence where $a_{4,k} = (2m-1)n_1 + 2k + 1$, $a_{5,k} = mn_1 + 2mk + 1$, and $a_{6,k} = 4mk + 1$, for all $1 \leq k \leq n_0$. Furthermore, let $\rho : NAPP(m, n) \rightarrow NAPP(m, n)$ (see Definition 24 for the definition of $NAPP(m, n)$) and $\sigma : NAPP(m, n) \rightarrow NAPP(m, n)$ be functions defined by

$$\begin{aligned} \rho(a_1, a_2, \dots, a_{n_0-1}, a_{n_0}) &= (a_{n_0}, a_{n_0-1}, \dots, a_2, a_1) \quad \text{and} \\ \sigma(a_1, a_2, \dots, a_{n_0-1}, a_{n_0}) &= (S - a_1, S - a_2, \dots, S - a_{n_0-1}, S - a_{n_0}). \end{aligned}$$

Suppose $A, A' \in NAPP(m, n)$ are sequences such that A is path partition equivalent to A' . By Lemma 26, either $A' = A$, $A' = \rho(A)$, $A' = \sigma(A)$, or $A' = (\sigma\rho)(A)$. Of these four possibilities for A' , let A' be the

minimal element in the set $\{A, \rho(A), \sigma(A), (\sigma\rho)(A)\}$ under the lexicographic order. Let

$$MNAPP(m, n) = \left\{ A \in NAPP(m, n) : A \text{ is the minimal element in the set} \right. \\ \left. \{A, \rho(A), \sigma(A), (\sigma\rho)(A)\} \text{ under the lexicographic order} \right\}$$

be the minimal set of sequences $A \in NAPP(m, n)$ needed in order to count all Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings on $\mathcal{K}_{m,n}$ in which A is not a palindrome. Let \tilde{A}_1 and \tilde{A}_2 be two distinct sequences in $MNAPP(m, n)$. For $i = 1, 2$, let X_i be a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X_i)$ is an \tilde{A}_i -admissible path partition of $G(\tilde{A}_i)$. Then, \tilde{A}_2 is not any of the pairs $\tilde{A}_1, \rho(\tilde{A}_1), \sigma(\tilde{A}_1)$, or $(\sigma\rho)(\tilde{A}_1)$. By Lemma 26, \tilde{A}_1 is not path partition equivalent to \tilde{A}_2 . Thus, X_1 and X_2 are Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings on $\mathcal{K}_{m,n}$. Let $N(A)$ be the number of distinct A -admissible path partitions of $G(A)$.

We consider the cases $n \equiv 0 \pmod{4}$ and $n \equiv 2 \pmod{4}$ individually.

Case 1. Suppose $n \equiv 2 \pmod{4}$. Then, n_1 is odd. Let $A = (a_1, a_2, \dots, a_{n_0})$ be a sequence such that there is an A -admissible path partition of $G(A)$. For the purposes of contradiction, assume A is a palindrome. Since $x_{1, \mp n_1} + x_{1, \mp n_1 + 1} = a_{\mp n_1} = a_{\mp n_1 + 1} = x_{1, \mp n_1 + 1} + x_{1, \mp n_1 + 2}$, we have $x_{1, \mp n_1} = x_{1, \mp n_1 + 2}$. This is a contradiction. Therefore, A is not a palindrome.

By Lemma 22, there are $N(A) \cdot (m-1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X)$ is an A -admissible path partition of $G(A)$. Thus, the number of distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings on $\mathcal{K}_{m,n}$, denoted by $NKBL(m, n)$, is

$$NKBL(m, n) = \sum_{A \in MNAPP(m, n)} N(A) \cdot (m-1)!$$

By Proposition 35, we have $N(A_i) \geq 2^m$ for all $1 \leq i \leq 3$. By Proposition 36, we have $N(A_i) \geq 2^m$ for all $4 \leq i \leq 6$. Thus, the number of distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings on $\mathcal{K}_{m,n}$ satisfies

$$NKBL(m, n) = \sum_{A \in MNAPP(m, n)} N(A) \cdot (m-1)! \\ \geq \sum_{i=1}^6 N(A_i) \cdot (m-1)! \geq (6 \cdot 2^m) \cdot (m-1)!$$

Case 2. Suppose $n \equiv 0 \pmod{4}$. Then, n_1 is even.

Let $A_7 = (a_{7,1}, a_{7,2}, \dots, a_{7,n_0})$ be the sequence where $a_{7,2k-1} = mn$ for all $1 \leq k \leq \mp n_1$, and $a_{7,2k} = mn+2$ for all $1 \leq k \leq \mp n_1 - 1$. Let $A_8 = (a_{8,1}, a_{8,2}, \dots, a_{8,n_0})$ be the sequence where $a_{8,2k-1} = mn-1$ for all $1 \leq k \leq \mp n_1$, and $a_{8,2k} = mn+3$ for all $1 \leq k \leq \mp n_1 - 1$. Let $A_9 = (a_{9,1}, a_{9,2}, \dots, a_{9,n_0})$ be the sequence where $a_{9,2k-1} = m(n-1)+1$ for all $1 \leq k \leq \mp n_1$, and $a_{9,2k} = m(n+1) = 1$ for all $1 \leq k \leq \mp n_1 - 1$. Let $A_{10} = (a_{10,1}, a_{10,2}, \dots, a_{10,n_0})$ be the sequence where $a_{10,2k-1} = m(n-2)+1$ for all $1 \leq k \leq \mp n_1$, and $a_{10,2k} = m(n+2)+1$ for all $1 \leq k \leq \mp n_1 - 1$.

Suppose $A, A' \in PAPP(m, n)$ (see Definition 24 for the definition of $PAPP(m, n)$) are sequences such that A is path partition equivalent to A' . Define $\sigma : PAPP(m, n) \rightarrow PAPP(m, n)$ by

$$\sigma(a_1, a_2, \dots, a_{n_0-1}, a_{n_0}) = (S - a_1, S - a_2, \dots, S - a_{n_0-1}, S - a_{n_0}).$$

By Lemma 26, either $A' = A$, or $A' = \sigma(A)$. Of these two possibilities for A' , let A' be the minimal element in the set $\{A, \sigma(A)\}$ under the lexicographic order. Let

$$MPAPP(m, n) = \left\{ A \in PAPP(m, n) : A \text{ is the minimal element in the set} \right. \\ \left. \{A, \sigma(A)\} \text{ under the lexicographic order} \right\}$$

be the minimal set of sequences $A \in PAPP(m, n)$ needed in order to count all Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings on $\mathcal{K}_{m,n}$ in which A is a palindrome. Let \tilde{A}_1 and \tilde{A}_2 be two distinct sequences in $MPAPP(m, n)$. For $i = 1, 2$, let X_i be a C_4 -face-magic Klein bottle labeling on $\mathcal{K}_{m,n}$ such

that $\mathcal{L}(X_i)$ is an \tilde{A}_i -admissible path partition of $G(\tilde{A}_i)$. Then, \tilde{A}_2 is not any of the pairs \tilde{A}_1 , or $\sigma(\tilde{A}_1)$. By Lemma 26, \tilde{A}_1 is not path partition equivalent to \tilde{A}_2 . Thus X_1 and X_2 are Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings on $\mathcal{K}_{m,n}$. Let $N(A)$ be the number of distinct A -admissible path partitions of $G(A)$. On one hand, if A is not a palindrome, by Lemma 22, there are $N(A) \cdot (m-1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X)$ is an A -admissible path partition of $G(A)$. On the other hand, if A is a palindrome, by Lemma 22, there are $N(A) \cdot 2^{m-1}(m-1)!$ distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings X on $\mathcal{K}_{m,n}$ such that $\mathcal{L}(X)$ is an A -admissible path partition of $G(A)$. Thus, the number of distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings on $\mathcal{K}_{m,n}$, denoted by $NKBL(m, n)$, is

$$NKBL(m, n) = \sum_{A \in MNAPP(m, n)} N(A) \cdot (m-1)! + \sum_{A \in MPAPP(m, n)} N(A) \cdot 2^{m-1}(m-1)!.$$

By Proposition 35, we have $N(A_i) \geq 2^m$ for all $1 \leq i \leq 3$. By Proposition 37, we have $N(A_i) \geq 1$ for all $7 \leq i \leq 10$. Thus, the number of distinct Klein bottle nonequivalent C_4 -face-magic Klein bottle labelings on $\mathcal{K}_{m,n}$ satisfies

$$\begin{aligned} NKBL(m, n) &= \sum_{A \in MNAPP(m, n)} N(A) \cdot (m-1)! + \sum_{A \in MPAPP(m, n)} N(A) \cdot 2^{m-1}(m-1)! \\ &\geq \sum_{i=1}^3 N(A_i) \cdot (m-1)! + \sum_{i=7}^{10} N(A_i) \cdot 2^{m-1}(m-1)! \geq (5 \cdot 2^m) \cdot (m-1)!. \end{aligned}$$

□

REFERENCES

- [1] M. Bača, On magic labelings of Mobius ladders. *J. Franklin Inst.* **326** (1989), 885–888, doi:10.1016/0016-0032(89)90010-0.
- [2] M. Bača, On magic labelings of honeycomb. *Discrete Math.* **105** (1992), 305–311, doi:10.1016/0012-365X(92)90153-7.
- [3] M. Bača, On magic labelings of grid graphs. *Ars Combin.* **33** (1992), 295–299.
- [4] J. A. Bondy, and U. S. R. Murty, *Graph Theory with Applications*, MacMillan, New York, (1976).
- [5] S. I. Butt, M. Numan, S. Ali and A. Semaničová-Feňovčíková, Face antimagic labelings of toroidal and Klein bottle grid graphs. *AKCE Int. J. Graphs Comb.* (2020), 1–9, doi:10.1016/j.akcej.2018.09.005.
- [6] S. J. Curran and R. M. Low, C_4 -face-magic torus labelings on $C_4 \times C_4$, *Cong. Numer.* **233** (2019), 79–94.
- [7] S. J. Curran, R. M. Low, and S. C. Locke, C_4 -face-magic toroidal labelings on $C_m \times C_n$, *Art Discrete Appl. Math.*, 4 (2021), no. 1, #P1.04, 33 pp., doi.org/10.26493/2590-9770.1368.f37.
- [8] J. A. Gallian, A dynamic survey of graph labeling. *Electron. J. Combin.* **24** (2021), #DS6, doi:10.37236/27.
- [9] J. Hsieh, S-M. Lee, P-J. Liang, and R. M. Low, On C_4 face-(1,0,0) magic polyomino graphs and their related graphs., manuscript.
- [10] A. Kasif, H. Muhammad, A. Ali, and M. Miller, Magic labeling of type (a, b, c) of families of wheels. *Math. Comput. Sci.* **7** (2013), 315–319, doi:10.1007/s11786-013-0162-9.
- [11] K. Kathiresan and S. Gokulakrishnan, On magic labelings of type $(1, 1, 1)$ for the special classes of plane graphs. *Util. Math.* **63** (2003), 25–32.
- [12] A. Kotzig and A. Rosa, Magic valuations of finite graphs. *Canad. Math. Bull.* **13** (1970), 451–461, doi:10.4153/CMB-1970-084-1.
- [13] K-W. Lih, On magic and consecutive labelings of plane graphs. *Util. Math.* **24** (1983), 165–197.
- [14] T-M. Wang, Toroidal grids are anti-magic. In: Wang L. (eds) *Computing and Combinatorics*. COCOON 2005. Lecture Notes in Computer Science, vol 3595. Springer, Berlin, Heidelberg, (2005), 671–679, doi:10.1007/11533719_68

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PITTSBURGH AT JOHNSTOWN, 450 SCHOOLHOUSE RD, JOHNSTOWN, PA 15904, USA

Email address: sjcurran@pitt.edu

DEPARTMENT OF MATHEMATICS, SAN JOSE STATE UNIVERSITY, 1 WASHINGTON SQ, SAN JOSE, CA 95192, USA

Email address: richard.low@sjsu.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, FLORIDA ATLANTIC UNIVERSITY, 777 GLADES RD, BOCA RATON, FL 33431, USA

Email address: lockes@fau.edu