

COMMON KINGS OF A CHAIN OF CYCLES IN A STRONG TOURNAMENT

LOGAN POST AND ZEYU ZHENG

ABSTRACT. It is known that every strong tournament has directed cycles of any length, and thereby strong subtournaments of any size. In this note, we prove that they also can share a common vertex which is a king of all of them. This common vertex can be any king in the whole tournament. Further, the Hamiltonian cycles in them can be recursively constructed by inserting an additional vertex to one directed edge.

1. INTRODUCTION

A **tournament** is a directed graph, obtained by assigning an orientation to each edge of a complete graph. A **king** is a vertex k such that each other vertex is reachable from k by a directed path of length at most 2 [1]. A **strong** tournament has the property that every vertex is reachable from every other by a directed path [2]. A **chain of cycles** is a sequence C_1, C_2, \dots, C_n such that for $i \in [1, n-1]$, the cycle C_{i+1} is obtained from C_i by replacing some edge (x, y) of C_i by two edges (x, z) and (z, y) where z is not on C_i .

It is known that every tournament has a king [1], and it is known that every strong tournament T_n has directed cycles of length $3, \dots, n$ [2]. The proof can be modified so that all cycles contain a common vertex which can be any king in the whole tournament, but not necessarily of the subtournaments determined by the vertices of the cycles.

In this note, we show that:

Theorem 1.1. *Given a strong tournament T_n on n vertices and any king k of T_n , there exist a chain of cycles C_3, C_4, \dots, C_n where C_i is of length i such that k is a king in all subtournaments induced by the vertex sets of the cycles in the chain.*

2. PROOF OF THEOREM 1.1

Let k be a king of T_n . We use A to denote $N^+(k)$ and use B to denote $N^-(k)$. A result in [3] shows that we can partition A into A_1, A_2, \dots, A_r such that A_i determines a strong subtournament for all integer $i \in [1, r]$, and for all $i < j$, all directed edges between A_i and A_j terminates in A_j .

Claim 1. There exists $a^* \in A_r$, $b^* \in B$ such that (a^*, b^*) is an edge in T_n .

For a chosen $a \in A_r$, as T_n is strong, there exists a directed path from a to k . Choose the shortest such path. This path will look like $a \rightarrow \dots \rightarrow a^* \rightarrow b^* \rightarrow k$ for some $b^* \in B$. $a^* \in A_r$ because A_r is fully beaten by A_i for all $i < r$, so this path resides in A_r and no other elements of B are included because this would imply a shorter path. This satisfies Claim 1.

Claim 2. There is a Hamiltonian path through A ending at a^* .

We see that the subtournament on $A_1 \cup A_2 \cup \dots \cup A_{r-1}$ has Hamiltonian path P_1 . Also, A_r is strong, so it has a Hamiltonian cycle, in which there is a Hamiltonian path P_2 ending at a^* . Concatenating P_1 and P_2 , this satisfies Claim 2. We denote this path by $P = a_1 \rightarrow a_2 \rightarrow \dots \rightarrow a_d = a^*$ where $d = \deg^+ k$.

Using Claim 1 and Claim 2, we can construct length- i directed cycles C_i of which k is the king for $i \in [3, n]$.

For any $i \in [1, d]$, we see that $C_{i+2} = k \rightarrow a_{d-i+1} \rightarrow \dots \rightarrow a_d \rightarrow b \rightarrow k$ is a directed cycle of length $i + 2$ and k is the king of the subtournament on $V(C_{i+2})$. This gives us cycles of length from 3 to $d + 2$.

Given a directed cycle C_{i-1} of length $i - 1$ where $i \in [d + 3, n]$ and $(A \cup \{k\}) \subset V(C_{i-1})$. We now show that we can always find a cycle of length i and its vertex set still contains $A \cup \{k\}$.

Since T_n is strong we know that there exists an edge $(w, z) \in T_n$ so that $w \in C_{i-1}$ and $z \in V(T_n) \setminus V(C_{i-1})$. Note that $(V(T_n) \setminus V(C_{i-1})) \subset B$, so (z, k) is an edge in T_n . Therefore, there exists an edge (x, y) on the cycle C_{i-1} such that both (x, z) and (z, y) are edges in T_n . We can insert z in the cycle C_{i-1} between x and y to get a directed cycle C_i of length i , and $(A \cup \{k\}) \subseteq V(C_i)$. This recursively gives us cycles of length from $d + 3$ to n .

Now we have created a collection of cycles C_3, C_4, \dots, C_n in T_n of which k is the king of subtournament induced by vertices of each of them. Note as an added fact that C_{i+1} can be obtained from C_i by replacing some edge (x, y) of C_i by two edges (x, z) and (z, y) where $z \in V(T_n) \setminus V(C_i)$ for all integer $i \in [3, n - 1]$, which finishes the proof.

ACKNOWLEDGEMENTS

The authors thank Professor András Gyárfás for proposing the research problem we solve here in his course Advanced Combinatorics and for his valuable suggestions. The work have been conducted under the auspices of the Budapest Semesters in Mathematics Program during the Spring of 2022.

Yaobin Chen, Professor Jie Ma, Professor Hehui Wu have provided different proofs to the main theorem. The second author would like to thank them for fruitful early discussions on this problem.

REFERENCES

- [1] HG Landau. “On dominance relations and the structure of animal societies: III The condition for a score structure”. In: *The bulletin of mathematical biophysics* 15.2 (1953), pp. 143–148.
- [2] JW Moon. “Topic on Tournaments, Holt, Rinehart and Winston”. In: *Inc., New York* (1968).
- [3] KB Reid and Lowell W Beineke. “Tournaments, Selected Topics in Graph Theory”. In: *Academic Press, London* (1978), pp. 169–204.

HAVERFORD COLLEGE, HAVERFORD, PA, 19041

Email address: `lpost@haverford.edu`

SCHOOL OF MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, SHANGHAI, CHINA 200433

Email address: `zeyuzheng19@fudan.edu.cn`