

Scales

Mathieu Helfter *

May 9, 2024

Abstract

We introduce the notion of *scale* to generalize and compare different invariants of metric spaces and their measures. Several versions of scales are introduced such as *Hausdorff*, *packing*, *box*, *local* and *quantization*. They moreover are defined for different *growth*, allowing in particular a refined study of infinite dimensional spaces. We prove general theorems comparing the different versions of scales. They are applied to describe geometries of ergodic decompositions, of the Wiener measure and of functional spaces. The first application solves a problem of Berger on the notions of emergence (2020); the second lies in the geometry of the Wiener measure and extends the work of Dereich-Lifshits (2005); the last refines Kolmogorov-Tikhomirov (1958) study on functional spaces.

Contents

1	Introduction and results	2
1.1	From dimension to scale	3
1.2	Results on comparisons of scales	5
1.3	Applications	7
2	Metric scales	10
2.1	Scalings	10
2.2	Box scales	13
2.3	Hausdorff scales	15
2.4	Packing scales	16
2.5	Properties and comparison of scales of metric spaces	19
3	Scales of measures	22
3.1	Hausdorff, packing and local scales of measures	22
3.2	Quantization and box scales of measures	23
3.3	Comparison between local and global scales of measures and proof of Theorem C	25
3.4	Proof of Theorem B	27

*Partially supported by the ERC project 818737 *Emergence on wild differentiable dynamical systems*

4 Applications	31
4.1 Scales of infinite products of finite sets	31
4.2 Functional spaces	34

1 Introduction and results

Dimension theory was popularized by Mandelbrot in the article *How long is the coast of Britain ?* [Man67] and shed light on the general problem of measuring how large a natural object is. The category of objects considered are metric spaces possibly endowed with a measure.

Dimension theory encompasses not only smooth spaces such as manifolds, but also wild spaces such as fractals, so that the dimension may be any non-negative real number. There are several notions of dimension: for instance Hausdorff [Hau18], packing [Tri82] or box dimensions; also when the space is endowed with a measure, there are moreover the local and the quantization dimensions. These different versions of dimension are bi-Lipschitz invariants. They are in general not equal, so that they reveal different aspects of the underlying space. The seminal works of Hausdorff, Frostman, Tricot, Fan, Tamashiro, Pötzelberger, Graf-Luschgy and Dereich-Lifshits described the relationship between these notions and gave conditions under which they coincide.

Obviously these invariants do not give much information on infinite dimensional spaces. However such spaces are subject to many studies. Most relevantly, Kolmogorov-Tikhomirov in [KT93] gave asymptotics of the covering numbers of functional spaces. Dereich-Lifshits gave asymptotics of the mass of the small balls for the Wiener measure and exhibited their relationship with the quantization problem, see [DFMS03, DL05, CM44, Chu47, BR92, KL93]. Also Berger and Bochi [Ber20] gave estimates on the covering number and quantization number of the ergodic decomposition of some smooth dynamical systems. See also [BR92, Klo15, BB21].

This leads to the following natural question:

Question. *Are there infinite dimensional counterparts of the different versions of dimension with the same relationships ?*

To answer this question, we introduce the notion of *scale*. The key idea is to consider a *scaling*, that is a one parameter family of gauge functions verifying some mild assumptions, that prescribes at which "scale" the size of space is studied. For instance the given for the *dimension* or the *order* given in Proposition 2.4 are scalings. Given by a scaling, different versions of scales are defined. In particular, the Hausdorff dimension, packing dimension or the box dimension are scales.

We will generalize comparison theorems between the different kind of dimensions to all the different growths of scales in **Theorem A**, **Theorem B** and **Theorem C**. The definition of scaling is tuned so that the proofs of **Theorem A** and **Theorem B** are almost direct generalization of the established case of dimension (see Section 1.2).

*The main difficulty will be then to prove **Theorem C** which enables to compare the quantization scales with both the local and the box scales. Also even for the specific case of dimension, new inequalities between quantization dimension of a measure and box dimension of the set of positive mass are proven in **Theorem C** (inequalities (f) and (h)).*

In the next Section 1.1 we recall usual definitions of dimension and introduce the notions of scaling and scales. The theorems comparing the different versions of scales are stated in Section 1.2. Precise definitions of the involved scales are given in Section 2 and in Section 3 when the space is endowed with a measure. Then Section 1.3 is dedicated to applications of the main results. In Section 1.3.1, a first application of **Theorem C** together with Dereich-Lifshits

estimate [DL05] implies the coincidence of local, Hausdorff, packing, quantization and box orders of the Wiener measure for the L^p -norm, for any $p \in [1, \infty]$. Then in Section 1.3.2, we apply **Theorem A** to show the coincidence of the box, Hausdorff and packing orders for finitely regular functional spaces; refining Kolmogorov-Tikhomirov study in [KT93, Thm XV]. Lastly in Section 1.3.3, a consequence of **Theorem C** is that the local order of the ergodic decomposition is at most its quantization order. This solves a problem set by Berger in [Ber20].

Thanks: Warmest thanks to Pierre Berger for his investment and advice; to Ai-Hua Fan for his interest, references and advice, François Ledrappier for answering my questions and giving references, Martin Leguil for his interest, and Camille Tardif and Nicolas de Saxcé for giving me references.

1.1 From dimension to scale

Let us first recall some classical definitions of dimension theory and see how they could be naturally extended to define finite invariants for infinite dimensional spaces. The Hausdorff, packing and box dimensions of a totally bounded metric space (X, d) are defined by looking at families of subsets of X . For the box dimensions. Given an error $\epsilon > 0$, recall that the covering number $\mathcal{N}_\epsilon(X)$ is the minimal cardinality of a covering of X by balls of radius ϵ . Then lower and upper box dimensions of (X, d) are given by:

$$\underline{\dim}_B X := \sup\{\alpha > 0 : \phi_\alpha(\epsilon) \cdot \mathcal{N}_\epsilon(X) \rightarrow +\infty\} \quad \text{and} \quad \overline{\dim}_B X := \inf\{\alpha > 0 : \phi_\alpha(\epsilon) \cdot \mathcal{N}_\epsilon(X) \rightarrow 0\},$$

where $(\phi_\alpha)_{\alpha > 0}$ is the family of functions on $(0, 1)$ given for $\alpha > 0$ by $\phi_\alpha : \epsilon \mapsto \epsilon^\alpha$.

In general upper and lower box dimensions do not coincide (see e.g. [FF97, Fal04]). However when X is a smooth manifold endowed with the euclidean metric these definitions coincide with the usual definition of dimension. Basic properties of box dimensions are revealed when looking at subsets of a metric space with the induced metric. Notably, box dimensions are non decreasing for the union of subsets and invariant by topological closure. In particular, in general they are not countable stable: the box dimensions of a countable union of subsets of a metric space are a priori not equal to the suprema of the corresponding dimensions of the subsets. The most popular version of dimension that enjoy the property of being countable stable is *Hausdorff dimension*. Let us recall its definition. Given an error $\epsilon > 0$, consider:

$$\mathcal{H}_\epsilon^\alpha(X) := \inf_{C \in \mathcal{C}_H(\epsilon)} \sum_{B(x, \delta) \in C} \phi_\alpha(\delta),$$

where $\mathcal{C}_H(\epsilon)$ is the set of countable coverings of X by disjoint balls of radius at most ϵ . Then, the *Hausdorff dimension* of (X, d) is given by:

$$\dim_H X = \sup \left\{ \alpha > 0 : \mathcal{H}_\epsilon^\alpha(X) \xrightarrow{\epsilon \rightarrow 0} +\infty \right\} = \inf \left\{ \alpha > 0 : \mathcal{H}_\epsilon^\alpha(X) \xrightarrow{\epsilon \rightarrow 0} 0 \right\}.$$

Lastly, another interesting dimension that enjoys countable stability is the *packing dimension*. Its construction is analogous to the one of Hausdorff dimension and was introduced by Tricot in his thesis [Tri82]. It is actually linked to upper box dimension by the following characterization that we will use for now as a definition:

$$\dim_P X := \inf \sup_{n \geq 1} \overline{\dim}_B E_n,$$

where the infimum is taken over countable coverings $(E_n)_{n \geq 1}$ by subsets of X . These four versions of dimension are bi-Lipschitz invariants; they quantify different aspects of the geometry of the studied metric space since they a priori do not coincide. However, it always holds:

$$\dim_H X \leq \underline{\dim}_B X \leq \overline{\dim}_B X \quad \text{and} \quad \dim_H X \leq \dim_P X \leq \overline{\dim}_B X .$$

See e.g. [FF97], [Fal04] for full proofs.

Let us now introduce *scales*. A simple observation is that all of the above versions of dimension imply a specific parameterized family $(\phi_\alpha)_{\alpha > 0} = (\epsilon \mapsto \epsilon^\alpha)_{\alpha > 0}$ of gauge functions with polynomial behaviour. Classically, a gauge function is a generalization of the measurement of diameters of balls that is used to refine the definition of Hausdorff measure for finite dimensional spaces. The idea here is totally different, instead of finding a refinement we will take functions with behaviour possibly far away from being polynomial. Let us precise the discussion. If a space (X, d) is infinite dimensional then its covering number $\mathcal{N}_\epsilon(X)$ grows faster than any polynomial in ϵ^{-1} as ϵ decreases to 0. Thus to hope defining finite invariants for infinite dimensional spaces we must allow other gauge functions that decreases faster than any polynomial when the radius of the involved balls decrease to 0. Consequently, we propose here to replace in all the above definitions of dimensions, the family $(\phi_\alpha)_{\alpha > 0} = (\epsilon \in (0, 1) \mapsto \epsilon^\alpha)_{\alpha > 0}$ by another family of gauge functions which encompasses the following examples of growth:

- Example 1.1.**
1. *The family $\text{dim} = (\epsilon \in (0, 1) \mapsto \epsilon^\alpha)_{\alpha > 0}$ which is used in the definitions of dimensions,*
 2. *the family $\text{ord} = (\epsilon \in (0, 1) \mapsto \exp(-\epsilon^{-\alpha}))_{\alpha > 0}$ which is called order. It fits with the growth of the covering number of spaces of finitely regular functions studied by Kolmogorov-Tikhomirov [KT93], see Theorem 1.11, or with the one of the space of ergodic measures spaces of dynamics by Berger-Bochi [Ber20], as we will see in Proposition 1.16,*
 3. *the family $(\epsilon \in (0, 1) \mapsto \exp(-(\log \epsilon^{-1})^\alpha))_{\alpha > 0}$ which fits with the growth of the covering number of holomorphic functions estimated by Kolmogorov-Tikhomirov [KT93], as we will see in Theorem 2.5.*

Yet to extend properly the definitions and comparison theorems between the scales, i.e. the generalization of box, Hausdorff and packing dimensions-, the family $(\phi_\alpha)_{\alpha > 0}$ must satisfy some mild assumptions, which leads to introduce *scalings*:

Definition 1.2 (Scaling). *A family $\text{scl} = (\text{scl}_\alpha)_{\alpha \geq 0}$ of positive non-decreasing functions on $(0, 1)$ is a scaling when for any $\alpha > \beta > 0$ and any $\lambda > 1$ close enough to 1, it holds:*

$$(*) \quad \text{scl}_\alpha(\epsilon) = o(\text{scl}_\beta(\epsilon^\lambda)) \quad \text{and} \quad \text{scl}_\alpha(\epsilon) = o(\text{scl}_\beta(\epsilon)^\lambda) \quad \text{when } \epsilon \rightarrow 0 .$$

Remark 1.3. *The left hand side condition is used in all the proof of our theorems represented on Fig. 1. The right hand side condition is only used to prove the equalities between packing and upper local scales in **Theorem B** and to compare upper local scales with upper box and upper quantization scales in **Theorem C** inequalities (c)&(g). It moreover allows to characterize packing scale with packing measure.*

Remark 1.4. *There are scalings that allows to study 0-dimensional spaces, e.g. $(\epsilon \in (0, 1) \mapsto \log(\epsilon^{-1})^{-\alpha})_{\alpha > 0}$.*

We will show in Proposition 2.4 that the families in Example 1.1 are scalings. Scalings allow to define *scales* which generalize packing dimension, Hausdorff dimension, box dimensions, quantization dimensions and local dimensions that are local counterparts for measures. For each scaling, the different kind of scales do not a priori coincide on a

generic space. Nevertheless in Section 1.3, as a direct application of our comparison theorems, we bring examples of metric spaces and measures where all those definitions coincide. In these examples, equalities between the different scales are linked to some underlying "homogeneity" of the space.

Now for a metric space (X, d) , replacing the specific family dim in the definition of box dimensions by a given scaling $\text{scl} = (\text{scl}_\alpha)_{\alpha>0}$ gives the following:

Definition 1.5 (Box scales). Lower and upper box scales of a metric space (X, d) are defined by:

$$\underline{\text{scl}}_B X = \sup \left\{ \alpha > 0 : \mathcal{N}_\epsilon(X) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} +\infty \right\} \quad \text{and} \quad \overline{\text{scl}}_B X = \inf \left\{ \alpha > 0 : \mathcal{N}_\epsilon(X) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} 0 \right\} .$$

Moreover we will generalize the notion Hausdorff and packing dimensions to the *Hausdorff scale* denoted $\text{scl}_H X$ (see Definition 2.13) and *packing scale* denoted $\text{scl}_P X$ (see Definition 2.14). The construction are fully detailed in the next section. Let us now state the main results on comparison of scales of metric spaces.

1.2 Results on comparisons of scales




Figure 1: Diagram presenting results of Theorems A, B and C. Each arrow is an inequality, the scale at the starting point of the arrow is at least the one at its ending point : " \rightarrow " = " \geq ". None of them is an equality in the general case. If there is no path between two scales scl_1 and scl_2 then there exist examples of spaces endowed with a measure such that both $\text{scl}_1 > \text{scl}_2$ and $\text{scl}_1 < \text{scl}_2$ can happen. See Example 4.6.

In this section, we introduce other kind of scales and Theorems A, B and C which state the inequalities between them as illustrated in Fig. 1. First, we bring the following generalization of classical inequalities comparing dimensions of metric spaces to the frame of scales:

Theorem A. Let (X, d) be a metric space and scl a scaling, the following inequalities hold:

$$\text{scl}_H X \leq \text{scl}_P X \leq \overline{\text{scl}}_B X \quad \text{and} \quad \text{scl}_H X \leq \underline{\text{scl}}_B X \leq \overline{\text{scl}}_B X .$$

In the specific case of dimension these inequalities are well known and redacted for instance by Tricot [Tri82] or Falconer [FF97, Fal04]. The proof of this theorem will be done in Section 2.5. The key part is to show that Hausdorff scales and packing scales are well defined quantities. Then we will follow the lines of Falconer's proof to show **Theorem A**.

When the metric space (X, d) is endowed with a measure μ , Frostman first studied the relationship between the Hausdorff dimension and the growth of the mass of the small balls. This has been intensively studied by Fan [FLR02, Fan94], Pötzelberger [Pöt99], Tamashiro [Tam95] as local dimension. Similarly we introduce *local scales* that extend the notion of local dimensions of a measure:

Definition 1.6 (Local scales). Let μ be Borel measure on a metric space (X, d) and scl a scaling. The lower and upper scales of μ are the functions that map a point $x \in X$ to:

$$\underline{\text{scl}}_{\text{loc}}\mu(x) = \sup \left\{ \alpha > 0 : \frac{\mu(B(x, \epsilon))}{\text{scl}_\alpha(\epsilon)} \xrightarrow{\epsilon \rightarrow 0} 0 \right\} \quad \text{and} \quad \overline{\text{scl}}_{\text{loc}}\mu(x) = \inf \left\{ \alpha > 0 : \frac{\mu(B(x, \epsilon))}{\text{scl}_\alpha(\epsilon)} \xrightarrow{\epsilon \rightarrow 0} +\infty \right\} .$$

As in dimension theory, we should not compare the local scales with the scales of X but to the ones of its subsets with positive mass. This observation leads to consider the following:

Definition 1.7 (Hausdorff, packing and box scales of a measure). Let scl be a scaling and μ a non-null Borel measure on a metric space (X, d) . For any $\text{scl}_\bullet \in \{\text{scl}_H, \text{scl}_P, \underline{\text{scl}}_B, \overline{\text{scl}}_B\}$ we define lower and upper scales of the measure μ by:

$$\text{scl}_\bullet\mu = \inf_{E \in \mathcal{B}} \{\text{scl}_\bullet E : \mu(E) > 0\} \quad \text{and} \quad \text{scl}_\bullet^*\mu = \inf_{E \in \mathcal{B}} \{\text{scl}_\bullet E : \mu(X \setminus E) = 0\} ,$$

where \mathcal{B} is the set of Borel subsets of X .

In the case of dimension, Frostman [Fro35], Tricot [Tri82], Fan [Fan94, FLR02] and Tamashiro [Tam95] exhibited the relationship between the Hausdorff and packing dimensions of measures and their local dimensions that we generalize as:

Theorem B. Let μ be a Borel measure on a metric space (X, d) , then for any scaling scl , Hausdorff and packing scales of μ are characterized by:

$$\text{scl}_H\mu = \text{ess inf } \underline{\text{scl}}_{\text{loc}}\mu, \quad \text{scl}_H^*\mu = \text{ess sup } \underline{\text{scl}}_{\text{loc}}\mu, \quad \text{scl}_P\mu = \text{ess inf } \overline{\text{scl}}_{\text{loc}}\mu, \quad \text{scl}_P^*\mu = \text{ess sup } \overline{\text{scl}}_{\text{loc}}\mu ,$$

where ess sup and ess inf denote the essential suprema and infima of a function.

The proof of the latter theorem is done in Section 3.1. The proof follows the lines of the one of the case of dimension from Fan in [Fan94, FLR02].

Let us introduce a last kind of scale, the *quantization scale*. It generalizes the quantization dimension which dragged much research interest [GL07, Pöt99, DFMS03, DL05, Ber17, BB21, Ber20].

Definition 1.8 (Quantization scales). Let (X, d) be a metric space and μ a Borel measure on X . The quantization number \mathcal{Q}_μ of μ is the function that maps $\epsilon > 0$ to the minimal cardinality of a set of points that is on average ϵ -close to any point in X :

$$\mathcal{Q}_\mu(\epsilon) = \inf \left\{ N \geq 0 : \exists \{c_i\}_{i=1, \dots, N} \subset X, \int_X d(x, \{c_i\}_{1 \leq i \leq N}) d\mu(x) \leq \epsilon \right\} .$$

Then lower and upper quantization scales of μ for a given scaling scl are defined by:

$$\underline{\text{scl}}_Q \mu = \sup \left\{ \alpha > 0 : \mathcal{Q}_\mu(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} +\infty \right\} \quad \text{and} \quad \overline{\text{scl}}_Q \mu = \inf \left\{ \alpha > 0 : \mathcal{Q}_\mu(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} 0 \right\} .$$

The following gives relationships between the remaining kind of introduced scales of measures:

Theorem C (Main). *Let (X, d) be a metric space. Let μ be a Borel measure on X . For any scaling scl , the following inequalities on the scales of μ hold:*

$$\text{ess inf } \underbrace{\underline{\text{scl}}_{\text{loc}} \mu}_{(a)} \leq \underbrace{\underline{\text{scl}}_B \mu}_{(b)} \leq \underline{\text{scl}}_Q \mu \quad ; \quad \text{ess inf } \underbrace{\overline{\text{scl}}_{\text{loc}} \mu}_{(c)} \leq \underbrace{\overline{\text{scl}}_B \mu}_{(d)} \leq \overline{\text{scl}}_Q \mu$$

and

$$\text{ess sup } \underbrace{\underline{\text{scl}}_{\text{loc}} \mu}_{(e)} \leq \underbrace{\underline{\text{scl}}_Q \mu}_{(f)} \leq \underbrace{\underline{\text{scl}}_B^* \mu}_{(g)} \quad ; \quad \text{ess sup } \underbrace{\overline{\text{scl}}_{\text{loc}} \mu}_{(h)} \leq \overline{\text{scl}}_Q \mu \leq \overline{\text{scl}}_B^* \mu .$$

The proof of inequalities (b) and (d) is done at Theorem 3.10 and relies mainly on the use of Borel-Cantelli lemma. Even in the specific case of dimension, these inequalities were not shown yet, as far as we know. The proof of inequalities (f) and (h) is straightforward, see Lemma 3.8. Inequalities (e) and (g) were shown by Pötzelberger in [Pöt99] for dimension and in $[0, 1]^d$. A new approach for the general case of scales of inequalities (e) and (g) is brought in Theorem 3.12. We deduce the inequality (a) from (e) and (f) and inequality (c) from (g) and (h). As a direct application, inequality (e) allows to answer to a problem set by Berger in [Ber20] (see Section 1.3.3). We will give in Section 4.1 examples of topological compact groups different versions of orders do not coincide. Moreover in that same section we show that for a metric group where the law is Lipschitz, the Hausdorff scale coincides with the lower box scale and the packing scale coincides with the upper box scale.

1.3 Applications

Let us see how our main theorems imply easily the coincidence of the scales of some natural infinite dimensional spaces.

1.3.1 Wiener measure

First example is the calculus of the orders of the Wiener measure W that describes uni-dimensional standard Brownian motion on $[0, 1]$. Recall that W is the law of a continuous process $(B_t)_{t \in [0, 1]}$ with independent increments. It is such that for any $t \geq s$ the law of the random variable $B_t - B_s$ is $\mathcal{N}(0, t - s)$. Computation of the local scales of the Wiener measure relies on small ball estimates which received much interest [CM44, Chu47, BR92, KL93]. These results gave asymptotics on the measure of small balls centered at 0 for L^p norms and Hölder norms. Moreover for a random ball the Dereich-Lifshits made the following estimate for L^p -norms:

Theorem 1.9 (Dereich-Lifshits [DL05][3.2, 5.1, 6.1, 6.3]). *For the Wiener measure on $C^0([0, 1], \mathbb{R})$ endowed with the L^p -norm, for $p \in [1, \infty]$, there exists¹ a constant $\kappa > 0$ such that for W -almost any $\omega \in C^0([0, 1], \mathbb{R})$:*

$$-\epsilon^2 \cdot \log W(B(\omega, \epsilon)) \rightarrow \kappa, \text{ when } \epsilon \rightarrow 0 \quad ,$$

and moreover the quantization number of W verifies:

$$\epsilon^2 \cdot \log \mathcal{Q}_W(\epsilon) \rightarrow \kappa, \text{ when } \epsilon \rightarrow 0 \quad .$$

¹Note that for $p < \infty$, the constant κ does not depend on the value of p .

As a direct consequence of **Theorem B** and **Theorem C** we get that the new invariants we introduced for a measure with growth given by ord all coincide:

Theorem D (Orders of the Wiener measure). *For the Wiener measure on $C^0([0, 1], \mathbb{R})$ endowed with the L^p -norm, for $p \in [1, \infty]$, verifies for W almost every $\omega \in C^0([0, 1])$:*

$$\underline{\text{ord}}_{\text{loc}}(\omega) = \text{ord}_H W = \text{ord}_H^* W = \overline{\text{ord}}_{\text{loc}}(\omega) = \text{ord}_P W = \text{ord}_P^* W = \underline{\text{ord}}_B W = \underline{\text{ord}}_Q W = \overline{\text{ord}}_B W = \overline{\text{ord}}_Q W = 2 .$$

Proof. By **Theorem 1.9**, for W -almost ω and for any $p \in [1, \infty]$, in the L^p -norm it holds:

$$\underline{\text{ord}}_{\text{loc}} W(\omega) = \overline{\text{ord}}_{\text{loc}} W(\omega) = 2 = \underline{\text{ord}}_Q W = \overline{\text{ord}}_Q W .$$

Now by **Theorem B**:

$$\text{ord}_H W = \underline{\text{ord}}_{\text{loc}} W(\omega) = \text{ord}_H^* W \quad \text{and} \quad \text{ord}_P W = \overline{\text{ord}}_{\text{loc}} W(\omega) = \text{ord}_P^* W .$$

Finally since by **Theorem C**:

$$\overline{\text{ord}}_Q W \geq \overline{\text{ord}}_B W \geq \underline{\text{ord}}_B W \geq \text{ord}_H W ,$$

the sought result comes by combining the three above lines of equalities and inequalities. \square

Remark 1.10. *Since $C^0([0, 1], \mathbb{R})$ and endowed with the L^p -norm is not totally bounded, as well as any of its subsets with total mass, it holds $\underline{\text{ord}}_B^* = +\infty$.*

1.3.2 Functional spaces endowed with the C^0 -norm

Let d be a positive integer. For any integer $k \geq 0$ and for any $\alpha \in [0, 1]$ denote:

$$\mathcal{F}^{d,k,\alpha} := \{ f \in C^k([0, 1]^d, [-1, 1]) : \|f\|_{C^k} \leq 1, \text{ and if } \alpha > 0, \text{ the map } D^k f \text{ is } \alpha\text{-H\"older with constant } 1 \} .$$

We endow this space with the C^0 norm. See **Section 4.2** for the definition of the C^k -norms.

Kolmogorov-Tikhomirov gave the following asymptotics:

Theorem 1.11 (Kolmogorov-Tikhomirov, [KT93][Thm XV]). *Let d be a positive integer. For any integer k and for any $\alpha \in [0, 1]$, there exist two constants $C_1 > C_2 > 0$ such that the covering number $\mathcal{N}_\epsilon(\mathcal{F}^{d,k,\alpha})$ of the space $(\mathcal{F}^{d,k,\alpha}, \|\cdot\|_\infty)$ verifies:*

$$C_1 \cdot \epsilon^{-\frac{d}{k+\alpha}} \geq \log \mathcal{N}_\epsilon(\mathcal{F}^{d,k,\alpha}) \geq C_2 \cdot \epsilon^{-\frac{d}{k+\alpha}} .$$

In **Section 4.2** by embedding a group whose Hausdorff order is bounded from below into $\mathcal{F}^{k,d,\alpha}$ (see **Section 4.2**), via an expanding map, we will prove:

Lemma 1.12. *Let d be a positive integer. For any integer k and for any $\alpha \in [0, 1]$, it holds:*

$$\text{ord}_H \mathcal{F}^{d,k,\alpha} \geq \frac{d}{k+\alpha} .$$

The above lemma together with **Theorem A** gives the following extension of Kolmogorov-Tikhomirov's Theorem:

Theorem E. *Let d be a positive integer. For any integer k and for any $\alpha \in [0, 1]$, it holds:*

$$\text{ord}_H \mathcal{F}^{d,k,\alpha} = \text{ord}_P \mathcal{F}^{d,k,\alpha} = \underline{\text{ord}}_B \mathcal{F}^{d,k,\alpha} = \overline{\text{ord}}_B \mathcal{F}^{d,k,\alpha} = \frac{d}{k + \alpha} .$$

Proof of Theorem E. First, by **Theorem A**, it holds:

$$\text{ord}_H \mathcal{F}^{d,k,\alpha} \leq \underline{\text{ord}}_B \mathcal{F}^{d,k,\alpha} \leq \overline{\text{ord}}_B \mathcal{F}^{d,k,\alpha} \quad \text{and} \quad \text{ord}_H \mathcal{F}^{d,k,\alpha} \leq \text{ord}_P \mathcal{F}^{d,k,\alpha} \leq \overline{\text{ord}}_B \mathcal{F}^{d,k,\alpha} .$$

From there by **Theorem 1.11** and **Lemma 1.12**, it holds:

$$\frac{d}{k + \alpha} \leq \text{ord}_H \mathcal{F}^{d,k,\alpha} \leq \underline{\text{ord}}_B \mathcal{F}^{d,k,\alpha} \leq \overline{\text{ord}}_B \mathcal{F}^{d,k,\alpha} = \frac{d}{k + \alpha} ,$$

and

$$\frac{d}{k + \alpha} \leq \text{ord}_H \mathcal{F}^{d,k,\alpha} \leq \text{ord}_P \mathcal{F}^{d,k,\alpha} \leq \overline{\text{ord}}_B \mathcal{F}^{d,k,\alpha} = \frac{d}{k + \alpha} .$$

From there, all of the above inequalities are indeed equalities, which gives the sought result. \square

1.3.3 Local and global emergence

The framework of scales moreover allow to answer to a problem set by Berger in **[Ber20]** on wild dynamical systems. We now consider a compact metric space (X, d) and a measurable map $f : X \rightarrow X$. We denote \mathcal{M} the set of probability Borel measures on X and \mathcal{M}_f the subset of \mathcal{M} of f -invariant measures. The space \mathcal{M} is endowed with the Wasserstein distance W_1 defined by:

$$W_1(\nu_1, \nu_2) = \sup_{\phi \in \text{Lip}^1(X)} \int \phi d(\nu_1 - \nu_2) ,$$

inducing the weak *- topology for which \mathcal{M} is compact. A way to measure the wildness of a dynamical system is to measure how far from being ergodic an invariant measure μ is. Then by Birkhoff's theorem given a measure $\mu \in \mathcal{M}_f$, for μ -almost every $x \in X$ the following measure is well defined:

$$e(x) := \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \delta_{f^k(x)} ,$$

and moreover the limit measure is ergodic. The definition of emergence, introduced by Berger, describes the size of the set of ergodic measures reachable by limits of empirical measures given an f -invariant probability measure on X .

Definition 1.13 (Emergence, **[Ber17, BB21]**). *The emergence of a measure $\mu \in \mathcal{M}_f$ at $\epsilon > 0$ is defined by:*

$$\mathcal{E}_\mu(\epsilon) = \min\{N \in \mathbb{N} : \exists \nu_1, \dots, \nu_N \in \mathcal{M}_f, \int_X W_1(e_f(x), \{\nu_i\}_{1 \leq i \leq N}) d\mu(x) \leq \epsilon\} .$$

The case of high emergence corresponds to dynamics where the considered measure is not ergodic at all. The following result shows us that the order is an adapted scaling in the study of the ergodic decomposition.

Theorem 1.14 (**[BGV07, Klo15, BB21]**). *Let (X, d) be a metric compact space of finite then:*

$$\underline{\dim}_B X \leq \underline{\text{ord}}_B(\mathcal{M}) \leq \overline{\text{ord}}_B(\mathcal{M}) \leq \overline{\dim}_B X .$$

For a given measure $\mu \in \mathcal{M}_f$ we define its *emergence order* by:

$$\overline{\text{ord}}\mathcal{E}_\mu := \limsup_{\epsilon \rightarrow 0} \frac{\log \log \mathcal{E}_\mu(\epsilon)}{-\log \epsilon} = \inf \left\{ \alpha > 0 : \mathcal{E}_\mu(\epsilon) \cdot \exp(-\epsilon^{-\alpha}) \xrightarrow{\epsilon \rightarrow 0} 0 \right\} .$$

We denote $\mu_{e_f} := e_f \star \mu$ the *ergodic decomposition* of μ ; it is the probability measure on \mathcal{M}_f equal to the push forward by e of μ . A local analogous local quantity to the emergence order is the local order of the ergodic decomposition of μ , for $\nu \in \mathcal{M}_f$ it is defined by:

$$\overline{\text{ord}}\mathcal{E}_\mu^{\text{loc}}(\nu) := \limsup_{\epsilon \rightarrow 0} \frac{\log -\log(\mu_{e_f}(B(\nu, \epsilon)))}{-\log \epsilon} .$$

Berger asked if the the following comparison between asymptotic behaviour of the mass of the balls of the ergodic decomposition of μ and the asymptotic behaviour of its quantization holds.

Problem 1.15 (Berger, [Ber20, Pbm 4.22]). *Let (X, d) be a compact metric space, $f : X \rightarrow X$ a measurable map and μ a Borel f -invariant measure on X . Does the following holds ?*

$$\int_{\mathcal{M}_f} \overline{\text{ord}}\mathcal{E}_\mu^{\text{loc}} d\mu_{e_f} \leq \overline{\text{ord}}\mathcal{E}_\mu .$$

We propose here a stronger result that answer to latter problem as a direct application of **Theorem C**:

Proposition 1.16. *Let (X, d) be a compact metric space, $f : X \rightarrow X$ a measurable map and μ a Borel f -invariant measure on X . For μ_{e_f} -almost every $\nu \in \mathcal{M}$, it holds:*

$$\overline{\text{ord}}\mathcal{E}_\mu^{\text{loc}}(\nu) \leq \overline{\text{ord}}\mathcal{E}_\mu .$$

Proof. Note that $\overline{\text{ord}}\mathcal{E}_\mu^{\text{loc}} = \overline{\text{ord}}_{\text{loc}}\mu_{e_f}$ and $\overline{\text{ord}}\mathcal{E}_\mu = \overline{\text{ord}}_Q\mu_{e_f}$. Now by **Theorem C**, it holds μ_{e_f} -almost surely that $\overline{\text{ord}}_{\text{loc}}\mu_{e_f} \leq \overline{\text{ord}}_Q\mu_{e_f}$ which is the sought result. \square

2 Metric scales

Before defining and comparing metric scales we show basic handful properties of scalings and present some relevant examples of scalings.

2.1 Scalings

We first recall that a scaling is a family $\text{scl} = (\text{scl}_\alpha)_{\alpha \geq 0}$ of positive non-decreasing functions on $(0, 1)$ is a *scaling* when for any $\alpha > \beta > 0$ and any $\lambda > 1$ close enough to 1, it holds:

$$(*) \quad \text{scl}_\alpha(\epsilon) = o(\text{scl}_\beta(\epsilon^\lambda)) \quad \text{and} \quad \text{scl}_\alpha(\epsilon) = o(\text{scl}_\beta(\epsilon)^\lambda) \quad \text{when } \epsilon \rightarrow 0 .$$

An immediate consequence of the latter definition is the following:

Fact 2.1. *Let scl be a scaling then for any $\alpha > \beta > 0$ and for any constant $C > 0$ it holds for $\epsilon > 0$ small enough:*

$$\text{scl}_\alpha(\epsilon) \leq \text{scl}_\beta(C \cdot \epsilon) .$$

A consequence of the latter fact is the following which compares scales of metric spaces and measures:

Lemma 2.2. *Let $f, g : \mathbb{R}_+^* \rightarrow \mathbb{R}_+^*$ be two functions defined such that $f \leq g$ near 0, thus for any constant $C > 0$:*

$$\inf \left\{ \alpha > 0 : f(C \cdot \epsilon) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} 0 \right\} \leq \inf \left\{ \alpha > 0 : g(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} 0 \right\}$$

and

$$\sup \left\{ \alpha > 0 : f(C \cdot \epsilon) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} +\infty \right\} \leq \sup \left\{ \alpha > 0 : g(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} +\infty \right\}.$$

Proof. It suffices to observe that, by **Fact 2.1**, for any $\alpha > \beta > 0$, and $\epsilon > 0$ small, it holds:

$$f(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \leq g(\epsilon) \cdot \text{scl}_\beta(C^{-1} \cdot \epsilon) = g(C \cdot \tilde{\epsilon}) \cdot \text{scl}_\beta(\tilde{\epsilon}),$$

with $\tilde{\epsilon} = C \cdot \epsilon$. □

The following gives a sequential characterization of scales.

Lemma 2.3 (Sequential characterization of scales). *Let scl be a scaling and $f : \mathbb{R}_+^* \rightarrow \mathbb{R}_+^*$ be a non increasing function.*

Let $(r_n)_{n \geq 1}$ be a positive sequence decreasing to 0 such that $\log r_{n+1} \sim \log r_n$ as $n \rightarrow +\infty$, then it holds:

$$\inf \left\{ \alpha > 0 : f(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} 0 \right\} = \inf \left\{ \alpha > 0 : f(r_n) \cdot \text{scl}_\alpha(r_n) \xrightarrow{n \rightarrow +\infty} 0 \right\}$$

and

$$\sup \left\{ \alpha > 0 : f(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} +\infty \right\} = \sup \left\{ \alpha > 0 : f(r_n) \cdot \text{scl}_\alpha(r_n) \xrightarrow{n \rightarrow +\infty} +\infty \right\}.$$

Proof. Consider $\alpha > 0$ and $\epsilon > 0$. If ϵ is small enough, there exists an integer $n > 0$ that verifies $r_{n+1} < \epsilon \leq r_n$, thus, since f is not increasing and scl_α is increasing, it holds:

$$f(r_n) \cdot \text{scl}_\alpha(r_{n+1}) \leq f(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \leq f(r_{n+1}) \cdot \text{scl}_\alpha(r_n).$$

Let β, γ be positive real numbers such that $0 < \beta < \alpha < \gamma$. For λ close to 1 and ϵ small enough, by **Definition 1.2** of scaling, it holds:

$$\text{scl}_\gamma(r_n) \leq \text{scl}_\alpha(r_n^\lambda) \quad \text{and} \quad \text{scl}_\alpha(r_n) \leq \text{scl}_\beta(r_n^\lambda).$$

Observe now that $r_{n+1} = r_n^{\frac{\log r_{n+1}}{\log r_n}}$, and since $\log r_{n+1} \sim \log r_n$. For n great enough, it holds then:

$$r_n^\lambda \leq r_{n+1}.$$

Since the functions of the scaling are increasing, it follows:

$$\text{scl}_\gamma(r_n) \leq \text{scl}_\alpha(r_{n+1}) \quad \text{and} \quad \text{scl}_\alpha(r_n) \leq \text{scl}_\beta(r_{n+1}).$$

We can now deduce from the latter and the first line of inequalities that:

$$f(r_n) \cdot \text{scl}_\gamma(r_n) \leq f(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \quad \text{and} \quad f(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \leq f(r_{n+1}) \cdot \text{scl}_\beta(r_{n+1}).$$

It follows:

$$\limsup_{n \rightarrow +\infty} f(r_n) \cdot \text{scl}_\gamma(r_n) \leq \limsup_{\epsilon \rightarrow 0} f(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \leq \limsup_{n \rightarrow +\infty} f(r_n) \cdot \text{scl}_\beta(r_n)$$

and

$$\liminf_{n \rightarrow +\infty} f(r_n) \cdot \text{scl}_\gamma(r_n) \leq \liminf_{\epsilon \rightarrow 0} f(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \leq \liminf_{n \rightarrow +\infty} f(r_n) \cdot \text{scl}_\beta(r_n).$$

Since this holds for every positive α and that β and γ can be taken arbitrarily close to α , we get the sought result. □

The following provides many scalings and shows in particular that the families brought in Example 1.1 are indeed scalings.

Proposition 2.4. *For any integers $p, q \geq 1$, the family $\text{scl}^{p,q} = (\text{scl}_\alpha^{p,q})_{\alpha>0}$ defined for any $\alpha > 0$ by:*

$$\text{scl}_\alpha^{p,q} : \epsilon \in (0, 1) \mapsto \frac{1}{\exp^{\circ p}(\alpha \cdot \log_+^{\circ q}(\epsilon^{-1}))}$$

is a scaling; where $\log_+ : t \in \mathbb{R} \mapsto \log(t) \cdot \mathbb{1}_{t>1}$ is the positive part of the logarithm.

We prove this proposition below. Now note in particular that $\text{scl}^{1,1} = \dim = (\epsilon \in (0, 1) \mapsto \epsilon^\alpha)_{\alpha>0}$ and $\text{scl}^{2,1} = \text{ord} = (\epsilon \in (0, 1) \mapsto \exp(-\epsilon^{-\alpha}))_{\alpha>0}$ are both scalings. Let us give an example of space which have finite box scales for the scaling $\text{scl}^{2,2}$ as defined in Proposition 2.4. Consider the space A of holomorphic functions on the disk $\mathbb{D}(R) \subset \mathbb{C}$ of radius $R > 1$ which are uniformly bounded by 1:

$$A = \left\{ \phi = \sum_{n \geq 0} a_n z^n \in C^\omega(\mathbb{D}(R), \mathbb{C}) : \sup_{\mathbb{D}(R)} |\phi| \leq 1 \right\} \text{ endowed with the norm } \|\phi\|_\infty := \sup_{z \in \mathbb{D}(1)} |\phi(z)| .$$

The following implies:

$$\text{scl}_B^{2,2} A = \overline{\text{scl}_B^{2,2} A} = 2 .$$

Theorem 2.5 (Kolmogorov, Tikhomirov [KT93][Equality 129]). *The following estimate on the covering number of $(A, \|\cdot\|_\infty)$ holds:*

$$\log \mathcal{N}_\epsilon(A) = (\log R)^{-1} \cdot |\log \epsilon|^2 + O(\log \epsilon^{-1} \cdot \log \log \epsilon^{-1}), \text{ when } \epsilon \text{ tends to } 0 .$$

Let us now prove Proposition 2.4:

Proof of Proposition 2.4. First it is clear that $\text{scl}^{p,q}$ is a family of non-decreasing functions. Moreover the family is non-increasing. We prove the following below:

Lemma 2.6. *For any $\gamma > 0$ and $\nu > 1$ close to 1, it holds for $\epsilon > 0$ small:*

$$\text{scl}_{\nu \cdot \gamma}^{p,q}(\epsilon) \leq \text{scl}_\gamma^{p,q}(\epsilon^\nu) \quad \text{and} \quad \text{scl}_{\nu \cdot \gamma}^{p,q}(\epsilon) \leq \text{scl}_\gamma^{p,q}(\epsilon)^\nu .$$

Let us show how this lemma implies condition (*) in Definition 1.2 of scaling and thus the result of the proposition. For $\alpha > \beta > 0$, consider $\lambda > 1$ such that $\alpha > \lambda^2 \cdot \beta$, since the family is non-increasing, it holds:

$$\text{scl}_\alpha^{p,q} \leq \text{scl}_{\lambda^2 \cdot \beta}^{p,q} .$$

Now by the above lemma, it holds for $\epsilon > 0$ small:

$$\text{scl}_{\lambda^2 \cdot \beta}^{p,q}(\epsilon) \leq \text{scl}_{\lambda \cdot \beta}^{p,q}(\epsilon^\lambda) \leq \left(\text{scl}_\beta^{p,q}(\epsilon^\lambda) \right)^\lambda \quad \text{and} \quad \text{scl}_{\lambda^2 \cdot \beta}^{p,q}(\epsilon) \leq \left(\text{scl}_\beta^{p,q}(\epsilon) \right)^{\lambda^2} .$$

Thus it comes:

$$\frac{\text{scl}_\alpha^{p,q}(\epsilon)}{\text{scl}_\beta^{p,q}(\epsilon^\lambda)} \leq \left(\text{scl}_\beta^{p,q}(\epsilon^\lambda) \right)^{\lambda-1} \xrightarrow{\epsilon \rightarrow 0} 0 \quad \text{and} \quad \frac{\text{scl}_\alpha^{p,q}(\epsilon)}{\text{scl}_\beta^{p,q}(\epsilon)^\lambda} \leq \left(\text{scl}_\beta^{p,q}(\epsilon) \right)^{\lambda(\lambda-1)} \xrightarrow{\epsilon \rightarrow 0} 0 ;$$

which allows to conclude the proof of the proposition. It remains to show the above lemma. First observe the following:

Fact 2.7. For every $\nu > 1$, for every $d \geq 1$ and for $y > 0$ great enough, it holds:

$$\log^{\circ d}(y^\nu) \leq \nu \cdot \log^{\circ d}(y) .$$

Proof. We prove this fact recursively on $d \geq 1$. For $d = 1$ note that for every $y > 0$, we have $\log(y^\nu) = \nu \cdot \log y$. Assume then that the inequality holds for $d \geq 1$, then for y great enough, it holds:

$$\log^{\circ(d+1)}(y^\nu) = \log(\log^{\circ d}(y^\nu)) \leq \log(\nu \log^{\circ d} y) = \log \nu + \log^{\circ(d+1)} y$$

Then, since $\nu > 1$, for y great enough we have $\log \nu + \log^{\circ(d+1)} y \leq \nu \cdot \log^{\circ(d+1)} y$ which allows to conclude. \square

Using **Fact 2.7** with $d = q$ and $y = \epsilon^{-1}$ for $\epsilon > 0$ small gives:

$$\gamma \cdot \log^{\circ q}(\epsilon^{-\nu}) \leq \lambda \cdot \gamma \cdot \log^{\circ q}(\epsilon^{-1}) .$$

Since $t \mapsto \exp^{\circ p}(t)^{-1}$ is decreasing, it comes:

$$\text{scl}_{\nu \cdot \gamma}^{p,q}(\epsilon) \leq \text{scl}_{\gamma}^{p,q}(\epsilon^\nu) ,$$

which gives the first inequality in the lemma.

Moreover by **Fact 2.7** with $d = p$ and y great enough, it holds:

$$\log^{\circ p}(y^\nu) \leq \nu \cdot \log^{\circ p}(y) .$$

Applying $\exp^{\circ p}$ to both sides gives:

$$y^\nu \leq \exp^{\circ p}(\nu \cdot \log^{\circ p}(y)) .$$

Now with $y^{-1} = \text{scl}_{\gamma}(\epsilon)$, we have:

$$\exp^{\circ p}(\nu \cdot \log^{\circ p}(y)) = (\text{scl}_{\nu \cdot \gamma}(\epsilon))^{-1} .$$

From there we obtain:

$$\text{scl}_{\nu \cdot \gamma}(\epsilon) \leq \text{scl}_{\gamma}(\epsilon)^\nu ,$$

which is the remaining inequality in the lemma. \square

2.2 Box scales

As we introduced in Definition 1.5, lower and upper box scales of a metric space (X, d) are defined by:

$$\underline{\text{scl}}_B X = \sup \left\{ \alpha > 0 : \mathcal{N}_\epsilon(X) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} +\infty \right\} \quad \text{and} \quad \overline{\text{scl}}_B X = \inf \left\{ \alpha > 0 : \mathcal{N}_\epsilon(X) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} 0 \right\} ,$$

where the covering number $\mathcal{N}_\epsilon(X)$ is the minimal cardinality of a covering of X by balls with radius $\epsilon > 0$.

In general, the upper and lower box scales must not coincide, we give new examples for order in Example 4.6. Now we give a few properties of box scales that are well known in the specific case of dimension.

Fact 2.8. Let (X, d) be a metric space. The following properties hold true:

1. if $\underline{\text{scl}}_B(X) < +\infty$, then (X, d) is totally bounded,
2. for every subset $E \subset X$ it holds $\underline{\text{scl}}_B E \leq \underline{\text{scl}}_B X$ and $\overline{\text{scl}}_B E \leq \overline{\text{scl}}_B X$,

3. for every subset E of X it holds $\underline{\text{scl}}_B E = \underline{\text{scl}}_B \text{cl}(E)$ and $\overline{\text{scl}}_B E = \overline{\text{scl}}_B \text{cl}(E)$.

1. and 2. are straightforward. To see 3. it is enough to observe that $\mathcal{N}_\epsilon(E) \leq \mathcal{N}_\epsilon(\text{cl}(E)) \leq \mathcal{N}_{\epsilon/2}(E)$ for every $\epsilon > 0$.
Box scales are sometimes easier to compare with other scales using packing number:

Definition 2.9 (Packing number). For $\epsilon > 0$ let $\tilde{\mathcal{N}}_\epsilon(X)$ be the packing number of the metric space (X, d) . It is the maximum cardinality of an ϵ -separated set of points in X for the distance d :

$$\tilde{\mathcal{N}}_\epsilon(X) = \sup\{N \geq 0 : \exists x_1, \dots, x_N \in X, d(x_i, x_j) \geq \epsilon \text{ for every } 1 \leq i < j \leq N\}.$$

A well know comparison between packing and covering number is the following:

Lemma 2.10. Let (X, d) be a metric space. For every $\epsilon > 0$, it holds:

$$\tilde{\mathcal{N}}_{2\epsilon}(X) \leq \mathcal{N}_\epsilon(X) \leq \tilde{\mathcal{N}}_\epsilon(X).$$

In virtue of basic properties of scalings, the covering number can be replaced by the packing number in the definitions of box scales:

Lemma 2.11. Let (X, d) be a metric space and scl a scaling, then box scales of X can be written as:

$$\underline{\text{scl}}_B X = \sup\left\{\alpha > 0 : \tilde{\mathcal{N}}_\epsilon(X) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} +\infty\right\} \quad \text{and} \quad \overline{\text{scl}}_B X = \inf\left\{\alpha > 0 : \tilde{\mathcal{N}}_\epsilon(X) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} 0\right\}.$$

Proof. Since for any $\epsilon > 0$ it holds by Lemma 2.10:

$$\tilde{\mathcal{N}}_{2\epsilon}(X) \leq \mathcal{N}_\epsilon(X) \leq \tilde{\mathcal{N}}_\epsilon(X),$$

we obtain the sought result by Lemma 2.2. □

Remark 2.12. Another property for the scaling $\text{scl}^{p,q}$ from Proposition 2.4, with $p, q \geq 1$, is that the upper and lower box scales for a metric space (X, d) can be written as:

$$\underline{\text{scl}}_B^{p,q}(X) = \liminf_{\epsilon \rightarrow 0} \frac{\log^{op}(\mathcal{N}_\epsilon(X))}{\log^{oq}(\epsilon^{-1})}$$

and

$$\overline{\text{scl}}_B^{p,q}(X) = \limsup_{\epsilon \rightarrow 0} \frac{\log^{op}(\mathcal{N}_\epsilon(X))}{\log^{oq}(\epsilon^{-1})}.$$

In particular, for dimension and order:

$$\underline{\text{dim}}_B(X) = \liminf_{\epsilon \rightarrow 0} \frac{\log(\mathcal{N}_\epsilon(X))}{\log(\epsilon^{-1})}, \quad \overline{\text{dim}}_B(X) = \limsup_{\epsilon \rightarrow 0} \frac{\log(\mathcal{N}_\epsilon(X))}{\log(\epsilon^{-1})}.$$

and

$$\underline{\text{ord}}_B(X) = \liminf_{\epsilon \rightarrow 0} \frac{\log \log(\mathcal{N}_\epsilon(X))}{\log(\epsilon^{-1})}, \quad \overline{\text{ord}}_B(X) = \limsup_{\epsilon \rightarrow 0} \frac{\log \log(\mathcal{N}_\epsilon(X))}{\log(\epsilon^{-1})}.$$

The above equalities coincide with the most usual definitions of box dimensions and orders.

2.3 Hausdorff scales

The definition of Hausdorff scales, generalizing Hausdorff dimension, is introduced here using the definition of Hausdorff outer measure as given by Tricot in [Tri82]. We still consider a metric space (X, d) . Given an increasing function $\phi \in C(\mathbb{R}_+^*, \mathbb{R}_+^*)$, such that $\phi(\epsilon) \rightarrow 0$ when $\epsilon \rightarrow 0$, we recall:

$$\mathcal{H}_\epsilon^\phi(X) := \inf_{J \text{ countable set}} \left\{ \sum_{j \in J} \phi(|B_j|) : X = \bigcup_{j \in J} B_j, \forall j \in J : |B_j| \leq \epsilon \right\},$$

where $|B|$ is the radius of a ball $B \subset X$. A countable family $(B_j)_{j \in J}$ of balls with radius at most $\epsilon > 0$ such that $X = \bigcup_{j \in J} B_j$ will be called an ϵ -cover of X .² Since the set of ϵ -cover is not decreasing for inclusion as ϵ decreases to 0, the following limit does exist:

$$\mathcal{H}^\phi(X) := \lim_{\epsilon \rightarrow 0} \mathcal{H}_\epsilon^\phi(X).$$

Now replacing (X, d) in the previous definitions by any subset E of X endowed with the same metric d , we observe that \mathcal{H}^ϕ defines an outer-measure on X . We now introduce the following:

Definition 2.13 (Hausdorff scale). *The Hausdorff scale of a metric space (X, d) is defined by:*

$$\text{scl}_H X = \sup \{ \alpha > 0 : \mathcal{H}^{\text{scl}_\alpha}(X) = +\infty \} = \inf \{ \alpha > 0 : \mathcal{H}^{\text{scl}_\alpha}(X) = 0 \}.$$

Note that the above definition gives us two quantities that are a priori not equal. However, the mild assumptions in the definition of scaling allow to verify that they indeed coincide and allows to use the machinery of Hausdorff outer measure to define metric invariants generalizing Hausdorff dimension.

Proof of the equality in Definition 2.13. It is clear from definition that $\alpha \mapsto \mathcal{H}^{\text{scl}_\alpha}(X)$ is non-increasing. It is then enough to check that if there exists $\alpha > 0$ such that $0 < \mathcal{H}^{\text{scl}_\alpha}(X) < +\infty$ then, for any positive $\delta < \alpha$, it holds:

$$\mathcal{H}^{\text{scl}_{\alpha+\delta}}(X) = 0 \quad \text{and} \quad \mathcal{H}^{\text{scl}_{\alpha-\delta}}(X) = +\infty.$$

Let us fix $\eta > 0$, by Definition 1.2, for $\epsilon > 0$ small it holds:

$$\text{scl}_{\alpha+\delta}(\epsilon) \leq \eta \cdot \text{scl}_\alpha(\epsilon) \quad \text{and} \quad \text{scl}_\alpha(\epsilon) \leq \eta \cdot \text{scl}_{\alpha-\delta}(\epsilon).$$

Since ϵ is small, it holds:

$$0 < \frac{1}{2} \mathcal{H}^{\text{scl}_\alpha}(X) \leq \mathcal{H}_\epsilon^{\text{scl}_\alpha}(X) \leq \mathcal{H}^{\text{scl}_\alpha}(X) < +\infty.$$

Given $(B_j)_{j \in J}$ an ϵ -cover of X , the following holds:

$$\frac{1}{2} \mathcal{H}^{\text{scl}_\alpha}(X) \leq \mathcal{H}_\epsilon^{\text{scl}_\alpha}(X) \leq \sum_{j \in J} \text{scl}_\alpha(|B_j|),$$

and then:

$$\frac{1}{2\eta} \mathcal{H}^{\text{scl}_\alpha}(X) \leq \frac{1}{\eta} \sum_{j \in J} \text{scl}_\alpha(|B_j|) \leq \sum_{j \in J} \text{scl}_{\alpha-\delta}(|B_j|).$$

²Note that the historical construction of the Hausdorff measures uses subsets of X with diameter at most ϵ instead of the balls with radius at most ϵ . However both these constructions lead to the same definitions of Hausdorff scales.

Since this holds for every ϵ -cover, the latter inequality leads to:

$$\frac{1}{2\eta} \mathcal{H}^{\text{scl}_\alpha}(X) \leq \mathcal{H}_\epsilon^{\text{scl}_{\alpha-\delta}}(X) ,$$

and so:

$$\frac{1}{2\eta} \mathcal{H}^{\text{scl}_\alpha}(X) \leq \mathcal{H}^{\text{scl}_{\alpha-\delta}}(X) .$$

On the other side, there exists an ϵ -cover $(B_j)_{j \in J}$ of E such that:

$$\sum_{j \in J} \text{scl}_\alpha(|B_j|) \leq 2\mathcal{H}_\epsilon^{\text{scl}_\alpha}(X) .$$

Now since $\mathcal{H}_\epsilon^{\text{scl}_\alpha}(X) \leq \mathcal{H}^{\text{scl}_\alpha}(X)$, this leads to:

$$\sum_{j \in J} \text{scl}_{\alpha+\delta}(|B_j|) \leq \eta \cdot \sum_j \text{scl}_\alpha(|B_j|) \leq 2\eta \cdot \mathcal{H}^{\text{scl}_\alpha}(X) .$$

From there:

$$\mathcal{H}_\epsilon^{\text{scl}_{\alpha+\delta}}(X) \leq 2\eta \cdot \mathcal{H}^{\text{scl}_\alpha}(X) ,$$

and this holds for every small ϵ . We have just shown:

$$\frac{1}{2\eta} \mathcal{H}^{\text{scl}_\alpha}(X) \leq \mathcal{H}^{\text{scl}_{\alpha-\delta}}(X) \quad \text{and} \quad \mathcal{H}_\epsilon^{\text{scl}_{\alpha+\delta}}(X) \leq 2\eta \cdot \mathcal{H}^{\text{scl}_\alpha}(X) .$$

Since η can be arbitrarily close to 0, it follows that $\mathcal{H}^{\text{scl}_{\alpha-\delta}}(X) = +\infty$ and $\mathcal{H}^{\text{scl}_{\alpha+\delta}}(X) = 0$, which concludes the proof. \square

As box scales, Hausdorff scales are increasing for inclusion. We show a stronger property of Hausdorff scales in Lemma 2.20.

2.4 Packing scales

2.4.1 Packing scales through modified box scales

The original construction of packing dimension relies on the packing measure introduced by Tricot in [Tri82]. We first define packing scales by modifying upper box scales and we show then later how it is related to packing measures.

Definition 2.14 (Packing scale). *Let (X, d) be a metric space and scl a scaling. The packing scale of X is defined by:*

$$\text{scl}_P X = \inf \left\{ \sup_{n \geq 1} \overline{\text{scl}}_B E_n : (E_n)_{n \geq 1} \subset X^{\mathbb{N}} \text{ s.t. } \bigcup_{n \geq 1} E_n = X \right\} .$$

The following comes directly from definition of packing scale:

Proposition 2.15. *Let (X, d) be a metric space and let scl be a scaling. It holds:*

$$\text{scl}_P X \leq \overline{\text{scl}}_B X .$$

2.4.2 Packing measures

In this paragraph we show the relationship between packing measures and packing scales. Let us first recall a few definitions.

Given $\epsilon > 0$, an ϵ -*pack* of a metric space (X, d) is a countable collection of disjoint balls of X with radii at most ϵ . As for Hausdorff outer measure, consider $\phi : \mathbb{R}_+^* \rightarrow \mathbb{R}_+^*$ an increasing function such that $\phi(\epsilon) \rightarrow 0$ when $\epsilon \rightarrow 0$. For $\epsilon > 0$, put:

$$\mathcal{P}_\epsilon^\phi(X) := \sup \left\{ \sum_{i \in I} \phi(|B_i|) : (B_i)_{i \in I} \text{ is an } \epsilon\text{-pack of } X \right\}.$$

Since $\mathcal{P}_\epsilon^\phi(X)$ is non-increasing when ϵ decreases to 0, the following quantity is well defined:

$$\mathcal{P}_0^\phi(X) := \lim_{\epsilon \rightarrow 0} \mathcal{P}_\epsilon^\phi(X).$$

The idea of Tricot is to build an outer measure from this quantity:

Definition 2.16 (Packing measure). *For every subset E of X endowed with the same metric d , the packing ϕ -measure of E is defined by:*

$$\mathcal{P}^\phi(E) = \inf \left\{ \sum_{n \geq 1} \mathcal{P}_0^\phi(E_n) : E = \bigcup_{n \geq 1} E_n \right\}.$$

Note that \mathcal{P}^ϕ is an outer-measure on X and can eventually be infinite or null. The following shows the equivalence of Tricot's counterpart definition of the packing scale; this will be useful to show the equality between upper local scale and packing scale of a measure given by **Theorem C** eq. (c&g).

Proposition 2.17. *The packing scale of a metric space (X, d) verifies:*

$$\sup \{ \alpha > 0 : \mathcal{P}^{\text{scl}_\alpha}(X) = +\infty \} = \text{scl}_P X = \inf \{ \alpha > 0 : \mathcal{P}^{\text{scl}_\alpha}(X) = 0 \}.$$

Proof. Let $(E_n)_{n \geq 1}$ be a family of subsets of X . Since each map $\alpha \mapsto \mathcal{P}_0^{\text{scl}_\alpha}(E_n)$ is not increasing and non negative, we have:

$$(2.1) \quad \inf \left\{ \alpha > 0 : \sum_{n \geq 1} \mathcal{P}_0^{\text{scl}_\alpha}(E_n) = 0 \right\} = \sup_{n \geq 1} \inf \left\{ \alpha > 0 : \mathcal{P}_0^{\text{scl}_\alpha}(E_n) = 0 \right\}.$$

We show below the following:

Lemma 2.18. *Given $\alpha > 0$, if $\mathcal{P}_0^{\text{scl}_\alpha}(E)$ is finite, then for every $\delta \in (0, \alpha)$, it holds:*

$$\mathcal{P}_0^{\text{scl}_{\alpha+\delta}}(E) = 0 \quad \text{and} \quad \mathcal{P}_0^{\text{scl}_{\alpha-\delta}}(E) = +\infty.$$

The right hand side equality of the latter lemma implies:

$$(2.2) \quad \sup \left\{ \alpha > 0 : \sum_{n \geq 1} \mathcal{P}_0^{\text{scl}_\alpha}(E_n) = +\infty \right\} = \sup_{n \geq 1} \sup \left\{ \alpha > 0 : \mathcal{P}_0^{\text{scl}_\alpha}(E_n) = +\infty \right\}.$$

We now compare the right hand term using the following shown below:

Lemma 2.19. For every $E \subset X$, it holds:

$$\sup \left\{ \alpha > 0 : \mathcal{P}_0^{\text{scl}_\alpha}(E) = +\infty \right\} = \overline{\text{scl}}_B E = \inf \left\{ \alpha > 0 : \mathcal{P}_0^{\text{scl}_\alpha}(E) = 0 \right\} .$$

Consequently by Eqs. (2.1) and (2.2) and Lemmas 2.18 and 2.19:

$$\sup \left\{ \alpha > 0 : \sum_{n \geq 1} \mathcal{P}_0^{\text{scl}_\alpha}(E_n) = +\infty \right\} = \sup_{n \geq 1} \overline{\text{scl}}_B E_n = \inf \left\{ \alpha > 0 : \sum_{n \geq 1} \mathcal{P}_0^{\text{scl}_\alpha}(E_n) = 0 \right\} .$$

Taking the infimum over families $(E_n)_{n \geq 1}$ which covers X we obtain the sought result. \square

Proof of Lemma 2.18. Given $\eta > 0$, by Definition 1.2 of scaling, for $\epsilon > 0$ small enough, it holds:

$$\text{scl}_{\alpha+\delta}(\epsilon) \leq \eta \cdot \text{scl}_\alpha(\epsilon) \quad \text{and} \quad \text{scl}_\alpha(\epsilon) \leq \eta \cdot \text{scl}_{\alpha-\delta}(\epsilon) .$$

Moreover there exists $(B_j)_{j \geq 1}$ an ϵ -pack of E such that:

$$\frac{1}{2} \mathcal{P}_\epsilon^{\text{scl}_\alpha}(E) \leq \sum_{j \geq 1} \text{scl}_\alpha(|B_j|) .$$

Combining the two inequalities above leads to:

$$\frac{1}{2} \mathcal{P}_\epsilon^{\text{scl}_\alpha}(E) \leq \eta^{-1} \cdot \sum_{j \geq 1} \text{scl}_{\alpha-\delta}(|B_j|) \leq \mathcal{P}_\epsilon^{\text{scl}_{\alpha-\delta}}(E) .$$

Taking the limit when ϵ tends to 0 gives $\frac{1}{2} \mathcal{P}_0^{\text{scl}_\alpha}(E) \leq \mathcal{P}_0^{\text{scl}_{\alpha-\delta}}(E)$. On the other side consider $(B_j)_{j \geq 1}$ an ϵ -pack of E . It holds:

$$\sum_{j \geq 1} \text{scl}_\alpha(|B_j|) \leq \mathcal{P}_\epsilon^{\text{scl}_\alpha}(E) \leq \mathcal{P}_0^{\text{scl}_\alpha}(E) ,$$

moreover it holds:

$$\eta \cdot \sum_{j \geq 1} \text{scl}_{\alpha+\delta}(|B_j|) \leq \sum_{j \geq 1} \text{scl}_\alpha(|B_j|) .$$

Since this holds true for any ϵ -cover and $\epsilon > 0$ arbitrary small, it follows:

$$\mathcal{P}_0^{\text{scl}_{\alpha+\delta}}(E) \leq \eta \cdot \mathcal{P}_0^{\text{scl}_\alpha}(E) .$$

By taking η arbitrarily small, it comes $\mathcal{P}_0^{\text{scl}_{\alpha-\delta}}(E) = +\infty$ and $\mathcal{P}_0^{\text{scl}_{\alpha+\delta}}(E) = 0$. \square

Proof of Lemma 2.19. By Lemma 2.18, it suffices to show that:

$$(2.3) \quad \sup \left\{ \alpha > 0 : \mathcal{P}_0^{\text{scl}_\alpha}(E) = +\infty \right\} \leq \overline{\text{scl}}_B E \leq \inf \left\{ \alpha > 0 : \mathcal{P}_0^{\text{scl}_\alpha}(E) = 0 \right\} ,$$

Consider $\alpha > 0$ such that $\mathcal{P}_0^{\text{scl}_\alpha}(E) = 0$. Then for $\epsilon > 0$ sufficiently small it holds $\mathcal{P}_\epsilon^{\text{scl}_\alpha}(E) \leq 1$. In particular the packing number (see def. 2.9) satisfies $\tilde{\mathcal{N}}_\epsilon(E) \cdot \text{scl}_\alpha(\epsilon) < 1$. Taking the limit when ϵ tends to 0 leads to $\overline{\text{scl}}_B E \leq \alpha$ by Lemma 2.11. This proves the right hand side of Eq. (2.3).

To show the left hand side inequality, it suffices to show that $\overline{\text{scl}}_B E$ is at least α for every $\alpha > 0$ such that $\mathcal{P}_0^{\text{scl}_\alpha}(E) = +\infty$. For such an α , given $\epsilon > 0$, there exists an ϵ -pack $(B_j)_{j \geq 1}$ such that:

$$\sum_{j \geq 1} \text{scl}_\alpha(|B_j|) > 1 .$$

For $k \geq 1$ an integer, put:

$$n_k := \text{Card} \left\{ j \geq 1 : 2^{-(k+1)} \leq \text{scl}_\alpha^{-1}(|B_j|) < 2^{-k} \right\} .$$

Thus, since scl_α is not decreasing, it holds:

$$\sum_{k \geq 1} n_k \cdot 2^{-k} > 1 .$$

Since B_j has radius at most δ , we have $n_k = 0$ for any $k < -\log_2 \text{scl}_\alpha(\delta)$. Then for $\delta > 0$ small, there exists an integer $j \geq 2$ such that $n_j > j^{-2}2^j$. In fact, otherwise we would have:

$$\sum_{k \geq 1} n_k \cdot 2^{-k} \leq \sum_{k \geq 2} \frac{1}{k^2} < 1 ,$$

which contradicts the above inequality. Then E contains the centers of n_j disjoint balls with radii at least $\text{scl}_\alpha^{-1}(2^{-(j+1)})$, in particular:

$$\tilde{\mathcal{N}}_{\text{scl}_\alpha^{-1}(2^{-(j+1)})}(E) \geq n_j > j^{-2}2^j ,$$

and moreover:

$$j \geq -\log_2 \text{scl}_\alpha(\delta) .$$

Since this inequality holds true for δ arbitrarily small, there exists an increasing sequence of integers $(j_n)_{n \geq 1}$ such that:

$$\tilde{\mathcal{N}}_{\epsilon_n}(E) > j_n^{-2}2^{j_n} ,$$

with $\epsilon_n = \text{scl}_\alpha^{-1}(2^{-(j_n+1)})$. Let us consider a positive number $\beta < \alpha$, by Definition 1.2 of scaling, for $\lambda > 1$ close to 1, it holds:

$$\text{scl}_\beta(\epsilon) \cdot (\text{scl}_\alpha(\epsilon))^{-\lambda^{-1}} \xrightarrow{\epsilon \rightarrow 0} +\infty .$$

On the other hand, given a such $\lambda > 1$, for n large enough, it holds:

$$j_n^{-2}2^{j_n} \geq 2^{\lambda^{-1}(j_n+1)} ,$$

it follows:

$$\tilde{\mathcal{N}}_{\epsilon_n}(E) \geq \left(2^{-(j_n+1)} \right)^{-\lambda^{-1}} = (\text{scl}_\alpha(\epsilon_n))^{-\lambda^{-1}} .$$

Thus we finally have:

$$\text{scl}_\beta(\epsilon_n) \cdot \tilde{\mathcal{N}}_{\epsilon_n}(E) > \text{scl}_\beta(\epsilon) \cdot (\text{scl}_\alpha(\epsilon))^{-\lambda^{-1}} \xrightarrow{\epsilon \rightarrow 0} +\infty .$$

By Lemma 2.11 we deduce $\overline{\text{scl}}_B E \geq \alpha$. Since this holds true for β arbitrary close to α , it follows $\overline{\text{scl}}_B E \geq \alpha$. \square

The following is similar to the proof below Definition 3.1 that Hausdorff scales are well defined.

2.5 Properties and comparison of scales of metric spaces

We first give a few basic properties of scales that would allow to compare them. Since both packing and Hausdorff scales are defined via measures, they both are countable stable as shown in the following:

Lemma 2.20 (Countable stability). *Let (X, d) be a metric space. Let I be a countable set and $(E_i)_{i \in I}$ a covering of X , then for any scaling scl :*

$$\text{scl}_H X = \sup_{i \in I} \text{scl}_H E_i \quad \text{and} \quad \text{scl}_P X = \sup_{i \in I} \text{scl}_P E_i .$$

Proof. The equality on packing scales is clear by definition. Let us prove the equality on Hausdorff scales. By monotonicity of the Hausdorff measure, it holds $\text{scl}_H X \geq \sup_{i \in I} \text{scl}_H E_i$. For the reverse inequality, consider $\alpha > \sup_{i \in I} \text{scl}_H E_i$, then for any $i \in I$ it holds $\mathcal{H}^{\text{scl}_\alpha}(E_i) = 0$. Thus:

$$\mathcal{H}^{\text{scl}_\alpha}(X) \leq \sum_{i \in I} \mathcal{H}^{\text{scl}_\alpha}(E_i) = 0 ,$$

and then $\text{scl}_H X \leq \alpha$. Since this is true for any $\alpha > \sup_{i \in I} \text{scl}_H E_i$, the sought result comes. \square

Note that countable stability is not a property of box scales. To see that, it suffices to consider a countable dense subset of a metric space (X, d) with positive box scales. This is actually a basic know fact for the specific case of dimension that naturally still holds there.

The following lemma shows in particular that the above scales are bi-Lipschitz invariants.

Lemma 2.21. *Let (X, d) and (Y, d) be two metric spaces such that there exists a Lipschitz map $f : (X, d) \rightarrow (Y, d)$. Then for any scaling scl , the scales of $f(X)$ are at most the ones of X :*

$$\text{scl}_H f(X) \leq \text{scl}_H X; \quad \text{scl}_P f(X) \leq \text{scl}_P X; \quad \underline{\text{scl}}_B f(X) \leq \underline{\text{scl}}_B X; \quad \overline{\text{scl}}_B f(X) \leq \overline{\text{scl}}_B X .$$

Remark the above lemma holds even for scalings that have sub-polynomial behaviours. We prove this lemma below. As a direct application, we obtain the following:

Corollary 2.22. *Let (X, d) and (Y, d) be two metric spaces. Assume that there exists an embedding $g : (Y, \delta) \rightarrow (X, d)$ such that g^{-1} is Lipschitz on $g(X)$. Then for every scaling scl , the scales of Y are at most the ones of X :*

$$\text{scl}_H Y \leq \text{scl}_H X; \quad \text{scl}_P Y \leq \text{scl}_P X; \quad \underline{\text{scl}}_B Y \leq \underline{\text{scl}}_B X; \quad \overline{\text{scl}}_B Y \leq \overline{\text{scl}}_B X .$$

Proof. By Lemma 2.21 we have $\text{scl}_\bullet Y \leq \text{scl}_\bullet g(Y)$ for any $\text{scl}_\bullet \in \{\text{scl}_H, \text{scl}_P, \underline{\text{scl}}_B, \overline{\text{scl}}_B\}$. As $g(Y) \subset X$, we have also $\text{scl}_\bullet g(Y) \leq \text{scl}_\bullet X$. \square

Proof of Lemma 2.21. Let us fix $\epsilon > 0$. Suppose that f is K -Lipschitz for a constant $K > 0$. We first show the inequalities on box and packing scales. Consider a finite covering by a collection of balls $(B(x_j, \epsilon))_{1 \leq j \leq N}$ where $x_j \in X$ for any $1 \leq j \leq N$ and $N = \mathcal{N}_\epsilon(X)$. Since $X = \bigcup_{j=1}^N B(x_j, \epsilon)$, it comes:

$$f(X) \subset f \left(\bigcup_{j=1}^N B(x_j, \epsilon_j) \right) \subset \bigcup_{j=1}^N B(f(x_j), K \cdot \epsilon_j) .$$

Then $(B(f(x_j), K \cdot \epsilon))_{1 \leq j \leq N}$ is a covering by $K \cdot \epsilon$ -balls of $f(X)$. Then $\mathcal{N}_{K \cdot \epsilon}(f(X)) \leq \mathcal{N}_\epsilon(X)$ and all the inequalities on the box and packing scales are immediately deduced. Now for Hausdorff scales, consider a countable set J and $\{B(x_j, \epsilon_j) : j \in J\}$ an ϵ -cover of X . Then it comes:

$$f(X) \subset \bigcup_{j \in J} B(f(x_j), K \cdot \epsilon_j) .$$

For any $\alpha > \beta > 0$ and $\delta > 0$ small enough, by **Fact 2.1**, it holds:

$$\text{scl}_\alpha(\delta) \leq \text{scl}_\beta(K^{-1} \cdot \delta) .$$

Hence for ϵ small, it holds:

$$\mathcal{H}_{K \cdot \epsilon}^{\text{scl}_\alpha}(f(X)) \leq \sum_{j \in J} \text{scl}_\alpha(K \cdot \epsilon_j) \leq \sum_{j \in J} \text{scl}_\beta(\epsilon_j) .$$

As $\beta > \text{scl}_H X$, the ϵ -cover $(B(x_j, \epsilon_j))_{j \in J}$ can be chosen such that $\sum_{j \in J} \text{scl}_\beta(\epsilon_j)$ is arbitrary small. Consequently, it holds $\mathcal{H}_{K \cdot \epsilon}^{\text{scl}_\alpha}(f(X)) = 0$, and so $\text{scl}_H f(X) \leq \alpha$. As α is arbitrary close to $\text{scl}_H X$, it holds:

$$\text{scl}_H f(X) \leq \text{scl}_H X .$$

□

The end of this section consists of comparing the different scales introduced and prove **Theorem A**. We start by comparing the Hausdorff with lower box scales. The following proposition generalizes well known facts on dimension. See e.g. [Fal04][(3.17)].

Proposition 2.23. *Let (X, d) be a metric space and scl a scaling, its Hausdorff scale is at most its lower box scale:*

$$\text{scl}_H X \leq \underline{\text{scl}}_B X .$$

Proof. We can assume without any loss that (X, d) is totally bounded. If $\text{scl}_H X = 0$ the inequality obviously holds, thus consider a positive number $\alpha < \text{scl}_H X$. For $\delta > 0$ small enough, $\mathcal{H}_\delta^{\text{scl}_\alpha}(X) > 1$. Thus for every δ -cover $(B_j)_{1 \leq j \leq \mathcal{N}_\delta(X)}$, it holds:

$$1 < \sum_{1 \leq j \leq \mathcal{N}_\delta(X)} \text{scl}_\alpha(|B_j|) = \mathcal{N}_\delta(X) \cdot \text{scl}_\alpha(\delta) .$$

From there, it holds $\underline{\text{scl}}_B X \geq \alpha$. We conclude by taking α arbitrarily close to $\text{scl}_H X$. □

We have compared Hausdorff and packing scales with their corresponding box scales. It remains to compare each other with the following:

Proposition 2.24. *Let (X, d) be a metric space and scl a scaling. It holds:*

$$\text{scl}_H X \leq \text{scl}_P X .$$

Proof. By Lemma 2.20, it holds:

$$\text{scl}_H X = \inf_{\bigcup_{n \geq 1} E_n = X} \sup_{n \geq 1} \text{scl}_H E_n ,$$

where the infimum is taken over countable coverings of X . Moreover by Proposition 2.23, we have:

$$\text{scl}_H E \leq \underline{\text{scl}}_B E \leq \overline{\text{scl}}_B E ,$$

for any subset E of X . It follows then:

$$\text{scl}_H X \leq \inf_{\bigcup_{n \geq 1} E_n = X} \sup_{n \geq 1} \overline{\text{scl}}_B E_n = \text{scl}_P X .$$

□

For the sake of completeness we will resume:

Proof of Theorem A. Let (X, d) be a metric space and scl a scaling. By Proposition 2.23, Proposition 2.24 and Proposition 2.15, it holds respectively:

$$\text{scl}_H X \leq \underline{\text{scl}}_B X, \quad \text{scl}_H X \leq \text{scl}_P X \quad \text{and} \quad \text{scl}_H X \leq \overline{\text{scl}}_B X .$$

Now since $\underline{\text{scl}}_B X \leq \overline{\text{scl}}_B X$ obviously holds, we deduce the sought result:

$$\text{scl}_H X \leq \text{scl}_P X \leq \overline{\text{scl}}_B X \quad \text{and} \quad \text{scl}_H X \leq \underline{\text{scl}}_B X \leq \overline{\text{scl}}_B X .$$

□

3 Scales of measures

In this section we recall the different versions of scales of measures we introduced and show the inequalities and equalities comparing them. In particular we provide proofs of **Theorem B** and **Theorem C**. They generalize known facts of dimension theory to any scaling and moreover bring new comparisons (see Theorem 3.10) between quantization and box scales that were not shown yet for even for the case of dimension.

3.1 Hausdorff, packing and local scales of measures

Let us recall the definition of local scales. Let μ be a Borel measure on a metric space (X, d) and scl a scaling. The *lower and upper scales* of μ are the functions that map a point $x \in X$ to:

$$\underline{\text{scl}}_{\text{loc}} \mu(x) = \sup \left\{ \alpha > 0 : \frac{\mu(B(x, \epsilon))}{\text{scl}_\alpha(\epsilon)} \xrightarrow{\epsilon \rightarrow 0} 0 \right\} \quad \text{and} \quad \overline{\text{scl}}_{\text{loc}} \mu(x) = \inf \left\{ \alpha > 0 : \frac{\mu(B(x, \epsilon))}{\text{scl}_\alpha(\epsilon)} \xrightarrow{\epsilon \rightarrow 0} +\infty \right\} .$$

We shall compare local scales with the followings:

Definition 3.1 (Hausdorff scales of a measure). *Let scl be a scaling and μ a non-null Borel measure on a metric space (X, d) . We define the Hausdorff and *-Hausdorff scales of the measure μ by:*

$$\text{scl}_H \mu = \inf_{E \in \mathcal{B}} \{ \text{scl}_H E : \mu(E) > 0 \} \quad \text{and} \quad \text{scl}_H^* \mu = \inf_{E \in \mathcal{B}} \{ \text{scl}_H E : \mu(X \setminus E) = 0 \} ,$$

where \mathcal{B} is the set of Borel subsets of X .

Definition 3.2 (Packing scales of a measure). *Let scl be a scaling and μ a non-null Borel measure on a metric space (X, d) . We define the packing and *-packing scales of μ by:*

$$\text{scl}_P \mu = \inf_{E \in \mathcal{B}} \{ \text{scl}_P E : \mu(E) > 0 \} \quad \text{and} \quad \text{scl}_P^* \mu = \inf_{E \in \mathcal{B}} \{ \text{scl}_P E : \mu(X \setminus E) = 0 \} .$$

Remark 3.3. *In order to avoid excluding the null measure 0, we set $\text{scl}_H 0 = \text{scl}_H^* 0 = \text{scl}_P 0 = \text{scl}_P^* 0 = 0$.*

The lemma below will allow to compare local scales with the other scales of measures.

Lemma 3.4. *Let μ be a Borel measure on X . Then for any Borel subset F of X such that $\mu(F) > 0$, the restriction σ of μ to F verifies:*

$$\text{ess inf } \overline{\text{scl}}_{\text{loc}} \mu \leq \text{ess inf } \overline{\text{scl}}_{\text{loc}} \sigma \quad \text{and} \quad \text{ess inf } \underline{\text{scl}}_{\text{loc}} \mu \leq \text{ess inf } \underline{\text{scl}}_{\text{loc}} \sigma .$$

Moreover, if there exists $\alpha > 0$ such that $F \subset \{x \in X : \overline{\text{scl}}_{\text{loc}}\mu(x) > \alpha\}$, it holds then:

$$\text{ess inf } \overline{\text{scl}}_{\text{loc}}\sigma \geq \alpha ,$$

and similarly if $F \subset \{x \in X : \underline{\text{scl}}_{\text{loc}}\mu(x) > \alpha\}$, it holds:

$$\text{ess inf } \underline{\text{scl}}_{\text{loc}}\sigma \geq \alpha .$$

Proof. Consider a point $x \in X$, then for any $\epsilon > 0$, one has $\sigma(B(x, \epsilon)) \leq \mu(B(x, \epsilon))$, thus by definition of local scales:

$$\overline{\text{scl}}_{\text{loc}}\mu \leq \overline{\text{scl}}_{\text{loc}}\sigma \quad \text{and} \quad \underline{\text{scl}}_{\text{loc}}\mu \leq \underline{\text{scl}}_{\text{loc}}\sigma .$$

Now if there exists $\alpha > 0$ such that $F \subset \{x \in X : \overline{\text{scl}}_{\text{loc}}\mu(x) > \alpha\}$, as $\overline{\text{scl}}_{\text{loc}}\mu(x) \geq \alpha$ for μ -almost every x in F , it comes by the above inequality that $\overline{\text{scl}}_{\text{loc}}\sigma(x) \geq \alpha$ for μ -almost every x in F , and thus for σ -almost every $x \in X$. It follows $\text{ess inf } \overline{\text{scl}}_{\text{loc}}\sigma \geq \alpha$. And the same holds for lower local scales. \square

The following is a first step in the proof of **Theorem B**. We prove this lemma later. We first use it to prove **Theorem C**.

Lemma 3.5. *Let (X, d) be a metric space and μ a Borel measure on X . Let scl be a scaling. The lower and upper local scales of μ are respectively not greater than the Hausdorff and packing scales of the space X :*

$$\text{ess sup } \underline{\text{scl}}_{\text{loc}}\mu \leq \text{scl}_H X \quad \text{and} \quad \text{ess sup } \overline{\text{scl}}_{\text{loc}}\mu \leq \text{scl}_P X .$$

Remark 3.6. *Note that in the above we can replace X by any subset of X with total mass, this leads to:*

$$\text{ess sup } \underline{\text{scl}}_{\text{loc}}\mu \leq \text{scl}_H^* \mu \quad \text{and} \quad \text{ess sup } \overline{\text{scl}}_{\text{loc}}\mu \leq \text{scl}_P^* \mu .$$

3.2 Quantization and box scales of measures

Let us first recall the definition of quantization scales. Let (X, d) be a metric space and μ a Borel measure on X . The *quantization number* \mathcal{Q}_μ of μ is the function that maps $\epsilon > 0$ to the minimal cardinality of a set of points that is on average ϵ -close to any point in X :

$$\mathcal{Q}_\mu(\epsilon) = \inf \left\{ N \geq 0 : \exists \{c_i\}_{i=1, \dots, N} \subset X, \int_X d(x, \{c_i\}_{1 \leq i \leq N}) d\mu(x) < \epsilon \right\} .$$

Then *lower and upper quantization scales* of μ for a given scaling scl are defined by:

$$\underline{\text{scl}}_Q \mu = \sup \left\{ \alpha > 0 : \mathcal{Q}_\mu(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} +\infty \right\} \quad \text{and} \quad \overline{\text{scl}}_Q \mu = \inf \left\{ \alpha > 0 : \mathcal{Q}_\mu(\epsilon) \cdot \text{scl}_\alpha(\epsilon) \xrightarrow{\epsilon \rightarrow 0} 0 \right\} .$$

Quantization scales of a measure are compared in **Theorem C** with box scales:

Definition 3.7 (Box scales of a measure). *Let scl be a scaling and μ a positive Borel measure on a metric space (X, d) . We define the lower box scale and the *-lower box scale of μ by:*

$$\underline{\text{scl}}_B \mu = \inf_{E \in \mathcal{B}} \{\underline{\text{scl}}_B E : \mu(E) > 0\} \quad \text{and} \quad \underline{\text{scl}}_B^* \mu = \inf_{E \in \mathcal{B}} \{\underline{\text{scl}}_B E : \mu(X \setminus E) = 0\} .$$

Similarly, we define the upper box scale and the *-upper box scale of μ by:

$$\overline{\text{scl}}_B \mu = \inf_{E \in \mathcal{B}} \{\overline{\text{scl}}_B E : \mu(E) > 0\} \quad \text{and} \quad \overline{\text{scl}}_B^* \mu = \inf_{E \in \mathcal{B}} \{\overline{\text{scl}}_B E : \mu(X \setminus E) = 0\} ,$$

where \mathcal{B} is the set of Borel subsets of X .

As for Hausdorff scales of measures we chose that all box scales of the null measure are equal to 0 as a convention. The following is straightforward:

Lemma 3.8. *Let (X, d) be a metric space and μ a Borel measure on X . Given scl a scaling, it holds:*

$$\underline{\text{scl}}_Q \mu \leq \underline{\text{scl}}_B^* \mu \quad \text{and} \quad \overline{\text{scl}}_Q \mu \leq \overline{\text{scl}}_B^* \mu .$$

Proof. We can assume without loss of generality that $\overline{\text{scl}}_B^* \mu$ and $\underline{\text{scl}}_B^* \mu$ are finite. Let E be a Borel set with total mass such that $\overline{\text{scl}}_B E$ is finite, then E is totally bounded by **Fact 2.8**. Now for $\epsilon > 0$, consider a covering by ϵ -balls centered at some points x_1, \dots, x_N in E . Since $\mu(X \setminus E) = 0$, it comes:

$$\int_X d(x, \{x_i\}_{1 \leq i \leq N}) d\mu(x) = \int_E d(x, \{x_i\}_{1 \leq i \leq N}) d\mu(x) < \epsilon .$$

Thus $\mathcal{Q}_\mu(\epsilon) \leq \mathcal{N}_\epsilon(E)$, and by **Lemma 2.2**:

$$\underline{\text{scl}}_Q \mu \leq \underline{\text{scl}}_B E \quad \text{and} \quad \overline{\text{scl}}_Q \mu \leq \overline{\text{scl}}_B E .$$

Since this holds true for any Borel set E with total mass, the sought results comes. \square

The following lemma will allow to compare quantization scales with box scales.

Lemma 3.9. *Let μ be a Borel measure on (X, d) such that $\mathcal{Q}_\mu(\epsilon) < +\infty$ for any $\epsilon > 0$. Let us fix $\epsilon > 0$ and an integer $N \geq \mathcal{Q}_\mu(\epsilon)$. Thus consider $x_1, \dots, x_N \in X$ such that:*

$$\int_X d(x, \{x_i\}_{1 \leq i \leq N}) d\mu(x) < \epsilon .$$

For any $r > 0$, with $E_r := \bigcup_{i=1}^N B(x_i, r)$, it holds:

$$\mu(X \setminus E_r) < \frac{\epsilon}{r} .$$

Proof. Since $X \setminus E_r$, the complement of E_r in X is the set of points with distance at most r from the set $\{x_1, \dots, x_n\}$, it holds:

$$r \cdot \mu(X \setminus E_r) \leq \int_{X \setminus E_r} d(x, \{x_i\}_{1 \leq i \leq N}) d\mu(x) < \epsilon ,$$

which gives the sought result by dividing both sides by r . \square

The following result exhibits the relationship between quantization scales and box scales. As far as we know, this result has not yet have been proved even for the specific case of dimension. It is a key element in the answer to **Problem 1.15**.

Theorem 3.10. *Let μ be a non null Borel measure on a metric space (X, d) . For any scaling scl, there exists a Borel set F with positive mass such that:*

$$\underline{\text{scl}}_B F \leq \underline{\text{scl}}_Q \mu \quad \text{and} \quad \overline{\text{scl}}_B F \leq \overline{\text{scl}}_Q \mu .$$

In particular, it holds:

$$\underline{\text{scl}}_B \mu \leq \underline{\text{scl}}_Q \mu \quad \text{and} \quad \overline{\text{scl}}_B \mu \leq \overline{\text{scl}}_Q \mu .$$

Proof. If $\mathcal{Q}_\mu(\epsilon)$ is not finite for any $\epsilon > 0$, then $F = X$ satisfies the sought properties. Let us suppose now that the quantization number of μ is finite. Given an integer $n \geq 0$, we set $\epsilon_n := \exp(-n)$ and $r_n := n^2 \cdot \exp(-n) = n^2 \cdot \epsilon_n$. We also consider a finite set of points $C_n \subset X$ that contains exactly $\mathcal{Q}_\mu(\epsilon_n)$ points and such that:

$$\int_X d(x, C_n) d\mu(x) < \epsilon_n .$$

We can then consider the following set:

$$E_n := \bigcup_{c \in C_n} B(c, r_n) ,$$

then by Lemma 3.9, it holds:

$$\mu(X \setminus E_n) < \frac{\epsilon_n}{r_n} = \frac{1}{n^2} .$$

Thus, it holds:

$$\sum_{n \geq 0} \mu(X \setminus E_n) < +\infty .$$

By Borell-Cantelli lemma, we obtain:

$$\mu \left(\bigcup_{m \geq 0} \bigcap_{n \geq m} E_n \right) = \mu(X) > 0 .$$

Thus there exists an integer $m \geq 0$ such that $\mu \left(\bigcap_{n \geq m} E_n \right) > 0$. We fix such an integer m and set $F := \bigcap_{n \geq m} E_n$. It remains to check that $\underline{\text{scl}}_B F \leq \underline{\text{scl}}_Q \mu$ and $\overline{\text{scl}}_B F \leq \overline{\text{scl}}_Q \mu$. By definition, one has $F \subset E_n$ for any $n \geq m$. Then since $F \subset E_n = \bigcup_{c \in C_n} B(c, r_n)$, it holds:

$$\mathcal{N}_{r_n}(F) \leq \text{Card } C_n = \mathcal{Q}_\mu(\epsilon_n)$$

Since this holds true for any n greater than m , and since $\log r_n \sim \log \epsilon_n = -n$, we finally have by Lemmas 2.2 and 2.3 that:

$$\underline{\text{scl}}_B F \leq \underline{\text{scl}}_Q \mu \quad \text{and} \quad \overline{\text{scl}}_B F \leq \overline{\text{scl}}_Q \mu .$$

□

As a corollary of the proof observe the following:

Remark 3.11. *If μ is positive finite, then by taking m large in the above we can have $\mu(F)$ arbitrarily close to $\mu(X)$.*

3.3 Comparison between local and global scales of measures and proof of Theorem C

By the latter theorem, to finish the proof of **Theorem C**, it remains only to show:

Theorem 3.12. *Let (X, d) be a separable metric space and μ a finite Borel measure on X . Let scl be a scaling. It holds:*

$$\text{ess sup } \underline{\text{scl}}_{\text{loc}} \mu \leq \underline{\text{scl}}_Q \mu \quad \text{and} \quad \text{ess sup } \overline{\text{scl}}_{\text{loc}} \mu \leq \overline{\text{scl}}_Q \mu .$$

Proof. We can suppose without any loss of generality that there exists $\alpha < \text{ess sup } \underline{\text{scl}}_{\text{loc}} \mu$ and $\beta < \text{ess sup } \overline{\text{scl}}_{\text{loc}} \mu$. We now set $E := \{x \in X : \underline{\text{scl}}_{\text{loc}} \mu(x) > \alpha \text{ and } \overline{\text{scl}}_{\text{loc}} \mu(x) > \beta\}$. By definition of essential suprema, we have

$\mu(E) > 0$. Thus the restriction σ of μ to E is a positive measure. Thus by Lemma 3.4 one has $\text{ess inf } \underline{\text{scl}}_{\text{loc}}\sigma \geq \alpha$ and $\text{ess inf } \overline{\text{scl}}_{\text{loc}}\sigma \geq \beta$. Moreover by Theorem 3.10, there is a Borel set $F \subset E$ with $\sigma(F) > 0$ and such that:

$$\underline{\text{scl}}_B F \leq \underline{\text{scl}}_Q \sigma \leq \underline{\text{scl}}_Q \mu \quad \text{and} \quad \overline{\text{scl}}_B F \leq \overline{\text{scl}}_Q \sigma \leq \overline{\text{scl}}_Q \mu .$$

Yet by Proposition 2.23 and Proposition 2.15, it holds respectively:

$$\text{scl}_H F \leq \underline{\text{scl}}_B F \quad \text{and} \quad \text{scl}_P F \leq \overline{\text{scl}}_B F .$$

Now, by setting τ the restriction of μ to F , Lemma 3.4 also gives:

$$\alpha \leq \text{ess inf } \underline{\text{scl}}_{\text{loc}}\sigma \leq \text{ess inf } \underline{\text{scl}}_{\text{loc}}\tau \quad \text{and} \quad \beta \leq \text{ess inf } \overline{\text{scl}}_{\text{loc}}\sigma \leq \text{ess inf } \overline{\text{scl}}_{\text{loc}}\tau .$$

By Lemma 3.5, it holds:

$$\text{ess inf } \underline{\text{scl}}_{\text{loc}}\tau \leq \text{scl}_H F \quad \text{and} \quad \text{ess inf } \overline{\text{scl}}_{\text{loc}}\tau \leq \text{scl}_P F .$$

Finally, combining all the above inequalities leads to:

$$\alpha \leq \text{ess inf } \underline{\text{scl}}_{\text{loc}}\tau \leq \text{scl}_H F \leq \underline{\text{scl}}_B F \leq \underline{\text{scl}}_Q \mu$$

and

$$\beta \leq \text{ess inf } \overline{\text{scl}}_{\text{loc}}\tau \leq \text{scl}_P F \leq \overline{\text{scl}}_B F \leq \overline{\text{scl}}_Q \mu .$$

Since this holds true for any α and β arbitrarily close to $\text{ess sup } \underline{\text{scl}}_{\text{loc}}\mu$ and $\text{ess sup } \overline{\text{scl}}_{\text{loc}}\mu$ we have the sought results. \square

We shall now prove:

Proof of Theorem C. By Theorem 3.10 and Lemma 3.8, it holds:

$$\underline{\text{scl}}_B \mu \leq \underline{\text{scl}}_Q \mu \leq \underline{\text{scl}}_B^* \mu \quad \text{and} \quad \overline{\text{scl}}_B \mu \leq \overline{\text{scl}}_Q \mu \leq \overline{\text{scl}}_B^* \mu .$$

By Theorem 3.12 it holds:

$$\text{ess sup } \underline{\text{scl}}_{\text{loc}}\mu \leq \underline{\text{scl}}_Q \mu \quad \text{and} \quad \text{ess sup } \overline{\text{scl}}_{\text{loc}}\mu \leq \overline{\text{scl}}_Q \mu .$$

Thus it remains only to show:

$$\text{ess inf } \underline{\text{scl}}_{\text{loc}}\mu \leq \underline{\text{scl}}_B \mu \quad \text{and} \quad \text{ess inf } \overline{\text{scl}}_{\text{loc}}\mu \leq \overline{\text{scl}}_B \mu .$$

Given E a subset of X with positive mass, we set σ the restriction of μ to E . By Lemma 3.4, it holds:

$$\text{ess inf } \underline{\text{scl}}_{\text{loc}}\mu \leq \text{ess inf } \underline{\text{scl}}_{\text{loc}}\sigma \quad \text{and} \quad \text{ess inf } \overline{\text{scl}}_{\text{loc}}\mu \leq \text{ess inf } \overline{\text{scl}}_{\text{loc}}\sigma .$$

By Theorem 3.12 it holds:

$$\text{ess sup } \underline{\text{scl}}_{\text{loc}}\sigma \leq \underline{\text{scl}}_Q \sigma \quad \text{and} \quad \text{ess sup } \overline{\text{scl}}_{\text{loc}}\sigma \leq \overline{\text{scl}}_Q \sigma .$$

Moreover by Lemma 3.8:

$$\underline{\text{scl}}_Q \sigma \leq \underline{\text{scl}}_B E \quad \text{and} \quad \overline{\text{scl}}_Q \sigma \leq \overline{\text{scl}}_B E .$$

Combining all of the above leads to:

$$\text{ess inf } \underline{\text{scl}}_{\text{loc}}\mu \leq \underline{\text{scl}}_B E \quad \text{and} \quad \text{ess inf } \overline{\text{scl}}_{\text{loc}}\mu \leq \overline{\text{scl}}_B E .$$

Taking the infima over such subsets $E \subset X$ with positive mass leads to the sought result. \square

3.4 Proof of Theorem B

This subsection contains the proof of **Theorem B**, we recall its statement below. We use Vitali's lemma [Vit08] to compare local scales with Hausdorff and packing scales as Fan and Tamashiro did in their proof for the case of dimension.

Lemma 3.13 (Vitali). *Let (X, d) be a separable metric space. Given $\delta > 0$, \mathcal{B} a family of open balls in X with radii at most δ and F the union of these balls. There exists a countable set J and a δ -pack $(B(x_j, r_j))_{j \in J} \subset \mathcal{B}$ of F such that:*

$$F \subset \bigcup_j B(x_j, 5r_j) .$$

We first prove Lemma 3.5 that was used to prove **Theorem C** in the previous paragraph.

Proof of Lemma 3.5. First we can assume that $\text{scl}_H X < +\infty$, $\text{scl}_P X < +\infty$, and that μ is not null, otherwise both inequalities immediately hold true. In particular, we can assume that X is separable.

LEFT HAND SIDE INEQUALITY: If $\text{ess sup } \underline{\text{scl}}_{\text{loc}} \mu = 0$ the inequality is obviously true. Suppose then that this quantity is positive and consider a positive $\alpha < \text{ess sup } \underline{\text{scl}}_{\text{loc}} \mu$. Thus, there exists $r_0 > 0$ such that the set:

$$A := \{x \in X : \mu(B(x, r)) \leq \text{scl}_\alpha(r), \forall r \in (0, r_0)\}$$

has positive measure. Consider $\delta \leq r_0$ and any δ -cover $(B_j)_{j \in J}$ of A . Then it holds:

$$0 < \mu(A) \leq \sum_{j \in J} \mu(B_j) \leq \sum_{j \in J} \text{scl}_\alpha(|B_j|) .$$

Since this holds true for any such cover, it follows:

$$0 < \mu(A) \leq \mathcal{H}_\delta^{\text{scl}_\alpha}(A) .$$

Taking δ arbitrarily close to 0 leads to:

$$0 < \mu(A) \leq \mathcal{H}^{\text{scl}_\alpha}(A) .$$

Finally since Hausdorff scale is non-decreasing for inclusion, it holds:

$$\text{scl}_H X \geq \text{scl}_H A \geq \alpha .$$

Note that since this holds true for any $\alpha < \text{ess sup } \underline{\text{scl}}_{\text{loc}} \mu$, we indeed have $\text{ess sup } \underline{\text{scl}}_{\text{loc}} \mu \leq \text{scl}_H(X)$.

RIGHT HAND SIDE INEQUALITY: Similarly, without any loss of generality we assume that there exists $0 < \alpha < \text{ess sup } \overline{\text{scl}}_{\text{loc}} \mu$ and put:

$$F = \{x \in X : \overline{\text{scl}}_{\text{loc}} \mu(x) > \alpha\} .$$

Let us fix a family of Borel subsets $(F_N)_{N \geq 1}$ of X such that $F = \bigcup_{N \geq 1} F_N$. For $0 < \beta < \alpha$, by **Fact 2.1**, there exists $\delta_0 > 0$ such that for any $r \leq \delta_0$, it holds:

$$\text{scl}_\alpha(5r) \leq \text{scl}_\beta(r) .$$

We fix $\delta \in (0, \delta_0)$ and an integer $N \geq 1$. For any x in F_N by Lemma 2.3 there exists an integer $n(x)$, minimal, such that $r(x) := \exp(-n(x)) \leq \delta$ and:

$$\mu(B(x, 5r(x))) \leq \text{scl}_\alpha(5r(x)) .$$

We now set:

$$\mathcal{F} = \{B(x, r(x)) : x \in F_N\} .$$

Thus by Vitali Lemma 3.13 there exists a countable set J and a δ -pack $\{B(x_j, r_j) : j \in J\} \subset \mathcal{F}$ of F_N such that $F_N \subset \bigcup_{j \in J} B(x_j, 5r_j)$. From there it holds:

$$\mu(F_N) \leq \sum_{j \in J} \mu(B(x_j, 5r_j)) \leq \sum_{j \in J} \text{scl}_\alpha(5r_j) \leq \sum_{j \in J} \text{scl}_\beta(r_j) .$$

Since this holds true for any pack, we have:

$$\mathcal{P}_\delta^{\text{scl}_\beta}(F_N) \geq \mu(F_N) ,$$

and then taking δ arbitrarily close to 0 leads to:

$$\mathcal{P}_0^{\text{scl}_\beta}(F_N) \geq \mu(F_N) .$$

By taking the sum over $N \geq 1$, it holds:

$$\sum_{N \geq 1} \mathcal{P}_0^{\text{scl}_\beta}(F_N) \geq \sum_{N \geq 1} \mu(F_N) \geq \mu(F) > 0 .$$

Recall that $(F_N)_{N \geq 1}$ is an arbitrary covering of Borel sets of F , thus:

$$\mathcal{P}^{\text{scl}_\beta}(F) \geq \mu(F) > 0 .$$

It holds then $\text{scl}_P F \geq \beta$ for any $\beta < \alpha < \text{ess sup ess sup } \overline{\text{scl}}_{\text{loc}} \mu$ which allows to conclude the proof. \square

We deduce then:

Proposition 3.14. *Let (X, d) be a metric space and μ a Borel measure on X , then:*

$$\text{ess inf } \underline{\text{scl}}_{\text{loc}} \mu \leq \text{scl}_H \mu \quad \text{and} \quad \text{ess inf } \overline{\text{scl}}_{\text{loc}} \mu \leq \text{scl}_P \mu ,$$

and

$$\text{ess sup } \underline{\text{scl}}_{\text{loc}} \mu \leq \text{scl}_H^* \mu \quad \text{and} \quad \text{ess sup } \overline{\text{scl}}_{\text{loc}} \mu \leq \text{scl}_P^* \mu .$$

Proof. The second line of inequalities are given by Remark 3.6. It remains to show the first line of inequalities. Let E be a Borel subset of X with μ positive mass. Thus with σ the restriction of μ to E , it holds by Lemma 3.5:

$$\text{ess sup } \underline{\text{scl}}_{\text{loc}} \sigma \leq \text{scl}_H E \quad \text{and} \quad \text{ess sup } \overline{\text{scl}}_{\text{loc}} \sigma \leq \text{scl}_P E .$$

By Lemma 3.4, it holds:

$$\text{ess inf } \underline{\text{scl}}_{\text{loc}} \mu \leq \text{scl}_H E \quad \text{and} \quad \text{ess inf } \overline{\text{scl}}_{\text{loc}} \mu \leq \text{scl}_P E .$$

Taking the infima over E with positive mass ends the proof. \square

Explicit links between packing scales, Hausdorff scales and local scales of measures can be now established by proving **Theorem B**. Let us first recall its statement: Let (X, d) be a metric space and μ a Borel measure on X , then:

$$\text{scl}_H \mu = \text{ess inf } \underline{\text{scl}}_{\text{loc}} \mu \leq \text{scl}_P \mu = \text{ess inf } \overline{\text{scl}}_{\text{loc}} \mu$$

and

$$\text{scl}_H^* \mu = \text{ess sup } \underline{\text{scl}}_{\text{loc}} \mu \leq \text{scl}_P^* \mu = \text{ess sup } \overline{\text{scl}}_{\text{loc}} \mu .$$

Proof of Theorem B. By Proposition 3.14 it remains only to show four inequalities. We first prove $\text{scl}_H \mu \leq \text{ess inf } \underline{\text{scl}}_{\text{loc}} \mu$. We can assume that $\text{ess inf } \underline{\text{scl}}_{\text{loc}} < +\infty$, otherwise the result immediately comes, and fix $\alpha > \text{ess inf } \underline{\text{scl}}_{\text{loc}}$. Consider $\beta > \alpha$, thus by definition of scaling, there exists $\delta > 0$ such that for any $r \in (0, \delta)$ one has $\text{scl}_\beta(5r) \leq \text{scl}_\alpha(r)$. Denote:

$$F := \{x \in X : \underline{\text{scl}}_{\text{loc}} \mu(x) < \alpha\},$$

then $\mu(F) > 0$ and by Lemma 2.3 for any x in F there exists an integer $n(x)$, minimal, such that $r(x) := \exp(-n(x))$ is at most δ and:

$$\mu(B(x, r(x))) \geq \text{scl}_\alpha(r(x)).$$

Now set:

$$\mathcal{F} := \{B(x, r(x)) : x \in F\}.$$

By Vitali Lemma 3.13, there exists a countable set J and a δ -pack $\{B(x_j, r_j)\}_{j \in J} \subset \mathcal{F}$ of F such that $F \subset \bigcup_{j \in J} B(x_j, 5r_j)$. Then, it holds:

$$\sum_{j \in J} \text{scl}_\beta(5r_j) \leq \sum_{j \in J} \text{scl}_\alpha(r_j) \leq \sum_{j \in J} \mu(B(x_j, r_j)) \leq \mu(F).$$

We then have $\mathcal{H}_\delta^{\text{scl}_\beta}(F) \leq \mu(F)$. Since this holds true δ as small as we want, we deduce $\mathcal{H}^{\text{scl}_\beta}(F) \leq \mu(F)$; and this holds true for any $\beta > \alpha$. We finally get $\text{scl}_H F \leq \alpha$ and then by taking α close to $\text{ess inf } \underline{\text{scl}}_{\text{loc}} \mu$, we indeed have $\text{scl}_H \mu \leq \text{scl}_H F \leq \text{ess inf } \underline{\text{scl}}_{\text{loc}} \mu$.

We prove now $\text{scl}_P \mu \leq \text{ess inf } \overline{\text{scl}}_{\text{loc}} \mu$. Similarly as for Hausdorff scales, we can assume $\text{ess inf } \overline{\text{scl}}_{\text{loc}} \mu < +\infty$. Consider then $\alpha > \text{ess sup } \overline{\text{scl}}_{\text{loc}} \mu$ and set $E = \{x \in X : \overline{\text{scl}}_{\text{loc}} \mu(x) < \alpha\}$, thus $\mu(X \setminus E) = 0$. Moreover it holds:

$$E := \bigcup_{i \geq 1} E_i \quad \text{where } E_i = \{x \in E : \forall r \leq 2^{-i}, \mu(B(x, r)) \geq \text{scl}_\alpha(r)\}.$$

By Lemma 2.20, it holds $\text{scl}_P E = \sup_{i \geq 1} \text{scl}_P E_i$. It is then enough to show that for any $i \geq 1$, we have $\text{scl}_P E_i \leq \alpha$. Indeed, then taking α arbitrarily close to $\text{ess sup } \overline{\text{scl}}_{\text{loc}} \mu$ allows to conclude. In that way let us fix $i \geq 1$. Fix $\delta \in (0, 2^{-i})$, and consider J a countable set and $(B_j)_{j \in J}$ a δ -pack of E_i . It follows:

$$\sum_{j \in J} \text{scl}_\alpha(|B_j|) \leq \sum_{j \in J} \mu(B_j) \leq 1.$$

Since this holds true for any δ -pack, we have:

$$\mathcal{P}_\delta^{\text{scl}_\alpha}(E_i) \leq 1.$$

Taking δ arbitrarily close to 0 leads to:

$$\mathcal{P}^{\text{scl}_\alpha}(E_i) \leq \mathcal{P}_0^{\text{scl}_\alpha}(E_i) \leq 1.$$

It follows that we indeed have $\text{scl}_P E_i \leq \alpha$.

Let us now prove $\text{scl}_H^* \mu \leq \text{ess inf } \overline{\text{scl}}_{\text{loc}} \mu$. We can assume that $\text{ess sup } \underline{\text{scl}}_{\text{loc}} < +\infty$ and fix a real positive number $\alpha > \text{ess sup } \underline{\text{scl}}_{\text{loc}}$. For $\beta > \alpha$, then consider $\delta > 0$ such that for any $r \in (0, \delta)$ we have $\text{scl}_\beta(5r) \leq \text{scl}_\alpha(r)$. Denote $F := \{x \in X : \underline{\text{scl}}_{\text{loc}} \mu(x) < \alpha\}$, thus F has total mass and by Lemma 2.3 for any x in F there exists an integer $n(x)$, minimal, such that $r(x) := \exp(-n(x)) \leq \delta$ and moreover:

$$\mu(B(x, r(x))) \geq \text{scl}_\alpha(r(x)).$$

Now put:

$$\mathcal{F} := \{B(x, r(x)) : x \in F\} .$$

By Vitali's Lemma 3.13, there exists a countable set J and a δ -pack $(B(x_j, r_j))_{j \in J} \subset \mathcal{F}$ of F such that F is included in $\bigcup_{j \in J} B(x_j, 5r_j)$. Thus, it holds:

$$\sum_{j \in J} \text{scl}_\beta(5r_j) \leq \sum_{j \in J} \text{scl}_\alpha(r_j) \leq \sum_{j \in J} \mu(B(x_j, r_j)) \leq \mu(F) .$$

Finally $\mathcal{H}_\delta^{\text{scl}_\beta}(F) \leq \mu(F)$. Since this holds true for δ arbitrarily close to 0, we deduce that $\mathcal{H}^{\text{scl}_\beta}(F) \leq \mu(F)$. Then, taking $\beta > \alpha$ close to α leads to $\text{scl}_H F \leq \alpha$, and thus by taking α close to $\text{ess sup scl}_{\text{loc}} \mu$, we end up with:

$$\text{scl}_H^* \mu \leq \text{scl}_H F \leq \text{ess sup scl}_{\text{loc}} \mu .$$

Now we shall prove $\text{scl}_P \mu \leq \text{ess inf scl}_{\text{loc}} \mu$. Let E be a Borel set with positive measure. Let σ be the restriction of μ to E , thus by Lemma 3.4:

$$\text{ess inf scl}_{\text{loc}} \mu \leq \text{ess inf scl}_{\text{loc}} \sigma ,$$

and then by Lemma 3.5, it holds:

$$\text{ess inf scl}_{\text{loc}} \mu \leq \text{ess inf scl}_{\text{loc}} \sigma \leq \text{ess sup scl}_{\text{loc}} \sigma \leq \text{scl}_P E .$$

This holds true for any E such that $\mu(E) > 0$, thus $\text{ess inf scl}_{\text{loc}} \mu \leq \text{scl}_P \mu$.

Finally, let us show $\text{scl}_P^* \mu \leq \text{ess sup scl}_{\text{loc}} \mu$. Put $\alpha > \text{ess inf scl}_{\text{loc}} \mu$ and set $F := \{x \in X : \text{scl}_{\text{loc}} \mu < \alpha\}$, then $\mu(F) > 0$, and denote:

$$E := \bigcup_{i \geq 1} E_i \quad \text{where } E_i = \{x \in E : \forall r \leq 2^{-i}, \mu(B(x, r)) \geq \text{scl}_\alpha(r)\} .$$

By Lemma 2.20, we have $\text{scl}_P E = \sup_{i \geq 1} \text{scl}_P E_i$, it is then enough to show that for any $i \geq 1$, we have $\text{scl}_P E_i \leq \alpha$. Indeed we can take α arbitrarily close to $\text{ess inf scl}_{\text{loc}} \mu$. We then fix $i \geq 1$. Fix $\delta \in (0, 2^{-i})$. We consider J a countable set and $(B_j)_{j \in J}$ a δ -pack of E_i . Then:

$$\sum_{j \in J} \text{scl}_\alpha(|B_j|) \leq \sum_{j \in J} \mu(B_j) \leq 1 .$$

Since this holds true for any δ -pack, it follows:

$$\mathcal{P}_\delta^{\text{scl}_\alpha}(E_i) \leq 1 .$$

When δ tends to 0, the latter inequality leads to:

$$\mathcal{P}^{\text{scl}_\alpha}(E_i) \leq \mathcal{P}_0^{\text{scl}_\alpha}(E_i) \leq 1 .$$

From there, we deduce $\text{scl}_P E_i \leq \alpha$, which concludes the proof of the last inequality and thus the one of **Theorem B**. \square

4 Applications

4.1 Scales of infinite products of finite sets

A natural toy model in the study of scales is given by a product $Z = \prod_{n \geq 1} Z_k$ of finite sets. To define the metric δ on this set, we fix a decreasing sequence $(\epsilon_n)_{n \geq 1}$ which verifies $\log \epsilon_{n+1} \sim \log \epsilon_n$ when $n \rightarrow +\infty$. We put for $\underline{x} = (x_n)_{n \geq 1} \in Z$ and $\underline{y} = (y_n)_{n \geq 1} \in Z$:

$$\delta(\underline{x}, \underline{y}) := \epsilon_m,$$

where $m = \nu(\underline{x}, \underline{y}) := \inf\{n \geq 1 : x_n \neq y_n\}$ is the minimal index such that the sequences \underline{x} and \underline{y} differ. Note that then if each Z_n is endowed with the discrete topology, then δ provides the product topology on Z .

A natural measure on Z is the following product measure:

$$\mu := \otimes_{n \geq 1} \mu_n,$$

where μ_n is the equidistributed measure on Z_n for $n \geq 1$. The scales of Z and μ are given by the following:

Proposition 4.1. *For any scaling scl, it holds for any $\underline{x} \in Z$:*

$$\underline{\text{scl}}_{\text{loc}} \mu(\underline{x}) = \underline{\text{scl}}_B Z = \sup \left\{ \alpha > 0 : \text{scl}_\alpha(\epsilon_n) \cdot \prod_{k=1}^n \frac{1}{\text{Card } Z_k} \xrightarrow{n \rightarrow +\infty} +\infty \right\}$$

and

$$\overline{\text{scl}}_{\text{loc}} \mu(\underline{x}) = \overline{\text{scl}}_B Z = \inf \left\{ \alpha > 0 : \text{scl}_\alpha(\epsilon_n) \cdot \prod_{k=1}^n \frac{1}{\text{Card } Z_k} \xrightarrow{n \rightarrow +\infty} 0 \right\}.$$

We shall prove this proposition below. A first corollary can be deduced directly from **Theorem A** and **C**:

Corollary 4.2. *For any scaling scl, it holds moreover:*

$$\text{scl}_H Z = \underline{\text{scl}}_Q \mu = \underline{\text{scl}}_B Z$$

and

$$\text{scl}_P Z = \overline{\text{scl}}_Q \mu = \overline{\text{scl}}_B Z.$$

For some particular choice of the sequence $(\epsilon_n)_{n \geq 1}$ and for $\text{scl} = \text{ord}$ we obtain moreover:

Corollary 4.3. *Suppose that $\frac{-\log \epsilon_n}{n}$ converges to $C > 1$ when $n \rightarrow +\infty$. Then for any scaling scl, it holds moreover:*

$$\text{ord}_H Z = \underline{\text{ord}}_B Z = \liminf_{n \rightarrow +\infty} \frac{1}{n \log C} \log \left(\sum_{k=1}^n \log(\text{Card } Z_k) \right)$$

and

$$\text{ord}_P Z = \overline{\text{ord}}_B Z = \limsup_{n \rightarrow +\infty} \frac{1}{n \log C} \log \left(\sum_{k=1}^n \log(\text{Card } Z_k) \right).$$

Note that $\log \epsilon_n \sim -C \cdot n$ implies $\log \epsilon_n \sim \log \epsilon_{n+1}$ when $n \rightarrow +\infty$.

The following lemma allows to prove both Proposition 4.1 and its corollaries:

Lemma 4.4. For any $n \geq 1$ the ϵ_n -covering number $\mathcal{N}_{\epsilon_n}(Z)$ verifies for any $\underline{z} \in Z$:

$$\mathcal{N}_{\epsilon_n}(Z) = \mu(B(\underline{z}, \epsilon_n))^{-1} = \prod_{k=1}^n \text{Card } Z_k .$$

Proof of Proposition 4.1. Since $\log \epsilon_{n+1} \sim \log \epsilon_n$ when $n \rightarrow +\infty$, we have by Lemma 2.3:

$$\underline{\text{scl}}_B Z = \sup \left\{ \alpha > 0 : \text{scl}_\alpha(\epsilon_n) \cdot \mathcal{N}_{\epsilon_n}(Z) \xrightarrow{n \rightarrow \infty} +\infty \right\} \quad \text{and} \quad \overline{\text{scl}}_B Z = \inf \left\{ \alpha > 0 : \text{scl}_\alpha(\epsilon_n) \cdot \mathcal{N}_{\epsilon_n}(Z) \xrightarrow{n \rightarrow \infty} 0 \right\} ,$$

and we have the same form for local scales. Then the sought results follow from Lemma 4.4. \square

Before proving the remaining lemma we first prove the second corollary:

Proof of Corollary 4.3. By Lemma 4.4, for any $n \geq 1$ we have $\mathcal{N}_{\epsilon_n}(Z) = \prod_{k=1}^n \text{Card } Z_k$, then by Remark 2.12 and Lemma 2.3, it holds:

$$\underline{\text{ord}}_B(X) = \liminf_{n \rightarrow +\infty} \frac{\log \log(\mathcal{N}_{\epsilon_n}(Z))}{\log(\epsilon_n^{-1})} = \liminf_{n \rightarrow +\infty} \frac{1}{n \log C} \log \left(\sum_{k=1}^n \log(\text{Card } Z_k) \right)$$

and

$$\overline{\text{ord}}_B(X) = \limsup_{\epsilon \rightarrow 0} \frac{\log \log(\mathcal{N}_{\epsilon_n}(Z))}{\log(\epsilon_n^{-1})} \limsup_{n \rightarrow +\infty} \frac{1}{n \log C} \log \left(\sum_{k=1}^n \log(\text{Card } Z_k) \right) .$$

This concludes the proof of the corollary. \square

Finally we provide the remaining:

Proof of Lemma 4.4. Note that for any $n \geq 1$ and for any $\underline{z} \in Z$:

$$B(\underline{z}, \epsilon_n) = \{ \underline{w} \in Z : w_1 = z_1, \dots, w_n = z_n \} .$$

Thus:

$$\mu(B(\underline{z}, \epsilon_n)) = \prod_{k=1}^n n_k^{-1} .$$

This shows the first equality, it remains to show that $\mathcal{N}_{\epsilon_n}(Z) = \prod_{k=1}^n n_k$. Let us consider $\{\underline{z}^1, \dots, \underline{z}^N\}$ a set of minimal cardinality such that:

$$Z = \bigcup_{j=1}^N B(\underline{z}^j, \epsilon_n) .$$

For $1 \leq j \leq N$, denote $\underline{z}^j = (z_k^j)_{k \geq 1}$, thus we have the following:

Fact 4.5. The map:

$$\phi : i \in \{1, \dots, N\} \mapsto (z_1^i, \dots, z_n^i) \in Z_1 \times \dots \times Z_n .$$

is a bijection.

Proof. We first start by showing ϕ injective. Let us assume that there exists $i \neq j$ such that $\phi(i) = \phi(j)$, then it holds $B(\underline{z}^i, \epsilon_n) = B(\underline{z}^j, \epsilon_n)$. It follows that there exists a covering of Z by $N - 1$ balls with radius ϵ_n , which contradicts the assumption on minimality of N . Thus ϕ is injective. We now show that ϕ is also surjective. Consider $\alpha \in Z_1 \times \dots \times Z_n$. Since Z_k is not empty for any $k \geq 1$, there exists $\underline{z} \in Z$ such that for any $1 \leq k \leq n$ it holds $z_k = \alpha_k$. Then there exists $i \in \{1, \dots, N\}$ such that $\underline{z} \in B(\underline{z}^i, \epsilon_n)$. Thus $\phi(i) = \alpha$ which gives us the surjectivity. \square

From there since ϕ is a bijection, we have:

$$\mathcal{N}_{\epsilon_n}(Z) = N = \text{Card } Z_1 \times \cdots \times Z_N = \prod_{k=1}^N n_k .$$

□

Such examples of products of groups allow to exhibit compact metric spaces with arbitrary high order:

Example 4.6. For any $\alpha \geq \beta > 0$, there exists compact metric probability space (Z, δ, μ) such that for any $z \in Z$:

$$\beta = \underline{\text{ord}}_{\text{loc}} \mu(z) = \text{ord}_H Z = \underline{\text{ord}}_Q \mu = \underline{\text{ord}}_B Z$$

and

$$\alpha = \overline{\text{ord}}_{\text{loc}} \mu(z) = \text{ord}_P Z = \overline{\text{ord}}_Q \mu = \overline{\text{ord}}_B Z .$$

In particular with $\alpha > \beta$ we obtain examples of metric spaces with finite order such that the Hausdorff and packing orders do not coincide. Moreover, for a countable dense subset F of X , it holds $\text{ord}_H F = \text{ord}_P F = 0$ and $\underline{\text{ord}}_B F = \beta < \alpha = \overline{\text{ord}}_B F$. It follows that none of the inequalities of **Theorem A** for the case of order in a equality in the general case. Moreover, using disjoint unions of such spaces allows to produce examples of metric spaces where either of the strict equality can happen between any pair of scales that are not compared in Fig. 1.

Proof. Let $(u_k)_{k \geq 0}$ be the sequence defined by:

$$u_k = \begin{cases} \lfloor \exp(\exp(\beta \cdot k)) \rfloor & \text{if } c^{2j} \leq k < c^{2j+1} \\ \lfloor \exp(\exp(\alpha \cdot k)) \rfloor & \text{if } c^{2j+1} \leq k < c^{2j+2} \end{cases}$$

where $c = \lfloor \frac{\alpha}{\beta} \rfloor + 1$. We denote $Z := \prod_{n \geq 1} \mathbb{Z}/u_n \mathbb{Z}$ endowed with the metric δ defined by:

$$\delta(\underline{z}, \underline{w}) := \exp(-\inf \{n \geq 1 : z_n \neq w_n\})$$

for $\underline{z} = (z_n)_{n \geq 1}$ and $\underline{w} = (w_n)_{n \geq 1}$ in Z . Let us denote $\lambda_n = \frac{1}{n} \log \sum_{k=1}^n \log u_k$. Thus by Corollary 4.3, it follows:

$$\text{ord}_H Z = \underline{\text{ord}}_B Z = \liminf_{n \rightarrow +\infty} \lambda_n$$

and

$$\text{ord}_P Z = \overline{\text{ord}}_B Z = \limsup_{n \rightarrow +\infty} \lambda_n .$$

It remains to show that $\lambda^- := \liminf_{n \rightarrow +\infty} \lambda_n = \beta$ and $\lambda^+ := \limsup_{n \rightarrow +\infty} \lambda_n = \alpha$ in order to show that (Z, δ) satisfies the sought properties. First notice that $\exp(\exp(\beta \cdot n)) \leq u_n \leq \exp(\exp(\alpha \cdot n))$ for every integer n . It follows that $\lambda^- \geq \beta$ and $\lambda^+ \leq \alpha$. Denote $n_j = c^{2j+1}$ and observe that:

$$\lambda_{n_j} \geq \frac{1}{n} \log \log(u_{n_j}) = \alpha .$$

Thus, taking $j \rightarrow +\infty$ leads to $\lambda^+ \geq \alpha$. Moreover, denote $m_j = c^{2j+1} - 1$. We have the following:

Lemma 4.7. For any $j \geq 1$ and for any $1 \leq k \leq m_j$, it holds:

$$u_k < u_{m_j} .$$

Proof. If $c^{2j} \leq k \leq m_j$, then $u_k = \exp(\exp(\beta \cdot k)) \leq \exp(\exp(\beta \cdot m_j)) = u_{m_j}$. Otherwise, we have $k < c^{2j}$, and then $u_k < \exp(\exp(\alpha \cdot c^{2j})) < \exp(\exp(\beta \cdot c^{2j+1})) = u_{m_j}$, since $\alpha < \beta \cdot c$. □

From the above lemma, we have:

$$\lambda_{m_j} \leq \frac{1}{m_j} \log m_j \log(u_{m_j}) \xrightarrow{j \rightarrow +\infty} \beta ,$$

and so $\lambda^- \leq \beta$ which concludes the proof of the proposition. □

4.2 Functional spaces

Metric spaces studied here are sub-spaces of differentiable spaces on compact subset of \mathbb{R}^d for d a positive integer. We denote by $\|\cdot\|_{C^k}$ the C^k -uniform norm on $C^k([0, 1]^d, \mathbb{R})$:

$$\|f\|_{C^k} := \sup_{0 \leq j \leq k} \|D^j f\|_{\infty} .$$

Definition 4.8. For $d \geq 1$ and k an integer, $\alpha \in [0, 1]$ let us define:

$$\mathcal{F}^{d,k,\alpha} := \{f \in C^k([0, 1]^d, [-1, 1]) : \|f\|_{C^k} \leq 1, \text{ and if } \alpha > 0, \text{ the map } D^k f \text{ is } \alpha\text{-H\"older with constant } 1 \} .$$

Recall that for $\alpha > 0$, the map $D^k f$ is α -H\"older with constant 1 if for any $x, y \in [0, 1]^d$ it holds:

$$\|D^k f(x) - D^k f(y)\|_{\infty} \leq \|x - y\|_{\infty}^{\alpha} .$$

In particular, $\mathcal{F}^{d,k,0}$ is the unit ball for the C^k -norm in $C^k([0, 1]^d, [-1, 1])$. Let us recall the asymptotic given by Kolmogorov-Tikhomirov [KT93][Thm XV] on the covering number of $(\mathcal{F}^{d,k,\alpha}, \|\cdot\|_{\infty})$, see Theorem 1.11:

$$C_1 \cdot \epsilon^{-\frac{d}{k+\alpha}} \geq \log \mathcal{N}_{\epsilon}(\mathcal{F}^{d,k,\alpha}) \geq C_2 \cdot \epsilon^{-\frac{d}{k+\alpha}} ,$$

where $C_1 > C_2 > 0$ are two constants depending on d, k and α . In order to prove **Theorem E** which states that box, packing and Hausdorff scales of $\mathcal{F}^{d,k,\alpha}$ are all equal to $\frac{d}{k+\alpha}$, by **Theorem B**, it remains to prove Lemma 1.12. The latter states:

$$\text{ord}_H \mathcal{F}^{d,k,\alpha} \geq \frac{d}{k+\alpha} .$$

Proof of Lemma 1.12. We first start with the case $\alpha > 0$. The case $\alpha = 0$ will be deduced from it. We consider the following set:

$$\Lambda = \prod_{n \geq 1} \Lambda_n .$$

where $\Lambda_n = \{-1, 0, +1\}^{R^{dn}}$ and with $R = \lfloor 5^{\frac{1}{k+\alpha}} \rfloor + 1$. We endow Λ with the metric δ defined by:

$$\delta(\underline{\lambda}, \underline{\lambda}') = \epsilon_m ,$$

with m the minimal index such that the sequences $\underline{\lambda}$ and $\underline{\lambda}'$ differ and with $(\epsilon_n)_{n \geq 0}$ is a decreasing sequence of positive real numbers such that $\frac{-\log \epsilon_n}{n} \rightarrow R^{k+\alpha}$ when $n \rightarrow +\infty$. We can choose $(\epsilon_n)_{n \geq 1}$ such that the following holds:

Lemma 4.9. *There exists an embedding $I : (\Lambda, \delta) \rightarrow (\mathcal{F}^{d,k,\alpha}, \|\cdot\|_\infty)$ such that for any $\lambda, \lambda' \in \Lambda$ it holds:*

$$\|I(\lambda) - I(\lambda')\|_\infty \geq \frac{1}{2} \delta(\lambda, \lambda') .$$

The above lemma allows to conclude the proof of Lemma 1.12. Indeed, since Λ is a product of finite sets endowed with a product metric, and since $\log \epsilon_{n+1} \sim \log \epsilon_n$, by Corollary 4.3 it holds:

$$\text{ord}_H \Lambda = \liminf_{n \rightarrow +\infty} \frac{1}{n \log R^{k+\alpha}} \log \left(\sum_{j=1}^n \log \text{Card } \Lambda_j \right) = \liminf_{n \rightarrow +\infty} \frac{1}{n \log R^{k+\alpha}} \log \left(\sum_{j=1}^n R^{dj} \cdot \log 3 \right) .$$

Now, for $n \geq 1$ it holds:

$$\frac{1}{n \log R^{k+\alpha}} \log \left(\sum_{j=1}^n R^{dj} \cdot \log 3 \right) = \frac{\log \log 3 + \log \frac{R^{d(n+1)} - R}{R-1}}{\log R^{k+\alpha}} \xrightarrow{n \rightarrow +\infty} \frac{d}{k+\alpha} .$$

It comes $\text{ord}_H \Lambda = \frac{d}{k+\alpha}$. Now since by assumption on I in Lemma 4.9, it holds by Corollary 2.22:

$$\text{ord}_H \mathcal{F}^{d,k,\alpha} \geq \text{ord}_H \Lambda = \frac{d}{k+\alpha} ,$$

which concludes the proof of Lemma 1.12. It remains to show:

Proof of Lemma 4.9. Let us denote $q := k + \alpha$ and recall that $R := \lceil 5^{\frac{1}{q}} \rceil + 1$. We consider the following map on \mathbb{R} :

$$\phi : t \in \mathbb{R} \mapsto (2t)^q (2 - 2t)^q \cdot \mathbb{1}_{0 < t < 1} .$$

Note that the function ϕ has its support in $[0, 1]$ and takes the value 1 at $\frac{1}{2}$. The k^{th} derivative of ϕ is non-constant. For $f \in \mathcal{F}^{d,k,\alpha}$, let $\|f\|_q$ be the infimum of the constants $C > 0$ such that for any $x, y \in \mathbb{R}$:

$$\|D^k f(x) - D^k f(y)\| \leq C \cdot \|x - y\|^q .$$

Note that $\|\cdot\|_q$ is a semi-norm on $\mathcal{F}^{d,k,\alpha}$ and moreover:

$$\mathcal{F}^{d,k,\alpha} = \{f \in \mathcal{F}^{d,k,0} : \|f\|_q \leq 1\} .$$

Observe that $\|\phi\|_q > 0$. Let $(x_j)_{1 \leq j \leq R^{dn}}$ be an exhaustive sequence of the set:

$$\left\{ \left(\frac{i_1}{R^n}, \dots, \frac{i_d}{R^n} \right) : i_1, \dots, i_d \in \{0, \dots, R^n - 1\} \right\} .$$

For any $\lambda = (\lambda_1, \dots, \lambda_{R^{dn}}) \in \Lambda_n = \{-1, 0, +1\}^{R^{dn}}$ we associate the following map:

$$f_\lambda : x = (x_1, \dots, x_d) \in [0, 1]^d \mapsto \epsilon_n \cdot \sum_{j=1}^{R^{dn}} \lambda_j \cdot \phi(R^n \cdot \|x - x_j\|) ,$$

with

$$\epsilon_n := \frac{6}{\pi^2 \cdot n^2 \cdot R^{qn} \cdot \|\phi\|_q} .$$

Let us denote \mathcal{S}_n the set of such maps:

$$\mathcal{S}_n = \{f_\lambda : \lambda \in \Lambda_n\} .$$

The sequence $(\epsilon_{n \geq 1})$ is chosen such as the following holds:

Lemma 4.10. *The distance between $f_\lambda, f_{\lambda'} \in \mathcal{S}_n$ is given by:*

$$\|f_\lambda - f_{\lambda'}\|_q = \frac{6}{\pi^2 \cdot n^2} \cdot \|\lambda - \lambda'\|_\infty \quad \text{and} \quad \|f_\lambda - f_{\lambda'}\|_\infty = \epsilon_n \cdot \|\lambda - \lambda'\|_\infty,$$

where $\|\lambda - \lambda'\|_\infty := \sup_{1 \leq i \leq R^{dn}} |\lambda_i - \lambda'_i|$.

Proof. For any $x \in [0, 1]^d$, there exists at most one value $j \in \{1, \dots, R^{dn}\}$ such that $\|x - x_j\| < R^{-n}$, thus the maps $x \mapsto \phi(\|x - x_j\| \cdot R^n)$ for $1 \leq j \leq R^{dn}$ have disjoint supports. It comes then:

$$\|f_\lambda - f_{\lambda'}\|_q = \epsilon_n \left\| \sum_{i=1}^{R^{dn}} |\lambda_i - \lambda'_i| \phi(R^n \|\cdot - x_i\|) \right\|_q = \epsilon_n \cdot \sup_{1 \leq i \leq R^{dn}} |\lambda_i - \lambda'_i| \cdot R^{qn} \|\phi\|_q = \frac{6}{\pi \cdot n^2} \cdot \|\lambda - \lambda'\|_\infty.$$

Now, for the C^0 -norm, it holds:

$$\|f_\lambda - f_{\lambda'}\|_\infty = \epsilon_n \left\| \sum_{i=1}^{R^{dn}} (\lambda - \lambda') \cdot \phi(R^n \|\cdot - x_i\|) \right\|_\infty = \epsilon_n \cdot \|\lambda - \lambda'\|_\infty.$$

□

Note in particular that since $0 \in \mathcal{S}_n$, for any $f_\lambda \in \mathcal{S}_n \setminus \{0\}$, it holds:

$$\|f_\lambda\|_q = \frac{6}{\pi^2 \cdot n^2} \quad \text{and} \quad \|f_\lambda\|_\infty = \epsilon_n.$$

We now embed $\prod_{n \geq 1} \mathcal{S}_n$ into $\mathcal{F}^{d,k,\alpha}$. For $\underline{\lambda} = (\lambda_n)_{n \geq 1} \in \Lambda$ we associate the formal series $\sum_{n \geq 1} f_{\lambda_n}$ where $f_{\lambda_n} \in \mathcal{S}_n$. Then we have the following:

Lemma 4.11. *For any $\underline{\lambda} = (\lambda_n)_{n \geq 1} \in \Lambda$ the function series $\sum_{n \geq 1} f_{\lambda_n}$ converges in $C^0([0, 1]^d, [-1, 1])$ and moreover its limit lies in $\mathcal{F}^{d,k,\alpha}$.*

Proof. By Lemma 4.10, it holds:

$$\sum_{n \geq 1} \|f_{\lambda_n}\|_\infty \leq \sum_{n \geq 1} \epsilon_n < +\infty.$$

It comes that the series $\sum_{n \geq 1} f_{\lambda_n}$ is normally convergent, thus it is also point-wise convergent and moreover the limit g is continuous. Now note that for any $n \geq 1$ and for any $1 \leq l \leq k$ it holds $D^l f_{\lambda_n}(0) = 0$, thus by Taylor integral formula, it holds:

$$\|D^l f_{\lambda_n}\|_\infty \leq \|D^k f_{\lambda_n}\|_\infty.$$

Moreover, still by Lemma 4.10, it holds:

$$\sum_{n \geq 1} \|f_{\lambda_n}\|_q \leq \sum_{n \geq 1} \frac{6}{\pi^2 n^2} = 1.$$

Now since $D^k f_{\lambda_n}$ is $\frac{6}{\pi^2 n^2}$ - α -Hölder and $D^k f_{\lambda_n}(0) = 0$, for any $n \geq 1$, it follows:

$$\sum_{n \geq 1} \|f_{\lambda_n}\|_{C^k} \leq \sum_{n \geq 1} \frac{6}{\pi^2 n^2} = 1.$$

Thus the partial sums lie in $\mathcal{F}^{d,k,\alpha}$ and so does g as a limit of elements of $\mathcal{F}^{d,k,\alpha}$, which is closed for the C^0 -norm. □

By Lemma 4.11, the following map is well defined:

$$I : \underline{\lambda} = (\lambda_n)_{n \geq 1} \in (\Lambda, \delta) \mapsto \lim_{n \rightarrow +\infty} \sum_{n \geq 1} f_{\lambda_n} \in (\mathcal{F}^{d,k,\alpha}, \|\cdot\|_\infty).$$

To conclude the proof, it remains to show that for any $\underline{\lambda}, \underline{\lambda}' \in \Lambda$:

$$\|I(\underline{\lambda}) - I(\underline{\lambda}')\|_\infty \geq \frac{1}{2} \delta(\underline{\lambda}, \underline{\lambda}'),$$

Consider $\underline{\lambda} = (\lambda_n)_{n \geq 1}, \underline{\lambda}' = (\lambda'_n)_{n \geq 1} \in \Lambda$. We denote $k := \nu(\underline{\lambda}, \underline{\lambda}')$. Then it holds:

$$\|I(\underline{\lambda}) - I(\underline{\lambda}')\|_\infty = \left\| \sum_{n \geq k} f_{\lambda_n} - f_{\lambda'_n} \right\|_\infty \geq \|f_{\lambda_k} - f_{\lambda'_k}\|_\infty - \sum_{n > k} \|f_{\lambda_n} - f_{\lambda'_n}\|_\infty.$$

Now by Lemma 4.10, it holds respectively:

$$\|f_{\lambda_k} - f_{\lambda'_k}\|_\infty \geq \epsilon_k \quad \text{and} \quad \sum_{n > k} \|f_{\lambda_n} - f_{\lambda'_n}\|_\infty \leq 2 \sum_{n > k} \epsilon_n.$$

Now recall that $\epsilon_n = \frac{6}{\pi^2 n^2 R^{qn}}$ for any $n \geq 1$, then:

$$\sum_{n > k} \epsilon_n \leq \sum_{n > k} \epsilon_k \cdot R^{-q(n-k)} = \epsilon_k \cdot \frac{1}{R^q - 1}.$$

Now since $R^q \geq 5$, it holds then $\frac{1}{R^q - 1} \leq \frac{1}{4}$ and it follows:

$$\|I(\underline{\lambda}) - I(\underline{\lambda}')\|_\infty \geq \frac{1}{2} \epsilon_{\nu(\underline{\lambda}, \underline{\lambda}')}. \quad \square$$

Now since $\epsilon_{\nu(\underline{\lambda}, \underline{\lambda}')} = \delta(\underline{\lambda}, \underline{\lambda}')$ by definition of δ , the sought result comes. □

This concludes the proof of Lemma 1.12 for the case $\alpha > 0$. It remains to deduce the case $\alpha = 0$ from that previous one. For any $\beta > 0$, it holds $\mathcal{F}^{d,k,\beta} \subset \mathcal{F}^{d,k,0}$. From there since Hausdorff scales are non decreasing for inclusion, it holds then $\text{ord}_H \mathcal{F}^{d,k,0} \geq \text{ord}_H \mathcal{F}^{d,k,\beta} \geq \frac{d}{k+\beta}$. Since we can take $\beta > 0$ arbitrary small, it indeed holds $\text{ord}_H \mathcal{F}^{d,k,0} \geq \frac{d}{k}$. □

References

- [ASY96] Kathleen T Alligood, Tim D Sauer, and James A Yorke. *Chaos*. Springer, 1996. (NOT CITED.)
- [BB21] Pierre Berger and Jairo Bochi. On emergence and complexity of ergodic decompositions. *Advances in Mathematics*, 390:107904, 2021. (Cited on pages 2, 6, and 9.)
- [Ber17] Pierre Berger. Emergence and non-typicality of the finiteness of the attractors in many topologies. *Proceedings of the Steklov Institute of Mathematics*, 297(1):1–27, 2017. (Cited on pages 6 and 9.)
- [Ber20] Pierre Berger. Complexities of differentiable dynamical systems. *Journal of Mathematical Physics*, 61(3):032702, 2020. (Cited on pages 2, 3, 4, 6, 7, 9, and 10.)
- [BGV07] François Bolley, Arnaud Guillin, and Cédric Villani. Quantitative concentration inequalities for empirical measures on non-compact spaces. *Probability Theory and Related Fields*, 137(3-4):541–593, 2007. (Cited on page 9.)

- [BR92] Paolo Baldi and Bernard Roynette. Some exact equivalents for the brownian motion in hölder norm. *Probability theory and related fields*, 93(4):457–484, 1992. (Cited on pages 2 and 7.)
- [Chu47] Kai Lai Chung. On the maximum partial sum of independent random variables. *Proceedings of the National Academy of Sciences of the United States of America*, 33(5):132, 1947. (Cited on pages 2 and 7.)
- [CM44] Robert H. Cameron and William Ted Martin. The wiener measure of hilbert neighborhoods in the space of real continuous functions. *Journal of Mathematics and Physics*, 23:195–209, 1944. (Cited on pages 2 and 7.)
- [DFMS03] Steffen Dereich, Franz Fehrer, Anis Matoussi, and Michael Scheutzow. On the link between small ball probabilities and the quantization problem for gaussian measures on banach spaces. *Journal of Theoretical Probability*, 16(1):249–265, 2003. (Cited on pages 2 and 6.)
- [DL05] S. Dereich and M. A. Lifshits. Probabilities of randomly centered small balls and quantization in Banach spaces. *The Annals of Probability*, 33(4):1397 – 1421, 2005. (Cited on pages 2, 3, 6, and 7.)
- [Fal04] Kenneth Falconer. *Fractal geometry: mathematical foundations and applications*. John Wiley and Sons, 2004. (Cited on pages 3, 4, 6, and 21.)
- [Fan94] Ai Hua Fan. Sur les dimensions de mesures. *Studia Mathematica*, 1(111):1–17, 1994. (Cited on page 6.)
- [FF97] Kenneth J Falconer and KJ Falconer. *Techniques in fractal geometry*, volume 3. Wiley Chichester, 1997. (Cited on pages 3, 4, and 6.)
- [FLR02] Ai-Hua Fan, Ka-Sing Lau, and Hui Rao. Relationships between different dimensions of a measure. *Monatshefte für Mathematik*, 135(3):191–201, 2002. (Cited on page 6.)
- [Fro35] Otto Frostman. *Potentiel d'équilibre et capacité des ensembles*. PhD thesis, Gleerup, 1935. (Cited on page 6.)
- [GL07] Siegfried Graf and Harald Luschgy. *Foundations of quantization for probability distributions*. Springer, 2007. (Cited on page 6.)
- [Hau18] Felix Hausdorff. Dimension und äußeres maß. *Mathematische Annalen*, 79(1-2):157–179, 1918. (Cited on page 2.)
- [KL93] James Kuelbs and Wenbo V Li. Metric entropy and the small ball problem for gaussian measures. *Journal of Functional Analysis*, 116(1):133–157, 1993. (Cited on pages 2 and 7.)
- [Klo15] Benoît R Kloeckner. A geometric study of wasserstein spaces: ultrametrics. *Mathematika*, 61(1):162–178, 2015. (Cited on pages 2 and 9.)
- [Kol30] Andrei Nikolaevitch Kolmogorov. Sur la loi forte des grands nombres. *Comptes rendus de l'Académie des Sciences*, 191:910–912, 1930. (NOT CITED.)
- [KT93] AN Kolmogorov and VM Tikhomirov. ε -entropy and ε -capacity of sets in functional spaces. In *Selected works of AN Kolmogorov*, pages 86–170. Springer, 1993. (Cited on pages 2, 3, 4, 8, 12, and 34.)
- [KZ⁺03] Marc Kesseböhmer, Sanguo Zhu, et al. Quantization dimension via quantization numbers. *Real Analysis Exchange*, 29(2):857–867, 2003. (NOT CITED.)

- [KZ07] Marc Kesseböhmer and Sanguo Zhu. Stability of quantization dimension and quantization for homogeneous cantor measures. *Mathematische Nachrichten*, 280(8):866–881, 2007. (NOT CITED.)
- [KZ15] Marc Kesseböhmer and Sanguo Zhu. Some recent developments in quantization of fractal measures. In *Fractal Geometry and Stochastics V*, pages 105–120. Springer, 2015. (NOT CITED.)
- [Man67] Benoit Mandelbrot. How long is the coast of britain? statistical self-similarity and fractional dimension. *science*, 156(3775):636–638, 1967. (Cited on page 2.)
- [Pöt99] Klaus Pötzelberger. The quantization dimension of distributions. 1999. (Cited on pages 6 and 7.)
- [Sma70] Steve Smale. Differentiable dynamical systems. *Uspekhi Matematicheskikh Nauk*, 25(1):113–185, 1970. (NOT CITED.)
- [Tam95] Masakazu Tamashiro. Dimensions in a separable metric space. *Kyushu Journal of Mathematics*, 49(1):143–162, 1995. (Cited on page 6.)
- [Tri82] Claude Tricot. Two definitions of fractional dimension. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 91, pages 57–74. Cambridge University Press, 1982. (Cited on pages 2, 3, 6, 15, and 16.)
- [Tri99] Claude Tricot. *Courbes et dimension fractale*. Springer Science & Business Media, 1999. (NOT CITED.)
- [Vit08] Giuseppe Vitali. Sui gruppi di punti e sulle funzioni di variabili reali. *Atti Accad. Sci. Torino*, 43:75–92, 1908. (Cited on page 27.)