THE ODD CHROMATIC NUMBER OF A TOROIDAL GRAPH IS AT MOST 9

FANGYU TIAN¹ YUXUE YIN²

¹Department of Mathematics, Central China Normal University, Wuhan, Hubei, China. ²Department of EE, Tsinghua University, Beijing, China.

ABSTRACT. It's well known that every planar graph is 4-colorable. A toroidal graph is a graph that can be embedded on a torus. It's proved that every toroidal graph is 7-colorable. A proper coloring of a graph is called *odd* if every non-isolated vertex has at least one color that appears an odd number of times in its neighborhood. The smallest number of colors that admits an odd coloring of a graph G is denoted by $\chi_o(G)$. In this paper, we prove that if G is tortoidal, then $\chi_o(G) \leq 9$; Note that K_7 is a toroidal graph, the upper bound is no less than 7.

1. INTRODUCTION

In this paper, all graphs are finite and simple, which means no parallel edges and no loops at their vertices. A proper k-coloring of a simple graph G is an assignment (or a special case of labeling) of k colors to the vertices of G so that no two adjacent vertices share the same color. We say a graph G is c-colorable if it admits a proper c-coloring. The chromatic number of a graph G is the minimum c such that G is c-colorable, and this minimum color is denoted by $\chi(G)$. It's well known that $\chi(G) \leq 4$ if G is planar. For a proper coloring, there exists at least one color that appears an odd number of times in the neighborhood of v, then we say v admits an odd coloring. We use $c_o(v)$ to denote the color and $C_o(v)$ to denote the set of the odd colors. An odd c-coloring of a graph is a proper c-coloring with the additional constraint that each vertex admits an odd coloring. A graph G is odd c-colorable if it has an odd c-coloring. The odd chromatic number of a graph G, denoted by $\chi_o(G)$, is the minimum c such that G has an odd c-coloring. Odd coloring has potential applications in many areas, for example, battery consumption aspects of sensor networks and in RFID protocols [1].

Odd coloring was introduced very recently by Petruševski and Škrekovski [2], who proved that planar graphs are odd 9-colorable. Note that a 5-cycle is a planar graph whose odd chromatic number is exactly 5, they further conjectured that planar graphs are odd 5-colorable. Petr and Portier [3] proved that planar graphs are odd 8-colorable. Fabrici[4] proved a strengthening version about planar graphs regarding similar coloring parameters. Cranston [5] studied the restriction of girth and proved that, $\chi_o(G) \leq 5$ if G is a planar graph with girth at least 7, and $\chi_o(G) \leq 6$ if G is a planar graph with girth at least

E-mail address: yinyuxue945@mail.tsinghua.edu.cn.

6. Eun-Kyung Cho [6] focused on a sparse graph and conjectured that, for $c \ge 4$, if G is a graph with $mad(G) \le \frac{4c-4}{c+1}$, then $\chi_o(G) \le c$. And proved that, if G is a graph with $mad(G) \le mad(K_{c+1}^*) = \frac{4c}{c+2}$, then $\chi_o(G) \le c$ for $c \ge 7$, unless G contains $K_{c_1}^*$ as a subgraph. They further proved that $\chi_o(G) \le 5$ if G is a planar graph with girth at least 6.

Suppose that G is a toroidal graph. The 7-color theorem [7] shows that $\chi(G) \leq 7$. Notice that K_7 is a toroidal graph, $\chi(G) = 7$. Note that $\chi_o(G) \geq \chi(G)$, if G is a toroidal graph, then $\chi_o(G) \geq 7$. We proved that,

Theorem 1.1. If G is a toroidal graph, then $\chi_o(G) \leq 9$

We prove Theorem 1.1 by reduction. In the construction of the minimal counterexample, we organize the constraints in a creative way, which simplifies our proof greatly and can be modified to settle other coloring problems. Moreover, we pay a lot attention to summarizing the complex situations in the proof and present it in brevity. More precisely, we force the most difficult part into the configuration as is shown in Figure 1 via discharging method, and simplify the analysis of this main reducible configuration by splitting it into Lemmas 2.4, 2.5 and 2.6, and then come to a conclusion in Lemma 2.7 based on the former tool Lemmas and Claim.

2. Proof

Let G be a counterexample to Theorem 1.1 with the minimum number of 4⁺-vertices, and subject to that, the number of 5⁺-neighbors of 5⁺-vertex G is minimized, and subject to these conditions |E(G)| is minimized.

Lemma 2.1. The minimum degree $\delta(G) \geq 5$

Proof. Suppose otherwise that there is a 4-vertex v in G. Let v_1, v_2, v_3, v_4 be the neighbors of v. Let G' be the graph obtained from G - v by adding $v_1x_1v_2, v_2x_2v_3, v_3x_3v_1$, where each of $x_i, i \in [3]$ is a new 2-vertex. By the minimality of G, G' has an odd 9-coloring c'. Then we can get an odd 9-coloring c of G by coloring each vertex other than v in G with the same color in G' and coloring v with $[9] \setminus \{c(v_1), c(v_2), c(v_3), c(v_4), c_o(v_1), c_o(v_2), c_o(v_3), c_o(v_4)\}$. Since each of x_i is a 2-vertex, $c(v_1) \neq (v_2) \neq (v_3)$. Then v has an odd coloring, a contradiction.

Lemma 2.2. The odd vertex is not adjacent to any odd vertex.

Proof. Suppose otherwise that there exist two odd adjacent vertices u and v. By Lemma 2.1, u and v are 5⁺-vertices. Let G' be the graph obtained from G by splitting edge uv with a 2-vertex w. Since 5⁺-vertex u and v have fewer 5⁺-neighbors in G', there is an odd 9-coloring c' of G' by the minimality of G. Note that $c'(u) \neq c'(v)$ since w is a 2-vertex. Let c(z) = c'(z) for $z \in v(G)$. Since u and v are odd vertices, u and v always admit an odd coloring. Then c is an odd coloring of G, a contradiction.

Lemma 2.3. Let u be a 5-vertex, u_1, u_2, \ldots, u_5 be the neighbors of u in clockwise order, each of u_2 and u_3 be a 6-vertex, $[uu_1u_2], [u_2uu_3], [u_3uu_4]$ be 3-faces. Then G has no such 5-vertex u.

Proof. Suppose otherwise that G has such a 5-vertex u satisfied these constraints in Lemma 2.3. Let $u'_2, u''_2, u''_2 \notin \{u, u_3, u_1\}$ be the neighbors of $u_2; u'_3, u''_3, u''_3 \notin \{u, u_2, u_4\}$ be the neighbors of u_3 . Let G' be the graph obtained from $G - \{u_2, u_3\}$ by adding edges between any two of u'_i, u''_i, u'''_i if they are not adjacent in G for $i \in \{2, 3\}$. Since G' has fewer 4⁺-vertices than G, G' has an odd 9-coloring of c' by the minimality of G. Let c(w) = c'(w) for $w \in V(G) - \{u_2, u_3\}$. Since c' is proper, $c(u'_i) \neq c(u''_i)$ for i = 2, 3.

If $c(u_1) \notin \{c(u'_2), c(u''_2), c(u'''_2)\}$, then u_2 must have an odd coloring regardless the colors of uand u_3 in G. Then color u_2 with the color in $[9] \setminus \{c(u'_2), c(u''_2), c_o(u''_2), c_o(u''_2), c_o(u'''_2), c(u'''_2)\}$. If either $\{c(u_2), c(u_4)\} \not\subseteq \{c(u'_3), c(u''_3), c(u'''_3)\}$ or $c(u_2) = c(u_4)$, then u_3 must have an odd color regardless the color of u in G. Then color u_3 with the color in $[9] \setminus \{c(u'_3), c(u''_3), c(u'''_3), c(u'''_3)\}$. $c_o(u'_3), c_o(u''_3), c_o(u'''_3), c(u_2), c(u_4)\}$. Recolor u with the color in $[9] \setminus \{c(u_1), c(u''_3), c(u'''_3)\}$.

 $c_o(u_1), c(u_2), c(u_3), c(u_4), c_o(u_4), c(u_5), c_o(u_5)\}$. Since u is a 5-vertex, u must have an odd color. Then G has an odd 9-coloring c, a contradiction. Thus, $\{c(u_2), c(u_4)\} \subseteq \{c(u'_3), c(u''_3), c(u'''_3)\}$ and $c(u_2) \neq c(u_4)$. We assume that $c(u_2) = c(u'_3), c(u_4) = c(u''_3)$. In this case, we first recolor u with the color in $[9] \setminus \{c(u_1), c_o(u_1), c(u_2), c(u_4), c(u_5), c_o(u_5), c(u''_3)\}$. Then u_3 has an odd color $c(u'''_3)$. Then color u_3 with the color in $[9] \setminus \{c(u'_3), c(u''_3), c(u''_3), c(u'''_3), c_o(u''_3), c_o(u''_3), c_o(u''_3), c_o(u''_3), c_o(u''_3), c_o(u''_3), c_o(u'''_3)\}$.

Thus, $c(u_1) \in \{c(u'_2), c(u''_2), c(u''_2)\}$. By symmetry, $c(u_4) \in \{c(u'_3), c(u''_3), c(u''_3)\}$. We assume that $c(u_1) = c(u'_2), c(u_4) = c(u'_3)$.

If $|\{c(u'_2), c(u''_2), c(u'''_2), c_o(u''_2), c_o(u''_2), c_o(u'''_2), c(u'''_3), c(u'''_3)\}| \leq 7$, then color u_3 with the color in $[9] \setminus \{c(u'_3), c(u''_3), c(u''_3), c_o(u''_3), c_o(u''_3), c(u'''_3), c(u'''_2)\}$. Then u_2 must have an odd color $c(u''_2)$ or $c(u'''_2)$ regardless the color of u in G. Then color u_2 with the color in $[9] \setminus \{c(u'_2), c(u''_2), c_o(u''_2), c_o(u''_2), c_o(u'''_2), c(u'''_3), c(u'''_3), c(u'''_3), c(u'''_3), c(u'''_3), c(u'''_3), c(u'''_2), c(u'''_2), c(u'''_2), c(u'''_2), c_o(u'''_2), c_o(u'''_2), c_o(u'''_2), c_o(u'''_3), c(u'''_3)\}| \leq 7$, u_2 has at least one color. Then u_3 must have an odd color $c(u''_3)$ or $c(u'''_3)$ regardless the color of u in G. Finally recolor u with the color in $[9] \setminus \{c(u_1), c_o(u_1), c(u_2), c(u_3), c(u_4), c_o(u_4), c(u_5), c_o(u_5)\}$, a contradiction. Thus, $|\{c(u'_2), c(u''_2), c(u'''_2), c_o(u''_2), c_o(u'''_2), c_o(u'''_2), c_o(u'''_2), c_o(u'''_2), c(u'''_3), c(u'''_3)\}| = 8$.

First color u_2 with the color in $[9] \setminus \{c(u'_2), c(u''_2), c_o(u''_2), c_o(u''_2), c_o(u'''_2), c(u'''_3), c(u'''_3)\}$. Then u_3 must have an odd color $c(u''_3)$ or $c(u'''_3)$ regardless the color of u in G. Let $\{c_1, c_2\} \in [9] \setminus \{c(u'_3), c(u''_3), c(u'''_3), c_o(u''_3), c_o(u'''_3), c_o(u'''_3), c(u_2)\}$. If one of c_1 and c_2 is not in $\{c(u''_2), c(u'''_2)\}$, then color u_3 with this color. Then u_2 must have an odd color $c(u''_2)$ or $c(u'''_2)$ regardless the color of u. Recolor u with the color in $[9] \setminus \{c(u_1), c_o(u_1), c(u_2), c(u_3), c(u_4), c_o(u_4), c(u_5), c_o(u_5)\}$, a contradiction. Thus, $\{c_1, c_2\} = \{c(u''_2), c(u'''_2)\}$. Then color u_3 with c_1 . If c_2 is not equal $[9] \setminus \{c(u_1), c_o(u_1), c(u_2), c(u_3), c(u_4), c_o(u_4), c(u_5), c_o(u_5)\}$, then recolor u with the color in $[9] \setminus \{c(u_1), c_o(u_1), c(u_2), c(u_3), c(u_4), c_o(u_4), c(u_5), c_o(u_5)\}$. Then u_2 has an odd color c_2 , a contradiction. Thus, c_2 is equal $[9] \setminus \{c(u_1), c_o(u_1), c(u_2), c(u_3), c(u_4), c_o(u_4), c(u_5), c_o(u_5)\}$. In this case, recolor u with $c(u_2)$. Since c_2 is equal to the only color in $[9] \setminus \{c(u_1), c_o(u_1), c(u_2), c(u_3)\}$, the color of u does not destroy the proper coloring of u_1, u_3, u_4, u_5 and the odd coloring of u_3, u_4, u_5 . Then we choose one of $c(u''_3)$ and $c(u'''_3)$ to color u_2 such that u_1 has an odd color. Then u_2 has an odd coloring c_2, u_3 has an odd coloring $c(u''_3)$ or $c(u''_3)$, a contradiction.

A 6-vertex v is special if v is incident with six 3-faces. A vertex u is free to the vertex v if u is adjacent to v and $|C_o(u)| \geq 3$. Especially, we use $\overline{c_o}(u)$ to denote the the only odd color of u if $|C_o(u)| = 1$. Note that if u is a 5-vertex, then $|C_o(u)| = 1$, 3 or 5.

In the following Lemmas 2.4-2.7, let u be a special 6-vertex, u_1, u_2, \ldots, u_6 be the neighbors of u in the clockwise order, $v_{12}, v_1, v_2 \notin \{u_6, u, u_2\}$ be the neighbors of $u_1; v_2, v_3, v_4 \notin u_6, u_7, u_8\}$ $\{u_1, u, u_3\}$ be the neighbors of u_2 ; $v_4, v_5, v_6 \notin \{u_2, u, u_4\}$ be the neighbors of u_3 ; $v_6, v_7, v_8 \notin \{u_1, u, u_3\}$ $\{u_3, u, u_5\}$ be the neighbors of $u_4; v_8, v_9, v_{10} \notin \{u_4, u, u_6\}$ be the neighbors of $u_5; v_{10}, v_{11}, v_{12} \notin \{u_4, u, u_6\}$ $\{u_5, u, u_1\}$ be the neighbors of u_6 ; each of $u, u_1, \ldots, u_6, v_1, v_2, \ldots, v_{12}$ is a special 6-vertex, as is depicted in Figure 1.



Figure 1. A cluster of special 6-vertices

In the following Lemmas 2.4-2.6, let c be an odd 9-coloring of G-u.

Lemma 2.4. Let $c(u_1) \neq c(u_2) \neq c(u_3)$, $c(u_2) = c(u_6)$, $c(u_3) = c(u_5)$, each of u_1, u_2 and u_6 has exactly one odd color and $\overline{c_o}(u_1) \neq \overline{c_o}(u_2) \neq \overline{c_o}(u_6), \overline{c_o}(u_i) \notin \{c(u_1), c(u_2), c(u_3), c(u_4)\}$ for each $i \in \{1, 2, 6\}$. Then u_1 can be recolored with one color in $[9] \setminus \{c(u_1), c(u_2), c(u_3), c(u_5)\}$ such that u_2 and u_6 are free to u.

Proof. Since $c(u_2) = c(u_6)$ and u_1 has exactly one odd color in G - u, $c(v_2) = c(v_{12})$, $c(v_1) = \overline{c_o}(u_1)$. Since c is proper, $c(v_{12}) = c(v_2) \neq c(u_1) \neq c(u_2)$. If $c(v_2) = c(v_{12}) \neq c(v_2)$ $c(u_3)$, then $\overline{c_o}(u_2) = c(v_2) = c(v_{12}) = \overline{c_o}(u_6)$, a contradiction. Thus, $c(v_2) = c(v_{12}) = c(v$ $c(u_3)$. Since each of u_2 and u_6 has exactly one odd color and $c(u_5) = c(u_3) \neq c(u_1)$, $\{c(v_3), c(v_4)\} = \{\overline{c_o}(u_2), c(u_1)\}$ and $\{c(v_{10}), c(v_{11})\} = \{\overline{c_o}(u_6), c(u_1)\}$. Let c_1 be the color in $[9] \setminus \{c(u_1), c(u_2), c(u_3), c(u_4), c(v_1), c_o(v_1), \overline{c_o}(u_2), \overline{c_o}(u_6)\}$. Recolor u_1 with c_1 . Since $c(u_3) =$ $c(v_2) = c(v_{12}), c_1 \neq c(v_2)$ and $c_1 \neq c(v_{12})$. Thus, each of v_1, v_2, v_{12} is proper. Observe the neighbors of v_2 , $c(v_1) \neq c(u_2) \neq c(v_3)$ since $c(v_3) \in \{c_o(u_2), c(u_1)\}$ and $c(v_1) = \overline{c_o}(u_1)$. Recall that $c_1 \notin \{c(v_1), c(u_2), c(u_1), \overline{c_o}(u_2)\}$. Thus, v_2 must have an odd coloring in the case of u_1 with color c_1 . By symmetry, v_{12} must have an odd coloring in the case of u_1 with color c_1 . Observe the neighbors of u_2 , $c(u_3) \neq c(v_3) \neq c(v_4) \neq c_1$. Then $|C_o(u_2)| \geq 3$ regardless of the color of u in G. Thus, u_2 is free to u. By symmetry, u_6 is free to u.

Lemma 2.5. If $c(u_6) = c(u_2) \neq c(u_1) \neq c(u_3) \neq c(u_5)$, $|C_o(u_1)| = 1$, $|C_o(u_2)| = 1$ and $|C_o(u_2)| = 1$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_6)\}$ occupies at most two different colors in [9] \setminus $\{c(u_1), c(u_2), c(u_3), c(u_5)\}$ together.

Proof. Suppose otherwise that $\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_6)\}$ occupies three colors in $[9] \setminus \{c(u_1), c(u_2), c($ $c(u_3), c(u_5)$ together. Thus, $\overline{c_o}(u_1) \neq \overline{c_o}(u_2) \neq \overline{c_o}(u_6)$. Since u_1 has an odd color and

 $c(u_2) = c(u_6), c(v_2) = c(v_{12}) \text{ and } c(v_1) = \overline{c_o}(v_1).$ If $c(v_2) = c(u_3)$, then $c(v_{12}) = c(u_3)$. Since $c(u_3) \neq c(u_5), u_6$ has an odd color $c(v_{12}) = c(u_3)$, a contradiction. If $c(v_2) = c(u_5)$, then u_2 has an odd color $c(v_2) = c(u_5)$ since $c(u_3) \neq c(u_5)$. Then $\overline{c_o}(u_2) = c(u_5)$, a contradiction. Thus, $c(v_2) \notin \{c(u_3), c(u_5)\}$. Then $\{c(v_3), c(v_4)\} = \{c(u_1), c(u_3)\}$ since u_2 has exactly one odd color and $c(u_1) \neq c(u_3)$. Since u_6 has exactly one odd color and $c(u_1) \neq c(u_5)$, $\{c(v_{10}), c(v_{11})\} = \{c(u_1), c(u_5)\}$. Then $\overline{c_o}(u_2) = c(v_2) = c(v_{12}) = \overline{c_o}(u_6)$, a contradiction. \Box

Lemma 2.6. If $c(u_1) \neq c(u_2) \neq c(u_3) \neq c(u_6)$, $|C_o(u_1)| = 1$ and $|C_o(u_2)| = 1$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_2)\}$ occupies at most one color in $[9] \setminus \{c(u_1), c(u_2), c(u_3), c(u_6)\}.$

Proof. Suppose otherwise that $\{\overline{c_o}(u_1), \overline{c_o}(u_2)\}$ occupies two colors in $[9] \setminus \{c(u_1), c(u_2), c(u_3), c(u_6)\}$. Thus, $\overline{c_o}(u_1) \neq \overline{c_o}(u_2)$. Since u_2 has an odd color and $c(u_3) \neq c(u_1)$, $\{\overline{c_o}(u_2), c(u_3), c(u_1)\} = \{c(v_2), c(v_3), c(v_4)\}$. Since c is proper and $v_2u_1 \in E(G - u)$, $c(v_2) = \overline{c_o}(u_2)$ or $c(v_2) = c(u_3)$. In the former case, since u_1 has exactly one odd color and $\overline{c_o}(u_2) \neq c(u_2) \neq c(u_6)$, $\{c(v_1), c(v_{12})\} = \{c(u_2), c(u_6)\}$. Then $\overline{c_o}(u_1) = c(v_2) = \overline{c_o}(u_2)$, a contradiction. In the latter case, $\{c(v_1), c(v_{12})\} = \{c(u_2), c(u_6)\}$ since u_1 has exactly one odd color and $c(u_3) \neq c(u_2) \neq c(u_2) \neq c(u_6)$. Then $\overline{c_o}(u_1) = c(v_2) = \overline{c_o}(u_2)$.

Lemma 2.7. G has no configure in Figure 1.

Proof. Suppose otherwise that G has the configure in Figure 1. Let G' be the graph obtained from G - u by adding paths $u_1x_1u_3, u_1x_2u_4, u_1x_3u_5$, where x_1, x_2, x_3 are 2-vertices. Since G' has fewer 6⁺-vertices than G, G' has an odd 9-coloring c'. Let c(z) = c'(z) for each vertex $z \in V(G) - u$. Since c' is an odd coloring, each of x_1, x_2 and x_3 has an odd color. Then $c(u_1) \neq c(u_3), c(u_1) \neq c(u_4), c(u_1) \neq c(u_5)$. Then all the color possibilities of u_1, u_2, \ldots, u_6 are shown as the following cases by symmetry.

We first establish the following claim:

Claim Let $|\{c(u_1), c(u_2), \ldots, c(u_6)\}| = k$. If one of the following statements hold, then u admits an odd coloring in G.

- (1) there exist at least k 2 neighbors u_i , u_i satisfies that u_i are free to u or $\overline{c_o}(u_i) \in \{c(u_1), c(u_2), \ldots, c(u_6)\};$
- (2) $|\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5), \overline{c_o}(u_6)\}| < 9 k.$

Case 1 $c(u_1) = 1, c(u_2) = c(u_6) = 2, c(u_3) = c(u_5) = 3, c(u_4) = 2.$

By Claim, $\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5), \overline{c_o}(u_6)\} = \{4, 5, 6, 7, 8, 9\}$. By Lemma 2.4, we can recolor u_1 with color not in $\{c(u_1), c(u_2), c(u_3), c(u_4)\}$ such that u_2 and u_6 are free to u. Then color u with a color in $[9] \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5), c(u_1), c(u_2), c(u_3), c(u_4)\}$.

Since $c(u_1) \neq c(u_4)$, u always admits an odd coloring in G. Thus, the odd coloring c' of G' can return back to G, a contradiction.

Case 2 $c(u_1) = 1, c(u_2) = c(u_6) = 2, c(u_3) = c(u_5) = 3, c(u_4) = 4.$ or $c(u_1) = 1, c(u_2) = c(u_5) = 3, c(u_6) = c(u_3) = 2, c(u_4) = 4.$

By claim, $\{5, 6, 7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5), \overline{c_o}(u_6)\} = \emptyset$. Since $c(u_1) \neq c(u_2) \neq c(u_3) \neq c(u_4), \{\overline{c_o}(u_2), \overline{c_o}(u_3)\}$ occupies at most one color in $\{5, \ldots, 9\}$ by Lemma 2.6. If $\overline{c_o}(u_3) \in \{5, \ldots, 9\}$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5), \overline{c_o}(u_6)\} = \{5, 6, 7, 8, 9\}$. If $\overline{c_o}(u_2) \in [1, 1]$

 $\{5,\ldots,9\}$, then $\{\overline{c_o}(u_1),\overline{c_o}(u_2),\overline{c_o}(u_4),\overline{c_o}(u_5),\overline{c_o}(u_6)\} = \{5,6,7,8,9\}$. If $\overline{c_o}(u_2),\overline{c_o}(u_3) \notin \{5,\ldots,9\}$, then u admits an odd coloring in G by Claim. In each case, $\overline{c_o}(u_5) \neq \overline{c_o}(u_6)$ and each of $\overline{c_o}(u_5)$ and $\overline{c_o}(u_6)$ be in $\{5,6,7,8,9\}$, which contradicts Lemma 2.6 since $c(u_1) \neq c(u_6) \neq c(u_5) \neq c(u_4)$.

Case 3 $c(u_1) = 1, c(u_2) = c(u_6) = c(u_4) = 2, c(u_3) = 3, c(u_5) = 4.$

By claim, $\{5, 6, 7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5), \overline{c_o}(u_6)\} = \emptyset$. Since $c(u_6) = c(u_2) \neq c(u_1) \neq c(u_3) \neq c(u_5)$,

 $\overline{c_o}(u_i)$ for $i \in \{1, 2, 6\}$ occupies at most two different colors in $\{5, 6, 7, 8, 9\}$ by Lemma 2.5. If $\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_6)\}$ occupies at most one color in $\{5, 6, 7, 8, 9\}$, then $\{5, 6, 7, 8, 9\} \setminus \{\overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5)\} \neq \emptyset$, it contradicts Claim. If $\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_6)\}$ occupies two colors in $\{5, 6, 7, 8, 9\}$, say 5 and 6, then $\{\overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \{7, 8, 9\}$. If $\{\overline{c_o}(u_2), \overline{c_o}(u_6)\} = \{5, 6\}$, then $\{\overline{c_o}(u_6), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \{5, 6, 7, 8, 9\}$. If $\{\overline{c_o}(u_1), \overline{c_o}(u_6)\} = \{5, 6\}$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_6), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \{5, 6, 7, 8, 9\}$. If $\{\overline{c_o}(u_1), \overline{c_o}(u_2)\} = \{5, 6\}$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \{5, 6, 7, 8, 9\}$. If $\{\overline{c_o}(u_1), \overline{c_o}(u_2)\} = \{5, 6\}$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \{5, 6, 7, 8, 9\}$. If $\{\overline{c_o}(u_1), \overline{c_o}(u_2)\} = \{5, 6\}$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \{5, 6, 7, 8, 9\}$. If $\{\overline{c_o}(u_1), \overline{c_o}(u_2)\} = \{5, 6\}$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_5)\} = \{5, 6, 7, 8, 9\}$. In the first two cases, $\overline{c_o}(u_4) \neq \overline{c_o}(u_5) \neq \overline{c_o}(u_6)$ and each of $\overline{c_o}(u_4), \overline{c_o}(u_5) \neq c(u_1) \neq c(u_3)$. In the last case, $\overline{c_o}(u_2) \neq \overline{c_o}(u_3) \neq \overline{c_o}(u_4)$ and each of $\overline{c_o}(u_2), \overline{c_o}(u_3)$ and $\overline{c_o}(u_4)$ is in $\{5, 6, 7, 8, 9\}$, which contradicts Lemma 2.5 since $c(u_6) = c(u_4) \neq c(u_5) \neq c(u_1) \neq c(u_3)$. In the last case, $\overline{c_o}(u_2) \neq \overline{c_o}(u_3) \neq \overline{c_o}(u_4)$ and each of $\overline{c_o}(u_2), \overline{c_o}(u_3)$ and $\overline{c_o}(u_4)$ is in $\{5, 6, 7, 8, 9\}$, which contradicts Lemma 2.5 since $c(u_2) = c(u_4) \neq c(u_1) \neq c(u_3) \neq c(u_5)$.

Case 4 $c(u_1) = 1, c(u_2) = c(u_4) = 3, c(u_6) = c(u_3) = 2, c(u_5) = 4.$

By claim, $\{5, 6, 7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5), \overline{c_o}(u_6)\} = \emptyset$. Since $c(u_1) \neq c(u_6) \neq c(u_5) \neq c(u_4), \{\overline{c_o}(u_5), \overline{c_o}(u_6)\}$ occupies at most one color in $\{5, \ldots, 9\}$ by Lemma 2.5. If $\overline{c_o}(u_6) \in \{5, \ldots, 9\}$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_6)\} = \{5, 6, 7, 8, 9\}$. If $\overline{c_o}(u_5) \in \{5, \ldots, 9\}$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \{5, 6, 7, 8, 9\}$. If $\overline{c_o}(u_5), \overline{c_o}(u_6) \notin \{5, \ldots, 9\}$, then u admits an odd coloring in G by Claim.

In each case, $\overline{c_o}(u_2) \neq \overline{c_o}(u_3) \neq \overline{c_o}(u_4)$ and each of $\overline{c_o}(u_2)$, $\overline{c_o}(u_3)$ and $\overline{c_o}(u_4)$ be in $\{5, 6, 7, 8, 9\}$, which contradicts Lemma 2.5 since $c(u_2) = c(u_4) \neq c(u_1) \neq c(u_3) \neq c(u_5)$.

Case 5 $c(u_1) = 1, c(u_2) = c(u_6) = 2, c(u_3) = 3, c(u_5) = 4, c(u_4) = 5.$

By claim, $\{6, 7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5), \overline{c_o}(u_6)\} = \emptyset$. By Lemma 2.6, $\{\overline{c_o}(u_2), \overline{c_o}(u_3)\}$ occupies at most one color in $\{6, 7, 8, 9\}$ due to $c(u_1) \neq c(u_2) \neq c(u_3) \neq c(u_4)$; $\{\overline{c_o}(u_5), \overline{c_o}(u_6)\}$ occupies at most one color in $\{6, 7, 8, 9\}$ due to $c(u_1) \neq c(u_6) \neq c(u_5) \neq c(u_4)$. If $\overline{c_o}(u_2), \overline{c_o}(u_5) \in \{6, 7, 8, 9\}$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \{6, 7, 8, 9\}$; if $\overline{c_o}(u_2), \overline{c_o}(u_6) \notin \{6, 7, 8, 9\}$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_4), \overline{c_o}(u_6)\} = \{6, 7, 8, 9\}$. They contradict Lemma 2.6 since $c(u_3) \neq c(u_4) \neq c(u_5) \neq c(u_6)$ and Lemma 2.5 since $c(u_6) = c(u_2) \neq c(u_1) \neq c(u_3) \neq c(u_5)$. If $\overline{c_o}(u_3), \overline{c_o}(u_5) \in \{6, 7, 8, 9\}$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \{6, 7, 8, 9\}$; if $\overline{c_o}(u_3), \overline{c_o}(u_6) \in \{6, 7, 8, 9\}$, then $\{\overline{c_o}(u_1), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \{6, 7, 8, 9\}$; They contradict Lemma 2.6 since $c(u_2) \neq c(u_3) \neq c(u_4) \neq c(u_5)$. If $\{\overline{c_o}(u_3), \overline{c_o}(u_5)\} = \{6, 7, 8, 9\}$; then $\{\overline{c_o}(u_1), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \{6, 7, 8, 9\}$; occupies at most one color in $\{6, 7, 8, 9\}$, then $\{\overline{c_o}(u_4) \neq c(u_5)$. If $\{\overline{c_o}(u_2), \overline{c_o}(u_5), \overline{c_o}(u_6)\}$ occupies at most one color in $\{6, 7, 8, 9\}$, then u admits an odd coloring by Claim.

Case 6 $c(u_1) = 1, c(u_2) = 3, c(u_6) = (u_3) = 2, c(u_5) = 4, c(u_4) = 5.$

In this case, $c(u_1) \neq c(u_2) \neq c(u_3) \neq c(u_4)$ and $c(u_1) \neq c(u_6) \neq c(u_5) \neq c(u_4)$. By same argument of Case 5, $\{6, 7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_4), \overline{c_o}(u_5)\}$ = \emptyset or $\{6, 7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_4), \overline{c_o}(u_6)\} = \emptyset$ or $\{6, 7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_4), \overline{c_o}(u_6)\}$ $\overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \emptyset \text{ or } \{6, 7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_6)\} = \emptyset. \text{ In the first case, we assume that } \overline{c_o}(u_1) = 6, \overline{c_o}(u_2) = 7, \overline{c_o}(u_4) = 8, \overline{c_o}(u_5) = 9. \text{ Since } \overline{c_o}(u_1) = 6 \text{ and } c(u_2) \neq c(u_6), \{6, 2, 3\} = \{c(v_{12}), c(v_1), c(v_2)\}. \text{ Then } c(v_2) = 6 \text{ or } 2. \text{ If } c(v_2) = 6, \text{ then } \overline{c_o}(u_2) = 6, \text{ a contradiction. Thus, } c(v_2) = 2. \text{ By symmetry, } c(v_8) = 2. \text{ We color } u \text{ with } 3, \text{ recolor } u_2 \in [9] \setminus \{1, 2, 3, 6, 7, \overline{c_o}(v_2), \overline{c_o}(v_3), \overline{c_o}(v_4)\}. \text{ Since } \overline{c_o}(u_2) = 7, c(v_4) = 1 \text{ or } 7. \text{ Since } \overline{c_o}(u_4) = 8, c(v_4) = 4 \text{ or } 8. \text{ Then } u_3 \text{ has an odd coloring. Since } \overline{c_o}(u_1) = 6, c(v_{12}) = 3 \text{ or } 6. \text{ Since } \overline{c_o}(u_4) = 9, c(v_{10}) = 5 \text{ or } 9. \text{ Then } u_6 \text{ has an odd coloring. Then } \overline{c_o}(u_1) = 6, \overline{c_o}(u_2) = 7, \overline{c_o}(u_4) = 8, \overline{c_o}(u_5) = 9. \text{ Thus, } G \text{ has a } 9\text{-odd coloring, a contradiction. In the second case, } \overline{c_o}(u_1) \neq \overline{c_o}(u_6) \text{ and each of } \overline{c_o}(u_1) \text{ and } \overline{c_o}(u_6) \text{ be in } \{6, 7, 8, 9\}, \text{ which contradicts Lemma 2.6 since } c(u_2) \neq c(u_3) \neq c(u_4) \neq c(u_5). \text{ and } \overline{c_o}(u_5) = 0 \text{ or } 10, 10 \text{ or } 10, 10$

Case 7 $c(u_1) = 1, c(u_2) = 3, c(u_6) = 2, c(u_3) = 4, c(u_5) = 4, c(u_4) = 5.$

In this case, $c(u_1) \neq c(u_2) \neq c(u_3) \neq c(u_4)$ and $c(u_1) \neq c(u_6) \neq c(u_5) \neq c(u_4)$. By same argument of Case 5, $\{6, 7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_4), \overline{c_o}(u_5)\}$

 $= \varnothing \text{ or } \{6,7,8,9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_4), \overline{c_o}(u_6)\} = \varnothing \text{ or } \{6,7,8,9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o$

 $\overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \emptyset \text{ or } \{6, 7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_6)\} = \emptyset. \text{ In first two cases, } \overline{c_o}(u_1) \neq \overline{c_o}(u_2) \text{ and each of } \overline{c_o}(u_1) \text{ and } \overline{c_o}(u_2) \text{ be in } \{6, 7, 8, 9\}, \text{ which contradicts Lemma 2.6 since } c(u_3) \neq c(u_2) \neq c(u_1) \neq c(u_6). \text{ In the third case, } \overline{c_o}(u_3) \neq \overline{c_o}(u_4) \neq \overline{c_o}(u_5) \text{ and each of } \overline{c_o}(u_3), \overline{c_o}(u_4) \text{ and } \overline{c_o}(u_5) \text{ be in } \{6, 7, 8, 9\}, \text{ which contradicts Lemma 2.5 since } c(u_5) = c(u_3) \neq c(u_2) \neq c(u_4) \neq c(u_6). \text{ In last case, } \overline{c_o}(u_1) \neq \overline{c_o}(u_6) \text{ and each of } \overline{c_o}(u_1) \text{ and } \overline{c_o}(u_6) \text{ be in } \{6, 7, 8, 9\}, \text{ which contradicts Lemma 2.5 since } c(u_6) \text{ be in } \{6, 7, 8, 9\}, \text{ which contradicts Lemma 2.6 since } c(u_2) \neq c(u_1) \neq c(u_6) \neq c(u_5).$

Case 8 $c(u_1) = 1, c(u_2) = 3, c(u_6) = c(u_4) = 2, c(u_3) = 4, c(u_5) = 5.$

By claim, $\{6, 7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5), \overline{c_o}(u_6)\} = \emptyset$. By Lemma 2.6, $\{\overline{c_o}(u_3), \overline{c_o}(u_4)\}$ occupies at most one color in $\{6, 7, 8, 9\}$ due to $c(u_2) \neq c(u_3) \neq c(u_4) \neq c(u_5)$; $\{\overline{c_o}(u_1)\overline{c_o}(u_6)\}$ occupies at most one color in $\{6, 7, 8, 9\}$ due to $c(u_1) \neq c(u_2) \neq c(u_5) \neq c(u_6)$. Thus, $\{6, 7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_5)\} = \emptyset$ or $\{6, 7, 8, 9\} \setminus \{\overline{c_o}(u_6), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_5)\}$ $= \emptyset$ or $\{6, 7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \emptyset$ or $\{6, 7, 8, 9\} \setminus \{\overline{c_o}(u_6), \overline{c_o}(u_6), \overline{c_o}(u_6), \overline{c_o}(u_6)\}$

 $\overline{c_o}(u_2), \overline{c_o}(u_4), \overline{c_o}(u_5)\} = \emptyset$. In the former two cases, they contradict Lemma 2.6 since $c(u_1) \neq c(u_2) \neq c(u_3) \neq c(u_4)$. In the latter two cases, they contradict Lemma 2.6 due to $c(u_6) \neq c(u_1) \neq c(u_2) \neq c(u_3)$ and Lemma 2.5 due to $c(u_6) = c(u_4) \neq c(u_3) \neq c(u_5) \neq c(u_1)$. Case.9 $c(u_1) = 1, c(u_2) = 2, c(u_6) = 6, c(u_3) = 3, c(u_5) = 5, c(u_4) = 4$.

By claim, $\{7, 8, 9\} \setminus \{\overline{c_o}(u_1), \overline{c_o}(u_2), \overline{c_o}(u_3), \overline{c_o}(u_4), \overline{c_o}(u_5), \overline{c_o}(u_6)\} = \emptyset$.

By Lemma 2.6, $\{u_i, u_{i+1}\}$ occupies at most one color in $\{7, 8, 9\}$, where $1 \leq i \leq 6$ and i+1=1 if i=6. Thus, $\{\overline{c_o}(u_1), \overline{c_o}(u_3), \overline{c_o}(u_5)\} = \{7, 8, 9\}$ by symmetry. We assume that, $\overline{c_o}(u_1) = 7, \overline{c_o}(u_3) = 8, \overline{c_o}(u_5) = 9$ and $c(u_1) \neq c(u_2) \neq \ldots \neq c(u_6)$. Then $c(v_{12}) = 2$ or 7, $c(v_2) = 6$ or 7, $c(v_4) = 4$ or 8, $c(v_6) = 2$ or 8, $c(v_8) = 9$ or 6, $c(v_{10}) = 9$ or 4. Then color u with 1. Recolor $u_1 \in [9] \setminus \{1, 2, 6, 7, c_o(v_1), c_o(v_2), c_o(v_{12})\}$. In each case, u_2, u_4 and u_6 has at least one odd coloring, $\overline{c_o}(u_1) = 1, \overline{c_o}(u_3) = 8, \overline{c_o}(u_5) = 9$, and u has the odd coloring, a contradiction.

Now we are ready to complete the proof of Theorem 1.1. Let each $v \in V(G)$ have an initial charge of $\mu(v) = d(v) - 6$, each $f \in \bigcup F(G)$ have an initial charge of $\mu(f) = 2d(f) - 6$. By Euler's Formula, $|V(G)| + |F(G)| - |E(G)| \ge 0$. Then $\sum_{v \in V} \mu(v) + \sum_{f \in F} \mu(f) = 0$.

Let $\mu^*(x)$ denote the final charge of $x \in V(G) \cup F(G)$ after the discharging procedure. To lead to a contradiction, we shall prove that $\sum_{x \in V(G) \cup F(G)} \mu^*(x)$

> 0. Since the total sum of charges is unchanged in the discharging procedure, this contradiction proves Theorem 1.1.

We use the following discharging rules:

- (R1) Every 4^+ -face sends 1 to each incident 5 -vertex.
- (R2) Every 8⁺-vertex u sends $\frac{3}{8}$ to each adjacent 5 -vertex v if both faces incident with the edge uv are 3-faces.

By Lemma 2.1, $\delta(G) \geq 5$. Then we check that the final charge of each 5⁺-vertex and each face is nonnegative.

- 1. Let f be a 3-face. By Rules. f is not involved in any discharging procedure. Thus, $\mu^*(f) = 2d(f) - 6 = 0.$
- 2. Let f be a 4⁺-face. By Lemma 2.2, 5-vertex is not adjacent to 5-vertex. Then f is incident with at most $\frac{d(f)}{2}$ 5-vertices. By (R1), f sends 1 to each incident 5-vertex. Thus, $\mu^*(f) = 2d(f) 6 \frac{d(f)}{2} \ge 0$. Note that $\mu^*(f) = 0$ if and only if d(f) = 4 and f is incident with two 5-vertices.
- 3. Let v be a 5-vertex. If v is incident with at least two 4⁺-faces, then each of 4-faces sends 1 to v by (R1). Thus, $\mu^*(v) = d(v) 6 + 2 > 0$. If v is incident with one 4-face, then v is incident with four 3-faces. Let v_1, v_2, \ldots, v_5 be the neighbors of v, the face incident with v_1vv_2 be a 4⁺-face. By Lemma 2.2, each of v_1, \ldots, v_5 is a 6⁺-vertex. By Lemma 2.3 and one of v_3 and v_4 is a 8⁺-vertex. By (R2), this 8⁺-vertex sends $\frac{3}{8}$ to v. Thus, $\mu^*(v) = d(v) 6 + 1 + \frac{3}{8} > 0$. Thus, each face incident with v is a 3-face. By Lemma 2.3, at least one of v_i and v_{i+1} is a 8⁺-vertex where $i \in \{1, 2, 3, 4, 5\}$ and i + 1 = 1 if i = 5. Thus, at least three of v_1, v_2, \ldots, v_5 are 8⁺-vertices. By (R2), $\mu^*(v) = d(v) 6 + \frac{3}{8} \times 3 > 0$.
- 4. Let v be a 6- or 7-vertex. By Rules. v is not involved in any discharging procedure. Thus, $\mu^*(v) = d(v) - 6 = 0$ if f(f) = 6, $\mu^*(v) = d(v) - 6 > 0$ if f(f) = 7.
- 5. Let v be a 8⁺-vertex. By (R2), v sends $\frac{3}{8}$ to adjacent 5-vertex v if uv is incident with two 3-faces. Let v_2 be the 5-neighbor of v and vv_2 is incident with two 5-faces $[vv_2v_1]$ and $[vv_2v_3]$. By Lemma 2.2, 5-vertex is not adjacent to 5-vertex. Then each of v_1 and v_3 are 6⁺-vertices. Thus, v sends charge to at most $\frac{d(v)}{2}$ 5-neighbors. Thus, $\mu^*(v) = d(v) 6 \frac{3}{8} \times \frac{d(v)}{2} > 0$.

From our hypothesis and the above discharging procedure, we know that the following configurations admit positive final charge and thus can not be contained in G. That's, if G has a 7⁺-vertex or a 5-vertex, then $\sum_{x \in V(G) \cup F(G)} \mu^*(x) > 0$; if G has a 5⁺-face or a 4-face incident with at most one 5-vertex, then $\sum_{x \in V(G) \cup F(G)} \mu^*(x) > 0$; if G has a 4-face incident with two 5-vertices, then these two 5-vertices has final charge more than 0, thus $\sum_{x \in V(G) \cup F(G)} \mu^*(x) > 0$. Thus, G only has 6-vertices and 3-faces. By Lemma 2.7, G has no such configurations, a contradiction.

References

- Smorodinsky, Shakhar. "Conflict-free coloring and its applications." Geometry—Intuitive, Discrete, and Convex. Springer, Berlin, Heidelberg, 2013. 331-389.
- [2] Petruševski, Mirko, and Riste Škrekovski. "Colorings with neighborhood parity condition." arXiv preprint arXiv:2112.13710 (2021).
- [3] Petr, Jan, and Julien Portier. "Odd chromatic number of planar graphs is at most 8." arXiv preprint arXiv:2201.12381 (2022).
- [4] Fabrici, Igor, et al. "Proper conflict-free and unique-maximum colorings of planar graphs with respect to neighborhoods." arXiv preprint arXiv:2202.02570 (2022).
- [5] Cranston, Daniel W. "Odd Colorings of Sparse Graphs." arXiv preprint arXiv:2201.01455 (2022).
- [6] Cho, Eun-Kyung, et al. "Odd coloring of sparse graphs and planar graphs." arXiv preprint arXiv:2202.11267 (2022).
- [7] Kauffman, Louis H. "Seven Knots and Knots in the Seven-Color Map." Homage to a Pied Puzzler (2009): 75.