

SOME SPECIAL SOLUTIONS OF A NONLINEAR SYSTEM OF 4 ORDINARY DIFFERENTIAL EQUATIONS RECENTLY INTRODUCED TO INVESTIGATE THE EVOLUTION OF HUMAN RESPIRATORY VIRUS EPIDEMICS

Francesco Calogero^{a,b*,†}, Andrea Giansanti^{a,b,‡§}, Farrin Payandeh^{c¶||}

June 15, 2022

^a *Physics Department, University of Rome "La Sapienza", Rome, Italy*

^b *INFN, Sezione di Roma 1*

^c *Department of Physics, Payame Noor University, PO BOX 19395-3697 Tehran, Iran*

Abstract

A system of 4 nonlinearly-coupled Ordinary Differential Equations has been recently introduced to investigate the evolution of human respiratory virus epidemics. In this paper we point out that some *explicit* solutions of that system can be obtained by *algebraic* operations, provided the parameters of the model satisfy certain *constraints*.

1 Introduction

The following system of 4 nonlinearly-coupled Ordinary Differential Equations (ODEs) has been recently introduced to investigate the evolution of human respiratory virus epidemics [1]:

$$\dot{\tilde{x}}_1 = -k_D \tilde{x}_1 + \alpha k_R (\tilde{x}_3 + \tilde{x}_4) , \quad (1a)$$

$$\dot{\tilde{x}}_2 = k_B \tilde{x}_1 + [k_B - k_D - f(\tilde{x}_1, \tilde{x}_2, \tilde{x}_3, \tilde{x}_4)] \tilde{x}_2 + [k_B + (1 - \alpha) k_R] (\tilde{x}_3 + \tilde{x}_4) , \quad (1b)$$

$$\dot{\tilde{x}}_3 = f(\tilde{x}_1, \tilde{x}_2, \tilde{x}_3, \tilde{x}_4) \tilde{x}_2 - (k_R + k_D + k_P) \tilde{x}_3 , \quad (1c)$$

$$\dot{\tilde{x}}_4 = k_P \tilde{x}_3 - (k_R + k_D + k_{DV}) \tilde{x}_4 , \quad (1d)$$

where

$$f(\tilde{x}_1, \tilde{x}_2, \tilde{x}_3, \tilde{x}_4) = \frac{k_I (\tilde{x}_3 + \beta \tilde{x}_4)}{\tilde{x}_1 + \tilde{x}_2 + \tilde{x}_3 + \beta \tilde{x}_4} . \quad (1e)$$

Notation. We maintained the original notation of [1], except for the following replacement of the 4 dependent variables $I(t)$ (number of *Immune* hosts), $S(t)$ (number of *Susceptible* hosts), $A(t)$ (number of *Asymptomatic* and infectious hosts), $C(t)$ (number of *Symptomatic* and infectious hosts) used there, and the use of a superimposed dot (instead of an appended prime) to denote differentiation with respect to the dependent variable t ("time"):

$$I(t) = \tilde{x}_1(t) , \quad S(t) = \tilde{x}_2(t) , \quad A(t) = \tilde{x}_3(t) , \quad C(t) = \tilde{x}_4(t) ; \quad \dot{\tilde{x}}(t) \equiv d\tilde{x}(t)/dt . \quad (2)$$

*e-mail: francesco.calogero@roma1.infn.it

†e-mail: francesco.calogero@uniroma1.it

‡e-mail: andrea.giansanti@uniroma1.it

§e-mail: andrea.giansanti@roma1.infn.it

¶e-mail: farrinpayandeh@yahoo.com

||e-mail: f_payandeh@pnu.ac.ir

Note that in the following we occasionally *omit* to indicate *explicitly* the t -dependence of the dependent variables: and see below **Remark 1.1** for our (notational) motivation of the tilde superimposed on these coordinates $\tilde{x}_n(t)$. ■

For the epidemiological significance of this model see [1], as well as for references to analogous models.

Remark 1.1. The system of 4 nonlinearly-coupled ODEs (1) features the 8 *a priori arbitrary* (of course time-independent) *parameters* $k_D, k_R, k_B, k_P, k_{DV}, k_I, \alpha, \beta$, for whose epidemiological significance we refer to [1]; in this paper we focus mainly on some mathematical properties of this system, so we generally assume that these are 8 *a priori arbitrary* (possibly even *complex*) *numbers*, although we shall comment occasionally on the relevance of such mathematical treatment on the *epidemic* model (when these numbers are *positive real* numbers).

One observation which is relevant for the mathematical discussion of this model (1)—which we think is reasonable to state at the very beginning of this paper—is to note that the parameter k_D plays a relatively trivial role in this system, because it can be altogether eliminated from it via the following very simple change of dependent variables:

$$\tilde{x}_n(t) = x_n(t) \exp(-k_D t) , \quad n = 1, 2, 3, 4 , \quad (3a)$$

implying of course

$$\tilde{x}_n(0) = x_n(0) , \quad n = 1, 2, 3, 4 ; \quad (3b)$$

indeed the system of ODEs satisfied by the 4 variables $x_n(t)$ is then *identical* with the original system (1), except for the elimination of the parameter k_D :

$$\dot{x}_1 = \alpha k_R (x_3 + x_4) , \quad (4a)$$

$$\dot{x}_2 = k_B x_1 + [k_B - f(x_1, x_2, x_3, x_4)] x_2 + [k_B + (1 - \alpha) k_R] (x_3 + x_4) , \quad (4b)$$

$$\dot{x}_3 = f(x_1, x_2, x_3, x_4) x_2 - (k_R + k_P) x_3 , \quad (4c)$$

$$\dot{x}_4 = k_P x_3 - (k_R + k_{DV}) x_4 , \quad (4d)$$

where of course now

$$f(x_1, x_2, x_3, x_4) = \frac{k_I (x_3 + \beta x_4)}{x_1 + x_2 + x_3 + \beta x_4} . \quad \blacksquare \quad (4e)$$

Hence hereafter we shall mainly deal with this, marginally simpler, system (4).

In the following **Section 2** we investigate a very simple solution of this model (4), characterized by the fact that the 4 components $x_n(t)$ of this solution all evolve proportionally to the same exponential function of time, $\exp(\mu t)$, with μ an appropriate parameter determined in terms of the parameters of the model; implying that the quantity $f(x_1, x_2, x_3, x_4)$ is *time-independent* (see (4e)), hence that the system (4)—for this class of solutions—reduces to a *linear* system of 4 ODEs.

In the subsequent **Section 3** we discuss the somewhat less simple solutions characterized by the presumably more interesting requirement that each of the 4 components $x_n(t)$ of the solution be *linear* combinations—with *time-independent* coefficients—of 2 exponential functions of time, $\exp(\mu_1 t)$ and $\exp(\mu_2 t)$, and moreover that the quantity $f(x_1, x_2, x_3, x_4)$, see (4e), be again *time-independent*. The related restrictions on the parameters of the model and the initial-data of this solution are also *explicitly* determined, up to *algebraic* operations.

The subsequent **Section 4** outlines the *analogous* treatments when the solution being identified is the sum of 3, or 4, exponentials.

A final **Section 5** concludes the paper, by mentioning its applicative relevance and possible further developments of the approach used in this paper.

2 A very simple solution

The right-hand sides of the 4 ODEs (4) are all *homogeneous* of degree 1 in the 4 dependent variables $x_n(t)$. This implies a well-known (see for instance [2]) consequence, which can be stated as the following

Proposition 2.1. The system of 4 ODEs (4) features the *simple explicit* solution

$$x_n(t) = x_n(0) \exp(\mu t) , \quad n = 1, 2, 3, 4 , \quad (5)$$

where $x_n(0)$ are clearly the 4 *initial values* of the 4 dependent variables $x_n(t)$ and μ is an *a priori arbitrary* time-independent parameter, provided these 5 quantities—i. e., $x_n(0)$ and μ , together with the 7 parameters of the model (4)—satisfy (as it were, *a posteriori*) the following 4 *algebraic* equations:

$$\mu x_1(0) = \alpha k_R [x_3(0) + x_4(0)] , \quad (6a)$$

$$\mu x_2(0) = k_B x_1(0) + [k_B - f(0)] x_2(0) + [k_B + (1 - \alpha) k_R] [x_3(0) + x_4(0)] , \quad (6b)$$

$$\mu x_3(0) = f(0) x_2(0) - (k_R + k_P) x_3(0) , \quad (6c)$$

$$\mu x_4(0) = k_P x_3(0) - (k_R + k_{DV}) x_4(0) , \quad (6d)$$

where of course (see (4e) and (5))

$$f(x_1, x_2, x_3, x_4) \equiv f(0) = \frac{k_I [x_3(0) + \beta x_4(0)]}{x_1(0) + x_2(0) + x_3(0) + \beta x_4(0)} . \blacksquare \quad (6e)$$

The validity of this **Proposition 2.1** can be easily verified by inserting the solution (5) in the system (4) and by then taking advantage of the conditions (6).

Somewhat less trivial is to ascertain which are the *constraints* on the 4 initial data $x_n(0)$ and on the parameter μ —by solving the system of algebraic equations (6)—when we consider the model (4) for an *arbitrary* assignment of its 7 parameters $k_R, k_B, k_P, k_{DV}, k_I, \alpha, \beta$. Remarkably, as we show below, this turns out to be *explicitly* doable by purely *algebraic* operations.

Since all the 5 eqs. (6) are invariant under a *common* rescaling of the 4 initial data $x_n(0)$, it is convenient to assume that one of them—say $x_4(0)$ —can be *arbitrarily assigned*, and to focus on the *ratios* of the other 3 to that one, hence on the 3 quantities

$$r_m = x_m(0)/x_4(0) , \quad x_m(0) = r_m x_4(0) , \quad m = 1, 2, 3 ; \quad (7)$$

thereby replacing the 5 eqs. (6) with the following 5 equations:

$$\mu r_1 = \alpha k_R (r_3 + 1) , \quad (8a)$$

$$\mu r_2 = k_B r_1 + [k_B - F(r_1, r_2, r_3)] r_2 + [k_B + (1 - \alpha) k_R] (r_3 + 1) , \quad (8b)$$

$$\mu r_3 = F(r_1, r_2, r_3) r_2 - (k_R + k_P) r_3 , \quad (8c)$$

$$\mu = k_P r_3 - (k_R + k_{DV}) , \quad (8d)$$

where of course (above and hereafter)

$$F(r_1, r_2, r_3) = k_I (r_3 + \beta) / (r_1 + r_2 + r_3 + \beta) . \quad (8e)$$

It is now convenient—in order to get rid of the *nonlinear* function $F(r_1, r_2, r_3)$ —to sum the 2 eqs. (8b) and (8c), getting thereby

$$\mu (r_2 + r_3) = k_B + (1 - \alpha) k_R + k_B r_1 + k_B r_2 + (k_B - k_P - \alpha k_R) r_3 . \quad (9)$$

The 3 eqs. (8a), (8d) and (9) constitute now a system of 3 *linear algebraic* equations for the 3 unknowns r_1, r_2, r_3 , which can be easily solved. Indeed from (8d) we get

$$r_3 = (\mu + k_{DV} + k_R) / k_P ; \quad (10a)$$

then from (8a) and (10a) we get

$$r_1 = \alpha k_R (\mu + k_{DV} + k_P + k_R) / (\mu k_P) ; \quad (10b)$$

and then from (9), (10a) and (10b) we get

$$r_2 = -\frac{\mu + k_{DV}}{k_P} - \frac{\mu - k_B + k_{DV}}{\mu - k_B} - \frac{[(1 + \alpha)\mu + \alpha(k_{DV} + k_P)] k_R + \alpha(k_R)^2}{\mu k_P} . \quad (10c)$$

Note that these are *explicit* expressions of the 3 parameters r_m in terms of the 6 parameters k_R , k_B , k_P , k_{DV} , k_I , α of the system (4), and moreover of the parameter μ featured by the solution (5) (where of course now $x_m(0) = r_m x_4(0)$ for $m = 1, 2, 3$, with $x_4(0)$ remaining as a *free* parameter).

Our remaining task in order to get the special solution (5) of the system (4) is to ascertain the permitted values of the parameter μ , as implied by inserting the following expression of $F(r_1, r_2, r_3)$ (obtained by inserting the 3 expressions (10) of r_1, r_2, r_3 in (8e)),

$$F(r_1, r_2, r_3) = \frac{-k_I(\mu - k_B)(\mu + k_{DV} + \beta k_P + k_R)}{k_P[(1 - \beta)(\mu - k_B) + k_{DV}]}, \quad (11)$$

into any one of the 2 eqs. (8b) or (8c). This yields the following *algebraic* equation of degree 4 (hence *explicitly solvable*) for the quantity μ :

$$\sum_{k=0}^4 (c_k \mu^k) = 0, \quad (12a)$$

with the following definitions of the 5 parameters c_k :

$$c_4 = k_I - k_P + \beta k_P, \quad (12b)$$

$$\begin{aligned} c_3 = & 2k_{DV}k_I - 2k_{DV}k_P + k_Ik_P - (k_P)^2 + 2k_Ik_R - 2k_Pk_R \\ & + \alpha k_Ik_R - k_B[k_I - (1 - \beta)k_P] + \beta k_P(k_{DV} + k_I + k_P + 2k_R), \end{aligned} \quad (12c)$$

$$\begin{aligned} c_2 = & (k_I - k_P) \left[(k_{DV})^2 + k_R(k_P + k_R) \right] + \alpha k_Ik_R(k_P + 2k_R) \\ & + \beta k_P \left[(k_I + k_R)(k_P + k_R) + \alpha k_Ik_R \right] \\ & + k_{DV} \{ (2 + \beta)k_Ik_P + 2k_I(k_R + \alpha k_R) - k_P[(2 - \beta)k_P + (3 - \beta)k_R] \} \\ & + k_B \{ (1 - \beta)k_P(k_P + 2k_R) + k_{DV}(-2k_I + k_P - \beta k_P) \\ & - k_I(k_P + 2k_R + \alpha k_R + \beta k_P) \}, \end{aligned} \quad (12d)$$

$$\begin{aligned} c_1 = & (k_{DV} + k_R) [k_{DV}k_P(k_I - k_P - k_R) + \alpha k_Ik_R(k_{DV} + k_P + k_R)] \\ & + \beta k_Ik_P [k_{DV}k_P + \alpha k_R(k_{DV} + k_P + k_R)] \\ & + k_B \{ -(k_{DV} + k_R) [k_{DV}k_I + (k_I - k_P)(k_P + k_R)] \\ & - \alpha k_Ik_R(2k_{DV} + k_P + 2k_R) \\ & - \beta k_P [(k_I + k_R)(k_P + k_R) + k_{DV}(k_I + k_P + k_R) + \alpha k_Ik_R] \}, \end{aligned} \quad (12e)$$

$$c_0 = -\alpha k_B k_I k_R (k_{DV} + k_P + k_R) (k_{DV} + k_R + \beta k_P). \quad (12f)$$

Remark 2.2. For completeness let us mention that the results reported just above require the validity of the following *inequalities*:

$$\mu \neq 0, \quad k_P \neq 0, \quad (1 - \beta)(k_B - \mu) - k_{DV} \neq 0. \quad \blacksquare \quad (13)$$

Some of these expressions of the 5 parameters c_k —see (12)—are rather cumbersome (albeit quite explicit), featuring the 7 *a priori arbitrary* parameters k_R , k_B , k_P , k_{DV} , k_I , α , β characterizing the system (4); and of course much more complicated are the—in principle easily available—*explicit* expressions of the 4 roots μ_n ($n = 1, 2, 3, 4$) of the fourth-degree equation (12a). We do not consider useful to report these formulas in this paper; since analogous—much more practical—formulas can be easily obtained from eq. (12a) in *applicative* contexts, whenever the 7 *a priori arbitrary* parameters k_R , k_B , k_P , k_{DV} , k_I , α , β have been assigned specific numerical values, entailing, via the explicit expressions of the parameters c_k written above (see (12)), the corresponding numerical values of these parameters c_k , to be then inserted in (12a) before the standard task of solving this *quartic* equation is performed.

Remark 2.3. Let us finally mention that clearly, by setting

$$\mu = k_D, \quad (14)$$

one is looking—see (3a) and (5)—at the *equilibrium* solution

$$\tilde{x}_n(t) = \bar{x}_n, \quad \dot{\tilde{x}}_n(t) = 0, \quad n = 1, 2, 3, 4, \quad (15)$$

of the original pandemic system (1), as given by the formulas (implied via (7) by (10))

$$\bar{x}_1 = \bar{x}_4 [\alpha k_R (k_D + k_{DV} + k_P + k_R) / (k_D k_P)], \quad (16a)$$

$$\bar{x}_2 = \bar{x}_4 \left\{ -\frac{k_D + k_{DV}}{k_P} - \frac{k_D - k_B + k_{DV}}{k_D - k_B} \right. \quad (16b)$$

$$\left. - \frac{[(1 + \alpha) k_D + \alpha (k_{DV} + k_P)] k_R + \alpha (k_R)^2}{k_P k_D} \right\}, \quad (16c)$$

$$\bar{x}_3 = \bar{x}_4 (k_D + k_{DV} + k_R) / k_P, \quad (16d)$$

where \bar{x}_4 is of course an *arbitrary* parameter. ■

Remark 2.4. Note that throughout this paper we assume that the 4 roots μ_n of the *quartic* algebraic equation (12a) are all different among themselves. ■

To conclude this **Section 2**, let us mention that the *special* solutions (5) are not very interesting in *applicative* contexts, since they imply that the 4 dependent variables $x_n(t)$ *all* evolve in the *same*, very *simple*, manner. But fortunately—as shown below—it is also possible to identify other—presumably more interesting—*explicit* solutions of the system of nonlinear ODEs (4).

3 A less simple solution: the linear combination of 2 exponentials

In this **Section 3** we investigate the following class of solutions of the system of ODEs (4):

$$x_n(t) = a_{n1} \exp(\mu_1 t) + a_{n2} \exp(\mu_2 t), \quad n = 1, 2, 3, 4, \quad (17a)$$

where μ_1 and μ_2 are 2 *different* roots of eq. (12); while corresponding values for the 8 time-independent parameters a_{n1} and a_{n2} are obtained below.

Remark 3.1. Since there are 4 (assumedly *different*) solutions μ of the fourth-order algebraic eq. (12a), there are $4 \cdot 3/2 = 6$ different assignments of the pair of values μ_1, μ_2 . Note moreover that, even if the 7 parameters $k_R, k_B, k_P, k_{DV}, k_I, \alpha, \beta$ of the system of ODEs (4) are *all real* numbers (as is of course the case in the pandemics case), the 4 solutions μ_n of the fourth-order algebraic eq. (12a) need *not* be *real* numbers; but if the 7 parameters of the system of ODEs (4) are *all real* numbers, then *non-real* solutions of the algebraic eq. (12a) must be present in *complex conjugate* pairs. ■

Remark 3.2. Note that we are again assuming, throughout this **Section 3**, that the quantity $f(x_1, x_2, x_3, x_4)$ in the system (4) is *time-independent*, hence equal to its value at the initial time $t = 0$ (see (6e)); although this property, which was obvious in the treatment of **Section 2** (see (4e) and (5))—and which is essential to justify the existence of the subclass of solutions (17a)—is now instead far from obvious: indeed conditions for it to hold—involving the *initial data* of these solutions, and also 1 *constraint* on the parameters of the system (4)—shall have to be ascertained, see below. ■

The 4 eqs. (17a) involve of course the following 4 relations among the 8 parameters a_{n1} and a_{n2} and the 4 initial data $x_n(0)$:

$$x_n(0) = a_{n1} + a_{n2}, \quad n = 1, 2, 3, 4. \quad (17b)$$

The assumption (see **Remark 3.2**) that the function $f(x_1, x_2, x_3, x_4)$ be *time-independent* implies that the system (4) is again a *linear* system of 4 ODEs with *time-independent* parameters; hence each of the 2 exponential functions in the right-hand side of the *ansatz* (17a) must satisfy (as it were, *separately*) the system (4). Therefore each of the 2 sets of 4 parameters a_{n1} and a_{n2} must satisfy the same requirements—see the 4 eqs. (6)—satisfied by the initial data $x_n(0)$ in the treatment of the previous **Section 2**; namely there must now hold the 8 relations

$$\mu_\ell a_{1\ell} = \alpha k_R (a_{3\ell} + a_{4\ell}), \quad \ell = 1, 2, \quad (18a)$$

$$\mu_\ell a_{2\ell} = k_B a_{1\ell} + [k_B - f(0)] a_{2\ell} + [k_B + (1 - \alpha) k_R] [a_{3\ell} + a_{4\ell}] , \quad \ell = 1, 2 , \quad (18b)$$

$$\mu_\ell a_{3\ell} = f(0) a_{2\ell} - (k_R + k_P) a_{3\ell} , \quad \ell = 1, 2 , \quad (18c)$$

$$\mu_\ell a_{4\ell} = k_P a_{3\ell} - (k_R + k_{DV}) a_{4\ell} , \quad \ell = 1, 2 ; \quad (18d)$$

where of course we again set

$$f(x_1, x_2, x_3, x_4) \equiv f(0) = \frac{k_I [x_3(0) + \beta x_4(0)]}{x_1(0) + x_2(0) + x_3(0) + \beta x_4(0)} ; \quad (18e)$$

but now with the 4 initial data $x_n(0)$ related to the 2 parameters a_{n1} and a_{n2} by the eq. (17b).

We can therefore now proceed in close analogy to the treatment of the previous **Section 2**, introducing 6 parameters $b_{m\ell}$ via the following position:

$$a_{m\ell} = b_{m\ell} a_{4\ell} , \quad b_{m\ell} = a_{m\ell} / a_{4\ell} , \quad m = 1, 2, 3, \quad \ell = 1, 2 . \quad (19)$$

These 6 parameters $b_{m\ell}$ ($m = 1, 2, 3, \ell = 1, 2$) are then *explicitly* expressed in terms of the parameters of the system (4) as follows (see (10)):

$$b_{1\ell} = \alpha k_R (\mu_\ell + k_{DV} + k_P + k_R) / (k_P \mu_\ell) , \quad \ell = 1, 2 , \quad (20a)$$

$$b_{2\ell} = -\frac{\mu_\ell + k_{DV}}{k_P} - \frac{\mu_\ell - k_B + k_{DV}}{\mu_\ell - k_B} - \frac{[(1 + \alpha) \mu_\ell + \alpha (k_{DV} + k_P)] k_R + \alpha (k_R)^2}{k_P \mu_\ell} , \quad \ell = 1, 2 , \quad (20b)$$

$$b_{3\ell} = (\mu_\ell + k_{DV} + k_R) / k_P , \quad \ell = 1, 2 ; \quad (20c)$$

with a_{41} and a_{42} remaining as 2 *free* parameters.

To complete the treatment of this case, it is necessary to identify the *constraints* on the parameters of the system (4) and on the parameters of the solution under present consideration, see (17a), which are necessary in order to comply with the requirement—essential for our treatment—that the quantity $h(t)$, related to the quantity $f(x_1, x_2, x_3, x_4)$, see (4e), by the simple relation

$$f(x_1, x_2, x_3, x_4) = k_I h(t) \quad (21a)$$

—hence reading as follows,

$$h(t) = \frac{x_3(t) + \beta x_4(t)}{x_1(t) + x_2(t) + x_3(t) + \beta x_4(t)} \quad (21b)$$

—be *time-independent*: therefore given in terms of the initial data as follows:

$$h(t) = h(0) = \frac{x_3(0) + \beta x_4(0)}{x_1(0) + x_2(0) + x_3(0) + \beta x_4(0)} . \quad (21c)$$

To fulfill this task, we now note that the solutions $x_n(t)$ under consideration in this **Section 3** are defined by the relations (17a), hence their insertion in the definition (21b) implies the following expression of $h(t)$:

$$h(t) = \frac{a_{31} + \beta a_{41} + (a_{32} + \beta a_{42}) \exp[(\mu_2 - \mu_1)t]}{a_{11} + a_{21} + a_{31} + \beta a_{41} + (a_{12} + a_{22} + a_{32} + \beta a_{42}) \exp[(\mu_2 - \mu_1)t]} . \quad (21d)$$

It is therefore easily seen that the requirement that this expression of $h(t)$ be *time-independent* implies that the 8 parameters $a_{n\ell}$ ($n = 1, 2, 3, 4; \ell = 1, 2$) satisfy the following *single constraint* on the 8 parameters $a_{n\ell}$:

$$(a_{12} + a_{22})(a_{31} + \beta a_{41}) - (a_{11} + a_{21})(a_{32} + \beta a_{42}) = 0 ; \quad (22)$$

entailing then that

$$h(t) = h(0) = \frac{a_{31} + \beta a_{41}}{a_{11} + a_{21} + a_{31} + \beta a_{41}} = \frac{a_{32} + \beta a_{42}}{a_{12} + a_{22} + a_{32} + \beta a_{42}} . \quad (23)$$

By inserting the formulas (19) in (22) we then get, for the 6 parameters $b_{m\ell}$ ($m = 1, 2, 3$; $\ell = 1, 2$), the following *single constraint*:

$$(b_{12} + b_{22})(b_{31} + \beta) - (b_{11} + b_{21})(b_{32} + \beta) = 0 ; \quad (24)$$

and, via (20), we finally get the following *single constraint* on the 7 parameters of the system (4) for the existence of the solution (17a):

$$(1 - \beta)(k_B)^2 - k_B k_{DV} + k_{DV}(k_{DV} + \beta k_P + k_R + \mu_1) + [k_{DV} + (1 - \beta)\mu_1]\mu_2 - (1 - \beta)k_B(\mu_1 + \mu_2) = 0 , \quad (25a)$$

provided there hold the following inequalities:

$$k_P \neq 0 , \quad k_B \neq \mu_1 , \quad k_B \neq \mu_2 , \quad \mu_1 \neq \mu_2 . \quad (25b)$$

Remark 3.3. Let us recall that there are in general 6 different versions of the *constraint* (25a) due to the 6 different possible selections of the 2 roots μ_1 and μ_2 (see **Remark 3.1**); and that the simplicity of this formula (25a) as providing a *constraint* on the 7 parameters of the system (4) is somewhat misleading, due to the (*explicit* but) *quite complicated* dependence on these parameters of the solutions μ_1 and μ_2 of the *fourth-degree* algebraic equation (12a). However—as already mentioned above—all these complicated relations (including those yielding the *initial data* of the class of solutions considered in this **Section 3**) become much more easily managed whenever any 6 of the 7 *a priori arbitrary* parameters featured by the system (4) are assigned *specific numerical* values, so that the remaining task left is to ascertain the values of the 7-th parameter implied by the constraint (25a) (as well as those characterizing the *initial data* $x_n(0)$ of the class of solutions considered in this **Section 3**), thereby identifying the corresponding class of systems (4) featuring the simple *explicit* solutions of type (17a). ■

4 Solutions which are the linear superposition of 3 or 4 exponentials

In this **Section 4** we treat the subclass of solutions of the system (4) whose time-evolution is a *linear superposition* of 3 or 4 exponentials.

Remark 4.1. Hence, throughout this **Section 4**, we assume that the quantity $f(x_1, x_2, x_3, x_4)$ in (4) is *time-independent*; although this property is—as in **Section 3**: see for instance **Remark 3.2**—far from obvious: indeed conditions for it to hold—involving the *initial data* of these subclass of solutions, as well as *constraints* on the parameters of the system (4)—shall have to be ascertained, see below. ■

4.1 Solutions which are the linear superposition of 3 exponentials

In this **Subsection 4.1** we investigate the following class of solutions of the system of ODEs (4):

$$x_n(t) = a_{n1} \exp(\mu_1 t) + a_{n2} \exp(\mu_2 t) + a_{n3} \exp(\mu_3 t) , \quad n = 1, 2, 3, 4 , \quad (26a)$$

where μ_1, μ_2, μ_3 are 3 different roots of the 4th-degree algebraic eq. (12a); while corresponding values for the 12 time-independent parameters a_{n1}, a_{n2}, a_{n3} are obtained below.

Of course these formulas (26a) imply the following relations among the 4 initial data $x_n(0)$ and the 12 parameters a_{nj} ($n = 1, 2, 3, 4$; $j = 1, 2, 3$):

$$x_n(0) = a_{n1} + a_{n2} + a_{n3} , \quad n = 1, 2, 3, 4 . \quad (26b)$$

Remark 4.1.1. Clearly symbols such as $x_n, \mu_{n\ell}, a_{n\ell}$ need not have the same significance nor the same values when appearing in different **Sections** or **Subsections** of this paper. But of course the statements made in **Remark 3.1** concerning the possibility that not all the 4 roots of the fourth-order algebraic eq. (12a) be *real* numbers are generally valid. ■

Remark 4.1.2. Since there are 4 (assumedly *different*) solutions μ of the fourth-order algebraic eq. (12a), there are 4 different selections—from the *quartet* of solutions μ_n of eq. (12a)—of the *trio* of values μ_1, μ_2, μ_3 in the *ansatz* (26a). ■

Let us now proceed in close analogy to the treatment provided in **Section 3**. Again we assume that the quantity f in the right-hand sides of the ODEs (4b) and (4c) is a *time-independent* parameter, up to identifying below conditions on the parameters of the system (4) and on the initial data of the solution (26) under consideration which are *sufficient* to guarantee—as it were, *a posteriori*—that this be the case; hence that the system of ODEs (4) be equivalent to a system of 4 *linear* ODEs, featuring independent solutions $a \exp(\mu t)$ each depending exponentially on the independent variable t (which can therefore be added without loosing the property to satisfy the system of ODEs (4)).

We thus obtain—in analogy to the 8 relations (18)—the following 12 relations:

$$\mu_j a_{1j} = \alpha k_R (a_{3j} + a_{4j}) , \quad j = 1, 2, 3 , \quad (27a)$$

$$\mu_j a_{2j} = k_B a_{1j} + [k_B - f(0)] a_{2j} + [k_B + (1 - \alpha) k_R] (a_{3j} + a_{4j}) , \quad j = 1, 2, 3 , \quad (27b)$$

$$\mu_j a_{3j} = f(0) a_{2j} - (k_R + k_P) a_{3j} , \quad j = 1, 2, 3 , \quad (27c)$$

$$\mu_j a_{4j} = k_P a_{3j} - (k_R + k_{DV}) a_{4j} , \quad j = 1, 2, 3 . \quad (27d)$$

Next we set (in analogy to (19))

$$a_{mj} = b_{mj} a_{4j} , \quad b_{mj} = a_{mj} / a_{4j} , \quad m = 1, 2, 3 , \quad j = 1, 2, 3 , \quad (28)$$

getting thereby the following 9 relations:

$$b_{1j} = \alpha k_R (\mu_j + k_{DV} + k_P + k_R) / (k_P \mu_j) , \quad j = 1, 2, 3 , \quad (29a)$$

$$b_{2j} = -\frac{\mu_j + k_{DV}}{k_P} - \frac{\mu_j - k_B + k_{DV}}{\mu_j - k_B} - \frac{[(1 + \alpha) \mu_j + \alpha (k_{DV} + k_P)] k_R + \alpha (k_R)^2}{k_P \mu_j} , \quad j = 1, 2, 3 , \quad (29b)$$

$$b_{3j} = (\mu_j + k_{DV} + k_R) / k_P , \quad j = 1, 2, 3 ; \quad (29c)$$

with a_{41} , a_{42} , a_{43} remaining as 3 *free* parameters.

We must now investigate the restrictions on the parameters a_{mj} implied by the requirement that the quantity $h(t)$ be *time-independent*. The analogous formula to (21d) now reads as follows:

$$h(t) = numh(t) / denh(t) , \quad (30a)$$

$$numh(t) = a_{31} + \beta a_{41} + (a_{32} + \beta a_{42}) \exp[(\mu_2 - \mu_1)t] + (a_{33} + \beta a_{43}) \exp[(\mu_3 - \mu_1)t] , \quad (30b)$$

$$denh(t) = a_{11} + a_{21} + a_{31} + \beta a_{41} + (a_{12} + a_{22} + a_{32} + \beta a_{42}) \exp[(\mu_2 - \mu_1)t] + (a_{13} + a_{23} + a_{33} + \beta a_{43}) \exp[(\mu_3 - \mu_1)t] . \quad (30c)$$

It is then easily seen that the condition (22) is now replaced by the following 2 restrictions:

$$(a_{11} + a_{21}) (a_{3k} + \beta a_{4k}) = (a_{1k} + a_{2k}) (a_{31} + \beta a_{41}) , \quad k = 2, 3 . \quad (31)$$

Hence, after the change of parameters (28), we get the following 2 *constraints*,

$$(b_{11} + b_{21}) (b_{3k} + \beta) = (b_{1k} + b_{2k}) (b_{31} + \beta) , \quad k = 2, 3 , \quad (32)$$

on the 9 parameters b_{mj} ($m = 1, 2, 3; j = 1, 2, 3$). Which, via the expressions (29) of these 9 parameters, entail the following 2 *constraints* on the original 7 parameters of the system (4):

$$(1 - \beta) (k_B)^2 + (k_{DV})^2 + (1 - \beta) \mu_1 \mu_2 + k_{DV} (\beta k_P + k_R + \mu_1 + \mu_2) - k_B [k_{DV} + (1 - \beta) (\mu_1 + \mu_2)] = 0 , \quad (33a)$$

$$\begin{aligned}
& k_{DV} (k_{DV} + \beta k_P + k_R + \mu_1) + [k_{DV} + (1 - \beta) \mu_1] \mu_3 \\
& - (1 - \beta) k_B (\mu_1 + \mu_3) + k_B [(1 - \beta) k_B - k_{DV}] = 0.
\end{aligned} \tag{33b}$$

While of course the initial data $x_n(0)$ of the solution under consideration in this **Subsection 4.1** are explicitly given by the formulas (26b) with (28) and (29).

4.2 Solutions which are the linear superposition of 4 exponentials

The treatment in this **Subsection 4.2** is quite terse, since it is quite analogous to that provided above in **Subsection 4.1**; hence we only report the key formulas which play an analogous role to the key formulas in **Subsection 4.1**.

Instead of (26) we now have

$$x_n(t) = \sum_{q=1}^4 [a_{nq} \exp(\mu_q t)] , \quad n = 1, 2, 3, 4 , \tag{34a}$$

$$x_n(0) = \sum_{q=1}^4 (a_{nq}) , \quad n = 1, 2, 3, 4 . \tag{34b}$$

In place of the 12 equations (27) we now have the following 16 equations:

$$\mu_q a_{1q} = \alpha k_R (a_{3q} + a_{4q}) , \quad q = 1, 2, 3, 4 , \tag{35a}$$

$$\begin{aligned}
\mu_q a_{2q} &= k_B a_{1q} + [k_B - f(0)] a_{2q} + [k_B + (1 - \alpha) k_R] (a_{3q} + a_{4q}) , \\
q &= 1, 2, 3, 4 ,
\end{aligned} \tag{35b}$$

$$\mu_q a_{3q} = f(0) a_{2q} - (k_R + k_P) a_{3q} , \quad q = 1, 2, 3, 4 , \tag{35c}$$

$$\mu_q a_{4q} = k_P a_{3q} - (k_R + k_{DV}) a_{4q} , \quad q = 1, 2, 3, 4 . \tag{35d}$$

Likewise, in place of the 9 equations (28), we now write the 12 relations

$$a_{jq} = b_{jq} a_{4j} , \quad b_{jq} = a_{jq} / a_{4j} , \quad j = 1, 2, 3 , \quad q = 1, 2, 3, 4 , \tag{36}$$

getting thereby, from (35), the following 12 relations (analogous to (29)):

$$b_{1q} = \alpha k_R (\mu_q + k_{DV} + k_P + k_R) / (k_P \mu_q) , \quad q = 1, 2, 3, 4 , \tag{37a}$$

$$\begin{aligned}
b_{2q} &= -\frac{\mu_q + k_{DV}}{k_P} - \frac{\mu_q - k_B + k_{DV}}{\mu_q - k_B} \\
&- \frac{[(1 + \alpha) \mu_q + \alpha (k_{DV} + k_P)] k_R + \alpha (k_R)^2}{k_P \mu_q} , \quad q = 1, 2, 3, 4 ,
\end{aligned} \tag{37b}$$

$$b_{3q} = (\mu_q + k_{DV} + k_R) / k_P , \quad q = 1, 2, 3, 4 ; \tag{37c}$$

with a_{41} , a_{42} , a_{43} , a_{44} remaining as 4 *free* parameters.

Next come the *constraints* on the parameters of the system (4) needed in order that the more general solution (34a), when inserted in the definition (4e) of the function $f(t)$, hence now reading

$$f(t) = \frac{k_I \sum_{q=1}^4 [(a_{3q} + \beta a_{4q}) \exp(\mu_q t)]}{\sum_{q=1}^4 [(a_{1q} + a_{2q} + a_{3q} + \beta a_{4q}) \exp(\mu_q t)]} , \tag{38a}$$

—or, equivalently, see (36)—

$$f(t) = \frac{k_I \sum_{q=1}^4 [(b_{3q} + 1) \exp(\mu_q t)]}{\sum_{q=1}^4 [(b_{1q} + b_{2q} + b_{3q} + \beta) \exp(\mu_q t)]} , \tag{38b}$$

be *time-independent*.

And since it is easily seen that

$$f(t) = \frac{k_I (b_{31} + 1) \varphi(t)}{(b_{11} + b_{21} + b_{31} + \beta)}, \quad (39a)$$

with

$$\varphi(t) = \frac{1 + \sum_{q=2}^4 \left\{ \left(\frac{b_{3q}+1}{b_{31}+1} \right) \exp [(\mu_q - \mu_1) t] \right\}}{1 + \sum_{q=2}^4 \left\{ \left(\frac{b_{1q}+b_{2q}+b_{3q}+\beta}{b_{11}+b_{21}+b_{31}+\beta} \right) \exp [(\mu_q - \mu_1) t] \right\}}, \quad (39b)$$

the requirement that $f(t)$ be *time-independent* amounts to the following 3 constraints:

$$\frac{b_{3q} + 1}{b_{31} + 1} = \frac{b_{1q} + b_{2q} + b_{3q} + \beta}{b_{11} + b_{21} + b_{31} + \beta}, \quad q = 2, 3, 4, \quad (40)$$

which clearly entail—via the expressions (29)—3 corresponding *constraints* on the parameters of the original model (4).

5 Concluding remarks

In this paper we have identified certain solutions of the *pandemic* model introduced in the paper [1]; these solutions—and the constraints on the parameters of the model required for their validity—are all identified by *algebraic* equations which can be *explicitly* solved; we did not report the corresponding explicit formulas because they are so complicated to be hardly useful when written for *a priori arbitrary* assignments of the parameters of the pandemic model, while they can instead be easily managed for any *specific numerical* assignment of these parameters. We therefore leave the utilization of these findings to the interested pandemics experts.

Additional solutions—more special but perhaps displaying more interesting evolutions—correspond to the special cases in which the algebraic quartic-equation (12) features 4 roots μ_n which are *not* all different among themselves. This case shall perhaps be eventually treated in a separate paper by ourselves or others.

Acknowledgments

We like to acknowledge with thanks 3 grants, facilitating our collaboration—mainly developed via e-mail exchanges—by making it possible for FP to visit three times the Department of Physics of the University of Rome "La Sapienza": two granted by that University, and one granted jointly by the Istituto Nazionale di Alta Matematica (INdAM) of that University and by the International Institute of Theoretical Physics (ICTP) in Trieste in the framework of the ICTP-INdAM "Research in Pairs" Programme. Finally, we also like to thank Fernanda Lupinacci who, in these difficult times—with extreme efficiency and kindness—facilitated all the arrangements necessary for the presence of FP with her family in Rome.

References

- [1] Nash Rochman, Yu. I. Wolf, Eugene V. Koonin, "Evolution of human respiratory virus epidemics", F1000Research 2021, **10:447**; <https://doi.org/19.12688/f1000research.53392.2>.
- [2] F. Calogero and F. Payandeh, "Explicitly solvable systems of first-order ordinary differential equations with homogeneous right-hand sides, and their periodic variants", arXiv:2106.06634v1 [math.DS] 11 Jun 2021; Theor. Math. Phys. (in press).