PARAMETER REGION FOR MULTISTATIONARITY IN n-SITE PHOSPHORYLATION NETWORKS

ELISENDA FELIU¹, NIDHI KAIHNSA², TIMO DE WOLFF³, OĞUZHAN YÜRÜK⁴

ABSTRACT. Multisite phosphorylation is a signaling mechanism well known to give rise to multiple steady states, a property termed multistationarity. When phosphorylation occurs in a sequential and distributive manner, we obtain a family of networks indexed by the number of phosphorylation sites n. This work addresses the problem of understanding the parameter region where this family of networks displays multistationarity, by focusing on the projection of this region onto the set of kinetic parameters. The problem is phrased in the context of real algebraic geometry and reduced to studying whether a polynomial, defined as the determinant of a parametric matrix of size three, attains negative values over the positive orthant. The coefficients of the polynomial are functions of the kinetic parameters. For any n, we provide sufficient conditions for the polynomial to be positive and hence, preclude multistationarity, and also sufficient conditions for it to attain negative values and hence, enable multistationarity. These conditions are derived by exploiting the structure of the polynomial, its Newton polytope, and employing circuit polynomials. A relevant consequence of our results is that the set of kinetic parameters that enable or preclude multistationarity are both connected for all n.

1. Introduction

Given a dynamical system, the notion of *multistationarity* refers to the existence of multiple steady states or equilibrium points. Often in biological processes, multistationarity has been associated with and found relevant for cellular decision making processes [28, 33, 46]. This has led to an extensive interest in developing methods to identify whether parametric models arising from reaction networks admit multistationarity, and to determine and understand the parameter region where this occurs, e.g. [1, 8, 9, 11, 12, 14–17, 32, 36, 45].

In this work, we focus on understanding the rate parameters that allow or restrict the multistationarity in a phosphorylation and dephosphorylation network where a substrate has n phosphorylation sites. For two sites, that is, for n = 2, we obtain the so-called dual phosphorylation cycle

(1)
$$E + S_0 \xrightarrow{\kappa_1} ES_0 \xrightarrow{\kappa_3} E + S_1 \xrightarrow{\kappa_7} ES_1 \xrightarrow{\kappa_9} E + S_2$$

$$F + S_2 \xrightarrow{\kappa_{10}} FS_2 \xrightarrow{\kappa_{12}} F + S_1 \xrightarrow{\kappa_4} FS_1 \xrightarrow{\kappa_6} F + S_0.$$

In this system, S, E, and F denote substrate, kinase, and phosphatase, respectively. Kinase and phosphatase catalyze the phosphorylation and dephosphorylation of the substrate. The subscripts 0, 1, and 2 of the phosphoforms S_0 , S_1 , and S_2 denote the number of phosphorylated sites of the substrate S.

Over the years, under the assumption of mass-action kinetics, the process of dual phosphorylation has been studied rigorously by means of polynomial ordinary differential equations (ODEs) [1, 7, 10, 17]. Besides its relevance in biology, this network has become testing grounds for developing various mathematical methods focused towards understanding multistationarity and dynamics of reaction networks in general. In [17], the authors studied the region of reaction rate constants (the $\kappa's$ in (1)) that yield multistationarity, giving a description of the boundary between the region of parameters that enable

¹Department of Mathematical Sciences, University of Copenhagen. efeliu@math.ku.dk

²Division of Applied Mathematics, Brown University. nidhi_kaihnsa@brown.edu

 $^{^3}$ Institute of Analysis and Algebra, TU Braunschweig. t.de-wolff@tu-braunschweig.de

⁴Chair of Discrete Mathematics/Geometry, Technische Universität Berlin. yuruk@math-tu.berlin.de

²⁰¹⁰ Mathematics Subject Classification. 92Bxx, 14Pxx, 37N25, 52B20, 90C26.

 $Key\ words\ and\ phrases.$ Phosphorylation network, Multistationarity, Parameter Estimation, Chemical reaction network, Real algebraic geometry, Circuit polynomial, Connectedness.

multistationarity and the region of parameters that preclude multistationarity. Moreover, it was shown that both regions are path connected.

In this article we explore the general n-phosphorylation network, i.e., the substrate has n sites available for phosphorylation, and the network takes the following form:

$$(2) \qquad E + S_0 \xrightarrow{\kappa_1} ES_0 \xrightarrow{\kappa_3} E + S_1 \cdots \longrightarrow E + S_i \xrightarrow{\kappa_{6i+1}} ES_i \xrightarrow{\kappa_{6i+3}} E + S_{i+1} \cdots \\ \cdots \longrightarrow E + S_{n-1} \xrightarrow{\kappa_{6n-5}} ES_{n-1} \xrightarrow{\kappa_{6n-3}} E + S_n \\ F + S_n \xrightarrow{\kappa_{6n-2}} FS_n \xrightarrow{\kappa_{6n}} F + S_{n-1} \cdots \longrightarrow F + S_{i+1} \xrightarrow{\kappa_{6i+4}} FS_{i+1} \xrightarrow{\kappa_{6i+6}} F + S_i \cdots \\ \cdots \longrightarrow F + S_1 \xrightarrow{\kappa_5} FS_1 \xrightarrow{\kappa_6} F + S_0.$$

Multisite phosphorylation, or more general, multisite posttranslational modification, is a ubiquitious process in cell signaling. It is believed that approximately 30% of all proteins in humans undergo phosphorylation [6], and there are proteins having more than 150 phosphorylation sites [24]. The order in which the sites are phosphorylated, or how many encounters with the enzyme are required for the phosphorylation of the multiple sites, give rise to different mechanisms [39]. Network (2) is a representative model where phosphorylation and dephosphorylation occur sequentially, that is, in a given order, and each encounter leads to the modification of one site, that is, it is distributive. For this mechanism it is known that the network has finitely many positive steady states (in a stoichiometric compatibility class) for any choice of reaction rate constants κ , specifically between 1 and 2n-1 [44], and that the possible number increases with n [42, 44]. It is conjectured that there exist parameter values for which the system has 2n-1 positive steady states, but this has only been established for $n \leq 4$ [21]. For general n, it has been proven that there exist parameter values for which the system admits n+1 (n even) or n (n odd) positive steady states with half plus one of them asymptotically stable [18, 44]. These properties illustrate that the dynamics of the system become qualitatively richer as n increases, but while much is known about the case n=2, very little has been established about the dependence of the dynamics on the choice of parameters for general n.

With this work we address the problem of determining for which reaction rate constants κ_i , $i \in \{1, \ldots, 6n\}$, the *n*-phosphorylation network can have multiple steady states, that is, there exist total amounts of kinase, phosphatase and substrate for which the system has at least two positive steady states. In other words, we study the projection of the parameter region of multistationarity onto the subset of parameters consisting of reaction rate constants. In [1] the authors study the projection of this region onto a different subset of parameters, including the three total amounts and some of the reaction rate constants. Their methods are similar in spirit to ours: they rely on connecting properties of the Newton polytope associated to a parametric system of three equations in three variables and certain sign combinations, to obtain a lower bound on the maximal number of positive solutions the system can attain; see also [2, 23].

Addressing this question for a fixed n requires challenging computations, since the number of parameters and variables become very large as n increases. However, for any n, the differential equation system associated with the network admits exactly three linear first integrals and the set of steady states admits a nice parametrization. These key facts allow us to reduce the study of multistationarity to studying whether a polynomial in three variables attains negative values over the positive orthant (Proposition 3.3). This question can in turn be addressed by exploring the Newton polytope associated with the polynomial (Theorem 3.5), and in the end, the multistationarity problem can be reduced to studying the signs that a bivariate polynomial attains (Corollary 3.9).

With this in place, we proceed similarly to the study of the case n=2 in [17], and relate the signs the polynomial attains to the signs of the coefficients, which depend on the κ 's, and to the point configuration of the exponents of the polynomial. Imposing that at least one of the coefficients corresponding to vertices of the Newton polytope is negative, we obtain sufficient conditions for multistationarity (Theorem 4.1). However, these are not necessary as Theorem 4.3 shows: multistationarity can arise even when the conditions in Theorem 4.1 are not satisfied and only points in the interior of the Newton polytope have negative coefficients.

To preclude multistationarity, we note that a polynomial can be nonnegative even if some points in the interior of its Newton polytope have negative coefficients. We use SONC (sums of nonnegative circuit) polynomials as a symbolic nonnegativity certificate in order to describe a non-empty region in the parameter space that ensures the monostationarity of the system, and where the polynomial has some of the coefficients negative (Theorem 5.3).

Our results find subregions of the region of multistationarity and of the region of monostationarity, and indicate that the set of rate parameters that enable multistationarity is not simply found by looking at the sign of the coefficients of the polynomial. For the parameters that fall outside these subregions, our tools cannot characterize what happens, and this asks for the development of new techniques.

Even if a full description of the projection of the parameter region of multistationarity of the n-phosphorylation network onto the set of reaction rate constants is out of reach, we can still conclude that it is path-connected. The path-connected components of the parameter region of multistationarity can be thought to represent different biological mechanisms that give rise to multistationarity, as it has recently been brought up in [32, 40]. For example, if the region of multistationarity is connected, then the set of such parameters cannot be classified into two groups corresponding to different mechanisms. This has been proven to be the case for the dual phosphorylation cycle in [40], but it remains open for n > 2.

We study connectivity in Section 6, and show in Theorem 6.2 and Corollary 6.4 that the projection of the full parameter region of multistationarity onto the set of reaction rate constants is path connected, and the same holds for the region of monostationarity. To do that, we use the results in the previous sections to identify subregions that are path connected, and then show that any other point is joined to a point in the subregion via a continuous path. It is our hope that knowing that these projections are path connected might give a route to show that the full regions are also path connected.

Throughout this work, we often view the relevant polynomial, whose coefficients depend on the κ 's, as a polynomial in some of the parameters as well, and then employ standard techniques involving the Newton polytope. This is for example the case in showing path connectivity. These ideas should apply to other systems for which the study of multistationarity is reduced to the study of the signs a polynomial attains (those in the setting of [7]). Hence, the methods brought forward in this manuscript may have consequences beyond the understanding of the system in play.

2. Preliminaries

2.1. The parametric system. We consider the reaction network (2) and describe the ODE system that governs evolution of the concentration of the species in time [1, 18, 44]. We then lay the groundwork to derive a polynomial for this system whose positivity will determine the monostationarity of network (2). We note that network (2) belongs to a general class of reaction networks called MESSI systems, see [35]. MESSI systems admit specific decompositions of the sets of species and reactions, and these can be exploited to guarantee that relevant properties, such as absence of boundary steady states or persistence, hold. We will use this fact later on.

The concentrations of the species in the network are denoted as follows:

$$e = [E], \quad f = [F], \quad s_i = [S_i] \quad \text{for } i = 0, \dots, n,$$

 $u_i = [FS_{i+1}], \quad y_i = [ES_i] \quad \text{for } i = 0, \dots, n-1.$

Under the assumption of mass-action kinetics, the ODE system in $\mathbb{R}^{3n+3}_{>0}$ is as follows:

$$\frac{de}{dt} = -\sum_{i=0}^{n-1} \kappa_{6i+1} s_i e + \sum_{i=0}^{n-1} \kappa_{6i+1} (\kappa_{6i+2} + \kappa_{6i+3}) y_i,$$

$$\frac{df}{dt} = -\sum_{i=0}^{n-1} \kappa_{6i+4} s_{i+1} f + \sum_{i=0}^{n-1} (\kappa_{6i+5} + \kappa_{6i+6}) u_i,$$

$$\frac{ds_i}{dt} = -\kappa_{6i+1} s_i e \kappa_{6i-3} y_{i-1} + \kappa_{6i+2} y_i - \kappa_{6i-2} s_i f + \kappa_{6i+6} u_i + \kappa_{6i-1} u_{i-1}, \quad \text{for } i = 0, \dots, n,$$

$$\frac{dy_i}{dt} = \kappa_{6i+1} s_i e - (\kappa_{6i+2} + \kappa_{6i+3}) y_i, \quad \text{for } i = 0, \dots, n-1,$$

$$\frac{du_i}{dt} = \kappa_{6i+4} s_{i+1} f - (\kappa_{6i+5} + \kappa_{6i+6}) u_i, \quad \text{for } i = 0, \dots, n-1,$$

with the convention that $\kappa_j = 0$ if j > 6n or j < 0 (this becomes relevant only for $\frac{ds_0}{dt}$ and $\frac{ds_n}{dt}$). This system of ODEs admits three linear first integrals, such that the trajectories are confined in the following level sets for some E_{tot} , F_{tot} , $S_{\text{tot}} \ge 0$:

(4)
$$e + \sum_{i=0}^{n-1} y_i = E_{\text{tot}}, \qquad f + \sum_{i=0}^{n-1} u_i = F_{\text{tot}}, \qquad s_0 + \sum_{i=1}^n s_i + \sum_{i=0}^{n-1} y_i + \sum_{i=0}^{n-1} u_i = S_{\text{tot}}.$$

It is straightforward to verify that these equations arise from linear first integrals, and there are no additional independent linear first integrals, as the rank of the coefficient matrix of the polynomials on the right-hand side of (3) is three for all κ 's; alternatively, see e.g. [35, Theorem 3.2]. Each of the equations in (4) is called a conservation law and E_{tot} , F_{tot} and S_{tot} are the total amounts of E, F, and S, respectively. These are, therefore, taken nonnegative. The intersection of a level set with the nonnegative orthant is called a stoichiometric compatibility class. In particular, the ODE system has 3n+3 variables and for a given initial condition, the trajectory lies in a 3n dimensional subspace determined by the three linear constraints in (4).

The steady states of the system are found by setting the left-hand side of (3) to zero. Using the notation in [1], we consider the inverses of Michaelis-Menten constants of the (de)phosphorylation events

(5)
$$K_i = \frac{\kappa_{6i+1}}{\kappa_{6i+2} + \kappa_{6i+3}}, \qquad L_i = \frac{\kappa_{6i+4}}{\kappa_{6i+5} + \kappa_{6i+6}}, \qquad i = 0, \dots, n-1,$$

and define

(6)
$$T_{i} = \prod_{j=0}^{i} \frac{\kappa_{6j+3} K_{j}}{\kappa_{6j+6} L_{j}} = T_{i-1} \frac{\kappa_{6i+3} K_{i}}{\kappa_{6i+6} L_{i}}, \quad i = 0, \dots, n-1, \qquad T_{-1} = 1.$$

We note that here, the definition of K_i is the inverse of that given in [17] for n=2, but it turns out to be more convenient to use the notation of [1] for general n. For later use, we define the following composition of surjective maps $\eta: \mathbb{R}^{6n}_{>0} \to \mathbb{R}^{4n}_{>0} \to \mathbb{R}^{3n}_{>0}$

(7)
$$\kappa = (\kappa_1, \dots, \kappa_{6n}) \mapsto (\kappa_3, \kappa_6, \dots, \kappa_{6n}, K_0, \dots, K_{n-1}, L_0, \dots, L_{n-1}) \\ \mapsto (T_0, \dots, T_{n-1}, K_0, \dots, K_{n-1}, L_0, \dots, L_{n-1}).$$

As a consequence of the three conservation laws, the equations corresponding to e, f, and s_0 are redundant as they are linearly dependent on the rest. The steady state equations for s_{i+1}, y_i, u_i with $i = 0, \ldots, n-1$ are linear in these variables, and have a unique solution, which is positive provided e, f, s_0 are positive:

$$s_{i+1} = T_i e^{i+1} f^{-(i+1)} s_0$$
 $y_i = K_i T_{i-1} e^{i+1} f^{-i} s_0$ $u_i = L_i T_i e^{i+1} f^{-i} s_0$.

This gives rise to a parametrization of the set of positive steady states in the variables e, f, s_0 ,

(8)
$$\varphi_{\kappa} \colon \mathbb{R}^{3}_{>0} \longrightarrow \mathbb{R}^{3n+3}_{>0},$$

that is, the image of φ_{κ} is precisely the set of positive steady states. For later use, we consider the polynomial function

(9)
$$\psi_{\kappa} \colon \mathbb{R}^{3n+3} \longrightarrow \mathbb{R}^{3n+3}.$$

whose first three entries are $(e + \sum_{i=0}^{n-1} y_i, f + \sum_{i=0}^{n-1} u_i, s_0 + \sum_{i=1}^n s_i + \sum_{i=0}^{n-1} y_i + \sum_{i=0}^{n-1} u_i)$, that is, the left-hand side of the three conservation laws listed in (4), and the entries 4 to 3n + 3 are the right-hand side of the equations for $\frac{ds_i}{dt}$ for $i = 1, \ldots, n$, $\frac{dy_i}{dt}$ for $i = 0, \ldots, n-1$, and $\frac{du_i}{dt}$ for $i = 0, \ldots, n-1$ in (3), in this order.

In a fixed stoichiometric compatibility class with positive total amounts, this system has finitely many positive steady states for any choice of reaction rate constants κ , specifically between 1 and 2n-1 [44], see the Introduction.

For a given vector of reaction rate constants κ , if there exist positive E_{tot} , F_{tot} , and S_{tot} such that some stoichiometric compatibility class has at least two positive steady states, then we say that κ enables multistationarity. If this is not the case, then κ is said to preclude multistationarity. For n=2, the set of reaction rate constants that enable multistationarity was explored in detail in [17], building on preliminary results by Conradi and Mincheva [10]. In this work we go beyond the case n=2 and

determine subsets of reaction rate constants that enable or preclude multistationarity for general n, that is, for the network (2).

2.2. Nonnegative Circuit Polynomials. Let f be a polynomial function on \mathbb{R}^m . Determining if f(x) is nonnegative over \mathbb{R}^m is a classical problem in real algebraic geometry that appears naturally in different contexts, and prominently in polynomial optimization. It has been studied since the 19th century and is the subject of Hilbert's 17th problem [25]. The critical result for the present work, Theorem 2.2 below, builds on pioneering by Reznick in 1989 [37] where the theorem was proven for a special case. Another special case was shown by Fidalgo and Kovacec in [20]. In its full generality, Theorem 2.2 was first shown in 2012 by Pantea, Craciun and Köppl in [34]. That approach is particularly interesting, since the result is motivated by the study of reaction networks rather than nonnegativity of real polynomials. In this article, we use Theorem 2.2 in the terminology introduced by Iliman and de Wolff in 2016 [26]. Specifically, we certify nonnegativity by decomposing f(x) as a sum of nonnegative circuit polynomials (SONCs), and we use that nonnegative circuit polynomials are characterized by means of Theorem 2.2. The theory of SONCs has been subsequently developed further in [13, 22, 27].

For a multivariate polynomial $f(x) = \sum_{\alpha} c_{\alpha} x_1^{\alpha_1} \cdots x_m^{\alpha_m}$ with $\alpha = (\alpha_1, \dots, \alpha_m) \in \mathbb{Z}_{\geq 0}^m$, the Newton polytope, denoted by N(f), is the convex hull of the exponent vectors of the monomials with non-zero coefficients, i.e. $N(f) := \text{conv}(\{\alpha \in \mathbb{Z}_{\geq 0}^m : c_{\alpha} \neq 0\})$. Often we will also use $\text{coeff}(f, \alpha)$ to denote the coefficient of x^{α} in the polynomial f(x). We next define circuit polynomials.

Definition 2.1. A polynomial $f \in \mathbb{R}[x_1, \dots, x_m]$ is called a *circuit polynomial* if

$$f(x) = c_{\beta} x^{\beta} + \sum_{j=0}^{r} c_{\alpha(j)} x^{\alpha(j)},$$

where $\beta, \alpha(j) \in \mathbb{Z}_{\geq 0}^m$, $c_{\alpha(j)} \in \mathbb{R}_{>0}$ for $j = 0, \ldots, r$, $c_{\beta} \in \mathbb{R}$, and N(f) is a simplex with vertices $\alpha(0), \ldots, \alpha(r)$ such that the exponent β is in the strict interior of N(f). That is, $\beta = \sum_{j=0}^r \lambda_j \alpha(j)$ with $\lambda_j > 0$ and $\sum_{j=0}^r \lambda_j = 1$. We define the *circuit number* as

$$\Theta_f := \prod_{j=0}^r \left(\frac{c_{\alpha(j)}}{\lambda_j}\right)^{\lambda_j}.$$

Note that the circuit number Θ_f can be computed by solving a system of linear equations. In this article, we will consider circuit polynomials restricted to the nonnegative orthant, as the variables will be concentrations and reaction rate constants. We are interested in polynomials that take nonnegative (resp. positive) values over the positive orthant, which we call respectively nonnegative (resp. positive) polynomials. Thus, as opposed to the definition of circuit polynomials given in [26] where the focus was on nonnegativity over the whole real space, we do not restrict $\alpha(j)$ in Definition 2.1 to have even entries. We can do this because in the positive orthant, we can redefine the polynomial by replacing every variable with its square. Then, the theorem below is a consequence of [34, Thm. 3.6. and Prop. 3] and a direct specialization of Theorem 3.8 in [26].

Theorem 2.2. With the notation in Definition 2.1, a circuit polynomial f is nonnegative on $\mathbb{R}^m_{\geq 0}$ if and only if $c_{\beta} \geq -\Theta_f$.

The Motzkin polynomial, $x^4y^2 + x^2y^4 - 3x^2y^2 + 1$, is a classical example of a nonnegative polynomial which is not a sum of squares [29]. This is a circuit polynomial with $\beta = (2,2)$, and $\lambda_j = 1/3$ for all j (see e.g., [26]). The circuit number is 3 and hence, the polynomial is nonnegative by Theorem 2.2.

We note that the scenario studied by Reznick in [37] was the case where $c_{\alpha(j)} = \lambda_j$ for all $j \in \{1, \dots, n\}$. In [34], the authors do not only consider the case of simplex Newton polytopes/circuits, but discuss also the case of non-unique barycentric coordinates and give nonnegativity certificates. These functions were later called AGE (Arithmetic-Geometric Exponential) functions by Chandrasekaran and Shah [4] in the context of signomial programming, and a theorem similar to Theorem 2.2 was given in the framework called SAGE. While it was not obvious that SONC and SAGE describe the same cone of functions, we know nowadays that this is the case. We summarize this fact in the following remark for later reference.

Remark 2.3. Let f be a polynomial whose support contains only a single exponent that is not a vertex of N(f), and let the coefficients of f at the vertices of N(f) be positive. Then f is nonnegative if and only if it is SONC, see [43, Theorem 3.10], also [30, Theorem 11]. Such polynomials are examples of AGE as defined in [4], and they yield an equivalent description of SONC polynomials.

We also point out that computationally a SONC/SAGE certificate can be found using relative entropy programs (REP). This is a convex optimization program, where the target function and the constraints are linear or a sum of entropy functions. In general, these problems can be solved effectively, for example using standard solvers like MOSEK; for further details on REPs see e.g. [5, 31]. In the context of reaction networks the situation is more involved though, as the polynomials, for which we aim to certify nonnegativity, come with *symbolic* coefficients, and this does not allow to simply compute a SONC/SAGE certificate via solving an REP in the usual way; see also Remark 5.6.

3. A polynomial for multistationarity and the Newton polytope

We follow the approach in [17] and study the space of reaction rate constants that enable multistationarity after first reducing the problem to understanding the positivity of a multivariate polynomial for given reaction rates. This section is devoted to computing the general expression of the polynomial and exploring the structure of the associated Newton polytope to determine whether it is positive on the positive orthant. To this end, we apply Theorem 1 in [7], and obtain the following proposition. Here $J_{\psi_{\kappa}}$ denotes the Jacobian matrix of the polynomial function defined in (9) in 3n + 3 variables.

Proposition 3.1. Let $\kappa \in \mathbb{R}^{6n}_{>0}$, ψ_{κ} be the polynomial function in (9) and φ_{κ} the parametrization of the set of positive steady states from (8). Define the following rational function

$$q_{\kappa}(e, f, s_0) := (-1)^{3n} \det J_{\psi_{\kappa}}(\varphi_{\kappa}(e, f, s_0)).$$

The following holds:

- (i) If $q_{\kappa}(e, f, s_0) > 0$ for all $(e, f, s_0) \in \mathbb{R}^3_{>0}$, then κ precludes multistationarity.
- (ii) If $q_{\kappa}(e^*, f^*, s_0^*) < 0$ for some $(e^*, f^*, s_0^*) \in \mathbb{R}^3_{>0}$, then κ enables multistationarity. In this case the stoichiometric compatibility class containing $\varphi_{\kappa}(e^*, f^*, s_0^*)$ contains at least two positive steady states.

Proof. The statement is exactly the conclusion of Theorem 1 in [7], as the dimension of the stoichiometric compatibility classes is 3n. So we need to check that the two hypotheses to apply Theorem 1 in [7] hold. First, the reaction network (1) is dissipative, as it is conservative (each concentration appears in at least one conservation law that has all coefficients nonnegative). Second, we have to verify that the ODE system (3) has no boundary steady states (steady states with some entry equal to zero) when total amounts are positive. This follows from Theorem 3.15 in [35], as network (1) is a MESSI system that satisfies the theorem's hypotheses.

Based on Proposition 3.1, a strategy to determine the set of reaction rate constants that enable multistationarity consists of studying the signs q_{κ} attains on the positive orthant. To compute q_{κ} , for all n, the first step is to reduce the computation of the determinant of a matrix of size 3n + 3, to that of the determinant of a 3×3 matrix. Let Φ be the linear map given by the left-hand side of (4). Then the composition $\Phi \circ \varphi_{\kappa}$ is a function from $\mathbb{R}^3_{>0}$ to $\mathbb{R}^3_{>0}$.

Proposition 3.2. For any $\kappa \in \mathbb{R}^{6n}_{>0}$ and $(e, f, s_0) \in \mathbb{R}^3_{>0}$, the sign of $q_{\kappa}(e, f, s_0)$ agrees with the sign of $\det J_{\Phi_{\Omega/2n}}(e, f, s_0)$.

Proof. By the chain rule, it holds

(10)
$$J_{\psi_{\kappa}\circ\varphi_{\kappa}}(e,f,s_{0}) = J_{\psi_{\kappa}}(\varphi_{\kappa}(e,f,s_{0})) \cdot J_{\varphi_{\kappa}}(e,f,s_{0}).$$

We write $J_{\psi_{\kappa}}(\varphi_{\kappa}(e, f, s_0))$ in block form as

$$J_{\psi_{\kappa}}(\varphi_{\kappa}(e, f, s_0)) = \begin{pmatrix} A' & A'' \\ B' & B'' \end{pmatrix}, \quad A' \in \mathbb{R}^{3 \times 3}, \quad B'' \in \mathbb{R}^{3n \times 3n}.$$

The first three components of $\psi_{\kappa} \circ \varphi_{\kappa}$ agree with $\Phi \circ \varphi_{\kappa}$ by construction, and the last 3n components are identically zero, as φ_{κ} is precisely found by solving the system given after setting the last 3n entries of


 ψ_{κ} to zero. Since φ_{κ} is the identity in the first three entries, for some $D \in \mathbb{R}^{3n \times 3}$ we write (10) in block form and obtain the consequent relations:

$$\begin{pmatrix} J_{\Phi \circ \varphi_{\kappa}}(e, f, s_0) \\ 0 \end{pmatrix} = \begin{pmatrix} A' & A'' \\ B' & B'' \end{pmatrix} \begin{pmatrix} Id_{3 \times 3} \\ D \end{pmatrix}$$
$$J_{\Phi \circ \varphi_{\kappa}}(e, f, s_0) = A' + A''D, \qquad B' + B''D = 0.$$

By adding to the first three columns of $J_{\psi_{\kappa}}(\varphi_{\kappa}(e, f, s_0))$ linear combinations of the remaining 3n columns with coefficients given by the three columns of D respectively, we obtain

$$\det J_{\psi_{\kappa}}(\varphi_{\kappa}(e, f, s_0)) = \det \begin{pmatrix} A' & A'' \\ B' & B'' \end{pmatrix} = \det \begin{pmatrix} A' + A''D & A'' \\ B' + B''D & B'' \end{pmatrix} = \det \begin{pmatrix} J_{\Phi \circ \varphi_{\kappa}}(e, f, s_0) & A'' \\ 0 & B'' \end{pmatrix}$$
$$= \det J_{\Phi \circ \varphi_{\kappa}}(e, f, s_0) \cdot \det B''.$$

All that remains is to show that the sign of $\det B''$ is $(-1)^{3n}$. The matrix B'' is the Jacobian of the last 3n rows of ψ_{κ} in the variables s_i, y_i, u_i . Note that it is precisely the coefficient matrix of the linear system that is solved to find φ_{κ} . To find the determinant, we use the theory developed in [19, 38], see also [41], where this determinant is found using the Matrix-Tree theorem on a suitable digraph. Consider the following digraph:



Consider the directed spanning trees rooted at the vertex * (that is, such that * is the only vertex without outgoing edges), and for each such tree, define its label as the product of the labels of its edges. By [38, Proposition 1], the determinant of B'' is $(-1)^{3n}$ times the sum of the labels of all spanning trees rooted at *. As there is at least one such tree, and all labels are positive, this gives that the sign of det B'' is $(-1)^{3n}$. This concludes the proof.

Proposition 3.2 is the key in this work, as it will allow us to find explicitly a polynomial whose signs determine whether or not a vector of reaction rate constants enables multistationarity. The proposition relies on some algebraic manipulations and the multivariate chain rule, and this would apply to any other system falling in the setting of Theorem 1 in [7] and for which a parametrization of the set of positive steady states exists. In the notation of the proof of Proposition 3.2, a critical aspect is that the sign of the determinant of the matrix B'' is constant and can be determined. This might seem very restrictive, but it occurs whenever the parametrization of the set of steady states arises from linear elimination of variables in the setting of [19, 38], see also [41].

In view of Propositions 3.1 and 3.2, multistationarity is established by considering the sign of the determinant of the Jacobian of $\Phi \circ \varphi_{\kappa}$. The entries of $\Phi \circ \varphi_{\kappa}$ are

$$e + \sum_{i=0}^{n-1} K_i T_{i-1} e^{i+1} f^{-i} s_0,$$

$$f + \sum_{i=0}^{n-1} L_i T_i e^{i+1} f^{-i} s_0,$$

$$s_0 + \sum_{i=1}^{n} T_{i-1} e^i f^{-i} s_0 + \sum_{i=0}^{n-1} (K_i T_{i-1} + L_i T_i) e^{i+1} f^{-i} s_0.$$

Hence, $\Phi \circ \varphi_{\kappa}$ depends on κ through the assembled parameters K_i, L_i, T_i . As we are only interested in the determinant of the Jacobian of $\Phi \circ \varphi_{\kappa}$, we subtract the first and second equations from the third, and

in this way the last equation is replaced by

$$-e - f + s_0 + \sum_{i=0}^{n-1} T_i e^{i+1} f^{-(i+1)} s_0.$$

For convenience, the parameters and variables are relabeled in the following way:

(11)
$$a_i = K_i T_{i-1}, \quad b_i = L_i T_i, \quad c_i = T_i, \quad \text{for } i = 0, \dots, n-1, \\ x_1 = e, \qquad x_2 = \frac{e}{f}, \qquad x_3 = s_0.$$

With this notation, an easy computation now gives that the determinant of the Jacobian of $\Phi \circ \varphi_{\kappa}$ is the determinant of the following matrix:

$$(12) \quad J := \begin{bmatrix} 1 + \sum_{i=0}^{n-1} (i+1) \, a_i \, x_2^i x_3 & - \sum_{i=0}^{n-1} i \, a_i \, x_2^{i+1} x_3 & \sum_{i=0}^{n-1} a_i \, x_1 x_2^i \\ \sum_{i=0}^{n-1} (i+1) \, b_i \, x_2^i x_3 & 1 - \sum_{i=0}^{n-1} i \, b_i \, x_2^{i+1} x_3 & \sum_{i=0}^{n-1} b_i \, x_1 x_2^i \\ -1 + \sum_{i=0}^{n-1} (i+1) \, c_i \, x_1^{-1} \, x_2^{i+1} x_3 & -1 - \sum_{i=0}^{n-1} (i+1) \, c_i \, x_1^{-1} x_2^{i+2} x_3 & 1 + \sum_{i=0}^{n-1} c_i \, x_2^{i+1} \end{bmatrix}.$$

The determinant of (12) is a polynomial in x_1, x_2, x_3 with coefficients depending on a_i, b_i, c_i , which in turn depend on $T_i, K_i, L_i, i = 0, ..., n-1$. We, therefore, view det J as a polynomial in x_1, x_2, x_3 with coefficients depending on a parameter vector in $\mathbb{R}^{3n}_{>0}$, and define:

(13)
$$p_{\eta}(x_1, x_2, x_3) := \det J, \qquad \eta = (T_0, \dots, T_{n-1}, K_0, \dots, K_{n-1}, L_0, \dots, L_{n-1}) \in \mathbb{R}^{3n}_{>0}.$$

The discussion above together with Propositions 3.1 and 3.2, shows that whether or not κ enables multistationarity depends only on the entries of the vector $\eta(\kappa)$ and yield the following proposition.

Proposition 3.3. Let $\kappa \in \mathbb{R}^{6n}_{>0}$, consider $\eta(\kappa) \in \mathbb{R}^{3n}_{>0}$ using (7), and the polynomial $p_{\eta(\kappa)}(x_1, x_2, x_3)$ in (13). It holds:

- (i) If $p_{\eta(\kappa)}(x_1, x_2, x_3) > 0$ for all $(x_1, x_2, x_3) \in \mathbb{R}^3_{>0}$, then κ precludes multistationarity.
- (ii) If $p_{\eta(\kappa)}(x_1^*, x_2^*, x_3^*) < 0$ for some $(x_1^*, x_2^*, x_3^*) \in \mathbb{R}^3_{>0}$, then κ enables multistationarity. In this case the stoichiometric compatibility class containing $\varphi_{\kappa}(x_1^*, \frac{x_1^*}{x_2^*}, x_3^*)$ contains at least two positive steady states.

In Appendix A we explicitly compute p_{η} as the determinant of the matrix J in (12). We now explore the structure of the polynomial $p_{\eta}(x_1, x_2, x_3)$ using techniques from real algebraic geometry, in particular, via studying its Newton polytope. The polynomial is quadratic in x_3 and can be written as:

(14)
$$p_{\eta}(x_1, x_2, x_3) = A_2(x_2)x_3^2 + \left(A_{10}(x_2) + A_{11}(x_2)x_1\right)x_3 + \left(A_{00}(x_2) + A_{01}(x_2)x_1\right),$$
 with

$$\begin{split} A_2(x_2) &= \left(1 + \sum_{i=0}^{n-1} c_i \, x_2^{i+1} \,\right) \left(\sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (i-j) \, a_i \, b_j \, x_2^{\ell+1} \right), \\ A_{11}(x_2) &= (1+x_2) \left(\sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (i-j) a_i \, b_j \, x_2^{\ell} \right), \\ A_{10}(x_2) &= \sum_{\ell=0}^{n-1} (\ell+1) \, a_\ell \, x_2^{\ell} - \sum_{\ell=1}^{n-1} \ell \, b_\ell \, x_2^{\ell+1} + \sum_{\ell=0}^{2n-2} \sum_{i+j=\ell} (j+1-i) \, b_i \, c_j \, x_2^{\ell+2} + \sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (i-j) \, a_i \, c_j \, x_2^{\ell+1}, \\ A_{01}(x_2) &= \sum_{i=0}^{n-1} (a_i + b_i) x_2^i, \qquad \text{and} \qquad A_{00}(x_2) = 1 + \sum_{i=0}^{n-1} c_i \, x_2^{i+1}. \end{split}$$

Lemma 3.4. For the polynomial p_{η} written as in (14), the polynomials $A_{01}(x_2)$ and $A_{00}(x_2)$ have only positive coefficients, while $A_2(x_2)$, $A_{10}(x_2)$, and $A_{11}(x_2)$ have both positive and negative coefficients. Moreover, $A_2(x_2)$ is nonnegative if and only if $A_{11}(x_2)$ is nonnegative.

Proof. The first part follows easily from inspection of the coefficients of the polynomials. The second statement is immediate as both A_2 and A_{11} are positive multiples of $\sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (i-j) a_i b_j x_2^{\ell}$.

Note that

(15)
$$\sum_{i+j=\ell} (i-j) a_i b_j x_2^{\ell+1} = \sum_{\substack{i>j\\i+j=\ell}} (i-j) (a_i b_j - a_j b_i) x_2^{\ell+1}.$$

Furthermore, recalling the definition of a_i, b_i , and c_i in (11), and of T_i in (6), we obtain that

(16)
$$a_{i}b_{j} - a_{j}b_{i} = K_{i}T_{i-1}L_{j}T_{j} - K_{j}T_{j-1}L_{i}T_{i} = T_{i-1}T_{j-1}\left(K_{i}L_{j}\frac{\kappa_{6j+3}K_{j}}{\kappa_{6j+6}L_{j}} - K_{j}L_{i}\frac{\kappa_{6i+3}K_{i}}{\kappa_{6i+6}L_{i}}\right) = T_{i-1}T_{j-1}K_{i}K_{j}\left(\frac{\kappa_{6j+3}}{\kappa_{6j+6}} - \frac{\kappa_{6i+3}}{\kappa_{6i+6}}\right).$$

Therefore, the signs of $a_ib_j - a_jb_i$ depend on the minors of the following matrix:

(17)
$$M_{\kappa} = \begin{bmatrix} \kappa_3 & \kappa_9 & \dots & \kappa_{6n-3} \\ \kappa_6 & \kappa_{12} & \dots & \kappa_{6n} \end{bmatrix}.$$

Newton Polytope of $p_{\eta}(x_1, x_2, x_3)$. To analyze the possible signs that $p_{\eta}(x_1, x_2, x_3)$ attains over the positive orthant, we begin by exploring the properties of its Newton polytope. The rest of this section focuses on the shape and structure of $N(p_{\eta})$. We start by noting that p_{η} is linear in x_1 , and hence can be expressed as

$$p_{\eta}(x_1, x_2, x_3) = P_0(x_2, x_3) + x_1 P_1(x_2, x_3)$$

with

(18)
$$P_0(x_2, x_3) = A_2(x_2)x_3^2 + A_{10}(x_2)x_3 + A_{00}(x_2)$$
 and $P_1(x_2, x_3) = A_{11}(x_2)x_3 + A_{01}(x_2)$.

Therefore, the two hyperplanes $H_0 := \{x \in \mathbb{R}^3 : x_1 = 0\}$ and $H_1 := \{x \in \mathbb{R}^3 : x_1 = 1\}$ contain all exponent vectors, and further

(19)
$$N(p_{\eta}) \cap H_0 = \{0\} \times N(P_0), \qquad N(p_{\eta}) \cap H_1 = \{1\} \times N(P_1).$$

Theorem 3.5. Let $n \ge 1$, and $\eta \in \mathbb{R}^{3n}_{>0}$ such that $a_1b_0 - a_0b_1 \ne 0$ and $a_{n-1}b_{n-2} - a_{n-2}b_{n-1} \ne 0$. Then the set of vertices of the Newton polytope $N(p_\eta)$ of p_η in (14) consists of the following 10 points:

$$\Big\{(0,0,0),(0,n,0),(0,0,1),(0,2n,1),(0,2,2),(0,3n-2,2),(1,0,0),(1,n-1,0),(1,1,1),(1,2n-2,1)\Big\}.$$

Furthermore, $N(P_0) = N(p_n) \cap H_0$ is a hexagon with set of vertices

$$\{(0,0),(n,0),(0,1),(2n,1),(2,2),(3n-2,2)\}.$$

Proof. The hypothesis implies that the polynomials A_2 , A_{11} , A_{10} , A_{01} , A_{00} in x_2 have degree 3n-2, 2n-2, 2n, n-1, n respectively, and lowest exponent 1, 1, 0, 0, 0 respectively.

Considering that the vertices of $N(p_{\eta})$ are contained in the union of hyperplanes $H_0 \cup H_1$, the set of vertices of $N(p_{\eta})$ is the union of the set of vertices of $N(p_{\eta}) \cap H_0$ and of $N(p_{\eta}) \cap H_1$. By (19), it is enough to find the vertices of $N(P_0)$ and $N(P_1)$, which are planar polytopes.

As P_0 is quadratic in x_3 , see (18), the vertices of $N(P_0)$ are on the lines $L_i := \{(x_2, x_3) \in \mathbb{R}^2 : x_3 = i\}$, for i = 0, 1, 2. The polytopes $N(P_0) \cap L_i$ for i = 0, 1, 2 correspond to the Newton polytopes of the univariate polynomials A_{00} , A_{10} , and A_2 respectively. Their vertices are simply the highest and lowest exponents of these polynomials. Therefore, the set of vertices of $N(P_0)$ is contained in $S_1 = \{(0,0),(n,0),(0,1),(2n,1),(2,2),(3n-2,2)\}$. To show that this is exactly the set of vertices of $N(P_0)$, it is enough to verify that (0,1),(2n,1) do not lie in the convex hull of the 4 remaining points. This is easily checked by considering the relative position of (0,1) with respect to the line joining (0,0) and (2,2), and similarly, that of (2n,1) with respect to the line joining (n,0) and (3n-2,2). The Newton polytope $N(P_0)$ is therefore the hexagon with vertex set S_1 .

For $N(P_1)$, we note that P_1 in (18) is linear in x_3 . Hence, as above, we consider first the vertices of $N(P_1) \cap \{(x_2, x_3) \in \mathbb{R}^2 : x_3 = i\}$ for i = 0, 1, which are 0, n - 1 and 1, 2n - 2 respectively. It follows that the set of vertices of $N(P_1)$ is $S_2 = \{(0, 0), (n - 1, 0), (1, 1), (2n - 2, 1)\}$. By (19), we conclude that the set of vertices of $N(p_\eta)$ is as given in the statement.

Remark 3.6. In the original parameter vector $\kappa \in \mathbb{R}^{6n}_{>0}$, the condition in Theorem 3.5 corresponds by (16) to $\kappa_3 \kappa_{12} - \kappa_6 \kappa_9 \neq 0$ and $\kappa_{6n-9} \kappa_{6n} - \kappa_{6n-3} \kappa_{6n-6} \neq 0$.

Given a polynomial $P \in \mathbb{R}[x_1, \dots, x_m]$, and a proper face F of N(P), we let P_F denote the restriction of P to F, that is, P_F is the sum of the terms $c_{\alpha}x_1^{\alpha_1}\cdots x_m^{\alpha_m}$ of P satisfying $(\alpha_1, \dots, \alpha_m) \in F$. In what follows we will use the following well-known result that gives a connection between the signs P attains over $\mathbb{R}^m_{>0}$, and the signs P_F attains.

Proposition 3.7. Let F be a proper face of the Newton polytope N(P) of a polynomial $P \in \mathbb{R}[x_1, \ldots, x_m]$ and let P_F be the restriction of P to F. Then for any $x \in \mathbb{R}^m_{>0}$ such that $P_F(x) \neq 0$, there exists $y \in \mathbb{R}^m_{>0}$ such that

$$sign(P(y)) = sign(P_F(x)).$$

In particular, for t > 0 large enough and $\omega \in \mathbb{R}^m$ in the open outer normal cone of F, $y = (x_1 t^{\omega_1}, \dots, x_m t^{\omega_m})$ generates the desired point.

A proof of this proposition can be found in [17, Prop. 2.3]. Recall that the open outer normal cone of F consists of the vectors ω such that $\omega \cdot x^\top > \omega \cdot y^\top$ for every $x \in F$ and $y \in \mathcal{N}(P) \backslash F$ where x^\top, y^\top denotes the transpose. We can easily deduce that if the coefficient of the monomial supported at one of the vertices is negative, then there exists some $x \in \mathbb{R}^m_{>0}$ such that the polynomial is negative.

Theorem 3.8. Fix $\eta \in \mathbb{R}^{3n}_{>0}$.

- (i) The polynomial p_{η} attains a negative value over $\mathbb{R}^3_{>0}$ if and only if P_0 attains negative values over $\mathbb{R}^2_{>0}$.
- (ii) Moreover, p_{η} is either positive over the positive orthant or p_{η} attains negative values over $\mathbb{R}^3_{>0}$. In other words, p_{η} has a zero in $\mathbb{R}^3_{>0}$ if and only if it takes negative values.

Proof. The two statements in the theorem follow from these two implications:

$$\begin{aligned} p_{\eta}(x_1, x_2, x_3) \leqslant 0 \text{ for some } (x_1, x_2, x_3) \in \mathbb{R}^3_{>0} & \Rightarrow & P_0(x_2, x_3) < 0 \text{ for some } (x_2, x_3) \in \mathbb{R}^2_{>0} \\ & \Rightarrow & p_{\eta}(x_1, x_2, x_3) < 0 \text{ for some } (x_1, x_2, x_3) \in \mathbb{R}^3_{>0}. \end{aligned}$$

To show the second implication, note that $N(p_{\eta}) \cap H_0$ and $N(p_{\eta}) \cap H_1$ are two facets of $N(p_{\eta})$, and p_{η} restricted to these facets is respectively P_0 and x_1P_1 . Therefore, if $P_0(x_2, x_3) < 0$ for some $(x_2, x_3) \in \mathbb{R}^2_{>0}$, then Proposition 3.7 readily gives that p_{η} also attains negative values.

For the first implication, $p_{\eta}(x_1, x_2, x_3) \leq 0$ implies that $P_0(x_2, x_3)$ or $P_1(x_2, x_3)$ are negative. If $P_1(x_2, x_3) < 0$, then by Lemma 3.4 necessarily $A_{11}(x_2) < 0$ and hence, by the same lemma, we also have $A_2(x_2) < 0$. As $A_2(x_2)$ is the leading coefficient of P_0 , there exists $x_3^* > 0$ such that $P_0(x_2, x_3^*) < 0$. This concludes the proof of the theorem.

As the signs p_{η} attain are completely determined by the signs of P_0 over $\mathbb{R}^2_{>0}$ by Theorem 3.8, we rewrite P_0 as P_{η} , to indicate dependence on the parameter vector $\eta \in \mathbb{R}^{3n}_{>0}$:

(20)
$$P_{\eta}(x_2, x_3) = A_2(x_2)x_3^2 + A_{10}(x_2)x_3 + A_{00}(x_2).$$

Theorem 3.8 tells us that the silent scenario in Proposition 3.3, namely when p_{η} is nonnegative over $\mathbb{R}^3_{>0}$, cannot occur. This gives the following characterization of multistationarity of the *n*-phosphorylation cycle.

Corollary 3.9. Let $\kappa \in \mathbb{R}^{6n}_{>0}$, consider $\eta(\kappa) \in \mathbb{R}^{3n}_{>0}$ from (7), and $P_{\eta(\kappa)}(x_2, x_3)$ as in (20). Then κ enables multistationarity if and only if $P_{\eta(\kappa)}(x_2^*, x_3^*) < 0$ for some $(x_2^*, x_3^*) \in \mathbb{R}^2_{>0}$.

4. Sufficient conditions for multistationarity

In this section we start by providing conditions on the parameters to ensure multistationarity using Corollary 3.9. Then, we describe subsets of the parameter space that preclude multistationarity. First, we make the observation that if the polynomial $A_2(x_2)$ in (14) is negative, then P_{η} in (20) attains negative values. This follows immediately from P_{η} being quadratic in x_3 with $A_2(x_2)$ as the leading coefficient. Recall that the sign of the coefficients of $A_2(x_2)$ are determined by (15).

Theorem 4.1. If $\kappa \in \mathbb{R}^{6n}_{>0}$ is such that

$$\kappa_3 \kappa_{12} - \kappa_6 \kappa_9 < 0$$
 or $\kappa_{6n-9} \kappa_{6n} - \kappa_{6n-3} \kappa_{6n-6} < 0$,

then $P_{n(\kappa)}$ attains negative values, and hence κ enables multistationarity.

Proof. By Theorem 3.5, the set of vertices of $N(P_{\eta(\kappa)})$ is $\{(0,0),(n,0),(0,1),(2n,1),(2,2),(3n-2,2)\}$, see Remark 3.6. The coefficient of the lowest degree term of $A_2(x_2)$ (i.e. $x_2^2x_3^2$) is a positive multiple of $a_1b_0 - b_0a_1$ by (15), and of $\kappa_3\kappa_{12} - \kappa_6\kappa_9$ by (16). Similarly, the coefficient of the highest degree term of $A_2(x_2)$ (i.e. $x_2^{3n-2}x_3^2$) is a positive multiple of $\kappa_{6n-9}\kappa_{6n} - \kappa_{6n-3}\kappa_{6n-6}$. If one of these coefficients is negative, then $P_{\eta(\kappa)}$ attains negative values by Proposition 3.7, and therefore, by Corollary 3.9, κ enables multistationarity. Note that the other coefficients of $P_{\eta(\kappa)}$ corresponding to vertices of $N(P_{\eta(\kappa)})$ are strictly positive.

The expressions in Theorem 4.1 correspond to the first and last maximal minors of the matrix M_{κ} in (17).

Proposition 4.2. Let $\kappa \in \mathbb{R}^{6n}_{>0}$ be such that the matrix M_{κ} in (17) has rank one. Then κ enables multistationarity if and only if the polynomial $A_{10}(x_2)$ is not nonnegative, that is, attains a negative value.

Proof. By hypothesis, all maximal minors of M_{κ} vanish, and therefore, A_2 is identically zero by (15) and (16). Therefore, $P_{\eta(\kappa)}(x_2, x_3) = A_{10}(x_2)x_3 + A_{00}(x_2)$. As A_{00} has only positive coefficients, $P_{\eta(\kappa)}(x_2, x_3)$ attains negative values if and only if $A_{10}(x_2)$ does. The statement now follows from Theorem 3.8.

In the next result we show that network (2) can be multistationary even if all the minors of the matrix M_{κ} are positive.

Theorem 4.3. Let $\kappa \in \mathbb{R}^{6n}_{>0}$ be such that all the minors of M_{κ} are positive. Then, by choosing $K_0 = \frac{\kappa_1}{\kappa_2 + \kappa_3}$ small enough by varying κ_1 and/or κ_2 and leaving the rest of the entries of κ fixed, we obtain a parameter vector that enables multistationarity.

Proof. Fix the value of the assembled parameters $K_1, \ldots, K_{n-1}, L_0, \ldots, L_{n-1}$ corresponding to κ , and construct $T_0(K_0), \ldots, T_{n-1}(K_0)$ as in (6) but with K_0 treated as a parameter. For the one parameter family of vectors $\xi(K_0) = (T_0(K_0), \ldots, T_{n-1}(K_0), K_0, \ldots, K_{n-1}, L_0, \ldots, L_{n-1}) \in \mathbb{R}^{3n}_{>0}$, the polynomial $P_{\xi(K_0)}(x_2, x_3)$ becomes a polynomial in K_0, x_2 , and x_3 , which we denote as $Q(K_0, x_2, x_3)$. We prove the theorem by studying the Newton polytope of $Q(K_0, x_2, x_3)$.

Note that K_0 is a factor with exponent 1 in each of a_i , b_i , and c_i . Consider the exponents of K_0 and x_2 in the polynomials A_{00} , A_{10} , and A_2 in (14). Close inspection of each of these show that N(Q) is contained in the convex hull of the following set of points:

$$\Omega := \{(0,0,0), (1,1,0), \dots, (1,n,0), (1,0,1), \dots, (1,n,1), (2,2,1), \dots, (2,2n,1), \dots, (2,2n,2), \dots, (2,2n-2,2), (3,3,2), \dots, (3,3n-2,2)\}.$$

Note that some points could appear with zero coefficient for certain parameter values.

Consider the point $(1, n, 1) \in \Omega$. The term of Q with monomial $K_0 x_2^n x_3$ is $-(n-1) b_{n-1} x_2^n x_3$. Hence, the coefficient is negative and $(1, n, 1) \in N(Q)$. With $\omega = (-2(n+1), 2, 4) \in \mathbb{R}^3$ and $\gamma = -1$, it holds

$$\omega \cdot (1, n, 1)^{\top} + \gamma > 0, \qquad \omega \cdot q^{\top} + \gamma < 0 \text{ for all } q \in \Omega \setminus \{(1, n, 1)\}.$$

Therefore, the hyperplane $\omega \cdot x^{\top} + \gamma = 0$ separates (1, n, 1) and $\operatorname{conv}(\Omega \setminus \{(1, n, 1)\})$. It follows that $(1, n, 1) \notin \operatorname{conv}(\Omega \setminus \{(1, n, 1)\})$, showing that (1, n, 1) is a vertex of N(Q).

By Proposition 3.7, Q attains negative values after appropriately choosing $K_0, x_2, x_3 > 0$. As ω belongs to the outer normal cone of (1, n, 1) by construction, the point $(t^{-2(n+1)}, t^2, t^4)$ makes Q negative for t large enough. This amounts to letting K_0 be small enough, and as κ_3 was fixed in the definition of $T_i(K_0)$, this must be achieved by varying κ_1 and/or κ_2 . This concludes the proof.

Remark 4.4. Theorem 4.3 implies that the conditions in Theorem 4.1 are not necessary for multistationarity. It gives, additionally, a way to construct parameters that enable multistationarity, while satisfying the condition that the negative coefficients of $N(P_{\eta(\kappa)})$ correspond to points in the interior of $N(P_{\eta(\kappa)})$.

Although we have shown the existence of values of K_0 for multistationarity when the rest of the parameters are fixed, the theorem remains true when all parameters but L_{n-1} are fixed. This is shown by considering $P_{\eta(\kappa)}$ as a polynomial in the variables L_{n-1}, x_2 and x_3 . Then, the point (0, n, 1) is a vertex of the corresponding polytope and the proof follows similar arguments as the proof of Theorem 4.3.

5. Sufficient Conditions for Monostationarity

In this section we present a systematic way to construct a sufficient condition for monostationarity for all n. To this end, we describe a region R in the parameter space such that the polynomial $P_{\eta(\kappa)}$ in (20) is positive for any $\kappa \in \mathbb{R}^{6n}_{>0}$, and contains parameter values for which $P_{\eta(\kappa)}$ has negative coefficients. The approach is a generalization of the methods that are used in [17, §2.2] for n = 2, and in [47] for n = 3.

For the particular method we use in this section, it is crucial that the coefficients of the terms that correspond to the vertices of $N(P_{\eta(\kappa)})$ have positive sign. If this is not the case, then Theorem 4.1 implies that multistationarity is enabled. Therefore, throughout this section we only consider parameters $\kappa \in \mathbb{R}^{6n}_{>0}$ where

(21)
$$\kappa_3 \kappa_{12} - \kappa_6 \kappa_9 > 0$$
 and $\kappa_{6n-9} \kappa_{6n} - \kappa_{6n-3} \kappa_{6n-6} > 0$.

For the cases n=2 and n=3, the conditions in (21) are sufficient for all minors of the matrix M_{κ} in (17) to be positive.

Remark 5.1. For n=2, assuming that the minors of the matrix M_{κ} are all positive, then P_{η} has only one term that can have negative coefficient. The exponent that corresponds to this particular term is contained in the interior of $N(P_{\eta})$. Therefore, as discussed in Remark 2.3, $P_{\eta(\kappa)}$ is nonnegative if and only if it is a SONC polynomial.

In order to find the region R, we will restrict to a scenario where all coefficients of $P_{\eta(\kappa)}$ corresponding to exponents at the boundary of $N(p_{\eta})$ are nonnegative, and hence, only points in the interior of $N(P_{\eta(\kappa)})$ can have negative coefficient. In particular, this holds when all minors of the matrix M_{κ} are nonnegative, as then all coefficients of $A_2(x_2)$ are nonnegative by (16) and only $A_{10}(x_2)$ contributes to negative terms to $P_{\eta(\kappa)}$. For n > 3, the conditions in (21) are not enough to guarantee that all minors of M_{κ} are positive. In Proposition 5.2, we give a sufficient condition on a_i, b_i, c_i , as defined in Equation (11), that ensures that all minors of M_{κ} are positive.

Proposition 5.2. Assume that for a given $\kappa \in \mathbb{R}^{6n}_{>0}$, it holds that $a_{i+1}b_i - a_ib_{i+1} > 0$ (resp. $\geqslant 0$) for all $i \in \{0, \ldots, n-1\}$. Then, all 2 by 2 minors of M_{κ} are positive (resp. nonnegative).

Proof. We need to show that $\kappa_{6i+3}\kappa_{6j+6} - \kappa_{6i+6}\kappa_{6j+3} > 0$, equivalently $\frac{\kappa_{6i+3}}{\kappa_{6i+6}} - \frac{\kappa_{6j+3}}{\kappa_{6j+6}} > 0$ for all j > i. Note that

$$\frac{\kappa_{6i+3}}{\kappa_{6i+6}} - \frac{\kappa_{6j+3}}{\kappa_{6j+6}} = \sum_{k=1}^{j-i} \frac{\kappa_{6(i+k-1)+3}}{\kappa_{6(i+k-1)+6}} - \frac{\kappa_{6(i+k)+3}}{\kappa_{6(i+k)+6}}.$$

By hypothesis and (16), all terms in the sum are positive and hence, so is their sum. The argument for nonnegativity is analogous. This concludes the proof.

In the rest of this section, we only consider parameter vectors $\kappa \in \mathbb{R}^{6n}_{>0}$ for which all 2 by 2 minors of M_{κ} are nonnegative and (21) holds. These yield to parameter vectors $\eta \in \mathbb{R}^{3n}_{>0}$ such that the set of exponents of the monomials of P_{η} that may have negative coefficient is

$$A_{-} := \{(2,1), (3,1), \dots, (2n-2,1)\},\$$

and we define

$$\iota_i := (i+1,1), \quad \text{for} \quad i \in \{1,\dots,2n-3\}.$$

Furthermore, we know from Theorem 3.5 that $N(P_{\eta})$ is a hexagon for $n \ge 2$, and let us label its vertices as follows:

$$\alpha_1 := (0,0), \quad \alpha_2 := (2,2), \quad \alpha_3 := (2n,1), \quad \alpha_4 := (0,1), \quad \alpha_5 := (n,0), \quad \alpha_6 := (3n-2,2).$$

Under the current assumptions, the coefficients corresponding to these vertices are nonnegative. The triangles

(22)
$$\Delta_1 := \operatorname{conv}(\{\alpha_1, \alpha_2, \alpha_3\}) \quad \text{and} \quad \Delta_2 := \operatorname{conv}(\{\alpha_4, \alpha_5, \alpha_6\}),$$

contain ι_i in the relative interior for all $i \in \{1, \ldots, 2n-3\}$. Hence, our goal now is to construct two circuit polynomials with supports $\Delta_1 \cup \{\iota_i\}$, $\Delta_2 \cup \{\iota_i\}$ for each exponent $\iota_i \in A_-$, such that we can apply Theorem 2.2 repeatedly to ensure nonnegativity of P_n .

Consider $\iota_i \in A_-$, $x = (x_2, x_3)$, and the two polynomials:

(23)
$$C_{1,i}(x) = \sum_{j=1}^{3} \frac{\operatorname{coeff}(P_{\eta,\alpha_{j}})}{2n-3} x^{\alpha_{j}} + \delta_{1,i} x^{\iota_{i}} \quad \text{and} \quad C_{2,i}(x) = \sum_{j=4}^{6} \frac{\operatorname{coeff}(P_{\eta,\alpha_{j}})}{2n-3} x^{\alpha_{j}} + \delta_{2,i} x^{\iota_{i}},$$

where $\delta_{1,i}, \delta_{2,i} \in \mathbb{R}$ are chosen such that $\delta_{1,i} + \delta_{2,i} = \operatorname{coeff}(P_{\eta}, \iota_i)$. By construction, $\operatorname{N}(C_{1,i})$ and $\operatorname{N}(C_{2,i})$ are respectively Δ_1, Δ_2 for all $i \in \{1, \ldots, 2n-3\}$ and hence $C_{1,i}$ and $C_{2,i}$ are circuit polynomials. We denote the circuit numbers associated with $C_{1,i}$ and $C_{2,i}$ with $\Theta_{1,i}$ and $\Theta_{2,i}$, respectively. Then,

for $i \in \{1, \dots, 2n-3\}$, it holds

$$\Theta_{1,i} = \left(\frac{\operatorname{coeff}(C_{1,i},\alpha_{1})}{\frac{2n-1-i}{4n-2}}\right)^{\frac{2n-1-i}{4n-2}} \left(\frac{\operatorname{coeff}(C_{1,i},\alpha_{2})}{\frac{2n-1-i}{4n-2}}\right)^{\frac{2n-1-i}{4n-2}} \left(\frac{\operatorname{coeff}(C_{1,i},\alpha_{3})}{\frac{2i}{4n-2}}\right)^{\frac{2i}{4n-2}} \\
= \left(\frac{\operatorname{coeff}(P_{\eta},\alpha_{1})}{(2n-1-i)}\right)^{\frac{2n-1-i}{4n-2}} \left(\frac{\operatorname{coeff}(P_{\eta},\alpha_{2})}{(2n-1-i)}\right)^{\frac{2n-1-i}{4n-2}} \left(\frac{\operatorname{coeff}(P_{\eta},\alpha_{3})}{2i}\right)^{\frac{2i}{4n-2}} \left(\frac{4n-2}{2n-3}\right),$$

and similarly.

(25)
$$\Theta_{2,i} = \left(\frac{\operatorname{coeff}(P_{\eta}, \alpha_4)}{(4n - 2i - 4)}\right)^{\frac{4n - 2i - 4}{4n - 2}} \left(\frac{\operatorname{coeff}(P_{\eta}, \alpha_5)}{(i + 1)}\right)^{\frac{i + 1}{4n - 2}} \left(\frac{\operatorname{coeff}(P_{\eta}, \alpha_6)}{i + 1}\right)^{\frac{i + 1}{4n - 2}} \left(\frac{4n - 2}{2n - 3}\right).$$

For each $i \in \{1, ..., 2n-3\}$ we define the region $R_i \subseteq \mathbb{R}^{6n}_{>0}$ as follows:

$$R_{i} := \{ \kappa \in \mathbb{R}^{6n}_{>0} : \operatorname{coeff}(P_{\eta(\kappa)}, \iota_{i}) \geqslant -\Theta_{1,i} - \Theta_{2,i}, \\ \kappa_{3}\kappa_{12} - \kappa_{6}\kappa_{9} > 0, \ \kappa_{6n-9}\kappa_{6n} - \kappa_{6n-3}\kappa_{6n-6} > 0, \text{ and} \\ \kappa_{6\ell+3}\kappa_{6j+6} - \kappa_{6\ell+6}\kappa_{6j+3} \geqslant 0 \text{ for all } \ell < j, \ \ell = 1, \dots, n-3 \},$$

where $\Theta_{1,i}, \Theta_{2,i}$ are defined using $\eta(\kappa)$. Note that if $\kappa \in R_i$, all minors of M_{κ} are nonnegative, and furthermore, the inequalities in (21) hold. We denote the intersection of these 2n-3 regions by

$$(26) R := \bigcap_{i=1}^{2n-3} R_i.$$

Theorem 5.3. With $R \subseteq \mathbb{R}^{6n}_{>0}$ as in (26), it holds:

- (i) There exists κ∈ R such that P_{η(κ)} has some negative coefficient. In particular, R≠ Ø.
 (ii) P_{η(κ)}(x₂, x₃) > 0 for any κ∈ R and (x₂, x₃) ∈ ℝ²_{>0}, and hence κ precludes multistationarity.

Proof. We start by proving (ii). Let $\kappa \in R$, and $\eta := \eta(\kappa)$. Let $A(P_{\eta})$ be the set of exponents of P_{η} . and define $A_{+} = A(P_{\eta}) \setminus (A_{-} \cup \Delta_{1} \cup \Delta_{2})$ (with Δ_{1}, Δ_{2} as in (22). We rewrite P_{η} using the circuit polynomials defined in (23) as follows:

(27)
$$P_{\eta}(x) = \sum_{j=1}^{2} \sum_{i=1}^{2n-3} C_{j,i}(x) + \sum_{\alpha \in A_{+}} \operatorname{coeff}(P_{\eta}, \alpha) x^{\alpha}.$$

For any $\alpha \in A_+$, it holds that $\operatorname{coeff}(P_{\eta}, \alpha) \geqslant 0$. Recall that the coefficient of x^{ι_i} in $C_{j,i}$ equals $\delta_{j,i}$, and these could be chosen arbitrarily as long as $\delta_{1,i} + \delta_{2,i} = \operatorname{coeff}(P_{\eta}, \iota_i)$. This equality holds if we pick

$$\delta_{j,i} = -\Theta_{j,i} + \frac{\operatorname{coeff}(P_{\eta}, \iota_i) + \Theta_{1,i} + \Theta_{2,i}}{2}, \qquad j = 1, 2.$$

Furthermore, as in R it holds $\operatorname{coeff}(P_{\eta}, \iota_i) \ge -\Theta_{1,i} - \Theta_{2,i}$, we find that $\delta_{j,i} \ge -\Theta_{j,i}$ for j = 1, 2 and hence $C_{1,i}$ and $C_{2,i}$ are nonnegative circuit polynomials by Theorem 2.2. By the decomposition (27), P_{η} is nonnegative as well, and combined with Corollary 3.9, we have shown (ii).

To prove (i) we construct explicitly a point satisfying the conditions. We consider the one parameter family of reaction rate constants $\kappa(k) \in \mathbb{R}^{6n}_{>0}$ with k>0 defined as

(28)
$$\kappa_{j} = \begin{cases}
n - i & \text{if } j = 6i + 3 \text{ and } i < n, \\
2 & \text{if } j = 6i + 4 \text{ and } i < n, \\
k & \text{if } j = 6n - 5 = 6(n - 1) + 1, \\
1 & \text{otherwise,}
\end{cases}$$

and let $\eta(k) := \eta(\kappa(k)) \in \mathbb{R}^{3n}_{>0}$. We show that there exists an interval $I \subset \mathbb{R}_{>0}$ such that for any $k \in I$, $\kappa(k) \in R$ and $P_{\eta(k)}$ contains a negative term. For any such $\kappa(k)$, we have $L_i = 1$ for any $i \in \{0, \ldots, n-1\}$, and hence, $b_i = c_i = T_i$ for all $i \in \{0, \ldots, n-1\}$. Similarly, we have

(29)
$$K_{i} = \begin{cases} \frac{1}{n-i+1} & \text{if } i \in \{0,\dots,n-2\}, \\ \frac{k}{2} & \text{if } i = n-1, \end{cases} \quad \text{and} \quad T_{i} = \begin{cases} \frac{n-i}{n+1} & \text{if } i \in \{-1,0,\dots,n-2\}, \\ \frac{k}{n+1} & \text{if } i = n-1. \end{cases}$$

We start by noting that for each $\kappa(k)$ and for any $i \in \{0, \dots, n-1\}$, it holds

$$\left(\frac{\kappa_{6i+3}}{\kappa_{6i+6}} - \frac{\kappa_{6(i+1)+3}}{\kappa_{6(i+1)+6}}\right) = (n-i) - (n-i-1) = 1 > 0.$$

Hence, all minors of $M_{\kappa(k)}$ are positive by Proposition 5.2, and so are the coefficients of $P_{\eta(k)}$ corresponding to the exponents $\alpha_1, \ldots, \alpha_6$. It follows from (24) and (25) that $\Theta_{1,i}(k) + \Theta_{2,i}(k) > 0$ for all $i \in \{1, \ldots, 2n-3\}$, where $\Theta_{1,i}(k)$, $\Theta_{2,i}(k)$ are the circuit numbers obtained for $\kappa(k)$.

Let $C_{\ell}(k) := \operatorname{coeff}(P_{\eta(k)}, \iota_{\ell-1})$, which is the coefficient of x_2^{ℓ} in the polynomial $A_{10}(x_2)$. We show in Lemma B.1 in the Appendix, that for any k > 0, $C_{\ell}(k) > 0$ when $\ell \in \{2, \ldots, 2n-2\}$ and $\ell \neq n$. Thus, $\kappa(k) \in R_{\ell}$ for any $\ell \neq n$.

We consider the coefficient of x_2^n in $A_{10}(x_2)$, see (14). As for $\kappa(k)$ we have $b_i c_j = T_i T_j$ and $\sum_{i+j=n-2} (j+1-i)T_i T_j = \sum_{i+j=n-2} T_i T_j$, we have

$$C_n(k) = -(n-1)b_{n-1} + \sum_{\substack{i+j=n-2\\i>j}} T_i T_j + \sum_{\substack{i+j=n-1\\i>j}} (i-j) (a_i c_j - a_j c_i).$$

Using the identities (11), (29), and (30) we find $a_{n-1}c_0 - a_0c_{n-1} = \frac{k(n-1)}{(n+1)^2}$ and $b_{n-1} = \frac{k}{n+1}$. This gives:

$$C_n(k) = k \left(\frac{n-1}{n+1}\right) \left(\frac{n-1}{n+1} - 1\right) + \sum_{\substack{i+j=n-2\\i>j, i\neq n-1}} T_i T_j + \sum_{\substack{i+j=n-1\\i>j, i\neq n-1}} (i-j) (a_i c_j - a_j c_i).$$

The variable k does not appear in the last two summands of $C_n(k)$ and hence, $C_n(k)$ is a linear function of k with a negative leading term and a positive constant term.

Let $k_0 > 0$ be the zero of $C_n(k)$, and consider the function $g(k) = C_n(k) + \Theta_{1,n-1}(k) + \Theta_{2,n-1}(k)$. Note that $g(k_0) > 0$ as $C_n(k_0) = 0$. As g is continuous in k, there exists $\epsilon_0 > 0$ such that for all $k \in (k_0, k_0 + \epsilon_0)$, g(k) > 0. Hence $\kappa(k) \in R_{n-1}$. By definition of R, it then follows that $\kappa(k) \in R$ and by construction $C_n(k) < 0$ for any such k, showing (i).


In the next example, we give the conditions to be in R explicitly when n = 3 and $\kappa(k)$ is the parameter vector constructed in (28) in the proof of Theorem 5.3(i).

Example 5.4. Let n = 3, $\kappa(k) = (1, 1, 3, 2, 1, 1, 1, 1, 2, 2, 1, 1, k, 1, 1, 2, 1, 1)$ be as in (28). The Newton polytope $N(P_{\eta(k)})$ is shown in the right panel of Figure 1. The circuit numbers from (24) and (25) are

The vectors $\kappa(k) \in \mathbb{R}^{18}_{>0}$ are in the regions R_1, R_2 and R_3 in the parameter space if they satisfy the following inequalities respectively

$$\begin{split} \frac{6k+1}{8} &\geqslant -2^{-\frac{32}{10}} \ 3^{-1} \ 5 \ k^{\frac{4}{10}} \ - \ 2^{-\frac{28}{10}} \ 3^{-\frac{16}{10}} \ 5 \ k^{\frac{6}{10}}, \\ \frac{3-k}{4} &\geqslant -2^{-\frac{26}{10}} \ 3^{-\frac{16}{10}} \ 5 \ k^{\frac{8}{10}} \ - \ 2^{-3} \ 3^{-\frac{16}{10}} \ 5 \ k^{\frac{9}{10}}, \\ \frac{7k+4}{16} &\geqslant -2^{-\frac{32}{10}} \ 3^{-\frac{16}{10}} \ 5 \ k^{\frac{12}{10}} \ - \ 2^{-\frac{44}{10}} \ 3^{-1} \ 5 \ k^{\frac{12}{10}}, \end{split}$$

respectively. As shown in the proof of Theorem 5.3, the first and the third inequalities hold for all k > 0. The second inequality holds if k < 10.03, and the coefficient corresponding to the term ι_{n-1} in $P_{\eta(k)}$ is






FIGURE 1. The support of P_{η} for n=2,3, black circles and red squares correspond to the exponents with positive and negative coefficients, respectively. On the left, blue triangles and green line segments denote the Newton polytopes of the circuit polynomials used in [17, Theorem 3.5]. On the right, blue and green triangles correspond to the simplices Δ_1 and Δ_2 from (22), respectively.

negative if k > 3. Therefore, for 10.03 > k > 3, $P_{\eta(k)}$ contains a negative term, but $P_{\eta(k)}$ is positive. Hence multistationarity is precluded for $\kappa(k)$.

We end this section with some remarks.

Remark 5.5. If the premise of Proposition 5.2 holds, then every term of $P_{\eta(\kappa)}$ with exponent in $A_+ = A(P_{\eta}) \setminus (A_- \cup \Delta_1 \cup \Delta_2)$ has a positive coefficient. This means that we can construct further circuit polynomials using the exponents in A_+ as vertices and in A_- as inner terms. For example, a similar approach was taken for the case n = 2 in [17, Theorem 3.5], where we considered two additional 1-dimensional circuits given by $\{(1,0),(2,1),(3,2)\}$ and $\{(1,1),(2,1),(3,1)\}$ (green line segments in the left pane in Figure 1). These were used to formulate a sufficient condition for monostationarity stronger than that of Theorem 5.3. In Theorem 5.3, we omit these extra polynomials in order to keep the notation tidy.

The region R in (26) is a carefully constructed set, and hence Theorem 5.3 might only address a limited part of the monostationarity region. The ideas presented in this section can be used when $P_{\eta(\kappa)}$ has an exponent with a negative coefficient that lies on a positive dimensional face F of its Newton polytope, by applying them to the polynomial restricted to F.

Remark 5.6. We note that if the values of the reaction rate constant κ is known, then we can find a lower bound for $P_{\eta(\kappa)}(x_2, x_3)$ by using various convex and polynomial optimization methods such as solving a relative entropy program induced by the SONC/SAGE decomposition [4, Example 3.1] or using semidefinite programming [3]. However, for the results presented here, the parameters are a priori symbolic. Hence, we instead use the method of choosing circuit covers to obtain inequalities on the parameters, which yields symbolic necessary conditions for multistationarity.

6. Connectivity

As motivated in the introduction, even in lack of a full description of the parameter region of multistationarity, it is of interest to understand the topological properties of the region. Here we show that the subset of parameters enabling multistationarity and that of those precluding multistationarity are both path connected in the space of reaction rate constants.

The strategy we follow, which is applicable to other networks, consists of two parts: first, we identify a subregion of the region of interest that is path connected. For multistationarity, it is the subregion given by Theorem 4.1, making that theorem critical for this section. For monostationarity, we consider the subregion characterized by P_{η} having all coefficients positive (non-emptiness of this region is a consequence of our results here). Afterwards, we show that any other point in the region can be joined to a point in the selected subregion via a continuous path. This is done by exploiting again properties of the Newton polytope of P_{η} , but now viewing the polynomial as a polynomial in two of the entries of η .

Let $X \subseteq \mathbb{R}^{6n}_{>0}$ be the set of parameter vectors κ that enable multistationarity. We prove that X is path connected in two steps: First we show in Lemma 6.1 that a subset Y of X is path connected, and then we construct a path in X from any η that enables multistationarity to the subset Y in Theorem 6.2.

Lemma 6.1. Let $Y \subseteq \mathbb{R}^{6n}_{>0}$ consist of the parameters $\kappa \in \mathbb{R}^{6n}_{>0}$ such that $\kappa_{6n-9}\kappa_{6n} - \kappa_{6n-6}\kappa_{6n-3} < 0$. Then, $Y \subseteq X$ is path connected.

Proof. From Theorem 4.1, we know that $Y \subseteq X$. The proof that Y is path connected is a straightforward adaptation of Lemma 5.1 and Remark 5.3 in [17].

Theorem 6.2. For all $n \ge 2$, the set $X \subseteq \mathbb{R}^{6n}_{>0}$ consisting on the parameters κ that enable multistationarity is path connected.

Proof. Let $\kappa \in X$. Then by Corollary 3.9, there exists $x_2, x_3 > 0$ such that $P_{\eta(\kappa)}(x_2, x_3) < 0$. We proceed similarly to the proof of Theorem 4.3 by treating some parameters as variables.

Fix the value of the assembled parameters $K_0, \ldots, K_{n-2}, L_0, \ldots, L_{n-1}, T_0, \ldots, T_{n-2}$ corresponding to κ , and construct $T_{n-1}(\kappa_{6n-3}, K_{n-1})$ as in (6) with κ_{6n-3}, K_{n-1} treated as parameters. We fix as well $x_2, x_3 > 0$, and consider the two parameter family of vectors

$$\xi(\kappa_{6n-3}, K_{n-1}) = (T_0, \dots, T_{n-2}, T_{n-1}(\kappa_{6n-3}, K_{n-1}), K_0, \dots, K_{n-1}, L_0, \dots, L_{n-1}) \in \mathbb{R}^{3n}$$

The polynomial $P_{\xi(\kappa_{6n-3},K_{n-1})}(x_2,x_3)$ becomes a polynomial in κ_{6n-3},K_{n-1} , denoted by $\overline{P}(\kappa_{6n-3},K_{n-1})$. Note that the parameters κ_{6n-3},K_{n-1} are only present in the terms involving a_{n-1},b_{n-1} and c_{n-1} , see (6) and (11). In particular, a_{n-1} depends only on K_{n-1} , while b_{n-1} and c_{n-1} depend on the product $\kappa_{6n-3}K_{n-1}$. Using (14), we express A_2,A_{10} and A_{00} in terms of κ_{6n-3} and K_{n-1} in the following way, where the coefficients $r_i^{(j)} \in \mathbb{R}$ appearing in the expressions are independent of κ_{6n-3} and K_{n-1} :

$$\begin{split} \overline{A}_2 &= \left(1 + \sum_{i=0}^{n-1} c_i \, x_2^{i+1} \,\right) \left(\sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (i-j) \, a_i \, b_j \, x_2^{\ell+1} \right) \\ &= (r_0^{(2)} + r_1^{(2)} \kappa_{6n-3} K_{n-1}) (r_2^{(2)} + r_3^{(2)} K_{n-1} + r_4^{(2)} \kappa_{6n-3} K_{n-1}) \qquad \text{with} \quad r_0^{(2)}, r_1^{(2)}, r_3^{(2)} > 0, \\ \overline{A}_{10} &= \sum_{\ell=0}^{n-1} (\ell+1) \, a_\ell \, x_2^\ell - \sum_{\ell=0}^{n-1} \ell \, b_\ell \, x_2^{\ell+1} + \sum_{\ell=0}^{2n-2} \sum_{i+j=\ell} (j+1-i) \, b_i \, c_j \, x_2^{\ell+2} + \sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (i-j) \, a_i \, c_j \, x_2^{\ell+1} \\ &= (r_0^{(10)} + r_1^{(10)} K_{n-1}) - (r_2^{(10)} + r_3^{(10)} \kappa_{6n-3} K_{n-1}) + (r_4^{(10)} + r_5^{(10)} \kappa_{6n-3} K_{n-1} + r_6^{(10)} \kappa_{6n-3}^2 K_{n-1}) \\ &+ (r_7^{(10)} + r_8^{(10)} K_{n-1} + r_9^{(10)} \kappa_{6n-3} K_{n-1}) \qquad \text{with} \quad r_1^{(10)}, r_8^{(10)} > 0, \\ \overline{A}_{00} &= 1 + \sum_{i=0}^{n-1} c_i \, x_2^{i+1} \, = (r_0^{(2)} + r_1^{(2)} \kappa_{6n-3} K_{n-1}). \end{split}$$

Collecting all the terms together we write \overline{P} as

$$\overline{P}(\kappa_{6n-3},K_{n-1}) = p_0 + p_1K_{n-1} + p_2\kappa_{6n-3}K_{n-1} + p_3\kappa_{6n-3}K_{n-1}^2 + p_4\kappa_{6n-3}^2K_{n-1}^2,$$

with $p_1 = r_0^{(2)} r_3^{(2)} x_3^2 + (r_1^{(10)} + r_8^{(10)}) x_3 > 0$ and $p_3 = r_1^{(2)} r_3^{(2)} x_3^2 > 0$. The Newton polytope of \overline{P} has vertices (0,0),(0,1),(1,2),(2,2). The terms of \overline{P} that can be negative are all supported on the edge F joining the vertices (0,0) and (2,2). The outer normal cone of this edge if generated by $v = (v_1, v_2) = (1,-1)$.

By the choice of κ , x_2 , x_3 , we know that \overline{P} attains negative values, namely by choosing κ_{6n-5} , κ_{6n-4} , κ_{6n-3} to be the entries of the given κ . To avoid confusion, we denote them by κ_{6n-5}^* , κ_{6n-4}^* , κ_{6n-3}^* and the corresponding assembled parameter as K_{n-1}^* . It follows that $\overline{P}_F(\kappa_{6n-3}^*, K_{n-1}^*) < 0$. We evaluate \overline{P} at $q_s = (\kappa_{6n-3}^* s^{v_1}, K_{n-1}^* s^{v_2}) = (\kappa_{6n-3}^* s, K_{n-1}^* s^{-1})$ and obtain

$$Q(s) := s \ \overline{P}(q_s) = p_1 K_{n-1}^* + p_3 \kappa_{6n-3}^* (K_{n-1}^*)^2 + s \ \overline{P}_F(\kappa_{6n-3}^*, K_{n-1}^*).$$

As for s=1, Q(s) is a linear function with negative coefficient and Q(1)<0, we have that Q(s)<0 for all $s \ge 1$. Hence, consider the parameter vector κ_s obtained from κ by replacing κ_{6n-4}^* , κ_{6n-3}^* by $\kappa_{6n-4} s$, $\kappa_{6n-3} s$ such that the new K_{n-1} is $K_{n-1}s^{-1}$. Then κ_s enables multistationarity for all $s \ge 1$. As s increases, so does $\kappa_{6n-3} s$, and eventually κ_s belongs to Y. The parameter vectors κ_s thus give a path connecting the point κ in X with a point in Y. This concludes the proof.

The rest of the section focuses on showing that the region of parameters precluding multistationarity is also path connected. We first show that the region of monostationarity is connected in the 3n dimensional

parameter space given by T_i, K_i, L_i for i = 0, ..., n-1. Written in terms of T_i, K_i, L_i instead of a_i, b_i, c_i , the terms of the polynomial P_{η} in (20) are

$$A_{2}(x_{2}) = \left(1 + \sum_{i=0}^{n-1} T_{i} x_{2}^{i+1}\right) \left(\sum_{\ell=1}^{2n-3} \sum_{1+j=\ell} (i-j) K_{i} L_{j} T_{i-1} T_{j} x_{2}^{\ell+1}\right),$$

$$A_{10}(x_{2}) = \sum_{\ell=0}^{n-1} (\ell+1) K_{\ell} T_{\ell-1} x_{2}^{\ell} - \sum_{\ell=0}^{n-1} \ell L_{\ell} T_{\ell} x_{2}^{\ell+1} + \sum_{\ell=0}^{2n-2} \sum_{i+j=\ell} (j+1-i) L_{i} T_{i} T_{j} x_{2}^{\ell+2}$$

$$+ \sum_{\ell=1}^{2n-3} \sum_{1+j=\ell} (i-j) K_{i} T_{i-1} T_{j} x_{2}^{\ell+1},$$

$$A_{00}(x_{2}) = 1 + \sum_{i=0}^{n-1} T_{i} x_{2}^{i+1}.$$

In Theorem 6.3 we show that the region $Z \subseteq \mathbb{R}^{3n}_{>0}$ of parameters for which P_{η} is positive is path connected. We prove this by showing that any point in Z is path connected to a point $\eta \in Z$ such that all the coefficients of P_{η} are positive. We then show that any two such points are also path connected in Z.

Theorem 6.3. For all $n \ge 1$, the set

$$Z := \{ \eta \in \mathbb{R}^{3n}_{>0} : P_{\eta}(x_2, x_3) > 0 \text{ for all } (x_2, x_3) \in \mathbb{R}^2_{>0} \}$$

is path connected.

Proof. Let $\eta \in Z$. Hence, for all $x_2, x_3 > 0$, $P_{\eta}(x_2, x_3) > 0$. Fix $(x_2, x_3) \in \mathbb{R}^2_{>0}$ and all parameters in η but K_{n-1}, L_0 , and view $P_{\eta}(x_2, x_3)$ as a polynomial in K_{n-1} and L_0 with real coefficients. Let P' denote this polynomial. It is straightforward to see using (16) that P' takes the form

$$P'(K_{n-1}, L_0) := r_3 K_{n-1} L_0 + r_2 K_{n-1} + r_1 L_0 + r_0,$$
 with $r_1, r_2, r_3 > 0$.

As in the proof of Theorem 6.2, we denote by K_{n-1}^*, L_0^* the corresponding parameter entries in η , such that $P'(K_{n-1}^*, L_0^*) > 0$.

We evaluate P' at $q_s = (K_{n-1}^* s, L_0^* s)$:

$$Q(s) := P'(q_s) = r_3 K_{n-1} L_0 s^2 + (r_2 K_{n-1} + r_1 L_0) s + r_0.$$

The polynomial Q is quadratic in s with positive leading term and Q(1) > 0. By Descartes' rule of signs, Q has at most one positive root and hence Q(s) > 0 for all $s \ge 1$.

As this holds for any fixed $(x_2, x_3) \in \mathbb{R}^2_{>0}$, it follows that

$$\eta_s := (T_0, \dots, T_{n-1}, K_0, \dots, K_{n-2}, K_{n-1}^* s, L_0^* s, L_1, \dots, L_{n-1}) \in \mathbb{Z}, \quad \text{ for all } s \geqslant 1.$$

We now show that for s large enough, all coefficients of P_{η_s} in x_2, x_3 are positive. The factor of $A_2(x_2)$ that can be negative is $\left(\sum_{\ell=1}^{2n-3}\sum_{i+j=\ell}(i-j)\,K_iL_jT_{i-1}\,T_j\,x_2^{\ell+1}\right)$. For the coefficient of x_2^k , seen as a polynomial in all parameters, the coefficients of all the monomials involving K_{n-1} and L_0 are positive, and there is at least one such monomial for every $2\leqslant k\leqslant 2n-2$. Hence, by letting K_{n-1} and L_0 be sufficiently large, all coefficients become positive. An analogous argument holds for $A_{10}(x_2)$ and therefore we conclude that, for s large enough, all coefficients of P_{η_s} in x_2, x_3 are positive.

All that is left is to show that any two parameter points η_1 , η_2 such that all coefficients of P_{η_i} are positive, are connected by a path in Z. First note that increasing the coordinates of K_{n-1} and L_0 in any of them leads to a parameter point also in Z with all coefficients of P_{η_i} positive. Second, consider the projection π outside the entries K_{n-1} and L_0 , and form the line segment $\eta'(s) := s\pi(\eta_1) + (1-s)\pi(\eta_2)$ for $s \in [0,1]$. Extend $\eta'(s)$ to a parameter vector $\eta(s,K_{n-1},L_0)$ with K_{n-1} and L_0 seen as variables. The coefficients of $P_{\eta(s,K_{n-1},L_0)}$ are polynomials in K_{n-1} , L_0 with coefficients in s, such that the coefficients of the monomials involving K_{n-1} , L_0 are positive continuous functions of s. As [0,1] is compact, there exist K_{n-1}^* and L_0^* such that for any $K_{n-1} \geq K_{n-1}^*$ and $L_0 \geq L_0^*$, all the coefficients of $P_{\eta(s,K_{n-1},L_0)}$ are positive for all $s \in [0,1]$. By picking K_{n-1}^* and L_0^* to be larger than the entries of η_1 and η_2 , the desired path is the composition of the line segments in Z joining the points η_1 , $\eta(0,K_{n-1}^*,L_0^*)$, $\eta(1,K_{n-1}^*,L_0^*)$ and η_2 in this order. This concludes the proof.

In Theorem 6.3 above we have shown that the region of monostationarity is path connected in the parameter space given by the parameters K_i , L_i , and T_i for all i = 0, ..., n - 1. Next, we extend this result to the 6n dimensional parameter space given by $\kappa \in \mathbb{R}^{6n}_{>0}$.

Corollary 6.4. The set of parameters $\kappa \in \mathbb{R}^{6n}_{>0}$ precluding multistationarity is path connected for all n.

Proof. Let $\kappa, \kappa' \in \mathbb{R}^{6n}_{>0}$ preclude multistationarity. We have that $\eta(\kappa), \eta(\kappa') \in Z$, with Z as in Theorem 6.3 and hence, there is a continuous path

$$\xi(s) = (T_0(s), \dots, T_{n-1}(s), K_0(s), \dots, K_{n-1}(s), L_0(s), \dots, L_{n-1}(s)), \quad s \in [0, 1]$$

connecting them in Z. Define the points $\kappa(s) \in \mathbb{R}^{6n}_{>0}$ for $i = 0, \dots, n-1$, as

$$\kappa_{6i+2}(s) := \kappa_{6i+2}, \qquad \kappa_{6i+3}(s) := \kappa_{6i+6} \frac{T_i(s)}{T_{i-1}(s)} \frac{L_i(s)}{K_i(s)}, \qquad \kappa_{6i+1}(s) := K_i(s) (\kappa_{6i+2} + \kappa_{6i+3}(s)),$$

$$\kappa_{6i+5}(s) := \kappa_{6i+5}, \qquad \kappa_{6i+6}(s) := \kappa_{6i+6}, \qquad \kappa_{6i+4}(s) := L_i(s) (\kappa_{6i+5} + \kappa_{6i+6}).$$

Using (6), $\frac{T_i(0)}{T_{i-1}(0)} \frac{L_i(0)}{K_i(0)} = \frac{\kappa_{6i+3}}{\kappa_{6i+6}}$ and hence $\kappa(0) = \kappa$. Furthermore, for all $s \in [0,1]$, $\eta(\kappa(s)) = \xi(s)$. Therefore, $\kappa(s)$ precludes multistationarity for all $s \in [0,1]$ and gives a path from κ to $\kappa(1)$. All that is left is to connect κ' with $\kappa(1)$. Note that for $i = 0, \ldots, n-1$,

$$(\kappa_{6i+1}(1), \ \kappa_{6i+2}(1), \ \kappa_{6i+3}(1), \ \kappa_{6i+4}(1), \ \kappa_{6i+5}(1), \ \kappa_{6i+6}(1)) =$$

$$\left(\frac{\kappa'_{6i+1}}{\kappa'_{6i+2} + \kappa'_{6i+3}} \left(\kappa_{6i+2} + \kappa_{6i+6} \frac{\kappa'_{6i+3}}{\kappa'_{6i+6}} \right), \kappa_{6i+2}, \kappa_{6i+6} \frac{\kappa'_{6i+3}}{\kappa'_{6i+6}}, \frac{\kappa'_{6i+4}}{\kappa'_{6i+5} + \kappa'_{6i+6}}, \kappa_{6i+5}, \kappa_{6i+6} \right)$$

To this end, define a path $\alpha(s)$ with, for $i = 0, \dots, n-1$,

$$\alpha_{6i+2}(s) := \kappa_{6i+2} \left(\frac{\kappa'_{6i+2}}{\kappa_{6i+2}}\right)^{s}, \quad \alpha_{6i+3}(s) := \kappa'_{6i+3} \left(\frac{\kappa_{6i+6}}{\kappa'_{6i+6}}\right)^{1-s}, \quad \alpha_{6i+1}(s) := \frac{\kappa'_{6i+1}}{\kappa'_{6i+2} + \kappa'_{6i+3}} \left(\alpha_{6i+2}(s) + \alpha_{6i+3}(s)\right)$$

$$\alpha_{6i+5}(s) := \kappa_{6i+5} \left(\frac{\kappa'_{6i+5}}{\kappa_{6i+5}}\right)^{s}, \quad \alpha_{6i+6}(s) := \kappa'_{6i+6} \left(\frac{\kappa_{6i+6}}{\kappa'_{6i+6}}\right)^{1-s}, \quad \alpha_{6i+4}(s) := \frac{\kappa'_{6i+4}}{\kappa'_{6i+5} + \kappa'_{6i+6}} \left(\alpha_{6i+5}(s) + \alpha_{6i+6}(s)\right).$$

We have $\alpha(0) = \xi(1)$ and $\alpha(1) = \kappa'$. Furthermore, $\eta(\alpha(s)) = \eta(\kappa')$ for all $s \in [0,1]$ and hence, $\alpha(s)$ precludes multistationarity. Thus, α gives the desired path and concludes the proof.

APPENDIX A.

In this appendix we compute the determinant of the matrix J in (12), which we rewrite here for convenience:

$$J := \begin{bmatrix} 1 + \sum_{i=0}^{n-1} (i+1) \, a_i \, x_2^i x_3 & - \sum_{i=0}^{n-1} i \, a_i \, x_2^{i+1} x_3 & \sum_{i=0}^{n-1} a_i \, x_1 x_2^i \\ \sum_{i=0}^{n-1} (i+1) \, b_i \, x_2^i x_3 & 1 - \sum_{i=0}^{n-1} i \, b_i \, x_2^{i+1} x_3 & \sum_{i=0}^{n-1} b_i \, x_1 x_2^i \\ -1 + \sum_{i=0}^{n-1} (i+1) \, c_i \, x_1^{-1} \, x_2^{i+1} x_3 & -1 - \sum_{i=0}^{n-1} (i+1) \, c_i \, x_1^{-1} x_2^{i+2} x_3 & 1 + \sum_{i=0}^{n-1} c_i \, x_2^{i+1} \end{bmatrix}.$$

Note that x_3 appears with degree one in each of the entries of columns 1 and 2 but the third column does not depend on x_3 . Therefore, the determinant of J can be written as a quadratic polynomial in x_3 :

$$P := \det J = A_2 x_3^2 + A_1 x_3 + A_0.$$

Specifically, after noticing that the first two columns can be written as a sum of two vectors, one independent of x_3 and the other a multiple of x_3 , we write the determinant of J as

$$P = \det J_1 + (\det J_2 + \det J_3)x_3 + (\det J_4)x_3^2,$$

where

$$J_{1} = \begin{bmatrix} 1 & 0 & \sum_{i=0}^{n-1} a_{i} x_{1} x_{2}^{i} \\ 0 & 1 & \sum_{i=0}^{n-1} b_{i} x_{1} x_{2}^{i} \\ -1 & -1 & 1 + \sum_{i=0}^{n-1} c_{i} x_{2}^{i+1} \end{bmatrix},$$

$$J_{2} = \begin{bmatrix} 1 & -\sum_{i=0}^{n-1} i a_{i} x_{2}^{i+1} & \sum_{i=0}^{n-1} a_{i} x_{1} x_{2}^{i} \\ 0 & -\sum_{i=0}^{n-1} i b_{i} x_{2}^{i+1} & \sum_{i=0}^{n-1} b_{i} x_{1} x_{2}^{i} \\ -1 & -\sum_{i=0}^{n-1} (i+1) c_{i} x_{1}^{-1} x_{2}^{i+2} & 1 + \sum_{i=0}^{n-1} c_{i} x_{2}^{i+1} \end{bmatrix},$$

$$J_{3} = \begin{bmatrix} \sum_{i=0}^{n-1} (i+1) \, a_{i} \, x_{2}^{i} & 0 & \sum_{i=0}^{n-1} \, a_{i} \, x_{1} x_{2}^{i} \\ \sum_{i=0}^{n-1} (i+1) \, b_{i} \, x_{2}^{i} & 1 & \sum_{i=0}^{n-1} \, b_{i} \, x_{1} x_{2}^{i} \\ \sum_{i=0}^{n-1} (i+1) \, c_{i} \, x_{1}^{-1} \, x_{2}^{i+1} & -1 & 1 + \sum_{i=0}^{n-1} \, c_{i} \, x_{2}^{i+1} \end{bmatrix},$$

$$J_{4} = \begin{bmatrix} \sum_{i=0}^{n-1} (i+1) \, a_{i} \, x_{2}^{i} & -\sum_{i=0}^{n-1} i \, a_{i} \, x_{2}^{i+1} & \sum_{i=0}^{n-1} a_{i} \, x_{1} x_{2}^{i} \\ \sum_{i=0}^{n-1} (i+1) \, b_{i} \, x_{2}^{i} & -\sum_{i=0}^{n-1} i \, b_{i} \, x_{2}^{i+1} & \sum_{i=0}^{n-1} b_{i} \, x_{1} x_{2}^{i} \\ \sum_{i=0}^{n-1} (i+1) \, c_{i} \, x_{1}^{-1} \, x_{2}^{i+1} & -\sum_{i=0}^{n-1} (i+1) \, c_{i} \, x_{1}^{-1} x_{2}^{i+2} & 1 + \sum_{i=0}^{n-1} c_{i} \, x_{2}^{i+1} \end{bmatrix}.$$

We proceed to compute each coefficient A_0, A_1, A_2 using the matrices J_i , i = 1, 2, 3, 4.

Computing A_0 . The determinant of J_1 is simply

$$A_0 = 1 + \sum_{i=0}^{n-1} c_i x_2^{i+1} + \sum_{i=0}^{n-1} b_i x_1 x_2^i + \sum_{i=0}^{n-1} a_i x_1 x_2^i.$$

Computing A_2 . We compute now the determinant of J_4 . After adding the first column multiplied by x_2 to the second column, det J_4 agrees with the determinant of

$$J_4' = \begin{bmatrix} \sum_{i=0}^{n-1} (i+1) a_i x_2^i & \sum_{i=0}^{n-1} a_i x_2^{i+1} & \sum_{i=0}^{n-1} a_i x_1 x_2^i \\ \sum_{i=0}^{n-1} (i+1) b_i x_2^i & \sum_{i=0}^{n-1} b_i x_2^{i+1} & \sum_{i=0}^{n-1} b_i x_1 x_2^i \\ \sum_{i=0}^{n-1} (i+1) c_i x_1^{-1} x_2^{i+1} & 0 & 1 + \sum_{i=0}^{n-1} c_i x_2^{i+1} \end{bmatrix}.$$

Subtraction of the second column of J'_4 multiplied by x_1/x_2 from the third column gives that det $J_4 = \det J'_4 = \det J''_4$ with

$$J_4'' = \begin{bmatrix} \sum_{i=0}^{n-1} (i+1) a_i x_2^i & \sum_{i=0}^{n-1} a_i x_2^{i+1} & 0\\ \sum_{i=0}^{n-1} (i+1) b_i x_2^i & \sum_{i=0}^{n-1} b_i x_2^{i+1} & 0\\ \sum_{i=0}^{n-1} (i+1) c_i x_1^{i-1} x_2^{i+1} & 0 & 1 + \sum_{i=0}^{n-1} c_i x_2^{i+1} \end{bmatrix}.$$

Hence, we compute the determinant of J_4'' . We start by finding the principal minor obtained by removing the last row and column of J_4'' :

$$M = \left(\sum_{i=0}^{n-1} (i+1) a_i x_2^i\right) \left(\sum_{i=0}^{n-1} b_i x_2^{i+1}\right) - \left(\sum_{i=0}^{n-1} a_i x_2^{i+1}\right) \left(\sum_{i=0}^{n-1} (i+1) b_i x_2^i\right)$$

$$= \left(\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (i+1) a_i b_j x_2^{i+j+1}\right) - \left(\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} a_i (j+1) b_j x_2^{i+j+1}\right)$$

$$= \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (i-j) a_i b_j x_2^{i+j+1} = \sum_{\ell=1}^{2n-2} \sum_{i+j=\ell} (i-j) a_i b_j x_2^{\ell+1} = \sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (i-j) a_i b_j x_2^{\ell+1},$$

where in the last equality we note that the term vanished for $\ell = 2n - 2$. From this, we obtain

$$A_2 = \det J_4 = \left(1 + \sum_{i=0}^{n-1} c_i \, x_2^{i+1} \right) \left(\sum_{\ell=1}^{2n-3} \sum_{i+i=\ell} (i-j) \, a_i \, b_j \, x_2^{\ell+1} \right).$$

Computing A_1 . We start by finding det J_2 .

$$\det J_2 = \det \begin{bmatrix} 1 & -\sum_{i=0}^{n-1} i \, a_i \, x_2^{i+1} & \sum_{i=0}^{n-1} a_i \, x_1 x_2^i \\ 0 & -\sum_{i=0}^{n-1} i \, b_i \, x_2^{i+1} & \sum_{i=0}^{n-1} b_i \, x_1 x_2^i \\ -1 & -\sum_{i=0}^{n-1} (i+1) \, c_i \, x_1^{-1} x_2^{i+2} & \sum_{i=0}^{n-1} c_i \, x_2^{i+1} \end{bmatrix} + \det \begin{bmatrix} 1 & -\sum_{i=0}^{n-1} i \, a_i \, x_2^{i+1} & 0 \\ 0 & -\sum_{i=0}^{n-1} i \, b_i \, x_2^{i+1} & 0 \\ -1 & -\sum_{i=0}^{n-1} (i+1) \, c_i \, x_1^{-1} x_2^{i+2} & 1 \end{bmatrix}.$$

The second determinant is simply

$$D_2 := -\sum_{i=0}^{n-1} i \, b_i \, x_2^{i+1}.$$

For the first determinant D_1 , adding the first row to the last gives

$$D_1 = \det \begin{bmatrix} 1 & -\sum_{i=0}^{n-1} i \, a_i \, x_2^{i+1} & \sum_{i=0}^{n-1} a_i \, x_1 x_2^i \\ 0 & -\sum_{i=0}^{n-1} i \, b_i \, x_2^{i+1} & \sum_{i=0}^{n-1} b_i \, x_1 x_2^i \\ 0 & -\sum_{i=0}^{n-1} (i \, a_i \, x_2^{i+1} + (i+1) \, c_i \, x_1^{-1} x_2^{i+2}) & \sum_{i=0}^{n-1} (a_i \, x_1 x_2^i + c_i \, x_2^{i+1}) \end{bmatrix},$$

and hence

$$D_{1} = -\left(\sum_{i=0}^{n-1} i \, b_{i} \, x_{2}^{i+1}\right) \left(\sum_{i=0}^{n-1} (a_{i} \, x_{1} x_{2}^{i} + c_{i} \, x_{2}^{i+1})\right) + \left(\sum_{i=0}^{n-1} b_{i} \, x_{1} x_{2}^{i}\right) \left(\sum_{i=0}^{n-1} (i \, a_{i} \, x_{2}^{i+1} + (i+1) \, c_{i} \, x_{1}^{-1} x_{2}^{i+2})\right)$$

$$= \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (j-i) \, a_{j} \, b_{i} \, x_{1} x_{2}^{i+j+1} + \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (j+1-i) \, b_{i} \, c_{j} \, x_{2}^{i+j+2}$$

$$= \sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (j-i) \, a_{j} \, b_{i} \, x_{1} x_{2}^{\ell+1} + \sum_{\ell=0}^{2n-2} \sum_{i+j=\ell} (j+1-i) \, b_{i} \, c_{j} \, x_{2}^{\ell+2}.$$

This gives $\det J_2 = D_1 + D_2$.

We proceed similarly with the computation of $\det J_3$:

$$\det J_3 = \det \begin{bmatrix} \sum_{i=0}^{n-1} (i+1) \, a_i \, x_2^i & 0 & \sum_{i=0}^{n-1} a_i \, x_1 x_2^i \\ \sum_{i=0}^{n-1} (i+1) \, b_i \, x_2^i & 1 & \sum_{i=0}^{n-1} b_i \, x_1 x_2^i \\ \sum_{i=0}^{n-1} (i+1) \, c_i \, x_1^{-1} \, x_2^{i+1} & -1 & \sum_{i=0}^{n-1} c_i \, x_2^{i+1} \end{bmatrix}$$

$$+ \det \begin{bmatrix} \sum_{i=0}^{n-1} (i+1) \, a_i \, x_2^i & 0 & 0 \\ \sum_{i=0}^{n-1} (i+1) \, b_i \, x_2^i & 1 & 0 \\ \sum_{i=0}^{n-1} (i+1) \, c_i \, x_1^{-1} \, x_2^{i+1} & -1 & 1 \end{bmatrix}.$$

The second determinant C_2 is

$$C_2 = \sum_{i=0}^{n-1} (i+1) a_i x_2^i.$$

For the first determinant C_1 , we have

$$C_{1} = \det \begin{bmatrix} \sum_{i=0}^{n-1} (i+1) a_{i} x_{2}^{i} & 0 & \sum_{i=0}^{n-1} a_{i} x_{1} x_{2}^{i} \\ \sum_{i=0}^{n-1} (i+1) b_{i} x_{2}^{i} & 1 & \sum_{i=0}^{n-1} b_{i} x_{1} x_{2}^{i} \\ \sum_{i=0}^{n-1} (i+1) (b_{i} x_{2}^{i} + c_{i} x_{1}^{-1} x_{2}^{i+1}) & 0 & \sum_{i=0}^{n-1} (b_{i} x_{1} x_{2}^{i} + c_{i} x_{2}^{i+1}) \end{bmatrix},$$

and hence

$$C_{1} = \left(\sum_{i=0}^{n-1} (i+1) a_{i} x_{2}^{i}\right) \left(\sum_{i=0}^{n-1} (b_{i} x_{1} x_{2}^{i} + c_{i} x_{2}^{i+1})\right) - \left(\sum_{i=0}^{n-1} a_{i} x_{1} x_{2}^{i}\right) \left(\sum_{i=0}^{n-1} (i+1) (b_{i} x_{2}^{i} + c_{i} x_{1}^{-1} x_{2}^{i+1})\right)$$

$$= \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (i-j) a_{i} b_{j} x_{1} x_{2}^{i+j} + \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (i-j) a_{i} c_{j} x_{2}^{i+j+1}$$

$$= \sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (i-j) a_{i} b_{j} x_{1} x_{2}^{\ell} + \sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (i-j) a_{i} c_{j} x_{2}^{\ell+1}.$$

Putting it all together, we get

$$\begin{split} A_1 &= C_2 + D_2 + C_1 + D_1 \\ &= \sum_{\ell=0}^{n-1} (\ell+1) \, a_\ell \, x_2^\ell - \sum_{\ell=0}^{n-1} \ell \, b_\ell \, x_2^{\ell+1} + \sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (j-i) \, a_j \, b_i \, x_1 x_2^{\ell+1} \\ &+ \sum_{\ell=0}^{2n-2} \sum_{i+j=\ell} (j+1-i) \, b_i \, c_j \, x_2^{\ell+2} + \sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (i-j) a_i \, b_j \, x_1 x_2^\ell + \sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (i-j) \, a_i \, c_j \, x_2^{\ell+1} \\ &= \sum_{\ell=0}^{n-1} (\ell+1) \, a_\ell \, x_2^\ell - \sum_{\ell=0}^{n-1} \ell \, b_\ell \, x_2^{\ell+1} + \sum_{\ell=0}^{2n-2} \sum_{i+j=\ell} (j+1-i) \, b_i \, c_j \, x_2^{\ell+2} + \sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (i-j) \, a_i \, c_j \, x_2^{\ell+1} \\ &+ x_1 (1+x_2) \left(\sum_{\ell=1}^{2n-3} \sum_{i+j=\ell} (j-i) \, a_j \, b_i \, x_2^\ell \right). \end{split}$$

Appendix B.

In this part of the appendix, we provide a supplementary lemma to prove Theorem 5.3.

Lemma B.1. Let $\kappa(k)$ as given in (28). If $\ell \in \{2, \ldots, 2n-2\}$, $\ell \neq n$, then the coefficient $\mathcal{C}_{\ell}(k)$ of x_2^{ℓ} in A_{10} is positive for any $k \in \mathbb{R}_{>0}$.

Proof. First, we consider the case $\ell \in \{n+1,\ldots,2n-2\}$, and note that in this case

$$\begin{split} \mathcal{C}_{\ell}(k) &= \sum_{i+j=\ell-2} \left(j+1-i\right) b_i \, c_j \ + \sum_{i+j=\ell-1} \left(i-j\right) a_i \, c_j \\ &= \sum_{i+j=\ell-2} T_i \, T_j \ + \sum_{\substack{i+j=\ell-1\\i>j}} \left(i-j\right) \left(\, a_i \, c_j - \, a_j \, c_i \right), \end{split}$$

as $b_i = c_i = T_i$ for any $i \in \{0, ..., n-1\}$ for $\kappa(k)$, and $\sum_{i+j=\ell-2} (j+1-i)T_i T_j = \sum_{i+j=\ell-2} T_i T_j$. The first summand is positive since each T_i is positive. Likewise, the last summand is also positive, because for any $i, j \in \{0, ..., n-1\}$ such that i > j, we find using (29):

(30)
$$a_i c_j - a_j c_i = K_i T_{i-1} T_j - K_j T_{j-1} T_i = \begin{cases} \frac{i-j}{(n+1)^2} & \text{if } i \in \{1 \dots, n-2\}, \\ \frac{k}{(n+1)^2} (n-j-1) & \text{if } i = n-1. \end{cases}$$

Therefore $C_{\ell}(k) > 0$ for $\ell \in \{n+1, \ldots, 2n-2\}$.

The coefficient $C_{\ell}(k)$ for $\ell \in \{2, \ldots, n-1\}$ is given by

$$C_{\ell}(k) = (\ell+1) a_{\ell} - (\ell-1) b_{\ell-1} + \sum_{i+j=\ell-2} T_i T_j + \sum_{\substack{i+j=\ell-1 \ i>j}} (i-j) (a_i c_j - a_j c_i)$$

$$= \left(-(\ell-1) b_{\ell-1} + \sum_{\substack{i+j=\ell-2 \ i\neq j=\ell-1}} T_i T_j \right) + (\ell+1) a_{\ell} + \sum_{\substack{i+j=\ell-1 \ i\neq j=\ell-1}} (i-j) (a_i c_j - a_j c_i),$$

where the second equality is simply a reordering of terms. The second and third terms are positive by (30). We will show that the first term is also positive.

We point out that T_{n-1} does not appear in any sum. For $\ell = 2$, the first term is $-(\ell - 1) T_{\ell-1} + T_0^2 = 0$ $T_0^2 > 0$ and hence, $C_2(k) > 0$. For $\ell > 2$, (29) gives $T_i T_j = \frac{(n-i)(n-j)}{(n+1)^2}$, and we obtain

$$(n+1)^2 \sum_{i+j=\ell-2} T_i T_j = \sum_{i+j=\ell-2} (n-i)(n-j) = \sum_{i=0}^{\ell-2} (n-i)(n-\ell+2+i)$$
$$= \sum_{i=0}^{\ell-2} \left((n+1) - (i+1) \right) \left((n-\ell+1) + (i+1) \right)$$
$$> (\ell-1)(n-\ell+1)(n+1) = (n+1)^2 (\ell-1)b_{\ell-1}.$$

This gives $C_{\ell}(k) > 0$ and concludes the proof.

Acknowledgements. We thank the anonymous referees for their helpful comments. EF acknowledges funding from the Independent Research Fund of Denmark. TdW is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Emmy Noether Programme, grant WO 2206/1-1. OY is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - The Berlin Mathematics Research Center MATH+ (EXC-2046/1, project ID 390685689, sub-project AA1-9).

References

- [1] F. Bihan, A. Dickenstein, and Giaroli M. Lower bounds for positive roots and regions of multistationarity in chemical reaction networks. J. Algebra, 542:367-411, 2020.
- [2] F. Bihan, F. Santos, and P.-J. Spaenlehauer. A polyhedral method for sparse systems with many positive solutions. SIAM J. Appl. Alg. Geom., 2(4):620-645, 2018.

- [3] G. Blekherman, P.A. Parrilo, and R.R. Thomas. Semidefinite Optimization and Convex Algebraic Geometry, volume 13 of MOS-SIAM Series on Optimization. SIAM and the Mathematical Optimization Society, Philadelphia, 2013.
- [4] V. Chandrasekaran and P. Shah. Relative entropy relaxations for signomial optimization. SIAM Journal on Optimization, 26(2):1147–1173, 2016.
- [5] V. Chandrasekaran and P. Shah. Relative entropy optimization and its applications. *Math. Program.*, 161(1-2, Ser. A):1–32, 2017.
- [6] P. Cohen. The structure and regulation of protein phosphatases. *Annu. Rev. Biochem.*, 58:453–508, Jan 1989.
- [7] C. Conradi, E. Feliu, M. Mincheva, and C. Wiuf. Identifying parameter regions for multistationarity. *PLoS Comput. Biol.*, 13(10):e1005751, 2017.
- [8] C. Conradi, D. Flockerzi, J. Raisch, and J. Stelling. Subnetwork analysis reveals dynamic features of complex (bio)chemical networks. *Proc. Nat. Acad. Sci.*, 104(49):19175–80, 2007.
- [9] C. Conradi, A. Iosif, and T. Kahle. Multistationarity in the space of total concentrations for systems that admit a monomial parametrization. *Bull. Math. Biol.*, 81(10):4174–4209, 2019.
- [10] C. Conradi and M. Mincheva. Catalytic constants enable the emergence of bistability in dual phosphorylation. J. R. S. Interface, 11(95), 2014.
- [11] G. Craciun, J. W. Helton, and R. J. Williams. Homotopy methods for counting reaction network equilibria. *Mathematical biosciences*, 216(2):140–149, 2008.
- [12] P. Donnell, M. Banaji, A. Marginean, and C. Pantea. Control: an open source framework for the analysis of chemical reaction networks. *Bioinformatics*, 30(11), 2014.
- [13] M. Dressler, S. Iliman, and T. de Wolff. A Positivstellensatz for Sums of Nonnegative Circuit Polynomials. SIAM J. Appl. Algebra Geom., 1(1):536–555, 2017.
- [14] P. Ellison, M. Feinberg, H. Ji, and D. Knight. Chemical reaction network toolbox, version 2.2. Available online at http://www.crnt.osu.edu/CRNTWin, 2012.
- [15] M. Feinberg. The existence and uniqueness of steady states for a class of chemical reaction networks. *Arch. Rational Mech. Anal.*, 132(4):311–370, 1995.
- [16] E. Feliu. Injectivity, multiple zeros, and multistationarity in reaction networks. *Proceedings of the Royal Society A*, doi:10.1098/rspa.2014.0530, 2014.
- [17] E. Feliu, N. Kaihnsa, T. de Wolff, and O. Yürük. The Kinetic Space of Multistationarity in Dual Phosphorylation. *J. Dyn. Diff. Equat.*, 34:825–852, 2022.
- [18] E. Feliu, A. D. Rendall, and C. Wiuf. A proof of unlimited multistability for phosphorylation cycles. Nonlinearity, 33(11):5629–5658, 2020.
- [19] E. Feliu and C. Wiuf. Variable elimination in chemical reaction networks with mass-action kinetics. SIAM J. Appl. Math., 72:959–981, 2012.
- [20] C. Fidalgo and A. Kovacec. Positive semidefinite diagonal minus tail forms are sums of squares. Math. Z., 269(3-4):629–645, 2011.
- [21] D. Flockerzi, K. Holstein, and C. Conradi. N-site Phosphorylation Systems with 2N-1 Steady States. Bull. Math. Biol., 76(8):1892–1916, 2014.
- [22] J. Forsgård and T. de Wolff. The algebraic boundary of the sonc cone, 2019. Preprint; see arXiv:1905.04776.
- [23] M. Giaroli, R. Rischter, M. P. Millán, and A. Dickenstein. Parameter regions that give rise to 2[n/2] +1 positive steady states in the n-site phosphorylation system. *Math Biosci Eng*, 16(6):7589–7615, 08 2019.
- [24] F. Gnad, S. Ren, J. Cox, J. V. Olsen, B. Macek, M. Oroshi, and M. Mann. PHOSIDA (phosphorylation site database): management, structural and evolutionary investigation, and prediction of phosphosites. *Genome Biol*, 8(11):R250, 2007.
- [25] D. Hilbert. Ueber die Darstellung definiter Formen als Summe von Formenquadraten. *Math. Ann.*, 32(3):342–350, 1888.
- [26] S. Iliman and T. de Wolff. Amoebas, nonnegative polynomials and sums of squares supported on circuits. *Res. Math. Sci.*, 3(9), 2016.
- [27] S. Iliman and T. de Wolff. Lower bounds for polynomials with simplex newton polytopes based on geometric programming. SIAM J. Optim., 26(2):1128–1146, 2016.

- [28] M. Laurent and N. Kellershohn. Multistability: a major means of differentiation and evolution in biological systems. Trends Biochem. Sciences, 24(11):418–422, 1999.
- [29] T.S. Motzkin. The arithmetic-geometric inequality. In *Inequalities: Proceedings, Volume 1*, chapter 10, pages 203–224. Academic Press, 1967.
- [30] R. Murray, V. Chandrasekaran, and A. Wiermann. Newton polytopes and relative entropy optimization. Foundations of Computational Mathematics, 21:1703–1737, 2021.
- [31] R. Murray, V. Chandrasekaran, and A. Wiermann. Signomial and polynomial optimization via relative entropy and partial dualization. *Math. Program. Comput.*, 13(2):257–295, 2021.
- [32] K. M. Nam, B. M. Gyori, S. V. Amethyst, D. J. Bates, and J. Gunawardena. Robustness and parameter geography in post-translational modification systems. *PLoS Comput. Biol.*, 16(5):e1007573, 05 2020.
- [33] E. M. Ozbudak, M. Thattai, H. N. Lim, B. I. Shraiman, and A. Van Oudenaarden. Multistability in the lactose utilization network of escherichia coli. *Nature*, 427(6976):737–740, 2004.
- [34] C. Pantea, H. Koeppl, and G. Craciun. Global injectivity and multiple equilibria in uni- and bimolecular reaction networks. *Discrete Contin. Dyn. Syst. Ser. B*, 17(6):2153–2170, 2012.
- [35] M. Pérez Millán and A. Dickenstein. The structure of MESSI biological systems. SIAM J. Appl. Dyn. Syst., 17:1650–1682, 2018.
- [36] M. Pérez Millán, A. Dickenstein, A. Shiu, and C. Conradi. Chemical reaction systems with toric steady states. *Bull. Math. Biol.*, 74:1027–1065, 2012.
- [37] B. Reznick. Forms derived from the arithmetic-geometric inequality. *Math. Ann.*, 283(3):431–464, 1989.
- [38] M. Sáez, E. Feliu, and C. Wiuf. Graphical criteria for positive solutions to linear systems. *Linear Algebra and its Applications*, 552:166 193, 2018.
- [39] C. Salazar and T. Hofer. Multisite protein phosphorylation–from molecular mechanisms to kinetic models. FEBS J., 276(12):3177–3198, Jun 2009.
- [40] M. L. Telek and E Feliu. Topological descriptors of the parameter region of multistationarity: deciding upon connectivity. arXiv, 2209.13936, 2022.
- [41] M. Thomson and J. Gunawardena. The rational parameterization theorem for multisite post-translational modification systems. *J. Theor. Biol.*, 261:626–636, 2009.
- [42] M. Thomson and J. Gunawardena. Unlimited multistability in multisite phosphorylation systems. Nature, 460(7252):274 – 277, 2009.
- [43] J. Wang. Nonnegative polynomials and circuit polynomials. SIAM J. Appl. Algebra Geom., 6(2):111–133, 2022.
- [44] L. Wang and E. D. Sontag. On the number of steady states in a multiple futile cycle. *J. Math. Biol.*, 57(1):29–52, 2008.
- [45] C. Wiuf and E. Feliu. Power-law kinetics and determinant criteria for the preclusion of multistationarity in networks of interacting species. SIAM J. Appl. Dyn. Syst., 12:1685–1721, 2013.
- [46] W. Xiong and J. E. Ferrell Jr. A positive-feedback-based bistable 'memory module' that governs a cell fate decision. *Nature*, 426(6965):460–465, 2003.
- [47] O. Yürük. On the Maximal Mediated Set Structure and the Applications of Nonnegative Circuit Polynomials. PhD thesis, TU Braunschweig, 2021.